STRUCTURE THEOREM FOR LOG DE RHAM-WITT SHEAVES WITH VANISHING

FEI REN

Abstract. We prove an elegant structure theorem for log de Rham-Witt sheaves with vanishing along an effective Cartier divisor D defined in [\[RR24\]](#page-27-0), answering a question of Shuji Saito during the Mainz conference and a question of Yigeng Zhao during a short visit of the author last summer. Our structural result for the log forms also lays the foundation for the study of Milnor K -theory with vanishing along D in the paper to come.

CONTENTS

1. INTRODUCTION

Let k be a perfect field of positive characteristic p. Let X be a smooth k-scheme (not necessarily proper) of pure dimension d and let D be an effective Cartier divisor such that the underlying reduced divisor D_{red} has simple normal crossings. Let U be the complement of D in X and $j: U \hookrightarrow X$ be the open immersion. In [\[RR24\]](#page-27-0), we proved a duality theorem for de Rham-Witt sheaves with certain pole- or zero-restrictions along D in the Nisnevich topology. The de Rham-Witt sheaf with zeros along D, defined by

$$
W_n\Omega_{(X,-D)}^q:=\mathrm{Ker}\left(W_n\Omega_X^q\rightarrow \bigoplus_i W_n\Omega_{D_i}^q\right)\quad\text{(where $D=\sum_i D_i$ with all $D_{i,\mathrm{red}}$ smooth)},
$$

is dual to the the corresponding sheaf with poles defined via the theory of sheaves with modulus [\[KMSY21a,](#page-27-2) [KMSY21b,](#page-27-3) [KMSY22\]](#page-27-4). This duality extends further to a duality theorem of Milne-Kato type. Namely, the \mathbb{Z}/p^n -coefficient motivic complex with zeros along D, defined by

$$
\mathbb{Z}/p^n(q)_{(X,-D)}:=\left(W_n\Omega_{(X,-D)}^q\xrightarrow{C^{-1}-1}\frac{W_n\Omega_{(X,-D)}^q}{dV^{n-1}\Omega_X^{q-1}\cap W_n\Omega_{(X,-D)}^q}\right)[-q],
$$

is dual to the corresponding motivic complex with poles in the étale topology. See $[RR24, The$ orem 9.3, Theorem 11.15] for the precise statements. Moreover, the complex $\mathbb{Z}/p^{n}(q)_{(X,-D)}$

is concentrated in degree q in the étale topology by $[RR24, Lemma 11.6]$. This only nonzero cohomology sheaf is denoted by $W_n \Omega_{(X,-D),\log}^q$, and it is the intersection of the usual log de Rham-Witt sheaf $W_n \Omega_{X,\log}^q$ with $W_n \Omega_{(0)}^q$ $(X, -D)$. We will refer to it as the *log de Rham-Witt* sheaf with vanishing along D or the log de Rham-Witt sheaf with zeros along D . Hence to study the \mathbb{Z}/p^n -motivic complex with zeros along D, it is essential to understand the structure of $\hat{W}_n \Omega^q_{(X, -D), \log}$. In this work, we prove the following fundamental structural result:

Theorem [1.1](#page-1-0). Let $D = D_0 + pD_1 + \cdots + p^n D_n$ be a p-divisibility decomposition (see §1.1 for the precise definition). For each i, set $\underline{D}_i = D_0 + pD_1 + \ldots p^i D_i$. Let $U_i = X \setminus \underline{D}_i \hookrightarrow X$ be the open immersion. Then

$$
W_n \Omega_{(X,-D),\log}^q = \sum_{i=0}^{n-1} \underline{p}^i \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\lceil \frac{D}{p^i} \rceil}^{\times}) \underbrace{\operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times} \dots \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times}}_{(q-1)\text{-times}}
$$

as subsheaves of $W_n \Omega^q$ X on X_{Nis} .

There are other definitions of log de Rham-Witt sheaves with zeros along D earlier in the literature. For instance, [\[GK21\]](#page-27-5) defined a sheaf

$$
\text{Ker}(W_n \Omega_X^q \to W_n \Omega_D^q) \cap W_n \Omega_{X,\text{log}}^q,
$$

while [\[JSZ18\]](#page-27-6) defined another sheaf

$$
\text{Im}\,\Big(\text{dlog}:\text{Ker}(\mathcal{O}_X^{\times}\to\mathcal{O}_D^{\times})\otimes_{\mathbb{Z}}\underline{j_*}\mathcal{O}_U^{\times}\otimes_{\mathbb{Z}}\cdots\otimes_{\mathbb{Z}}j_*\mathcal{O}_U^{\times}\to W_n\Omega_X^q\Big).
$$

As n varies, all the three definitions form pro-systems, and these three pro-systems are isomorphic. It remained as a question how much they differ as sheaves. As a corollary of our structural theorem, we are able to give a complete answer to this question. See Remark [2.6.](#page-5-0)

This structural result for log forms with vanishing also laid the foundation for the study of Milnor K-theory in the paper to come. To shed some light, we end this introduction session by providing our definition of the Milnor K -sheaf with vanishing along D . Suppose $D = D_0 + pD_1 + \cdots + p^L D_L$ is the *p*-divisibility decomposition of *D* of maximal length (see §[1.1](#page-1-0) for the precise definition), then we define

$$
K_{q,(X,D)}^M:=\sum_{i=0}^L\operatorname{Im}\left(p^i\operatorname{Ker}({\cal O}_X^{\times}\to{\cal O}_{\lceil\frac{D}{p^i}\rceil}^{\times})\otimes_{{\mathbb Z}} j_{i*}K_{q-1,U_i}^M\to j_*K_{q,U}^M\right)
$$

as an abelian sheaf on X_{Nis} . With this definition, we arrive at a Bloch-Gabber-Kato theorem with vanishing along D . Namely, the dlog map induces an isomorphism of sheaves

$$
\mathrm{dlog} : K_{q,(X,D)}^M/pK_{q,X}^M \cap K_{q,(X,D)}^M \xrightarrow{\simeq} W_n \Omega_{(X,-D),\mathrm{log}}^q.
$$

1.1. **Notations and conventions.** Throughout the article, k denotes a perfect field of positive characteristic p. For an effective Cartier divisor A with A_{red} being a simple normal crossing divisor (SNCD), we denote $\Omega_X^q(\log A) := \Omega_X^q(\log A_{\text{red}})$.

If E is any divisor on X and $m \in \mathbb{Z} \setminus \{0\}$, then we write

$$
m \mid E \iff m
$$
 divides the multiplicity of every irreducible component of E

and

 $m \nmid E \iff m$ does not divide the multiplicity of any irreducible component of E.

Given an increasing sequence of natural numbers $1 \leq r_1 < \ldots < r_s$ we say

$$
E=E'+p^{r_1}E_1+\ldots+p^{r_s}E_s
$$

is a p-divisibility decomposition of E (with respect to $r_1 < \ldots < r_s$), if $p^{r_1} \nmid E'$ and $p^{r_i} \nmid$ $p^{r_{i-1}}E_{i-1}$, for $i=2,\ldots,s$, and $E'_{\text{red}}+\sum_{i=1}^{s}E_{i,\text{red}}$ is a reduced divisor. Note that $p \mid E_s$ is allowed and that a p-divisibility decomposition of E always exists and is uniquely determined by the sequence $r_1 < \ldots < r_s$. When $p \nmid E_s$, we say that this is the *p*-divisibility decomposition of E of maximal length. It is apparently uniquely determined by E itself.

For example if $E = \sum_i n_i \mathcal{E}_i$ with \mathcal{E}_i the irreducible components of $E, \mathcal{E}_i \neq \mathcal{E}_j$, for $i \neq j$, then to say that

$$
E = E' + p^r E_r
$$

is a *p*-divisibility decomposition of E means that

$$
E' = \sum_{i, p^r \nmid n_i} n_i \mathcal{E}_i \quad \text{and} \quad E_r = \sum_{i, p^r \mid n_i} \frac{n_i}{p^r} \mathcal{E}_i.
$$

Finally for E and m as above we set

$$
\lceil E/m \rceil := \sum_i \lceil n_i/m \rceil \mathcal{E}_i \quad \text{and} \quad \lfloor E/m \rfloor := \sum_i \lfloor n_i/m \rfloor \mathcal{E}_i,
$$

where $\lceil - \rceil$ (resp. $\lceil - \rceil$) denotes the round-up (resp. round-down).

Acknowledgement. The author thanks Kay Rülling for various discussions and for checking an earlier draft of this article. Gratitude also goes to Shuji Saito and Yigeng Zhao for raising questions that steered the exploration in this direction.

2. The structure of log de Rham-Witt differentials

Let k be a perfect field of positive characteristic p. Let X be a separated smooth k-scheme of finite type, and D be an effective Cartier divisor with D_{red} being a SNCD.

Definition 2.1 ([\[RR24,](#page-27-0) Proposition 9.3]). Define $W_n \Omega_{(X, -D), \log}^q$ to be the abelian subsheaf of $W_n \Omega_{X,\log}^q$ on X_{Nis} such that the following sequence is exact

$$
0 \to W_n\Omega_{(X,-D),\log}^q \to W_n\Omega_{(X,-D)}^q \xrightarrow{C^{-1}-1} \frac{W_n\Omega_{(X,-D)}^q}{dV^{n-1}\Omega_X^{q-1} \cap W_n\Omega_{(X,-D)}^q}.
$$

We write $\Omega_{(X, -D), \log}^q := W_1 \Omega_{(X, -D), \log}^q$.

Theorem 2.2. Let $A, B \ge 0$ be Cartier divisors with A and $(A + B)_{\text{red}}$ being SNCDs. Write $B = B' + B''$ with $B'_{red} \leq A$ and $B'_{red} + B''_{red}$ being SNCD. Let $B'' = B''_0 + pB''_1$ be a p-divisibility decomposition. Denote by $j: U = X \setminus (A + B) \hookrightarrow X$ the open immersion. For $q \geq 1$, let

$$
\Omega_X^q(\log A)(-B)_\mathrm{log}:=j_*\Omega^q_{U,\mathrm{log}}\cap \Omega_X^q(\log A)(-B).
$$

Then we have the following identity of Nisnevich subsheaves of $\Omega_X^q(\log(A+B))$

$$
\Omega_X^q(\log A)(-B)_{\log} = \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\widetilde{B}}^{\times}) \wedge \underbrace{\operatorname{dlog} j_{0*} \mathcal{O}_V^{\times} \wedge \cdots \wedge \operatorname{dlog} j_{0*} \mathcal{O}_V^{\times}}_{(q-1) \ times}.
$$

where $\widetilde{B}=B+B''_{0,\text{red}}$, and $j_0: V:=X\setminus (A+B''_0)\hookrightarrow X$ is the open immersion. In particular, since $\Omega_{(X, -D)}^q = \Omega_X^q(\log D_0)(-D)$ by [\[RR24,](#page-27-0) Lemma 8.4],

$$
\Omega^q_{(X,-D),\log} = \text{dlog Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_D^{\times}) \wedge \underbrace{\text{dlog } j_{0*} \mathcal{O}_{U_0}^{\times} \wedge \cdots \wedge \text{dlog } j_{0*} \mathcal{O}_{U_0}^{\times}}_{(q-1) \text{ times}}
$$

where $j_0 : U_0 := X \setminus D_0 \hookrightarrow X$ is the open immersion.

Proof of Theorem [2.2.](#page-2-2) Since X is of finite type over k, it suffices to check that the two subsheaves have the same Nisnevich stalk at each closed point of X . Let R be the henselization of a local ring at a closed point of X. The residue field of R is a finite extension of the base field k, and is hence perfect. Since we will be only concerned with the identification of two subsheaves on the stalk at R , to avoid complications of notations, we suppose R has residue field k. Hence we can choose a system T_1, \ldots, T_d of regular parameters of R such that

 $A = Div(T_1 ... T_e), \quad B = Div(T_1^{r_1} ... T_g^{r_g})$

where e, f, g are integers such that $0 \le e \le f \le g \le d$ and r_1, \ldots, r_g are nonnegative integers such that at least one $r_i \geq 1$, and

$$
p \nmid r_j \ge 1 \text{ for all } j \in [e+1, f], \quad p \mid r_j \ge 1 \text{ for all } j \in [f+1, g].
$$

Put $r_{g+1} = \cdots = r_d = 0$ and set

$$
\underline{r} := (r_1, \ldots, r_d) \in \mathbb{N}^d.
$$

Set

$$
G_R^{\underline{r}}:=\Omega_R^q(\log A)(-B),\quad G_{R,\log}^{\underline{r}}:=\Omega_R^q(\log A)(-B)\cap \Omega_{\mathrm{Frac}\,R,\log}^q.
$$

Theorem [2.2](#page-2-2) now follows from Proposition [2.3.](#page-3-0)

Proposition 2.3. Define

$$
\widetilde{r}_i = \begin{cases} r_i + 1 & i \in [e+1, f]; \\ r_i & i \notin [e+1, f]. \end{cases}
$$

Set
$$
\tilde{\underline{r}} := (\tilde{r}_1, ..., \tilde{r}_d)
$$
. Then
\n
$$
G_{R, \log}^{\underline{r}} = \Big\{ \sum_j^{\text{finite}} \text{dlog } x_{1,j} \wedge \text{dlog } x_{2,j} \wedge \cdots \wedge \text{dlog } x_{q,j} \Big|
$$
\n
$$
x_{1,j} \in 1 + (T_1^{\tilde{r}_1} ... T_g^{\tilde{r}_g}) \cdot R, x_{2,j}, \dots, x_{q,j} \in R[\frac{1}{T_1 ... T_f}]^{\times} \Big\}.
$$

Proposition [2.3](#page-3-0) is the main technical part of this work. Its proof is long and tedious and we postpone it to the next session.

Theorem 2.4. Let $D = D_0 + pD_1 + \cdots + p^n D_n$ be a p-divisibility decomposition. For each i, set $\underline{D}_i = D_0 + pD_1 + \dots p^i D_i$. Let $U_i = X \setminus \underline{D}_i \hookrightarrow X$ be the open immersion. Then

$$
W_n \Omega_{(X,-D),\log}^q = \sum_{i=0}^{n-1} \underline{p}^i \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\lceil \frac{D}{p^i} \rceil}^{\times}) \underbrace{\operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times} \dots \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times}}_{(q-1)\text{-times}}
$$

as subsheaves of $W_n \Omega_X^q$ on X_{Nis} . Moreover, the restriction map

$$
R: W_{n+1}\Omega_{(X,-D),\log}^q \to W_n\Omega_{(X,-D),\log}^q
$$

is surjective.

Proof. Once we have the first statement, the surjectivity of R clearly follows from the formula. Hence our aim is to show the equality

$$
W_n \Omega_{(X,-D)}^q \cap W_n \Omega_{X,\log}^q = \sum_{i=0}^{n-1} \underline{p}^i \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\lceil \frac{D}{p^i} \rceil}^{\times}) \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times} \dots \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times}.
$$

When $n = 1$, this is done in Theorem [2.2.](#page-2-2) Now we assume $n \geq 2$ and denote the right hand side of the formula above by M_n .

We first show the direction ⊃, that is

(2.4.1)
$$
\sum_{i=1}^{n-1} \underline{p}^i \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\lceil \frac{D}{p^i} \rceil}^{\times}) \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times} \dots \operatorname{dlog} j_{i*} \mathcal{O}_{U_i}^{\times} \subset W_n \Omega_{(X, -D)}^q.
$$

Since $W_n \Omega_{\ell}^q$ $\binom{q}{(X,-D)}$ is Cohen-Macaulay (see Example [4.2\(](#page-24-1)[3\)](#page-24-2)), it suffices to check [\(2.4.1\)](#page-4-0) locally around all codimension 1 points by Lemma [4.1.](#page-23-2) Let R be the henselization of the local ring of a codimension 1 point x of X , \mathfrak{m} be the maximal ideal of R , and T be a uniformizer of R. When D does not pass through the point x, $(2.4.1)$ trivially holds. Now we suppose $D = Div(T^r)$ in a neighborhood of x with $r \ge 1$. Suppose $r = p^j r'$ with $(p, r') = 1$. The Nisnevich stalk of the left hand side of $(2.4.1)$ is (2.4.2)

$$
\sum_{i=0}^{j-1} p^i \operatorname{dlog}(1 + \mathfrak{m}^{r/p^i}) \operatorname{dlog} R^\times \dots \operatorname{dlog} R^\times + \sum_{i=j}^{n-1} p^i \operatorname{dlog}(1 + \mathfrak{m}^{\lceil r/p^i \rceil}) \operatorname{dlog} R[\frac{1}{T}]^\times \dots \operatorname{dlog} R[\frac{1}{T}]^\times.
$$

(When $j > n-1$, the second summand above is 0.) This is clearly contained in the Nisnevich stalk of the right hand side of $(2.4.1)$ at x.

We show the direction ⊂. The proof below is adapted from the proof of [\[JSZ18,](#page-27-6) Theorem 2.3.1]. Do induction on n. Let $x \in W_n \Omega_0^q$ $(X_{(X,-D)}^q \cap \hat{W}_n \Omega_{X,\log}^q$ be a local section. Then $Rx \in$ $W_{n-1}\Omega_{\ell}^q$ $(X_{(X,-D)}^q \cap W_{n-1}\Omega_{X,\log}^q$, hence it lies in M_{n-1} by induction hypothesis. Consider the following commutative diagram

$$
M_n \xrightarrow{R} M_{n-1} \longrightarrow 0
$$

$$
0 \longrightarrow \Omega_{X,\log}^q \xrightarrow{p^{n-1}} W_n \Omega_{X,\log}^q \xrightarrow{R} W_{n-1} \Omega_{X,\log}^q \longrightarrow 0.
$$

Furthermore, the following claim is direct.

Claim 2.4.1. $R: M_n \to M_{n-1}$ is surjective.

Let $y \in M_n$ such that $R(y) = R(x)$. By the exactness of the second row (see [\[CTSS83,](#page-27-7) Lemme 3]), $x - y = p^{n-1}(z)$ for some $z \in \Omega_{X,\log}^q$. Note that $p^{n-1}(z) = x - y \in W_n \Omega_q^q$ $(X,-D)$ [.] Hence by [\[RR24,](#page-27-0) Lemma 6.11], we have

$$
z \in \Omega_{n-1}^q(-D', -pD_n) := \Omega_X^q(\log D')(-\lceil \frac{D}{p^{n-1}} \rceil).
$$

In particular, $z \in \Omega^q_{X, \log} \cap \Omega^q_n$ $_{n-1}^{q}(-D', -pD_n)$. By Theorem [2.2\(](#page-2-2)1), we have

$$
z \in \mathrm{dlog}\,\mathrm{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_{\lceil \frac{D}{p^{n-1}} \rceil}^{\times}) \,\mathrm{dlog}\,\mathcal{O}_{U_{n-1}}^{\times} \dots \mathrm{dlog}\,\mathcal{O}_{U_{n-1}}^{\times}
$$

where $U_{n-1} = X \setminus D'$. That is, $p^{n-1}(z) \in M_n$.

The following corollary is direct from the explicit formula for $W_n \Omega_{(X,-D),\log}^q$ given in Theorem [2.2.](#page-2-2) This is a filtered version of [\[CTSS83,](#page-27-7) Lemme 3]. Similar short exact sequences but for pro-systems were studied earlier for other filtrations, e.g. [\[GK21,](#page-27-5) Proposition 5.9] and [\[Mor19,](#page-27-8) Theorem 4.6]

Corollary 2.5. The following sequence is exact

$$
0 \to W_{n-1}\Omega_{(X,-D),\log}^q \xrightarrow{p} W_n\Omega_{(X,-D),\log}^q \xrightarrow{R^{n-1}} \Omega_{(X,-D),\log}^q \to 0.
$$

Remark 2.6. Our structural result Theorem [2.2](#page-2-2) answers a remaining question in [\[RR24,](#page-27-0) $11.5(4)$] and a question in [\[RR24,](#page-27-0) Remark 11.7(1)]. In [\[GK21\]](#page-27-5) and [\[JSZ18\]](#page-27-6) the following subsheaves of $W_n \Omega_{\mathcal{I}}^q$ $X(\log D)$ are introduced:

$$
\begin{aligned} W_n[\mathrm{GK}]^q_{(X,-D)}&:=\mathrm{Ker}(W_n\Omega_X^q\rightarrow W_n\Omega_D^q),\quad W_n[\mathrm{GK}]^q_{(X,-D),\log}:=W_n[\mathrm{GK}]^q_{(X,-D)}\cap W_n\Omega_{X,\log}^q;\\ W_n[\mathrm{JSZ}]^q_{(X,-D)}&:=W_n\mathcal{O}_X(-D)\cdot W_n\Omega_X^q(\log D),\quad W_n[\mathrm{JSZ}]^q_{(X,-D),\log}:=W_n[\mathrm{JSZ}]^q_{(X,-D)}\cap W_n\Omega_{X,\log}^q. \end{aligned}
$$

Here $W_n\mathcal{O}_X(-D) \subset W_n\mathcal{O}_X$ refers to the ideal sheaf which is a free line bundle of $W_n\mathcal{O}_X$. modules generated by the Teichmüller lifts of local sections of the effective Cartier divisor D. Then

(2.6.1)
$$
W_n[\text{GK}]_{(X, -D)}^q \subset W_n\Omega_{(X, -D)}^q \subset W_n[\text{JSZ}]_{(X, -D)}^q;
$$

(2.6.2)
$$
W_n[\text{GK}]_{(X,-D),\text{log}}^q \subset W_n\Omega_{(X,-D),\text{log}}^q \subset W_n[\text{JSZ}]_{(X,-D),\text{log}}^q.
$$

When $n = 1$, we have the following more explicit expressions (we use the notations from Theorem [2.2\)](#page-2-2):

(2.6.3)
$$
W_1[\text{GK}]_{(X,-D)}^q = \text{Im}(\mathcal{O}_X(-D) \cdot \Omega_X^q + d\mathcal{O}_X(-D) \cdot \Omega_X^{q-1} \hookrightarrow \Omega_X^q),
$$

$$
(2.6.4) \qquad \Omega_{(X,-D)}^q = \Omega_X^q(\log D_0)(-D), \qquad \Omega_{(X,-D),\log}^q = \text{dlog}\,\text{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_D^{\times}) \wedge j_{0*} K_{q-1,U_0}^M,
$$

$$
(2.6.5)
$$

$$
W_1[\operatorname{JSZ}]_{(X,-D)}^q = \Omega_X^q(\log D)(-D), \quad W_1[\operatorname{JSZ}]_{(X,-D)}^q = \operatorname{dlog} \operatorname{Ker}(\mathcal{O}_X^{\times} \to \mathcal{O}_D^{\times}) \wedge \operatorname{dlog} j_* K_{q-1,U}^M,
$$

The equation [\(2.6.3\)](#page-5-1) follows from the fact that $W_n[\text{GK}]^*_{(X, -D)}$ is the differential graded ideal of $W_n \Omega_X^*$ generated by $W_n \mathcal{O}_X(-D)$, see e.g. [Rül07, Lemma 1.20]. The equations [\(2.6.4\)](#page-5-2) follow from $[RR24, \text{ Lemma } 8.4]$ and Theorem [2.2.](#page-2-2) The second equation of $(2.6.5)$ follows from [\[JSZ18,](#page-27-6) Theorem 1.2.1] (or our Theorem [2.2\)](#page-2-2).

(1) Both inclusions in $(2.6.1)$ can be strict, this answers $[RR24, 11.5(4)]$. We illustrate this by the following explicit example when $n = 1$. Let $X = \text{Spec } A, D = \text{Div}(f^r g^s h^t)$ with $p \nmid rs, p \mid t$, and $D_{\text{red}} = Div(f) + Div(g) + Div(h)$ being an SNCD. Then

$$
\Omega_X^q(-D) \subsetneqq \text{Im}(d\mathcal{O}_X(-D) \cdot \Omega_X^{q-1} \hookrightarrow \Omega_X^q) \subsetneqq W_1[\text{GK}]_{(X, -D)}^q \subsetneqq \Omega_{(X, -D)}^q \subsetneqq W_1[\text{JSZ}]_{(X, -D)}^q
$$

where $M(-D) = M \otimes_{\mathcal{O}_X} \mathcal{O}_X(-D)$ for any \mathcal{O}_X -module M. Concretely, let $\omega \in \Omega_X^{q-1}$ X be any regular differential, then

$$
d(f^r g^s h^t) \wedge \omega \in \operatorname{Im} (d\mathcal{O}_X(-D) \cdot \Omega_X^{q-1} \to \Omega_X^q) \setminus \Omega_X^q(-D),
$$

$$
f^r h^t d(g^s) \wedge \omega \in \Omega_{(X, -D)}^q \setminus W_1[\operatorname{GK}]_{(X, -D)}^q,
$$

$$
f^r g^s h^{t-1} dh \wedge \omega \in W_1[\operatorname{JSZ}]_{(X, -D)}^q \setminus \Omega_{(X, -D)}^q.
$$

(2) While $\Omega_{(X,-D)}^q$ and $W_1[JSZ]_{(X,-D)}^q$ are both locally free, $W_1[GK]_{(X,-D)}^q$ is not a Cohen-Macaulay \mathcal{O}_X -module.

We use Lemma [4.1](#page-23-2) to see this. In fact, if it is Cohen-Macaulay, then any rational section which is integral at all codimension 1 points is an integral section. Take any $\alpha \in \Gamma(X, \Omega^q$ $\langle (X,-D) \rangle \backslash \Gamma(X,W_1[\text{GK}]_{(X,-D)}^q)$. In particular, $\alpha \in \Omega^q(\mathbb{R}^d)$ $_{(X,-D),x}^{q}$ for all $x \in X^{(1)}$. Note that the codimension 1 stalks of $\Omega_{(X, -D)}^q$ and of $W_1[\text{GK}]_{(X, -D)}^q$ are the same. We

have hence find a section α which lies in $W_1[\text{GK}]_{(X,-D),x}^q$ for all $x \in X^{(1)}$ but cannot be extended to a global section. Therefore $W_1[\text{GK}]_{(X,-D)}^{\check{q}}$ is not Cohen-Macaulay.

(3) Both inclusions in $(2.6.2)$ can be strict, this answers [\[RR24,](#page-27-0) Remark 11.7(1)]. If we take the example from (1), then

$$
\mathrm{dlog}(1 + f^r g^s h^t) \wedge \mathrm{dlog} h \in W_1[\mathrm{JSZ}]^2_{(X, -D), \mathrm{log}} \setminus \Omega^2_{(X, -D), \mathrm{log}}.
$$

$$
\mathrm{dlog}(1 + f^r g^s h^t) \wedge \mathrm{dlog} g \in \Omega^2_{(X, -D), \mathrm{log}} \setminus W_1[\mathrm{GK}]^2_{(X, -D), \mathrm{log}}.
$$

We can also see the second strict inclusion of $(2.6.2)$ directly from Theorem [2.2.](#page-2-2)

3. Proof of Proposition [2.3](#page-3-0)

This section is completely devoted to the proof of Proposition [2.3.](#page-3-0) The proof is inspired by the proofs of [\[JSZ18,](#page-27-6) Proposition 1.2.3] and of [\[Kat82,](#page-27-10) Proposition 1]. By Artin approximation, any finitely generated R-subalgebra A of R has an R-homomorphism $A \to R$. Hence it suffices to show the statement with R replaced by $k[[T_1, \ldots, T_d]]$.

Now we prove the lemma for $R = k[[T_1, \ldots, T_d]]$. We first setup some notations. For each $0 \leq i \leq d$, write

$$
R_i = k[[T_1, \ldots, \widehat{T}_i, \ldots, T_d]]
$$

so that $R = R_i[[T_i]]$. Denote

$$
\widetilde{A} = \text{Div}(T_1 \dots T_f), \quad \widetilde{B} = \text{Div}(T_1^{\widetilde{r}_1} \dots T_g^{\widetilde{r}_g}),
$$

and for each $i \in [1, g]$, set

$$
\widetilde{A}_i = \text{Div}(T_1 \dots \widehat{T}_i \dots T_f), \quad \widetilde{B}_i = \text{Div}(T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_g^{\widetilde{r}_g}),
$$

$$
A_i = \text{Div}(T_1 \dots \widehat{T}_i \dots T_e), \quad B_i = \text{Div}(T_1^{r_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_g^{r_g}).
$$

For each $i \in [g+1, d]$, set $A_i = A, B_i = B$ and similarly $\widetilde{A}_i = \widetilde{A}$ and $\widetilde{B}_i = \widetilde{B}$. (We use $\widehat{(-)}$ denotes that the corresponding element does not appear in the string.) Let I_i^q $\frac{q}{i}$ be the set of strictly increasing functions $\{1, \ldots, q\} \to \{1, \ldots, \hat{i}, \ldots, d\}$. For $s \in I_i^q$ i^q , set

$$
e(s) := \max_{1 \le j \le q} \{ s(j) \mid s(j) \le e \}, \quad f(s) := \min_{1 \le j \le q} \{ s(j) \mid s(j) \le f \}.
$$

Denote

$$
\omega_{i,s} = \operatorname{dlog} T_{s(1)} \wedge \cdots \wedge \operatorname{dlog} T_{e(s)} \wedge \operatorname{dlog} (1 + T_{e(s)+1}) \wedge \cdots \wedge \operatorname{dlog} (1 + T_{s(q)}) \in \Omega_{R_i}^q(\log A_i),
$$

 $\widetilde{\omega}_{i,s} = \text{dlog} T_{s(1)} \wedge \cdots \wedge \text{dlog} T_{f(s)} \wedge \text{dlog}(1 + T_{f(s)+1}) \wedge \cdots \wedge \text{dlog}(1 + T_{s(q)}) \in \Omega_f^q$ $R_i^q(\log A_i).$ Then $\{\omega_{i,s} \mid s \in I_i^q\}$ ^q} form a basis of $\Omega_{R_i}^q(\log A_i)$ as a free R_i -module. Similarly, $\{\widetilde{\omega}_{i,s} \mid s \in I_i^q\}$ $_i^q$ form a basis of $\Omega_{R_i}^q(\log \tilde{A}_i)$ as a free R_i -module.

The direction "⊃" of Proposition [2.3](#page-3-0) is straightforward. Now we show the reverse direction. Consider the isomorphism of R_i -modules

$$
(3.0.1)
$$

\n
$$
\left(R \otimes_{R_i} \Omega_{R_i}^q (\log A_i)\right) \oplus \left(R \otimes_{R_i} \Omega_{R_i}^{q-1} (\log A_i)\right) \xrightarrow{\simeq} \Omega_R^q (\log A),
$$

\n
$$
(a \otimes w, b \otimes v) \mapsto \begin{cases} aw + bv \wedge d \log T_i, & \text{if } 1 \leq i \leq e; \\ aw + bv \wedge d \log(1 + T_i), & \text{if } e + 1 \leq i \leq d. \end{cases}
$$

With these notations, we have the following easy description of the graded piece of $G_{\overline{f}}^r$ $\frac{r}{R}$: **Lemma 3.1.** Fix $i \in [1, d]$. Then [\(3.0.1\)](#page-6-1) induces the following isomorphism of R_i -modules

$$
(T_1^{r_1} \dots \widehat{T}_i^{r_i} \dots T_d^{r_d}) \cdot \Omega_{R_i}^q (\log A_i) \oplus (T_1^{r_1} \dots \widehat{T}_i^{r_i} \dots T_d^{r_d}) \cdot \Omega_{R_i}^{q-1} (\log A_i)
$$

\n
$$
\xrightarrow{\simeq} G_R^r / G_R^{(r_1, \dots, r_{i-1}, r_i+1, r_{i+1}, \dots, r_d)}
$$

\n
$$
(w, v) \mapsto \begin{cases} T_i^{r_i} w + T_i^{r_i} v \wedge \text{dlog} T_i & i \in [1, e], \\ T_i^{r_i} w + T_i^{r_i} v \wedge \text{dlog}(1 + T_i) & i \in [e+1, d]. \end{cases}
$$

For any $i \in [1, d]$ and $l \geq 0$, define

 $V_i^l =$ the image via [\(3.0.1\)](#page-6-1) of $(T_i^l) \otimes_{R_i} \Omega_f^q$ $\mathcal{C}_{R_i}^q(\log A_i) \oplus (T_i^l) \otimes_{R_i} \Omega_{R_i}^{q-1}$ $q_{R_i}^{q-1}(\log A_i),$

For each i, $\{V_i^l\}_{l\geq 0}$ form a decreasing filtration for $\Omega_R^q(\log A)$ which is exhaustive. In particular, $V_i^{r_i} \supset G_I^{\dot{r}}$ R_R^{r} for any *i*. Recall that $\Omega_R^q(\log A)_{\log} := \Omega_R^q(\log A) \cap \Omega_{\text{Frac }R,\log}^q$. Set

$$
U_i^l := V_i^l \cap \Omega_R^q(\log A)_{\log}.
$$

We will repeatedly use the following fact below:

(3.1.1)
$$
if C, D \in \Omega_R^q(\log A)_{\log} \text{ and } C \equiv D \mod V_i^h, \text{ then } C \equiv D \mod U_i^h.
$$

Lemma 3.2. Let $h > 0$ be a positive integer.

(1) If $i \in [1, e]$, any q-form in U_i^h/U_i^{h+1} is of the form

$$
\sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}T_i^h) \wedge w_{i,s} + \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}T_i^h) \wedge w'_{i,t} \wedge \text{dlog} T_i
$$

where $a_{i,s}, b_{i,t} \in R_i, w_{i,s}$, and $w_{i,s} \in \Omega^{q-1}_{R_i}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $q_{R_i}^{q-2}(\log A_i)_{\log}.$ (2) If $i \in [e+1,g]$, p | h, then any q-form in U_i^h/U_i^{h+1} is of the form

$$
\sum_{s=1}^{\text{finite}} \left(\mathrm{dlog}(1 + a'_{i,s} T_i^{h+1}) \wedge w_{i,s} + \mathrm{dlog}(1 + a_{i,s} T_i^h) \wedge w_{i,s} \right) + \sum_{t=1}^{\text{finite}} \mathrm{dlog}(1 + b_{i,t} T_i^h) \wedge w'_{i,t} \wedge \mathrm{dlog}(1 + T_i),
$$

where $a'_{i,s}, a_{i,s}, b_{i,t} \in R_i$, and $w_{i,s} \in \Omega_{R_i}^{q-1}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}.$ (3) If $i \in [e+1,g]$, $p \nmid h(h+1)$, then any q-form in U_i^h/U_i^{h+1} is of the form

$$
\sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}T_i^{h+1}) \wedge w_{i,s} + \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}T_i^h) \wedge w'_{i,t} \wedge \text{dlog}(1 + T_i),
$$

where $a_{i,s}, b_{i,t} \in R_i$, and $w_{i,s} \in \Omega_{R_i}^{q-1}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}.$ (4) If $i \in [e+1,g]$, p $|(h+1)$, then any q-form in $v \in U_i^h/U_i^{h+1}$ is of the form

$$
\sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t} T_i^h) \wedge w'_{i,t} \wedge \text{dlog}(1 + T_i),
$$

where $b_{i,t} \in R_i$, and $w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}.$

Proof. Every element in $R = R_i[[T_i]]$ which is a unit can be written uniquely as a infinite product

$$
u \prod_{l \ge 0} (1 - a_l T_i^l), \quad u \in k^{\times}, a_l \in R_i.
$$

Therefore $\Omega^1_{R,\text{log}}$ is generated by convergent power series of

$$
d\log(1 + a_l T_i^l)
$$

with $a_l \in R_i$ and $l \geq 0$. Moreover, for any $j \in [1, d]$, the natural map

$$
R^{\times} \oplus \mathbb{Z}^{j} \to R[\frac{1}{T_1 \dots T_j}]^{\times}, \quad (f, (g_1, \dots, g_j)) \mapsto f \cdot T_1^{g_1} \dots T_j^{g_j}
$$

is an isomorphism. In particular, $\Omega^1_{R[\frac{1}{T_1...T_d}],\log}$ is generated by convergent power series of

$$
d\log(1 + a_l T_i^l), \, d\log T_1, \, \dots, \, d\log T_d
$$

with $a_l \in R_i$, $l \geq 0$. An element of $\Omega^1_{R[\frac{1}{T_1...T_d}],\log}$ lies in $\Omega^1_R(\log A)_{\log}$ if its residue in terms of T_i lies in $\Omega_{R_i}^{q-1}(\log A_i)$, for all $i = e+1, \ldots, d$. Therefore $\Omega_R^1(\log A)_{\log}$ is generated by convergent power series of

(3.2.1) $\text{dlog}(1 + a_l T_i^l), \quad (a_l \in R_i, l \ge 0)$

(3.2.2) dlog T1, . . . , dlog Te.

(Note that $d\log(1+T_{e+1}), \ldots, d\log(1+T_d)$ are among these generators.)

Observation 3.2.1. Let $w \in \Omega_{R_1}^{q-1}$ $_{R_i}^{q-1}(\log A_i), w' \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)$, and $a \in R_i$. (1) If $i \in [1, e]$ and $h \geq 1$, then

$$
\mathrm{dlog}(1 + aT_i^h) \wedge w = \frac{haT_i^h \mathrm{dlog} T_i + T_i^h da}{1 + aT_i^h} \wedge w \in V_i^h,
$$

$$
\operatorname{dlog}(1 + aT_i^h) \wedge w' \wedge \operatorname{dlog} T_i = \frac{T_i^h da}{1 + aT_i^h} \wedge w' \wedge \operatorname{dlog} T_i \in V_i^h.
$$

(2) If $i \in [e+1, q], p \mid h > 1, then$

$$
d\log(1 + aT_i^h) \wedge w = \frac{T_i^h da}{1 + aT_i^h} \wedge w \in V_i^h,
$$

$$
\mathrm{dlog}(1+aT_i^h) \wedge w' \wedge \mathrm{dlog}(1+T_i) = \frac{T_i^h da}{1+aT_i^h} \wedge w' \wedge \mathrm{dlog}(1+T_i) \in V_i^h.
$$

(3) But if $i \in [e+1, f]$ and $p \nmid h \geq 1$, then

$$
\operatorname{dlog}(1 + aT_i^h) \wedge w = \frac{haT_i^{h-1}(1 + T_i) \operatorname{dlog}(1 + T_i) + T_i^h da}{1 + aT_i^h} \wedge w \in V_i^{h-1},
$$

$$
\operatorname{dlog}(1 + aT_i^h) \wedge w' \wedge \operatorname{dlog}(1 + T_i) = \frac{T_i^h da}{1 + aT_i^h} \wedge w' \wedge \operatorname{dlog}(1 + T_i) \in V_i^h.
$$

Observation 3.2.2. By [\[RS18,](#page-27-11) Lemma 2.7(1)], for $l_1, l_2 \ge 1$ and $a, b \in R_i$,

$$
\{1 + aT_i^{l_1}, 1 + bT_i^{l_2}\} = -\{1 + \frac{ab}{1 + aT_i^{l_1}}T_i^{l_1+l_2}, -a(1 + bT_i^{l_2}) \cdot T_i^{l_1}\} \text{ in } K_2^M(\text{Frac }R).
$$

Hence if there are more than one factor of the shape [\(3.2.1\)](#page-8-0) with $l \geq 1$ in a wedge product, one can always reorganize them so that only one such factor appears.

We continue the proof of Lemma [3.2.](#page-7-0) [\(1\)](#page-7-1) is straightforward. It remains to prove $(2)-(4)$ $(2)-(4)$. Let $h_1 \in p\mathbb{Z}$ be the unique integer with $h \in [h_1, h_1 + p)$. Write $h = h_0 + h_1$ with $h_0 \in [0, p)$. By Observation [3.2.1](#page-8-1) and Observation [3.2.2,](#page-8-2) a general element $v \in U_i^{h_1}/U_i^{h_1+p}$ is of the form

$$
(3.2.3)\sum_{l=0}^{p-1}\sum_{s=1}^{\text{finite}}\text{dlog}(1+a_{i,s}^{(l)}T_i^{h_1+l})\wedge w_{i,s}+\sum_{l=0}^{p-1}\sum_{t=1}^{\text{finite}}\text{dlog}(1+b_{i,t}^{(l)}T_i^{h_1+l})\wedge w_{i,t}'\wedge \text{dlog}(1+T_i).
$$

where all $a_{i,s}^{(l)}, b_{i,t}^{(l)} \in R_i$, and $w_{i,s} \in \Omega_{R_i}^{q-1}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}$. [\(2\)](#page-7-2) then follows directly from this expression. It remains to discuss (3) and (4) . Obviously, it will be sufficient to prove the following claim.

Claim 3.2.1. If $i \in [e+1,g]$, $p \nmid h$, then any q-form in U_i^h/U_i^{h+1} is of the form

$$
\sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}T_i^{h+1}) \wedge w_{i,s} + \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}T_i^h) \wedge w'_{i,t} \wedge \text{dlog}(1 + T_i),
$$

where $a_{i,s}, b_{i,t} \in R_i$, and $w_{i,s} \in \Omega_{R_i}^{q-1}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}.$

Case 1. $h_1 \geq p$. In this case, we have

$$
(a_{i,s}^{(l)}T_i^{h_1+l})^{l'}d(a_{i,s}^{(l)}T_i^{h_1+l}) \wedge w_{i,s} \in V_i^{h_1+p},
$$

$$
(b_{i,t}^{(l)}T_i^{h_1+l})^{l'} \cdot T_i^{h_1+l}db_{i,t}^{(l)} \wedge w_{i,t} \wedge d\log(1+T_i) \in V_i^{h_1+p}
$$

for all $l \in [0, p-1]$, $l' \geq 1$. These two relations imply that

$$
d \log(1 + a_{i,s}^{(l)} T_i^{h_1 + l}) \wedge w_{i,s} \equiv d(a_{i,s}^{(l)} T_i^{h_1 + l}) \wedge w_{i,s} \mod V_i^{h_1 + p},
$$

$$
\operatorname{dlog}(1+b_{i,t}^{(l)}T_i^{h_1+l}) \wedge w_{i,t} \wedge \operatorname{dlog}(1+T_i) \equiv T_i^{h_1+l} db_{i,t}^{(l)} \wedge w_{i,t} \wedge \operatorname{dlog}(1+T_i) \mod V_i^{h_1+p}.
$$

We will repeatedly use these congrence relations below.

Now let $v \in U_i^{h_1+h_0}/U_i^{h_1+h_0+1}$, and we prove $(3)(4)$ by decreasing induction on h_0 . We start from $h_0 = p - 1$, i.e., $v \in U_i^{h_1 + p - 1}$ $i^{h_1+p-1}/U_i^{h_1+p}$. A direct computation from [\(3.2.3\)](#page-8-3) shows

$$
v = \sum_{l=0}^{p-1} \sum_{s=1}^{\text{finite}} T_i^{h_1+l} da_{i,s}^{(l)} \wedge w_{i,s}
$$

+
$$
\sum_{l=0}^{p-2} T_i^{h_1+l} \left(\sum_{s=1}^{\text{finite}} (-1)^{q-1} (h_1 + l + 1) a_{i,s}^{(l+1)} w_{i,s} + \sum_{t=1}^{\text{finite}} \left(\sum_{l'=0}^l (-1)^{l-l'} db_{i,t}^{(l')} \right) \wedge w_{i,t} \right) \wedge dT_i
$$

+
$$
T_i^{h_1+p-1} \left(\sum_{t=1}^{\text{finite}} (db_{i,t}^{(p-1)} - db_{i,t}^{(p-2)} + \dots + (-1)^{p-1} db_{i,t}^{(0)}) \wedge w_{i,t} \right) \wedge dT_i \qquad \text{mod } V_i^{h_1+p}.
$$

The condition $v \in V_i^{h_1+p-1}$ forces

$$
\sum_{s=1}^{\text{finite}} da_{i,s}^{(l)} \wedge w_{i,s} = 0, \text{ for all } l \in [0, p-2]; \text{ and}
$$

$$
\sum_{s=1}^{\text{finite}} (-1)^{q-1} (h_1 + l + 1) a_{i,s}^{(l+1)} w_{i,s} + \sum_{t=1}^{\text{finite}} \left(\sum_{l'=0}^l (-1)^{l-l'} db_{i,t}^{(l')} \right) \wedge w_{i,t} = 0, \text{ for all } l \in [0, p-2].
$$

Taking $l = n-2$ in the second vanishing condition above and differentiating it, we get

Taking $l = p - 2$ in the second vanishing condition above and differentiating it, we get

$$
\sum_{s=1}^{\text{finite}} da_{i,s}^{(p-1)} \wedge w_{i,s} = 0.
$$

Therefore

$$
v \equiv T_i^{h_1+p-1} \left(\sum_{t=1}^{\text{finite}} (db_{i,t}^{(p-1)} - db_{i,t}^{(p-2)} + \dots + (-1)^{p-1} db_{i,t}^{(0)}) \wedge w_{i,t} \right) \wedge dT_i \mod V_i^{h_1+p}
$$

$$
\equiv \sum_{t=1}^{\text{finite}} \text{dlog}(1 + \left(\sum_{l=0}^{p-1} (-1)^{p-1-l} b_{i,t}^{(l)}\right) T_i^{h_1+p-1}) \wedge w_{i,t} \wedge \text{dlog}(1+T_i) \mod V_i^{h_1+p}.
$$

Since both sides of the congruence equation above lie in $U_i^{h_1+p-1}$, their difference lies in $U_i^{h_1+p-1} \cap V_i^{h_1+p} = U_i^{h_1+p}$ by [\(3.1.1\)](#page-7-5). In other words, any general element $v \in U_i^{h_1+p-1}/U_i^{h_1+p}$ is of the form

(3.2.4)
$$
\sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t} T_i^{h_1+p-1}) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i)
$$

for some $b_{i,t} \in R_i$. In particular, Claim [3.2.1](#page-9-0) holds in this case.

For the induction step, suppose that any element in $U_i^{h_1+h_0}/U_i^{h_1+h_0+1}$ $(h_0 \leq p-1)$ is of the form

$$
(3.2.5) \quad \sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s} T_i^{h_1 + h_0 + 1}) \wedge w_{i,s} + \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t} T_i^{h_1 + h_0}) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i)
$$

for some $a_{i,s}, b_{i,t} \in R_i$. Let $v \in U_i^{h_1+h_0-1}/U_i^{h_1+h_0}$. Following the general form $(3.2.3)$, we can assume

$$
v \equiv \sum_{l=0}^{h_0} \sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}^{(l)} T_i^{h_1+l}) \wedge w_{i,s} + \sum_{l=0}^{h_0-1} \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}^{(l)} T_i^{h_1+l}) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i)
$$

mod $V_i^{h_1+h_0}$.

A direct computation shows

$$
v \equiv \sum_{l=0}^{h_0-1} \sum_{s=1}^{\text{finite}} T_i^{h_1+l} da_{i,s}^{(l)} \wedge w_{i,s}
$$

+
$$
\sum_{l=0}^{h_0-2} T_i^{h_1+l} \left(\sum_{s=1}^{\text{finite}} (-1)^{q-1} (h_1 + l + 1) a_{i,s}^{(l+1)} w_{i,s} + \sum_{t=1}^{\text{finite}} \left(\sum_{l'=0}^l (-1)^{l-l'} db_{i,t}^{(l')} \right) \wedge w_{i,t} \right) \wedge dT_i
$$

+
$$
T_i^{h_1+h_0-1} \left(\sum_{s=1}^{\text{finite}} (-1)^{q-1} (h_1 + h_0) a_{i,s}^{(h)} w_{i,s} + \sum_{t=1}^{\text{finite}} \left(\sum_{l'=0}^{h_0-1} (-1)^{h_0-1-l'} db_{i,t}^{(l')} \right) \wedge w_{i,t} \right) \wedge dT_i
$$

mod $V_i^{h_1+h_0}$.

The condition $v \in V_i^{h_1+h_0-1}$ forces

$$
\sum_{s=1}^{\text{finite}} da_{i,s}^{(l)} \wedge w_{i,s} = 0, \text{ for all } l \in [0, h_0 - 2]; \text{ and}
$$

$$
\sum_{s=1}^{\text{finite}} a_{i,s}^{(l+1)} w_{i,s} + \sum_{t=1}^{\text{finite}} (db_{i,t}^{(l)} + \dots + db_{i,t}^{(0)}) \wedge w_{i,t} = 0, \text{ for all } l \in [0, h_0 - 2].
$$

Taking $l = p - 2$ in the second vanishing condition above and differentiating it, we get

$$
\sum_{s=1}^{\text{finite}} da_{i,s}^{(h-1)} \wedge w_{i,s} = 0,
$$

and therefore

$$
v \equiv T_i^{h_1 + h_0 - 1} \left(\sum_{s=1}^{\text{finite}} (-1)^{q-1} (h_1 + h_0) a_{i,s}^{(h)} w_{i,s} + \sum_{t=1}^{\text{finite}} \left(\sum_{l=0}^{h_0 - 1} (-1)^{h_0 - 1 - l} db_{i,t}^{(l)} \right) \wedge w_{i,t} \right) \wedge dT_i
$$

\nmod $V_i^{h_1 + h_0}$
\n
$$
\equiv \sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}^{(h)} T_i^{h_1 + h_0}) \wedge w_{i,s}
$$

\n
$$
+ \sum_{t=1}^{\text{finite}} \text{dlog}(1 + \left(\sum_{l=0}^{h_0 - 1} (-1)^{h_0 - 1 - l} b_{i,t}^{(l)} \right) T_i^{h_1 + h_0 - 1}) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i) \mod V_i^{h_1 + h_0}.
$$

Since both sides of the congruence equation above lies in $U_i^{h_1+h_0-1}$, their subtraction lies in $U_i^{h_1+h_0-1} \cap V_i^{h_1+h_0} = U_i^{h_1+h_0}$ by [\(3.1.1\)](#page-7-5). This finishes the induction step.

Case 2. $h \in [1, p)$. A general element U_i^0/U_i^p is of the form

$$
(3.2.6) \qquad \sum_{l=0}^{p-1} \sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}^{(l)} T_i^l) \wedge w_{i,s} + \sum_{l=0}^{p-1} \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}^{(l)} T_i^l) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i).
$$

(Due to formatting reasons, below we will freely omit some indices which should be clear from the context.) We prove by decreasing induction on h the general form of an element $v \in U_i^h/U_i^{h+1}$. First let $h = p - 1$, i.e., $v \in U_i^{p-1}$ i^{p-1}/U_i^p . [\(3.2.6\)](#page-11-0) implies

$$
v = \sum_{s} \text{dlog}(1 + a^{(0)}) \wedge w_{i,s}
$$

+
$$
\sum_{l=1}^{p-1} \sum_{s} \left(\sum_{l'=0}^{\infty} (-a^{(l)}T^{l})^{l'} \cdot (T^{l}da^{(l)} + la^{(l)}T^{l-1}dT) \right) \wedge w_{i,s}
$$

+
$$
\sum_{t} \text{dlog}(1 + b^{(0)}) \wedge w_{i,t} \wedge \left(\sum_{l'' \geq 0}^{\infty} (-T)^{l''} \right) dT
$$

+
$$
\sum_{l=1}^{p-1} \sum_{t} \left(\sum_{l' \geq 0}^{\infty} (-b^{(l)}T^{l})^{l'} \cdot T^{l}db^{(l)} \right) \wedge w_{i,t} \wedge \left(\sum_{l'' \geq 0}^{\infty} (-T)^{l''} \right) dT
$$

=
$$
\sum_{s} \text{dlog}(1 + a^{(0)}) \wedge w_{i,s}
$$

+
$$
\sum_{l=1}^{p-1} \sum_{s} T^{l} \left(\sum_{l'|l} (-a^{(l')})^{l/l' - 1} da^{(l')} \right) \wedge w_{i,s}
$$

+
$$
\sum_{l=0}^{p-1} T^{l} \left(\sum_{s} \left(\sum_{l'|l+1} l'a^{(l')} \cdot (-a^{(l')})^{(l+1)/l' - 1} \right) \cdot w_{i,s}
$$

+
$$
\sum_{t} \left((-1)^{l} \text{dlog}(1 + b^{(0)}) + \sum_{l'=1}^{l} \sum_{l'' \geq 0} (-1)^{l - l'(l''+1)} (-b^{(l')})^{l''} db^{(l')} \right) \wedge w_{i,t} \right) \wedge dT
$$

 $\mod V_i^p$ i^p . The condition that $v \in V_i^{p-1}$ i^{p-1} forces

(3.2.7)
$$
\sum_{s} \mathrm{dlog}(1 + a^{(0)}) \wedge w_{i,s} = 0;
$$

(3.2.8)
$$
\sum_{s} \left(\sum_{l'|l} (-a^{(l')})^{l/l'-1} da^{(l')} \right) \wedge w_{i,s} = 0 \text{ for all } l \in [1, p-2];
$$

$$
(3.2.9) \qquad \sum_{s} \left(\sum_{l' \mid (l+1)} (-1)^{(l+1)/l'-1} l'(a^{(l')})^{(l+1)/l'} \right) \cdot w_{i,s} + \sum_{t} \left((-1)^{t} \operatorname{dlog}(1+b^{(0)}) + \sum_{l'=1}^{l} \sum_{\substack{l'' \geq 0 \\ l'(l''+1) \leq l}} (-1)^{l-l'(l''+1)} (-b^{(l')})^{l''} db^{(l')} \right) \wedge w_{i,t} = 0
$$

for all $l \in [0, p-2]$.

Taking $l = p - 2$ in [\(3.2.9\)](#page-12-0), differentiating it and dividing it by $(p - 1)$, we deduce that

$$
\sum_{s} \left(\sum_{l' \mid (p-1)} (-a^{(l')})^{(p-1)/l'-1} da^{(l')} \right) \wedge w_{i,s} = 0.
$$

That is, $(3.2.8)$ holds for all $l \in [1, p-1]$. As a result, (3.2.10)

$$
v \equiv T^{p-1} \Bigg(\sum_{s} \Bigg(\sum_{l'|p} l' a^{(l')} \cdot (-a^{(l')})^{p/l'-1} \Bigg) \cdot w_{i,s} + \sum_{t} \Bigg((-1)^{p-1} \operatorname{dlog}(1+b^{(0)})
$$

+
$$
\sum_{l'=1}^{p-1} \sum_{\substack{l'' \geq 0 \\ l''(l''+1) \leq p-1}} (-1)^{p-1-l'(l''+1)+l''} (b^{(l')})^{l''} db^{(l')}\Bigg) \wedge w_{i,t} \Bigg) \wedge dT \mod V_i^p
$$

=
$$
T^{p-1} \sum_{s} \Bigg((-1)^{p-1} \cdot (a^{(1)})^p \Bigg) \cdot w_{i,s} \wedge dT
$$

+
$$
T^{p-1} \Bigg(\sum_{t} \Bigg((-1)^{p-1} \operatorname{dlog}(1+b^{(0)})
$$

+
$$
\sum_{l'=1}^{p-1} \sum_{\substack{l'' \geq 0 \\ l''(l''+1) \leq p-1}} (-1)^{p-1-l'(l''+1)+l''} (b^{(l')})^{l''} db^{(l')}\Bigg) \wedge w_{i,t} \Bigg) \wedge \operatorname{dlog}(1+T) \mod V_i^p
$$

=
$$
\sum_{s} (-1)^{p-1} C^{-1} (w_{i,s} \wedge a^{(1)} dT_i)
$$

+
$$
\sum_{t} \sum_{l'=1}^{p-1} \sum_{l'' \geq 0} (-1)^{p-1} \frac{(-1)^{l'l''+l'+l''}}{l''+1} \operatorname{dlog}(1+b^{(l')})^{l''+1} T_i^{p-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1+T_i) \mod V_i^p
$$

 i^p .

In the second congruence relation, we have used that $pa^{(p)} = 0$ in R_i . Note that

(3.2.11)
$$
C^{-1}(a^{(1)}dT \wedge w_{i,s}) \equiv -C^{-1}(Td(a^{(1)}w_{i,s})) \mod B_2
$$

$$
\equiv 0 \mod V_i^p.
$$

In the meanwhile,

$$
(3.2.12) \t C^{-1}(\text{dlog}(1+b^{(0)}) \wedge w_{i,t} \wedge dT) = (-1)^{q-1}C^{-1}(d(T \text{dlog}(1+b^{(0)}) \wedge w_{i,t})) \in \mathcal{B}_2.
$$

Here $\mathcal{B}_n := C^{-n} d\Omega_X^{q-1}$ $\chi^{q-1}(\log A)$ for any $n \geq 1$. Let $\mathcal{B}_{\infty} := \bigcup_{n \geq 1} \mathcal{B}_n$. By Lemma [4.4,](#page-26-1) the natural map

(3.2.13)
$$
\Omega_R^q(\log A)_{\log} \to \Omega_R^q(\log A)/\mathcal{B}_{\infty}
$$

is an injection. In particular,

$$
\Omega_R^q(\log A)_{\log} \simeq \text{Image}\left(\Omega_R^q(\log A)_{\log} \hookrightarrow \frac{\Omega_R^q(\log A)}{\mathcal{B}_{\infty}}\right) = \text{Image}\left(\Omega_R^q(\log A)_{\log} + \mathcal{B}_{\infty} \rightarrow \frac{\Omega_R^q(\log A)}{\mathcal{B}_{\infty}}\right).
$$

Moreover, for any l, since $\Omega_R^q(\log A)_{\log} \cap (V_i^l + \mathcal{B}_{\infty})$ is the preimage of the image of $\Omega_R^q(\log A)_{\log} \cap$ V_i^l under the injective map $(3.2.13)$, we have

(3.2.14)
$$
\Omega_R^q(\log A)_{\log} \cap V_i^l = \Omega_R^q(\log A)_{\log} \cap (V_i^l + \mathcal{B}_{\infty}).
$$

Now [\(3.2.10\)](#page-12-2), [\(3.2.11\)](#page-13-1), [\(3.2.12\)](#page-13-2) and [\(3.2.14\)](#page-13-3) together imply

$$
v \equiv \sum_{t} \sum_{l'=1}^{p-1} \sum_{l'' \ge 0}^{\frac{p-1}{l'}-1} (-1)^{p-1} \frac{(-1)^{l'l''+l'+l''}}{l''+1} \operatorname{dlog}(1+(b^{(l')})^{l''+1} T_i^{p-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1+T_i) \mod U_i^p.
$$

Taking into account that

$$
\begin{aligned} \operatorname{dlog}(1 + bT_i^{p-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1 + T_i) + \operatorname{dlog}(1 + b'T_i^{p-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1 + T_i) \\ &\equiv \operatorname{dlog}(1 + (b+b')T_i^{p-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1 + T_i) \mod U_i^p \end{aligned}
$$

for any $b, b' \in R_i$, we deduce that any $v \in U_i^{p-1}$ i^{p-1}/U_i^p is of the form

$$
\sum_{t} d \log(1 + b T_i^{p-1}) \wedge w_{i,t} \wedge d \log(1 + T_i), \quad b \in R_i.
$$

In particular, Claim [3.2.1](#page-9-0) holds in this case.

Now we prove the induction step. Suppose we have proven that any general element in U_i^h/U_i^{h+1} (with $h \leq p-1$) is of the form

$$
\sum_{s=1}^{\text{finite}} d\log(1 + a_{i,s}^{(h+1)}T_i^{h+1}) \wedge w_{i,s} + \sum_{t=1}^{\text{finite}} d\log(1 + b_{i,t}^{(h)}T_i^h) \wedge w_{i,t} \wedge d\log(1 + T_i).
$$

Let $v \in U_i^{h-1}/U_i^h$. By [\(3.2.6\)](#page-11-0), we can write

$$
v = \sum_{l=0}^{h} \sum_{s=1}^{\text{finite}} \text{dlog}(1 + a_{i,s}^{(l)} T_i^l) \wedge w_{i,s} + \sum_{l=0}^{h-1} \sum_{t=1}^{\text{finite}} \text{dlog}(1 + b_{i,t}^{(l)} T_i^l) \wedge w_{i,t} \wedge \text{dlog}(1 + T_i).
$$

Similar as the calculations before, the condition that $v \in V_i^{p-1}$ i^{p-1} forces

(3.2.15)
$$
\sum_{s} \text{dlog}(1 + a^{(0)}) \wedge w_{i,s} = 0
$$

(3.2.16)
$$
\sum_{s} \left(\sum_{l'|l} (-a^{(l')})^{l/l'-1} da^{(l')} \right) \wedge w_{i,s} = 0 \text{ for all } l \in [1, h-2],
$$

$$
(3.2.17) \sum_{s} \left(\sum_{l' \mid (l+1)} (-1)^{(l+1)/l'-1} l'(a^{(l')})^{(l+1)/l'} \right) \cdot w_{i,s} + \sum_{t} \left((-1)^{t} \operatorname{dlog}(1+b^{(0)}) + \sum_{l'=1}^{l} \sum_{\substack{l'' \geq 0 \\ l'(l''+1) \leq l}} (-1)^{l-l'(l''+1)} (-b^{(l')})^{l''} db^{(l')} \right) \wedge w_{i,t} = 0
$$

for all $l \in [0, h-2]$.

Taking $l = h - 2$ in [\(3.2.17\)](#page-14-0), differentiating it and dividing the result by $(h - 1)$, we deduce that $\overline{ }$

$$
\sum_{s} \left(\sum_{l' | (h-1)} (-a^{(l')})^{(p-1)/l'-1} da^{(l')} \right) \wedge w_{i,s} = 0.
$$
for all $l \in [1, h-1]$. Hence

 \mathbb{R}^2

That is, [\(3.2.16\)](#page-14-1) holds for all $l \in [1, h-1].$ Hence

$$
v \equiv T^{h-1} \Bigg(\sum_{s} \Bigg(\sum_{l'|h} l' a^{(l')} \cdot (-a^{(l')})^{h/l'-1} \Bigg) \cdot w_{i,s} + \sum_{t} \Bigg((-1)^{h-1} \operatorname{dlog}(1+b^{(0)})
$$

+
$$
\sum_{l'=1}^{h-1} \sum_{l'' \atop l'''' \ge 0} (-1)^{h-1-l'(l''+1)+l''} (b^{(l')})^{l''} db^{(l')} \Bigg) \wedge w_{i,t} \Bigg) \wedge dT \mod V_i^h
$$

=
$$
T^{h-1} \Bigg(\sum_{s} \sum_{l'|h} (-1)^{h/l'-1} l' \cdot (a^{(1)})^{h/l'} \cdot w_{i,s} \Bigg) \wedge \operatorname{dlog}(1+T)
$$

+
$$
T^{h-1} \Bigg(\sum_{t} \Bigg((-1)^{h-1} \operatorname{dlog}(1+b^{(0)})
$$

+
$$
\sum_{l'=1}^{h-1} \sum_{l'' \ge 0 \atop l''(l''+1) \le h-1} (-1)^{h-1-l'(l''+1)+l''} (b^{(l')})^{l''} db^{(l')} \Bigg) \wedge w_{i,t} \Bigg) \wedge \operatorname{dlog}(1+T) \mod V_i^h
$$

=
$$
\sum_{s} \sum_{l'|h} \frac{(-1)^{h/l'-1} l'}{h} \cdot w_{i,s} \wedge \operatorname{dlog}(1+(a^{(1)})^{h/l'} T^h)
$$

+
$$
\sum_{t} (-1)^{h-1} \operatorname{dlog}(1+b^{(0)}) \wedge w_{i,t} \wedge T^{h-1} dT
$$

+
$$
\sum_{t} \sum_{l'' \ge 0 \atop l''(l''+1) \le h-1} \frac{(-1)^{p-1+l'(l''+1)+l''}}{l''+1} \operatorname{dlog}(1+(b^{(l')})^{l''+1} T^{h-1}) \wedge w_{i,t} \wedge \operatorname{dlog}(1+T)
$$

 $\mod V_i^h$.

Note that

(3.2.18) dlog(1 + b (0)) ∧ wi,t ∧ T h−1 dT = (−1)q−¹ h d(T ^h dlog(1 + b (0)) ∧ wi,t) ∈ B1.

[\(3.2.14\)](#page-13-3) implies

$$
v \equiv \sum_{s} \sum_{l'|h} \frac{(-1)^{h/l'-1}l'}{h} \cdot w_{i,s} \wedge \text{dlog}(1 + (a^{(1)})^{h/l'}T^h)
$$

+
$$
\sum_{t} \sum_{l'=1}^{h-1} \sum_{\substack{l'' \geq 0 \\ l''(l''+1) \leq h-1}} \frac{(-1)^{p-1+l'(l''+1)+l''}}{l''+1} \text{dlog}(1 + (b^{(l')})^{l''+1}T^{h-1}) \wedge w_{i,t} \wedge \text{dlog}(1+T) \mod U_i^h.
$$

This last expression and the additivity of such forms at the positions $(a^{(1)})^{h/l'}$ and $(b^{(l')})^{l''+1}$ give the induction step. \Box

Lemma 3.3. Set

$$
\mathcal{Z}^q := \operatorname{Ker}(d : \Omega^q_{R_i}(\log A_i) \to \Omega^{q+1}_{R_i}(\log A_i)).
$$

As a convention, we set $\Omega^{-1} = 0$, and

$$
\Omega_{R_i}^0(\log A_i) = R_i.
$$

Let $h \geq 0$ be an integer.

(1) If $e > 0$, $i \in [1, e]$ and $p \nmid h$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \xrightarrow{\simeq} U_i^h/U_i^{h+1}
$$

$$
\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s} \mapsto \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s}T_i^h) \wedge \omega_{i,s},
$$

where $a_{i,s} \in R_i$. Moreover,

$$
0 \to \frac{\Omega_{R_i}^{q-2}(\log A_i)}{\mathcal{Z}^{q-2}} \xrightarrow{\beta} U_i^h / U_i^{h+1} \xrightarrow{\alpha} \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \to 0
$$

is a short exact sequence, where β is defined by

(3.3.1)
$$
\beta(\sum_{t} b_{i,t} \omega_{i,t}) = \sum_{t} \text{dlog}(1 + b_{i,t} T_i^h) \wedge \omega_{i,t} \wedge \text{dlog} T_i
$$

and α is defined by the inverse of $\rho_{i,h}$ followed by the projection.

(2) If $e > 0$, $i \in [1, e]$ and $p \mid h \ge 1$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h} : \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \oplus \frac{\Omega_{R_i}^{q-2}(\log A_i)}{\mathcal{Z}^{q-2}} \xrightarrow{\simeq} U_i^h / U_i^{h+1}
$$
\n
$$
\left(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}\right) \mapsto \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s} T_i^h) \wedge \omega_{i,s} + \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^h) \wedge \omega_{i,t} \wedge \text{dlog} T_i,
$$

where $a_{i,s}, b_{i,t} \in R_i$.

(3) If $e > 0$, $i \in [1, e]$ and $h = 0$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h}: \Omega_{R_i}^q(\log A_i)_{\log} \oplus \Omega_{R_i}^{q-1}(\log A_i)_{\log} \xrightarrow{\simeq} U_i^0/U_i^1
$$

$$
(w, w') \mapsto w + w' \wedge \text{dlog } T_i.
$$

(4) If $i \in [e+1, g]$ and $p \mid h \ge 1$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \oplus \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \xrightarrow{\simeq} U_i^h / U_i^{h+1},
$$

$$
\Big(\sum_{s \in I_i^{q-1}} a'_{i,s} \omega_{i,s}, \sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \Big) \mapsto \sum_{s \in I_i^{q-1}} \mathrm{dlog}(1 + a'_{i,s} T_i^{h+1}) \wedge \omega_{i,s} + \sum_{s \in I_i^{q-1}} \mathrm{dlog}(1 + a_{i,s} T_i^h) \wedge \omega_{i,s},
$$

where $a'_{i,s}, a_{i,s} \in R_i$. Moreover,

$$
0 \to \frac{\Omega_{R_i}^{q-2}(\log A_i)}{\mathcal{Z}^{q-2}} \xrightarrow{\beta} U_i^h/U_i^{h+1} \xrightarrow{\alpha} \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \oplus \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \to 0
$$

is a short exact sequence, where β is defined by

(3.3.2)
$$
\beta(\sum_t b_{i,t}\omega_{i,t}) = \sum_t \text{dlog}(1 + b_{i,t}T_i^h) \wedge \omega_{i,t} \wedge \text{dlog}(1 + T_i),
$$

and α is defined by the inverse of $\rho_{i,h}$ followed by the projection.

(5) If $i \in [e+1,g]$ and $p \nmid h(h+1)$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \to U_i^h/U_i^{h+1}
$$

$$
\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \mapsto \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s}T_i^{h+1}) \wedge \omega_{i,s}.
$$

where $a_{i,s} \in R_i$. Moreover,

$$
0 \to \frac{\Omega_{R_i}^{q-2}(\log A_i)}{\mathcal{Z}^{q-2}} \xrightarrow{\beta} U_i^h/U_i^{h+1} \xrightarrow{\alpha} \frac{\Omega_{R_i}^{q-1}(\log A_i)}{\mathcal{Z}^{q-1}} \to 0
$$

is a short exact sequence, where β is defined by [\(3.3.2\)](#page-16-0), and α is defined by the inverse of $\rho_{i,h}$ followed by the projection.

(6) If $i \in [e+1,g]$ and $p \mid (h+1)$, then we have the following isomorphism of abelian groups

$$
\rho_{i,h}: \frac{\Omega_{R_i}^{q-2}(\log A_i)}{\mathcal{Z}^{q-2}} \to U_i^h/U_i^{h+1}
$$

$$
\sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} \mapsto \sum_{t \in I_i^{q-2}} d\log(1 + b_{i,t}T_i^h) \wedge \omega_{i,t} \wedge d\log(1 + T_i).
$$

where $b_{i,t} \in R_i$. (In particular, when $q = 1$, we have $U_i^h = U_i^{h+1}$.)

(7) If
$$
i \in [g+1, d]
$$
 (i.e., $h = 0$ *), then we have the following isomorphism of abelian groups*
\n
$$
\rho_{i,h}: \Omega_{R_i}^q(\log A_i)_{\log} \oplus \Omega_{R_i}^{q-1}(\log A_i)_{\log} \xrightarrow{\simeq} U_i^0/U_i^1.
$$
\n
$$
(w, w') \mapsto w + w' \wedge \text{dlog}(1 + T_i).
$$

Proof of Lemma [3.3.](#page-15-0) The first statements in $(1)(2)$ are already contained in the proof of [\[Kat82,](#page-27-10) p. 224, Lemma 4]. We include the proof here for the convenience of the reader.

(a) If $i \in [1, e]$ and $h \geq 1$, consider the map

$$
(3.3.3) \qquad \rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \oplus \Omega_{R_i}^{q-2}(\log A_i) \to U_i^h/U_i^{h+1},
$$
\n
$$
\Big(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}\Big) \mapsto \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s} T_i^h) \wedge \omega_{i,s}
$$
\n
$$
+ \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^h) \wedge \omega_{i,t} \wedge \text{dlog} T_i.
$$

This map is clearly well-defined and is a map of abelian groups. It is also surjective by Lemma $3.2(1)$ $3.2(1)$. In fact, take any general element

(3.3.4)
$$
\sum_{s \in I_i^{q-1}} d \log(1 + a'_{i,s} T_i^h) \wedge w_{i,s} + \sum_{t \in I_i^{q-2}} d \log(1 + b'_{i,t} T_i^h) \wedge w'_{i,t} \wedge d \log T_i
$$

from U_i^h , where $a'_{i,s}, b'_{i,t} \in R_i$, $w_{i,s} \in \Omega_{R_i}^{q-1}$ $_{R_i}^{q-1}(\log A_i)_{\log}, w'_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i)_{\log}$. Suppose $a_{i,s}, b_{i,t} \in R_i$ satisfy

$$
\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s} = \sum_{s \in I_i^{q-1}} a'_{i,s} w_{i,s}, \quad \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} = \sum_{t \in I_i^{q-2}} b'_{i,t} w'_{i,t}.
$$

Then as an element in the graded piece U_i^h / U_i^{h+1} ,

$$
(3.3.4) = \sum_{s \in I_i^{q-1}} T_i^h da'_{i,s} \wedge w_{i,s} + T_i^h \left((-1)^{q-1} \sum_{s \in I_i^{q-1}} h a'_{i,s} w_{i,s} + \sum_{t \in I_i^{q-2}} d b'_{i,t} \wedge w'_{i,t} \right) \wedge \text{dlog } T_i
$$

$$
= \sum_{s \in I_i^{q-1}} T_i^h da_{i,s} \wedge \omega_{i,s} + T_i^h \left((-1)^{q-1} \sum_{s \in I_i^{q-1}} h a_{i,s} \omega_{i,s} + \sum_{t \in I_i^{q-2}} d b_{i,t} \wedge \omega_{i,t} \right) \wedge \text{dlog } T_i
$$

$$
= \rho_{i,h} \Big(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} \Big)
$$

This proves the surjectivity.

An element $(\sum a_{i,s}\omega_{i,s}, \sum b_{i,t}\omega_{i,t})$ lies in the kernel of $(3.3.3)$ if and only if

$$
(3.3.5)\sum_{s\in I_i^{q-1}} T_i^h \cdot da_{i,s} \wedge \omega_{i,s} + T_i^h \cdot \left((-1)^{q-1} \sum_{s\in I_i^{q-1}} ha_{i,s} \omega_{i,s} + \sum_{t\in I_i^{q-2}} db_{i,t} \wedge \omega_{i,t} \right) \wedge \text{dlog} T_i = 0.
$$

If $p \nmid h$, this gives

(3.3.6)
$$
\text{Ker}(3.3.3) = \Big\{ \Big(\frac{(-1)^q}{h} \sum_{t \in I_i^{q-2}} db_{i,t} \omega_{i,t}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} \Big) \mid b_{i,t} \in R_i \Big\},\
$$

which is canonically isomorphic to $\Omega_{R_i}^{q-2}(\log A_i)$. This proves the first statement in (1). If p | h, then [\(3.3.5\)](#page-17-2) implies $d(\sum a_{i,s}\omega_{i,s})=0, d(\sum b_{i,t}\omega_{i,t})=0$, and therefore $\text{Ker (3.3.3)} = \mathcal{Z}^{q-1} \oplus \mathcal{Z}^{q-2}$ $\text{Ker (3.3.3)} = \mathcal{Z}^{q-1} \oplus \mathcal{Z}^{q-2}$ $\text{Ker (3.3.3)} = \mathcal{Z}^{q-1} \oplus \mathcal{Z}^{q-2}$

This proves (2).

For the "moreover" part of (1), the surjectivity of the map α is clear. [\(3.3.6\)](#page-17-3) gives the exactness in the middle. Suppose $\beta(\sum_{t} b_{i,t} \omega_{i,t}) = 0$ in U_i^h / U_i^{h+1} . Note that

$$
\beta(\sum_{t} b_{i,t}\omega_{i,t}) = \sum_{t} \text{dlog}(1 + b_{i,t}T_i^h) \wedge \omega_{i,t} \wedge \text{dlog} T_i \stackrel{(3.3.6)}{=} \sum_{s} \text{dlog}(1 + a_{i,s}T_i^h) \wedge \omega_{i,s}
$$

for some $a_{i,s}$ with $\sum_s a_{i,s} \omega_{i,s} = -\frac{(-1)^q}{h}$ $\frac{(1)^q}{h} \sum_t db_{i,t} \omega_{i,t}$ in $\Omega_{R_i}^{q-1}(\log A_i)$. Since we have proven that $\rho_{i,h}$ is an isomorphism, $\beta(\sum_t b_{i,t} \omega_{i,t}) = 0$ if and only if $\sum_s a_{i,s} \omega_{i,s} = 0$. That is, $\sum_{t} b_{i,t} \omega_{i,t} \in \text{Ker } d$. This proves the remaining part of (1). (b) If $i \in [1, e]$ and $h = 0$, any q-form in U_i^0/U_i^1 is of the form

$$
w + w' \wedge \mathrm{dlog} T_i,
$$

where $w \in \Omega^q$ $_{R_i}^q(\log A_i)_{\log}, w' \in \Omega_{R_i}^{q-1}$ $R_i^{q-1}(\log A_i)_{\log}$. This proves (3).

(c) Assume that $i \in [e+1,g]$ and $p \mid h$. Based on Lemma [3.2\(](#page-7-0)[2\)](#page-7-2), a similar argument as in (a) shows that any q-form in U_i^h/U_i^{h+1} is of the form

$$
\sum_{s \in I_i^{q-1}} \text{dlog}(1 + a'_{i,s} T_i^{h+1}) \wedge \omega_{i,s} + \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s} T_i^h) \wedge \omega_{i,s} + \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^h) \wedge \omega_{i,t} \wedge \text{dlog}(1 + T_i),
$$

where $a'_{i,s}, a_{i,s}, b_{i,t} \in R_i$. Consider the map

$$
(3.3.7)
$$
\n
$$
\rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \oplus \Omega_{R_i}^{q-1}(\log A_i) \oplus \Omega_{R_i}^{q-2}(\log A_i) \to U_i^h/U_i^{h+1}
$$
\n
$$
\left(\sum_{s \in I_i^{q-1}} a'_{i,s}\omega_{i,s}, \sum_{s \in I_i^{q-1}} a_{i,s}\omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t}\omega_{i,t}\right) \mapsto \sum_{s \in I_i^{q-1}} d\log(1 + a'_{i,s}T_i^{h+1}) \wedge \omega_{i,s}
$$
\n
$$
+ \sum_{s \in I_i^{q-1}} d\log(1 + a_{i,s}T_i^h) \wedge \omega_{i,s}
$$
\n
$$
+ \sum_{t \in I_i^{q-2}} d\log(1 + b_{i,t}T_i^h) \wedge \omega_{i,t} \wedge d\log(1 + T_i)
$$

This map is then surjective. An element $(\sum a'_{i,s}\omega_{i,s}, \sum a_{i,s}\omega_{i,s} \sum b_{i,t}\omega_{i,t})$ lies in the kernel of [\(3.3.7\)](#page-18-0) if and only if

$$
\sum_{s \in I_i^{q-1}} T_i^h \cdot da_{i,s} \wedge \omega_{i,s} + T_i^h \cdot \left((-1)^{q-1} \sum_{s \in I_i^{q-1}} (h+1) a'_{i,s} \omega_{i,s} + \sum_{t \in I_i^{q-2}} db_{i,t} \wedge \omega_{i,t} \right) \wedge \mathrm{dlog}(1+T_i) = 0.
$$

Note that in this step, we have used the assumption that $h \geq 2$ and hence $a_{i,s}T_i^h d(a_{i,s}T_i^h) \in$ V_i^{h+1} . As a result,

(3.3.8)

$$
\text{Ker}\,(3.3.7) = \Big\{ \Big(\frac{(-1)^q}{h+1} \sum_{t \in I_i^{q-2}} db_{i,t} \wedge \omega_{i,t}, \sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} \Big) \mid d(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}) = 0 \Big\}
$$

This proves the first half of (4). The "moreover" part of (4) is proven analogously as the "moreover" part of (1).

(d) Assume that $i \in [e+1,g]$ and $p \nmid h \geq p$. Consider the map

$$
(3.3.9) \quad \rho_{i,h}: \Omega_{R_i}^{q-1}(\log A_i) \oplus \Omega_{R_i}^{q-2}(\log A_i) \to U_i^h / U_i^{h+1}
$$
\n
$$
\Big(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}\Big) \mapsto \sum_{s \in I_i^{q-1}} \text{dlog}(1 + a_{i,s} T_i^{h+1}) \wedge \omega_{i,s}
$$
\n
$$
+ \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^h) \wedge \omega_{i,t} \wedge \text{dlog}(1 + T_i)
$$

This map is surjective by a similar argument as in (a) based on Lemma $3.2(3)(4)$ $3.2(3)(4)$ $3.2(3)(4)$. An element $(\sum a_{i,s}\omega_{i,s}, \sum b_{i,t}\omega_{i,t})$ lies in the kernel of $(3.3.9)$ if and only if

$$
(3.3.10) \t T_i^h \cdot \left((-1)^{q-1} \sum_{s \in I_i^{q-1}} (h+1) a_{i,s} \omega_{i,s} + \sum_{t \in I_i^{q-2}} db_{i,t} \wedge \omega_{i,t} \right) \wedge \text{dlog}(1+T_i) = 0.
$$

If $p \nmid (h + 1)$, $(3.3.10)$ is equivalent to

$$
\text{Ker}(3.3.9) = \Big\{ \Big(\frac{(-1)^q}{h+1} \sum_{t \in I_i^{q-2}} db_{i,t} \wedge \omega_{i,t}, \sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t} \Big) \Big\}.
$$

This gives the first half of (5). The "moreover" part of (5) is proven analogously as the "moreover" part of (1). If $p \mid (h+1)$, [\(3.3.10\)](#page-19-1) is equivalent to $d(\sum b_{i,t}\omega_{i,t}) = 0$, and hence

$$
Ker (3.3.9) = \Omega_{R_i}^{q-1}(\log A_i) \oplus \mathcal{Z}^{q-2}
$$

This proves (6).

(e) If $i \in [g+1, d]$, then $h = 0$ by our assumptions. Any q-form in U_i^0/U_i^1 is of the form $(3.3.11)$ $w + w' \wedge \text{dlog}(1 + T_i)$,

where
$$
w \in \Omega_{R_i}^q(\log A)_{\log}, w' \in \Omega_{R_i}^{q-1}(\log A)_{\log}
$$
. This finishes (7).

We hence finished the proof of Lemma [3.3.](#page-15-0) \Box

Lemma 3.4. $Let\ w = \sum_{s\in I_i^{q-1}} a_{i,s} \omega_{i,s} \in \Omega_{R_i}^{q-1}$ $R_i^{q-1}(\log A_i), w' = \sum_{s \in I_i^{q-1}} a'_{i,s} \omega_{i,s} \in \Omega_{R_i}^{q-1}$ $q_{R_i}^{q-1}(\log A_i),$ $w'' = \sum_{s \in I_i^{q-2}} b_{i,t} \omega_{i,t} \in \Omega_{R_i}^{q-2}$ $R_i^{q-2}(\log A_i).$

(1) Suppose $e > 0$ and $i \in [1, e], p \nmid r_i$. If $\rho_{i,r_i}(w) \in G_R^{\underline{r}}$ mod $V_i^{r_i+1}$, then there exist $\widetilde{a}_{i,s},\widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \ldots \widehat{T}_i^{\widetilde{r}_i} \ldots T_d^{\widetilde{r}_d})R_i$

for all $s \in I_i^{q-1}$ $a_i^{q-1}, t \in I_i^{q-2}$ i^{q-2} , such that

$$
\rho_{i,r_i}(w) \equiv \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}_{i,s} T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} \text{dlog}(1 + \widetilde{b}_{i,t} T_i^{r_i}) \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i \mod U_i^{r_i+1}.
$$

(2) Suppose $e > 0$ and $i \in [1, e], p \mid r_i$. If $\rho_{i,r_i}(w, w'') \in G_R^r \mod V_i^{r_i+1}$, then there exist $\widetilde{a}_{i,s},\widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \ldots \widehat{T}_i^{\widetilde{r}_i} \ldots T_d^{\widetilde{r}_d})R_i$

such that

$$
\rho_{i,r_i}(w,w'') \equiv \sum_{s \in I_i^{q-1}} d\log(1 + \widetilde{a}_{i,s}T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} d\log(1 + \widetilde{b}_{i,t}T_i^{r_i}) \wedge \widetilde{\omega}_{i,t} \wedge d\log T_i \mod U_i^{r_i+1}.
$$

(3) Suppose
$$
i \in [e+1, g]
$$
 and $p | r_i \ge 1$. If $\rho_{i,r_i}(w', w) \in G_R^{\underline{r}}$ mod $V_i^{r_i+1}$, then there exist
\n
$$
\widetilde{a}'_{i,s}, \widetilde{a}_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T_i^{\widetilde{r}_i}} \dots T_d^{\widetilde{r}_d}) R_i
$$

for all
$$
s \in I_i^{q-1}
$$
, such that
\n
$$
\rho_{i,r_i}(w', w) \equiv \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \tilde{a}'_{i,s} T_i^{r_i+1}) \wedge \tilde{\omega}_{i,s} + \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \tilde{a}_{i,s} T_i^{r_i}) \wedge \tilde{\omega}_{i,s} \mod U_i^{r_i+1}
$$
\n(4) Suppose $i \in [e+1, g]$, $p \nmid r_i(r_i+1)$. If $\rho_{i,r_i}(w') \in G_R^{\mathcal{L}}$ mod $V_i^{r_i+1}$, then there exist

$$
\widetilde{a}_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) R_i
$$

for all $s \in I_i^{q-1}$ i^{q-1} , such that

$$
\rho_{i,r_i}(w') \equiv \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}'_{i,s} T_i^{r_i+1}) \wedge \widetilde{\omega}_{i,s} \mod U_i^{r_i+1}.
$$

(5) Suppose $i \in [e+1,g], p \nmid r_i, p \mid (r_i+1)$. If $\rho_{i,r_i}(w'') \in G_R^{\underline{r}}$ mod $V_i^{r_i+1}$, then there exist

$$
\widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) R_i
$$

for all $t \in I_i^{q-2}$ i^{q-2} , such that

$$
\rho_{i,r_i}(w'') \equiv \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^{r_i+1}) \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i \mod U_i^{r_i+1}.
$$

Proof of Lemma [3.4.](#page-19-2) If $i \in [1, e]$ and $p \nmid r_i$,

$$
\rho_{i,r_i}(w) \equiv \sum_{s \in I_i^{q-1}} T_i^{r_i} \left(da_{i,s} \wedge \omega_{i,s} + (-1)^{q-1} r_i a_{i,s} \omega_{i,s} \wedge \text{dlog} T_i \right) \mod V_i^{r_i+1}.
$$

By Lemma [3.1,](#page-6-2) the condition $\rho_{i,r_i}(w) \in G_R^{\underline{r}}$ mod $V_i^{r_i+1}$ gives

$$
d(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}) \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^q(\log A_i), \sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s} \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^{q-1}(\log A_i).
$$

By Lemma [4.3,](#page-24-3) the first relation implies that there exist

$$
\widetilde{a}_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T_i^{\widetilde{r}_i}} \dots T_d^{\widetilde{r}_d}) R_i
$$

for all s, such that $d(\sum_s a_{i,s}\omega_{i,s}) = d(\sum_s \tilde{a}_{i,s}\tilde{\omega}_{i,s})$. The short exact sequence in Lemma [3.3\(](#page-15-0)1) implies

$$
\rho_{i,r_i}(w) - \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}_{i,s}T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} = \sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t}T_i^{r_i}) \wedge \omega_{i,s} \wedge \text{dlog} T_i
$$

for some $b_{i,t} \in R_i$. Since $\sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}_{i,s}T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} \in G_{\overline{I}}^{\underline{r}}$ $\frac{r}{R}$, we have

$$
\sum_{t \in I_i^{q-2}} \text{dlog}(1 + b_{i,t} T_i^{r_i}) \wedge \omega_{i,t} \wedge \text{dlog} T_i \in G_R^{\underline{r}} \mod V_i^{r_i+1}.
$$

This is equivalent to $\sum_t db_{i,t} \wedge \omega_{i,t} \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^{q-1}(\log A_i)$, which after applying again Lemma [4.3](#page-24-3) gives

$$
\widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) R_i.
$$

This together with $(3.1.1)$ proves (1) .

If $i \in [1, e]$ and $p | r_i \geq 1$, $\rho_{i,r_i}(w,w'') \equiv \sum$ $s \in I_i^{q-1}$ $T_i^{r_i}da_{i,s}\wedge\omega_{i,s}+\sum\limits_{s}$ $t \in I_i^{q-2}$ $T_i^{r_i}db_{i,t} \wedge \omega_{i,t} \wedge \text{dlog } T_i \mod V_i^{r_i+1},$

Lemma [3.1](#page-6-2) implies that

$$
d(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}) \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^q(\log A_i), \quad d(\sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}) \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^{q-1}(\log A_i).
$$

Lemma [4.3](#page-24-3) implies that there exist

$$
\widetilde{a}_{i,s}, \widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \dots \widehat{T_i^{\widetilde{r}_i}} \dots T_d^{\widetilde{r}_d}) R_i
$$
\nsuch that $d(\sum_s a_{i,s} \omega_{i,s}) = d(\sum_s \widetilde{a}_{i,s} \widetilde{\omega}_{i,s}), d(\sum_t b_{i,t} \omega_{i,t}) = d(\sum_t \widetilde{b}_{i,t} \widetilde{\omega}_{i,t}).$ Hence\n
$$
\rho_{i,r_i}(w, w'') \equiv \sum_{s \in I_i^{q-1}} T_i^{r_i} d\widetilde{a}_{i,s} \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} T_i^{r_i} d\widetilde{b}_{i,t} \wedge \widetilde{\omega}_{i,t} \wedge d\log T_i \mod V_i^{r_i+1}
$$
\n
$$
\equiv \sum_{s \in I_i^{q-1}} d\log(1 + \widetilde{a}_{i,s} T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} d\log(1 + \widetilde{b}_{i,t} T_i^{r_i}) \wedge \widetilde{\omega}_{i,t} \wedge d\log T_i \mod V_i^{r_i+1}.
$$

This together with $(3.1.1)$ proves (2) .

If
$$
i \in [e+1, g]
$$
 and $p | r_i$,
\n
$$
\rho_{i,r_i}(w', w) \equiv (-1)^{q-1} \sum_{s \in I_i^{q-1}} (r_i + 1) a'_{i,s} T_i^{r_i} \omega_{i,s} \wedge \text{dlog}(1 + T_i) + \sum_{s \in I_i^{q-1}} T_i^{r_i} d a_{i,s} \wedge \omega_{i,s} \mod V_i^{r_i+1}.
$$

Lemma [3.1](#page-6-2) implies that

$$
\sum_{s \in I_i^{q-1}} a'_{i,s} \omega_{i,s} \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^{q-1}(\log A_i), \quad d(\sum_{s \in I_i^{q-1}} a_{i,s} \omega_{i,s}) \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^q(\log A_i).
$$

By Lemma [4.3,](#page-24-3) the second relation implies $d(\sum_s a_{i,s}\omega_{i,s}) = d(\sum_s \tilde{a}_{i,s}\tilde{\omega}_{i,s})$ for some

$$
\widetilde{a}_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) R_i.
$$

In particular,

$$
\sum_{s\in I_i^{q-1}}\mathrm{dlog}(1+a'_{i,s}T_i^{r_i+1})\wedge\omega_{i,s}\equiv\rho_{i,r_i}(w',w)-\sum_{s\in I_i^{q-1}}\mathrm{dlog}(1+\widetilde{a}_{i,s}T_i^{r_i})\wedge\widetilde{\omega}_{i,s}\in G_R^{\underline{r}}\mod U_i^{r_i+1}.
$$

This yields

$$
d(\sum_{s\in I_i^{q-1}}a'_{i,s}T_i^{r_i+1}\omega_{i,s})\in G_R^{\underline{r}}\mod V_i^{r_i+1}.
$$

Applying Lemma [4.3](#page-24-3) again, there exist

$$
\widetilde{a}'_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T_i^{\widetilde{r}_i}} \dots T_d^{\widetilde{r}_d}) R_i
$$

for all s, such that $d(\sum_s a'_{i,s}T_i^{r_i+1}\omega_{i,s}) = d(\sum_s \tilde{a}'_{i,s}T_i^{r_i+1}\tilde{\omega}_{i,s})$. Altogether,

$$
\rho_{i,r_i}(w',w) \equiv \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}'_{i,s} T_i^{r_i+1}) \wedge \widetilde{\omega}_{i,s} + \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}_{i,s} T_i^{r_i}) \wedge \widetilde{\omega}_{i,s} \mod V_i^{r_i+1}.
$$

This together with $(3.1.1)$ proves (3) .

If $p \nmid r_i(r_i+1), \rho_{i,r_i}(w') \in G_R^{\underline{r}} \mod V_i^{r_i+1}$ implies $d(\sum_{s \in I_i^{q-1}} a'_{i,s} T_i^{r_i+1} \omega_{i,s}) \in G_R^{\underline{r}} \mod V_i^{r_i+1}$ for some $\widetilde{a}'_{i,s} \in (T_1^{\widetilde{r}_1} \dots \widehat{T_i^{\widetilde{r}_i}} \dots T_d^{\widetilde{r}_d})R_i$

$$
22\quad
$$

for all s, according to Lemma [4.3.](#page-24-3) Hence

$$
\rho_{i,r_i}(w') \equiv \sum_{s \in I_i^{q-1}} \text{dlog}(1 + \widetilde{a}'_{i,s} T_i^{r_i+1}) \wedge \widetilde{\omega}_{i,s} \mod V_i^{r_i+1}.
$$

This together with $(3.1.1)$ proves (4) .

If $p \nmid r_i$ and $p \mid (r_i + 1)$,

$$
\rho_{i,r_i}(w'') \equiv \sum_{t \in I_i^{q-2}} T_i^{r_i} db_{i,t} \wedge \omega_{i,t} \wedge \text{dlog}(1+T_i) \mod V_i^{r_i+1},
$$

Lemma [3.1](#page-6-2) gives

$$
d(\sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}) \in (T_1^{r_1} \dots \hat{T}_i^{r_i} \dots T_d^{r_d}) \Omega_{R_i}^{q-1}(\log A_i).
$$

Lemma [4.3](#page-24-3) implies that $d(\sum_{t \in I_i^{q-2}} b_{i,t} \omega_{i,t}) = d(\sum_{t \in I_i^{q-2}} \tilde{b}_{i,t} \tilde{\omega}_{i,t})$ for some $\widetilde{b}_{i,t} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) R_i.$

Moreover,

$$
\rho_{i,r_i}(w'') \equiv \sum_{t \in I_i^{q-2}} T_i^{r_i} d(b_{i,t} \wedge \omega_{i,t}) \wedge \text{dlog}(1 + T_i) \mod V_i^{r_i+1}
$$

$$
\equiv \sum_{t \in I_i^{q-2}} T_i^{r_i+1} d(\widetilde{b}_{i,t} \wedge \widetilde{\omega}_{i,t}) \wedge \text{dlog} T_i \mod V_i^{r_i+1}
$$

$$
\equiv \sum_{t \in I_i^{q-2}} \text{dlog}(1 + \widetilde{b}_{i,t} T_i^{r_i+1}) \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i \mod V_i^{r_i+1}.
$$

This together with $(3.1.1)$ proves (5) .

We continue the proof of Proposition [2.3.](#page-3-0) If $e > 0$, pick an $i \in [1, e]$. Take $w \in G_{R, \log}^{\mathcal{L}}$. Then $w \in G_{R,\log}^{\mathcal{L}} \subset U_i^{r_i}$ and hence we can write w as an infinite sum

$$
w = w_{i,r_i} + w_{i,r_i+1} + w_{i,r_i+2} + \dots
$$

where the first term $w_{i,r_i} \in U_i^{r_i} \cap G_{R,\log}^{\mathcal{L}}$ is any chosen lift of the residue class of w in $U_i^{r_i}/U_i^{r_i+1}$, and for each $l \geq r_i + 1$, $w_{i,l} \in U_i^l \cap G_{R,\log}^{\underline{\mathcal{P}}}$ is any chosen lift of the residue class of $w - w_{i,r_i}$ $\cdots - w_{i,l-1}$ in U_i^l / U_i^{l+1} . By Lemma [3.4](#page-19-2)[\(1\)](#page-19-3)(2) and Lemma [3.3\(](#page-15-0)1)(2), there exists

$$
a_{i,s,l} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) \cdot R_i, \quad \text{for every } s \in I_i^{q-1} \text{ and every integer } l \ge r_i
$$

and

 $b_{i,t,pm} \in (T_1^{\tilde{r}_1} \dots \hat{T}_i^{\tilde{r}_i} \dots T_d^{\tilde{r}_d}) \cdot R_i$, for every $t \in I_i^{q-2}$ i^{q-2} and every integer $m \geq r_i/p$, such that

$$
v_{i,l} = \sum_{s \in I_i^{q-1}} a_{i,s,l} \widetilde{\omega}_{i,s}, \quad \text{when } p \nmid l,
$$

$$
v_{i,l} = \left(\sum_{s \in I_i^{q-1}} a_{i,s,l} \widetilde{\omega}_{i,s}, \sum_{t \in I_i^{q-2}} b_{i,t,pm} \widetilde{\omega}_{i,t} \right), \quad \text{when } l = pm \text{ for some integer } m,
$$

satisfy

$$
w_{i,l} = \rho_{i,l}(v_{i,l}) \quad \text{for each } l \ge r_i
$$

.

$$
\Box
$$

Altogether,

$$
w = \sum_{s \in I_i^{q-1}} \sum_{l \ge r_i} \text{dlog}(1 + a_{i,s,l} T_i^l) \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} \sum_{\{m \mid pm \ge r_i\}} \text{dlog}(1 + b_{i,t,pm} T_i^{pm}) \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i
$$

=
$$
\sum_{s \in I_i^{q-1}} \text{dlog} \, x \wedge \widetilde{\omega}_{i,s} + \sum_{t \in I_i^{q-2}} \text{dlog} \, y \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i
$$

with

$$
x = \prod_{l \ge r_i} (1 + a_{i,s,l} T_i^l), \quad y = \prod_{\{m | pm \ge r_i\}} (1 + b_{i,t,pm} T_i^{pm}).
$$

(When $q = 1$, we have $y = 1$ as a convention.) Then both x and y converge in R, and more precisely,

$$
x \in 1 + (T_1^{\widetilde{r}_1} \dots T_d^{\widetilde{r}_d}), \quad y \in 1 + (T_1^{\widetilde{r}_1} \dots T_i^{l_0} \dots T_d^{\widetilde{r}_d}).
$$

where l_0 is the minimum integer $l \geq r_i$ such that $p \mid l$. This completes the proof of Proposition [2.3](#page-3-0) in this case.

If $e = 0$, we choose an $i \in [1, g]$. Set

$$
l_0 = \begin{cases} r_i, & \text{if } p \mid r_i; \\ r_i + 1, & \text{if } p \nmid r_i. \end{cases}
$$

By Lemma $3.4(3)(4)(5)$ $3.4(3)(4)(5)$ and Lemma $3.3(4)(5)(6)$ $3.3(4)(5)(6)$, there exist

$$
a_{i,s,l} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) \cdot R_i
$$
, for every $s \in I_i^{q-1}$ and every $l \geq r_i$,

$$
b_{i,t,l} \in (T_1^{\widetilde{r}_1} \dots \widehat{T}_i^{\widetilde{r}_i} \dots T_d^{\widetilde{r}_d}) \cdot R_i, \quad \text{for every } t \in I_i^{q-2} \text{ and every } p \mid (l+1), \ l \geq r_i,
$$

such that

$$
w = \sum_{s \in I_i^{q-1}} \sum_{l \ge l_0} \text{dlog}(1 + a_{i,s,l}T_i^{l+1}) \wedge \omega_{i,s} + \sum_{t \in I_i^{q-2}} \sum_{\substack{l \ge r_i \\ p \mid (l+1)}} \text{dlog}(1 + b_{i,t,l}T_i^{l+1}) \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i
$$

=
$$
\sum_{s \in I_i^{q-1}} \text{dlog} x \wedge \omega_{i,s} + \sum_{t \in I_i^{q-2}} \text{dlog} y \wedge \widetilde{\omega}_{i,t} \wedge \text{dlog} T_i
$$

with

$$
x = \prod_{l \ge l_0} (1 + a_{i,s,l} T_i^{l+1}), \quad y = \prod_{\substack{l \ge r_i \\ p | (l+1)}} (1 + b_{i,t,l} T_i^{l+1}).
$$

This finishes the proof of Proposition [2.3.](#page-3-0)

4. Appendix

4.1. A Hartog's type lemma for Cohen-Macaulay modules.

Lemma 4.1. Let X be a locally noetherian equidimensional scheme, and F be a Cohen-Macaulay \mathcal{O}_X -module (see [\[Har66,](#page-27-12) Remark on p.238]), and U be an open dense subset of X. Then for any section $\alpha \in \Gamma(U, F)$ such that $\alpha \in F_x$ for all $x \in X^{(1)}$, we have $\alpha \in \Gamma(X, F)$.

Proof. If I^{\bullet} is an injective resolution of F , then

$$
0 \to \underline{\Gamma_{\overline{\{x\}}}}(I^{\bullet}) \to I^{\bullet} \to j_* I^{\bullet} \to 0,
$$

is a short exact of complexes, where $j : X \setminus \{x\} \hookrightarrow X$ is the open immersion. The long exact sequence it induces gives a four-term exact sequence of sheaves

$$
0 \to \underline{\Gamma_{\overline{\{x\}}}(F)} \to F \to j_*F \stackrel{\delta}{\to} \underline{H^1_{\overline{\{x\}}}(F)} \to 0.
$$

Taking stalks at x , we get a canonical isomorphism

(4.1.1)
$$
H_x^1(F) \simeq \bigoplus_{\substack{\eta \in X^{(0)} \\ x \in \{\eta\}}} F_\eta/F_x
$$

for every $x \in X^{(1)}$. Since F is Cohen-Macauley, the Cousin complex

$$
0 \to \bigoplus_{\eta \in X^{(0)}} i_{\eta *} F_{\eta} \to \bigoplus_{x \in X^{(1)}} i_{x *} H_x^1(F) \to \dots
$$

is a flasque resolution of the Zariski sheaf F . Bringing $(4.1.1)$ into this sequence, we deduce that

$$
0 \to F \to \bigoplus_{\eta \in X^{(0)}} i_{\eta *} F_{\eta} \to \bigoplus_{x \in X^{(1)}} i_{x *} \Big(\bigoplus_{\substack{\eta \in X^{(0)} \\ x \in \{\eta\}}} F_{\eta}/F_x \Big)
$$

is exact. Applying the functor $\Gamma(X, -)$, we get a left exact sequence

$$
0 \to \Gamma(X, F) \to \bigoplus_{\eta \in X^{(0)}} \Gamma(X_{\eta}, F_{\eta}) \to \bigoplus_{x \in X^{(1)}} \Big(\bigoplus_{\substack{\eta \in X^{(0)} \\ x \in \overline{\{\eta\}}} } F_{\eta}/F_x \Big).
$$

Hence if $\alpha \in F_x$ for all $x \in X^{(1)}$, then the left exactness of the above sequence gives $\alpha \in$ $\Gamma(X,F)$.

Example 4.2. Let k be a perfect field of positive characteristic. The followings are examples of Cohen-Macaulay sheaves:

- (1) Locally free sheaves of finite rank on a smooth k-scheme are Cohen-Macaulay.([\[Har66,](#page-27-12) p.239, Example])
- (2) For any $n \geq 1$ and $q \geq 0$, $W_n \Omega_X^q$ is a Cohen-Macaulay $W_n \mathcal{O}_X$ -module when X is smooth over k by [\[Ill79,](#page-27-13) I. Corollaire 3.9]. It is not a locally free $W_n \mathcal{O}_X$ -module.
- (3) More generally, $W_n \Omega^q$ $\mathcal{L}_{(X,\pm D)}^{q}$ as defined in [\[RR24\]](#page-27-0) is a Cohen-Macaulay $W_n\mathcal{O}_X$ -module when X is separated smooth of finite type over k . This follows from [\[RR24,](#page-27-0) Theorem 6.4] and the injectivity of p for the pair (X, D) , and from the duality statement Theorem 9.3 in loc. cit. for the pair $(X, -D)$.

4.2. A generalization of [\[RR24,](#page-27-0) Lemma 8.10].

Lemma 4.3. Let X be a smooth k-scheme with étale coordinates $T_1, \ldots, T_d \in \Gamma(X, \mathcal{O}_X)$. Set

$$
\widetilde{A} = \text{Div}(T_1 \dots T_e T_{e+1} \dots T_f), \quad \widetilde{B} = \text{Div}(T_1^{\widetilde{r}_1} \dots T_d^{\widetilde{r}_d}),
$$

where $\underline{r}' = (\widetilde{r}_1, \ldots, \widetilde{r}_d)$ is defined by

(4.3.1) $\widetilde{r}_i = r_i$ for all $j \notin [e + 1, f]$, and $\widetilde{r}_i = r_i + 1$ for all $j \in [e + 1, f]$.

Then $dF_{X*}\left(\Omega_X^{q-1}\right)$ $\left(\frac{q-1}{X}(\log \widetilde{A})(-\widetilde{B})\right)$ is a locally free sheaf of finite rank, and we have an identification of Nisnevich sheaves

$$
\left(\Omega_X^q(\log A)(-B)\right)\cap d\Omega_X^{q-1}(\log A)=d\left(\Omega_X^{q-1}(\log \widetilde{A})(-\widetilde{B})\right)
$$

on X_{Nis} .

Proof of Lemma [4.3.](#page-24-3) This claim is a generalization of $[RR24, Lemma 8.10]$. The direction "⊃" is clear. Now we show "⊂". Since X is smooth, the sheaf $dF_{X*}(\Omega_R^{q-1}(\log \tilde{A})(-\tilde{B}))$ is locally free. In fact, define \mathcal{Z}^{q-1} by the following exact sequence

$$
0 \to \mathcal{Z}^{q-1} \to \Omega_X^{q-1}(\log \widetilde{A})(-\widetilde{B}) \stackrel{d}{\to} d\Omega_X^{q-1}(\log \widetilde{A})(-\widetilde{B}) \to 0
$$

and consider this sequence together with

$$
0 \to d\Omega_X^{q-1}(\log A)(-\widetilde{B}) \to \mathcal{Z}^q \xrightarrow{C} \Omega_X^q(\log \widetilde{A})(-\widetilde{B}) \to 0.
$$

Then a decreasing induction argument on q gives the local freeness. By Lemma [4.1,](#page-23-2) it suffices to show the inclusion for the Nisnevich stalk at any codimension 1 point x . The statement is only nontrivial when x is the generic point of the reduced divisors cutting out by T_i for $j \in [1, g]$. If $j \in [1, e]$, or $j \in [f + 1, g]$, the lemma follows from the part (a) and part (b) of the proof of [\[RR24,](#page-27-0) Lemma 10.8]. Now we suppose x is the generic point of $Div(T_i)$ for some $j \in [e+1, f]$. Write $T := T_j$ and $r = r_j$. By replacing X with a Nisnevich neighborhood Spec R such that $R_0 := R/T$ is isomorphic to a subring of R ([\[BRS22,](#page-27-14) Lemma 7.14]), we can assume that X is of the form $Spec R_0[T]$. Hence as free R_0 -modules,

$$
\Omega_X^{q-1}(\log A) = \Omega_{R_0[T]}^{q-1} = \bigoplus_{i \ge 0} \Omega_{R_0}^{q-1} \cdot T^i \oplus \bigoplus_{i \ge 1} \Omega_{R_0}^{q-2} \cdot T^i \operatorname{dlog} T,
$$

$$
\Omega_X^{q-1}(\log \widetilde{A}) = \Omega_{R_0[T]}^{q-1}(\log T) = \bigoplus_{i \ge 0} \Omega_{R_0}^{q-1} \cdot T^i \oplus \bigoplus_{i \ge 0} \Omega_{R_0}^{q-2} \cdot T^i \operatorname{dlog} T.
$$

Now suppose $\alpha \in \Omega_X^{q-1}$ $\chi^{q-1}(\log A)$. Then there exists $a_i \in \Omega_{R_0}^{q-1}$ $\frac{q-1}{R_0}, b_i \in \Omega_{R_0}^{q-2}$ $\frac{q-2}{R_0}$, among which only finitely many are nonzero, such that

$$
\alpha = a_0 + \sum_{i \ge 1} (a_i T^i + b_i T^i \operatorname{dlog} T).
$$

Hence

$$
d\alpha = \sum_{i \ge 0} T^i da_i + \sum_{i \ge 1} ((-1)^{q-1} i a_i + db_i) \cdot T^i \operatorname{dlog} T.
$$

The condition $d\alpha \in T^r \Omega_{R_0[T]}^q = \bigoplus_{i \geq r} \Omega_{I}^q$ $R_0^q \cdot T^i \oplus \bigoplus_{i \geq r+1} \Omega_{R_0}^{q-1}$ $_{R_0}^{q-1} \cdot T^i \operatorname{dlog} T$ gives

$$
da_i = 0
$$
 for all $i \in [0, r-1]$, and $(-1)^{q-1}ia_i + db_i = 0$ for all $i \in [1, r]$.

In particular, the second equation above implies $db_i = 0$ when $p | i \in [1, r]$; and $da_r = 0$ (since $p \nmid r$ by our assumption). Let

$$
\epsilon := a_0 + \sum_{p|i \in [1,r]} a_i T^i + \sum_{i \in [1,r]} b_i T^i \operatorname{dlog} T - \sum_{p|i \in [1,r]} \frac{(-1)^{q-1}}{i} T^i db_i.
$$

It is direct to check that

$$
d\epsilon = da_0 + \sum_{p|i \in [1,r]} T^i (da_i + db_i dlog T) = 0.
$$

Moreover,

$$
\alpha - \epsilon = \sum_{i \ge r+1} (a_i T^i + b_i T^i \operatorname{dlog} T) \in T^{r+1} \cdot \Omega_X^{q-1}(\log \widetilde{A}).
$$

 \Box

4.3. Log forms as the Frobenius invariant in $W_n \Omega_1^q$ $\int_X^q (\log A)/B_{\infty}$. The following lemma originates from a lemma by Kay Rülling. Below we formulate and prove a log version. We only need the $n = 1$ version of it, but nevertheless we record a proof for general n.

Lemma 4.4. Let X be a connected smooth k -scheme and A be a simple normal crossing divisor on X. Let $j: U := X \setminus A \hookrightarrow X$ be the open immersion. Let $W_n \Omega_X^* (\log A)$ be the de Rham-Witt complex associated to the log structure on X associated to A (relative to the trivial log structure on the base k) defined by Hyodo-Kato. More explicitly (see $[HK94,$ Proposition 4.6]),

$$
W_n \Omega_X^q(\log A) = W_n \mathcal{O}_X \cdot j_* \operatorname{dlog} K_{q,U}^M + dW_n \mathcal{O}_X \cdot j_* \operatorname{dlog} K_{q-1,U}^M \subset j_* W_n \Omega_U^q,
$$

where $K_{q,U}^M$ is the étale sheaf of Milnor K-theory on U. Let

$$
W_n \Omega_X^q (\log A)_{\log} := W_n \Omega_X^q (\log A) \cap j_* W_n \Omega_{U, \log}^q.
$$

Then the sequence

$$
0 \to W_n \Omega_X^q(\log A)_{\log} \to W_n \Omega_X^q(\log A) / \mathcal{B}_{\infty} \xrightarrow{\overline{F}-1} W_n \Omega_X^q(\log A) / \mathcal{B}_{\infty} \to 0
$$

is exact on $X_{\text{\'et}}$. Here $\mathcal{B}_{\infty} := \bigcup_{j \geq 0} F^j(dW_{n+j}\Omega_U^{q-1})$ $\binom{q-1}{U}(\log A)$, and F is induced from F via the restriction map.

Proof. We first show that the map \overline{F} is well-defined. By [\[HK94,](#page-27-15) Theorem 4.4],

$$
\operatorname{Ker}(R:W_{n+1}\Omega_X^q(\log A)\to W_n\Omega_X^q(\log A))=V^n(\Omega_X^q(\log A))+dV^n(\Omega_X^{q-1}(\log A)).
$$

Hence $F: W_{n+1}\Omega_U^q \to W_n\Omega_U^q$ U induces a well-defined map

$$
W_n \Omega_X^q(\log A) \to W_n \Omega_X^q(\log A) / dV^{n-1} \Omega_X^{q-1}(\log A).
$$

Since $dV^{n-1}\Omega_X^q(\log A) \subset dW_n\Omega_X^q(\log A) \subset \mathcal{B}_{\infty}$, the map F further induces a well-defined map

 $\overline{F}:W_n\Omega^q_{\overline{X}}$ $X^q \left(\log A \right) / \mathcal{B}_{\infty} \to W_n \Omega_{\mathcal{X}}^q$ $X^q \left(\log A \right) / \mathcal{B}_{\infty}.$

Clearly the sequence is a complex. When $A = \emptyset$, the surjectivity of $\overline{F} - 1$ follows from [\[Ill79,](#page-27-13) I, Proposition 3.26]. In general, we can run a similar (but way easier) argument as in the proof in loc. cit.. For any $\alpha \in W_n \mathcal{O}_X \cdot j_* \text{dlog } K_{q-1,U}^M$, we have $\sum_{i=1}^n dV^i(R^{i-1}\alpha) \in W_{n+1}\Omega_X^q(\log A)$ and it satisfies

$$
d\alpha = (\overline{F} - 1)(R\Big(\sum_{i=1}^n dV^i(R^{i-1}\alpha)\Big)).
$$

It remains to show $W_n \mathcal{O}_X \cdot j_* \text{dlog } K_{q,U}^M \subset (\overline{F} - 1)(W_n \Omega_X^q(\log A))$ étale locally. Since $\overline{F} - 1$: $W_n\mathcal{O}_X \to W_n\mathcal{O}_X$ is surjective on $X_{\text{\'et}}$ and log forms are invariant under the endomorphism \overline{F} , it follows that $W_n \mathcal{O}_X \cdot j_* \text{dlog } K_{q,U}^M \subset (\overline{F} - 1)(W_n \mathcal{O}_X \cdot j_* \text{dlog } K_{q,U}^M)$, and in particular, it lies in $(\overline{F} - 1)(W_n \Omega_{\mathcal{I}}^q)$ $_X^q(\log A))$ as well.

We continue to prove the injectivity. Let $\alpha \in W_n \Omega_2^q$ $\frac{q}{X}(\log A)_{\log}$ be zero in $W_n \Omega^q$ $X^q \left(\log A \right) / \mathcal{B}_{\infty}.$ Then by the definition of \mathcal{B}_{∞} , $\alpha = F^m d\beta$ for some $\beta \in W_{n+m} \Omega_X^{q-1}$ $X^{q-1}(\log A)$. It follows that $V^{n+m}(\alpha) = 0$. Let $\widetilde{\alpha} \in W_{2n+m}\Omega_1^q$ $X^q_X(\log A)_{\log}$ be a lift of α . Then $0 = V^{n+m}(\alpha) =$ $V^{n+m}(F^{n+m}(\tilde{\alpha})) = p^{n+m}(\alpha)$, where in the second equality we have used that $\tilde{\alpha}$ is a log form and hence $F(\tilde{\alpha}) = R(\tilde{\alpha})$. By the injectivity of p, we deduce that $\alpha = 0$ as desired.

Next we prove the exactness in the middle. Let $\alpha \in W_n \Omega_2^q$ X^q (log A) and assume $F(\alpha) - \alpha =$ $F^m d\beta$ for some $\beta \in W_{n+m} \Omega_X^{q-1}$ $\chi^{q-1}(\log A)$. Let $\gamma = \sum_{i=1}^n dV^i(R^{i-1}(\beta)) \in W_{n+m+1}\Omega_j^q$ X^q (log A). Then $d\beta = (\overline{F} - 1)(R\gamma)$, and

$$
\overline{F}(\alpha - F^m R(\gamma)) - (\alpha - F^m R(\gamma)) = F^m d\beta - \overline{F}(F^m R \gamma) + F^m R(\gamma) = 0.
$$

By [\[CTSS83,](#page-27-7) Lemme 2], $\alpha - F^m R(\gamma)$ is a log form.

REFERENCES

- [BRS22] Federico Binda, Kay Rülling, and Shuji Saito. On the cohomology of reciprocity sheaves. Forum Math. Sigma, 10:Paper No. e72, 111, 2022.
- [CTSS83] Jean-Louis Colliot-Thélène, Jean-Jacques Sansuc, and Christophe Soulé. Torsion dans le groupe de Chow de codimension deux. Duke Math. J., 50(3):763–801, 1983.
- [GK21] Rahul Gupta and Amalendu Krishna. Ramified class field theory and duality over finite fields. Preprint, <https://arxiv.org/abs/2104.03029>, 2021.
- [Har66] Robin Hartshorne. Residues and duality, volume No. 20 of Lecture Notes in Mathematics. Springer-Verlag, Berlin-New York, 1966. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64, With an appendix by P. Deligne.
- [HK94] Osamu Hyodo and Kazuya Kato. Semi-stable reduction and crystalline cohomology with logarithmic poles. $Ast\acute{e}risque$, (223):221–268, 1994. Périodes p-adiques (Bures-sur-Yvette, 1988).
- [Ill79] Luc Illusie. Complexe de de Rham-Witt et cohomologie cristalline. Annales Scientifiques de l'Ecole ´ Normale Supérieure. Quatrième Série, 12(4):501–661, 1979.
- [JSZ18] Uwe Jannsen, Shuji Saito, and Yigeng Zhao. Duality for relative logarithmic de Rham–Witt sheaves and wildly ramified class field theory over finite fields. Compos. Math., 154(6):1306–1331, 2018.
- [Kat82] Kazuya Kato. Galois cohomology of complete discrete valuation fields. In Algebraic K-theory, Part II (Oberwolfach, 1980), volume 967 of Lecture Notes in Math., pages 215–238. Springer, Berlin-New York, 1982.
- [KMSY21a] Bruno Kahn, Hiroyasu Miyazaki, Shuji Saito, and Takao Yamazaki. Motives with modulus, I: Modulus sheaves with transfers for non-proper modulus pairs. Epijournal Géom. Algébrique, 5:Art. 1, 46, 2021.
- [KMSY21b] Bruno Kahn, Hiroyasu Miyazaki, Shuji Saito, and Takao Yamazaki. Motives with modulus, II: Modulus sheaves with transfers for proper modulus pairs. Epijournal Géom. Algébrique, 5:Art. 2, 31, 2021.
- [KMSY22] Bruno Kahn, Hiroyasu Miyazaki, Shuji Saito, and Takao Yamazaki. Motives with modulus, III: The categories of motives. Ann. K-Theory, 7(1):119–178, 2022.
- [Mor19] Matthew Morrow. K-theory and logarithmic Hodge-Witt sheaves of formal schemes in characteristic p. Ann. Sci. Éc. Norm. Supér. (4) , 52 (6) :1537-1601, 2019.
- [RR24] Fei Ren and Kay R¨ulling. Duality for hodge-witt cohomology with modulus, 2024. <https://arxiv.org/abs/2403.18763v1>.
- [RS18] Kay Rülling and Shuji Saito. Higher Chow groups with modulus and relative Milnor K-theory. Trans. Amer. Math. Soc., 370(2):987–1043, 2018.
- [Rül07] Kay Rülling. The generalized de Rham-Witt complex over a field is a complex of zero-cycles. J. Algebraic Geom., 16(1):109–169, 2007.

BERGISCHE UNIVERSITÄT WUPPERTAL, GAUSSSTR. 20, D-42119 WUPPERTAL, GERMANY Email address: renfei@uni-wuppertal.de