

Doubly Robust Inference on Causal Derivative Effects for Continuous Treatments

Yikun Zhang* and Yen-Chi Chen†

Department of Statistics, University of Washington

*yikun@uw.edu †yenchic@uw.edu

January 14, 2025

Abstract

Statistical methods for causal inference with continuous treatments mainly focus on estimating the mean potential outcome function, commonly known as the dose-response curve. However, it is often not the dose-response curve but its derivative function that signals the treatment effect. In this paper, we investigate nonparametric inference on the derivative of the dose-response curve with and without the positivity condition. Under the positivity and other regularity conditions, we propose a doubly robust (DR) inference method for estimating the derivative of the dose-response curve using kernel smoothing. When the positivity condition is violated, we demonstrate the inconsistency of conventional inverse probability weighting (IPW) and DR estimators, and introduce novel bias-corrected IPW and DR estimators. In all settings, our DR estimator achieves asymptotic normality at the standard nonparametric rate of convergence. Additionally, our approach reveals an interesting connection to nonparametric support and level set estimation problems. Finally, we demonstrate the applicability of our proposed estimators through simulations and a case study of evaluating a job training program.

Keywords: Causal inference; dose-response curve; derivative estimation; positivity; kernel smoothing.

1 Introduction

This paper investigates the construction of a doubly robust estimator for the derivative of the continuous treatment effect using kernel smoothing. The analysis considers scenarios both with and without the positivity condition. Specifically, positivity (Assumption A2) requires that every individual has a nonzero probability of being exposed to any treatment level $T = t$ across all possible values of the covariate vector $\mathbf{S} \in \mathcal{S} \subset \mathbb{R}^d$. Let $Y(t)$ be the potential outcome (Rubin, 1974) that would have been observed under treatment level $T = t$. The focus of this work is the

(causal) derivative effect curve $t \mapsto \theta(t) := \frac{d}{dt} \mathbb{E}[Y(t)]$, where $t \mapsto m(t) := \mathbb{E}[Y(t)]$ represents the (causal) dose-response curve.

Valid inference on $\theta(t)$ is essential for understanding how the outcome of interest $Y \in \mathcal{Y}$ changes with treatment t , offering insights beyond the expected value $\mathbb{E}[Y(t)] = m(t)$ of the potential outcome across the population. In some sense, the derivative effect curve $\theta(t)$ can be viewed as continuous-treatment counterpart to the average treatment effect $\mathbb{E}[Y(1)] - \mathbb{E}[Y(0)]$. Despite the importance of estimating $\theta(t)$, the current research for continuous treatments has largely focused on inferring $m(t)$ (Díaz and van der Laan, 2013; Kennedy et al., 2017; Bonvini and Kennedy, 2022; Takatsu and Westling, 2024), with few effectors directed at $\theta(t)$. The only exceptions are Colangelo and Lee (2020) and Bong and Lee (2023), who approximated $\theta(t)$ by the finite difference of the estimated dose-response curve or a closely related matching method.

To achieve precise inference on $\theta(t)$ without numerical approximation, a straightforward approach is to impose structural assumptions on the conditional mean outcome function $\mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s})$ or directly on the dose-response curve $m(t)$, known as the marginal structural modeling (Robins et al., 2000; Neugebauer and van der Laan, 2007). Although this approach can easily construct an estimator of $\theta(t)$ via a standard differentiation on the estimated dose-response curve, those structural assumptions are difficult to verify in practice. Alternatively, existing methods for derivative estimation (Gasser and Müller, 1984; Mack and Müller, 1989; Zhou and Wolfe, 2000), combined with the inverse probability weighting (IPW) technique (Hirano and Imbens, 2004; Imai and van Dyk, 2004), can define an estimator of $\theta(t)$; see, *e.g.*, our proposed IPW estimator in Section 3. Yet, this approach requires correct specification of the conditional density model of T given \mathbf{S} . The sensitivity of these approaches to model misspecification motivates us to propose a doubly robust (DR) inference procedures for $\theta(t)$, accommodating misspecification in either the outcome regression or the conditional density models (Robins, 1986; van der Laan and Robins, 2003; Bang and Robins, 2005).

The existing inference methods for $m(t)$ and the above discussion on $\theta(t)$ relies on the positivity condition (Assumption A2), which may be violated in observational studies with continuous treatments (Cole and Hernán, 2008; Westreich and Cole, 2010). When positivity fails, the identifications of both $m(t)$ and $\theta(t)$ become infeasible without structural assumptions; see Section 4 for details. Zhang et al. (2024) address this problem without positivity by imposing an assumption on the potential outcome model that can be satisfied by additive confounding models and proposing a regression adjustment (RA) estimator of $\theta(t)$. We extend their identification and estimation strategies to propose IPW and DR estimators of $\theta(t)$ under additive confounding models. This extension not only advances the field but also reveals novel connections between the derivative effect curve inference and classical support estimation problems (Cuevas and Fraiman, 1997; Cuevas, 2009).

1.1 Contributions and Outline of the Paper

1. Identification and Estimation: Under the positivity and other regularity conditions that are stated in Section 2, we propose our IPW and DR estimators of $\theta(t)$ using kernel smoothing in

Section 3. In particular, we demonstrate that naively combining the RA estimator with the IPW one only leads to a singly robust estimator. Our proposal leverages a local polynomial approximation to the outcome variable and is robust to the misspecification of either the outcome regression or the conditional density models.

2. Challenges and Remedies Under Violations of Positivity: When the positivity condition is violated, we demonstrate that the conventional approaches exhibit two types of bias due to lack of identification and support discrepancy in [Section 4](#). To resolve these issues, we adopt the additive structural assumption to maintain identification and utilize nonparametric set estimation techniques to develop our bias-corrected IPW and DR estimators of $\theta(t)$ in [Section 5.1](#).

3. Asymptotic Theory: We establish the consistency and asymptotic properties of RA, IPW, and DR estimators of $\theta(t)$ when the nuisance functions are nonparametrically estimated under cross-fitting; see [Section 3.1](#) with positivity and [Section 5.2](#) without positivity. Specifically, our proposed DR estimators are asymptotically normal and can be used to conduct valid inference on $\theta(t)$.

4. Numerical Experiments: We showcase the finite-sample performances of our proposed estimators of $\theta(t)$ with and without the positivity condition through simulations and a case study of the Job Corps program in the United States in [Section 6](#) and [Section B](#). All the codes for our experiments are available at <https://github.com/zhangyk8/npDRDeriv>, and we provide some practical considerations for implementing our proposed estimators in [Section A](#).

1.2 Other Related Works

The dose-response curve $m(t)$ and its derivative $\theta(t)$ are non-regular target parameters, as they lack unique Gâteaux derivatives and Riesz representers, depending on how the treatment distribution is localized at t ([van der Vaart, 1991](#); [Carone et al., 2019](#); [Ichimura and Newey, 2022](#)). As one of the key ingredients in this paper, kernel-based localization is a common approach in the literature, which has been used to construct IPW or DR estimators of $m(t)$ ([Kallus and Zhou, 2018](#); [Su et al., 2019](#); [Huber et al., 2020](#); [Colangelo and Lee, 2020](#); [Klosin, 2021](#)). An alternative localization method is through the basis approach or series estimator ([Chen et al., 2014](#); [Chen and Liao, 2014](#); [Luedtke and Chung, 2024](#)). Additionally, a general form of the IPW estimator of $m(t)$ was studied by [Galvao and Wang \(2015\)](#). Under the positivity condition, the RA or G-computation ([Robins, 1986](#)) estimators of $\theta(t)$ have been explored by [Gill and Robins \(2001\)](#); [Flores \(2007\)](#); [Lee \(2018\)](#).

Although inference methods for $\theta(t)$ are less developed, related works exist for the estimation of average derivative effects $\mathbb{E}[\theta(T)] = \mathbb{E}\left[\frac{\partial}{\partial t}\mathbb{E}(Y|T, \mathbf{S})\right]$ under some regularity conditions ([Härdle and Stoker, 1989](#); [Powell et al., 1989](#); [Newey and Stoker, 1993](#); [Cattaneo et al., 2010](#); [Hirshberg and Wager, 2020](#); [Hines et al., 2023](#)). These average derivative effects are also connected to incremental treatment effects ([Kennedy, 2019](#)), linking them to our estimand $\theta(t)$ of interest ([Rothenhäusler and Yu, 2019](#)).

Growing interest in relaxing the positivity condition has led to new developments in causal inference. For continuous treatments, [Branson et al. \(2023\)](#) studied a smoothed causal effect with

trimmed conditional densities, while [Schindl et al. \(2024\)](#) examined stochastic interventions via exponentially tilted treatment distributions. Notably, dynamic stochastic interventions with continuous treatments can be robust to the violation of positivity ([Díaz and Hejazi, 2020](#); [Bonvini et al., 2023](#); [McClellan et al., 2024](#)). To our knowledge, no existing works directly consider nonparametric inference on $\theta(t)$ without positivity, and our work takes an initial step to fill in this gap.

1.3 Notations

Throughout this paper, we consider an outcome variable $Y \in \mathcal{Y} \subset \mathbb{R}$, univariate continuous treatment $T \in \mathcal{T} \subset \mathbb{R}$, and a vector of confounding variables or covariates $\mathbf{S} = (S_1, \dots, S_d) \in \mathcal{S} \subset \mathbb{R}^d$ with a fixed dimension d . We write $Y \perp\!\!\!\perp \mathbf{X}$ when the random variables Y, \mathbf{X} are independent. The common distribution and expectation of $\mathbf{U} = (Y, T, \mathbf{S})$ are denoted by \mathbb{P} and \mathbb{E} respectively, whose Lebesgue density is $p(y, t, \mathbf{s}) = p_{Y|T, \mathbf{S}}(y|t, \mathbf{s}) \cdot p_{T|\mathbf{S}}(t|\mathbf{s}) \cdot p_{\mathbf{S}}(\mathbf{s})$. Here, $p_T(t)$ and $p_{\mathbf{S}}(\mathbf{s})$ are the marginal densities of T and \mathbf{S} , respectively, and $p_{T|\mathbf{S}}(t|\mathbf{s}) = \frac{\partial}{\partial t} \mathbb{P}(T \leq t | \mathbf{S} = \mathbf{s})$ is the conditional density of T given covariates $\mathbf{S} = \mathbf{s}$. We also denote the joint density of (T, \mathbf{S}) by $p(t, \mathbf{s}) = p_{T|\mathbf{S}}(t|\mathbf{s}) \cdot p_{\mathbf{S}}(\mathbf{s}) = p_{\mathbf{S}|T}(\mathbf{s}|t) \cdot p_T(t)$ and the support of $p_{\mathbf{S}|T}(\mathbf{s}|t)$ by $\mathcal{S}(t)$ for $t \in \mathcal{T}$. For any real-valued \mathbb{P} -integrable function f , we write $\mathbb{P}f = \int f(\mathbf{u}) d\mathbb{P}(\mathbf{u})$ and denote the $L_p(\mathbb{P})$ -norm of f by $\|f\|_{L_p} := \left(\int |f(\mathbf{u})|^p d\mathbb{P}(\mathbf{u}) \right)^{\frac{1}{p}}$. If \hat{f} is estimated on an independent data sample, then $\|\hat{f}\|_{L_p} := \left(\int |\hat{f}(\mathbf{u})|^p d\mathbb{P}(\mathbf{u}) \right)^{\frac{1}{p}}$. Additionally, we let \mathbb{P}_n denote the empirical measure so that $\mathbb{P}_n f = \frac{1}{n} \sum_{i=1}^n f(\mathbf{U}_i) = \int f(\mathbf{u}) d\mathbb{P}_n(\mathbf{u})$ and $\mathbb{G}_n(f) = \sqrt{n}(\mathbb{P}_n - \mathbb{P})f$. Finally, we use $\mathbb{1}_A$ to denote the indicator function of a set A . The big- O notation $h_n = O(g_n)$ means that $|h_n|$ is upper bounded by a positive constant multiple of $g_n > 0$ when n is sufficiently large. In contrast, $h_n = o(g_n)$ when $\lim_{n \rightarrow \infty} \frac{|h_n|}{g_n} = 0$. For random variables, $o_P(1)$ is short for a sequence of random variables converging to zero in probability, while $O_P(1)$ denotes the sequence that is bounded in probability.

2 Basic Framework

Suppose that the data sample consists of independent and identically distributed (i.i.d.) observations $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$ generated from the following structural equation model

$$Y = \mu(T, \mathbf{S}) + \epsilon, \tag{1}$$

where $\epsilon \in \mathbb{R}$ is an exogenous noise variable with $\epsilon \perp\!\!\!\perp \mathbf{S}$ and $\mathbb{E}(\epsilon) = 0$, $\text{Var}(\epsilon) = \sigma^2 > 0$, $\mathbb{E}|\epsilon^{2+c_1}| < \infty$ for some absolute constant $c_1 \geq 1$. Under model (1), the potential outcome model becomes $Y(t) = \mu(t, \mathbf{S}) + \epsilon$ for any hypothetical treatment value t . Since the main estimands of interest $\theta(t) = \frac{d}{dt} \mathbb{E}[Y(t)]$ and $m(t) = \mathbb{E}[Y(t)]$ are defined by potential outcomes that are not directly observable, we introduce some basic identification conditions for identifying $m(t)$ and $\theta(t)$ with observed data.

Assumption A1 (Basic identification conditions).

- (a) (Consistency) $T = t$ implies that $Y(t) = Y$ for any $t \in \mathcal{T}$.
- (b) (Ignorability or unconfoundedness) $Y(t) \perp\!\!\!\perp T \mid \mathbf{S}$ for all $t \in \mathcal{T}$.
- (c) (Treatment variation) The conditional variance of T given $\mathbf{S} = \mathbf{s}$ is strictly positive for all $\mathbf{s} \in \mathcal{S}$, i.e., $\text{Var}(T \mid \mathbf{S} = \mathbf{s}) > 0$.
- (d) (Interchangeability) The equality $\frac{d}{dt} \mathbb{E}[\mu(t, \mathbf{S})] = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right]$ holds true under model (1).

Assumption A1(a,b) are standard identification conditions for causal dose-response curves (Gill and Robins, 2001; Kennedy et al., 2017), while Example 1 in Zhang et al. (2024) demonstrates the necessity of imposing Assumption A1(c) for identifiability. In particular, Assumption A1(c) ensures that the distribution of (T, \mathbf{S}) has a nontrivial support in $\mathcal{T} \times \mathcal{S}$. Finally, Assumption A1(d) only requires the interchangeability of the expectation and (partial) differentiation under model (1). It is a mild condition and can be satisfied when $\left| \frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right|$ is upper bounded by an integrable function with respect to the distribution of \mathbf{S} ; see Theorem 1.1 and Example 1.8 in Shao (2003).

We emphasize that Assumption A1(d) is imposed under model (1), because the conditional mean outcome (or regression) function $\mu(t, \mathbf{s}) = \mathbb{E}(Y \mid T = t, \mathbf{S} = \mathbf{s})$ is not well-defined outside the support of the joint density $p(t, \mathbf{s})$. To resolve this definition issue, the following positivity condition is often imposed.

Assumption A2 (Positivity). *The conditional density $p_{T \mid \mathbf{S}}(t \mid \mathbf{s})$ is bounded away from 0 for all $(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}$, i.e., there exist $p_{\min} > 0$ such that $p_{T \mid \mathbf{S}}(t \mid \mathbf{s}) \geq p_{\min}$.*

Under Assumptions A1 and A2, the dose-response curve $m(t)$ and its derivative $\theta(t)$ are identifiable as:

$$m(t) = \mathbb{E}[\mu(t, \mathbf{S})] \quad \text{and} \quad \theta(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right],$$

respectively. In Section 2.1 and Section 3, we first study nonparametric inference on $m(t)$ and $\theta(t)$ this positivity condition. Later in Section 4 and Section 3, we examine inference without positivity.

2.1 Nonparametric Estimation on $m(t)$ With Positivity

Before discussing our estimation strategy on the derivative effect curve $\theta(t)$, we first review the existing approaches for estimating the dose-response curve $m(t)$ under the positivity condition. Specifically, under Assumptions A1 and A2, there are three major estimation strategies for $t \mapsto m(t) = \mathbb{E}[Y(t)]$ with observed data $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$ listed as follows.

- **Regression Adjustment (RA) Estimator:** Since $m(t)$ coincides with the form $\mathbb{E}[\mu(t, \mathbf{S})]$ under Assumptions A1 and A2, it leads to a plug-in estimator as:

$$\hat{m}_{\text{RA}}(t) = \frac{1}{n} \sum_{i=1}^n \hat{\mu}(t, \mathbf{S}_i), \tag{2}$$

where $\hat{\mu}(t, \mathbf{s})$ is a (consistent) estimator of the conditional mean outcome function $\mu(t, \mathbf{s})$.

• **Inverse Probability Weighting (IPW) Estimator:** The IPW estimator follows from the rationale that $m(t) = \mathbb{E} \left[\frac{Y \cdot \mathbb{1}_{\{T=t\}}}{p_{T|\mathbf{S}}(t|\mathbf{S})} \right]$ under Assumptions A1 and A2. In the context of continuous treatment $T \in \mathcal{T}$, we smooth the indicator function by a kernel function $K : \mathbb{R} \rightarrow [0, \infty)$ and obtain that

$$\widehat{m}_{\text{IPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i, \quad (3)$$

where $h > 0$ is a smoothing bandwidth and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ is a (consistent) estimator of the conditional density $p_{T|\mathbf{S}}(t|\mathbf{s})$. In practice, without loss of its consistency, one can implement a self-normalized IPW estimator (29) of $m(t)$ as shown in Section A.1 to reduce the variance of (3).

• **Doubly Robust (DR) Estimator:** The above RA estimator (2) can be combined with the IPW estimator (3) to obtain the following DR estimator as:

$$\widehat{m}_{\text{DR}}(t) = \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot [Y_i - \widehat{\mu}(t, \mathbf{S}_i)] + h \cdot \widehat{\mu}(t, \mathbf{S}_i) \right\}, \quad (4)$$

where $\widehat{\mu}(t, \mathbf{s})$ and $\widehat{p}_{T|\mathbf{S}}(t, \mathbf{s})$ are (consistent) estimators of $\mu(t, \mathbf{s})$ and $p_{T|\mathbf{S}}(t, \mathbf{s})$ respectively. For completeness, we state and prove the asymptotic properties of the above estimators in Section E.

Remark 1. *There exists a slightly different formulation of the IPW estimator of $m(t)$ in the literature (Colangelo and Lee, 2020; Klosin, 2021) as:*

$$\widehat{m}_{\text{IPW},2}(t) = \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}_i)} \cdot Y_i, \quad (5)$$

in which the (estimated) inverse probability weight $\frac{1}{\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}_i)}$ is evaluated at query point t conditioning on each \mathbf{S}_i . We demonstrate in Section D.1 that the asymptotic difference between the oracle versions of (3) and (5) will be of order $O(h^2) + O_P\left(\sqrt{\frac{h}{n}}\right)$ under some regularity conditions, which thus shrinks to 0 as $h \rightarrow 0$ and $n \rightarrow \infty$. In practice, we recommend using the form (3) for the IPW estimator of $m(t)$, because the estimated conditional density $\widehat{p}_{T|\mathbf{S}}$ is more likely to be positive at sample points $(T_i, \mathbf{S}_i), i = 1, \dots, n$ than at the (query) points $(t, \mathbf{S}_i), i = 1, \dots, n$.

3 Nonparametric Inference on $\theta(t)$ With Positivity

In this section, analogous to the estimation of $m(t)$ in Section 2.1, we study three different methods for estimating the derivative effect curve $t \mapsto \theta(t) = \frac{d}{dt} \mathbb{E}[Y(t)]$ with kernel smoothing under Assumptions A1 and A2. Notably, both the IPW and DR estimators of $\theta(t)$ are novel contribution to the existing literature and exhibit distinct insights.

• **Regression Adjustment (RA) Estimator:** Assumption A1(d), together with other conditions in A1 and A2, guarantees the identification of $\theta(t)$ as $\mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right]$ and provides a natural

RA estimator as:

$$\widehat{\theta}_{\text{RA}}(t) = \frac{1}{n} \sum_{i=1}^n \widehat{\beta}(t, \mathbf{S}_i), \quad (6)$$

where $\widehat{\beta}(t, \mathbf{s})$ is a (consistent) estimator of $\beta(t, \mathbf{s}) = \frac{\partial}{\partial t} \mu(t, \mathbf{s})$.

• **Inverse Probability Weighting (IPW) Estimator:** Inspired by the nonparametric derivative estimator in [Mack and Müller \(1989\)](#), we propose the following IPW estimator of $\theta(t)$ as:

$$\widehat{\theta}_{\text{IPW}}(t) = \frac{1}{nh^2} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right)}{\kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}(T_i | \mathbf{S}_i)}, \quad (7)$$

where $K : \mathbb{R} \rightarrow [0, \infty)$ is a kernel function with $\kappa_2 = \int u^2 K(u) du$, $h > 0$ is a smoothing bandwidth, and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ is a (consistent) estimator of the conditional density $p_{T|\mathbf{S}}(t|\mathbf{s})$. One can implement the self-normalized IPW estimator (30) of $\theta(t)$ in [Section A.1](#) to reduce the variance of (7).

Remark 2. One might define the IPW estimator by evaluating the estimated inverse probability weights at points (t, \mathbf{S}_i) , $i = 1, \dots, n$ as:

$$\widehat{\theta}_{\text{IPW},2}(t) = \frac{1}{nh^2} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right)}{\kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}(t | \mathbf{S}_i)}. \quad (8)$$

However, different from (7) in [Remark 1](#), this IPW estimator $\widehat{\theta}_{\text{IPW},2}(t)$ of $\theta(t)$ is (asymptotically) biased even when $h \rightarrow 0$ and $n \rightarrow \infty$; see [Section D.2](#) for details. Hence, our proposed IPW form (7) is preferable not only due to the practical reason as stated in [Remark 1](#) but also because of its statistical consistency as justified in [Theorem 1](#) below.

• **Doubly Robust (DR) Estimator:** Naively, one may combine the RA estimator (6) with the IPW estimator (7) to derive an augmented IPW (AIPW) estimator of $\theta(t)$ with the following (or other similar) form as:

$$\widehat{\theta}_{\text{AIPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{K \left(\frac{T_i - t}{h} \right)}{\widehat{p}_{T|\mathbf{S}}(T_i | \mathbf{S}_i)} \left[\frac{Y_i \left(\frac{T_i - t}{h^2} \right) - \widehat{\beta}(t, \mathbf{S}_i)}{\kappa_2} \right] + h \cdot \widehat{\beta}(t, \mathbf{S}_i) \right\}. \quad (9)$$

However, this naive AIPW estimator is not doubly robust—it is only robust to the misspecification of the limiting quantity of $\widehat{\beta}(t, \mathbf{s})$. In other words, $\widehat{\theta}_{\text{AIPW}}(t)$ will be asymptotically unbiased only when the estimated conditional density $\widehat{p}_{T|\mathbf{S}}$ converges to the true conditional density $p_{T|\mathbf{S}}$ in a certain rate; see [Proposition F.1](#) in [Section F.4](#) for details.

To achieve the doubly robust property like $\widehat{m}_{\text{DR}}(t)$ in (4) (see also [Section E](#)), we propose the following DR estimator of $\theta(t)$ as:

$$\widehat{\theta}_{\text{DR}}(t) = \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{\left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right)}{h \cdot \kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}(T_i | \mathbf{S}_i)} \left[Y_i - \widehat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \widehat{\beta}(t, \mathbf{S}_i) \right] + h \cdot \widehat{\beta}(t, \mathbf{S}_i) \right\}, \quad (10)$$

where $\widehat{\mu}(t, \mathbf{s}), \widehat{\beta}(t, \mathbf{s}), \widehat{p}_{T|\mathbf{S}}(t, \mathbf{s})$ are (consistent) estimators of $\mu(t, \mathbf{s}), \beta(t, \mathbf{s}), p_{T|\mathbf{S}}(t, \mathbf{s})$, respectively. We discuss how these nuisance functions can be estimated in [Section A.3](#). The key insight of why $\widehat{\theta}_{\text{DR}}(t)$ in (10) embraces the doubly robust property is that we leverage a local polynomial approximation ([Fan and Gijbels, 1996](#)) to push the residual of the IPW component in (10) to at least second order before combining with the RA component. In other words, it can be shown that the Neyman orthogonality holds as $h \rightarrow 0$ ([Neyman, 1959, 1979; Chernozhukov et al., 2018](#)). As pointed out in [Remark 2](#), we need to compute the inverse probability weights at the sample points as $\frac{1}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}, i = 1, \dots, n$ for the above DR estimator (10). If we otherwise compute the inverse probability weights at the (query) points as $\frac{1}{\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s}_i)}$ for $i = 1, \dots, n$, then the resulting $\widehat{\theta}_{\text{DR}}(t)$ will be asymptotically biased even when both of the conditional density model $p_{T|\mathbf{S}}$ and the outcome model μ, β are correctly specified. Finally, we also outline a self-normalized version of (10) in [Section A.1](#) for stabilizing its variance.

3.1 Asymptotic Theory

We introduce some regularity conditions under model (1) for our subsequent theoretical analysis. Let $\mathcal{J} \subset \mathcal{T} \times \mathcal{S}$ be the support of the joint density $p(t, \mathbf{s})$, \mathcal{J}° be the interior of \mathcal{J} , and $\partial\mathcal{J}$ be the boundary of \mathcal{J} .

Assumption A3 (Differentiability of the conditional mean outcome function). *For any $(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}$, it holds under model (1) that*

- (a) $\mu(t, \mathbf{s})$ is at least four times continuously differentiable with respect to t .
- (b) $\mu(t, \mathbf{s})$ and all of its partial derivatives are uniformly bounded on $\mathcal{T} \times \mathcal{S}$.

Assumption A4 (Differentiability of the density functions). *For any $(t, \mathbf{s}) \in \mathcal{J}$, it holds that*

- (a) The joint density $p(t, \mathbf{s})$ and the conditional density $p_{T|\mathbf{S}}(t|\mathbf{s})$ are at least three times continuously differentiable with respect to t .
- (b) $p(t, \mathbf{s}), p_{T|\mathbf{S}}(t|\mathbf{s}), p_{\mathbf{S}|T}(\mathbf{s}|t)$, as well as all of the partial derivatives of $p(t, \mathbf{s})$ and $p_{T|\mathbf{S}}(t|\mathbf{s})$ are bounded and continuous up to the boundary $\partial\mathcal{J}$.
- (c) The support \mathcal{T} of the marginal density $p_T(t)$ is compact and $p_T(t)$ is uniformly bounded away from 0 within \mathcal{T} .

Assumption A5 (Regular kernel conditions). *A kernel function $K : \mathbb{R} \rightarrow [0, \infty)$ is bounded and compactly supported on $[-1, 1]$ with $\int_{\mathbb{R}} K(t) dt = 1$ and $K(t) = K(-t)$. In addition, it holds that*

- (a) $\kappa_j := \int_{\mathbb{R}} u^j K(u) du < \infty$ and $\nu_j := \int_{\mathbb{R}} u^j K^2(u) du < \infty$ for all $j = 1, 2, \dots$.
- (b) K is a second-order kernel, i.e., $\kappa_1 = 0$ and $\kappa_2 > 0$.
- (c) $\mathcal{K} = \left\{ t' \mapsto \left(\frac{t'-t}{h} \right)^{k_1} K \left(\frac{t'-t}{h} \right) : t \in \mathcal{T}, h > 0, k_1 = 0, 1 \right\}$ is a bounded VC-type class of measurable functions on \mathbb{R} .

Assumptions A3 and A4 are common smoothness conditions for derivative estimation with kernel smoothing methods (Gasser and Müller, 1984; Mack and Müller, 1989; Wand and Jones, 1994; Wasserman, 2006). These assumptions can be relaxed by the Hölder continuity condition. The uniform lower bound on $p_T(t)$ within its support \mathcal{T} in Assumption A4(c) is only needed when we establish the uniform consistency of our proposed estimators and identify the derivative effect curve $\theta(t)$ when the positivity condition is violated. Assumption A5(a,b) are more like properties than regularity conditions on those commonly used kernel functions, such as the triangular kernel $K(u) = (1 - |u|) \mathbb{1}_{\{|u| \leq 1\}}$ and Epanechnikov kernel $K(u) = \frac{3}{4}(1 - |u|) \mathbb{1}_{\{|u| \leq 1\}}$. Finally, the VC-type condition in Assumption A5(c) is only required when we are interested in the uniform consistency of our proposed estimators over \mathcal{T} .

The following theorem summarizes the consistency and asymptotic results of all the estimators of $\theta(t)$, whose proof is in Section F. Notice that our results in Theorem 1 hold true with a non-separable noise in the potential outcome model as $Y(t) = f(t, \mathbf{S}, \epsilon)$ for some deterministic function f , under which $\mu(t, \mathbf{s})$ stands for $\mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s})$ exclusively.

Theorem 1 (Consistency of estimating $\theta(t)$ under positivity). *Suppose that Assumptions A1, A3, A4, A5, and A2 hold under the general model (1) and $\widehat{\mu}, \widehat{\beta}, \widehat{p}_{T|\mathbf{S}}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s}), \bar{\beta}(t, \mathbf{s})$, and $\bar{p}_{T|\mathbf{S}}(t|\mathbf{s})$ be fixed bounded functions to which $\widehat{\mu}(t, \mathbf{s}), \widehat{\beta}(t, \mathbf{s})$ and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ converge under the rates of convergence as:*

$$\begin{aligned} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} &= O_P(\Upsilon_{1,n}), \quad \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{3,n}), \\ \text{and} \quad \sup_{|u-t| \leq h} \left\| \widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} &= O_P(\Upsilon_{2,n}), \end{aligned}$$

where $\Upsilon_{1,n}, \Upsilon_{3,n}, \Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we have that

$$\begin{aligned} \widehat{\theta}_{\text{RA}}(t) - \theta(t) &= O_P \left(\Upsilon_{3,n} + \left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} + \frac{1}{\sqrt{n}} \right), \\ \widehat{\theta}_{\text{IPW}}(t) - \theta(t) &= O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{2,n} + \sup_{|u-t| \leq h} \left\| \bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} \right). \end{aligned}$$

If, in addition, we assume that

(a) $\bar{p}_{T|\mathbf{S}}$ satisfies Assumptions A4 and A2;

(b) either (i) “ $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ ” with only $h \cdot \Upsilon_{3,n} \rightarrow 0$ or (ii) “ $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$ ”;

(c) $\sqrt{nh} \sup_{|u-t| \leq h} \left\| \widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} \left[\left\| \widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} + h \left\| \widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right] = o_P(1)$,

then

$$\sqrt{nh^3} \left[\widehat{\theta}_{\text{DR}}(t) - \theta(t) \right] = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) + \sqrt{h^3} [\bar{\beta}(t, \mathbf{S}_i) - \mathbb{E}[\beta(t, \mathbf{S})]] \right\} + o_P(1)$$

when $nh^7 \rightarrow c_3$ for some finite number $c_3 \geq 0$, where

$$\phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})].$$

Furthermore,

$$\sqrt{nh^3} \left[\hat{\theta}_{\text{DR}}(t) - \theta(t) - h^2 B_\theta(t) \right] \xrightarrow{d} \mathcal{N}(0, V_\theta(t))$$

with $V_\theta(t) = \mathbb{E} \left[\phi_{h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right]$ and

$$B_\theta(t) = \begin{cases} \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{3 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + p_{T|\mathbf{S}}(t|\mathbf{S}) \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) - 3 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right\} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) \right] & \text{when } \bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}. \end{cases}$$

As established by [Theorem 1](#), the proposed estimator $\hat{\theta}_{\text{DR}}(t)$ achieves doubly robust consistency for $\theta(t)$, provided that either the conditional density model $\bar{p}_{T|\mathbf{S}}$ or the outcome model $\bar{\mu}, \bar{\beta}$ is correctly specified. Unlike the DR estimator $\hat{m}_{\text{DR}}(t)$ of the dose-response curve $m(t)$, which only requires the specification of $\mu(t, \mathbf{s})$ in the outcome model, the DR estimator $\hat{\theta}_{\text{DR}}(t)$ of the derivative effect $\theta(t)$ necessitates specifying both $\mu(t, \mathbf{s})$ and its partial derivative $\beta(t, \mathbf{s}) = \frac{\partial}{\partial t} \mu(t, \mathbf{s})$ in the outcome model. This added complexity is essential for accurately estimating derivatives.

We require in [Theorem 1](#) and other subsequent results that $\hat{\mu}, \hat{\beta}, \hat{p}_{T|\mathbf{S}}$ are obtained from a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. This requirement avoids the need for uniform entropy conditions on $\bar{\mu}, \bar{p}_{T|\mathbf{S}}$ imposed by [Kennedy et al. \(2017\)](#). When no additional data sample is available, these nuisance function estimators $\hat{\mu}, \hat{\beta}, \hat{p}_{T|\mathbf{S}}$ can still be estimated using cross-fitting techniques, allowing for valid construction of the associated estimators of $\theta(t)$; see [Section A.2](#) for the detailed procedures. Importantly, the established rates of convergence in [Theorem 1](#) remain unchanged for the cross-fitted estimators.

3.2 Statistical Inference on $\theta(t)$

To leverage the asymptotic normality of $\hat{\theta}_{\text{DR}}(t)$ for pointwise inference or hypothesis testing on $\theta(t)$ in practice, we need to address two additional challenges: (i) estimate the asymptotic variance $V_\theta(t)$; and (ii) select a proper bandwidth parameter $h > 0$.

For challenge (i), we estimate $V_\theta(t)$ in [Theorem 1](#) by the sample variance of the influence function $\phi_{h,t}$ or the asymptotic linear form as:

$$\hat{V}_\theta(t) = \frac{1}{n} \sum_{i=1}^n \left\{ \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \hat{\mu}, \hat{\beta}, \hat{p}_{T|\mathbf{S}}) + \sqrt{h^3} \left[\hat{\beta}(t, \mathbf{S}_i) - \hat{\theta}_{\text{DR}}(t) \right] \right\}^2. \quad (11)$$

The cross-fitted version of $\hat{V}_\theta(t)$ can be found in (37) of [Section A.2](#). Notice that the second part $\sqrt{h^3} \left[\hat{\beta}(t, \mathbf{S}_i) - \hat{\theta}_{\text{DR}}(t) \right]$ in (11) is asymptotically negligible. We keep this part mainly for a more

conservative estimate of the asymptotic variance $V_\theta(t)$ to guarantee a better empirical coverage of the resulting pointwise confidence interval.

For challenge (ii), the optimal bandwidth that minimizes the asymptotic mean squared error of $\widehat{\theta}_{\text{DR}}(t)$ is of order $O\left(n^{-\frac{1}{7}}\right)$. However, to construct a valid Wald-type confidence interval, an undersmoothing bandwidth h is typically required for the first-order bias of $\widehat{\theta}_{\text{DR}}(t)$ to be asymptotically negligible, *i.e.*, $h^2\sqrt{nh^3} = o(1)$ (Wasserman, 2006, Section 5.7). Therefore, we recommend choosing the bandwidth h to be of order $O\left(n^{-\frac{1}{5}}\right)$, aligning with the outputs of standard bandwidth selection methods for nonparametric regression (Wand and Jones, 1994; Li and Racine, 2004).

Finally, the $(1 - \tau)$ -level confidence interval for $\theta(t)$ is thus given by $\left[\widehat{\theta}_{\text{DR}}(t) \pm q_{1-\frac{\tau}{2}} \sqrt{\frac{\widehat{V}_\theta(t)}{nh^3}}\right]$, where $q_{1-\frac{\tau}{2}}$ is the $(1 - \frac{\tau}{2})$ quantile of the standard normal distribution $\mathcal{N}(0, 1)$.

Remark 3 (Uniform inference via multiplier bootstrap). *It is also statistically valid to conduct uniform inference on $\theta(t)$ over $t \in \mathcal{T}$ via multiplier bootstrap under our regularity conditions in Theorem 1. Specifically, let $\{Z_i\}_{i=1}^n$ be a sequence of i.i.d. random variables independent of the observed data $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$ with $\mathbb{E}(Z_i) = \text{Var}(Z_i) = 1$ and sub-exponential tails. Then, we sample B different i.i.d. datasets $\{Z_i^{(b)}\}_{i=1}^n, b = 1, \dots, B$ and compute the bootstrap DR estimators of $\theta(t)$ as:*

$$\widehat{\theta}_{\text{DR}}^{(b)*}(t) = \frac{1}{nh} \sum_{i=1}^n Z_i^{(b)} \left\{ \frac{\left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{h \cdot \kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \left[Y_i - \widehat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \widehat{\beta}(t, \mathbf{S}_i) \right] + h \cdot \widehat{\beta}(t, \mathbf{S}_i) \right\}$$

for $b = 1, \dots, B$. If $\widehat{Q}(1-\tau)$ is the $(1-\tau)$ quantile of the sequence $\left\{ \sup_{t \in \mathcal{T}} \sqrt{nh^3} \left| \frac{\widehat{\theta}_{\text{DR}}^{(b)*}(t) - \widehat{\theta}_{\text{DR}}(t)}{\sqrt{\widehat{V}_\theta(t)}} \right| \right\}_{b=1}^B$, then the $(1 - \tau)$ uniform confidence band of $\theta(t)$ is given by $\left[\widehat{\theta}_{\text{DR}}(t) \pm \widehat{Q}(1 - \tau) \sqrt{\frac{\widehat{V}_\theta(t)}{nh^3}} \right]$. The asymptotic validity of this confidence band under cross-fitting follows from Theorem 4.2 in Fan et al. (2022); see also Section S4 in Colangelo and Lee (2020).

4 Identification and Inconsistency Issues Without Positivity

This section discusses the general identification issue on the dose-response curve $t \mapsto m(t) = \mathbb{E}[Y(t)]$ and its derivative effect curve $t \mapsto \theta(t) = \frac{d}{dt} \mathbb{E}[Y(t)]$ when the positivity condition (Assumption A2) is violated. We propose an additive structural assumption on the outcome model in (13) to address the identification issue. However, even under this additive confounding model (13), the IPW and DR estimators of $m(t)$ and $\theta(t)$ remain inconsistent without the positivity condition due to the support discrepancy. To resolve this inconsistency, we leverage techniques from nonparametric set estimation to propose our bias-corrected IPW and DR estimators.

4.1 Identification Issue Under the General Model (1)

When the positivity condition (Assumption A2) fails to hold, the conditional mean outcome (or regression) function $\mu(t, \mathbf{s}) = \mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s})$ under model (1) is not well-defined in those regions of $\mathcal{T} \times \mathcal{S}$ that lie outside the support \mathcal{J} of the joint density $p(t, \mathbf{s})$. Hence, the G-computation formulae $\mathbb{E}[\mu(t, \mathbf{S})]$ and $\mathbb{E}\left[\frac{\partial}{\partial t}\mu(t, \mathbf{S})\right]$ are ill-defined and cannot be used to identify $m(t)$ and $\theta(t)$, respectively.

Similarly, identifying $m(t)$ and $\theta(t)$ through the IPW formulae requires the positivity condition as well, because we demonstrate in the proofs of Theorem 1 and Proposition E.1 that

$$\lim_{h \rightarrow 0} \mathbb{E} \left[\frac{Y \cdot K\left(\frac{T-t}{h}\right)}{h \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \right] = \mathbb{E}[\mu(t, \mathbf{S})] \quad \text{and} \quad \lim_{h \rightarrow 0} \mathbb{E} \left[\frac{Y \left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\kappa_2 h^2 \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \right] = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right]. \quad (12)$$

Therefore, it is impossible in general to identify the causal dose-response curve $t \mapsto m(t) = \mathbb{E}[Y(t)]$ and its derivative effect curve $t \mapsto \theta(t) = \frac{d}{dt}\mathbb{E}[Y(t)]$ without further identification or structural assumptions when the positivity condition is violated.

4.1.1 Remedy: Identification Under an Additive Structural Model

While the identifications of $m(t)$ and $\theta(t)$ are infeasible under the general confounding model (1), they are indeed identifiable under an additive structural assumption on the potential outcome model as $Y(t) = \bar{m}(t) + \eta(\mathbf{S}) + \epsilon$ for any $t \in \mathcal{T}$ (Zhang et al., 2024), which, under the consistency condition (Assumption A1(a)), is equivalent to the following additive confounding model

$$Y = \bar{m}(T) + \eta(\mathbf{S}) + \epsilon, \quad (13)$$

where the constraint on ϵ remains the same as in model (1). Such an additive model is a common working model in the context of spatial statistics (Paciorek, 2010; Schnell and Papadogeorgou, 2020), where the covariates $\mathbf{S} \in \mathcal{S} \subset \mathbb{R}^d$ consist of spatial locations or other spatially correlated confounding variables. More broadly, it also appears in the literature of nonparametric (Stone, 1985) and high-dimensional statistics (Meier et al., 2009; Guo et al., 2019).

Under model (13), the dose-response curve $m(t)$ and its derivative $\theta(t)$ become

$$m(t) = \mathbb{E}[Y(t)] = \bar{m}(t) + \mathbb{E}[\eta(\mathbf{S})] \quad \text{and} \quad \theta(t) = m'(t) = \bar{m}'(t). \quad (14)$$

They are identifiable from the observable data through the formulas

$$\begin{aligned} \theta(t) &= \bar{m}'(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = t \right], \\ m(t) &= \mathbb{E} \left[Y + \int_T^t \theta(\tilde{t}) d\tilde{t} \right] = \mathbb{E} \left\{ Y + \int_T^t \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = \tilde{t} \right] d\tilde{t} \right\}. \end{aligned} \quad (15)$$

For completeness, we also summarize this identification theory as Proposition C.1 in Section C. As

a result, the RA estimator of $\theta(t)$ under model (13) without assuming the positivity condition is given by

$$\widehat{\theta}_{C,RA}(t) = \int \widehat{\beta}(t, \mathbf{s}) d\widehat{F}_{\mathbf{S}|T}(\mathbf{s}|t), \quad (16)$$

where $\widehat{\beta}(t, \mathbf{s})$ and $\widehat{F}_{\mathbf{S}|T}(\mathbf{s}|t)$ are (consistent) estimators of $\beta(t, \mathbf{s}) = \frac{\partial}{\partial t}\mu(t, \mathbf{S})$ and the conditional cumulative distribution function (CDF) $P_{\mathbf{S}|T}(\mathbf{s}|t) := F_{\mathbf{S}|T}(\mathbf{s}|t)$, respectively. By (15), the integral RA estimator of $m(t)$ under model (13) can be written as:

$$\widehat{m}_{C,RA}(t) = \frac{1}{n} \sum_{i=1}^n \left[Y_i + \int_{\widetilde{T}_i}^{\widetilde{t}=t} \widehat{\theta}_{C,RA}(\widetilde{t}) d\widetilde{t} \right]. \quad (17)$$

Both estimators (16) and (17) are consistent even when the positivity condition is violated (Zhang et al., 2024); see also Theorem 5 and Section I. In the sequel, we will discuss both the challenges and solutions for extending these RA estimators to IPW and DR estimators of $\theta(t)$ and $m(t)$ under model (13).

4.2 Estimation Issues of IPW Estimators Under the Additive Confounding Model (13)

Although the causal quantities $m(t)$ and $\theta(t)$ are identifiable under the additive confounding model (13), the IPW formulae (12) are indeed biased without positivity due to the support discrepancy between the conditional density $p_{\mathbf{S}|T}(\mathbf{s}|t)$ for $t \in \mathcal{T}$ and the marginal density $p_{\mathbf{S}}(\mathbf{s})$. To examine these biases, we can equivalently analyze the following oracle IPW estimators of $m(t)$ and $\theta(t)$ defined as:

$$\widetilde{m}_{IPW}(t) = \frac{1}{nh} \sum_{i=1}^n \frac{Y_i \cdot K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \quad \text{and} \quad \widetilde{\theta}_{IPW}(t) = \frac{1}{nh^2} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}, \quad (18)$$

where the estimated conditional density $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ is replaced by the true one $p_{T|\mathbf{S}}(t|\mathbf{s})$.

Proposition 2 (Inconsistency of IPW estimators). *Suppose that Assumptions A1(a-c), A3, A4(c), and A5(a-b) hold under the additive confounding model (13). Assume also that when the bandwidth h is small, the Lebesgue measure of the symmetric difference set satisfies*

$$|\mathcal{S}(t+uh) \Delta \mathcal{S}(t)| = |[\mathcal{S}(t+uh) \setminus \mathcal{S}(t)] \cup [\mathcal{S}(t) \setminus \mathcal{S}(t+uh)]| = o(1)$$

for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$. Then, when h is small, the expectation of $\widetilde{m}_{IPW}(t)$ in (18) is given by

$$\mathbb{E}[\widetilde{m}_{IPW}(t)] = \bar{m}(t) \cdot \rho(t) + \omega(t) + o(1),$$

where $\rho(t) = P(\mathbf{S} \in \mathcal{S}(t))$ and $\omega(t) = \mathbb{E}[\eta(\mathbf{S}) \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t)\}}]$. If, in addition, there exists a constant

$A_h > 0$ depending on h such that

$$\int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t) + \eta(\mathbf{S})] [\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}}] \right\} u \cdot K(u) du = O(A_h) \quad (19)$$

for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$ when h is small, then the expectation of $\tilde{\theta}_{\text{IPW}}(t)$ in (12) is given by

$$\mathbb{E} \left[\tilde{\theta}_{\text{IPW}}(t) \right] = \bar{m}'(t) \cdot \rho(t) + O\left(\frac{A_h}{h}\right).$$

The proof of Proposition 2 is in Section G.1. We emphasize that the IPW estimators in (18) have two layers of bias. First, if $\frac{A_h}{h} \rightarrow 0$ as $h \rightarrow 0$ (see also Remark 4 below), then the results in Proposition 2 will imply that

$$\begin{aligned} \lim_{h \rightarrow 0} \mathbb{E} [\tilde{m}_{\text{IPW}}(t)] &= \lim_{h \rightarrow 0} \mathbb{E} \left[\frac{Y \cdot K\left(\frac{T-t}{h}\right)}{h \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \right] = \bar{m}(t) \cdot \rho(t) + \omega(t) \neq m(t), \\ \lim_{h \rightarrow 0} \mathbb{E} [\tilde{\theta}_{\text{IPW}}(t)] &= \lim_{h \rightarrow 0} \mathbb{E} \left[\frac{Y \left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \cdot \kappa_2 \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \right] = \bar{m}'(t) \cdot \rho(t) \neq \theta(t), \end{aligned}$$

where we recall that $m(t) = \bar{m}(t) + \mathbb{E}[\eta(\mathbf{S})]$ and $\theta(t) = \bar{m}'(t)$ from (14). Second, if $\frac{A_h}{h}$ does not converge to 0, then the bias of $\tilde{\theta}_{\text{IPW}}(t)$ will be larger or even diverging to infinity as $h \rightarrow 0$. In reality, the estimation biases or inconsistencies of IPW estimators in (18) are due to the discrepancy between the conditional support $\mathcal{S}(t)$ of $p_{\mathbf{S}|T}(\mathbf{s}|t)$ and the marginal support \mathcal{S} of $p_{\mathbf{S}}(\mathbf{s})$. To correct for the bias of IPW estimators, it is necessary to address the geometric discrepancy, a solution to which will be elaborated upon in Section 5.

Finally, since both RA and IPW estimators cannot be used to identify and estimate $m(t)$ and $\theta(t)$ due to identification and inconsistency issues, the previously studied DR estimators (4) and (10) will be pointless without the positivity condition.

Remark 4. *The regularity condition (19) is indeed not an assumption but rather a natural property. This is because as $h \rightarrow 0$, the differences between two sets $\mathcal{S}(t+uh) \setminus \mathcal{S}(t)$ and $\mathcal{S}(t) \setminus \mathcal{S}(t+uh)$ shrink to 0 for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$. Additionally, when the expectation in (19) is independent of u , one can deduce by the second-order kernel property of K that the left-hand side of (19) is 0. Hence, as $h \rightarrow 0$, the left-hand side of (19) should converge to 0 in a certain rate depending on h .*

5 Nonparametric Inference on $\theta(t)$ Without Positivity

In this section, we present our solution for addressing the estimation biases of IPW estimators for the dose-response curve $m(t)$ and its derivative $\theta(t)$, as described in Section 4.2, when the positivity condition (Assumption A2) is violated. Specifically, our proposed IPW and DR estimators for $\theta(t)$ under the additive confounding model (13) rely on a consistent estimation of the interior region of the support of the conditional density $p_{\mathbf{S}|T}(\mathbf{s}|t)$. Our approach establishes a connection between

the classical support estimation problem and a contemporary causal inference challenge, namely the dose-response curve estimation problem.

5.1 Bias-Corrected IPW and DR Estimators of $\theta(t)$

Recall from (18) and Proposition 2 that the oracle IPW estimator of $\theta(t)$ is the sample average of the IPW quantity $\Xi_t(Y, T, \mathbf{S}) = \frac{Y \left(\frac{T-t}{h}\right) K \left(\frac{T-t}{h}\right)}{h^2 \cdot \kappa_2 \cdot p_{T|\mathbf{S}}(T|\mathbf{S})}$, and it is biased for estimating the quantity of interest $\theta(t) = \bar{m}'(t)$ even under model (13). In particular, $\mathbb{E}[\Xi_t(Y, T, \mathbf{S})]$ converges to $\bar{m}'(t) \cdot \rho(t)$ as $h \rightarrow 0$ under some mild regularity conditions, where $\rho(t) = P(\mathbf{S} \in \mathcal{S}(t))$ for any $t \in \mathcal{T}$. The first step toward removing the bias of $\mathbb{E}[\Xi_t(Y, T, \mathbf{S})]$ is to decouple the quantity of interest $\theta(t) = \bar{m}'(t)$ from the nuisance function $\rho(t)$. To this end, we consider a modified IPW quantity defined as:

$$\tilde{\Xi}_t(Y, T, \mathbf{S}) = \Xi_t(Y, T, \mathbf{S}) \cdot \frac{p_{\mathbf{S}|T}(\mathbf{S}|t)}{p_{\mathbf{S}}(\mathbf{S})} = \frac{Y \left(\frac{T-t}{h}\right) K \left(\frac{T-t}{h}\right) p_{\mathbf{S}|T}(\mathbf{S}|t)}{h^2 \cdot \kappa_2 \cdot p(T, \mathbf{S})}, \quad (20)$$

in which we multiply the original IPW quantity $\Xi_t(Y, T, \mathbf{S})$ by a density ratio $\frac{p_{\mathbf{S}|T}(\mathbf{S}|t)}{p_{\mathbf{S}}(\mathbf{S})}$. The following proposition demonstrates that the remaining bias in $\mathbb{E}[\tilde{\Xi}_t(Y, T, \mathbf{S})]$ can be disentangled from the quantity of interest $\theta(t) = \bar{m}'(t)$ in an additive form.

Proposition 3. *Suppose that Assumptions A1(a-c), A3, A4(c), and A5(a-b) hold under the additive confounding model (13). Then, when the bandwidth h is small, the expectation of the modified IPW quantity (20) is given by*

$$\begin{aligned} \mathbb{E}[\tilde{\Xi}_t(Y, T, \mathbf{S})] &= \bar{m}'(t) + O(h^2) \\ &+ \int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t+uh) + \eta(\mathbf{S})] [\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}}] \Big| T = t \right\} u \cdot K(u) du. \end{aligned}$$


Remark 5. *Different from Remarks 1 and 2, the conditional density $p_{\mathbf{S}|T}$ should be evaluated at the (query) point (t, \mathbf{S}) instead of the sample point (T, \mathbf{S}) in the modified IPW quantity (20). Otherwise, the expectation of (20) will have an asymptotically non-vanishing additive bias; see the proof of Proposition 3 in Section G.2 for details.*

Proposition 3 reveals that the estimation bias of the modified IPW quantity (20) results from the support discrepancy between $\mathcal{S}(t)$ and the integration range $\mathcal{S}(t+uh)$ for a given integration variable $u \in \mathbb{R}$; see Figure 1 for an illustration. As shown in Proposition 2, this additive bias may not always shrink at the rate $O(h^2)$ as $h \rightarrow 0$. To further reduce the bias of the modified IPW quantity (20) to $O(h^2)$ without assuming positivity, we address the support discrepancy of (20) by restricting the conditional density $p_{\mathbf{S}|T}(\mathbf{s}|t)$ to its interior region, defining it as $p_{\zeta}(\mathbf{s}|t)$, and refining (20) as:

$$\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) = \frac{Y \left(\frac{T-t}{h}\right) K \left(\frac{T-t}{h}\right) p_{\zeta}(\mathbf{S}|t)}{h^2 \cdot \kappa_2 \cdot p(T, \mathbf{S})}. \quad (21)$$

Essentially, the only requirement for defining the ζ -interior conditional density $p_{\zeta}(\mathbf{s}|t)$ is that its

Support shrinking approach



Level set approach




Figure 1: Graphical illustrations of the support discrepancy between $\mathcal{S}(t)$ and $\mathcal{S}(t + \delta)$ for $t \in \mathcal{T}$ as well as Assumption A6, where δ can take its value as $uh \in \mathbb{R}$.

support satisfies the following condition:

$$\{\mathbf{s} \in \mathcal{S}(t) : p_{\zeta}(\mathbf{s}|t) > 0\} \subset \mathcal{S}(t + \delta) \quad \text{for any } \delta \in [-h, h]. \quad (22)$$

Here, we propose two approaches for defining $p_{\zeta}(\mathbf{s}|t)$ and leave other options to interested reader.

1. Support Shrinking Approach: Let $\mathcal{S}(t) \ominus \zeta = \{\mathbf{s} \in \mathcal{S}(t) : \inf_{\mathbf{x} \in \partial \mathcal{S}(t)} \|\mathbf{s} - \mathbf{x}\|_2 \geq \zeta\}$ denote the set of interior points of $\mathcal{S}(t)$ that are at least a distance ζ away from the boundary $\mathcal{S}(t)$. Then, we define the ζ -interior conditional density with $\zeta > 0$ being a tuning parameter as:

$$p_{\zeta}(\mathbf{s}|t) = \frac{p_{\mathcal{S}|T}(\mathbf{s}|t) \cdot \mathbb{1}_{\{\mathbf{s} \in \mathcal{S}(t) \ominus \zeta\}}}{\int_{\mathcal{S}(t) \ominus \zeta} p_{\mathcal{S}|T}(\mathbf{s}_1|t) d\mathbf{s}_1} \propto p_{\mathcal{S}|T}(\mathbf{s}|t) \cdot \mathbb{1}_{\{\mathbf{s} \in \mathcal{S}(t) \ominus \zeta\}}. \quad (23)$$

This interior density is indeed the conditional density $p_{\mathcal{S}|T}(\mathbf{s}|t)$ restricted to the interior of its support $\mathcal{S}(t)$. Its estimator $\hat{p}_{\zeta}(\mathbf{s}|t)$ can be constructed using a support estimator $\hat{\mathcal{S}}(t)$ and constraining the conditional density estimator $\hat{p}_{\mathcal{S}|T}(\mathbf{s}|t)$ within the region $\hat{\mathcal{S}}(t) \ominus \zeta$.

2. Level Set Approach: Let $\mathcal{L}_{\zeta}(t) = \{\mathbf{s} \in \mathcal{S}(t) : p_{\mathcal{S}|T}(\mathbf{s}|t) \geq \zeta\}$ be the ζ -upper level set of the conditional density $p_{\mathcal{S}|T}(\mathbf{s}|t)$. Then, we define the ζ -interior conditional density as:

$$p_{\zeta}(\mathbf{s}|t) = \frac{p_{\mathcal{S}|T}(\mathbf{s}|t) \cdot \mathbb{1}_{\{\mathbf{s} \in \mathcal{L}_{\zeta}(t)\}}}{\int_{\mathcal{L}_{\zeta}(t)} p_{\mathcal{S}|T}(\mathbf{s}_1|t) d\mathbf{s}_1} \propto p_{\mathcal{S}|T}(\mathbf{s}|t) \cdot \mathbb{1}_{\{\mathbf{s} \in \mathcal{L}_{\zeta}(t)\}}. \quad (24)$$

The level set approach restricts the conditional density $p_{\mathcal{S}|T}(\mathbf{s}|t)$ to the high-density region, which is generally located away from the support boundary. We may construct the estimator $\hat{p}_{\zeta}(\mathbf{s}|t)$ using a level set estimator $\hat{\mathcal{L}}_{\zeta}(t) = \{\mathbf{s} \in \mathcal{S}(t) : \hat{p}_{\mathcal{S}|T}(\mathbf{s}|t) \geq \zeta\}$ and constraining $\hat{p}_{\mathcal{S}|T}(\mathbf{s}|t)$ to $\hat{\mathcal{L}}_{\zeta}(t)$.

We further specialize condition (22) for the above two approaches by introducing the following smoothness condition on the conditional support $\mathcal{S}(t)$.

Assumption A6 (Smoothness condition on $\mathcal{S}(t)$). *For any $\delta \in \mathbb{R}$ and $t \in \mathcal{T}$, there exists an*

absolute constant $A_0 > 0$ such that either (i) “ $\mathcal{S}(t) \ominus (A_0|\delta|) \subset \mathcal{S}(t + \delta)$ ” for the support shrinking approach or (ii) “ $\mathcal{L}_{A_0|\delta|}(t) \subset \mathcal{S}(t + \delta)$ ” for the level set approach.

To some extent, Assumption A6 can be viewed as a Lipschitz condition of the conditional support $\mathcal{S}(t)$. It can be satisfied when the Euclidean norm of the gradient $\|\nabla_{\mathbf{s}} p_{\mathcal{S}|T}(\mathbf{s}|t)\|_2$ is bounded away from 0 at the boundary of $\mathcal{S}(t)$ (Cadre, 2006). This assumption allows us to ignore the boundary discrepancy as long as we do not evaluate our IPW quantity (21) near the boundary; see Figure 1 for a graphical illustration.

Proposition 4. *Suppose that Assumptions A1(a-c), A3, A4(c), A5(a-b), and A6 hold under the additive confounding model (13). Then, when the bandwidth $h > 0$ is small, the expectation of the modified IPW quantity (21) is given by*

$$\mathbb{E} \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] = \bar{m}'(t) + \frac{h^2 \kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) + O(h^3).$$

The proof of Proposition 4 is in Section G.3. This result demonstrates that the expectation of our newly modified IPW quantity $\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S})$ in (21) converges to the quantity of interest $\theta(t) = \bar{m}'(t)$ in the standard order $O(h^2)$ as $h \rightarrow 0$ under the additive confounding model (13). Notice that the tuning parameter $\zeta = \zeta_n > 0$ in (23) is allowed to converge to 0 as $n \rightarrow \infty$, as long as the condition $h = h_n < \frac{\zeta_n}{A_0}$ holds under Assumption A6.

Given this newly modified IPW quantity (21), we propose the bias-corrected IPW estimator of $\theta(t)$ without the positivity condition as:

$$\hat{\theta}_{\text{C,IPW}}(t) = \frac{1}{nh^2} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right) \hat{p}_{\zeta}(\mathbf{S}_i|t)}{\kappa_2 \cdot \hat{p}(T_i, \mathbf{S}_i)}, \quad (25)$$

where $\hat{p}(t, \mathbf{s})$ is a consistent estimator of the joint density $p(t, \mathbf{s})$ and $\hat{p}_{\zeta}(\mathbf{s}|t)$ is an estimated ζ -interior conditional density.

Finally, we combine the modified RA estimator (16) with our bias-corrected IPW estimator (25) to propose our bias-corrected DR estimator of $\theta(t)$ as:

$$\hat{\theta}_{\text{C,DR}}(t) = \frac{1}{nh^2} \sum_{i=1}^n \frac{\left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right) \hat{p}_{\zeta}(\mathbf{S}_i|t)}{\kappa_2 \cdot \hat{p}(T_i, \mathbf{S}_i)} \left[Y_i - \hat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \hat{\beta}(t, \mathbf{S}_i) \right] + \int \hat{\beta}(t, \mathbf{s}) \cdot \hat{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s}. \quad (26)$$

Notice that for the RA component of $\hat{\theta}_{\text{C,DR}}(t)$, we replace the original conditional CDF estimator $\hat{F}_{\mathcal{S}|T}$ in (16) with the estimated ζ -interior conditional density \hat{p}_{ζ} . This modification is necessary because the IPW component of $\hat{\theta}_{\text{C,DR}}(t)$ is defined through \hat{p}_{ζ} . Both the RA and IPW components need to match up with each other in the definition of $\hat{\theta}_{\text{C,DR}}(t)$ for its consistency.

Remark 6. *While both the support shrinking and level set approaches are valid, we recommend the level set approach in practice, because support estimation is a notoriously challenging problem in nonparametric statistics (Devroye and Wise, 1980). Additionally, selecting an appropriate ζ*

for the support shrinking method is nontrivial. In contrast, level set estimation has been studied over decades (Cuevas and Fraiman, 1997; Cadre, 2006), and the threshold can be set as $\zeta = 0.5 \cdot \max \{\widehat{p}_{\mathbf{S}|T}(\mathbf{S}_i|t) : i = 1, \dots, n\}$. Notice that users may adjust the multiplier 0.5 in this rule, where a smaller value generally increases the effective sample size but also raises the risk of violating condition (22).

5.2 Asymptotic Theory

The following theorem summarizes the consistency and asymptotic results of our RA (16), IPW (25), and DR (26) estimators of $\theta(t)$ under the additive confounding model (13) without assuming the positivity condition.

Theorem 5 (Consistency of estimating $\theta(t)$ without positivity). *Suppose that Assumptions A1(a-c), A3, A4, A5, and A6 hold under the additive confounding model (13), and the support $\mathcal{S} \subset \mathbb{R}^d$ of the marginal density $p_{\mathbf{S}}$ is compact. In addition, $\widehat{\mu}, \widehat{\beta}, \widehat{p}_{\zeta}, \widehat{p}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s}), \bar{\beta}(t, \mathbf{s}), \bar{p}_{\zeta}(\mathbf{s}|t)$, and $\bar{p}(t, \mathbf{s})$ be fixed bounded functions to which $\widehat{\mu}(t, \mathbf{s}), \widehat{\beta}(t, \mathbf{s}), \widehat{p}_{\zeta}(\mathbf{s}|t)$, and $\widehat{p}(t, \mathbf{s})$ converge under the rates of convergence as:*

$$\begin{aligned} \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} &= O_P(\Upsilon_{3,n}), \quad \sup_{\mathbf{s} \in \mathcal{S}} \left| \widehat{F}_{\mathbf{S}|T}(\mathbf{s}|t) - F_{\mathbf{S}|T}(\mathbf{s}|t) \right| = O_P(\Upsilon_{4,n}), \\ \left\| \widehat{p}_{\zeta}(\mathbf{S}|t) - \bar{p}_{\zeta}(\mathbf{S}|t) \right\|_{L_2} &= O_P(\Upsilon_{5,n}), \quad \text{and} \quad \sup_{|u-t| \leq h} \left\| \widehat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{6,n}), \end{aligned}$$

where $\Upsilon_{3,n}, \Upsilon_{4,n}, \Upsilon_{5,n}, \Upsilon_{6,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we have that

$$\begin{aligned} \widehat{\theta}_{\text{C,RA}}(t) - \theta(t) &= O_P\left(\Upsilon_{3,n} + \Upsilon_{4,n} + \left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2}\right), \\ \widehat{\theta}_{\text{C,IPW}}(t) - \theta(t) &= O(h^2) + O_P\left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{5,n} + \Upsilon_{6,n} + \sup_{|u-t| \leq h} \left\| \bar{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2}\right). \end{aligned}$$

If, in addition, we assume that

(a) \bar{p}, \bar{p}_{ζ} satisfy Assumptions A4 and A6 as well as $\sqrt{nh^3} \cdot \Upsilon_{5,n} = o(1)$;

(b) either (i) “ $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ ” or (ii) “ $\bar{p} = p$ ”;

(c) $\sqrt{nh} \left[\left\| \widehat{p}_{\zeta}(\mathbf{S}|t) - \bar{p}_{\zeta}(\mathbf{S}|t) \right\|_{L_2} + \sup_{|u-t| \leq h} \left\| \widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2} \right] \left[\left\| \widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} + h \left\| \widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right] = o_P(1)$,

then

$$\begin{aligned} &\sqrt{nh^3} \left[\widehat{\theta}_{\text{C,DR}}(t) - \theta(t) \right] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{\text{C},h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_{\zeta}) + \sqrt{h^3} \left[\int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} - \theta(t) \right] \right\} + o_P(1) \end{aligned}$$

when $nh^7 \rightarrow c_3$ for some finite number $c_3 \geq 0$, where

$$\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})].$$

Furthermore,

$$\sqrt{nh^3} \left[\hat{\theta}_{C,DR}(t) - \theta(t) - h^2 B_{C,\theta}(t) \right] \xrightarrow{d} \mathcal{N}(0, V_{C,\theta}(t))$$

with $V_{C,\theta}(t) = \mathbb{E} \left[\phi_{C,h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right]$ and

$$B_{C,\theta}(t) = \begin{cases} \frac{\kappa_4}{6\kappa_2} \int \left\{ \frac{3 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \bar{m}''(t) + p(t, \mathbf{s}) [\bar{m}^{(3)}(t) - 3 \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \cdot \bar{m}''(t)]}{\bar{p}(t, \mathbf{s})} \right\} \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) & \text{when } \bar{p} = p. \end{cases}$$

The proof of [Theorem 5](#) is in [Section H](#). Similar to our discussion after [Theorem 1](#) in [Section 3.2](#), we can estimate the asymptotic variance $V_{C,\theta}(t)$ in [Theorem 5](#) by

$$\hat{V}_{C,\theta}(t) = \frac{1}{n} \sum_{i=1}^n \left\{ \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \hat{\mu}, \hat{\beta}, \hat{p}, \hat{p}_\zeta) + \sqrt{h^3} \left[\int \hat{\beta}(t, \mathbf{s}) \cdot \hat{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \hat{\theta}_{C,DR}(t) \right] \right\}^2$$

and choose the bandwidth h to be of order $O\left(n^{-\frac{1}{5}}\right)$ to ensure valid inference. As a corollary, we can plug either IPW ([25](#)) or DR ([26](#)) estimators into our integral formula ([17](#)) to obtain the integral IPW or DR estimators of the dose-response curve $m(t)$ under model ([13](#)). We establish the asymptotic theory for these integral estimators in [Corollary I.1](#) of [Section I](#).

Remark 7. Under usual regularity conditions and the setting of nonparametric estimation ([Wasserman, 2006](#)), the rates of convergence $\Upsilon_{1,n}, \Upsilon_{2,n}, \Upsilon_{6,n}$ for $\hat{\mu}, \hat{p}_{T|\mathbf{S}}, \hat{p}$ in [Theorem 1](#) and [Theorem 5](#) would be of order $O\left(n^{-\frac{2}{4+d}}\right)$ up to some possible $\log n$ factors, while the rate of convergence $\Upsilon_{3,n}$ for $\hat{\beta}$ is of order $O\left(n^{-\frac{2}{6+d}}\right)$. As shown by [Farrell et al. \(2021\)](#); [Colangelo and Lee \(2020\)](#), these rates are attainable by neural network models. Additionally, the rate of convergence $\Upsilon_{4,n}$ for $\hat{F}_{\mathbf{S}|T}$ can be dimensionally independent and of order $O\left(\left(\frac{\log n}{n}\right)^{\frac{2}{5}}\right)$ ([Einmahl and Mason, 2005](#)), while $\Upsilon_{5,n}$ for \hat{p}_ζ is typically of order $O\left(n^{-\frac{2}{5+d}}\right)$ ([Cuevas and Fraiman, 1997](#); [Tsybakov, 1997](#)). Faster rates of convergence are possible if we assume higher-order smoothness conditions and use higher-order kernel functions.

6 Numerical Experiments

In this section, we evaluate the finite-sample performances of our proposed estimators of $\theta(t) = \frac{d}{dt} \mathbb{E}[Y(t)]$ in [Section 3](#) and compare them with the finite-difference approach in [Colangelo and Lee \(2020\)](#) under the positivity condition through simulation studies and an analysis of the Job Corps program in the United States. Furthermore, we compare the bias-corrected estimators of $\theta(t)$ in

Section 5.1 with their counterparts via simulation studies when the positivity condition is violated.

6.1 Simulation Studies With Positivity

We generate i.i.d. observations $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$ from the following data-generating model as in Colangelo and Lee (2020); Klosin (2021):

$$\begin{aligned} Y &= 1.2T + T^2 + TS_1 + 1.2\boldsymbol{\xi}^T \mathbf{S} + \epsilon \cdot \sqrt{0.5 + F_{\mathcal{N}(0,1)}(S_1)}, \quad \epsilon \sim \mathcal{N}(0, 1), \\ T &= F_{\mathcal{N}(0,1)}(3\boldsymbol{\xi}^T \mathbf{S}) - 0.5 + 0.75E, \quad \mathbf{S} = (S_1, \dots, S_d)^T \sim \mathcal{N}_d(\mathbf{0}, \Sigma), \quad E \sim \mathcal{N}(0, 1), \end{aligned} \tag{27}$$

where $F_{\mathcal{N}(0,1)}$ is the CDF of $\mathcal{N}(0, 1)$, $\boldsymbol{\xi} = (\xi_1, \dots, \xi_d)^T \in \mathbb{R}^d$ has its entry $\xi_j = \frac{1}{j^2}$ for $j = 1, \dots, d$ as well as $\Sigma_{ii} = 1$, $\Sigma_{ij} = 0.5$ when $|i - j| = 1$, and $\Sigma_{ij} = 0$ when $|i - j| > 1$ for $i, j = 1, \dots, d$. Here, $d = 20$ unless stated otherwise. The dose-response curve is thus given by $m(t) = 1.2t + t^2$, and our parameter of interest is the derivative effect curve $\theta(t) = 1.2 + 2t$.

We evaluate our proposed estimators of $\theta(t)$ in Section 3 alongside the finite-difference estimator by Colangelo and Lee (2020) with 5-fold cross-fitting. In particular, we replicate their finite-difference estimators using their neural network (NN) and kernel neural network (KNN) models for estimating the nuisance functions $\mu(t, \mathbf{s})$ and $p_{T|\mathbf{S}}(t|\mathbf{s})$, which yield their best performances. Additionally, similar to the setups in Colangelo and Lee (2020); Klosin (2021), we use the Epanechnikov kernel $K(u) = \frac{3}{4}(1 - |u|) \mathbb{1}_{\{|u| \leq 1\}}$ under a bandwidth choice $h = 1.25 \hat{\sigma}_T \cdot n^{-\frac{1}{5}}$, where $\hat{\sigma}_T$ is the sample standard deviation of $\{T_1, \dots, T_n\}$. Furthermore, for our proposed estimators, the nuisance functions $\mu(t, \mathbf{s})$ and $\beta(t, \mathbf{s})$ are estimated by neural network models as well, while $p_{T|\mathbf{S}}(t|\mathbf{s})$ is estimated by either the method of kernel density estimation (KDE) on residuals or the approach of regressing kernel-smoothed outcomes (RKS); see Section A.3 for details. To prevent division by zero, all estimated conditional density values $\hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i), i = 1, \dots, n$ smaller than 0.001 are set to this value. For comparison, we also implement our proposed DR estimator of $\theta(t)$ under the true conditional density (“True”). All our DR estimators are self-normalized as described in Section A.1 to reduce their variances. The nominal levels of all the yielded pointwise confidence intervals are set to 95%.

The simulation results are shown in Figure 2 for various sample sizes, where the estimation biases, root mean square errors (RMSEs), and coverage rates of confidence intervals are calculated by averaging over 1000 Monte Carlo replications. Additional results when the bandwidth parameter varies or cross-fitting is not employed are in Section B.1 and Section B.2. Unlike prior studies in Colangelo and Lee (2020); Klosin (2021), which focus solely on $t = 0$, our comparative simulations evaluate 81 treatment values across $t \in [-2, 2]$. Overall, our proposed DR estimators, using either true or KDE-estimated conditional density values, outperform the finite-difference methods of Colangelo and Lee (2020) in terms of estimation bias while maintaining comparable RMSE. When it comes to statistical inference, the confidence intervals from our DR estimators consistently show better empirical coverages than those from Colangelo and Lee (2020). These performance advantages of our DR estimators arise from directly estimating and inferring $\theta(t)$ without requiring a




Figure 2: Comparisons between our proposed estimators and the finite-difference approaches by Colangelo and Lee (2020) (“CL20”) under positivity and with 5-fold cross-fitting across various sample sizes. Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different values for n .

step-size parameter for finite-difference approximations.

6.2 Simulation Studies Without Positivity

We now assess the finite-sample performances of our bias-corrected IPW and DR estimators of $\theta(t)$ in Section 5.1 and compare them with those counterparts in Section 3 when the positivity condition is violated. To this end, we generate i.i.d. data $\{(Y_i, T_i, S_i)\}_{i=1}^n$ from the following data-generating model

$$Y = T^3 + T^2 + 10S + \epsilon, \quad T = \sin(\pi S) + E, \quad S \sim \text{Uniform}[-1, 1] \subset \mathbb{R}, \quad (28)$$

where $E \sim \text{Uniform}[-0.3, 0.3]$ is an independent treatment variation and $\epsilon \sim \mathcal{N}(0, 1)$ is an independent noise variable. The marginal supports of T and S are $\mathcal{T} = [-1.3, 1.3]$ and $\mathcal{S} = [-1, 1]$ respectively, while the joint support of (T, S) only covers a thin band region of the product space $\mathcal{T} \times \mathcal{S}$; see Figure 1 in Zhang et al. (2024) for illustration. The true derivative effect curve is thus given by $\theta(t) = 3t^2 + 2t$.

We evaluate our bias-corrected estimators of $\theta(t)$ in Section 5.1 on the simulated dataset, alongside those estimators from Section 3 that assumes the positivity condition. All these estimators are




Figure 3: Comparisons between our bias-corrected estimators (NP) in Section 5.1 and their counterparts (P) in Section 3 under the violation of positivity and with 5-fold cross-fitting across different sample sizes. Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different values for n .

assessed with 5-fold cross-fitting. Again, we use the Epanechnikov kernel $K(u) = \frac{3}{4}(1 - |u|) \mathbb{1}_{\{|u| \leq 1\}}$ under a bandwidth choice $h = 2\hat{\sigma}_T \cdot n^{-\frac{1}{5}}$. For those estimators assuming positivity, we estimate the nuisance functions $\mu(t, s)$ and $\beta(t, s)$ by neural network models in Section A.3 and utilize the true conditional density function $p_{T|S}$ evaluated at the observations $\{(T_i, S_i)\}_{i=1}^n$. For the bias-corrected estimators, we estimate the joint density $p(t, s)$ and conditional density $p_{S|T}(s|t)$ using kernel density estimation with a Gaussian kernel $K(u) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right)$. The estimated interior densities $\hat{p}_\zeta(S_i|t), i = 1, \dots, n$ are computed via the trimming method outlined in Remark 6. All the estimators are self-normalized as described in Section A.1 to reduce their variances, and the nominal levels of all the yielded pointwise confidence intervals are set to 95%.

The simulation results for different sample sizes are presented in Figure 3, where the estimation biases, root mean square errors (RMSEs), and coverage rates of confidence intervals are calculated by averaging over 1000 Monte Carlo replications. Additional results when the bandwidth parameter varies or cross-fitting is not employed are in Section B.3 and Section B.4. The bias-corrected IPW estimator (25) effectively reduces the estimation biases of the standard IPW estimator (7) of $\theta(t)$ across when the positivity condition is violated. Furthermore, the bias-corrected DR estimator (26) achieves comparable biases and RMSEs to its standard counterpart (10), even when (10) uses the oracle conditional density $p_{T|S}$. Notably, the confidence intervals yielded by the bias-corrected




Figure 4: Estimated derivative effect curves with 95% confidence intervals using our proposed estimators and the finite-difference approaches by Colangelo and Lee (2020) (“CL20”) under 5-fold cross-fitting. The vertical red dotted lines mark the original treatment range [320, 1840] analyzed in Colangelo and Lee (2020).

DR estimator (26) exhibit better coverage probabilities compared to its counterpart (10). These findings support the theoretical properties of our proposed bias-corrected IPW and DR estimators in Section 5.1. Nonetheless, the bias-corrected RA estimator (16) remains the preferred choice when it comes to estimation accuracy due to its simplicity under violations of the positivity condition.

6.3 Case Study: An Analysis of the Job Corps Program

We demonstrate the applicability of our proposed DR estimators for $\theta(t)$ by extending the analysis of Colangelo and Lee (2020) on the Job Corps program in the United States (U.S.). This program aims at providing academic and vocational training to U.S. legal residents aged 16–24 who come from low-income households (Schochet et al., 2001). The data used in our analysis originated from the National Job Corps Study, which conducted some randomized experiments on first-time applicants in the 48 contiguous states and the District of Columbia between November 1994 and February 1996 (Schochet et al., 2008).

Numerous studies have examined the causal effects of the Job Corps program from various angles (Flores and Flores-Lagunes, 2009; Flores et al., 2012; Huber, 2014; Lee, 2018; Huber et al., 2020; Lee and Liu, 2024). Following Colangelo and Lee (2020), we analyze the relationship between employment outcomes and the duration of academic and vocational training, focusing on the derivative effect curve $\theta(t) = \frac{d}{dt} \mathbb{E}[Y(t)]$. The data sample includes 4,024 individuals who received at least 40 hours of training. The outcome variable Y represents the proportions of weeks employed in the second year following the program assignment, and the treatment variable T is the total hours of academic and vocational training received. The covariate vector \mathbf{S} , comprising 49 socioeconomic characteristics, ensures the validity of the ignorability assumption (Flores et al., 2012); see Table 4 in Huber et al. (2020) for detailed descriptions of the covariates. Before applying derivative effect estimation methods, categorical covariates were converted to dummy variables, and all variables were standardized to have mean 0 and variance 1.

We apply our proposed DR estimator (10) with the same setup as in Section 6.1 to the standardized data, extending the range of queried treatment values from [320, 1840] to [40, 4000]. For consistency, we use the same bandwidth parameter $h = 223$ and apply the neural network model for conditional density estimation as in Colangelo and Lee (2020). The estimated derivative effect curves with 95% confidence intervals under 5-fold cross-fitting are shown in Figure 4. Overall, our DR estimator produces similar patterns to the finite-difference estimates from Colangelo and Lee (2020). However, our confidence intervals are more conservative and include 0 for nearly all treatment values, suggesting insufficient evidence to confirm the program’s effectiveness. Additional results when cross-fitting is not employed are shown in Section B.5.

7 Discussion

In summary, this paper studies nonparametric DR inference methods for the derivative function of the dose-response curve with and without the positivity condition. We establish the asymptotic properties of our proposed estimators under mild conditions, permitting the use of machine learning methods for nuisance function estimation with cross-fitting. Furthermore, our identification theory and refinements of IPW and DR estimators without positivity open up a novel link between the dose-response curve inference challenge and the nonparametric set estimation problem. Simulation studies and empirical applications demonstrate the advantages of our DR estimator over the existing finite-difference method for derivative effect inference. This work also highlights several avenues for future research.

1. Efficiency theory for DR estimators: As discussed in Section 1.2, the derivative effect curve $\theta(t)$ is not pathwise differentiable relative to a nonparametric model (Bickel, 1982; Bickel et al., 1998). A promising direction for future work is to define a sequence of smoothed parameters approximating $\theta(t)$ through kernel smoothing, each pathwise differentiable (van der Laan et al., 2018), and derive their efficient influence functions. Investigating the alignment of these kernel-smoothed efficient influence functions with the asymptotic linear forms derived in Theorem 1 and Theorem 5 would provide deeper theoretical insights.

2. Bias correction for DR estimators: As shown in Theorem 1 and Theorem 5, our DR estimators of $\theta(t)$ contain bias terms of order $O(h^2)$. These biases become asymptotically negligible when the bandwidth is chosen as $h \asymp n^{-\frac{1}{5}}$ that matches up the standard rate of convergence for nonparametric regression. To guarantee valid inference, an alternative approach is to explicitly estimate and correct these bias terms, as demonstrated by Calonico et al. (2018); Cheng and Chen (2019); Takatsu and Westling (2024). A rigorous investigation of this bias-corrected approach for our DR estimators would be a valuable direction for future research.

3. Derivative estimation in other causal contexts: Our proposed DR inference methods for $\theta(t)$ can be naturally extended to conduct inference on other causal estimands of interest, such as the instantaneous causal effect $\frac{d}{dt}\mathbb{E}[Y(t)|\mathbf{S} = \mathbf{s}]$ (Stolzenberg, 1980; Ratkovic and Tingley, 2017) or the marginal direct and indirect effects in causal mediation analysis (Huber et al., 2020).

Acknowledgement

We thank Alex Luedtke and Jon A. Wellner for their helpful comments. YZ is supported in part by YC’s NSF grant DMS-2141808. YC is supported by NSF grants DMS-1952781, 2112907, 2141808, and NIH U24-AG07212.

References

- H. Bang and J. M. Robins. Doubly robust estimation in missing data and causal inference models. *Biometrics*, 61(4):962–973, 2005.
- A. G. Baydin, B. A. Pearlmutter, A. A. Radul, and J. M. Siskind. Automatic differentiation in machine learning: a survey. *Journal of machine learning research*, 18(153):1–43, 2018.
- P. Bickel, C. Klaassen, Y. Ritov, and J. Wellner. *Efficient and Adaptive Estimation for Semiparametric Models*. Springer New York, 1998.
- P. J. Bickel. On adaptive estimation. *The Annals of Statistics*, 10(3):647–671, 1982.
- M. Blondel and V. Roulet. The elements of differentiable programming. *arXiv preprint arXiv:2403.14606*, 2024.
- S. Bong and K. Lee. Local causal effects with continuous exposures: A matching estimator for the average causal derivative effect. *arXiv preprint arXiv:2311.18532*, 2023.
- M. Bonvini and E. H. Kennedy. Fast convergence rates for dose-response estimation. *arXiv preprint arXiv:2207.11825*, 2022.
- M. Bonvini, A. McClean, Z. Branson, and E. H. Kennedy. Incremental causal effects: an introduction and review. In *Handbook of matching and weighting adjustments for causal inference*, pages 349–372. Chapman and Hall/CRC, 2023.
- Z. Branson, E. H. Kennedy, S. Balakrishnan, and L. Wasserman. Causal effect estimation after propensity score trimming with continuous treatments. *arXiv preprint arXiv:2309.00706*, 2023.
- B. Cadre. Kernel estimation of density level sets. *Journal of Multivariate Analysis*, 97(4):999–1023, 2006.
- S. Calonico, M. D. Cattaneo, and M. H. Farrell. On the effect of bias estimation on coverage accuracy in nonparametric inference. *Journal of the American Statistical Association*, 113(522):767–779, 2018.
- M. Carone, A. R. Luedtke, and M. J. van der Laan. Toward computerized efficient estimation in infinite-dimensional models. *Journal of the American Statistical Association*, 114(527):1174–1190, 2019.
- M. D. Cattaneo, R. K. Crump, and M. Jansson. Robust data-driven inference for density-weighted average derivatives. *Journal of the American Statistical Association*, 105(491):1070–1083, 2010.
- X. Chen and Z. Liao. Sieve m inference on irregular parameters. *Journal of Econometrics*, 182(1):70–86, 2014.

- X. Chen, Z. Liao, and Y. Sun. Sieve inference on possibly misspecified semi-nonparametric time series models. *Journal of Econometrics*, 178:639–658, 2014.
- G. Cheng and Y.-C. Chen. Nonparametric inference via bootstrapping the debiased estimator. *Electronic Journal of Statistics*, 13(1):2194 – 2256, 2019.
- V. Chernozhukov, D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins. Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68, 01 2018.
- V. Chernozhukov, W. K. Newey, and R. Singh. Automatic debiased machine learning of causal and structural effects. *Econometrica*, 90(3):967–1027, 2022.
- K. Colangelo and Y.-Y. Lee. Double debiased machine learning nonparametric inference with continuous treatments. *arXiv preprint arXiv:2004.03036*, 2020.
- S. R. Cole and M. A. Hernán. Constructing inverse probability weights for marginal structural models. *American Journal of Epidemiology*, 168(6):656–664, 2008.
- A. Cuevas. Set estimation: Another bridge between statistics and geometry. *Boletín de Estadística e Investigación Operativa*, 25(2):71–85, 2009.
- A. Cuevas and R. Fraiman. A plug-in approach to support estimation. *The Annals of Statistics*, 25(6):2300 – 2312, 1997.
- L. Devroye and G. L. Wise. Detection of abnormal behavior via nonparametric estimation of the support. *SIAM Journal on Applied Mathematics*, 38(3):480–488, 1980.
- I. Díaz and N. S. Hejazi. Causal mediation analysis for stochastic interventions. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82(3):661–683, 2020.
- I. Díaz and M. J. van der Laan. Targeted data adaptive estimation of the causal dose–response curve. *Journal of Causal Inference*, 1(2):171–192, 2013.
- U. Einmahl and D. M. Mason. Uniform in bandwidth consistency of kernel-type function estimators. *The Annals of Statistics*, 33(3):1380 – 1403, 2005.
- J. Fan and I. Gijbels. *Local polynomial modelling and its applications*, volume 66. Chapman & Hall/CRC, 1996.
- Q. Fan, Y.-C. Hsu, R. P. Lieli, and Y. Zhang. Estimation of conditional average treatment effects with high-dimensional data. *Journal of Business & Economic Statistics*, 40(1):313–327, 2022.
- M. H. Farrell, T. Liang, and S. Misra. Deep neural networks for estimation and inference. *Econometrica*, 89(1):181–213, 2021.
- C. Flores. Estimation of dose-response functions and optimal doses with a continuous treatment. Technical report, Department of Economics, University of Miami, 2007. URL <https://core.ac.uk/download/pdf/7169663.pdf>.
- C. A. Flores and A. Flores-Lagunes. Identification and estimation of causal mechanisms and net effects of a treatment under unconfoundedness. IZA Discussion Papers 4237, Institute of Labor Economics (IZA), 2009.

- C. A. Flores, A. Flores-Lagunes, A. Gonzalez, and T. C. Neumann. Estimating the effects of length of exposure to instruction in a training program: The case of job corps. *Review of Economics and Statistics*, 94(1):153–171, 2012.
- A. F. Galvao and L. Wang. Uniformly semiparametric efficient estimation of treatment effects with a continuous treatment. *Journal of the American Statistical Association*, 110(512):1528–1542, 2015.
- T. Gasser and H.-G. Müller. Estimating regression functions and their derivatives by the kernel method. *Scandinavian Journal of Statistics*, pages 171–185, 1984.
- R. D. Gill and J. M. Robins. Causal inference for complex longitudinal data: the continuous case. *Annals of Statistics*, 29(6):1785–1811, 2001.
- V. Godambe and V. Joshi. Admissibility and bayes estimation in sampling finite populations. i. *The Annals of Mathematical Statistics*, 36(6):1707–1722, 1965.
- Z. Guo, W. Yuan, and C.-H. Zhang. Decorrelated local linear estimator: Inference for non-linear effects in high-dimensional additive models. *arXiv preprint arXiv:1907.12732*, 2019.
- W. Härdle and T. M. Stoker. Investigating smooth multiple regression by the method of average derivatives. *Journal of the American statistical Association*, 84(408):986–995, 1989.
- J. D. Hart and P. Vieu. Data-driven bandwidth choice for density estimation based on dependent data. *The Annals of Statistics*, pages 873–890, 1990.
- O. Hines, K. Diaz-Ordaz, and S. Vansteelandt. Optimally weighted average derivative effects. *arXiv preprint arXiv:2308.05456*, 2023.
- K. Hirano and G. W. Imbens. *The Propensity Score with Continuous Treatments*, chapter 7, pages 73–84. John Wiley & Sons, Ltd, 2004.
- D. A. Hirshberg and S. Wager. Debiased inference of average partial effects in single-index models: Comment on wooldridge and zhu. *Journal of Business & Economic Statistics*, 38(1):19–24, 2020.
- M. Huber. Identifying causal mechanisms (primarily) based on inverse probability weighting. *Journal of Applied Econometrics*, 29(6):920–943, 2014.
- M. Huber, Y.-C. Hsu, Y.-Y. Lee, and L. Lettry. Direct and indirect effects of continuous treatments based on generalized propensity score weighting. *Journal of Applied Econometrics*, 35(7):814–840, 2020.
- H. Ichimura and W. K. Newey. The influence function of semiparametric estimators. *Quantitative Economics*, 13(1):29–61, 2022.
- K. Imai and D. A. van Dyk. Causal inference with general treatment regimes: Generalizing the propensity score. *Journal of the American Statistical Association*, 99(467):854–866, 2004.
- N. Kallus and A. Zhou. Policy evaluation and optimization with continuous treatments. In *International Conference on Artificial Intelligence and Statistics*, pages 1243–1251. PMLR, 2018.
- E. H. Kennedy. Nonparametric causal effects based on incremental propensity score interventions. *Journal of the American Statistical Association*, 114(526):645–656, 2019.

- E. H. Kennedy, Z. Ma, M. D. McHugh, and D. S. Small. Nonparametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 79(4):1229–1245, 2017.
- S. Klosin. Automatic double machine learning for continuous treatment effects. *arXiv preprint arXiv:2104.10334*, 2021.
- D. S. Lee. Training, wages, and sample selection: Estimating sharp bounds on treatment effects. *The Review of Economic Studies*, 76(3):1071–1102, 2009.
- Y.-Y. Lee. Partial mean processes with generated regressors: Continuous treatment effects and nonseparable models. *arXiv preprint arXiv:1811.00157*, 2018.
- Y.-Y. Lee and C.-A. Liu. Lee bounds with a continuous treatment in sample selection. *arXiv preprint arXiv:2411.04312*, 2024.
- E. L. Lehmann. *Elements of large-sample theory*. Springer, 1999.
- Q. Li and J. Racine. Cross-validated local linear nonparametric regression. *Statistica Sinica*, 14: 485–512, 2004.
- A. Luedtke. Simplifying debiased inference via automatic differentiation and probabilistic programming. *arXiv preprint arXiv:2405.08675*, 2024.
- A. Luedtke and I. Chung. One-step estimation of differentiable hilbert-valued parameters. *The Annals of Statistics*, 52(4):1534–1563, 2024.
- Y. Mack and H.-G. Müller. Derivative estimation in nonparametric regression with random predictor variable. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 59–72, 1989.
- A. McClean, Y. Li, S. Bae, M. A. McAdams-DeMarco, I. Díaz, and W. Wu. Fair comparisons of causal parameters with many treatments and positivity violations. *arXiv preprint arXiv:2410.13522*, 2024.
- L. Meier, S. van de Geer, and P. Bühlmann. High-dimensional additive modeling. *The Annals of Statistics*, 37(6B):3779 – 3821, 2009.
- J. Meloche. Asymptotic behaviour of the mean integrated squared error of kernel density estimators for dependent observations. *The Canadian Journal of Statistics/La Revue Canadienne de Statistique*, pages 205–211, 1990.
- R. Neugebauer and M. van der Laan. Nonparametric causal effects based on marginal structural models. *Journal of Statistical Planning and Inference*, 137(2):419–434, 2007.
- W. K. Newey and J. R. Robins. Cross-fitting and fast remainder rates for semiparametric estimation. *arXiv preprint arXiv:1801.09138*, 2018.
- W. K. Newey and T. M. Stoker. Efficiency of weighted average derivative estimators and index models. *Econometrica*, 61(5):1199–1223, 1993.
- J. Neyman. Optimal asymptotic tests of composite hypotheses. *Probability and Statistics*, pages 213–234, 1959.

- J. Neyman. $C(\alpha)$ tests and their use. *Sankhyā: The Indian Journal of Statistics, Series A*, 41(1/2): 1–21, 1979.
- C. J. Paciorek. The importance of scale for spatial-confounding bias and precision of spatial regression estimators. *Statistical Science*, 25(1):107–125, 2010.
- A. Paszke, S. Gross, S. Chintala, G. Chanan, E. Yang, Z. DeVito, Z. Lin, A. Desmaison, L. Antiga, and A. Lerer. Automatic differentiation in pytorch. In *NIPS 2017 Workshop on Autodiff*, 2017.
- A. Paszke, S. Gross, F. Massa, A. Lerer, J. Bradbury, G. Chanan, T. Killeen, Z. Lin, N. Gimeshain, L. Antiga, A. Desmaison, A. Kopf, E. Yang, Z. DeVito, M. Raison, A. Tejani, S. Chilamkurthy, B. Steiner, L. Fang, J. Bai, and S. Chintala. Pytorch: An imperative style, high-performance deep learning library. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.
- J. L. Powell, J. H. Stock, and T. M. Stoker. Semiparametric estimation of index coefficients. *Econometrica*, pages 1403–1430, 1989.
- M. Ratkovic and D. Tingley. Causal inference through the method of direct estimation. *arXiv preprint arXiv:1703.05849*, 2017.
- J. Robins. A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Mathematical Modelling*, 7(9-12):1393–1512, 1986.
- J. M. Robins, M. A. Hernan, and B. Brumback. Marginal structural models and causal inference in epidemiology. *Epidemiology*, 11(5):550–560, 2000.
- D. Rothenhäusler and B. Yu. Incremental causal effects. *arXiv preprint arXiv:1907.13258*, 2019.
- D. B. Rubin. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688–701, 1974.
- A. Schick. On Asymptotically Efficient Estimation in Semiparametric Models. *The Annals of Statistics*, 14(3):1139 – 1151, 1986.
- K. Schindl, S. Shen, and E. H. Kennedy. Incremental effects for continuous exposures. *arXiv preprint arXiv:2409.11967*, 2024.
- P. Schnell and G. Papadogeorgou. Mitigating unobserved spatial confounding when estimating the effect of supermarket access on cardiovascular disease deaths. *Annals of Applied Statistics*, 14: 2069–2095, 12 2020.
- P. Z. Schochet, J. Burghardt, and S. Glazerman. National job corps study: The impacts of job corps on participants' employment and related outcomes. Mathematica policy research reports, Mathematica Policy Research, 2001.
- P. Z. Schochet, J. Burghardt, and S. McConnell. Does job corps work? impact findings from the national job corps study. *American Economic Review*, 98(5):1864–1886, 2008.
- J. Shao. *Mathematical Statistics*. Springer Science & Business Media, 2003.

- R. M. Stolzenberg. The measurement and decomposition of causal effects in nonlinear and nonadditive models. *Sociological Methodology*, 11:459–488, 1980.
- C. J. Stone. Additive regression and other nonparametric models. *The Annals of Statistics*, 13(2):689–705, 1985.
- L. Su, T. Ura, and Y. Zhang. Non-separable models with high-dimensional data. *Journal of Econometrics*, 212(2):646–677, 2019.
- A. Swaminathan and T. Joachims. The self-normalized estimator for counterfactual learning. In C. Cortes, N. Lawrence, D. Lee, M. Sugiyama, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 28, 2015.
- K. Takatsu and T. Westling. Debiased inference for a covariate-adjusted regression function. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, page qkae041, 2024.
- H. F. Trotter and J. W. Tukey. Conditional monte carlo for normal samples. In *Symposium on Monte Carlo Methods*, pages 64–79. John Wiley and Sons, 1956.
- A. B. Tsybakov. On nonparametric estimation of density level sets. *The Annals of Statistics*, 25(3):948–969, 1997.
- M. J. van der Laan and J. M. Robins. *Unified methods for censored longitudinal data and causality*. Springer, 2003.
- M. J. van der Laan, A. Bibaut, and A. R. Luedtke. Cv-tml for nonpathwise differentiable target parameters. In M. J. van der Laan and S. Rose, editors, *Targeted Learning in Data Science: Causal Inference for Complex Longitudinal Studies*, pages 455–481. Springer, 2018.
- A. van der Vaart. On differentiable functionals. *The Annals of Statistics*, 19(1):178–204, 1991.
- A. W. van der Vaart. *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 1998.
- M. P. Wand and M. C. Jones. *Kernel Smoothing*. CRC press, 1994.
- L. Wasserman. *All of nonparametric statistics*. Springer Science & Business Media, 2006.
- D. Westreich and S. R. Cole. Invited commentary: positivity in practice. *American Journal of Epidemiology*, 171(6):674–677, 2010.
- X. Wu, F. Mealli, M.-A. Kioumourtzoglou, F. Dominici, and D. Braun. Matching on generalized propensity scores with continuous exposures. *Journal of the American Statistical Association*, 119(545):757–772, 2024.
- Y. Zhang, Y.-C. Chen, and A. Giessing. Nonparametric inference on dose-response curves without the positivity condition. *arXiv preprint arXiv:2405.09003*, 2024.
- S. Zhou and D. A. Wolfe. On derivative estimation in spline regression. *Statistica Sinica*, 10(1):93–108, 2000.

Supplementary Materials to “Nonparametric Doubly Robust Inference on Derivative of Dose-Response Curve: With and Without Positivity”

Contents

A Practical Considerations	2
A.1 Self-Normalized IPW Estimators	2
A.2 Implementation of Proposed Estimators in Section 3 with Cross-Fitting	4
A.3 Nuisance Function Estimation	5
B Additional Simulation Results	7
B.1 Simulation Studies With Positivity Across Different Bandwidth Choices	7
B.2 Simulation Studies With Positivity and No Cross-Fitting	7
B.3 Simulation Studies Without Positivity Across Different Bandwidth Choices	8
B.4 Simulation Studies Without Positivity and No Cross-Fitting	9
B.5 Analysis of the Job Corps Program With No Cross-fitted Estimators	10
C Identification of $m(t)$ and $\theta(t)$ Under the Additive Confounding Model (13)	11
D Asymptotic Differences Between Two Variants of IPW Estimators	12
D.1 Asymptotic Difference Between IPW Estimators (3) and (5) of $m(t)$	13
D.2 Asymptotic Difference Between IPW Estimators (7) and (8) of $\theta(t)$	14
E Consistency of Estimating $m(t)$ Under Positivity	17
E.1 Rate of Convergence of $\hat{m}_{\text{RA}}(t)$	18
E.2 Rate of Convergence of $\hat{m}_{\text{IPW}}(t)$	19
E.3 Asymptotic Properties and Double Robustness of $\hat{m}_{\text{DR}}(t)$	21
F Proof of Theorem 1	30
F.1 Rate of Convergence of $\hat{\theta}_{\text{RA}}(t)$	31
F.2 Rate of Convergence of $\hat{\theta}_{\text{IPW}}(t)$	32
F.3 Asymptotic Properties of $\hat{\theta}_{\text{DR}}(t)$	35
F.4 Asymptotic Bias of $\hat{\theta}_{\text{AIPW}}(t)$ in (9)	45
G Proofs of Propositions 2, 3, and 4	47
G.1 Proof of Proposition 2	47
G.2 Proof of Proposition 3	49
G.3 Proof of Proposition 4	51

H	Proof of Theorem 5	52
H.1	Rate of Convergence of $\widehat{\theta}_{C,RA}(t)$	53
H.2	Rate of Convergence of $\widehat{\theta}_{C,IPW}(t)$	54
H.3	Asymptotic Properties of $\widehat{\theta}_{C,DR}(t)$	57
I	Asymptotic Theory of Estimating $m(t)$ Without Positivity	67
I.1	Uniform Rate of Convergence of $\widehat{m}_{C,RA}(t)$	69
I.2	Uniform Rate of Convergence of $\widehat{m}_{C,IPW}(t)$	70
I.3	Asymptotic Properties of $\widehat{m}_{C,DR}(t)$	70

A Practical Considerations

In this section, we outline some practical aspects involved in implementing our proposed estimators of dose-response and derivative effect curves.

A.1 Self-Normalized IPW Estimators

The classical IPW estimators of $m(t)$ and $\theta(t)$ suffer from the variance blowup when some estimated conditional densities $\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i), i = 1, \dots, n$ are close to 0. While truncating these estimates in a threshold value can reduce the instability of IPW estimators (Branson et al., 2023; Wu et al., 2024), determining the appropriate threshold value in practice is not straightforward. Alternatively, one can reduce the variances of IPW estimators and maintain its consistency by implementing the self-normalized version of IPW estimators (Swaminathan and Joachims, 2015; Kallus and Zhou, 2018). This idea was originally from the importance sampling literature (Trotter and Tukey, 1956) and also known as Hájek estimator (Godambe and Joshi, 1965).

• **Estimators Under Positivity:** For the IPW estimator (3) of $m(t)$, its self-normalized version takes the form

$$\widehat{m}_{IPW}^{\text{norm}}(t) = \frac{\widehat{m}_{IPW}(t)}{\frac{1}{nh} \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}} = \frac{\sum_{i=1}^n \frac{Y_i \cdot K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}}{\sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}}. \quad (29)$$

The self-normalized IPW estimator (29) maintains the consistency of the original IPW estimator (3), because $\widehat{p}_{T|\mathbf{S}}$ is a consistent estimator of $p_{T|\mathbf{S}}$ and the (oracle) denominator of (29) has its expectation as:

$$\begin{aligned} \mathbb{E} \left[\frac{1}{nh} \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{p_{T|\mathbf{S}}(T_j|\mathbf{S}_j)} \right] &= \frac{1}{h} \int_{\mathcal{T} \times \mathcal{S}} \frac{K\left(\frac{t_1-t}{h}\right)}{p_{T|\mathbf{S}}(t_1|\mathbf{s}_1)} \cdot p(t_1, \mathbf{s}_1) d\mathbf{s}_1 dt_1 \\ &= \int_{\mathbb{R} \times \mathcal{S}} K(u) \cdot p_{\mathbf{S}}(\mathbf{s}_1) d\mathbf{s}_1 du = 1 \end{aligned}$$

under Assumption A5.

Similarly, for the IPW estimator (7) of $\theta(t)$, its self-normalized version can be written as:

$$\hat{\theta}_{\text{IPW}}^{\text{norm}}(t) = \frac{\hat{\theta}_{\text{IPW}}(t)}{\frac{1}{nh} \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}} = \frac{\sum_{i=1}^n \frac{Y_i \left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}}{\kappa_2 h \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}}. \quad (30)$$

This self-normalized technique can also be applied to the IPW component of the DR estimators (4) and (10) to stabilize their variances, leading to self-normalized DR estimators of $m(t)$ and $\theta(t)$ as:

$$\hat{m}_{\text{DR}}^{\text{norm}}(t) = \frac{\sum_{i=1}^n \frac{[Y_i - \hat{\mu}(t, \mathbf{S}_i)] K\left(\frac{T_i-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}}{\sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}} + \frac{1}{n} \sum_{i=1}^n \hat{\mu}(t, \mathbf{S}_i) \quad (31)$$

and

$$\hat{\theta}_{\text{DR}}^{\text{norm}}(t) = \frac{\sum_{i=1}^n \frac{[Y_i - \hat{\mu}(t, \mathbf{S}_i) - (T_i-t) \cdot \hat{\beta}(t, \mathbf{S}_i)] \left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}}{\kappa_2 h \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}} + \frac{1}{n} \sum_{i=1}^n \hat{\beta}(t, \mathbf{S}_i), \quad (32)$$

respectively. Compared to (11), the estimated asymptotic variance of the self-normalized DR estimator (32) thus becomes

$$\hat{V}_{\theta}(t) = \frac{1}{n} \sum_{i=1}^n \left\{ \frac{nh \cdot \phi_{h,t} \left(Y_i, T_i, \mathbf{S}_i; \hat{\mu}, \hat{\beta}, \hat{p}_{T|\mathbf{S}} \right)}{\sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right)}{\hat{p}_{T|\mathbf{S}}(T_j|\mathbf{S}_j)}} + \sqrt{h^3} \left[\hat{\beta}(t, \mathbf{S}_i) - \hat{\theta}_{\text{DR}}^{\text{norm}}(t) \right] \right\}^2. \quad (33)$$

• **Estimators Without Positivity:** For the bias-corrected IPW estimator (25) of $\theta(t)$, we also adopt the self-normalized technique to deduce that

$$\hat{\theta}_{\text{C,IPW}}^{\text{norm}}(t) = \frac{\hat{\theta}_{\text{C,IPW}}(t)}{\frac{1}{nh} \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right) \cdot \hat{p}_{\zeta}(\mathbf{S}_j|t)}{\hat{p}(T_j, \mathbf{S}_j)}} = \frac{\sum_{i=1}^n \frac{Y_i \left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right) \cdot \hat{p}_{\zeta}(\mathbf{S}_i|t)}{\hat{p}(T_i, \mathbf{S}_i)}}{\kappa_2 h \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right) \cdot \hat{p}_{\zeta}(\mathbf{S}_j|t)}{\hat{p}(T_j, \mathbf{S}_j)}}. \quad (34)$$

The self-normalized IPW estimator (34) again maintains the consistency of the original IPW estimator (25), because \hat{p}, \hat{p}_{ζ} are consistent estimators of p, \bar{p}_{ζ} respectively and the (oracle) denominator of (34) has its expectation as:

$$\begin{aligned} \mathbb{E} \left[\frac{1}{nh} \sum_{j=1}^n \frac{K\left(\frac{T_j-t}{h}\right) \cdot \bar{p}_{\zeta}(\mathbf{S}_j|t)}{p(T_j, \mathbf{S}_j)} \right] &= \frac{1}{h} \int_{\mathcal{T} \times \mathcal{S}} K\left(\frac{t_1-t}{h}\right) \cdot \bar{p}_{\zeta}(\mathbf{s}_1|t) d\mathbf{s}_1 dt_1 \\ &= \int_{\mathbb{R} \times \mathcal{S}} K(u) \cdot \bar{p}_{\zeta}(\mathbf{s}_1|t) d\mathbf{s}_1 du = 1 \end{aligned}$$

under Assumption A5.

Analogously, the self-normalized bias-corrected DR estimator of $\theta(t)$ is given by

$$\widehat{\theta}_{\text{C,DR}}^{\text{norm}}(t) = \frac{\sum_{i=1}^n \frac{[Y_i - \widehat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \widehat{\beta}(t, \mathbf{S}_i)] \left(\frac{T_i - t}{h}\right) K\left(\frac{T_i - t}{h}\right) \cdot \widehat{p}_{\zeta}(\mathbf{S}_i | t)}{\widehat{p}(T_i, \mathbf{S}_i)}}{\kappa_2 h \sum_{j=1}^n \frac{K\left(\frac{T_j - t}{h}\right) \cdot \widehat{p}_{\zeta}(\mathbf{S}_j | t)}{\widehat{p}(T_j, \mathbf{S}_j)}} + \int \widehat{\beta}(t, \mathbf{s}) \cdot \widehat{p}_{\zeta}(\mathbf{s} | t) d\mathbf{s}, \quad (35)$$

whose estimated asymptotic variance becomes

$$\widehat{V}_{C,\theta}(t) = \frac{1}{n} \sum_{i=1}^n \left\{ \frac{nh \cdot \phi_{C,h,t} \left(Y_i, T_i, \mathbf{S}_i; \widehat{\mu}, \widehat{\beta}, \widehat{p}, \widehat{p}_{\zeta} \right)}{\sum_{j=1}^n \frac{K\left(\frac{T_j - t}{h}\right) \cdot \widehat{p}_{\zeta}(\mathbf{S}_j | t)}{\widehat{p}(T_j, \mathbf{S}_j)}} + \sqrt{h^3} \left[\int \widehat{\beta}(t, \mathbf{s}) \cdot \widehat{p}_{\zeta}(\mathbf{s} | t) d\mathbf{s} - \widehat{\theta}_{\text{C,DR}}(t) \right] \right\}^2.$$

A.2 Implementation of Proposed Estimators in Section 3 with Cross-Fitting

We explain the implementation details for our proposed estimators of $\theta(t)$ in Section 3 with cross-fitting (Schick, 1986; Newey and Robins, 2018; Chernozhukov et al., 2018) as follows. The same procedures can be applied to the estimators of $m(t)$ in Section 2.1 as well.

1. Partitioning the Data: The observed data $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$ are partitioned into L distinct subsets of approximately equal size. Commonly, the 5-fold ($L = 5$) or 10-fold ($L = 10$) cross-fitting is applied in practice, and no cross-fitting is used when $L = 1$ by convention. Let $I_{\ell}, \ell = 1, \dots, L$ be the index sets of such a partition so that $\cup_{\ell=1}^L I_{\ell} = \{1, \dots, n\}$.

2. Estimating the Nuisance Functions: For each index set I_{ℓ} , we estimate the nuisance functions $\mu, \beta, p_{T|\mathbf{S}}$ using the observations that are *not* in I_{ℓ} ; see Section A.3 for details. The estimated nuisance functions are denoted by $\widehat{\mu}^{(\ell)}, \widehat{\beta}^{(\ell)}, \widehat{p}_{T|\mathbf{S}}^{(\ell)}$, respectively, for $\ell = 1, \dots, L$. Recall that $\mu(t, \mathbf{s})$ is the conditional mean outcome function, $\beta(t, \mathbf{s})$ is the partial derivative of $\mu(t, \mathbf{s})$ with respect to t , and $p_{T|\mathbf{S}}(t|\mathbf{s})$ is the conditional density function of T given $\mathbf{S} = \mathbf{s}$.

3. Constructing the Final Estimators: The RA, IPW, and DR estimators of $\theta(t)$ under cross-fitting are given by

$$\begin{aligned} \widehat{\theta}_{\text{RA}}(t) &= \frac{1}{n} \sum_{\ell=1}^L \sum_{i \in I_{\ell}} \widehat{\beta}^{(\ell)}(t, \mathbf{S}_i), \\ \widehat{\theta}_{\text{IPW}}(t) &= \frac{1}{nh^2} \sum_{\ell=1}^L \sum_{i \in I_{\ell}} \frac{Y_i \left(\frac{T_i - t}{h}\right) K\left(\frac{T_i - t}{h}\right)}{\kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}^{(\ell)}(T_i | \mathbf{S}_i)}, \\ \widehat{\theta}_{\text{DR}}(t) &= \frac{1}{nh} \sum_{\ell=1}^L \sum_{i \in I_{\ell}} \left\{ \frac{\left(\frac{T_i - t}{h}\right) K\left(\frac{T_i - t}{h}\right)}{h \cdot \kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}^{(\ell)}(T_i | \mathbf{S}_i)} \left[Y_i - \widehat{\mu}^{(\ell)}(t, \mathbf{S}_i) - (T_i - t) \cdot \widehat{\beta}^{(\ell)}(t, \mathbf{S}_i) \right] + h \cdot \widehat{\beta}^{(\ell)}(t, \mathbf{S}_i) \right\}, \end{aligned} \quad (36)$$

where the bandwidth parameter $h > 0$ is chosen beforehand. The estimated asymptotic variance of $\widehat{\theta}_{\text{DR}}(t)$ under cross-fitting is given by

$$\widehat{V}_{\theta}(t) = \frac{1}{n} \sum_{\ell=1}^L \sum_{i \in I_{\ell}} \left\{ \phi_{h,t} \left(Y_i, T_i, \mathbf{S}_i; \widehat{\mu}^{(\ell)}, \widehat{\beta}^{(\ell)}, \widehat{p}_{T|\mathbf{S}}^{(\ell)} \right) + \sqrt{h^3} \left[\widehat{\beta}^{(\ell)}(t, \mathbf{S}_i) - \widehat{\theta}_{\text{DR}}(t) \right] \right\}^2. \quad (37)$$

The self-normalized technique in [Section A.1](#) can be applied to these cross-fitted estimators accordingly.

A.3 Nuisance Function Estimation

The implementation of the DR estimator (10) of $\theta(t)$ in [Section 3](#) requires the estimation of three nuisance functions: (i) the conditional mean outcome function $\mu(t, \mathbf{s}) = \mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s})$; (ii) the partial derivative function $\beta(t, \mathbf{s}) = \frac{\partial}{\partial t} \mu(t, \mathbf{s})$; and (iii) the conditional density $p_{T|\mathbf{S}}(t|\mathbf{s})$. Below, we discuss how these nuisance functions are estimated in our numerical experiments of [Section 6](#).

• **Estimations of $\mu(t, \mathbf{s})$ and $\beta(t, \mathbf{s})$:** We apply a fully connected neural network model with two hidden layers of size 100×50 and use the sigmoid linear unit function $u \mapsto \frac{u}{1+e^{-u}}$ as the activation function to ensure the smoothness of resulting estimators $\widehat{\mu}(t, \mathbf{s})$ and $\widehat{\beta}(t, \mathbf{s})$. Other choices of the neural network architectures and activation functions also works for our proposed DR estimator (10). Theoretically, Theorem 1 in [Farrell et al. \(2021\)](#) and Section 3.1 in [Colangelo and Lee \(2020\)](#) discuss some regularity conditions under which our requirements on the rates of convergence are satisfied by neural network models. Practically, our neural network model is implemented via PyTorch ([Paszke et al., 2019](#)), and we use its automatic differentiation engine ([Paszke et al., 2017](#)) to compute the estimated partial derivative $\widehat{\beta}(t, \mathbf{s})$ from the fitted conditional mean outcome function $\widehat{\mu}(t, \mathbf{s})$. In contrast to numerical differentiation, automatic differentiation offers the key advantage of being hyperparameter-free and inherently accurate to working precision ([Baydin et al., 2018](#); [Blondel and Roulet, 2024](#)). Recently, it has been employed to compute the semi-parametric or non-parametric efficient influence function for any statistical functional ([Luedtke, 2024](#)).

• **Estimation of $p_{T|\mathbf{S}}(t|\mathbf{s})$:** Given the data-generating model (27), we consider two different methods for estimating the conditional density $p_{T|\mathbf{S}}(t|\mathbf{s})$ with kernel smoothing techniques.

1. *Method 1 (Kernel density estimation (KDE) on residuals):* Notice that the relationship between the covariate vector \mathbf{S} and the treatment variation variable E is additive in model (27), *i.e.*,

$$T = g_{\mathbf{S}}(\mathbf{S}) + g_E(E) \quad \text{with} \quad g_{\mathbf{S}}(\mathbf{S}) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{3\xi^T \mathbf{S}} \exp\left(-\frac{u^2}{2}\right) du - 0.5 \quad \text{and} \quad g_E(E) = 0.75E.$$

In addition, the regression function of T against \mathbf{S} is given by $\mathbb{E}(T|\mathbf{S} = \mathbf{s}) = g_{\mathbf{S}}(\mathbf{s})$. To estimate the regression function $g_{\mathbf{S}}$, we can apply any machine learning method to the data $\{(T_i, \mathbf{S}_i)\}_{i=1}^n$ using cross-fitting. In the actual implementation, we use a neural network model

with one hidden layer of size 20 and rectified linear unit function $u \mapsto \max\{0, u\}$ as the activation function. Based on the fitted regression function $\hat{g}_{\mathbf{S}}$, we construct an estimator of $p_{T|\mathbf{S}}(t|\mathbf{s})$ as:

$$\hat{p}_{T|\mathbf{S}}(t|\mathbf{s}) = \frac{1}{nh_e} \sum_{i=1}^n K_e \left[\frac{t - \hat{g}_{\mathbf{S}}(\mathbf{s}) - (T_i - \hat{g}_{\mathbf{S}}(\mathbf{S}_i))}{h_e} \right],$$

where the kernel function K_e and bandwidth $h_e > 0$ may differ from those in our DR estimator (10). It is worth noting that the observations $\{T_1 - \hat{g}_{\mathbf{S}}(\mathbf{S}_1), \dots, T_n - \hat{g}_{\mathbf{S}}(\mathbf{S}_n)\}$ are not i.i.d., necessitating additional analysis for asymptotic theory and bandwidth selection (Hart and Vieu, 1990; Meloche, 1990). For simplicity, we use the Epanechnikov kernel $K_e(u) = \frac{3}{4}(1 - |u|) \mathbb{1}_{\{|u| \leq 1\}}$ and choose the bandwidth via Silverman's rule of thumb as $\hat{h}_e = \left(\frac{4}{3}\right)^{\frac{1}{5}} \hat{\sigma}_e n^{-\frac{1}{5}}$, where $\hat{\sigma}_e$ is the sample standard deviation of $\{T_1 - \hat{g}_{\mathbf{S}}(\mathbf{S}_1), \dots, T_n - \hat{g}_{\mathbf{S}}(\mathbf{S}_n)\}$.

2. *Method 2 (Regression on kernel-smoothed outcomes (RKS))* The validity of Method 1 relies on the additive relation between \mathbf{S} and E in the model for T . Since this additive structure may not hold in general, we consider another kernel smoothing method for estimating $p_{T|\mathbf{S}}(t|\mathbf{s})$. Specifically, we estimate a kernel-smoothed regression function $g(t, \mathbf{s}) = \mathbb{E} \left[K_r \left(\frac{T-t}{h_r} \right) \mid \mathbf{S} = \mathbf{s} \right]$ by regressing kernel-smoothed outcomes $\left\{ K_r \left(\frac{T_i-t}{h_r} \right) \right\}_{i=1}^n$ against the covariate vectors $\{\mathbf{S}_i\}_{i=1}^n$ via any machine learning method. The fitted kernel-smoothed regression function $\hat{g}(t, \mathbf{s})$ is a consistent estimator of $p_{T|\mathbf{S}}(t|\mathbf{s})$ when the regression method is accurate, because

$$\begin{aligned} g(t, \mathbf{s}) &= \mathbb{E} \left[K_r \left(\frac{T-t}{h_r} \right) \mid \mathbf{S} = \mathbf{s} \right] \\ &= \int_{\mathcal{T}} K_r \left(\frac{t_1-t}{h_r} \right) p_{T|\mathbf{S}}(t_1|\mathbf{s}) dt_1 \\ &= \int_{\mathbb{R}} K_r(u) \cdot p_{T|\mathbf{S}}(t + uh_r|\mathbf{s}) du \\ &= \int_{\mathbb{R}} K_r(u) \left[p_{T|\mathbf{S}}(t|\mathbf{s}) + uh_r \cdot p'_{T|\mathbf{S}}(t|\mathbf{s}) + \frac{u^2 h_r^2}{2} \cdot p''_{T|\mathbf{S}}(t|\mathbf{s}) + o(h_r^3) \right] du \\ &= p_{T|\mathbf{S}}(t|\mathbf{s}) + O(h_r^2) \\ &\rightarrow p_{T|\mathbf{S}}(t|\mathbf{s}) \end{aligned}$$

as $h_r \rightarrow 0$ under Assumptions A4 and A5(a-b). In the actual implementation, we again use a neural network model with one hidden layer of size 20 and rectified linear unit function $u \mapsto \max\{0, u\}$ as the activation function. Here, the kernel function K_r and bandwidth $h_r > 0$ can be different from those in our DR estimator (10). To ensure a relatively large effective sample size for fitting g , we use the Gaussian kernel $K_r(u) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right)$ and choose the bandwidth by Silverman's rule of thumb as $\hat{h}_r = \left(\frac{4}{3}\right)^{\frac{1}{5}} \hat{\sigma}_T n^{-\frac{1}{5}}$, where $\hat{\sigma}_T$ is the sample standard deviation of $\{T_1, \dots, T_n\}$.

Besides that, Klosin (2021) proposed another method with kernel smoothing that directly esti-

mates the reciprocal $\frac{1}{p_{T|\mathcal{S}}(t|\mathbf{s})}$ of the conditional density using a minimum distance Lasso approach (Chernozhukov et al., 2022). This method employs polynomial basis functions of the covariate vector \mathbf{S} and a kernel-smoothed L_2 loss function. We briefly experimented with this approach and found that its performance and computational efficiency are inferior to the two methods above. In addition, this approach is very sensitive to the choice of its tuning parameter as shown in Section 6 of Klosin (2021). Thus, we choose not to report its results.

B Additional Simulation Results

This section provides supplementary simulation results assessing the impact of varying the bandwidth parameter on the performance of our proposed estimators of $\theta(t)$ and the finite-difference method by Colangelo and Lee (2020). Furthermore, we evaluate the finite-sample performances of our proposed estimators without cross-fitting in both simulation studies and the empirical analysis of the U.S. Job Corps Program dataset.

B.1 Simulation Studies With Positivity Across Different Bandwidth Choices

We follow the same data-generating process and experimental setup in Section 6.1 to evaluate the performances of different estimators of $\theta(t)$ under the positivity condition, varying the bandwidth parameter h . In line with the bandwidth choices in Colangelo and Lee (2020); Klosin (2021), we examine four scaling factors for the bandwidth parameter as $h = C_h \cdot \hat{\sigma}_T \cdot n^{-\frac{1}{5}}$ with $C_T \in \{0.75, 1, 1.25, 1.5\}$, where $\hat{\sigma}_T$ is the sample standard deviation of $\{T_1, \dots, T_n\}$. For supplementary purposes, we only present the simulation results with 5-fold cross-fitting when the sample size is $n = 4000$ in Figure 5. The results are mostly consistent with our findings in Section 6.1. Our proposed DR estimators, leveraging either true or KDE-estimated conditional densities, demonstrate lower estimation biases and superior empirical coverage probabilities for their confidence intervals compared to the finite-difference method of Colangelo and Lee (2020). At the same time, they maintain RMSEs that are comparable to the finite-difference method. This additional results further demonstrate the robustness of our proposed DR estimator (10) to variations in its bandwidth parameter.

B.2 Simulation Studies With Positivity and No Cross-Fitting

For exploratory purposes, we conduct additional simulations to compare the performances of our proposed estimators of $\theta(t)$ in Section 3 with the finite-difference method by Colangelo and Lee (2020) when cross-fitting is not employed.

We replicate the experimental setup in Section 6.1 to generate the simulation results shown in Figure 6 across various sample sizes without using any cross-fitting. When the sample size is small, our proposed DR estimators without cross-fitting exhibit lower RMSEs but higher estimation biases than the finite-difference method by Colangelo and Lee (2020), resulting in inferior empirical coverage probabilities for the associated confidence intervals. However, as the sample size increases,




Figure 5: Comparisons between our proposed estimators and the finite-difference approaches by Colangelo and Lee (2020) (“CL20”) under positivity and with 5-fold cross-fitting across different bandwidth values (h). Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different scaling factors for h .

the estimation biases of our DR estimators diminish, and the empirical coverage probabilities of their confidence intervals improve, ultimately surpassing the finite-sample performance of the finite-difference method by Colangelo and Lee (2020).

These results without cross-fitting again highlight the practical utility of our proposed estimators in Section 3 under the positivity condition. However, developing rigorous theoretical guarantees for these estimators without cross-fitting is beyond the scope of this paper and will be addressed in future work.

B.3 Simulation Studies Without Positivity Across Different Bandwidth Choices

We adopt the same data-generating process and experimental setup in Section 6.2 to evaluate the performances of different estimators of $\theta(t)$ under various choices of the bandwidth parameter without assuming the positivity condition. Specifically, we test four scaling factors for the bandwidth parameter as $h = C_h \cdot \hat{\sigma}_T \cdot n^{-\frac{1}{5}}$ with $C_T \in \{0.75, 1, 1.5, 2\}$, where $\hat{\sigma}_T$ is the sample standard deviation of $\{T_1, \dots, T_n\}$. For supplementary purposes, we only present the simulation results with 5-fold cross-fitting when the sample size is $n = 2000$ in Figure 7. Again, our proposed bias-corrected estimators of $\theta(t)$ demonstrate significant improvements by reducing bias and enhancing the em-




Figure 6: Comparisons between our proposed estimators and the finite-difference approaches by Colangelo and Lee (2020) (“CL20”) under positivity and without cross-fitting across various sample sizes. Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different values for n . This figure follows an identical simulation setup as Figure 2 but without using any cross-fitting.

empirical coverage probabilities of the resulting confidence intervals compared to their counterparts in Section 3 across various bandwidth parameter choices.

B.4 Simulation Studies Without Positivity and No Cross-Fitting

For exploratory purposes, we conduct additional simulations for our bias-corrected estimators of $\theta(t)$ in Section 5.1 when the positivity condition is violated and cross-fitting is not employed.

Using the same experimental setup described in Section 6.2, we generate simulation results, shown in Figure 8, across various sample sizes without using any cross-fitting. As expected, the estimation biases and RMSEs of our bias-corrected estimators improve as the sample size increases, consistently outperforming their standard counterparts. However, in comparison to the results obtained with 5-fold cross-fitting in Figure 3, the performance of our bias-corrected estimators without cross-fitting deteriorates, particularly in terms of the empirical coverage probabilities of the resulting confidence intervals. These results consolidate the need of cross-fitting for constructing our bias-corrected estimators as Theorem 5 suggests.




Figure 7: Comparisons between our bias-corrected estimators (NP) in Section 5.1 and their counterparts (P) under the violation of positivity and with 5-fold cross-fitting ($L = 5$) across different bandwidth values (h). Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different scaling factors for h .

B.5 Analysis of the Job Corps Program With No Cross-fitted Estimators

Finally, we explore the behaviors of our proposed DR estimator (10) and the finite-difference method by Colangelo and Lee (2020) when cross-fitting is not employed. Following the same analysis pipeline described in Section 6.3, but without employing cross-fitting, we produce the results shown in Figure 9. Interestingly, our DR estimator without cross-fitting reveals some distinct trends compared to its counterpart with 5-fold cross-fitting. Specifically, it suggests a positive impact on employment during the first 20 weeks (~ 800 hours), diminishing benefits after 23 weeks (~ 920 hours), and statistically significant negative effects beyond 43 weeks (~ 1720 hours). These trends align with prior research (*e.g.*, Figure 2 of Lee 2009), which documented short-term negative impacts of the program on employment propensities (104 weeks after the program assignment). However, it is worth mentioning that the analysis in Lee (2009) was based on a binary treatment variable of being in the program or not. Since we do not establish any theoretical guarantees for our DR estimator of $\theta(t)$ when cross-fitting is not applied in this paper, more thorough investigations are necessary in the future to substantiate these short-term negative impacts of the Job Corps program.




Figure 8: Comparisons between our bias-corrected estimators (NP) in Section 5.1 and their counterparts (P) in Section 3 under the violation of positivity and without cross-fitting across different sample sizes. Rows present estimation bias, RMSE, and coverage probability for each estimator of $\theta(t)$, while columns correspond to different values for n . This figure follows an identical simulation setup as in Figure 3 but without using any cross-fitting.

C Identification of $m(t)$ and $\theta(t)$ Under the Additive Confounding Model (13)

Proposition C.1 (Identifications of $m(t)$ and $\theta(t)$). *Suppose that Assumptions A1(a-c) and A4(c) holds under model (13). Then, for any $t \in \mathcal{T}$ with $p_{\mathcal{S}|T}(s|t) > 0$ for some $s \in \mathcal{S}$, we have that*

$$\theta(t) = \bar{m}'(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = t \right],$$

where $\mu(t, \mathbf{s}) = \mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s})$. If, in addition, the marginal support \mathcal{T} of $p_T(t)$ is connected, then

$$m(t) = \mathbb{E} \left[Y + \int_T^t \theta(\tilde{t}) d\tilde{t} \right] = \mathbb{E} \left\{ Y + \int_T^t \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = \tilde{t} \right] d\tilde{t} \right\}.$$

Proof of Proposition C.1. We first study the identification of $\theta(t)$. By (14), $\theta(t) = \bar{m}'(t)$, and the conditional mean outcome function $\mu(t, \mathbf{s}) = \mathbb{E}(Y|T = t, \mathbf{S} = \mathbf{s}) = \bar{m}(t) + \eta(\mathbf{s})$ is well-defined within the support \mathcal{J} of the joint density $p(t, \mathbf{s})$. In particular, for any $t \in \mathcal{T}$ with $p_{\mathcal{S}|T}(\tilde{\mathbf{s}}|t) > 0$ for some $\tilde{\mathbf{s}} \in \mathcal{S}$, we know that $p(t, \tilde{\mathbf{s}}) > 0$ so that $\frac{\partial}{\partial t} \mu(t, \tilde{\mathbf{s}}) = \bar{m}'(t)$ is also well-defined for these




Figure 9: Estimated derivative effect curves with 95% confidence intervals using our proposed estimators and the finite-difference approaches by Colangelo and Lee (2020) (“CL20”) without cross-fitting. The vertical red dotted lines mark the original treatment range [320, 1840] analyzed in Colangelo and Lee (2020). This figure follows an identical analysis pipeline as in Figure 4 but without using any cross-fitting.

$\tilde{\mathbf{s}} \in \mathcal{S}(t)$. Furthermore, under Assumption A4(c), the support $\mathcal{S}(t)$ of the conditional distribution $p_{\mathbf{S}|T}(\mathbf{s}|t)$ is non-degenerate (*i.e.*, has nonzero Lebesgue measure). Thus,

$$\theta(t) = \bar{m}'(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = t \right]$$

is valid.

As for the identification of $m(t)$, we apply the fundamental theorem of calculus and argue that

$$m(t) = m(T) + \int_T^t \theta(\tilde{t}) d\tilde{t}.$$

Taking the expectation over T yields that

$$m(t) = \mathbb{E} \left[m(T) + \int_T^t \theta(\tilde{t}) d\tilde{t} \right] = \mathbb{E} \left\{ Y + \int_T^t \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \Big| T = \tilde{t} \right] d\tilde{t} \right\},$$

where the second equality follows from the fact that $\mathbb{E}[m(T)] = \mathbb{E}[\bar{m}(T)] + \mathbb{E}[\eta(\mathbf{S})] = \mathbb{E}(Y)$ by (14). Here, the connectedness of \mathcal{T} ensures that the integration of $\theta(t)$ is only over the region where it is identifiable. When \mathcal{T} has multiple connected components, the integral formula (15) as well as the observations should be restricted to the connected component in which the point of interest $t \in \mathcal{T}$ lies. \square

D Asymptotic Differences Between Two Variants of IPW Estimators

In this section, we study the asymptotic differences between the IPW estimators when the inverse probability weights are evaluated at the sample points (T_i, \mathbf{S}_i) , $i = 1, \dots, n$ or at the (query) points

$(t, \mathbf{S}_i), i = 1, \dots, n$. Specifically, for estimating the dose-response curve $m(t)$, we have two variants of the IPW estimators as (3) and (5). Similarly, for estimating the derivative effect $\theta(t) = m'(t)$, we also consider two different versions of the IPW estimators as (7) and (8). For the sake of illustrations, we assume that the conditional density $p_{T|\mathbf{S}}$ is known and only consider the oracle IPW estimators.

D.1 Asymptotic Difference Between IPW Estimators (3) and (5) of $m(t)$

We define the difference between two oracle IPW estimators of $m(t)$ as:

$$\tilde{\Delta}_{\text{IPW},m}(t) = \tilde{m}_{\text{IPW},2}(t) - \tilde{m}_{\text{IPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S}_i)} - \frac{1}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \right] Y_i \cdot K \left(\frac{T_i - t}{h} \right). \quad (38)$$

Proposition D.1. *Suppose that Assumptions A3, A4, A5, and A2 hold under the general model (1). Then, for any fixed $t \in \mathcal{T}$, we have that*

$$\begin{aligned} \tilde{\Delta}_{\text{IPW},m}(t) &= h^2 \kappa_2 \cdot \mathbb{E} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{2p_{T|\mathbf{S}}(t|\mathbf{S})} \right] + O(h^3) + O_P \left(\sqrt{\frac{h}{n}} \right) \\ &= O(h^2) + O_P \left(\sqrt{\frac{h}{n}} \right) \end{aligned}$$

as $h \rightarrow 0$ and $n \rightarrow \infty$.

Proof of Proposition D.1. By Chebyshev's inequality, we know that

$$\begin{aligned} \tilde{\Delta}_{\text{IPW},m}(t) &= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] + \tilde{\Delta}_{\text{IPW},m}(t) - \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] \\ &= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] + O_P \left(\sqrt{\text{Var} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right]} \right). \end{aligned}$$

On one hand, we calculate that

$$\begin{aligned} &\mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] \\ &= \mathbb{E} \left\{ \frac{1}{h} \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right] Y \cdot K \left(\frac{T-t}{h} \right) \right\} \\ &= \mathbb{E} \left\{ \int_{\mathcal{T}} \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(t_1|\mathbf{S})} \right] \mu(t_1, \mathbf{S}) \cdot K \left(\frac{t_1-t}{h} \right) \cdot p_{T|\mathbf{S}}(t_1|\mathbf{S}) dt_1 \right\} \\ &\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \left[\frac{p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \mu(t+uh, \mathbf{S}) \cdot K(u) du \right\} \\ &\stackrel{(ii)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \left[\frac{uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + O(h^3) \right] \right. \\ &\quad \left. \times \left[\mu(t, \mathbf{S}) + uh \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S}) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + O(h^3) \right] K(u) du \right\} \end{aligned}$$

$$= h^2 \kappa_2 \cdot \mathbb{E} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{2p_{T|\mathbf{S}}(t|\mathbf{S})} \right] + O(h^3)$$

where (i) uses a change of variable $u = \frac{t_1 - t}{h}$ while (ii) applies Taylor's expansions on $p_{T|\mathbf{S}}$ and μ under Assumptions A3 and A4. On the other hand, we also compute that

$$\begin{aligned} & \text{Var} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] \\ &= \frac{1}{nh^2} \cdot \text{Var} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right] Y \cdot K \left(\frac{T-t}{h} \right) \right\} \\ &\stackrel{\text{(iii)}}{=} \frac{1}{nh^2} \cdot \mathbb{E} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right]^2 Y^2 K^2 \left(\frac{T-t}{h} \right) \right\} + O \left(\frac{h^4}{n} \right) \\ &\stackrel{\text{(iv)}}{=} \frac{1}{nh} \cdot \mathbb{E} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(t+uh|\mathbf{S})} \right]^2 [\mu(t+uh, \mathbf{S})^2 + \sigma^2] K^2(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} \\ &\stackrel{\text{(v)}}{=} \frac{1}{nh} \cdot \mathbb{E} \left\{ \frac{\left[uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^3) \right]^2}{p_{T|\mathbf{S}}^2(t|\mathbf{S}) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})} \cdot [\mu(t+uh, \mathbf{S})^2 + \sigma^2] K^2(u) du \right\} \\ &= \frac{h}{n} \cdot \nu_2 \cdot \mathbb{E} \left\{ \frac{\left[\frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 [\mu(t, \mathbf{S})^2 + \sigma^2]}{p_{T|\mathbf{S}}(t|\mathbf{S})} \right\} + O \left(\frac{h^2}{n} \right) \end{aligned}$$

where (iii) leverages our above calculation on $\mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] = O(h^2)$, (iv) applies a change of variable under model (1), and (v) utilizes Taylor's expansion on $p_{T|\mathbf{S}}$ under Assumption A4.

In total, we conclude that

$$\begin{aligned} \tilde{\Delta}_{\text{IPW},m}(t) &= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right] + O_P \left(\sqrt{\text{Var} \left[\tilde{\Delta}_{\text{IPW},m}(t) \right]} \right) \\ &= h^2 \kappa_2 \cdot \mathbb{E} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{2p_{T|\mathbf{S}}(t|\mathbf{S})} \right] + O(h^3) + O_P \left(\sqrt{\frac{h}{n}} \right) \\ &= O(h^2) + O_P \left(\sqrt{\frac{h}{n}} \right) \end{aligned}$$

as $h \rightarrow 0$ and $n \rightarrow \infty$. The result follows. \square

D.2 Asymptotic Difference Between IPW Estimators (7) and (8) of $\theta(t)$

We define the difference between two oracle IPW estimators of $\theta(t)$ as:

$$\tilde{\Delta}_{\text{IPW},\theta}(t) = \tilde{\theta}_{\text{IPW},2}(t) - \tilde{\theta}_{\text{IPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S}_i)} - \frac{1}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \right] \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right)}{\kappa_2}. \quad (39)$$

Proposition D.2. *Suppose that Assumptions A3, A4, A5, and A2 hold under the general model*

(1). Then, for any fixed $t \in \mathcal{T}$, we have that

$$\begin{aligned}
\tilde{\Delta}_{\text{IPW},\theta}(t) &= \mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right] \\
&\quad + \frac{h^2 \kappa_4}{2\kappa_2} \cdot \mathbb{E} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S})}{3p_{T|\mathbf{S}}(t|\mathbf{S})} \right] \\
&\quad + O(h^3) + O_P \left(\sqrt{\frac{1}{nh}} \right) \\
&= \mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right] + O(h^2) + O_P \left(\sqrt{\frac{1}{nh}} \right)
\end{aligned}$$

as $h \rightarrow 0$ and $nh \rightarrow \infty$.

Proof of Proposition D.2. By Chebyshev's inequality, we know that

$$\begin{aligned}
\tilde{\Delta}_{\text{IPW},\theta}(t) &= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] + \tilde{\Delta}_{\text{IPW},\theta}(t) - \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] \\
&= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] + O_P \left(\sqrt{\text{Var} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right]} \right).
\end{aligned}$$

On one hand, we calculate that

$$\begin{aligned}
&\mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] \\
&= \mathbb{E} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right] \frac{Y \left(\frac{T-t}{h} \right) K \left(\frac{T-t}{h} \right)}{h \cdot \kappa_2} \right\} \\
&= \mathbb{E} \left\{ \int_{\mathcal{T}} \left[\frac{p_{T|\mathbf{S}}(t_1|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \frac{\mu(t_1, \mathbf{S}) \left(\frac{t_1-t}{h} \right) K \left(\frac{t_1-t}{h} \right)}{h \cdot \kappa_2} dt_1 \right\} \\
&\stackrel{\text{(i)}}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \left[\frac{p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \frac{\mu(t+uh, \mathbf{S}) \cdot u \cdot K(u)}{h \cdot \kappa_2} du \right\} \\
&\stackrel{\text{(ii)}}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \left[\frac{uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4)}{p_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right. \\
&\quad \left. \times \left[\mu(t, \mathbf{S}) + uh \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S}) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + O(h^3) \right] \frac{u \cdot K(u)}{h \cdot \kappa_2} du \right\} \\
&= \mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right] \\
&\quad + \frac{h^2 \kappa_4}{2\kappa_2} \cdot \mathbb{E} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S})}{3p_{T|\mathbf{S}}(t|\mathbf{S})} \right] + O(h^3),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ while (ii) applies Taylor's expansions on $p_{T|\mathbf{S}}$ and μ

under Assumptions A3 and A4. On the other hand, we also compute that

$$\begin{aligned}
& \text{Var} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] \\
&= \frac{1}{nh^2} \cdot \text{Var} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right] \frac{Y \left(\frac{T-t}{h^2} \right) K \left(\frac{T-t}{h} \right)}{\kappa_2} \right\} \\
&= \frac{1}{nh^4} \cdot \mathbb{E} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right]^2 \frac{Y^2 \left(\frac{T-t}{h} \right)^2 K^2 \left(\frac{T-t}{h} \right)}{\kappa_2^2} \right\} + O \left(\frac{1}{n} \right) \\
&\stackrel{\text{(iii)}}{=} \frac{1}{nh^4} \cdot \mathbb{E} \left\{ \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(T|\mathbf{S})} \right]^2 \frac{[\mu(T, \mathbf{S})^2 + \sigma^2] \left(\frac{T-t}{h} \right)^2 K^2 \left(\frac{T-t}{h} \right)}{\kappa_2^2} \right\} + O \left(\frac{1}{n} \right) \\
&\stackrel{\text{(iv)}}{=} \frac{1}{nh^3} \cdot \mathbb{E} \left\{ \int_{\mathbb{R}} \left[\frac{1}{p_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{1}{p_{T|\mathbf{S}}(t+uh|\mathbf{S})} \right]^2 \frac{[\mu(t+uh, \mathbf{S})^2 + \sigma^2] u^2 K^2(u)}{\kappa_2^2} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} + O \left(\frac{1}{n} \right) \\
&\stackrel{\text{(v)}}{=} \frac{1}{nh^3} \cdot \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{\left[uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right]^2}{p_{T|\mathbf{S}}^2(t|\mathbf{S}) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})} \right. \\
&\quad \left. \times \frac{[\mu(t+uh, \mathbf{S})^2 + \sigma^2] u^2 K^2(u)}{\kappa_2^2} du \right\} + O \left(\frac{1}{n} \right) \\
&= \frac{\nu_4}{nh \cdot \kappa_2^2} \cdot \mathbb{E} \left\{ \frac{\left[\frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 [\mu(t, \mathbf{S})^2 + \sigma^2]}{p_{T|\mathbf{S}}(t|\mathbf{S})} \right\} + O \left(\frac{1}{n} \right),
\end{aligned}$$

where (iii) leverages our above calculation on $\mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] = O(1)$, (iv) applies a change of variable under model (1), and (v) utilizes Taylor's expansion on $p_{T|\mathbf{S}}$ under Assumption A4.

In total, we conclude that

$$\begin{aligned}
\tilde{\Delta}_{\text{IPW},\theta}(t) &= \mathbb{E} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right] + O_P \left(\sqrt{\text{Var} \left[\tilde{\Delta}_{\text{IPW},\theta}(t) \right]} \right) \\
&= \mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right] \\
&\quad + \frac{h^2 \kappa_4}{2\kappa_2} \cdot \mathbb{E} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S})}{p_{T|\mathbf{S}}(t|\mathbf{S})} + \frac{\mu(t, \mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S})}{3p_{T|\mathbf{S}}(t|\mathbf{S})} \right] \\
&\quad + O(h^3) + O_P \left(\sqrt{\frac{1}{nh}} \right) \\
&= \mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right] + O(h^2) + O_P \left(\sqrt{\frac{1}{nh}} \right)
\end{aligned}$$

as $h \rightarrow 0$ and $nh \rightarrow \infty$. The result follows. \square

Remark D.1. Given our convergence analysis for $\hat{\theta}_{\text{IPW}}(t)$ in Theorem 1, one can easily calculate that the non-vanishing bias term $\mathbb{E} \left[\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{S}) \right]$ for $\tilde{\Delta}_{\text{IPW},\theta}(t)$ in Proposition D.2 results from the IPW estimator $\tilde{\theta}_{\text{IPW},2}(t)$ in (8). Therefore, unless $\mu(t, \mathbf{s}) = 0$ or $\frac{\partial}{\partial t} \log p_{T|\mathbf{S}}(t|\mathbf{s}) = 0$,

the IPW estimator (8) of $\theta(t)$ is asymptotically biased and inconsistent.

E Consistency of Estimating $m(t)$ Under Positivity

In this section, we review and prove the consistency results of $\widehat{m}_{\text{RA}}(t)$, $\widehat{m}_{\text{IPW}}(t)$, and $\widehat{m}_{\text{DR}}(t)$ in (2), (3), (4) for estimating the dose-response curve $t \mapsto m(t) = \mathbb{E}[Y(t)]$ under the positivity condition.

Proposition E.1 (Consistency of Estimating $m(t)$ Under Positivity). *Suppose that Assumptions A1, A3, A4, A5, and A2 hold under the general model (1) and $\widehat{\mu}, \widehat{p}_{T|\mathbf{S}}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s})$ and $\bar{p}_{T|\mathbf{S}}(t|\mathbf{s})$ be fixed bounded functions to which $\widehat{\mu}(t, \mathbf{s})$ and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ converge under the rates of convergence as:*

$$\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n}) \quad \text{and} \quad \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n}),$$

where $\Upsilon_{1,n}, \Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh \rightarrow \infty$, we have that

$$\begin{aligned} \widehat{m}_{\text{RA}}(t) - m(t) &= O_P\left(\Upsilon_{1,n} + \|\bar{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} + \frac{1}{\sqrt{n}}\right), \\ \widehat{m}_{\text{IPW}}(t) - m(t) &= O(h^2) + O_P\left(\sqrt{\frac{1}{nh}} + \Upsilon_{2,n} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right). \end{aligned}$$

If, in addition, we assume that

- (a) $\bar{p}_{T|\mathbf{S}}$ satisfies Assumptions A4 and A2;
- (b) either (i) “ $\bar{\mu} = \mu$ ” or “ $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$ ” almost surely;
- (c) $\sqrt{nh} \|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = o_P(1)$,

then

$$\sqrt{nh} [\widehat{m}_{\text{DR}}(t) - m(t)] = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \psi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) + \sqrt{h} [\bar{\mu}(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S})]] \right\} + o_P(1)$$

when $nh^5 \rightarrow c_2$ for some finite number $c_2 \geq 0$, where $\psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) = \frac{K(\frac{T-t}{h})}{\sqrt{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})}} [Y - \bar{\mu}(t, \mathbf{S})]$ and

$$\sqrt{nh} [\widehat{m}_{\text{DR}}(t) - m(t) - h^2 B_m(t)] \xrightarrow{d} \mathcal{N}(0, V_m(t))$$

with $V_m(t) = \mathbb{E} \left[\psi_{h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) \right]$ and

$$B_m(t) = \begin{cases} \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{2 \frac{\partial}{\partial t} \mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + p_{T|\mathbf{S}}(t|\mathbf{S}) \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) - 2 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right\} & \text{when } \bar{\mu} = \mu, \\ \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \right] & \text{when } \bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}. \end{cases}$$

Remark E.1 (Uniform asymptotic theory for estimating $m(t)$). *If we assume that*

$$\begin{cases} \sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n}), \\ \sup_{t \in \mathcal{T}} \|\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n}), \\ \sqrt{nh} \sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \|\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}) - p_{T|\mathbf{S}}(t|\mathbf{S})\|_{L_2} = o_P(1), \end{cases}$$

then the pointwise convergence results in Proposition E.1 can be strengthened to the uniform ones; see our side notes in the proof below.

Proof of Proposition E.1. We derive the rates of convergence of $\widehat{m}_{\text{RA}}(t)$ given by (2) and $\widehat{m}_{\text{IPW}}(t)$ given by (3) in Section E.1 and Section E.2, respectively. We also prove the asymptotic linearity, double robustness, and asymptotic normality of $\widehat{m}_{\text{DR}}(t)$ given by (4) in Section E.3.

E.1 Rate of Convergence of $\widehat{m}_{\text{RA}}(t)$

Firstly, we derive the rate of convergence for $\widehat{m}_{\text{RA}}(t)$ in (2). Notice that

$$\begin{aligned} \widehat{m}_{\text{RA}}(t) - m(t) &= \frac{1}{n} \sum_{i=1}^n \widehat{\mu}(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &= \underbrace{\frac{1}{n} \sum_{i=1}^n [\widehat{\mu}(t, \mathbf{S}_i) - \bar{\mu}(t, \mathbf{S}_i)]}_{\text{Term I}} + \underbrace{\frac{1}{n} \sum_{i=1}^n [\bar{\mu}(t, \mathbf{S}_i) - \mu(t, \mathbf{S}_i)]}_{\text{Term II}} + \underbrace{\frac{1}{n} \sum_{i=1}^n \{\mu(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S}_i)]\}}_{\text{Term III}}. \end{aligned}$$

- **Term I:** By Markov's inequality (and Hölder's inequality), we know that

$$\begin{aligned} \text{Term I} &\leq \frac{1}{n} \sum_{i=1}^n |\widehat{\mu}(t, \mathbf{S}_i) - \bar{\mu}(t, \mathbf{S}_i)| \\ &= O_P(\mathbb{E}|\widehat{\mu}(t, \mathbf{S}_1) - \bar{\mu}(t, \mathbf{S}_1)|) = O_P(\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}) = O_P(\Upsilon_{1,n}). \end{aligned}$$

- **Term II:** We similarly derive that

$$\text{Term II} \leq \frac{1}{n} \sum_{i=1}^n |\bar{\mu}(t, \mathbf{S}_i) - \mu(t, \mathbf{S}_i)| = O_P(\|\bar{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2}).$$

- **Term III:** By the central limit theorem and the boundedness of $\mu(t, \mathbf{s})$ on $\mathcal{T} \times \mathcal{S}$ under Assumption A3, we know that

$$\text{Term III} = \frac{1}{n} \sum_{i=1}^n \{\mu(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S}_i)]\} = O_P\left(\frac{1}{\sqrt{n}}\right).$$

As a side note, under Assumption A3, we know that $|\mu(t_1, \mathbf{s}) - \mu(t_2, \mathbf{s})| \leq A_1|t_1 - t_2|$ for some absolute constant $A_1 > 0$. Together with the compactness of \mathcal{T} and Example 19.7 in van der Vaart

(1998), we also deduce that

$$\sup_{t \in \mathcal{T}} \left| \frac{1}{n} \sum_{i=1}^n \{\mu(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S}_i)]\} \right| = O_P\left(\frac{1}{\sqrt{n}}\right).$$

In summary, we conclude that

$$\widehat{m}_{\text{RA}}(t) - m(t) = O_P\left(\Upsilon_{1,n} + \|\bar{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} + \frac{1}{\sqrt{n}}\right).$$

E.2 Rate of Convergence of $\widehat{m}_{\text{IPW}}(t)$

Secondly, we derive the rate of convergence for $\widehat{m}_{\text{IPW}}(t)$ in (3). Note that

$$\begin{aligned} & \widehat{m}_{\text{IPW}}(t) - m(t) \\ &= \widetilde{m}_{\text{IPW}}(t) - m(t) + \widehat{m}_{\text{IPW}}(t) - \widetilde{m}_{\text{IPW}}(t) \\ &= \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i - \mathbb{E}[\mu(t, \mathbf{S}_1)]}_{\text{Term IV}} + \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i - \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i}_{\text{Term V}}, \end{aligned}$$

where $\widetilde{m}_{\text{IPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} Y_i$ is the oracle IPW estimator of $m(t)$ defined in (18). We shall handle **Term IV** and **Term V** in Section E.2.1 and Section E.2.2, respectively.

E.2.1 Rate of Convergence of Term IV for $\widehat{m}_{\text{IPW}}(t)$

Under model (1) as well as Assumptions A3 and A5, we calculate the bias of $\widetilde{m}_{\text{IPW}}(t)$ as:

$$\begin{aligned} & \mathbb{E}[\widetilde{m}_{\text{IPW}}(t)] - m(t) \\ &= \mathbb{E}\left[\frac{1}{h} \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i\right] - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &= \frac{1}{h} \int_{\mathcal{S} \times \mathcal{T}} \frac{K\left(\frac{t_1-t}{h}\right)}{p_{T|\mathbf{S}}(t_1|\mathbf{s}_1)} \cdot \mu(t_1, \mathbf{s}_1) \cdot p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &\stackrel{(i)}{=} \int_{\mathcal{S}} \int_{\mathbb{R}} K(u) \cdot \mu(t + uh, \mathbf{s}_1) \cdot p_{\mathcal{S}}(\mathbf{s}_1) duds_1 - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &\stackrel{(ii)}{=} \int_{\mathcal{S}} \int_{\mathbb{R}} K(u) \left[\mu(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(\tilde{t}, \mathbf{s}_1) \right] p_{\mathcal{S}}(\mathbf{s}_1) duds_1 - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &\stackrel{(iii)}{=} \int_{\mathcal{S}} \mu(t, \mathbf{s}_1) \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 - \mathbb{E}[\mu(t, \mathbf{S}_1)] + \int_{\mathcal{S}} \int_{\mathbb{R}} K(u) \cdot \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(\tilde{t}, \mathbf{s}_1) \cdot p_{\mathcal{S}}(\mathbf{s}_1) duds_1 \\ &= O(h^2), \end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$, (ii) applies Taylor's expansion with some \tilde{t} that lies between t and $t + uh$, and (iii) utilizes the properties of the second-order kernel function K . Similarly, we compute the variance of $\tilde{m}_{\text{IPW}}(t)$ as:

$$\begin{aligned}
& \text{Var} [\tilde{m}_{\text{IPW}}(t)] \\
&= \frac{1}{nh^2} \cdot \text{Var} \left[\frac{K \left(\frac{T_i-t}{h} \right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} Y_i \right] \\
&= \frac{1}{nh^2} \cdot \mathbb{E} \left[\frac{K^2 \left(\frac{T_i-t}{h} \right)}{[p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)]^2} \cdot Y_i^2 \right] - \frac{1}{nh^2} \left\{ \mathbb{E} \left[\frac{K \left(\frac{T_i-t}{h} \right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \right] \right\}^2 \\
&= \frac{1}{nh^2} \int_{\mathcal{S} \times \mathcal{T}} \frac{K^2 \left(\frac{t_1-t}{h} \right)}{[p_{T|\mathbf{S}}(t_1|\mathbf{s}_1)]^2} \cdot [\mu(t_1, \mathbf{s}_1)^2 + \sigma^2] \cdot p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 - \frac{\{\mathbb{E}[\mu(t, \mathbf{S}_1)]\}^2}{n} + O\left(\frac{h^2}{n}\right) \\
&\stackrel{\text{(i)}}{=} \frac{1}{nh} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K^2(u)}{p_{T|\mathbf{S}}(t+uh|\mathbf{s}_1)} \cdot [\mu(t+uh, \mathbf{s}_1)^2 + \sigma^2] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{\text{(ii)}}{=} \frac{1}{nh} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K^2(u)}{p_{T|\mathbf{S}}(t|\mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t'|\mathbf{s}_1)} \left[\mu(t, \mathbf{s}_1)^2 + 2uh \cdot \mu(t'', \mathbf{s}_1) \cdot \frac{\partial}{\partial t} \mu(t'', \mathbf{s}_1) + \sigma^2 \right] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 \\
&\quad + O\left(\frac{1}{n}\right) \\
&\stackrel{\text{(iii)}}{=} \frac{1}{nh} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K^2(u)}{p_{T|\mathbf{S}}(t|\mathbf{s}_1)} \cdot [\mu(t, \mathbf{s}_1)^2 + \sigma^2] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{\text{(iv)}}{=} O\left(\frac{1}{nh}\right),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ and the boundedness of $\mu(t, \mathbf{s})$, (ii) applies the Taylor's expansion under Assumptions [A3](#) and [A4](#) with t', t'' being two points between t and $t + uh$, (iii) absorbs the higher order terms to $O\left(\frac{1}{n}\right)$, and (iv) utilizes the properties of K under Assumption [A5](#) and the positivity condition (Assumption [A2](#)). Now, by Chebyshev's inequality and our above calculations, we obtain that

$$\begin{aligned}
\tilde{m}_{\text{IPW}}(t) - m(t) &= \tilde{m}_{\text{IPW}}(t) - \mathbb{E}[\tilde{m}_{\text{IPW}}(t)] + \mathbb{E}[\tilde{m}_{\text{IPW}}(t)] - m(t) \\
&= O_P\left(\sqrt{\text{Var}[\tilde{m}_{\text{IPW}}(t)]}\right) + O(h^2) \\
&= O_P\left(\sqrt{\frac{1}{nh}}\right) + O(h^2)
\end{aligned}$$

as $h \rightarrow 0$ and $nh \rightarrow \infty$. As a side note, under the VC-type condition on K (Assumption [A5\(c\)](#)), we can apply Theorem 2 in [Einmahl and Mason \(2005\)](#) to strengthen the above pointwise rate of convergence to the uniform one as:

$$\sup_{t \in \mathcal{T}} |\tilde{m}_{\text{IPW}}(t) - m(t)| = O_P\left(\sqrt{\frac{|\log h|}{nh}}\right) + O(h^2).$$

E.2.2 Rate of Convergence of Term V for $\widehat{m}_{\text{IPW}}(t)$

By direct calculations, we have that

Term V

$$\begin{aligned}
&= \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \left[\frac{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \right] \\
&= \frac{1}{nh} \sum_{i=1}^n \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \left\{ \frac{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) + \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - [p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)] - [\bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)]} \right\} \\
&\stackrel{(i)}{=} \left\{ \mathbb{E}[\mu(t, \mathbf{S})] + O(h^2) + O_P\left(\sqrt{\frac{1}{nh}}\right) \right\} \\
&\quad \times \frac{O_P\left(\sup_{|u-t|\leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} + \sup_{|u-t|\leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right)}{\inf_{(t,\mathbf{s})\in\mathcal{T}\times\mathcal{S}} p_{T|\mathbf{S}}(t|\mathbf{s}) - O_P\left(\sup_{|u-t|\leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} + \sup_{|u-t|\leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right)} \\
&= O_P\left(\Upsilon_{2,n} + \sup_{|u-t|\leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right) \left[O(1+h^2) + O_P\left(\sqrt{\frac{1}{nh}}\right) \right] \\
&= O_P\left(\Upsilon_{2,n} + \sup_{|u-t|\leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right)
\end{aligned}$$

as $h \rightarrow 0$ and $nh \rightarrow \infty$, where (i) utilizes our results for **Term IV** and Markov's inequality.

Combining our results for **Term IV** and **Term V** in [Section E.2.1](#) and [Section E.2.2](#), we conclude that

$$\widehat{m}_{\text{IPW}}(t) - m(t) = O(h^2) + O_P\left(\sqrt{\frac{1}{nh}} + \Upsilon_{2,n} + \sup_{|u-t|\leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right).$$

E.3 Asymptotic Properties and Double Robustness of $\widehat{m}_{\text{DR}}(t)$

Finally, we establish the double robustness and asymptotic properties of $\widehat{m}_{\text{DR}}(t)$ in (4). Some parts of the following proof are inspired by the proof of Theorem 3.1 in [Colangelo and Lee \(2020\)](#). Notice that under Assumption [A1](#),

$$\begin{aligned}
&\widehat{m}_{\text{DR}}(t) - m(t) \\
&= \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot [Y_i - \widehat{\mu}(t, \mathbf{S}_i)] + h \cdot \widehat{\mu}(t, \mathbf{S}_i) \right\} - \mathbb{E}[\mu(t, \mathbf{S})] \\
&= \mathbb{P}_n \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) - \mathbb{E}[\mu(t, \mathbf{S})] + \mathbb{P}_n [\Psi_{h,t}(Y, T, \mathbf{S}; \widehat{\mu}, \widehat{p}_{T|\mathbf{S}}) - \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}})]
\end{aligned}$$

$$\begin{aligned}
&= \underbrace{\mathbb{P}_n \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) - \mathbb{E}[\mu(t, \mathbf{S})]}_{\text{Term VI}} \\
&\quad + \underbrace{(\mathbb{P}_n - \mathbb{P})[\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})]}_{\text{Term VII}} + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S})] \right\}}_{\text{Term VIII}} \\
&\quad + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\}}_{\text{Term IX}} \\
&\quad + \underbrace{\mathbb{P}_n \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\}}_{\text{Term X}} \\
&\quad + \underbrace{\mathbb{P} \left\{ \left[1 - \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} + \mathbb{P} \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S})] \right\}}_{\text{Term XI}},
\end{aligned}$$

where $\Psi_{h,t}(Y, T, \mathbf{S}; \mu, p_{T|\mathbf{S}}) = \frac{K\left(\frac{T-t}{h}\right)}{h \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \mu(t, \mathbf{S})] + \mu(t, \mathbf{S})$. It remains to show that the dominating **Term VI** is of order $O(h^2) + O_P\left(\sqrt{\frac{1}{nh}}\right)$ in [Section E.3.1](#) and the remainder terms are of order $o_P\left(\sqrt{\frac{1}{nh}}\right)$ for any $t \in \mathcal{T}$ in [Section E.3.2](#), [Section E.3.3](#), [Section E.3.4](#), and [Section E.3.5](#). We shall also derive the asymptotic normality of $\hat{m}_{\text{DR}}(t)$ in [Section E.3.6](#).

E.3.1 Analysis of Term VI for $\hat{m}_{\text{DR}}(t)$

We analyze the variance and bias of **Term VI** separately as follows. Notice that

$$\begin{aligned}
&\text{Var}[\text{Term VI}] \\
&= \text{Var}[\mathbb{P}_n \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}})] \\
&= \frac{1}{nh^2} \text{Var} \left[\frac{K\left(\frac{T-t}{h}\right)}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S})] + h \cdot \bar{\mu}(t, \mathbf{S}) \right] \\
&\stackrel{(i)}{\lesssim} \frac{1}{nh^2} \text{Var} \left[\frac{K\left(\frac{T-t}{h}\right)}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S})] \right] + \frac{1}{n} \text{Var}[\bar{\mu}(t, \mathbf{S})] \\
&\stackrel{(ii)}{=} \frac{1}{nh^2} \cdot \mathbb{E} \left[\frac{K^2\left(\frac{T-t}{h}\right)}{\bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S})]^2 \right] + O\left(\frac{1}{n}\right) \\
&= \frac{1}{nh^2} \int_{\mathcal{S}} \int_{\mathcal{T}} \frac{K^2\left(\frac{t_1-t}{h}\right)}{\bar{p}_{T|\mathbf{S}}^2(t_1|\mathbf{s}_1)} \left\{ [\mu(t_1, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)]^2 + \sigma^2 \right\} p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{(iii)}{=} \frac{1}{nh} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K^2(u)}{\bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{s}_1)} \left\{ [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)]^2 + \sigma^2 \right\} p(t+uh, \mathbf{s}_1) dud\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{(iv)}{=} \frac{1}{nh} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K^2(u)}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{s}_1) + O(h^2)} \left\{ [\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)]^2 + O(h^2) + \sigma^2 \right\} [p(t, \mathbf{s}_1) + O(h)] dud\mathbf{s}_1 + O\left(\frac{1}{n}\right)
\end{aligned}$$

$$= O\left(\frac{1}{nh}\right),$$

where (i) uses Cauchy-Schwarz inequality on the covariance, (ii) uses the boundedness of $\bar{\mu}$ under Assumption A3 to derive the term $O\left(\frac{1}{n}\right)$, (iii) leverages a change of variable $u = \frac{t_1-t}{h}$, and (iv) applies the Taylor's expansion under Assumptions A3 and A4. In the above calculations, we also note from the line (i) that the second part $\bar{\mu}(t, \mathbf{S})$ of $\Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S})$ is of smaller order than the first term $\frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|S}(T|\mathbf{S})} [Y - \bar{\mu}(t, \mathbf{S})]$. Thus, we can only keep the first term in the final asymptotically linear form of $\hat{m}_{\text{DR}}(t)$. Now, by Chebyshev's inequality, we conclude that

$$\begin{aligned} (\mathbb{P}_n - \mathbb{P}) \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S}) &= O_P\left(\sqrt{\text{Var}\left[\mathbb{P}_n \Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S})\right]}\right) \\ &= O_P\left(\sqrt{\text{Var}\left[\frac{1}{\sqrt{h}} \cdot \mathbb{P}_n \psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S})\right]}\right) = O_P\left(\sqrt{\frac{1}{nh}}\right), \end{aligned}$$

where $\psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S}) = \frac{K\left(\frac{T-t}{h}\right)}{\sqrt{h \cdot \bar{p}_{T|S}(T|\mathbf{S})}} [Y - \bar{\mu}(t, \mathbf{S})]$. In addition, by direct calculations and Taylor's expansions, we derive that

Bias [Term VI]

$$\begin{aligned} &= \mathbb{P}\left[\Psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|S})\right] - \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &= \mathbb{E}\left[\frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|S}(T|\mathbf{S})} [Y - \bar{\mu}(t, \mathbf{S})]\right] + \mathbb{E}[\bar{\mu}(t, \mathbf{S}_1) - \mu(t, \mathbf{S}_1)] \\ &= \int_{\mathcal{S}} \int_{\mathcal{T}} \frac{K\left(\frac{t_1-t}{h}\right)}{h \cdot \bar{p}_{T|S}(t_1|\mathbf{s}_1)} [\mu(t_1, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)] p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 + \mathbb{E}[\bar{\mu}(t, \mathbf{S}_1) - \mu(t, \mathbf{S}_1)] \\ &= \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K(u)}{\bar{p}_{T|S}(t+uh|\mathbf{s}_1)} [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)] p(t+uh, \mathbf{s}_1) dud\mathbf{s}_1 + \mathbb{E}[\bar{\mu}(t, \mathbf{S}_1) - \mu(t, \mathbf{S}_1)] \\ &= \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{K(u) \left[\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) + uh \frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + O(h^3) \right]}{\bar{p}_{T|S}(t|\mathbf{s}_1) + uh \frac{\partial}{\partial t} \bar{p}_{T|S}(t|\mathbf{s}_1) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|S}(t|\mathbf{s}_1) + O(h^3)} \\ &\quad \times \left[p(t, \mathbf{s}_1) + uh \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}_1) + O(h^3) \right] dud\mathbf{s}_1 + \mathbb{E}[\bar{\mu}(t, \mathbf{S}_1) - \mu(t, \mathbf{S}_1)] \\ &= \int_{\mathcal{S}} \int_{\mathbb{R}} K(u) \left[\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) + uh \frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + O(h^3) \right] \\ &\quad \times \left[\frac{1}{\bar{p}_{T|S}(t|\mathbf{s}_1)} - \frac{uh \frac{\partial}{\partial t} \bar{p}_{T|S}(t|\mathbf{s}_1)}{\bar{p}_{T|S}^2(t|\mathbf{s}_1)} - \frac{u^2 h^2 \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|S}(t|\mathbf{s}_1)}{2\bar{p}_{T|S}^3(t|\mathbf{s}_1)} + \frac{u^2 h^2 \left[\frac{\partial}{\partial t} \bar{p}_{T|S}(t|\mathbf{s}_1) \right]^2}{\bar{p}_{T|S}^3(t|\mathbf{s}_1)} + O(h^3) \right] \\ &\quad \times \left[p(t, \mathbf{s}_1) + uh \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}_1) + O(h^3) \right] dud\mathbf{s}_1 + \mathbb{E}[\bar{\mu}(t, \mathbf{S}_1) - \mu(t, \mathbf{S}_1)] \\ &= \int_{\mathcal{S}} \frac{[\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)]}{\bar{p}_{T|S}(t|\mathbf{s}_1)} \cdot p(t, \mathbf{s}_1) d\mathbf{s}_1 + \int_{\mathcal{S}} [\bar{\mu}(t, \mathbf{s}_1) - \mu(t, \mathbf{s}_1)] p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 \end{aligned}$$

$$\begin{aligned}
& + \frac{h^2 \kappa_2}{2} \int_{\mathcal{S}} \left\{ \frac{\frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1)}{\bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1)} \cdot p(t, \mathbf{s}_1) + \frac{2 \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1) \right]^2 / \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1) - \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1)}{\bar{p}_{T|\mathcal{S}}^2(t|\mathbf{s}_1)} [\bar{\mu}(t, \mathbf{s}_1) - \mu(t, \mathbf{s}_1)] p(t, \mathbf{s}_1) \right. \\
& + \frac{[\bar{\mu}(t, \mathbf{s}_1) - \mu(t, \mathbf{s}_1)]}{\bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1)} \cdot \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}_1) - \frac{2 \left[\frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) \right] \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1) \right]}{\bar{p}_{T|\mathcal{S}}^2(t|\mathbf{s}_1)} \cdot p(t, \mathbf{s}_1) + \frac{2 \left[\frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) \right] \left[\frac{\partial}{\partial t} p(t, \mathbf{s}_1) \right]}{\bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1)} \\
& \left. - \frac{2 \left[\frac{\partial}{\partial t} p(t, \mathbf{s}_1) \right] \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1) \right] [\bar{\mu}(t, \mathbf{s}_1) - \mu(t, \mathbf{s}_1)]}{\bar{p}_{T|\mathcal{S}}^2(t|\mathbf{s}_1)} \right\} d\mathbf{s}_1 + o(h^2) \\
& = \int_{\mathcal{S}} \frac{[\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)] [p_{T|\mathcal{S}}(t|\mathbf{s}_1) - \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1)]}{\bar{p}_{T|\mathcal{S}}(t|\mathbf{s}_1) \cdot p_{T|\mathcal{S}}(t|\mathbf{s}_1)} \cdot p(t, \mathbf{s}_1) d\mathbf{s}_1 + h^2 B_m(t) + o(h^2) \\
& = \mathbb{E}_{\mathcal{S}} \left\{ \frac{[\mu(t, \mathcal{S}) - \bar{\mu}(t, \mathcal{S})] [p_{T|\mathcal{S}}(t|\mathcal{S}) - \bar{p}_{T|\mathcal{S}}(t|\mathcal{S})]}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right\} + h^2 B_m(t) + o(h^2),
\end{aligned}$$

where the complicated bias term $B_m(t)$ is given by

$$\begin{aligned}
B_m(t) & = \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left\{ \frac{[\mu(t, \mathcal{S}) - \bar{\mu}(t, \mathcal{S})]}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \left[\frac{2 \frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \cdot \frac{\partial}{\partial t} \log \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) - \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathcal{S}}(t|\mathcal{S})}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \cdot p_{T|\mathcal{S}}(t|\mathcal{S}) + \frac{\partial^2}{\partial t^2} p_{T|\mathcal{S}}(t|\mathcal{S}) \right. \right. \\
& \quad \left. \left. - \frac{2 \left[\frac{\partial}{\partial t} p_{T|\mathcal{S}}(t|\mathcal{S}) \right] \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \right]}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right] \right\} \\
& + \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left\{ \frac{\left[\bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathcal{S}) - 2 \left[\frac{\partial}{\partial t} \mu(t, \mathcal{S}) \right] \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \right] \right]}{\bar{p}_{T|\mathcal{S}}^2(t|\mathcal{S})} \cdot p_{T|\mathcal{S}}(t|\mathcal{S}) \right\} \\
& + \kappa_2 \cdot \mathbb{E}_{\mathcal{S}} \left[\frac{\frac{\partial}{\partial t} \mu(t, \mathcal{S}) \cdot \frac{\partial}{\partial t} p_{T|\mathcal{S}}(t|\mathcal{S})}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right]
\end{aligned}$$

and $\bar{p}(t, \mathbf{s}) = \bar{p}_{T|\mathcal{S}}(t|\mathbf{s}) \cdot p_{\mathcal{S}}(\mathbf{s})$. Under the condition that either $\bar{\mu} = \mu$ or $\bar{p}_{T|\mathcal{S}} = p_{T|\mathcal{S}}$, we have that

$$\mathbb{E}_{\mathcal{S}} \left\{ \frac{[\mu(t, \mathcal{S}) - \bar{\mu}(t, \mathcal{S})] [p_{T|\mathcal{S}}(t|\mathcal{S}) - \bar{p}_{T|\mathcal{S}}(t|\mathcal{S})]}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right\} = 0$$

and

$$\begin{aligned}
B_m(t) & = \begin{cases} \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left\{ \frac{\left[\bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathcal{S}) - 2 \left[\frac{\partial}{\partial t} \mu(t, \mathcal{S}) \right] \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \right] \right]}{\bar{p}_{T|\mathcal{S}}^2(t|\mathcal{S})} \cdot p_{T|\mathcal{S}}(t|\mathcal{S}) + \frac{2 \frac{\partial}{\partial t} \mu(t, \mathcal{S}) \cdot \frac{\partial}{\partial t} p_{T|\mathcal{S}}(t|\mathcal{S})}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right\} & \text{when } \bar{\mu} = \mu, \\ \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathcal{S}) \right] & \text{when } \bar{p}_{T|\mathcal{S}} = p_{T|\mathcal{S}}, \end{cases} \\
& = \begin{cases} \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left\{ \frac{2 \frac{\partial}{\partial t} \mu(t, \mathcal{S}) \cdot \frac{\partial}{\partial t} p_{T|\mathcal{S}}(t|\mathcal{S}) + p_{T|\mathcal{S}}(t|\mathcal{S}) \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathcal{S}) - 2 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathcal{S}}(t|\mathcal{S}) \cdot \frac{\partial}{\partial t} \mu(t, \mathcal{S}) \right]}{\bar{p}_{T|\mathcal{S}}(t|\mathcal{S})} \right\} & \text{when } \bar{\mu} = \mu, \\ \frac{\kappa_2}{2} \cdot \mathbb{E}_{\mathcal{S}} \left[\frac{\partial^2}{\partial t^2} \mu(t, \mathcal{S}) \right] & \text{when } \bar{p}_{T|\mathcal{S}} = p_{T|\mathcal{S}}. \end{cases}
\end{aligned}$$

As a result, as $h \rightarrow 0$ and $nh \rightarrow \infty$, we have that

$$\mathbf{Term VI} = \mathbb{P}_n \Psi_{h,t}(Y, T, \mathcal{S}; \bar{\mu}, \bar{p}_{T|\mathcal{S}}) - \mathbb{E}[\mu(t, \mathcal{S})]$$

$$\begin{aligned}
&= h^2 B_m(t) + o(h^2) + O_P \left(\sqrt{\frac{1}{nh}} \right) \\
&= O(h^2) + O_P \left(\sqrt{\frac{1}{nh}} \right).
\end{aligned}$$

As a side note, under some VC-type condition on the kernel function K (Einmahl and Mason, 2005), we can strengthen the above pointwise rate of convergence to the following uniform one as:

$$\sup_{t \in \mathcal{T}} |\mathbf{Term VI}| = O(h^2) + O_P \left(\sqrt{\frac{|\log h|}{nh}} \right);$$

see Theorem 4 in Einmahl and Mason (2005) for details.

E.3.2 Analysis of Term VII for $\hat{m}_{\text{DR}}(t)$

By Markov's inequality, we know that

$$\begin{aligned}
\sqrt{nh} \cdot \mathbf{Term VII} &= \sqrt{h} \cdot \mathbb{G}_n [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \\
&= O_P \left(\sqrt{h} \cdot \Upsilon_{1,n} \right) = o_P(1)
\end{aligned}$$

because $\mathbb{E} \left\{ h \cdot [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})]^2 \right\} = h \cdot \|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}^2 = O_P(h \cdot \Upsilon_{1,n}^2)$ and $\Upsilon_{1,n} \rightarrow 0$ as $n \rightarrow \infty$. As a side note, under Assumption A3 on $\bar{\mu}$ and $\hat{\mu}$, we know that the function $\mathbf{s} \mapsto \hat{\mu}(t, \mathbf{s}) - \bar{\mu}(t, \mathbf{s})$ is Lipschitz continuous with respect to $t \in \mathcal{T}$. Together with the compactness of \mathcal{T} and Example 19.7 in van der Vaart (1998), we can also deduce that

$$\sup_{t \in \mathcal{T}} \left| \sqrt{h} \cdot \mathbb{G}_n [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right| = O_P \left(\sqrt{h} \cdot \sup_{t \in \mathcal{T}} \|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} \right),$$

which will be $o_P(1)$ as well if $\sup_{t \in \mathcal{T}} \|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = o_P(1)$.

E.3.3 Analyses of Term VIII and Term IX for $\hat{m}_{\text{DR}}(t)$

The argument for showing **Term VIII** and **Term IX** to be $o_P \left(\sqrt{\frac{1}{nh}} \right)$ will be similar to the one for **Term VII** above. By Markov's inequality, we know that

$$\begin{aligned}
\sqrt{nh} \cdot \mathbf{Term VIII} &= \mathbb{G}_n \left[\frac{K \left(\frac{T-t}{h} \right)}{\sqrt{h}} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S})] \right] \\
&= O_P(\Upsilon_{2,n}) = o_P(1)
\end{aligned}$$

because

$$\mathbb{E} \left\{ \frac{K^2 \left(\frac{T-t}{h} \right)}{h} \cdot \frac{[\hat{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{\hat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S})]^2 \right\}$$

$$\begin{aligned}
&= \mathbb{E} \left\{ \frac{K^2 \left(\frac{T-t}{h} \right)}{h} \cdot \frac{[\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{\widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [(\mu(T, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}))^2 + \sigma^2] \right\} \\
&\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} K^2(u) \cdot \frac{[\widehat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [(\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}))^2 + \sigma^2] \right\} \\
&\stackrel{(ii)}{\lesssim} \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}^2 \\
&\stackrel{(iii)}{=} O_P(\Upsilon_{2,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{T-t}{h}$ in the integration, (ii) leverages the boundedness of $\mu, \bar{\mu}$ under Assumption A3 and the positivity condition (Assumption A2) on $\bar{p}_{T|\mathbf{S}}$, as well as (iii) applies $\sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n})$ with $\Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$. As a side note again, under the VC-type condition on the kernel function K (Einmahl and Mason, 2005) and $\sup_{t \in \mathcal{T}} \|\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n}) = o_P(1)$, we can strengthen the above pointwise rate of convergence to the following uniform result as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left[\frac{K \left(\frac{T-t}{h} \right)}{\sqrt{h}} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S})] \right] \right| = o_P(1).$$

Similarly, by Markov's inequality, we have that

$$\sqrt{nh} \cdot \mathbf{Term IX} = \mathbb{G}_n \left\{ \frac{K \left(\frac{T-t}{h} \right)}{\sqrt{h} \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S})] \right\} = O_P(\Upsilon_{1,n}) = o_P(1)$$

because

$$\begin{aligned}
\mathbb{E} \left\{ \frac{K^2 \left(\frac{T-t}{h} \right)}{h \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S})]^2 \right\} &= \mathbb{E} \left\{ \int_{\mathcal{T}} \frac{K^2 \left(\frac{t_1-t}{h} \right) \cdot p_{T|\mathbf{S}}(t_1|\mathbf{S})}{h \cdot \bar{p}_{T|\mathbf{S}}^2(t_1|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S})]^2 dt_1 \right\} \\
&\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K^2(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S})]^2 du \right\} \\
&\stackrel{(ii)}{\lesssim} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}^2 \\
&= O_P(\Upsilon_{1,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{t_1-t}{h}$ and (ii) leverages the boundedness of $p_{T|\mathbf{S}}$ under Assumption A3, the positivity condition (Assumption A2) on $\bar{p}_{T|\mathbf{S}}$, the boundedness condition on K under Assumption A5, as well as $\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n})$ with $\Upsilon_{1,n} \rightarrow 0$ as $n \rightarrow \infty$. In addition, if $\sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = o_P(1)$, then the above pointwise rate of convergence

can be strengthened to the uniform one as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\} \right| = o_P(1).$$

E.3.4 Analysis of Term X for $\hat{m}_{\text{DR}}(t)$

We first calculate that

$$\begin{aligned} & \mathbb{E} \left| \sqrt{nh} \cdot \mathbf{Term X} \right| \\ & \mathbb{E} \left| \sqrt{\frac{n}{h}} \cdot K\left(\frac{T-t}{h}\right) \cdot \frac{[\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(T|\mathbf{S})]}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right| \\ & \stackrel{(i)}{\leq} \sqrt{nh} \cdot \sqrt{\mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right) [\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{h \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \right\} \cdot \mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \cdot [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})]^2 \right\}} \\ & = \sqrt{nh} \cdot \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]^2}{\hat{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S})} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\ & \quad \times \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} K(u) \cdot [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\ & \lesssim \sqrt{nh} \sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \|\hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \\ & \stackrel{(ii)}{=} o_P(1), \end{aligned}$$

where (i) uses Cauchy-Schwarz inequality and (ii) leverages our assumption (c) on the doubly robust rate of convergence in the proposition statement. As a result, by Markov's inequality, we obtain that

$$\sqrt{nh} \cdot \mathbf{Term X} = \sqrt{\frac{n}{h}} \cdot \mathbb{P}_n \left\{ K\left(\frac{T-t}{h}\right) \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\} = o_P(1).$$

E.3.5 Analysis of Term XI for $\hat{m}_{\text{DR}}(t)$

By direct calculations under model (1) with some change of variable, we have that

Term XI

$$\begin{aligned} & = \mathbb{E} \left\{ \left[1 - \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} + \mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S})] \right\} \\ & = \mathbb{E} \left\{ \mathbb{E} \left[1 - \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \middle| \mathbf{S} \right] [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} \\ & \quad + \mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right) [\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(T|\mathbf{S})]}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\mu(T, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} \end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left\{ \underbrace{\left[1 - \int_{\mathbb{R}} \frac{K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} du \right]}_{\text{Term XIa}} [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} \\
&+ \mathbb{E} \left\{ \underbrace{\int_{\mathbb{R}} \frac{K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du}_{\text{Term XIb}} \right\}.
\end{aligned}$$

On one hand, when $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$, we know from Assumption A5 that **Term XIa** = 0 and

$$\begin{aligned}
\text{Term XIb} &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [p_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{\hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] du \right\} \\
&\lesssim \sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \\
&= o_P \left(\sqrt{\frac{1}{nh}} \right)
\end{aligned}$$

by the boundedness of $\mu, \bar{\mu}$ under Assumption A3, the positivity condition (Assumption A2), and our assumption (c) on the doubly robust rate of convergence in the proposition statement. Specifically, since $\|\hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} = O_P(1)$ when $\bar{\mu} \neq \mu$, our assumption (c) entails that $\sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$.

On the other hand, when $\bar{\mu} = \mu$, we know from Assumption A2 on $\bar{p}_{T|\mathbf{S}}$ and the boundedness of $p_{T|\mathbf{S}}$ by Assumption A4 that

$$\begin{aligned}
\text{Term XIa} &= \mathbb{E} \left\{ \left[1 - \int_{\mathbb{R}} \frac{K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} du \right] [\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} \\
&\lesssim \|\hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \\
&= o_P \left(\sqrt{\frac{1}{nh}} \right),
\end{aligned}$$

where we again argue from our assumption (c) on the doubly robust rate of convergence in the proposition statement that $\|\hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$ if $\bar{p}_{T|\mathbf{S}} \neq p_{T|\mathbf{S}}$ and

$$\sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = O_P(1).$$

In addition, we have that

$$\begin{aligned}
\text{Term XIb} &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \mu(t, \mathbf{S})] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} \\
&\leq \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}}
\end{aligned}$$

$$\begin{aligned}
& \times \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [\mu(t+uh, \mathbf{S}) - \mu(t, \mathbf{S})]^2}{\bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S})} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\
& \stackrel{(i)}{=} O_P \left(\sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \right) \\
& \times \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [uh \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{S}) + O(h^2)]^2 [p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^2)]^2}{\left[\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S}) + 2uh \cdot \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^2) \right]^2 \left[1 + O_P(\Upsilon_{1,n}^2) \right]} du \right\}} \\
& = O_P \left(h^2 \cdot \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \right) \\
& = O_P(h^2 \cdot \Upsilon_{2,n}) \\
& \stackrel{(ii)}{=} o_P \left(\sqrt{\frac{1}{nh}} \right),
\end{aligned}$$

where (i) applies Taylor's expansion and uses the fact that the difference between $\bar{p}_{T|\mathbf{S}}$ and $\widehat{p}_{T|\mathbf{S}}$ is small when $\sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n})$ as well as (i) leverages the arguments that $\sqrt{nh} \cdot h^2 = \sqrt{nh^5} \rightarrow \sqrt{c_2} \in [0, \infty)$ and $\Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$.

E.3.6 Asymptotic Normality of $\widehat{m}_{\text{DR}}(t)$

For the asymptotic normality of $\widehat{m}_{\text{DR}}(t)$, it follows from the Lyapunov central limit theorem. Specifically, we already show in [Section E.3.1](#) and subsequent subsections that

$$\begin{aligned}
\sqrt{nh} [\widehat{m}_{\text{DR}}(t) - m(t)] &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \psi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) + \sqrt{h} [\bar{\mu}(t, \mathbf{S}_i) - \mathbb{E}[\mu(t, \mathbf{S})]] \right\} + o_P(1) \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \psi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) + o_P(1)
\end{aligned}$$

with $\psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) = \frac{K(\frac{T-t}{h})}{\sqrt{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})}} [Y - \bar{\mu}(t, \mathbf{S})]$ and $V_m(t) = \mathbb{E} \left[\psi_{h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) \right] = O(1)$ by our calculation in [Term VI](#). Then, $\sum_{i=1}^n \text{Var} \left[\frac{1}{\sqrt{n}} \cdot \psi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) \right] = O(1)$ and

$$\begin{aligned}
& \sum_{i=1}^n \mathbb{E} \left| \frac{1}{\sqrt{n}} \cdot \psi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{p}_{T|\mathbf{S}}) \right|^{2+c_1} \\
&= \mathbb{E} \left| \frac{K^{2+c_1} \left(\frac{T-t}{h} \right) \cdot [Y - \bar{\mu}(t, \mathbf{S})]^{2+c_1}}{n^{\frac{c_1}{2}} h^{1+\frac{c_1}{2}} \cdot \bar{p}_{T|\mathbf{S}}^{2+c_1}(T|\mathbf{S})} \right| \\
&\lesssim \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K^{2+c_1}(u) \cdot \left[[\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})]^{2+c_1} + \mathbb{E}|\epsilon|^{2+c_1} \right]}{\sqrt{(nh)^{c_1}} \cdot \bar{p}_{T|\mathbf{S}}^{2+c_1}(t+uh|\mathbf{S})} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}
\end{aligned}$$

$$= O\left(\sqrt{\frac{1}{(nh)^{c_1}}}\right) = o(1)$$

by the boundedness of $\mu, \bar{\mu}, p_{T|\mathbf{S}}$, the positivity condition on $\bar{p}_{T|\mathbf{S}}$, the assumption that $\mathbb{E}|\epsilon|^{2+c_1} < \infty$, and the requirement that $nh \rightarrow \infty$ as $n \rightarrow \infty$. Hence, the Lyapunov condition holds, and we have that

$$\sqrt{nh} [\widehat{m}_{\text{DR}}(t) - m(t) - h^2 B_m(t)] \xrightarrow{d} \mathcal{N}(0, V_m(t))$$

after subtracting the dominating bias term $h^2 B_m(t)$ of $\psi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{p}_{T|\mathbf{S}})$ that we have computed in **Term VI**. The proof is thus completed. \square

F Proof of [Theorem 1](#)

Theorem 1 (Consistency of estimating $\theta(t)$ under positivity). *Suppose that Assumptions [A1](#), [A3](#), [A4](#), [A5](#), and [A2](#) hold under the general model (1) and $\widehat{\mu}, \widehat{\beta}, \widehat{p}_{T|\mathbf{S}}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s})$, $\bar{\beta}(t, \mathbf{s})$, and $\bar{p}_{T|\mathbf{S}}(t|\mathbf{s})$ be fixed bounded functions to which $\widehat{\mu}(t, \mathbf{s})$, $\widehat{\beta}(t, \mathbf{s})$ and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ converge under the rates of convergence as:*

$$\begin{aligned} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} &= O_P(\Upsilon_{1,n}), \quad \|\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{3,n}), \\ \text{and} \quad \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} &= O_P(\Upsilon_{2,n}), \end{aligned}$$

where $\Upsilon_{1,n}, \Upsilon_{3,n}, \Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we have that

$$\begin{aligned} \widehat{\theta}_{\text{RA}}(t) - \theta(t) &= O_P\left(\Upsilon_{3,n} + \|\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2} + \frac{1}{\sqrt{n}}\right), \\ \widehat{\theta}_{\text{IPW}}(t) - \theta(t) &= O(h^2) + O_P\left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{2,n} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right). \end{aligned}$$

If, in addition, we assume that

(a) $\bar{p}_{T|\mathbf{S}}$ satisfies Assumptions [A4](#) and [A2](#);

(b) either (i) “ $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ ” with only $h \cdot \Upsilon_{3,n} \rightarrow 0$ or (ii) “ $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$ ”;

(c) $\sqrt{nh} \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \left[\|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} + h \|\widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2} \right] = o_P(1)$,

then

$$\sqrt{nh^3} [\widehat{\theta}_{\text{DR}}(t) - \theta(t)] = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) + \sqrt{h^3} [\bar{\beta}(t, \mathbf{S}_i) - \mathbb{E}[\beta(t, \mathbf{S})]] \right\} + o_P(1)$$

when $nh^7 \rightarrow c_3$ for some finite number $c_3 \geq 0$, where

$$\phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})].$$

Furthermore,

$$\sqrt{nh^3} \left[\widehat{\theta}_{\text{DR}}(t) - \theta(t) - h^2 B_\theta(t) \right] \xrightarrow{d} \mathcal{N}(0, V_\theta(t))$$

with $V_\theta(t) = \mathbb{E} \left[\phi_{h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right]$ and

$$B_\theta(t) = \begin{cases} \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{3 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + p_{T|\mathbf{S}}(t|\mathbf{S}) \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) - 3 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right\} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) \right] & \text{when } \bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}. \end{cases}$$

Remark F.1 (Uniform asymptotic theory for estimating $\theta(t)$). *If we assume that*

$$\begin{cases} \sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n}), \\ \sup_{t \in \mathcal{T}} \|\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n}), \\ \sup_{t \in \mathcal{T}} \|\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{3,n}), \\ \sqrt{nh} \sup_{t \in \mathcal{T}} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \left[\|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} + h \|\widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2} \right] = o_P(1), \end{cases}$$

then the pointwise convergence results in [Theorem 1](#) can be strengthened to the uniform ones; see our side notes in the proof below.

Proof of [Theorem 1](#). We derive the rates of convergence of $\widehat{\theta}_{\text{RA}}(t)$ given by [\(6\)](#) and $\widehat{\theta}_{\text{IPW}}(t)$ given by [\(7\)](#) in [Section F.1](#) and [Section F.2](#), respectively. We also prove the asymptotic linearity, double robustness, and asymptotic normality of $\widehat{\theta}_{\text{DR}}(t)$ given by [\(10\)](#) in [Section F.3](#).

F.1 Rate of Convergence of $\widehat{\theta}_{\text{RA}}(t)$

Firstly, we derive the rate of convergence for $\widehat{\theta}_{\text{RA}}(t)$ in [\(6\)](#). Under Assumption [A1](#)(d), we have that

$$\begin{aligned} \widehat{\theta}_{\text{RA}}(t) - \theta(t) &= \frac{1}{n} \sum_{i=1}^n \widehat{\beta}(t, \mathbf{S}_i) - \frac{d}{dt} \mathbb{E}[\mu(t, \mathbf{S}_1)] \\ &= \underbrace{\frac{1}{n} \sum_{i=1}^n [\widehat{\beta}(t, \mathbf{S}_i) - \bar{\beta}(t, \mathbf{S}_i)]}_{\text{Term I}} + \underbrace{\frac{1}{n} \sum_{i=1}^n [\bar{\beta}(t, \mathbf{S}_i) - \beta(t, \mathbf{S}_i)]}_{\text{Term II}} + \underbrace{\frac{1}{n} \sum_{i=1}^n \{\beta(t, \mathbf{S}_i) - \mathbb{E}[\beta(t, \mathbf{S}_i)]\}}_{\text{Term III}}. \end{aligned}$$

- **Term I:** By Markov's inequality (and Hölder's inequality), we know that

$$\text{Term I} \leq \frac{1}{n} \sum_{i=1}^n \left| \widehat{\beta}(t, \mathbf{S}_i) - \bar{\beta}(t, \mathbf{S}_i) \right|$$

$$= O_P \left(\mathbb{E} \left| \widehat{\beta}(t, \mathbf{S}_1) - \bar{\beta}(t, \mathbf{S}_1) \right| \right) = O_P \left(\left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right) = O_P(\Upsilon_{3,n}).$$

• **Term II:** Analogously, we derive that

$$\mathbf{Term II} \leq \frac{1}{n} \sum_{i=1}^n |\bar{\beta}(t, \mathbf{S}_i) - \beta(t, \mathbf{S}_i)| = O_P \left(\left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right).$$

• **Term III:** By the central limit theorem and the boundedness of $\beta(t, \mathbf{s})$ on $\mathcal{T} \times \mathcal{S}$ under Assumption A3, we know that

$$\mathbf{Term III} = \frac{1}{n} \sum_{i=1}^n \{\beta(t, \mathbf{S}_i) - \mathbb{E}[\beta(t, \mathbf{S}_i)]\} = O_P \left(\frac{1}{\sqrt{n}} \right).$$

As a side note, under Assumption A3, we know that $|\beta(t_1, \mathbf{s}) - \beta(t_2, \mathbf{s})| \leq A_2|t_1 - t_2|$ for some absolute constant $A_2 > 0$. Together with the compactness of \mathcal{T} and Example 19.7 in van der Vaart (1998), we also deduce that

$$\sup_{t \in \mathcal{T}} \left| \frac{1}{n} \sum_{i=1}^n \{\beta(t, \mathbf{S}_i) - \mathbb{E}[\beta(t, \mathbf{S}_i)]\} \right| = O_P \left(\frac{1}{\sqrt{n}} \right).$$

In summary, we conclude that

$$\widehat{\theta}_{\text{RA}}(t) - \theta(t) = O_P \left(\Upsilon_{3,n} + \left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} + \frac{1}{\sqrt{n}} \right).$$

F.2 Rate of Convergence of $\widehat{\theta}_{\text{IPW}}(t)$

Secondly, we derive the rate of convergence for $\widehat{\theta}_{\text{IPW}}(t)$ in (7). Note that

$$\begin{aligned} & \widehat{\theta}_{\text{IPW}}(t) - \theta(t) \\ &= \widetilde{\theta}_{\text{IPW}}(t) - \theta(t) + \widehat{\theta}_{\text{IPW}}(t) - \widetilde{\theta}_{\text{IPW}}(t) \\ &= \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i - \mathbb{E}[\beta(t, \mathbf{S})]}_{\mathbf{Term IV}} + \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot \widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i - \frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i}_{\mathbf{Term V}}, \end{aligned}$$

where $\widetilde{\theta}_{\text{IPW}}(t) = \frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i$ is the oracle IPW estimator of $\theta(t)$ defined in (18) and $\beta(t, \mathbf{s}) = \frac{\partial}{\partial t} \mu(t, \mathbf{s})$. We shall handle **Term IV** and **Term V** in Section F.2.1 and Section F.2.2, respectively.

F.2.1 Rate of Convergence of Term IV for $\widehat{\theta}_{\text{IPW}}(t)$

Under model (1) as well as Assumptions A3 and A5, we calculate the bias of $\widetilde{\theta}_{\text{IPW}}(t)$ as:

$$\begin{aligned}
& \mathbb{E} \left[\widetilde{\theta}_{\text{IPW}}(t) \right] - \theta(t) \\
&= \mathbb{E} \left[\frac{1}{h} \frac{\left(\frac{T_i-t}{h^2} \right) K \left(\frac{T_i-t}{h} \right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \right] - \mathbb{E} [\beta(t, \mathbf{S})] \\
&= \frac{1}{h} \int_{\mathcal{S} \times \mathcal{T}} \frac{\left(\frac{t_1-t}{h^2} \right) K \left(\frac{t_1-t}{h} \right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(t_1|\mathbf{s}_1)} \cdot \mu(t_1, \mathbf{s}_1) \cdot p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 - \mathbb{E} [\beta(t, \mathbf{S})] \\
&\stackrel{(i)}{=} \frac{1}{h \cdot \kappa_2} \int_{\mathcal{S}} \int_{\mathbb{R}} u K(u) \cdot \mu(t+uh, \mathbf{s}_1) \cdot p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 - \mathbb{E} [\beta(t, \mathbf{S})] \\
&\stackrel{(ii)}{=} \frac{1}{h \cdot \kappa_2} \int_{\mathcal{S}} \int_{\mathbb{R}} u K(u) \left[\mu(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \frac{\partial^3}{\partial t^3} \mu(\tilde{t}, \mathbf{s}_1) \right] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 \\
&\quad - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] \\
&\stackrel{(iii)}{=} \int_{\mathcal{S}} \frac{\partial}{\partial t} \mu(t, \mathbf{s}_1) \cdot p_{\mathbf{S}}(\mathbf{s}_1) d\mathbf{s}_1 - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] + \int_{\mathcal{S}} \int_{\mathbb{R}} K(u) \cdot \frac{u^3 h^2}{6 \kappa_2} \cdot \frac{\partial^3}{\partial t^3} \mu(\tilde{t}, \mathbf{s}_1) \cdot p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 \\
&= O(h^2),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$, (ii) applies Taylor's expansion with some \tilde{t} that lies between t and $t+uh$, and (iii) utilizes the properties of the second-order symmetric kernel function K . Similarly, we compute the variance of $\widetilde{\theta}_{\text{IPW}}(t)$ as:

$$\begin{aligned}
& \text{Var} \left[\widetilde{\theta}_{\text{IPW}}(t) \right] \\
&= \frac{1}{nh^2 \kappa_2^2} \cdot \text{Var} \left[\frac{\left(\frac{T_i-t}{h^2} \right) K \left(\frac{T_i-t}{h} \right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} Y_i \right] \\
&= \frac{1}{nh^2 \kappa_2^2} \cdot \mathbb{E} \left[\frac{\left(\frac{T_i-t}{h^2} \right)^2 K^2 \left(\frac{T_i-t}{h} \right)}{\left[p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) \right]^2} \cdot Y_i^2 \right] - \frac{1}{nh^2 \kappa_2^2} \left\{ \mathbb{E} \left[\frac{\left(\frac{T_i-t}{h^2} \right) K \left(\frac{T_i-t}{h} \right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \right] \right\}^2 \\
&= \frac{1}{nh^4 \kappa_2^2} \int_{\mathcal{S} \times \mathcal{T}} \frac{\left(\frac{t_1-t}{h} \right)^2 K^2 \left(\frac{t_1-t}{h} \right)}{\left[p_{T|\mathbf{S}}(t_1|\mathbf{s}_1) \right]^2} \cdot [\mu(t_1, \mathbf{s}_1)^2 + \sigma^2] p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 - \frac{\{\mathbb{E} [\beta(t, \mathbf{S})]\}^2}{n} + O \left(\frac{h^2}{n} \right) \\
&\stackrel{(i)}{=} \frac{1}{nh^3 \kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u)}{p_{T|\mathbf{S}}(t+uh|\mathbf{s}_1)} \cdot [\mu(t+uh, \mathbf{s}_1)^2 + \sigma^2] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 + O \left(\frac{1}{n} \right) \\
&\stackrel{(ii)}{=} \frac{1}{nh^3 \kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u)}{p_{T|\mathbf{S}}(t|\mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t'|\mathbf{s}_1)} \left[\mu(t, \mathbf{s}_1)^2 + 2uh \cdot \mu(t'', \mathbf{s}_1) \cdot \frac{\partial}{\partial t} \mu(t'', \mathbf{s}_1) + \sigma^2 \right] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 \\
&\quad + O \left(\frac{1}{n} \right) \\
&\stackrel{(iii)}{=} \frac{1}{nh^3 \kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u)}{p_{T|\mathbf{S}}(t|\mathbf{s}_1)} \cdot [\mu(t, \mathbf{s}_1)^2 + \sigma^2] p_{\mathbf{S}}(\mathbf{s}_1) dud\mathbf{s}_1 + O \left(\frac{1}{n} \right)
\end{aligned}$$

$$\stackrel{\text{(iv)}}{=} O\left(\frac{1}{nh^3}\right),$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ and the boundedness of $\beta(t, \mathbf{s})$, (ii) applies the Taylor's expansion under Assumptions A3 and A4 with t', t'' being two points between t and $t + uh$, (iii) absorbs the higher order terms to $O\left(\frac{1}{n}\right)$, and (iv) utilizes the properties of K under Assumption A5 and the positivity condition (Assumption A2). Now, by Chebyshev's inequality and our above calculations, we obtain that

$$\begin{aligned} \tilde{\theta}_{\text{IPW}}(t) - \theta(t) &= \tilde{\theta}_{\text{IPW}}(t) - \mathbb{E}\left[\tilde{\theta}_{\text{IPW}}(t)\right] + \mathbb{E}\left[\tilde{\theta}_{\text{IPW}}(t)\right] - \theta(t) \\ &= O_P\left(\sqrt{\text{Var}\left[\tilde{\theta}_{\text{IPW}}(t)\right]}\right) + O(h^2) \\ &= O_P\left(\sqrt{\frac{1}{nh^3}}\right) + O(h^2) \end{aligned}$$

as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$. As a side note, under the VC-type condition on K (Assumption A5(c)), we can apply Theorem 2 in Einmahl and Mason (2005) to strengthen the above pointwise rate of convergence to the uniform one as:

$$\sup_{t \in \mathcal{T}} \left| \tilde{\theta}_{\text{IPW}}(t) - \theta(t) \right| = O_P\left(\sqrt{\frac{|\log h|}{nh^3}}\right) + O(h^2).$$

F.2.2 Rate of Convergence of Term V for $\hat{\theta}_{\text{IPW}}(t)$

By direct calculations, we have that

Term V

$$\begin{aligned} &= \frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \left[\frac{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}{\hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \right] \\ &= \frac{1}{nh} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h^2}\right) K\left(\frac{T_i-t}{h}\right)}{\kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \left\{ \frac{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) + \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - [p_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)] - [\bar{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i) - \hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)]} \right\} \\ &\stackrel{\text{(i)}}{=} \left\{ \mathbb{E}[\beta(t, \mathbf{S})] + O(h^2) + O_P\left(\sqrt{\frac{1}{nh^3}}\right) \right\} \\ &\quad \times \frac{O_P\left(\sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right)}{\inf_{(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}} p_{T|\mathbf{S}}(t|\mathbf{s}) - O_P\left(\sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right)} \\ &= O_P\left(\Upsilon_{2,n} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}\right) \left[O(1+h^2) + O_P\left(\sqrt{\frac{1}{nh^3}}\right) \right] \end{aligned}$$

$$= O_P \left(\Upsilon_{2,n} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \right)$$

as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, where (i) utilizes our results for **Term IV** and Markov's inequality.

Combining our results for **Term IV** and **Term V**, we conclude that

$$\hat{\theta}_{\text{IPW}}(t) - \theta(t) = O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{2,n} + \sup_{|u-t| \leq h} \|\bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \right).$$

F.3 Asymptotic Properties of $\hat{\theta}_{\text{DR}}(t)$

Finally, using the similar arguments to [Section E.3](#), we establish the asymptotic properties of $\hat{\theta}_{\text{DR}}(t)$ in (10). Under Assumption [A1](#), we have that

$$\begin{aligned} & \hat{\theta}_{\text{DR}}(t) - \theta(t) \\ &= \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{\left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{h \cdot \kappa_2 \cdot \hat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \left[Y_i - \hat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \hat{\beta}(t, \mathbf{S}_i) \right] + h \cdot \hat{\beta}(t, \mathbf{S}_i) \right\} - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] \\ &= \mathbb{P}_n \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] + \mathbb{P}_n \left[\Phi_{h,t}(Y, T, \mathbf{S}; \hat{\mu}, \hat{\beta}, \hat{p}_{T|\mathbf{S}}) - \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right] \\ &= \underbrace{\mathbb{P}_n \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \mathbb{E}[\beta(t, \mathbf{S})]}_{\text{Term VI}} + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left[\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right]}_{\text{Term VII}} \\ &+ \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\}}_{\text{Term VIII}} \\ &+ \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) \left[\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S}) \right] \right] \right\}}_{\text{Term IX}} \\ &+ \underbrace{\mathbb{P}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) \left[\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S}) \right] \right] \right\}}_{\text{Term X}} \\ &+ \underbrace{\mathbb{P} \left\{ \left[1 - \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] \left[\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \right\}}_{\text{Term XIa}} + \underbrace{\mathbb{P} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\}}_{\text{Term XIb}} \\ &+ \underbrace{\mathbb{P} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\}}_{\text{Term XIc}}, \end{aligned}$$

where $\Phi_{h,t}(Y, T, \mathbf{S}; \mu, \beta, p_{T|S}) = \frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{h^2\kappa_2 \cdot \bar{p}_{T|S}(T|\mathbf{S})} \cdot [Y - \mu(t, \mathbf{S}) - (T-t) \cdot \beta(t, \mathbf{S})] + \beta(t, \mathbf{S})$. It remains to show that the dominating **Term VI** is of order $O(h^2) + O_P\left(\sqrt{\frac{1}{nh^3}}\right)$ in [Section F.3.1](#) and the remainder terms are of order $o_P\left(\sqrt{\frac{1}{nh^3}}\right)$ for any fixed $t \in \mathcal{T}$ in [Section F.3.2](#), [Section F.3.3](#), and [Section F.3.4](#), and [Section F.3.5](#). We shall also derive the asymptotic normality of $\widehat{\theta}_{\text{DR}}(t)$ in [Section F.3.6](#).

F.3.1 Analysis of Term VI for $\widehat{\theta}_{\text{DR}}(t)$

We analyze the variance and bias of **Term VI** separately as follows. By direct calculations, we have that

$$\begin{aligned}
& \text{Var}[\mathbf{Term VI}] \\
&= \text{Var}[\mathbb{P}_n \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|S})] \\
&= \frac{1}{nh^2} \cdot \text{Var} \left\{ \frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{h \cdot \kappa_2 \cdot \bar{p}_{T|S}(T|\mathbf{S})} [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] + h \cdot \bar{\beta}(t, \mathbf{S}) \right\} \\
&\stackrel{(i)}{\lesssim} \frac{1}{nh^2} \cdot \text{Var} \left\{ \frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{h \cdot \kappa_2 \cdot \bar{p}_{T|S}(T|\mathbf{S})} [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} + \frac{1}{n} \cdot \text{Var}[\bar{\beta}(t, \mathbf{S})] \\
&\stackrel{(ii)}{=} \frac{1}{nh^2} \cdot \mathbb{E} \left\{ \frac{(\frac{T-t}{h})^2 K^2(\frac{T-t}{h})}{h^2 \kappa_2^2 \cdot \bar{p}_{T|S}^2(T|\mathbf{S})} [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^2 \right\} + O\left(\frac{1}{n}\right) \\
&= \frac{1}{nh^2} \int_{\mathcal{S}} \int_{\mathcal{T}} \frac{(\frac{t_1-t}{h})^2 K^2(\frac{t_1-t}{h})}{h^2 \kappa_2^2 \cdot \bar{p}_{T|S}^2(t_1|\mathbf{s}_1)} \left\{ \sigma^2 + [\mu(t_1, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - (t_1-t) \cdot \bar{\beta}(t, \mathbf{s}_1)]^2 \right\} p(t_1, \mathbf{s}_1) dt_1 d\mathbf{s}_1 \\
&\quad + O\left(\frac{1}{n}\right) \\
&\stackrel{(iii)}{=} \frac{1}{nh^3} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 \cdot K^2(u)}{\kappa_2^2 \cdot \bar{p}_{T|S}^2(t+uh|\mathbf{s}_1)} \cdot \left\{ \sigma^2 + [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)]^2 \right\} p(t+uh, \mathbf{s}_1) du d\mathbf{s}_1 \\
&\quad + O\left(\frac{1}{n}\right) \\
&= \frac{1}{nh^3} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 \cdot K^2(u)}{\kappa_2^2 \cdot \bar{p}_{T|S}^2(t|\mathbf{s}_1) + O(h^2)} \cdot \left\{ \sigma^2 + [\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)]^2 + O(h^2) \right\} [p(t, \mathbf{s}_1) + O(h)] du d\mathbf{s}_1 \\
&\quad + O\left(\frac{1}{n}\right) \\
&\stackrel{(iv)}{=} O\left(\frac{1}{nh^3}\right)
\end{aligned}$$

where (i) uses Cauchy-Schwarz inequality on the covariance, (ii) leverages the boundedness of $\bar{\beta}$ under [Assumption A3](#) to derive the term $O\left(\frac{1}{n}\right)$, (iii) applies a change of variable $u = \frac{t_1-t}{h}$, as well as (iv) utilizes the boundedness of $\mu, \bar{\beta}$ under [Assumption A3](#) and the positivity condition ([Assumption A2](#)) on $\bar{p}_{T|S}$. In the above calculation, we also note from the line (i) that the sec-

ond part $\bar{\beta}(t, \mathbf{S})$ of $\Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}})$ is of smaller order than the first term $\frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{h^2\kappa_2\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$. Thus, we can only keep the first term in the final asymptotically linear form of $\hat{\theta}_{\text{DR}}(t)$. Now, by Chebyshev's inequality, we conclude that

$$\begin{aligned} (\mathbb{P}_n - \mathbb{P}) \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) &= O_P \left(\sqrt{\text{Var} [\mathbb{P}_n \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}})]} \right) \\ &= O_P \left(\sqrt{\text{Var} \left[\frac{1}{\sqrt{h^3}} \cdot \mathbb{P}_n \phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right]} \right) \\ &= O_P \left(\sqrt{\frac{1}{nh^3}} \right), \end{aligned}$$

where $\phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) = \frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{\sqrt{h\kappa_2\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})}} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$. In addition, by direct calculations and Taylor's expansions, we derive that

Bias **[Term VI]**

$$\begin{aligned} &= \mathbb{P} [\Phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}})] - \mathbb{E} [\beta(t, \mathbf{S})] \\ &= \mathbb{E} \left\{ \frac{(\frac{T-t}{h})K(\frac{T-t}{h})}{h^2\kappa_2\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\ &= \mathbb{E} \left\{ \int_{\mathcal{T}} \frac{(\frac{t_1-t}{h})K(\frac{t_1-t}{h})}{h^2\kappa_2\bar{p}_{T|\mathbf{S}}(t_1|\mathbf{S})} \cdot [\mu(t_1, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - (t_1-t) \cdot \bar{\beta}(t, \mathbf{S})] p_{T|\mathbf{S}}(t_1|\mathbf{S}) dt_1 \right\} + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\ &\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u)}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} \cdot [\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S})] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} \\ &\quad + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\ &\stackrel{(ii)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u) \left[p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right]}{h \cdot \kappa_2 \left[\bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right]} \right. \\ &\quad \times \left. \left[(\mu(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})) + hu (\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) + O(h^4) \right] du \right\} \\ &\quad + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\ &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u)}{h \cdot \kappa_2} \left[p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right] \right. \\ &\quad \times \left[\frac{1}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{uh \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} - \frac{u^2 h^2 \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{2\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} + \frac{u^2 h^2 \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2}{\bar{p}_{T|\mathbf{S}}^3(t|\mathbf{S})} - \frac{u^3 h^3 \cdot \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{6\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} \right. \\ &\quad \left. \left. + \frac{u^3 h^3 \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right] \left[\frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}^3(t|\mathbf{S})} + O(h^4) \right] \right. \\ &\quad \left. \times \left[(\mu(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})) + hu (\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) + O(h^4) \right] du \right\} \end{aligned}$$

$$\begin{aligned}
& + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\
= & \mathbb{E}_{\mathbf{S}} \left\{ \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} \right] [\mu(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})] \right\} \\
& + \mathbb{E}_{\mathbf{S}} \left\{ [\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \right\} + h^2 B_{\theta}(t) + O(h^3),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ and (ii) applies Taylor's expansion. Here, the complicated bias term $B_{\theta}(t)$ is given by

$$\begin{aligned}
B_{\theta}(t) = & \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{[\mu(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{s})]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right. \\
& \times \left[\frac{1}{3} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) - \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + 2 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \left[\frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 \right. \\
& \left. \left. + \frac{6 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) - 3 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{3 \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} \\
& + \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{[\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{s})]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \left[\frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) - 2 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right. \right. \\
& \left. \left. + 2 p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \left[\frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} \\
& + \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} + \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S})}{3 \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right].
\end{aligned}$$

Under the condition that either $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ or $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$, we have that

$$\mathbb{E}_{\mathbf{S}} \left\{ \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) [\mu(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \cdot \frac{\partial}{\partial t} \log \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} + \mathbb{E}_{\mathbf{S}} \left\{ [\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \right\} = 0$$

and

$$\begin{aligned}
B_{\theta}(t) = & \begin{cases} \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} + \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S})}{3 \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) \right] & \text{when } \bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}, \end{cases} \\
= & \begin{cases} \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{3 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + p_{T|\mathbf{S}}(t|\mathbf{S}) \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) - 3 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right\} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) \right] & \text{when } \bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}. \end{cases}
\end{aligned}$$

As a result, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we know that

$$\mathbf{Term VI} = \mathbb{P}_n \Phi_{h,t} (Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \mathbb{E} [\beta(t, \mathbf{S})]$$

$$\begin{aligned}
&= h^2 B_\theta(t) + o(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right) \\
&= O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right).
\end{aligned}$$

As a side note, under the VC-type condition on the kernel function K (Einmahl and Mason, 2005) (Assumption A5(c)), we can strengthen the above pointwise rate of convergence to the following uniform one as:

$$\sup_{t \in \mathcal{T}} |\mathbf{Term VI}| = O(h^2) + O_P \left(\sqrt{\frac{|\log h|}{nh^3}} \right);$$

see Theorem 4 in Einmahl and Mason (2005) for details.

F.3.2 Analysis of Term VII for $\widehat{\theta}_{\text{DR}}(t)$

By Markov's inequality, we know that

$$\begin{aligned}
\sqrt{nh^3} \cdot \mathbf{Term VII} &= \sqrt{h^3} \cdot \mathbb{G}_n \left[\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \\
&= O_P \left(\sqrt{h^3} \cdot \Upsilon_{3,n} \right) = o_P(1)
\end{aligned}$$

because $\mathbb{E} \left\{ h^3 \cdot \left[\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right]^2 \right\} = h^3 \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2}^2 = O_P \left(h^3 \cdot \Upsilon_{3,n}^2 \right)$ and $h \cdot \Upsilon_{3,n} \rightarrow 0$ as $n \rightarrow \infty$. As a side note, under Assumption A3 on $\bar{\beta}$ and $\widehat{\beta}$, we know that the function $\mathbf{s} \mapsto \widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})$ is Lipschitz continuous with respect to $t \in \mathcal{T}$. Together with the compactness of \mathcal{T} and Example 19.7 in van der Vaart (1998), we can also deduce that

$$\sup_{t \in \mathcal{T}} \left| \sqrt{h^3} \cdot \mathbb{G}_n \left[\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \right| = O_P \left(\sqrt{h^3} \cdot \sup_{t \in \mathcal{T}} \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right),$$

which will be $o_P(1)$ as well if $\sup_{t \in \mathcal{T}} \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = o_P \left(\frac{1}{h} \right)$.

F.3.3 Analyses of Term VIII and Term IX for $\widehat{\theta}_{\text{DR}}(t)$

The argument for showing **Term VIII** and **Term IX** to be $o_P \left(\sqrt{\frac{1}{nh^3}} \right)$ will be similar to the one for **Term VII** above. By Markov's inequality, we know that

$$\begin{aligned}
\sqrt{nh^3} \cdot \mathbf{Term VIII} &= \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h} \right) K \left(\frac{T-t}{h} \right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \\
&= O_P \left(\Upsilon_{2,n} \right) = o_P(1)
\end{aligned}$$

because

$$\begin{aligned}
& \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \cdot \frac{[\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{\widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^2 \right\} \\
&= \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \cdot \frac{[\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{\widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \cdot [(\mu(T, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S}))^2 + \sigma^2] \right\} \\
&\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u^2 K^2(u) [\widehat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\kappa_2^2 \cdot \widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \right. \\
&\quad \left. \times [(\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S}))^2 + \sigma^2] \right\} \\
&\stackrel{(ii)}{\lesssim} \sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2}^2 \\
&\stackrel{(iii)}{=} O_P(\Upsilon_{2,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{T-t}{h}$ in the integration, (ii) leverages the boundedness of $\mu, \bar{\mu}, \bar{\beta}$ under Assumption A3 and the positivity condition (Assumption A2) on $\bar{p}_{T|\mathbf{S}}$, as well as (iii) applies $\sup_{|u-t| \leq h} \|\widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = O_P(\Upsilon_{2,n})$ with $\Upsilon_{2,n} \rightarrow 0$ as $n \rightarrow \infty$. As a side note again, under the VC-type condition on the kernel function K (Einmahl and Mason, 2005) and $\sup_{t \in \mathcal{T}} \|\widehat{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})\|_{L_2} = o_P(1)$, we can strengthen the above pointwise rate of convergence to the following uniform result as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \right| = o_P(1).$$

Similarly, by Markov's inequality, we have that

$$\begin{aligned}
\sqrt{nh^3} \cdot \mathbf{Term IX} &= \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right] \right\} \\
&= O_P(\max\{\Upsilon_{1,n}, h \cdot \Upsilon_{3,n}\}) = o_P(1)
\end{aligned}$$

because

$$\begin{aligned}
& \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2 \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 \right\} \\
&= \mathbb{E} \left\{ \int_{\mathcal{T}} \frac{\left(\frac{t_1-t}{h}\right)^2 K^2\left(\frac{t_1-t}{h}\right) \cdot p_{T|\mathbf{S}}(t_1|\mathbf{S})}{h \cdot \kappa_2^2 \cdot \bar{p}_{T|\mathbf{S}}^2(t_1|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (t_1-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 dt_1 \right\} \\
&\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u^2 K^2(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\kappa_2^2 \cdot \bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + hu \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 du \right\}
\end{aligned}$$

$$\begin{aligned}
& \stackrel{\text{(ii)}}{\lesssim} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}^2 + h^2 \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2}^2 \\
& = O_P(\Upsilon_{1,n}^2 + h^2 \Upsilon_{3,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{t_1-t}{h}$ and (ii) leverages the boundedness of $p_{T|\mathbf{S}}$ under Assumption A3, the positivity condition (Assumption A2) on $\bar{p}_{T|\mathbf{S}}$, the boundedness condition on K under Assumption A5, as well as $\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n})$ and $h \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(h \cdot \Upsilon_{3,n})$ with $\Upsilon_{1,n}, h \cdot \Upsilon_{3,n} \rightarrow 0$ as $n \rightarrow \infty$. In addition, if $\sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = o_P(1)$ and $\sup_{t \in \mathcal{T}} \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = o_P(\frac{1}{h})$, then the above pointwise rate of convergence can be strengthened to the uniform one as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right] \right\} \right| = o_P(1).$$

F.3.4 Analysis of Term X for $\widehat{\theta}_{\text{DR}}(t)$

We first calculate that

$$\begin{aligned}
& \mathbb{E} \left| \sqrt{nh^3} \cdot \mathbf{Term X} \right| \\
& \mathbb{E} \left| \sqrt{\frac{n}{h}} \cdot \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\kappa_2} \cdot \frac{[\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})]}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right] \right| \\
& \stackrel{\text{(i)}}{\leq} \sqrt{nh} \cdot \sqrt{\mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right) [\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})]^2}{h \cdot \widehat{p}_{T|\mathbf{S}}^2(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(T|\mathbf{S})} \right\}} \\
& \quad \times \sqrt{\mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 \right\}} \\
& = \sqrt{nh} \cdot \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \widehat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]^2}{\widehat{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}^2(t+uh|\mathbf{S})} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\
& \quad \times \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u^2 K(u)}{\kappa_2^2} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + hu \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\
& \lesssim \sqrt{nh} \sup_{|u-t| \leq h} \left\| \widehat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} \left[\left\| \widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) \right\|_{L_2} + h \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right] \\
& \stackrel{\text{(ii)}}{=} o_P(1),
\end{aligned}$$

where (i) uses Cauchy-Schwarz inequality and (ii) leverages our assumption (c) on the doubly robust rate of convergence in the theorem statement. As a result, by Markov's inequality, we obtain that

$$\sqrt{nh^3} \cdot \mathbf{Term X}$$

$$\begin{aligned}
&= \sqrt{\frac{n}{h}} \cdot \mathbb{P}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\kappa_2} \cdot \frac{[\bar{p}_{T|\mathbf{S}}(T|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(T|\mathbf{S})]}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S}) \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) (\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S})) \right] \right\} \\
&= o_P(1).
\end{aligned}$$

F.3.5 Analysis of Term XI for $\hat{\theta}_{\text{DR}}(t)$

By direct calculations under model (1) with some change of variable, we have that

Term XI

$$\begin{aligned}
&= \mathbb{E} \left\{ \left[1 - \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \right\} + \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\} \\
&+ \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \cdot \kappa_2} \left[\frac{1}{\hat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \\
&= \mathbb{E} \left\{ \underbrace{\left[1 - \int_{\mathbb{R}} \frac{u^2 \cdot K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} du \right]}_{\text{Term XIa}} [\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \right\} \\
&+ \mathbb{E} \left\{ \underbrace{\int_{\mathbb{R}} \frac{u \cdot K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] du}_{\text{Term XIb}} \right\} \\
&+ \mathbb{E} \left\{ \underbrace{\int_{\mathbb{R}} \frac{u \cdot K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S})] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du}_{\text{Term XIc}} \right\}
\end{aligned}$$

On one hand, when $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$, we know from Assumption A5 that **Term XIa** = **Term XIb** = 0 and

$$\begin{aligned}
\text{Term XIc} &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u) [p_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{h \cdot \kappa_2 \cdot \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S})] du \right\} \\
&\lesssim \frac{1}{h} \cdot \sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} \\
&= o_P \left(\sqrt{\frac{1}{nh^3}} \right)
\end{aligned}$$

by the boundedness of $\mu, \bar{\mu}$ under Assumption A3, the positivity condition (Assumption A2), and our assumption (c) on the doubly robust rate of convergence in the theorem statement. Specifically, since $\|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} + h \|\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})\|_{L_2} = o_P(1)$ when $\bar{\mu} \neq \mu$ and $\bar{\beta} \neq \beta$, our assumption (c) ensures that $\sup_{|u-t| \leq h} \|\hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$.

On the other hand, when $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$, we know from Assumption A2 on $\bar{p}_{T|\mathbf{S}}$ and the

boundedness of $p_{T|\mathbf{S}}$ by Assumption A4 that

$$\begin{aligned} \text{Term XIa} &= \mathbb{E} \left\{ \left[1 - \int_{\mathbb{R}} \frac{u^2 \cdot K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{\kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} du \right] \left[\hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \right\} \\ &\lesssim \left\| \hat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \\ &= o_P \left(\sqrt{\frac{1}{nh^3}} \right), \end{aligned}$$

where we argue from our assumption (c) on the doubly robust rate of convergence in the theorem statement that $\left\| \hat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh^3}} \right)$ if $\bar{p}_{T|\mathbf{S}} \neq p_{T|\mathbf{S}}$ and $\sup_{|u-t| \leq h} \left\| \hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} = O_P(1)$. In addition, we also have that

$$\begin{aligned} \text{Term XIb} &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u) \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S})}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] du \right\} \\ &\lesssim \frac{1}{h} \left\| \hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} \\ &= o_P \left(\sqrt{\frac{1}{nh^3}} \right), \end{aligned}$$

where we again argue from our assumption (c) on the doubly robust rate of convergence in the theorem statement that $\left\| \hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$ if $\bar{p}_{T|\mathbf{S}} \neq p_{T|\mathbf{S}}$ and $\sup_{|u-t| \leq h} \left\| \hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} = O_P(1)$. Finally, we also derive that

Term XIc

$$\begin{aligned} &= \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u \cdot K(u) [\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) - \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})]}{h \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S}) \cdot \hat{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} [\mu(t+uh, \mathbf{S}) - \mu(t, \mathbf{S}) - hu \cdot \beta(t, \mathbf{S})] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} \\ &\stackrel{(i)}{=} \mathbb{E} \left\{ \sup_{|u-t| \leq h} \left| \hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right| \right. \\ &\quad \times \left. \int_{\mathbb{R}} \frac{uK(u) \left[\frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + O(h^3) \right] [p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^2)]}{h \cdot \kappa_2 \left[\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S}) + 2uh \cdot \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S}) + O(h^2) \right] [1 + O_P(\Upsilon_{1,n})]} du \right\} \\ &= O_P \left(h^2 \sup_{|u-t| \leq h} \left\| \hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - p_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} \right) \\ &= O_P(h^2 \cdot \Upsilon_{2,n}) \\ &\stackrel{(ii)}{=} o_P \left(\sqrt{\frac{1}{nh^3}} \right), \end{aligned}$$

where (i) applies Taylor's expansion and mean-value theorem for integrals as well as uses the fact that the difference between $\bar{p}_{T|\mathbf{S}}$ and $\hat{p}_{T|\mathbf{S}}$ is small when $\sup_{|u-t| \leq h} \left\| \hat{p}_{T|\mathbf{S}}(u|\mathbf{S}) - \bar{p}_{T|\mathbf{S}}(u|\mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{2,n})$, while (ii) leverages the arguments that $\sqrt{nh^3} \cdot h^2 = \sqrt{nh^7} \rightarrow \sqrt{c_3} \in [0, \infty)$ and $\Upsilon_{2,n} \rightarrow 0$

as $n \rightarrow \infty$.

F.3.6 Asymptotic Normality of $\widehat{\theta}_{\text{DR}}(t)$

For the asymptotic normality of $\widehat{\theta}_{\text{DR}}(t)$, it follows from the Lyapunov central limit theorem. Specifically, we already show in [Section F.3.1](#) and subsequent subsections that

$$\begin{aligned} \sqrt{nh^3} \left[\widehat{\theta}_{\text{DR}}(t) - \theta(t) \right] &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) + \sqrt{h^3} [\bar{\beta}(t, \mathbf{S}_i) - \mathbb{E}[\bar{\beta}(t, \mathbf{S})]] \right\} + o_P(1) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) + o_P(1) \end{aligned}$$

with

$$\phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$$

and $V_\theta(t) = \mathbb{E} \left[\phi_{h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right] = O(1)$ by our calculation in [Term VI](#). Then,

$$\sum_{i=1}^n \text{Var} \left[\frac{1}{\sqrt{n}} \cdot \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right] = O(1)$$

and

$$\begin{aligned} &\sum_{i=1}^n \mathbb{E} \left| \frac{1}{\sqrt{n}} \cdot \phi_{h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right|^{2+c_1} \\ &= \mathbb{E} \left| \frac{\left(\frac{T-t}{h}\right)^{2+c_1} K^{2+c_1}\left(\frac{T-t}{h}\right) \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^{2+c_1}}{n^{\frac{c_1}{2}} h^{1+\frac{c_1}{2}} \cdot \kappa_2^{2+c_1} \cdot \bar{p}_{T|\mathbf{S}}^{2+c_1}(T|\mathbf{S})} \right| \\ &\lesssim \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u^{2+c_1} K^{2+c_1}(u) \cdot \left[\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S}) \right]^{2+c_1} + \mathbb{E}|\epsilon|^{2+c_1}}{\sqrt{(nh)^{c_1}} \cdot \bar{p}_{T|\mathbf{S}}^{2+c_1}(t+uh|\mathbf{S})} \cdot p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} \\ &= O \left(\sqrt{\frac{1}{(nh)^{c_1}}} \right) = o(1) \end{aligned}$$

by the boundedness of $\mu, \bar{\mu}, p_{T|\mathbf{S}}$, the positivity condition on $\bar{p}_{T|\mathbf{S}}$, the assumption that $\mathbb{E}|\epsilon|^{2+c_1} < \infty$, and the requirement that $nh^3 \rightarrow \infty$ as $n \rightarrow \infty$. Hence, the Lyapunov condition holds, and we have that

$$\sqrt{nh^3} \left[\widehat{\theta}_{\text{DR}}(t) - \theta(t) - h^2 B_\theta(t) \right] \xrightarrow{d} \mathcal{N}(0, V_\theta(t))$$

after subtracting the dominating bias term $h^2 B_\theta(t)$ of $\phi_{h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}_{T|\mathbf{S}})$ that we have computed in [Term VI](#). The proof of [Theorem 1](#) is thus completed. \square

F.4 Asymptotic Bias of $\widehat{\theta}_{\text{AIPW}}(t)$ in (9)

Analogous to our calculations in Section F.3, we can decompose $\widehat{\theta}_{\text{AIPW}}(t) - \theta(t)$ under Assumption A1 as:

$$\begin{aligned}
& \widehat{\theta}_{\text{AIPW}}(t) - \theta(t) \\
&= \frac{1}{nh} \sum_{i=1}^n \left\{ \frac{K\left(\frac{T_i-t}{h}\right)}{\widehat{p}_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \left[\frac{Y_i}{\kappa_2} \left(\frac{T_i-t}{h^2}\right) - \widehat{\beta}(t, \mathbf{S}_i) \right] + h \cdot \widehat{\beta}(t, \mathbf{S}_i) \right\} - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] \\
&= \mathbb{P}_n \widetilde{\Phi}_{h,t}(Y, T, \mathbf{S}; \widehat{\beta}, \widehat{p}_{T|\mathbf{S}}) - \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \right] + \mathbb{P}_n \left[\Phi_{h,t}(Y, T, \mathbf{S}; \widehat{\beta}, \widehat{p}_{T|\mathbf{S}}) - \Phi_{h,t}(Y, T, \mathbf{S}; \bar{\beta}, \bar{p}_{T|\mathbf{S}}) \right] \\
&= \underbrace{\mathbb{P}_n \widetilde{\Phi}_{h,t}(Y, T, \mathbf{S}; \widehat{\beta}, \widehat{p}_{T|\mathbf{S}}) - \mathbb{E} [\beta(t, \mathbf{S})]}_{\text{Dominating Term}} + (\mathbb{P}_n - \mathbb{P}) \left[\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \\
&\quad + (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] \left[\frac{Y}{\kappa_2} \left(\frac{T-t}{h^2}\right) - \bar{\beta}(t, \mathbf{S}) \right] \right\} \\
&\quad + (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \left[\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right] \right\} \\
&\quad + \mathbb{P}_n \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] \left[\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right] \right\} \\
&\quad + \mathbb{P} \left\{ \left[1 - \frac{K\left(\frac{T-t}{h}\right)}{h \cdot \bar{p}(T|\mathbf{S})} \right] \left[\widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] + \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{1}{\widehat{p}_{T|\mathbf{S}}(T|\mathbf{S})} - \frac{1}{\bar{p}_{T|\mathbf{S}}(T|\mathbf{S})} \right] \left[\frac{Y}{\kappa_2} \left(\frac{T-t}{h^2}\right) - \bar{\beta}(t, \mathbf{S}) \right] \right\},
\end{aligned}$$

where $\widetilde{\Phi}_{h,t}(Y, T, \mathbf{S}; \beta, p_{T|\mathbf{S}}) = \frac{K\left(\frac{T-t}{h}\right)}{h \cdot p_{T|\mathbf{S}}(T|\mathbf{S})} \left[\frac{Y}{\kappa_2} \left(\frac{T-t}{h^2}\right) - \beta(t, \mathbf{S}) \right] + \beta(t, \mathbf{S})$. Thus, in order to study the asymptotically dominating bias of $\widehat{\theta}_{\text{AIPW}}(t)$ in (9), it suffices to compute $\mathbb{E} \left[\widetilde{\Phi}_{h,t}(Y, T, \mathbf{S}; \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \beta(t, \mathbf{S}) \right]$ as in Proposition F.1 below.

Proposition F.1 (Asymptotically dominating bias of $\widehat{\theta}_{\text{AIPW}}(t)$). *Suppose that Assumptions A1, A3, A4, A5, and A2 hold under the general model (1) and $\widehat{\beta}, \widehat{p}_{T|\mathbf{S}}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s})$, $\bar{\beta}(t, \mathbf{s})$, and $\bar{p}_{T|\mathbf{S}}(t|\mathbf{s})$ be fixed bounded functions to which $\widehat{\mu}(t, \mathbf{s})$, $\widehat{\beta}(t, \mathbf{s})$ and $\widehat{p}_{T|\mathbf{S}}(t|\mathbf{s})$ converge. Assume also that $\bar{p}_{T|\mathbf{S}}$ satisfies Assumptions A4 and A2. Then, the asymptotically dominating bias of $\widehat{\theta}_{\text{AIPW}}(t)$ is given by*

$$\begin{aligned}
& \mathbb{E} \left[\widetilde{\Phi}_{h,t}(Y, T, \mathbf{S}; \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \beta(t, \mathbf{S}) \right] \\
&= \mathbb{E}_{\mathbf{S}} \left\{ \frac{\mu(t, \mathbf{S}) \cdot p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \cdot \frac{\partial}{\partial t} \log \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} + \mathbb{E}_{\mathbf{S}} \left\{ \left[\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right] \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \right\} + O(h^2)
\end{aligned}$$

when $h \rightarrow 0$ and $n \rightarrow \infty$.

Proof of Proposition F.1. By direct calculations under model (1) and Assumption A1, we derive

that

$$\begin{aligned}
& \mathbb{E} \left[\tilde{\Phi}_{h,t} (Y, T, \mathbf{S}; \bar{\beta}, \bar{p}_{T|\mathbf{S}}) - \beta(t, \mathbf{S}) \right] \\
&= \mathbb{E} \left\{ \int_{\mathcal{T}} \frac{K\left(\frac{t_1-t}{h}\right)}{h \cdot \bar{p}_{T|\mathbf{S}}(t_1|\mathbf{S})} \cdot \left[\frac{\mu(t_1, \mathbf{S})}{\kappa_2} \left(\frac{t_1-t}{h^2} \right) - \bar{\beta}(t, \mathbf{S}) \right] p_{T|\mathbf{S}}(t_1|\mathbf{S}) dt_1 \right\} + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\
&\stackrel{(i)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u)}{\bar{p}_{T|\mathbf{S}}(t+uh|\mathbf{S})} \cdot \left[\frac{u \cdot \mu(t+uh, \mathbf{S})}{h \cdot \kappa_2} - \bar{\beta}(t, \mathbf{S}) \right] p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\} + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\
&\stackrel{(ii)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} \frac{K(u) \left[p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4)} \right. \\
&\quad \times \left. \left[\frac{u \cdot \mu(t, \mathbf{S})}{h \cdot \kappa_2} + \frac{u^2 \beta(t, \mathbf{S})}{\kappa_2} - \bar{\beta}(t, \mathbf{S}) + \frac{u^3 h}{2\kappa_2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + \frac{u^4 h^2}{6\kappa_2} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) + O(h^3) \right] du \right\} \\
&\quad + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\
&\stackrel{(iii)}{=} \mathbb{E} \left\{ \int_{\mathbb{R}} K(u) \left[p_{T|\mathbf{S}}(t|\mathbf{S}) + uh \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) + O(h^4) \right] \right. \\
&\quad \times \left[\frac{1}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{uh \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} - \frac{u^2 h^2 \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{2\bar{p}_{T|\mathbf{S}}^3(t|\mathbf{S})} + \frac{u^2 h^2 \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2}{\bar{p}_{T|\mathbf{S}}^3(t|\mathbf{S})} - \frac{u^3 h^3 \cdot \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{6\bar{p}_{T|\mathbf{S}}^4(t|\mathbf{S})} \right. \\
&\quad \left. \left. + \frac{u^3 h^3 \left[\frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right] \left[\frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}^3(t|\mathbf{S})} + O(h^4) \right] \right. \\
&\quad \times \left. \left[\frac{u \cdot \mu(t, \mathbf{S})}{h \cdot \kappa_2} + \frac{u^2 \beta(t, \mathbf{S})}{\kappa_2} - \bar{\beta}(t, \mathbf{S}) + \frac{u^3 h}{2\kappa_2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S}) + \frac{u^4 h^2}{6\kappa_2} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) + O(h^3) \right] du \right\} \\
&\quad + \mathbb{E} [\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})] \\
&= \mathbb{E} \left[\frac{\mu(t, \mathbf{S}) \cdot \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{\mu(t, \mathbf{S}) \cdot p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} \right] + \mathbb{E} \left\{ [\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \right\} \\
&\quad + h^2 \tilde{B}_\theta(t) + O(h^3),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ while (ii) and (iii) apply Taylor's expansions under Assumptions A4 and A3. Here, the complicated bias term $\tilde{B}_\theta(t)$ is given by

$$\begin{aligned}
\tilde{B}_\theta(t) &= \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{\mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \left[\frac{1}{3} \cdot \frac{\partial^3}{\partial t^3} p_{T|\mathbf{S}}(t|\mathbf{S}) - \frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right. \right. \\
&\quad \left. \left. + 2 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \left[\frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 \right. \right. \\
&\quad \left. \left. + \frac{6 \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) - \frac{\partial^3}{\partial t^3} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) - 3 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{3\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} \\
&\quad + \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left\{ \frac{\left[\frac{\kappa_4}{\kappa_2} \cdot \beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right]}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \left[\frac{\partial^2}{\partial t^2} p_{T|\mathbf{S}}(t|\mathbf{S}) - 2 \frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right. \right.
\end{aligned}$$

$$\begin{aligned}
& + 2p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \left[\frac{\partial}{\partial t} \log \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \right]^2 - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \Bigg\} \\
& + \frac{\kappa_4}{2\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\frac{\partial}{\partial t} p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial}{\partial t} \bar{p}_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{S})}{\bar{p}_{T|\mathbf{S}}^2(t|\mathbf{S})} + \frac{p_{T|\mathbf{S}}(t|\mathbf{S}) \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S})}{3\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right].
\end{aligned}$$

When $\bar{p}_{T|\mathbf{S}} = p_{T|\mathbf{S}}$, we have that

$$\mathbb{E}_{\mathbf{S}} \left\{ \frac{\mu(t, \mathbf{S}) \cdot p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \cdot \frac{\partial}{\partial t} \log \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} + \mathbb{E}_{\mathbf{S}} \left\{ [\beta(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S})] \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} - 1 \right] \right\} = 0$$

and $\tilde{B}_\theta(t) = \frac{\kappa_4}{6\kappa_2} \cdot \mathbb{E}_{\mathbf{S}} \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{S}) \right]$. In this case, the dominating bias term is $h^2 \tilde{B}_\theta(t)$, which tends to 0 as $h \rightarrow 0$ and $n \rightarrow \infty$.

However, when $\bar{\beta} = \beta$ (and $\bar{\mu} = \mu$), the dominating bias is equal to

$$\mathbb{E}_{\mathbf{S}} \left\{ \frac{\mu(t, \mathbf{S}) \cdot p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \cdot \frac{\partial}{\partial t} \log \left[\frac{p_{T|\mathbf{S}}(t|\mathbf{S})}{\bar{p}_{T|\mathbf{S}}(t|\mathbf{S})} \right] \right\} + h^2 \tilde{B}_\theta(t),$$

which is not necessarily 0.

This also shows that the naive AIPW estimator (9) is not doubly robust. \square

G Proofs of Propositions 2, 3, and 4

G.1 Proof of Proposition 2

Proposition 3 (Inconsistency of IPW estimators). *Suppose that Assumptions A1(a-c), A3, A4(c), and A5(a-b) hold under the additive confounding model (13). Assume also that when the bandwidth h is small, the Lebesgue measure of the symmetric difference set satisfies*

$$|\mathcal{S}(t+uh) \Delta \mathcal{S}(t)| = |[\mathcal{S}(t+uh) \setminus \mathcal{S}(t)] \cup [\mathcal{S}(t) \setminus \mathcal{S}(t+uh)]| = o(1)$$

for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$. Then, when h is small, the expectation of $\tilde{m}_{\text{IPW}}(t)$ in (18) is given by

$$\mathbb{E}[\tilde{m}_{\text{IPW}}(t)] = \bar{m}(t) \cdot \rho(t) + \omega(t) + o(1),$$

where $\rho(t) = \mathbb{P}(\mathbf{S} \in \mathcal{S}(t))$ and $\omega(t) = \mathbb{E}[\eta(\mathbf{S}) \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t)\}}]$. If, in addition, there exists a constant $A_h > 0$ depending on h such that

$$\int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t) + \eta(\mathbf{S})] [\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}}] \right\} u \cdot K(u) du = O(A_h)$$

for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$ when h is small, then the expectation of $\tilde{\theta}_{\text{IPW}}(t)$ in (12) is given by

$$\mathbb{E}[\tilde{\theta}_{\text{IPW}}(t)] = \bar{m}'(t) \cdot \rho(t) + O\left(\frac{A_h}{h}\right).$$

Proof of Proposition 2. Notice that the conditional density support $\mathcal{S}(t)$ depends on t when the positivity condition is violated. Under the additive confounding model (13), we have that

$$\begin{aligned}
\mathbb{E} [\tilde{m}_{\text{IPW}}(t)] &= \mathbb{E} \left[\frac{1}{h} \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \cdot Y_i \right] \\
&= \mathbb{E} \left\{ \frac{1}{h} \frac{K\left(\frac{T_i-t}{h}\right)}{p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} [\bar{m}(T_i) + \eta(\mathbf{S}_i)] \right\} \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(\tilde{t})} \frac{1}{h} \frac{K\left(\frac{\tilde{t}-t}{h}\right)}{p_{T|\mathbf{S}}(\tilde{t}|\mathbf{s})} [\bar{m}(\tilde{t}) + \eta(\mathbf{s})] p(\tilde{t}, \mathbf{s}) d\mathbf{s} d\tilde{t} \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(\tilde{t})} \frac{1}{h} K\left(\frac{\tilde{t}-t}{h}\right) [\bar{m}(\tilde{t}) + \eta(\mathbf{s})] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} d\tilde{t} \\
&\stackrel{\text{(i)}}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} K(u) [\bar{m}(t+uh) + \eta(\mathbf{s})] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} du \\
&\stackrel{\text{(ii)}}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} K(u) [\bar{m}(t) + \eta(\mathbf{s}) + uh \cdot \bar{m}'(t) + O(h^2)] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} du \\
&= \int_{\mathbb{R}} \int_{\mathcal{S}(t)} K(u) [\bar{m}(t) + \eta(\mathbf{s}) + uh \cdot \bar{m}'(t) + O(h^2)] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} du \\
&\quad + \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} K(u) [\bar{m}(t) + \eta(\mathbf{s}) + uh \cdot \bar{m}'(t) + O(h^2)] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} du \\
&\quad - \int_{\mathbb{R}} \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} K(u) [\bar{m}(t) + \eta(\mathbf{s}) + uh \cdot \bar{m}'(t) + O(h^2)] p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} du \\
&\stackrel{\text{(iii)}}{=} \int_{\mathcal{S}(t)} [\bar{m}(t) + \eta(\mathbf{s})] \cdot p_{\mathcal{S}}(\mathbf{s}) d\mathbf{s} + O(h^2) + o(1) \\
&= \bar{m}(t) \cdot \mathbb{P}(\mathbf{S} \in \mathcal{S}(t)) + \mathbb{E} [\eta(\mathbf{S}) \cdot \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t)\}}] + o(1) \\
&= \bar{m}(t) \cdot \rho(t) + \omega(t) + o(1),
\end{aligned}$$

where (i) follows from a change of variable $u = \frac{\tilde{t}-t}{h}$, (ii) is due to Taylor's expansion under Assumption A3, and (iii) relies on our assumption on the Lebesgue measure $|\mathcal{S}(t+uh) \Delta \mathcal{S}(t)| = o(1)$ for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$.

Similarly, we can also derive that

$$\begin{aligned}
&\mathbb{E} [\tilde{\theta}_{\text{IPW}}(t)] \\
&= \mathbb{E} \left[\frac{[\bar{m}(T_i) + \eta(\mathbf{S}_i)] \left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right)}{h^2 \cdot \kappa_2 \cdot p_{T|\mathbf{S}}(T_i|\mathbf{S}_i)} \right] \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right)}{h^2 \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 dt_1 \\
&\stackrel{\text{(iv)}}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u)}{h \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 du
\end{aligned}$$

$$\begin{aligned}
&= \int_{\mathbb{R}} \int_{\mathcal{S}(t)} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u)}{h \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 du \\
&\quad + \int_{\mathbb{R}} \left[\int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} - \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} \right] \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u)}{h \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 du \\
&\stackrel{(v)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t)} \frac{[\bar{m}(t) + \eta(\mathbf{s}_1) + uh \cdot \bar{m}'(t) + O(h^2)] u \cdot K(u)}{h \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 du \\
&\quad + \int_{\mathbb{R}} \left[\int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} - \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} \right] \frac{[\bar{m}(t) + \eta(\mathbf{s}_1) + uh \cdot \bar{m}'(t) + O(h^2)] u \cdot K(u)}{h \cdot \kappa_2} \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 du \\
&\stackrel{(vi)}{=} \int_{\mathcal{S}(t)} \bar{m}'(t) \cdot p_{\mathcal{S}}(\mathbf{s}_1) d\mathbf{s}_1 + O(h) \\
&\quad + \frac{1}{h \cdot \kappa_2} \int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t) + \eta(\mathbf{S})] \left[\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}} \right] \right\} u \cdot K(u) du \\
&\quad + \frac{1}{\kappa_2} \int_{\mathbb{R}} \mathbb{E} \left\{ \bar{m}'(t) \left[\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}} \right] \right\} u^2 K(u) du \\
&\stackrel{(vii)}{=} \bar{m}'(t) \cdot \rho(t) + O\left(\frac{A_h}{h}\right),
\end{aligned}$$

where (iv) follows from a change of variable $u = \frac{\tilde{t}-t}{h}$, (v) is due to Taylor's expansion under Assumption A3, (vi) leverages the property of the second-order kernel K , and (vii) relies on our assumptions on the Lebesgue measure $|\mathcal{S}(t+uh) \Delta \mathcal{S}(t)| = o(1)$ and (19) for any $t \in \mathcal{T}$ and $u \in \mathbb{R}$. The results follow. \square

G.2 Proof of Proposition 3

Proposition 4. *Suppose that Assumptions A1(a-c), A3, A4(c), and A5(a-b) hold under the additive confounding model (13). Then, when the bandwidth h is small, the expectation of the modified IPW quantity (20) is given by*

$$\begin{aligned}
\mathbb{E} \left[\tilde{\Xi}_t(Y, T, \mathbf{S}) \right] &= \bar{m}'(t) + O(h^2) \\
&\quad + \int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t+uh) + \eta(\mathbf{S})] \left[\mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathbf{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}} \right] \middle| T = t \right\} u \cdot K(u) du.
\end{aligned}$$

Proof of Proposition 3. Recall that the conditional density support $\mathcal{S}(t)$ depends on t when the positivity condition fails to hold. Under the additive confounding model (13), we have that

$$\begin{aligned}
&\mathbb{E} \left[\tilde{\Xi}_t(Y, T, \mathbf{S}) \right] \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h^2 \cdot \kappa_2} d\mathbf{s}_1 dt_1 \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(t)} \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h^2 \cdot \kappa_2} d\mathbf{s}_1 dt_1
\end{aligned}$$

$$\begin{aligned}
& + \left\{ \left[\int_{\mathcal{T}} \int_{\mathcal{S}(t_1) \setminus \mathcal{S}(t)} - \int_{\mathcal{T}} \int_{\mathcal{S}(t) \setminus \mathcal{S}(t_1)} \right] \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h^2 \cdot \kappa_2} d\mathbf{s}_1 dt_1 \right\} \\
& \stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t)} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u) \cdot p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\
& + \left\{ \int_{\mathbb{R}} \left[\int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} - \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} \right] \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h \cdot \kappa_2} d\mathbf{s}_1 du \right\} \\
& \stackrel{(ii)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t)} \frac{\left[\bar{m}(t) + \eta(\mathbf{s}_1) + uh \cdot \bar{m}'(t) + \frac{u^2 h^2}{2} \cdot \bar{m}''(t) + O(h^3) \right] u \cdot K(u) \cdot p(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\
& + \left\{ \int_{\mathbb{R}} \left[\int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} - \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} \right] \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h \cdot \kappa_2} d\mathbf{s}_1 du \right\} \\
& \stackrel{(iii)}{=} \bar{m}'(t) + O(h^2) \\
& + \left\{ \int_{\mathbb{R}} \left[\int_{\mathcal{S}(t+uh) \setminus \mathcal{S}(t)} - \int_{\mathcal{S}(t) \setminus \mathcal{S}(t+uh)} \right] \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u) p_{\mathcal{S}|T}(\mathbf{s}_1|t)}{h \cdot \kappa_2} d\mathbf{s}_1 du \right\} \\
& = \bar{m}'(t) + O(h^2) \\
& + \int_{\mathbb{R}} \mathbb{E} \left\{ [\bar{m}(t+uh) + \eta(\mathbf{S})] [\mathbb{1}_{\{\mathcal{S} \in \mathcal{S}(t+uh) \setminus \mathcal{S}(t)\}} - \mathbb{1}_{\{\mathcal{S} \in \mathcal{S}(t) \setminus \mathcal{S}(t+uh)\}}] \Big| T = t \right\} u \cdot K(u) du,
\end{aligned}$$

where (i) follows from a change of variable $u = \frac{t_1-t}{h}$, (ii) is due to Taylor's expansion, and (iii) utilizes the fact that K is a second-order kernel function by Assumption A5(b). The result thus follows.

As stated in Remark 5, we should evaluate the conditional density $p_{\mathcal{S}|T}$ at the (query) point (t, \mathbf{S}) instead of the sample point (T, \mathbf{S}) in the modified IPW quantity (20). To see this, we consider the alternative modified IPW quantity

$$\tilde{\Xi}_{t,2}(Y, T, \mathbf{S}) = \frac{Y \left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) p_{\mathcal{S}|T}(\mathbf{S}|T)}{h^2 \cdot \kappa_2 \cdot p(T, \mathbf{S})}$$

and compute its expectation as:

$$\begin{aligned}
& \mathbb{E} \left[\tilde{\Xi}_{t,2}(Y, T, \mathbf{S}) \right] \\
& = \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) \cdot p_{\mathcal{S}|T}(\mathbf{s}_1|t_1)}{h^2 \cdot \kappa_2} d\mathbf{s}_1 dt_1 \\
& \stackrel{(iv)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] u \cdot K(u) \cdot p_{\mathcal{S}|T}(\mathbf{s}_1|t+uh)}{h \cdot \kappa_2} d\mathbf{s}_1 du \\
& \stackrel{(v)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{\left[\bar{m}(t) + \eta(\mathbf{s}_1) + uh \cdot \bar{m}'(t) + \frac{u^2 h^2}{2} \cdot \bar{m}''(t) + O(h^3) \right] u \cdot K(u) \cdot p_{\mathcal{S}|T}(\mathbf{s}_1|t+uh)}{h \cdot \kappa_2} d\mathbf{s}_1 du \\
& \stackrel{(vi)}{=} \bar{m}'(t) + O(h^2) + \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{\eta(\mathbf{s}_1) \cdot u \cdot K(u) \cdot p_{\mathcal{S}|T}(\mathbf{s}_1|t+uh)}{h \cdot \kappa_2} d\mathbf{s}_1 du
\end{aligned}$$

$$= \bar{m}'(t) + O(h^2) + \int_{\mathcal{T}} \frac{\left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right)}{h^2 \cdot \kappa_2} \cdot \mathbb{E}[\eta(\mathbf{S})|T = t_1] dt_1,$$

where (i) follows from a change of variable $u = \frac{t_1-t}{h}$, (ii) is due to Taylor's expansion, and (iii) utilizes the fact that K is a second-order kernel function by Assumption A5(b). Hence, it is unclear how we can eliminate the additional bias term $\int_{\mathcal{T}} \frac{\left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right)}{h^2 \cdot \kappa_2} \cdot \mathbb{E}[\eta(\mathbf{S})|T = t_1] dt_1$ from $\mathbb{E}[\tilde{\Xi}_{t,2}(Y, T, \mathbf{S})]$ unless $\mathbb{E}[\eta(\mathbf{S})|T = t] = 0$, which is not true in general. \square

G.3 Proof of Proposition 4

Proposition 5. *Suppose that Assumptions A1(a-c), A3, A4(c), A5(a-b), and A6 hold under the additive confounding model (13). Then, when the bandwidth $h > 0$ is small, the expectation of the modified IPW quantity (21) is given by*

$$\mathbb{E}[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S})] = \bar{m}'(t) + \frac{h^2 \kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) + O(h^3).$$

Proof of Proposition 4. Since the kernel function K has a compact support under Assumption A5, we can assume, without loss of generality, that it is supported on $[-1, 1]$. Then, when $h < \frac{\zeta}{A_0}$ (or, equivalently $A_0|uh| < \zeta$ for any $u \in [-1, 1]$), we have that

$$\mathcal{S}(t) \ominus \zeta \subset \mathcal{S}(t) \ominus (A_0|uh|) \subset \mathcal{S}(t + uh) \quad \text{and} \quad \mathcal{L}_\zeta(t) \subset \mathcal{L}_{A_0|\delta|}(t) \subset \mathcal{S}(t + \delta)$$

by Assumption A6. Then, under model (13) and the support shrinking approach for $p_\zeta(\mathbf{s}|t)$, the expectation of $\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S})$ is given by

$$\begin{aligned} & \mathbb{E}[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S})] \\ &= \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{[\bar{m}(t_1) + \eta(\mathbf{s}_1)] \left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\ &= \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] \cdot u \cdot K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\ &\stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{[\bar{m}(t+uh) + \eta(\mathbf{s}_1)] \cdot u \cdot K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\ &= \int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{\left[\bar{m}(t) + \eta(\mathbf{s}_1) + uh \cdot \bar{m}'(t) + \frac{u^2 h^2}{2} \cdot \bar{m}''(t) + \frac{u^3 h^3}{6} \cdot \bar{m}^{(3)}(t) + O(h^4)\right] u \cdot K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du \\ &\stackrel{(ii)}{=} \underbrace{\int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{[\bar{m}(t) + \eta(\mathbf{s}_1)] \cdot u \cdot K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2 \cdot h} d\mathbf{s}_1 du}_{=0} + \int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{\bar{m}'(t) \cdot u^2 K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{\kappa_2} d\mathbf{s}_1 du \\ &\quad + \underbrace{\int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{h \cdot \bar{m}''(t) \cdot u^3 K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{2\kappa_2} d\mathbf{s}_1 du}_{=0} + \int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{h^2 \cdot \bar{m}^{(3)}(t) \cdot u^4 K(u) \cdot p_\zeta(\mathbf{s}_1|t)}{6\kappa_2} d\mathbf{s}_1 du \end{aligned}$$

$$\begin{aligned}
& + O(h^3) \\
& = \bar{m}'(t) + \frac{h^2 \kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) + O(h^3) \\
& = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(t, \mathbf{S}) \Big| T = t \right] + \frac{h^2 \kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) + O(h^3),
\end{aligned}$$

where (i) uses the definition of the ζ -interior conditional density (23) and (ii) follows from the fact that K is a second-order kernel function under Assumption A5(b). The result under the level set approach for $p_\zeta(\mathbf{s}|t)$ follows from almost identical arguments. \square

H Proof of Theorem 5

Theorem 6 (Consistency of estimating $\theta(t)$ without positivity). *Suppose that Assumptions A1(a-c), A3, A4, A5, and A6 hold under the additive confounding model (13), and the support $\mathcal{S} \subset \mathbb{R}^d$ of the marginal density $p_{\mathbf{S}}$ is compact. In addition, $\hat{\mu}, \hat{\beta}, \hat{p}_\zeta, \hat{p}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s}), \bar{\beta}(t, \mathbf{s}), \bar{p}_\zeta(\mathbf{s}|t)$, and $\bar{p}(t, \mathbf{s})$ be fixed bounded functions to which $\hat{\mu}(t, \mathbf{s}), \hat{\beta}(t, \mathbf{s}), \hat{p}_\zeta(\mathbf{s}|t)$, and $\hat{p}(t, \mathbf{s})$ converge under the rates of convergence as:*

$$\begin{aligned}
& \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{3,n}), \quad \sup_{\mathbf{s} \in \mathcal{S}} \left| \hat{F}_{\mathbf{S}|T}(\mathbf{s}|t) - F_{\mathbf{S}|T}(\mathbf{s}|t) \right| = O_P(\Upsilon_{4,n}), \\
& \left\| \hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t) \right\|_{L_2} = O_P(\Upsilon_{5,n}), \quad \text{and} \quad \sup_{|u-t| \leq h} \left\| \hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{6,n}),
\end{aligned}$$

where $\Upsilon_{3,n}, \Upsilon_{4,n}, \Upsilon_{5,n}, \Upsilon_{6,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we have that

$$\begin{aligned}
& \hat{\theta}_{\text{C,RA}}(t) - \theta(t) = O_P \left(\Upsilon_{3,n} + \Upsilon_{4,n} + \left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right), \\
& \hat{\theta}_{\text{C,IPW}}(t) - \theta(t) = O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{5,n} + \Upsilon_{6,n} + \sup_{|u-t| \leq h} \left\| \bar{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2} \right).
\end{aligned}$$

If, in addition, we assume that

(a) \bar{p}, \bar{p}_ζ satisfy Assumptions A4 and A6 as well as $\sqrt{nh^3} \cdot \Upsilon_{5,n} = o(1)$;

(b) either (i) “ $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ ” or (ii) “ $\bar{p} = p$ ”;

(c) $\sqrt{nh} \left[\left\| \hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t) \right\|_{L_2} + \sup_{|u-t| \leq h} \left\| \hat{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2} \right] \left[\left\| \hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} + h \left\| \hat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right] = o_P(1)$,

then

$$\sqrt{nh^3} \left[\hat{\theta}_{\text{C,DR}}(t) - \theta(t) \right]$$

$$= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + \sqrt{h^3} \left[\int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \theta(t) \right] \right\} + o_P(1)$$

when $nh^7 \rightarrow c_3$ for some finite number $c_3 \geq 0$, where

$$\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})].$$

Furthermore,

$$\sqrt{nh^3} \left[\hat{\theta}_{C,DR}(t) - \theta(t) - h^2 B_{C,\theta}(t) \right] \xrightarrow{d} \mathcal{N}(0, V_{C,\theta}(t))$$

with $V_{C,\theta}(t) = \mathbb{E} \left[\phi_{C,h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right]$ and

$$B_{C,\theta}(t) = \begin{cases} \frac{\kappa_4}{6\kappa_2} \int \left\{ \frac{3 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \bar{m}''(t) + p(t, \mathbf{s}) [\bar{m}^{(3)}(t) - 3 \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \cdot \bar{m}''(t)]}{\bar{p}(t, \mathbf{s})} \right\} \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) & \text{when } \bar{p} = p. \end{cases}$$

Proof of Theorem 5. We derive the rates of convergence of $\hat{\theta}_{C,RA}(t)$ given by (16) and $\hat{\theta}_{IPW}(t)$ given by (25) in Section H.1 and Section H.2, respectively. We also prove the asymptotic linearity, double robustness, and asymptotic normality of $\hat{\theta}_{C,DR}(t)$ given by (26) in Section H.3.

H.1 Rate of Convergence of $\hat{\theta}_{C,RA}(t)$

Firstly, we derive the rate of convergence for $\hat{\theta}_{C,RA}(t)$ in (16). By Proposition C.1, we know that $\theta(t) = \bar{m}'(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) \middle| T = t \right] = \mathbb{E} [\beta(T, \mathbf{S}) \middle| T = t]$ and

$$\begin{aligned} & \hat{\theta}_{C,RA}(t) - \theta(t) \\ &= \int \hat{\beta}(t, \mathbf{s}) d\hat{F}_{\mathbf{S}|T}(\mathbf{s}|t) - \int \beta(t, \mathbf{s}) dF_{\mathbf{S}|T}(\mathbf{s}|t) \\ &= \underbrace{\int [\hat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})] d\hat{F}_{\mathbf{S}|T}(\mathbf{s}|t)}_{\text{Term I}} + \underbrace{\int \bar{\beta}(t, \mathbf{s}) d[\hat{F}_{\mathbf{S}|T}(\mathbf{s}|t) - F_{\mathbf{S}|T}(\mathbf{s}|t)]}_{\text{Term II}} + \underbrace{\int [\bar{\beta}(t, \mathbf{s}) - \beta(t, \mathbf{s})] dF_{\mathbf{S}|T}(\mathbf{s}|t)}_{\text{Term III}}. \end{aligned}$$

- **Term I:** By Markov's inequality (and Hölder's inequality), we know that

$$\left| \hat{\beta}(t, \mathbf{S}_1) - \bar{\beta}(t, \mathbf{S}_1) \right| = O_P \left(\mathbb{E} \left| \hat{\beta}(t, \mathbf{S}_1) - \bar{\beta}(t, \mathbf{S}_1) \right| \right) = O_P \left(\left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right) = O_P(\Upsilon_{3,n})$$

for any random vector \mathbf{S}_1 supported on $\mathcal{S} \subset \mathbb{R}^d$. Thus,

$$\text{Term I} \leq \int \left| \hat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right| d\hat{F}_{\mathbf{S}|T}(\mathbf{s}|t) = O_P \left(\left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right) = O_P(\Upsilon_{3,n}).$$

- **Term II:** By the compactness of \mathcal{S} and the fact that $\mathcal{S}(t) \subset \mathcal{S}$, we know that the Lebesgue

measure $|\mathcal{S}(t)|$ satisfies $|\mathcal{S}(t)| \leq |\mathcal{S}| < \infty$ for any $t \in \mathcal{T}$ and thus,

$$\begin{aligned} \mathbf{Term II} &\leq \sup_{\mathbf{s} \in \mathcal{S}} |\bar{\beta}(t, \mathbf{s})| \cdot \left\| \widehat{F}_{\mathcal{S}|T}(\cdot|t) - F_{\mathcal{S}|T}(\cdot|t) \right\|_{\text{TV}} \\ &\leq \sup_{\mathbf{s} \in \mathcal{S}} |\bar{\beta}(t, \mathbf{s})| \cdot \sup_{\mathbf{s} \in \mathcal{S}} \left| \widehat{F}_{\mathcal{S}|T}(\mathbf{s}|t) - F_{\mathcal{S}|T}(\mathbf{s}|t) \right| \cdot |\mathcal{S}| \\ &= O_P(\Upsilon_{4,n}) \end{aligned}$$

under Assumption [A3](#) and the condition that $\sup_{\mathbf{s} \in \mathcal{S}} \left| \widehat{F}_{\mathcal{S}|T}(\mathbf{s}|t) - F_{\mathcal{S}|T}(\mathbf{s}|t) \right| = O_P(\Upsilon_{4,n})$, where $\left\| \widehat{F}_{\mathcal{S}|T}(\cdot|t) - F_{\mathcal{S}|T}(\cdot|t) \right\|_{\text{TV}}$ is the total variation distance between the probability measures associated with $\widehat{F}_{\mathcal{S}|T}(\cdot|t)$ and $F_{\mathcal{S}|T}(\cdot|t)$. Notice that $\widehat{F}_{\mathcal{S}|T}(\cdot|t)$ can be constructed on the same data sample $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$.

• **Term III:** Similar to the argument for **Term I**, we have that

$$\mathbf{Term III} \leq \int |\bar{\beta}(t, \mathbf{s}) - \beta(t, \mathbf{s})| d\widehat{F}_{\mathcal{S}|T}(\mathbf{s}|t) = O_P\left(\|\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2}\right).$$

In summary, we conclude that

$$\widehat{\theta}_{\text{C,RA}}(t) - \theta(t) = O_P\left(\Upsilon_{3,n} + \Upsilon_{4,n} + \|\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2}\right).$$

H.2 Rate of Convergence of $\widehat{\theta}_{\text{C,IPW}}(t)$

Secondly, we derive the rate of convergence for $\widehat{\theta}_{\text{C,IPW}}(t)$ in [\(25\)](#). Recall from [\(21\)](#) that

$$\begin{aligned} \widehat{\theta}_{\text{C,IPW}}(t) - \theta(t) &= \mathbb{P}_n \left[\widetilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \theta(t) + \widehat{\theta}_{\text{C,IPW}}(t) - \mathbb{P}_n \left[\widetilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] \\ &= \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i-t}{h} \right) K \left(\frac{T_i-t}{h} \right) \bar{p}_\zeta(\mathbf{S}_i|t)}{h \cdot \kappa_2 \cdot p(T_i, \mathbf{S}_i)} - \mathbb{E} [\beta(T, \mathbf{S})|T=t]}_{\mathbf{Term IV}} \\ &\quad + \underbrace{\frac{1}{nh} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i-t}{h} \right) K \left(\frac{T_i-t}{h} \right)}{h \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}_i|t)}{\widehat{p}(T_i, \mathbf{S}_i)} - \frac{\bar{p}_\zeta(\mathbf{S}_i|t)}{p(T_i, \mathbf{S}_i)} \right]}_{\mathbf{Term V}}. \end{aligned}$$

We shall handle **Term IV** and **Term V** in [Section H.2.1](#) and [Section H.2.2](#), respectively.

H.2.1 Rate of Convergence of **Term IV** for $\widehat{\theta}_{\text{C,IPW}}(t)$

We already computed in [Proposition 4](#) that

$$\mathbb{E} \left[\widetilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \theta(t) = \frac{h^2 \kappa_4}{6 \kappa_2} \cdot \bar{m}^{(3)}(t) + O(h^3)$$

under model (13). In particular, the above equality holds true even when we replace the true ζ -interior conditional density p_ζ with the limiting ζ -interior conditional density \bar{p}_ζ because \bar{p}_ζ also satisfies Assumptions A4 and A6. Additionally, we calculate the variance of $\mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right]$ under \bar{p}_ζ as:

$$\begin{aligned}
& \text{Var} \left\{ \mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] \right\} \\
&= \frac{1}{nh^4\kappa_2^2} \cdot \text{Var} \left[\frac{Y_i \left(\frac{T_i-t}{h} \right) K \left(\frac{T_i-t}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_i|t)}{p(T_i, \mathbf{S}_i)} \right] \\
&= \frac{1}{nh^4\kappa_2^2} \cdot \mathbb{E} \left[\frac{Y_i^2 \left(\frac{T_i-t}{h} \right)^2 K^2 \left(\frac{T_i-t}{h} \right) \cdot \bar{p}_\zeta^2(\mathbf{S}_i|t)}{p^2(T_i, \mathbf{S}_i)} \right] - \frac{1}{nh^4\kappa_2^2} \left\{ \mathbb{E} \left[\frac{Y_i \left(\frac{T_i-t}{h} \right) K \left(\frac{T_i-t}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_i|t)}{p(T_i, \mathbf{S}_i)} \right] \right\}^2 \\
&\stackrel{(i)}{=} \frac{1}{nh^4\kappa_2^2} \int_{\mathcal{S} \times \mathcal{T}} \frac{\left(\frac{t_1-t}{h} \right)^2 K^2 \left(\frac{t_1-t}{h} \right) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t)}{p(t_1, \mathbf{s}_1)} \cdot [\mu(t_1, \mathbf{s}_1)^2 + \sigma^2] dt_1 d\mathbf{s}_1 - \frac{\left\{ \mathbb{E} [\beta(T, \mathbf{S})|T=t] \right\}^2}{n} \\
&\quad + O\left(\frac{h^2}{n}\right) \\
&\stackrel{(ii)}{=} \frac{1}{nh^3\kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t)}{p(t+uh, \mathbf{s}_1)} \cdot [\mu(t+uh, \mathbf{s}_1)^2 + \sigma^2] dud\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{(iii)}{=} \frac{1}{nh^3\kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t)}{p(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} p(t', \mathbf{s}_1)} \left[\mu(t, \mathbf{s}_1)^2 + 2uh \cdot \mu(t'', \mathbf{s}_1) \cdot \frac{\partial}{\partial t} \mu(t'', \mathbf{s}_1) + \sigma^2 \right] dud\mathbf{s}_1 \\
&\quad + O\left(\frac{1}{n}\right) \\
&\stackrel{(iv)}{=} \frac{1}{nh^3\kappa_2^2} \int_{\mathcal{S}} \int_{\mathbb{R}} \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t)}{p(t, \mathbf{s}_1)} [\mu(t, \mathbf{s}_1)^2 + \sigma^2] dud\mathbf{s}_1 + O\left(\frac{1}{n}\right) \\
&\stackrel{(v)}{=} O\left(\frac{1}{nh^3}\right)
\end{aligned}$$

with $\mu(t, \mathbf{s}) = \bar{m}(t) + \eta(\mathbf{s})$ under model (13), where (i) utilizes our result in Proposition 4 for $\mathbb{E} \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right]$, (ii) uses a change of variable $u = \frac{t_1-t}{h}$ and the boundedness of $\beta(t, \mathbf{s})$, (iii) applies the Taylor's expansion under Assumptions A3 and A4 with t', t'' being two points between t and $t+uh$, (iv) absorbs the higher order terms to $O\left(\frac{1}{n}\right)$, and (v) utilizes the properties of K under Assumption A5 and the positivity of $\frac{\bar{p}_\zeta^2(\mathbf{s}|t)}{p(t, \mathbf{s})}$ within the support \mathcal{J} of $p(t, \mathbf{s})$. Now, by Chebyshev's inequality and our above calculations, we obtain that

$$\begin{aligned}
\mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \theta(t) &= \mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \mathbb{E} \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] + \mathbb{E} \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \theta(t) \\
&= O_P \left(\sqrt{\text{Var} \left\{ \mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] \right\}} \right) + O(h^2) \\
&= O_P \left(\sqrt{\frac{1}{nh^3}} \right) + O(h^2)
\end{aligned}$$

as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$. As a side note, under the VC-type condition on K (Assumption A5(c)), we can apply Theorem 2 in Einmahl and Mason (2005) to strengthen the above pointwise rate of convergence to the uniform one as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{P}_n \left[\tilde{\Xi}_{t,\zeta}(Y, T, \mathbf{S}) \right] - \theta(t) \right| = O_P \left(\sqrt{\frac{|\log h|}{nh^3}} \right) + O(h^2).$$

H.2.2 Rate of Convergence of Term V for $\hat{\theta}_{C,IPW}(t)$

By direct calculations, we have that

Term V

$$\begin{aligned} &= \frac{1}{nh} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_i | t)}{h \cdot \kappa_2 \cdot p(T_i, \mathbf{S}_i)} \left[\frac{\hat{p}_\zeta(\mathbf{S}_i | t) \cdot p(T_i, \mathbf{S}_i) - \bar{p}_\zeta(\mathbf{S}_i | t) \cdot \hat{p}(T_i, \mathbf{S}_i)}{\hat{p}(T_i, \mathbf{S}_i) \cdot \bar{p}_\zeta(\mathbf{S}_i | t)} \right] \\ &= \frac{1}{nh} \sum_{i=1}^n \frac{Y_i \left(\frac{T_i - t}{h} \right) K \left(\frac{T_i - t}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_i | t)}{h \cdot \kappa_2 \cdot p(T_i, \mathbf{S}_i)} \left\{ \frac{[\hat{p}_\zeta(\mathbf{S}_i | t) - \bar{p}_\zeta(\mathbf{S}_i | t)] p(T_i, \mathbf{S}_i) - \bar{p}_\zeta(\mathbf{S}_i | t) [\hat{p}(T_i, \mathbf{S}_i) - p(T_i, \mathbf{S}_i)]}{\hat{p}(T_i, \mathbf{S}_i) \cdot \bar{p}_\zeta(\mathbf{S}_i | t)} \right\} \\ &\stackrel{(i)}{=} \left\{ \mathbb{E} [\beta(T, \mathbf{S}) | T = t] + O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right) \right\} \\ &\quad \times \left[\frac{O_P \left(\|\hat{p}_\zeta(\mathbf{S} | t) - \bar{p}_\zeta(\mathbf{S} | t)\|_{L_2} \right)}{\inf_{(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}} p(t, \mathbf{s}) - O_P \left(\sup_{|u-t| \leq h} [\|\hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} + \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2}] \right)} \right. \\ &\quad \left. + \frac{O_P \left(\sup_{|u-t| \leq h} [\|\hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} + \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2}] \right)}{\inf_{(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}} p(t, \mathbf{s}) - O_P \left(\sup_{|u-t| \leq h} [\|\hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} + \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2}] \right)} \right] \\ &= \left[O(1 + h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right) \right] \cdot O_P \left(\Upsilon_{5,n} + \Upsilon_{6,n} + \sup_{|u-t| \leq h} \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right) \\ &= O_P \left(\Upsilon_{5,n} + \Upsilon_{6,n} + \sup_{|u-t| \leq h} \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right) \end{aligned}$$

as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, where (i) utilizes our results for **Term IV** and Markov's inequality.

Combining our results for **Term IV** and **Term V**, we conclude that

$$\hat{\theta}_{C,IPW}(t) - \theta(t) = O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} + \Upsilon_{5,n} + \Upsilon_{6,n} + \sup_{|u-t| \leq h} \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right).$$

H.3 Asymptotic Properties of $\widehat{\theta}_{\text{C,DR}}(t)$

Finally, using some similar arguments to [Section F.3](#), we shall establish the asymptotic properties of $\widehat{\theta}_{\text{C,DR}}(t)$ in [\(26\)](#). By [Proposition C.1](#), we have that

$$\theta(t) = \mathbb{E} \left[\frac{\partial}{\partial t} \mu(T, \mathbf{S}) | T = t \right] = \int \beta(t, \mathbf{s}) dF_{\mathbf{S}|T}(\mathbf{s}|t) = \int \beta(t, \mathbf{s}) \cdot p_{\zeta}(\mathbf{s}|t) d\mathbf{s} = \int \beta(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s}$$

where $\mu(t, \mathbf{s}) = \bar{m}(t) + \eta(\mathbf{s})$. Therefore,

$$\begin{aligned} & \widehat{\theta}_{\text{C,DR}}(t) - \theta(t) \\ &= \frac{1}{n} \sum_{i=1}^n \frac{\left(\frac{T_i-t}{h}\right) K\left(\frac{T_i-t}{h}\right) \cdot \widehat{p}_{\zeta}(\mathbf{S}_i|t)}{h^2 \cdot \kappa_2 \cdot \widehat{p}(T_i, \mathbf{S}_i)} \left[Y_i - \widehat{\mu}(t, \mathbf{S}_i) - (T_i - t) \cdot \widehat{\beta}(t, \mathbf{S}_i) \right] + \int \widehat{\beta}(t, \mathbf{s}) \cdot \widehat{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} \\ & \quad - \int \beta(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} \\ &= \mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{\text{C},h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_{\zeta}) \right] + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} \\ & \quad + \int \widehat{\beta}(t, \mathbf{s}) \cdot \widehat{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} - \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} \\ & \quad + \mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{\text{C},h,t}(Y, T, \mathbf{S}; \widehat{\mu}, \widehat{\beta}, \widehat{p}, \widehat{p}_{\zeta}) - \frac{1}{\sqrt{h^3}} \cdot \phi_{\text{C},h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_{\zeta}) \right] \\ &= \underbrace{\mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{\text{C},h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_{\zeta}) \right] + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s}}_{\text{Term VI}} \\ & \quad + \underbrace{\int \widehat{\beta}(t, \mathbf{s}) [\widehat{p}_{\zeta}(\mathbf{s}|t) - \bar{p}_{\zeta}(\mathbf{s}|t)] d\mathbf{s}}_{\text{Term VII}} \\ & \quad + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_{\zeta}(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_{\zeta}(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\}}_{\text{Term VIII}} \\ & \quad + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_{\zeta}(\mathbf{S}|t)}{h^2 \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) [\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S})]] \right\}}_{\text{Term IX}} \\ & \quad + \underbrace{\mathbb{P}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_{\zeta}(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_{\zeta}(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] [\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) [\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S})]] \right\}}_{\text{Term X}} \\ & \quad + \underbrace{\mathbb{P} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right) \cdot \bar{p}_{\zeta}(\mathbf{S}|t)}{h \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S})] \right\} + \int [\widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})] \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s}}_{\text{Term XIa}} \end{aligned}$$

$$\begin{aligned}
& + \underbrace{\text{P} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S})] \right\}}_{\text{Term XIb}} \\
& + \underbrace{\text{P} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{\hat{p}_\zeta(\mathbf{S}|t)}{\hat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\}}_{\text{Term XIc}},
\end{aligned}$$

where $\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$. It remains to show that the dominating **Term VI** is of order $O(h^2) + O_P\left(\sqrt{\frac{1}{nh^3}}\right)$ in [Section H.3.1](#) and the remainder terms are of order $o_P\left(\sqrt{\frac{1}{nh^3}}\right)$ for any fixed $t \in \mathcal{T}$ in [Section H.3.2](#), [Section H.3.3](#), and [Section H.3.4](#), and [Section H.3.5](#). We shall also derive the asymptotic normality of $\hat{\theta}_{C,DR}(t)$ in [Section H.3.6](#).

H.3.1 Analysis of Term VI for $\hat{\theta}_{C,DR}(t)$

We analyze the variance and bias of **Term VI** separately as follows. By direct calculations, we have that

$$\begin{aligned}
\text{Var}[\text{Term VI}] &= \text{Var} \left\{ \mathbb{P}_n \left[\frac{1}{\sqrt{nh^3}} \cdot \phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] \right\} \\
&= \frac{1}{nh^4} \cdot \text{Var} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \\
&= \frac{1}{nh^4} \cdot \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta^2(\mathbf{S}|t)}{\kappa_2^2 \cdot \bar{p}^2(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^2 \right\} \\
&\quad - \frac{1}{nh^4} \left\{ \mathbb{E} \left[\frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right]^2 \right\} \\
&\stackrel{(i)}{\lesssim} \frac{1}{nh^3} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{\kappa_2^2 \cdot \bar{p}^2(t+uh, \mathbf{s}_1)} \\
&\quad \times \left\{ [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)]^2 + \sigma^2 \right\} d\mathbf{s}_1 du \\
&\stackrel{(ii)}{=} \frac{1}{nh^3} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t) \cdot [p(t, \mathbf{s}_1) + O(h)]}{\kappa_2^2 \cdot [\bar{p}^2(t, \mathbf{s}_1) + O(h^2)]} \\
&\quad \times \left\{ [\mu(t, \mathbf{s}_1) + O(h) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)]^2 + \sigma^2 \right\} d\mathbf{s}_1 du \\
&\stackrel{(iii)}{=} O\left(\frac{1}{nh^3}\right),
\end{aligned}$$

where (i) uses a change of variable and only keeps the dominating first term, (ii) leverages Taylor's expansions, and (iii) utilizes the upper boundedness of $\mu, \bar{\mu}, \bar{\beta}, \bar{p}_\zeta$ under [Assumption A3](#) as well as the fact that \bar{p} is lower bounded away from 0 around the support \mathcal{J} . Now, by Chebyshev's

inequality, we conclude that

$$\begin{aligned}
(\mathbb{P}_n - \mathbb{P}) \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] &= O_P \left(\sqrt{\text{Var} \left\{ \mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] \right\}} \right) \\
&= O_P \left(\sqrt{\frac{1}{nh^3}} \right).
\end{aligned}$$

In addition, we calculate the bias term as:

Bias **[Term VI]**

$$\begin{aligned}
&= \mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \\
&\quad + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{\left(\frac{t_1-t}{h}\right) K\left(\frac{t_1-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{s}_1|t)}{h^2 \kappa_2 \cdot \bar{p}(t_1, \mathbf{s}_1)} \cdot [\mu(t_1, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - (t_1-t) \cdot \bar{\beta}(t, \mathbf{s}_1)] p(t_1, \mathbf{s}_1) d\mathbf{s}_1 dt_1 \\
&\quad + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t)}{h \cdot \kappa_2 \cdot \bar{p}(t+uh, \mathbf{s}_1)} \cdot [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)] p(t+uh, \mathbf{s}_1) d\mathbf{s}_1 du \\
&\quad + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\stackrel{(ii)}{=} \int_{\mathbb{R}} \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \frac{u \cdot K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t) \left[p(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p(t, \mathbf{s}_1) + O(h^4) \right]}{h \cdot \kappa_2 \left[\bar{p}(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \bar{p}(t, \mathbf{s}_1) + O(h^4) \right]} \\
&\quad \times \left[(\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)) + hu (\beta(t, \mathbf{s}_1) - \bar{\beta}(t, \mathbf{s}_1)) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{s}_1) + O(h^4) \right] d\mathbf{s}_1 du \\
&\quad + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \int_{\mathbb{R}} \frac{u \cdot K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t)}{h \cdot \kappa_2} \left[p(t, \mathbf{s}_1) + uh \cdot \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} p(t, \mathbf{s}_1) + O(h^4) \right] \\
&\quad \times \left[\frac{1}{\bar{p}(t, \mathbf{s}_1)} - \frac{uh \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1)}{\bar{p}^2(t, \mathbf{s}_1)} - \frac{u^2 h^2 \cdot \frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s}_1)}{2\bar{p}^2(t, \mathbf{s}_1)} + \frac{u^2 h^2 \left[\frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1) \right]^2}{\bar{p}^3(t, \mathbf{s}_1)} - \frac{u^3 h^3 \cdot \frac{\partial^3}{\partial t^3} \bar{p}(t, \mathbf{s}_1)}{6\bar{p}^2(t, \mathbf{s}_1)} \right. \\
&\quad \left. + \frac{u^3 h^3 \left[\frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1) \right] \left[\frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s}_1) \right]}{\bar{p}^3(t, \mathbf{s}_1)} + O(h^4) \right] \\
&\quad \times \left[(\mu(t, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1)) + hu (\beta(t, \mathbf{s}_1) - \bar{\beta}(t, \mathbf{s}_1)) + \frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + \frac{u^3 h^3}{6} \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{s}_1) + O(h^4) \right] d\mathbf{s}_1 du \\
&\quad + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s}
\end{aligned}$$

$$\begin{aligned}
&= \int \left[\frac{\frac{\partial}{\partial t} p(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - \frac{p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s})}{\bar{p}^2(t, \mathbf{s})} \right] [\mu(t, \mathbf{s}) - \bar{\mu}(t, \mathbf{s})] \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\quad + \int [\beta(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})] \left[\frac{p(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - 1 \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} + h^2 \cdot B_{C,\theta}(t) + O(h^3),
\end{aligned}$$

where (i) uses a change of variable $u = \frac{t_1-t}{h}$ and (ii) applies Taylor's expansions. Here, the complicated bias term $B_{C,\theta}(t)$ is given by

$$\begin{aligned}
B_{C,\theta}(t) &= \frac{\kappa_4}{2\kappa_2} \int \frac{[\mu(t, \mathbf{s}) - \bar{\mu}(t, \mathbf{s})]}{\bar{p}(t, \mathbf{s})} \\
&\quad \times \left[\frac{1}{3} \cdot \frac{\partial^3}{\partial t^3} p(t, \mathbf{s}) - \frac{\partial^2}{\partial t^2} p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) + 2 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \left[\frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \right]^2 \right. \\
&\quad \left. + \frac{6 \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s}) - \frac{\partial^3}{\partial t^3} \bar{p}(t, \mathbf{s}) - 3 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s})}{3\bar{p}(t, \mathbf{s})} \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\quad + \frac{\kappa_4}{2\kappa_2} \int \frac{[\beta(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})]}{\bar{p}(t, \mathbf{s})} \left[\frac{\partial^2}{\partial t^2} p(t, \mathbf{s}) - 2 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \right. \\
&\quad \left. + 2p(t, \mathbf{s}) \cdot \left[\frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \right]^2 - \frac{p(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \bar{p}(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\quad + \frac{\kappa_4}{2\kappa_2} \int \left[\frac{\frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - \frac{p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s})}{\bar{p}^2(t, \mathbf{s})} + \frac{p(t, \mathbf{s}) \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{s})}{3\bar{p}(t, \mathbf{s})} \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s}.
\end{aligned}$$

Under the condition that either $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ or $\bar{p} = p$, we have that

$$\begin{aligned}
&\int \left[\frac{\frac{\partial}{\partial t} p(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - \frac{p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s})}{\bar{p}^2(t, \mathbf{s})} \right] [\mu(t, \mathbf{s}) - \bar{\mu}(t, \mathbf{s})] \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&\quad + \int [\beta(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s})] \left[\frac{p(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - 1 \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= 0
\end{aligned}$$

and

$$\begin{aligned}
B_{C,\theta}(t) &= \begin{cases} \frac{\kappa_4}{2\kappa_2} \int \left[\frac{\frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s})}{\bar{p}(t, \mathbf{s})} - \frac{p(t, \mathbf{s}) \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}) \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s})}{\bar{p}^2(t, \mathbf{s})} + \frac{p(t, \mathbf{s}) \cdot \frac{\partial^3}{\partial t^3} \mu(t, \mathbf{s})}{3\bar{p}(t, \mathbf{s})} \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \int \left[\frac{\partial^3}{\partial t^3} \mu(t, \mathbf{s}) \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{p} = p, \end{cases} \\
&= \begin{cases} \frac{\kappa_4}{6\kappa_2} \int \left\{ \frac{3 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \bar{m}''(t) + p(t, \mathbf{s}) [\bar{m}^{(3)}(t) - 3 \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \cdot \bar{m}''(t)]}{\bar{p}(t, \mathbf{s})} \right\} \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) & \text{when } \bar{p} = p. \end{cases}
\end{aligned}$$

As a result, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we know that

$$\begin{aligned}
\mathbf{Term VI} &= \mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] + \int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \int \beta(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= h^2 B_{C,\theta}(t) + o(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right) \\
&= O(h^2) + O_P \left(\sqrt{\frac{1}{nh^3}} \right).
\end{aligned}$$

As a side note, under some VC-type condition on the kernel function K (Einmahl and Mason, 2005), we can strengthen the above pointwise rate of convergence to the following uniform one as:

$$\sup_{t \in \mathcal{T}} |\mathbf{Term VI}| = O(h^2) + O_P \left(\sqrt{\frac{|\log h|}{nh^3}} \right);$$

see Theorem 4 in Einmahl and Mason (2005) for details.

H.3.2 Analysis of Term VII for $\hat{\theta}_{C,DR}(t)$

Notice that

$$\begin{aligned}
\mathbf{Term VII} &= \int \hat{\beta}(t, \mathbf{s}) [\hat{p}_\zeta(\mathbf{s}|t) - \bar{p}_\zeta(\mathbf{s}|t)] d\mathbf{s} \\
&\leq \int \left| \bar{\beta}(t, \mathbf{s}) + \hat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right| |\hat{p}_\zeta(\mathbf{s}|t) - \bar{p}_\zeta(\mathbf{s}|t)| d\mathbf{s} \\
&\stackrel{(i)}{\lesssim} \int \left[|\bar{\beta}(t, \mathbf{s})| + \left| \hat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right| \right] |\hat{p}_\zeta(\mathbf{s}|t) - \bar{p}_\zeta(\mathbf{s}|t)| p_S(\mathbf{s}) d\mathbf{s} \\
&\stackrel{(ii)}{=} O_P \left(\|\hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} \right) \\
&= O_P(\Upsilon_{5,n}),
\end{aligned}$$

where (i) uses the fact that the marginal density p_S is lower bounded away from 0 within the union set $(\mathcal{S}(t) \ominus \zeta) \cup (\hat{\mathcal{S}}(t) \ominus \zeta)$ and (ii) leverages the boundedness of $\bar{\beta}$ under Assumption A3 as well as $\left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(\Upsilon_{3,n})$ with $\Upsilon_{3,n} \rightarrow 0$ as $n \rightarrow \infty$. Hence, $\sqrt{nh^3} \cdot \mathbf{Term VII} = o_P(1)$ when $\sqrt{nh^3} \cdot \Upsilon_{5,n} = o(1)$.

H.3.3 Analyses of Term VIII and Term IX for $\hat{\theta}_{C,DR}(t)$

By Markov's inequality, we know that

$$\begin{aligned}
\sqrt{nh^3} \cdot \mathbf{Term VIII} &= \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\hat{p}_\zeta(\mathbf{S}|t)}{\hat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \\
&= O_P(\Upsilon_{5,n} + \Upsilon_{6,n}) = o_P(1)
\end{aligned}$$

because

$$\begin{aligned}
& \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right]^2 [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^2 \right\} \\
& \lesssim \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \cdot \frac{[\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)]^2 \bar{p}^2(T, \mathbf{S}) + [\widehat{p}(T, \mathbf{S}) - \bar{p}(T, \mathbf{S})] \bar{p}_\zeta^2(\mathbf{S}|t)}{\widehat{p}^2(T, \mathbf{S}) \cdot \bar{p}^2(T, \mathbf{S})} \right. \\
& \quad \left. \times [(\mu(T, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S}))^2 + \sigma^2] \right\} \\
& \stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{u^2 K^2(u)}{\kappa_2^2} \cdot \frac{[\widehat{p}_\zeta(\mathbf{s}_1|t) - \bar{p}_\zeta(\mathbf{s}_1|t)]^2 \bar{p}^2(t+uh, \mathbf{s}_1) + [\widehat{p}(t+uh, \mathbf{s}_1) - \bar{p}(t+uh, \mathbf{s}_1)] \bar{p}_\zeta^2(\mathbf{s}_1|t)}{\widehat{p}^2(t+uh, \mathbf{s}_1) \cdot \bar{p}^2(t+uh, \mathbf{s}_1)} \\
& \quad \times [(\mu(t+uh, \mathbf{S}) - \bar{\mu}(t, \mathbf{S}) - hu \cdot \bar{\beta}(t, \mathbf{S}))^2 + \sigma^2] p(t+uh, \mathbf{s}_1) d\mathbf{s}_1 du \\
& \stackrel{(ii)}{\lesssim} \|\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2}^2 + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2}^2 \\
& = O_P(\Upsilon_{5,n}^2 + \Upsilon_{6,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{T-t}{h}$ in the integration as well as (ii) leverages the upper boundedness of $\mu, \bar{\mu}, \bar{\beta}$ under Assumption A3 and the lower boundedness on $\frac{\bar{p}_\zeta}{\bar{p}}, \widehat{p}$ away from 0 around the support \mathcal{J} by definition. As a side note, under the VC-type condition on the kernel function K (Einmahl and Mason, 2005) and

$$\sup_{t \in \mathcal{T}} \left[\|\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} \right] = o_P(1),$$

we can strengthen the above pointwise rate of convergence to the following uniform result as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \right| = o_P(1).$$

Similarly, by Markov's inequality, we have that

$$\begin{aligned}
\sqrt{nh^3} \cdot \mathbf{Term IX} &= \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right] \right\} \\
&= O_P(\max\{\Upsilon_{1,n}, h \cdot \Upsilon_{3,n}\}) = o_P(1)
\end{aligned}$$

because

$$\begin{aligned}
& \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta^2(\mathbf{S}|t)}{h \cdot \kappa_2^2 \cdot \bar{p}^2(T, \mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right]^2 \right\} \\
&= \int_{\mathcal{T}} \int_{\mathcal{S}(t)} \frac{\left(\frac{t_1-t}{h}\right)^2 K^2\left(\frac{t_1-t}{h}\right) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t) \cdot p(t_1, \mathbf{s}_1)}{h \cdot \kappa_2^2 \cdot \bar{p}^2(t_1, \mathbf{s}_1)} \left[\bar{\mu}(t, \mathbf{s}_1) - \widehat{\mu}(t, \mathbf{s}_1) + (t_1-t) \left(\bar{\beta}(t, \mathbf{s}_1) - \widehat{\beta}(t, \mathbf{s}_1) \right) \right]^2 d\mathbf{s}_1 dt_1
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{u^2 K^2(u) \cdot \bar{p}_{\zeta}^2(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2^2 \cdot \bar{p}^2(t+uh, \mathbf{s}_1)} \left[\bar{\mu}(t, \mathbf{s}_1) - \hat{\mu}(t, \mathbf{s}_1) + hu \cdot (\bar{\beta}(t, \mathbf{s}_1) - \hat{\beta}(t, \mathbf{s}_1)) \right]^2 d\mathbf{s}_1 du \\
&\stackrel{(ii)}{\lesssim} \|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}^2 + h^2 \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2}^2 \\
&= O_P(\Upsilon_{1,n}^2 + h^2 \Upsilon_{3,n}^2) = o_P(1),
\end{aligned}$$

where (i) uses the change of variable $u = \frac{t_1-t}{h}$ and (ii) leverages the boundedness of p, p_{ζ} under Assumption A3, the lower bound on $\frac{\bar{p}_{\zeta}}{\bar{p}}$ away from 0 around the support \mathcal{J} by definition, the boundedness condition on K under Assumption A5, as well as $\|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{1,n})$ and $h \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(h \cdot \Upsilon_{3,n})$ with $\Upsilon_{1,n}, h \cdot \Upsilon_{3,n} \rightarrow 0$ as $n \rightarrow \infty$. In addition, if $\sup_{t \in \mathcal{T}} \|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} = o_P(1)$ and $\sup_{t \in \mathcal{T}} \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = o_P\left(\frac{1}{h}\right)$, then the above pointwise rate of convergence can be strengthened to the uniform one as:

$$\sup_{t \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_{\zeta}(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T|\mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) (\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S})) \right] \right\} \right| = o_P(1).$$

H.3.4 Analysis of Term X for $\hat{\theta}_{C,DR}(t)$

We first calculate that

$$\begin{aligned}
&\mathbb{E} \left| \sqrt{nh^3} \cdot \text{Term X} \right| \\
&\mathbb{E} \left| \sqrt{\frac{n}{h}} \cdot \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\kappa_2} \left[\frac{\hat{p}_{\zeta}(\mathbf{S}|t)}{\hat{p}(T, \mathbf{S})} - \frac{\bar{p}_{\zeta}(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) (\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S})) \right] \right| \\
&\stackrel{(i)}{\lesssim} \sqrt{nh} \cdot \sqrt{\mathbb{E} \left\{ \frac{K\left(\frac{T-t}{h}\right)}{h} \left[\frac{[\bar{p}_{\zeta}(\mathbf{S}|t) - \hat{p}_{\zeta}(\mathbf{S}|t)]^2}{\hat{p}^2(T, \mathbf{S})} + \frac{[\hat{p}(T, \mathbf{S}) - \bar{p}(T, \mathbf{S})]^2 \bar{p}_{\zeta}^2(\mathbf{S}|t)}{\bar{p}^2(T, \mathbf{S})} \right] \right\}} \\
&\quad \times \sqrt{\mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right)}{h \cdot \kappa_2^2} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + (T-t) (\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S})) \right]^2 \right\}} \\
&= \sqrt{nh} \cdot \sqrt{\int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} K(u) \left[\frac{[\bar{p}_{\zeta}(\mathbf{s}_1|t) - \hat{p}_{\zeta}(\mathbf{s}_1|t)]^2}{\hat{p}^2(t+uh, \mathbf{s}_1)} + \frac{[\hat{p}(t+uh, \mathbf{s}_1) - \bar{p}(t+uh, \mathbf{s}_1)]^2 \bar{p}_{\zeta}^2(\mathbf{s}_1|t)}{\bar{p}^2(t+uh, \mathbf{s}_1)} \right] p(t+uh, \mathbf{s}_1) d\mathbf{s}_1 du} \\
&\quad \times \sqrt{\mathbb{E} \left\{ \int_{\mathbb{R}} \frac{u^2 K(u)}{\kappa_2^2} \cdot \left[\bar{\mu}(t, \mathbf{S}) - \hat{\mu}(t, \mathbf{S}) + hu (\bar{\beta}(t, \mathbf{S}) - \hat{\beta}(t, \mathbf{S})) \right]^2 p_{T|\mathbf{S}}(t+uh|\mathbf{S}) du \right\}} \\
&\lesssim \sqrt{nh} \left[\|\hat{p}_{\zeta}(\mathbf{S}|t) - \bar{p}_{\zeta}(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} \right] \left[\|\hat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} + h \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} \right] \\
&\stackrel{(ii)}{=} o_P(1),
\end{aligned}$$

where (i) uses Cauchy-Schwarz inequality and (ii) leverages our assumption (c) on the doubly robust rate of convergence in the theorem statement. As a result, by Markov's inequality, we

obtain that

$$\begin{aligned}
& \sqrt{nh^3} \cdot \mathbf{Term X} \\
&= \sqrt{\frac{n}{h}} \cdot \mathbb{P}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{\kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T-t) \left(\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right) \right] \right\} \\
&= o_P(1).
\end{aligned}$$

H.3.5 Analysis of Term XI for $\widehat{\theta}_{C,DR}(t)$

By direct calculations under model (1) with some change of variable, we have that

Term XI

$$\begin{aligned}
&= \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{h \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \left[\bar{\beta}(t, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right] \right\} + \int \left[\widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&+ \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \left[\bar{\mu}(t, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) \right] \right\} \\
&+ \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|t)}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|t)}{\bar{p}(T, \mathbf{S})} \right] \left[Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S}) \right] \right\} \\
&= \underbrace{\int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{u^2 K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{\kappa_2 \cdot \bar{p}(t+uh, \mathbf{s}_1)} \left[\bar{\beta}(t, \mathbf{s}_1) - \widehat{\beta}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 du + \int \left[\widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s}}_{\mathbf{Term XIa}} \\
&+ \underbrace{\int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2 \cdot \bar{p}(t+uh, \mathbf{s}_1)} \left[\bar{\mu}(t, \mathbf{s}_1) - \widehat{\mu}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 du}_{\mathbf{Term XIb}} \\
&+ \underbrace{\int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{s}_1|t)}{\widehat{p}(t+uh, \mathbf{s}_1)} - \frac{\bar{p}_\zeta(\mathbf{s}_1|t)}{\bar{p}(t+uh, \mathbf{s}_1)} \right] \left[\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 du}_{\mathbf{Term XIc}}.
\end{aligned}$$

On one hand, when $\bar{p}(t, \mathbf{s}) = p(t, \mathbf{s})$, we know that **Term XIb** = 0, and under Assumption A6 and model (13),

$$\begin{aligned}
\mathbf{Term XIa} &= \int_{\mathbb{R}} \int_{\mathcal{S}(t) \ominus \zeta} \frac{u^2 K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t)}{\kappa_2} \left[\bar{\beta}(t, \mathbf{s}_1) - \widehat{\beta}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 du + \int \left[\widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= \int_{\mathcal{S}(t) \ominus \zeta} p_\zeta(\mathbf{s}_1|t) \left[\bar{\beta}(t, \mathbf{s}_1) - \widehat{\beta}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 + \int \left[\widehat{\beta}(t, \mathbf{s}) - \bar{\beta}(t, \mathbf{s}) \right] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} \\
&= 0.
\end{aligned}$$

In addition, we also have that

Term XIc

$$= \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot [p(t+uh, \mathbf{s}_1) - \widehat{p}(t+uh, \mathbf{s}_1)] \cdot \bar{p}_\zeta(\mathbf{s}_1|t)}{h \cdot \kappa_2 \cdot \widehat{p}(t+uh, \mathbf{s}_1)} \left[\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1) \right] d\mathbf{s}_1 du$$

$$\begin{aligned}
& + \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot [\widehat{p}_{\zeta}(\mathbf{s}_1|t) - \bar{p}_{\zeta}(\mathbf{s}_1|t)] \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2 \cdot \widehat{p}(t+uh, \mathbf{s}_1)} [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)] d\mathbf{s}_1 du \\
& \lesssim \frac{1}{h} \left[\|\widehat{p}_{\zeta}(\mathbf{S}|t) - p_{\zeta}(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right] \\
& = o_P \left(\sqrt{\frac{1}{nh^3}} \right)
\end{aligned}$$

by the upper boundedness of $\mu, \bar{\mu}$ under Assumption A3, the lower boundedness of \widehat{p} away from 0 around the support \mathcal{J} , and our assumption (c) on the doubly robust rate of convergence in the theorem statement. Specifically, since $\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} + h \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(1)$ when $\bar{\mu} \neq \mu$ and $\bar{\beta} \neq \beta$, our assumption (c) ensures that $\|\widehat{p}_{\zeta}(\mathbf{S}|t) - \bar{p}_{\zeta}(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$.

On the other hand, when $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$, we know from Assumptions A4 and A6 on \bar{p}, \bar{p}_{ζ} that $\frac{\bar{p}_{\zeta}}{\bar{p}}$ is bounded away from 0 within the support and thus,

Term XIa

$$\begin{aligned}
& = \int_{\mathbb{R}} \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \frac{u^2 K(u) \cdot \bar{p}_{\zeta}(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{\kappa_2 \cdot \bar{p}(t+uh, \mathbf{s}_1)} [\beta(t, \mathbf{s}_1) - \widehat{\beta}(t, \mathbf{s}_1)] d\mathbf{s}_1 du + \int [\widehat{\beta}(t, \mathbf{s}) - \beta(t, \mathbf{s})] \bar{p}_{\zeta}(\mathbf{s}|t) d\mathbf{s} \\
& \stackrel{(i)}{\lesssim} \left\| \widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \\
& \stackrel{(ii)}{=} o_P \left(\sqrt{\frac{1}{nh^3}} \right),
\end{aligned}$$

where (i) uses the facts that \bar{p}_{ζ} is upper bounded while the marginal density $p_{\mathcal{S}}$ is lower bounded away from 0 within the ζ -interior conditional support $\mathcal{S}(t) \ominus \zeta$ and (ii) utilizes our assumption (c) on the doubly robust rate of convergence in the theorem statement to argue that $\left\| \widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh^3}} \right)$ if $\bar{p} \neq p, \bar{p}_{\zeta} \neq p_{\zeta}$ and $\|\widehat{p}_{\zeta}(\mathbf{S}|t) - p_{\zeta}(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} = O_P(1)$. In addition, we also have that

$$\begin{aligned}
\text{Term XIb} & = \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot \bar{p}_{\zeta}(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2 \cdot \bar{p}(t+uh, \mathbf{s}_1)} [\mu(t, \mathbf{s}_1) - \widehat{\mu}(t, \mathbf{s}_1)] d\mathbf{s}_1 du \\
& \lesssim \frac{1}{h} \|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \\
& = o_P \left(\sqrt{\frac{1}{nh^3}} \right),
\end{aligned}$$

where we again argue from our assumption (c) on the doubly robust rate of convergence in the theorem statement that $\|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$ if $\bar{p} \neq p, \bar{p}_{\zeta} \neq p_{\zeta}$ and $\|\widehat{p}_{\zeta}(\mathbf{S}|t) - p_{\zeta}(\mathbf{S}|t)\|_{L_2} +$

$\sup_{|u-t|\leq h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} = O_P(1)$. Finally, we also derive that

Term XIc

$$\begin{aligned}
&= \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot p(t+uh, \mathbf{s}_1)}{h \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{s}_1|t)}{\widehat{p}(t+uh, \mathbf{s}_1)} - \frac{\bar{p}_\zeta(\mathbf{s}_1|t)}{\bar{p}(t+uh, \mathbf{s}_1)} \right] [\mu(t+uh, \mathbf{s}_1) - \mu(t, \mathbf{s}_1) - hu \cdot \beta(t, \mathbf{s}_1)] d\mathbf{s}_1 du \\
&= \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) \cdot p(t+uh, \mathbf{s}_1) [\widehat{p}_\zeta(\mathbf{s}_1|t) - \bar{p}_\zeta(\mathbf{s}_1|t)]}{h \cdot \kappa_2 \cdot \widehat{p}(t+uh, \mathbf{s}_1)} [\mu(t+uh, \mathbf{s}_1) - \mu(t, \mathbf{s}_1) - hu \cdot \beta(t, \mathbf{s}_1)] d\mathbf{s}_1 du \\
&\quad + \int_{\mathbb{R}} \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \frac{uK(u) \cdot p(t+uh, \mathbf{s}_1) \cdot \bar{p}_\zeta(\mathbf{s}_1|t) [\bar{p}(t+uh, \mathbf{s}_1) - \widehat{p}(t+uh, \mathbf{s}_1)]}{h \cdot \kappa_2 \cdot \widehat{p}(t+uh, \mathbf{s}_1) \cdot \bar{p}(t+uh, \mathbf{s}_1)} \\
&\quad \times [\mu(t+uh, \mathbf{s}_1) - \mu(t, \mathbf{s}_1) - hu \cdot \beta(t, \mathbf{s}_1)] d\mathbf{s}_1 du \\
&\stackrel{(i)}{=} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} \frac{uK(u) [\widehat{p}_\zeta(\mathbf{s}_1|t) - \bar{p}_\zeta(\mathbf{s}_1|t)] \left[\frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + O(h^3) \right] \left[p(t, \mathbf{s}_1) + 2uh \cdot \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + O(h^2) \right]}{h \cdot \kappa_2 [\bar{p}(t, \mathbf{s}_1) + 2uh \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1) + O(h^2)] [1 + O_P(\Upsilon_{6,n})]} d\mathbf{s}_1 du \\
&\quad + \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \sup_{|u-t|\leq h} |\bar{p}(u, \mathbf{s}_1) - \widehat{p}(u, \mathbf{s}_1)| \\
&\quad \times \left| \int_{\mathbb{R}} \frac{u \cdot K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|t) \left[\frac{u^2 h^2}{2} \cdot \frac{\partial^2}{\partial t^2} \mu(t, \mathbf{s}_1) + O(h^3) \right] \left[p(t, \mathbf{s}_1) + 2uh \cdot \frac{\partial}{\partial t} p(t, \mathbf{s}_1) + O(h^2) \right]}{h \cdot \kappa_2 [\bar{p}^2(t, \mathbf{s}_1) + 2uh \cdot \bar{p}(t, \mathbf{s}_1) \cdot \frac{\partial}{\partial t} \bar{p}(t, \mathbf{s}_1) + O(h^2)] [1 + O_P(\Upsilon_{6,n}^2)]} du \right| d\mathbf{s}_1 \\
&= O_P \left(h^2 \|\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} \right) + O_P \left(h^2 \sup_{|u-t|\leq h} \|\bar{p}(u, \mathbf{S}) - \widehat{p}(u, \mathbf{S})\|_{L_2} \right) \\
&= O_P \left(h^2 [\Upsilon_{5,n} + \Upsilon_{6,n}] \right) \\
&\stackrel{(ii)}{=} o_P \left(\sqrt{\frac{1}{nh^3}} \right),
\end{aligned}$$

where (i) applies Taylor's expansion and mean-value theorem for integrals as well as uses the fact that the difference between \bar{p} and \widehat{p} is small when $\sup_{|u-t|\leq h} \|\widehat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2} = O_P(\Upsilon_{6,n})$, while (ii) leverages the arguments that $\sqrt{nh^3} \cdot h^2 = \sqrt{nh^7} \rightarrow \sqrt{c_3} \in [0, \infty)$ and $\Upsilon_{5,n}, \Upsilon_{6,n} \rightarrow 0$ as $n \rightarrow \infty$.

H.3.6 Asymptotic Normality of $\widehat{\theta}_{C,DR}(t)$

For the asymptotic normality of $\widehat{\theta}_{C,DR}(t)$, it follows from the Lyapunov central limit theorem. Specifically, we already show in [Section H.3.1](#) and subsequent subsections that

$$\begin{aligned}
&\sqrt{nh^3} \left[\widehat{\theta}_{C,DR}(t) - \theta(t) \right] \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + \sqrt{h^3} \left[\int \bar{\beta}(t, \mathbf{s}) \cdot \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} - \theta(t) \right] \right\} + o_P(1) \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + o_P(1)
\end{aligned}$$

with

$$\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$$

and $V_{C,\theta}(t) = \mathbb{E} \left[\phi_{C,h,t}^2(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] = O(1)$ by our calculation in **Term VI**. Then,

$$\sum_{i=1}^n \text{Var} \left[\frac{1}{\sqrt{n}} \cdot \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] = O(1)$$

and

$$\begin{aligned} & \sum_{i=1}^n \mathbb{E} \left| \frac{1}{\sqrt{n}} \cdot \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right|^{2+c_1} \\ &= \mathbb{E} \left| \frac{\left(\frac{T-t}{h}\right)^{2+c_1} K^{2+c_1}\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta^{2+c_1}(\mathbf{S}|t) \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]^{2+c_1}}{n^{\frac{c_1}{2}} h^{1+\frac{c_1}{2}} \cdot \kappa_2^{2+c_1} \cdot \bar{p}^{2+c_1}(T, \mathbf{S})} \right| \\ &\lesssim \int_{\mathbb{R}} \int_{\bar{\mathcal{S}}(t) \ominus \zeta} \frac{u^{2+c_1} K^{2+c_1}(u) \cdot \bar{p}_\zeta^{2+c_1}(\mathbf{s}_1|t) \left[[\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)]^{2+c_1} + \mathbb{E}|\epsilon|^{2+c_1} \right]}{\sqrt{(nh)^{c_1}} \bar{p}^{2+c_1}(t+uh, \mathbf{s}_1)} \\ &\quad \times p(t+uh, \mathbf{s}_1) d\mathbf{s}_1 du \\ &= O \left(\sqrt{\frac{1}{(nh)^{c_1}}} \right) = o(1) \end{aligned}$$

by the upper boundedness of $\mu, \bar{\mu}, p$ under Assumptions [A3](#) and [A4](#), the lower boundedness of \bar{p} away from 0 around the support \mathcal{J} , the assumption that $\mathbb{E}|\epsilon|^{2+c_1} < \infty$, and the requirement that $nh^3 \rightarrow \infty$ as $n \rightarrow \infty$. Hence, the Lyapunov condition holds, and we have that

$$\sqrt{nh^3} \left[\widehat{\theta}_{C,DR}(t) - \theta(t) - h^2 B_{C,\theta}(t) \right] \xrightarrow{d} \mathcal{N}(0, V_{C,\theta}(t))$$

after subtracting the dominating bias term $h^2 B_{C,\theta}(t)$ of $\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)$ that we have computed in **Term VI**. The proof is thus completed. \square

I Asymptotic Theory of Estimating $m(t)$ Without Positivity

Under the additive confounding model [\(13\)](#), Proposition [C.1](#) implies that we can define the integral RA estimator [\(17\)](#) as well as the integral IPW and DR estimators of $m(t)$ based on [\(26\)](#) and [\(25\)](#) as:

$$\widehat{m}_{C,IPW}(t) = \frac{1}{n} \sum_{i=1}^n \left[Y_i + \int_{\tilde{t}=T_i}^{\tilde{t}=t} \widehat{\theta}_{C,IPW}(\tilde{t}) d\tilde{t} \right] \quad \text{and} \quad \widehat{m}_{C,DR}(t) = \frac{1}{n} \sum_{i=1}^n \left[Y_i + \int_{\tilde{t}=T_i}^{\tilde{t}=t} \widehat{\theta}_{C,DR}(\tilde{t}) d\tilde{t} \right].$$

We prove in the following corollary that these integral estimators are consistent to $m(t)$ without assuming the positivity condition.

Corollary I.1 (Consistency of estimating $m(t)$ without positivity). *Suppose that Assumptions A1, A3, A4, A5, and A6 are valid under the additive confounding model (13). In addition, $\hat{\mu}, \hat{\beta}, \hat{p}_\zeta, \hat{p}$ are constructed on a data sample independent of $\{(Y_i, T_i, \mathbf{S}_i)\}_{i=1}^n$. For any fixed $t \in \mathcal{T}$, we let $\bar{\mu}(t, \mathbf{s}), \bar{\beta}(t, \mathbf{s}), \bar{p}_\zeta(\mathbf{s}|t)$, and $\bar{p}(t, \mathbf{s})$ be fixed bounded functions to which $\hat{\mu}(t, \mathbf{s}), \hat{\beta}(t, \mathbf{s}), \hat{p}_\zeta(\mathbf{s}|t)$, and $\hat{p}(t, \mathbf{s})$ converge under the rates of convergence as:*

$$\begin{aligned} \sup_{t \in \mathcal{T}} \left\| \hat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} &= O_P(\tilde{\Upsilon}_{3,n}), \quad \sup_{t \in \mathcal{T}} \sup_{\mathbf{s} \in \mathcal{S}} \left| \hat{F}_{\mathbf{S}|T}(\mathbf{s}|t) - F_{\mathbf{S}|T}(\mathbf{s}|t) \right| = O_P(\tilde{\Upsilon}_{4,n}), \\ \sup_{t \in \mathcal{T}} \left\| \hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t) \right\|_{L_2} &= O_P(\tilde{\Upsilon}_{5,n}), \quad \text{and} \quad \sup_{u \in \mathcal{T} \oplus h} \left\| \hat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S}) \right\|_{L_2} = O_P(\tilde{\Upsilon}_{6,n}), \end{aligned}$$

where $\mathcal{T} \oplus h = \{u \in \mathbb{R} : \inf_{t \in \mathcal{T}} |u - t| \leq h\}$ and $\tilde{\Upsilon}_{3,n}, \tilde{\Upsilon}_{4,n}, \tilde{\Upsilon}_{5,n}, \tilde{\Upsilon}_{6,n} \rightarrow 0$ as $n \rightarrow \infty$. Then, as $h \rightarrow 0$ and $nh^3 \rightarrow \infty$, we have that

$$\begin{aligned} \sup_{t \in \mathcal{T}} |\hat{m}_{C,RA}(t) - m(t)| &= O_P \left(\tilde{\Upsilon}_{3,n} + \tilde{\Upsilon}_{4,n} + \sup_{t \in \mathcal{T}} \left\| \bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} + \frac{1}{\sqrt{n}} \right), \\ \sup_{t \in \mathcal{T}} |\hat{m}_{C,IPW}(t) - m(t)| &= O(h^2) + O_P \left(\sqrt{\frac{|\log h|}{nh^3}} + \tilde{\Upsilon}_{5,n} + \tilde{\Upsilon}_{6,n} + \sup_{u \in \mathcal{T} \oplus h} \left\| \bar{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2} \right). \end{aligned}$$

If, in addition, we assume that

(a) \bar{p}, \bar{p}_ζ satisfy Assumptions A4 and A6 as well as $\sqrt{nh^3} \cdot \tilde{\Upsilon}_{5,n} = o(1)$;

(b) either $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$ or $\bar{p} = p$;

(c) $\sup_{t \in \mathcal{T}} \left[\left\| \hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t) \right\|_{L_2} + \sup_{|u-t| \leq h} \left\| \hat{p}(u, \mathbf{S}) - p(u, \mathbf{S}) \right\|_{L_2} \right] \left[\left\| \hat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S}) \right\|_{L_2} + h \left\| \hat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right] = o_P \left(\frac{1}{\sqrt{nh}} \right)$,

then

$$\begin{aligned} &\sqrt{nh^3} [\hat{m}_{C,DR}(t) - m(t)] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \left\{ \phi_{C,h,\tilde{t}}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + \sqrt{h^3} \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) d\mathbf{s} \right\} d\tilde{t} \right] \right\} + o_P(1) \end{aligned}$$

when $nh^7 \rightarrow c_3$ for some finite number $c_3 \geq 0$, where $\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)$ is defined in Theorem 5. Furthermore,

$$\sqrt{nh^3} \left\{ \hat{m}_{C,DR}(t) - m(t) - h^2 \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t B_{C,\theta}(\tilde{t}) d\tilde{t} \right] \right\} \xrightarrow{d} \mathcal{N}(0, V_{C,m}(t))$$

with $V_{C,m}(t) = \mathbb{E} \left[\left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right] \right\}^2 \right]$ with $B_{C,\theta}(t)$ defined in Theorem 5.

Proof of Corollary I.1. Recall from (17) that our integral estimator of $m(t)$ is defined as:

$$\widehat{m}_C(t) = \frac{1}{n} \sum_{i=1}^n \left[Y_i + \int_{\tilde{T}_i}^{\tilde{T}=t} \widehat{\theta}_C(\tilde{t}) d\tilde{t} \right],$$

where $\widehat{\theta}_C(t)$ can be either RA (16), IPW (25), or DR (26) estimators. By (15) in Proposition C.1, we have that

$$\widehat{m}_C(t) - m(t) = \frac{1}{n} \sum_{i=1}^n Y_i - \mathbb{E}(Y) + \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \theta(\tilde{t}) d\tilde{t} - \mathbb{E} \left[\int_T^t \theta(\tilde{t}) d\tilde{t} \right] + \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \left[\widehat{\theta}_C(\tilde{t}) - \theta(\tilde{t}) \right] d\tilde{t}. \quad (40)$$

Under Assumption A3 and the condition $\mathbb{E}|\epsilon^{2+c_1}| < \infty$ for some constant $c_1 > 0$, it is valid that $\text{Var}(Y) < \infty$ and $\text{Var} \left[\int_T^t \theta(\tilde{t}) d\tilde{t} \right] < \infty$. Thus, by Chebyshev's inequality, we know that

$$\frac{1}{n} \sum_{i=1}^n Y_i - \mathbb{E}(Y) = O_P \left(\frac{1}{\sqrt{n}} \right) \quad \text{and} \quad \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \theta(\tilde{t}) d\tilde{t} - \mathbb{E} \left[\int_T^t \theta(\tilde{t}) d\tilde{t} \right] = O_P \left(\frac{1}{\sqrt{n}} \right).$$

Furthermore, under Assumption A3, $\left| \int_T^{t_1} \theta(\tilde{t}) d\tilde{t} - \int_T^{t_2} \theta(\tilde{t}) d\tilde{t} \right| \leq \sup_{t \in \mathcal{T}} |\theta(t)| \cdot |t_1 - t_2|$. Together with the compactness of \mathcal{T} and Example 19.7 in van der Vaart (1998), we also deduce that

$$\sup_{t \in \mathcal{T}} \left| \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \theta(\tilde{t}) d\tilde{t} - \mathbb{E} \left[\int_T^t \theta(\tilde{t}) d\tilde{t} \right] \right| = O_P \left(\frac{1}{\sqrt{n}} \right).$$

Therefore, plugging the above rates of convergence back into (40), we conclude that

$$\widehat{m}_C(t) - m(t) = \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \left[\widehat{\theta}_C(\tilde{t}) - \theta(\tilde{t}) \right] d\tilde{t} + O_P \left(\frac{1}{\sqrt{n}} \right). \quad (41)$$

Now, we derive the uniform rates of convergence for $\widehat{m}_{C,RA}(t)$ and $\widehat{m}_{C,IPW}(t)$ when $\widehat{\theta}_C(t)$ are given by $\widehat{\theta}_{C,RA}(t)$ in (16) and $\widehat{\theta}_{C,IPW}(t)$ in (25) respectively in Section I.1 and Section I.2. We also prove the asymptotic linearity, double robustness, and asymptotic normality of $\widehat{m}_{C,DR}(t)$ when $\widehat{\theta}_C(t)$ is given by $\widehat{\theta}_{C,DR}(t)$ in (26) in Section I.3.

I.1 Uniform Rate of Convergence of $\widehat{m}_{C,RA}(t)$

Firstly, we derive the rate of convergence for $\widehat{m}_{C,RA}(t)$ when the derivative estimator is given by $\widehat{\theta}_{C,RA}(t)$ in (16). By (41), we have that

$$\sup_{t \in \mathcal{T}} |\widehat{m}_{C,RA}(t) - m(t)| \leq \sup_{t \in \mathcal{T}} \left| \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \left[\widehat{\theta}_{C,RA}(\tilde{t}) - \theta(\tilde{t}) \right] d\tilde{t} \right| + O_P \left(\frac{1}{\sqrt{n}} \right)$$

$$\leq \sup_{t \in \mathcal{T}} \left[\left| \widehat{\theta}_{C,RA}(t) - \theta(t) \right| \cdot \frac{1}{n} \sum_{i=1}^n |t - T_i| \right] + O_P \left(\frac{1}{\sqrt{n}} \right).$$

The compactness of \mathcal{T} by Assumption A3 and Markov's inequality imply that $\sup_{t \in \mathcal{T}} \frac{1}{n} \sum_{i=1}^n |t - T_i| = O_P(1)$. Therefore, by Theorem 5, we conclude that

$$\begin{aligned} \sup_{t \in \mathcal{T}} |\widehat{m}_{C,RA}(t) - m(t)| &\leq \sup_{t \in \mathcal{T}} \left| \widehat{\theta}_{C,RA}(t) - \theta(t) \right| \cdot \sup_{t \in \mathcal{T}} \left[\frac{1}{n} \sum_{i=1}^n |t - T_i| \right] + O_P \left(\frac{1}{\sqrt{n}} \right) \\ &= O_P \left(\bar{\Upsilon}_{3,n} + \bar{\Upsilon}_{4,n} + \sup_{t \in \mathcal{T}} \|\bar{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2} + \frac{1}{\sqrt{n}} \right). \end{aligned}$$

I.2 Uniform Rate of Convergence of $\widehat{m}_{C,IPW}(t)$

Secondly, we derive the rate of convergence for $\widehat{m}_{C,IPW}(t)$ when the derivative estimator is given by $\widehat{\theta}_{C,IPW}(t)$ in (25). Similar to our arguments in Section I.1, we derive from our results in the proof of Theorem 5 (Section H.2) that

$$\begin{aligned} \sup_{t \in \mathcal{T}} |\widehat{m}_{C,IPW}(t) - m(t)| &\leq \sup_{t \in \mathcal{T}} \left| \widehat{\theta}_{C,IPW}(t) - \theta(t) \right| \cdot \sup_{t \in \mathcal{T}} \left[\frac{1}{n} \sum_{i=1}^n |t - T_i| \right] + O_P \left(\frac{1}{\sqrt{n}} \right) \\ &= O(h^2) + O_P \left(\sqrt{\frac{|\log h|}{nh^3}} + \bar{\Upsilon}_{5,n} + \bar{\Upsilon}_{6,n} + \sup_{u \in \mathcal{T} \oplus h} \|\bar{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right). \end{aligned}$$

I.3 Asymptotic Properties of $\widehat{m}_{C,DR}(t)$

Finally, using the asymptotic properties of $\widehat{\theta}_{C,DR}(t)$ in Theorem 5, we shall establish the asymptotic properties of $\widehat{m}_{C,DR}(t)$ when the derivative function is given by $\widehat{\theta}_{C,DR}(t)$ in (26). Recall from the proof of Theorem 5 (specifically, Section H.3) that

$$\begin{aligned} \widehat{\theta}_{C,DR}(t) - \theta(t) &= \mathbb{P}_n \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,t}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] + \int [\bar{\beta}(t, \mathbf{s}) - \beta(t, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} + A_{n,h}(t), \end{aligned}$$

where $\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{(T-t)K(\frac{T-t}{h}) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})}} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})]$ and $A_{n,h}(t) = o_P \left(\sqrt{\frac{1}{nh^3}} \right)$ consists of **Term VII–XI** for $\widehat{\theta}_{C,DR}(t)$ in Section H.3. Then, by (41) and model (13), we know that

$$\begin{aligned} \widehat{m}_{C,DR}(t) - m(t) &= \frac{1}{n} \sum_{i_1=1}^n \int_{T_{i_1}}^t \left\{ \frac{1}{n} \sum_{i_2=1}^n \frac{\left(\frac{T_{i_2}-\tilde{t}}{h} \right) K \left(\frac{T_{i_2}-\tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_{i_2}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T_{i_2}, \mathbf{S}_{i_2})} \cdot [Y_{i_2} - \bar{\mu}(\tilde{t}, \mathbf{S}_{i_2}) - (T_{i_2} - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S}_{i_2})] \right\} d\tilde{t} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} + \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t A_{n,h}(\tilde{t}) d\tilde{t} + O_P\left(\frac{1}{\sqrt{n}}\right) \\
= & \underbrace{\frac{1}{n^2} \sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y_{i_2}, T_{i_2}, \mathbf{S}_{i_2}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} + \mathbb{E} \left\{ \int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right\}}_{\text{Term I}} \\
& + \underbrace{\frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} - \mathbb{E} \left\{ \int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right\}}_{\text{Term II}} \\
& + \underbrace{\frac{1}{n} \sum_{i=1}^n \int_{T_i}^t A_{n,h}(\tilde{t}) d\tilde{t} + O_P\left(\frac{1}{\sqrt{n}}\right)}_{\text{Term III}}.
\end{aligned}$$

As for **Term II**, we know from Assumptions [A3](#) and [A4](#) that $\mathbb{E} \left\{ \left[\int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right]^2 \right\} < \infty$ for any $t \in \mathcal{T}$. By the central limit theorem,

$$\begin{aligned}
\text{Term II} &= \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} - \mathbb{E} \left\{ \int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right\} \\
&= O_P\left(\frac{1}{\sqrt{n}}\right).
\end{aligned}$$

Thus, it remains to derive the asymptotic linearity of $\widehat{m}_{C,DR}(t) - m(t)$ from **Term I** in [Section I.3.1](#) and argue that **Term III** = $o_P\left(\sqrt{\frac{1}{nh^3}}\right)$ in [Section I.3.2](#).

I.3.1 Analysis of Term I for $\widehat{m}_{C,DR}(t)$

Notice that the first term in **Term I** takes a form of V-statistics with a symmetric ‘kernel’ defined as:

$$\begin{aligned}
\Lambda_t(\mathbf{U}_{i_1}, \mathbf{U}_{i_2}) &= \frac{1}{2} \int_{T_{i_1}}^t \frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y_{i_2}, T_{i_2}, \mathbf{S}_{i_2}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \\
&\quad + \frac{1}{2} \int_{T_{i_2}}^t \frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y_{i_1}, T_{i_1}, \mathbf{S}_{i_1}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t},
\end{aligned}$$

where $\mathbf{U}_i \equiv (Y_i, T_i, \mathbf{S}_i)$ for $i = 1, \dots, n$. By Pascal’s rule, we know that

$$\begin{aligned}
\text{Term I} &= \mathbb{P}_n^2 \Lambda_t + \mathbb{E} \left\{ \int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right\} \\
&= 2(\mathbb{P}_n - \mathbb{P}) \mathbb{P} \Lambda_t + (\mathbb{P}_n - \mathbb{P})^2 \Lambda_t + \mathbb{P}^2 \Lambda_t + \mathbb{E} \left\{ \int_T^t \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds d\tilde{t} \right\} \\
&= \underbrace{\mathbb{E}_{T_1} \left\{ \int_{T_1}^t (\mathbb{P}_n - \mathbb{P}) \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] d\tilde{t} \right\}}_{\text{Term Ia}}
\end{aligned}$$

$$\begin{aligned}
& + \underbrace{(\mathbb{P}_n - \mathbb{P}) \left\{ \int_T^t \mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] d\tilde{t} \right\}}_{\text{Term Ib}} \\
& + \underbrace{(\mathbb{P}_n - \mathbb{P})^2 \left\{ \int_{T_{i_1}}^t \frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y_{i_2}, T_{i_2}, \mathbf{S}_{i_2}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right\}}_{\text{Term Ic}} \\
& + \underbrace{\mathbb{E}_{T_1} \left\{ \int_{T_1}^t \left(\mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] + \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) d\mathbf{s} \right) d\tilde{t} \right\}}_{\text{Term Id}},
\end{aligned}$$

where we use the shorthand notation $\text{P}\Lambda_t$ referring to the function $\mathbf{U}_{i_1} \mapsto \int \Lambda_t(\mathbf{U}_{i_1}, \mathbf{u}_{i_2}) d\text{P}(\mathbf{u}_{i_2})$ and $\text{P}^2\Lambda_t = \int \int \Lambda_t(\mathbf{u}_{i_1}, \mathbf{u}_{i_2}) d\text{P}(\mathbf{u}_{i_1})d\text{P}(\mathbf{u}_{i_2})$.

We shall show that the dominating terms **Term Ia** and **Term Id** are of orders $O_P\left(\sqrt{\frac{1}{nh^3}}\right)$ and $O(h^2)$ respectively, and the remainder terms **Term Ib** and **Term Ic** are of order $o_P\left(\sqrt{\frac{1}{nh^3}}\right)$ as follows.

- **Term Ia:** By our calculations in [Section H.3.1](#), we know that

Var [**Term Ia**]

$$\begin{aligned}
& = \text{Var} \left[(\mathbb{P}_n - \mathbb{P}) \left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] d\tilde{t} \right] \right\} \right] \\
& = \frac{1}{n} \cdot \text{Var} \left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] d\tilde{t} \right] \right\} \\
& = \frac{1}{n} \cdot \mathbb{E} \left[\left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] d\tilde{t} \right] \right\}^2 \right] - \frac{1}{n} \cdot \text{P}^2\Lambda_t \\
& \stackrel{(i)}{\leq} \frac{1}{nh^4} \cdot \mathbb{E}_{T_1} \left[|t - T_1| \int_{T_1}^t \mathbb{E} \left\{ \frac{\left(\frac{T-t}{h}\right)^2 K^2\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta^2(\mathbf{S}|\tilde{t})}{\kappa_2^2 \cdot \bar{p}^2(T, \mathbf{S})} \cdot [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})]^2 \right\} d\tilde{t} \right] - \frac{1}{n} \cdot \text{P}^2\Lambda_t \\
& \stackrel{(ii)}{\sim} \frac{1}{nh^3} \int_{\mathcal{T}} \int_{\mathbb{R}} \int_{\mathcal{S}(t+uh)} |t - t_1| \cdot p_T(t_1) \cdot \frac{u^2 K^2(u) \cdot \bar{p}_\zeta^2(\mathbf{s}_1|t) \cdot p(t+uh, \mathbf{s}_1)}{\kappa_2^2 \cdot \bar{p}^2(t+uh, \mathbf{s}_1)} \\
& \quad \times \left\{ [\mu(t+uh, \mathbf{s}_1) - \bar{\mu}(t, \mathbf{s}_1) - hu \cdot \bar{\beta}(t, \mathbf{s}_1)]^2 + \sigma^2 \right\} d\mathbf{s}_1 dudt_1 \\
& \stackrel{(iii)}{=} O_P\left(\frac{1}{nh^3}\right),
\end{aligned}$$

where (i) follows from Jensen's inequality on the squared function and Cauchy-Schwarz inequality as $\left[\int_{T_1}^t g(\tilde{t}) d\tilde{t}\right]^2 \leq \left[\int_{T_1}^t g(\tilde{t}) d\tilde{t}\right] \cdot |t - T_1|$ for the function $g(\tilde{t}) = \mathbb{E} \left[\phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right]$, (ii) uses a change of variable and only keeps the dominating first term, and (iii) leverages our arguments in [Section H.3.1](#). Moreover, Var $\left[\sqrt{nh^3} \cdot \text{Term Ia}\right]$ is strictly positive as long as Var $(\epsilon) = \sigma^2 > 0$.

Then, by Chebyshev's inequality, we obtain that

$$\mathbf{Term Ia} = O_P \left(\sqrt{\text{Var} [\mathbf{Term Ia}]} \right) = O_P \left(\sqrt{\frac{1}{nh^3}} \right).$$

- **Term Id:** By our calculation of the bias term in [Section H.3.1](#), we know that

$$\mathbf{Term Id} = h^2 \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t B_{C,\theta}(\tilde{t}) dt \right] + O(h^3) = O(h^2),$$

where $B_{C,\theta}(t)$ is defined in [Theorem 5](#) as:

$$B_{C,\theta}(t) = \begin{cases} \frac{\kappa_4}{6\kappa_2} \int \left\{ \frac{3 \frac{\partial}{\partial t} p(t, \mathbf{s}) \cdot \bar{m}''(t) + p(t, \mathbf{s}) [\bar{m}^{(3)}(t) - 3 \frac{\partial}{\partial t} \log \bar{p}(t, \mathbf{s}) \cdot \bar{m}''(t)]}{\bar{p}(t, \mathbf{s})} \right\} \bar{p}_\zeta(\mathbf{s}|t) d\mathbf{s} & \text{when } \bar{\mu} = \mu \text{ and } \bar{\beta} = \beta, \\ \frac{\kappa_4}{6\kappa_2} \cdot \bar{m}^{(3)}(t) & \text{when } \bar{p} = p. \end{cases}$$

- **Term Ib:** By Chebyshev's inequality, we have that

$$\begin{aligned} \mathbf{Term Ib} &= (\mathbb{P}_n - \mathbb{P}) \left\{ \int_T^t \mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] dt \right\} \\ &= O_P \left(\sqrt{\frac{\text{Var} \left\{ \int_T^t \mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] dt \right\}}{n}} \right) \\ &\stackrel{(i)}{=} O_P \left(\sqrt{\frac{1}{nh^2}} \right) = o_P \left(\sqrt{\frac{1}{nh^3}} \right). \end{aligned}$$

Here, the equality (i) above follows from the calculation that

$$\begin{aligned} &\text{Var} \left\{ \int_T^t \mathbb{E} \left[\frac{1}{\sqrt{h^3}} \cdot \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) \right] dt \right\} \\ &= \int_{\mathcal{T}} \left\{ \int_{t_2}^t \int_{\mathcal{T}} \int_{\mathcal{S}(t_1)} \frac{\left(\frac{t_1 - \tilde{t}}{h} \right) K \left(\frac{t_1 - \tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{s}_1|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(t_1, \mathbf{s}_1)} \cdot [\mu(t_1, \mathbf{s}_1) - \bar{\mu}(\tilde{t}, \mathbf{s}_1) - (t_1 - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{s}_1)] d\mathbf{s}_1 dt_1 d\tilde{t} \right\}^2 p_T(t_2) dt_2 \\ &\stackrel{(ii)}{=} \int_{\mathcal{T}} \left\{ \int_{t_2}^t \int_{\mathbb{R}} \int_{\mathcal{S}(\tilde{t}+uh)} \frac{u \cdot K(u) \cdot \bar{p}_\zeta(\mathbf{s}_1|\tilde{t})}{h \cdot \kappa_2 \cdot \bar{p}(\tilde{t} + uh, \mathbf{s}_1)} \cdot [\mu(\tilde{t} + uh, \mathbf{s}_1) - \bar{\mu}(\tilde{t}, \mathbf{s}_1) - uh \cdot \bar{\beta}(\tilde{t}, \mathbf{s}_1)] d\mathbf{s}_1 du d\tilde{t} \right\}^2 p_T(t_2) dt_2 \\ &\stackrel{(iii)}{=} O \left(\frac{1}{h^2} \right), \end{aligned}$$

where (ii) utilizes a change of variable and (iii) utilizes the upper boundedness of $\mu, \bar{\mu}, \bar{\beta}, \bar{p}_\zeta, p_T$ under Assumptions [A3](#) and [A4](#) as well as the fact that \bar{p} is lower bounded away from 0 around the support \mathcal{J} .

- **Term Ic:** Recall from Assumption [A5\(c\)](#) that $\mathcal{K} = \left\{ t' \mapsto \left(\frac{t' - t}{h} \right)^{k_1} K \left(\frac{t' - t}{h} \right) : t \in \mathcal{T}, h > 0, k_1 = 0, 1 \right\}$

is a bounded VC-type class of measurable functions on \mathbb{R} . Under Assumption A3 and the condition that $\mathbb{E}|e^{2+c_1}| < \infty$, we deduce by Theorem 4 in Einmahl and Mason (2005) that with probability 1,

$$\begin{aligned} & \sup_{t \in \mathcal{T}} |\mathbb{G}_n [\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)]| \\ &= \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{G}_n \left\{ \frac{\left(\frac{T-t}{h}\right) K\left(\frac{T-t}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T-t) \cdot \bar{\beta}(t, \mathbf{S})] \right\} \right| \\ &= O\left(\sqrt{|\log h|}\right) \end{aligned}$$

when $\frac{|\log h|}{\log \log n} \rightarrow \infty$. Thus, by Chebyshev's inequality,

$$\begin{aligned} \sqrt{nh^3} \cdot \mathbf{Term Ic} &= (\mathbb{P}_n - \mathbb{P}) \left\{ \int_{T_{i_1}}^t \mathbb{G}_n [\phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)] d\tilde{t} \right\} \\ &= O_P \left(\sqrt{\frac{1}{n} \cdot \mathbb{E} \left\{ \left[\int_{T_{i_1}}^t \mathbb{G}_n [\phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)] d\tilde{t} \right]^2 \right\}} \right) \\ &= O_P \left(\sqrt{\frac{|\log h|}{n}} \right) = o_P \left(\sqrt{\frac{1}{nh^3}} \right). \end{aligned}$$

Here, the last equality follows from the calculation that

$$\begin{aligned} \mathbb{E} \left\{ \left[\int_{T_{i_1}}^t \mathbb{G}_n [\phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)] d\tilde{t} \right]^2 \right\} &\stackrel{(i)}{\leq} \mathbb{E} \left[|t - T_{i_1}|^2 \cdot \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{G}_n [\phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta)] \right|^2 \right] \\ &= O(|\log h|), \end{aligned}$$

where (i) applies the mean-value theorem for integrals.

As a summary for this subsection, we conclude that

$$\begin{aligned} & \sqrt{nh^3} \cdot \mathbf{Term I} \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \left\{ \phi_{C,h,\tilde{t}}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + \sqrt{h^3} \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) d\mathbf{s} \right\} d\tilde{t} \right] \right\} + o_P(1). \end{aligned}$$

I.3.2 Analysis of Term III for $\hat{m}_{C,DR}(t)$

Recall from those **Term VII–XI** for $\hat{\theta}_{C,DR}(t)$ in Section H.3 that **Term III** for $\hat{m}_{C,DR}(t)$ here is given by

Term III

$$\begin{aligned}
&= \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t A_{n,h}(\tilde{t}) d\tilde{t} \\
&= \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \int \widehat{\beta}(\tilde{t}, \mathbf{s}) [\widehat{p}_\zeta(\mathbf{s}|\tilde{t}) - \bar{p}_\zeta(\mathbf{s}|\tilde{t})] ds d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIa}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] \right\} d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIb}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(\tilde{t}, \mathbf{S}) + (T - \tilde{t}) [\bar{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(\tilde{t}, \mathbf{S})]] \right\} d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIc}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \mathbb{P}_n \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [\bar{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(\tilde{t}, \mathbf{S}) + (T - \tilde{t}) [\bar{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(\tilde{t}, \mathbf{S})]] \right\} d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIId}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \left(\mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right)^2 K\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(\tilde{t}, \mathbf{S})] \right\} + \int [\widehat{\beta}(\tilde{t}, \mathbf{s}) - \bar{\beta}(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds \right) d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIe}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\bar{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(\tilde{t}, \mathbf{S})] \right\} d\tilde{t} \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIf}} \\
&+ \frac{1}{n} \sum_{i=1}^n \int_{T_i}^t \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] \right\} d\tilde{t}. \\
&\quad \underbrace{\hspace{10em}}_{\text{Term IIIg}}
\end{aligned}$$

We shall argue that all these terms above are of order $o_P\left(\sqrt{\frac{1}{nh^3}}\right)$ respectively as follows.

- **Term IIIa:** By direct calculations, we know that

$$\begin{aligned}
\sqrt{nh^3} \cdot \text{Term IIIa} &= \sqrt{\frac{h^3}{n}} \sum_{i=1}^n \int_{T_i}^t \int \widehat{\beta}(\tilde{t}, \mathbf{s}) [\widehat{p}_\zeta(\mathbf{s}|\tilde{t}) - \bar{p}_\zeta(\mathbf{s}|\tilde{t})] ds d\tilde{t} \\
&\leq \left(\sqrt{\frac{h^3}{n}} \sum_{i=1}^n |t - T_i| \right) \sup_{\tilde{t} \in \mathcal{T}} \left| \int \widehat{\beta}(\tilde{t}, \mathbf{s}) [\widehat{p}_\zeta(\mathbf{s}|\tilde{t}) - \bar{p}_\zeta(\mathbf{s}|\tilde{t})] ds \right| \\
&= O_P\left(\sqrt{nh^3} \cdot \bar{\Upsilon}_{5,n}\right) = o_P(1),
\end{aligned}$$

where in the last equality, we use the compactness of the marginal support $\mathcal{T} \subset \mathbb{R}$ to argue that $\frac{1}{n} \sum_{i=1}^n |t - T_i| = O_P(1)$ for any $t \in \mathcal{T}$ and utilize our derivations in [Section H.3.2](#) to obtain that

$$\sup_{\tilde{t} \in \mathcal{T}} \left| \int \hat{\beta}(\tilde{t}, \mathbf{s}) [\hat{p}_\zeta(\mathbf{s}|\tilde{t}) - \bar{p}_\zeta(\mathbf{s}|\tilde{t})] d\mathbf{s} \right| = O_P \left(\|\hat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} \right) = O_P(\tilde{\Upsilon}_{5,n}).$$

• **Term IIIb:** Notice that **Term IIIb** can be written in a form of V-statistics. Specifically,

$\sqrt{nh^3} \cdot$ **Term IIIb**

$$\begin{aligned} &= \frac{1}{\sqrt{n}} \sum_{i_1=1}^n \int_{T_{i_1}}^t (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\hat{p}_\zeta(\mathbf{S}|\tilde{t})}{\hat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] \right\} d\tilde{t} \\ &:= \frac{1}{n^{\frac{3}{2}}} \sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} d\tilde{t} \end{aligned}$$

with $\mathbf{Z}_i(\tilde{t}) = \frac{\left(\frac{T_i-\tilde{t}}{h}\right) K\left(\frac{T_i-\tilde{t}}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\hat{p}_\zeta(\mathbf{S}_i|\tilde{t})}{\hat{p}(T_i, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}_i|\tilde{t})}{\bar{p}(T_i, \mathbf{S}_i)} \right] [Y_i - \bar{\mu}(\tilde{t}, \mathbf{S}_i) - (T_i - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S}_i)]$. Note that the random variables $\int_{T_i}^t \{ \mathbf{Z}_j(\tilde{t}) - \mathbb{E}[\mathbf{Z}_j(\tilde{t})] \} d\tilde{t}$ and $\int_{T_k}^t \{ \mathbf{Z}_\ell(\tilde{t}) - \mathbb{E}[\mathbf{Z}_\ell(\tilde{t})] \} d\tilde{t}$ are dependent and have a nonzero covariance when any of i, j, k, ℓ coincides. Thus, the variance of

$$\sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} d\tilde{t}$$

involves a sum of $O(n^3)$ nonzero terms; see, *e.g.*, Section 6.1 in [Lehmann \(1999\)](#) for detailed explanations. By Chebyshev's inequality,

$$\begin{aligned} \sqrt{nh^3} \cdot \mathbf{Term IIIb} &= \frac{1}{n^{\frac{3}{2}}} \sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} \\ &= O_P \left(\text{Var} \left[\int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} \right] \right) \\ &= O_P \left(\sqrt{\mathbb{E} \left[\left(\int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} \right)^2 \right]} \right) \\ &= o_P(1). \end{aligned}$$

Here, the last equality follows from the calculations that

$$\begin{aligned} &\mathbb{E} \left[\left(\int_{T_{i_1}}^t \{ \mathbf{Z}_{i_2}(\tilde{t}) - \mathbb{E}[\mathbf{Z}_{i_2}(\tilde{t})] \} \right)^2 \right] \\ &\stackrel{(i)}{\leq} \mathbb{E} \left[(t - T_{i_1})^2 |\mathbf{Z}_{i_2}(t') - \mathbb{E}[\mathbf{Z}_{i_2}(t')]|^2 \right] \\ &= \sup_{t' \in \mathcal{T}} \mathbb{E} \left[(t - T_{i_1})^2 |\mathbf{Z}_{i_2}(t')|^2 \right] \end{aligned}$$

$$\begin{aligned}
& \stackrel{\text{(ii)}}{\lesssim} \sup_{t' \in \mathcal{T}} \mathbb{E} \left\{ \frac{\left(\frac{T_i - t'}{h} \right)^2 K^2 \left(\frac{T_i - t'}{h} \right)}{h \cdot \kappa_2^2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}_i | t')}{\widehat{p}(T_i, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}_i | t')}{\bar{p}(T_i, \mathbf{S}_i)} \right]^2 [Y_i - \bar{\mu}(t', \mathbf{S}_i) - (T_i - t') \cdot \bar{\beta}(t', \mathbf{S}_i)]^2 \right\} \\
& \stackrel{\text{(iii)}}{\lesssim} \sup_{t \in \mathcal{T}} \|\widehat{p}_\zeta(\mathbf{S} | t) - \bar{p}_\zeta(\mathbf{S} | t)\|_{L_2}^2 + \sup_{u \in \mathcal{T} \oplus h} \|\widehat{p}(u, \mathbf{S}) - \bar{p}(u, \mathbf{S})\|_{L_2}^2 \\
& = O_P(\bar{\Upsilon}_{5,n}^2 + \bar{\Upsilon}_{6,n}^2) = o_P(1),
\end{aligned}$$

where (i) applies the mean-value theorem for integrals with t' lying between $t, T_{i_1} \in \mathcal{T}$, (ii) uses the compactness of $\mathcal{T} \subset \mathbb{R}$, and (iii) utilizes our derivations in [Section H.3.3](#).

• **Term IIIc:** Analogous to our arguments for **Term IIIb**, we write **Term IIIc** in terms of V-statistics and deduce that

$$\begin{aligned}
& \sqrt{nh^3} \cdot \mathbf{Term IIIc} \\
& = \frac{1}{\sqrt{n}} \sum_{i=1}^n \int_{T_i}^t (\mathbb{P}_n - \mathbb{P}) \left\{ \frac{\left(\frac{T_i - \tilde{t}}{h} \right) K \left(\frac{T_i - \tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S} | \tilde{t})}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T_i, \mathbf{S})} \left[\bar{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(\tilde{t}, \mathbf{S}) + (T_i - \tilde{t}) \left[\bar{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(\tilde{t}, \mathbf{S}) \right] \right] \right\} d\tilde{t} \\
& := \frac{1}{n^{\frac{3}{2}}} \sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \{ \bar{\mathbf{Z}}_{i_2}(\tilde{t}) - \mathbb{E}[\bar{\mathbf{Z}}_{i_2}(\tilde{t})] \} d\tilde{t} \\
& = O_P \left(\sqrt{\mathbb{E} \left[\left(\int_{T_{i_1}}^t \{ \bar{\mathbf{Z}}_{i_2}(\tilde{t}) - \mathbb{E}[\bar{\mathbf{Z}}_{i_2}(\tilde{t})] \} \right)^2 \right]} \right) \\
& = o_P(1).
\end{aligned}$$

Here, $\bar{\mathbf{Z}}_i(\tilde{t}) = \frac{\left(\frac{T_i - \tilde{t}}{h} \right) K \left(\frac{T_i - \tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}_i | \tilde{t})}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T_i, \mathbf{S}_i)} \left[\bar{\mu}(\tilde{t}, \mathbf{S}_i) - \widehat{\mu}(\tilde{t}, \mathbf{S}_i) + (T_i - \tilde{t}) \left[\bar{\beta}(\tilde{t}, \mathbf{S}_i) - \widehat{\beta}(\tilde{t}, \mathbf{S}_i) \right] \right]$, and the last equality above follows from some similar calculations as:

$$\begin{aligned}
& \mathbb{E} \left[\left(\int_{T_{i_1}}^t \{ \bar{\mathbf{Z}}_{i_2}(\tilde{t}) - \mathbb{E}[\bar{\mathbf{Z}}_{i_2}(\tilde{t})] \} \right)^2 \right] \\
& \leq \mathbb{E} \left[(t - T_{i_1})^2 \left| \bar{\mathbf{Z}}_{i_2}(t) - \mathbb{E}[\bar{\mathbf{Z}}_{i_2}(t)] \right|^2 \right] \\
& = \sup_{t' \in \mathcal{T}} \mathbb{E} \left[(t - T_{i_1})^2 \left| \bar{\mathbf{Z}}_{i_2}(t') \right|^2 \right] \\
& \lesssim \sup_{t' \in \mathcal{T}} \mathbb{E} \left\{ \frac{\left(\frac{T_i - \tilde{t}}{h} \right)^2 K^2 \left(\frac{T_i - \tilde{t}}{h} \right) \cdot \bar{p}_\zeta^2(\mathbf{S}_i | \tilde{t})}{h \cdot \kappa_2^2 \cdot \bar{p}^2(T_i, \mathbf{S}_i)} \left[\bar{\mu}(\tilde{t}, \mathbf{S}_i) - \widehat{\mu}(\tilde{t}, \mathbf{S}_i) + (T_i - \tilde{t}) \left[\bar{\beta}(\tilde{t}, \mathbf{S}_i) - \widehat{\beta}(\tilde{t}, \mathbf{S}_i) \right] \right]^2 \right\} \\
& \stackrel{\text{(iv)}}{\lesssim} \sup_{t \in \mathcal{T}} \left[\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2}^2 + h^2 \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2}^2 \right] \\
& = O_P(\bar{\Upsilon}_{1,n}^2 + h^2 \bar{\Upsilon}_{3,n}^2) = o_P(1),
\end{aligned}$$

where (iv) again leverages our derivations in [Section H.3.3](#).

• **Term IIIId:** Similar to our arguments for **Term IIIB**, we also write **Term IIIId** in terms of V-statistics and utilize Markov's inequality to deduce that

$$\begin{aligned}
& \sqrt{nh^3} \cdot \mathbf{Term\ IIIId} \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \int_{T_i}^t \mathbb{P}_n \left\{ \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] \left[\widehat{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T - \tilde{t}) \left[\widehat{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right] \right] \right\} d\tilde{t} \\
&:= \frac{1}{n^{\frac{3}{2}}} \sum_{i_1=1}^n \sum_{i_2=1}^n \int_{T_{i_1}}^t \mathbf{V}_{i_2}(\tilde{t}) d\tilde{t} \\
&= O_P \left(\sqrt{n} \cdot \mathbb{E} \left| \int_{T_{i_1}}^t \mathbf{V}_{i_2}(\tilde{t}) d\tilde{t} \right| \right) \\
&= o_P(1).
\end{aligned}$$

Here, $\mathbf{V}_i(\tilde{t}) = \frac{\left(\frac{T_i-\tilde{t}}{h}\right) K\left(\frac{T_i-\tilde{t}}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}_i|\tilde{t})}{\widehat{p}(T_i, \mathbf{S}_i)} - \frac{\bar{p}_\zeta(\mathbf{S}_i|\tilde{t})}{\bar{p}(T_i, \mathbf{S}_i)} \right] \left[\widehat{\mu}(\tilde{t}, \mathbf{S}_i) - \widehat{\mu}(t, \mathbf{S}_i) + (T_i - \tilde{t}) \left[\widehat{\beta}(\tilde{t}, \mathbf{S}_i) - \widehat{\beta}(t, \mathbf{S}_i) \right] \right]$, and the last equality above follows from the following calculation as:

$$\begin{aligned}
& \sqrt{n} \cdot \mathbb{E} \left| \int_{T_{i_1}}^t \mathbf{V}_{i_2}(\tilde{t}) d\tilde{t} \right| \\
&\stackrel{(i)}{\leq} \sqrt{n} \cdot \mathbb{E} [|t - T_{i_1}| \cdot |\mathbf{V}_{i_2}(t')|] \\
&\stackrel{(ii)}{\lesssim} \sqrt{n} \cdot \sup_{\tilde{t} \in \mathcal{T}} \mathbb{E} \left| \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right)}{\sqrt{h} \cdot \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] \left[\widehat{\mu}(\tilde{t}, \mathbf{S}) - \widehat{\mu}(t, \mathbf{S}) + (T - \tilde{t}) \left[\widehat{\beta}(\tilde{t}, \mathbf{S}) - \widehat{\beta}(t, \mathbf{S}) \right] \right] \right| \\
&\stackrel{(iii)}{\lesssim} \sqrt{nh} \cdot \sup_{t \in \mathcal{T}} \left\{ \left[\|\widehat{p}_\zeta(\mathbf{S}|t) - p_\zeta(\mathbf{S}|t)\|_{L_2} + \sup_{|u-t| \leq h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right] \right. \\
&\quad \left. \times \left[\|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} + h \|\widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S})\|_{L_2} \right] \right\} \\
&\stackrel{(iv)}{=} o_P(1),
\end{aligned}$$

where applies the mean-value theorem for integrals with t' lying between $t, T_{i_1} \in \mathcal{T}$, (ii) uses the compactness of $\mathcal{T} \subset \mathbb{R}$, (iii) utilizes our derivations in [Section H.3.4](#), and (iv) leverages our assumption (c) on the doubly robust rate of convergence in the corollary statement.

• **Term IIIe, Term IIIf, and Term IIIg:** On one hand, when $\bar{p}(t, \mathbf{s}) = p(t, \mathbf{s})$ for all $(t, \mathbf{s}) \in \mathcal{T} \times \mathcal{S}$, we know from our calculations in [Section H.3.5](#) that **Term IIIe** = **Term IIIf** = 0

and

Term IIIg

$$\begin{aligned}
& \stackrel{(i)}{\leq} \left(\frac{1}{n} \sum_{i=1}^n |T_i - t| \right) \cdot \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h} \right) K \left(\frac{T-\tilde{t}}{h} \right)}{h^2 \kappa_2} \left[\frac{\widehat{p}_\zeta(\mathbf{S}|\tilde{t})}{\widehat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] \right\} \right| \\
& \stackrel{(ii)}{\lesssim} \left(\frac{1}{n} \sum_{i=1}^n |T_i - t| \right) \cdot \frac{1}{h} \left[\sup_{t \in \mathcal{T}} \|\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} + \sup_{u \in \mathcal{T} \oplus h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right] \\
& \stackrel{(iii)}{=} O_P \left(\frac{1}{h} (\bar{\Upsilon}_{5,n} + \bar{\Upsilon}_{6,n}) \right) \\
& = o_P \left(\sqrt{\frac{1}{nh^3}} \right),
\end{aligned}$$

where (i) utilizes the mean-value theorem for integrals, (ii) follows from our derivations in [Section H.3.5](#), and (iii) applies Markov's inequality to $\frac{1}{n} \sum_{i=1}^n |T_i - t|$ as well as leverages our assumption (c) on the doubly robust rate of convergence in the corollary statement. Specifically, since $\|\widehat{\mu}(t, \mathbf{S}) - \bar{\mu}(t, \mathbf{S})\|_{L_2} + h \left\| \widehat{\beta}(t, \mathbf{S}) - \bar{\beta}(t, \mathbf{S}) \right\|_{L_2} = O_P(1)$ for any $t \in \mathcal{T}$ when $\bar{\mu} \neq \mu$ and $\bar{\beta} \neq \beta$, our assumption (c) ensures that $\sup_{t \in \mathcal{T}} \|\widehat{p}_\zeta(\mathbf{S}|t) - \bar{p}_\zeta(\mathbf{S}|t)\|_{L_2} + \sup_{u \in \mathcal{T} \oplus h} \|\widehat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} = o_P \left(\sqrt{\frac{1}{nh}} \right)$.

On the other hand, when $\bar{\mu} = \mu$ and $\bar{\beta} = \beta$, we again know from our calculations in [Section H.3.5](#) that

Term IIIe

$$\begin{aligned}
& \leq \left(\frac{1}{n} \sum_{i=1}^n |T_i - t| \right) \\
& \quad \times \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h} \right)^2 K \left(\frac{T-\tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\beta(\tilde{t}, \mathbf{S}) - \widehat{\beta}(\tilde{t}, \mathbf{S})] \right\} + \int [\widehat{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) d\mathbf{s} \right| \\
& = O_P \left(\sup_{t \in \mathcal{T}} \left\| \widehat{\beta}(t, \mathbf{S}) - \beta(t, \mathbf{S}) \right\|_{L_2} \right) \\
& = o_P \left(\sqrt{\frac{1}{nh^3}} \right),
\end{aligned}$$

$$\begin{aligned}
\text{Term IIIf} & \leq \left(\frac{1}{n} \sum_{i=1}^n |T_i - t| \right) \cdot \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h} \right) K \left(\frac{T-\tilde{t}}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{h^2 \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} [\mu(\tilde{t}, \mathbf{S}) - \widehat{\mu}(\tilde{t}, \mathbf{S})] \right\} \right| \\
& = O_P \left(\frac{1}{h} \cdot \sup_{t \in \mathcal{T}} \|\widehat{\mu}(t, \mathbf{S}) - \mu(t, \mathbf{S})\|_{L_2} \right)
\end{aligned}$$

$$= o_P \left(\sqrt{\frac{1}{nh^3}} \right),$$

and

Term IIIg

$$\begin{aligned} &\leq \left(\frac{1}{n} \sum_{i=1}^n |T_i - t| \right) \cdot \sup_{\tilde{t} \in \mathcal{T}} \left| \mathbb{P} \left\{ \frac{\left(\frac{T-\tilde{t}}{h} \right) K \left(\frac{T-\tilde{t}}{h} \right)}{h^2 \kappa_2} \left[\frac{\hat{p}_\zeta(\mathbf{S}|\tilde{t})}{\hat{p}(T, \mathbf{S})} - \frac{\bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\bar{p}(T, \mathbf{S})} \right] [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] \right\} \right| \\ &= O_P \left(h^2 \left[\sup_{t \in \mathcal{T}} \|\hat{p}_\zeta(\mathbf{S}|t) - p_\zeta(\mathbf{S}|t)\|_{L_2} + \sup_{u \in \mathcal{T} \oplus h} \|\hat{p}(u, \mathbf{S}) - p(u, \mathbf{S})\|_{L_2} \right] \right) \\ &= o_P \left(\sqrt{\frac{1}{nh^3}} \right). \end{aligned}$$

As a summary for this subsection, we conclude that $\sqrt{nh^3} \cdot \mathbf{Term III} = o_P(1)$.

I.3.3 Asymptotic Normality of $\hat{m}_{C,DR}(t)$

For the asymptotic normality of $\hat{m}_{C,DR}(t)$, it follows from the Lyapunov central limit theorem. Specifically, we already show in [Section I.3.1](#) and [Section I.3.2](#) that

$$\begin{aligned} &\sqrt{nh^3} [\hat{m}_{C,DR}(t) - m(t)] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \left\{ \phi_{C,h,\tilde{t}}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) + \sqrt{h^3} \int [\bar{\beta}(\tilde{t}, \mathbf{s}) - \beta(\tilde{t}, \mathbf{s})] \bar{p}_\zeta(\mathbf{s}|\tilde{t}) ds \right\} d\tilde{t} \right] \right\} + o_P(1) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbb{E}_{T_1} \left[\int_{T_1}^t \phi_{C,h,\tilde{t}}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right] + o_P(1) \end{aligned}$$

with

$$\phi_{C,h,t}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) = \frac{\left(\frac{T-t}{h} \right) K \left(\frac{T-t}{h} \right) \cdot \bar{p}_\zeta(\mathbf{S}|t)}{\sqrt{h} \cdot \kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(t, \mathbf{S}) - (T - t) \cdot \bar{\beta}(t, \mathbf{S})]$$

and $V_{C,m}(t) = \mathbb{E} \left[\left\{ \mathbb{E}_{T_1} \left[\int_{T_1}^t \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right] \right\}^2 \right] = O(1)$ by our calculation of **Term Ia** in [Section I.3.1](#). Then,

$$\sum_{i=1}^n \text{Var} \left\{ \frac{1}{\sqrt{n}} \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t \phi_{C,h,\tilde{t}}(Y_i, T_i, \mathbf{S}_i; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right] \right\} = O(1)$$

and

$$\sum_{i=1}^n \mathbb{E} \left| \frac{1}{\sqrt{n}} \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t \phi_{C,h,\tilde{t}}(Y, T, \mathbf{S}; \bar{\mu}, \bar{\beta}, \bar{p}, \bar{p}_\zeta) d\tilde{t} \right] \right|^{2+c_4}$$

$$\begin{aligned}
&\leq \mathbb{E} \left[\frac{1}{\sqrt{n^{c_4} h^{c_4+2}}} \cdot \mathbb{E}_{T_1} \left| \int_{T_1}^t \frac{\left(\frac{T-\tilde{t}}{h}\right) K\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta(\mathbf{S}|\tilde{t})}{\kappa_2 \cdot \bar{p}(T, \mathbf{S})} \cdot [Y - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})] d\tilde{t} \right|^{2+c_4} \right] \\
&= \mathbb{E} \left\{ \frac{1}{\sqrt{n^{c_4} h^{c_4+2}}} \mathbb{E}_{T_1} \left[|t - T_1|^{2+c_4} \right. \right. \\
&\quad \times \sup_{\tilde{t} \in \mathcal{T}} \frac{\left(\frac{T-\tilde{t}}{h}\right)^{2+c_4} K^{2+c_4}\left(\frac{T-\tilde{t}}{h}\right) \cdot \bar{p}_\zeta^{2+c_4}(\mathbf{S}|\tilde{t})}{\kappa_2^{2+c_4} \cdot \bar{p}^{2+c_4}(T, \mathbf{S})} \cdot [\mu(T, \mathbf{S}) + \epsilon - \bar{\mu}(\tilde{t}, \mathbf{S}) - (T - \tilde{t}) \cdot \bar{\beta}(\tilde{t}, \mathbf{S})]^{2+c_4} d\tilde{t} \left. \left. \right] \right\} \\
&= O \left(\sqrt{\frac{1}{n^{c_4} h^{2+c_4}}} \right) = o(1)
\end{aligned}$$

by the upper boundedness of $\mu, \bar{\mu}, p$ under Assumptions [A3](#) and [A4](#), the upper boundedness of the kernel function under Assumption [A5\(c\)](#), the lower boundedness of \bar{p} away from 0 around the support \mathcal{J} , the assumption that $\mathbb{E}|\epsilon|^{2+c_1} < \infty$ for some constant $c_1 \geq 1$, and the requirement that $nh^3 \rightarrow \infty$ as $n \rightarrow \infty$. Hence, the Lyapunov condition holds, and we have that

$$\sqrt{nh^3} \left\{ \widehat{m}_{C,DR}(t) - m(t) - h^2 \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t B_{C,\theta}(\tilde{t}) d\tilde{t} \right] \right\} \xrightarrow{d} \mathcal{N}(0, V_{C,m}(t))$$

after subtracting the dominating bias term $h^2 \cdot \mathbb{E}_{T_1} \left[\int_{T_1}^t B_{C,\theta}(\tilde{t}) d\tilde{t} \right]$ that we have computed in **Term Id** in [Section I.3.1](#). The proof is thus completed. \square