

SPECIES OF ROTA-BAXTER ALGEBRAS BY ROOTED TREES, TWISTED BIALGEBRAS AND FOCK FUNCTORS

LOÏC FOISSY, LI GUO, XIAO-SONG PENG, YUNZHOU XIE, AND YI ZHANG

ABSTRACT. As a fundamental and ubiquitous combinatorial notion, species has attracted sustained interest, generalizing from set-theoretical combinatorial to algebraic combinatorial and beyond. The Rota-Baxter algebra is one of the algebraic structures with broad applications from Renormalization of quantum field theory to integrable systems and multiple zeta values. Its interpretation in terms of monoidal categories has also recently appeared. This paper studies species of Rota-Baxter algebras, making use of the combinatorial construction of free Rota-Baxter algebras in terms of angularly decorated trees and forests. The notion of simple angularly decorated forests is introduced for this purpose and the resulting Rota-Baxter species is shown to be free. Furthermore, a twisted bialgebra structure, as the bialgebra for species, is established on this free Rota-Baxter species. Finally, through the Fock functor, another proof of the bialgebra structure on free Rota-Baxter algebras is obtained.

CONTENTS

1. Introduction	1
1.1. Species and monoidal categories	2
1.2. Species and Rota-Baxter algebras	2
2. Rota-Baxter algebras on species	3
2.1. Species and Rota-Baxter species	3
2.2. Species of simple angularly decorated forests	6
2.3. Free Rota-Baxter species on the species of simple angularly decorated forests	7
3. Twisted bialgebras on the species of simple angularly decorated forests	11
3.1. Rota-Baxter structures on tensor powers	11
3.2. The twisted bialgebra structure	13
4. Fock functors and colored Fock functors	17
References	20

1. INTRODUCTION

This paper introduces the species of Rota-Baxter algebras, by equipping the species of angularly decorated rooted trees with the operations for a Rota-Baxter algebra, which is further enriched to a twisted bialgebra structure. Through the Fock functors, we obtain another proof of the bialgebra structure on free Rota-Baxter algebras on rooted trees.

Date: January 14, 2025.

2010 Mathematics Subject Classification. 18M80, 17B38, 18D10, 05C05, 16S10, 16T10, 08B20.

Key words and phrases. Species, monoidal category, Rota-Baxter algebra, rooted tree, bialgebra, Fock functor.

1.1. Species and monoidal categories. The concept of species was introduced by A. Joyal in his seminal paper [30], formulated as functors from the category of finite sets with bijective maps to the category of sets, providing a way to categorize combinatorial objects based on their structures. The theory was further developed as a fundamental tool in combinatorics and several other areas of mathematics, providing deep insights into the nature of combinatorial objects and their relationships [7, 14, 21, 24, 34].

Vector species can be viewed as objects in a category, with morphisms given by natural transformations. This category is endowed with a symmetric monoidal structure, which gives rise to various interesting notions, such as monoids, comonoids, bimonoids, and Hopf monoids within the context of species. A general structured theory of these algebraic structures, internal to the functor category of species, has been developed by Aguiar and Mahajan [3, 4]. Furthermore, there are natural functors that transform vector species into graded vector spaces, and Hopf monoids in species into graded Hopf algebras. The constructions and properties of Hopf algebras, when lifted to the species level, are often more accessible and easier to generalize through the use of these functors. In the other direction, through Fock functors, these enriched species with additional structures can be regarded as lifts of these structures in their original (graded) form.

As further enrichment and extensions of the classical species theory, species with additional structures, such as the associative algebra and bialgebra, were studied recently [16, 22, 32, 33, 35, 41, 44]. In particular, [22] studied bialgebras in the category of linear species under the name of twisted bialgebras and gave a species version of the chromatic symmetric series and chromatic polynomials. Later, Tamaroff [41] gave a Koszul duality theory between twisted algebras and coalgebras on the twisted coalgebra that gives rise to their cohomology theory to give a new alternative description of it. At the same time, Carlier [16] showed that Schmitt's hereditary species induce monoidal decomposition spaces and exemplify Schmitt's bialgebra construction as a case of the general bialgebra construction on these spaces. Furthermore, Cohen-Macaulay Hopf monoids in the category of species was studied in [44].

Species theory has found applications in various areas, including graph theory, algebra, and combinatorial enumeration, providing deep insights into the nature of combinatorial objects and their relationships. Recently, Norledge [35] developed an algebraic framework for perturbative quantum field theory (pQFT) based on Joyal's combinatorial species. He demonstrated that fundamental pQFT structures can be viewed as algebraic structures within species, constructed using the Cauchy monoidal product.

1.2. Species and Rota-Baxter algebras. Rota-Baxter algebra was introduced by G. Baxter from fluctuation theory in probability [12]. Afterwards, it attracted attentions of prominent mathematicians such as Atkinson, Cartier and Rota [17, 37, 38]. In the early 1980s, Rota-Baxter operators on Lie algebras were independently discovered by Semenov-Tian-Shansky in [39] as the operator form of the classical Yang-Baxter equation [13], named after the physicists C. N. Yang and R. J. Baxter.

Over the past two decades, the study of Rota-Baxter algebras has undergone an extraordinary renaissance thanks to broad applications and connections such as Renormalization of quantum field theory [18], Yang-Baxter equations [8, 11, 15, 25, 26], operads [1, 9, 43], combinatorics [5, 45], Lie groups [29, 31], deformation theories [19, 42] and Hopf algebras [6, 10, 46]. See [27] for an introduction.

The category of species is a monoidal category [3]. Thus monoids and Hopf monoids can be defined for species. On the other hand, Rota-Baxter operators have been defined for monoidal categories in [2, 40].

In this paper, we combine species and Rota-Baxter algebras to define Rota-Baxter species by the Cauchy product, yielding a species with a Rota-Baxter algebra structure. In particular, we equipped angularly decorated forests with a free Rota-Baxter species, making use of the combinatorial constructions of free Rota-Baxter algebras given by rooted trees and angularly decorated forests [5, 20, 27, 46, 47]. A twisted bialgebra on the free Rota-Baxter species is also obtained, which recovers the biaglebra structure on free Rota-Baxter algebras. In a related work [23], the rings of various species (set species, linear species and localized set species) are equipped with the structures of an integro-differential ring [36] or a modified integro-differential ring [28].

Here is an outline of the paper. In Section 2, we recall the needed preliminaries on linear species. Then we use the Rota-Baxter operators on monoidal categories to define Rota-Baxter species and give the notion of free Rota-Baxter species. We construct a Rota-Baxter species on what we call simple angularly decorated forests and show that it is the free Rota-Baxter species.

In Section 3, we show that a species of bialgebras, namely a twisted bialgebra, can be given on the above-obtained free Rota-Baxter species. This is accomplished by making use of the universal property of the free Rota-Baxter species, which allows us to obtain the existence of the coproduct and its coassociativity.

In Section 4, we show that the (Bosonic) Fock functor from a species to its associated graded space can be enrich to a functor from a Rota-Baxter species to a Rota-Baxter algebra on the associated grades space. As a special case, the Fock functor applied to the free Rota-Baxter species recovers the construction of free Rota-Baxter algebras previously obtained by a direct construction [20, 27, 47].

Notations. In this paper, we fix a base field \mathbf{k} of characteristic zero. All linear spaces, linear maps and tensor products are taken on this base field.

2. ROTA-BAXTER ALGEBRAS ON SPECIES

This section introduces Rota-Baxter species and constructs free Rota-Baxter species by angularly decorated trees.

2.1. Species and Rota-Baxter species. First we recall basic notions and facts about species that will be used later. For more details, see [4, 14, 30].

Definition 2.1. Let \mathbf{Set} be the category whose objects are finite sets and whose morphisms are bijections, and let \mathbf{Vect} be the category whose objects are vector spaces and whose morphisms are linear maps.

(a) A **species** is a functor

$$\mathbf{p} : \mathbf{Set} \longrightarrow \mathbf{Vect}.$$

(b) For a species \mathbf{p} , its value on a finite set X is denoted by $\mathbf{p}[X]$. Its value on a bijection $\sigma : X \rightarrow Y$ is given by

$$\mathbf{p}[\sigma] : \mathbf{p}[X] \rightarrow \mathbf{p}[Y].$$

(c) For any finite set X , we have $\mathbf{p}[\text{id}_X] = \text{id}_{\mathbf{p}[X]}$. For any bijections $\sigma : X \rightarrow Y$ and $\tau : Y \rightarrow Z$, we have

$$\mathbf{p}[\tau \circ \sigma] = \mathbf{p}[\tau] \circ \mathbf{p}[\sigma].$$

(d) A **morphism between species** is a natural transformation of the functors for the two species. More precisely, for two species \mathbf{p} and \mathbf{q} , a **morphism $f : \mathbf{p} \rightarrow \mathbf{q}$ of species** is a collection of linear maps

$$f_X : \mathbf{p}[X] \rightarrow \mathbf{q}[X],$$

one for each finite set X such that, for each bijection $\sigma : X \rightarrow Y$ of finite sets, the following diagram commutes.

$$\begin{array}{ccc} \mathbf{p}[X] & \xrightarrow{f_X} & \mathbf{q}[X] \\ \mathbf{p}[\sigma] \downarrow & & \downarrow \mathbf{q}[\sigma] \\ \mathbf{p}[Y] & \xrightarrow{f_Y} & \mathbf{q}[Y] \end{array}$$

- (e) If the linear maps $f_X : \mathbf{p}[X] \rightarrow \mathbf{q}[X]$ in a morphism between species are bijections, then the morphism is called an **isomorphism between species**.
- (f) For any element $k \in \mathbf{k}$ and any morphisms of species $f, g : \mathbf{p} \rightarrow \mathbf{q}$, define the morphisms $kf, f + g : \mathbf{p} \rightarrow \mathbf{q}$ by

$$(kf)_X := kf_X \quad \text{and} \quad (f + g)_X := f_X + g_X,$$

for each finite set X . Then the category of species is a \mathbf{k} -linear category.

- (g) Given two species \mathbf{p} and \mathbf{q} , the **Cauchy product** $\mathbf{p} \otimes \mathbf{q}$ is defined by

$$\mathbf{p} \otimes \mathbf{q}[X] := \bigoplus_{X_1 \sqcup X_2 = X} \mathbf{p}[X_1] \otimes \mathbf{q}[X_2],$$

for any finite set X .

- (h) Denote by $\mathbf{1}_{\mathbf{k}}$ the species defined by

$$(1) \quad \mathbf{1}_{\mathbf{k}}[X] := \begin{cases} \mathbf{k}, & \text{if } X = \emptyset, \\ 0, & \text{otherwise.} \end{cases}$$

Then for any species \mathbf{p} , there is

$$\mathbf{p} \otimes \mathbf{1}_{\mathbf{k}} = \mathbf{p} = \mathbf{1}_{\mathbf{k}} \otimes \mathbf{p}.$$

Moreover, the category of species is a monoidal category under the Cauchy product, with the unit object $\mathbf{1}_{\mathbf{k}}$ [4].

Example 2.2. Here are two examples of species [22].

- (a) The species Com is defined by $\text{Com}[X] = \mathbf{k}$ for any finite set X and $\text{Com}[\sigma] = \text{id}_{\mathbf{k}}$ for any bijection $\sigma : X \rightarrow Y$.
- (b) For any finite set X , a set composition of X is a finite sequence (X_1, \dots, X_k) of nonempty subset of X such that $X_1 \sqcup \dots \sqcup X_k = X$. Denote by $\text{Comp}[X]$ the set of set compositions of X . Then define the species Comp as follows. For any finite set X , define $\text{Comp}[X]$ to be the space generated by $\text{Comp}[X]$; while for any set bijection $\sigma : X \rightarrow Y$ and $(X_1, \dots, X_k) \in \text{Comp}[X]$, define

$$\text{Comp}[\sigma](X_1, \dots, X_k) = (\sigma(X_1), \dots, \sigma(X_k)) \in \text{Comp}[Y].$$

As the category of species is a monoidal category, the notion of algebras is defined in the category of species as follows [22].

Definition 2.3. [22] An **algebra in the category of species**, or **twisted algebra**, is a triple (\mathbf{p}, m, ℓ) , where \mathbf{p} is a species, $m : \mathbf{p} \otimes \mathbf{p} \rightarrow \mathbf{p}$ and $\ell : \mathbf{1}_{\mathbf{k}} \rightarrow \mathbf{p}$ are morphisms of species such that

$$m \circ (m \otimes \text{id}_{\mathbf{p}}) = m \circ (\text{id}_{\mathbf{p}} \otimes m) \quad \text{and} \quad m \circ (\ell \otimes \text{id}_{\mathbf{p}}) = \text{id}_{\mathbf{p}} = m \circ (\text{id}_{\mathbf{p}} \otimes \ell).$$

Equivalently, a twisted algebra \mathbf{p} satisfies that for any two finite sets X and Y , there is a map $m_{X,Y} : \mathbf{p}[X] \otimes \mathbf{p}[Y] \rightarrow \mathbf{p}[X \sqcup Y]$ such that

(a) For any two bijections $\sigma : X \rightarrow X'$ and $\tau : Y \rightarrow Y'$ of finite sets,

$$m_{X',Y'} \circ (\mathbf{p}[\sigma] \otimes \mathbf{p}[\tau]) = \mathbf{p}[\sigma \sqcup \tau] \circ m_{X,Y} : \mathbf{p}[X] \otimes \mathbf{p}[Y] \rightarrow \mathbf{p}[X' \sqcup Y'],$$

where $\sigma \sqcup \tau : X \sqcup Y \rightarrow X' \sqcup Y'$ is the bijection induced by σ and τ .

(b) For finite sets X, Y, Z ,

$$m_{X \sqcup Y, Z} \circ (m_{X,Y} \otimes \text{id}_{\mathbf{p}[Z]}) = m_{X,Y \sqcup Z} \circ (\text{id}_{\mathbf{p}[X]} \otimes m_{Y,Z}).$$

(c) There is an element $1_{\mathbf{p}} \in \mathbf{p}[\emptyset]$ such that for any finite set X and any element $u \in \mathbf{p}[X]$,

$$m_{0,X}(1_{\mathbf{p}} \otimes u) = x = m_{X,0}(u \otimes 1_{\mathbf{p}}).$$

Remark 2.4. [22] Let \mathbf{p} be a twisted algebra and X be a finite set. For $(X_1, \dots, X_k) \in \text{Comp}[X]$, there is a map m_{X_1, \dots, X_k} from $\mathbf{p}[X_1] \otimes \dots \otimes \mathbf{p}[X_k]$ to $\mathbf{p}[X]$ inductively defined by

$$m_X = \text{id}_{\mathbf{p}[X]},$$

$$m_{X_1, \dots, X_k} = m_{X_1 \sqcup X_2, X_3, \dots, X_k} \circ (m_{X_1, X_2} \otimes \text{id}_{\mathbf{p}[X_3]} \otimes \dots \otimes \text{id}_{\mathbf{p}[X_k]}).$$

By associativity, if $(X_1, \dots, X_{k+\ell}) \in \text{Comp}[X]$, then

$$m_{X_1 \sqcup \dots \sqcup X_k, X_{k+1} \sqcup \dots \sqcup X_{k+\ell}} \circ (m_{X_1, \dots, X_k} \otimes m_{X_{k+1}, \dots, X_{k+\ell}}) = m_{X_1, \dots, X_{k+\ell}}.$$

Example 2.5. The species $\mathbf{1}_k$ is a twisted algebra, where for $k_1 \in \mathbf{1}_k[\emptyset] = \mathbf{k}$ and $k_2 \in \mathbf{1}_k[\emptyset] = \mathbf{k}$,

$$m_{0,0}(k_1 \otimes k_2) = k_1 k_2 \in \mathbf{1}_k[\emptyset] = \mathbf{k}.$$

The unit of $\mathbf{1}_k$ is $1_{\mathbf{k}} \in \mathbf{k} = \mathbf{1}_k[\emptyset]$.

Rota-Baxter operators have been defined for monoidal categories in [2, 40]. As the category of species is a monoidal category, we give the concept of Rota-Baxter species as follows.

Definition 2.6. Fix a scalar λ . A **Rota-Baxter algebra of weight λ in the category of species**, or **Rota-Baxter species of weight λ** , is a quadruple (\mathbf{p}, m, ℓ, R) , where (\mathbf{p}, m, ℓ) is a twisted algebra and $R : \mathbf{p} \rightarrow \mathbf{p}$ is a morphism of species such that the following diagram commutes:

$$\begin{array}{ccccc} \mathbf{p} \otimes \mathbf{p} & \xrightarrow{R \otimes \text{id}_{\mathbf{p}} + \text{id}_{\mathbf{p}} \otimes R + \lambda \text{id}_{\mathbf{p}} \otimes \text{id}_{\mathbf{p}}} & \mathbf{p} \otimes \mathbf{p} & \xrightarrow{m} & \mathbf{p} \\ R \otimes R \downarrow & & & & \downarrow R \\ \mathbf{p} \otimes \mathbf{p} & \xrightarrow{m} & & & \mathbf{p} \end{array}$$

In more detail, the commutative diagram means that, for any two finite sets X, Y and any elements $x \in \mathbf{p}[X], y \in \mathbf{p}[Y]$,

$$m_{X,Y} \circ (R_X \otimes R_Y)(x \otimes y) = R_{X \sqcup Y} \circ m_{X,Y} \circ (R_X \otimes \text{id}_{\mathbf{p}[Y]} + \text{id}_{\mathbf{p}[X]} \otimes R_Y + \lambda \text{id}_{\mathbf{p}[X]} \otimes \text{id}_{\mathbf{p}[Y]})(x \otimes y).$$

Definition 2.7. Let (\mathbf{p}, m, ℓ, R) and $(\mathbf{p}', m', \ell', R')$ be two Rota-Baxter species of the same weight, and let $f : \mathbf{p} \rightarrow \mathbf{p}'$ be a morphisms of species. We shall say that f is a **morphism of Rota-Baxter species** if

$$f \circ m = m' \circ (f \otimes f), \quad \ell' = f \circ \ell, \quad f \circ R = R' \circ f.$$

Inspired by the concept of free Rota-Baxter algebras [27], we also give the concept of free Rota-Baxter species as follows.

Definition 2.8. Let P be a species. Let (F, m, ℓ, R) be a Rota-Baxter species and $i : P \rightarrow F$ be a morphism of species. We say that F with the morphism $i : P \rightarrow F$ is a **free Rota-Baxter species** on P if for any Rota-Baxter species $(Q, m, 1_Q, R')$ and any morphism $g : P \rightarrow Q$ of species, there is a unique morphism of Rota-Baxter species $\bar{g} : F \rightarrow Q$ such that $g = \bar{g} \circ i$.


Remark 2.9. If it exists, the pair (F, i) is unique up to an isomorphism. Indeed, if (F, i) and (G, j) are two free Rota-Baxter species on P , using the species morphism $j : P \rightarrow G$, we obtain a morphism of Rota-Baxter species $\bar{j} : F \rightarrow G$ such that $\bar{j} \circ i = j$. Using the species morphism $i : P \rightarrow F$, we obtain a morphism of Rota-Baxter species $\bar{i} : G \rightarrow F$ such that $\bar{i} \circ j = i$. Then $\bar{i} \circ \bar{j}$ is an endomorphism of Rota-Baxter species of F such that $\bar{i} \circ \bar{j} \circ i = i$. By unicity of such an endomorphism, $\bar{i} \circ \bar{j} = \text{id}_F$. Similarly, $\bar{j} \circ \bar{i} = \text{id}_G$, so \bar{i} and \bar{j} are two isomorphisms, inverse one from the other.

2.2. Species of simple angularly decorated forests. In this subsection, we first recall the concept of angularly decorated forests [20, 27, 47]. Then as an example of species, we give the species of simple angularly decorated forests.


Definition 2.10. Let X be a set.

- (a) An **angularly X -decorated rooted tree** is a planar rooted tree together with a decoration of the angles between adjacent leaves by elements of X .
- (b) For a rooted forest $F = T_1 \cdots T_n$, as the concatenation of rooted trees T_1, \dots, T_n , an **angle of the forest F** is defined to be either the angle between two leaves of a rooted tree, or the space between two adjacent rooted trees T_i and T_{i+1} , $1 \leq i \leq n - 1$.
- (c) An **angularly X -decorated rooted forest** is a planar rooted forest F together with a decoration of the angles of F by elements of X . Let $\mathcal{F}(X)$ denote the set of angularly X -decorated rooted forests.

The one leaf tree \bullet is trivially an angularly X -decorated rooted tree with no angle to decorate. Here are some examples of angularly X -decorated trees



and here are some examples of angularly X -decorated forests



In general, an angularly X -decorated rooted forest is of the form

$$T_1 x_1 T_2 x_2 \cdots x_{n-1} T_n,$$

where $x_1, \dots, x_{n-1} \in X$ and T_1, \dots, T_n are angularly X -decorated rooted trees.

Remark 2.11. Note that the number of angles of a rooted forest F is precisely $\ell - 1$ for the number ℓ of leaves of F .

With this remark in mind, the following description of angularly X -decorated forests from [27] will be convenient to apply. For an angularly X -decorated forest F , let $u(F)$ be the underlying

forest of F and let $d(F) = (x_1, \dots, x_{\ell-1}) \in X^{\ell-1}$ denote the angular decorations of F . Then F can be written in the form

$$(2) \quad F = (u(F), d(F)).$$

Conversely, every pair (S, \vec{x}) consisting of a forest S with ℓ leaves and a vector $\vec{x} \in X^{\ell-1}$ uniquely determines an angular X -decorated rooted forest $F = F_{(S, \vec{x})}$, by decorating the angles of S with the entries of \vec{x} in the order from the left to the right.

Let X be a finite set. An angularly X -decorated rooted forest $F \in \mathcal{F}(X)$ is called **simple** if the angles of F are decorated by all the distinct elements of X . Since the number of angles of F is $\ell - 1$ if the number of leaves F is ℓ , this means that $|X| = \ell - 1$ and each element of X appears exactly once in the angles of F . Denote by $\mathcal{PF}(X)$ the set of all simple angularly X -decorated rooted forests. Thus for a fixed ordering \vec{x} of the elements of X , we have

$$\mathcal{PF} = \{F = (u(F), \tau(\vec{x})) \mid \tau \in S_{\ell-1}\}$$

for the permutation group $S_{\ell-1}$ on $\ell - 1$ letters.

Define a functor $\text{ADF} : \mathbf{Set} \rightarrow \mathbf{Vect}$ by taking

$$\text{ADF}[X] := \mathbf{k}\mathcal{PF}(X)$$

for any finite set X , and for any bijection $\sigma : X \rightarrow X'$ of finite sets, defining

$$\text{ADF}[\sigma] : \text{ADF}[X] \rightarrow \text{ADF}[Y], \quad F \mapsto \text{ADF}[\sigma](F),$$

where F is a simple angularly X -decorated rooted forest and $\text{ADF}[\sigma](F)$ is the simple angularly X' -decorated rooted forest obtained by replacing the decoration $x \in X$ for any given angle of F by the decoration $\sigma(x)$ for the same angle.

In terms of the description of angular decorated forests in Eq. (2), let $F = (u(F), d(F))$ with the underlying rooted forest $u(F)$ of F and the angular decoration vector $\vec{x} = (x_1, \dots, x_{\ell-1}) \in X^{\ell-1}$. Then we simply have

$$(3) \quad \text{ADF}[\sigma](F) = (u(F), \sigma(\vec{x})),$$

where $\sigma(\vec{x}) = (\sigma(x_1), \dots, \sigma(x_{\ell-1}))$.

By the definition of ADF , we have

$$\text{ADF}[\text{id}_X] = \text{id}_{\text{ADF}[X]}, \quad \text{ADF}[\sigma \circ \tau] = \text{ADF}[\sigma] \circ \text{ADF}[\tau],$$

for the identity set map $\text{id}_X : X \rightarrow X$, and bijective maps $\sigma : Y \rightarrow Z, \tau : X \rightarrow Y$. Hence $\text{ADF} : \mathbf{Set} \rightarrow \mathbf{Vect}$ is a species.

2.3. Free Rota-Baxter species on the species of simple angularly decorated forests. In this subsection, we construct a Rota-Baxter species on the species of simple angularly decorated forests and show that this Rota-Baxter species is a free Rota-Baxter species. First we recall the Rota-Baxter algebra structure on angularly decorated forests [20, 27].

For an angularly decorated rooted tree T , define the **depth** $\text{dep}(T)$ of T to be the maximal length of the paths from the root to the leaves of the tree. For an angularly decorated forest $F = T_1 x_1 T_2 \cdots x_n T_n$ with rooted trees T_1, \dots, T_n , define the **depth** $\text{dep}(F)$ of F to be the maximum of the depths of the trees T_1, \dots, T_n .

There is a **grafting operator** $B^+ : \mathcal{F}(X) \rightarrow \mathcal{F}(X)$ which sends an angularly X -decorated rooted forest $T_1 x_1 T_2 x_2 \cdots x_{n-1} T_n$ to a new angularly X -decorated rooted tree that is obtained by adding

a new root, and then adding an edge from the new root to the root of each of the trees T_1, \dots, T_n . For example,

$$B^+ \left(\bullet x \bullet \right) = \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ x \end{array} \quad \text{and} \quad B^+ \left(\bullet x_1 \bullet x_2 \bullet \right) = \begin{array}{c} \bullet \quad \bullet \quad \bullet \\ \diagdown \quad | \quad \diagup \\ x_1 \quad x_2 \end{array}.$$

In terms of the representation $F = (u(F), d(F))$ in Eq. (2), we have $B^+(F) = (B^+(u(F)), d(F))$.

Let $\mathbf{k}\mathcal{F}(X)$ be the free \mathbf{k} -module generated by $\mathcal{F}(X)$. To equip $\mathbf{k}\mathcal{F}(X)$ with a Rota-Baxter algebra structure, a multiplication \diamond is defined on $\mathbf{k}\mathcal{F}(X)$ as follows [20, 27]. For $F, F' \in \mathbf{k}\mathcal{F}(X)$, $F \diamond F'$ is defined by induction on $\text{dep} = \text{dep}(F) + \text{dep}(F')$. If $\text{dep} = 0$, then F, F' are of the form

$$F = \bullet x_1 \bullet \cdots \bullet x_m \bullet \quad \text{and} \quad F' = \bullet y_1 \bullet \cdots \bullet y_n \bullet,$$

with the convention that $F = \bullet$ if $m = 0$ and $F' = \bullet$ if $n = 0$. Then define

$$F \diamond F' := \bullet x_1 \bullet \cdots \bullet x_m \bullet y_1 \bullet \cdots \bullet y_n \bullet.$$

For the induction step of $\text{dep} > 0$, let

$$(4) \quad F = T_1 x_1 T_2 \cdots x_m T_{m+1} \quad \text{and} \quad F' = T'_1 y_1 T'_2 \cdots y_n T'_{n+1}.$$

First consider the case of $F = T_1, F' = T'_1$ being angularly decorated trees. Then define

$$(5) \quad F \diamond F' := \begin{cases} \bullet, & \text{if } T_1 = T'_1 = \bullet, \\ T_1, & \text{if } T'_1 = \bullet, \\ T'_1, & \text{if } T_1 = \bullet, \\ B^+(T_1 \diamond \overline{T}'_1) + B^+(\overline{T}_1 \diamond T'_1) + \lambda B^+(\overline{T}_1 \diamond \overline{T}'_1), & \text{if } T_1 = B^+(\overline{T}_1) \text{ and } T'_1 = B^+(\overline{T}'_1). \end{cases}$$

Here in the last case, each term on the right hand side is well defined by the induction hypothesis on the sum of depths. In the general case of F and F' as in Eq. (4), define

$$F \diamond F' := T_1 x_1 T_2 \cdots x_m (T_{m+1} \diamond T'_1) y_1 T'_2 \cdots y_n T'_{n+1},$$

where the product in the middle is given by Eq. (5) and the rest by concatenation. Equivalently, one can define a multiplication \diamond_u on the underlying forests by the same procedure, then for $F = (u(F), d(F))$ and $F' = (u(F'), d(F'))$, define

$$F \diamond F' = \left(u(F) \diamond_u u(F'), (d(F), d(F')) \right).$$

Here $(d(F), d(F'))$ denotes the vector concatenation of $d(F)$ and $d(F')$.

Finally expand the operations \diamond and B^+ to $\mathbf{k}\mathcal{F}(X)$ by (bi-)linearity. Then the following result is obtained.

Theorem 2.12. [20, 27] *Let X be a set. Let $i_x : X \rightarrow \mathbf{k}\mathcal{F}(X)$, $x \mapsto \bullet x \bullet$ be the set map. The quadruple $(\mathbf{k}\mathcal{F}(X), \diamond, B^+, i_x)$ is the free non-commutative unitary Rota-Baxter algebra generated by X .*

To construct the twisted algebra on the species ADF, we first prove the following lemma.

Lemma 2.13. *Let X, Y be two finite sets. Then for $F \in \text{ADF}[X]$ and $G \in \text{ADF}[Y]$, we have*

$$F \diamond G \in \text{ADF}[X \sqcup Y],$$

where the product \diamond is the one defined in Eq. (5).

Proof. Let

$$(6) \quad F = T_1 x_1 T_2 \cdots x_m T_{m+1} \in \text{ADF}[X], \quad G = T'_1 y_1 T'_2 \cdots y_n T'_{n+1} \in \text{ADF}[Y]$$

for some $m, n \geq 0$ as in Eq. (4). As in the definition of \diamond in Eq. (5), we prove the result by induction on $\text{dep}(F) + \text{dep}(G)$. If $\text{dep}(F) + \text{dep}(G) = 0$, then

$$F = \bullet x_1 \bullet \cdots x_m \bullet, \quad G = \bullet y_1 \bullet \cdots y_n \bullet,$$

and so the result holds trivially. For the induction step of $\text{dep}(F) + \text{dep}(G) > 0$, we prove the result by first considering $F = T_1, G = T'_1$ for simple angularly X -decorated tree T_1 and simple angularly Y -decorated tree T'_1 . If $T_1 = \bullet$ or $T'_1 = \bullet$, then the result holds directly. In the remaining case, we may write $T_1 = B^+(\bar{T}_1)$ and $T'_1 = B^+(\bar{T}'_1)$ for some $\bar{T}_1 \in \text{ADF}[X], \bar{T}'_1 \in \text{ADF}[Y]$, then

$$F \diamond G = B^+(\bar{T}_1) \diamond B^+(\bar{T}'_1) = B^+(T_1 \diamond \bar{T}'_1) + B^+(\bar{T}_1 \diamond T'_1) + \lambda B^+(\bar{T}_1 \diamond \bar{T}'_1).$$

By the induction hypothesis on the depth, $T_1 \diamond \bar{T}'_1, \bar{T}_1 \diamond T'_1$ and $\bar{T}_1 \diamond \bar{T}'_1$ are in $\text{ADF}[X \sqcup Y]$. Then $F \diamond G$ is in $\text{ADF}[X \sqcup Y]$. For general F and F' in Eq. (6), we have $T_{m+1} \in \text{ADF}[X']$ and $T'_1 \in \text{ADF}[Y']$, for some $X' \subseteq X$ and $Y' \subseteq Y$. By the definition of \diamond ,

$$F \diamond G = T_1 x_1 T_2 \cdots x_m (T_{m+1} \diamond T'_1) y_1 T'_2 \cdots y_n T'_{n+1}.$$

Since $T_{m+1} \diamond T'_1 \in \text{ADF}[X' \sqcup Y']$ by the previous case and other decorations are intact, it follows that $F \diamond G$ is in $\text{ADF}[X \sqcup Y]$. \square

For finite sets X, Y , and elements $F \in \text{ADF}[X], G \in \text{ADF}[Y]$, define

$$(7) \quad m_{X,Y}(F, G) := F \diamond G.$$

Then we have the following proposition.

Proposition 2.14. *The species ADF is a twisted algebra when it is equipped with the multiplication $m_{X,Y}$ defined by Eq. (7) and the unit $\bullet \in \text{ADF}[\emptyset]$.*

Proof. By the definition of ADF, we observe that for any two bijections $\sigma : X \rightarrow X'$ and $\tau : Y \rightarrow Y'$ of finite sets, there is

$$m_{X',Y'} \circ (\text{ADF}[\sigma] \otimes \text{ADF}[\tau]) = \text{ADF}[\sigma \sqcup \tau] \circ m_{X,Y}.$$

Also, for finite sets X, Y, Z and $F \in \text{ADF}[X], G \in \text{ADF}[Y], H \in \text{ADF}[Z]$, we have

$$\begin{aligned} & m_{X \sqcup Y, Z} \circ (m_{X,Y} \otimes \text{id}_{\text{ADF}[Z]})(F \otimes G \otimes H) \\ &= (F \diamond G) \diamond H \\ &= F \diamond (G \diamond H) \\ &= m_{X,Y \sqcup Z} \circ (\text{id}_{\text{ADF}[X]} \otimes m_{Y,Z})(F \otimes G \otimes H). \end{aligned}$$

Finally, for a finite set X and $F \in \text{ADF}[X]$,

$$m_{\emptyset, X}(\bullet \otimes F) = \bullet \diamond F = F = F \diamond \bullet = m_{X, \emptyset}(F \otimes \bullet).$$

Thus the required conditions are verified. \square

Now for any finite set X , define a map

$$(8) \quad R_X : \text{ADF}[X] \rightarrow \text{ADF}[X], \quad F \mapsto B^+(F),$$

where B^+ is the grafting operator recalled above. Then for any bijection $\sigma : X \rightarrow X'$ and $F \in \text{ADF}[X]$, there is

$$\text{ADF}[\sigma] \circ R_X(F) = \text{ADF}[\sigma](B^+(F)) = B^+(\text{ADF}[\sigma](F)) = R_{X'} \circ \text{ADF}[\sigma](F).$$

Hence $R : \text{ADF} \rightarrow \text{ADF}$ is a morphism of species. Moreover, we have

Proposition 2.15. *The twisted algebra (ADF, m, \bullet) , with the morphism R of species, is a Rota-Baxter species.*

Proof. For any finite sets X, Y and any elements $F \in \text{ADF}[X], G \in \text{ADF}[Y]$, we have

$$\begin{aligned} & m_{X,Y} \circ (R_X \circ R_Y)(F \otimes G) \\ &= m_{X,Y}(B^+(F) \otimes B^+(G)) \\ &= B^+(F) \diamond B^+(G) \\ &= B^+(B^+(F) \diamond G + F \diamond B^+(G) + \lambda F \diamond G) \text{ (by the definition of } \diamond) \\ &= R_{X \sqcup Y} \circ m_{X,Y} \circ (R_X \otimes \text{id}_{\text{ADF}[Y]} + \text{id}_{\text{ADF}[X]} \otimes R_Y + \lambda \text{id}_{\text{ADF}[X]} \otimes \text{id}_{\text{ADF}[Y]})(F \otimes G), \end{aligned}$$

as required. \square

We define a species O by

$$(9) \quad O[X] := \begin{cases} \mathbf{k}\{\bullet x \bullet\}, & \text{if } X = \{x\} \text{ is a singleton set,} \\ 0, & \text{otherwise.} \end{cases}$$

If $\sigma : X = \{x\} \rightarrow Y = \{y\}$ is a bijection, then

$$O[\sigma] : \begin{cases} O[X] \longrightarrow O[Y] \\ \bullet x \bullet \longmapsto \bullet y \bullet. \end{cases}$$

There is a morphism $i_O : O \rightarrow \text{ADF}$ of species, where $(i_O)_X : O[X] \rightarrow \text{ADF}[X]$ is the natural inclusion if X is a singleton set and otherwise $i_O = 0$.

Now we show that the Rota-Baxter species ADF with the morphism $i := i_O : O \rightarrow \text{ADF}$ is a free Rota-Baxter species on O .

Theorem 2.16. *The Rota-Baxter species ADF together with the morphism $i : O \rightarrow \text{ADF}$ of species is the free Rota-Baxter species on O .*

Proof. For the proof, we prove that for any Rota-Baxter species $(Q, m, 1_Q, R')$ and any morphism $g : O \rightarrow Q$ of species, there is a unique morphism of Rota-Baxter species $\bar{g} : \text{ADF} \rightarrow Q$ such that $g = \bar{g} \circ i$.

For a finite set X , define $\bar{g}_X : \text{ADF}[X] \rightarrow Q[X]$ by induction on $|X|$. If $|X| = 0$, then $X = \emptyset$. For a rooted forest $F \in \text{ADF}[\emptyset]$, we define $\bar{g}_\emptyset(F)$ by induction on $\text{dep}(F)$. If $\text{dep}(F) = 0$, then $F = \bullet$ and define

$$\bar{g}_\emptyset(\bullet) := 1_Q.$$

For the induction step of $\text{dep}(F) > 0$, note that $F \in \text{ADF}[\emptyset]$ and so $F = B^+(\bar{F})$ for some $\bar{F} \in \text{ADF}[\emptyset]$. Then define

$$\bar{g}_\emptyset(F) := R'_\emptyset(\bar{g}(\bar{F})).$$

Hence $\bar{g}_\emptyset(F)$ are defined for all $F \in \text{ADF}[\emptyset]$. For the induction step of $|X| > 0$, take a rooted forest $F \in \text{ADF}[X]$ and define $\bar{g}_X(F)$ by a second induction on $\text{dep}(F)$. If $\text{dep}(F) = 0$, then F is of the form $\bullet x_1 \bullet \cdots \bullet x_m \bullet$, where $X = \{x_1, \dots, x_m\}$. If $m = 1$ and $F = \bullet x_1 \bullet$, then define

$$\bar{g}_X(F) := g_{\{x_1\}}(\bullet x_1 \bullet).$$

If $m > 1$, then define

$$\bar{g}_X(F) := m_{\{x_1\}, \{x_2\}, \dots, \{x_m\}}(g_{\{x_1\}}(\bullet x_1 \bullet) \otimes \cdots \otimes g_{\{x_m\}}(\bullet x_m \bullet)),$$

where $m_{\{x_1, \{x_2\}, \dots, \{x_m\}}$ is the map from $Q[\{x_1\}] \otimes \dots \otimes Q[\{x_m\}]$ into $Q[X]$ inductively defined as in Remark 2.4.

For the induction step of $\text{dep}(F) > 0$, let $F = T_1 x_1 T_2 \cdots T_m x_m T_{m+1} \in \text{ADF}[X]$, we define $\bar{g}_X(F)$ by induction on m . If $m = 0$, then $F = B^+(\bar{F})$ for some $\bar{F} \in \text{ADF}[X]$. Then define

$$\bar{g}_X(F) = R'_X(\bar{g}_X(\bar{F})),$$

where $\bar{g}_X(\bar{F})$ is defined by induction hypothesis.

For the induction step of $m > 0$, assume $T_1 \in \text{ADF}[X_1], \dots, T_m \in \text{ADF}[X_m], T_{m+1} \in \text{ADF}[X_{m+1}]$ and $X_1 \sqcup \dots \sqcup X_{m+1} \sqcup \{x_1, \dots, x_{m+1}\} = X$. Then define

$$\bar{g}_X(F) = m_{X_1, \{x_1\}, \dots, X_m, \{x_m\}, X_{m+1}}(\bar{g}_{X_1}(T_1) \otimes \bar{g}_{\{x_1\}}(\bullet x_1 \bullet) \otimes \dots \otimes \bar{g}_{\{x_m\}}(\bullet x_m \bullet) \otimes \bar{g}_{X_{m+1}}(T_{m+1})),$$

where $m_{X_1, \{x_1\}, \dots, X_m, \{x_m\}, X_{m+1}}$ is the map from $Q[X_1] \otimes Q[\{x_1\}] \otimes \dots \otimes Q[\{x_m\}] \otimes Q[X_{m+1}]$ into $Q[X]$ inductively defined as in Remark 2.4.

By the construction of \bar{g} , it is the unique morphism of twisted Rota-Baxter algebras such that $g = \bar{g} \circ i$. \square

3. TWISTED BIALGEBRAS ON THE SPECIES OF SIMPLE ANGULARLY DECORATED FORESTS

In this section, applying the universal property of the free Rota-Baxter species, a bialgebra structure is given to the free Rota-Baxter species on simple angularly decorated forests.

3.1. Rota-Baxter structures on tensor powers. We first construct an augmentation on ADF which will be the counit in the bialgebra construction.

The species $1_{\mathbf{k}}$ is a Rota-Baxter species, with the product of \mathbf{k} and the null Rota-Baxter operator. Thus the universal property of ADF gives a unique morphism of Rota-Baxter species

$$\epsilon : \text{ADF} \longrightarrow 1_{\mathbf{k}}$$

such that for any singleton set $\{x\}$, $\epsilon_{\{x\}}(\bullet x \bullet) = 0$.

Lemma 3.1. *The species morphism ϵ thus defined is an augmentation: for any finite set X and $F \in \text{ADF}[X]$, we have*

$$(10) \quad \epsilon_X(F) := \begin{cases} 1_{\mathbf{k}}, & \text{if } X = \emptyset \text{ and } F = \bullet, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. As ϵ is a morphism of Rota-Baxter species, $\epsilon_{\emptyset}(\bullet) = 1_{\mathbf{k}}$. Its kernel is an ideal of the twisted algebra ADF, containing the elements $\bullet x \bullet$, and the image of R as $\epsilon \circ R = 0$. An easy induction on the depth shows that this kernel contains all the simple angularly decorated forests except \bullet . \square

For later applications, we give some properties of the morphism ϵ as follows.

Lemma 3.2. *For any finite sets X, Y and any $F \in \text{ADF}[X], G \in \text{ADF}[Y]$, we have*

- (a) $\epsilon_{X \sqcup Y}(R_X(F) \diamond G) = \epsilon_{X \sqcup Y}(F \diamond R_Y(G)) = 0$;
- (b) $R_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = \epsilon_X(F) \bullet \diamond R_Y(G)$, $R_{X \sqcup Y}(F \diamond \epsilon_Y(G) \bullet) = R_X(F) \diamond \epsilon_Y(G) \bullet$;
- (c) $\epsilon_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = \epsilon_{X \sqcup Y}(F \diamond \epsilon_Y(G) \bullet) = \epsilon_X(F) \epsilon_Y(G)$.

Proof. (a) By the definition of \diamond , we see that the expansions of $R_X(F) \diamond G$ and $F \diamond R_Y(G)$ do not include the term \bullet . Hence $\epsilon_{X \sqcup Y}(R_X(F) \diamond G) = \epsilon_{X \sqcup Y}(F \diamond R_Y(G)) = 0$.

(b) If $X = \emptyset$ and $F = \bullet$, then

$$R_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = R_Y(\bullet \diamond G) = R_Y(G) = \bullet \diamond R_Y(G) = \epsilon_X(F) \bullet \diamond R_Y(G).$$

If not, then

$$R_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = R_{X \sqcup Y}(0 \diamond G) = 0 = \epsilon_X(F) \bullet \diamond R_Y(G).$$

Similarly, we have

$$R_{X \sqcup Y}(F \diamond \epsilon_Y(G) \bullet) = R_X(F) \diamond \epsilon_Y(G) \bullet.$$

(c) If $X = \emptyset$ and $F = \bullet$, then

$$\epsilon_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = \epsilon_Y(\bullet \diamond G) = \epsilon_Y(G) = \epsilon_X(F) \epsilon_Y(G).$$

If not, then

$$\epsilon_{X \sqcup Y}(\epsilon_X(F) \bullet \diamond G) = \epsilon_Y(0 \diamond G) = 0 = \epsilon_X(F) \epsilon_Y(G).$$

Similarly, we have

$$\epsilon_{X \sqcup Y}(F \diamond \epsilon_Y(G) \bullet) = \epsilon_X(F) \epsilon_Y(G). \quad \square$$

Note that [22] if $(\mathbf{p}, m_{\mathbf{p}})$ and $(\mathbf{q}, m_{\mathbf{q}})$ are two twisted algebras, then $\mathbf{p} \otimes \mathbf{q}$ is also a twisted algebra with the product

$$m_{\mathbf{p} \otimes \mathbf{q}} := (m_{\mathbf{p}} \otimes m_{\mathbf{q}}) \circ (\text{id}_{\mathbf{p}} \otimes c_{\mathbf{p}, \mathbf{q}} \otimes \text{id}_{\mathbf{q}}),$$

where $c_{\mathbf{p}, \mathbf{q}} : \mathbf{p} \otimes \mathbf{q} \rightarrow \mathbf{q} \otimes \mathbf{p}$ is the isomorphism of species

$$(c_{\mathbf{p}, \mathbf{q}})_X : \begin{cases} \mathbf{p} \otimes \mathbf{q}[X] & \rightarrow \mathbf{q} \otimes \mathbf{p}[X], \\ x \otimes y & \mapsto y \otimes x. \end{cases}$$

Since ADF is a twisted algebra, $\text{ADF} \otimes \text{ADF}$ is also a twisted algebra. We further have:

Proposition 3.3. *The twisted algebra $\text{ADF} \otimes \text{ADF}$ is a Rota-Baxter species, where the Rota-Baxter operator $R^{(2)} : \text{ADF} \otimes \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$ is defined by*

$$R_X^{(2)} : \begin{cases} \text{ADF} \otimes \text{ADF}[X] & \rightarrow \text{ADF} \otimes \text{ADF}[X], \\ \text{ADF}[X_1] \otimes \text{ADF}[X \setminus X_1] \ni F \otimes G & \mapsto R_{X_1}(F) \otimes G + \epsilon_{X_1}(F) \bullet \otimes R_{X \setminus X_1}(G). \end{cases}$$

Proof. If $X_1 = \emptyset$ and for $F \otimes G \in \text{ADF}[X_1] \otimes \text{ADF}[X \setminus X_1]$ with $F = \bullet$,

$$R_X^{(2)}(F \otimes G) = R_{\emptyset}(F) \otimes G + \epsilon_{\emptyset}(F) \bullet \otimes R_X(G) = B^+(F) \otimes G + \bullet \otimes B^+(G) \in \text{ADF}[\emptyset] \otimes \text{ADF}[X].$$

Otherwise, $\epsilon_{X_1}(F) = 0$ and so

$$R_X^{(2)}(F \otimes G) = R_{X_1}(F) \otimes G + \epsilon_{X_1}(F) \bullet \otimes R_{X \setminus X_1}(G) = B^+(F) \otimes G \in \text{ADF}[X_1] \otimes \text{ADF}[X \setminus X_1].$$

For any bijection $\sigma : X \rightarrow X'$ and $F \otimes G \in \text{ADF} \otimes \text{ADF}[X]$, by the definition of $R^{(2)}$, we find that

$$(\text{ADF} \otimes \text{ADF}[\sigma]) \circ R_X^{(2)}(F \otimes G) = R_{X'}^{(2)} \circ (\text{ADF} \otimes \text{ADF}[\sigma])(F \otimes G).$$

Hence $R^{(2)} : \text{ADF} \otimes \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$ is a morphism of species. Now for $F_1 \otimes G_1 \in \text{ADF}[X_1] \otimes \text{ADF}[X \setminus X_1] \subset \text{ADF} \otimes \text{ADF}[X]$ and $F_2 \otimes G_2 \in \text{ADF}[Y_1] \otimes \text{ADF}[Y \setminus Y_1] \subset \text{ADF} \otimes \text{ADF}[Y]$, there is

$$\begin{aligned} & R_{X \sqcup Y}^{(2)} \circ (m_{\text{ADF} \otimes \text{ADF}})_{X, Y} \circ (R_X^{(2)} \otimes \text{id} + \text{id} \otimes R_Y^{(2)} + \lambda \text{id} \otimes \text{id})((F_1 \otimes G_1) \otimes (F_2 \otimes G_2)) \\ &= R_{X \sqcup Y}^{(2)} \circ (m_{\text{ADF} \otimes \text{ADF}})_{X, Y} (R_X^{(2)}(F_1 \otimes G_1) \otimes (F_2 \otimes G_2) + (F_1 \otimes G_1) \otimes R_Y^{(2)}(F_2 \otimes G_2) \\ &\quad + \lambda(F_1 \otimes G_1) \otimes (F_2 \otimes G_2)) \\ &= R_{X \sqcup Y}^{(2)} \circ (m_{\text{ADF} \otimes \text{ADF}})_{X, Y} ((R_{X_1}(F_1) \otimes G_1 + \epsilon_{X_1}(F_1) \bullet \otimes R_{X \setminus X_1}(G_1)) \otimes (F_2 \otimes G_2) \\ &\quad + (F_1 \otimes G_1) \otimes (R_{Y_1}(F_2) \otimes G_2 + \epsilon_{Y_1}(F_2) \bullet \otimes R_{Y \setminus Y_1}(G_2)) + \lambda(F_1 \otimes G_1) \otimes (F_2 \otimes G_2)) \\ &= R_{X \sqcup Y}^{(2)} ((R_{X_1}(F_1) \diamond F_2) \otimes (G_1 \diamond G_2) + (\epsilon_{X_1}(F_1) \bullet \diamond F_2) \otimes (R_{X \setminus X_1}(G_1) \diamond G_2)) \end{aligned}$$

$$\begin{aligned}
& + (F_1 \diamond R_{Y_1}(F_2)) \otimes (G_1 \diamond G_2) + ((F_1 \diamond \epsilon_{Y_1}(F_2)) \bullet) \otimes (G_1 \diamond R_{Y \setminus Y_1}(G_2)) + \lambda(F_1 \diamond F_2) \otimes (G_1 \diamond G_2) \\
= & R_{X_1 \sqcup Y_1}(R_{X_1}(F_1) \diamond F_2) \otimes (G_1 \diamond G_2) + \epsilon_{X_1 \sqcup Y_1}(R_{X_1}(F_1) \diamond F_2) \bullet \otimes R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond G_2) \\
& + R_{X_1 \sqcup Y_1}(\epsilon_{X_1}(F_1) \bullet \diamond F_2) \otimes (R_{X \setminus X_1}(G_1) \diamond G_2) \\
& + \epsilon_{X_1 \sqcup Y_1}((\epsilon_{X_1}(F_1) \bullet) \diamond F_2) \bullet \otimes R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(R_{X \setminus X_1}(G_1) \diamond G_2) \\
& + R_{X_1 \sqcup Y_1}(F_1 \diamond R_{Y_1}(F_2)) \otimes (G_1 \diamond G_2) + \epsilon_{X_1 \sqcup Y_1}(F_1 \diamond R_{Y_1}(F_2)) \bullet \otimes R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond G_2) \\
& + R_{X_1 \sqcup Y_1}(F_1 \diamond (\epsilon_{Y_1}(F_2) \bullet)) \otimes (G_1 \diamond R_{Y \setminus Y_1}(G_2)) \\
& + \epsilon_{X_1 \sqcup Y_1}(F_1 \diamond (\epsilon_{Y_1}(F_2) \bullet)) \bullet \otimes R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond R_{Y \setminus Y_1}(G_2)) \\
& + \lambda R_{X_1 \sqcup Y_1}(F_1 \diamond F_2) \otimes (G_1 \diamond G_2) + \lambda \epsilon_{X_1 \sqcup Y_1}(F_1 \diamond F_2) \bullet \otimes R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond G_2) \\
= & (R_{X_1 \sqcup Y_1}(R_{X_1}(F_1) \diamond F_2) + R_{X_1 \sqcup Y_1}(F_1 \diamond R_{Y_1}(F_2)) + \lambda R_{X_1 \sqcup Y_1}(F_1 \diamond F_2)) \otimes (G_1 \diamond G_2) \\
& + (\epsilon_{X_1}(F_1) \bullet \diamond R_{Y_1}(F_2)) \otimes (R_{X \setminus X_1}(G_1) \diamond G_2) + (R_{X_1}(F_1) \diamond \epsilon_{Y_1}(F_2) \bullet) \otimes (G_1 \diamond R_{Y \setminus Y_1}(G_2)) \\
& + (\epsilon_{X_1}(F_1) \epsilon_{Y_1}(F_2)) \bullet \otimes (R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(R_{X \setminus X_1}(G_1) \diamond G_2) + R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond R_{Y \setminus Y_1}(G_2)) \\
& + \lambda R_{(X \sqcup Y) \setminus (X_1 \sqcup Y_1)}(G_1 \diamond G_2)) \quad (\text{by Lemmas 3.1 and 3.2}) \\
= & (R_{X_1}(F_1) \diamond R_{Y_1}(F_2)) \otimes (G_1 \diamond G_2) + (\epsilon_{X_1}(F_1) \bullet \diamond R_{Y_1}(F_2)) \otimes (R_{X \setminus X_1}(G_1) \diamond G_2) \\
& + (R_{X_1}(F_1) \diamond \epsilon_{Y_1}(F_2) \bullet) \otimes (G_1 \diamond R_{Y \setminus Y_1}(G_2)) + (\epsilon_{X_1}(F_1) \epsilon_{Y_1}(F_2)) \bullet \otimes (R_{X \setminus X_1}(G_1) \diamond R_{Y \setminus Y_1}(G_2)) \\
= & (m_{\text{ADF} \otimes \text{ADF}})_{X,Y} \left((R_{X_1}(F_1) \otimes G_1 + \epsilon_{X_1}(F_1) \bullet \otimes R_{X \setminus X_1}(G_1)) \otimes (R_{Y_1}(F_2) \otimes G_2 + \epsilon_{Y_1}(F_2) \bullet \otimes R_{Y \setminus Y_1}(G_2)) \right) \\
= & (m_{\text{ADF} \otimes \text{ADF}})_{X,Y} \circ (R_X^{(2)} \otimes R_Y^{(2)})((F_1 \otimes G_1) \otimes (F_2 \otimes G_2)),
\end{aligned}$$

and hence $R^{(2)}$ satisfies the Rota-Baxter identity. \square

Based on ADF being a Rota-Baxter species with R , and $\text{ADF} \otimes \text{ADF}$ being a Rota-Baxter species with $R^{(2)}$, we define $R^{(3)} : \text{ADF} \otimes \text{ADF} \otimes \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$ by

$$\begin{aligned}
R_{X_1, X_2, X \setminus (X_1 \sqcup X_2)}^{(3)}(F \otimes G \otimes H) & := R_{X_1}(F) \otimes G \otimes H + \epsilon_{X_1}(F) \bullet \otimes R_{X \setminus X_1}^{(2)}(G \otimes H) \\
& = R_{X_1}(F) \otimes G \otimes H + \epsilon_{X_1}(F) \bullet \otimes R_{X_2}(G) \otimes H + \epsilon_{X_1}(F) \bullet \otimes \epsilon_{X_2}(G) \bullet \otimes R_{X \setminus (X_1 \sqcup X_2)}(H),
\end{aligned}$$

for $F \otimes G \otimes H \in \text{ADF}[X_1] \otimes \text{ADF}[X_2] \otimes \text{ADF}[X \setminus (X_1 \sqcup X_2)]$. Then by a similar argument for the proof of Proposition 3.3, we obtain

Proposition 3.4. *The species $\text{ADF} \otimes \text{ADF} \otimes \text{ADF}$ equipped with the operator $R^{(3)}$ is a Rota-Baxter species.*

3.2. The twisted bialgebra structure. We now apply the universal property of the free Rota-Baxter species ADF and the Rota-Baxter structure on the tensor powers of this algebra to obtain a bialgebra structure on ADF.

Definition 3.5. [22] A **coalgebra in the category of species**, or **twisted coalgebra**, is a triple $(\mathbf{p}, \Delta, \varepsilon)$, where \mathbf{p} is a species, $\Delta : \mathbf{p} \rightarrow \mathbf{p} \otimes \mathbf{p}$ and $\varepsilon : \mathbf{p} \rightarrow \mathbf{1}_k$ are morphisms of species such that

$$(\Delta \otimes \text{id}_{\mathbf{p}}) \circ \Delta = (\text{id}_{\mathbf{p}} \otimes \Delta) \circ \Delta \quad \text{and} \quad (\varepsilon \otimes \text{id}_{\mathbf{p}}) \circ \Delta = \text{id}_{\mathbf{p}} = (\text{id}_{\mathbf{p}} \otimes \varepsilon) \circ \Delta.$$

Equivalently, a twisted coalgebra \mathbf{p} satisfies that for any two finite sets X and Y , there is a map $\Delta_{X,Y} : \mathbf{p}[X \sqcup Y] \rightarrow \mathbf{p}[X] \otimes \mathbf{p}[Y]$ such that

(a) For any two bijections $\sigma : X \rightarrow X'$ and $\tau : Y \rightarrow Y'$ of finite sets,

$$(\mathbf{p}[\sigma] \otimes \mathbf{p}[\tau]) \circ \Delta_{X,Y} = \Delta_{X',Y'} \circ \mathbf{p}[\sigma \sqcup \tau] : \mathbf{p}[X \sqcup Y] \rightarrow \mathbf{p}[X'] \otimes \mathbf{p}[Y'].$$

(b) For finite sets X, Y, Z ,

$$(\Delta_{X,Y} \otimes \text{id}_{\mathbf{p}[Z]}) \circ \Delta_{X \sqcup Y, Z} = (\text{id}_{\mathbf{p}[X]} \otimes \Delta_{Y,Z}) \circ \Delta_{X, Y \sqcup Z}.$$

(c) There is a linear map $\varepsilon_\emptyset : \mathbf{p}[\emptyset] \rightarrow \mathbf{k}$ such that for any finite set X ,

$$(\varepsilon_\emptyset \otimes \text{id}_{\mathbf{p}[X]}) \circ \Delta_{\emptyset, X} = \text{id}_{\mathbf{p}[X]} = (\text{id}_{\mathbf{p}[X]} \otimes \varepsilon_\emptyset) \circ \Delta_{X, \emptyset}.$$

Remark 3.6. (a) The species $\mathbf{1}_k$ is a twisted coalgebra, where for $k \in \mathbf{1}_k[\emptyset]$,

$$\Delta_{\emptyset, \emptyset}(k) = k(1_k \otimes 1_k) \in \mathbf{1}_k[\emptyset] \otimes \mathbf{1}_k[\emptyset].$$

(b) [22] If \mathbf{p} and \mathbf{q} are twisted coalgebras, then $\mathbf{p} \otimes \mathbf{q}$ is also a twisted coalgebra.

Definition 3.7. A **bialgebra in the category of species**, or **twisted bialgebra**, is a quintuple $(\mathbf{p}, m, \ell, \Delta, \varepsilon)$, where (\mathbf{p}, m, ℓ) is a twisted algebra, $(\mathbf{p}, \Delta, \varepsilon)$ is a twisted coalgebra, and one of the following equivalent conditions holds.

(a) $\varepsilon : \mathbf{p} \rightarrow \mathbf{1}_k$ and $\Delta : \mathbf{p} \rightarrow \mathbf{p} \otimes \mathbf{p}$ are algebra morphisms.

(b) $\ell : \mathbf{1}_k \rightarrow \mathbf{p}$ and $m : \mathbf{p} \otimes \mathbf{p} \rightarrow \mathbf{p}$ are coalgebra morphisms.

To construct the twisted coalgebra on ADF, we define a morphism of species

$$g : \mathcal{O} \rightarrow \text{ADF} \otimes \text{ADF}$$

as follows. If $X = \{x\}$ is a singleton set, then $g_X(\bullet x \bullet) = \bullet x \bullet \otimes \bullet + \bullet \otimes \bullet x \bullet$. Otherwise, $g_X = 0$. Then by Theorem 2.16 and Proposition 3.3, there is a unique morphism

$$\Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$$

of Rota-Baxter species such that $g = \Delta \circ i$.

Lemma 3.8. For any finite set X and any $F \in \text{ADF}[X]$,

$$(11) \quad (\text{id} \otimes \varepsilon_\emptyset) \circ \Delta_{X, \emptyset}(F) = (\varepsilon_\emptyset \otimes \text{id}) \circ \Delta_{\emptyset, X}(F) = F.$$

Proof. Let us consider the morphism of species $\rho : \text{ADF} \rightarrow \text{ADF}$ defined as follows: for any finite set X , for any $F \in \text{ADF}[X]$,

$$\rho_X(F) = (\varepsilon_\emptyset \circ \text{id}) \circ \Delta_{\emptyset, X}(F).$$

By composition, it is an algebra morphism. Let us prove that it is a Rota-Baxter algebra endomorphism of ADF. Let X be a finite set and $F \in \text{ADF}[X]$. We put $\Delta_{\emptyset, X}(F) = \sum F^{(1)} \otimes F^{(2)}$.

$$\begin{aligned} \rho_X \circ R_X(F) &= (\varepsilon_\emptyset \circ \text{id}) \circ \Delta_{\emptyset, X} \circ R_X(F) \\ &= (\varepsilon_\emptyset \circ R_\emptyset \otimes \text{id}) \circ \Delta_{\emptyset, X}(F) + \varepsilon_\emptyset(F^{(1)})\varepsilon_\emptyset(\bullet)R_X(F^{(2)}) \\ &= 0 + R_X(\varepsilon_\emptyset(F^{(1)})F^{(2)}) \\ &= R_X \circ \rho_X(F), \end{aligned}$$

as $\varepsilon \circ R = 0$. So ρ is a morphism of Rota-Baxter species. Moreover, for any singleton set $X = \{x\}$,

$$\Delta_{\emptyset, X}(\bullet x \bullet) = \bullet \otimes \bullet x \bullet,$$

so $\rho_X(\bullet x \bullet) = x \bullet x \bullet$. In other words, $\rho \circ i = \text{id}_{\text{ADF}} \circ i$. By the unicity part of the universal property, $\rho = \text{id}_{\text{ADF}}$. Consequently, for any finite set X and any $F \in \text{ADF}[X]$,

$$(\text{id} \otimes \varepsilon_\emptyset) \circ \Delta_{X, \emptyset}(F) = F.$$

We now consider the morphism of species $\rho' : \text{ADF} \rightarrow \text{ADF}$ defined as follows: for any finite set X , for any $F \in \text{ADF}[X]$,

$$\rho'_X(F) = (\text{id} \circ \epsilon_0) \circ \Delta_{X,0}(F).$$

By composition, it is an algebra morphism. Let us prove that it is a Rota-Baxter algebra endomorphism of ADF. Let X be a finite set and $F \in \text{ADF}[X]$. We put $\Delta_{X,0}(F) = \sum F_{(1)} \otimes F_{(2)}$.

$$\begin{aligned} \rho'_X \circ R_X(F) &= (\text{id} \otimes \epsilon_X) \circ \Delta_{X,0} \circ R_X(F) \\ &= R_X(F_{(1)})\epsilon_0(F_{(2)}) + \epsilon_X(F_{(1)}) \bullet \epsilon_0 \circ R_0(F_{(2)}) \\ &= R_X(F_{(1)})\epsilon_0(F_{(2)}) + 0 \\ &= R_X \circ \rho'_X(F), \end{aligned}$$

as $\epsilon \circ R = 0$. So ρ' is a Rota-Baxter species. As $\Delta_{X,0}(\bullet x \bullet) = \bullet x \bullet \otimes \bullet$, we conclude as for ρ that $\rho' = \text{id}_{\text{ADF}}$, which gives the missing equality. \square

Lemma 3.9. *For any finite set X with $X_1 \sqcup X_2 = X$ and any $F \in \text{ADF}[X]$, assume $\Delta_{X_1, X_2}(F) = \sum_{(F)} F_{(1)} \otimes F_{(2)}$, then*

$$(12) \quad \Delta_{X_1, X_2}(\epsilon_X(F) \bullet) = \sum_{(F)} \epsilon_{X_1}(F_{(1)}) \bullet \otimes \epsilon_{X_2}(F_{(2)}) \bullet.$$

Proof. By the proof of Theorem 2.16, $\Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$ is defined by induction on $|X|$.

If $|X| = 0$, then $X = \emptyset$. For a rooted forest $F \in \text{ADF}[\emptyset]$, $\Delta_{0,0}(F)$ is defined by induction on $\text{dep}(F)$. If $\text{dep}(F) = 0$, then $F = \bullet$ and

$$\Delta_{0,0}(\bullet) = \bullet \otimes \bullet.$$

Thus $\Delta_{0,0}(\epsilon_0(\bullet) \bullet) = \Delta_{0,0}(\bullet) = \bullet \otimes \bullet = \epsilon_0(\bullet) \bullet \otimes \epsilon_0(\bullet) \bullet$, Eq. (12) holds.

For the induction step of $\text{dep}(F) > 0$, assume $F = B^+(\bar{F})$ for some $\bar{F} \in \text{ADF}[\emptyset]$. Assume $\Delta_{0,0}(\bar{F}) = \sum_{(\bar{F})} \bar{F}_{(1)} \otimes \bar{F}_{(2)}$, then

$$\Delta_{0,0}(F) = R_0^{(2)}(\Delta_{0,0}(\bar{F})) = \sum_{(\bar{F})} (R_0(\bar{F}_{(1)}) \otimes \bar{F}_{(2)} + \epsilon_0(\bar{F}_{(1)}) \bullet \otimes R_0(\bar{F}_{(2)})).$$

Thus both sides of Eq. (12) equal to zero.

If $|X| > 0$, then at least one of X_1, X_2 is nonempty and so $\epsilon_{X_1}(\bar{F}_{(1)}) = 0$ or $\epsilon_{X_2}(\bar{F}_{(2)}) = 0$. Thus both sides of Eq. (12) equal to zero. \square

Proposition 3.10. *The triple $(\text{ADF}, \Delta, \epsilon)$ is a twisted coalgebra.*

Proof. Define another morphism of species

$$g' : \mathcal{O} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF},$$

$$g'_X(F) := \begin{cases} \bullet x \bullet \otimes \bullet \otimes \bullet + \bullet \otimes \bullet x \bullet \otimes \bullet + \bullet \otimes \bullet \otimes \bullet x \bullet, & X = \{x\}, F = \bullet x \bullet, \\ 0, & \text{otherwise.} \end{cases}$$

Then by Theorem 2.16 and Proposition 3.4, there is a unique morphism

$$\Delta' : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$$

of Rota-Baxter species such that $g' = \Delta' \circ i$.

As $\Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$ is a morphism of Rota-Baxter species,

$$(\Delta \otimes \text{id}) \circ \Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$$

and

$$(\text{id} \otimes \Delta) \circ \Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$$

are morphisms of twisted algebras. For $F \in \text{ADF}[X]$ with $X_1 \sqcup X_2 \sqcup X_3 = X$, we have

$$\begin{aligned} & (\Delta_{X_1, X_2} \otimes \text{id}) \circ \Delta_{X_1 \sqcup X_2, X_3} \circ R_X(F) \\ &= (\Delta_{X_1, X_2} \otimes \text{id}) \circ R_X^{(2)} \circ \Delta_{X_1 \sqcup X_2, X_3}(F) \quad (\text{by the proof of Theorem 2.16}) \\ &= (\Delta_{X_1, X_2} \otimes \text{id}) \circ (R_{X_1 \sqcup X_2} \otimes \text{id} + \bullet \epsilon_{X_1 \sqcup X_2} \otimes R_{X_3}) \circ \Delta_{X_1 \sqcup X_2, X_3}(F) \\ &= ((\Delta_{X_1, X_2} \circ R_{X_1 \sqcup X_2}) \otimes \text{id} + (\Delta_{X_1, X_2} \circ (\bullet \epsilon_{X_1 \sqcup X_2}) \otimes R_{X_3})) \circ \Delta_{X_1 \sqcup X_2, X_3}(F) \\ &= \left((R_{X_1 \sqcup X_2}^{(2)} \circ \Delta_{X_1, X_2}) \otimes \text{id} + (\bullet \epsilon_{X_1} \otimes \bullet \epsilon_{X_2} \otimes R_{X_3}) \circ (\Delta_{X_1, X_2} \otimes \text{id}) \right) \circ \Delta_{X_1 \sqcup X_2, X_3}(F) \quad (\text{by Lemma 3.9}) \\ &= (R_{X_1} \otimes \text{id} \otimes \text{id} + \bullet \epsilon_{X_1} \otimes R_{X_2} \otimes \text{id} + \bullet \epsilon_{X_1} \otimes \bullet \epsilon_{X_2} \otimes R_{X_3}) \circ (\Delta_{X_1, X_2} \otimes \text{id}) \circ \Delta_{X_1 \sqcup X_2, X_3}(F) \\ &= R_X^{(3)} p \circ (\Delta_{X_1, X_2} \otimes \text{id}) \circ \Delta_{X_1 \sqcup X_2, X_3}(F). \end{aligned}$$

Hence $(\Delta \otimes \text{id}) \circ \Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$ is a morphism of Rota-Baxter species. Similarly, $(\text{id} \otimes \Delta) \circ \Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF} \otimes \text{ADF}$ is also a morphism of Rota-Baxter species. Moreover, for any singleton set $X = \{x\}$ with $X_1 \sqcup X_2 \sqcup X_3 = X$, there is

$$(\Delta_{X_1, X_2} \otimes \text{id}) \circ \Delta_{X_1 \sqcup X_2, X_3}(\bullet x \bullet) = (\text{id} \otimes \Delta_{X_2, X_3}) \circ \Delta_{X_1, X_2 \sqcup X_3}(\bullet x \bullet),$$

which implies that

$$(\Delta \otimes \text{id}) \circ \Delta \circ i = g' = (\text{id} \otimes \Delta) \circ \Delta \circ i.$$

By the uniqueness of Δ' such that $\Delta' \circ i = g'$, we conclude $(\Delta \otimes \text{id}) \circ \Delta = \Delta' = (\text{id} \otimes \Delta) \circ \Delta$. Then by Lemma 3.8, $(\text{ADF}, \Delta, \epsilon)$ is a twisted coalgebra. \square

Theorem 3.11. *The quintuple $(\text{ADF}, m, \bullet, \Delta, \epsilon)$ is a twisted bialgebra.*

Proof. By the construction of $\Delta : \text{ADF} \rightarrow \text{ADF} \otimes \text{ADF}$, it is a morphism of twisted algebras. By Lemma 3.1, $\epsilon : \text{ADF} \rightarrow \mathbf{1}_k$ is a morphism of twisted algebras. By Propositions 2.14 and 3.10, we get that $(\text{ADF}, m, \bullet, \Delta, \epsilon)$ is a twisted bialgebra. \square

Remark 3.12. (a) By [3, §8.4.1], if a twisted bialgebra \mathbf{p} is connected in the sense that $\mathbf{p}[\emptyset]$ is one-dimensional over the base field, then \mathbf{p} is a twisted Hopf algebra. In our case, the species ADF is not connected since $\text{ADF}[\emptyset]$ contains all trees with one leaf. In particular,



\bullet , \bullet are both in $\text{ADF}[\emptyset]$. Thus one cannot apply that result and conclude that ADF is a twisted Hopf algebra. In fact, note that $\text{ADF}[\emptyset] = \mathbf{k}\{R^n(\bullet), n \geq 0\}$ and an easy induction proves that for any $n \geq 0$,

$$\Delta_{\emptyset, \emptyset}(R^n(\bullet)) = \sum_{k=0}^n R^k(\bullet) \otimes R^{n-k}(\bullet).$$

(b) On the other hand, as shown in [46, 47], a filtration and coproduct can be given to free Rota-Baxter algebras such that the resulting filtered bialgebra is connected, thereby yielding a Hopf algebra structure on free Rota-Baxter algebras. It would be interesting to lift this construction to species and show that the new Rota-Baxter species is a twisted Hopf algebra.

4. FOCK FUNCTORS AND COLORED FOCK FUNCTORS

In this section, we define the Fock functor from Rota-Baxter species to Rota-Baxter graded algebras. As consequence of the bialgebra structure on the free Rota-Baxter species in the previous section, we recover the construction of the free Rota-Baxter algebra [20, 27, 47] by direct construction.

We first recall the concept of Fock functors [3, 22]. For each $n \geq 0$, denote the set $\{1, \dots, n\}$ by \underline{n} with the convention $\underline{0} = \emptyset$. For a left \mathfrak{S}_n -module V , denote the space of coinvariants of V by $\text{Coinv}(V)$, i.e.,

$$\text{Coinv}(V) := V/\text{Vect}(x - \alpha \cdot x, x \in V, \alpha \in \mathfrak{S}_n).$$

Definition 4.1. [22]

(a) **(Full Fock functor)** Let \mathbf{p} be a species. Define $\mathcal{F}(\mathbf{p}) := \bigoplus_{n=0}^{\infty} \mathbf{p}[\underline{n}]$. If $f : \mathbf{p} \rightarrow \mathbf{q}$ is a morphism of specie, define

$$\mathcal{F}(f) : \begin{cases} \mathcal{F}(\mathbf{p}) & \rightarrow \mathcal{F}(\mathbf{q}), \\ x \in \mathbf{p}[\underline{n}] & \mapsto f_{\underline{n}}(x) \in \mathbf{q}[\underline{n}]. \end{cases}$$

Then \mathcal{F} is a functor from the category of species to the category of graded vector spaces.

(b) **(Bosonic Fock functor)** Let \mathbf{p} be a species. Define $\widehat{\mathcal{F}}(\mathbf{p}) := \bigoplus_{n=0}^{\infty} \text{Coinv}(\mathbf{p}[\underline{n}])$. If $f : \mathbf{p} \rightarrow \mathbf{q}$ is a morphism of species. Define

$$\widehat{\mathcal{F}}(f) : \begin{cases} \widehat{\mathcal{F}}(\mathbf{p}) & \rightarrow \widehat{\mathcal{F}}(\mathbf{q}), \\ \bar{x} \in \text{Coinv}(\mathbf{p}[\underline{n}]) & \mapsto \overline{f_{\underline{n}}(x)} \in \text{Coinv}(\mathbf{q}[\underline{n}]). \end{cases}$$

Then $\widehat{\mathcal{F}}$ is a functor from the category of species to the category of graded vector spaces.

For $m, n \in \mathbb{N}$, denote by $\sigma_{m,n} : \underline{m} \sqcup \underline{n} \rightarrow \underline{m+n}$ the bijection

$$\sigma_{m,n}(i) = \begin{cases} i & \text{if } i \in \underline{m}, \\ i + m & \text{if } i \in \underline{n}. \end{cases}$$

Then

Lemma 4.2. [22] *Let \mathbf{p} be a twisted algebra (resp. coalgebra, bialgebra).*

(a) $\mathcal{F}(\mathbf{p})$ is a graded bialgebra, in which

(i) the product is given by

$$x \cdot y = \mathbf{p}[\sigma_{m,n}] \circ m_{\mathbf{p}}(x \otimes y), \quad x \in \mathbf{p}[\underline{m}], y \in \mathbf{p}[\underline{n}], m, n \in \mathbb{N};$$

(ii) the unit is $1_{\mathbf{p}} \in \mathbf{p}[\underline{0}]$;

(iii) the coproduct is given by

$$\Delta(x) = \sum_{i=0}^m \mathbf{p}[\sigma_{i,m-i}^{-1}] \circ \Delta_{\mathbf{p}}(x), \quad x \in \mathbf{p}[\underline{m}], m \in \mathbb{N};$$

(iv) the counit is given by

$$\varepsilon(x) = \varepsilon_{\mathbf{p}}(x), \quad x \in \mathcal{F}(\mathbf{p}).$$

(b) $\widehat{\mathcal{F}}(\mathbf{p})$ is a graded bialgebra.

Definition 4.3. A **graded Rota-Baxter algebra** is a graded algebra $A = \bigoplus_{n=0}^{\infty} A_n$ with a Rota-Baxter operator $P : A \rightarrow A$ such that $P(A_n) \subseteq A_n$ for each $n \geq 0$.

Lemma 4.4. Let (\mathbf{p}, m, ℓ, R) be a Rota-Baxter species of weight λ . Then

- (a) $\mathcal{F}(\mathbf{p})$ is a graded Rota-Baxter algebra of the same weight, with the Rota-Baxter operator $P_{\mathcal{F}}$ defined by

$$P_{\mathcal{F}}(x) := R_{\underline{m}}(x), \quad x \in \mathbf{p}[\underline{m}], m \in \mathbb{N},$$

for the grafting operator $R_{\underline{m}}$ defined in Eq. (8).

- (b) $\widehat{\mathcal{F}}(\mathbf{p})$ is a graded Rota-Baxter algebra of the same weight, with the Rota-Baxter operator $P_{\widehat{\mathcal{F}}}$ defined by

$$P_{\widehat{\mathcal{F}}}(x) := \overline{R_{\underline{m}}(x)}, \quad \bar{x} \in \text{Coinv}(\mathbf{p}[\underline{m}]), m \in \mathbb{N}.$$

Proof. ((a)). By Lemma 4.2 (a), $\mathcal{F}(\mathbf{p})$ is a graded algebra. For $m, n \in \mathbb{N}$ and $x \in \mathbf{p}[\underline{m}], y \in \mathbf{p}[\underline{n}]$, we have

$$\begin{aligned} P_{\mathcal{F}}(x) \cdot P_{\mathcal{F}}(y) &= \mathbf{p}[\sigma_{m,n}] \circ m_{\underline{m}, \underline{n}}(R_{\underline{m}}(x) \otimes R_{\underline{n}}(y)) \\ &= \mathbf{p}[\sigma_{m,n}](R_{\underline{m} \sqcup \underline{n}}(R_{\underline{m}}(x) \cdot y + x \cdot R_{\underline{n}}(y) + \lambda x \cdot y)) \\ &= P_{\mathcal{F}}(P_{\mathcal{F}}(x) \cdot y + x \cdot P_{\mathcal{F}}(y) + \lambda x \cdot y). \end{aligned}$$

Hence $\mathcal{F}(\mathbf{p})$ is a graded Rota-Baxter algebra of weight λ .

((b)). Let I be the subspace of $\mathcal{F}(\mathbf{p})$ generated by the elements $x - \mathbf{p}[\alpha](x)$, where $x \in \mathbf{p}[\underline{n}]$, $\alpha \in S_n$ with $n \geq 0$. Then $\widehat{\mathcal{F}}(\mathbf{p}) = \mathcal{F}(\mathbf{p})/I$. By [22], I is an ideal of the algebra $\mathcal{F}(\mathbf{p})$. So we only need to show that I is a Rota-Baxter ideal, i. e., $P_{\mathcal{F}}(I) \subseteq I$. For any $x \in \mathbf{p}[\underline{n}], \alpha \in S_n$,

$$P_{\mathcal{F}}(x - \mathbf{p}[\alpha](x)) = R_{\underline{n}}(x - \mathbf{p}[\alpha](x)) = R_{\underline{n}}(x) - \mathbf{p}[\alpha](R_{\underline{n}}(x)) \in I.$$

Then $\widehat{\mathcal{F}}(\mathbf{p})$ is a graded Rota-Baxter algebra. □

Denote by \mathcal{H} the vector space of simple angularly decorated rooted forests whose angles are decorated by $\{1, 2, \dots, n\}$, where n is the number of angles. Denoted by $\widehat{\mathcal{H}}$ the vector space of all angularly decorated forests whose angles are decorated by one element. Then

Proposition 4.5. The vector spaces \mathcal{H} and $\widehat{\mathcal{H}}$ are both graded bialgebras and graded Rota-Baxter algebras.

Proof. By Theorems 2.16 and 3.11, the species ADF is a twisted Rota-Baxter algebra and a twisted bialgebra. Then by Lemmas 4.2 and 4.4, $\mathcal{F}(\text{ADF})$ and $\widehat{\mathcal{F}}(\text{ADF})$ are graded bialgebras and graded Rota-Baxter algebras. Then by Definition 4.1, $\mathcal{H} \cong \mathcal{F}(\text{ADF})$ and $\widehat{\mathcal{H}} \cong \widehat{\mathcal{F}}(\text{ADF})$ as graded vector spaces. Hence \mathcal{H} and $\widehat{\mathcal{H}}$ are both graded bialgebras and graded Rota-Baxter algebras. □

Remark 4.6. It is easy to check the multiplication and Rota-Baxter operator of $\widehat{\mathcal{H}}$ correspond to \diamond and B^+ , hence $\widehat{\mathcal{H}}$ agrees with the free non-commutative unitary Rota-Baxter algebra generated by one element given in [20, 27].

Let E be a given set. We define a specie S_E in the following way:

- For any finite set X , $S_E[X]$ is the vector space generated by sequences $(e_x)_{x \in X} \in E^X$ of elements of E indexed by X . The empty sequence is denoted by $\mathbf{1} \in S_E[\emptyset]$.
- For any bijection $\sigma : X \rightarrow Y$ of finite sets and any $(e_x)_{x \in X} \in E^X$, define

$$P[\sigma]((e_x)_{x \in X}) = (e_{\sigma^{-1}(y)})_{y \in Y}.$$

This species is a twisted bialgebra, where

- $m_{X,Y}((e_x)_{x \in X} \otimes (f_y)_{y \in Y}) = (g_z)_{z \in X \sqcup Y}$, with

$$g_z = \begin{cases} e_z & \text{if } z \in X, \\ f_z & \text{if } z \in Y. \end{cases}$$

The unit is the empty sequence $\mathbf{1}$.

- $\Delta_{X,Y}((e_x)_{x \in X \sqcup Y}) = (e_x)_{x \in X} \otimes (e_y)_{y \in Y}$. The counit sends the empty sequence $\mathbf{1}$ to 1.

Next we recall the Hadamard product of species [4, 22].

Definition 4.7. Let \mathbf{p} and \mathbf{q} be species. The **Hadamard product of \mathbf{p} and \mathbf{q}** is defined by

$$(\mathbf{p} \boxtimes \mathbf{q})[X] := \mathbf{p}[X] \otimes \mathbf{q}[X], \text{ for any finite set } X$$

$$(\mathbf{p} \boxtimes \mathbf{q})[\sigma] := \mathbf{p}[\sigma] \otimes \mathbf{q}[\sigma], \text{ for any bijection } \sigma : X \rightarrow Y.$$

Remark 4.8. If \mathbf{p} and \mathbf{q} are twisted algebras (respectively coalgebras, bialgebras), then $\mathbf{p} \boxtimes \mathbf{q}$ is also a twisted algebra (respectively coalgebra, bialgebra) [4], with componently defined product and coproduct.

If \mathbf{p} and \mathbf{q} are two species, for any $n \in \mathbb{N}$,

$$\begin{aligned} \text{Coinv}(\mathbf{p} \boxtimes \mathbf{q})[\underline{n}] &= \frac{\mathbf{p}[\underline{n}] \otimes \mathbf{q}[\underline{n}]}{\langle \mathbf{p}[\sigma](p) \otimes \mathbf{q}[\sigma](q) - p \otimes q \mid p \in \mathbf{p}[\underline{n}], q \in \mathbf{q}[\underline{n}], \sigma \in \mathfrak{S}_n \rangle} \\ &= \frac{\mathbf{p}[\underline{n}] \otimes \mathbf{q}[\underline{n}]}{\langle \mathbf{p}[\sigma^{-1}](p) \otimes q - p \otimes \mathbf{q}[\sigma](q) \mid p \in \mathbf{p}[\underline{n}], q \in \mathbf{q}[\underline{n}], \sigma \in \mathfrak{S}_n \rangle} \\ &= \frac{\mathbf{p}[\underline{n}] \otimes \mathbf{q}[\underline{n}]}{\langle p \cdot \sigma \otimes q - p \otimes \sigma \cdot q \mid p \in \mathbf{p}[\underline{n}], q \in \mathbf{q}[\underline{n}], \sigma \in \mathfrak{S}_n \rangle} \\ &= \mathbf{p}[\underline{n}] \otimes_{\mathfrak{S}_n} \mathbf{q}[\underline{n}], \end{aligned}$$

where the right action of \mathfrak{S}_n on $\mathbf{p}[\underline{n}]$ is given by

$$p \cdot \sigma = \mathbf{p}[\sigma^{-1}](p),$$

whereas the left action of \mathfrak{S}_n on $\mathbf{q}[\underline{n}]$ is given by

$$\sigma \cdot q = \mathbf{q}[\sigma](q).$$

The E -colored Fock functor $\widehat{\mathcal{F}}_E$ sends a species \mathbf{p} to

$$\widehat{\mathcal{F}}_E(\mathbf{p}) = \widehat{\mathcal{F}}(\mathbf{p} \boxtimes S_E) = \bigoplus_{n \in \mathbb{N}} \mathbf{p}[\underline{n}] \otimes_{\mathfrak{S}_n} S_E[\underline{n}].$$

If $f : \mathbf{p} \rightarrow \mathbf{q}$ is a species morphism, for any $\overline{p \otimes (e_i)_{i \in \underline{n}}} \in \mathbf{p}[\underline{n}] \otimes_{\mathfrak{S}_n} S_E[\underline{n}]$,

$$\widehat{\mathcal{F}}_E(f)(\overline{p \otimes (e_i)_{i \in \underline{n}}}) = \overline{f(p) \otimes (e_i)_{i \in \underline{n}}} \in \widehat{\mathcal{F}}_E(\mathbf{q}).$$

If E is a singleton, then $S_E[\underline{n}]$ is a trivial \mathfrak{S}_n -module of dimension 1 for any $n \in \mathbb{N}$, and $\widehat{\mathcal{F}}_E$ is naturally isomorphic to $\widehat{\mathcal{F}}$.

As S_E is a twisted bialgebra, we have

- If \mathbf{p} is a twisted algebra, then $\mathbf{p} \boxtimes S_E$ is a twisted algebra, so $\widehat{\mathcal{F}}_E(\mathbf{p})$ is an algebra.
- If \mathbf{p} is a twisted coalgebra, then $\mathbf{p} \boxtimes S_E$ is a twisted coalgebra, so $\widehat{\mathcal{F}}_E(\mathbf{p})$ is a coalgebra.
- If \mathbf{p} is a twisted bialgebra, then $\mathbf{p} \boxtimes S_E$ is a twisted bialgebra, so $\widehat{\mathcal{F}}_E(\mathbf{p})$ is a bialgebra.

Note that for ADF, $\widehat{\mathcal{F}}_E(\text{ADF})$ is just the vector space of angularly E -decorated forests. Hence, by Proposition 3.11, we recover the bialgebra structure of angularly decorated forests.

Corollary 4.9. *The vector space $\widehat{\mathcal{F}}_E(\text{ADF})$ of angularly E -decorated forests, equipped with the operations defined above, is a bialgebra.*

For $\overline{x \otimes (e_i)_{i \in \underline{m}}} \in \text{ADF}[\underline{m}] \otimes_{\varepsilon_m} S_E[\underline{m}]$, define

$$P(\overline{x \otimes (e_i)_{i \in \underline{m}}}) = \overline{R_m(x) \otimes (e_i)_{i \in \underline{m}}}.$$

Then the Rota-Baxter algebra structure of angularly decorated forests is recovered:

Proposition 4.10. *Equipped with the above operator P , the vector space $\widehat{\mathcal{F}}_E(\text{ADF})$ of angularly E -decorated forests is a Rota-Baxter algebra.*

Proof. We need to prove that P is well-defined. For any bijection $\sigma : \underline{m} \rightarrow \underline{m}$ such that $\text{ADF}[\sigma^{-1}](x) = x'$ and $\text{ADF}[\sigma]((e_i)_{i \in \underline{m}}) = (e'_i)_{i \in \underline{m}}$. By the definition of R_m , we have $\text{ADF}[\sigma^{-1}](R_m(x)) = R_m(\text{ADF}[\sigma^{-1}](x)) = R_m(x')$. Hence

$$P(\overline{x \otimes (e_i)_{i \in \underline{m}}}) = \overline{R_m(x) \otimes (e_i)_{i \in \underline{m}}} = \overline{R_m(x') \otimes (e'_i)_{i \in \underline{m}}} = P(\overline{x' \otimes (e'_i)_{i \in \underline{m}}}).$$

Then by the multiplication of twisted algebras and ADF being a Rota-Baxter species, $\widehat{\mathcal{F}}_E(\text{ADF})$ is a Rota-Baxter algebra. \square

Remark 4.11. Note that the elements of $\widehat{\mathcal{F}}_E(\text{ADF})$ are just the linear combinations of angularly E -decorated forests, and the multiplication and Rota-Baxter operator of $\widehat{\mathcal{F}}_E(\text{ADF})$ correspond to the product \diamond defined in Eq. (5) and the grafting operator B^+ . Hence $\widehat{\mathcal{F}}_E(\text{ADF})$ corresponds the free noncommutative unitary Rota-Baxter algebra generated by the set E as in [20, 27].

Acknowledgments. This research is supported by NNSFC (12301025 and 12101316).

Declaration of interests. The authors have no conflicts of interest to disclose.

Data availability. Data sharing is not applicable as no new data were created or analyzed.

REFERENCES

- [1] M. Aguiar, On the associative analog of Lie bialgebras, *J. Algebra* **244** (2001), 492–532. [2](#)
- [2] M. Aguiar, Dendriform algebras relative to a semigroup, *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), 15 pp. [2](#), [5](#)
- [3] M. Aguiar and S. Mahajan, Monoidal functors, species and Hopf algebras, CRM Monograph Series, vol. 29, American Mathematical Society, Providence, RI, 2010. [2](#), [16](#), [17](#)
- [4] M. Aguiar and S. Mahajan, Hopf monoids in the category of species, *Contemporary Mathematics* **585** (2013), 17–124. [2](#), [3](#), [4](#), [19](#)
- [5] M. Aguiar and W. Moreira, Combinatorics of the free Baxter algebra, *Electronic J. Combinator.* **13** (2006), (R17):1. [2](#), [3](#)
- [6] G. Andrew, L. Guo, W. Keigher and K. One, Baxter algebras and Hopf algebras, *Trans. Amer. Math. Soc.* **355** (2003), 4639–4656. [2](#)
- [7] P. Auger, G. Labelle and P. Leroux, Generalized binomial coefficients for molecular species, In memory of Gian-Carlo Rota, *J. Combin. Theory Ser. A* **91** (2000), 15–48. [2](#)
- [8] C. Bai, A unified algebraic approach to the classical Yang-Baxter equations, *J. Phys. A: Math. Theor.* **40** (2007), 11073–11082. [2](#)

- [9] C. Bai, O. Bellier, L. Guo and X. Ni, Splitting of operations, Manin products and Rota-Baxter operators, *Int. Math. Res. Not. IMRN* (2013), 485-524. [2](#)
- [10] C. Bai, L. Guo and T. Ma, Bialgebras, Frobenius algebras and associative Yang-Baxter equations for Rota-Baxter algebras, *Asian J. Math.* **28** (2024), 411-436 [2](#)
- [11] V. G. Bardakov and V. Gubarev, Rota-Baxter groups, skew left braces, and the Yang-Baxter equation, *J. Algebra* **596** (2022), 328-351. [2](#)
- [12] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* **10** (1960), 731-742. [2](#)
- [13] A. A. Belavin and V. G. Drinfeld, Solutions of the classical Yang-Baxter equation for simple Lie algebras, *Funct. Anal. Appl.* **16** (1982), 159-180. [2](#)
- [14] F. Bergeron, G. Labelle and P. Leroux, Combinatorial Species and Tree-Like Structures, Cambridge University Press, 1998. [2](#), [3](#)
- [15] M. Bordemann, Generalized Lax pairs, the modified classical Yang-Baxter equation, and affine geometry of Lie groups. *Comm. Math. Phys.* **135** (1990), 201-216. [2](#)
- [16] L. Carlier, Hereditary species as monoidal decomposition spaces, comodule bialgebras, and operadic categories, *Int. Math. Res. Not. IMRN* (2022), no. 8, 5745-5780. [2](#)
- [17] P. Cartier, On the structure of free Baxter algebras, *Adv. Math.* **9** (1972), 253-265. [2](#)
- [18] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem I: the Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* **210** (2000), 249-273. [2](#)
- [19] A. Das, Deformations of associative Rota-Baxter operators, *J. Algebra* **560** (2020), 144-180. [2](#)
- [20] K. Ebrahimi-Fard, L. Guo, Free Rota-Baxter algebras and rooted trees, *J. Algebra Appl.* **7** (2008), 167-194. [3](#), [6](#), [7](#), [8](#), [17](#), [18](#), [20](#)
- [21] M. Fiore, N. Gambino, M. Hyland, G. Winskel, The Cartesian closed bicategory of generalised species of structures, *J. Lond. Math. Soc. (2)* **77** (2008), 203-220. [2](#)
- [22] L. Foissy, Twisted bialgebras, cofreeness and cointeraction, arXiv:1905.10199 [2](#), [4](#), [5](#), [12](#), [13](#), [14](#), [17](#), [18](#), [19](#)
- [23] X. Gao, L. Guo, M. Rosenkranz, H. Zhang and S. Zhang, Integro-differential rings on species and derived structures, arXiv:2501.05540, 2025. [3](#)
- [24] I. M. Gessel, Counting tanglegrams with species, *J. Combin. Theory Ser. A* **184** (2021), Paper No. 105498, 15 pp. [2](#)
- [25] I. Z. Golubchik and V. V. Sokolov, Generalized perator Yang-Baxter equations, integrable ODEs and nonassociative algebras, *J. Nonlinear Math. Phys.* **7** (2000), 184-197. [2](#)
- [26] M. Goncharov and V. Gubarev, Double Lie algebras of a nonzero weight, *Adv. Math.* **409** (2022), part B, Paper No. 108680, 30 pp. [2](#)
- [27] L. Guo, An Introduction to Rota-Baxter Algebra, International Press, 2012. [2](#), [3](#), [5](#), [6](#), [7](#), [8](#), [17](#), [18](#), [20](#)
- [28] L. Guo, R. Gustavson and Y. Li, Generalized Reynolds algebras from Volterra integrals and their free construction by complete shuffle product, 2411.02633. [3](#)
- [29] L. Guo, H. Lang and Y. Sheng, Integration and geometrization of Rota-Baxter Lie algebras, *Adv. Math.* **387** (2021), 107834. [2](#)
- [30] A. Joyal, Une théorie combinatoire des séries formelles, *Adv. in Math.* **42** (1981), 1-82. [2](#), [3](#)
- [31] H. Lang and Y. Sheng, Factorizable Lie bialgebras, quadratic Rota-Baxter Lie algebras and Rota-Baxter Lie bialgebras, *Commun. Math. Phys.* **397** (2023), 763-791. [2](#)
- [32] E. Marberg, Strong forms of linearization for Hopf monoids in species, *J. Algebraic Combin.* **42** (2015), 391-428. [2](#)
- [33] E. Marberg, Strong forms of self-duality for Hopf monoids in species, *Trans. Amer. Math. Soc.* **368** (2016), 5433-5473. [2](#)
- [34] M. Maia and M. Méndez, On the arithmetic product of combinatorial species, *Discrete Math.* **308** (2008), 5407-5427. [2](#)
- [35] W. Norledge, Species-theoretic foundations of perturbative quantum field theory, arXiv:2009.09969. [2](#)
- [36] M. Rosenkranz and G. Regensburger, Solving and factoring boundary problems for linear ordinary differential equations in differential algebras, *J. Symbolic Computation* **43** (2008), 515-544. [3](#)
- [37] G.-C. Rota, Baxter algebras and combinatorial identities I, II, *Bull. Amer. Math. Soc.* **75** (1969), 325-329, 330-334. [2](#)
- [38] G.-C. Rota, Baxter operators, an introduction, in: *Gian-Carlo Rota on Combinatorics, Introductory Papers and Commentaries* (1995), 504-512, Birkhäuser. [2](#)

- [39] M. A. Semenov-Tian-Shansky, What is a classical r-matrix? *Funct. Anal. Appl.* **17** 254-272 (1983). [2](#)
- [40] R. Street, Weighted tensor products of Joyal species, graphs, and charades, *SIGMA* **12** (2016), 005, 20 pages. [2](#), [5](#)
- [41] P. Tamaroff, The cohomology of coalgebras in species, *Comm. Algebra* **50** (2022), 2811-2830. [2](#)
- [42] R. Tang, C. Bai, L. Guo and Y. Sheng, Deformations and their controlling cohomologies of \mathcal{O} -operators. *Commun. Math. Phys.* **368** (2019), 665-700. [2](#)
- [43] K. Wang and G. Zhou, The minimal model of Rota-Baxter operad with arbitrary weight, *Selecta Math.* **30** (2024), no. 5, Paper No. 99, 44 pp. [2](#)
- [44] J. A. White, On Cohen-Macaulay Hopf monoids in species, *Sem. Lothar. Combin.* 84B (2020), Art. 84, 12 pp. [2](#)
- [45] H. Yu, L. Guo and J.-Y. Thibon, Weak quasi-symmetric functions, Rota-Baxter algebras and Hopf algebras, *Adv. Math.* **344** (2019), 1-34. [2](#)
- [46] T. J. Zhang, X. Gao and L. Guo, Hopf algebras, cocycles and Rota-Baxter algebras. *J. Math. Phys.* **57** (2016), 101701. [2](#), [3](#), [16](#)
- [47] X. G. Zhang, A. Q. Xu and L. Guo, Hopf algebra structure on free Rota-Baxter algebras by angularly decorated rooted trees, *J. Algebr Combinat.* **55** (2022), 1331–1349. [3](#), [6](#), [16](#), [17](#)

UNIV. LITTORAL CÔTE D'OPALE, UR 2597 LMPA, LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES JOSEPH LIOUVILLE F-62100 CALAIS, FRANCE

Email address: loic.foissy@univ-littoral.fr

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, RUTGERS UNIVERSITY, NEWARK, NJ 07102, UNITED STATES

Email address: liguo@rutgers.edu

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGSU NORMAL UNIVERSITY, XUZHOU, JIANGSU 221116, P. R. CHINA

Email address: pengxiaosong3@163.com

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, LONDON SW7 2AZ, UK

Email address: yx3021@ic.ac.uk

SCHOOL OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF INFORMATION SCIENCE & TECHNOLOGY, NANJING, JIANGSU 210044, P. R. CHINA

Email address: zhangy2016@nuist.edu.cn