

Magnetically arrested disk flux eruption events to describe SgrA* flares

Eleni Antonopoulou^{1,2}, Argyrios Loules¹, Antonios Nathanail²

¹ Department of Physics, National and Kapodistrian University of Athens, University Campus, GR 15784 Zografos, Greece

² Research Center for Astronomy and Applied Mathematics, Academy of Athens, Soranou Efessiou 4, 115 27 Athens, Greece

Received Month Day, 2024; accepted Month Day, 2024

ABSTRACT

Context. Magnetically arrested disks are among the most suitable candidates for describing the gas accretion and observed emission in the vicinity of supermassive black holes.

Aims. This work aims to provide a direct correlation between the quasi-periodic *flux eruption events*, characteristic of MAD accretion disk simulations, and the observed flaring behavior in the Galactic center.

Methods. We employ a MAD accretion disk with a distinct counter-clockwise rotation and investigate the evolution of magnetized flux tubes generated during a prominent flux eruption event. Although these flux tubes demonstrate a clockwise pattern, they experience significant dragging by the accretion disk's rotation. This study models the motion of hot spots, formed on the disk's equatorial plane due to magnetic reconnection, as they travel along the magnetized flux tubes at a fraction of the speed of light.

Results. Hot spots with a relativistic ejection velocity are able to balance out the counter-clockwise dragging of the flux tube's foot-point on the disk and demonstrate a clockwise motion in the sky, that is in good agreement with the NIR flares in the Galactic center. In addition, our flare models favor face-on inclinations in the range $[0^\circ, 34^\circ]$ and $[163^\circ, 180^\circ]$ for SgrA*.

Conclusions. The flux eruption events that arise naturally in the MAD accretion state provide a promising framework for reproducing the observed flaring behavior in the vicinity of SgrA*.

Key words. black hole physics – magnetohydrodynamics (MHD) – radiative transfer – Galaxy:center – relativistic processes

1. Introduction

Sagittarius A*, the supermassive black hole in the Galactic center, demonstrates intense flaring events in the near-infrared (Genzel et al. 2003; Ghez et al. 2004) and X-ray (Baganoff et al. 2001) wavelengths several times a day. Recent observations of the GRAVITY instrument at the Very Large Telescope have shed light on both the astrometry and the polarimetric signature of bright NIR flares (GRAVITY Collaboration et al. 2018, 2020b, 2023). The most prominent events exhibit relativistic clockwise motion a few gravitational radii from the supermassive black hole, accompanied by a matching evolution in polarization angle over a period of approximately one hour. These profound observations provide significant constraints on the orbital motion of hot spots in the vicinity of SgrA* (GRAVITY Collaboration et al. 2020b) and suggest the existence of strong poloidal magnetic fields in the emitting region of our Galactic center (GRAVITY Collaboration et al. 2020a).

The origin of Galactic center hot spots was first investigated in Matsumoto et al. (2020), who pointed out the super-Keplerian pattern speed of the underlying accretion flow. Since then, several models have been employed to investigate the dynamics of the highly polarized NIR emission and reproduce the observed flaring behavior in the vicinity of SgrA* (Tursunov et al. 2020; Ball et al. 2021; Vos et al. 2022; Vincent et al. 2023; Yfantis et al. 2023; Lin et al. 2023; Aimar et al. 2023; Huang et al. 2024; Kocherlakota et al. 2024; Lin & Yuan 2024; Antonopoulou & Nathanail 2024). The aforementioned studies have provided significant insight into the parametric properties of such flares, favoring super-Keplerian motion and nearly face-on observation

angles, denoting the impact of considering ejected hot spot configurations and electromagnetic interaction within the inner magnetosphere, as well as highlighting the non-thermal properties of the emitting electron population. Nevertheless, the precise mechanism governing the ejection and energetic emission of hot spots in the Galactic center is still unknown.

On the other hand, magnetically arrested disk (MAD) models are among the most suitable candidates to describe active galactic nuclei that exhibit strong jets (Bisnovatyi-Kogan & Ruzmaikin 1974; Narayan et al. 2003; Suková et al. 2021; Cruz-Orsorio et al. 2022; Fromm, Christian M. et al. 2022). In addition, large-scale simulations of stellar mass feeding in the Galactic center have also successfully reproduced MAD disks (Ressler et al. 2019). In the MAD state, poloidal magnetic flux is advected toward the black hole by the accreting gas, until the magnetic pressure becomes strong enough to balance the ram pressure of the disk (Igumenshchev et al. 2003; Igumenshchev 2008; Tchekhovskoy et al. 2011). When the two competing forces reach equipartition, gas accretion is significantly hindered and can only occur via three-dimensional non-axisymmetric processes, such as the magnetic Rayleigh - Taylor instability (Papadopoulos & Contopoulos 2018).

General-relativistic magnetohydrodynamics (GRMHD) simulations of MAD accretion disks reveal a quasi-periodic cycle of magnetic flux accumulation and ejection, a process we define as a *flux eruption event*. In each flux eruption event, magnetized bundles or hot spots are expelled into the disk. One of the main energy-dissipation mechanisms responsible for hot spot generation close to the event horizon of an accreting black hole is magnetic reconnection (Ripperda et al. 2022; Nathanail et al. 2022b).




Fig. 1. Evolution of quantities on the equatorial plane throughout the selected flux eruption event. The first column illustrates the ratio of vertical magnetic field strength to gas pressure, the second column depicts the magnetization, and the third column shows their product $\sigma B_z^2/P_g$ used for the identification of magnetic reconnection sites on the equatorial plane. The white and black contours on the first two columns outline the magnetic reconnection sites illustrated in the third column.

Turbulence and instabilities in the accretion flows of MAD models generate large current sheets with alternating magnetic field polarity, which are highly efficient sites for particle acceleration (Kagan et al. 2015). Magnetic reconnection in these sheets creates chains of magnetized hot spots, each a few gravitational radii in size (Loureiro et al. 2007; Fermo et al. 2010; Uzdensky et al. 2010; Huang & Bhattacharjee 2012; Loureiro et al. 2012; Takamoto 2013; Nathanail et al. 2020; Chatterjee et al. 2021; Suková et al. 2021). While these hot spots have minimal impact on the overall accretion flow variability, they are linked to observed variability in NIR and X-ray wavelengths and episodic flares near our Galactic center (Dimitropoulos et al. 2024).

Consequently, MAD models are well-suited for flux tube formation and energetic flare emission, and provide promising candidates for reproducing the observed flares in the vicinity of SgrA*. The magnetic reconnection events that naturally arise in the MAD state have been shown to produce episodic flares that demonstrate the continuous rotation and typical timescales of NIR flares in the Galactic center (Dexter et al. 2020), as well as the polarization signature and timescales of flares at sub-mm wavelengths (Najafi-Ziyazi et al. 2023). Although the magnetic

energy associated with such events is capable of powering the observed IR and X-ray flares, the motion of the generated flux tubes within the accretion flow is substantially sub-Keplerian, in contrast with the GRAVITY observations (Porth et al. 2021). Most recently, (Lin & Yuan 2024) identified flux ropes consistent with the theoretical 'coronal mass ejection' model (Yuan et al. 2009) within GRMHD simulations of black hole accretion disks and semi-analytically reconstructed their evolved trajectories to fit the July 2018 flare.

This study extends previous research on MAD models by conducting a three-dimensional GRMHD simulation, focused on isolating flux tubes from energetic flux eruption events and modeling the ejected hot spot emission, that is in accordance with the most accurate astrophysical models for SgrA*. Although Event Horizon Telescope Collaboration et al. (2022) has not been able to identify a preferred Position Angle for the source, they have determined a cluster of strongly magnetized (MAD) models with positive spin and low inclination, that successfully pass all but two of the observational constraints for SgrA* across a broad frequency range. We employ a highly spinning Kerr black hole with parameter $a = 0.94$ and a counter-clockwise rotation (on the

plane of the sky) for the magnetized accretion disk, in agreement with one of the most promising MAD models. In this configuration, the generated flux bundles experience significant bending due to rotation and demonstrate a distinct clockwise pattern during the flux eruption events. The present analysis emphasizes the role of MAD configurations in modeling gas accretion near SgrA* and directly links simulated hot spot ejection to the observed flaring activity in our Galactic center. Our findings impose strict constraints on disk dynamics, revealing that a counter-clockwise rotation of the disk plasma, coupled with the clockwise motion of hot spots atop the flux bundle, is capable of replicating the observed NIR flares.

This letter is organized as follows: Section 2 outlines the methodology for identifying promising flux eruption events and modeling flares in MAD simulations (details on the modeling process can be found in Appendix C). Section 3 compares the flare models to the observations of GRAVITY Collaboration et al. (2018) and imposes constraints on the inclination of SgrA* (detailed discussion in Appendix E). Section 4 contains the conclusions of our analysis. The numerical setup for the GRMHD simulation and GRRT calculations is described in Appendix A, whereas the time series of the mass accretion rate and normalized magnetic flux are given in Appendix B. Appendix D focuses on replicating the first observational data points and Appendix F briefly investigates the scenario of clockwise disk rotation.

2. Methodology

This study investigates the evolution of a 3D MAD accretion disk simulation, focusing on the generation and development of energetic flux tubes during flux eruption events. These events are initiated by intense magnetic reconnection on the disk's equatorial plane, producing bundles of energized particles. We assume that the produced bundles travel along the magnetized flux tubes. Flux tubes in this context are typically dominated by coherent, large-scale vertical magnetic fields, resulting in significant magnetic field strength. Their foot-points on the disk are also highly magnetized¹. Notably, magnetic flux from previous flux eruption events remains within the disk, continuing to evolve (move around) and potentially influence subsequent eruptions, creating regions with "stagnating" magnetic flux. Therefore, we initially shift our focus to the accretion disk's equator and search for regions with a high value of magnetization, $\sigma \geq 0.4$, and a large ratio of vertical magnetic field strength over thermal pressure, $B_z^2/P_g \geq 0.8$, that have been produced by newly generated flares.

Our analysis centers on a specific flux eruption event occurring at $32228 M$ and originating approximately at $4.7 r_g$ from the supermassive black hole. Fig. 1 illustrates this event by displaying the ratio of vertical magnetic field strength over gas pressure (first column), the magnetization (second column), and their product (third column) evaluated on the equatorial plane of the accretion disk during the selected flux eruption event. Initially confined, the active region quickly expands in both azimuthal directions. As shown in Fig. 1, this region exhibits both an orbital motion due to the accretion disk's rotation, and an outward motion toward larger orbital radii, remaining active for over $90 M$ (Porth et al. 2021). This activity makes it an ideal candidate to model the observed flaring behavior in our Galactic center. In addition to this eruption, other flux eruption events appear throughout the simulation. These events tend to stagnate within the disk, moving passively with the flow. They can be identified




Fig. 2. Three-dimensional representation of the accretion flow and magnetic field topology during a flux eruption event. The color in the equatorial plane corresponds to the product of the magnetization times the vertical magnetic field strength, divided by the gas pressure ($\sigma \times B_z^2/P_g$), highlighting newly generated flux bundles (in red) created by the flux eruption event. The red field lines are drawn from the peak of the quantity plotted in color, corresponding to the most energetic part of the flux bundle. The grey lines represent the magnetic field lines of the accretion disk, while the orange lines show the magnetized funnel field lines, originating from the black hole and previously connected to the flux bundle before the event.

by their strong vertical magnetic field and characteristic peaks in B_z^2/P_g (first column of Fig. 1). Over time, their magnetization decreases, indicating that only recently generated flux tubes from fresh eruptions maintain high magnetization (second column of Fig. 1). The product of these quantities, shown in the third column of Fig. 1, marks where newly generated flux tubes are introduced into the disk via a flux eruption event.

After tracking the evolution of the magnetic reconnection site on the equatorial plane, we examine the three-dimensional morphology of the energetic flux tubes generated during the selected flux eruption event. Fig. 2 presents a 3D view of the magnetic field lines, showing in red the newly generated flux bundles from the flux eruption event initially identified on the equatorial plane. The disk's magnetic field lines are shown in grey, while the magnetized funnel field lines —connected to the red flux bundle prior to the eruption and originating from the black hole's event horizon— are highlighted in orange. This visualization captures the shape and dynamics of the magnetized flux tube originating from the active region identified in the third panel of Fig. 1, displaying its evolution at several points throughout the prominent flux eruption event. This approach allows us to track the morphology of magnetic field lines within the three-dimensional space-time of the MAD accretion disk, providing the framework for analyzing the evolution of the hot spot trajectory.

Notably, each hot spot model corresponds to a fixed flux tube structure captured at a specific timestamp in the MAD simulation, while the overall evolution of the flux tube's shape throughout the flux eruption event is represented by different models (see the left panel of Fig. 3). This study considers spherical hot spots with an effective emission radius of $1M$, that are generated at

¹ $\sigma = B_{co}^2/\rho$, where B_{co} is the magnetic field in the co-moving frame and ρ is the density of the accretion disk.




Fig. 3. Hot spot trajectories along the magnetized flux tubes generated during the prominent flaring event overlapped with the observations of July 22 2018 (grey and colorful circles, respectively). The direction of motion is indicated by blue arrows. The position of SgrA* in the sky is denoted by a black cross. *Left Panel:* Flare models for hot spot ejection velocities between $0.5c - 0.8c$ and foot-point orbital velocities ranging from $0.1u_K - 0.4u_K$. The best-fit black hole position, which is slightly different for each model, is illustrated by a blue rectangle. *Right Panel:* The flare model providing the best fit with the GRAVITY observations. The position of the black hole is depicted by a black diamond.

the equatorial plane of the disk and propagate along the magnetized flux tube with a constant ejection velocity a fraction of the speed of light. In addition, we track the position of the flux tube's foot-point on the accretion disk during the selected flux eruption event and implement both the outward motion and sub-Keplerian orbital rotation in the flare models, assuming solid-body rotation for the flux bundle. Note that other models, which reconstruct trajectories from the toroidal velocity from a numerical simulation, suggest that the hot spot moves in the same direction as the rotation of the flow (Nathanail et al. 2020; Lin & Yuan 2024). In contrast, our model predicts that the disk plasma and the hot spot, which is anchored to the flux bundle, move in the opposite direction. More details on the modeling procedure can be found in Appendix C.

3. Results

According to the analysis outlined in Section 2, the prominent flux eruption event illustrated in Fig. 1 constitutes a promising candidate for replicating the observed flaring behavior in the vicinity of SgrA*. In the following paragraphs, we track the trajectory of the hot spot along the magnetized flux tubes generated during the selected flux eruption event and calculate the emitted radiation for individual models. Unless stated otherwise, observer inclination is face-on throughout our calculations. We note that the focal point of this study is the most investigated flare of July 22 2018. The observed flare positions represent the average of the "Waisberg" and "Pfuhl" analyses, thus incorporating the individual differences between the GRAVITY Collaboration et al. (2018) reduction methodologies (a more detailed discussion is given in Appendix D).

The left panel of Fig. 3 illustrates the flaring behavior for various hot spot models –and therefore several flux tube configurations throughout the selected flux eruption event– overlapped with the observations of July 22 2018. Each flare model corresponds to a different ejection for the hot spot, ranging from $0.5c$ to $0.8c$, and orbital velocity for the flux tube's foot-point on the disk, between $0.1u_K$ to $0.4u_K$ based on the simulation (for more details see Appendix C). Even though the MAD accretion disk possesses a distinct counter-clockwise rotation, the observed hot spot trajectories demonstrate a clockwise motion in the sky, due to the relativistic velocity of the ejected hot spots. In particular, the flares modeled throughout the flux eruption event demonstrate a clear trend, whose general characteristics are in good agreement with the observed flaring behavior in our Galactic center.

The right panel of Fig. 3 depicts a flare model yielding very good agreement with the observations of July 22 2018. In this particular model, the ejected hot spot traverses along the magnetized flux tube with a relativistic velocity of $0.8c$, whereas the flux tube's foot-point on the accretion disk is rotating with an orbital velocity equal to $0.1u_K$. The illustrated trajectory demonstrates good correlation with the general trend of the GRAVITY observations and falls within the associated error bars for all but one of the observed flaring positions (see purple point at $t = 27.03\text{min}$). Although this particular flare model does not capture the kinematics of the first data points (dark red point at $t = 1.03\text{min}$), it still falls within the corresponding error bars, since the uncertainty for the first flaring position is significantly big. It is important to note, however, that a number of models illustrated in Fig. 3 are capable of reproducing the first data points of the July 22 flare. A more detailed discussion on the topic is given in Appendix D.

The final point of our analysis is to calculate the image of the best fit flare model (see right panel of Fig. 3) for several observer inclinations. We consider an evenly spaced range of observation angles up to 85° , along with their supplementary angles. At edge-on inclinations, the orbits exhibit significant deformation, either shrinking or stretching in the plane of the sky, which pushes the hot spot trajectories outside the bounds of the observed flaring emission (see Fig. D.2). Our analysis highlights the strong preference the GRAVITY observations demonstrate towards face-on inclinations and constrains the inclination of SgrA* to the range between $[0^\circ, 34^\circ]$ and $[163^\circ, 180^\circ]$. A more detailed discussion on the effect of inclination on the resulting flare emission is given in Appendix E.

4. Conclusions and Discussion

This study explored the dynamics of flux eruption events in MAD accretion flows, with a particular focus on the formation, evolution, and motion of hot spots generated from magnetic reconnection during a flux eruption event. We have performed a three-dimensional GRMHD simulation –motivated by the best-bet models of Event Horizon Telescope Collaboration et al. (2022)–, in order to identify energetic flux tubes and track their morphology and evolution within the disk's highly magnetized environment. Our calculations take into account both the orbital motion of the flux tube's foot-point on the disk and the outward radial motion identified in the simulation, while maintaining a constant ejection velocity for the hot spots. The focal point of this study is the incorporation of a counter-clockwise

motion for the MAD accretion disk, in contrast to the clockwise NIR flares observed with GRAVITY. This distinct sense of rotation is consistent with the assumption that the Rotation Measure can be entirely attributed to an internal Faraday rotation in the polarimetric analysis of [Event Horizon Telescope Collaboration et al. \(2024\)](#).

By modeling specific flux eruption events, we impose constraints on disk plasma rotation and examine the role of magnetic field geometry in shaping the flare trajectories. The flux eruption events observed in our simulation closely align with flaring activity seen near SgrA*, providing a promising framework for understanding high-energy phenomena in the vicinity of supermassive black holes. This study highlights the distinct counter-clockwise flow of the flux bundles and the clockwise motion of the energetic particles within them, driven by the rotational dynamics of the disk, as well as the bending of the magnetized tube. Our findings further highlight that MAD configurations offer a robust description of the variable environment of SgrA*, while at the same time providing a direct link between the multi-wavelength observations of the Galactic center and the episodic flares observed in the vicinity of the supermassive black hole.

However, to differentiate this model from other competing scenarios, specific ‘smoking gun’ observations are required. Key distinguishing features include the absence of a steady-state jet and the presence of an intermittent flow transitioning between quiescence and flaring activity. Crucially, observations at 43 and 86 GHz ([Nathanail et al. 2022a](#)) could reveal a significant signature of these behaviors, particularly through the detection or non-detection of a structure near the black hole’s base from a well-defined jet. A structure that would differ from what has been observed at the base of a jet, for example, in M87, would provide direct evidence for the unique magnetic field geometry proposed.

The quasi-periodic flux eruption events –characteristic of MAD simulations– and the flare model presented in this work offer valuable insights into the energetic phenomena near supermassive black holes beyond the Galactic center. Consequently, an interesting application of our flare model would be the quasi-periodic outflows (QPOs) observed in extragalactic nuclei. Most recently, [Pasham et al. \(2024\)](#) reported the discovery of an ultra-fast QPO from a low luminosity active galactic nucleus, which they attributed to a binary black hole system. In particular, they suggest that the intermediate-mass black hole secondary is orbiting the supermassive black hole primary and expelling blobs of gas into the magnetized funnel, in a mechanism closely related to our hot spot ejection model.

Last but not least, this study imposes strict constraints on the inclination of SgrA*, which should lie on the face-on range between $[0^\circ, 34^\circ]$ and $[163^\circ, 180^\circ]$. These findings are in agreement with the results of [Event Horizon Telescope Collaboration et al. \(2022\)](#), that has identified two promising MAD models with positive spin and low inclination to describe the supermassive black hole in our Galactic center. Specifically, the parameters for the model associated with our GRMHD simulation read: $(a, i) = (0.94, 30^\circ)$, and align perfectly with the results of our analysis. These insights contribute to a more detailed understanding of accretion-driven magnetic activity and offer a basis for future observational comparisons, particularly with near-infrared (NIR) and X-ray variability linked to episodic flares around supermassive black holes.

Acknowledgements

The authors thank O. Porth and I. Contopoulos for fruitful discussions and comments on this work, A. Liberatori and F. Zanias for support in the calculations, and the anonymous referee for constructive comments on this work. This work was supported by computational time granted from the National Infrastructures for Research and Technology S.A. (GRNET S.A.) in the National HPC facility - ARIS - under project ID 16033. AL acknowledges financial support by the State Scholarships Foundation (IKY) scholarship program from the proceeds of the “Nic. D. Chrysovergis” bequest.

References

- Aimar, N., Dmytriiev, A., Vincent, F. H., et al. 2023, *A&A*, 672, A62
 Antonopoulou & Nathanail. 2024, *A&A*, 690, A240
 Baganoff, F. K., Bautz, M. W., Brandt, W. N., et al. 2001, *Nature*, 413, 45
 Ball, D., Özel, F., Christian, P., Chan, C.-K., & Psaltis, D. 2021, *The Astrophysical Journal*, 917, 8
 Bisnovatyi-Kogan, G. S. & Ruzmaikin, A. A. 1974, *Ap&SS*, 28, 45
 Chatterjee, K., Markoff, S., Neilsen, J., et al. 2021, *MNRAS*, 507, 5281
 Cruz-Orsio, A., Fromm, C., Mizuno, Y., et al. 2022, *Nature Astronomy*, 6
 Dexter, J., Tchekhovskoy, A., Jiménez-Rosales, A., et al. 2020, *Monthly Notices of the Royal Astronomical Society*, 497, 4999
 Dimitropoulos, I., Nathanail, A., Petropoulou, M., Contopoulos, I., & Fromm, C. M. 2024, arXiv e-prints, arXiv:2407.14312
 Event Horizon Telescope Collaboration et al. 2019, *ApJL*, 875, L5
 Event Horizon Telescope Collaboration et al. 2022, *ApJL*, 930, L16
 Event Horizon Telescope Collaboration et al. 2024, *ApJ*, 964, L26
 Fermo, R. L., Drake, J. F., & Swisdak, M. 2010, *Physics of Plasmas*, 17, 010702
 Fishbone, L. G. & Moncrief, V. 1976, *ApJ*, 207, 962
 Fromm, Christian M., Cruz-Orsio, Alejandro, Mizuno, Yosuke, et al. 2022, *A&A*, 660, A107
 Genzel, R., Schödel, R., Ott, T., et al. 2003, *Nature*, 425, 934
 Ghez, A. M., Wright, S. A., Matthews, K., et al. 2004, *The Astrophysical Journal*, 601, L159
 Ghisellini, G. 2013, *Radiative Processes in High Energy Astrophysics* (Springer International Publishing)
 GRAVITY Collaboration et al. 2018, *A&A*, 618, L10
 GRAVITY Collaboration et al. 2020a, *A&A*, 643, A56
 GRAVITY Collaboration et al. 2020b, *A&A*, 635, A143
 GRAVITY Collaboration et al. 2023, *A&A*, 677, L10
 Huang, J., Zhang, Z., & Guo, Minyong & Chen, B. 2024, arXiv e-prints, arXiv:2402.16293
 Huang, Y.-M. & Bhattacharjee, A. 2012, *Phys. Rev. Lett.*, 109, 265002
 Igumenshchev, I. V. 2008, *The Astrophysical Journal*, 677, 317
 Igumenshchev, I. V., Narayan, R., & Abramowicz, M. A. 2003, *The Astrophysical Journal*, 592, 1042
 Kagan, D., Sironi, L., Cerutti, B., & Giannios, D. 2015, *Space Sci. Rev.*, 191, 545
 Kocherlakota, P., Rezzolla, L., Roy, R., & Wielgus, M. 2024, arXiv e-prints, arXiv:2403.08862
 Lin, X., Li, Y.-P., & Yuan, F. 2023, *MNRAS*, 520, 1271
 Lin, X. & Yuan, F. 2024, *MNRAS*, 531, 3136
 Loureiro, N. F., Samtaney, R., Schekochihin, A. A., & Uzdensky, D. A. 2012, *Physics of Plasmas*, 19, 042303
 Loureiro, N. F., Schekochihin, A. A., & Cowley, S. C. 2007, *Physics of Plasmas*, 14, 100703
 Matsumoto, T., Chan, C.-H., & Piran, T. 2020, *Monthly Notices of the Royal Astronomical Society*, 497, 2385
 McKinney, J. C. & Gammie, C. F. 2004, *The Astrophysical Journal*, 611, 977
 Najafi-Ziyazi, M., Davelaar, J., Mizuno, Y., & Porth, O. 2023, *Flares in the Galactic Centre II: polarisation signatures of flares at mm-wavelengths*
 Narayan, R., Chael, A., Chatterjee, K., Ricarte, A., & Curd, B. 2022, *MNRAS*, 511, 3795
 Narayan, R., Igumenshchev, I. V., & Abramowicz, M. A. 2003, *PASJ*, 55, L69
 Nathanail, A., Dhing, P., & Fromm, C. M. 2022a, *Mon. Not. R. Astron. Soc.*, 513, 5204
 Nathanail, A., Fromm, C. M., Porth, O., et al. 2020, *MNRAS*, 495, 1549
 Nathanail, A., Mpisketzis, V., Porth, O., Fromm, C. M., & Rezzolla, L. 2022b, *MNRAS*, 513, 4267
 Olivares, H., Porth, O., Davelaar, J., et al. 2019, *A&A*, 629, A61
 Pandya, A., Zhang, Z., Chandra, M., & Gammie, C. F. 2016, *The Astrophysical Journal*, 822, 34

- Papadopoulos, D. B. & Contopoulos, I. 2018, *Monthly Notices of the Royal Astronomical Society*, 483, 2325
- Pasham, D. R., Tombesi, F., Suková, P., et al. 2024, *Science Advances*, 10, eadj8898
- Porth, O., Mizuno, Y., Younsi, Z., & Fromm, C. M. 2021, *Monthly Notices of the Royal Astronomical Society*, 502, 2023
- Porth, O., Olivares, H., Mizuno, Y., et al. 2017, *Computational Astrophysics and Cosmology*, 4, 1
- Ressler, S. M., Quataert, E., & Stone, J. M. 2019, *Monthly Notices of the Royal Astronomical Society*, 492, 3272
- Ripperda, B., Liska, M., Chatterjee, K., et al. 2022, *The Astrophysical Journal Letters*, 924, L32
- Rybicki, G. B. & Lightman, A. P. 1986, *Radiative Processes in Astrophysics*
- Suková, P., Zajaček, M., Witzany, V., & Karas, V. 2021, *ApJ*, 917, 43
- Takamoto, M. 2013, *The Astrophysical Journal*, 775, 50
- Tchekhovskoy, A., Narayan, R., & McKinney, J. C. 2011, *Monthly Notices of the Royal Astronomical Society: Letters*, 418, L79
- Tursunov, A., Zajaček, M., Eckart, A., et al. 2020, *ApJ*, 897, 99
- Uzdensky, D. A., Loureiro, N. F., & Schekochihin, A. A. 2010, *Phys. Rev. Lett.*, 105, 235002
- Van der Laan, H. 1966, *Nature*, 211, 1131
- Vincent, F. H., Wielgus, M., Aimar, N., Paumard, T., & Perrin, G. 2023, *arXiv e-prints*, arXiv:2309.10053
- Vos, J., Mościbrodzka, M. A., & Wielgus, M. 2022, *A&A*, 668, A185
- Wong, G. N., Du, Y., Prather, B. S., & Gammie, C. F. 2021, *ApJ*, 914, 55
- Yfantis, A. I., Mościbrodzka, M. A., Wielgus, M., Vos, J. T., & Jimenez-Rosales, A. 2023, *arXiv e-prints*, arXiv:2310.07762
- Yuan, F., Lin, J., Wu, K., & Ho, L. C. 2009, *Monthly Notices of the Royal Astronomical Society*, 395, 2183
- Zhang, G. Q., Bégué, D., Pe'er, A., & Zhang, B. B. 2024, *ApJ*, 962, 135

Appendix A: Numerical Setup

This study investigates a MAD simulation, motivated by the best-bet models of [Event Horizon Telescope Collaboration et al. \(2022\)](#). The simulation is performed in three spatial dimensions with BHAC ([Porth et al. 2017](#); [Olivares et al. 2019](#))², using modified Kerr-Schild coordinates ([McKinney & Gammie 2004](#)) and 2–3 levels of adaptive mesh refinement. We employ units where $c = G = 1$, which corresponds to a length unit equal to the black hole mass $r_g = M$, and a base resolution of $384 \times 192 \times 192$ grid cells in the r -, θ -, and ϕ -directions, respectively. The initial conditions consist of a [Fishbone & Moncrief \(1976\)](#) torus in hydrodynamic equilibrium, with a constant specific angular momentum of $l = 6.76$, orbiting a Kerr black hole with a dimensionless spin parameter of $a = 0.94$. The inner edge of the disk lies at $r_{in} = 20M$, the density maximum is located at $r_{max} = 40M$, and the equation of state is ideal, with an adiabatic index of $\hat{\gamma} = 4/3$. Furthermore, the magnetic field is initialized as a nested loop described by the vector potential:

$$A_\phi = \max\left(\left(\frac{\rho}{\rho_{max}} \left(\frac{r}{r_{in}}\right)^3 \sin^3 \theta \exp\left(\frac{-r}{400}\right)\right) - 0.2, 0\right), \quad (\text{A.1})$$

where ρ_{max} is the maximum rest-mass density in the torus ([Event Horizon Telescope Collaboration et al. 2019](#); [Wong et al. 2021](#); [Cruz-Ororio et al. 2022](#); [Fromm, Christian M. et al. 2022](#); [Narayan et al. 2022](#); [Event Horizon Telescope Collaboration et al. 2022](#); [Zhang et al. 2024](#)). The field strength is set such that $2P_{max}/B_{max}^2 = 100$, where the global maxima of pressure P_{max} and magnetic field strength B_{max}^2 do not necessarily coincide. With this choice of initial magnetic field geometry and strength, the simulation progresses according to the MAD regime (see [Appendix B](#)). MAD models are characterized by high values of magnetic flux at the event horizon $\phi_{BH} \sim \phi_{crit} \sim 50$ and demonstrate a quasi-periodic cycle of flux eruption events. In particular, magnetic flux accretes onto the black hole until it reaches the saturation value $\phi_{BH} \geq \phi_{crit}$. Thereafter, accretion of additional flux leads to flux expulsion events, during which ϕ_{BH} rapidly decreases. In contrast, SANE models are characterized by low values of magnetic flux $\phi_{BH} \ll \phi_{crit}$ and do not demonstrate the characteristic flux eruption events of the MAD accretion state ([Cruz-Ororio et al. 2022](#); [Fromm, Christian M. et al. 2022](#)). As a result, this study does not investigate SANE accretion disk simulations.

Radiative transfer calculations are performed in Boyer-Lindquist coordinates with the Black Hole Imaging code, first presented in [Antonopoulou & Nathanail \(2024\)](#). A Kerr black hole, with dimensionless spin parameter $a = 0.94$, rests at the center of the coordinate system, and a grid of 600×600 photons is initialized at a distance of $500M$ from the center. Unless clearly stated otherwise, observer inclination is face-on $-i = 5^\circ$ to eliminate issues on the polar axis—throughout our calculations. In the following calculations, hot spot emission is modeled by a power-law electron distribution ([Pandya et al. 2016](#)), while synchrotron absorption is neglected as a first-order approximation.




Fig. B.1. Time series of the mass accretion rate (top panel) and the normalized magnetic flux (bottom panel) throughout the simulation. The dashed vertical lines indicate the times displayed in [Fig. 1](#).

Appendix B: MAD accretion state

We define the mass accretion rate and magnetic flux over the surface of the black hole’s horizon, as follows

$$\dot{M} \equiv - \iint \rho u^r \sqrt{-g} d\theta d\phi, \quad (\text{B.1})$$

$$\Phi_{BH} \equiv \frac{1}{2} \iint |B^r| \sqrt{-g} d\theta d\phi. \quad (\text{B.2})$$

where, ρ , u^r , g and B^r are the mass density, radial four-velocity, metric determinant, and radial component of the magnetic field as seen by the local fiducial observers, respectively. The dimensionless normalized magnetic flux at the horizon can then be defined as

$$\phi_{BH} \equiv \sqrt{4\pi} \frac{\Phi_{BH}}{\sqrt{\dot{M}}}. \quad (\text{B.3})$$

[Fig. B.1](#) shows the time series of the mass accretion rate and the dimensionless normalized magnetic flux during the selected flux eruption event. At time $30000M$, we can say that the simulation has already reached a quasi-stationary MAD state and quasi-periodic dips in the normalized magnetic flux can be identified throughout the bottom part of the figure. These dips correspond to flux eruption events in the accretion disk, where the magnetic flux is expelled and matter accretion is significantly halted. They occur when the parameter ϕ_{BH} reaches its saturation limit ($\phi_{BH} \sim 50$, [Tchekhovskoy et al. \(2011\)](#)). The dashed vertical lines indicate the snapshots of the flaring event illustrated in [Fig. 1](#).

Appendix C: Modeling MAD flares

Our basic assumptions in this work are the following: during magnetic reconnection, the released magnetic energy generates energetic particles, forming hot spots on the accretion disk’s equatorial plane, spanning a fraction of the gravitational radius. These hot spots are then ejected from the disk plane and travel along highly magnetized flux tubes at a fraction of the speed of light. It is clear from the evolution of the simulation that as the flux bundle moves counterclockwise with the disk’s accretion flow, it experiences bending in the opposite direction due to rotation. Consequently, hot spots trapped within these flux bundles are observed to move in a clockwise manner, opposite to the general flow of the accretion disk. The focus of [Section 2](#) was set on identifying a promising flux eruption event candidate, capable of replicating the flaring observations in the vicinity of SgrA*, as well as tracking the evolution of both the magnetic reconnection site on the disk’s equator and the magnetic field lines comprising the energetic flux tube. The following paragraphs outline the

² <https://www.bhac.science>

Model	a	Resolution	Duration (M)	$\hat{\gamma}$	$r_{in}(M)$	$r_{max}(M)$	$r_{out}(M)$
MAD	0.94	$384 \times 192 \times 192$	100.000	4/3	20	40	2.500

Table 1. GRMHD simulation parameters: Model specifies the accretion disk model, a is the dimensionless black hole spin, Resolution denotes the number of grid zones along each direction as $N_r \times N_\theta \times N_\phi$. Duration denotes the total duration of the simulation, γ is the fluid element adiabatic index, r_{in} and r_{max} are the inner and pressure maximum radii of the FM torus, and r_{out} is the outer radial boundary of the simulation domain.

modeling procedure adopted for the selected flux eruption event, illustrated in Fig. 1.

This model considers spherical hot spots with a constant radius of $r_{spot} = 0.5M$ and a radiative sphere of radius $r_{rad} = 1M$, respectively. In particular, the interior of the hot spot is considered to be optically thick, therefore the total non-thermal synchrotron emission originates from the radiative sphere surrounding it. Although the shape of the hot spot is certainly subjected to continuous alterations throughout the course of its outward trajectory (due to shearing, for instance), the precise mechanism behind this process strongly depends on the morphology of the highly magnetized surrounding plasma and has yet to be investigated in a concise manner. Due to a lack of a concise analytic prescription, a constant spherical hot spot shape is maintained throughout the radiative transfer calculations, with an effective emission radius of $1M$.

The hot spots are in principal generated at the equatorial plane of the disk, fueled by the magnetic reconnection process, and ejected outward along the magnetized flux tube. As a first approximation, we assume that the generated hot spot gains a large amount of energy, enough to traverse along the flux tube with a relativistic velocity a fraction of the speed of light. We further assume that the energetic hot spot effectively maintains a constant ejection velocity throughout its trajectory. In the present analysis, the ejection velocity ranges from 0.5 up to 0.8 times the speed of light for individual models.

Another important parameter to be taken into account is the motion of the active magnetic reconnection site on the equatorial plane of the disk. As discussed in section 2, the active region exhibits two distinct motion patterns throughout the flaring event. Firstly, one must consider the outward motion, toward larger orbital radii, by carefully tracking the position of the flux tube's foot-point on the equator of the accretion disk. For the flaring event illustrated in Fig. 1, this corresponds to an average radial velocity of $0.07c$. Secondly, one must account for the orbital motion of the active region, as a result of the accretion disk's rotation. However, the selection of magnetic field lines and foot-points on the equatorial plane introduces some deviations in the flux tube's rotation. In particular, the flux tube continuously changes shape during the flux eruption event, rendering the measurement of the precise orbital velocity for each foot-point a difficult task (Porth et al. 2021). This analysis assumes a solid-body rotation for the flux tube and investigates sub-Keplerian orbital frequencies ranging from $0.1u_K - 0.4u_K$ for individual models, based on the simulation.

Appendix D: Reproducing the first data points

The GRAVITY Collaboration et al. (2018) has employed four independent codes to fit the signal of SgrA*. They report that all four codes agree on the main features and results of the observed flares, despite their individual differences (fitting approach, number of free parameters, etc.). The flare positions investigated throughout the paper represent the average of the "Waisberg" and "Pfuhl" analyses, in agreement with previous




Fig. D.1. One of the flare models illustrated in Fig. 3 capable of capturing the kinematics of the first data points. *Top panel:* A close-up view of the selected flare model overlapped with the first four observed flaring positions. *Bottom panel:* A three-dimensional visualization of the corresponding flux tube. The hot spot positions for the first four data points are color-coordinated with the associated points in the Top panel.

studies. This methodology effectively incorporates the individual differences between the unique codes and reflects the systematic uncertainty of the GRAVITY observations. For instance, the "Waisberg" analysis of the July 22 flare results in a significantly less elongated loop in the y-direction than both the "Pfuhl" analysis and the averaged data (Fig. B.1. of GRAVITY Collaboration et al. (2018)). As a result, investigating the "Waisberg" flare positions would allow for configurations that produce less extended orbits to fit the GRAVITY observations. However, since the key




Fig. D.2. The best fit flare model of Fig. 3 for various inclinations, denoted on the bottom right legend of each panel, overlapped with the observations of July 22 2018 (grey circles). The direction of motion is indicated by blue arrows. The black hole position for each inclination is color-coordinated with the corresponding hot spot trajectory and illustrated by a diamond. *Left panel:* inclinations between $[0^\circ, 90^\circ]$, *Right panel:* inclinations between $[90^\circ, 180^\circ]$.

flare features are present in both analyses –such as the kinematics of the first data points –, we do not expect this choice to significantly affect the results of our work.

The focus of this Appendix is set on reproducing the first data points of the July 22 2018 flare. While several models have attempted to replicate the prominent flare of July 22, none has successfully captured the assumed kinematics of the first data point (notice the dark red point at $t = 1.03\text{min}$). Although the flare demonstrates a clear clockwise motion in the sky plane, the first flaring position exhibits a sudden push in the opposite direction (notice the bright red point at $t = 3.59\text{min}$), before continuing along the observed clockwise pattern.

Fig. D.1 depicts close-up view of the first four observations for the July 22 flare, overlapped with one of the hot spot models illustrated in Fig. 3 (left panel), as well as a three-dimensional visualization of the corresponding flux tube (right panel). This flare model corresponds to an orbital frequency of $0.4\omega_K$ for the foot-point on the disk, and an ejection velocity of $0.8c$ for the hot spot. The position of the hot spot along the flux tube for the first four GRAVITY observations is color-coordinated in both panels of Fig. D.1. Notice that the shape of the magnetized flux tube demonstrates a continuous clockwise loop and an upward trend, as well as a distinct counter-clockwise motion due to the rotation of the accretion disk. Even though the hot spot is continuously gaining height, it appears to be passing very close to its initial position (notice the dark red and orange points) due to the projection of the flux tube’s shape on the sky plane. The resulting hot spot motion is in great agreement with the observed flaring positions at the beginning of the flaring event, as is evident in Fig. D.1. Consequently, our analysis naturally recovers the observed flaring behavior of GRAVITY Collaboration et al. (2018).

Appendix E: Constraining the inclination of SgrA*

The trajectories illustrated in Fig. 3 correspond to realistic flux tube configurations that are self-consistently generated throughout the flux eruption events observed in a MAD accretion disk simulation. The following analysis investigates the image of the best fit hot spot trajectory (see the right panel of Fig. 3) for a distant observer at various inclinations.

Fig. D.2 illustrates the best fit hot spot model for an evenly spaced range of observation angles up to 85° (left panel), as well as their supplementary angles (right panel). We note that the best fit trajectory of Fig. 3 is depicted in red in both panels. It is evident that larger observation angles produce increasingly deformed hot spot orbits. Specifically, observation angles between $[0^\circ, 90^\circ]$ result in oblate trajectories that are shrunk in the

y -direction, whereas observation angles in the range $[90^\circ, 180^\circ]$ exhibit the opposite behavior, producing increasingly elongated orbits. The GRAVITY observations demonstrate a strong preference towards face-on inclinations, as illustrated in Fig. D.2. Observing the flares at an edge-on inclination introduces large deformations in the shape of the orbit and pushes the hot spot trajectory beyond the bounds of the observed flaring emission in the Galactic center. In particular, the present analysis constrains the inclination of SgrA* to the face-on range between $[0^\circ, 34^\circ]$ and $[163^\circ, 180^\circ]$. On the other hand, hot spot trajectories observed at an inclination higher than 34° and lower than 163° are incompatible with the observed flaring behavior due to their large degree of deformation.

Appendix F: Clockwise disk rotation

Throughout the paper, we have investigated the quasi-periodic flux eruption events of a MAD accretion disk with a distinct counter-clockwise rotation. The generated flux tubes in this configuration demonstrate a clockwise pattern on the sky, as illustrated in Fig. D.1. Section 3 demonstrated that flare models with a small sub-Keplerian disk rotation and a relativistic hot spot velocity are in good agreement with the observed flaring behavior in the Galactic center. On the contrary, if the accretion disk was rotating in a clockwise manner, the magnetized flux tubes would exhibit a counter-clockwise pattern on the sky. This section briefly investigates the scenario of clockwise accretion disk rotation.

To consider a clockwise rotation in our MAD simulation, one has to observe the accretion disk from below ($\theta_{obs} = \pi$). When viewed from below, the disk is indeed rotating in a clockwise manner with respect to the observer and the flux tubes demonstrate a distinct counter-clockwise pattern on the sky. We study the evolution of the selected flux eruption event (see Fig. 1) and model the motion of hot spots along the generated flux tubes, following the formulation of Appendix C. Fig. F.1 illustrates a selection of flare models (red lines) overlapped with the observations of July 22 2018 (grey circles). The depicted models correspond to various flux tube configurations during the selected flux eruption event, a hot spot ejection velocity equal to $0.1c$, and a clockwise disk rotation ranging from $0.8\omega_K$ to $0.9\omega_K$. When the flux tube is rotating in a clockwise manner -in accordance with the observed flares-, the motion of the ejected hot spot along the counter-clockwise shaped magnetic field lines is opposed to the clockwise evolution of the observed flare on the sky. Fig. F.1 demonstrates that flare models with a nearly Keplerian disk rotation are able to balance out the ejected hot spot velocity and exhibit a distinct clockwise evolution that is in good agreement




Fig. F.1. Flare models for a hot spot ejection velocity of $0.1c$ and foot-point orbital velocities between $0.8u_K - 0.9u_K$ (red lines) overlapped with the observations of July 22 2018 (grey circles). The direction of motion is indicated by blue arrows. The position of SgrA* in the sky is denoted by a black cross, whereas the best-fit black hole position, which is slightly different for each model, is illustrated by a blue rectangle.

with the observed flares in the vicinity of SgrA*. However, the rotation of the accretion disk in MAD simulations is well below the Keplerian limit and does not typically reach the values necessary to account for the flare models illustrated in Fig. F.1 (Porth et al. 2021).