vv, Number ii, pp. 1–18

disco: Distributional Synthetic Controls

Florian Gunsilius Emory University Atlanta, Georgia/US fgunsil@emory.edu David Van Dijcke University of Michigan Ann Arbor, Michigan/US dvdijcke@umich.edu

Abstract. The method of synthetic controls is widely used for evaluating causal effects of policy changes in settings with observational data. Often, researchers aim to estimate the causal impact of policy interventions on a treated unit at an aggregate level while also possessing data at a finer granularity. In this article, we introduce the new disco command, which implements the Distributional Synthetic Controls method introduced in Gunsilius (2023). This command allows researchers to construct entire synthetic distributions for the treated unit based on an optimally weighted average of the distributions of the control units. Several aggregation schemes are provided to facilitate clear reporting of the distributional effects of the treatment. The package offers both quantile-based and CDF-based approaches, comprehensive inference procedures via bootstrap and permutation methods, and visualization capabilities. We empirically illustrate the use of the package by replicating the results in Van Dijcke et al. (2024).

Keywords: synthetic controls, treatment effects, distributional analysis, causal inference, disco, optimal transport

1 Introduction

Synthetic control methods, pioneered by Abadie and Gardeazabal (2003) and formalized in Abadie et al. (2010), have revolutionized policy evaluation with aggregate data. These methods have become a cornerstone of modern causal inference, particularly in settings where traditional regression approaches are infeasible due to a limited number of treated units or the lack of a good counterfactual control group. While these methods have proven invaluable for estimating average treatment effects, they traditionally focus on matching means, potentially overlooking rich heterogeneous effects across the distribution of outcomes that are often of primary interest to researchers and policymakers.

The Distributional Synthetic Controls (DiSCo) method (Gunsilius 2023) addresses this limitation by extending the synthetic control framework to match entire outcome distributions rather than just means. This extension is particularly valuable in empirical settings where researchers have access to repeated cross-sectional data within aggregate units but cannot track individual units over time. Such scenarios are common in policy evaluation, where interventions occur at an aggregate level (e.g., states, counties, or firms) but researchers have access to individual-level or more granular data within these units. In these contexts, treatment effects may vary substantially across the distribution of outcomes, making that distribution a crucial object of study.

This article introduces the disco command in Stata, which implements the Distribu-

 \bigodot yyyy StataCorp LLC

tional Synthetic Controls method. We lay out the command syntax and usage, provide a practical overview of the underlying methodology, and illustrate the command's use in an empirical application. A companion R package is available at Van Dijcke et al. (2024). Replication code and data can be found at https://tinyurl.com/msz9ct2e. When using the command, please cite this paper.

2 The disco Command

2.1 Syntax

disco depvar idvar timevar *if* [in], <u>idtarget(varname)</u> <u>t</u>O(#) [options]

depvar is the outcome variable (numeric).

idvar is the unit identifier variable (numeric).

timevar is the time-period variable (numeric).

2.2 Options

Required Options

idtarget(varname) specifies the ID value of the treated unit in *idvar*.

t0(#) specifies the first time period of treatment in *timevar*. Note the slight difference in notation with Abadie and Gardeazabal (2003), who use capital T_0 to denote the last pre-treatment period.

Model Options

- m(#) specifies the number of grid points for the approximation of the integral in (4).See the explanation there. The default is 1000.
- g(#) specifies the number of grid points for the evaluation of the quantile functions and CDFs, i.e., for the vectors/functions that are returned by the command. The default is g(100).
- mixture requests the CDF-based approach using the 1-Wasserstein metric in (5) instead of the default quantile-based approach using the 2-Wasserstein metric in (3). This is recommended for categorical outcomes or when distributions are likely mixtures.
- **nosimplex** removes the constraint that weights must lie in the unit simplex, allowing for negative weights while maintaining that the weights sum to one. This enables extrapolation beyond the convex hull of control outcomes.

qmin(#) specifies the minimum quantile for the estimation range; the default is qmin(0).

qmax(#) specifies the maximum quantile for the estimation range; the default is qmax(1).

$\mathbf{2}$

Inference Options

- ci computes bootstrap confidence intervals for all distributional effects. Confidence intervals account for estimation uncertainty in both the weights and empirical distributions.
- boots(#) specifies the number of bootstrap replications for confidence intervals. The default is boots(300). More replications provide more stable intervals but increase computation time linearly.
- cl(#) sets the confidence level as a percentage for intervals. The default is cl(0.95). Must be between 0 and 1.
- permutation performs a permutation test by applying the estimation to each control unit as if it were treated, following Abadie et al. (2010); Van Dijcke et al. (2024). Reports a p-value for the null of no effect.
- seed(#) sets the random-number seed for reproducibility of bootstrap and permutation
 inference.
- **nouniform** requests pointwise rather than uniform confidence bands. Uniform bands control for multiple testing across quantiles but are wider than pointwise bands. Use of this option is discouraged since inference on functional objects should generally be uniform.

agg specifies the type of aggregation for summary statistics and plots. One of

- "quantile": summarize estimated quantile functions
- "cdf": summarize estimated CDFs
- "quantileDiff": summarize differences in quantiles between treated and synthetic
- "cdfDiff": summarize differences in CDFs between treated and synthetic
- samples(numlist) specifies quantile or CDF points to aggregate over for summary statistics. For quantiles, these are in [0,1]. For CDFs, these are values of the outcome variable. If not specified, the default is to partition the support (either [0,1] or the range of the outcome variable) into 5 equally spaced points ([0, 0.25, 0.5, 0.75, 1]) and aggregate the treatment effects within those intervals.

2.3 Stored Results

~ .	
Scalars	
e(amin)	minimum observed outcome value
e(amax)	maximum observed outcome value
e(m)	number of samples to approximate integrals in estimation step
e(g)	number of grid points to evaluate quantile function / CDF on
e(t_max)	last time period in the dataset
e(N)	number of observations
e(pval)	p-value from permutation test
e(doci)	indicates whether CIs were requested
e(t0)	first treatment period
e(cl)	confidence level (e.g., 0.95)
Macros	
e(cmd)	disco
e(agg)	aggregation type (e.g., "quantile")
e(cmdline)	command as typed
Matrices	
e(cids)	control IDs $(1 \times J)$
e(weights)	synthetic control weights $(J \times 1)$
$e(cdf_t)$	treated unit CDFs (G \times T)
e(cdf_synth)	synthetic CDFs $(G \times T)$
$e(quantile_t)$	treated unit quantiles $(G \times T)$
e(quantile_synth)	synthetic quantiles $(G \times T)$
e(cdf_diff)	CDF differences $(G \times T)$
e(quantile_diff)	quantile differences $(G \times T)$
e(summary_stats)	summary stats matrix (length of samples option - 1×7)

2.4 Post-estimation Commands

disco_estat

The disco_estat command displays summary statistics after disco estimation. It provides a detailed summary of aggregated distributional treatment effects, including point estimates, standard errors, and confidence intervals across different time periods. Aggregation is done along the partition of the support specified in the samples option in the disco command. Syntax:

disco_estat <u>sum</u>mary

Important: disco_estat requires that disco has been run previously with a specific agg and samples option, which cannot be altered post-estimation. This limitation is imposed to avoid the command having to pass large bootstrap matrices back to Stata when the ci option is specified.

disco_plot

The disco_plot command creates visualizations after disco estimation. It displays quantile functions, CDFs, and their differences over time, with optional confidence intervals. Syntax:

disco_plot [, options]

Plot types are automatically adapted to the aggregation method used (e.g., quantileDiff, cdfDiff). Confidence intervals appear if ci was specified in disco. The options for disco_plot are:

agg(string) specifies if the outcome variable is categorical and a CDF plot is requested in agg. If specified, a bar plot is created instead of a line plot.

categorical specifies if the outcome variable is categorical and a CDF plot is requested in agg. If specified, a bar plot is created instead of a line plot.

title(string) specifies a custom title for the graph. Defaults vary by plot type.

ytitle(string) specifies a custom y-axis title. Defaults vary by plot type.

xtitle(string) specifies a custom x-axis title. Defaults vary by plot type.

color1(string) specifies the color for the first series (default "blue").

color2(string) specifies the color for the second series in level plots (default "red").

cicolor(string) specifies the color for confidence intervals (default "gs12").

lwidth(string) specifies the line width (default "medium").

lpattern(string) specifies the line pattern for the second series (default "dash").

- legend(string) specifies legend options, passed directly to Stata's graph commands; see [R] legend_options.
- byopts(string) specifies options for small multiples using by(), such as rows(2) or ytitle; see [R] by_option.
- plotregion(string) specifies options for the plot region in Stata graphs; see [R] region_options.

graphregion(string) specifies options for the overall graph region; see [R] region_options.

scheme(string) specifies a scheme name for the graph; see [R] scheme_option.

hline(real) y coordinate for a dashed grey horizontal line; see [R] added_line_options.

vline(real) x coordinate for a dashed grey vertical line; see [R] added_line_options.

xrange(numlist) numlist of size 2 to set the x range; see [R] scale_options.

yrange(numlist) numlist of size 2 to set the y range; see [R] scale_options.

disco_weight

The disco_weight command displays and stores synthetic control weights after disco estimation, matching numeric unit IDs (id_var) with their corresponding string names (name_var). The results are stored in a new frame for further analysis or easy

export to tables.

```
disco_weight id_var name_var [, options]
```

The command requires that disco has been run previously. The options for disco_weight are:

n(#) specifies the number of top weights to display and store. The default is 5.

format(string) specifies the display format for weights. The default is %12.4f.

frame(string) specifies the name of the frame where results will be stored. The default
 is "disco_weights". If a frame with this name already exists, it will be replaced.

round (#) specifies the rounding precision for weights. The default is 0.0001.

The command stores the following results:

```
Scalars

r(n) number of top weights displayed

Macros

r(cmd) disco_weight

r(format) display format used

r(frame) name of frame where results are

stored
```

3 Methodological Background

3.1 Overview and Notation

Consider a panel of J + 1 units (for example, firms) observed over T periods. One unit (conventionally labeled as unit 1) receives treatment after period T_0 (or equivalently in period $t_0 = T_0 + 1$), while the remaining J units serve as potential controls. For each unit j and time period t, we observe individual-level outcomes Y_{ijt} (for example, employee wages). From these, we can construct the cumulative distribution function (CDF) $F_{Y_{it}}$

$$F_{Y_{jt}}(y) = P(Y_{jt} \le y)$$

which gives the probability P that the outcome of unit j at time t, Y_{jt} , is less than or equal to y, and its corresponding quantile function $F_{Y_{it}}^{-1}$,

$$F_{Y_{it}}^{-1}(q) = \inf\{x \in \mathbb{R} : q \le F(x)\},\tag{1}$$

which captures the value in the support of Y_{jt} that is larger than $q \times 100$ percent of the distribution. These functions capture the entire distribution of outcomes within each aggregate unit, preserving information about heterogeneity that would be lost by focusing only on means.

The goal is to estimate what the distribution of outcomes in the treated unit would have been in the absence of treatment. We denote this counterfactual distribution's

CDF as $F_{Y_{1t,N}}$ and its quantile function as $F_{Y_{1t,N}}^{-1}$ for periods $t > T_0$. The DiSCo method constructs entire synthetic distributions for the treated unit using optimally chosen weights on the control distributions.

3.2 Quantile-Based Approach

The quantile-based approach constructs the counterfactual by finding optimal weights λ_j that create a synthetic control distribution via a weighted average of control unit quantile functions:

$$F_{Y_{1t,N}}^{-1}(q) = \sum_{j=2}^{J+1} \lambda_j^* F_{Y_{jt}}^{-1}(q) \quad \text{for } q \in (0,1).$$
(2)

These weights are chosen to minimize the squared distance between the treated unit's pre-treatment quantile function and the weighted combination of control unit quantile functions, corresponding to minimizing the 2-Wasserstein distance,

$$\vec{\lambda}_{t}^{*} = \operatorname{argmin}_{\lambda \in \Delta^{J}} \int_{\operatorname{qmin}}^{\operatorname{qmax}} \left| \sum_{j=2}^{J+1} \lambda_{j} F_{Y_{jt}}^{-1}(q) - F_{Y_{1t}}^{-1}(q) \right| dq,$$
(3)

for $t \leq T_0$, where $(\vec{\lambda}_t^*) = (\lambda_{1t}^*, \ldots, \lambda_{jt}^*)$ is the vector of J weights (one for each control unit), and Δ^J is the J-dimensional unit simplex, which constrains the weights to satisfy $\sum_{j=2}^{J+1} \lambda_j = 1$ and $\lambda_j \geq 0$ for all j > 1. Then, the overall synthetic control weights are computed as a simple average over the pre-treatment weights,

$$\lambda_{j}^{*} = \frac{1}{T_{0}} \sum_{t=1}^{T_{0}} \lambda_{jt}^{*}.$$

Here, qmin and qmax default to 0 and 1, respectively, matching the entire distribution. In practice, researchers may sometimes wish to match or conduct inference on specific parts of the distribution. To accommodate this, the command allows users to specify the qmin and qmax options, restricting the integral to an interval $[a, b] \subset [0, 1]$.

To relax the positivity constraint and allow for extrapolation as in Doudchenko and Imbens (2016), the user can specify the **nosimplex** option in the main command. The primary motivation for choosing this particular objective function is that it has the intuitive interpretation of a "regression of quantile functions." Alternatively, one can interpret the problem as finding the set of weights that makes the treated unit's quantile function as similar as possible to the weighted average of the control units' quantile functions (Gunsilius 2023, p. 1109). This aligns with the interpretation of the weighted average quantile function, $\sum_{j=2}^{J+1} \lambda_j^* F_{Y_{jt}}^{-1}(q)$, as a Wasserstein barycenter (Agueh and Carlier 2011).

Note that problem (3) is "distributional" in the sense that it assigns a single weight to each quantile function in its entirety, which can be seen from the outer integration

over y. Thus, the synthetic control is truly a weighted average of distributions, rather than a pointwise average of quantiles as in (Chen 2020). The benefits of working with full distributions as the fundamental objects of estimation are: (1) it corresponds to the classical synthetic controls approach of assigning one single weight to every unit j; and (2) it allows for individuals to drop in and out of a unit, meaning one need not track individuals over time. This accommodates settings with entry and exit (e.g., employees leaving firms, as in Van Dijcke et al. (2024)), and settings where the researcher may only have a repeated cross-section rather than a balanced panel of observations.

In practice, the integral in (3) must be simulated, resulting in the empirical problem,

$$\vec{\lambda}_{i}^{*} = \underset{\bar{\lambda}\in\Delta^{J}}{\operatorname{argmin}} \frac{1}{G} \sum_{g=1}^{G} \sum_{j=2}^{J+1} \left| \lambda_{jt} F_{Y_{j_{t}}^{-1}}^{-1} \left(V_{g} \right) - F_{Y_{1t}}^{-1} \left(V_{g} \right) \right|^{2}, \tag{4}$$

where V_g are G independent draws from the uniform distribution on [0, 1], with G large. In practice, the command simply uses uniformly spaced draws from [0, 1]. We also use a single parameter G as both the number of simulation draws for the integral and the number of grid points on which we evaluate the synthetic control distributions in practice.

3.3 CDF-Based Approach

As an alternative, one can match the cumulative distribution functions directly, minimizing the 1-Wasserstein distance:

$$\vec{\lambda}_t^* = \operatorname{argmin}_{\lambda \in \Delta^J} \int_{-\infty}^{\infty} \left| \sum_{j=2}^{J+1} \lambda_j F_{Y_{jt}}(y) - F_{Y_{1t}}(y) \right| \, dy.$$
(5)

The synthetic control distribution function is then estimated analogously to (2) as

$$F_{Y_{1t,N}}(y) = \sum_{j=2}^{J+1} \lambda_j^* F_{Y_{jt}}(y) \quad \text{for } y \in \text{supp}(Y),$$

with $\operatorname{supp}(Y)$ the support of the outcome variable Y. Of course, one can easily obtain a synthetic control quantile function from this by using the pseudo-inverse in (1), and vice versa for (2). Indeed, the command always computes both synthetic quantile functions and CDFs, regardless of whether the mixture option is specified, and stores them in $e(quantile_synth)$ and $e(cdf_synth)$.

The CDF-based approach is often advantageous for discrete outcomes or settings where preserving specific portions of the distribution is critical (Gunsilius 2023, §4.3). To have the command solve (5) instead of (3), one can specify the **mixture** option. A prominent example where the mixture (CDF-based) approach is useful is when the outcome is a categorical variable. In that case, a weighted average of quantile functions, as computed in (3), will interpolate between the support points, which is undesirable.

A "mixture" of distribution functions, however, restricts the weighted average to the same support points as the "donor" distribution functions (see Figure 1 in Gunsilius (2023)). We further illustrate this with a simple simulation in Stata: our target (treated) distribution is a discrete uniform distribution with equal probability mass on $\{1, 2, 3, 4\}$, i.e.,

$$Pr(Y_t = i) = \frac{1}{4}$$
 for $i = 1, \dots, 4$.

Our donor distributions take this target distribution but place more mass at one of the support points each, i.e.,

$$Pr(Y_j = i) = \begin{cases} 1/5 & \text{if } i \neq j, \\ 2/5 & \text{if } i = j, \end{cases}$$

for i, j = 1, ..., 4. The "raw" control and target quantile functions corresponding to these random variables are plotted together in Figure 1a. The synthetic CDFs and quantile functions, constructed by either solving (3) (dashed) or (5) (dash-dot), are plotted against the target distribution (solid) in panels 1b and 1c. As clearly shown, the CDF-based synthetic controls correctly preserve the support of the control (donor) functions, while the quantile-based approach undesirably interpolates between support points. Of course, such interpolation *would* be desirable for a continuous outcome variable.

Finally, note that when your variable is categorical, you should specify the g and m options to reflect the number of support points. For instance, if your variable takes values on the integers from 1 to 10, you would specify g(10) and m(10). This ensures that the integral in (5) correctly sums over the support points. Note that the code does expect evenly spaced support points for this to work. If your variable is not evenly spaced, you can normalize it without affecting the estimates. Furthermore, when creating graphs for categorical variables with disco_plot, you may want to specify the categorical option to change the graph type to a bar plot instead of a line plot.

3.4 Inference

Permutation Test

When the **permutation** option is specified, the command carries out the distributional version of the classical permutation test for synthetic controls proposed in Gunsilius (2023) and further developed in Van Dijcke et al. (2024). The idea is intuitive: recompute the DiSCo estimator anew for each of the J control units, while pretending that the given control unit was, in fact, treated. If the treatment effect on the treated unit is not driven by statistical noise, the estimated treatment effect on the true treated unit should be an extreme value of the resulting distribution of "placebo" treatment effects.

The only complication in the distributional setting is that each single permutation delivers an entire distribution of treatment effects, so we need a way to aggregate these into a single statistic that we can rank. Echoing the use of the root mean squared prediction error in Abadie et al. (2010), Gunsilius (2023) proposed using the 2-Wasserstein







(c) Target and Synthetic Quantile Functions

Figure 1: Illustrative Simulation of mixture Option

distance. Then, an analogous permutation test to Abadie et al. (2010) is constructed by normalizing the average post-treatment prediction error by the average pre-treatment one (Van Dijcke et al. 2024, p.10),

$$r_{j} = \frac{R_{j} \left(T^{*} + 1, T\right)}{R_{j} \left(1, T^{*}\right)},$$

where

$$R_j(t_1, t_2) = \left(\frac{1}{t_2 - t_1 + 1} \sum_{t=t_1}^{t_2} d_t^2\right)^{1/2},$$

the root mean squared prediction error in the 2-Wasserstein distance at time t,

$$d_t^2 = \int_{q_{\min}}^{q_{\max}} \left| \sum_{j=1}^J \lambda_j^* F_{jt}^{-1}(q) - F_{0t}^{-1}(q) \right|^2 \mathrm{d}q,$$

with λ_j^* the optimal DiSCo weights. As explained above, one can test for effects in a restricted part of the distribution by setting qmin > 0 or qmax < 1. Then we calculate the p-value of the permutation test as

$$p = \frac{1}{J+1} \sum_{i=0}^{J} 1\left\{r_{j} \ge r_{0}\right\},\,$$

which calculates how frequently the normalized placebo effects exceed the true effect on the treated. The null hypothesis that the effect on the treated is drawn from the placebo distribution can be rejected at standard significance levels, for instance $p \leq 0.05$. This p-value is stored in the e(pval) scalar by the disco command when the permutation option is specified.

Bootstrapped Confidence Intervals

The permutation test described above provides an intuitive, finite-sample approach to inference that has become standard in synthetic controls. In addition to the permutation test, we also provide a bootstrap procedure to compute confidence intervals that account for the randomness in the estimation of the distributions. It can be requested by specifying the ci option. Note that this can be computationally intensive, depending on the number of bootstrap replications requested via the **boots** option. If ci is specified, the disco_plot and disco_estat commands will automatically include confidence intervals and/or bootstrapped standard errors.

In Van Dijcke et al. (2024, Theorem 1), we prove that the bootstrap is uniformly valid for the DiSCo method up to a negligible set, meaning that the bootstrapped empirical process,

$$\tilde{\mathbb{G}}_{n} = \sqrt{n} \left(\tilde{F}_{0tn,N}^{-1} - \hat{F}_{0tn,N}^{-1} \right) = \sqrt{n} \left(\sum_{j=1}^{J} \tilde{\lambda}_{jn}^{*} \tilde{F}_{jtn}^{-1} - \sum_{j=1}^{J} \hat{\lambda}_{jn}^{*} \hat{F}_{jtn}^{-1} \right)$$
(6)

converges uniformly to the "true" empirical process,

$$\mathbb{G}_n = \sqrt{n} \left(\hat{F}_{0tn,N}^{-1} - F_{0tn}^{-1} \right) = \sqrt{n} \left(\sum_{j=1}^J \hat{\lambda}_{jn}^* \hat{F}_{jtn}^{-1} - \sum_{j=1}^J \lambda_j^* F_{jtn}^{-1} \right),$$

where $\tilde{\lambda}_{jn}$ and \tilde{F}_{jtn}^{-1} are the bootstrapped optimal weights and quantile functions.

In practice, the following pseudo-code provides a high-level overview of how the bootstrapped confidence intervals are computed. For a more detailed exposition, see Algorithm 1 in Van Dijcke et al. (2024).

- 1. Estimate the main DiSCo weights λ_j^* on the original pre-treatment data by solving (the empirical version of) (3) or (5).
- 2. Construct the main post-treatment estimate the synthetic quantile function in (2).

3. For b from 1 to boots:

- a. Resample each firm's data (pre + post) to create a bootstrap sample.
- b. Re-estimate weights on the bootstrapped pre-treatment data.
- c. Construct the estimate for each bootstrapped post-treatment $t > T_0$, using (1) the new weights calculated with the resampled pre-treatment quantile functions and (2) the resampled post-treatment quantile functions: $\sum_{j=1}^{J} \tilde{\lambda}_j \tilde{F}_{jt}^{-1}$. d. Compute the bootstrap "gap" in (6).
- 4. Form confidence intervals/bands using the empirical distribution of the bootstrap "gaps."

3.5 Relation to Classical Synthetic Controls

The DiSCo method reduces to the classical synthetic controls method of Abadie and Gardeazabal (2003) when only aggregate data are given — i.e., when each unit j has a single aggregate observation rather than an entire distribution of observations (Gunsilius 2023, §4.1). As such, it complements the classical method for settings where researchers have access to sub-aggregate data *within* each unit j and want to (1) use all available data to inform estimation rather than collapsing it down to a single point for each unit and/or (2) estimate distributional treatment effects.

3.6 Implementation Details

The command uses a C++ plugin for the constrained quadratic optimization in (3) to improve performance, based on the **quadprog** code by Luca Di Gaspero. This code implements the Goldfarb–Idani active-set dual method (Goldfarb and Idnani 1983). The C++ plugin is compatible with Linux/Unix, macOS (both Apple Silicon and Intel), and Windows system architectures. For the optimization in (5), the command relies on the LinearProgram class in Mata.

Memory requirements scale with the number of units (J), time periods (T), grid points (G), and bootstrap replications (B).

4 Empirical Illustration: Van Dijcke et al. (2024)

To contextualize the commands, we now use them to reproduce the main analysis in Van Dijcke et al. (2024). The code and data for this exercise are available at https://github.com/Davidvandijcke/disco_stata_journal.

4.1 Data and Methods

Data Overview. In the paper, we use quarterly résumé data from People Data Labs,¹ a large provider of individual-level employment records that covers over half of the total

^{1.} https://www.peopledatalabs.com/

headcount for most major tech firms (see Van Dijcke et al. 2024 for details). Each firm-quarter includes a repeated cross-section of employees, from which we observe job tenure and a seniority rank (ranging from "unpaid" to "CXO"). We merge in information on return-to-office (RTO) mandates gathered from public announcements, employee forums, and the Flex Index by Scoop Technologies, as well as layoff records from layoffs.fyi. Our main estimation sample focuses on the return to office at Microsoft in April 2022, plus a control pool of firms never adopting RTO within the observation window. For more details on the data, see Van Dijcke et al. (2024). In practice, we use a perturbed version of the original data for confidentiality reasons, where we add uniform noise to the outcome variables. This preserves the shape of the distributions while keeping the dataset shareable.

Empirical Approach. Using these data, we then apply a distributional synthetic controls design that computes synthetic control weights to closely match Microsoft's distribution of employee tenure in the two quarters before its RTO. The post-RTO gap between the observed and synthetic distributions isolates the policy's causal effect on workforce composition.

In the notation from Section 3, *i* refers to a single employee at firm *j* (with j = 1 corresponding to Microsoft), *t* refers to a quarter of the year, and Y_{ijt} is one of two main outcomes:

- 1. $Tenure_{ijt}$, the number of days employee *i* has been employed at company *j*.
- 2. $Title_{ijt}$, an integer classification of employee *i*'s job title at firm *j*, ranging from 1 (trainee) to 10 (C-suite). See Van Dijcke et al. (2024, Table 1) for the full classification.

We solve (3) for $Tenure_{ijt}$ (as it is continuous) and (5) for the categorical $Title_{ijt}$. Put differently, it makes sense to interpolate over the support of tenure (e.g., 5 days, 5.5 days), but it generally does not make sense to talk about an employee being "25% junior associate and 75% manager."

4.2 Results: Quantile Approach

First, we reproduce the main Figure 4 in the paper. To that end, we start by loading and inspecting the (perturbed) data:

. use "tenure_anonymized.dta", clear

. list in 1/5, ab(20)

	time_col	id_col	company_name	y_col
1.	2	17	oracle	1682.25
2.	3	1	deloitte	375.783
з.	2	72	Зm	5276.48
4.	3	2	microsoft	957.55
5.	2	2	microsoft	2745.96

The data structure shows the essential columns required for estimation: (1) a *time column*, indicating the time period (quarters of 2022); (2) a numeric ID column indicating the ID of the aggregate unit (firms, e.g. ID "2" for Oracle); (3) a numeric outcome variable (days the employee at Oracle had been working there at the start of quarter 2). There is also an optional string column indicating the name of the aggregate unit ("Oracle"), which can be used in the disco_weight command to inspect synthetic control weights by name.

With these data, we can replicate the figure with:

Here, we evaluate the quantile functions on 10 grid points using g(10), but use 100 points for estimation (m(100)) to approximate the integral in (4). In the paper, we used 1,000 samples; we lower that to 100 here for replication speed. Similarly, we compute confidence intervals via the ci option with only 300 bootstrap repetitions (boots(300)) instead of 1,000 as in the paper.

After estimation, we first examine the top 5 weights by company name:

```
. disco_weight id_col company_name, n(5)
Top 5 weights:
```

name	weight
amazon	.2203
autodesk	.1271
cisco	.1066
dell technologies	.0991
slalom consulting	.0962
slalom consulting	.0962

These weights are very close to those in Van Dijcke et al. (2024, Table 3), despite the data perturbation. As noted there, firms like Amazon, Cisco, and Dell receive high weights because they are highly similar to Microsoft (the treated firm).

We then directly plot the quantile effects, defined as the differences between the treated and synthetic quantile functions at each of the 10 deciles:

This command creates Figure 2, which closely resembles Figure 4 in the paper.

As discussed in the paper, panel 3 in the figure suggests a statistically significant decrease in tenure (about 50 days) for employees in the top 2–3 deciles of the tenure distribution, but no such decrease at the lower deciles. In other words, the RTO caused longer-tenured employees to leave Microsoft, but not those with shorter tenures. Because these effects are calculated as the difference between the observed and the synthetic quantile function, they have a counterfactual interpretation — i.e., the drop in tenure at the top deciles is relative to a "synthetic" Microsoft that did not return to the office.



Figure 2: Replication of Tenure Result

Finally, panels 1 and 2 in the figure show the quantile differences for the two quarters before the RTO. Since 0 is included in the 95% confidence bands there, we see that the synthetic control replicated Microsoft's tenure distribution well, supporting its use for constructing post-treatment counterfactuals.

Although not shown in the paper, a researcher may be interested in comparing the synthetic and observed quantile functions side by side. This can be done with:

```
. disco_plot, title(" ") ytitle("Tenure (Days)") agg("quantile")
```

which yields Figure 3.

We observe similar patterns here: a strong pre-treatment fit, with the two quantile functions overlapping, and a downward shift in the synthetic quantile function at the top deciles after the RTO. Additionally, it clarifies the absolute magnitudes in the distribution: the top 3 deciles, where the largest effects occur, correspond to employees who have been at Microsoft for at least 1,500 days (about 4 years).

Finally, a researcher may want to study the treatment effects in aggregate parts of the distribution. To do so, we can run:

. disco_estat summary



Figure 3: Replication of Tenure Result

Time period	Range		Effect	Std. Err.	[.95% Conf. In	Interval]
3	0.00-0.25	-6.659	26.779	-59.147	45.828	
3	0.25-0.50	-21.443	26.779	-73.931	31.045	
3	0.50-0.75	-50.509	26.779	-102.997	1.978	
3	0.75-1.00	-54.751	26.779	-107.239	-2.263*	

Summary of quantile effects

* denotes significance at .95% confidence level

By default, the command aggregates the treatment effects into four buckets corresponding to the four quartiles. We see a statistically significant effect in the top quartile only, corresponding to a decrease of about 55 days in employee tenure.

4.3 Results: CDF Approach

An analogous exercise can be carried out for the categorical Title variable, except that we now want the command to solve (5), as this variable takes integer values from 1 to 10. Hence, we specify the **mixture** option and force the command to use a grid of 10 points by setting g(10) and m(10). Concretely:

- . use "titles_anonymized.dta", clear
- . list in 1/5, ab(20)

	time_col	id_col	company_name	y_col
1.	3	17	oracle	8
2.	2	1	deloitte	5
з.	3	1	deloitte	6
4.	1	1	deloitte	5
5.	2	245	splunk	4

. disco y_col id_col time_col, idtarget(2) t0(3) agg("cdfDiff") seed(12143) ///
> mixture g(10) m(10) ci boots(300)

. disco_plot, title(" ") ytitle("Change in CDF") hline(0) categorical ///
> scheme("stsj") color("bluishgray")



Figure 4: Replication of Titles Result

Since y_col is clearly categorical, specifying categorical in the disco_plot command generates a bar plot rather than a line plot. The resulting figure, shown in Figure 4, closely resembles Figure 5 in Van Dijcke et al. (2024), though the data perturbation slightly widens the confidence intervals.

This figure can be interpreted as showing a statistically significant increase in the

mass under the CDF for employees with titles below the (senior) manager level (4), but no such increase for higher-ranked employees. By the nature of cumulative distribution functions, this means Microsoft's workforce rebalanced toward lower-ranked employees after the RTO, likely due to an outflow of senior talent.

5 Conclusion

This article introduced a new Stata command, disco, which implements the Distributional Synthetic Controls method of Gunsilius (2023). A practically oriented methodological section provided background, and we illustrated the command's usage through examples from Van Dijcke et al. (2024). A companion R package is available at Van Dijcke et al. (2024).

6 References

- Abadie, A., A. Diamond, and J. Hainmueller. 2010. Synthetic control methods for comparative case studies: Estimating the effect of California's tobacco control program. *Journal of the American statistical Association* 105(490): 493–505.
- Abadie, A., and J. Gardeazabal. 2003. The economic costs of conflict: A case study of the Basque Country. *American economic review* 93(1): 113–132.
- Agueh, M., and G. Carlier. 2011. Barycenters in the Wasserstein space. SIAM Journal on Mathematical Analysis 43(2): 904–924.
- Chen, Y.-T. 2020. A distributional synthetic control method for policy evaluation. Journal of Applied Econometrics 35(5): 505–525.
- Doudchenko, N., and G. W. Imbens. 2016. Balancing, regression, difference-indifferences and synthetic control methods: A synthesis. Technical report, National Bureau of Economic Research.
- Goldfarb, D., and A. Idnani. 1983. A numerically stable dual method for solving strictly convex quadratic programs. *Mathematical programming* 27(1): 1–33.
- Gunsilius, F. F. 2023. Distributional synthetic controls. *Econometrica* 91(3): 1105–1117.
- Van Dijcke, D., F. Gunsilius, and S. He. 2024. DiSCos: Distributional Synthetic Controls Estimation. http://www.davidvandijcke.com/DiSCos.
- Van Dijcke, D., F. Gunsilius, and A. Wright. 2024. Return to Office and the Tenure Distribution. arXiv preprint arXiv:2405.04352.

About the authors

Florian Gunsilius is an Associate Professor in the Department of Economics at Emory University. His research interests are nonparametric approaches for statistical identification, estimation, and inference. His current focus is on statistical optimal transport theory, mean field estimation, causal inference, and free discontinuity problems.

David Van Dijcke is a Ph.D. candidate in Economics at the University of Michigan. His research interests include econometrics, industrial organization, and political economy.