The regular singular inverse problem in differential Galois theory

Thomas Serafini and Michael Wibmer^{*}

January 15, 2025

Abstract

We show that every linear algebraic group over an algebraically closed field of characteristic zero is the differential Galois group of a regular singular linear differential equation with rational function coefficients.

1 Introduction

Let k be an algebraically closed field of characteristic zero. We consider linear differential equations over k(z), the field of rational functions with coefficients in k, equipped with the usual derivation $\frac{d}{dz}$. The solution of the inverse problem in differential Galois theory states that every linear algebraic group over k is the differential Galois group of a linear differential equation with coefficients in k(z). This was first proved in [Har05], based on previous work of several authors.

There are various directions into which the inverse problem can be generalized. For example, instead of trying to realize linear algebraic groups as differential Galois groups, one can try to realize surjective morphisms of linear algebraic groups. This idea can be formalized through differential embedding problems: Recall that a *differential embedding problem* for k(z) is given by a Picard-Vessiot extension L of k(z) with differential Galois group H together with a surjective morphism $G \to H$ of linear algebraic groups. A *solution* consists of a Picard-Vessiot extension M of k(z), containing L, with differential Galois group G such that the restriction map $G \to H$ of differential Galois groups corresponds to the given morphism $G \to H$. Note that the special case H = 1 recovers the inverse problem. Building on previous work ([Bac+18], [Bac+21a], [Bac+21b], [Bac+21c]), it was recently shown in [FW24] that every differential embedding problem for k(z) has a solution.

Another way to sharpen the inverse problem is to restrict the class of differential equations via which one hopes to realize a given linear algebraic group as a differential Galois group. As the class of *regular singular differential equations* is one of the best studied classes of linear differential equations, it is natural to wonder "Is every linear algebraic group the Galois group of a regular singular linear differential equation?" Our main result is that the answer is yes. In fact, we offer a somewhat more precise statement.

Theorem (Theorem 2.16). Let $G \leq \operatorname{GL}_{n,k}$ be a linear algebraic group over k that can be generated by d elements (as an algebraic group) and let S be a subset of $\mathbb{P}^1(k)$ with d + 1 elements. Then there exists a matrix $A \in k(z)^{n \times n}$ such that the differential equation $\partial(y) = Ay$ is regular singular with singularities contained in S and has differential Galois group G.

^{*}Supported by the Lise Meitner grant M-2582-N32 of the Austrian Science Fund FWF.

Mathematics Subject Classification Codes: 34M50, 12H05. Key words and phrases: Differential Galois theory, regular singular differential equations, Riemann-Hilbert correspondence, free proalgebraic groups.

The above theorem is reminiscent of Schlesinger's density theorem: For the field $k = \mathbb{C}$ of complex numbers, the differential Galois group of a regular singular differential equation with d + 1 singularities is the Zariski closure of the monodromy group, which is generated by d elements. Moreover, the weak solution of the Riemann-Hilbert problem states that every subgroup of $\operatorname{GL}_n(\mathbb{C})$ that can be generated by d elements, occurs as the monodromy group of a regular singular differential equation over $\mathbb{C}(z)$ of order n with the d + 1 (potential) singularities being fixed a priori. Therefore, the above theorem holds for $k = \mathbb{C}$. This is well known and, in fact, this is exactly how the inverse problem was first solved over $\mathbb{C}(z)$ in [TT79].

Our strategy to prove the above theorem is to deduce it from the case $k = \mathbb{C}$ via a specialization argument, based on a specialization theorem from [FW24].

For $k = \mathbb{C}$ it is an immediate consequence of the Riemann-Hilbert correspondence that the differential Galois group of the family of all regular singular differential equations with singularities contained in S, is the free proalgebraic group on d generators (where |S| = d + 1). It seems natural to expect that this statement is true in full generality, i.e., for any algebraically closed field k of characteristic zero. This question was already raised in [Wib24] and the above theorem can be seen as a small step towards its resolution. Our theorem implies that every linear algebraic group over k that is a quotient of the free proalgebraic group on d generators also is a quotient of the differential Galois group of the family of all regular singular differential equations with singularities in S (Corollary 2.18).

2 The regular singular inverse problem

Throughout this article k is an algebraically closed field of characteristic zero and all fields are assumed to have characteristic zero. All rings are assumed to be commutative. For the sake of clarity, we use t for the local variable and z for the global variable. In other words, we write k((t)) for the field of formal Laurent series over k and k(z) for the field of rational functions over k. For brevity, we use the term "algebraic group (over k)" for "affine group scheme of finite type (over k)". In other words, all algebraic groups are assumed to be linear. A "proalgebraic group (over k)" is an affine group scheme (over k). A "closed subgroup" of a proalgebraic group is a closed subgroup scheme. For a morphism $R \to R'$ of rings and a scheme X over R we denote with $X_{R'}$ the scheme over R' obtained from X by base change via $R \to R'$. For an affine scheme X over a ring R, we denote with R[X] the R-algebra representing X. In particular, for an algebraic group G over k its coordinate ring is denoted with k[G].

In this section we prove the results announced in the introduction. After recalling the basics of differential Galois theory and the theory of regular singular differential equations, we study regular singular differential equations under an extension of the constants and under specialization. These preparatory results are then combined to yield the proof of the main theorem.

2.1 Differential Galois theory

In this section we recall a few key definitions and facts from differential Galois theory. For more background see any of the textbooks [PS03], [Mag94], [CH11] or [Sau16].

Let (K, ∂) be a differential field with field of constants $K^{\partial} = \{a \in K | \ \partial(a) = 0\}$ equal to k. A differential module (M, ∇) over (K, ∂) is a finite dimensional K-vector space together with an additive map $\nabla \colon M \to M$ satisfying $\nabla(am) = \partial(a)m + a\nabla(m)$ for $a \in K$ and $m \in M$. To a differential equation $\partial(y) = Ay$ with $A \in K^{n \times n}$ we can associate the differential module (K^n, ∇_A) , where $\nabla_A = \partial - A \colon K^n \to K^n$ is given by $\nabla_A(\xi) = \partial(\xi) - A\xi$.

Conversely, if (M, ∇) is a differential module and $\underline{e} = (e_1, \ldots, e_n)$ is a K-basis of M, then we can write $\nabla(\underline{e}) = \underline{e}(-A)$ for some matrix $A \in K^{n \times n}$ so that $\nabla(\underline{e}\xi) = \underline{e}\partial(\xi) + \nabla(\underline{e})\xi = \underline{e}(\partial(\xi) - A\xi)$ for all $\xi \in K^n$. Thus, up to the choice of a basis, (M, ∇) is of the form (K^n, ∇_A) . The choice of

another basis \underline{e}' , say $\underline{e} = \underline{e}'F$ with $F \in \operatorname{GL}_n(K)$, leads to another matrix $A' \in K^{n \times n}$ satisfying

$$A' = FAF^{-1} + \partial(F)F^{-1} \tag{2.1}$$

Two matrices $A, A' \in K^{n \times n}$ are said to be *Gauge equivalent* (over K) if there exists a matrix $F \in GL_n(K)$ such that (2.1) is satisfied.

Base change for differential modules works as expected: If (K', ∂') is a differential field extension of (K, ∂) and (M, ∇) is a differential module over (K, ∂) , then $M' = M \otimes_K K'$ becomes as differential module over (K', ∂') via $\nabla'(m \otimes a) = \nabla(m) \otimes a + m \otimes \partial'(a)$ for $m \in M$ and $a \in K'$. If $(M, \nabla) = (K^n, \nabla_A)$ for some $A \in K^{n \times n}$, then $(M', \nabla') = (K'^n, \nabla_A)$. I.e., in terms of differential equations, this construction simply means that a differential equation $\partial(y) = Ay$ over K is considered as a differential equation over K'.

Definition 2.1. For $A \in K^{n \times n}$, a differential K-algebra R is a Picard-Vessiot ring for $\partial(y) = Ay$ if

- (i) R is ∂ -simple (i.e., it has no ∂ -stable ideals other than $\{0\}$ and R),
- (ii) there exists a fundamental solution matrix for $\partial(y) = Ay$ in R, i.e., there exists $Y \in GL_n(R)$ such that $\partial(Y) = AY$,
- (iii) as a K-algebra R is generated by the entries of Y and the inverse of its determinant, i.e., $R = K[Y_{i,j}, \frac{1}{\det(Y)}].$

Given $A \in K^{n \times n}$, there exists a unique (up to K- ∂ -isomorphisms) Picard-Vessiot ring R/Kfor $\partial(y) = Ay$. The differential Galois group of $\partial(y) = Ay$ or of R/K is then defined as the group of differential automorphisms of R/K. More formally, we define $\operatorname{Gal}(R/K)$ as the functor, from the category of k-algebras to the category of groups, that associates to a k-algebra T the group of differential automorphisms of $R \otimes_k T$ over $K \otimes_k T$, where T is considered as a constant differential ring. The functor $\operatorname{Gal}(R/K)$ is representable by a finitely generated k-algebra, i.e., $\operatorname{Gal}(R/K)$ is an algebraic group (over k).

Definition 2.1 and the definition of the differential Galois group can be generalized to (potentially infinite) families $(\partial(y) = A_i y)_{i \in I}$ of linear differential equations over K. For example, for K = k(z) and S a subset of $\mathbb{P}^1(k)$, one can consider the differential Galois of the family of all regular singular differential equations over k(z) with singularities contained in S. In condition (ii) one requires that for every member of the family there exists a fundamental solution matrix in R and in condition (iii) one allows all entries and the inverse of the determinant of all fundamental solution matrices.

For families $\operatorname{Gal}(R/K)$ need not be an algebraic group, however, it is still a proalgebraic group (over k). Also note that the Picard-Vessiot ring for a family $(\partial(y) = A_i y)_{i \in I}$ is generated by the Picard-Vessiot rings of its members $\partial(y) = A_i y$.

2.2 Regular singular differential equations

In this section we recall the basic definitions regarding regular singular differential equations and observe that the relevant properties are preserved under an extension of the base field k. We also establish a key result (Lemma 2.9) for the proof of our main theorem (Theorem 2.16): We show that if a matrix $A \in k(t)^{n \times n}$ is Gauge equivalent over k((t)) to a matrix A' such that $tA' \in k[[t]]^{n \times n}$, then A is already Gauge equivalent over k(t) to such a matrix. More background on regular singular differential equations can be found in [PS03, Sections 5 and 6], [Sau16] or [MS16, Part I].

We first recall the basic definitions regarding formal differential modules, i.e., differential modules over the differential field $(k(t)), \frac{d}{dt})$ of formal Laurent series with coefficients in k.

A k[[t]]-lattice Λ in a finite dimensional k((t))-vector space M is a k[[t]]-submodule of the form $\Lambda = k[[t]]e_1 + \ldots + k[[t]]e_n$, where e_1, \ldots, e_n is a k((t))-basis of M. A differential module (M, ∇) over $(k((t)), \frac{d}{dt})$ is

- regular if there exists a k[[t]]-lattice Λ in M such that $\nabla(\Lambda) \subseteq \Lambda$;
- regular singular if there exists a k[[t]]-lattice Λ in M such that $\nabla(\Lambda) \subseteq \frac{1}{t}\Lambda$.

For a matrix $A \in k((t))^{n \times n}$, the differential equation $\partial(y) = Ay$ is regular (or regular singular) if the corresponding differential module $(k((t))^n, \nabla_A)$ is regular (or regular singular). Thus $\partial(y) = Ay$ is regular if and only if A is Gauge equivalent over k((t)) to a matrix $A' \in k[[t]]^{n \times n}$ and $\partial(y) = Ay$ is regular singular if and only if A is Gauge equivalent over k((t)) to a matrix $A' \in k((t))^{n \times n}$ such that $tA' \in k[[t]]^{n \times n}$.

We next discuss our main topic of interest, linear differential equations over k(z), the field of rational functions with coefficients in k. Throughout this article we consider k(z) as a differential field with respect to the standard derivation $\frac{d}{dz}$.

A differential module (M, ∇) over k(z) defines for every $p \in \mathbb{P}^1(k) = k \cup \{\infty\}$ a formal differential module (M_p, ∇_p) . In detail: For $p \in k$ we set $\widehat{k(z)}^p = k((z-p))$. Then, writing t = z - p so that $\widehat{k(z)}^p = k((t))$, we see that $(k(z), \frac{d}{dz})$ is a differential subfield of $(k((t)), \frac{d}{dt})$. We define the differential module (M_p, ∇_p) over $\widehat{k(z)}^p = k((t))$ as the base change of (M, ∇) via the inclusion $k(z) \subseteq \widehat{k(z)}^p$.

For $p = \infty$, we set $\widehat{k(z)}^p = k((z^{-1}))$. Then, writing $t = z^{-1}$ so that $\widehat{k(z)}^p = k((t))$, we see that $(k(z), \frac{d}{dz})$ is a differential subfield of $(k((t)), -t^2 \frac{d}{dt})$. Thus, via base change from $(k(z), \frac{d}{dz})$ to $(k((t)), -t^2 \frac{d}{dt})$ we obtain from (M, ∇) a differential module (M_p, ∇') over $(k((t)), -t^2 \frac{d}{dt})$. Set $\nabla_p = -\frac{1}{t^2} \nabla' \colon M_p \to M_p$. Then (M_p, ∇_p) is a differential module over $(k((t)), \frac{d}{dt})$.

A point $p \in \mathbb{P}^1(k)$ is regular for (M, ∇) if (M_p, ∇_p) is regular. Otherwise it is singular. A point $p \in \mathbb{P}^1(k)$ is regular singular for (M, ∇) if (M_p, ∇_p) is regular singular. Finally, (M, ∇) is regular singular if all points $p \in \mathbb{P}^1(k)$ are regular singular for (M, ∇) . As regular points are regular singular, this is equivalent to saying that all singular points are regular singular.

For $A \in k(z)^{n \times n}$, a point $p \in \mathbb{P}^1(k)$ is regular (or regular singular) for $\partial(y) = Ay$ if p is a regular (or regular singular) point of the corresponding differential module $(k(z)^n, \nabla_A)$. So, explicitly, a point $p \in k$ is regular (or regular singular) for $\partial(y) = Ay$ if and only if $\partial(y) =$ A(t+p)y is regular (or regular singular). On the other hand, $p = \infty$ is a regular (or regular singular) point for $\partial(y) = Ay$ if and only if $\partial(y) = -\frac{1}{t^2}A(\frac{1}{t})y$ is regular (or regular singular).

The singularities or singular points of $\partial(y) = Ay$ are the points of $\mathbb{P}^1(k)$ that are not regular for $\partial(y) = Ay$. If $p \in k$ is not a pole of one of the entries of A, then $A \in k[[t]]^{n \times n}$ for t = z - pand so p is a regular point for $\partial(y) = Ay$. Thus the singularities of $\partial(y) = Ay$ are among the poles of the entries of A and ∞ . On the other hand, a pole of an entry of A need not be a singularity of $\partial(y) = Ay$. For example, p = 0 is a regular point for $\partial(y) = \frac{1}{z}y$ (with solution y(z) = z). Indeed, for F = -z, we have $F\frac{1}{z}F^{-1} + \partial(F)F^{-1} = 0$.

The differential equation $\partial(y) = Ay$ is regular singular if all points $p \in \mathbb{P}^1(k)$ are regular singular for $\partial(y) = Ay$, i.e., if the differential module $(k(z)^n, \nabla_A)$ is regular singular.

For $k = \mathbb{C}$, the condition of being regular singular is usually defined via a growth condition on the solutions (see e.g., [Sau16, Def. 9.11] or [MS16, Part I, Def. 1.24]). However, this growth condition is equivalent to the existence of a local meromorphic gauge transformation yielding a transformed matrix that has at most a pole of order one ([Sau16, Theorem 9.17]). The existence of such a local meromorphic gauge transformation is equivalent to the existence of a local formal gauge transformation with the same effect ([PS03, Prop. 3.12 and Lemma 3.42]). Thus the above definition of a regular singular differential equation $\partial(y) = Ay$ with $A \in k(z)^{n \times n}$ is equivalent to the classical definition for $k = \mathbb{C}$.

The following three lemmas show that being regular singular is preserved under an extension of the base field k.

Lemma 2.2. Let $k \subseteq k'$ be an inclusion of algebraically closed fields and let (M, ∇) be a differential module over $(k((t)), \frac{d}{dt})$. If M is regular (or regular singular) then $M' = M \otimes_{k((t))} k'((t))$ is regular (or regular singular).

Proof. If Λ is a lattice in M with $\nabla(\Lambda) \subseteq \Lambda$ (or $\nabla(\Lambda) \subseteq \frac{1}{t}\Lambda$), then $\Lambda' = k'[[t]]\Lambda$ is a lattice in M' with $\nabla'(\Lambda') \subseteq \Lambda'$ (or $\nabla'(\Lambda') \subseteq \frac{1}{t}\Lambda'$). Thus if M is regular (or regular singular), so is M'. \Box

Lemma 2.3. Let $k \subseteq k'$ be an inclusion of algebraically closed fields and let (M, ∇) be a differential module over k(z). If M is regular singular with singular points in $S \subseteq \mathbb{P}^1(k)$, then $M' = M \otimes_{k(z)} k'(z)$ is regular singular with singular points in S.

Proof. We may assume that $(M, \nabla) = (k(z)^n, \nabla_A)$ for some $A \in k(z)^{n \times n}$. Let $p \in \mathbb{P}^1(k')$ be a singular point for M'. We have to show that p is regular singular for M' and that $p \in S$. All points in k' that are not poles of an entry of A are regular for M'. Thus p must be a pole of an entry of A or $p = \infty$. In particular, $p \in \mathbb{P}^1(k)$. By assumption, p is a regular singular point for M. So p is a regular singular point for M' by Lemma 2.2. Also, by the same lemma, p cannot be a regular point for M because then it would be a regular point for M'. Thus p is a singular point for M and therefore lies in S.

Lemma 2.4. Let $k \subseteq k'$ be an inclusion of algebraically closed fields. If $A \in k(z)^{n \times n}$ is such that $\partial(y) = Ay$ is regular singular with singularities in $S \subseteq \mathbb{P}^1(k)$ and differential Galois group G, then $\partial(y) = Ay$, considered as a differential equation over k'(z), is regular singular with singularities in $S \subseteq \mathbb{P}^1(k')$ and differential Galois group $G_{k'}$.

Proof. The statement about the singularities follows from Lemma 2.3. The statement about the differential Galois group follows from [FW24, Lemma 3.2]. \Box

Our next goal is to show that if a matrix $A \in k(t)^{n \times n}$ is Gauge equivalent over k((t)) to a matrix A' such that $tA' \in k[[t]]^{n \times n}$, then A is already Gauge equivalent over k(t) to such a matrix. For the sake of clarity of the exposition, we separate the proof into a series of lemmas.

Lemma 2.5. If $\Lambda \subseteq k((t))^n$ is a k[[t]]-lattice and E is a finite subset of $k((t))^n$, then there exists a non-zero $h \in k((t))$ such that $he \in \Lambda$ for all $e \in E$.

Proof. Let e_1, \ldots, e_n be a k((t))-basis of $k((t))^n$ such that $\Lambda = k[[t]]e_1 + \ldots + k[[t]]e_n$. Fix $e \in E$ and write $e = h_1e_1 + \ldots + h_ne_n$ with $h_1, \ldots, h_n \in k((t))$. Let $m_e \in \mathbb{N}$ be such that $t^{m_e}h_i \in k[[t]]$ for $i = 1, \ldots, n$. Then $t^{m_e}e \in \Lambda$. Let m be the maximum of all m_e 's. Then $h = t^m$ has the required property.

Lemma 2.6. Let (M, ∇) be a differential module over $(k((t)), \frac{d}{dt})$ and $\Lambda \subseteq M$ a k[[t]]-lattice such that $\nabla(\Lambda) \subseteq \frac{1}{t}\Lambda$. Then $h\Lambda \subseteq M$ is a k[[t]]-lattice with $\nabla(h\Lambda) \subseteq \frac{1}{t}h\Lambda$ for any non-zero $h \in k((t))$.

Proof. Clearly $h\Lambda$ is a k[[t]]-lattice in M. Furthermore, $\nabla(h\Lambda) \subseteq \frac{d}{dt}(h)\Lambda + h\nabla(\Lambda) \subseteq \frac{d}{dt}(h)\Lambda + \frac{1}{t}h\Lambda$. So it suffices to show that $\frac{d}{dt}(h)\Lambda \subseteq \frac{1}{t}h\Lambda$. But $\frac{t\frac{d}{dt}(h)}{h} \in k[[t]]$ because the order of $t\frac{d}{dt}(h)$ is the order of h unless $h \in k$. Thus $\frac{d}{dt}(h)\Lambda = \frac{h}{t}\frac{t\frac{d}{dt}(h)}{h}\Lambda \subseteq \frac{h}{t}\Lambda = \frac{1}{t}h\Lambda$.

As usual, we denote with $k[t]_{(t)}$ the localization of the ring k[t] at the prime ideal (t), i.e., $k[t]_{(t)} = \left\{ \frac{h_1}{h_2} | \ h_1, h_2 \in k[t], \ t \text{ does not divide } h_2 \right\}.$

Lemma 2.7. Let e_1, \ldots, e_n be a k(t)-basis of $k(t)^n$. For $\Lambda = k[[t]]e_1 + \ldots + k[[t]]e_n \subseteq k((t))^n$ we have $\Lambda \cap k(t)^n = k[t]_{(t)}e_1 + \ldots + k[t]_{(t)}e_n$. Proof. The inclusion $k[t]_{(t)}e_1 + \ldots + k[t]_{(t)}e_n \subseteq \Lambda \cap k(t)^n$ follows from $k[t]_{(t)} \subseteq k[[t]]$. For the reverse inclusion, consider $m \in \Lambda \cap k(t)^n$. As e_1, \ldots, e_n is a k((t))-basis of $k((t))^n$, we have $m = a_1e_1 + \ldots + a_ne_n$ for uniquely determined $a_1, \ldots, a_n \in k((t))$. As $m \in \Lambda$, it follows that $a_i \in k[[t]]$ and as $m \in k(t)^n$ is follows that $a_i \in k(t)$. Thus $a_i \in k[[t]] \cap k(t) = k[t]_{(t)}$ for $i = 1, \ldots, n$ and so $m \in k[t]_{(t)}e_1 + \ldots + k[t]_{(t)}e_n$.

The following lemma is a variant of Nakayama's Lemma. For a proof see e.g., [Lan02, Chapter X, Theorem 4.4].

Lemma 2.8. Let R be a local ring with maximal ideal \mathfrak{m} and residue field $k = R/\mathfrak{m}$. If M is a free R-module of finite rank, then a tuple of elements from M is an R-basis of M if and only if their images are a k-basis of $M/\mathfrak{m}M$.

Recall that the condition of being regular singular is characterized via local formal gauge transformations. The following lemma shows that in fact it is enough to consider rational gauge transformations. We believe this lemma is well-known but we could not find a suitable reference. For statements of a similar spirit in the analytic context see [Sau16, Lemma 12.1] and [PS03, Lemma 3.42].

Lemma 2.9. Let $A \in k(t)^{n \times n} \subseteq k((t))^{n \times n}$ be such $\partial(y) = Ay$ is regular singular. Then there exists a matrix $F \in \operatorname{GL}_n(k(t))$ such that $A' = FAF^{-1} + \partial(F)F^{-1}$ satisfies $tA' \in k[t]_{(t)}^{n \times n}$. In other words, if A is Gauge equivalent over k((t)) to a matrix A' satisfying $tA' \in k[[t]]^{n \times n}$, then A is already Gauge equivalent over k(t) to such a matrix.

Proof. As $\partial(y) = Ay$ is regular singular, there exists a k[[t]]-lattice Λ in $k((t))^n$ such that $\nabla_A(\Lambda) \subseteq \frac{1}{t}\Lambda$, where $\nabla_A = \frac{d}{dt} - A$.

Let $\underline{e} = (e_1, \ldots, e_n)$ be the standard basis of $k(t)^n$. By Lemma 2.5 there exists a non-zero $h \in k((t))$ such that $he_i \in \Lambda$ for $i = 1, \ldots, n$. Then $\frac{1}{h}\Lambda$ is a k[[t]]-lattice in $k((t)))^n$ with $\nabla_A(\frac{1}{h}\Lambda) \subseteq \frac{1}{t}\frac{1}{h}\Lambda$ (Lemma 2.6) and $e_1, \ldots, e_n \in \frac{1}{h}\Lambda$. Replacing Λ with $\frac{1}{h}\Lambda$, we can thus assume that $e_1, \ldots, e_n \in \Lambda$.

Let Λ' be the k[[t]]-submodule of Λ generated by all $(t\nabla_A)^i(e_j)$, where $i \geq 0$, and $j \in \{1, \ldots, n\}$. Note that, by construction, $t\nabla_A(\Lambda') \subseteq \Lambda'$.

Recall (e.g. from [Lan02, Chapter III, Theorem 7.1]) that a submodule of a free module over a principal ideal domain is free of rank bounded by the rank of the ambient module. As k[[t]]is a principal ideal domain and Λ is a free k[[t]]-module of rank n, it follows that Λ' is a free k[[t]]-module of rank at most n. As Λ' contains the standard basis \underline{e} , we see that in fact Λ' is free of rank n. So $\Lambda'/t\Lambda'$ is a k-vector space of dimension n. Since the $(t\nabla_A)^i(e_j)$'s generate Λ' as a k[[t]]-module, their images generate $\Lambda'/t\Lambda'$ as a k-vector space. We can select n of these generators, say e'_1, \ldots, e'_n , such that their images are k-basis of $\Lambda'/t\Lambda'$. It follows from Lemma 2.8 that e'_1, \ldots, e'_n are a k[[t]]-basis of L'. In particular, e'_1, \ldots, e'_n are k[[t]]-linearly independent and they lie in $k(t)^n$. Thus e'_1, \ldots, e'_n are a k(t)-basis of $k(t)^n$.

As the $(t\nabla_A)^i(e_j)$'s generate Λ' as a k[[t]]-module, Lemma 2.8 shows that we can select n of these generators, say e'_1, \ldots, e'_n , such that they form a k[[t]]-basis of Λ' . In particular, e'_1, \ldots, e'_n are k[[t]]-linearly independent and they lie in $k(t)^n$. Thus $\underline{e} = (e'_1, \ldots, e'_n)$ is a k(t)-basis of $k(t)^n$.

Lemma 2.7 applied to e'_1, \ldots, e'_n and $\Lambda' = k[[t]]e'_1 + \ldots + k[[t]]e'_n$ shows that $\Lambda' \cap k(t)^n = k[t]_{(t)}e'_1 + \ldots + k[t]_{(t)}e'_n$. As Λ' and $k(t)^n$ are stable under $t\nabla_A$, it follows that also $k[t]_{(t)}e'_1 + \ldots + k[t]_{(t)}e'_n$ is stable under $t\nabla_A$. That is, if we write $\nabla_A(\underline{e}') = \underline{e'}(-A')$, then $tA' \in k[t]_{(t)}^{n \times n}$. For $F \in \operatorname{GL}_n(k(t))$ with $\underline{e} = \underline{e'}F$, we have $A' = FAF^{-1} + \partial(F)F^{-1}$ as desired.

2.3 Generating algebraic groups

In this short section we consider algebraic groups generated by finitely many elements and how this property behaves under base change. Let G be an algebraic group over k and $g_1, \ldots, g_d \in G(k)$. The intersection of all closed subgroups H of G such that $g_1, \ldots, g_d \in H(k)$, is a closed subgroup of G that also satisfies this property. Thus there exists a smallest closed subgroup of G whose k-points contain g_1, \ldots, g_d . If this closed subgroup agrees with G, we say that g_1, \ldots, g_d generated G (as an algebraic group).

Let Γ be the (abstract) subgroup of G(k) generated by g_1, \ldots, g_d . As the Zariski closure of Γ is a subgroup of G(k) ([Spr98, Lemma 2.2.4]), we see that g_1, \ldots, g_d generate G if and only if Γ is Zariski dense in G. This means that g_1, \ldots, g_d generate G if and only if $f(\gamma) = 0$ for all $\gamma \in \Gamma$ implies f = 0 for $f \in k[G]$.

Lemma 2.10. Let $k \subseteq k'$ be an inclusion of algebraically closed fields, G an algebraic group over k and $g_1, \ldots, g_d \in G(k)$. Then g_1, \ldots, g_d generate G if and only if g_1, \ldots, g_d generate $G_{k'}$.

Proof. Let Γ be the (abstract) subgroup of $G(k) \subseteq G(k') = G_{k'}(k')$ generated by g_1, \ldots, g_d . First assume that g_1, \ldots, g_d generate G. We have to show that $f'(\gamma) = 0$ for all $\gamma \in \Gamma$ implies f' = 0 for $f' \in k'[G_{k'}] = k[G] \otimes_k k'$. Let $(\lambda_i)_{i \in I}$ be k-basis of k'. Then f' can be written as $f' = \sum f_i \otimes \lambda_i$ with $f_i \in k[G]$. For $\gamma \in \Gamma$, we have $0 = f'(\gamma) = \sum f_i(\gamma)\lambda_i$. As the λ_i are k-linearly independent and $f_i(\gamma_i) \in k$, it follows that $f_i(\gamma) = 0$ for all $\gamma \in \Gamma$ and $i \in I$. Because g_1, \ldots, g_d generate G, this implies that $f_i = 0$. So f' = 0 as desired.

Conversely, assume that g_1, \ldots, g_d generate $G_{k'}$. Then an $f' \in k'[G_{k'}] = k[G] \otimes_k k'$ that vanishes on all of Γ must be zero. In particular, this holds for $f \in k[G]$ and therefore g_1, \ldots, g_d generate G.

2.4 Specializations

The proof strategy for our main result is to deduce it from the already known special case $k = \mathbb{C}$ via a specialization argument. In this section we recall the specialization theorem from [FW24] required for this approach. We also show that the property of being regular singular is preserved under specialization in an appropriate sense.

We begin by recalling some terminology from [FW24]. Let \mathcal{B} be a k-algebra (considered as a constant differential ring). The specializations we will be considering are morphisms $c \in$ $\operatorname{Hom}_k(\mathcal{B}, k)$, i.e., $c \colon \mathcal{B} \to k$ is a morphism of k-algebras. We consider $\mathcal{B}[z]$ as a differential ring with respect to the derivation $\frac{d}{dz}$. For a monic polynomial $f \in \mathcal{B}[z]$ a specialization $c \colon \mathcal{B} \to k$ extends to a morphism $c \colon \mathcal{B}[z]_f \to k(z)$ of differential k-algebras via c(z) = z. For a matrix $\mathcal{A} \in$ $\mathcal{B}[z]_f^{n \times n}$ we denote with $A^c \in k(z)^{n \times n}$ the matrix obtained from \mathcal{A} by applying $c \colon \mathcal{B}[z]_f \to k(z)$ to the entries of \mathcal{A} . Similarly, for $a \in \mathcal{B}$ we will write a^c rather than c(a).

Assuming that \mathcal{B} is an integral domain, so that we can consider the algebraic closure K of the field of fractions of \mathcal{B} , the specialization problem is about comparing the differential Galois group of the generic equation $\partial(y) = \mathcal{A}y$ (over K(z)) with the differential Galois group of the specialized equations $\partial(y) = A^c y$ (over k(z)). To capture the idea of specializing differential Galois groups, we need to consider group schemes over \mathcal{B} and we also need some more terminology.

Let \mathcal{B} be a k-algebra and let \mathcal{Q} be a differential \mathcal{B} -algebra. (The most relevant case for us is $\mathcal{Q} = \mathcal{B}[z]_{f}$.) For a differential \mathcal{Q} -algebra \mathcal{R} , we define $\underline{\operatorname{Aut}}(\mathcal{R}/\mathcal{Q})$ as the functor form the category of \mathcal{B} -algebras to the category of groups given by $\underline{\operatorname{Aut}}(\mathcal{R}/\mathcal{Q})(\mathcal{T}) = \operatorname{Aut}^{\partial}(\mathcal{R} \otimes_{\mathcal{B}} \mathcal{T}/\mathcal{Q} \otimes_{\mathcal{B}} \mathcal{T})$, where \mathcal{T} is considered as a constant differential ring. On morphisms, $\underline{\operatorname{Aut}}(\mathcal{R}/\mathcal{Q})$ is defined by base change. An action of an affine group scheme \mathcal{G} over \mathcal{B} on \mathcal{R}/\mathcal{Q} is a morphism of group functors $\mathcal{G} \to \underline{\operatorname{Aut}}(\mathcal{R}/\mathcal{Q})$.

Let $\mathcal{B}[\mathcal{G}]$ denote the coordinate ring of \mathcal{G} , i.e., the \mathcal{B} -algebra representing \mathcal{G} . By [FW24, Lemma 3.3], to specify an action of \mathcal{G} on \mathcal{R}/\mathcal{Q} is equivalent to specifying a morphism (the

coaction) $\rho \colon \mathcal{R} \to \mathcal{R} \otimes_{\mathcal{B}} \mathcal{B}[\mathcal{G}]$ of \mathcal{Q} -algebras such that



commute, where Δ and ε denote the comultiplication and counit of the Hopf-algebra $\mathcal{B}[\mathcal{G}]$.

Definition 2.11. Assume, as above, that \mathcal{G} acts \mathcal{R}/\mathcal{Q} . Then \mathcal{R}/\mathcal{Q} is a differential \mathcal{G} -torsor if the morphism $\mathcal{R} \otimes_{\mathcal{Q}} \mathcal{R} \to \mathcal{R} \otimes_{\mathcal{B}} \mathcal{B}[\mathcal{G}], a \otimes b \mapsto (a \otimes 1) \cdot \rho(b)$ is an isomorphism. Moreover, for $\mathcal{A} \in \mathcal{Q}^{n \times n}$, if there exists a matrix $\mathcal{Y} \in \operatorname{GL}_n(\mathcal{R})$ such that

- $\partial(\mathcal{Y}) = \mathcal{AY},$
- $\mathcal{R} = \mathcal{Q}\left[\mathcal{Y}_{i,j}, \frac{1}{\det(\mathcal{Y})}\right]$, and
- for every \mathcal{B} -algebra \mathcal{T} and every $g \in \mathcal{G}(\mathcal{T})$, there exists a matrix $M_g \in \mathrm{GL}_n(\mathcal{T})$ such that $g(\mathcal{Y} \otimes 1) = (\mathcal{Y} \otimes 1)(1 \otimes M_g)$,

then \mathcal{R}/\mathcal{Q} is a differential \mathcal{G} -torsor for $\partial(y) = \mathcal{A}y$.

The definition of a differential torsor can be interpreted geometrically: With $\mathcal{Z} = \operatorname{Spec}(\mathcal{R})$, the coaction $\rho \colon \mathcal{R} \to \mathcal{R} \otimes_{\mathcal{B}} \mathcal{B}[\mathcal{G}] = \mathcal{R} \otimes_{\mathcal{Q}} \mathcal{Q} \otimes_{\mathcal{B}} \mathcal{B}[\mathcal{G}]$ induces a group action $\mathcal{Z} \times_{\mathcal{Q}} \mathcal{G}_{\mathcal{Q}} \to \mathcal{Z}$ (in the category of \mathcal{Q} -schemes) and \mathcal{R}/\mathcal{Q} is a differential \mathcal{G} -torsor if and only if $\mathcal{Z} \times_{\mathcal{Q}} \mathcal{G}_{\mathcal{Q}} \to \mathcal{Z} \times_{\mathcal{Q}} \mathcal{Z}$ is an isomorphism.

Definition 2.11 mimics and generalizes the definition of a Picard-Vessiot ring. In particular, if R/K is a Picard-Vessiot ring for $\partial(y) = Ay$ (Definition 2.1), then R/K is a differential $\operatorname{Gal}(R/K)$ -torsor for $\partial(y) = Ay$.

The following specialization theorem is the key to our specialization argument.

Theorem 2.12 ([FW24], Theorem 4.26). Assume that

- \mathcal{B} is a finitely generated k-algebra and an integral domain,
- K is the algebraic closure of the field of fractions of \mathcal{B} ,
- \mathcal{G} is an affine group scheme of finite type over \mathcal{B} ,
- $f \in \mathcal{B}[z]$ is a monic polynomial,
- $\mathcal{A} \in \mathcal{B}[z]_f^{n \times n}$,
- $\mathcal{R}/\mathcal{B}[z]_f$ is a differential \mathcal{G} -torsor for $\partial(y) = \mathcal{A}y$ such that \mathcal{R} is flat over $\mathcal{B}[z]_f$,
- $\mathcal{R} \otimes_{\mathcal{B}[z]_f} K(z)$ is ∂ -simple.

Then there exists a $c \in \operatorname{Hom}_k(\mathcal{B}, k)$ such that $R^c = \mathcal{R} \otimes_{\mathcal{B}[z]_f} k(z)$ is a Picard-Vessiot ring for $\partial(y) = A^c y$ with differential Galois group \mathcal{G}_k , where the involved base change is $c \colon \mathcal{B} \to k$.

Note that the assumptions in Theorem 2.12 imply that $R^{\text{gen}} = \mathcal{R} \otimes_{\mathcal{B}[z]_f} K(z)$ is a Picard-Vessiot ring for $\partial(y) = \mathcal{A}y$ (over K(z)) with differential Galois group \mathcal{G}_K . The following lemma shows that, starting from an inclusion $k \subseteq k'$ of algebraically closed fields and a differential equation $\partial(y) = Ay$ over k'(z), the technical conditions of Theorem 2.12 can always be achieved.

Lemma 2.13 ([FW24], Lemma 3.11). Let $k \subseteq k'$ be an inclusion of algebraically closed fields and let $A \in k'(z)^{n \times n}$. Then there exist

- a finitely generated k-subalgebra \mathcal{B} of k',
- a monic polynomial $f \in \mathcal{B}[z]$,
- an affine group scheme \mathcal{G} of finite type over \mathcal{B} , and
- a differential \mathcal{G} -torsor $\mathcal{R}/\mathcal{B}[z]_f$,

such that

- $A \in \mathcal{B}[z]_f^{n \times n}$,
- \mathcal{R} is flat over $\mathcal{B}[z]_f$ and $\mathcal{R}/\mathcal{B}[z]_f$ is a differential \mathcal{G} -torsor for $\partial(y) = Ay$
- $\mathcal{R} \otimes_{\mathcal{B}[z]_{\ell}} K(z)$ is ∂ -simple, where K the algebraic closure of the field of fractions of \mathcal{B} .

Moreover, for a fixed finitely generated k-algebra \mathcal{B}_0 and monic $f_0 \in \mathcal{B}_0[z]$, \mathcal{B} and f can be chosen such that $\mathcal{B}_0 \subseteq \mathcal{B}$ and f_0 divides f. Furthermore, if there exists an algebraic group G over k such that the differential Galois group of $\partial(y) = Ay$ (over k'(z)) is of the form $G_{k'}$, then \mathcal{G} can be chosen to be $G_{\mathcal{B}}$.

We next study regular singular differential equations under specialization, first locally, then globally.

Lemma 2.14. Let $k \subseteq k'$ be an inclusion of algebraically closed fields and let $A \in k'(t)^{n \times n} \subseteq k((t))^{n \times n}$ be such that $\partial(y) = Ay$ is regular singular. Then there exists a finitely generated k-subalgebra \mathcal{B} of k' and a monic polynomial $f \in \mathcal{B}[t]$ such that $A \in \mathcal{B}[t]_f^{n \times n}$ and $\partial(y) = A^c y$ is regular singular for every $c \in \operatorname{Hom}_k(\mathcal{B}, k)$.

Proof. By Lemma 2.9, there exists a matrix $F \in \operatorname{GL}_n(k'(t))$ such that the matrix $t(FAF^{-1} + \partial(F)F^{-1})$ has entries in $k'[t]_{(t)}$. We may choose a monic polynomial $f \in k'[t]$ such that fA and fF have entries in k'[t]. Let \mathcal{B} be the k-subalgebra of k' generated by the coefficients of f and all coefficients of all entries of fA and fF. Then $A, F \in \mathcal{B}[t]_f^{n \times n}$. As $\det(F) \in \mathcal{B}[t]_f$ is non-zero, we may write $\det(f) = \frac{h}{f^m}$ for some non-zero $h \in \mathcal{B}[t]$. Adding the inverse of the leading coefficient of h to \mathcal{B} and replacing f with the product of f with h and the inverse of the leading coefficient of h, we may assume that $A, F, F^{-1} \in \mathcal{B}[t]_f^{n \times n}$.

As $t(FAF^{-1} + \partial(F)F^{-1})$ has entries in $k'[t]_{(t)}$, there exist non-zero elements $a_1, \ldots, a_m \in k'$ and a matrix $B \in k'[t]$ such that

$$(t_1 - a_1) \dots (t - a_m)t(FAF^{-1} + \partial(F)F^{-1}) = B.$$
(2.2)

Replacing \mathcal{B} with the k-subalgebra of k' generated by \mathcal{B} , the coefficients of the entries of Band $a_1, \ldots, a_m, a_1^{-1}, \ldots, a_m^{-1}$, identity (2.2) becomes an identity in $\mathcal{B}[t]_f$. Thus, for any $c \in$ $\operatorname{Hom}_k(\mathcal{B}, k)$, we can apply the morphism $c: \mathcal{B}[t]_f \to k(t)$ of differential k-algebras to (2.2) to obtain

$$(t_1 - a_1^c) \dots (t - a_m^c) t(F^c A^c (F^c)^{-1} + \partial (F^c) (F^c)^{-1}) = B^c,$$

an identity in k(t). As $B^c \in k[t]^{n \times n}$ and a_1^c, \ldots, a_m^c are non-zero, this shows that $\partial(y) = A^c y$ is regular singular.

Let $A \in k(z)^{n \times n}$. Recall that a pole of an entry of A need not be a singularity of $\partial(y) = Ay$. Such points are sometimes called *apparent singularities*. Note, however, that they are not singularities of $\partial(y) = Ay$ according to our terminology. To have an unambiguous notation we make the following definition. A point $p \in \mathbb{P}^1(k)$ is a *pole of* $\partial(y) = Ay$ if

• $p \in k$ and p is a pole of one of the entries of A or

• $p = \infty$ and z = 0 is a pole of one of the entries of $-\frac{1}{z^2}A(\frac{1}{z})$.

Thus the singularities of $\partial(y) = Ay$ are contained in the poles of $\partial(y) = Ay$.

Lemma 2.15. Let $k \subseteq k'$ be an inclusion of algebraically closed fields and let S be a finite subset of $\mathbb{P}^1(k) \subseteq \mathbb{P}^1(k')$. Furthermore, let $A \in k'(z)^{n \times n}$ be such that $\partial(y) = Ay$ is regular singular with poles contained in S. Then there exists a finitely generated k-subalgebra $\mathcal{B} \subseteq k'$, and a monic polynomial $f \in \mathcal{B}[z]$ such that $A \in \mathcal{B}[z]_f^{n \times n}$ and for every $c \in \operatorname{Hom}_k(\mathcal{B}, k)$ the differential equation $\partial(y) = A^c y$ is regular singular with poles contained in S.

Proof. As the poles of $\partial(y) = Ay$ are contained in S, there exist (not necessarily distinct) $a_1, \ldots, a_m \in S$ such that $B = (z - a_1) \ldots (z_1 - a_m)A$ has entries in k'[z]. Let \mathcal{B}_0 be the k-subalgebra of k' generated by the coefficients of all entries of B and set $f_0 = (z - a_1) \ldots (z_1 - a_m)$. Then $A \in \mathcal{B}_0[z]_{f_0}^{n \times n}$ and for any $c \in \operatorname{Hom}_k(\mathcal{B}_0, k)$ the poles in k of the differential equation $\partial(y) = A^c y$ are contained in $\{a_1, \ldots, a_m\} \subseteq S$.

For a non-zero rational function $h \in \mathcal{B}_0[z]_{f_0} \subseteq k'(z)$, when written as $h = \frac{h_1}{f_0^c}$, with $h_1 \in \mathcal{B}_0[z]$, the rational function $-\frac{1}{z^2}h(\frac{1}{z}) \in k'(z)$ has a pole at z = 0 if and only if $\deg(h_1) - em + 2 > 0$. Thus, if $-\frac{1}{z^2}h(\frac{1}{z})$ does not have a pole at z = 0, for any specialization $c \in \operatorname{Hom}_k(\mathcal{B}_0, k)$, the rational function $-\frac{1}{z^2}h^c(\frac{1}{z})$ does not have a pole at z = 0. So if $p = \infty$ is not a pole of $\partial(y) = Ay$, then $p = \infty$ is also not a pole of $\partial(y) = A^c y$ for any $c \in \operatorname{Hom}_k(\mathcal{B}_0, k)$. In summary, for any $c \in \operatorname{Hom}_k(\mathcal{B}_0, k)$, the poles of $\partial(y) = A^c y$ are contained in S.

For i = 1, ..., m, the differential equation $\partial(y) = A_i y$, where $A_i = A(t + a_i) \in k'(t)^{n \times n}$ is regular singular. Thus, by Lemma 2.14, there exists a finitely generated k-subalgebra \mathcal{B}_i of k' and a monic polynomial $f_i \in \mathcal{B}_i[t]$ such that $A_i \in \mathcal{B}_i[t]_{f_i}^{n \times n}$ and $\partial(y) = A_i^c y$ is regular singular for all $c \in \operatorname{Hom}_k(\mathcal{B}_i, k)$. Enlarging \mathcal{B}_i if necessary, we may assume that $\mathcal{B}_0 \subseteq \mathcal{B}_i$. Then $A_i^c = A(t + a_i)^c = A^c(t + a_i)$ for every $c \in \operatorname{Hom}_k(\mathcal{B}_i, k)$. As $\partial(y) = A_i^c y$ is regular singular, this shows that $p = a_i$ is a regular singular point for $\partial(y) = A^c y$.

For the point $p = \infty$ we proceed similarly. As $\partial(y) = -\frac{1}{t^2}A(\frac{1}{t})y$ is regular singular, there exists, by Lemma 2.14, a finitely generated k-subalgebra \mathcal{B}_{∞} of k' and a monic polynomial $f_{\infty} \in \mathcal{B}_{\infty}[t]$ such that $-\frac{1}{t^2}A(\frac{1}{t}) \in \mathcal{B}_{\infty}[t]_{f_{\infty}}^{n \times n}$ and $\partial(y) = -\frac{1}{t^2}A(\frac{1}{t})^c y$ is regular singular for every $c \in \operatorname{Hom}_k(\mathcal{B}_{\infty}, k)$. We may assume that $\mathcal{B}_0 \subseteq \mathcal{B}_{\infty}$. Then $-\frac{1}{t^2}A(\frac{1}{t})^c = -\frac{1}{t^2}A^c(\frac{1}{t})$ and we see that $p = \infty$ is a regular singular point for $\partial(y) = A^c y$ for all $c \in \operatorname{Hom}_k(\mathcal{B}_{\infty}, k)$.

Let \mathcal{B} be the k-subalgebra of k' generated by $\mathcal{B}_1, \ldots, \mathcal{B}_m, \mathcal{B}_\infty$ and set $f = f_0 f_1 \ldots f_m f_\infty \in \mathcal{B}[z]$. We claim that \mathcal{B} and f have the desired properties.

Clearly $A \in \mathcal{B}[z]_f^{n \times n}$. Let $c \in \operatorname{Hom}_k(\mathcal{B}, k)$. As $\mathcal{B}_0 \subseteq \mathcal{B}$ (so that c restricts to a morphism $\mathcal{B}_0 \to k$), it follows from the first two paragraphs, that the poles of $\partial(y) = Ay$ are contained in S. Similarly, as c restricts to morphisms on $\mathcal{B}_1, \ldots, \mathcal{B}_m, \mathcal{B}_\infty$, it follows from the above that a_1, \ldots, a_m, ∞ are regular singular points for $\partial(y) = A^c y$. As the poles of $\partial(y) = A^c y$ are contained in $\{a_1, \ldots, a_m, \infty\}$, and all these points are regular singular, it follows that $\partial(y) = A^c y$ is regular singular.

2.5 Proof of the main result

We are now prepared to prove our main theorem.

Theorem 2.16. Let G be a closed subgroup of $\operatorname{GL}_{n,k}$ that can be generated by d elements and let S be a subset of $\mathbb{P}^1(k)$ with d+1 elements. Then there exists a matrix $A \in k(z)^{n \times n}$ such that the differential equation $\partial(y) = Ay$ is regular singular with singularities contained in S and has differential Galois group G.

Proof. We roughly follow the short solution to the inverse problem from [FW24, Section 5.2]. The algebraic group G (together with the closed embedding into $\operatorname{GL}_{n,k}$) descends to a finitely generated field k_0 , i.e., there exist a subfield k_0 of k, finitely generated over \mathbb{Q} , and a closed subgroup G_0 of $\operatorname{GL}_{n,k_0}$ such that $(G_0)_k = G$ (as closed subgroups of $\operatorname{GL}_{n,k}$).

Let $g_1, \ldots, g_d \in G(k)$ generate G. Enlarging k_0 if necessary, we can assume that $g_1, \ldots, g_d \in G_0(k_0) \subseteq G_0(k) = G(k)$ and $S \subseteq \mathbb{P}^1(k_0)$.

Let $k_1 \subseteq k$ be the algebraic closure of k_0 and set $G_1 = (G_0)_{k_1}$. Then G_1 is a closed subgroup of $\operatorname{GL}_{n,k_1}$. As k_1 is countable, there exists an embedding $k_1 \hookrightarrow \mathbb{C}$ (that we fix).

Because $g_1, \ldots, g_d \in G_0(k_0) \subseteq G_0(k_1) = G_1(k_1)$ generate $G = (G_1)_k$, it follows from Lemma 2.10 that g_1, \ldots, g_d generate G_1 . The same lemma, but applied in the reverse direction, shows that $(G_1)_{\mathbb{C}}$ is generated by g_1, \ldots, g_d . In particular, $(G_1)_{\mathbb{C}}$ is a closed subgroup of $\operatorname{GL}_{n,\mathbb{C}}$ that can be generated by d elements.

As noted in the introduction, Theorem 2.16 holds for $k = \mathbb{C}$ as a consequence of the (weak) solution of the Riemann-Hilbert problem ([PS03, Theorem 5.15]) and Schlesinger's density theorem ([PS03, Theorem 5.8]). In fact, the (weak) solution of the Riemann-Hilbert problem yields a slightly stronger statement than Theorem 2.16 in the case $k = \mathbb{C}$: The linear differential equation can be chosen such that its poles are contained in S. (See e.g., [MS16, Part I, Theorem 4.49] or [Sau16, Theorem 12.4].) Thus, there exists a matrix $A \in \mathbb{C}(z)^{n \times n}$ such that $\partial(y) = Ay$ is regular singular with poles contained in S and differential Galois group $(G_1)_{\mathbb{C}}$.

Applying Lemma 2.15 to the inclusion $k_1 \subseteq \mathbb{C}$ and the differential equation $\partial(y) = Ay$ yields a finitely generated k_1 -subalgebra \mathcal{B}_0 of \mathbb{C} and a monic $f_0 \in \mathcal{B}_0[z]$ such that $A \in \mathcal{B}_0[z]_{f_0}^{n \times n}$ and $\partial(y) = A^c y$ is regular singular with singularities contained in S for all $c \in \operatorname{Hom}_{k_1}(\mathcal{B}_0, k_1)$. By Lemma 2.13 there exists a finitely generated k_1 -subalgebra \mathcal{B} of \mathbb{C} containing \mathcal{B}_0 , a monic polynomial $f \in \mathcal{B}[z]$ such that f_0 divides f, a differential $(G_1)_{\mathcal{B}}$ -torsor $\mathcal{R}/\mathcal{B}[z]_f$ for $\partial(y) = Ay$ such that \mathcal{R} is flat over $\mathcal{B}[z]_f$ and $\mathcal{R} \otimes_{\mathcal{B}[z]_f} K(z)$ is ∂ -simple, where K is the algebraic closure of the field of fractions of \mathcal{B} . By Theorem 2.12, there exists a $c \in \operatorname{Hom}_{k_1}(\mathcal{B}, k_1)$ such that $R^c = \mathcal{R} \otimes_{\mathcal{B}[z]_f} k_1(z)$ is a Picard-Vessiot ring for $\partial(y) = A^c y$ (over $k_1(z)$) with differential Galois group $((G_1)_{\mathcal{B}})_{k_1} = G_1$. By construction of \mathcal{B}_0 and f_0 , and using that $A \in \mathcal{B}_0[z]_{f_0}^{n \times n} \subseteq \mathcal{B}[z]_f^{n \times n}$ we see that $\partial(y) = A^c y$ is regular singular with singularities contained in S.

Finally, by Lemma 2.4, the equation $\partial(y) = A^c y$, considered as a differential equation over k(z), has the required properties, i.e., $\partial(y) = A^c y$ (over k(z)) is regular singular with singularities contained in S and has differential Galois group $(G_1)_k = G$.

Corollary 2.17. Any algebraic group over k is the differential Galois group of a regular singular differential equation $\partial(y) = Ay$ over k(z).

Proof. As any (affine) algebraic group is isomorphic to a closed subgroup of some general linear group ([Mil17, Cor. 4.10]) and every algebraic group (in characteristic zero) is finitely generated ([PS03], Lemma 5.13), this follows from Theorem 2.16. \Box

2.6 An interpretation in terms of proalgebraic groups

In this final section we offer an interpretation of our main result (Theorem 2.16) in terms of proalgebraic groups.

Recall that the *free proalgebraic group* Γ_d on d generators (over k) is characterized by the following universal property of the map $\iota: X \to \Gamma_d(k)$ from a d-element set X into the k-points of Γ_d : For every proalgebraic group G and every map $\varphi: X \to G(k)$ there exists a unique morphism $\phi: \Gamma_d \to G$ of proalgebraic groups such that



commutes. The proalgebraic group Γ_d can be constructed as the fundamental group of the neutral tannakian category of all (finite dimensional, k-linear) representations of the (abstract)

free group F_d on d generators. In other words, Γ_d is the proalgebraic completion (or proalgebraic hull) of F_d . (See [Wib20] for more background on free proalgebraic groups.)

Let S be a subset of $\mathbb{P}^1(k)$ with d + 1 elements and let Γ_S be the differential Galois group of the family of all regular singular differential equations over k(z) with singularities contained in S. For $k = \mathbb{C}$, it is an immediate corollary of the Riemann-Hilbert correspondence ([PS03, Theorem 6.15]) that Γ_S is isomorphic to Γ_d . It therefore seems natural to expect that Γ_S is isomorphic to Γ_d in general. This question was already raised in [Wib24, Section 4.1]. While we are far from being able to show that indeed Γ_S is isomorphic to Γ_d , Theorem 2.16 can be seen as a small step into the right direction. It implies that every algebraic quotient of Γ_d is also a quotient of Γ_S .

Corollary 2.18. Let G be an algebraic group over k. If G is a quotient of Γ_d , then G is also a quotient of Γ_S .

Proof. If G is a quotient of Γ_d , then G can be generated by d elements ([Wib20, Lemma 2.16]). After embedding G into some $\operatorname{GL}_{n,k}$, we can apply Theorem 2.16 to find a regular singular differential equation $\partial(y) = Ay$ with differential Galois group G. The Picard-Vessiot ring R of $\partial(y) = Ay$ is canonically embedded in the Picard-Vessiot ring R_S of the family of all regular singular differential equations with singularities in S. By the second fundamental theorem of differential Galois theory (see e.g., [AMT09, Theroem 2.11]), the differential Galois group of R is a quotient of the differential Galois group of R_S , i.e., G is a quotient of Γ_S as desired.

References

[AMT09]	Katsutoshi Amano, Akira Masuoka, and Mitsuhiro Takeuchi. "Hopf algebraic approach to Picard-Vessiot theory". In: vol. 6. Handb. Algebr. Elsevier/North-Holland, Amsterdam, 2009, pp. 127–171. ISBN: 978-0-444-53257-2. DOI: 10.1016/S1570-7954(08)00204-0. URL: https://doi.org/10.1016/S1570-7954(08)00204-0.
[Bac+18]	Annette Bachmayr, David Harbater, Julia Hartmann, and Michael Wibmer. "Differ- ential embedding problems over complex function fields". In: <i>Doc. Math.</i> 23 (2018), pp. 241–291. ISSN: 1431-0635.
[Bac+21a]	Annette Bachmayr, David Harbater, Julia Hartmann, and Florian Pop. "Large fields in differential Galois theory". In: J. Inst. Math. Jussieu 20.6 (2021), pp. 1931–1946. ISSN: 1474-7480,1475-3030. DOI: 10.1017/S1474748020000018. URL: https://doi.org/10.1017/S
[Bac+21b]	Annette Bachmayr, David Harbater, Julia Hartmann, and Michael Wibmer. "Free differential Galois groups". In: <i>Trans. Amer. Math. Soc.</i> 374.6 (2021), pp. 4293–4308. ISSN: 0002-9947,1088-6850. DOI: 10.1090/tran/8352. URL: https://doi.org/10.1090/tran/8352
[Bac+21c]	Annette Bachmayr, David Harbater, Julia Hartmann, and Michael Wibmer. "The differential Galois group of the rational function field". In: <i>Adv. Math.</i> 381 (2021), Paper No. 107605, 27 pp. ISSN: 0001-8708. DOI: 10.1016/j.aim.2021.107605. URL: https://doi.org/10.1016/j.aim.2021.107605.
[CH11]	Teresa Crespo and Zbigniew Hajto. Algebraic groups and differential Galois theory. Vol. 122. Graduate Studies in Mathematics. American Mathematical Society, Provi- dence, RI, 2011, pp. xiv+225. ISBN: 978-0-8218-5318-4. DOI: 10.1090/gsm/122. URL: https://doi.org/10.1090/gsm/122.
[FW24]	Ruyong Feng and Michael Wibmer. "Differential Galois groups, specializations and Matzat's conjecture". In: to appear in Mem. Amer. Math. Soc. (2024). arXiv: 2209.01581 [math.AG]
[Har05]	Julia Hartmann. "On the inverse problem in differential Galois theory". In: <i>De</i> <i>Gruuter</i> 2005,586 (Sept. 2005), pp. 21–44, ISSN: 1435-5345, DOI: 10.1515/crll.2005.2005.586.21.

[Lan02]	Serge Lang. <i>Algebra</i> . third. Vol. 211. Graduate Texts in Mathematics. Springer- Verlag, New York, 2002, pp. xvi+914. ISBN: 0-387-95385-X. DOI: 10.1007/978-1-4613-0041-0. URL: https://doi.org/10.1007/978-1-4613-0041-0.
[Mag94]	Andy R. Magid. <i>Lectures on differential Galois theory</i> . Vol. 7. University Lecture Series. American Mathematical Society, Providence, RI, 1994, pp. xiv+105. ISBN: 0-8218-7004-1. DOI: 10.1090/ulect/007. URL: https://doi.org/10.1090/ulect/007.
[Mil17]	J. S. Milne. <i>Algebraic groups</i> . Vol. 170. Cambridge Studies in Advanced Mathemat- ics. The theory of group schemes of finite type over a field. Cambridge University Press, Cambridge, 2017, pp. xvi+644. ISBN: 978-1-107-16748-3. DOI: 10.1017/9781316711736. URL: https://doi.org/10.1017/9781316711736.
[MS16]	Claude Mitschi and David Sauzin. <i>Divergent series, summability and resurgence. I.</i> Vol. 2153. Lecture Notes in Mathematics. Monodromy and resurgence, With a fore- word by Jean-Pierre Ramis and a preface by Éric Delabaere, Michèle Loday-Richaud, Claude Mitschi and David Sauzin. Springer, [Cham], 2016, pp. xxi+298. ISBN: 978-3- 319-28735-5; 978-3-319-28736-2. DOI: 10.1007/978-3-319-28736-2. URL: https://doi.org/10.1007/978-3-319-28736-2.
[PS03]	Marius van der Put and Michael F. Singer. Galois theory of linear differential equations. Vol. 328. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2003, pp. xviii+438. ISBN: 3-540-44228-6. DOI: 10.1007/978-3-642-55750-7. URL: https://doi.org/10.1007/978-3-
[Sau16]	Jacques Sauloy. Differential Galois theory through Riemann-Hilbert correspondence. Vol. 177. Graduate Studies in Mathematics. An elementary introduction, With a foreword by Jean-Pierre Ramis. American Mathematical Society, Providence, RI, 2016, pp. xxiii+275. ISBN: 978-1-4704-3095-5. DOI: 10.1090/gsm/177. URL: https://doi.org/10.10
[Spr98]	 T. A. Springer. <i>Linear algebraic groups</i>. Second. Vol. 9. Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1998, pp. xiv+334. ISBN: 0-8176-4021-5. DOI: 10.1007/978-0-8176-4840-4. URL: https://doi.org/10.1007/978-0-8176-4840-4.
[TT79]	Carol Tretkoff and Marvin Tretkoff. "Solution of the inverse problem of differential Galois theory in the classical case". In: <i>Amer. J. Math.</i> 101.6 (1979), pp. 1327–1332. ISSN: 0002-9327,1080-6377. DOI: 10.2307/2374143. URL: https://doi.org/10.2307/2374143.
[Wib20]	Michael Wibmer. "Free Proalgebraic Groups". In: Épijournal de Géométrie Algébrique Volume 4 (Feb. 2020). DOI: 10.46298/epiga.2020.volume4.5733. URL: https://doi.org/10.4629
[Wib24]	Michael Wibmer. "Regular singular differential equations and free proalgebraic groups". In: <i>Bull. Lond. Math. Soc.</i> 56.7 (2024), pp. 2568–2583. ISSN: 0024-6093,1469-2120.

Thomas Serafini, Sorbonne Université, IMJ-PRG, 4 place Jussieu, 75005 Paris, France, $\tt tserafini@dma.ens.fr$

Michael Wibmer, School of Mathematics, University of Leeds, LS2 9JT, Leeds, United Kingdom, m.wibmer@leeds.ac.uk