# Multiplayer Federated Learning: Reaching Equilibrium with Less Communication

TaeHo Yoon	Sayantan Choudhury	Nicolas Loizou
tyoon7@jhu.edu	schoudh8@jhu.edu	nloizou@jhu.edu

Mathematical Institute for Data Science (MINDS), Johns Hopkins University Department of Applied Mathematics & Statistics, Johns Hopkins University

#### Abstract

Traditional Federated Learning (FL) approaches assume collaborative clients with aligned objectives working towards a shared global model. However, in many real-world scenarios, clients act as rational players with individual objectives and strategic behaviors, a concept that existing FL frameworks are not equipped to adequately address. To bridge this gap, we introduce *Multiplayer Federated Learning (MpFL)*, a novel framework that models the clients in the FL environment as players in a game-theoretic context, aiming to reach an equilibrium. In this scenario, each player tries to optimize their own utility function, which may not align with the collective goal. Within MpFL, we propose *Per-Player Local Stochastic Gradient Descent* (PEARL-SGD), an algorithm in which each player/client performs local updates independently and periodically communicates with other players. We theoretically analyze PEARL-SGD and prove that it reaches a neighborhood of equilibrium with less communication in the stochastic setup compared to its non-local counterpart. Finally, we verify our theoretical findings through numerical experiments.

## 1 Introduction

Federated Learning (FL) has emerged as a powerful collaborative learning paradigm where multiple clients jointly train a machine learning model without sharing their local data. In the classical FL setting, a central server coordinates multiple clients (e.g., mobile devices, edge devices) to collaboratively learn a shared global model without exchanging their own training data [48, 54, 79, 64]. In this scenario, each client performs local computations on its private data and periodically communicates model updates to the server, which aggregates them to update the global model. This collaborative approach has been successfully applied in various domains, including natural language processing [69, 43], computer vision [70, 63], and healthcare [4, 116].

Despite their success, traditional FL frameworks rely on the key assumption that all participants are fully cooperative and share aligned objectives, collectively working towards optimizing the performance of a shared global model (e.g., minimizing the average of individual loss functions). This assumption overlooks situations where participants have individual objectives, or competitive interests that may not align with the collective goal. Diverse examples of such scenarios have been extensively considered in the game theory literature, including Cournot competition in economics [2], optical networks [91], electricity markets [98], energy consumption control in smart grid [120], or mobile robot control [49]. Despite their relevance, these applications have yet to be associated with FL, presenting an unexplored opportunity to bridge game theory and FL for more robust and realistic frameworks.

To address these limitations of classical FL approaches, we propose a novel framework called *Multiplayer Federated Learning (MpFL)*, which models the FL process as a game among rational

players with individual utility functions. In MpFL, each participant is considered a player who aims to optimize their own objective while interacting strategically with other players in the network via a central server. This game-theoretic perspective acknowledges that participants may act in their self-interest, have conflicting goals, or be unwilling to fully cooperate. By incorporating these dynamics, MpFL provides a more realistic and flexible foundation for FL in competitive and heterogeneous environments.

In the literature, there are multiple strategies that aim to incorporate a personalization approach into classical FL, including multi-task learning [106, 82], transfer learning [52], and mixing of the local and global models [40, 41], to name a few. However, to the best of our knowledge, none of them is able to formulate the behaviour of the clients/players in a non-cooperative environment. This gap is precisely what Multiplayer Federated Learning (MpFL) aims to address.

### 1.1 Main contributions

In this work, we make the following main contributions:

- Introducing Multiplayer Federated Learning. We develop a novel framework of *Multiplayer Federated Learning (MpFL)*, which models the FL process as a game among rational players with individual utility functions. In MpFL, each client within the FL environment is viewed as a player of the game, and their local models are viewed as their actions. Each player constantly adjusts their model (action) to optimize their own objective function, and the MpFL framework aims for each player to reach to a Nash equilibrium by collaboratively training their model under the orchestration of a central server (e.g., service provider), while keeping the training data decentralized. That is, MpFL extends the scope of FL to scenarios where clients are allowed to have more general, diversified, possibly competing objectives.
- **Design and analysis of Per-Player Local SGD.** To handle the Multiplayer Federated Learning framework, we introduce *Per-Player Local SGD* (PEARL-SGD), a new algorithm inspired by the stochastic gradient descent ascent method in minimax optimization, that is able to handle the competitive nature of the players/clients. In PEARL-SGD, each player performs local SGD steps independently on their own actions/strategies (keeping the strategies of the other players fixed), and the udpated actions/models are periodically communicated with the other players of the network via a central server.
- Convergence guarantees for PEARL-SGD on heterogeneous data. We provide tight convergence guarantees for PEARL-SGD, in both deterministic and stochastic regimes with heterogeneous data (see Table 1 for a summary of our results).
  - **Deterministic setting:** For the full-batch (deterministic) variant of PEARL-SGD, we prove that under suitable assumptions, PEARL-SGD converges linearly to an equilibrium for any communication period  $\tau > 1$ , provided that the constant step-size  $\gamma$  is sufficiently small (see Theorem 3.3).
  - Stochastic setting: In its more general version, PEARL-SGD assumes that each player uses an unbiased estimator of its gradient in the update rule. For this setting, we provide two Theorems based on two different step-size choices:
    - \* Constant step-size: We show that under the same assumptions as in the deterministic case, PEARL-SGD converges linearly to a neighborhood of equilibrium (see Theorem 3.4). In Corollary 3.5, we show that with appropriate step-size depending on the total number of local SGD iterations T, PEARL-SGD achieves  $\tilde{\mathcal{O}}(1/T)$  convergence rate with improved communication complexity when T is sufficiently large.

- \* Decreasing step-size rule: We prove that PEARL-SGD converges to an exact equilibrium (without neighborhood of convergence) with sublinear convergence (see Theorem 3.6). In this scenario, the asymptotic rate and communication complexity are essentially the same as in Corollary 3.5, but this result does not require the step-sizes to depend on T.
- Numerical Evaluation: We provide extensive numerical experiments verifying our theoretical results and show the benefits in terms of communications of PEARL-SGD over its non-local counterpart in the MpFL settings.

Table 1: Summary of theoretical results for PEARL-SGD. Theorem 3.3 considers the full-batch (deterministic) scenario. Theorem 3.4 and Theorem 3.6 both considers the general stochastic case. These results differ in the step-size choice; the former uses a constant step-size, while the latter uses decreasing step-sizes. In the *Convergence* column, "Linear" and "Sublinear" indicates the convergence rate, "Exact" refers to convergence to an equilibrium, and "Neighborhood" refers to convergence to a neighborhood of an equilibrium.

Theorem	Setting	Step-size	Convergence
Theorem 3.3	Deterministic	Constant	Linear+Exact
Theorem 3.4	Stochastic	Constant	Linear+Neighborhood
Theorem 3.6	Stochastic	Decreasing	Sublinear+Exact

## 2 Multiplayer Federated Learning: Definition and Related Work

In this section, we introduce the Multiplayer Federated Learning (MpFL) framework and explain its main differences compared to the classical FL [48], federated minimax optimization [19, 104, 124] and personalized FL [26, 111].

### 2.1 Definition of MpFL

Multiplayer Federated Learning (MpFL) is a machine learning setting that combines the benefits of a game-theoretic formulation with classical federated learning. In this setting, the problem is an *n*-player game in which multiple players/clients (e.g. mobile devices or whole organizations) communicate with each other via a central server (e.g. service provider) to reach equilibrium. That is, reach a set of strategies where no player can unilaterally deviate from their strategy to achieve a better payoff, given the strategies chosen by all other players.

In classical *n*-player games, communication between players was assumed to be cheap, easy, and straightforward, mainly because all players were in close proximity and had direct access to one another. This assumption made communication an insignificant concern in typical game theory analysis. However, with the advent of new large-scale machine learning applications, this is no longer the case. Communication between players can be expensive and challenging, especially in distributed systems where the clients/players are geographically dispersed or operate under communication constraints. Addressing communication costs and designing communication-efficient algorithms for *n*-player games have become increasingly important, and this is precisely the challenge that Multiplayer Federated Learning aims to address.

**Equilibrium in** *n***-player game.** Let  $x^i \in \mathbb{R}^{d_i}$  denote the action of player  $i = 1, \ldots, n$  and let  $\mathbf{x} = (x^1, \ldots, x^n) \in \mathbb{R}^D = \mathbb{R}^{d_1 + \cdots + d_n}$  be the joint action/strategy vector of all players. Let  $f_i(x^1, \ldots, x^n) : \mathbb{R}^{d_1 + \cdots + d_n} \to \mathbb{R}$  be the function of the player *i* (which player *i* prefers to minimize in  $x^i$ ) and let  $x^{-i} = (x^1, \ldots, x^{i-1}, x^{i+1}, \ldots, x^n) \in \mathbb{R}^{D-d_i}$  be the vector containing all players' actions except that of player *i*. With this notation in place, the goal in a *n* player game is to find an *equilibrium*, a joint action  $\mathbf{x}_{\star} = (x_{\star}^1, \ldots, x_{\star}^n) \in \mathbb{R}^D$ , formally expressed as:

$$\inf_{\mathbf{x}_{\star}=(x_{\star}^{1},\ldots,x_{\star}^{n})\in\mathbb{R}^{D}} \quad f_{i}(x_{\star}^{i};x_{\star}^{-i}) \leq f_{i}(x^{i};x_{\star}^{-i}), \quad \forall x^{i}\in\mathbb{R}^{d_{i}}, \quad \forall i\in[n],$$
(1)

where  $f_i(x^i; x^{-i}) = f_i(x^1, ..., x^n)$ .

**MpFL.** As mentioned above, in the setting of interest of this work, we focus on an n-player game in which multiple players communicate via a central server to reach an equilibrium. In this setting, each player of the n-player game represents a client to the system (see Figure 1). Mathematically, the problem is formulated as solving (1) with

$$f_i(x^1,\ldots,x^n) = \mathbb{E}_{\xi^i \sim \mathcal{D}_i} \left[ f_{i,\xi^i}(x^1,\ldots,x^n) \right].$$

Here  $\mathcal{D}_i$  denotes the data distribution of the *i*th player,  $f_{i,\xi^i}$  is the loss of the *i*-th player for a data point  $\xi^i$  sampled from  $\mathcal{D}_i$ .

In our proposed FL environment, each client/player uses the strategies of all players to execute local updates. In particular, each player keeps the other players' strategies fixed



 $f_1(x^1, x^2, ..., x^n) \quad f_2(x^1, x^2, ..., x^n) \qquad f_n(x^1, x^2, ..., x^n)$ 

Figure 1: Illustration of MpFL for heterogeneous functions  $f_i$ . The goal is for each player to reach the equilibrium  $\mathbf{x}_{\star} = (x_{\star}^1, \ldots, x_{\star}^n)$  (see (1)) with as little communication as possible.

player keeps the other players' strategies fixed and updates their own value, which is later shared with the master server, which concatenates all new strategies and sends them back to all players. Later, in Section 3 we introduce and analyze Algorithm 1, named *Per-Player Local SGD* (PEARL-SGD), which formalizes the above setting.

Similarly to the classical FL regime, our setting focuses on *heterogeneous data* (non-i.i.d.) as we do not make any restrictive assumption on the data distribution  $\mathcal{D}_i$  or the similarity between the functions of the players.

Assumptions on multiplayer game. Let us present the main assumptions on the functions of the multiplayer game, which we later use to provide the convergence analysis for the proposed *Per-Player Local SGD*. In our work, we make two main assumptions on the functions  $f_i$  of each player  $i \in [n]$ . We assume that the function is convex and smooth.

Throughout this work, we denote the gradient of  $f_i$  (function of player  $i \in [n]$ ) with respect to  $x^i$  by:

$$\nabla_{x^i} f_i(x^1, \dots, x^n) = \nabla f_i(x^i; x^{-i}).$$

This convention allows us to remove the cumbersome subscript  $x^i$  from the  $\nabla$  notation; we only differentiate  $f_i$  with respect to  $x^i$  but never with  $x^{-i}$ .

Assumption 2.1 (*Convex (CVX)*). For  $i \in [n]$ , for any  $x^{-i} \in \mathbb{R}^{D-d_i}$ , the local function  $f_i(\cdot; x^{-i}) \colon \mathbb{R}^{d_i} \to \mathbb{R}$  is convex. That is, for any  $x^i, y^i \in \mathbb{R}^{d_i}$  and  $x^{-i} \in \mathbb{R}^{D-d_i}$ ,

$$f_i(y^i; x^{-i}) \ge f_i(x^i; x^{-i}) + \left\langle \nabla f_i(x^i; x^{-i}), y^i - x^i \right\rangle$$

Assumption 2.2 (*Smoothness (SM*)). For  $i \in [n]$ , for any  $x^{-i} \in \mathbb{R}^{D-d_i}$ , the local function  $f_i(\cdot; x^{-i}) \colon \mathbb{R}^{d_i} \to \mathbb{R}$  is  $L_i$ -smooth. That is, for any  $x^i, y^i \in \mathbb{R}^{d_i}$  and  $x^{-i} \in \mathbb{R}^{D-d_i}$ ,

$$\left\| \nabla f_i(x^i; x^{-i}) - \nabla f_i(y^i; x^{-i}) \right\| \le L_i \left\| x^i - y^i \right\|.$$

As in (1), in the stochastic regime of MpFL we have  $f_i(x^1, \ldots, x^n) = \mathbb{E}_{\xi^i \sim \mathcal{D}_i} \left[ f_{i,\xi^i}(x^1, \ldots, x^n) \right]$ . To obtain convergence guarantees for PEARL-SGD in this scenario, we need the following assumption of bounded variance of the gradient oracle, commonly used in stochastic optimization.

Assumption 2.3 (Bounded Variance (BV)). Let  $\sigma_i \ge 0, \forall i \in [n]$ . For each i = 1, ..., n,  $\mathbb{E}_{\xi^i \sim \mathcal{D}_i} \left[ \left\| \nabla f_{i,\xi^i}(x^i; x^{-i}) - \nabla f_i(x^i; x^{-i}) \right\|^2 \right] \le \sigma_i^2, \quad \forall x^i \in \mathbb{R}^{d_i}, x^{-i} \in \mathbb{R}^{D-d_i}.$ 

### 2.2 Comparison with closely related FL frameworks

Having presented the MpFL setting, let us now provide a concise survey of related setups: classical FL, federated minimax optimization and personalized FL. We compare each of them with our proposed MpFL. An additional list of related work is provided in Appendix A.

Federated learning (FL). In its basic formulation, classical federated learning can be expressed as the minimization of the objective function [48],

$$\underset{x \in \mathbb{R}^d}{\text{minimize}} \quad f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x) \quad \text{where} \quad f_i(x) = \mathbb{E}_{\xi^i \sim \mathcal{D}_i}[F_i(x,\xi^i)].$$

Here,  $x \in \mathbb{R}^d$  represents the parameter for the global model,  $f_i$  denotes the local objective function at client *i*, and  $\mathcal{D}_i$  denotes the data distribution of client *i*. The local loss functions  $F_i(x,\xi^i)$  are often the same across all clients, but the local data distribution  $\mathcal{D}_i$  will often vary, capturing data heterogeneity. The foundational communication-efficient algorithm for this setup is FedAvg (Local SGD), proposed and massively popularized by McMahan et al. [78]. Despite its simplicity, Local SGD has shown empirical success in terms of convergence speed and communication frequency, and many works have provided theoretical explanation for this performance [109, 23, 110, 51].

In these works, clients work in a fully cooperative manner to find  $x_{\star} = \operatorname{argmin}_{x \in \mathbb{R}^d} f(x)$ , unlike our proposed MpFL where the clients who now serve as players of the game seek an equilibrium among possibly competing (non-cooperative) objectives.

**Federated minimax optimization.** Federated minimax optimization is a more recent concept proposed as a federated extension of minimax optimization problems appearing in many machine learning applications. In that scenario, the main problem can be expressed as:

$$\underset{x \in \mathbb{R}^{d_x}}{\text{minimize}} \quad \mathcal{L}(x) = \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(x, y) \quad \text{where} \quad \mathcal{L}_i(x, y) = \mathbb{E}_{\xi^i \sim \mathcal{D}_i}[\phi_i(x, y, \xi)].$$

Here *n* is the number of clients, and  $\mathcal{L}_i(x, y)$  represents the local loss function at client *i* that depends on both *x* and *y*, defined as  $\mathcal{L}_i(x, y) = \mathbb{E}_{\xi^i \sim \mathcal{D}_i}[\phi_i(x, y, \xi)]$ . Note that in that scenario, each client has access to the information of both players *x* and *y*.  $\phi_i(x, y, \xi)$  denotes the loss for the data point  $\xi$ , sampled from the local data distribution  $\mathcal{D}_i$  at client *i*. Based on the properties of the model, the functions  $\mathcal{L}_i(x, y)$  can be smooth/non-smooth, convex/non-convex with respect to player *x*, and concave/non-concave with respect to player *y*. The extension of Local SGD for solving this problem are Local Stochastic Gradient Descent-Ascent (SGDA) [19, 104] or Local Stochastic Extragradient (SEG) [6, 7] algorithms. More recently there was also an approach based on primal-dual updates [14].

While this line of work also studied federated learning in the context of minimax optimization and games, it is totally different from MpFL. As we explained above, the setup in this scenario assumes that each FL client has access to both players of the minimax game, and they do not take the *multiplayer* aspect into account. In contrast, in our setting, we assume that each client is a player of a large-scale multiplayer game who only has access to their objective  $f_i$  and its gradient and only updates their action  $x^i$ . In our work, we design the novel PEARL-SGD algorithm, which is suitable for the MpFL setting, a task not possible using the existing Local SGDA and Local SEG methods.

**Personalized federated learning.** In personalized FL [26, 111, 41, 40, 20, 112], clients aim to learn models tailored to each local data distribution, while generalizing well over the data from other clients [42]. One way of formulating personalized FL is [41, 42]:

$$\underset{\mathbf{x}=(x^1,\dots,x^n)\in\mathbb{R}^{nd}}{\text{minimize}} \quad \frac{1}{n}\sum_{i=1}^n h_i(x^i) + \frac{\lambda}{2n}\sum_{i=1}^n \left\|x^i - \overline{x}\right\|^2$$

where  $x^1, \ldots, x^n \in \mathbb{R}^d$  are local models of each player,  $h_1, \ldots, h_n$  are the local losses following the each player's distributions,  $\overline{x} = \frac{1}{n} \sum_{i=1}^{n} x^i$  is the average model and  $\lambda > 0$  is regularization strength. Given that each  $h_i$  is convex, the first-order optimality condition for the above is  $x^i_{\star} - \overline{x}_{\star} + \frac{1}{\lambda} \nabla h_i(x^i_{\star}) = 0$  [41]. This is equivalent to the equilibrium condition for the *n*-player game (MpFL) where each player has the objective function  $f_i(x^i; x^{-i}) = h_i(x^i) + \frac{\lambda}{2} \sum_{i=1}^n ||x^i - \overline{x}||^2$ . In this respect, personalized FL can be viewed as a particular instance of MpFL.

## **3** PEARL-SGD: Algorithm and Convergence Guarantees

In this section, we introduce and analyze Algorithm 1, named *Per-Player Local SGD* (PEARL-SGD), which is suitable for the MpFL setting we described in Section 2.

### 3.1 Algorithm and assumptions

PEARL-SGD works by having the clients/players of the game run SGD independently in parallel for updating their strategy (keeping the strategies  $x^{-i}$  of the other players fixed) and concatenate the strategies of all players only once in a while (via a central server). In more detail, in every round of PEARL-SGD, each player  $i \in [n]$  runs  $\tau$  iterations of stochastic gradient descent (SGD) with respect to  $f_i(\cdot, x^{-i})$ , having the  $x^{-i}$  fixed to be the information of the other players' actions obtained from the previous synchronization step. Once each player completes  $\tau$  iterations of SGD (local updates), a synchronization occurs. In each synchronization, the central server collects the actions of all players, and then the concatenation of all updated strategies/actions is distributed to all clients/players. We emphasize that PEARL-SGD and its convergence guarantees hold without any assumption on players' data distributions  $\mathcal{D}_i$ . That is, functions  $f_i$  can be very different between players, and the setting is fully heterogeneous.

### Algorithm 1 Per-Player Local SGD (PEARL-SGD)

**Input:** Step-sizes  $\gamma_k > 0$ , Synchronization interval  $\tau \ge 1$ , Number of synchronization/local update rounds  $R \ge 1$  **for**  $p = 0, \ldots, R - 1$  **do** Master server collects  $x_{\tau p}^i$  from players  $i = 1, \ldots, n$  and forms  $\mathbf{x}_{\tau p} = (x_{\tau p}^1, \ldots, x_{\tau p}^n)$ Master server distributes  $\mathbf{x}_{\tau p}$  back to players  $i = 1, \ldots, n$  **for each players**  $i = 1, \ldots, n$  **in parallel do for**  $k = \tau p, \ldots, \tau(p+1) - 1$  **do** Draw  $\xi_k^i \sim \mathcal{D}_i$   $g_k^i \leftarrow \nabla f_{i,\xi_k^i}(x_k^i; x_{\tau p}^{-i})$   $x_{k+1}^i \leftarrow x_k^i - \gamma_k g_k^i$ end for end for **output:**  $\mathbf{x}_{\tau R} \in \mathbb{R}^D$ 

Let us note that the synchronization step in PEARL-SGD involves transferring a  $D = (d_1 + \cdots + d_n)$ -dimensional vector  $\mathbf{x}_{\tau p}$  from the master server to the players. This is an important difference compared to the classical FL (minimization problem), where the dimension of the communication vectors is the same from client to master and from master to client, and it does not scale with n. While PEARL-SGD aims to reduce this overhead compared to its distributed variant ( $\tau = 1$ ) by communicating less frequently (with  $\tau > 1$ ), the high complexity of the synchronization step makes MpFL (and distributed n-player games in general) more suitable for cross-silo FL setups with relatively small number of organizations and more reliable communication bandwidths. We expect that the potentially expensive communication of high-dimensional vectors  $\mathbf{x}_{\tau p}$  could be addressed by incorporating additional techniques such as gradient compression [3, 8]. This is an orthogonal approach to our proposed local methods, and we leave it for future work.

Assumptions on the joint gradient operator. We require some definitions and additional assumptions in order to carry out the theory. Define the joint gradient operator  $\mathbb{F} \colon \mathbb{R}^D \to \mathbb{R}^D$  as

 $\mathbb{F}(\mathbf{x}) = \left(\nabla f_1(x^1; x^{-1}), \dots, \nabla f_n(x^n; x^{-n})\right).$ 

Assumption 3.1 (*Quasi-strong monotonicity (QSM*)). There exists a unique equilibrium  $\mathbf{x}_{\star} = (x_{\star}^{1}, \ldots, x_{\star}^{n}) \in \mathbb{R}^{D}$ , for which  $\mathbb{F}(\mathbf{x}_{\star}) = 0$ , and  $\mu > 0$  such that for any  $\mathbf{x} \in \mathbb{R}^{D}$ ,  $\langle \mathbb{F}(\mathbf{x}), \mathbf{x} - \mathbf{x}_{\star} \rangle \geq \mu \| \mathbf{x} - \mathbf{x}_{\star} \|^{2}$ .

(QSM) is a concept extending quasi-strong convexity [37, 36] to the context of variational inequality problems (VIPs). This condition has been referred to as different names in the literature, such as strong coherent VIPs [108], VIPs with strong stability condition [80], or the strong Minty variational inequality [22]. It generalizes strong monotonicity, capturing some non-monotone problems. In [75], it was proposed and utilized as an assumption ensuring the convergence of SGDA dynamics in minimax games without the well-known issues of cycling or

diverging [81, 16]. Later, it was also used in the analysis of stochastic extragradient method [33] and its single-call variants (optimistic and past stochastic etragradient) [13].

Assumption 3.2 (*Star-cocoercivity (SCO*)).  $\mathbb{F}$  is  $\frac{1}{\ell}$ -star-cocoercive, i.e., there is  $\ell > 0$  such that for any  $\mathbf{x} \in \mathbb{R}^D$ ,  $\langle \mathbb{F}(\mathbf{x}), \mathbf{x} - \mathbf{x}_{\star} \rangle \geq \frac{1}{\ell} \| \mathbb{F}(\mathbf{x}) \|^2$ .

(SCO) generalizes the class of coercive operators and, interestingly, can hold for non-Lipschitz operators [75]. This has also been taken as minimal assumption for SGDA analysis in prior work [8]. Note that (QSM) and (SCO) together imply  $\mu \|\mathbf{x} - \mathbf{x}_{\star}\| \leq \|\mathbf{F}(\mathbf{x})\| \leq \ell \|\mathbf{x} - \mathbf{x}_{\star}\|$ for any  $\mathbf{x} \in \mathbb{R}^{D}$ , which implies  $\mu \leq \ell$ . We call  $\kappa = \ell/\mu \geq 1$  the condition number of the problem. In Appendix D, we provide a detailed discussion on the set of our theoretical assumptions and explain connections to other commonly assumed properties in the literature such as cocoercivity, Lipschitzness and monotonicity.

### 3.2 Convergence of PEARL-SGD: Deterministic setup

First, we provide the convergence result for PEARL-SGD with constant step-size  $\gamma_k \equiv \gamma$  in the full-batch (deterministic) scenario, where there is no noise in the gradient computation. While this is recovered as a special case of Theorem 3.4, we state it separately because the deterministic case provides several points of discussion that are worth emphasizing on their own.

**Theorem 3.3.** Assume *(CVX)*, *(SM)*, *(QSM)* and *(SCO)*. Let  $L_{\max} = \max\{L_1, \ldots, L_n\}$ ,  $0 < \gamma_k \equiv \gamma \leq \frac{1}{\ell\tau + 2(\tau-1)L_{\max}\sqrt{\kappa}}$ , and let  $\kappa = \ell/\mu$  be the condition number. Then the Deterministic PEARL-SGD (Algorithm 1 with full-batch) converges with the rate

$$\|\mathbf{x}_{\tau R} - \mathbf{x}_{\star}\|^{2} \leq (1 - \gamma \tau \mu \zeta)^{R} \|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2}$$

where  $\zeta = 2 - \gamma \ell \tau - 2(\tau - 1)\gamma L_{\max} \sqrt{\kappa/3} > 0$  (by the choice of  $\gamma$ ).

Theorem 3.3 shows that deterministic PEARL-SGD converges linearly to an equilibrium. This distinguishes our result from the analyses of local gradient descent for finite sum minimization in heterogeneous data setups, where one has convergence to a neighborhood of optimum even when there is no noise [51], unless further correction mechanism is used [84]. In addition, note that when  $\tau = 1$ , the step-size constraint and the convergence rate of Theorem 3.3 coincide with those from the analysis of the gradient descent-ascent (GDA) under the *(QSM)* and *(SCO)* assumptions from [75], showing the tightness of our analysis.

Player drift and step-size constraint. If  $\gamma$  does not appropriately scale down with  $\tau$ , then at each round, players' actions (SGD iterates) converge to minimizers of local functions. We call this phenomenon *player drift*, analogous to client drift in classical FL [50], enforcing the  $\mathcal{O}(1/\tau)$ step-size. In our setting, note that the local minimizers  $x_{\star}^{i}(x_{\tau p}^{-i}) := \operatorname{argmin}_{x^{i} \in \mathbb{R}^{d_{i}}} f_{i}(x^{i}; x_{\tau p}^{-i})$ depend on other players' strategies  $x_{\tau p}^{-i}$ . Due to this dependence, under extreme player drift, PEARL-SGD may display undesirable dynamics such as diverging to infinity<sup>1</sup>. As these features are not typically observed in client drift in classical FL, player drift represents a distinct phenomenon despite some conceptual similarities. Therefore, we consider understanding and mitigating player drift an intriguing direction for future work in MpFL, which may necessitate novel insights that differ from existing approaches to client drift [50, 84].

<sup>&</sup>lt;sup>1</sup>This can be checked with simple examples such as the two-player quadratic minimax game  $\min_{u \in \mathbb{R}} \max_{v \in \mathbb{R}} \frac{\mu}{2} u^2 + uv - \frac{\mu}{2} v^2$  with  $\mu < 1$ .

### 3.3 Convergence of PEARL-SGD: Stochastic setup

We now discuss the convergence of PEARL-SGD with stochastic gradients. We first present the convergence of PEARL-SGD to a neighborhood of an equilibrium  $\mathbf{x}_{\star}$  given constant step-sizes  $\gamma_k \equiv \gamma$ , and then discuss the communication complexity gain we achieve. Then we present the convergence result using a decreasing step-size selection, showing sublinear convergence to the exact equilibrium  $\mathbf{x}_{\star}$  rather than its neighborhood. While we defer the details of the proofs to Appendix B, we provide a proof outline for Theorem 3.4 in Section 3.4.

**Theorem 3.4.** Assume *(CVX)*, *(SM)*, *(BV)*, *(QSM)* and *(SCO)* hold. Let  $0 < \gamma_k \equiv \gamma \leq \frac{1}{\ell\tau + 2(\tau-1)L_{\max}\sqrt{\kappa}}$  and denote  $q = L_{\max}/\sqrt{\ell\mu}$ . Then PEARL-SGD exhibits the rate:

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau R}-\mathbf{x}_{\star}\right\|^{2}\right] \leq \left(1-\gamma\tau\mu\zeta\right)^{R}\|\mathbf{x}_{0}-\mathbf{x}_{\star}\|^{2}+\left(1+\left(\tau-1\right)\left(\left(4+\sqrt{3}q\right)\gamma\tau L_{\max}+\frac{q}{2\tau}\right)\right)\frac{\gamma\sigma^{2}}{\mu\zeta}.$$

where  $\sigma^2 = \sum_{i=1}^n \sigma_i^2$  and  $\zeta = 2 - \gamma \ell \tau - 2(\tau - 1)\gamma L_{\max} \sqrt{\kappa/3} > 0$  by the choice of  $\gamma$ .

When  $\tau = 1$ , with  $\gamma \leq 1/\ell$ , the above rate becomes  $\mathbb{E}\left[\|\mathbf{x}_R - \mathbf{x}_\star\|^2\right] \leq (1 - \gamma \mu)^R \|\mathbf{x}_0 - \mathbf{x}_\star\|^2 + \gamma^{\sigma^2/\mu}$ , which is consistent with the classical analysis of the stochastic gradient descent-ascent (SGDA). In the result, note that  $\sigma^2$  is the sum of  $\sigma_i^2$ 's, the (upper bounds on) playerwise gradient variances  $(\sigma_i^2 \geq \mathbb{E}_{\xi^i \sim D_i} \left[ \|\nabla f_{i,\xi^i}(x^i; x^{-i}) - \nabla f_i(x^i; x^{-i})\|^2 \right]$ ). Hence,  $\sigma^2$  represents the upper bound on the variance in estimating the joint gradient operator  $\mathbb{F}(\cdot)$ .

**Remark.** If we use the largest possible step-size  $\gamma = \frac{1}{\ell\tau + 2(\tau-1)L_{\max}\sqrt{\kappa}}$  allowed in Theorem 3.4, then the right-hand side of the bound does not scale down indefinitely with  $\tau$ . In fact, with this choice of  $\gamma$ , one can expect the communication gain by a factor of approximately  $\frac{L_{\max}}{\ell}$  (when  $L_{\max} \ll \ell$ ). More precisely, suppose  $q \leq 1$  (equivalently  $L_{\max} \leq \sqrt{\ell\mu}$ —refer to Appendix D for the explanation that this is a common parameter regime). Then we have  $\gamma = \Theta\left(\frac{1}{\ell\tau}\right)$  and

$$\mathbb{E}\left[\|\mathbf{x}_{\tau R} - \mathbf{x}_{\star}\|^{2}\right] \leq (1 - \gamma \tau \mu)^{R} \|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2} + \mathcal{O}\left(\frac{1}{\tau} + \frac{L_{\max}}{\ell}\right) \frac{\sigma^{2}}{\ell \mu}.$$

The first linear convergence term is essentially unaffected by  $\tau$ , as the effect of using smaller  $\gamma = \Theta\left(\frac{1}{\ell\tau}\right)$  is canceled out by the factor  $\tau$  within  $(1 - \gamma\tau\mu)^R$ . In the second term (which is usually dominant), we see that the size of the convergence neighborhood is reduced by the factor  $\mathcal{O}\left(\frac{1}{\tau} + \frac{L_{\max}}{\ell}\right) = \mathcal{O}\left(\frac{1}{\tau} + \frac{1}{\sqrt{\kappa}}\right)$ . Therefore, we see that with  $\tau = \Omega(\sqrt{\kappa})$ , PEARL-SGD reaches about  $\sqrt{\kappa}$  times smaller neighborhood within the same number of communication rounds R (compared to the case  $\tau = 1$ ).

In Corollary 3.5, we analyze the convergence and communication gain of PEARL-SGD in the regime where the total number of iterations  $T = \tau R$  is large. In this setting, we use a step-size that depends on the total number of iterations T.

**Corollary 3.5.** Suppose the assumptions of Theorem 3.4 hold, and let  $\tau \ge 1$  be fixed. Let  $q = L_{\max}/\sqrt{\ell\mu}$ . Then PEARL-SGD with  $\gamma_k \equiv \gamma = \frac{1}{\mu\eta(1+2q)}$  exhibits the rate

$$\mathbb{E}\left[\left\|\mathbf{x}_{T} - \mathbf{x}_{\star}\right\|^{2}\right] = \tilde{\mathcal{O}}\left(\frac{(1+q)^{2} \left\|\mathbf{x}_{0} - \mathbf{x}_{\star}\right\|^{2}}{T^{2}} + \frac{(1+q)\sigma^{2}}{\mu^{2}T} + \frac{(1+q)\tau^{2}L_{\max}\sigma^{2}}{\mu^{3}T^{2}}\right)$$

where  $\eta$  is selected so that  $T = 2(1+2q)\eta \log \eta$ , provided that T is large enough so that  $\eta > \kappa \tau$ . Here the  $\tilde{\mathcal{O}}$ -notation hides polylogarithmic terms in T and constant factors.

Reduction of communication complexity. Note that the  $\tilde{\mathcal{O}}\left(\frac{(1+q)^2 \|\mathbf{x}_0 - \mathbf{x}_\star\|^2}{T^2}\right)$  term decays fast in Corollary 3.5 (as T grows) and the terms proportional to  $\sigma^2$  become dominant. The order of convergence is not slower than the  $\tilde{\mathcal{O}}(1/T)$  rate of the fully communicating case  $\tau = 1$ , provided that  $\tau^2 L_{\max} \sigma^2 / \mu^3 T^2 = \mathcal{O}(\sigma^2 / \mu^2 T) \iff \tau = \mathcal{O}\left(\sqrt{\mu T / L_{\max}}\right)$ . Therefore, as long as we select  $\tau = \mathcal{O}\left(\sqrt{\mu T / L_{\max}}\right)$ , in PEARL-SGD the communication cost is reduced by the factor of  $\tau$ (because the total number of communications is  $T/\tau$ ). With the largest possible  $\tau$ , the resulting communication complexity is  $T/\tau = \Theta\left(\sqrt{T L_{\max} / \mu}\right) = \Theta\left(\sqrt{T}\right)$ .

Convergence to equilibrium via decreasing step-sizes. We conclude the section with convergence result for PEARL-SGD using a decreasing step-size selection. While showing a similar convergence rate in terms of T as in Corollary 3.5, Theorem 3.6 has the advantage of not requiring to fix T in advance to determine the step-sizes.

**Theorem 3.6.** Under the assumptions of Theorem 3.4, let  $q = L_{\max}/\sqrt{\ell\mu}$ , and choose the step-sizes  $\gamma_k = \begin{cases} \frac{1}{\ell\tau(1+2q)} & \text{if } p < 2(1+2q)\kappa \\ \frac{1}{\tau\mu}\frac{2p+1}{(p+1)^2} & \text{if } p \geq 2(1+2q)\kappa \end{cases}$  for  $\tau p \leq k \leq \tau(p+1) - 1$ ,  $p = 0, \ldots, R - 1$ . Then PEARL-SGD converges with the rate

$$\mathbb{E}\left[\left\|\mathbf{x}_{T} - \mathbf{x}_{\star}\right\|^{2}\right] \leq \frac{4(1+2q)^{2}\kappa^{2}\tau^{2} \|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2}}{eT^{2}} + \frac{4(1+q)\sigma^{2}}{\mu^{2}T} + \frac{4(1+2q)^{2}\kappa\tau\sigma^{2}}{\mu^{2}T^{2}}\left(1 + \frac{2\tau}{\sqrt{\kappa}}\right) + \frac{32(1+q)\tau^{2}L_{\max}\sigma^{2}\log T}{\mu^{3}T^{2}}$$

where  $T = \tau R$  is the total number of iterations.

### 3.4 Proof outline

In this section, we provide a proof outline for Theorem 3.4. The key components of the proof are as follows: (i) a round of local SGD in PEARL-SGD behaves like a large single descent step with respect to the joint gradient operator  $\mathbb{F}$  except for *local error* terms caused by running multiple SGD steps locally (Lemma 3.7), and (ii) we bound these local error terms (Lemma 3.8).

**Lemma 3.7.** Assume *(SM)*, and let  $L_{\max} = \max\{L_1, \ldots, L_n\}$ . Let  $0 \le p \le R-1$  be a fixed round index in PEARL-SGD and suppose  $\gamma_k \equiv \gamma > 0$  for  $k = \tau p, \ldots, \tau(p+1) - 1$ . Then for arbitrary  $\alpha > 0$ , we have

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] \leq \left(1 + (\tau - 1)\alpha\gamma\right) \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} - 2\gamma\tau \left\langle \mathbb{F}(\mathbf{x}_{\tau p}), \mathbf{x}_{\tau p} - \mathbf{x}_{\star} \right\rangle \\ + \frac{\gamma L_{\max}^{2}}{\alpha} \sum_{j=\tau p+1}^{\tau p+\tau-1} \mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{j}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] + \mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\tau(p+1)}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right]\right].$$

Local error bound. The right hand side of the bound in Lemma 3.7 involves the quantities

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{j}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] = \sum_{i=1}^{n} \mathbb{E}\left[\left\|x_{\tau p}^{i} - x_{j}^{i}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right]\right]$$
(2)

for  $j = \tau p + 1, \ldots, \tau (p + 1)$ . We further bound (2) using the following result.

**Lemma 3.8.** Suppose Assumptions *(CVX)*, *(SM)* and *(BV)* hold. For a fixed  $i \in [n]$  and a fixed communication round p in PEARL-SGD, suppose  $\gamma_k \equiv \gamma$  for  $k = \tau p, \ldots, \tau (p+1) - 1$ , where  $0 < \gamma \leq \frac{1}{L_i} \min \left\{ 1, \frac{1}{\tau - 1} \right\}$ . Then for  $t = 0, \ldots, \tau$ ,  $\mathbb{E} \left[ \left\| x_{\tau p}^i - x_{\tau p+t}^i \right\|^2 \, \left| \mathbf{x}_{\tau p} \right] \leq \gamma^2 t^2 \left\| \nabla f(x_{\tau p}^i; x_{\tau p}^{-i}) \right\|^2 + \gamma^2 t \left( 1 + 2(t-1)(t+1)\gamma L_i \right) \sigma_i^2.$ 

Here we sketch the proof of Lemma 3.8 and clarify the role of Assumption (CVX). By assuming that each  $f_i(\cdot; x_{\tau p}^{-i})$  is convex and  $L_i$ -smooth, we can prove Lemma 3.9, showing that the expectation of squared gradient norm is "almost" nonincreasing along the local SGD steps, except for some additional term due to stochasticity. Then, we rewrite each summand in (2) as

$$\mathbb{E}\left[\left\|x_{\tau p}^{i}-x_{k}^{i}\right\|^{2}\left\|\mathbf{x}_{\tau p}\right]=\mathbb{E}\left[\gamma^{2}\left\|\sum_{j=\tau p}^{k-1}g_{j}^{i}\right\|^{2}\left\|\mathbf{x}_{\tau p}\right]=\mathbb{E}\left[\gamma^{2}\left\|\sum_{j=\tau p}^{k-1}\nabla f_{i,\xi_{j}^{i}}(x_{j}^{i};x_{\tau p}^{-i})\right\|^{2}\left\|\mathbf{x}_{\tau p}\right]$$
(3)

and use Lemma 3.9 to bound (3).

**Lemma 3.9.** Under the assumptions of Lemma 3.8, for 
$$j = \tau p + 1, \ldots, \tau (p+1)$$
,  

$$\mathbb{E} \left[ \left\| \nabla f_i(x_j^i; x_{\tau p}^{-i}) \right\|^2 \, \left| \, \mathbf{x}_{\tau p} \right] \leq \left\| \nabla f_i(x_{\tau p}^i; x_{\tau p}^{-i}) \right\|^2 + 2(j - \tau p) \gamma L_i \sigma_i^2.$$

**Remark.** Given (3), it is tempting to apply Jensen's inequality to the rightmost quantity and then apply Lemma 3.9. However, this results in a bound that is looser than our Lemma 3.8. We need more careful arguments regarding the expectations, which we detail throughout Appendix B.

Proof outline for Theorem 3.4. We combine Lemmas 3.7 and 3.8, and then apply (SCO) to eliminate the  $\|\mathbf{F}(\mathbf{x}_{\tau p})\|^2$  terms to obtain

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] \leq \left(1 + (\tau - 1)\alpha\gamma\right) \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} + (\text{terms proportional to }\sigma^{2}) - \underbrace{\left(2\gamma\tau - \gamma^{2}\tau^{2}\ell - \frac{\gamma^{3}L_{\max}^{2}\tau^{2}(\tau - 1)\ell}{3\alpha}\right)}_{:=C} \langle \mathbb{F}(\mathbf{x}_{\tau p}), \mathbf{x}_{\tau p} - \mathbf{x}_{\star} \rangle \qquad (4)$$

Provided that  $C \geq 0$ , we can upper bound the second line of (4) by  $-C\mu \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^2$  using *(QSM)*. Then we choose  $\alpha = \gamma \tau L_{\max} \sqrt{\frac{\ell \mu}{3}}$  which minimizes the resulting coefficient of  $\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^2$ , and rewrite it in the form  $1 - \gamma \tau \mu \zeta$ . Finally, take expectation over  $\mathbf{x}_{\tau p}$  in (4) and unroll the recursion.

## 4 Numerical Experiments

In this section, we conduct experiments to assess the empirical performance of PEARL-SGD and verify our theory. We focus on three different setups: a minimax game (where n = 2), a multiplayer game with n = 5 players, and a distributed mobile robot control problem with n = 5. Details of all experiments are provided in Appendix C.

### 4.1 Quadratic minimax game

Consider the minimax game  $\min_{u \in \mathbb{R}^d} \max_{v \in \mathbb{R}^d} \mathcal{L}(u, v) = \frac{1}{M} \sum_{m=1}^M \mathcal{L}_m(u, v)$  where  $\mathcal{L}_m(u, v)$  is as below  $(\mathbf{A}_m, \mathbf{B}_m, \mathbf{C}_m \text{ are matrices and } a_m, c_m \text{ are vectors})$ . In this two-player zero-sum game, we have n = 2 with  $f_1(x^1; x^2) = \mathcal{L}(x^1, x^2)$  and  $f_2(x^2; x^1) = -\mathcal{L}(x^1, x^2)$ .

$$\mathcal{L}_m(u,v) := \frac{1}{2} \langle u, \mathbf{A}_m u \rangle + \langle u, \mathbf{B}_m v \rangle - \frac{1}{2} \langle v, \mathbf{C}_m v \rangle + \langle a_m, u \rangle - \langle c_m, v \rangle.$$
(5)

PEARL-SGD with theoretical step-size. Figures 2a and 2b demonstrates the performance of PEARL-SGD using the theoretical step-size  $\gamma = 1/(\ell\tau + 2(\tau-1)L_{\max}\sqrt{\kappa})$  from Theorems 3.3 and 3.4 with  $\tau \in \{1, 2, 4, 5, 8, 20\}$ . Figure 2a shows results from Deterministic PEARL-SGD, while Figure 2b shows the stochastic case, where we repeat each experiment 5 times and plot the mean relative error with standard deviation (shaded region). In the deterministic case, as  $\gamma$ scales down with  $\tau$ , we observe similar linear convergence pattern for all values of  $\tau$ . On the other hand, in the stochastic case, we observe a clear benefit of using larger  $\tau$ ; it reaches smaller relative error within the same number of communication rounds. This is consistent with our theory, which predicts reduced communication cost in the stochastic case.

PEARL-SGD with tuned step-size. This experiment simulates the scenario where we do not know the precise theoretical parameters in advance. For each  $\tau \in \{1, 2, 4, 5, 8, 20\}$ , we tune  $\gamma$  by running PEARL-SGD with each  $\gamma \in \{10^{-1}, 10^{-2}, \ldots, 10^{-6}\}$ , and plot the best relative error  $\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2}/\|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2}$  (y-axis) versus the communication round index p (x-axis). Figure 2c presents results from Deterministic PEARL-SGD. We observe that performance improves as  $\tau$ is increased from 1 to 5, and then degrades. Figure 2d presents results under stochasticity, imposed by mini-batching from the finite sum. Here we observe the lowest relative errors with large values of  $\tau$ , once again demonstrating the advantage of larger synchronization intervals in PEARL-SGD given stochastic gradients.



Figure 2: Performance of PEARL-SGD on quadratic minimax game (5) with different values of  $\tau$ . Figures 2a (deterministic) and 2b (stochastic) show the performance of PEARL-SGD with tight theoretical step-sizes, and Figures 2c (deterministic) and 2d (stochastic) show its performance with empirically tuned step-sizes.

**Performance of PEARL-SGD for different**  $(\gamma, \tau)$ **pairs.** Figure 3 displays the heatmap of relative errors (log-scale) after 100 communication rounds of Deterministic PEARL-SGD on a quadratic minimax game. White and yellow regions indicate divergence/poor performance; darker regions indicate lower relative errors.

Figure 3 reveals a trend: for a fixed  $\gamma$ , PEARL-SGD's performance improves as  $\tau$  increases up to certain threshold, after which it declines and finally diverges. Another key observation is that the dark region of the heatmap (signifying the best performance) takes the shape of a hyperbola. This is consistent with our Theorem 3.3, showing the relationship  $\gamma_{\tau} \propto 1/\tau$  where  $\gamma_{\tau}$  is the optimal step-size choice given  $\tau$  (providing fastest convergence).



Figure 3: Heatmap of relative errors in logarithmic scale.

## 4.2 Quadratic *n*-player game

In our second experiment, we consider an n-player game where the local function of the i-th player is given by

$$f_i(x^i; x^{-i}) := \frac{1}{M} \sum_{m=1}^M f_{i,m}(x^i; x^{-i}), \tag{6}$$

for i = 1, ..., n (with  $d_1 = \cdots = d_n = d$ ). In this setting, each  $f_{i,m}$  takes the form:

$$f_{i,m}(x^i; x^{-i}) = \frac{1}{2} \langle x^i, \mathbf{A}_{i,m} x^i \rangle + \sum_{1 \le j \le n, j \ne i} \langle x^i, \mathbf{B}_{i,j,m} x^j \rangle + \langle a_{i,m}, x^i \rangle,$$

where  $\mathbf{A}_{i,m}, \mathbf{B}_{i,j,m} \in \mathbb{R}^{d \times d}$  and  $a_{i,m} \in \mathbb{R}^d$  for  $m = 1, \ldots, M$ . We set the number of players to n = 5.

**Connection to game & control theory literature.** The above *n*-player game formulation has been often considered in game theory and the closely related control theory literature. It can been used to model applications including the Cournot competition in economics [2], optical networks [91], electricity markets [98], energy consumption control in smart grid [120], mobile robot control [49], etc., and has been utilized in recent works on distributed game optimization (Nash equilibrium search) [99, 119, 67, 113]. In connection with this literature, in Section 4.3, we demonstrate an experiment on a concrete robot control setup from [49].



Figure 4: Performance of PEARL-SGD on the *n*-player game defined by (6), with different  $\tau$ . Figures 4a (deterministic) and 4b (stochastic) show the performance of PEARL-SGD with tight theoretical step-sizes, and Figures 4c (deterministic) and 4d (stochastic) show its performance with empirically tuned step-sizes.

PEARL-SGD with theoretical step-size. Again, we run PEARL-SGD with the theoretical step-size  $\gamma = 1/(\ell \tau + 2(\tau - 1)L_{\max}\sqrt{\kappa})$  of Theorems 3.3 and 3.4 with  $\tau \in \{1, 2, 4, 5, 8, 20\}$ . We set the cocoercivity parameter to  $\ell = L^2/\mu$  following [25], where L and  $\mu$  are explicitly computed Lipschitz constant and strong monotonicity parameter of  $\mathbb{F}$ . Figure 4a displays the results from Deterministic PEARL-SGD; similarly as in Section 4.1, we observe that all values of  $\tau$  produce indistinguishable performance plots. On the other hand, Figure 4b demonstrates that in the stochastic setting, PEARL-SGD with larger synchronization interval  $\tau$  provides a clear benefit of achieving smaller relative error using the same number of communication rounds.

PEARL-SGD with tuned step-size. In this experiment we use tuned step-size for each choice of synchronization interval  $\tau \in \{1, 2, 4, 5, 8, 20\}$ . We use the same  $\gamma$ -grid  $\{10^{-1}, 10^{-2}, \cdots, 10^{-6}\}$  as in Section 4.1 and proceed similarly. Figure 4c shows results from the deterministic setting; the choices  $\tau = 2$  and  $\tau = 20$  outperform the fully communicating case  $\tau = 1$  with step-size tuning. Figure 4d shows results from the stochastic setting, indicating that using larger values of  $\tau$  could lead to higher accuracy levels.

### 4.3 Distributed mobile robot control

Here, we consider a distributed control problem of mobile robots from [49]. This is a multiagent system where each robot has its own objective, depending on the positions  $x^i \in \mathbb{R}^d$ (corresponding to action/strategy in our formulation of multiplayer game) of each *i*-th robot. Specifically, the objective function of the robot *i* is:

$$f_i(\mathbf{x}) = \underbrace{\frac{a_i}{2} \|x^i - x_{\text{anc}}^i\|^2}_{:=J_{i1}(x^i)} + \underbrace{\frac{b_i}{2} \sum_{j=1}^N \|x^i - x^j - h_{ij}\|^2}_{:=J_{i2}(x^i;x^{-i})}$$
(7)

where  $J_{i1}(x^i)$  represents the cost penalizing the distance of agent *i* from the anchor point  $x_{anc}^i \in \mathbb{R}^d$ , and  $J_{i2}(x^i; x^{-i})$  is the cost associated with the relative displacement between the robots' positions. The control problem finds an equilibrium of the *n*-player game, which is the concatenation of all robots' position vectors, ensuring that each robot stays close to  $x_{anc}^i$  while maintaining designated displacement from other robots.



Figure 5: Performance of PEARL-SGD on the distributed mobile robot control problem.

We implement PEARL-SGD with synchronization intervals  $\tau \in \{1, 2, 4, 5, 8, 20\}$  and the theoretical step-size  $\gamma = \frac{1}{\ell \tau + L_{\max}(\tau - 1)\sqrt{\kappa}}$ . Figure 5a shows that with larger values of  $\tau$ , PEARL-SGD achieves better accuracy (in terms of distance to  $\mathbf{x}_{\star}$ ) within a given number of communication rounds. This highlights the potential benefit of using local update steps in solving real-data problems formulated as multiplayer games. Figure 5b displays how the local objective values  $f_i$ are reduced by PEARL-SGD, in the case  $\tau = 5$ .

## 5 Conclusion

In this paper, we introduce Multiplayer Federated Learning (MpFL), a FL framework under setups where clients, strategically acting in their own interests, collaborate through a central server to train models (actions) with the goal of reaching an equilibrium. We propose the PEARL-SGD algorithm handling MpFL, and provide its tight convergence guarantees under heterogeneous setups where each player has distinct objectives and data distributions. We show that PEARL-SGD provides improved communication complexity, reducing the primary overhead in large-scale applications.

Our work offers a number of potential extensions by incorporating the ideas such as Extragradient [56, 34], asynchronous updates [17, 109], gradient compression [3], gradient tracking [90] and algorithmic correction for drifts [50, 84]. We anticipate that our initiation of the study of MpFL will lead to interesting future work including but not limited to these topics.

## Acknowledgments

Sayantan Choudhury acknowledges support from the Chen Family Dissertation Fellowship. Nicolas Loizou acknowledges support from CISCO Research.

### References

- J. Abernethy, K. A. Lai, and A. Wibisono. Last-iterate convergence rates for min-max optimization: Convergence of hamiltonian gradient descent and consensus optimization. *international Conference on Algorithmic Learning Theory*, 2021.
- [2] E. Ahmed and H. Agiza. Dynamics of a cournot game with n-competitors. Chaos, Solitons & Fractals, 9(9):1513-1517, 1998.
- [3] D. Alistarh, D. Grubic, J. Li, R. Tomioka, and M. Vojnovic. QSGD: Communicationefficient SGD via gradient quantization and encoding. *Neural Information Processing* Systems, 2017.
- [4] R. S. Antunes, C. André da Costa, A. Küderle, I. A. Yari, and B. Eskofier. Federated learning for healthcare: Systematic review and architecture proposal. ACM Transactions on Intelligent Systems and Technology (TIST), 13(4):1–23, 2022.
- [5] W. Azizian, I. Mitliagkas, S. Lacoste-Julien, and G. Gidel. A tight and unified analysis of gradient-based methods for a whole spectrum of differentiable games. *International Conference on Artificial Intelligence and Statistics*, 2020.
- [6] A. Beznosikov, V. Samokhin, and A. Gasnikov. Distributed saddle-point problems: Lower bounds, near-optimal and robust algorithms. arXiv preprint arXiv:2010.13112, 2020.
- [7] A. Beznosikov, P. Dvurechenskii, A. Koloskova, V. Samokhin, S. U. Stich, and A. Gasnikov. Decentralized local stochastic extra-gradient for variational inequalities. *Neural Information Processing Systems*, 2022.
- [8] A. Beznosikov, E. Gorbunov, H. Berard, and N. Loizou. Stochastic gradient descentascent: Unified theory and new efficient methods. *International Conference on Artificial Intelligence and Statistics*, 2023.
- [9] R. I. Boţ, E. R. Csetnek, and D.-K. Nguyen. Fast Optimistic Gradient Descent Ascent (OGDA) Method in Continuous and Discrete Time. Foundations of Computational Mathematics, 2023.
- [10] G. W. Brown. Some Notes on Computation of Games Solutions. Rand Corporation, 1949.
- [11] Y. Cai and W. Zheng. Accelerated single-call methods for constrained min-max optimization. International conference on Learning Representations, 2023.
- [12] T. Chavdarova, G. Gidel, F. Fleuret, and S. Lacoste-Julien. Reducing noise in GAN training with variance reduced extragradient. *Neural Information Processing Systems*, 2019.
- [13] S. Choudhury, E. Gorbunov, and N. Loizou. Single-call stochastic extragradient methods for structured non-monotone variational inequalities: Improved analysis under weaker conditions. *Neural Information Processing Systems*, 2024.
- [14] L. Condat and P. Richtárik. RandProx: Primal-dual optimization algorithms with randomized proximal updates. *NeurIPS OPT 2022 Workshop*, 2022.
- [15] C. Daskalakis and I. Panageas. The limit points of (optimistic) gradient descent in minmax optimization. Neural Information Processing Systems, 2018.

- [16] C. Daskalakis, A. Ilyas, V. Syrgkanis, and H. Zeng. Training GANs with optimism. International Conference on Learning Representations, 2018.
- [17] J. Dean, G. Corrado, R. Monga, K. Chen, M. Devin, M. Mao, M. Ranzato, A. Senior, P. Tucker, K. Yang, et al. Large scale distributed deep networks. *Neural Information Processing Systems*, 2012.
- [18] V. Dem'yanov and A. Pevnyi. Numerical methods for finding saddle points. USSR Computational Mathematics and Mathematical Physics, 12(5):11–52, 1972.
- [19] Y. Deng and M. Mahdavi. Local stochastic gradient descent ascent: Convergence analysis and communication efficiency. *International Conference on Artificial Intelligence and Statistics*, 2021.
- [20] Y. Deng, M. M. Kamani, and M. Mahdavi. Adaptive personalized federated learning. arXiv:2003.13461, 2020.
- [21] J. Diakonikolas. Halpern iteration for near-optimal and parameter-free monotone inclusion and strong solutions to variational inequalities. *Conference on Learning Theory*, 2020.
- [22] J. Diakonikolas, C. Daskalakis, and M. I. Jordan. Efficient methods for structured nonconvex-nonconcave min-max optimization. *International Conference on Artificial Intelligence and Statistics*, 2021.
- [23] A. Dieuleveut and K. K. Patel. Communication trade-offs for Local-SGD with large step size. Neural Information Processing Systems, 2019.
- [24] K. Emmanouilidis, R. Vidal, and N. Loizou. Stochastic extragradient with random reshuffling: Improved convergence for variational inequalities. *International Conference on Artificial Intelligence and Statistics*, 2024.
- [25] F. Facchinei and J.-S. Pang. Finite-Dimensional Variational Inequalities and Complementarity Problems. Springer, 2003.
- [26] A. Fallah, A. Mokhtari, and A. Ozdaglar. Personalized federated learning with theoretical guarantees: A model-agnostic meta-learning approach. *Neural Information Processing* Systems, 2020.
- [27] S. Feng, B. Li, H. Yu, Y. Liu, and Q. Yang. Semi-supervised federated heterogeneous transfer learning. *Knowledge-Based Systems*, 252:109384, 2022.
- [28] T. Fiez and L. J. Ratliff. Local convergence analysis of gradient descent ascent with finite timescale separation. *International Conference on Learning Representations*, 2021.
- [29] I. Gemp, R. Patel, Y. Bachrach, M. Lanctot, V. Dasagi, L. Marris, G. Piliouras, S. Liu, and K. Tuyls. Steering language models with game-theoretic solvers. *ICML Agentic Markets Workshop*, 2024.
- [30] G. Gidel, H. Berard, G. Vignoud, P. Vincent, and S. Lacoste-Julien. A variational inequality perspective on generative adversarial networks. *International conference on Learning Representations*, 2019.
- [31] I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial nets. *Neural Information Processing Systems*, 2014.

- [32] E. Gorbunov, F. Hanzely, and P. Richtarik. Local SGD: Unified theory and new efficient methods. *International Conference on Artificial Intelligence and Statistics*, 2021.
- [33] E. Gorbunov, H. Berard, G. Gidel, and N. Loizou. Stochastic extragradient: General analysis and improved rates. *International Conference on Artificial Intelligence and Statistics*, 2022.
- [34] E. Gorbunov, N. Loizou, and G. Gidel. Extragradient method: O (1/k) last-iterate convergence for monotone variational inequalities and connections with cocoercivity. *International Conference on Artificial Intelligence and Statistics*, 2022.
- [35] E. Gorbunov, A. Taylor, and G. Gidel. Last-iterate convergence of optimistic gradient method for monotone variational inequalities. *Neural Information Processing Systems*, 2022.
- [36] R. Gower, O. Sebbouh, and N. Loizou. SGD for structured nonconvex functions: Learning rates, minibatching and interpolation. *International Conference on Artificial Intelligence* and Statistics, 2021.
- [37] R. M. Gower, N. Loizou, X. Qian, A. Sailanbayev, E. Shulgin, and P. Richtárik. Sgd: General analysis and improved rates. *International Conference on Machine Learning*, 2019.
- [38] M. Grudzień, G. Malinovsky, and P. Richtarik. Can 5th generation local training methods support client sampling? Yes! International Conference on Artificial Intelligence and Statistics, 2023.
- [39] F. Haddadpour and M. Mahdavi. On the convergence of local descent methods in federated learning. arXiv:1910.14425, 2019.
- [40] F. Hanzely and P. Richtárik. Federated learning of a mixture of global and local models. arXiv:2002.05516, 2020.
- [41] F. Hanzely, S. Hanzely, S. Horváth, and P. Richtarik. Lower bounds and optimal algorithms for personalized federated learning. *Neural Information Processing Systems*, 2020.
- [42] F. Hanzely, B. Zhao, and m. kolar. Personalized federated learning: A unified framework and universal optimization techniques. *Transactions on Machine Learning Research*, 2023.
- [43] A. Hard, K. Rao, R. Mathews, S. Ramaswamy, F. Beaufays, S. Augenstein, H. Eichner, C. Kiddon, and D. Ramage. Federated learning for mobile keyboard prediction. arXiv preprint arXiv:1811.03604, 2018.
- [44] J. C. Harsanyi and R. Selten. A general theory of equilibrium selection in games. MIT Press Books, 1, 1988.
- [45] Z. Hu and H. Huang. Tighter analysis for ProxSkip. International Conference on Machine Learning, 2023.
- [46] A. P. Jacob, Y. Shen, G. Farina, and J. Andreas. The consensus game: Language model generation via equilibrium search. *International Conference on Learning Representations*, 2024.
- [47] J. Jordan. Three problems in learning mixed-strategy nash equilibria. Games and Economic Behavior, 5(3):368–386, 1993.

- [48] P. Kairouz, H. B. McMahan, B. Avent, A. Bellet, M. Bennis, A. N. Bhagoji, K. Bonawitz, Z. Charles, G. Cormode, R. Cummings, et al. Advances and open problems in federated learning. *Foundations and trends (R) in machine learning*, 14(1–2):1–210, 2021.
- [49] D. Kalyva and H. E. Psillakis. Distributed control of a mobile robot multi-agent system for Nash equilibrium seeking with sampled neighbor information. *Automatica*, 166:111712, 2024.
- [50] S. P. Karimireddy, S. Kale, M. Mohri, S. Reddi, S. Stich, and A. T. Suresh. SCAF-FOLD: Stochastic controlled averaging for federated learning. *International Conference* on Machine Learning, 2020.
- [51] A. Khaled, K. Mishchenko, and P. Richtarik. Tighter theory for local SGD on identical and heterogeneous data. *International Conference on Artificial Intelligence and Statistics*, 2020.
- [52] M. Khodak, M.-F. F. Balcan, and A. S. Talwalkar. Adaptive gradient-based meta-learning methods. *Neural Information Processing Systems*, 2019.
- [53] A. Koloskova, N. Loizou, S. Boreiri, M. Jaggi, and S. Stich. A unified theory of decentralized SGD with changing topology and local updates. *International Conference on Machine Learning*, 2020.
- [54] J. Konečný, H. Brendan McMahan, F. X. Yu, P. Richtárik, A. Theertha Suresh, and D. Bacon. Federated learning: Strategies for improving communication efficiency. arXiv e-prints, pages arXiv-1610, 2016.
- [55] J. Konečný, H. B. McMahan, D. Ramage, and P. Richtárik. Federated optimization: Distributed machine learning for on-device intelligence. arXiv:1610.02527, 2016.
- [56] G. M. Korpelevich. The extragradient method for finding saddle points and other problems. *Ekonomika i Matematicheskie Metody*, 12(4):747–756, 1976.
- [57] D. M. Kreps. Game Theory and Economic Modelling. Oxford University Press, 1990.
- [58] D. M. Kreps, P. Milgrom, J. Roberts, and R. Wilson. Rational cooperation in the finitely repeated prisoners' dilemma. *Journal of Economic theory*, 27(2):245–252, 1982.
- [59] M. Lanctot, V. Zambaldi, A. Gruslys, A. Lazaridou, K. Tuyls, J. Pérolat, D. Silver, and T. Graepel. A unified game-theoretic approach to multiagent reinforcement learning. *Neural Information Processing Systems*, 2017.
- [60] J. Lee, H. Cho, and C. Yun. Fundamental benefit of alternating updates in minimax optimization. *International Conference on Machine Learning*, 2024.
- [61] S. Lee and D. Kim. Fast extra gradient methods for smooth structured nonconvexnonconcave minimax problems. *Neural Information Processing Systems*, 2021.
- [62] C. J. Li, Y. Yu, N. Loizou, G. Gidel, Y. Ma, N. Le Roux, and M. Jordan. On the convergence of stochastic extragradient for bilinear games using restarted iteration averaging. *International Conference on Artificial Intelligence and Statistics*, 2022.
- [63] Q. Li, Z. Wen, Z. Wu, S. Hu, N. Wang, Y. Li, X. Liu, and B. He. A survey on federated learning systems: Vision, hype and reality for data privacy and protection. *IEEE Transactions on Knowledge and Data Engineering*, 35(4):3347–3366, 2021.

- [64] T. Li, A. K. Sahu, A. Talwalkar, and V. Smith. Federated learning: Challenges, methods, and future directions. *IEEE signal processing magazine*, 37(3):50–60, 2020.
- [65] T. Li, A. K. Sahu, M. Zaheer, M. Sanjabi, A. Talwalkar, and V. Smith. Federated optimization in heterogeneous networks. *Annual Conference on Machine Learning and* Systems, 2020.
- [66] T. Li, K. Zhu, N. C. Luong, D. Niyato, Q. Wu, Y. Zhang, and B. Chen. Applications of multi-agent reinforcement learning in future Internet: A comprehensive survey. *IEEE Communications Surveys & Tutorials*, 24(2):1240–1279, 2022.
- [67] Z. Li and Z. Ding. Distributed nash equilibrium searching via fixed-time consensus-based algorithms. In 2019 American Control Conference (ACC), pages 2765–2770, 2019.
- [68] T. Lin, C. Jin, and M. Jordan. On gradient descent ascent for nonconvex-concave minimax problems. *International Conference on Machine Learning*, 2020.
- [69] M. Liu, S. Ho, M. Wang, L. Gao, Y. Jin, and H. Zhang. Federated learning meets natural language processing: A survey. arXiv:2107.12603, 2021.
- [70] Y. Liu, A. Huang, Y. Luo, H. Huang, Y. Liu, Y. Chen, L. Feng, T. Chen, H. Yu, and Q. Yang. FedVision: An online visual object detection platform powered by federated learning. AAAI Conference on Artificial Intelligence, 2020.
- [71] Y. Liu, Y. Kang, C. Xing, T. Chen, and Q. Yang. A secure federated transfer learning framework. *IEEE Intelligent Systems*, 35(4):70–82, 2020.
- [72] Y. Liu, X. Zhang, Y. Kang, L. Li, T. Chen, M. Hong, and Q. Yang. FedBCD: A communication-efficient collaborative learning framework for distributed features. *IEEE Transactions on Signal Processing*, 70:4277–4290, 2022.
- [73] Y. Liu, Y. Kang, T. Zou, Y. Pu, Y. He, X. Ye, Y. Ouyang, Y.-Q. Zhang, and Q. Yang. Vertical federated learning: Concepts, advances, and challenges. *IEEE Transactions on Knowledge and Data Engineering*, 36(7):3615–3634, 2024.
- [74] N. Loizou. Distributionally robust games with risk-averse players. International Conference on Operations Research and Enterprise Systems, 2016.
- [75] N. Loizou, H. Berard, G. Gidel, I. Mitliagkas, and S. Lacoste-Julien. Stochastic gradient descent-ascent and consensus optimization for smooth games: Convergence analysis under expected co-coercivity. *Neural Information Processing Systems*, 2021.
- [76] R. D. Luce and H. Raiffa. Games and Decisions: Introduction and Critical Survey. Courier Corporation, 1989.
- [77] O. Marfoq, G. Neglia, A. Bellet, L. Kameni, and R. Vidal. Federated multi-task learning under a mixture of distributions. *Neural Information Processing Systems*, 2021.
- [78] B. McMahan, E. Moore, D. Ramage, S. Hampson, and B. A. y Arcas. Communicationefficient learning of deep networks from decentralized data. *International Conference on Artificial Intelligence and Statistics*, 2017.
- [79] B. McMahan, E. Moore, D. Ramage, S. Hampson, and B. A. y Arcas. Communicationefficient learning of deep networks from decentralized data. *International Conference on Artificial Intelligence and Statistics*, 2017.

- [80] P. Mertikopoulos and Z. Zhou. Learning in games with continuous action sets and unknown payoff functions. *Mathematical Programming*, 173:465–507, 2019.
- [81] L. Mescheder, S. Nowozin, and A. Geiger. The numerics of GANs. Neural Information Processing Systems, 2017.
- [82] J. Mills, J. Hu, and G. Min. Multi-task federated learning for personalised deep neural networks in edge computing. *IEEE Transactions on Parallel and Distributed Systems*, 33 (3):630–641, 2021.
- [83] K. Mishchenko, D. Kovalev, E. Shulgin, P. Richtárik, and Y. Malitsky. Revisiting stochastic extragradient. *International Conference on Artificial Intelligence and Statistics*, 2020.
- [84] K. Mishchenko, G. Malinovsky, S. Stich, and P. Richtarik. ProxSkip: Yes! Local gradient steps provably lead to communication acceleration! Finally! *International Conference on Machine Learning*, 2022.
- [85] A. Mitra, R. Jaafar, G. J. Pappas, and H. Hassani. Linear convergence in federated learning: Tackling client heterogeneity and sparse gradients. *Neural Information Processing Systems*, 2021.
- [86] A. Mokhtari, A. Ozdaglar, and S. Pattathil. A unified analysis of extra-gradient and optimistic gradient methods for saddle point problems: Proximal point approach. *International Conference on Artificial Intelligence and Statistics*, 2020.
- [87] D. Monderer and L. S. Shapley. Fictitious play property for games with identical interests. Journal of economic theory, 68(1):258–265, 1996.
- [88] J. Nash. Non-cooperative games. Annals of Mathematics, 54(2):286–295, 1951.
- [89] J. F. Nash Jr. Equilibrium points in n-person games. Proceedings of the national academy of sciences, 36(1):48–49, 1950.
- [90] A. Nedic, A. Olshevsky, and W. Shi. Achieving geometric convergence for distributed optimization over time-varying graphs. SIAM Journal on Optimization, 27(4):2597–2633, 2017.
- [91] Y. Pan and L. Pavel. Global convergence of an iterative gradient algorithm for the nash equilibrium in an extended OSNR game. *IEEE INFOCOM*, 2007.
- [92] L. D. Popov. A modification of the Arrow-Hurwicz method for search of saddle points. Mathematical notes of the Academy of Sciences of the USSR, 28(5):845–848, 1980.
- [93] A. Rakhlin and K. Sridharan. Online learning with predictable sequences. Conference on Learning Theory, 2013.
- [94] S. Rakhlin and K. Sridharan. Optimization, learning, and games with predictable sequences. Neural Information Processing Systems, 2013.
- [95] J. Robinson. An iterative method of solving a game. Annals of Mathematics, 54(2): 296–301, 1951.
- [96] E. K. Ryu and W. Yin. Large-scale Convex Optimization: Algorithms & Analyses via Monotone Operators. Cambridge University Press, 2022.

- [97] E. K. Ryu, R. Hannah, and W. Yin. Scaled relative graphs: Nonexpansive operators via 2d euclidean geometry. *Mathematical Programming*, 194(1):569–619, 2022.
- [98] W. Saad, Z. Han, H. V. Poor, and T. Basar. Game-theoretic methods for the smart grid: An overview of microgrid systems, demand-side management, and smart grid communications. *IEEE Signal Processing Magazine*, 29(5):86–105, 2012.
- [99] F. Salehisadaghiani and L. Pavel. Distributed Nash equilibrium seeking: A gossip-based algorithm. Automatica, 72:209–216, 2016.
- [100] T. C. Schelling. The Strategy of Conflict: With a New Preface by the Author. Harvard university press, 1980.
- [101] J. Shamma and G. Arslan. Dynamic fictitious play, dynamic gradient play, and distributed convergence to Nash equilibria. *IEEE Transactions on Automatic Control*, 50(3):312–327, 2005.
- [102] L. S. Shapley. Stochastic games. Proceedings of the national academy of sciences, 39(10): 1095–1100, 1953.
- [103] L. S. Shapley et al. Some topics in two-person games. Rand Corporation, 1963.
- [104] P. Sharma, R. Panda, G. Joshi, and P. Varshney. Federated minimax optimization: Improved convergence analyses and algorithms. *International Conference on Machine Learn*ing, 2022.
- [105] S. Sharma, C. Xing, Y. Liu, and Y. Kang. Secure and efficient federated transfer learning. IEEE International Conference on Big Data, 2019.
- [106] V. Smith, C.-K. Chiang, M. Sanjabi, and A. S. Talwalkar. Federated multi-task learning. *Neural Information Processing Systems*, 2017.
- [107] S. Sokota, R. D'Orazio, J. Z. Kolter, N. Loizou, M. Lanctot, I. Mitliagkas, N. Brown, and C. Kroer. A unified approach to reinforcement learning, quantal response equilibria, and two-player zero-sum games. *International Conference on Learning Representations*, 2023.
- [108] C. Song, Z. Zhou, Y. Zhou, Y. Jiang, and Y. Ma. Optimistic dual extrapolation for coherent non-monotone variational inequalities. *Neural Information Processing Systems*, 2020.
- [109] S. U. Stich. Local SGD converges fast and communicates little. *International Conference* on Learning Representations, 2019.
- [110] S. U. Stich and S. P. Karimireddy. The error-feedback framework: SGD with delayed gradients. Journal of Machine Learning Research, 21(237):1–36, 2020.
- [111] C. T. Dinh, N. Tran, and J. Nguyen. Personalized federated learning with moreau envelopes. *Neural Information Processing Systems*, 2020.
- [112] A. Z. Tan, H. Yu, L. Cui, and Q. Yang. Towards personalized federated learning. IEEE Transactions on Neural Networks and Learning Systems, 34(12):9587–9603, 2023.
- [113] T. Tatarenko, W. Shi, and A. Nedić. Geometric convergence of gradient play algorithms for distributed nash equilibrium seeking. *IEEE Transactions on Automatic Control*, 66 (11):5342–5353, 2021.

- [114] J. Von Neumann and O. Morgenstern. Theory of games and economic behavior: 60th anniversary commemorative edition. In *Theory of Games and Economic Behavior*. Princeton university press, 2007.
- [115] S. Wang, T. Tuor, T. Salonidis, K. K. Leung, C. Makaya, T. He, and K. Chan. Adaptive federated learning in resource constrained edge computing systems. *IEEE Journal on Selected Areas in Communications*, 37(6):1205–1221, 2019.
- [116] J. Xu, B. S. Glicksberg, C. Su, P. Walker, J. Bian, and F. Wang. Federated learning for healthcare informatics. *Journal of healthcare informatics research*, 5:1–19, 2021.
- [117] J. Yang, N. Kiyavash, and N. He. Global convergence and variance reduction for a class of nonconvex-nonconcave minimax problems. *Neural Information Processing Systems*, 2020.
- [118] Q. Yang, Y. Liu, T. Chen, and Y. Tong. Federated machine learning: Concept and applications. ACM Transactions on Intelligent Systems and Technology, 10(2), 2019.
- [119] M. Ye and G. Hu. Distributed Nash equilibrium seeking by a consensus based approach. *IEEE Transactions on Automatic Control*, 62(9):4811–4818, 2017.
- [120] M. Ye and G. Hu. Game design and analysis for price-based demand response: An aggregate game approach. *IEEE Transactions on Cybernetics*, 47(3):720–730, 2017.
- [121] T. Yoon and E. K. Ryu. Accelerated algorithms for smooth convex-concave minimax problems with O(1/k<sup>2</sup>) rate on squared gradient norm. *International Conference on Machine Learning*, 2021.
- [122] T. Yoon, J. Kim, J. J. Suh, and E. K. Ryu. Optimal acceleration for minimax and fixed-point problems is not unique. *International Conference on Machine Learning*, 2024.
- [123] H. Yu, S. Yang, and S. Zhu. Parallel restarted SGD with faster convergence and less communication: Demystifying why model averaging works for deep learning. AAAI Conference on Artificial Intelligence, 2019.
- [124] S. Zhang, S. Choudhury, S. U. Stich, and N. Loizou. Communication-efficient gradient descent-accent methods for distributed variational inequalities: Unified analysis and local updates. *International Conference on Learning Representations*, 2024.
- [125] Y. Zhao, M. Li, L. Lai, N. Suda, D. Civin, and V. Chandra. Federated learning with non-iid data. arXiv:1806.00582, 2018.
- [126] T. Zheng, N. Loizou, P. You, and E. Mallada. Dissipative gradient descent ascent method: A control theory inspired algorithm for min-max optimization. *IEEE Control Systems Letters*, 2024.

# **Supplementary Material**

We organize the Supplementary Material as follows: Section A provides an additional survey of related work. Section B presents the proofs of theoretical results omitted from the main text. Section C provides the details of the experiments omitted from the main paper. Section D provides detailed explanation and interpretation on the theoretical assumptions made in the paper.

## Contents

1	Introduction	1
	1.1 Main contributions	2
2	Multiplayer Federated Learning: Definition and Related Work         2.1       Definition of MpFL         2.2       Comparison with closely related FL frameworks	<b>3</b> 3 5
		5
3	PEARL-SGD: Algorithm and Convergence Guarantees         3.1       Algorithm and assumptions         3.2       Convergence of PEARL-SGD: Deterministic setup         3.3       Convergence of PEARL-SGD: Stochastic setup         3.4       Proof outline	6 6 8 9 10
4	Numerical Experiments         4.1 Quadratic minimax game         4.2 Quadratic n-player game         4.3 Distributed mobile robot control	<b>12</b> 12 13 14
5	Conclusion	15
A	Additional related work	<b>25</b>
в	Omitted proofs for Per-Player Local SGD (PEARL-SGD)B.1Proof of Lemma 3.7B.2General properties and bounds for SGDB.3Proofs of Lemmas 3.9 and 3.8B.4Remaining details in proof of Theorem 3.4B.5Proof of Corollary 3.5B.6Proof of Theorem 3.6	27 28 33 33 35 36
С	Details of Numerical ExperimentsC.1Quadratic minimax gameC.2Quadratic n-player gameC.3Distributed mobile robot control	<b>38</b> 38 38 39
D	<b>Discussion on Theoretical Assumptions</b> D.1 Possible simplification of assumptions: Assuming cocoercivity of $\mathbb{F}$	<b>40</b> 40

## A Additional related work

Game Theory & Equilibrium Computation. Multiplayer games, where multiple players each minimize their own cost function that is affected by the actions of the others, are a long-studied fundamental topic in mathematics and economics [89, 88, 102, 100, 58, 44, 76, 57, 74, 114]. More recently, there has been an increasing interest in the ML community in game-theoretic problems with motivating applications, including adversarial learning [31, 16], multi-agent reinforcement learning (MARL) [59, 66, 107], and language models [29, 46]. This incoming stream of applications has led to the development of novel analyses and insights regarding classical equilibrium-searching algorithms including gradient descent-ascent [18, 68, 117, 28, 126, 75, 60], extragradient [56, 12, 24, 62, 86, 83, 33, 34], optimistic gradient [92, 93, 94, 15, 30, 35] and consensus optimization/Hamiltonian gradient method [81, 5, 1, 75], and even the discovery of new accelerated algorithms for games [21, 121, 61, 11, 9, 122].

Heterogeneity and client drift. One fundamental challenge for theory of Local SGD (FedAvg) is heterogeneity, i.e., varying  $f_i$ 's due to differences in local data distributions [55, 65]. Under such setup, Local SGD is prone to client drift [125, 50] where local descent trajectories head toward distinct minima (of local objectives), and convergence theories require either additional assumptions [115, 123, 39, 65] or technical analyses [51, 53] to control this drift. Some papers, based on theoretical insights, introduced or analyzed correction mechanisms for Local SGD to mitigate client drift [50, 32, 85, 84, 45, 38]. Extension of such ideas to federated minimax optimization was explored in [124].

We note that the *n*-player game setup of MpFL is also fully heterogeneous as each player has distinct (possibly even conflicting) objective functions, and consequently, we have the analogous concept of *player drift*. We refer the readers interested in this topic to the discussion at the end of Section 3.2.

**FL frameworks with individual models.** There are several distinct contexts for FL frameworks (other than personalized FL) where each client learns an individual model. In Vertical FL [118, 72, 73] scenarios, multiple organizations hold distinct features from the common set of samples and they collaborate to train their each local model. In Federated Transfer Learning [105, 71, 27], the participating organizations similarly keep and train local models, but their datasets have heterogeneity over both sample and feature spaces with limited overlaps. Federated Multi-Task Learning [106, 77, 82] extends FL to cases where each client solves different, but related tasks.

Fictitious play. The Fictitious Play (FP) is a classical algorithm, originally proposed by [10] to solve minimax games where each player has a finite action space and plays mixed (randomized) strategies. In FP, each player selects an action that minimizes their expected loss (best response), assuming that the other player plays the empirical (historical) strategy, which is a uniform random mixture of their previously played actions. The convergence of FP to a Nash equilibrium for minimax games was established in [95], but FP fails to converge for general *n*-player (with n > 2) or non-zero-sum games [103, 47], except for particular cases such as all players having identical objectives [87].

While it may appear that PEARL-SGD is conceptually similar to FP (as each player performing multiple local SGD steps can be interpreted as seeking a local approximate best response to others' strategies) the connection is opaque due to some fundamental differences. First, in PEARL-SGD, players make their updates based on only the most recent strategies of other players (not the entire history as in FP). Second, in PEARL-SGD, local SGD steps are not run until players converge to local optima—this results in player drift as we discuss at the end of Section 3.2, and is rather avoided by using step-sizes scaling down with the number of local steps. Third, in the FP setting players are assumed to have finite action spaces and mixed strategies (corresponding to points on a probability simplex), while the MpFL setting deals with continuous action spaces with pure (non-random) strategies. However, despite distinctions, as FP has been previously studied in the distributed n-player game setup [101], exploring the further connection between MpFL and FP could be an interesting direction.

# B Omitted proofs for Per-Player Local SGD (PEARL-SGD)

### B.1 Proof of Lemma 3.7

Note that for  $k = \tau p + 1, \dots, \tau (p + 1)$  (iterations between p-th and (p + 1)-th communications), we have

$$\|\mathbf{x}_{k} - \mathbf{x}_{\star}\|^{2} = \sum_{i=1}^{n} \|x_{k}^{i} - x_{\star}^{i}\|^{2}$$

$$= \sum_{i=1}^{n} \|x_{\tau p}^{i} - x_{\star}^{i} - (x_{\tau p}^{i} - x_{k}^{i})\|^{2}$$

$$= \sum_{i=1}^{n} \left[ \|x_{\tau p}^{i} - x_{\star}^{i}\|^{2} - 2\langle x_{\tau p}^{i} - x_{\star}^{i}, x_{\tau p}^{i} - x_{k}^{i}\rangle + \|x_{\tau p}^{i} - x_{k}^{i}\|^{2} \right]$$

$$= \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} - 2\gamma \sum_{i=1}^{n} \sum_{j=\tau p}^{k-1} \langle x_{\tau p}^{i} - x_{\star}^{i}, g_{j}^{i}\rangle + \sum_{i=1}^{n} \|x_{\tau p}^{i} - x_{k}^{i}\|^{2}, \qquad (8)$$

where for the last equality, we use

$$g_j^i = \nabla f_{i,\xi_j^i}(x_j^i; x_{\tau p}^{-i}), \quad j = \tau p, \dots, k-1, \quad i = 1, \dots, n$$

and

$$x_{j+1}^{i} = x_{j}^{i} - \gamma g_{j}^{i}, \quad j = \tau p, \dots, k-1, \quad i = 1, \dots, n$$

to rewrite  $x_{\tau p}^i - x_k^i = \gamma \sum_{j=\tau p}^{k-1} g_j^i$ . Note that we have

$$\mathbb{E}\left[-\left\langle x_{\tau p}^{i}-x_{\star}^{i},g_{\tau p}^{i}\right\rangle \left|\mathbf{x}_{\tau p}\right]=-\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})\right\rangle,$$

while for the other indices  $j = \tau p + 1, \ldots, k - 1$ , we have the upper bound

$$\begin{split} & \mathbb{E}\left[-\left\langle x_{\tau p}^{i}-x_{\star}^{i},g_{j}^{i}\right\rangle \left|x_{j}^{i}\right]\right] \\ &=-\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{j}^{i};x_{\tau p}^{-i})\right\rangle \\ &=-\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})\right\rangle +\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})-\nabla f_{i}(x_{j}^{i};x_{\tau p}^{-i})\right\rangle \\ &\leq-\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})\right\rangle +\frac{\alpha}{2}\left\|x_{\tau p}^{i}-x_{\star}^{i}\right\|^{2}+\frac{1}{2\alpha}\left\|\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})-\nabla f_{i}(x_{j}^{i};x_{\tau p}^{-i})\right\|^{2} \\ &\leq-\left\langle x_{\tau p}^{i}-x_{\star}^{i},\nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})\right\rangle +\frac{\alpha}{2}\left\|x_{\tau p}^{i}-x_{\star}^{i}\right\|^{2}+\frac{L_{i}^{2}}{2\alpha}\left\|x_{\tau p}^{i}-x_{j}^{i}\right\|^{2} \end{split}$$

where in the fourth line, we use Young's inequality with an arbitrary  $\alpha > 0$  that we determine later. Take expectations of the both sides in (8) (conditioned on  $\mathbf{x}_{\tau p}$ ), and apply the above bound with the tower rule to obtain

$$\mathbb{E}\left[\left\|\mathbf{x}_{k}-\mathbf{x}_{\star}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] \leq \left\|\mathbf{x}_{\tau p}-\mathbf{x}_{\star}\right\|^{2}-2\gamma \sum_{i=1}^{n} \sum_{j=\tau p}^{k-1} \left\langle x_{\tau p}^{i}-x_{\star}^{i}, \nabla f_{i}(x_{\tau p}^{i};x_{\tau p}^{-i})\right\rangle+2\gamma \sum_{i=1}^{n} \sum_{j=\tau p+1}^{k-1} \frac{\alpha}{2} \left\|x_{\tau p}^{i}-x_{\star}^{i}\right\|^{2} +2\gamma \sum_{i=1}^{n} \sum_{j=\tau p+1}^{k-1} \mathbb{E}\left[\frac{L_{i}^{2}}{2\alpha} \left\|x_{\tau p}^{i}-x_{j}^{i}\right\|^{2}\right] \mathbf{x}_{\tau p}\right]+\sum_{i=1}^{n} \mathbb{E}\left[\left\|x_{\tau p}^{i}-x_{k}^{i}\right\|^{2}\right] \mathbf{x}_{\tau p}\right].$$
(9)

Now we apply the identities

$$\sum_{i=1}^{n} \left\langle x_{\tau p}^{i} - x_{\star}^{i}, \nabla f_{i}(x_{\tau p}^{i}; x_{\tau p}^{-i}) \right\rangle = \left\langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p}) \right\rangle, \quad \sum_{i=1}^{n} \left\| x_{\tau p}^{i} - x_{\star}^{i} \right\|^{2} = \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2}$$
$$\sum_{i=1}^{n} \mathbb{E} \left[ \left\| x_{\tau p}^{i} - x_{k}^{i} \right\|^{2} \left\| \mathbf{x}_{\tau p} \right] = \mathbb{E} \left[ \left\| \mathbf{x}_{\tau p} - \mathbf{x}_{k} \right\|^{2} \left\| \mathbf{x}_{\tau p} \right] \right]$$

and the inequality

$$\sum_{i=1}^{n} \sum_{j=\tau p+1}^{k-1} \mathbb{E}\left[\frac{L_i^2}{2\alpha} \left\|x_{\tau p}^i - x_j^i\right\|^2 \left\|\mathbf{x}_{\tau p}\right\|\right] \le \frac{L_{\max}^2}{2\alpha} \sum_{j=\tau p+1}^{k-1} \sum_{i=1}^{n} \mathbb{E}\left[\left\|x_{\tau p}^i - x_j^i\right\|^2 \left\|\mathbf{x}_{\tau p}\right\|\right]$$
$$= \frac{L_{\max}^2}{2\alpha} \sum_{j=\tau p+1}^{k-1} \mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_j\right\|^2 \left\|\mathbf{x}_{\tau p}\right\|\right]$$

to (9) and plug in  $k = \tau(p+1)$ , which gives the desired result.

### B.2 General properties and bounds for SGD

In this section, we present some general properties of stochastic gradient descent (SGD) for an *L*-smooth, convex function  $f \colon \mathbb{R}^m \to \mathbb{R}$ . Suppose that we have a stochastic oracle  $\nabla f_{\xi}(\cdot)$  for the gradient operator  $\nabla f(\cdot)$ , satisfying

$$\mathbb{E}_{\xi}[\nabla f_{\xi}(x)] = \nabla f(x), \quad \mathbb{E}_{\xi}\left[\|\nabla f_{\xi}(x) - \nabla f(x)\|^{2}\right] \le \rho^{2}, \quad \forall x \in \mathbb{R}^{m}$$
(10)

where  $\mathbb{E}_{\xi}$  denotes the expectation with respect to randomness in  $\xi$ . This setup and the subsequent results are the abstractions of intermediate results that we need for the proofs of Lemma 3.9 and Lemma 3.8. Specifically, we will later use the results of this section with

$$f(\cdot) = f_i(\cdot; x_{\tau p}^{-i}), \qquad \rho^2 = \sigma_i^2$$

for each i = 1, ..., n. We make this abstraction to simplify notations and to more effectively convey the key intuitions underlying the analyses.

**Lemma B.1.** Let  $f : \mathbb{R}^m \to \mathbb{R}$  be convex and *L*-smooth. Suppose that a stochastic gradient oracle  $\nabla f_{\xi}(\cdot)$  satisfies (10). Let  $y = x - \gamma \nabla f_{\xi}(x)$ , where  $0 < \gamma \leq \frac{2}{L}$ . Then we have

$$\mathbb{E}_{\xi}\left[\left\|\nabla f(y)\right\|^{2}\right] \leq \left\|\nabla f(x)\right\|^{2} + 2\gamma L\rho^{2}.$$

*Proof.* It is well-known that if f is convex and L-smooth, then  $\nabla f$  is  $\frac{1}{L}$ -cocoercive, i.e., for any  $x, y \in \mathbb{R}^m$ ,

$$\langle x - y, \nabla f(x) - \nabla f(y) \rangle \ge \frac{1}{L} \|\nabla f(x) - \nabla f(y)\|^2$$

By cocoercivity and the step-size condition  $\gamma \leq \frac{2}{L}$ , we have

$$\begin{split} &\frac{\gamma}{2} \|\nabla f(x) - \nabla f(y)\|^2 \\ &\leq \frac{1}{L} \|\nabla f(x) - \nabla f(y)\|^2 \\ &\leq \langle x - y, \nabla f(x) - \nabla f(y) \rangle \\ &= \langle \gamma \nabla f_{\xi}(x), \nabla f(x) - \nabla f(y) \rangle \\ &= \gamma \left( \langle \nabla f_{\xi}(x), \nabla f(x) \rangle - \langle \nabla f(x), \nabla f(y) \rangle + \langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) \rangle \right). \end{split}$$

Taking expectation of the both sides, we obtain

$$\mathbb{E}_{\xi} \left[ \frac{\gamma}{2} \| \nabla f(x) - \nabla f(y) \|^{2} \right]$$
  

$$\leq \mathbb{E}_{\xi} \left[ \gamma \left\langle \nabla f_{\xi}(x), \nabla f(x) \right\rangle - \gamma \left\langle \nabla f(x), \nabla f(y) \right\rangle + \gamma \left\langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) \right\rangle \right]$$
  

$$= \gamma \| \nabla f(x) \|^{2} - \gamma \mathbb{E}_{\xi} \left[ \left\langle \nabla f(x), \nabla f(y) \right\rangle \right] + \gamma \mathbb{E}_{\xi} \left[ \left\langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) \right\rangle \right].$$

Cancelling out the terms and dividing both sides by  $\frac{\gamma}{2}$ , we then have

$$\mathbb{E}_{\xi}\left[\left\|\nabla f(y)\right\|^{2}\right] \leq \left\|\nabla f(x)\right\|^{2} + 2\mathbb{E}_{\xi}\left[\left\langle\nabla f(x) - \nabla f_{\xi}(x), \nabla f(y)\right\rangle\right].$$
(11)

Now observe that

$$\mathbb{E}_{\xi}\left[\langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) \rangle\right] = \mathbb{E}_{\xi}\left[\langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) - \nabla f(x - \gamma \nabla f(x)) \rangle\right]$$

because  $\nabla f(x - \gamma \nabla f(x))$  is a non-random quantity and  $\mathbb{E}_{\xi}[\nabla f(x) - \nabla f_{\xi}(x)] = 0$ . Then

$$\begin{split} & \mathbb{E}_{\xi} \left[ \langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(y) - \nabla f(x - \gamma \nabla f(x)) \rangle \right] \\ &= \mathbb{E}_{\xi} \left[ \langle \nabla f(x) - \nabla f_{\xi}(x), \nabla f(x - \gamma \nabla f_{\xi}(x)) - \nabla f(x - \gamma \nabla f(x)) \rangle \right] \\ &\leq \mathbb{E}_{\xi} \left[ \| \nabla f(x) - \nabla f_{\xi}(x) \| \| \nabla f(x - \gamma \nabla f_{\xi}(x)) - \nabla f(x - \gamma \nabla f(x)) \| \right] \\ &\leq \mathbb{E}_{\xi} \left[ \| \nabla f(x) - \nabla f_{\xi}(x) \| L \| (x - \gamma \nabla f_{\xi}(x)) - (x - \gamma \nabla f(x)) \| \right] \\ &= \gamma L \mathbb{E}_{\xi} \left[ \| \nabla f(x) - \nabla f_{\xi}(x) \|^{2} \right] \\ &= \gamma L \rho^{2}, \end{split}$$

and plugging this into (11) completes the proof.

**Lemma B.2.** Let  $f : \mathbb{R}^m \to \mathbb{R}$  be convex and *L*-smooth and let the stochastic gradient oracle  $\nabla f_{\xi}(\cdot)$  satisfy (10). Let  $x_0 \in \mathbb{R}^m$  be any initial point,  $0 < \gamma \leq \frac{2}{L}$ , and  $x_1, \ldots, x_t$  be a sequence generated by the stochastic gradient descent algorithm

$$x_{s+1} = x_s - \gamma \nabla f_{\xi_s}(x_s)$$

for  $s = 0, \ldots, t - 1$ . Then we have

$$\mathbb{E}\left[\left\|\nabla f(x_s)\right\|^2\right] \le \left\|\nabla f(x_0)\right\|^2 + 2s\gamma L\rho^2$$

for s = 0, ..., t - 1.

*Proof.* Apply Lemma B.1 recursively and use the tower rule (law of total expectation).

**Lemma B.3.** Let  $f: \mathbb{R}^m \to \mathbb{R}$  be *L*-smooth and let  $x_0, \ldots, x_t$  be a sequence generated by stochastic gradient descent

$$x_{s+1} = x_s - \gamma \nabla f_{\xi_s}(x_s)$$

where the stochastic gradient oracle satisfies (10). Let  $\hat{x}_0, \ldots, \hat{x}_t$  be generated via *deterministic* gradient descent

$$\hat{x}_{s+1} = \hat{x}_s - \gamma \nabla f(\hat{x}_s)$$

where  $\hat{x}_0 = x_0$ . Then, provided that  $0 < \gamma \leq \frac{1}{L(t-1)}$ , we have

$$||x_t - \hat{x}_t|| \le 3\gamma \sum_{s=0}^{t-1} ||\nabla f_{\xi_s}(x_s) - \nabla f(x_s)||.$$

**Remark.** This result only assumes *L*-smoothness of f (which is *L*-Lipschitz continuity of  $\nabla f$ ) and does not require convexity.

*Proof.* When t = 1, we have  $||x_t - \hat{x}_t|| = \gamma ||\nabla f_{\xi_0}(x_0) - \nabla f(x_0)||$  as  $x_0 = \hat{x}_0$ . Now assume t > 1. Observe that

$$\begin{aligned} x_t - \hat{x}_t &= (x_{t-1} - \hat{x}_{t-1}) - \gamma \left( \nabla f_{\xi_{t-1}}(x_{t-1}) - \nabla f(\hat{x}_{t-1}) \right) \\ &= (x_{t-1} - \hat{x}_{t-1}) - \gamma \left( \nabla f_{\xi_{t-1}}(x_{t-1}) - \nabla f(x_{t-1}) \right) - \gamma \left( \nabla f(x_{t-1}) - \nabla f(\hat{x}_{t-1}) \right) \end{aligned}$$

and therefore,

$$\begin{aligned} \|x_t - \hat{x}_t\| &\leq \|x_{t-1} - \hat{x}_{t-1}\| + \gamma \left\| \nabla f_{\xi_{t-1}}(x_{t-1}) - \nabla f(x_{t-1}) \right\| + \gamma \left\| \nabla f(x_{t-1}) - \nabla f(\hat{x}_{t-1}) \right\| \\ &\leq (1 + \gamma L) \left\| x_{t-1} - \hat{x}_{t-1} \right\| + \gamma \left\| \nabla f_{\xi_{t-1}}(x_{t-1}) - \nabla f(x_{t-1}) \right\| \end{aligned}$$

where the last inequality uses the L-smoothness assumption. Now unrolling the recursion and using the fact  $||x_0 - \hat{x}_0|| = 0$  we obtain

$$\begin{aligned} \|x_t - \hat{x}_t\| &\leq \sum_{s=0}^{t-1} \gamma (1 + \gamma L)^{t-s-1} \|\nabla f_{\xi_s}(x_s) - \nabla f(x_s)\| \\ &\leq \gamma \left( 1 + \frac{1}{t-1} \right)^{t-1} \sum_{s=0}^{t-1} \|\nabla f_{\xi_s}(x_s) - \nabla f(x_s)\| \\ &\leq 3\gamma \sum_{s=0}^{t-1} \|\nabla f_{\xi_s}(x_s) - \nabla f(x_s)\|. \end{aligned}$$

Lemma B.4. Under the assumptions of Lemma B.3, we have

$$\mathbb{E}\left[\left\langle \nabla f_{\xi_0}(x_0) - \nabla f(x_0), \nabla f(x_t)\right\rangle\right] \le 3t\gamma L\rho^2.$$

*Proof.* Observe that because  $\hat{x}_t$  as defined in Lemma B.3 is a non-random quantity and

$$\mathbb{E}\left[\nabla f_{\xi_0}(x_0) - \nabla f(x_0)\right] = 0,$$

we have

$$\mathbb{E}\left[\left\langle \nabla f_{\xi_0}(x_0) - \nabla f(x_0), \nabla f(\hat{x}_t)\right\rangle\right] = 0.$$

Using this, we can proceed as in the following to obtain the desired bound:

$$\begin{split} & \mathbb{E} \left[ \langle \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}), \nabla f(x_{t}) \rangle \right] \\ &= \mathbb{E} \left[ \langle \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}), \nabla f(x_{t}) - \nabla f(\hat{x}_{t}) \rangle \right] \\ &\leq \mathbb{E} \left[ \| \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}) \| \| \nabla f(x_{t}) - \nabla f(\hat{x}_{t}) \| \right] \\ &\leq \mathbb{E} \left[ \| \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}) \| L \| x_{t} - \hat{x}_{t} \| \right] \\ &\leq 3\gamma L \, \mathbb{E} \left[ \left\| \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}) \right\| \sum_{s=0}^{t-1} \| \nabla f_{\xi_{s}}(x_{s}) - \nabla f(x_{s}) \| \right] \\ &\leq 3\gamma L \, \mathbb{E} \left[ \sum_{s=0}^{t-1} \left( \frac{\| \nabla f_{\xi_{0}}(x_{0}) - \nabla f(x_{0}) \|^{2}}{2} + \frac{\| \nabla f_{\xi_{s}}(x_{s}) - \nabla f(x_{s}) \|^{2}}{2} \right) \right] \\ &\leq 3t \gamma L \rho^{2}. \end{split}$$

Lemma B.5. Under the assumptions of Lemma B.3, we have

$$\mathbb{E}\left[\|x_0 - x_t\|^2\right] \le \gamma^2 \mathbb{E}\left[\left\|\sum_{s=0}^{t-1} \nabla f(x_s)\right\|^2\right] + \gamma^2 t \rho^2 + (t-1)t(t+1)\gamma^3 L \rho^2.$$

*Proof.* In the case t = 1, we have

$$\mathbb{E}\left[\|x_0 - x_1\|^2\right] = \gamma^2 \mathbb{E}_{\xi_0}\left[\|\nabla f_{\xi_0}(x_0)\|^2\right] \le \gamma^2 \rho^2 + \gamma^2 \|\nabla f(x_0)\|^2,$$

which is the desired statement. Now we use induction on t. Suppose that the result holds for any initial point and t steps of SGD. Consider a sequence  $x_0, \ldots, x_{t+1}$  generated via SGD with initial point  $x_0$  and step-size  $\gamma > 0$ . Observe that

$$\mathbb{E}\left[\left\|x_{0}-x_{t+1}\right\|^{2}\right]$$

$$=\gamma^{2}\mathbb{E}\left[\left\|\sum_{s=0}^{t}\nabla f_{\xi_{s}}(x_{s})\right\|^{2}\right]$$

$$=\gamma^{2}\mathbb{E}\left[\left\|\sum_{s=0}^{t-1}\nabla f_{\xi_{s}}(x_{s})\right\|^{2}+\mathbb{E}_{\xi_{t}}\left[2\left\langle\nabla f_{\xi_{t}}(x_{t}),\sum_{s=0}^{t-1}\nabla f_{\xi_{s}}(x_{s})\right\rangle+\|\nabla f_{\xi_{t}}(x_{t})\|^{2}\left|x_{t}\right]\right]$$

$$\leq\mathbb{E}\left[\left\|x_{0}-x_{t}\right\|^{2}\right]+\gamma^{2}\mathbb{E}\left[2\left\langle\nabla f(x_{t}),\sum_{s=0}^{t-1}\nabla f_{\xi_{s}}(x_{s})\right\rangle+\|\nabla f(x_{t})\|^{2}+\rho^{2}\right]$$
(12)

where the third line uses the tower rule. Now observe that for  $s = 0, \ldots, t - 1$ ,

$$\mathbb{E}\left[\langle \nabla f(x_t), \nabla f_{\xi_s}(x_s) \rangle\right] = \mathbb{E}\left[\langle \nabla f(x_t), \nabla f(x_s) \rangle\right] + \mathbb{E}\left[\langle \nabla f(x_t), \nabla f_{\xi_s}(x_s) - \nabla f(x_s) \rangle\right]$$
$$= \mathbb{E}\left[\langle \nabla f(x_t), \nabla f(x_s) \rangle\right] + \mathbb{E}\left[\mathbb{E}\left[\langle \nabla f_{\xi_s}(x_s) - \nabla f(x_s), \nabla f(x_t) \rangle \mid x_s\right]\right]$$
$$\leq \mathbb{E}\left[\langle \nabla f(x_t), \nabla f(x_s) \rangle\right] + 3(t-s)\gamma L\rho^2$$

where the last inequality uses Lemma B.4 (with  $x_s$  regarded as initial point of the stochastic gradient descent). Now we apply this inequality and the induction hypothesis to (12):

$$\begin{split} & \mathbb{E}\left[ \|x_{0} - x_{t+1}\|^{2} \right] \\ & \leq \gamma^{2} \mathbb{E}\left[ \left\| \sum_{s=0}^{t-1} \nabla f(x_{s}) \right\|^{2} + t\rho^{2} + (t-1)t(t+1)\gamma L\rho^{2} \\ & \quad + \sum_{s=0}^{t-1} \left( 2 \left\langle \nabla f(x_{t}), \nabla f(x_{s}) \right\rangle + 6(t-s)\gamma L\rho^{2} \right) + \|\nabla f(x_{t})\|^{2} + \rho^{2} \right] \\ & = \gamma^{2} \left( t\rho^{2} + (t-1)t(t+1)\gamma L\rho^{2} + 3t(t+1)\gamma L\rho^{2} + \rho^{2} \right) \\ & \quad + \gamma^{2} \mathbb{E}\left[ \left\| \sum_{s=0}^{t-1} \nabla f(x_{s}) \right\|^{2} + 2 \left\langle \sum_{s=0}^{t-1} \nabla f(x_{s}), \nabla f(x_{t}) \right\rangle + \|\nabla f(x_{t})\|^{2} \right] \\ & = \gamma^{2} (t+1)\rho^{2} + t(t+1)(t+2)\gamma^{3}L\rho^{2} + \gamma^{2} \mathbb{E}\left[ \left\| \sum_{s=0}^{t} \nabla f(x_{s}) \right\|^{2} \right] \end{split}$$

where for the first equality we use  $\sum_{s=0}^{t-1} 6(t-s) = 3t(t+1)$ . This completes the induction.

**Lemma B.6.** Let  $f : \mathbb{R}^m \to \mathbb{R}$  be convex and *L*-smooth, and let  $x_0 \in \mathbb{R}^m$  be any initial point. Let  $x_1, \ldots, x_t$  be generated by stochastic gradient descent

$$x_{s+1} = x_s - \gamma \nabla f_{\xi_s}(x_s)$$

with  $0 < \gamma \leq \frac{1}{L} \min \left\{ 1, \frac{1}{t-1} \right\}$ . Then  $\mathbb{E} \left[ \|x_0 - x_t\|^2 \right] \leq \gamma^2 t^2 \|\nabla f(x_0)\|^2 + \gamma^2 t (1 + 2(t-1)(t+1)\gamma L)\rho^2.$ 

*Proof.* Lemma B.5 gives

$$\mathbb{E}\left[\|x_0 - x_t\|^2\right] \le \gamma^2 \mathbb{E}\left[\left\|\sum_{s=0}^{t-1} \nabla f(x_s)\right\|^2\right] + \gamma^2 t \rho^2 + (t-1)t(t+1)\gamma^3 L \rho^2.$$
(13)

Next, by Jensen's inequality and Lemma B.2,

$$\mathbb{E}\left[\left\|\sum_{s=0}^{t-1} \nabla f(x_s)\right\|^2\right] \le t \sum_{s=0}^{t-1} \mathbb{E}\left[\left\|\nabla f(x_s)\right\|^2\right]$$
$$\le t \sum_{s=0}^{t-1} \left(\left\|\nabla f(x_0)\right\|^2 + 2s\gamma L\rho^2\right)$$
$$\le t^2 \left\|\nabla f(x_0)\right\|^2 + (t-1)t(t+1)\gamma L\rho^2$$

where the last inequality uses  $\sum_{s=0}^{t-1} 2s = t(t-1) \leq (t-1)(t+1)$ . Applying the above inequality to (13) we obtain the desired result.

### B.3 Proofs of Lemmas 3.9 and 3.8

Proof of Lemma 3.9. Observe that given  $\mathbf{x}_{\tau p}$ , the sequence  $x_{\tau p}^i, \ldots, x_{\tau(p+1)}^i$  is a sequence generated via stochastic gradient descent

$$x_{j+1}^i = x_j^i - \gamma \nabla f_{i,\xi_j^i}(x_j^i; x_{\tau p}^{-i})$$

for the  $L_i$ -smooth convex function  $f_i(\cdot; x_{\tau p}^{-i})$ , with  $x_{\tau p}^i$  as initial point, using the stochastic oracle  $\nabla f_{i,\xi^i}(\cdot; x_{\tau p}^{-i})$  satisfying **(BV)** (unbiased estimator of  $\nabla f_i(\cdot; x_{\tau p}^{-i})$  with variance at most  $\sigma_i^2$ ). Therefore, we can apply Lemma B.2 with

$$f(\cdot) = f_i(\cdot; x_{\tau p}^{-i}), \qquad \rho^2 = \sigma_i^2, \qquad x_0 = x_{\tau p}^i, \qquad x_s = x_j^i, \qquad L = L_i$$

and this immediately proves the desired statement. (Note that s is replaced with  $j - \tau p$  because  $x_j^i$  is obtained by  $j - \tau p$  steps of SGD from  $x_{\tau p}^i$ .)

*Proof of Lemma 3.8.* This is a direct consequence of Lemma B.6 with same choice of  $f, \rho^2, x_0$  and L as in the proof of Lemma 3.9 and  $x_s = x_{\tau p+t}^i$ .

### B.4 Remaining details in proof of Theorem 3.4

Note that the step-size condition of Lemma 3.8 is satisfied by our step-size selection, as  $\gamma < \frac{2}{\ell\tau+2(\tau-1)L_{\max}\sqrt{\kappa}} \leq \frac{1}{L_{\max}(\tau-1)}$  (because  $\kappa \geq 1$ ). Now combine Lemmas 3.7 and 3.8 to obtain  $\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2} \middle| \mathbf{x}_{\tau p} \right]$   $\leq (1 + \alpha\gamma(\tau-1)) \left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2} - 2\gamma\tau \left\langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p}) \right\rangle$   $+ \sum_{j=\tau p+1}^{\tau(p+1)-1} \sum_{i=1}^{n} \frac{\gamma L_{i}^{2}}{\alpha} \left(\gamma^{2}(j-\tau p)^{2} \left\|\nabla f(x_{\tau p}^{i}; x_{\tau p}^{-i})\right\|^{2} + \gamma^{2}(j-\tau p) \left(1 + 2(j-\tau p-1)(j-\tau p+1)\gamma L_{i}\right)\sigma_{i}^{2}\right)$   $+ \sum_{i=1}^{n} \left(\gamma^{2}\tau^{2} \left\|\nabla f(x_{\tau p}^{i}; x_{\tau p}^{-i})\right\|^{2} + \gamma^{2}\tau \left(1 + 2(\tau-1)(\tau+1)\gamma L_{i}\right)\sigma_{i}^{2}\right)$   $\leq (1 + \alpha\gamma(\tau-1)) \left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2} - 2\gamma\tau \left\langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p})\right\rangle + \left(\gamma^{2}\tau^{2} + \frac{\gamma^{3}L_{\max}^{2}\tau^{2}(\tau-1)}{3\alpha}\right) \left\|\mathbb{F}(\mathbf{x}_{\tau p})\right\|^{2}$   $+ \gamma^{2}\tau \left(1 + (\tau-1)\gamma L_{\max}\left(2(\tau+1) + \frac{L_{\max}}{2\alpha} + \frac{\gamma L_{\max}^{2}}{2\alpha}(\tau+1)^{2}\right)\right)\sigma^{2}$ (14)

where for the last inequality, we replace all occurrences of  $L_i$ 's by  $L_{\max} = \max\{L_1, \ldots, L_n\}$  and use the identities

$$\sigma^{2} = \sum_{i=1}^{n} \sigma_{i}^{2}, \quad \|\mathbb{F}(\mathbf{x}_{\tau p})\|^{2} = \sum_{i=1}^{n} \|\nabla f_{i}(x_{\tau p}^{i}; x_{\tau p}^{-i})\|^{2}$$

to eliminate the summations  $\sum_{i=1}^{n}$  and use the following elementary summation results:

$$\sum_{\substack{j=\tau p+1\\j=\tau p+1}}^{\tau(p+1)-1} (j-\tau p)^2 = \frac{(\tau-1)\tau(2\tau-1)}{6} \le \frac{(\tau-1)\tau^2}{3}$$
$$\sum_{\substack{j=\tau p+1\\2}}^{\tau(p+1)-1} (j-\tau p) = \frac{(\tau-1)\tau}{2}$$

and

$$\sum_{j=\tau p+1}^{\tau(p+1)-1} (j-\tau p-1)(j-\tau p)(j-\tau p+1) = \frac{(\tau-2)(\tau-1)\tau(\tau+1)}{2} \le \frac{(\tau-1)\tau(\tau+1)^2}{2}.$$

Now in (14), we use the assumption (SCO) to bound

$$-2\gamma\tau \langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p}) \rangle + \left(\gamma^{2}\tau^{2} + \frac{\gamma^{3}L_{\max}^{2}\tau^{2}(\tau-1)}{3\alpha}\right) \|\mathbb{F}(\mathbf{x}_{\tau p})\|^{2}$$

$$\leq -\left(2\gamma\tau - \ell\left(\gamma^{2}\tau^{2} + \frac{\gamma^{3}L_{\max}^{2}\tau^{2}(\tau-1)}{3\alpha}\right)\right) \langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p}) \rangle$$

$$= -\gamma\tau\left(2 - \gamma\ell\tau - \frac{\gamma^{2}\ell L_{\max}^{2}\tau(\tau-1)}{3\alpha}\right) \langle \mathbf{x}_{\tau p} - \mathbf{x}_{\star}, \mathbb{F}(\mathbf{x}_{\tau p}) \rangle.$$
(15)

Provided that

$$2 - \gamma \ell \tau - \frac{\gamma^2 \ell L_{\max}^2 \tau(\tau - 1)}{3\alpha} \ge 0, \tag{16}$$

we can again upper bound (15) using the assumption (QSM):

$$-\gamma\tau\left(2-\gamma\ell\tau-\frac{\gamma^{2}\ell L_{\max}^{2}\tau(\tau-1)}{3\alpha}\right)\langle\mathbf{x}_{\tau p}-\mathbf{x}_{\star},\mathbb{F}(\mathbf{x}_{\tau p})\rangle$$
  
$$\leq-\gamma\tau\left(2-\gamma\ell\tau-\frac{\gamma^{2}\ell L_{\max}^{2}\tau(\tau-1)}{3\alpha}\right)\mu\|\mathbf{x}_{\tau p}-\mathbf{x}_{\star}\|^{2}.$$

We plug this into (14) and rearrange the terms to obtain

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2} \left\|\mathbf{x}_{\tau p}\right] \leq \left(1 + \alpha\gamma(\tau - 1) - \gamma\tau\left(2 - \gamma\tau\ell - \frac{\gamma^{2}\ell L_{\max}^{2}\tau(\tau - 1)}{3\alpha}\right)\mu\right) \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} + \gamma^{2}\tau\left(1 + (\tau - 1)\gamma L_{\max}\left(2(\tau + 1) + \frac{L_{\max}}{2\alpha} + \frac{\gamma L_{\max}^{2}}{2\alpha}(\tau + 1)^{2}\right)\right)\sigma^{2}.$$
(17)

Now, we optimize the coefficient of the  $\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^2$  term in (17) by taking

$$\alpha = \operatorname*{argmin}_{\alpha>0} \alpha \gamma(\tau-1) + \frac{\gamma^3 \ell L_{\max}^2 \tau^2(\tau-1)\mu}{3\alpha} = \gamma \tau L_{\max} \sqrt{\frac{\ell \mu}{3}}.$$

With this choice of  $\alpha$ , the bound (17) becomes

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2} \left|\mathbf{x}_{\tau p}\right] \\
\leq \left(1 - \gamma\tau\mu\left(2 - \gamma\ell\tau - 2(\tau - 1)\gamma L_{\max}\sqrt{\frac{\ell}{3\mu}}\right)\right) \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} \\
+ \gamma^{2}\tau\left(1 + (\tau - 1)\gamma L_{\max}\left(2(\tau + 1) + \frac{1}{2\gamma\tau\sqrt{\ell\mu/3}} + \frac{L_{\max}(\tau + 1)^{2}}{2\tau\sqrt{\ell\mu/3}}\right)\right)\sigma^{2} \\
\leq (1 - \gamma\tau\mu\zeta) \|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^{2} + \gamma^{2}\tau\sigma^{2}\left(1 + (\tau - 1)\left(4\gamma\tau L_{\max} + \frac{L_{\max}}{2\tau\sqrt{\ell\mu/3}} + \frac{\gamma\tau L_{\max}^{2}}{\sqrt{\ell\mu/3}}\right)\right) (18)$$

where for the last inequality, we use  $\tau + 1 \leq 2\tau$  and make the substitution

$$\zeta = 2 - \gamma \ell \tau - 2(\tau - 1)\gamma L_{\max} \sqrt{\frac{\ell}{3\mu}} = 2 - \gamma \ell \tau - 2(\tau - 1)\gamma L_{\max} \sqrt{\kappa/3}.$$

Note that with our choice  $\alpha = \gamma \tau L_{\max} \sqrt{\frac{\ell \mu}{3}}$  and  $0 < \gamma < \frac{2}{\ell \tau + 2(\tau - 1)L_{\max}\sqrt{\kappa}}$ , the condition (16) is satisfied because

$$2 - \gamma \ell \tau - \frac{\gamma^2 \ell L_{\max}^2 \tau(\tau - 1)}{3\alpha} \ge 2 - \gamma \ell \tau - \frac{\gamma^2 \ell L_{\max}^2 \tau(\tau - 1)}{3\alpha}$$
$$= 2 - \gamma \ell \tau - (\tau - 1) \gamma L_{\max} \sqrt{\frac{\ell}{3\mu}}$$
$$\ge 2 - \gamma \left(\ell \tau + (\tau - 1) L_{\max} \sqrt{\kappa}\right) > 0.$$

Finally, unrolling the recursion (18) using the following simple lemma, with  $a_p = \mathbb{E}\left[\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\|^2\right]$ ,  $A = \tau \mu \zeta$  and

$$B = \tau \sigma^2 \left( 1 + (\tau - 1) \left( 4\gamma \tau L_{\max} + \frac{L_{\max}}{2\tau \sqrt{\ell\mu/3}} + \frac{\gamma \tau L_{\max}^2}{\sqrt{\ell\mu/3}} \right) \right)$$

gives the desired rate. (Note that  $\gamma A = \gamma \tau \mu \zeta \leq \gamma \tau \mu (2 - \gamma \ell \tau) \leq \gamma \ell \tau (2 - \gamma \ell \tau) \leq 1.$ )

**Lemma B.7.** Let  $\gamma, A, B > 0$  with  $\gamma A \leq 1$ . If a sequence  $a_0, \ldots, a_R \in \mathbb{R}$  satisfies

$$a_{p+1} \leq (1 - \gamma A)a_p + \gamma^2 B$$
  
for  $p = 0, \dots, R - 1$ , then  $a_R \leq (1 - \gamma A)^R a_0 + \frac{\gamma B}{A}$ .

Proof of Lemma B.7. As there is nothing to prove if  $\gamma A = 1$ , suppose  $\gamma A < 1$ . Recursively applying the given inequality we have

$$a_R \le (1 - \gamma A)a_{R-1} + \gamma^2 B \le \dots \le (1 - \gamma A)^R a_0 + \gamma^2 B \sum_{p=0}^{R-1} (1 - \gamma A)^p.$$

Now apply the bound  $\sum_{p=0}^{R-1} (1 - \gamma A)^p \leq \sum_{p=0}^{\infty} (1 - \gamma A)^p = \frac{1}{1 - (1 - \gamma A)} = \frac{1}{\gamma A}$  to the above inequality.

### B.5 Proof of Corollary 3.5

First, because  $\eta > \kappa \tau$ , we have

$$\gamma < \frac{1}{\mu\kappa\tau \left(1 + \frac{2L_{\max}}{\sqrt{\ell\mu}}\right)} = \frac{1}{\ell\tau \left(1 + \frac{2L_{\max}}{\sqrt{\ell\mu}}\right)} \le \frac{1}{\ell\tau + 2(\tau - 1)L_{\max}\sqrt{\frac{\ell}{\mu}}} = \frac{1}{\ell\tau + 2(\tau - 1)L_{\max}\sqrt{\kappa}}.$$

Hence we can apply Theorem 3.4. Now observe that  $\zeta > 2 - \gamma \left(\ell \tau + 2(\tau - 1)L_{\max}\sqrt{\kappa}\right) > 1$ , and  $(1-u)^R \leq e^{-uR}$  for u < 1, so

$$(1 - \gamma \tau \mu \zeta)^R \le e^{-\gamma \mu \zeta \tau R} \le e^{-\gamma \mu T} = e^{-2\log\eta} = \frac{1}{\eta^2} = \frac{4(\log\eta)^2 (1 + 2q)^2}{T^2} = \tilde{\mathcal{O}}\left(\frac{(1+q)^2}{T^2}\right)$$

where we use  $T = 2(1+2q)\eta \log \eta$  and remove the factor  $\log \eta < \log T$  within the  $\tilde{\mathcal{O}}$  notation. Next, for the terms proportional to  $\sigma^2$ , we have

$$\begin{split} &\left(1 + (\tau - 1)\left(4\gamma\tau L_{\max} + \frac{L_{\max}}{2\tau\sqrt{\ell\mu/3}} + \frac{\gamma\tau L_{\max}^2}{\sqrt{\ell\mu/3}}\right)\right)\frac{\gamma\sigma^2}{\mu\zeta} \\ &\leq \frac{\gamma\sigma^2}{\mu}\left(1 + \tau\left(4\gamma\tau L_{\max} + \frac{\sqrt{3}q}{2\tau} + \sqrt{3}\gamma\tau L_{\max}q\right)\right)\right) \\ &\leq \frac{\gamma\sigma^2}{\mu}\left(1 + \frac{\sqrt{3}q}{2}\right) + \frac{\gamma^2\tau^2 L_{\max}\sigma^2}{\mu}(4 + \sqrt{3}q) \\ &= \frac{\sigma^2(1 + \sqrt{3}q/2)}{\mu^2\eta(1 + 2q)} + \frac{\tau^2 L_{\max}\sigma^2(4 + \sqrt{3}q)}{\mu^3\eta^2(1 + 2q)^2} \\ &= \tilde{\mathcal{O}}\left(\frac{(1 + q)\sigma^2}{\mu^2T} + \frac{(1 + q)\tau^2 L_{\max}\sigma^2}{\mu^3T^2}\right). \end{split}$$

Combining these with Theorem 3.4 we arrive at the desired conclusion.

### B.6 Proof of Theorem 3.6

Note that we use constant step-size  $\gamma_k \equiv \gamma_{\tau p}$  within each communication round p, i.e., for  $\tau p \leq k \leq \tau (p+1) - 1$ , so we can apply the bound (18) from the proof of Theorem 3.4, provided that

$$\gamma_{\tau p} \le \frac{1}{\ell \tau + 2(\tau - 1)L_{\max}\sqrt{\kappa}}$$

This clearly holds true when  $p < 2(1+2q)\kappa - 1$ , and when  $p \ge 2(1+2q)\kappa - 1$  then

$$\gamma_{\tau p} = \frac{1}{\tau \mu} \frac{2p+1}{(p+1)^2} < \frac{1}{\tau \mu} \frac{2}{p+1} \le \frac{1}{\tau \mu} \frac{1}{(1+2q)\kappa} = \frac{1}{\ell \tau + 2\tau L_{\max} \sqrt{\kappa}}$$

so we see that the step-size condition is satisfied. Furthermore we have

$$\zeta_{\tau p} = 2 - \gamma_{\tau p} \ell \tau - 2(\tau - 1) \gamma_{\tau p} L_{\max} \sqrt{\kappa/3} > 1,$$

so (18), with  $q = \frac{L_{\text{max}}}{\sqrt{\ell\mu}}$  and taking expectation with respect to  $\mathbf{x}_{\tau p}$ , gives

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2}\right] \leq (1 - \gamma_{\tau p}\tau\mu\zeta_{\tau p})\mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2}\right] \\ + \gamma_{\tau p}^{2}\tau\sigma^{2}\left(1 + (\tau - 1)\left(\gamma_{\tau p}\tau L_{\max}(4 + \sqrt{3}q) + \frac{\sqrt{3}}{2\tau q}\right)\right) \\ \leq (1 - \gamma_{\tau p}\tau\mu)\mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2}\right] + (1 + q)\gamma_{\tau p}^{2}\tau\sigma^{2} + 4(1 + q)\gamma_{\tau p}^{3}\tau^{2}(\tau - 1)L_{\max}\sigma^{2}.$$

$$\tag{19}$$

For  $p \ge 2(1+2q)\kappa - 1$ , plugging in  $\gamma_{\tau p} = \frac{1}{\tau \mu} \frac{2p+1}{(p+1)^2}$  we obtain

$$\mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2}\right] \leq \frac{p^{2}}{(p+1)^{2}} \mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2}\right] + \frac{(2p+1)^{2}\sigma^{2}(1+q)}{\tau\mu^{2}(p+1)^{4}} \left(1 + \frac{4(\tau-1)L_{\max}(2p+1)}{\mu(p+1)^{2}}\right).$$

Multiplying  $\tau^2(p+1)^2$  to both sides and upper-bounding  $\frac{2p+1}{p+1} \leq 2$ , we obtain

$$(\tau(p+1))^{2} \mathbb{E}\left[\left\|\mathbf{x}_{\tau(p+1)} - \mathbf{x}_{\star}\right\|^{2}\right] \leq (\tau p)^{2} \mathbb{E}\left[\left\|\mathbf{x}_{\tau p} - \mathbf{x}_{\star}\right\|^{2}\right] + \frac{4(1+q)\tau\sigma^{2}}{\mu^{2}}\left(1 + \frac{8(\tau-1)L_{\max}}{\mu(p+1)}\right).$$

Let  $p_0 = \lfloor 2(1+2q)\kappa - 1 \rfloor$ . Chaining the above inequality for  $p = p_0, \ldots, R-1$  gives

$$\begin{aligned} &(\tau R)^{2} \mathbb{E} \left[ \| \mathbf{x}_{\tau R} - \mathbf{x}_{\star} \|^{2} \right] \\ &\leq (\tau p_{0})^{2} \mathbb{E} \left[ \| \mathbf{x}_{\tau p_{0}} - \mathbf{x}_{\star} \|^{2} \right] + \frac{4(1+q)\tau (R-p_{0})\sigma^{2}}{\mu^{2}} + \frac{32(1+q)\tau (\tau-1)L_{\max}\sigma^{2}}{\mu^{3}} \sum_{p=p_{0}}^{R-1} \frac{1}{p+1} \\ &\leq (\tau p_{0})^{2} \mathbb{E} \left[ \| \mathbf{x}_{\tau p_{0}} - \mathbf{x}_{\star} \|^{2} \right] + \frac{4(1+q)\tau (R-p_{0})\sigma^{2}}{\mu^{2}} + \frac{32(1+q)\tau^{2}L_{\max}\sigma^{2}\log(R/p_{0})}{\mu^{3}} \end{aligned}$$

where we use  $\sum_{p=p_0}^{R-1} \frac{1}{p+1} \leq \int_{p_0}^R \frac{dp}{p} = \log \frac{R}{p_0}$ . Now substitute  $T = \tau R$  using the upper bounds  $\tau(R-p_0) \leq \tau R = T$  and  $\log(R/p_0) \leq \log T$ , we can write

$$T^{2}\mathbb{E}\left[\|\mathbf{x}_{T} - \mathbf{x}_{\star}\|^{2}\right] \leq (\tau p_{0})^{2}\mathbb{E}\left[\|\mathbf{x}_{\tau p_{0}} - \mathbf{x}_{\star}\|^{2}\right] + \frac{4(1+q)T\sigma^{2}}{\mu^{2}} + \frac{32(1+q)\tau^{2}L_{\max}\sigma^{2}\log T}{\mu^{3}}.$$
 (20)

As  $\gamma_k$  is constantly  $\gamma_0 = \frac{1}{\ell \tau (1+2q)}$  over rounds  $p = 0, \ldots, p_0 - 1$ , we can directly apply Theorem 3.4 with  $R = p_0$  and similar simplification of the  $\sigma^2$ -terms as in (19) to bound

$$\mathbb{E}\left[\|\mathbf{x}_{\tau p_{0}} - \mathbf{x}_{\star}\|^{2}\right] \leq \left(1 - \frac{\mu}{\ell(1+2q)}\right)^{p_{0}} \|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2} + \frac{(1+q)\gamma_{0}\sigma^{2}}{\mu}\left(1 + 4\gamma_{0}\tau(\tau-1)L_{\max}\right)$$
$$\leq \left(1 - \frac{1}{\kappa(1+2q)}\right)^{\kappa(1+2q)} \|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2} + \frac{\sigma^{2}}{\ell\mu\tau}\left(1 + \frac{4(\tau-1)L_{\max}}{\ell(1+2q)}\right)$$
$$\leq \frac{\|\mathbf{x}_{0} - \mathbf{x}_{\star}\|^{2}}{e} + \frac{\sigma^{2}}{\ell\mu\tau}\left(1 + \frac{2\tau}{\sqrt{\kappa}}\right),$$

where the second line uses  $p_0 \ge 2(1+2q)\kappa - 1 \ge \kappa(1+2q)$ , and the third line uses the bound  $(1-\frac{1}{t})^t \le \frac{1}{e}$  for t > 1 and  $\frac{4(\tau-1)L_{\max}}{\ell(1+2q)} \le \frac{4q\tau\sqrt{\ell\mu}}{\ell(1+2q)} \le 2\tau\sqrt{\frac{\mu}{\ell}} = \frac{2\tau}{\sqrt{\kappa}}$ . Now plugging this into (20) and dividing both sides by  $T^2$  we obtain

$$\mathbb{E}\left[\left\|\mathbf{x}_{T} - \mathbf{x}_{\star}\right\|^{2}\right]$$

$$\leq \frac{p_{0}^{2}\tau^{2}\left\|\mathbf{x}_{0} - \mathbf{x}_{\star}\right\|^{2}}{eT^{2}} + \frac{\tau p_{0}^{2}\sigma^{2}}{\ell\mu T^{2}}\left(1 + \frac{2\tau}{\sqrt{\kappa}}\right) + \frac{4(1+q)\sigma^{2}}{\mu^{2}T} + \frac{32(1+q)\tau^{2}L_{\max}\sigma^{2}\log T}{\mu^{3}T^{2}}$$

$$\leq \frac{4(1+2q)^{2}\kappa^{2}\tau^{2}\left\|\mathbf{x}_{0} - \mathbf{x}_{\star}\right\|^{2}}{eT^{2}} + \frac{4(1+q)\sigma^{2}}{\mu^{2}T} + \frac{4(1+2q)^{2}\kappa\tau\sigma^{2}}{\mu^{2}T^{2}}\left(1 + \frac{2\tau}{\sqrt{\kappa}}\right) + \frac{32(1+q)\tau^{2}L_{\max}\sigma^{2}\log T}{\mu^{3}T^{2}}$$

which is the desired result.

## C Details of Numerical Experiments

### C.1 Quadratic minimax game

To ensure that the quadratic objective (5) is strongly-convex-strongly-concave and smooth, we randomly generate the symmetric matrices  $\mathbf{A}_m, \mathbf{B}_m, \mathbf{C}_m \in \mathbb{R}^{d \times d}$  to respectively have eigenvalues within the interval  $[\mu_A, L_A]$ ,  $[0, L_B]$  and  $[\mu_C, L_C]$ , respectively, following [75]. We generate the vectors  $a_m, c_m \in \mathbb{R}^d$  to be uniformly random. In all our experiments, we set d = 10 and M = 100, where M represents the size of dataset (number of functions in the finite sum structure). We implement PEARL-SGD with two computational nodes, one corresponding to the  $x^1$ -variable and the other to the  $x^2$ -variable.

## C.2 Quadratic *n*-player game

As before, we set d = 10 and M = 100. The matrices  $\mathbf{A}_{i,m}$  are generated randomly with their eigenvalues in the range  $[\mu_{\mathbf{A}}, L_{\mathbf{A}}]$   $(0 < \mu_{\mathbf{A}} < L_{\mathbf{A}})$ . Similarly, for  $1 \le i < j \le n$ , we generate the matrices  $\mathbf{B}_{i,j,m}$  randomly with their eigenvalues in  $[0, L_{\mathbf{B}}]$ . Notably, we set  $\mathbf{B}_{j,i,m} = -\mathbf{B}_{i,j,m}^{\mathsf{T}}$  for  $1 \le j < i \le n$ . With this condition, we can ensure that the *n*-player game (6) satisfies the (QSM) assumption, regardless of the values of  $\mu_{\mathbf{A}}, L_{\mathbf{A}}$  and  $L_{\mathbf{B}}$ . We show below why this is the case.

Recall that we have

$$f_i(x^1, \dots, x^n) = \frac{1}{2} \left\langle x^i, \mathbf{A}_i x^i \right\rangle + \left\langle a_i, x^i \right\rangle + \sum_{j \neq i} \left\langle x^i, \mathbf{B}_{i,j} x^j \right\rangle$$

for i = 1, ..., n. Differentiating  $f_i$  with respect to  $x^i$ , we get

$$\nabla f_i(x^i; x^{-i}) = \mathbf{A}_i x^i + a_i + \sum_{j \neq i} \mathbf{B}_{i,j} x^j$$

and thus

$$\nabla f_i(x^i; x^{-i}) - \nabla f_i(x^i_\star; x^{-i}_\star) = \left( \mathbf{A}_i x^i + a_i + \sum_{j \neq i} \mathbf{B}_{i,j} x^j \right) - \left( \mathbf{A}_i x^i_\star + a_i + \sum_{j \neq i} \mathbf{B}_{i,j} x^j_\star \right)$$
$$= \mathbf{A}_i(x^i - x^i_\star) + \sum_{j \neq i} \mathbf{B}_{i,j}(x^j - x^j_\star)$$

and

$$\langle \mathbb{F}(\mathbf{x}) - \mathbb{F}(\mathbf{x}_{\star}), \mathbf{x} - \mathbf{x}_{\star} \rangle = \sum_{i=1}^{n} \left\langle \nabla f_{i}(x^{i}; x^{-i}) - \nabla f_{i}(x^{i}_{\star}; x^{-i}_{\star}), x^{i} - x^{i}_{\star} \right\rangle$$

$$= \sum_{i=1}^{n} \left\langle x^{i} - x^{i}_{\star}, \mathbf{A}_{i}(x^{i} - x^{i}_{\star}) \right\rangle + \sum_{i=1}^{n} \sum_{j \neq i} \left\langle x^{i} - x^{i}_{\star}, \mathbf{B}_{i,j}(x^{j} - x^{j}_{\star}) \right\rangle.$$

Now, the double summation term vanishes because for any  $i \neq j$ ,

$$\left\langle x^{i} - x^{i}_{\star}, \mathbf{B}_{i,j}(x^{j} - x^{j}_{\star}) \right\rangle + \left\langle x^{j} - x^{j}_{\star}, \mathbf{B}_{j,i}(x^{i} - x^{i}_{\star}) \right\rangle = 0$$

due to the condition  $\mathbf{B}_{j,i} = -\mathbf{B}_{i,j}^{\mathsf{T}}$ . Therefore, provided that each  $\mathbf{A}_i \succeq \mu I$  we see that  $\mathbb{F}$  satisfies (QSM) (in fact, the same argument with arbitrary  $\mathbf{y}$  in place of  $\mathbf{x}_{\star}$  shows that  $\mathbb{F}$  is  $\mu$ -strongly monotone).

## C.3 Distributed mobile robot control

We follow the same choice of parameter values  $a_i, b_i, x_{anc}^i, h_{ij}$  within (7) from [49]:  $n = 5, d = 1, a_i = 10 + i/6, b_i = i/6,$ 

$$(x_{\text{anc}}^1, x_{\text{anc}}^2, x_{\text{anc}}^3, x_{\text{anc}}^4, x_{\text{anc}}^5) = (1, -4, 8, -9, 13)$$

and

$$(h_{ij})_{\substack{1 \le i \le 5\\1 \le j \le 5}} = \begin{pmatrix} 0 & 5 & -7 & 9 & -8\\ -5 & 0 & -6 & 2 & -9\\ 7 & 6 & 0 & 7 & -4\\ -9 & -2 & -7 & 0 & -2\\ 8 & 9 & 4 & 2 & 0 \end{pmatrix}.$$

We add Gaussian noise with  $\sigma^2 = 100$  to the gradients to simulate stochasticity. In this setup, all our theoretical assumptions are satisfied.

## D Discussion on Theoretical Assumptions

### D.1 Possible simplification of assumptions: Assuming cocoercivity of $\mathbb{F}$

In fact, the convergence of PEARL-SGD can still be proved even if the three assumptions (CVX), (SM) and (SCO) are replaced with the single assumption that  $\mathbb{F} \colon \mathbb{R}^D \to \mathbb{R}^D$  is  $\frac{1}{\ell}$ -cocoercive, i.e.,

$$\langle \mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle \ge \frac{1}{\ell} \|\mathbf{x} - \mathbf{y}\|^2, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^D.$$
 (COCO)

In the subsequent paragraphs, we explain in detail why this is the case. However, we emphasize here that if we derived all convergence theory using (*COCO*) in place of (*CVX*), (*SM*) and (*SCO*) and did not distinguish the role of  $L_i$ 's (the local Lipschitzness parameters from (*SM*)) from that of  $\ell$ , then the resulting convergence rates would have become much more pessimistic (worse) in many cases. Therefore, in our work, we choose to use the current set of assumptions. It allows us to more clearly present the tight dependency of convergence rates to  $L_i$ 's. Also note that assuming (*CVX*), (*SM*) and (*SCO*) is strictly more general than assuming (*COCO*), as we illustrate in Appendix D.2.

(*COCO*) implies (*CVX*), (*SM*) and (*SCO*). Trivially, (*COCO*) implies (*SCO*). Furthermore, if  $\mathbb{F}$  is  $\frac{1}{\ell}$ -cocoercive, then  $\mathbb{F}$  is monotone:

$$\langle \mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle \ge 0, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^{D},$$
 (21)

and  $\ell$ -Lipschitz continuous:

$$\|\mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y})\| \le \ell \|\mathbf{x} - \mathbf{y}\|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^{D}.$$
(22)

In particular, for each i = 1, ..., n, we can take

$$\mathbf{x} = (x^1, \dots, x^{i-1}, x^i, x^{i+1}, \dots, x^n), \quad \mathbf{y} = (x^1, \dots, x^{i-1}, y^i, x^{i+1}, \dots, x^n)$$
(23)

in (21), which gives

$$\left\langle \nabla f_i(x^i; x^{-i}) - \nabla f_i(y^i; x^{-i}), x^i - y^i \right\rangle \ge 0$$

for any  $x^i, y^i \in \mathbb{R}^{d_i}$  and  $x^{-i} \in \mathbb{R}^{D-d_i}$ . That is, the gradient of  $f_i(\cdot; x^{-i}) \colon \mathbb{R}^{d_i} \to \mathbb{R}$  is a monotone operator on  $\mathbb{R}^{d_i}$ , and this implies that  $f_i(\cdot; x^{-i})$  is convex, i.e., *(CVX)* holds. Similarly, plugging the choice (23) into (22) we obtain

$$\left\|\nabla f_i(x^i; x^{-i}) - \nabla f_i(y^i; x^{-i})\right\| \le \ell \left\|x^i - y^i\right\|,$$

showing that (SM) holds, with  $L_i = \ell$ . Therefore, all theorems from the main paper hold under the assumptions (QSM), (COCO), and (BV), with  $\ell$  in place of  $L_{\max}$  in step-size restrictions and convergence rates.

What do we lose by replacing  $L_{\text{max}}$  with  $\ell$ ? The previous discussion shows that we can assume (*COCO*) and replace all occurrences of  $L_{\text{max}}$  with  $\ell$  within the theory. In this case, however, the step-size conditions in Theorems 3.3 and 3.4 become

$$\gamma \le \frac{1}{\ell(\tau + 2(\tau - 1)\sqrt{\kappa})} = \mathcal{O}\left(\frac{1}{\ell\tau\sqrt{\kappa}}\right),\tag{24}$$

and the  $\sqrt{\kappa}$  factor in the denominator is undesirable as it significantly restricts the range of step-size one can use if  $\kappa$  is large. Furthermore, in Corollary 3.5 and Theorem 3.6, the factor q becomes  $\sqrt{\frac{\ell}{\mu}} = \sqrt{\kappa}$ , causing the constant factors in the convergence bounds to potentially become large.

However, there are many cases where  $L_{\text{max}} \ll \ell$ , showing why it is beneficial to keep the dependency on  $L_{\text{max}}$  tight as we do. As an abstract example, when  $\mathbb{F}$  is a generic  $\mu$ -strongly monotone and M-Lipschitz continuous operator, the tight (smallest) cocoercivity parameter one can guarantee on  $\mathbb{F}$  is  $\ell = M^2/\mu$  [25] (tightness can be shown using, e.g., the scaled relative graph theory in [97], [96, Chapter 13]). On the other hand, we have

$$L_{\max} \leq \max_{i=1,\dots,n} \sup_{\mathbf{x}=(x^{i},x^{-i}),\mathbf{y}=(y^{i},x^{-i})} \frac{\|\mathbb{F}(\mathbf{x}) - \mathbb{F}(\mathbf{y})\|}{\|\mathbf{x} - \mathbf{y}\|} \leq \sup_{\mathbf{x}\neq\mathbf{y}} \frac{\|\mathbb{F}(\mathbf{x}) - \mathbb{F}(\mathbf{y})\|}{\|\mathbf{x} - \mathbf{y}\|} = M,$$

i.e., M is an upper bound on  $L_{\max}$  (better than  $\ell$ ). Therefore,  $\ell$  is at least  $\frac{\ell}{M} = \frac{\ell}{\sqrt{\ell\mu}} = \sqrt{\kappa}$  times larger than  $L_{\max}$ , and the largest step-size allowed in Theorems 3.3 and 3.4 is

$$\frac{1}{\ell\tau + 2(\tau - 1)L_{\max}\sqrt{\kappa}} = \Omega\left(\frac{1}{\ell\tau}\right)$$

which is in contrast with (24) where we used  $\ell$  in place of  $L_{\max}$  and obtained  $\sqrt{\kappa}$  times smaller step-size range. Additionally, note that in this case  $q = \frac{L_{\max}}{\sqrt{\ell\mu}} = \frac{L_{\max}}{M} \leq 1$  in Corollary 3.5 and Theorem 3.6, so we can avoid the  $\kappa$ -dependent factors appearing in the convergence results.

We demonstrate another problem class for which  $L_{\max} \ll \ell$ . Consider a two-player matrix game, regularized by adding strongly convex (resp. strongly concave) quadratic terms in x (resp. y):

$$\underset{u \in \mathbb{R}^m}{\text{minimize}} \underset{v \in \mathbb{R}^m}{\text{minimize}} \mathcal{L}(u, v) = \frac{\mu}{2} \|u\|^2 + g^{\mathsf{T}}u + u^{\mathsf{T}}\mathbf{B}v - h^{\mathsf{T}}v - \frac{\mu}{2} \|v\|^2$$
(25)

where  $\mathbf{B} \in \mathbb{R}^{m \times m}, g, h \in \mathbb{R}^{m}$ . In our *n*-player game notation, the first and second players respectively use the objective function  $f_1(x^1; x^2) = \mathcal{L}(x^1, x^2)$  and  $f_2(x^2; x^1) = -\mathcal{L}(x^1, x^2)$ . In this case, the operator  $\mathbb{F}$  is  $\mu$ -strongly monotone with  $\mu$  and *M*-Lipschitz continuous with parameter  $M \geq \sqrt{\|\mathbf{B}\|_2^2 + \mu^2} \geq \|\mathbf{B}\|_2$ . Note that the cocoercivity parameter  $\ell$  is at least *M* (and at most  $M^2/\mu$ ). On the other hand,

$$\nabla f_1(x^1; x^2) = \mu x^1 + g + \mathbf{B} x^2, \quad \nabla f_2(x^2; x^1) = \mu x^2 + h - \mathbf{B}^{\mathsf{T}} x^1,$$

so the Lipschitz constant for  $\nabla f_1$  with  $x^2$  fixed (resp.  $\nabla f_2$  with  $x^1$  fixed) is  $\mu$ , i.e.,  $L_{\max} = \mu$ . Therefore, we have  $L_{\max} \ll \ell$  in this scenario, as strength of regularization  $\mu$  is usually small compared to the smoothness parameter M. The same principle applies to the *n*-player analogue of this setup we use in Section 4.2, where each player has the objective function

$$f_i(x^i; x^{-i}) = \frac{1}{2} \left\langle x^i, \mathbf{A}_i x^i \right\rangle + \left\langle a_i, x^i \right\rangle + \sum_{\substack{1 \le j \le n \\ j \ne i}} \left\langle x^i, \mathbf{B}_{i,j} x^j \right\rangle$$

with  $\mathbf{B}_{j,i} = -\mathbf{B}_{i,j}^{\mathsf{T}}$ . If the quadratic terms are the small regularization terms introduced to induce convergence, so that  $\mathbf{A}_i = \mu \mathbf{I}$  with  $\mu \ll ||\mathbf{B}_{i,j}||_2$ , then we have  $L_{\max} = \mu \ll \max_{i \neq j} ||\mathbf{B}_{i,j}||_2 \leq \ell$ .

## D.2 Example of non-cocoercive $\mathbb{F}$ satisfying (CVX), (SM), (QSM) and (SCO)

Consider the two-player game where two players have the objectives

$$f_1(u;v) = \frac{u^2}{2}\varphi(v)$$
$$f_2(v;u) = \frac{v^2}{2}\varphi(u)$$

where  $\varphi \colon \mathbb{R} \to \mathbb{R}$  is defined by

$$\varphi(t) = \left(\mu + (\ell - \mu)\sin^2 t\right).$$

Here  $0 < \mu < \ell$ , and we use the notation  $\mathbf{x} = (u, v) \in \mathbb{R} \times \mathbb{R}$  instead of  $\mathbf{x} = (x^1, x^2)$  for better readability. Note that because  $\varphi$  satisfies

$$0 < \mu \le \varphi(t) \le \ell, \quad \forall t \in \mathbb{R},$$

 $f_1(\cdot, v) \colon \mathbb{R} \to \mathbb{R}$  is convex (quadratic) for any  $v \in \mathbb{R}$ , and so is  $f_2(u, \cdot)$  for any  $u \in \mathbb{R}$ . Therefore, this game satisfies *(CVX)*. For any  $\mathbf{x} = (u, v)$ , we have

$$\mathbb{F}(\mathbf{x}) = \left(\nabla_u f_1(u; v), \nabla_v f_2(v; u)\right) = \left(u\varphi(v), v\varphi(u)\right).$$

Therefore, the unique equilibrium of the game is  $\mathbf{x}_{\star} = (u_{\star}, v_{\star}) = (0, 0)$ . Additionally, observe that

$$\nabla_{uu} f_1(u; v) = \varphi(v) \in [\mu, \ell], \quad \nabla_{vv} f_2(v; u) = \varphi(u) \in [\mu, \ell].$$

In particular, the both second derivatives are bounded, so (SM) is satisfied. Next, we have

$$\langle \mathbb{F}(\mathbf{x}), \mathbf{x} - \mathbf{x}_{\star} \rangle = u^2 \varphi(v) + v^2 \varphi(u) \ge \mu(u^2 + v^2) = \mu \|\mathbf{x} - \mathbf{x}_{\star}\|^2$$

i.e.,  $\mathbb{F}$  satisfies **(QSM)**. Finally, we have

$$\|\mathbf{F}(\mathbf{x})\|^2 = u^2 \varphi(v)^2 + v^2 \varphi(u)^2 \le \max\{\varphi(v), \varphi(u)\} \left(u^2 \varphi(v) + v^2 \varphi(u)\right) \le \ell \left\langle \mathbf{F}(\mathbf{x}), \mathbf{x} - \mathbf{x}_\star \right\rangle,$$

showing that  $\mathbb{F}$  satisfies (SCO).

On the other hand,  $\mathbb{F}$  is not cocoercive with respect to any parameter; in fact, it is not even Lipschitz continuous nor monotone. Observe that the cross-derivatives

$$\nabla_{uv} f_1(u;v) = (\ell - \mu) u \sin(2v), \quad \nabla_{vu} f_2(u;v) = (\ell - \mu) v \sin(2u)$$

are unbounded over  $\mathbb{R} \times \mathbb{R}$ , so  $\mathbb{F}$  is cannot be Lipschitz continuous with any fixed parameter. Furthermore, we have

$$(D\mathbb{F} + D\mathbb{F}^{\mathsf{T}})(u, v) = \begin{bmatrix} 2\varphi(v) & (\ell - \mu)(u\sin(2v) + v\sin(2u)) \\ (\ell - \mu)(u\sin(2v) + v\sin(2u)) & 2\varphi(u) \end{bmatrix}$$

so with  $u = v = \left(2N + \frac{1}{2}\right)\pi$ , we have

$$det (D\mathbb{F} + D\mathbb{F}^{\mathsf{T}}) (u, v) = 4\varphi^2 \left( \left( 2N + \frac{1}{2} \right) \pi \right) - 4(\ell - \mu)^2 \left( 2N + \frac{1}{2} \right)^2 \pi^2$$
  
=  $4\ell^2 - 4(\ell - \mu)^2 \left( 2N + \frac{1}{2} \right)^2 \pi^2$   
<  $0$ 

provided that N is sufficiently large. As a differentiable operator  $\mathbb{F}$  is monotone if and only if  $D\mathbb{F} + D\mathbb{F}^{\intercal} \succeq 0$  everywhere [96], this shows that  $\mathbb{F}$  is not monotone.

Note that while we provided a two-player example for simplicity, one can easily use the essentially same ideas to construct a non-coccercive *n*-player game satisfying our assumptions with any n > 2. For example, we can choose  $f_i(x^i; x^{-i}) = \frac{(x^i)^2}{2}\varphi(x^{i+1})$  where we identify  $x^{n+1} = x^1$ .