

INTERMINGLED OPEN AND CLOSED MAGNETIC FIELD LINES NEAR THE RADIAL ORIGIN OF THE HELIOSPHERIC
CURRENT SHEET

Forrest Mozer^{1,2}, Andrii Voshchepynets⁴, Oleksiy Agapitov¹, Kyung-Eu Choi¹, Richard Sydora³,

1 Space Sciences Laboratory, University of California, Berkeley, California, USA

2 Physics Department, University of California, Berkeley, California, USA

3. Physics Department, University of Alberta, Edmonton, Canada, T6G 2E1

4. Uzhhorod National University, Uzhhorod, Ukraine

Aims. To investigate the magnetic field geometry and waves in the region near the Sun where the heliospheric current sheet is formed.

Methods. Many Parker Solar Probe heliospheric current sheet crossings were searched for an example below 15 solar radii having apparent open and closed magnetic field lines. One good example was found and its fields and plasmas were analyzed.

Results. The radial component of the magnetic field (the Z-component) measured on the Parker Solar Probe (PSP) changed sign between 12:00 and 13:00 UT on March 30, 2024, when the spacecraft was at 13 solar radii. This sign change may have occurred because the spacecraft crossed the heliospheric current sheet on long open magnetic field lines or it may have occurred because the spacecraft crossed from one side of the equator to the other on much shorter closed magnetic field lines. During this crossing, two distinct regions having different magnetic field geometries, strahl flows, plasma densities, and electric field spectra were observed and identified as regions with open and closed magnetic field lines, respectively. The two regions intermingled on time scales less than 100 milliseconds to create a complex magnetic field geometry. The waves observed in both regions were electrostatic and composed of wide band signatures (in the open field lines regions) and well-structured frequency harmonics (in both the open and closed field lines regions). These harmonic frequencies correlated with the proton plasma frequency, f_{pp} , with the lowest frequency at $\sim 0.1f_{pp}$. This result plus the field aligned electric field perturbations and plasma density fluctuations, require that the observed intense electrostatic mode and associated harmonics were ion acoustic waves. The absence of broadband electrostatic signals in closed field line regions is explained by the lower (than in open field line regions) hot and core electron density and higher ratio of the electron plasma frequency to the electron gyrofrequency, suppressing wave generation.

INTRODUCTION

Maxwell's equations show that magnetic field lines do not end. In the absence of currents in the plasma, such field lines near the Sun are dipole-like and are called closed. Further from the Sun, in the presence of mostly axial currents (called the heliospheric current sheet) the field lines become much longer to form a tail-like structure of so-called open field lines (although they close at distances far from the Sun) that point mostly in the radial direction [Wilcox and Ness, 1965]. Near the Sun, the field lines may have both topologies and lengths, so the plasmas in the two regions may be different because of the processes that do or do not occur, depending on the length of the field line.

The radial component of the magnetic field measured on the Parker Solar Probe (PSP) can reverse sign either because the spacecraft was on shorter closed magnetic field lines as it moved from one side of the equatorial region to the other or because it was on longer open magnetic field lines and it moved across the current sheet. A search has been undertaken to find an example that displays both geometries in a local region. The FIELDS instrument on PSP measured the electric and magnetic field waveforms at frequencies below about one kHz and collected short bursts of data at frequencies to more than one MHz during an event of interest [Bale et al, 2016]. The electron data in this paper were produced by the SWEAP instrument on the same spacecraft [Whittlesey et al, 2020]. All data in this paper are presented in the spacecraft reference frame whose X and Y directions are perpendicular to the Sun-satellite line.

DATA

A magnetic field reversal occurred on Parker Solar Probe orbit 19 when the spacecraft was 13 solar radii from the center of the sun, as illustrated by the radial (Z-component) magnetic field change during the one-hour interval given in Figure 1a. Figure 1b shows that the electric field spectra consisted of several lines plus a broadband signal that disappeared twice during the crossing (the electron cyclotron frequency was about 10,000 Hz at this time). Figures 1e and 1f present the electric and magnetic field spectra obtained to 1 MHz from 17-millisecond bursts that occurred during each of these intervals. The spectrum of Figure 1e was obtained in the region containing the broadband waves while that of Figure 1f was obtained in the absence of such waves. In addition to the larger power at frequencies below a few kHz in Figure 1e as compared to Figure 1f, there are much more complex higher frequency waves in the latter case, in part due to the harmonics. In both cases, no magnetic field signature was observed below about 100 kHz. This may be due to either the magnetic field strength being below the threshold of detection or the high frequency waves being electrostatic. In these figures, the black vertical dashed lines denote the electron cyclotron frequency and the red dashed lines give the proton plasma frequency. It is noted that some of the waves are above and some below the electron cyclotron frequency, as has been previously reported [Shi, et al, 2022].

As seen in the plasma density plot of Figure 1c, the dropouts in the broadband electric field occurred when the plasma density became large, as large as $25,000 \text{ cm}^{-3}$. The coordinated sharp boundaries in the electric field spectra and the plasma density suggest the possibility that the spacecraft passed from one region to another

at these boundary times. If such boundary crossings meant the passage from open to closed magnetic field lines, the magnetic field should change from near-radial on the open field lines to inclined with respect to the radial direction on the closed field lines. Figure 1d, a plot of the angle between the magnetic field and the radial direction, shows that the magnetic field acquired a closed-field-line-like geometry when the density increased and the broadband electric field signal disappeared.




Figure 1. A one-hour interval during which the radial component of the magnetic field in panel 1a changed sign, the electric field spectrum in panel 1b had two different characteristics, the plasma density in panel 1c changed rapidly between two different values, and the magnetic field of panel 1d varied in direction synchronously with the changes of the plasma density and electric field spectrum. Panel 1e presents spectra during the time interval having a broadband electric field and relatively low plasma density while panel 1f provides spectra obtained in the region of large density and missing broadband electric field. The spectra in the latter region are much more complex although they contain less power at frequencies below about 1 kHz. The absence of magnetic field waves (the black curves in panels (1e) and (1f) indicate that all the observed waves are electrostatic. The black dashed vertical line in both figures gives the electron cyclotron frequency while the dashed red line gives the proton plasma frequency.

The interpretation that the spacecraft passed between closed and open magnetic field lines may be further tested by considering the flow direction of the strahl. Figure 2a gives the pitch angle distribution of 314 eV electrons, which are a major component of the strahl. It shows that, at times of large plasma density and missing broadband electric field spectra (12:10-12:20 and 12:42-12:55), the near-zero and near-180-degree pitch angle electron fluxes were both large, indicating that the strahl was propagating in both directions along closed

magnetic field lines. The increases of the electron flux near 90-degree pitch angles in Figure 2a further support the conclusion that the spacecraft was on closed magnetic field lines at these times. Figure 2b shows the changes of the proton spectra at such times.

Figures 2c, 2d, and 2e give 30 second averaged electron velocity distribution functions at the times of the vertical dashed lines in Figure 2a or 2b. In each figure, the four dashed curves give the core, halo, and two strahl distributions. They indicate, respectively, strahl in the minus direction at 12:05 on open field lines, strahl in both directions at 12:43 on closed field lines, and strahl in the plus direction at 12:59 on open field lines having a reversed magnetic field. Panels 2f, 2g, and 2h, provide the proton distributions at these same times.




Figure 2. Panel 2a gives the 314-eV electron pitch angle distribution during the hour of interest, which shows the bi-directional flow of the strahl at 12:10-12:20 and 12:42-12:55, indicating that the spacecraft was on closed magnetic field lines during these times. The enhanced fluxes near 90-degree pitch angles during these times support this interpretation. Panel 2b displays the proton energy distribution while panels 2c, 2d, and 2e display one-minute velocity distributions broken-down to display the core, halo, and two strahl distributions at the times of the vertical dashed lines in panel 2a or 2b. They indicate, respectively, strahl in the minus direction, strahl in both directions and strahl in the plus direction. Panels 2f, 2g, and 2h, provide the proton distributions at these same times.

The interrelation between open and closed magnetic field lines on small spatial scales is studied in Figure 3, during which the spacecraft traveled from a closed-like to an open-like magnetic field geometry in eight seconds because the magnetic field direction in Figure 3a changed from about 45 degrees to nearly zero with respect to the radial direction. In Figure 3b the plasma density is shown to decrease in correlation with the magnetic field change, showing again that the large plasma density was a property of the closed, not open, field lines. Figure 3c plots one component of the electric field collected in a burst that lasted about

3.5 seconds and that is expanded in Figure 3d. During this burst the electric field changed amplitude and character abruptly on time scales shorter than 100 milliseconds, which is a spatial scale about 1000 times the core electron gyroradius. The electric field spectrum to three kHz is shown in Figure 3e while the spectrum of the fluctuations of the plasma density is shown in Figure 3f [Mozer et al, 2022]. Both spectra change on time scales less than 100 milliseconds, showing that the waves were electrostatic. This also suggests that the open and closed magnetic field lines wind around each other on very short spatial scales that show turbulent rather than well-defined static geometries.




Figure 3. An eight-second- interval during which the magnetic field direction of panel 3a showed the spacecraft passage from a closed field-line-like geometry to an open geometry. The density of panel 3b decreased synchronously with the changing magnetic field direction and there was a 3.5 second burst of high-rate electric field measurements shown in panel 3c. Panel 3d shows one component of the electric field during this burst while the electric field spectrum from zero to 3000 Hz is displayed in panel 3e and the plasma density fluctuations from zero to 3000 Hz are illustrated in panel 3f. The variations of the spectra on time scales shorter than 100 milliseconds suggest that the spacecraft crossed the intermingled open and closed magnetic field lines of a complex magnetic field geometry near the radial origin of the heliospheric current sheet. The similarity of the electric field and density spectra again shows that all the observed waves were electrostatic.

Figure 4 provides the relationship between the proton plasma frequency and the electric field harmonics over a three-hour interval surrounding the reversal of the radial magnetic field. The harmonics were visible for the full three-hour interval and their correlation with the proton plasma frequency (the white curve)

ACKNOWLEDGEMENTS

This work was supported by NASA contract NNN06AA01C. The authors acknowledge the extraordinary contributions of the Parker Solar Probe spacecraft engineering team at the Applied Physics Laboratory at Johns Hopkins University. The FIELDS experiment on the Parker Solar Probe was designed and developed under NASA contract NNN06AA01C. Our sincere thanks to J.W. Bonnell, M. Moncuquet, T. Quinn, M. Pulupa, and P. Harvey for providing data analysis material and for managing the spacecraft commanding and data processing, which have become heavy loads thanks to the complexity of the instruments and the orbit.

is seen by the three black curves at 0.1fpp, 0.2fpp, and 0.3fpp which line up nicely with the three lowest harmonics of the electric field. This is the correlation expected for ion acoustic waves. It is noted that the electron gyrofrequency, f_{ce} , the red curve in Figure 4, happens to be roughly anti-correlated with the fpp, so the harmonics do not correlate at all with f_{ce} .




Figure 4. The electric field spectra during the three-hour interval surrounding the reversal of the radial magnetic field. The white curve gives the proton plasma frequency, f_{pp} , while the three black curves display 0.1fpp, 0.2fpp, and 0.3fpp, respectively. The correlation between the harmonic frequencies and the fractional proton plasma frequency is emphasized because it identifies the waves as ion acoustic waves.

SUMMARY

The above observations of two distinct regions containing widely different plasma densities, electric field spectra, X-Y components of the magnetic field and strahl electron flows suggest that the spacecraft passed through both open and closed magnetic field lines in close proximity. The two regions intermingled on time scales less than 100 milliseconds to create a complex magnetic field geometry near the radial origin of the heliospheric current sheet. The waves observed in both regions were electrostatic and composed of a wideband signal (in the open field lines regions) and well-structured harmonics of the main mode (in both the open and closed field lines regions). The harmonic frequencies correlated with the ion plasma frequency with the main frequency of ~ 0.1 fpp. These waves had field aligned polarizations and they appeared also as perturbations of the plasma density. These properties suggest that the observed electrostatic mode and associated harmonics are ion acoustic waves. The absence of broadband electrostatic noise in the regions of closed field lines is explained by the lower (than in open field lines regions) ratio of hot and core electron density and higher w_{pe}/w_{ce} ratio suppressing wave generation.

REFERENCES

Bale, S. D., Goetz, K., Harvey, P. R., et al. 2016, *SSRv*, 204,

Mozer, F.S., Bale, S.D., Kellogg, P.J., Larson, D.E., Livi, R., and Romeo, O., 2022, "An Improved Technique for Measuring Plasma Density to High Frequencies on the Parker Solar Probe", *The Astrophysical Journal*, 926:220, 2022, <https://doi.org/10.3847/1538-4357/ac4f42>

Whittlesey, P. L., Larson, D. E., Kasper, J. C., Halekas, J., Abatcha, M., Abiad, R., Berthomier, M., Case, A. W., Chen, J., Curtis, D. W. et al., 2020, The solar probe analyzers—Electrons on the Parker Solar Probe, *Astrophys. J. Suppl. Ser.* 246(2), 74

Wilcox, J. M., & Ness, N. F. (1965). *Quasi-stationary corotating structure in the interplanetary medium*. *Journal of Geophysical Research*, 70(23), 5793-5805

Shi, C., Zhao, J., Malaspina, D.M., Bale, S.D., Dong, X., Wang, T. and Wu, D., (2022), *The Astrophysical Journal Letters*, 926:L3 <https://doi.org/10.3847/2041-8213/ac4d37>