arXiv:2501.09199v2 [math.CA] 30 Jan 2025

WEIGHTED EQUILIBRIUM AND THE FLOW OF DERIVATIVES OF POLYNOMIALS

ANDREI MARTÍNEZ-FINKELSHTEIN AND EVGENII A. RAKHMANOV

Dedicated to Ed Saff on the occasion of his 80th birthday, in recognition of his profound contributions to approximation theory and his enduring inspiration as a mentor, colleague, and friend.

ABSTRACT. Given a sequence of polynomials Q_n of degree n with zeros on [-1, 1], we consider the triangular table of derivatives $Q_{n,k}(x) = d^k Q_n(x)/dx^k$. Under the assumption that the sequence $\{Q_n\}$ has a weak* limiting zero distribution (an empirical distribution of zeros) given by the arcsine law, we show that as $n, k \to \infty$ such that $k/n \to t \in [0, 1)$, the zero-counting measure of the polynomials $Q_{n,k}$ converges to an explicitly given measure μ_t . This measure is the equilibrium measure of [-1, 1] of size 1 - t in an external field given by two mass points of size t/2 fixed at ± 1 . The main goal of this paper is to provide a direct potential theory proof of this fact.

1. INTRODUCTION

Given a sequence of (monic) polynomials Q_n , deg $Q_n = n$, $n \in \mathbb{N}$, with zeros on the interval [-1, 1], we consider the triangular table of their iterated derivatives

$$Q_{n,k}(x) \stackrel{\text{def}}{=} \frac{(n-k)!}{n!} \frac{d^k}{dx^k} Q_n(x) = x^{n-k} + \text{lower degree terms}, \quad k = 0, 1, \dots, n-1.$$
(1.1)

The main result of this paper is the following theorem:

Theorem 1.1. Let Q_n be a sequence of polynomials, deg $Q_n = n$, whose zeros belong to the interval $[-1,1] \subset \mathbb{R}$, and such that the sequence of probability zero-counting measures¹

$$\sigma_{n,0} = \frac{1}{n} \chi\left(Q_n\right), \quad \chi(Q_n) \stackrel{\text{def}}{=} \sum_{Q_n(x)=0} \delta_x,$$

has a weak-* limit μ_0 , given by the arc-sine distribution, namely,

$$d\mu_0(x) = \frac{dx}{\pi\sqrt{1-x^2}}, \quad x \in (-1,1).$$
(1.2)

Date: January 31, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary: 30C15; Secondary: 30C10; 31A05.

Key words and phrases. Polynomials; Zeros; Asymptotic distribution of zeros; Weak convergence; Equilibrium measure; Weighted equilibrium .

¹ In all zero-counting measures the zeros are considered with account of their multiplicity.

Let

$$\sigma_{n,k} \stackrel{\text{def}}{=} \frac{1}{n} \chi\left(Q_{n,k}\right) = \frac{1}{n} \sum_{Q_{n,k}(x)=0} \delta_x.$$
(1.3)

Then for any $t \in [0,1)$ and any sequence k_n of natural numbers with $k_n/n \to t$, the sequence σ_{n,k_n} has a weak-* limit μ_t , where μ_t is absolutely continuous and

$$d\mu_t(x) = \frac{1}{\pi} \frac{\sqrt{1 - t^2 - x^2}}{1 - x^2} dx, \qquad x \in \left[-\sqrt{1 - t^2}, \sqrt{1 - t^2}\right],\tag{1.4}$$

A simple illustration of this result is given by the sequence $Q_n = P_n^{(\alpha,\beta)}$, $\alpha, \beta > -1$, of the (monic) Jacobi polynomials, see [2, §18.3]. The well-known property [2, §18.9]

$$Q'_{n}(x) = nP_{n-1}^{(\alpha+1,\beta+1)}(x)$$

allows for the explicit computation of the weak-* limit μ_t of the sequence (1.3), as $k/n \to t$, in this case. This yields the expression in (1.4). Theorem 1.1 shows that this behavior is not intrinsic to Jacobi polynomials, but it is generic for any sequence of polynomials whose initial distribution is given by the arc-sine law.

Theorem 1.1 can be derived, in particular, from the results contained in the recent paper [3], where the description of the Cauchy transform of the limiting measure μ_t (under very general assumptions) is done in terms of solutions of the inviscid Burgers equation. The main contribution of this paper is the potential-theoretic approach, which characterizes the flow in terms of the equilibrium measure in an external field generated by two mass points. Since one of the main steps in our analysis is an estimate of the Cauchy integral expressing the polynomials $Q_{n,k}$, it is worth mentioning a recent work of [1] on zeros of polynomials generated by a Rodrigues-type formula.

2. Proof of Theorem 1.1

Let us start by setting up the following notation: for $0 < \tau \leq 1$, denote $\mathcal{M}_{\tau}(F)$ the set of Borel measures of total mass τ supported on $F \subset \mathbb{R}$.

Given a finite compactly supported Borel measure μ on \mathbb{R} , we define its *logarithmic potential* as

$$U^{\mu}(z) \stackrel{\text{def}}{=} \int \log \frac{1}{|z-y|} d\mu(y).$$
(2.1)

For our purposes, it will be sufficient to consider measures on an interval of the real line, without loss of generality, $\Delta \stackrel{\text{def}}{=} [-1,1] \subset \mathbb{R}$. Recall the following weighted equilibrium problem that we state in its simplified form, see e.g. [4]: given $\tau \in (0,1]$ and a continuous function $\varphi : (-1,1) \to \mathbb{R}$ (the *external field*), there is the unique equilibrium measure $\lambda = \lambda(\tau, \varphi, \Delta) \in \mathcal{M}_{\tau}(\Delta)$ minimizing the weighted energy

$$\int U^{\mu} d\mu + 2 \int \varphi \, d\mu$$

in the class $\mathcal{M}_{\tau}(\Delta)^2$. It is known that there is a constant $m_{\tau,\varphi}$ such that

$$U^{\lambda}(x) + \varphi(x) \begin{cases} = m_{\tau,\varphi}, & x \in \operatorname{supp}(\lambda), \\ \geqslant m_{\tau,\varphi}, & x \in \Delta. \end{cases}$$
(2.2)

The equilibrium measure λ is also characterized by the following property:

$$m_{\tau,\varphi} = \min_{x \in \Delta} \left(U^{\lambda}(x) + \varphi(x) \right) = \max_{\mu \in \mathcal{M}_{\tau}(\Delta)} \min_{x \in \Delta} \left(U^{\mu}(x) + \varphi(x) \right).$$
(2.3)

In other words, for any $\mu \in \mathcal{M}_{\tau}(\Delta)$,

$$\min_{x \in \Delta} \left(U^{\mu}(x) + \varphi(x) \right) \ge m_{\tau,\varphi} \text{ quasi-everywhere } \Rightarrow \quad \mu = \lambda(\tau,\varphi,\Delta).$$
(2.4)

This will be the key property we will use to prove Theorem 1.1, taking into account the following well-known fact:

Lemma 2.1. For $t \in [0, 1)$, measure μ_t in (1.4) is the equilibrium measure $\lambda(1-t, \varphi, \Delta) \in \mathcal{M}_{1-t}(\Delta)$ in the external field

$$\varphi(x) = \varphi_t(x) \stackrel{\text{def}}{=} \frac{t}{2} \log \frac{1}{|x^2 - 1|}, \quad x \in (-1, 1).$$
 (2.5)

For the proof, see e.g. [4, Examples IV.1.18 and IV.6.2].

Lemma 2.2. Let φ be a continuous external field on Δ , $\tau \in (0,1]$, $\lambda = \lambda(\tau, \varphi, \Delta) \in \mathcal{M}_{\tau}(\Delta)$ the equilibrium measure. If the support $S \stackrel{\text{def}}{=} \operatorname{supp}(\lambda_{\tau,\varphi}) \subset \Delta$ is an interval then

$$m_{\tau,\varphi} = u_{\varphi}(\infty) + \tau \log \frac{1}{\operatorname{cap}(S)},$$
(2.6)

where u_{φ} is the solution of the Dirichlet problem in $\overline{\mathbb{C}} \setminus S$ with the boundary values φ , and $\operatorname{cap}(S) = \operatorname{length}(S)/4$ is the logarithmic capacity of S.

In particular, with $\varphi = \varphi_t$ given in (2.5),

$$m_{1-t,\varphi} = \log(2) - \frac{1+t}{2}\log(1+t) - \frac{1-t}{2}\log(1-t).$$
(2.7)

Proof. Let Ψ be the conformal mapping of $\overline{\mathbb{C}} \setminus S$ onto the exterior of the unit circle, such that $\Psi(\infty) = \infty$, and normalized by $\Psi'(\infty) > 0$. Then $\Psi'(\infty) = 1/\operatorname{cap}(S)$. Consider the function

$$H(z) \stackrel{\text{def}}{=} U^{\lambda}(z) + \tau \log |\Psi(z)| + u_{\varphi}(z).$$

² We could clearly reformulate the problem in the class $\mathcal{M}_1(\Delta)$; the parameter τ is introduced here for convenience.

By (2.2), H is harmonic in $\overline{\mathbb{C}} \setminus S$ and identically equal to $m_{\tau,\varphi}$ on S, which implies that $H(z) = m_{\tau,\varphi}$ for all $z \in \overline{\mathbb{C}} \setminus S$. In particular,

$$m_{\tau,\varphi} = \lim_{z \to \infty} \left(U^{\lambda}(z) + \tau \log |\Psi(z)| + u_{\varphi}(z) \right)$$
$$= \lim_{z \to \infty} \left(U^{\lambda}(z) + \tau \log |z| \right) + \lim_{z \to \infty} \tau \log \left| \frac{\Psi(z)}{z} \right| + u_{\varphi}(\infty)$$

and (2.6) follows from the well-known properties of the logarithmic potential and the conformal mapping Ψ .

By Lemma 2.1 and formula (1.4), for $\varphi = \varphi_t$, we have $\tau = 1 - t$, S = [-s, s] with $s = \sqrt{1 - t^2}$. In this case, $\Psi(z) = \Phi(z/s)$, where $\Phi(z)$ is the inverse Zhukovski map,

$$\Phi(z) = z + (z^2 - 1)^{1/2}, \qquad (2.8)$$

and we take the single-valued branch of the square root in $\mathbb{C}\backslash\Delta$ fixed by $(z^2 - 1)^{1/2} > 0$ for z > 1. On the other hand,

$$u_{\varphi}(z) = \operatorname{Re}\left(\frac{\sqrt{z^2 - s^2}}{\pi} \int_{-s}^{s} \frac{\varphi_t(x)}{\sqrt{s^2 - x^2}} \frac{dx}{z - x}\right),$$

so that

$$\begin{aligned} u_{\varphi}(\infty) &= \frac{1}{\pi} \int_{-s}^{s} \frac{\varphi_{t}(x)}{\sqrt{s^{2} - x^{2}}} dx \\ &= \frac{t}{2} \left(\frac{1}{\pi} \int_{-s}^{s} \log \frac{1}{1 - x} \frac{dx}{\sqrt{s^{2} - x^{2}}} + \frac{1}{\pi} \int_{-s}^{s} \log \frac{1}{1 + x} \frac{dx}{\sqrt{s^{2} - x^{2}}} \right) \\ &= \frac{t}{\pi} \int_{-s}^{s} \log \frac{1}{1 - x} \frac{dx}{\sqrt{s^{2} - x^{2}}} = t U^{\lambda}(1), \end{aligned}$$

where $\lambda = \lambda(1, 0, [-s, s])$ is the equilibrium (Robin) measure of the interval [-s, s]. Since

$$U^{\lambda}(z) = -\log\left(\exp([-s,s])\right) - \log|\Phi(z/s)| = \log\left(\frac{2}{s}\right) - \log|\Phi(z/s)|,$$

we get that

$$u_{\varphi}(\infty) = t\left(\log\left(\frac{2}{s}\right) - \log|\Phi(1/s)|\right) = t\left(\log\left(\frac{2}{s}\right) - \frac{1}{2}\log(1+t) + \frac{1}{2}\log(1-t)\right).$$

In consequence,

$$m_{\tau,\varphi} = -t \log |\Phi(1/s)| + \log \left(\frac{2}{s}\right)$$

= $\log(2) - \frac{1+t}{2} \log(1+t) - \frac{1-t}{2} \log(1-t).$

Remark 2.3. Clearly, formula (2.6) is valid under much milder assumptions on φ . We have stated the restricted version here, sufficient for our purposes, for the sake of simplicity.

Now we turn to the sequence of zero-counting measures $\{\sigma_{n,k_n}\}_{n\in\Lambda}$, defined in (1.3), with the assumption that

$$\sigma_{n,0} \xrightarrow{*} \mu_0 \in \mathcal{M}_1(\Delta), \quad n \to \infty.$$
(2.9)

The following lemma is valid even if μ_0 is not the arc-sine distribution on Δ :

Lemma 2.4. Let $\{k_n\}, n \in \mathbb{N}$, be a sequence such that

$$\lim_{n} \frac{k_n}{n} = t \in [0, 1) \tag{2.10}$$

exists, and let $\Lambda \subseteq \mathbb{N}$ be such that the following weak-* limit exists:

$$\sigma_{n,k_n} \xrightarrow{*} \nu \in \mathcal{M}_{1-t}(\Delta), \quad n \in \Lambda.$$
 (2.11)

Then, for any $\zeta \in \Delta$ where U^{ν} is continuous, and quasi-everywhere on Δ , for any Jordan contour Γ enclosing Δ ,

$$U^{\nu}(\zeta) \ge -c_t + \min_{z \in \Gamma} \left(U^{\mu_0}(z) + t \log |z - \zeta| \right),$$
(2.12)

where

$$c_t \stackrel{\text{def}}{=\!\!=} t \log t + (1-t) \log(1-t). \tag{2.13}$$

Proof. Let first ζ be in the bounded complement of $\mathbb{C} \setminus (\Gamma \cup \Delta)$. We have the integral representation

$$Q_{n,k}(\zeta) = \binom{n}{k}^{-1} \frac{1}{2\pi i} \oint_{\Gamma} \frac{Q_n(z)}{(z-\zeta)^{k+1}} dz,$$

from which we get that

$$U^{\sigma_{n,k}}(\zeta) = -\frac{1}{n} \log |Q_{n,k}(\zeta)| = \frac{1}{n} \log \binom{n}{k} + \frac{1}{n} \log(2\pi) - \frac{1}{n} \log \left| \oint_{\Gamma} \frac{Q_n(z)}{(z-\zeta)^{k+1}} dz \right|.$$

By Stirling's formula,

$$\lim_{n} \frac{1}{n} \log \binom{n}{k_n} = -c_t$$

On the other hand,

$$\begin{aligned} \frac{1}{n} \log \left| \oint_{\Gamma} \frac{Q_n(z)}{(z-\zeta)^{k+1}} dz \right| &\leq \frac{1}{n} \log \left(\max_{z \in \Gamma} \left| \frac{Q_n(z)}{(z-\zeta)^{k+1}} \right| \cdot \operatorname{length}(\Gamma) \right) \\ &= \max_{z \in \Gamma} \left(-U^{\sigma_{n,0}}(z) - \frac{k+1}{n} \log |z-\zeta| \right) + \frac{1}{n} \log \left(\operatorname{length}(\Gamma) \right) \\ &= -\min_{z \in \Gamma} \left(U^{\sigma_{n,0}}(z) + \frac{k+1}{n} \log |z-\zeta| \right) + \frac{1}{n} \log \left(\operatorname{length}(\Gamma) \right). \end{aligned}$$

Thus, we conclude that

$$U^{\sigma_{n,k_n}}(\zeta) \ge -c_t + \min_{z \in \Gamma} \left(U^{\sigma_{n,0}}(z) + \frac{k_n + 1}{n} \log|z - \zeta| \right) + o(1), \quad n \to \infty.$$

$$(2.14)$$

Taking limits in both sides and using assumptions (2.9) and (2.11) we conclude that (2.12) is valid for ζ in the bounded complement of $\mathbb{C} \setminus (\Gamma \cup \Delta)$. By the the Continuity Theorem [4, Theorem A.IV.4.6], we can extend this inequality to any $\zeta \in \Delta$ where the potential U^{ν} is continuous. Moreover, by the Lower Envelope Theorem [4, Theorem A.IV.4.8], it holds quasi-everywhere on Δ .

We need to estimate the right-hand side of (2.12) in the case when μ_0 is the arcsine distribution given in (1.2). In this case,

$$U^{\mu_0}(z) = \log(2) - \log|\Phi(z)|, \qquad (2.15)$$

with Φ given in (2.8).

Also, following [4], we use the notation

$$\inf_{z \in \Delta} h(z)$$

"

for essential infimum, that is, the largest number L such that on Δ the real function h takes values smaller than L only on a set of zero capacity.

Lemma 2.5. With the assumptions of Lemma 2.4,

$$\lim_{x \in [-1,1]} (U^{\nu} + \varphi)(x) \ge m_{1-t,\varphi},$$
(2.16)

where $m_{1-t,\varphi}$ is given in (2.7).

Proof. Fix $\zeta \in (-1, 1)$ where the potential U^{ν} is continuous; as a contour Γ in (2.12) we take the level curve of Φ , namely

$$\Gamma = \Gamma_A \stackrel{\text{def}}{=} \left\{ z \in \mathbb{C} \backslash \Delta : |\Phi(z)| = \Phi(A) \right\}, \quad A > 1.$$

It is easy to see that Γ_A is an ellipse whose equation in Cartesian coordinates on the (x, y) plane is

$$\frac{x^2}{A^2} + \frac{y^2}{(A^2 - 1)} = 1.$$

By (2.15), it is also a level curve of U^{μ_0} . Thus, the problem of minimizing $\log |z - \zeta|$ with $z \in \Gamma_A$ is equivalent to the real-valued constrained optimization problem

minimize
$$(x - \zeta)^2 + y^2$$

subject to $\frac{x^2}{A^2} + \frac{y^2}{A^2 - 1} = 1$

Assume w.o.l.g. that $0 \leq \zeta \leq 1$. Standard methods show that the only possible points of minimum are (x, y) = (A, 0) and, if $\zeta A^2 \leq 1$,

$$(x,y) = \left(\zeta A^2, \pm \sqrt{(A^2 - 1)(1 - \zeta)}\right).$$

Hence, the global minimum is

$$\begin{cases} (A - \zeta)^2, & \text{if } \zeta A^2 \ge 1, \\ \min\left\{ (A - \zeta)^2, (A^2 - 1)(1 - \zeta^2) \right\}, & \text{if } \zeta A^2 \le 1. \end{cases}$$

Since

$$(A - \zeta)^2 = (A^2 - 1)(1 - \zeta^2) + (\zeta A - 1)^2 \ge (A^2 - 1)(1 - \zeta^2),$$

for $-1 \leq \zeta \leq 1$ and $z \in \Gamma_A$,

$$\log|z-\zeta| \ge \frac{1}{2}\log(A^2-1) + \frac{1}{2}\log(1-\zeta^2),$$

so that, combining it with (2.15),

$$U^{\mu_0}(z) + t \log|z - \zeta| \ge \log(2) - \log|\Phi(A)| + \frac{t}{2}\log(A^2 - 1) + \frac{t}{2}\log(1 - \zeta^2).$$

Using it in (2.12), we see that for $-1 \leq \zeta \leq 1$,

$$U^{\mu_t}(\zeta) \ge -c_t + \log(2) - \log|\Phi(A)| + \frac{t}{2}\log(A^2 - 1) + \frac{t}{2}\log(1 - \zeta^2).$$
(2.17)

,

With φ as in (2.5) we can rewrite it as

$$(U^{\mu_t} + \varphi)(\zeta) \ge -c_t + \log(2) - \log|\Phi(A)| + \frac{t}{2}\log(A^2 - 1),$$

and we conclude that

$$\min_{x \in [-1,1]} \left(U^{\mu_0} + \varphi \right)(x) \ge -c_t + \log(2) - \log|\Phi(A)| + \frac{t}{2}\log(A^2 - 1).$$
(2.18)

The global maximum of the right-hand side over A > 1 is attained at

$$A = \frac{1}{\sqrt{1-t^2}} = \frac{1}{s}$$

and is equal to

$$-c_t + \log(2) - \log|\Phi(1/s)| + \frac{t}{2}\log(s^{-2} - 1) = -c_t + \log(2) - \log|\Phi(1/s)| + t\log(t/s).$$

Simplifying, we obtain that

$$\min_{x \in [-1,1]} \left(U^{\mu_t} + \varphi \right)(x) \ge -c_t + \log(2) + (1-t)\log(1-t) + \left(\frac{t}{2} - 1\right)\log(2-t) + \frac{t}{2}\log(t)$$
2.16) follows.

and (2.16) follows.

The assertion of Theorem 1.1 follows from Lemma 2.5 and the characterization (2.4).

Acknowledgments

The first author was partially supported by Simons Foundation Collaboration Grants for Mathematicians (grant 710499). He also acknowledges the support of the project PID2021-124472NB-I00, funded by MCIN/AEI/10.13039/501100011033 and by "ERDF A way of making Europe", as well as the support of Junta de Andalucía (research group FQM-229 and Instituto Interuniversitario Carlos I de Física Teórica y Computacional).

A. MARTÍNEZ-FINKELSHTEIN AND E. A. RAKHMANOV

References

- R. Bøgvad, C. Hägg, and B. Shapiro. Rodrigues' descendants of a polynomial and Boutroux curves. Constr. Approx., 59(3):737-798, 2024.
- [2] NIST Digital Library of Mathematical Functions. https://dlmf.nist.gov/, Release 1.2.3 of 2024-12-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [3] A. Martínez-Finkelshtein and E. A. Rakhmanov. Flow of the zeros of polynomials under iterated differentiation. Preprint arXiv:2408.13851, 2024.
- [4] E. B. Saff and V. Totik. Logarithmic Potentials with External Fields, volume 316 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin, second edition, 2024.

(AMF) DEPARTMENT OF MATHEMATICS, BAYLOR UNIVERSITY, WACO, TX 76706, USA, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALMERÍA, ALMERÍA, SPAIN

Email address: A_Martinez-Finkelshtein@baylor.edu

(ER) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FL 33620, USA

Email address: rakhmano@usf.edu