A Revisit to Rate-distortion Theory via Optimal Weak Transport

Jiayang Zou, Luyao Fan, Jiayang Gao and Jia Wang Department of Electronic Engineering Shanghai Jiao Tong University Shanghai, China

Email: qiudao@sjtu.edu.cn, fanluyao@sjtu.edu.cn, gjy0515@sjtu.edu.cn, jiawang@sjtu.edu.cn

Abstract—This paper revisits the rate-distortion theory from the perspective of optimal weak transport, as recently introduced by Gozlan et al. While the conditions for optimality and the existence of solutions are well-understood in the case of discrete alphabets, the extension to abstract alphabets requires more intricate analysis. Within the framework of weak transport problems, we derive a parametric representation of the rate-distortion function, thereby connecting the rate-distortion function with the Schrödinger bridge problem, and establish necessary conditions for its optimality. As a byproduct of our analysis, we reproduce K. Rose's conclusions regarding the achievability of Shannon lower bound concisely, without reliance on variational calculus.

I. Introduction

Rate-distortion (RD) theory is a fundamental branch of information theory, primarily concerned with the problem of compressing data at the minimum coding rate subject to a given distortion constraint. Initially proposed by Claude E. Shannon in his groundbreaking works in 1948 [1] and 1959 [2], this theory has since become a cornerstone of modern data compression and communication system design. RD problems focus on designing a coding scheme that minimizes the coding rate, while satisfying a specific distortion level for a given source and distortion measure.

The classical rate-distortion problems for finite-alphabet sources and continuous sources defined in Euclidean spaces have been extensively studied in [3]-[6]. As research advanced, scholars began to explore more complex problems involving abstract alphabets. A key contribution was made by I. Csiszár, who extended RD theory to abstract alphabets [7]. He rigorously derived the parametric representation of the RD function, laying a solid theoretical foundation for numerical algorithms [4], [8]. Subsequently, F. Rezaei et al. [9] significantly advanced Csiszár's results by employing techniques such as Stone-Cěch compactification and fixed-point methods in locally convex topological vector spaces. These approaches eliminated the assumption of compactness for the target space. Moreover, they provided a comprehensive characterization of the optimal reconstruction under these generalized conditions. Further contributions were made by E. Riegler et al. [10], [11], who focused on lossy compression theory for sources defined on compact manifolds and fractal sets. By leveraging results related to the rate-distortion dimension [12], they derived the Shannon lower bound (SLB) in more general settings, thereby extending the applicability of RD theory to more complex

source structures. In terms of the parametric representation of the RD function, V. Kostina and E. Tuncel [13] avoided reliance on the Karush-Kuhn-Tucker (KKT) optimality conditions. Instead, they streamlined I. Csiszár's argument by utilizing the Donsker-Varadhan characterization of minimum relative entropy [14]. Their work significantly advanced the theory of successive refinement for abstract sources, thereby generalizing the earlier results of Equitz and Cover [15].

In 2013, Robert M. Gray highlighted the connections between optimal transport (OT) and information theory [16], offering a geometric view of source coding and a brief survey of optimal transport from a quantization perspective. Since then, the relationship between OT and RD theory has gradually garnered attention. However, conventional optimal transport theory is challenging to apply directly to RD problems due to fundamental differences in their problem settings. Conventional optimal transport minimizes transportation cost between probability distributions with marginal constraints, while RD theory balances coding rate and distortion under a disintegration framework. To bridge these differences, an entropy regularization term have been introduced to establish equivalence within the framework of entropic optimal transport in [17]. In fact, optimal weak transport (OWT) theory, as a generalization of OT, naturally links to RD problems through optimizing the disintegration of the target measure with respect to the source measure. This framework enables a more concise reproduction of several key results in RD theory.

In this paper we conduct a study of revisiting the RD theory via OWT, which was recently introduced by Gozlan et al. [18]. We derive a parametric representation that connects the RD function with the Schrödinger problem [19] within the framework of weak transport problems. Additionally, we significantly streamline the proof of the necessary optimality conditions for the RD function using the existence theorem of OWT problems [20], [21]. This approach provides a concise method to reproduce K. Rose's conclusions regarding the achievability of the Shannon lower bound [22], avoiding reliance on variational calculus and circumventing the potential risks of a lack of rigor that may arise in the use of variational methods, which typically entails more intricate and complex analysis to ensure rigor.

II. BACKGROUND AND PROBLEM SETUP

A. Fundamentals of Weak Transport Theory

Optimal transport theory studies the transportation of mass between two probability measures. Given two probability measures μ and ν on Polish spaces (complete separable metric spaces) $(\mathcal{X}, d_{\mathcal{X}}), (\mathcal{Y}, d_{\mathcal{Y}})$, the optimal transport problem is to find a transport plan π that minimizes the transportation cost

$$\inf_{\pi \in \Pi(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x,y) d\pi(x,y) \tag{1}$$

where $\Pi(\mu,\nu)$ is the set of all transport plans between μ and ν , and c(x,y) is the cost of transporting mass from x to y. The optimal transport problem has been widely studied in mathematics, and has applications in various fields, such as machine learning [23]–[26], statistics [27]–[29], and information theory [30]–[33]. Readers are referred to [34] for background on optimal transport.

Weak transport theory was initiated by Gozlan et al. [18] motivated by applications in geometric inequalities. Their contribution sparked extensive research activity among various groups and led to applications in numerous fields in [20], [21], [35]–[39]. Throughout $\mathcal X$ and $\mathcal Y$ denote Polish spaces. Given probability measures $\mu \in \mathcal P(\mathcal X), \nu \in \mathcal P(\mathcal Y)$ we write (μ, ν) for the set of all couplings on $\mathcal X \times \mathcal Y$ with marginals μ and ν . Given a coupling π on $\mathcal X \times \mathcal Y$, we denote a regular disintegration with respect to the first marginal by $(\pi_x), x \in \mathcal X$. We consider cost functionals of the form

$$C: \mathcal{X} \times \mathcal{P}(\mathcal{Y}) \to \mathbb{R} \cup \{+\infty\} \tag{2}$$

where C is lower bounded and lower semicontinuous, and $C(x,\cdot)$ is assumed to be convex on $\mathcal{P}(\mathcal{Y})$ for all $x\in\mathcal{X}$. The weak transport problem is to find a coupling π that minimizes the cost functional

$$V_C(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X}} C(x, \pi_x) d\mu(dx).$$
 (3)

The classical transport problem is included via $C(x,p) = \int c(x,y)dp(y)$ for some cost function $c: \mathcal{X} \times \mathcal{Y} \to \mathbb{R} \cup \{+\infty\}$. For $t \geq 1$, $\mathcal{P}^t_{d_{\mathcal{Y}}}(\mathcal{Y})$, which, for convenience, can be abbreviated as $\mathcal{P}_t(\mathcal{Y})$, denotes the set of Borel probability measures with finite t-th moment for some fixed metric $d_{\mathcal{Y}}$ (compatible with the topology on \mathcal{Y}), that is, a Borel probability measure ν is in $\mathcal{P}_t(\mathcal{Y})$ if and only if for some $y_0 \in \mathcal{Y}$ we have

$$\int_{\mathcal{Y}} dy(y, y_0)^t \nu(dy) < \infty.$$

The set of continuous functions on $\mathcal Y$ which are dominated by a multiple of $1+d_{\mathcal Y}(y,y_0)^t$, is denoted by $\Phi_t(\mathcal Y)$. We equip the set of probability measures $\mathcal P_t(\mathcal Y)$ with the t-th Wasserstein topology. Specifically, a sequence $(\nu_k)_{k\in\mathbb N}$ converges to $\nu\in\mathcal P_t(\mathcal Y)$ if $\nu_k(f):=\int fd\nu_k$ converges to $\nu(f)$ for all $f\in\Phi_t(\mathcal Y)$. The space $\mathcal P(\mathcal Y)$ itself is equipped with the usual weak topology. The same conventions apply to $\mathcal X$ instead of $\mathcal Y$.

Definition 1. (A) A cost functional $C: \mathcal{X} \times \mathcal{P}_t(\mathcal{Y}) \to \mathbb{R} \cup \{+\infty\}$ satisfies property (A) if and only if

- 1) C is lower semicontinuous with respect to the product topology on $\mathcal{X} \times \mathcal{P}_t(\mathcal{Y})$,
- 2) C is bounded from below,
- 3) the map $p \mapsto C(x, p)$ is convex, that is, for all $x \in \mathcal{X}$ and all $p, q \in \mathcal{P}_t(\mathcal{Y}), \ \lambda \in [0, 1]$, we have

$$C(x, \lambda p + (1 - \lambda)q) \le \lambda C(x, p) + (1 - \lambda)C(x, q).$$

Theorem 1. [Theorem 1.1 in [20]] Assume that $C: \mathcal{X} \times \mathcal{P}(\mathcal{Y}) \to \mathbb{R} \cup \{+\infty\}$ is jointly lower semicontinuous, bounded from below and convex in the second argument. Then the problem

$$\inf_{\pi \in \Pi(\mu,\nu)} \int_{\mathcal{X}} C(x,\pi_x) \mu(dx)$$

admits a minimizer.

However, the cost function C may not be lower semicontinuous with respect to weak convergence in many cases, such as $\rho(x,y) = \|x-y\|^2$ for $\mathcal{X} = \mathcal{Y} = \mathbb{R}^n$, hence we will need to employ a refined version of Existence Theorem 1 to carry out applications in general cases.

Theorem 2. (Existence and semicontinuity)[Theorem 3.2 in [21]] If the cost function C satisfies property (A), the infimum in (3) is attained and the value $V_C(\mu, \nu)$ depends in a lower semicontinuous way on the marginals $(\mu, \nu) \in \mathcal{P}(\mathcal{X}) \times \mathcal{P}_t(\mathcal{Y})$.

B. A Sketch of The Schödinger Problem

The (static) Schödinger problem (SP), also called the Schödinger bridge problem, was addressed in [40] in 1931. While the problem is historically old, it has attracted significant attention in the past decades due to its deep connection with various fields, especially (entropic) optimal transport theory. In modern terminology, SP can be expressed as follows. For two Polish spaces \mathcal{X}, \mathcal{Y} with given probability measures $\mu \in \mathcal{P}(\mathcal{X}), \nu \in \mathcal{P}(\mathcal{Y})$, and a given reference probability measure $\gamma \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$. We denote by $\Pi(\mu, \nu) \subset \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ the set of couplings; namely, the set of all $\pi \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ satisfying

$$\int_{\mathcal{X}\times\mathcal{Y}} f(x)\pi(dx,dy) = \int_{\mathcal{X}} f(x)\mu(dx)$$

$$\int_{\mathcal{X}\times\mathcal{Y}} g(y)\pi(dx,dy) = \int_{\mathcal{Y}} g(y)\nu(dy)$$
(4)

for all bounded measurable functions f,g. The Schrödinger problem is to find a coupling $\pi \in \Pi(\mu,\nu)$ that minimizes the relative entropy with respect to the reference measure γ :

$$H(\pi|\gamma) = \begin{cases} \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi}{d\gamma} d\pi, & \text{if } \pi \ll \gamma \\ +\infty, & \text{otherwise} \end{cases}$$
 (5)

This problem can be seen as a special case of OWT problems. Assume that the value $\inf_{\pi\in\Pi(\mu,\nu)} H(\pi|\gamma)$ is finite, then OWT guarantees the existence of minimizer and provides the

Lemma 1. [Corollary 4.3 in [21]] Let $\mu \in \mathcal{P}(\mathcal{X})$, $\nu \in \mathcal{P}(\mathcal{Y})$ and $\gamma \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ a probability measure equivalent to $\mu \times \nu$.

structure of optimizers in the SP.

Assume that the value of the corresponding entropic optimal transport problem is finite, that is

$$\inf_{\pi \in \Pi(\mu,\nu)} H(\pi|\gamma) < +\infty. \tag{6}$$

A coupling π^* minimizes the above problem if and only if $H(\pi^*|\gamma) < \infty$ and there exist measurable functions f,g such that

$$\frac{d\pi^*}{d\gamma}(x,y) = f(x)g(y), \gamma - a.e.$$

In fact, f(x) and g(y) in the lemma above are nonnegative and unique up to a multiplicative constant [41]. In the case where $d\gamma = Ke^{-\beta\rho(x,y)}d\mu \times d\nu$ for a nonnegative measurable function $\rho(x,y)$, since $\pi^\star \in \Pi(\mu,\nu)$, we have $\int_{\mathcal{Y}} d\pi^\star = d\mu$, $\int_{\mathcal{X}} d\pi^\star = d\nu$, which can be written as Schrödinger equations [41]

$$1 = Kf(x) \int_{\mathcal{Y}} g(y)e^{-\beta\rho(x,y)}d\nu$$

$$= Kg(y) \int_{\mathcal{X}} f(x)e^{-\beta\rho(x,y)}d\mu$$

$$= \int_{\mathcal{X}} \frac{g(y')e^{-\beta\rho(x,y')}}{\int_{\mathcal{Y}} g(y)e^{-\beta\rho(x,y)}d\nu}d\mu, \quad \forall y' \in \mathcal{Y}.$$
(7)

where

$$K = \left(\int_{\mathcal{X} \times \mathcal{Y}} e^{-\beta \rho(x,y)} d\mu d\nu\right)^{-1}$$
$$= \left(\int_{\mathcal{X} \times \mathcal{Y}} f(x)g(y)e^{-\beta \rho(x,y)} d\mu d\nu\right)^{-1}$$
(8)

In the following sections, we will recapitulate the RD theory, and investigate its connection to the Schrödinger problem within the framework of weak transport theory.

C. A Recap of Rate-distortion Theory

Given a source X with probability distribution p(x), and a distortion measure $\rho(x,y)$, the RD function R(D) is defined as the minimum rate required to achieve a distortion level D,

$$R(D) = \inf_{Y} I(X;Y) \tag{9}$$

subject to the constraint

$$\mathbb{E}[\rho(X,Y)] \leqslant D \tag{10}$$

where I(X;Y) is the mutual information between X and Y, and Y is the reconstruction. I. Csiszár proved the following basic properties about the rate-distortion function and its parametric representation under sufficiently weak assumptions in [7], providing a solid theoretical foundation for the RD theory:

Assumption 1. Let $\rho: \mathcal{X} \times \mathcal{Y} \to [0, \infty]$ be a given measurable function, called a loss function, with

$$\inf_{y} \rho(x, y) = 0, \forall x \in \mathcal{X}. \tag{11}$$

Lemma 2. [Lemma 1.1 in [7]] Under Assumption 1, the ratedistortion function R(D) is a non-increasing convex function of D and one may write

$$R(D) = \inf_{\mathbb{E}_{\pi}(\rho) = D} I(X; Y) \tag{12}$$

where $D \leq D_1$, which is the smallest non-negative number such that R(D) is constant in $(D_1, +\infty)$.

Assumption 2. We assume the existence of random variables ξ, η with $\xi \sim \mu$ such that

$$I(\xi;\eta) < \infty, \mathbb{E}\rho(\xi,\eta) < \infty.$$
 (13)

This assumption can exclude the contingency that ${\cal R}(D)$ is identically infinite.

Assumption 3. Suppose that there exists a finite set $B \subset \mathcal{Y}$ such that

$$\int \rho(x,B)\mu(dx) < \infty \quad \textit{where} \quad \rho(x,B) \coloneqq \min_{y \in B} \rho(x,y).$$

Define $D_{\min} := \inf\{D \ge 0 | R(D) < \infty\}.$

Theorem 3. [Theorem 2.1 in [7]] Under Assumption 1 and 2, suppose that to any $\epsilon > 0$ there exists measurable subsets $A_1 \subset A_2 \subset \cdots$ of \mathcal{X} and finite subsets $B_1 \subset B_2 \subset \cdots$ of \mathcal{Y} such that $\mu(A_n) \to 1$ as $n \to \infty$ and

$$\min_{y \in B_n} \rho(x, y) < \epsilon, \forall x \in A_n, n = 1, 2, \cdots$$
 (14)

Then R(D) is finite for all D > 0, namely, $D_{\min} = 0$.

The hypothesis of this theorem is trivially fulfilled if \mathcal{Y} is at most countable. It is also fulfilled if \mathcal{X} is Polish and $\rho^{(K)}(x,y) := \min\{\rho(x,y),K\}$ is upper semicontinuous for some fixed K>0. In this case the A_i 's may be chosen as an increasing sequence of compact subsets of \mathcal{X} , since on a Polish space every probability distribution is tight [7].

Theorem 4. [Theorem 2.1 in [7]] Under Assumption 1 and 2, if there exist A_n 's as in Theorem 3 and $y_n \in \mathcal{Y}$ with

$$\max_{x \in A_n} \rho(x, y_n) < \infty, n = 1, 2, \cdots$$
 (15)

then $R(D) \to 0$ as $D \to \infty$. Furthermore, setting

$$D_{\max} = \lim_{K \to \infty} \left(\inf_{y} \int \rho^{(K)}(x, y) \mu(dx) \right). \tag{16}$$

Then R(D) is positive for $D < D_{\max}$ and if $D_{\max} < \infty$ then R(D) = 0 for $D > D_{\max}$.

Theorem 5. [Theorem 2.3 in [7]] Under Assumption 1, 2 and 3, for each $D > D_{\min}$ holds

$$R(D) = \max_{\beta, \alpha(x)} \left\{ \int \log \alpha(x) \mu(dx) - \beta D \right\}$$
 (17)

where the maximization refers to $\alpha(x) \geqslant 1$ and $\beta \geqslant 0$ under the constraints

$$\int \alpha(x)e^{-\beta\rho(x,y)}\mu(dx) \leqslant 1, \forall y \in \mathcal{Y}.$$
 (18)

The maximizing β values are exact those associated with D by

$$R(D') + \beta D' \geqslant R(D) + \beta D, \forall D'.$$
 (19)

III. OWT MEETS RATE-DISTORTION THEORY

In this section, we will revisit rate-distortion theory using the framework of weak transport theory. We begin by transforming the original RD problem, which is focused on optimizing the disintegration, into an OWT problem of optimizing the joint distribution under the condition of a given target measure (or reconstruction). This is followed by seeking the infimum over the target measure. We will utilize the existence theorem for weak transport problems, along with its application in the Schrödinger problem, to provide a new parametric representation of the RD function. Furthermore, under the condition of the existence of an optimal reconstruction, we will prove the necessary optimality conditions for the RD function, thereby offering a concise proof of the SLB achievability as proposed by K. Rose [22]. Our main contributions are summarized from Theorem 6 to Theorem 8. We provide detailed proofs of them in Appendix.

Lemma 3. Under Assumption 1 and 2, the rate-distortion function R(D) can be represented as

$$\begin{split} \inf_{\pi_x: \mathbb{E}_\pi \rho \leqslant D} I(X;Y) &= \inf_{\nu \in \mathcal{P}(\mathcal{Y})} \inf_{\pi \in \Pi(\mu,\nu): \mathbb{E}_\pi \rho \leqslant D} I(X;Y) \\ &= \inf_{(\nu,\pi) \in \mathcal{P}(\mathcal{Y}) \times \Pi(\mu,\nu): \mathbb{E}_\pi \rho \leqslant D} I(X;Y). \end{split}$$

Proof. See Appendix A.

Remark 1. Note that $I(X;Y) = \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi}{d\mu d\nu} d\pi = \int_{\mathcal{X}} d\mu(x) \int_{\mathcal{Y}} \log \frac{d\pi_x}{d\nu} d\pi_x$, and $\pi \ll \mu \times \nu \iff \pi_x \ll \nu$. Hence we always assume that $\pi_x \ll \nu$ so that the infimum is proper.

Consider the (weak) dual problem of rate-distortion problem

$$R(D) = \inf_{\nu \in \mathcal{P}(\mathcal{Y})} \inf_{\pi \in \Pi(\mu,\nu): \mathbb{E}_{\pi} \rho \leqslant D} I(X;Y)$$

$$\geqslant \inf_{\nu \in \mathcal{P}(\mathcal{Y})} \sup_{\beta \geqslant 0} \inf_{\pi \in \Pi(\mu,\nu)} I(X;Y) + \beta(\mathbb{E}_{\pi} \rho - D).$$
(20)

Define

$$J(\nu,\beta) \coloneqq \inf_{\pi \in \Pi(\mu,\nu)} I(X;Y) + \beta (\mathbb{E}_{\pi}\rho - D)$$

$$= \int_{\mathcal{X}} d\mu(x) \int_{\mathcal{Y}} \left[\log \frac{d\pi_x}{d\nu} + \beta \rho(x,y) - \beta D \right] d\pi_x$$

$$d\gamma(x,y) \coloneqq K e^{-\beta \rho(x,y)} d\mu(x) d\nu(y)$$
where $K = \left(\int_{\mathcal{X} \times \mathcal{Y}} e^{-\beta \rho(x,y)} d\mu d\nu \right)^{-1}$ such that $\gamma \ll \mu \times \nu$ is a probability measure on $\mathcal{X} \times \mathcal{Y}$ (note that γ is indeed

equivalent to $\mu \times \nu$). We have

$$I(X;Y) + \beta(\mathbb{E}_{\pi}\rho - D)$$

$$= \int_{\mathcal{X}} D_{KL}(\pi_x || \gamma_x) d\mu + \int_{\mathcal{X}} \log K_x d\mu(x) - \beta D$$

$$= \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi_x}{d\gamma_x} d\pi - \int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu(x) - \beta D$$

$$= \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi}{d\gamma} d\pi - \int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu(x)$$

$$- \beta D + \int_{\mathcal{X}} \log \frac{d\gamma_0}{d\mu} d\mu$$

where γ_0 is the X-marginal of γ and

$$d\gamma_x = K_x e^{-\beta \rho(x,y)} d\nu, K_x = \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right)^{-1}.$$

Hence we only need to consider the infimum of $D_{KL}(\pi \| \gamma)$ over $\pi \in \Pi(\mu, \nu)$.

Remark 2. One might be concerned about the finiteness of K, K_x and integrability of $\log K_x$, while they always hold indeed. It is noticeable that $K = +\infty$ if and only if $\rho(x,y) = +\infty, \mu \times \nu$ -a.e., implying that $\rho = +\infty, \pi$ -a.e. for each $\pi \in \Pi(\mu, \nu)$ with $\pi_x \ll \nu$. In this case, $\mathbb{E}_\pi \rho = +\infty$ and $J(\nu, \beta) = +\infty$. Hence we only need to consider the case that $K < +\infty$. Similarly, $K_x = +\infty$ if and only if $\rho(x,y) = +\infty, \nu$ -a.e. for given $x \in \mathcal{X}$, implying that $\rho = +\infty, \pi_x$ -a.e. with $\pi_x \ll \nu$. Then we have $K_x < +\infty, \mu$ -a.e. otherwise $J(\nu, \beta)$ becomes infinite. Hence the case of $K_x = +\infty$ can be ignored. Moreover, if $\int_{\mathcal{X}} \log K_x d\mu(x) = +\infty$, we also have $J(\nu, \beta) = +\infty$, which can be excluded.

We consider the existence of the minimizer of $J(\nu,\beta)$. Define

$$C(x,p) := \int_{\mathcal{Y}} \left[\log \frac{dp}{d\nu} + \beta \rho(x,y) - \beta D \right] dp, p \in \mathcal{P}(\mathcal{Y}).$$
(22)

Assumption 4. $\int_{\mathcal{Y}} \rho(x,y) dp$ is jointly lower semicontinuous in $(x,p) \in \mathcal{X} \times \mathcal{P}(\mathcal{Y})$.

This assumption can hold if \mathcal{X}, \mathcal{Y} are Polish spaces and $\rho \in C_b(\mathcal{X} \times \mathcal{Y})$. It can be checked that $(x,p) \to \int_{\mathcal{Y}} \rho(x,y) dp$ is indeed bounded and jointly continuous. In addition, if $\rho(x,y)$ is jointly lower semicontinuous with an upper bound (the lower bound is 0), then it can be increasingly converged by a sequence of bounded k-Lipschitz functions, which implies that C(x,p) is jointly lower semicontinuous with respect to the product topology on $\mathcal{X} \times \mathcal{P}(\mathcal{Y})$. One way to verify these assertions is to use *Theorem A.3.12* in [43].

Lemma 4. Under Assumption 4, the cost function C(x, p) is jointly lower semicontinuous, bounded from below and convex in the second argument.

Corollary 1. Under the Assumption 4, $J(\nu, \beta)$ admits a minimizer, that is, there exists $\pi^* \in \Pi(\mu, \nu)$ such that

$$J(\nu,\beta) = \int_{\mathcal{X}} d\mu(x) \int_{\mathcal{Y}} \left[\log \frac{d\pi_x^*}{d\nu} + \beta \rho(x,y) - \beta D \right] d\pi_x^*$$
$$= \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi^*}{d\gamma} d\pi^* - \int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu(x)$$
$$-\beta D + \int_{\mathcal{X}} \log \frac{d\gamma_0}{d\mu} d\mu.$$

However, Assumption 4 may not hold in many cases, such as $\rho(x,y) = \|x-y\|^2$ for $\mathcal{X} = \mathcal{Y} = \mathbb{R}^n$, which has been mentioned below Theorem 1. Hence we need to utilize the Refined Existence heorem 2 to carry out applications in general cases. Before this, we need to impose some additional conditions on $\rho(x,y)$ and the source μ .

Assumption 5. $\int_{\mathcal{Y}} \rho(x,y) dp$ is jointly lower semicontinuous in $(x,p) \in \mathcal{X} \times \mathcal{P}_t(\mathcal{Y})$ for given metric $d_{\mathcal{Y}}$ on space \mathcal{Y} .

This assumption is weaker than Assumption 4, as convergence (or topology) in the space $\mathcal{P}_t(\mathcal{Y})$ is much stronger than in $\mathcal{P}(\mathcal{Y})$. This extension broadens the applicable scenarios, such as the case where the cost function takes the form $\rho(x,y) = \|x-y\|^t$ with $t \geqslant 1$ and $\mathcal{X} = \mathcal{Y} = \mathbb{R}^n$ under the Euclidean metric. We can also demonstrate that the cost function C(x,p) satisfies property (A).

Theorem 6. Under Assumption 5, the cost function C(x,p) satisfies property (A). Assume that $\nu \in \mathcal{P}_t(\mathcal{Y})$. Then $J(\nu,\beta)$ admits a minimizer π^* .

In order to restrict the reconstruction to the space $\mathcal{P}_t(\mathcal{Y})$, we introduce the following assumption, which allows us to apply the Refined Existence Theorem 2 of OWT to guarantee the existence of π^* . Although this assumption is stronger than those in the works of I. Csiszár [7] and F. Rezaei [9] et al., it remains reasonable and widely applicable in practical scenarios.

Assumption 6. $\mathcal{X} = \mathcal{Y}$ are Polish spaces and the source $\mu \in \mathcal{P}_t(\mathcal{X})$. The loss function $\rho(x, y)$ satisfies that

$$\rho(x,y) \geqslant c \cdot d_{\mathcal{X}}(x,y)^t$$
 for some constant $c > 0$.

Proposition 1. The reconstruction distribution $\nu \in \mathcal{P}_t(\mathcal{Y})$ for any $D > D_{\min}$ under Assumption 1, 2 and 6.

By Proposition 1 and Lemma 3, we can obtain that

$$R(D) = \inf_{\pi_{x}: \mathbb{E}_{\pi} \rho \leqslant D} I(X; Y)$$

$$= \inf_{\nu \in \mathcal{P}_{t}(\mathcal{Y})} \inf_{\pi \in \Pi(\mu, \nu): \mathbb{E}_{\pi} \rho \leqslant D} I(X; Y)$$

$$\geqslant \inf_{\nu \in \mathcal{P}_{t}(\mathcal{Y})} \sup_{\beta \geqslant 0} \inf_{\pi \in \Pi(\mu, \nu)} I(X; Y) + \beta(\mathbb{E}_{\pi} \rho - D)$$

$$= \inf_{\nu \in \mathcal{P}_{t}(\mathcal{Y})} \sup_{\beta \geqslant 0} J(\nu, \beta).$$
(23)

In fact, by applying the existence theorem of OWT and relevant results from the Schrödinger problem [19], [41], we can directly obtain the expression for $J(\nu,\beta)$, which in turn provides a new parametric representation for R(D). By Lemma 1, we can obtain the parametric representation of R(D) and a series of equivalent forms of the RD function.

Theorem 7. Under Assumption 1-6, the RD function R(D) can be represented as

$$\begin{split} R(D) &= \inf_{\pi_x: \mathbb{E}_\pi \rho \leqslant D} I(X;Y) \\ &= \inf_{\nu \in \mathcal{P}_t(\mathcal{Y})} \sup_{\beta \geqslant 0} \inf_{\pi \in \Pi(\mu,\nu)} I(X;Y) + \beta \big(\mathbb{E}_\pi \rho - D\big) \\ &= \inf_{\nu} \max_{\beta \in \partial R(D)} \left\{ -\int \! \log \left(\int \! e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D \! + \! L(\nu,\beta) \right\} \\ &= \max_{\beta \in \partial R(D)} \inf_{\nu} \left\{ -\int \! \log \left(\int \! e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D \! + \! L(\nu,\beta) \right\} \\ &= \inf_{\nu} \left\{ -\int \! \log \left(\int \! e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D \! + \! L(\nu,\beta) \right\}, \forall \beta \in \! \partial R(D) \end{split}$$

where

$$L(\nu,\beta) = \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu.$$

with g(y) satisfying the Schrödinger equations (7)

If we further assume the existence of the optimal reconstruction ν^* , $L(\nu, \beta)$ equals 0 so that the parametric representation (25) can be reproduced.

Theorem 8. Under the set of Theorem 7, if additionally, there exists an optimal reconstruction ν^* , then the corresponding optimal joint distribution satisfies that

$$d\pi^{\star} = \frac{e^{-\beta\rho(x,y)}}{\int_{\mathcal{V}} e^{-\beta\rho(x,y)} d\nu^{\star}} d\mu \times d\nu^{\star}$$
 (24)

which is equivalent to that g(y) in Lemma 1 is a constant, namely, $L(\nu^*,\beta)=0$ in Theorem 7. Then the RD function R(D) can be represented as

$$R(D) = \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D \right\}$$

=
$$-\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu^{\star} \right) d\mu - \beta D$$
 (25)

holding for any $\beta \in \partial R(D)$ and ν^* does not change even though β can vary in $\partial R(D)$.

By (24) along with K. Rose's methods of using the completeness of Hermite polynomials, we can reproduce the following conclusion.

Corollary 2. [Theorem 2 and Corollary 2 in [22]] For $\mathcal{X} = \mathcal{Y} = \mathbb{R}^n$ and $\rho(x,y) = \|x-y\|^2$, if the support of the optimal

reproduction random variable has an accumulation point, then the RD function coincides with the SLB.

If the SLB does not hold with equality, then the support of the optimal reproduction random variable consists of isolated singularities. Further, if this support is bounded, then Y is discrete and finite.

IV. DISCUSSION AND FUTURE WORK

In this paper, we revisit RD theory from the perspective of OWT. Within the framework of OWT, we derive a parametric representation of the RD function based on the existence theorem of OWT, thereby linking the RD function to the Schrödinger bridge problem. Building upon this, we provide a concise proof of the optimality condition (24) under the existence of an optimal reconstruction. Potential directions for future research include relaxing the current assumptions and exploring novel numerical algorithms to solve the RD problem. This can be inspired by the numerical techniques developed for solving the Schrödinger bridge problem, as discussed in [44]–[46].

REFERENCES

- [1] C. E. Shannon, "A mathematical theory of communication," *The Bell System Technical Journal*, vol. 27, no. 3, pp. 379–423, 1948.
- [2] —, "Coding theorems for a discrete source with a fidelity criterion," International Convention Record, vol. 7, pp. 325–350, 1959.
- [3] T. Berger, Rate-distortion theory: A mathematical basis for data compression. Englewood Cliffs: Prentice-Hall, 1971.
- [4] R. Blahut, "Computation of channel capacity and rate-distortion functions," *IEEE Trans. Inf. Theory*, vol. 18, no. 4, pp. 460–473, 1972.
- [5] R. E. Blahut, Principles and practice of information theory. Addison-Wesley Longman Publishing Co., Inc., 1987.
- [6] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. New York, NY, USA: Wiley, 2006.
- [7] I. Csiszár, "On an extremum problem of information theory," *Studia Scientiarum Mathematicarum Hungarica*, vol. 9, no. 1, pp. 57–71, 1974.
- [8] S. Arimoto, "An algorithm for computing the capacity of arbitrary discrete memoryless channels," *IEEE Trans. Inf. Theory*, vol. 18, no. 1, pp. 14–20, 1972.
- [9] F. Rezaei, N. Ahmed, and C. D. Charalambous, "Rate distortion theory for general sources with potential application to image compression," *International Journal of Applied Mathematical Sciences*, vol. 3, no. 2, pp. 141–165, 2006.
- [10] E. Riegler, H. Bölcskei, and G. Koliander, "Rate-distortion theory for general sets and measures," in *Proc. IEEE Int. Symp. Inf. Theory (ISIT)*, Vail, CO, USA, 2018, pp. 101–105.
- [11] E. Riegler, G. Koliander, and H. Bölcskei, "Lossy compression of general random variables," *Information and Inference: A Journal of the IMA*, vol. 12, no. 3, pp. 1759–1829, 2023.
- [12] T. Kawabata and A. Dembo, "The rate-distortion dimension of sets and measures," *IEEE Trans. Inf. Theory*, vol. 40, no. 5, pp. 1564–1572, 1994.
- [13] V. Kostina and E. Tuncel, "Successive refinement of abstract sources," IEEE Trans. Inf. Theory, vol. 65, no. 10, pp. 6385–6398, 2019.
- [14] M. D. Donsker and S. S. Varadhan, "Asymptotic evaluation of certain Markov process expectations for large time, I," *Commun. Pure Appl. Math.*, vol. 28, no. 1, pp. 1–47, 1975.
- [15] W. H. Equitz and T. M. Cover, "Successive refinement of information," IEEE Trans. Inf. Theory, vol. 37, no. 2, pp. 269–275, 1991.
- [16] R. M. Gray, "Transportation distance, shannon information, and source coding," in GRETSI Symposium on Signal and Image Processing, 2013, https://ee.stanford.edu/ gray/gretsi.pdf.
- [17] Y. Yang, S. Eckstein, M. Nutz, and S. Mandt, "Estimating the ratedistortion function by Wasserstein gradient descent," Advances in Neural Information Processing Systems, vol. 36, 2024.
- [18] N. Gozlan, C. Roberto, P.-M. Samson, and P. Tetali, "Kantorovich duality for general transport costs and applications," *J. Funct. Anal.*, vol. 273, no. 11, pp. 3327–3405, 2017.

- [19] C. Léonard, "A survey of the Schrödinger problem and some of its connections with optimal transport," arXiv:1308.0215, 2013.
- [20] J. Backhoff-Veraguas, M. Beiglböck, and G. Pammer, "Existence, duality, and cyclical monotonicity for weak transport costs," *Calculus of Variations and Partial Differential Equations*, vol. 58, no. 203, 2019.
- [21] J. Backhoff-Veraguas and G. Pammer, "Applications of weak transport theory," *Bernoulli*, vol. 28, no. 1, pp. 370–394, 2022.
- [22] K. Rose, "A mapping approach to rate-distortion computation and analysis," *IEEE Trans. Inf. Theory*, vol. 40, no. 6, pp. 1939–1952, 1994.
- [23] N. Courty, R. Flamary, D. Tuia, and A. Rakotomamonjy, "Optimal transport for domain adaptation," *IEEE Trans. Pattern Analysis and Machine Intelligence*, vol. 39, no. 9, pp. 1853–1865, 2016.
- [24] M. E. Hamri and Y. Bennani, "Regularized optimal transport for dynamic semi-supervised learning," arXiv:2103.11937, 2021.
- [25] M. Arjovsky, S. Chintala, and L. Bottou, "Wasserstein generative adversarial networks," in *International Conference on Machine Learning*. PMLR, 2017, pp. 214–223.
- [26] L. Chen, Y. Song, Y. Cai, J. Lu, Y. Li, Y. Xie, C. Wang, and G. He, "Multi-prototype space learning for commonsense-based scene graph generation," in *Proceedings of the AAAI Conference on Artificial Intelligence*, vol. 38, no. 2, 2024, pp. 1129–1137.
- [27] M. Talagrand, "Transportation cost for Gaussian and other product measures," *Geometric & Functional Analysis (GAFA)*, vol. 6, no. 3, pp. 587–600, 1996.
- [28] H. Cao, X. Guo, and M. Laurière, "Connecting GANs, mean-field games, and optimal transport," SIAM Journal on Applied Mathematics, vol. 84, no. 4, pp. 1255–1287, 2024.
- [29] F. Gunsilius and Y. Xu, "Matching for causal effects via multimarginal unbalanced optimal transport," arXiv:2112.04398, 2021.
- [30] A. Wibisono and V. Jog, "Convexity of mutual information along the Ornstein-Uhlenbeck flow," in *International Symposium on Information Theory and Its Applications (ISITA)*, Singapore, 2018, pp. 55–59.
- [31] T. A. Courtade, "Transportation proof of an inequality by Anantharam, Jog and Nair," *arXiv:1901.10893*, 2019.
- [32] Y. Bai, "Optimal transport meets information science: from measure concentration, to information theory, to machine learning," PhD Thesis, University of Delaware, 2022.
- [33] Y. Bai, X. Wu, and A. Özgür, "Information constrained optimal transport: From Talagrand, to Marton, to Cover," *IEEE Trans. Inf. Theory*, vol. 69, no. 4, pp. 2059–2073, 2023.
- [34] C. Villani, Optimal transport: old and new, ser. Grundlehren der mathematischen Wissenschaften. Springer, 2009.
- [35] N. Gozlan, C. Roberto, P.-M. Samson, Y. Shu, and P. Tetali, "Characterization of a class of weak transport-entropy inequalities on the line," Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, vol. 54, no. 3, pp. 1667 – 1693, 2018.
- [36] N. Gozlan and N. Juillet, "On a mixture of Brenier and Strassen theorems," Proc. Lond. Math. Soc., vol. 120, no. 3, pp. 434–463, 2020.
- [37] J. Backhoff-Veraguas, M. Beiglböck, and G. Pammer, "Weak monotone rearrangement on the line," *Electronic Communications in Probability*, vol. 25, pp. 1–16, 2020.
- [38] J. Backhoff-Veraguas and G. Pammer, "Stability of martingale optimal transport and weak optimal transport," *Ann. Appl. Probab.*, vol. 32, no. 1, pp. 721–752, 2022.
- [39] B. Acciaio, M. Beiglböck, and G. Pammer, "Weak transport for non-convex costs and model-independence in a fixed-income market," *Math. Finance*, vol. 31, no. 4, pp. 1423–1453, 2021.
- [40] E. Schrödinger, "Über die umkehrung der naturgesetze," Sitzungsberichte der Preussischen Akademie der Wissenschaften. Physikalisch-Mathematische Klasse, vol. 144, pp. 144–153, 1931.
- [41] M. Nutz, "Introduction to entropic optimal transport," Lecture notes, Columbia University, 2021.
- [42] J. Zou, L. Fan, J. Gao, and J. Wang, "A revisit to rate-distortion theory via optimal weak transport," arXiv:2501.09362, 2025.
- [43] P. Dupuis and R. S. Ellis, A weak convergence approach to the theory of large deviations. John Wiley & Sons, 2011.
- [44] A.-A. Pooladian and J. Niles-Weed, "Plug-in estimation of Schrödinger bridges," arXiv:2408.11686, 2024.
- [45] K. Caluya and A. Halder, "Wasserstein proximal algorithms for the Schrödinger bridge problem: Density control with nonlinear drift," *IEEE Trans. Automatic Control*, vol. 67, no. 3, pp. 1163–1178, 2021.
- [46] S. Marino and A. Gerolin, "An optimal transport approach for the Schrödinger bridge problem and convergence of sinkhorn algorithm," J. Sci. Comput., vol. 85(2), no. 27, 2020.

- [47] E. Posner, "Random coding strategies for minimum entropy," *IEEE Trans. Inf. Theory*, vol. 21, no. 4, pp. 388–391, 1975.
 [48] Y. Polyanskiy and Y. Wu, *Information Theory: From Coding to Learning*. Cambridge University Press, 2025.

APPENDIX A PROOF OF LEMMA 3

For any fixed $\pi_x \in \mathcal{P}(\mathcal{Y})$ with $\mathbb{E}_{\pi} \rho \leq D$, one can always find the corresponding $\nu \in \mathcal{P}(\mathcal{Y})$ and $\pi \in \Pi(\mu, \nu)$, where for any Borel measurable function $f : \mathcal{X} \times \mathcal{Y} \to [0, \infty]$

$$\int_{\mathcal{X}\times\mathcal{Y}} f(x,y)d\pi(x,y) = \int_{\mathcal{X}} d\mu(x) \int_{\mathcal{Y}} f(x,y)d\pi_x(y)$$
$$d\nu(y) = \int_{\mathcal{X}} d\pi(x,y).$$

Hence we can obtain that

$$|I(X;Y)|_{\pi_x} \geqslant \inf_{\nu,\pi:\mathbb{E}_{\pi},\rho\leqslant D} I(X;Y)$$

that is,

$$\inf_{\pi_x: \mathbb{E}_{\pi} \rho \leqslant D} I(X; Y) \geqslant \inf_{\nu, \pi: \mathbb{E}_{\pi} \rho \leqslant D} I(X; Y).$$

Additionally, for all fixed $\nu, \pi : \mathbb{E}_{\pi} \rho \leq D$, one can also determine the disintegration π_x of π , and we have

$$I(X;Y)|_{\nu,\pi} \geqslant \inf_{\pi_x: \mathbb{E}_{\pi}\rho \leqslant D} I(X;Y)$$

that is,

$$\inf_{\nu,\pi:\mathbb{E}_\pi\rho\leqslant D}I(X;Y)\geqslant \inf_{\pi_x:\mathbb{E}_\pi\rho\leqslant D}I(X;Y) \Longrightarrow \inf_{\nu,\pi:\mathbb{E}_\pi\rho\leqslant D}I(X;Y) = \inf_{\pi_x:\mathbb{E}_\pi\rho\leqslant D}I(X;Y).$$

Furthermore, we take the infimum sequence $\{\nu_k, \pi_k\}$ of $\inf_{\nu, \pi: \mathbb{E}_{\pi} \rho \leqslant D} I(X; Y)$, then

$$\inf_{\nu} \inf_{\pi \in \Pi(\mu,\nu): \mathbb{E}_{\pi}\rho \leqslant D} I(X;Y) \leqslant \inf_{\nu} \inf_{\pi \in \Pi(\mu,\nu_k): \mathbb{E}_{\pi}\rho \leqslant D} I(X;Y)$$

$$= \inf_{\pi \in \Pi(\mu,\nu_k): \mathbb{E}_{\pi}\rho \leqslant D} I(X;Y)$$

$$\leqslant I(X;Y)|_{(\mu,\pi_k)}.$$

Hence

$$\inf_{\nu}\inf_{\pi\in\Pi(\mu,\nu):\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y)\leqslant \lim_{k\to\infty}I(X;Y)|_{(\nu_k,\pi_k)}=\inf_{\nu,\pi:\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y).$$

Conversely, since for any fixed $\tilde{\nu} \in \mathcal{P}(\mathcal{Y})$,

$$\inf_{\nu,\pi:\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y)\leqslant\inf_{\pi\in\Pi(\mu,\bar{\nu}):\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y)$$

we take the infimum on both sides over $\tilde{\nu}$, and we have

$$\inf_{\nu,\pi:\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y)\leqslant\inf_{\tilde{\nu}}\inf_{\pi\in\Pi(\mu,\tilde{\nu}):\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y)=\inf_{\nu}\inf_{\pi\in\Pi(\mu,\nu):\mathbb{E}_{\pi}\rho\leqslant D}I(X;Y).$$

APPENDIX B PROOF OF LEMMA 4

Step 1. Jointly lower semi-continuous:

For $(x_k, p_k) \to (x, p)$ in the sense of product topology, i.e. $x_n \to x \in \mathcal{X}$ and $p_n \to p \in \mathcal{P}(\mathcal{Y})$, we need to verify that $C(x, p) \leq \liminf_{k \to \infty} C(x_k, p_k)$. On the one hand, we have

$$\int_{\mathcal{Y}} \log \frac{dp}{d\nu} dp \leqslant \liminf_{k \to \infty} \int_{\mathcal{Y}} \log \frac{dp_k}{d\nu} dp_k$$

by *Theorem 1* in [47] or *Theorem 4.9* in [48]. On the other hand, $\beta \int_{\mathcal{Y}} (\rho(x,y) - D) dp$ is lower semicontinuous in (x,p) by the Assumption 4. Hence C(x,p) is jointly lower semicontinuous in this case.

Step 2. Lower bounded:

$$C(x,p) := \int_{\mathcal{Y}} \left[\log \frac{dp}{d\nu} + \beta(\rho(x,y) - D) \right] dp$$

$$\geqslant \int_{\mathcal{Y}} \log \frac{dp}{d\nu} dp - \beta D$$

$$\geqslant -\beta D.$$

Note that the relative entropy is always non-negative since

$$\int_{\mathcal{Y}} \log \frac{dp}{d\nu} dp = \int_{\mathcal{Y}} \varphi(\frac{dp}{d\nu}) d\nu$$

$$\geqslant \varphi\left(\int_{\mathcal{Y}} \frac{dp}{d\nu} d\nu\right)$$

$$= \varphi(1) = 0.$$

Here $\varphi(x) = x \log x$ is a convex function over $(0, +\infty)$.

Step 3. Convex in the second argument:

For any $p_1, p_2 \in \mathcal{P}(\mathcal{Y})$ and $\lambda \in [0, 1]$, we have

$$C(x, \lambda p_1 + (1 - \lambda) p_2) - \lambda C(x, p_1) + (1 - \lambda) C(x, p_2)$$

$$= \int_{\mathcal{Y}} \left[\log \frac{\lambda p_1 + (1 - \lambda p_2)}{d\nu} + \beta \rho(x, y) \right] (\lambda dp_1 + (1 - \lambda) dp_2)$$

$$- \lambda \int_{Y} \left[\log \frac{dp_1}{d\nu} + \beta \rho(x, y) \right] dp_1 + (1 - \lambda) \int_{Y} \left[\log \frac{dp_2}{d\nu} + \beta \rho(x, y) \right] dp_2$$

$$= \int_{Y} \log \frac{\lambda dp_1 + (1 - \lambda) dp_2}{d\nu} (\lambda dp_1 + (1 - \lambda) dp_2) - \lambda \int_{Y} \log \frac{dp_1}{d\nu} dp_1 + (1 - \lambda) \int_{Y} \log \frac{dp_2}{d\nu} dp_2$$

$$= \int_{\mathcal{Y}} \varphi(\lambda \frac{dp_1}{d\nu} + (1 - \lambda) \frac{dp_2}{d\nu}) d\nu - \lambda \int_{\mathcal{Y}} \varphi(\frac{dp_1}{d\nu}) d\nu - (1 - \lambda) \int_{\mathcal{Y}} \varphi(\frac{dp_2}{d\nu}) d\nu$$

$$\leq 0.$$

APPENDIX C PROOF OF THEOREM 6

The lower bounded property and the convexity in the second argument of C(x,p) over $\mathcal{P}_t(\mathcal{Y})$ can be proved by the same method as in Lemma 4. Hence we only need to prove the jointly lowe semi-continuity of C(x,p) over $\mathcal{X} \times \mathcal{P}_t(\mathcal{Y})$. On the one hand, we have

$$\int_{\mathcal{Y}} \log \frac{dp}{d\nu} dp \leqslant \liminf_{k \to \infty} \int_{\mathcal{Y}} \log \frac{dp_k}{d\nu} dp_k$$

by Theorem 1 in [47] or Theorem 4.9 in [48]. On the other hand, $\beta \int_{\mathcal{Y}} (\rho(x,y) - D) dp$ is lower semicontinuous in $(x,p) \in \mathcal{X} \times \mathcal{P}_t(\mathcal{Y})$ by the Assumption 5. Hence C(x,p) is jointly lower semicontinuous in this case.

Furthermore, by Theorem 2, we know that the minimizer π^* of $J(\nu, \beta)$ exists.

APPENDIX D PROOF OF PROPOSITION 1

By the property of Wasserstein distance, we have

$$W_t(\nu, \delta_{y_0}) = \left(\int_{\mathcal{X}} d_{\mathcal{X}}(x, y_0)^t\right)^{1/t}$$

$$\leqslant W_t(\mu, \delta_{y_0}) + W_t(\mu, \nu).$$

Note that

$$W_t(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \left[\mathbb{E}_{\pi}(d_X^t) \right]^{1/t}$$

$$\leqslant c^{-1/t} \inf_{\pi \in \Pi(\mu, \nu)} \left[\mathbb{E}_{\pi} \rho \right]^{1/t}$$

$$\leqslant c^{-1/t} D^{1/t}.$$

Hence we have

$$W_t(\nu, \delta_{y_0}) \leqslant c^{-1/t} D^{1/t} + W_t(\mu, \delta_{y_0}) < +\infty.$$

APPENDIX E

PROOF OF THEOREM 7

By Lemma 1, we know that the minimizers π_x^{\star} , π^{\star} exist and there exist nonnegative measurable functions f, g such that

$$\frac{d\pi^*}{d\gamma} = f(x)g(y), \gamma - a.e..$$

Namely,

$$d\pi^* = Kf(x)g(y)e^{-\beta\rho(x,y)}d\mu(x)d\nu(y).$$

Then we have:

$$d\pi_x^{\star} = Kf(x)g(y)e^{-\beta\rho(x,y)}d\nu = \frac{K}{K_x}f(x)g(y)d\gamma_x$$
$$\frac{d\pi_x^{\star}}{d\gamma_x} = \frac{Kf(x)g(y)}{K_x}.$$

Since $\pi^{\star} \in \Pi(\mu, \nu)$, we have $\int_{\mathcal{Y}} d\pi^{\star} = d\mu$, $\int_{\mathcal{X}} d\pi^{\star} = d\nu$, which can be written as Schrödinger equations [41]

$$K = \left(\int_{\mathcal{X} \times \mathcal{Y}} e^{-\beta \rho(x,y)} d\mu d\nu\right)^{-1} = \left(\int_{\mathcal{X} \times \mathcal{Y}} f(x)g(y)e^{-\beta \rho(x,y)} d\mu d\nu\right)^{-1}$$

$$1 = Kf(x) \int_{\mathcal{Y}} g(y)e^{-\beta \rho(x,y)} d\nu = Kg(y) \int_{\mathcal{X}} f(x)e^{-\beta \rho(x,y)} d\mu$$

$$\implies \int_{\mathcal{X}} \frac{g(y')e^{-\beta \rho(x,y')}}{\int_{\mathcal{Y}} g(y)e^{-\beta \rho(x,y)} d\nu} d\mu = 1, \quad \forall y' \in \mathcal{Y}.$$
(26)

Then we have

$$J(\nu,\beta) = \int_{\mathcal{X}\times\mathcal{Y}} \log \frac{d\pi^*}{d\gamma} d\pi^* + \log K - \beta D$$

$$= \int_{\mathcal{X}\times\mathcal{Y}} \log(f(x)g(y)) d\pi^* + \log K - \beta D$$

$$= \int_{\mathcal{X}} \log(Kf(x)) d\mu + \int_{\mathcal{Y}} \log(g(y)) d\nu - \beta D$$

$$= -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu - \beta D$$

$$= -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu$$
(27)

$$\begin{split} R(D) &= \inf_{\nu} \inf_{\pi \in \Pi(\mu,\nu): \mathbb{E}_{\pi} \rho \leqslant D} I(X;Y) \\ &\geqslant \inf_{\nu} \sup_{\beta \geqslant 0} \inf_{\pi \in \Pi(\mu,\nu)} I(X;Y) + \beta (\mathbb{E}_{\pi} \rho - D) \\ &= \inf_{\nu} \sup_{\beta \geqslant 0} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\} \\ &\geqslant \sup_{\beta \geqslant 0} \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}. \end{split}$$

Define

$$L(\beta) = \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}.$$

Then for any $\beta \geqslant 0$

$$R(D) \geqslant L(\beta) - \beta D$$

$$R(D) + \beta D \geqslant L(\beta).$$
(28)

It is well known that R(D) is a non-increasing convex function of D, hence it has second derivatives almost everywhere, and for any $D \geqslant D_{\min}$ there exists subgradient of R(D), denoted by $\partial R(D)$, which is always a closed, convex and bounded set for convex function R(D). Then for each $-\beta_D \in \partial R(D)$, we have

$$R(D') \geqslant R(D) - \beta_D(D' - D), \ \forall D' \in (D_{\min}, D_{\max})$$

that is,

$$R(D') + \beta_D D' \geqslant R(D) + \beta_D D \geqslant L(\beta_D).$$

Additionally, by the definition of $L(\beta)$, for any sufficiently small $\epsilon > 0$, there exists corresponding $\nu^{\epsilon} \in \mathcal{P}(\mathcal{Y})$, such that

$$-\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta_D \rho(x,y)} d\nu^{\epsilon} \right) d\mu + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta_D \rho(x,y)} d\nu^{\epsilon}}{\int_{\mathcal{Y}} g(y) e^{-\beta_D \rho(x,y)} d\nu^{\epsilon}} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu^{\epsilon} < L(\beta_D) + \epsilon.$$

On the other hand,

$$-\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta_{D}\rho(x,y)} d\nu^{\epsilon} \right) d\mu + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta_{D}\rho(x,y)} d\nu^{\epsilon}}{\int_{\mathcal{Y}} g(y) e^{-\beta_{D}\rho(x,y)} d\nu^{\epsilon}} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu^{\epsilon}$$

$$= (I(X;Y) + \beta_{D} \mathbb{E}_{\pi} \rho)|_{\pi_{x} = \pi_{x}^{\star}(\nu^{\epsilon})}$$

$$\geqslant R(\mathbb{E}_{\pi^{\star}(\nu^{\epsilon})} \rho) + \beta_{D} \mathbb{E}_{\pi^{\star}(\nu^{\epsilon})} \rho$$

$$\geqslant R(D) + \beta_{D} D.$$
(29)

By equations above, we have

$$R(D) + \beta_D D - \epsilon < L(\beta_D) \leq R(D) + \beta_D D, \quad \forall \epsilon > 0.$$

Furthermore,

$$L(\beta_D) = R(D) + \beta_D D$$

$$R(D) \geqslant \sup_{\beta \geqslant 0} (L(\beta) - \beta D)$$

$$\geqslant \sup_{\beta_D \in \partial R(D)} (L(\beta_D) - \beta_D D)$$

$$= R(D).$$
(30)

Above all, we have proved that

$$R(D) = \inf_{\pi_x : \mathbb{E}_{\pi} \rho \leqslant D} I(X;Y) = \inf_{\nu} \inf_{\pi \in \Pi(\mu,\nu) : \mathbb{E}_{\pi} \rho \leqslant D} I(X;Y) = \inf_{\nu,\pi \in \Pi(\mu,\nu) : \mathbb{E}_{\pi} \rho \leqslant D} I(X;Y)$$

$$= \inf_{\nu} \sup_{\beta \geqslant 0} \inf_{\pi \in \Pi(\mu,\nu)} I(X;Y) + \beta(\mathbb{E}_{\pi} \rho - D)$$

$$= \inf_{\nu} \sup_{\beta \geqslant 0} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}$$

$$= \sup_{\beta \geqslant 0} \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}$$

$$= \inf_{\nu} \max_{\beta \in \partial R(D)} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}$$

$$= \max_{\beta \in \partial R(D)} \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu \right\}$$

$$= \inf_{\nu} \left\{ -\int_{\mathcal{X}} \log \left(\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu \right) d\mu - \beta D + L(\nu, \beta) \right\}, \quad \forall \beta \in \partial R(D)$$

where

$$L(\nu,\beta) = \int_{\mathcal{X}} \log \left(\frac{\int_{\mathcal{Y}} e^{-\beta \rho(x,y)} d\nu}{\int_{\mathcal{Y}} g(y) e^{-\beta \rho(x,y)} d\nu} \right) d\mu + \int_{\mathcal{Y}} \log g(y) d\nu.$$

APPENDIX F
PROOF OF THEOREM 8

We have known that for each $D > D_{\min}$ and each $\beta \in \partial R(D)$

$$R(D) = \inf_{\nu \in \mathcal{P}_t(\mathcal{Y})} \inf_{\pi \in \Pi(\mu,\nu)} \left\{ \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi}{d\mu \times d\nu} d\pi + \beta \int_{\mathcal{X} \times \mathcal{Y}} \rho(x,y) d\pi \right\} - \beta D.$$
 (32)

Define

$$\tilde{J}(\pi, \nu, \beta) = D_{KL}(\pi \| \mu \times \nu) + \beta \mathbb{E}_{\pi}(\rho) = \int_{\mathcal{X} \times \mathcal{Y}} \log \frac{d\pi}{d\mu \times d\nu} d\pi + \beta \int_{\mathcal{X} \times \mathcal{Y}} \rho(x, y) d\pi$$

$$f_{\nu, \beta}(x) = \left(\int_{\mathcal{Y}} e^{-\beta \rho(x, y)} d\nu \right)^{-1}$$

$$\pi_{\nu, \beta}(dx, dy) = f_{\nu, \beta}(x) e^{-\beta \rho(x, y)} \mu(dx) \times \nu(dy).$$
(33)

Then we obtain

$$R(D) = \inf_{\nu} \inf_{\pi \in \Pi(\mu,\nu)} \tilde{J}(\pi,\nu,\beta) - \beta D$$

= $\inf_{\pi \in \Pi(\mu,\nu)} \tilde{J}(\pi,\pi_1,\beta) - \beta D$, where π_1 is the Y-marginal of π . (34)

Assume that the optimal reconstruction ν^* exists, then we have

$$R(D) = \inf_{\pi \in \Pi(\mu, \nu^{\star})} \tilde{J}(\pi, \nu^{\star}, \beta) - \beta D$$

$$\stackrel{OWT}{=} \tilde{J}(\pi^{\star}, \nu^{\star}, \beta) - \beta D.$$
(35)

On the other hand,

$$R(D) = \inf_{\pi \in \Pi(\mu, \cdot)} \tilde{J}(\pi, \pi_1, \beta) - \beta D, \text{ where } \pi_1 \text{ is the } Y \text{-marginal of } \pi.$$
(36)

If $\inf_{\pi \in \Pi(\mu,\cdot)} \tilde{J}(\pi,\pi_1,\beta)$ can not be achieved, then there exists $\tilde{\pi}$, such that

$$\tilde{J}(\tilde{\pi}, \tilde{\pi}_1, \beta) < \tilde{J}(\pi^*, \nu^*, \beta). \tag{37}$$

Then we have

$$R(D) + \beta D = \tilde{J}(\pi^*, \nu^*, \beta) > \tilde{J}(\tilde{\pi}, \tilde{\pi}_1, \beta)$$

$$\geq \inf_{\pi \in \Pi(\mu, \cdot)} \tilde{J}(\pi, \pi_1, \beta)$$

$$= R(D) + \beta D.$$
(38)

It is a contradiction. Hence $\inf_{\pi \in \Pi(\mu, \cdot)} \tilde{J}(\pi, \pi_1, \beta)$ can be achieved at π^*, ν^* if the optimal reconstruction ν^* exists. Furthermore, by Lemma 1.4 in [7], we have

$$d\pi^{\star} = \alpha(x)e^{-\beta\rho(x,y)}d\mu \times d\nu^{\star}$$

$$= \frac{e^{-\beta\rho(x,y)}}{\int_{\mathcal{V}} e^{-\beta\rho(x,y)}d\nu^{\star}}d\mu \times d\nu^{\star}.$$
(39)