

## ELLIPSECT: A SURFACE BRIGHTNESS ANALYSIS TOOL FOR GALFIT 3

C. Añorve,<sup>1</sup> O. U. Reyes-Amador,<sup>2</sup> E. Ríos-López,<sup>3,4,5</sup> D. de Ramón Tadeo,<sup>6</sup>  
and O. López-Cruz<sup>3</sup>

*Draft version: January 20, 2025*

### RESUMEN

**Ellipsect** es una herramienta de análisis en Python fácil de utilizar que genera figuras de calidad para publicación a partir de los archivos de salida de **GALFIT 3**, un algoritmo popular para el ajuste 2D del brillo superficial de galaxias. **Ellipsect** permite el análisis directo de modelos preliminares de **GALFIT 3**, para guiar la selección óptima de parámetros. Además, **Ellipsect** genera medidas no paramétricas de algunos parámetros útiles que no son proporcionados por **GALFIT 3**, como el radio efectivo resultante de ajustes de multicomponentes, radio de cúspide y radio de Petrosian. Este artículo proporciona ejemplos y una guía rápida para **Ellipsect**.

### ABSTRACT

**EllipSect** is a user-friendly Python analysis tool that generates publication-quality figures from the output of **GALFIT 3**, a popular galaxy surface brightness 2D-fitting algorithm. **EllipSect** allows the direct analysis of **GALFIT 3** preliminary modeling, to guide optimal parameter selection. Furthermore, **EllipSect** generates non-parametric measurements of some useful parameters that are not provided by **GALFIT 3**, such as the effective radius resulting from multi-component fits, cusp radius, and the Petrosian radius. This paper provides examples and a quick guide for **EllipSect**.

*Key Words:* Galaxy surface brightness — Python analysis tool — photometric analysis

---

<sup>1</sup>Facultad de Ciencias de la Tierra y el Espacio, Universidad Autónoma de Sinaloa, Blvd. de la Américas y Av. Universitarios S/N, Ciudad Universitaria, C.P. 80010 Culiacán, Sinaloa, México.

<sup>2</sup>Instituto de Radioastronomía y Astrofísica, UNAM, Campus Morelia, AP 3-72, C.P. 58089, México.

<sup>3</sup>Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Luis Enrique Erro 1, Apartado Postal 51 y 216, C.P. 72840 Puebla, México.

<sup>4</sup>Instituto de Astrofísica de Canarias (IAC), Vía Láctea s/n, 38205, La Laguna, Tenerife, España.

<sup>5</sup>Universidad Pedagógica del Estado de Sinaloa (UPES), Castiza s/n, 80027, Culiacán, Sinaloa, México.

<sup>6</sup>Facultad de Ciencias de la Computación, Benemérita Universidad Autónoma de Puebla (FCC-BUAP), Av. San Claudio y 14 Sur, Ciudad Universitaria, C. P. 72570, Puebla, Puebla, México.

## 1. INTRODUCTION

**GALFIT 3**<sup>7</sup> (Peng et al. 2002, 2010) is a widely used stand-alone-two-dimensional (2D) fitting code designed to model the surface brightness (SB) distribution of astronomical sources. **GALFIT 3** applies 2D-parametric models to account for galaxies' substructures and individual stars; this means more information is used than the 1D approach. Each subcomponent has its independent orientation; hence, the separation of components is more reliable. A common misconception suggests that 1D approaches work better in low signal-to-noise (S/N) regions or the presence of low-SB features; indeed, there isn't any advantage from 1D over the 2D approaches due to the Poisson behavior of the signal<sup>8</sup>. The user provides the point spread function (PSF), and **GALFIT 3** has built-in functions to model the PSF. **GALFIT 3** convolves the PSF with the chosen profile to fit a given galaxy. This differs from most previous approaches that deconvolved the galaxy's image, generated the isophotes, and then fit a 1D profile (Jedrzejewski 1987).

**GALFIT 3** built-in functions or profiles are the following<sup>9</sup>:

1. **Sérsic (Sérsic 1968)**: This profile is a generalization of the de Vaucouleurs profile and is typically used to fit bulges, bars, and elliptical galaxies by allowing the concentration index  $n$  to vary:

$$I(r) = I_e \exp \left[ -\kappa_n \left( \left( \frac{r}{r_e} \right)^{\frac{1}{n}} - 1 \right) \right], \quad (1)$$

where  $I_e$  is the SB at the effective radius,  $r_e$ , which encloses half of the luminosity. The parameter  $\kappa_n$  is related to  $n$ , so  $r_e$  becomes the half-light radius; for  $n \in \mathbb{R}$ ,  $\kappa_n = 2n - \frac{1}{3} + \frac{4}{405n} + \frac{46}{25515n^2} + O(n^{-3})$  gives an approximation (Ciotti & Bertin 1999). See, for example, **GALFIT 3** user's manual and Graham & Driver (2005) for a full discussion on the properties of the Sérsic profile.

2. **de Vaucouleurs (de Vaucouleurs 1948)**: A particular case of the Sérsic profile (index  $n = 4$ ,  $\kappa_4 = -7.66925$ ), provides an excellent approximation to the SB profile of giant ellipticals. This profile has the following expression:

$$I(r) = I_e \exp \left[ -7.66925 \left( 1 - \left( \frac{r}{r_e} \right)^{1/4} \right) \right] \quad (2)$$

3. **Exponential (Patterson 1940)**: This popular profile is used to fit the disks of spiral galaxies; it's a particular case of the Sérsic profile with

<sup>7</sup>For installation, usage, and general help visit <https://users.obs.carnegiescience.edu/peng/work/galfit/galfit.html>

<sup>8</sup>See a more detailed discussion at <https://users.obs.carnegiescience.edu/peng/work/galfit/TFAQ.html#misconception1> on 1D vs. 2D modeling.

<sup>9</sup>See **GALFIT 3** user's manual, downloadable from <https://users.obs.carnegiescience.edu/peng/work/galfit/galfit.html>

$n = 1$ :

$$I(r) = I_0 \exp\left(-\frac{r}{r_s}\right), \quad (3)$$

where  $I_0$  is the central SB and  $r_s$  is called the scale length, which is related to the effective radius by  $r_e = 1.67835 r_s$  for  $n = 1$ . Freeman (1970) proposed a combination of a de Vaucouleurs bulge and an exponential disk to fit late-type galaxies.

4. **Nuker (Lauer et al. 1995)**: This is a general profile that includes three power-law slopes, namely:  $\gamma$  (inner),  $\alpha$  (sharpness of the transition from inner to outer slopes), and  $\beta$  (outer), they are free to vary and help to account for the large cores of very luminous early-type galaxies, including cD galaxies. This profile is expressed as follows:

$$I(r) = I_b 2^{(\frac{\beta-\gamma}{\alpha})} \left(\frac{r_b}{r}\right)^\gamma \left[1 + \left(\frac{r}{r_b}\right)^\alpha\right]^{(\frac{\gamma-\beta}{\alpha})}, \quad (4)$$

where  $r_b$  is called the break radius, and it's the place where the curvature of the profile reaches its maximum and  $I_b$  is the SB at  $r_b$ .

Recently, Baes (2020) showed that the Sérsic profile is a unique subgroup of the Nuker function; hence, the de Vaucouleurs profile, the exponential profile, and the Gaussian profile discussed above become specific cases of the Nuker profile, as well.

5. **Edge-on disk (van der Kruit & Searle 1981)**: This profile is used to fit the vertical structure of edge-on spiral galaxies. It results after considering an locally isothermal thin disk (van der Kruit & Searle 1981):

$$I(r, z) = I_0 \left(\frac{r}{r_s}\right) K_1\left(\frac{r}{r_s}\right) \operatorname{sech}^2\left(\frac{z}{z_h}\right), \quad (5)$$

where  $I_0 = I(0, 0)$  is the central SB,  $r_s$  is the scale length on the disk's major axis,  $z_s$  is the scale high on the  $z$  axis perpendicular to the disk, and  $K_1$  is the modified Bessel function of the second kind.

6. **Ferrers**<sup>10</sup>: This model is named after the triaxial potential introduced by Norman Macleod Ferrers (Ferrers 1877; Binney & Tremaine 1987, pp. 61-62). The modified Ferrer profile programmed in `GALFIT 3` is given by the following expression:

$$I(r) = I_0 \left[1 - \left(\frac{r}{r_{out}}\right)^{(2-\beta)}\right]^\alpha, \quad (6)$$

which shows an almost flat core and an outer truncation, and whose sharpness of the truncation is set by  $\alpha$ , while the central slope is determined by  $\beta$ ,  $I_0$  is the central SB. This profile is only valid for  $r \leq r_{out}$ , i.e.,  $I(r \geq r_{out}) \equiv 0$ .

<sup>10</sup>Peng et al. (2010) called it modified Ferrer profile.

This profile has been used to fit bars and lenses; it has eight free parameters. Notwithstanding, a Sérsic profile whose index is  $n < 0.5$  approximates the Ferrers profile; therefore, either profile can be used to model the SB of bars. Besides, the results are indistinguishable (e.g., Kim et al. 2015).

7. **Empirical King (Elson 1999):** It is given as follows:

$$I(r) = I_0 \left[ 1 - \frac{1}{\left(1 - \left(\frac{r_t}{r_c}\right)^2\right)^{\frac{1}{\alpha}}} \right]^{-\alpha} \left[ \frac{1}{\left(1 - \left(\frac{r}{r_c}\right)^2\right)^{\frac{1}{\alpha}}} - \frac{1}{\left(1 - \left(\frac{r_t}{r_c}\right)^2\right)^{\frac{1}{\alpha}}} \right]^{\alpha}, \quad (7)$$

where the core radius  $r_c$ , the truncation radius  $r_t$ , and the power law index  $\alpha$  are free parameters; however, in the standard modified King profile,  $\alpha = 2$  is fixed. This profile fits the SB of globular clusters.

8. **Gaussian profile** is used to fit PSF profiles, the contribution of active galactic nuclei and bars; it's a special case of the Sérsic profile when  $n = 0.5$ . It is described by the following equation:

$$I(r) = I_0 \exp\left(\frac{-r^2}{2\sigma^2}\right), \quad (8)$$

the scale is parametrized by the full width at half maximum  $FWHM = 2\sqrt{2 \ln 2} \approx 2.355 \sigma$ , instead of  $r_e$ .

9. **Moffat (Moffat 1969)** is a profile used to fit PSF profiles or the emission of active nuclei of galaxies. The following equation describes it:

$$I(r) = \frac{I_0}{\left(1 + \left(\frac{r}{r_d}\right)^2\right)^n}, \quad (9)$$

where  $n$  is the concentration index; as for the Gaussian profile, the scale is given by  $FWHM = 2 \left(2^{\frac{1}{n}} - 1\right)^{\frac{1}{2}} r_d$ . Nevertheless, the Gaussian profile is more widely used than the Moffat.

10. The user could provide the **PSF** as an image. However, care should be taken when selecting the stellar image's centroid, intensity, and sampling (see **GALFIT 3** user's manual).
11. **The background sky** is modeled as a plane, with an optional gradient applied along the  $x$  and  $y$  directions.
12. **Spiral modes** are used to model the spiral arms or asymmetrical galaxies; see Peng et al. (2010) and **GALFIT 3** user's manual for more details.

With **GALFIT 3**, the user can apply multiple profiles to fit the following components: bulges, disks, bars, spiral arms, and asymmetries to one or several galaxies simultaneously. This gives a substantial advantage in crowded galaxy fields, such as compact groups or clusters of galaxies. Galaxies being extended objects are modeled better when the galaxy of interest is fitted together with its closely neighboring galaxies instead of masking. This has been shown through simulations (Häussler et al. 2007; Añorve 2012).

**GALFIT 3** applies a  $\chi^2_\nu$  optimization scheme for parameter selection, including the background, generating parameter uncertainties, and produces a FITS (Flexible Image Transport System) image cube containing the image of the input object, the model image resulting after the fit, and the residual image. However, **GALFIT 3** does not include a graphical tool to compare the input image with its resulting model image. Users commonly applied IRAF's *ellipse*; nevertheless, this suite of programs became difficult to access after IRAF's discontinuation in 2013; fortunately, IRAF's *ellipse* is implemented in the Astropy library and new releases through. In any case, the main difficulty is that one needs to adequate **GALFIT 3**'s output format to the format admitted by external codes. However, this process could be time-consuming and prone to mistakes in handling.

Additionally, in crowded fields, neighboring galaxies near the target galaxy can be fitted simultaneously using **GALFIT 3**, resulting in more accurate outcomes. Häussler et al. (2007) demonstrated through simulations that masking galaxies in such fields leads to inaccurate fits for both the target galaxy and its neighbors. Improved fits were achieved when galaxies nearby were fitted simultaneously. However, this method is not feasible with IRAF's *ellipse* task, which processes only one galaxy at a time.

Hence, we are introducing **EllipSect**, a new Python tool that generates SB profiles and extracts complementary information directly from **GALFIT 3**'s output. We provide an easy-to-use tool to help users select and improve their models by introducing additional components or removing redundant components. **EllipSect**'s output includes data from the observed galaxy, and its model fits in the form of one-dimensional profiles. For a more straightforward assessment of the results, **EllipSect** can handle individual model components for a detailed analysis.

Additionally, **EllipSect** complements **GALFIT 3** analysis by calculating integrated magnitudes, mean SB at  $r_e$  ( $\langle\mu\rangle_e$ ), the SB at  $r_e$  ( $\mu_e$ ), the radius at 90% of total light, Kron radius (Kron 1980), Petrosian radius (Petrosian 1976), bulge-to-total ratio, and the total effective radius ( $r_e^t$ ) of a galaxy when multi-components are used.

This paper is organized as follows: §1 presents a general introduction to **GALFIT 3** and the built-in profiles, §2 explains **EllipSect**'s implemented algorithms. §3 shows applications and examples, §4 and §5 presents a discussion and conclusions, respectively. The Appendices show how to install and run **EllipSect**.

## 2. ELLIPSECT

This section explains how `EllipSect` works. We have divided it into two parts: first, we explain the main algorithm, and then, we explain the photometric measurements. Throughout this article, we refer to the individual profiles that make a SB model the *components*, while the set of SB components is called the *model*.

### 2.1. The Algorithm

`EllipSect` was designed to be user-friendly for most users. It omits any direct interaction with the code or translation of `GALFIT 3`'s data format. The program is efficiently executed via the command line. It only requires the latest `GALFIT 3`'s model output files.

`GALFIT 3`'s output files consist of a data cube in FITS format, which includes the original image but chopped into the fit region, the model image, and residual image (i.e., galaxy image – model image), a file named `galfit.nn` (where `nn` indicates the number of attempts run by `GALFIT 3`). This last file contains the parameters that fit each component of the model. Additionally, `GALFIT 3` creates a log file including the model parameter values and their errors. `EllipSect` uses the `GALFIT` output file (`galfit.nn`) to parse the input magnitude zero point, plate scale, mask file name, output file, and model parameters.

The output `GALFIT 3` file may contain information about the simultaneous fit of neighboring galaxies to the target galaxy. Using the distances among model components, `EllipSect` can discern which components belong to each galaxy.

`EllipSect` uses the same mask employed by `GALFIT` to remove the undesired pixels from the image. Once this is done, `EllipSect` reads the sky component from the `GALFIT` output file and subtracts it from the output galaxy and model images. The program divides the galaxy and its model into sectors equally spaced in angle around their centers. This is accomplished using the `sectors_photometry` routine from the `mgefit`<sup>11</sup> Python library by Cappellari (2002). The routine also calculates the counts within each sector, extending outward testing the galaxy's center until a specified minimum count level is reached. This is set by the `MINLEVEL` parameter of the routine. The default value is zero for this parameter. The ellipse is divided into sectors from the major axis to the minor axis, with the number of sectors set to a default value of 19, resulting in sectors spanning an angle of 5°. Sectors around the galaxy are built using the position angle and ellipticity found in the model component of the output `GALFIT` file. If multiple components are fitted to the galaxy, the geometric information is taken from the last component in the model, as it is identified as the galaxy's outermost component. This component is typically an exponential model, although this may vary depending on the specific galaxy model used. The user can change all these default parameters.

<sup>11</sup>`mgefit` is a software to obtain Multi-Gaussian Expansion (MGE) fits galaxy images (Cappellari 2002)

**EllipSect** generates publication-quality graphs. It produces two basic graphs in its simplest mode: one showing the averaged surface brightness along the major axis and another displaying the SB for various azimuthal angles across multiple plots (see examples in the Results section). Additionally, this mode generates a *png* image featuring the galaxy, model, and residual. **EllipSect** can also export a file containing the data points for each plot, allowing users to create custom visualizations with other plotting routines. Optionally, the SB profile for each plot component can include the galaxy (via the `--comp` flag). This feature allows users to visualize the behavior of model components and identify irrelevant components or redundancies (see, for example, Figure 3).

Based on the *mgefit* manual, the SB is calculated according to the following formula<sup>12</sup>:

$$\mu_I = zpt + 5 \times \log(Pl_{scc}) + 2.5 \times \log(t_{exp}) - 2.5 \times \log(C_0) + 0.1, \quad (10)$$

where  $zpt$  is the magnitude zero point,  $C_0$  are the  $counts \times pixel^{-1}$ ,  $Pl_{scc}$  is the plate scale,  $t_{exp}$  is the exposition time, and 0.1 is a correction for infinite aperture.

## 2.2. Output photometry

In addition to generating surface brightness plots, **EllipSect** calculates additional photometric variables from the models beyond those provided by **GALFIT 3**. These variables are stored in an output file and are derived after **EllipSect** extracts the count values for each sector. This section describes these variables and explains their computation.

**EllipSect** connects to NED through the command `curl` to download a *xml* file to extract the photometric corrections and compute the absolute magnitude according to the appropriate band. In the case of luminosity, it is calculated using the absolute magnitude of the Sun taken from Willmer (2018). This file extracts the distance modulus and the galactic extinction. It also obtains a conversion parameter used to compute the radius in Kpc units from angular units. SB dimming also corrects the mean SB at effective radius  $\langle \mu \rangle_e$  and SB at the effective radius  $\mu_e$ . The only correction not applied is the k-correction since it needs the color information that is not always available<sup>13</sup>. However, correction through galactic morphology can be used as well. It is important to note that corrections are applied only to the quantities in the output file; the single and multiple plots mentioned earlier are not adjusted for these corrections. However, users can apply a correction factor as a command-line argument or generate plots using the data from the **EllipSect** output files.

<sup>12</sup>this does not include the extinction parameter

<sup>13</sup>see <http://kcor.sai.msu.ru/>

NED provides two distance modulus measurements: one derived from cosmological models and another obtained directly from the galaxy. By default, `EllipSect` uses the redshift-independent measurement to calculate the luminosity, if available. Otherwise, it uses the distance modulus based on cosmological models.

Most output photometric variables are calculated within the ellipse corresponding to the radius enclosing 90% of the total light. Key variables include Bumpiness (Blakeslee et al. 2006), tidal parameter (Tal et al. 2009), reduced chi-squared ( $\chi^2_\nu$ ) within this ellipse (as opposed to the value provided by `GALFIT 3`, which applies to the entire fitted image section), signal-to-noise ratio (S/N), residual sum of squares, number of degrees of freedom, number of free parameters, absolute magnitude, the bulge-to-total ratio, and luminosity.

The Bumpiness parameter quantifies the irregular structure of a galaxy by measuring deviations in its SB from a smooth SB model. Blakeslee et al. (2006) employed this parameter and the Sérsic index to classify galaxies into elliptical, S0, spiral, and irregular types.

The Bumpiness parameter is computed as follows:

$$B = \frac{\left(\frac{1}{N} \sum_{r_{i,j} < 2r_e} [I_{i,j} - S(x_i, y_j | r_e, n)]^2 - \sigma_s^2\right)^{\frac{1}{2}}}{\frac{1}{N} \sum_{r_{i,j} < 2r_e} S(x_i, y_j | r_e, n)}, \quad (11)$$

The sum is performed over two effective radii ( $2r_e$ ), but in this case, it is applied within the ellipse region defined above. Here,  $N$  represents the number of unmasked pixels included in the computation,  $I_{i,j}$  is the galaxy intensity at pixel  $(i, j)$ ,  $S$  is the convolved surface brightness (SB) model, and  $\sigma_s^2$  is the estimated photometric error, which is taken directly from the sigma image produced by `GALFIT 3`.

Another useful parameter is the `Tidal` (Tal et al. 2009) parameter is defined as:

$$T_{gal} = \left[ \left( \frac{I_{(x,y)}}{M_{(x,y)}} \right) - 1 \right], \quad (12)$$

where  $I_{(x,y)}$  are the galaxy pixel intensity values in  $(x, y)$  and  $M_{(x,y)}$  represent the intensity values of the model.

This parameter measures the differences between the galaxy image and its corresponding model. Tal et al. (2009) used this to measure the tidal disturbances in elliptical galaxies and explored its relation with environmental properties, AGN activity, and color.

Additionally, suppose the galaxy has been fitted with multiple Sérsic components (or variants such as Gaussian and exponential models). In that case, the program gives  $r_e^t$ . While `GALFIT 3` provides  $r_e$  for each component, the program determines  $r_e^t$  by summing the surface brightness model of all components, calculating half of the total flux, and identifying the corresponding radius using the bisection method.

The Akaike Information Criterion (AIC) (Akaike 1974) and the Bayesian



Information Criterion (BIC) (Schwarz 1978) determines which model best fits the galaxy among a given set. Both criteria penalize adding new parameters by increasing the estimate of  $\chi_\nu$ . The model with the lowest AIC or BIC value is considered the best fit.

`EllipSect` computes both criteria and also calculates an alternative Bayesian Information Criterion (BIC) using the number of resolution elements instead of the number of pixels defined by  $N_{res} = \frac{n_{pix}}{\pi(\text{FWHM})^2}$ .

Finally, additional output parameters are computed for each model component, including the fraction of light (such as the bulge-to-total ratio for each component),  $\langle\mu\rangle_e$ ,  $\mu_e$ , absolute magnitude, luminosity radius at 90% of the total light, Kron radius (Kron 1980), and Petrosian radius (Petrosian 1976).

### 2.2.1. Sky computation

An incorrect sky background estimation can lead to inaccurate surface brightness model parameters, especially the Sérsic index. An appropriate strategy is determining the sky background and fixing it during model fitting. This reduces the number of free parameters and prevents parameter degeneracy.

`EllipSect` provides two methods for computing the sky background: The Sky gradient method and the random box method. These methods are not used to estimate the photometric variables discussed above. Instead, they enable a comparison with the sky value computed by `GALFIT 3`, allowing the user to decide whether to refine the model using the newly derived sky value from `EllipSect`.

**Sky gradient method.** This method follows Barden et al. (2012). In this approach, `EllipSect` constructs concentric rings using the same angular parameters and ellipticity employed for the elliptical sectors described in 2.1. The user specifies the ring width, and `EllipSect` enlarges the rings' major axes outward from the galaxy's center. A mask file is used to exclude undesired pixels. Optionally, `EllipSect` can remove the top 80% and bottom 20% of sky values in each ring. Finally, it computes the gradient based on the mean sky values of each ring.

A negative gradient arises in the galaxy's inner regions because the central counts exceed those in the outskirts. Beyond this zone, the gradient fluctuates randomly between positive and negative values (see Figure 5 in Barden et al. (2012)). `EllipSect` generates a fit image with sky rings to illustrate this gradient. It begins calculating the gradient from an initial radius and stops when it becomes positive for the second time, following the recommendation by Barden et al. (2012). At that point, `EllipSect` returns the sky value determined for this radius. Figure 1 shows an example of an elliptical ring used to compute the sky.

### 2.2.2. Random box method

For this method, `EllipSect` calculates the sky using randomly placed boxes around the galaxy. To achieve this, it first creates an elliptical mask




Fig. 1. Example of sky computation using the gradient method. The ring around the bright cluster member of Abell 2029 indicates the radius at which the sky was computed for this galaxy. For visualization purposes, all pixels in that elliptical ring are assigned the same numerical value corresponding to the ellipse’s major-axis length.

that covers the entire galaxy and excludes pixels in the original input mask file. By default, the geometry for this elliptical mask comes from the `GALFIT 3` output file, although the user may modify it.

Next, the program positions boxes randomly between this elliptical mask and a specified maximum radius (often at the image boundary). It can also optionally remove the top 80% and bottom 20% of pixel values in these boxes. The final sky value is derived by computing the mean, median, and standard deviation of all pixels in the boxes. The default settings place 20 boxes, each measuring 20 pixels across. This method follows the approach described in Gao & Ho (2017).




Fig. 2. Example of cube image generated by `EllipSect`. The left panel shows the galaxy, the middle panel is the model, and the right is the residual. The user can change the color palette, brightness, and contrast.

### 3. RESULTS: EXAMPLES OF SB PLOTS

`EllipSect` has been used to analyze galaxy images from 2MASS (Two Micron All Sky Survey) in which estimations of morphological and structural parameters have been obtained through photometric decompositions using `GALFIT 3` (Ríos-López et al. 2021). Moreover, `EllipSect` was used in a study of dust nuclear structures of a sample of Active Galactic Nuclei (AGN) in the local Universe through the Hubble Space Telescope (HST) images, SB models, and radiative transfer simulations (Reyes-Amador 2021). The code has also been used in reports, conference proceedings, professional papers, and undergraduate and graduate theses (Ríos-López 2021, Valerdi M. et al. paper in progress; Ríos-López et al. paper in progress).

Here, we present examples of surface brightness profile plots for three images: Holm 15A with Kitt Peak National Observatory (KPNO), M61 with 2MASS, and NGC4261 with the Hubble Space Telescope (HST).

#### 3.1. Kitt Peak: Holm 15A

This example shows the SB profile plots of Abell 85 brightest cluster galaxy (BCG), the cD galaxy Holm 15A. Here, `GALFIT 3` was used to fit a multi-gaussian component to model its SB; another analysis using the Nuker profiles has already been published (López-Cruz et al. 2014). This frame was taken with KPNO 0.9m Telescope, part of the catalog of LOCOS (López-Cruz 1997). Details on the selection criteria, observations, and data reduction can be found in López-Cruz (1997), López-Cruz et al. (2004), and López-Cruz et al. (2025, in preparation). In Figure 2, we show the cube image generated by `EllipSect` for this galaxy.

In Figure 3, we present the single-plot and multiplot of the surface brightness for the brightest cluster member of Abell 85. This galaxy was modeled using seven Gaussians to capture its full SB profile. The multiplot displays the error, while the single-plot shows the percent error. These plots were generated automatically by `EllipSect`, which computes the SB profiles of the galaxy, model, and individual components similar to IRAF's `ellipse` task.




Fig. 3. Example of `EllipSect` output for the elliptical galaxy Holm 15a and its model. This was fitted using seven Gaussian components. In both panels, red represents the galaxy and blue represents the `GALFIT 3` model. The color for each Gaussian component is shown in the top right inset box. These colors represent the same for both panels. Left panel: average SB vs. radius along the major axis. The residual percentage is shown in the bottom left plot. The model errors depend on the average over the ellipse for the model in that radius. Right panel: Multiple plots of galaxy SB and model at different angles from the major axis (the major axis is the one with 0 deg). The error is shown on the right side of the multiplot. The angle in red at the top right indicates the angle measured from the Y-axis of the image. On the other hand, the subtle grey angle at the bottom left indicates the angle measured from the major axis starting from the top with the major axis and the bottom plot with the minor axis




Fig. 4. Example of cube image generated by `EllipSect` for M 61 galaxy. Details for this image are the same as the caption of Figure 2.

### 3.2. *2MASS: M61*

In Figure 4, we show the galaxy, model, and residual for galaxy M 61 in the *Ks* band of 2MASS.

Figure 5 shows the multiplot of the fitting of M 61. This model consists of a series of Sérsic and Exponential components. As can be seen, deviations from the model can be appreciated in the region where the model fails to reproduce the SB model of the galaxy.




Fig. 5. Example of `EllipSect` output for M 61 galaxy and its model. The description of the image is the same as in the right panel of Figure 3




Fig. 6. Example of cube image generated by `EllipSect` for NGC4261 galaxy. Details for this image are the same as the caption of Figure 2.

### 3.3. *HST*: NGC4261

In the last example, we show an image from *HST*; this is NGC4261 and in Figure 6 is shown the galaxy, model, and residual.

Figure 7 shows the galaxy NGC4261 taken with the *HST* using filter F547M. This model used a single Sersic component to fit the galaxy's SB.

## 4. DISCUSSION

`EllipSect` was developed as an extension to `GALFIT 3` to generate SB profile plots and offer additional photometric information beyond `GALFIT 3`. Its




Fig. 7. Example of `EllipSect` output for NGC4261 galaxy and its model. The description of the image is the same as the right panel of Figure 3.

primary purpose is to test the reliability of `GALFIT 3`'s model at various radii and angles for a given galaxy. In addition, `EllipSect` allows users to include individual components of the `GALFIT 3` models in the plots, enabling verification that these components behave as expected. This feature streamlines the analysis process by eliminating the need to switch between different program formats, making it practical to quickly confirm that the `GALFIT` model is appropriate for the galaxy. `EllipSect` aids `GALFIT 3` users with a readily accessible and easy-to-use tool for generating accurate SB measurements and publication-quality plots.

Using the *sectors-photometry* routine, `EllipSect` creates evenly spaced angular sectors to compute surface brightness at various radii. It divides an ellipse, centered on the galaxy, into multiple sectors according to the geometric information from the `GALFIT 3` output file. By default, `EllipSect` derives the angular orientation and ellipticity from the last component of the `GALFIT 3` output file (see A.2).

In the conventional approach to generating SB profiles, one typically uses IRAF's *ellipse*. This routine fits an ellipse to each isophote at every radius, from the central region to the galaxy's outskirts. As the radius changes, the ellipticity and position angle of the ellipse may also vary.

One disadvantage of `EllipSect` is that it maintains a fixed ellipticity and angular position values throughout the entire galaxy, which may hinder studying how these parameters change with radius. However, this fixed parameter

approach is advantageous when dealing with galaxies simultaneously fitted by `GALFIT 3`. In such cases, masking or removing the galaxy is insufficient, as the galaxies are blended in the line of sight, and it is challenging to distinguish where one galaxy ends and the other begins (Haussler et al. 2007). In addition, `EllipSect` avoids the need for "double" fitting, i.e., two-dimensional and one-dimensional fitting, which IRAF's *ellipse* requires. One-dimensional routines may not be capable of simultaneous fitting of isophotes for two neighboring galaxies, making simultaneous fitting difficult or impossible. While `EllipSect` is not a complete substitute for IRAF's *ellipse*, it offers distinct advantages depending on the scientific question being addressed.

Despite the widespread use of IRAF's *ellipse*, alternative techniques such as `EllipSect` offer unique advantages for SB analysis, which may prove to be beneficial for astronomers seeking to enhance their understanding of galactic structures.

An important output parameter from `EllipSect` is  $r_e^t$ , the effective radius of the entire galaxy, which can be used to parameterize metallicity gradients and other relevant information (e.g., Garduño et al. 2023). If the galaxy has only a single Sérsic component,  $r_e$  from that component can represent the whole galaxy. However, no single component's  $r_e$  can accurately describe the entire galaxy when multiple components are present, because galaxies do not perfectly follow the surface models.

To resolve this, `EllipSect` employs a numerical method that adds together the models of all components and then locates the radius where the total luminosity reaches half of its overall value ( $r_e^t$ ). This procedure can also be applied to determine the radius corresponding to any percentage of the total light.

Determining which aperture to use for a galaxy's photometric variables can be somewhat arbitrary. Different photometric quantities require a suitable radius for proper evaluation, and standardizing this across various galaxies is difficult. In the case of `GALFIT 3`, the reduced chi-squared ( $\chi_\nu^2$ ) is computed using all pixels within the fitting region. However, changing the size of this region can alter  $\chi_\nu^2$  and may include pixels from other objects, introducing additional noise. Consequently, the  $\chi_\nu^2$  reflects the fit quality for every object within that region.

In contrast, `EllipSect` uses the radius containing 90% of the total light to compute various photometric variables. This radius is found by applying the same procedure used to determine the effective radius but at 90% of the total light. Focusing on this region ensures that only pixels belonging to the galaxy are included, minimizing contamination from external objects. Within this radius, `EllipSect` calculates several photometric variables, such as the tidal parameter, Bumpiness, AIC, BIC, and a local  $\chi_\nu^2$ .

`EllipSect` provides corrections for galaxy extinction and surface brightness (SB) dimming, but does not perform k-corrections at this time. Users can still apply morphological-based corrections to the galaxy, although this may be imprecise for cluster galaxies whose colors can differ from those of

field galaxies. In such situations, manual correction is recommended. Note that k-corrections can be generated using online tools such as <http://kcor.sai.msu.ru/>.

`EllipSect` provides a helpful tool for wrapping codes aimed at automating the `GALFIT 3` fitting process. It can efficiently compute the necessary photometric variables required by these codes, thereby allowing for rapid analysis of model outputs for multiple galaxies, which would otherwise be time-consuming. In fact, `EllipSect` has been specifically designed to be used in conjunction with the Driver for `GALFIT` on Cluster Galaxies (DGCG) code (Añorve et al., in preparation), which automates `GALFIT 3` fitting for galaxies in cluster images.

## 5. CONCLUSIONS AND FUTURE WORK

The paper introduces `EllipSect`, a Python tool for analyzing `GALFIT 3` output files. It generates surface brightness profiles for galaxies, models, and individual components, offering single-plot and multiplot visualizations. Additionally, `EllipSect` computes various photometric parameters, including light fractions for each component,  $r_e^t$ ,  $\langle\mu\rangle_e$ ,  $\mu_e$ , absolute magnitudes, the luminosity radius 90%, the effective radius of the galaxy, the Kron radius, and the Petrosian radius. It also calculates tidal and Bumpiness parameters, signal-to-noise ratios, and local  $\chi_\nu^2$  values to assess differences between models and observed galaxies.

`EllipSect` offers two methods for computing the sky background and enables users to distinguish among different models for galaxies by calculating the Akaike and Bayesian information criteria. The code can be easily adapted to other `GALFIT 3` wrapping codes and will be integrated with DGCG in future work. This project aims to apply `EllipSect` to analyze data images containing hundreds of galaxies, such as galaxy clusters.

`EllipSect` is a robust tool for analyzing `GALFIT 3` output and extracting extensive photometric information for galaxies. Its flexibility and versatility make it an essential tool for studying galaxy properties and morphology. Future development will focus on expanding its capacity to handle larger datasets and integrate it with other `GALFIT 3` wrapping codes.

## APPENDICES

### A. HOW TO INSTALL AND RUN ELLIPSECT

#### A.1. *Installing EllipSect*

The code is written for *Python 3*. It can be downloaded from the site: <https://github.com/canorve/EllipSect>

##### A.1.1. *Quick installation*

The quickest way to install `EllipSect` is just by typing this into the command line:

```
1 pip install ellipse
```



### A.1.2. Step-by-step installation

The following libraries are required to install `EllipSect` (v3.4.4) along with Python  $\geq$  v3.8:

- `numpy`  $\geq$  v1.20.3
- `astropy`  $\geq$  v5.1
- `scipy`  $\geq$  v1.5.2
- `matplotlib`  $\geq$  v3.5.2
- `mgefit`  $\geq$  v5.0.13
- `sphinx`  $\geq$  v3.2.1

After installing the libraries, download the latest release from the link shown above, and install it via:

```
1 cd EllipSect
2 pip install .
```

or

```
1 cd EllipSect
2 python setup.py install
```

To check that everything is working properly, run the automated tests:

```
1 python -m pytest
```

Although `GALFIT` is not strictly required, it will require to create the model components and sigma image. Ensure you can call `GALFIT` from the command line. Otherwise, the automated tests will fail.

### A.2. Running *EllipSect*

**Easy run:** Once installed, run the command `EllipSect` in the same directory that you run `GALFIT`. The easiest way to run the program is:

```
1 EllipSect GALFIT.01
```

**Note:** `EllipSect` takes the `GALFIT` output files (`GALFIT.XX`) where `XX` is the output file number you want to plot.

**More options:** As explained in the paper, `EllipSect` uses an ellipsoid-based grid to compute the SB. The information of this ellipse is obtained from the `GALFIT` output file, but if the user wants to provide their ellipse parameters, this is done via:

```
1 EllipSect GALFIT.01 -q 0.5 -pa 80
```

In this last example, an ellipse with a 0.5 axis ratio and position angular of 80 (measured from Y-axis) was used to create the plots.

In case your model contains two or more components, the `comp` (or `cp`) option will plot each component (for an example of this check figure 3):

```
1 EllipSect GALFIT.01 -cp
```

There are many options to change the format of the plots such as: `logx`, `ranx`, `rany`, `pix`, `grid`, `dpi`, `galax`, but if the user prefers to make their plots the option `sabout` will create a file (or files) containing the points of the the SB of the galaxy and model at each radius.

To compute the photometry of the galaxy and model such as `snr`, `BIC`, `AIC`, `absolute Magnitude`, `Luminosity`, `Bumpiness`, `Tidal` and other photometric variables the option `phot` creates a file with the computed variables:

```
1 EllipSect GALFIT.01 --phot
```

`EllipSect` has the option to compute the sky to compare with or provide to `GALFIT`. As explained above, the program uses two methods to compute it: The gradient method and the random box method. To compute the gradient method:

```
1 EllipSect GALFIT.01 -gsky
```

And to compute using the random box method:

```
1 EllipSect GALFIT.01 -rsky
```

The ring's width and box's size can be modified from their default options. For more options and to see how to use them, type:

```
1 EllipSect --help
```

#### A.2.1. Running *EllipSect* as a script

If you want to use `EllipSect` inside your Python script, you can call it in the following way:

```
1 from ellipsect.inout.read import ArgParsing
2 from ellipsect.sectors.sect import SectorsGalfit
3
4 args=['GALFIT.01', '--logx', '--phot', '--noplot']
5 parser_args = ArgParsing(args)
6 photapi = SectorsGalfit(parser_args)
7
8 print("Akaike Criterion: ", photapi.AICrit)
9 print("Bulge to Total: ", photapi.BulgeToTotal)
```

Listing 1: Example Python Code with `EllipSect`

A manual and more details can be found in <https://github.com/canorve/EllipSect>

## REFERENCES

- Añorve, C. 2012, PhD thesis, INAOE
- Akaike, H. 1974, *IEEE Transactions on Automatic Control*, 19, 716
- Baes, M. 2020, *A&A*, 634, A109
- Barden, M., Häußler, B., Peng, C. Y., McIntosh, D. H., & Guo, Y. 2012, *MNRAS*, 422, 449
- Binney, J. & Tremaine, S. 1987, *Galactic dynamics*
- Blakeslee, J. P., Holden, B. P., Franx, M., Rosati, P., Bouwens, R. J., Demarco, R., Ford, H. C., Homeier, N. L., Illingworth, G. D., Jee, M. J., Mei, S., Menanteau, F., Meurer, G. R., Postman, M., & Tran, K.-V. H. 2006, *ApJ*, 644, 30
- Cappellari, M. 2002, *MNRAS*, 333, 400
- Ciotti, L. & Bertin, G. 1999, *A&A*, 352, 447
- de Vaucouleurs, G. 1948, *Annales d’Astrophysique*, 11, 247
- Elson, R. A. W. 1999, in *Globular Clusters*, 209–248
- Ferrers, N. M. 1877, *Quart. J. Pure and Appl. Math.*, 14, 1
- Freeman, K. C. 1970, *ApJ*, 160, 811
- Gao, H. & Ho, L. C. 2017, *ApJ*, 845, 114
- Garduño, L. E., Zaragoza-Cardiel, J., Lara-López, M. A., Zinchenko, I. A., Zerbo, M. C., De Rossi, M. E., Fritz, J., Dib, S., Pilyugin, L. S., Sánchez-Cruces, M., Heesen, V., O’Sullivan, S. P., López-Cruz, O., Valerdi, M., & Rosado, M. 2023, *MNRAS*, 526, 2479
- Graham, A. W. & Driver, S. P. 2005, *Publ. Astron. Soc. Aust.*, 22, 118
- Häussler, B., McIntosh, D. H., Barden, M., Bell, E. F., Rix, H.-W., Borch, A., Beckwith, S. V. W., Caldwell, J. A. R., Heymans, C., Jahnke, K., Jogee, S., Koposov, S. E., Meisenheimer, K., Sánchez, S. F., Somerville, R. S., Wisotzki, L., & Wolf, C. 2007, *ApJS*, 172, 615
- Jedrzejewski, R. I. 1987, *MNRAS*, 226, 747
- Kim, T., Sheth, K., Gadotti, D. A., Lee, M. G., Zaritsky, D., Elmegreen, B. G., Athanassoula, E., Bosma, A., Holwerda, B., Ho, L. C., Comerón, S., Knapen, J. H., Hinz, J. L., Muñoz-Mateos, J.-C., Erroz-Ferrer, S., Buta, R. J., Kim, M., Laurikainen, E., Salo, H., Madore, B. F., Laine, J., Menéndez-Delmestre, K., Regan, M. W., de Swardt, B., Gil de Paz, A., Seibert, M., & Mizusawa, T. 2015, *ApJ*, 799, 99
- Kron, R. G. 1980, *ApJS*, 43, 305
- Lauer, T. R., Ajhar, E. A., Byun, Y. I., Dressler, A., Faber, S. M., Grillmair, C., Kormendy, J., Richstone, D., & Tremaine, S. 1995, *AJ*, 110, 2622
- López-Cruz, O. 1997, PhD thesis, University of Toronto, Canada
- López-Cruz, O., Añorve, C., Birkinshaw, M., Worrall, D. M., Ibarra-Medel, H. J., Barkhouse, W. A., Torres-Papaqui, J. P., & Motta, V. 2014, *ApJ*, 795, L31
- López-Cruz, O., Barkhouse, W. A., & Yee, H. K. C. 2004, *ApJ*, 614, 679
- Moffat, A. F. J. 1969, *A&A*, 3, 455
- Patterson, F. S. 1940, *Harvard College Observatory Bulletin*, 914, 9
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, *AJ*, 124, 266
- . 2010, *AJ*, 139, 2097
- Petrosian, V. 1976, *ApJ*, 209, L1
- Reyes-Amador, C. 2021, Master’s thesis, Instituto de Radioastronomía y Astrofísica
- Ríos-López, E. 2021, PhD thesis, INAOE
- Ríos-López, E., Añorve, C., Ibarra-Medel, H. J., López-Cruz, O., Alvira-Enríquez,

- J., Iacobuta, G., & Valerdi, M. 2021, MNRAS, 507, 5952  
Schwarz, G. 1978, Annals of Statistics, 6, 461  
Sérsic, J. L. 1968, Atlas de Galaxias Australes  
Tal, T., van Dokkum, P. G., Nelan, J., & Bezanson, R. 2009, AJ, 138, 1417  
van der Kruit, P. C. & Searle, L. 1981, A&A, 95, 105  
Willmer, C. N. A. 2018, ApJS, 236, 47

Christopher Añorve: Facultad de Ciencias de la Tierra y el Espacio, Universidad Autónoma de Sinaloa, Blvd. de la Américas y Av. Universitarios S/N, Ciudad Universitaria, C.P. 80010 Culiacán, Sinaloa, México (canorve@uas.edu.mx).

Omar Ulises Reyes-Amador: Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apartado Postal 3-72, 58090 Morelia, Michoacán, México (o.reyes@irya.unam.mx).

Emmanuel Ríos-López: Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Luis Enrique Erro 1, Apartado Postal 51 y 216, C.P. 72840 Puebla, México; Instituto de Astrofísica de Canarias (IAC), Vía Láctea s/n, 38205, La Laguna, Tenerife, España; Universidad Pedagógica del Estado de Sinaloa (UPES), Castiza s/n, 80027, Culiacán, Sinaloa, México (riloemm@inaoep.mx).

Omar López-Cruz: Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE), Luis Enrique Erro 1, Apartado Postal 51 y 216, C.P. 72840 Puebla, México (omarlx@inaoep.mx).

Diego de Ramón Tadeo: Facultad de Ciencias de la Computación, Benemérita Universidad Autónoma de Puebla, Av. San Claudio y 14 Sur, Ciudad Universitaria, C. P. 72570, Puebla, Puebla, México (deramontadeo@gmail.com).