DISTINGUISHED FILTRATIONS OF THE 0-HECKE MODULES FOR DUAL IMMACULATE QUASISYMMETRIC FUNCTIONS

SO-YEON LEE AND YOUNG-TAK OH

ABSTRACT. Let α range over the set of compositions. Dual immaculate quasisymmetric functions \mathfrak{S}^*_{α} , introduced by Berg, Bergeron, Saliola, Serrano, and Zabrocki, provide a quasisymmetric analogue of Schur functions. They also constructed an indecomposable 0-Hecke module \mathcal{V}_{α} whose image under the quasisymmetric characteristic is \mathfrak{S}^*_{α} . In this paper, we prove that \mathcal{V}_{α} admits a distinguished filtration with respect to the basis of Young quasisymmetric Schur functions. This result offers a novel representation-theoretic interpretation of the positive expansion of \mathfrak{S}^*_{α} in the basis of Young quasisymmetric Schur functions. A key tool in our proof is Mason's analogue of the Robinson-Schensted-Knuth algorithm, for which we establish a version of Green's theorem. As an unexpected byproduct of our investigation, we construct an indecomposable 0-Hecke module \mathbf{Y}_{α} whose image under the quasisymmetric characteristic is the Young quasisymmetric Schur function $\hat{\mathscr{T}}_{\alpha}$. Further properties of this module are also investigated. And, by applying a suitable automorphism twist to this module, we obtain an indecomposable 0-Hecke module whose image under the quasisymmetric characteristic is the quasisymmetric Schur function \mathscr{T}_{α} .

CONTENTS

| 2 |
|----|
| 5 |
| 5 |
| 6 |
| 7 |
| 8 |
| 11 |
| |
| 12 |
| 12 |
| 17 |
| 24 |
| |

Date: January 22, 2025.

²⁰²⁰ Mathematics Subject Classification. 20C08, 06A07, 05E10, 05E05.

Key words and phrases. 0-Hecke algebra, Representation, Filtration, Quasisymmetric function, Schur function.

S.-Y. LEE AND Y.-T. OH

| 4.1. Distinguished filtrations of \mathcal{V}_{α} | 25 |
|--|----|
| 4.2. Distinguished filtrations of X_{α} | 30 |
| 5. Indecomposable 0-Hecke modules for Young quasisymmetric Schur functions | |
| and quasisymmetric Schur functions | 33 |
| 5.1. Indecomposable $H_n(0)$ -modules \mathbf{Y}_{α} and $\boldsymbol{\phi}[\mathbf{Y}_{\alpha}]$ | 33 |
| 5.2. A surjection series containing \mathbf{Y}_{α} | 35 |
| 5.3. Weak Bruhat interval module structure for \mathbf{Y}_{α} | 37 |
| 6. Remarks concerned with future research | 45 |
| Appendix A. Examples of $H_n(0)$ -modules that have no distinguished filtrations | |
| with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ or $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$. | 46 |
| References | 49 |
| | |

1. INTRODUCTION

Let n be a nonnegative integer. The 0-Hecke algebra $H_n(0)$ is defined by setting q = 0in the generic Hecke algebra $H_n(q)$. However, the representation theory of $H_n(0)$ differs significantly from that of the generic Hecke algebra. In particular, $H_n(0)$ is neither semisimple nor representation-finite for $n \ge 4$; it is of tame type when n = 4 and of wild type for n > 4 (see [13, 14]).

The representation theory of 0-Hecke algebras has deep connections with quasisymmetric functions. Let $\mathcal{G}_0(H_n(0)\operatorname{-\mathbf{mod}})$ denote the Grothendieck group of the category $H_n(0)\operatorname{-\mathbf{mod}}$ of finitely generated left $H_n(0)\operatorname{-mod}$ used and let QSym be the ring of quasisymmetric functions. In [15, 6], it has been shown that the direct sum $\bigoplus_{n\geq 0} \mathcal{G}_0(H_n(0)\operatorname{-\mathbf{mod}})$, equipped with a Hopf algebra structure defined by the induction product and restriction, is isomorphic to QSym under the map

ch :
$$\bigoplus_{n \ge 0} \mathcal{G}_0(H_n(0)\operatorname{-\mathbf{mod}}) \to \operatorname{QSym}, \quad [\mathbf{F}_\alpha] \mapsto F_\alpha,$$

known as quasisymmetric characteristic. Here, α is a composition, \mathbf{F}_{α} is the irreducible $H_{|\alpha|}(0)$ -module indexed by α , and F_{α} is the fundamental quasisymmetric function indexed by α (see Section 2.3).

Given an index set I, let $\mathcal{B} = \{\mathcal{B}_{\alpha} \mid \alpha \in I\}$ be a linearly independent subset of the *n*th homogeneous component QSym_n of QSym , where each \mathcal{B}_{α} has a positive expansion in the basis $\{F_{\alpha} \mid \alpha \models n\}$. For an $H_n(0)$ -module M, a distinguished filtration of M with respect to \mathcal{B} is defined to be an $H_n(0)$ -submodule series

$$0 =: M_0 \subsetneq M_1 \subsetneq M_2 \subsetneq \cdots \subsetneq M_l := M,$$

such that for each $1 \leq k \leq l$, there exists $\alpha \in I$ satisfying $ch([M_k/M_{k-1}]) = \mathcal{B}_{\alpha}$. It is important to observe that such a distinguished filtration of M with respect to \mathcal{B} may not exist, even if ch([M]) admits a positive expansion in terms of \mathcal{B} . The concept of

 $\mathbf{2}$

distinguished filtrations was introduced in [22, Section 6] in the context of studying 0-Hecke modules associated with regular Schur labeled posets. Roughly speaking, a regular Schur labeled skew shape poset P is a poset where the set of linear extensions forms a weak Bruhat interval and the P-partition generating function is symmetric. It was shown therein that for any regular Schur labeled poset P with underlying set $[n] := \{1, 2, \ldots, n\}$, the associated $H_n(0)$ -module M_P admits a distinguished filtration with respect to the Schur basis $\{s_{\lambda} \mid \lambda \vdash n\}$.

In this paper, we consider two bases for QSym_n : the basis $\mathcal{S} := \{\mathscr{S}_\alpha \mid \alpha \models n\}$ consisting of quasisymmetric Schur functions, and the basis $\hat{\mathcal{S}} := \{\mathscr{S}_\alpha \mid \alpha \models n\}$ consisting of Young quasisymmetric Schur functions. The quasisymmetric Schur functions are widely viewed as a natural quasisymmetric analogue of the Schur functions (see [19, 8, 20, 24]). Meanwhile, the Young quasisymmetric Schur functions are derived from the quasisymmetric Schur functions through the automorphism ρ : QSym \rightarrow QSym, defined by $F_\alpha \mapsto F_{\alpha^r}$, where α^r denotes the reverse of the composition α . With these bases, we study distinguished filtrations of the $H_n(0)$ -modules for the dual immaculate quasisymmetric functions and the extended Schur functions.

The dual immaculate quasisymmetric functions \mathfrak{S}^*_{α} , introduced by Berg, Bergeron, Saliola, Serrano, and Zabrocki [3], serve as quasisymmetric analogues of the immaculate noncommutative symmetric functions. The set { $\mathfrak{S}^*_{\alpha} \mid \alpha \models n$ } forms a basis for QSym_n, which has been extensively studied in the context of quasisymmetric functions and the representation theory of $H_n(0)$ (for instance, see [4, 7, 5, 11, 12]). In particular, in [4], the authors constructed an indecomposable $H_n(0)$ -module \mathcal{V}_{α} and showed that its image under the quasisymmetric characteristic is \mathfrak{S}^*_{α} . Meanwhile, Allen, Hallam, and Mason [1] demonstrated that the dual immaculate quasisymmetric functions expand positively in $\hat{\mathcal{S}}$. Therefore, it is natural to ask whether the module \mathcal{V}_{α} admits a distinguished filtration with respect to $\hat{\mathcal{S}}$.

The extended Schur functions \mathcal{E}_{α} , introduced by Assaf and Searles [2] as stable limits of lock polynomials, also constitute a basis for QSym. In [31, Theorems 3.10 and 3.13], Searles constructed an indecomposable $H_n(0)$ -module X_{α} and showed that its image under the quasisymmetric characteristic is \mathcal{E}_{α} . Recently, Marcum and Niese [25, Theorem 3.5] showed that if α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then \mathcal{E}_{α} expands positively in $\hat{\mathcal{S}}$. Analogous to the case of \mathcal{V}_{α} , this naturally leads to the question of whether X_{α} admits a distinguished filtration with respect to $\hat{\mathcal{S}}$.

The primary objective of this paper is to address the aforementioned questions. The main results obtained in this work are outlined below.

In Section 3, we deal with Mason's analogue of the Robinson-Schensted-Knuth algorithm, which is the most important tool in our study. In Subsection 3.1, following [26, Procedure 4.2], we introduce an algorithm that, given a two-line array w, produces a pair of Young composition tableaux ($\hat{P}(w), \hat{Q}(w)$) (Algorithm 3.3). We then investigate the similarities and differences between this algorithm and the Robinson-Schensted-Knuth

S.-Y. LEE AND Y.-T. OH

algorithm. In Subsection 3.2, we present a theorem that determines the shapes of the insertion tableaux $\hat{\mathsf{P}}(w)$ (Theorem 3.15). In the case of the Robinson-Schensted-Knuth algorithm, this problem was resolved by Greene [18]. For our purpose, we define the concept of the *initial entries set* of a longest k-increasing subsequence of w (see Definition 3.10).

In Section 4, we present answers to the aforementioned questions. It was shown in [21, Theorem 4.4] that both \mathcal{V}_{α} and X_{α} are equipped with the structures of weak Bruhat interval modules. Specifically,

$$\mathcal{V}_{\alpha} \cong \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})) \quad \text{and} \quad X_{\alpha} \cong \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathsf{T}'_{\alpha}))$$

(for details, see Subsection 2.5). In Subsection 4.1, we first define an equivalence relation \simeq_M on \mathfrak{S}_n , which refines the dual Knuth equivalence relation. We then show that $[\mathsf{w}_r(\mathscr{T}_\alpha), \mathsf{w}_r(\mathscr{T}'_\alpha)]_L \cdot w_0$ is closed under \simeq_M , and a suitable order relation exists among the shapes of $\hat{\mathsf{P}}(\sigma)$ for all $\sigma \in [\mathsf{w}_r(\mathscr{T}_\alpha), \mathsf{w}_r(\mathscr{T}'_\alpha)]_L \cdot w_0$. Finally, using these results, we prove that for any composition α of n, the module \mathcal{V}_α admits a distinguished filtration with respect to $\hat{\mathcal{S}}$ (Theorem 4.7). In Subsection 4.2, we restrict α to a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$. We first show that $[\mathsf{w}_r(\mathsf{T}_\alpha), \mathsf{w}_r(\mathsf{T}'_\alpha)]_L \cdot w_0$ is closed under \simeq_M . Then, following the approach used in the proof of Theorem 4.7, we prove that X_α admits a distinguished filtration with respect to $\hat{\mathcal{S}}$ (Theorem 4.10). It is worth noting that if λ is a partition of n, then $ch([X_\lambda]) = s_\lambda$. However, X_λ may not necessarily admit a distinguished filtration with respect to \mathcal{S} (see Appendix A and FIGURE A.1).

Section 5 concerns a significant open question in the representation theory of 0-Hecke algebras: whether there exists an indecomposable $H_n(0)$ -module M whose image under the quasisymmetric characteristic equals \mathscr{S}_{α} or $\hat{\mathscr{S}}_{\alpha}$. In 2015, Tewari and van Willigenburg [34] constructed an $H_n(0)$ -module \mathbf{S}_{α} that satisfies $ch([\mathbf{S}_{\alpha}]) = \mathscr{S}_{\alpha}$. However, \mathbf{S}_{α} is not generally indecomposable. In Subsection 5.1, we provide an indecomposable $H_n(0)$ -module \mathbf{Y}_{α} whose image under the quasisymmetric characteristic is $\hat{\mathscr{S}}_{\alpha}$ (Theorem 5.1). To be precise, for any distinguished filtration

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{l-1} \subsetneq M_l = \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha}))$$

of $\mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha}))$, as constructed in the proof of Theorem 4.7, we define \mathbf{Y}_{α} as the quotient M_l/M_{l-1} . It turns out that this module is independent of the choice of the filtration. We then provide a basis $\mathcal{K}_{\alpha} \subseteq [\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_L$ for \mathbf{Y}_{α} (see (5.2)). In Subsection 5.2, we construct a surjection series

$$\overline{\mathbf{P}}_{\alpha^{\mathrm{c}}} \twoheadrightarrow \mathcal{V}_{\alpha} \twoheadrightarrow \mathbf{Y}_{\alpha} \twoheadrightarrow \hat{\mathbf{S}}_{\alpha,C}$$

(Proposition 5.3). Additionally, we show that this series can be extended when α is obtained by shuffling a partition with (1^k) for some $k \ge 0$:

$$\overline{\mathbf{P}}_{\alpha^{\mathrm{c}}} \twoheadrightarrow \mathcal{V}_{\alpha} \twoheadrightarrow X_{\alpha} \twoheadrightarrow \mathbf{Y}_{\alpha} \twoheadrightarrow \hat{\mathbf{S}}_{\alpha,C}$$

(Proposition 5.5). In Subsection 5.3, we show that \mathcal{K}_{α} is a weak Bruhat interval when α is a composition obtained by shuffling a partition and (1^k) for some $k \geq 0$. To achieve this,

we first introduce the concept of southeast decreasing fillings (see Definition 5.6). Using this concept, we characterize the elements in \mathcal{K}_{α} . We then construct a specific filling τ'_{α} and show that $\mathcal{K}_{\alpha} = [\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\tau'_{\alpha})]_{L}$ (Theorem 5.8 and Theorem 5.12).

2. Preliminaries

For integers m and n, we define [m, n] and [n] to be the intervals $\{t \in \mathbb{Z} \mid m \leq t \leq n\}$ and $\{t \in \mathbb{Z} \mid 1 \leq t \leq n\}$, respectively. Throughout this paper, n will denote a nonnegative integer unless otherwise stated.

2.1. Compositions and their diagrams. A composition α of n, denoted by $\alpha \models n$, is a finite ordered list of positive integers $(\alpha_1, \alpha_2, \ldots, \alpha_k)$ satisfying $\sum_{i=1}^k \alpha_i = n$. We call α_i $(1 \le i \le k)$ a part of $\alpha, k =: \ell(\alpha)$ the length of α , and $n =: |\alpha|$ the size of α . And, we define the empty composition \emptyset to be the unique composition of size and length 0. Whenever necessary, we set $\alpha_i = 0$ for all $i > \ell(\alpha)$. We represent each composition visually using its composition diagram. The composition diagram $cd(\alpha)$ of α is the array of left-justified cells, with α_i cells in the *i*th row from the bottom for $1 \le i \le k$. If $\alpha = (3, 1, 2)$, then



Each cell is described by its row and column indices, with the coordinate (i, j) denoting the cell in the *i*th row from the bottom and the *j*th column from the left. We often use the set of coordinates of the cells in $cd(\alpha)$ to denote $cd(\alpha)$ itself. For example,

 $\mathsf{cd}((3,1,2)) = \{(1,1), (1,2), (1,3), (2,1), (3,1), (3,2)\}.$

For a composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \models n$ and a set $I = \{i_1 < i_2 < \dots < i_l\} \subseteq [n-1]$, let

set
$$(\alpha) := \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{k-1}\}, \text{ and}$$

comp $(I) := (i_1, i_2 - i_1, i_3 - i_2, \dots, n - i_l).$

The set of all compositions of n is in bijection with the set of all subsets of [n-1] under the correspondence $\alpha \mapsto \operatorname{set}(\alpha)$ (or $I \mapsto \operatorname{comp}(I)$). The reverse composition α^r of α is defined to be the composition $(\alpha_k, \alpha_{k-1}, \ldots, \alpha_1)$ and the complement α^c of α is defined to be the unique composition satisfying $\operatorname{set}(\alpha^c) = [n-1] \setminus \operatorname{set}(\alpha)$.

If a composition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \models n$ satisfies $\lambda_1 \ge \lambda_2 \ge \dots \ge \lambda_k$, then it is called a partition of n and denoted as $\lambda \vdash n$. Given two partitions λ and μ with $\ell(\lambda) \ge \ell(\mu)$, we write $\lambda \supseteq \mu$ if $\lambda_i \ge \mu_i$ for all $1 \le i \le \ell(\mu)$. A skew partition λ/μ is a pair (λ, μ) of partitions with $\lambda \supseteq \mu$. We call $|\lambda/\mu| := |\lambda| - |\mu|$ the size of λ/μ . When λ is a partition, we typically use $yd(\lambda)$ to denote $cd(\lambda)$ and refer to it as the Young diagram of λ (in French notation). For a skew partition λ/μ , the Young diagram $yd(\lambda/\mu)$ of λ/μ is defined as the Young diagram $yd(\lambda)$ with the cells corresponding to $yd(\mu)$ removed. We denote by $SYT(\lambda/\mu)$ the set of all standard Young tableaux of shape λ/μ . And, we let $\operatorname{SYT}_n := \bigsqcup_{\lambda \vdash n} \operatorname{SYT}(\lambda)$. Given a composition α , let $\lambda(\alpha)$ be the partition obtained by rearranging the entries of α in decreasing order.

Convention. Let D be a composition diagram or a Young diagram of skew shape. In this paper, we regard a filling T of D with positive integers as a map

$$T: D \to \mathbb{Z}_{>0}, \quad (i,j) \mapsto T(i,j),$$

where T(i, j) denotes the entry in the cell (i, j) of T. Given a filling T of D, we primarily utilize two reading words, $w_r(T)$ and $w_c(T)$, defined as follows:

- $w_r(T)$ is the word obtained by reading the entries of T from right to left starting with the bottom-most row.
- $w_c(T)$ is the word obtained by reading the entries of T from bottom to top starting with the rightmost column.

Unless otherwise stated, the *i*th row of T refers to the *i*th row from the bottom and the *j*th column of T refers to the *j*th column from the left.

2.2. The left weak Bruhat order on the symmetric group. Let \mathfrak{S}_n denote the symmetric group on [n]. Every permutation $\sigma \in \mathfrak{S}_n$ can be expressed as a product of simple transpositions $s_i := (i, i + 1)$ for $1 \leq i \leq n - 1$. A reduced expression for σ is an expression that represents σ in the shortest possible length and the length $\ell(\sigma)$ of σ is the number of simple transpositions in any reduced expression for σ . The left weak Bruhat order \preceq_L on \mathfrak{S}_n is the partial order on \mathfrak{S}_n whose covering relation \preceq_L^c is given as follows:

$$\sigma \preceq^{c}_{L} s_i \sigma$$
 if $\ell(\sigma) < \ell(s_i \sigma)$.

Let

$$\operatorname{Des}_{L}(\sigma) := \{ 1 \le i \le n-1 \mid \ell(s_{i}\sigma) < \ell(\sigma) \}.$$

It is well known that if $\sigma = w_1 w_2 \dots w_n$ in one-line notation, then

 $\operatorname{Des}_{L}(\sigma) = \{ 1 \le i \le n-1 \mid i \text{ is right of } i+1 \text{ in } w_1 w_2 \dots w_n \}.$

For each $\gamma \in \mathfrak{S}_n$, let

 $\operatorname{Inv}_{L}(\gamma) := \{(i, j) \mid 1 \le i < j \le n \text{ and } \gamma(i) > \gamma(j)\}.$

Then, for $\sigma, \rho \in \mathfrak{S}_n$,

 $\sigma \preceq_L \rho$ if and only if $\operatorname{Inv}_L(\sigma) \subseteq \operatorname{Inv}_L(\rho)$.

Given $\sigma, \rho \in \mathfrak{S}_n$, the *left weak Bruhat interval* $[\sigma, \rho]_L$ denotes the closed interval $\{\gamma \in \mathfrak{S}_n \mid \sigma \preceq_L \gamma \preceq_L \rho\}$ with respect to the left weak Bruhat order \preceq_L .

We collect notations associated with the symmetric group that will be used throughout the paper. For any subset $I \subseteq [n-1]$, let \mathfrak{S}_I denote the parabolic subgroup of \mathfrak{S}_n generated by $\{s_i \mid i \in I\}$ and let $w_0(I)$ denote the longest element in \mathfrak{S}_I . When I = [n-1], we simply write w_0 to denote the longest element in \mathfrak{S}_n . For $\alpha \models n$, we define

$$w_0(\alpha) := w_0(\operatorname{set}(\alpha)).$$

Additionally, for any $\sigma \in \mathfrak{S}_n$, we define $\sigma^{w_0} := w_0 \sigma w_0$. Finally, for $S \subseteq \mathfrak{S}_n$ and $\xi \in \mathfrak{S}_n$, we define

$$S \cdot \xi := \{ \gamma \xi \mid \gamma \in S \} \text{ and } \xi \cdot S := \{ \xi \gamma \mid \gamma \in S \}.$$

2.3. The 0-Hecke algebra and its representation theory. The 0-Hecke algebra $H_n(0)$ is the associative \mathbb{C} -algebra with 1 generated by $\pi_1, \pi_2, \ldots, \pi_{n-1}$ subject to the following relations:

$$\pi_i^2 = \pi_i \quad \text{for } 1 \le i \le n - 1, \\ \pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1} \quad \text{for } 1 \le i \le n - 2, \\ \pi_i \pi_j = \pi_j \pi_i \quad \text{if } |i - j| \ge 2.$$

For any reduced expression $s_{i_1}s_{i_2}\cdots s_{i_p}$ for $\sigma \in \mathfrak{S}_n$, let $\pi_{\sigma} := \pi_{i_1}\pi_{i_2}\cdots \pi_{i_p}$. It is well known that these elements are independent of the choices of reduced expressions and $\{\pi_{\sigma} \mid \sigma \in \mathfrak{S}_n\}$ is a \mathbb{C} -bases for $H_n(0)$.

Let us briefly review the representation theory of 0-Hecke algebras. Unless stated otherwise, all $H_n(0)$ -modules are finitely generated and carry a left action. By [28], there are 2^{n-1} pairwise nonisomorphic irreducible $H_n(0)$ -modules, naturally indexed by compositions of n. To be precise, for each $\alpha \models n$, there exists an irreducible $H_n(0)$ -module $\mathbf{F}_{\alpha} := \mathbb{C}v_{\alpha}$ endowed with the $H_n(0)$ -action defined as follows: for each $1 \le i \le n-1$,

$$\pi_i \cdot v_{\alpha} = \begin{cases} 0 & i \in \operatorname{set}(\alpha), \\ v_{\alpha} & i \notin \operatorname{set}(\alpha). \end{cases}$$

Let $H_n(0)$ -mod be the category of finitely generated left $H_n(0)$ -modules and $\mathcal{R}(H_n(0))$ the Z-span of the set of (representatives of) isomorphism classes of modules in $H_n(0)$ -mod. We denote by [M] the isomorphism class corresponding to an $H_n(0)$ -module M. The Grothendieck group $\mathcal{G}_0(H_n(0))$ of $H_n(0)$ -mod is the quotient of $\mathcal{R}(H_n(0))$ modulo the relations [M] = [M'] + [M''] whenever there exists a short exact sequence $0 \to M' \to$ $M \to M'' \to 0$. The equivalence classes of the irreducible $H_n(0)$ -modules form a Z-basis for $\mathcal{G}_0(H_n(0))$. Let

$$\mathcal{G} := \bigoplus_{n \ge 0} \mathcal{G}_0(H_n(0)).$$

Let us review the connection between \mathcal{G} and the ring QSym of quasisymmetric functions. For the definition of quasisymmetric functions, see [32, Section 7.19]. For a composition α , the fundamental quasisymmetric function F_{α} , which was firstly introduced in [17], is defined by

$$F_{\varnothing} = 1$$
 and $F_{\alpha} = \sum_{\substack{1 \le i_1 \le i_2 \le \dots \le i_n \\ i_j < i_{j+1} \text{ if } j \in \text{set}(\alpha)}} x_{i_1} x_{i_2} \cdots x_{i_n}$ if $\alpha \neq \emptyset$.

It is known that $\{F_{\alpha} \mid \alpha \text{ is a composition}\}$ is a \mathbb{Z} -basis for QSym. When M is an $H_m(0)$ module and N is an $H_n(0)$ -module, we write $M \boxtimes N$ for the induction product of M and N, that is,

$$M \boxtimes N := M \otimes N \uparrow_{H_m(0) \otimes H_n(0)}^{H_{m+n}(0)}$$

Here, $H_m(0) \otimes H_n(0)$ is viewed as the subalgebra of $H_{m+n}(0)$ generated by $\{\pi_i \mid i \in [m+n-1] \setminus \{m\}\}$. The induction product induces a multiplication on \mathcal{G} . It was shown in [15] that the linear map

$$ch: \mathcal{G} \to QSym, \quad [\mathbf{F}_{\alpha}] \mapsto F_{\alpha}$$

called quasisymmetric characteristic, is a ring isomorphism.

2.4. Bases for QSym and related 0-Hecke modules. In this paper, we deal with the following quasi-analogues of Schur functions: quasisymmetric Schur functions, Young quasisymmetric Schur functions, dual immaculate quasisymmetric functions, and extended Schur functions. We provide a brief introduction to each of these functions. Furthermore, for the latter two, we introduce indecomposable 0-Hecke modules that have these functions as images under the quasisymmetric characteristic.

2.4.1. Quasisymmetric Schur functions, Young quasisymmetric Schur functions, and related 0-Hecke modules. Quasisymmetric Schur functions were first introduced in [19, Definition 5.1] and arise from the specialization of nonsymmetric Macdonald polynomials to standard bases, commonly known as Demazure atoms. Moreover, their F-expansions, where F denotes the basis of fundamental quasisymmetric functions, are also provided. Here, we introduce this expansion.

Definition 2.1. Let α be a composition of n. A composition tableau of shape α is a filling T of $cd(\alpha)$ with entries in $\mathbb{Z}_{>0}$ satisfying the following conditions:

- (1) The entries in each row of T weakly decrease from left to right.
- (2) The entries in the leftmost column of T strictly increase from top to bottom.
- (3) Triple rule: for any cells $(i, k+1), (j, k) \in cd(\alpha)$ with i < j,

if
$$T(i, k+1) \le T(j, k)$$
, then $(j, k+1) \in cd(\alpha)$ and $T(i, k+1) < T(j, k+1)$.

Let $\alpha \models n$. A standard composition tableau (SCT) of shape α is a composition tableau in which each number in [n] appears exactly once. Let $SCT(\alpha)$ denote the set of all standard composition tableaux of shape α . For each $T \in SCT(\alpha)$, define

 $Des_{\mathscr{S}}(T) := \{1 \le i \le n-1 \mid i+1 \text{ appears weakly to the right of } i \text{ in } T\}.$

It was shown in [19, Section 5] that

$$\mathscr{S}_{\alpha} = \sum_{T \in \mathrm{SCT}(\alpha^{\mathrm{r}})} F_{\mathrm{comp}(\mathrm{Des}_{\mathscr{S}}(T))}$$

and $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ forms a basis for QSym_{n} .

In 2015, Tewari and van Willigenberg [34] introduced an $H_n(0)$ -action on the C-span of SCT($\alpha^{\rm r}$). The resulting $H_n(0)$ -module, denoted by \mathbf{S}_{α} , satisfies

$$\operatorname{ch}([\mathbf{S}_{\alpha}]) = \mathscr{S}_{\alpha}$$

This module is not indecomposable in general. For its decomposition into indecomposable modules, see [23, 10]. Define

 $SCT(\alpha^{r}; C) := \{T \in SCT(\alpha^{r}) \mid \text{ each column of } T \text{ increases from top to bottom} \}.$

The C-span of $SCT(\alpha^r; C)$ forms an indecompsable $H_n(0)$ -submodule of \mathbf{S}_{α} , which is denoted by $\mathbf{S}_{\alpha,C}$ and referred to as the *canonical submodule* of \mathbf{S}_{α} .

On the other hand, the Young quasisymmetric Schur function $\hat{\mathscr{S}}_{\alpha}$ is defined as $\rho(\mathscr{S}_{\alpha^r})$, where $\rho : \operatorname{QSym} \to \operatorname{QSym}$ is the automorphism defined by $F_{\alpha} \mapsto F_{\alpha^r}$. These functions exhibit many properties analogous to those of quasisymmetric Schur functions. Additionally, the Young quasisymmetric function $\hat{\mathscr{S}}_{\alpha}$ can be expressed using the combinatorial object known as Young composition tableaux.

Definition 2.2. Let α be a composition of n. A Young composition tableau of shape α is a filling T of $cd(\alpha)$ with entries in $\mathbb{Z}_{>0}$ satisfying the following conditions:

- (1) The entries in each row weakly increase from left to right,
- (2) The entries in the first column strictly increase from bottom to top, and
- (3) Young triple rule: for any cells $(i, k+1), (j, k) \in cd(\alpha)$ with i < j,

if $T(i, k+1) \ge T(j, k)$, then $(j, k+1) \in cd(\alpha)$ and T(i, k+1) > T(j, k+1).

A standard Young composition tableau of shape α is a Young composition tableau of shape α in which each of the numbers in [n] appears exactly once. We denote the set of all standard Young composition tableaux of shape α by SYCT (α) . For each $T \in SYCT(\alpha)$, we define

$$\operatorname{Des}_{\hat{\mathscr{S}}}(T) := \{ 1 \le i \le n-1 \mid i \text{ appears weakly right of } i+1 \text{ in } T \}.$$
(2.1)

It was shown in [24, Proposition 5.2.2] that

$$\hat{\mathscr{P}}_{\alpha} = \sum_{T \in \text{SYCT}(\alpha)} F_{\text{comp}(\text{Des}_{\hat{\mathscr{S}}}(T))}$$

(see [24, Section 4 and Section 5]).

In 2022, Choi, Kim, Nam, and Oh introduced an $H_n(0)$ -action on the \mathbb{C} -span of SYCT (α) . The resulting $H_n(0)$ -module, denoted by $\hat{\mathbf{S}}_{\alpha}$, satisfies that $ch([\hat{\mathbf{S}}_{\alpha}]) = \hat{\mathscr{I}}_{\alpha}$. This module is also not indecomposable in general. Define

 $SYCT(\alpha; C) := \{T \in SYCT(\alpha) \mid each column of T increases from bottom to top\}.$

The C-span of SYCT($\alpha; C$) forms an indecomposable submodule of $\hat{\mathbf{S}}_{\alpha}$, denoted by $\hat{\mathbf{S}}_{\alpha,C}$, which is referred to as the *canonical submodule* of $\hat{\mathbf{S}}_{\alpha}$ (see [12, Subsection 4.2]).

2.4.2. Dual immaculate quasisymmetric functions and related 0-Hecke modules. For each composition α , the dual immaculate quasisymmetric function \mathfrak{S}^*_{α} was initially introduced as the quasisymmetric dual of the corresponding immaculate noncommutative symmetric function in [3, Section 3.7]. Similar to \mathscr{S}_{α} and $\hat{\mathscr{S}}_{\alpha}$, it has a positive *F*-expansion, which can be expressed in terms of tableaux.

Definition 2.3. ([3, Definition 3.9]) Let α be a composition of n. A standard immaculate tableau of shape α is a filling \mathscr{T} of $cd(\alpha)$ with the entries $1, 2, \ldots, n$ such that the entries are all distinct, the entries in each row increase from left to right, and the entries in the first column increase from bottom to top.

Let $SIT(\alpha)$ denote the set of all standard immaculate tableaux of shape α . It was shown in [3, Proposition 3.48] that

$$\mathfrak{S}^*_{\alpha} = \sum_{T \in \operatorname{SIT}(\alpha)} F_{\operatorname{comp}(\operatorname{Des}_L(\mathsf{w}_{\mathsf{r}}(T)))^c}.$$

Then, Berg *et al.* [4] constructed an $H_n(0)$ -module structure for the dual immaculate quasisymmetric function \mathfrak{S}^*_{α} . Define an $H_n(0)$ -action on the \mathbb{C} -span of $SIT(\alpha)$ by

 $\pi_i \cdot \mathscr{T} = \begin{cases} \mathscr{T} & \text{if } i \text{ appears weakly above of } i+1 \text{ in } \mathscr{T}, \\ 0 & \text{if } i \text{ and } i+1 \text{ are in the first column of } \mathscr{T}, \\ s_i \cdot \mathscr{T} & \text{otherwise} \end{cases}$

for $1 \leq i \leq n-1$ and $\mathscr{T} \in \operatorname{SIT}(\alpha)$. Here, $s_i \cdot \mathscr{T}$ is obtained from \mathscr{T} by swapping *i* and i+1. The resulting module is denoted by \mathcal{V}_{α} .

Theorem 2.4. ([4, Theorem 3.5]) For a composition α of n, \mathcal{V}_{α} is an indecomposable $H_n(0)$ -module whose image under quasisymmetric characteristic is \mathfrak{S}^*_{α} .

2.4.3. Extended quasisymmetric Schur functions and related 0-Hecke modules. The extended Schur functions \mathcal{E}_{α} were introduced by Assaf and Searles in [2] as stable limits of lock polynomials. They form a basis for QSym, and their *F*-expansions can be expressed in terms of standard extended tableaux.

Definition 2.5. ([2, Definition 6.18]) Let α be a composition of n. A standard extended tableau of shape α is a filling T of $cd(\alpha)$ with the entries $1, 2, \ldots, n$ such that the entries are all distinct, the entries in each row increase from left to right, and the entries in each column increase from bottom to top.

We denote the set of all standard extended tableaux of shape α by SET(α). It was shown in [2, Theorem 6.19] that

$$\mathcal{E}_{\alpha} = \sum_{T \in \operatorname{SET}(\alpha)} F_{\operatorname{comp}(\operatorname{Des}_{L}(\mathsf{w}_{\mathsf{r}}(T)))^{c}}.$$

Subsequently, Searles [31] constructed an $H_n(0)$ -module for the extended Schur functions \mathcal{E}_{α} . Define an $H_n(0)$ -action on the C-span of SET(α) by

$$\pi_i \cdot T = \begin{cases} T & \text{if } i \text{ is strictly left of } i+1 \text{ in } T, \\ 0 & \text{if } i \text{ and } i+1 \text{ are in the same column of } T, \\ s_i \cdot T & \text{if } i \text{ is strictly right of } i+1 \text{ in } T \end{cases}$$

for $1 \leq i \leq n-1$ and $T \in SET(\alpha)$. Here, $s_i \cdot T$ is obtained from T by swapping i and i+1. The resulting module is denoted by X_{α} .

Theorem 2.6. ([31, Theorem 3.10, 3.13]) For $\alpha \models n$, X_{α} is an indecomposable $H_n(0)$ -module whose image under quasisymmetric characteristic is \mathcal{E}_{α} .

2.5. Weak Bruhat interval modules. Finally, we introduce 0-Hecke modules known as weak Bruhat interval modules, which were introduced by Jung, Kim, Lee, and Oh in [21] to provide a unified approach for studying 0-Hecke modules constructed using tableaux.

Definition 2.7. ([21, Definition 1]) For each left weak Bruhat interval $[\sigma, \rho]_L$ in \mathfrak{S}_n , define $\mathsf{B}([\sigma, \rho]_L)$ (simply, $\mathsf{B}(\sigma, \rho)$) to be the $H_n(0)$ -module with $\mathbb{C}[\sigma, \rho]_L$ as the underlying space and with the $H_n(0)$ -action defined by

$$\pi_i \cdot \gamma := \begin{cases} \gamma & \text{if } i \in \text{Des}_L(\gamma), \\ 0 & \text{if } i \notin \text{Des}_L(\gamma) \text{ and } s_i \gamma \notin [\sigma, \rho]_L, \\ s_i \gamma & \text{if } i \notin \text{Des}_L(\gamma) \text{ and } s_i \gamma \in [\sigma, \rho]_L. \end{cases}$$

The $H_n(0)$ -module $\mathsf{B}(\sigma, \rho)$ is called the *weak Bruhat interval module associated to* $[\sigma, \rho]_L$. Considering the above $H_n(0)$ -action on $\mathsf{B}(\sigma, \rho)$, one can derive that

$$\operatorname{ch}([\mathsf{B}(\sigma,\rho)]) = \sum_{\gamma \in [\sigma,\rho]_L} F_{\operatorname{comp}(\operatorname{Des}_L(\gamma))^c}.$$
(2.2)

The authors proved that \mathcal{V}_{α} and X_{α} are equipped with weak Bruhat interval module structures. Let us provide a brief explanation of this result.

We first define two special standard immaculate tableaux of shape α . Let \mathscr{T}_{α} be the filling of $\mathsf{cd}(\alpha)$ with entries $1, 2, \ldots, n$ from left to right starting with the bottommost row. On the other hand, let \mathscr{T}'_{α} be the filling of $\mathsf{cd}(\alpha)$ defined by the following steps:

- (1) Fill the first column with entries $1, 2, \ldots, \ell(\alpha)$ from bottom to top.
- (2) Fill the remaining cells with entries $\ell(\alpha) + 1, \ell(\alpha) + 2, ..., n$ from left to right starting with the topmost row.

In a similar manner, we define two standard extended tableaux of shape α . Let T_{α} be the standard extended tableau obtained by filling $\mathsf{cd}(\alpha)$ with the entries $1, 2, \ldots, n$ from left to right starting with the bottommost row and T'_{α} be the standard extended tableau obtained by filling $\mathsf{cd}(\alpha)$ with the entries $1, 2, \ldots, n$ from bottom to top starting with the leftmost column.

Theorem 2.8. ([21, Theorem 4.4]) Let α be a composition of n.

(1) The \mathbb{C} -linear isomorphism

$$\Theta_V: \mathcal{V}_{\alpha} \to \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})), \ \mathscr{T} \mapsto \mathsf{w}_{\mathsf{r}}(\mathscr{T}) \quad (\mathscr{T} \in \operatorname{SIT}(\alpha))$$

is an $H_n(0)$ -module isomorphism.

(2) The \mathbb{C} -linear isomorphism

$$\Theta_X : X_\alpha \to \mathsf{B}(\mathsf{w}_\mathsf{r}(\mathsf{T}_\alpha), \mathsf{w}_\mathsf{r}(\mathsf{T}'_\alpha)), \ \mathsf{T} \mapsto \mathsf{w}_\mathsf{r}(\mathsf{T}) \qquad (\mathsf{T} \in \operatorname{SET}(\alpha))$$

is an $H_n(0)$ -module isomorphism.

3. An analogue of the Robinson-Schensted-Knuth Algorithm and its Greene's Theorem

In this section, we introduce an analogue of the Robinson-Schensted-Knuth algorithm and examine how the shapes of the insertion tableaux are determined. In the case of the Robinson-Schensted-Knuth algorithm, this issue was resolved by Greene [18]. The result we establish here will be essential for proving the main theorems of Section 4.

3.1. An analogue of the Robinson-Schensted-Knuth Algorithm. Schensted's insertion is a procedure for inserting a positive integer into a semistandard Young tableau, which is the fundamental operation of the Robinson-Schensted-Knuth algorithm. Using this algorithm, for each $\sigma \in \mathfrak{S}_n$, we associate a pair of standard Young tableaux $(\mathsf{P}(\sigma), \mathsf{Q}(\sigma))$ with the same shape. It is well known that the map

$$\mathfrak{S}_n \to \bigsqcup_{\lambda \vdash n} \operatorname{SYT}(\lambda) \times \operatorname{SYT}(\lambda), \quad \sigma \mapsto (\mathsf{P}(\sigma), \mathsf{Q}(\sigma))$$

is bijective.

In 2006, Mason [26] introduced a procedure for inserting a nonnegative integer into a semi-skyline augmented filling. In 2011, Haglund, Luoto, Mason, and van Willigenburg [19, pages 480-481] rewrote this as a procedure for inserting a nonnegative integer into a composition tableau through the weight-preserving bijection, provided in [19, Lemma 4.2].

Convention. In this paper, a semi-skyline augmented filling is identified with the corresponding composition tableau through the weight-preserving bijection in [19, Lemma 4.2]. Therefore, we use composition tableaux instead of semi-skyline augmented fillings. With this identification, all the results from [26], originally described in terms of semi-skyline augmented fillings, are interpreted as results involving composition tableaux. In particular, Mason's insertion procedure is viewed as the procedure in [19, pages 480-481], and the algorithm in [26, Procedure 4.2] is understood as one that produces a pair of composition tableaux from a two-line array.

In 2018, Allen, Hallam, and Mason [1, Procedure 3.1] introduces a related procedure for inserting a nonnegative integer into a Young composition tableau. In this paper, we primarily focus on this insertion procedure.

Procedure 3.1. ([1, Procedure 3.1]) Let T be a Young composition tableau of shape $\alpha \models n$ and k a positive integer. Let $cd(\alpha) = \{(c_1, d_1), (c_2, d_2), \ldots, (c_n, d_n)\}$, where each pair (c_i, d_i) satisfies the following lexicographic order: For indices $1 \le i < j \le n$,

$$d_i > d_j$$
 or $d_i = d_j$ and $c_i > c_j$.

Set i := 1 and $k_0 := k$.

- **P1.** If $i \leq n$, then go to **P2**. Otherwise, go to **P3**.
- **P2.** We proceed by considering the following three cases. Case 1: $(c_i, d_i + 1) \notin cd(\alpha)$ and $T(c_i, d_i) \leq k_0$. Place k_0 in the cell $(c_i, d_i + 1)$ and then terminate the procedure. Case 2: $(c_i, d_i + 1) \in cd(\alpha)$ and $T(c_i, d_i) \leq k_0 < T(c_i, d_i + 1)$. Set $k_1 := T(c_i, d_i + 1)$, and update $T(c_i, d_i + 1)$ to k_0 and k_0 to k_1 . Then, update i to i + 1 and go to **P1**. Case 3: If neither of the above applies, update i to i + 1 and go to **P1**.
- **P3.** Create a new row with a single cell containing k_0 and place the row so that the first column strictly increases from bottom to top. Then, terminate the procedure.

We denote the Young composition tableau obtained from Algorithm 3.3 as $k \to T$. The set of cells that are modified during the insertion process, including the final cell added is referred to as the *insertion sequence*. The final cell added to the tableau is called the *new cell*.

Example 3.2. Let

$$T = \begin{bmatrix} 5 \\ 2 & 2 & 3 \\ 1 & 4 \end{bmatrix} \quad \text{and} \quad k = 2.$$

Then, $k \to T$ is given by

with the insertion sequence $\{(2,3), (1,2), (3,1)\}$ and the new cell (3,1).

Let $\widetilde{\mathcal{W}}_n$ be the set of all two-line arrays

$$w = \begin{pmatrix} i_1 & i_2 & \dots & i_n \\ j_1 & j_2 & \dots & j_n \end{pmatrix}$$

of positive integers such that for $1 \le r \le n-1$,

$$i_r < i_{r+1}$$
 or $i_r = i_{r+1}$ and $j_r \le j_{r+1}$.

Each word $w = w_1 w_2 \dots w_n$ consisting of positive integers, especially a permutation, is identified with the two-line array

$$\left(\begin{array}{ccc}1&2&\ldots&n\\w_1&w_2&\ldots&w_n\end{array}\right)\in\widetilde{\mathcal{W}}_n.$$

By modifying [26, Procedure 4.2], we present an algorithm that produces a pair of fillings of composition diagrams from a two-line array.

Algorithm 3.3. Let

$$w = \left(\begin{array}{ccc} i_1 & i_2 & \dots & i_n \\ j_1 & j_2 & \dots & j_n \end{array}\right) \in \widetilde{\mathcal{W}}_n$$

Let $\hat{P} = \emptyset$, $\hat{Q} = \emptyset$, and k = 1.

- **P1.** Say the new cell created during $j_k \to \hat{P}$ is located in the *c*th column of $j_k \to \hat{P}$. Update \hat{P} to $j_k \to \hat{P}$.
- **P2.** If c = 1, then create a new row with a single cell containing i_k and place the row so that the first column of \hat{Q} increases from bottom to top. Otherwise, denote the topmost row of \hat{Q} with c 1 cells by the *r*th row of \hat{Q} and update $\hat{Q}(r, c) = i_k$.
- **P3.** If k < n, update k to k + 1 and go back to the step **P1**. Otherwise, terminate the process.

We denote the pair (\hat{P}, \hat{Q}) of fillings obtained by applying Algorithm 3.3 to w as $(\hat{\mathsf{P}}(w), \hat{\mathsf{Q}}(w))$. It is evident that $\hat{\mathsf{P}}(w)$ is a Young composition tableau. Furthermore, from Lemma 3.6, it follows that $\hat{\mathsf{Q}}(w)$ is also a Young composition tableau.

Example 3.4. Let

$$w = \left(\begin{array}{rrrr} 1 & 2 & 3 & 3 & 4 \\ 3 & 1 & 2 & 5 & 4 \end{array}\right) \in \widetilde{\mathcal{W}}_5.$$

Then, we see that

$$\begin{pmatrix} 3 & 1 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 3 & 2 \\ 1 & 2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 3 & 2 & 3 \\ 1 & 2 & 1 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 3 & 2 & 3 & 3 \\ 1 & 2 & 5 & 1 \end{pmatrix}$$
$$\rightsquigarrow \begin{pmatrix} 3 & 5 & 2 & 3 & 3 \\ 1 & 2 & 4 & 1 \end{pmatrix} = (\hat{\mathsf{P}}(w), \hat{\mathsf{Q}}(w)).$$

We will derive important properties of Algorithm 3.3 from those of [26, Procedure 4.2]. As a first step, we examine the relationship between Procedure 3.1 and Mason's insertion procedure. For a finite subset $A = \{a_1 < a_2 < \cdots < a_k\} \subseteq \mathbb{Z}_{>0}$, consider the bijection

$$\mathbf{r}_A: A \to A, \quad a_i \mapsto a_{k+1-i}.$$

For any filling T of a composition diagram, we denote by ent(T) the set of all entries in T. With these notations, we have a bijection

 $\mathsf{rev}: \{\text{composition tableaux}\} \to \{\text{Young composition tableaux}\}, \quad T \mapsto \mathsf{r}_{\mathsf{ent}(T)} \circ T.$

For each composition α , this map induces a bijection

$$\operatorname{rev}|_{\operatorname{SCT}(\alpha)} : \operatorname{SCT}(\alpha) \to \operatorname{SYCT}(\alpha),$$

which satisfies, for any $T \in SCT(\alpha)$, the relation

$$\operatorname{Des}_{\mathscr{S}}(T) = \{ |\alpha| - i \mid i \in \operatorname{Des}_{\widehat{\mathscr{S}}}(\operatorname{rev}(T)) \}.$$

For clarity, we denote the composition tableau obtained by inserting k into the composition tableau T using [19, pages 480-481] as $k \stackrel{\text{CT}}{\to} T$. The following lemma is easily derived.

Lemma 3.5. (cf. [1, page 78]) Let T be a Young composition tableau and k a positive integer. Then, the following relations hold.

- (1) $k \to T = \operatorname{rev}(\mathsf{r}_{\operatorname{ent}(k \to T)}(k) \xrightarrow{\operatorname{CT}} \mathsf{r}_{\operatorname{ent}(k \to T)} \circ T).$ (2) The new cells created by inserting k into T using Procedure 3.1 and by inserting $\mathsf{r}_{\mathsf{ent}(k\to T)}(k)$ into $\mathsf{r}_{\mathsf{ent}(k\to T)} \circ T$ using the procedure described in [19, pages 480-481] are same.

Now, given a two-line array $w \in \widetilde{\mathcal{W}}_n$, we denote by (F(w), G(w)) the pair of composition tableaux obtained by applying [26, Procedure 4.2] to w. For $w = \begin{pmatrix} i_1 & i_2 & \dots & i_n \\ j_1 & j_2 & \dots & j_n \end{pmatrix} \in$ $\widetilde{\mathcal{W}}_n$, define

$$w[s:t] := \begin{pmatrix} i_s & i_{s+1} & \dots & i_t \\ j_s & j_{s+1} & \dots & j_t \end{pmatrix} \quad \text{for } 1 \le s \le t \le n,$$
$$w^{\text{conj}} := \begin{pmatrix} \mathsf{r}_X(i_n) & \mathsf{r}_X(i_{n-1}) & \dots & \mathsf{r}_X(i_1) \\ \mathsf{r}_Y(j_n) & \mathsf{r}_Y(j_{n-1}) & \dots & \mathsf{r}_Y(j_1) \end{pmatrix},$$

where $X = \{i_t \mid 1 \le t \le n\}$ and $Y = \{j_t \mid 1 \le t \le n\}$. The second notation is introduced because, if $w \in \mathfrak{S}_n$, then $w^{\text{conj}} = w^{w_0}$. It is straightforward to see that $w[s:t] \in \widetilde{\mathcal{W}}_{t-s+1}$, $w^{\operatorname{conj}} \in \widetilde{\mathcal{W}}_n$ and the map

$$\widetilde{\mathcal{W}}_n \to \widetilde{\mathcal{W}}_n, \quad w \mapsto w^{\operatorname{conj}}$$

is a bijection. Using these notations, we can derive the following lemma from Lemma 3.5.

Lemma 3.6. Let
$$w = \begin{pmatrix} i_1 & i_2 & \dots & i_n \\ j_1 & j_2 & \dots & j_n \end{pmatrix} \in \widetilde{\mathcal{W}}_n$$
. Then, we have
 $\hat{\mathsf{P}}(w) = \mathsf{rev}(F(w^{\text{conj}}))$ and $\hat{\mathsf{Q}}(w) = \mathsf{rev}(G(w^{\text{conj}}))$

Proof. We first prove that $\hat{\mathsf{P}}(w) = \mathsf{rev}(F(w^{\text{conj}}))$. We achieve our purpose by applying mathematical induction on n. Since it is trivial for n = 1, we may assume that n > 1. Let w' := w[1: n-1] and let

$$Y := \{j_t \mid 1 \le t \le n\}, \ Y' := \{j_t \mid 1 \le t \le n-1\}, \ \text{and} \ Y'_1 := \mathsf{r}_Y(Y').$$

Given any two finite subsets $A, B \subseteq \mathbb{Z}_{>0}$ with |A| = |B|, let $\iota_{A,B} : A \to B$ denote the order-preserving bijection such that for any $a_1, a_2 \in A$,

 $a_1 < a_2$ if and only if $\iota_{A,B}(a_1) < \iota_{A,B}(a_2)$.

Then one sees that $F(w'^{\text{conj}}) = \iota_{Y'_1,Y'} \circ F(w^{\text{conj}}[2:n])$. Combining this equality with the induction hypothesis, we derive that

$$\hat{\mathsf{P}}(w') = \mathsf{rev}(F(w'^{\text{conj}})) = \mathsf{r}_{Y'} \circ \iota_{Y'_1,Y'} \circ F(w^{\text{conj}}[2:n]) = \mathsf{r}_Y \circ F(w^{\text{conj}}[2:n]).$$
(3.1)

Here, the last equality follows from

$$\mathbf{r}_{Y'} \circ \iota_{Y'_1,Y'}(y_k) = \mathbf{r}_{Y'}(\mathbf{r}_Y(y_{m+1-k})) = \mathbf{r}_Y(y_k),$$

when we write $Y'_1 = \{y_1 < y_2 < \cdots < y_m\}$. Therefore, we have

$$\begin{split} \hat{\mathsf{P}}(w) &= j_n \to \hat{\mathsf{P}}(w') \\ &= \mathsf{rev}(\mathsf{r}_Y(j_n) \xrightarrow{\mathrm{CT}} \mathsf{r}_Y \circ \hat{\mathsf{P}}(w')) \quad \text{(by Lemma 3.5(1))} \\ &= \mathsf{rev}(\mathsf{r}_Y(j_n) \xrightarrow{\mathrm{CT}} F(w^{\operatorname{conj}}[2:n])) \quad \text{(by (3.1))} \\ &= \mathsf{rev}(F(w^{\operatorname{conj}})). \end{split}$$

Next, we prove that $\hat{Q}(w) = \text{rev}(G(w^{\text{conj}}))$ by applying mathematical induction on n. Since it is trivial for n = 1, we may assume that n > 1. Let

$$X := \{i_t \mid 1 \le t \le n\}, \ X' := \{i_t \mid 1 \le t \le n-1\}, \ \text{and} \ X'_1 := \mathsf{r}_X(X').$$

One sees that $G(w'^{\text{conj}}) = \iota_{X'_1,X'} \circ G(w^{\text{conj}}[2 : n])$. Combining this equality with the induction hypothesis, we derive that

$$\hat{\mathsf{Q}}(w') = \mathsf{rev}(G(w'^{\text{ conj}})) = \iota_{X'_1,X'} \circ G(w^{\text{conj}}[2:n]).$$
(3.2)

This implies that $\hat{\mathsf{Q}}(w')$ and $G(w^{\text{conj}}[2:n])$ have the same shape, denoted by α . And, by Lemma 3.5(2), the new cells created during $j_n \to \hat{\mathsf{P}}(w')$ and $\mathsf{r}_Y(j_n) \to F(w^{\text{conj}}[2:n])$ are same. Denote this cell by (r, c). Therefore, it follows that

- $\hat{Q}(w)$ is obtained from $\hat{Q}(w')$ by filling (k, c) with i_n , and
- $G(w^{\text{conj}})$ is obtained from $G(w^{\text{conj}}[2:n])$ by filling (k,c) with $\mathsf{r}_X(i_n)$,

where $k = \max\{1 \le t \le \ell(\alpha) + 1 \mid \alpha_t = c - 1\}$. Now, the desired result can be derived by combining these properties with (3.2).

Consider the correspondence

$$\mathfrak{S}_{n} \to \bigsqcup_{\substack{\alpha,\beta \models n \\ \lambda(\alpha) = \lambda(\beta)}} \operatorname{SYCT}(\alpha) \times \operatorname{SYCT}(\beta), \quad w \mapsto (\hat{\mathsf{P}}(w), \hat{\mathsf{Q}}(w)).$$
(3.3)

By combining the properties in [26, Section 4] with Lemma 3.6, we derive the following properties:

- The correspondence given in (3.3) is a bijection.
- For $\sigma \in \mathfrak{S}_n$,

$$\hat{\mathsf{P}}(\sigma^{-1}) = \hat{\mathsf{Q}}(\sigma) \text{ and } \hat{\mathsf{Q}}(\sigma^{-1}) = \hat{\mathsf{P}}(\sigma).$$
 (3.4)

• For $\sigma, \rho \in \mathfrak{S}_n$,

$$\mathsf{P}(\sigma) = \mathsf{P}(\rho)$$
 if and only if $\hat{\mathsf{P}}(\sigma) = \hat{\mathsf{P}}(\rho)$. (3.5)

Another important property is that the mapping $\sigma \mapsto \hat{\mathsf{P}}(\sigma)$ preserves descents, as shown in the following proposition:

Proposition 3.7. For $\sigma \in \mathfrak{S}_n$, we have

$$\operatorname{Des}_{L}(\sigma) = \operatorname{Des}_{\hat{\mathscr{S}}}(\hat{\mathsf{P}}(\sigma)).$$

For the definition of $\text{Des}_{\hat{\mathscr{S}}}(\hat{\mathsf{P}}(\sigma))$, see (2.1).

Proof. We first prove our assertion in the case where σ is equal to $\mathbf{w}_{\mathbf{r}}(T)w_0$ for some $T \in \text{SYT}_n$. Then, it holds that $\mathsf{P}(\mathsf{w}_{\mathbf{r}}(T)w_0) = T$. Moreover, since $\mathsf{w}_{\mathbf{r}}(T)w_0$ is the immaculate reading word of T defined in [1, Definition 2.7], it follows from [1, Lemma 3.11, Lemma 3.12] that $\text{Des}_L(\mathsf{w}_{\mathbf{r}}(T)w_0) = \text{Des}_{\hat{\mathscr{S}}}(\hat{\mathsf{P}}(\mathsf{w}_{\mathbf{r}}(T)w_0))$.

Now, consider the case where σ is an arbitrary permutation in \mathfrak{S}_n . Let $T = \mathsf{P}(\sigma)$. Since $\mathsf{P}(\mathsf{w}_{\mathsf{r}}(T)w_0) = \mathsf{P}(\sigma)$, it follows that $\mathrm{Des}_L(\sigma) = \mathrm{Des}_L(\mathsf{w}_{\mathsf{r}}(T)w_0)$ and $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(T)w_0) = \hat{\mathsf{P}}(\sigma)$ by (3.5). Therefore, the desired result is obtained from the above discussion.

However, unlike the Robinson-Schensted-Knuth algorithm, the shapes of $\hat{\mathsf{P}}(\sigma)$ and $\hat{\mathsf{Q}}(\sigma)$ are not necessarily the same. In fact, it holds that

$$\lambda(\operatorname{sh}(\mathsf{P}(\sigma))) = \lambda(\operatorname{sh}(\mathsf{Q}(\sigma))) = \operatorname{sh}(\mathsf{P}(\sigma))$$

(for further details, see [26, Section 3.3, Corollary 4.4, Corollary 4.5]).

3.2. The corresponding Greene's theorem. We begin by introducing Greene's theorem, which provides a characterization of how the shapes of insertion tableaux in the Robinson-Schensted-Knuth algorithm are determined.

Let $\sigma \in \mathfrak{S}_n$ be fixed throughout this subsection. In 1961, Schensted [30] discovered that the size of the first row of the insertion tableau $\mathsf{P}(\sigma)$ is equal to the length of a longest increasing subsequence of σ . In 1974, Greene [18] extended this result by fully characterizing the shape of $\mathsf{P}(\sigma)$ in terms of *k*-increasing subsequences.

For each positive integer k, we introduce the notion of k-increasing subsequences. Consider a sequence $w = w_1 w_2 \dots w_n$ of positive integers. By abuse of notation, we use $m \in w$ to denote that m is an entry of w and $m \notin w$ otherwise. Let $\ell(w) := n$ and refer to it as the *length* of w. By convention, we denote \emptyset as the unique sequence of length 0. We say that w is *increasing* if $\ell(w) = 0$, or $\ell(w) > 1$ and $w_1 < w_2 < \dots < w_n$. For $1 \leq k \leq n$, a subsequence w' of w is called k-increasing (with a hyphen) if w' contains no decreasing subsequence of length k+1, Note that w' is k-increasing if and only if it can be expressed

as a disjoint union of k increasing subsequences of w. Let $i_k(w)$ denote the length of a longest k-increasing subsequence of w.

Example 3.8. Let w = 943156827. The subsequence 4156827 of w is 2-increasing since it is a union of 2 increasing subsequences 4568 and 127 of w. Also, one can easily check that it is a longest 2-increasing subsequence of w and thus $i_2(w) = 7$.

From now on, we write each permutation in one-line notation, viewing it as a sequence of positive integers. The following result is due to Greene.

Theorem 3.9. ([18]) Let $\sigma \in \mathfrak{S}_n$ and $\lambda = \operatorname{sh}(\mathsf{P}(\sigma))$. For $1 \leq k \leq n$, we have

$$\mathfrak{i}_k(\sigma) = \lambda_1 + \lambda_2 + \dots + \lambda_k.$$

The purpose of this subsection is to present an analogue of Theorem 3.9 for $\operatorname{sh}(\hat{\mathsf{P}}(\sigma))$. For a sequence $w = w_1 w_2 \dots w_n$ of distinct positive integers and $1 \leq k \leq n$, let $\operatorname{Inc}_k(w)$ be the set of all k-tuples $u = (u^{(i)})_{1 \leq i \leq k}$ of increasing subsequences of w such that

- if $i \neq j$, then $u^{(i)}$ and $u^{(j)}$ have no common entries and
- $\sum_{1 \le i \le k} \ell(u^{(i)}) = \mathfrak{i}_k(w).$

For $u = (u^{(i)})_{1 \le i \le k} \in \operatorname{Inc}_k(w)$, we denote by the boldfaced **u** the subsequence of w that consists of all entries in the subsequence $u^{(i)}$ for all *i*'s. For example, if $w = 943156827 \in \mathfrak{S}_9$ and $u = (4568, 127) \in \operatorname{Inc}_2(w)$, then $\mathbf{u} = 4156827$. One sees that **u** is a longest *k*-increasing subsequence of w and $\{\mathbf{u} \mid u \in \operatorname{Inc}_k(w)\}$ is the set of all longest *k*-increasing subsequences of w.

We introduce a total order \leq_{set} on the set of all finite subsets of $\mathbb{Z}_{>0}$ by defining $A \leq_{set} B$ if one of the following conditions holds:

• |A| < |B|,

•
$$A = B$$
,

• |A| = |B| and there exists $1 \le i \le |A|$ such that $a_j = b_j$ for all $1 \le j < i$ and $a_i < b_i$, where $A = \{a_1 < a_2 < \cdots < a_{|A|}\}$ and $B = \{b_1 < b_2 < \cdots < b_{|B|}\}$.

Definition 3.10. Let $\sigma \in \mathfrak{S}_n$ and $1 \leq k \leq n$.

(1) For $u = (u^{(i)})_{1 \le i \le k} \in \operatorname{Inc}_k(\sigma)$, we define the *initial entries set of* u by

$$\mathsf{IES}(u) := \{ u_1^{(i)} \mid 1 \le i \le k \text{ with } \ell(u^{(i)}) > 0 \}.$$

(2) With respect to the order \leq_{set} , we define

$$\underline{\mathsf{IES}}_k(\sigma) := \max\{\mathsf{IES}(u) \mid u \in \mathrm{Inc}_k(\sigma)\}$$

By convention, we set $\underline{\mathsf{IES}}_0(\sigma) := \emptyset$. From the definition, it follows that $|\underline{\mathsf{IES}}_k(\sigma)| = k$.

Example 3.11. Let $\sigma = 637$. Then $\text{Inc}_1(\sigma) = \{(67), (37)\}$, $\text{Inc}_2(\sigma) = \{(67, 3), (37, 6)\}$, and $\text{Inc}_3(\sigma) = \{(67, 3, \emptyset), (37, 6, \emptyset), (7, 6, 3)\}$. For these, we have

$$\begin{aligned} \mathsf{IES}((67)) &= \{6\}, \quad \mathsf{IES}((37)) &= \{3\}, \\ \mathsf{IES}(((67,3)) &= \{3,6\}, \quad \mathsf{IES}((37,6)) &= \{3,6\}, \text{ and} \\ \mathsf{IES}((67,3,\emptyset)) &= \{3,6\}, \quad \mathsf{IES}((37,6,\emptyset)) &= \{3,6\}, \quad \mathsf{IES}((3,6,7)) &= \{3,6,7\}. \end{aligned}$$

It follows that $\underline{\mathsf{IES}}_1(\sigma) = \{6\}$, $\underline{\mathsf{IES}}_2(\sigma) = \{3, 6\}$, and $\underline{\mathsf{IES}}_3(\sigma) = \{3, 6, 7\}$.

We begin by proving that $\underline{\mathsf{IES}}_k(\cdot)$ is constant on the Knuth equivalence classes. Let us briefly recall the Knuth equivalence relation. For $\sigma \in \mathfrak{S}_n$ and 1 < i < n, we write $\sigma \stackrel{1}{\cong} \sigma s_i$ if

$$\sigma(i+1) < \sigma(i-1) < \sigma(i) \quad \text{or} \quad \sigma(i) < \sigma(i-1) < \sigma(i+1).$$

We write $\sigma \stackrel{2}{\cong} \sigma s_{i-1}$ if

$$\sigma(i) < \sigma(i+1) < \sigma(i-1) \quad \text{or} \quad \sigma(i-1) < \sigma(i+1) < \sigma(i).$$

The Knuth equivalence is an equivalence relation $\stackrel{K}{\cong}$ on \mathfrak{S}_n defined by $\sigma \stackrel{K}{\cong} \rho$ if and only if there exist $\gamma_1, \gamma_2, \ldots, \gamma_k \in \mathfrak{S}_n$ such that

$$\sigma = \gamma_1 \stackrel{a_1}{\cong} \gamma_2 \stackrel{a_2}{\cong} \cdots \stackrel{a_{k-1}}{\cong} \gamma_k = \rho,$$

where $a_1, a_2, \ldots, a_{k-1} \in \{1, 2\}$. Note that $\sigma \stackrel{K}{\cong} \rho$ if and only if $\mathsf{P}(\sigma) = \mathsf{P}(\rho)$. For more information, see [9, 16, 29, 32].

Lemma 3.12. Let $\sigma, \rho \in \mathfrak{S}_n$. If σ and ρ are Knuth equivalent, then $\underline{\mathsf{IES}}_k(\sigma) = \underline{\mathsf{IES}}_k(\rho)$ for all $1 \leq k \leq n$.

Proof. Assume that σ and ρ are Knuth equivalent and let $1 \leq k \leq n$. To prove the assertion, it suffices to consider the situation where $\sigma \stackrel{a}{\cong} \rho$ with $a \in \{1, 2\}$ and $\ell(\sigma) > \ell(\rho)$. The following cases are possible:

Case 1.
$$\sigma \cong \rho$$
, that is, $\sigma = \dots yzx \dots$ and $\rho = \dots yxz \dots$, where $x < y < z$.
Case 2. $\sigma \cong \rho$, that is, $\sigma = \dots zxy \dots$ and $\rho = \dots xzy \dots$, where $x < y < z$.

We deal with only Case 1, as Case 2 can be handled in a similar manner.

Since $\operatorname{Inc}_k(\sigma) \subseteq \operatorname{Inc}_k(\rho)$, it immediately follows that $\underline{\operatorname{\mathsf{IES}}}_k(\sigma) \leq_{\operatorname{set}} \underline{\operatorname{\mathsf{IES}}}_k(\rho)$. Therefore, it remains to show that $\underline{\operatorname{\mathsf{IES}}}_k(\sigma) \geq_{\operatorname{set}} \underline{\operatorname{\mathsf{IES}}}_k(\rho)$. Choose $u = (u^{(i)})_{1 \leq i \leq k} \in \operatorname{Inc}_k(\rho)$ such that $\operatorname{\mathsf{IES}}(u) = \underline{\operatorname{\mathsf{IES}}}_k(\rho)$. If there is no index $1 \leq t \leq k$ with $x, z \in u^{(t)}$, then $u \in \operatorname{Inc}_k(\sigma)$, and thus $\underline{\operatorname{\mathsf{IES}}}_k(\sigma) \geq_{\operatorname{set}} \underline{\operatorname{\mathsf{IES}}}_k(\rho)$. Otherwise, let $1 \leq t \leq k$ be the index such that $x, z \in u^{(t)}$. Write $u^{(t)} = A xz B$. There are two subcases.

• Suppose $y \in u^{(s)}$ for some $1 \le s \le k$. In this case, $s \ne t$. Write $u^{(s)} = C y D$ and define $v = (v^{(i)})_{1 \le i \le k}$ as follows:

$$v^{(i)} := \begin{cases} A \, x \, D & \text{if } i = t, \\ C \, yz \, B & \text{if } i = s, \\ u^{(i)} & \text{otherwise} \end{cases}$$

By Theorem 3.9, one sees that $v \in \text{Inc}_k(\sigma)$; in addition, $\mathsf{IES}(v) = \mathsf{IES}(u) = \underline{\mathsf{IES}}_k(\rho)$. Therefore, $\underline{\mathsf{IES}}_k(\sigma) \ge_{\text{set}} \underline{\mathsf{IES}}_k(\rho)$, as required.

• Suppose $y \notin u^{(s)}$ for any $1 \leq s \leq k$. Let $v = (v^{(i)})_{1 \leq i \leq k}$ be defined as:

$$v^{(i)} := \begin{cases} A \, yz \, B & \text{if } i = t, \\ u^{(i)} & \text{otherwise.} \end{cases}$$

By Theorem 3.9, one sees that $v \in \text{Inc}_k(\sigma)$; in addition, $\mathsf{IES}(v) \ge_{\text{set}} \mathsf{IES}(u) = \underline{\mathsf{IES}}(\rho)$. Therefore, we obtain that $\underline{\mathsf{IES}}_k(\sigma) \ge_{\text{set}} \underline{\mathsf{IES}}_k(\rho)$.

Lemma 3.13. For $\sigma \in \mathfrak{S}_n$, $\underline{\mathsf{IES}}_l(\sigma)$ is equal to the set of all entries in the first column of $\hat{\mathsf{P}}(\sigma)$, where l denotes the length of $\operatorname{sh}(\hat{\mathsf{P}}(\sigma))$.

Proof. First, consider the case where $\sigma = w_r(T)w_0$ for some $T \in \text{SYT}(\lambda)$, with λ being a partition of n. Let $l = \ell(\lambda)$. In this setting, $\mathsf{P}(\sigma) = T$ and the set of all entries in the first column of $\hat{\mathsf{P}}(\sigma)$ is precisely the set of entries in the first column of T. Thus, to prove the assertion, it suffices to verify that $\underline{\mathsf{IES}}_l(\sigma) = \{T(1,1), T(2,1), \ldots, T(l,1)\}$. This follows from the observation that

$$T(i, 1) \in \mathsf{IES}(u)$$
 for all $u \in \mathrm{Inc}_l(\sigma)$ and $1 \le i \le l$.

Otherwise, there exist $u \in \text{Inc}_l(\sigma)$ and $1 \leq i \leq l$ such that $T(i, 1) \notin \text{IES}(u)$. Since T(j,k) > T(i,1) for all $i < j \leq l$ and $1 \leq k \leq \lambda_j$, it follows that $T(i,1) \notin \mathbf{u}$. Consequently, $\mathbf{u} \neq \sigma$, which contradicts the assumption that $u \in \text{Inc}_l(\sigma)$.

Let σ be an arbitrary permutation in \mathfrak{S}_n and let $T := \mathsf{P}(\sigma)$. Then,

$$\dot{\mathsf{P}}(\sigma) = \dot{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(T)w_0) \text{ and } \underline{\mathsf{IES}}_l(\sigma) = \underline{\mathsf{IES}}_l(\mathsf{w}_{\mathsf{r}}(T)w_0),$$

where the first equality follows from (3.5), while the second is a consequence of Lemma 3.12. Now, the desired assertion can be obtained by combining these equalities with the preceding discussion.

For a partition λ , we denote the skew partition whose Young diagram is obtained by rotating $yd(\lambda)$ by 180° by λ °.

Lemma 3.14. Let $\lambda = (\lambda_1, \lambda_2, ..., \lambda_l)$ be a partition of n and let $T \in SYT(\lambda^\circ)$. For $1 \le k \le l$ with $\lambda_k = \lambda_1$, the following hold:

- (1) For u = (u⁽ⁱ⁾)_{1≤i≤k} ∈ Inc_k(w_c(T)), IES(u) consists of the entries in the first row of T.
 (2) IES_k(w_c(T)) = {T(1, j) | l − k + 1 ≤ j ≤ l}.
- *Proof.* It is well known in the literature that the shape of $\mathsf{P}(\mathsf{w}_{\mathsf{c}}(T))$ is λ (for instance, see

[16]). (1) By Theorem 3.9, an increasing subsequence of $w_c(T)$ has length at most λ_1 and the

(1) By Theorem 5.9, an increasing subsequence of $w_c(T)$ has length at most λ_1 and the length of a longest k-increasing subsequence of $w_c(T)$ is $k\lambda_1$. Combining these properties, we deduce that the length of $u^{(i)}$ is equal to λ_1 for every $1 \le i \le k$. On the other hand, if an increasing subsequence of $w_c(T)$ has length λ_1 , then it starts with an entry in the first row of T. Therefore, we obtain the desired result.

(2) Let X denote the set of all entries in the first row of T. By (1), we have

$$\{\mathsf{IES}(u) \mid u \in \mathsf{Inc}(\mathsf{w}_{\mathsf{c}}(T))\} \subseteq \{S \subseteq X \mid |S| = k\}.$$

Combining this inclusion with

$$\{T(1,j) \mid l-k+1 \le j \le l\} = \max\{S \subseteq X \mid |S| = k\} \text{ (with respect to } \ge_{\text{set}}),\$$

it suffices to show that there exists $u \in \text{Inc}_k(\mathsf{w}_{\mathsf{c}}(T))$ such that

$$\mathsf{IES}(u) = \{ T(1,j) \mid l - k + 1 \le j \le l \}.$$

The existence of u can be shown by considering a k-tuple $v = (v^{(i)})_{1 \le i \le k}$ of subsequences of $w_c(T)$, where $v^{(i)}$ is the increasing subsequence of $w_c(T)$ consisting of all entries in the (l-k+i)th column of T.

We are now ready to present the main result of this section.

Theorem 3.15. Let $\sigma \in \mathfrak{S}_n$ and let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l) = \operatorname{sh}(\mathsf{P}(\sigma))$.

- (1) $\emptyset = \underline{\mathsf{IES}}_0(\sigma) \subsetneq \underline{\mathsf{IES}}_1(\sigma) \subsetneq \cdots \subsetneq \underline{\mathsf{IES}}_l(\sigma).$
- (2) Let $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = (\alpha_1, \alpha_2, \dots, \alpha_l)$ and let $\underline{\mathsf{IES}}_l(\sigma) = \{x_1 < x_2 < \dots < x_l\}$. Then, for each $1 \leq k \leq l$, $\alpha_k = \lambda_{i_k}$, where i_k is the smallest index $1 \leq t \leq l$ such that $x_k \in \underline{\mathsf{IES}}_t(\sigma)$.

Proof. Before proving the assertions, we recall that there exists $T \in \text{SYT}(\lambda^{\circ})$ such that $\mathsf{P}(\mathsf{w}_{\mathsf{c}}(T)) = \mathsf{P}(\sigma)$. For $1 \leq k \leq l$, let $T^{(k)}$ denote the subfilling of T consisting of the first k columns from the right. Clearly, $T^{(k)}$ is a Young tableau of shape $(\lambda_1, \lambda_2, \ldots, \lambda_k)^{\circ}$.

(1) By Lemma 3.12, it suffices to prove the assertion for $\sigma = w_c(T)$. To begin with, we will show that for every $1 \le k \le l$,

$$\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T)) = \underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T^{(k)})). \tag{3.6}$$

Assume first that $\lambda_k = \lambda_1$. Now, applying Lemma 3.14(2) to both $w_c(T)$ and $w_c(T')$ shows that

 $\underline{\mathsf{IES}}_k(\mathsf{w_c}(T)) = \{T(1,j) \mid l-k+1 \le j \le l\} = \underline{\mathsf{IES}}_k(\mathsf{w_c}(T^{(k)})).$

From now on, assume that $\lambda_k < \lambda_1$. Since $\operatorname{Inc}_k(\mathsf{w}_{\mathsf{c}}(T)) \supseteq \operatorname{Inc}_k(\mathsf{w}_{\mathsf{c}}(T^{(k)}))$, it follows that

$$\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T)) \geq_{\text{set}} \underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T^{(k)})).$$

Therefore, it suffices to verify that $\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T)) \leq_{\text{set}} \underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T^{(k)}))$. This proceeds in three steps as follows:

Step 1. Let $u = (u^{(i)})_{1 \le i \le k} \in \operatorname{Inc}_k(\mathsf{w}_{\mathsf{c}}(T))$ be given. We show that each $u^{(i)}$ contains at least one element from the (m+1)st row in T, where $m = \lambda_1 - \lambda_k$. Let X be the set of entries in the first m rows of T, that is,

$$X = \{T(i,j) \mid (i,j) \in \operatorname{yd}(\lambda^\circ) \text{ with } 1 \le i \le m\}.$$

Let T' be the subfilling of T obtained by removing all elements in X. Consider the k-tuple $\hat{u} = (\hat{u}^{(i)})_{1 \leq i \leq k}$, where $\hat{u}^{(i)}$ is the sequence obtained from $u^{(i)}$ by removing all elements in X. By its definition, $\hat{\mathbf{u}}$ is a k-increasing subsequence of $w_{\mathsf{c}}(T')$ and

$$\ell(\hat{\mathbf{u}}) \ge \ell(\mathbf{u}) - |X| = k\lambda_k = \mathfrak{i}_k(\mathsf{w}_\mathsf{c}(T')).$$

Combining this inequality with Theorem 3.9 yields that $\ell(\hat{\mathbf{u}}) = i_k(\mathsf{w}_{\mathsf{c}}(T'))$, that is, $\hat{u} \in \operatorname{Inc}_k(\mathsf{w}_{\mathsf{c}}(T'))$. Therefore, by Lemma 3.14(1),

$$\mathsf{IES}(\hat{u}) \subseteq \{ T(m+1,j) \mid (m+1,j) \in \mathsf{yd}(\lambda^\circ) \}.$$

As a consequence, we have

$$u^{(i)} = A^{(i)} T(m+1, c_i) B^{(i)} \quad (1 \le i \le k),$$

where $T(m+1, c_i)$ is the initial entry of $\hat{u}^{(i)}$. By suitably rearranging the indices, we may assume that $c_1 < c_2 < \cdots < c_k$.

Step 2. We show that

$$X = \{ x \in A^{(i)} \mid 1 \le i \le k \}.$$
(3.7)

To prove this equality, it suffices to show that for each $1 \leq i \leq k$, $A^{(i)}$ consists of elements that belong to X, while $B^{(i)}$ contains no elements from X. If the cell $C \in \mathrm{yd}(\lambda^{\circ})$ is strictly right and weakly above of $(m+1, c_i)$, then $T(C) > T(m+1, c_i)$, and thus $T(C) \notin A^{(i)}$. This shows that $A^{(i)}$ consists only of elements in X. On the other hand, if the cell $C \in \mathrm{yd}(\lambda^{\circ})$ is strictly left and strictly below of $(m+1, c_i)$, then $T(C) < T(m+1, c_i)$, and thus $T(C) \notin B^{(i)}$. This shows that $B^{(i)}$ does not contain any element in X.

Step 3. Define a k-tuple $v = (v^{(i)})_{1 \le i \le k}$ of sequences by

$$v^{(i)} := A^{(i)} T(m+1, l-k+i) T(m+2, l-k+i) \dots T(\lambda_1, l-k+i).$$

Since $c_i \leq l - k + i$, we have

$$T(m+1, l-k+i) \ge T(m+1, c_i),$$

which implies that $v^{(i)}$ is an increasing sequence. And, **v** is equal to $w_c(T^{(k)})$ by (3.7). Consequently, $v \in \text{Inc}_k(w_c(T^{(k)}))$. Moreover, it is easy to see that $\mathsf{IES}(u) \leq_{\text{set}} \mathsf{IES}(v)$. Since $u \in \text{Inc}_k(w_c(T))$ was chosen arbitrarily, we conclude that

$$\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T)) \leq_{\text{set}} \underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T^{(k)})).$$

Now we are ready to prove the assertion. For a given positive integer $2 \leq k \leq l$, every entry in the first column of $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{c}}(T^{(k-1)}))$ is also in the first column of $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{c}}(T^{(k)}))$. Combining this with Lemma 3.13, we have

$$\underline{\mathsf{IES}}_{k-1}(\mathsf{w}_{\mathsf{c}}(T^{(k-1)})) \subseteq \underline{\mathsf{IES}}_{k}(\mathsf{w}_{\mathsf{c}}(T^{(k)})).$$

Finally, by (3.6), we conclude that

$$\underline{\mathsf{IES}}_{k-1}(\mathsf{w}_{\mathsf{c}}(T)) \subseteq \underline{\mathsf{IES}}_{k}(\mathsf{w}_{\mathsf{c}}(T)).$$

(2) By (1), the set of entries in the first column of $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{c}}(T))$ is given by $\{y_1, y_2, \ldots, y_l\}$, where y_k denotes a unique element in $\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{c}}(T)) \setminus \underline{\mathsf{IES}}_{k-1}(\mathsf{w}_{\mathsf{c}}(T))$. For the assertion, it suffices to show that the number of cells in the row starting with y_k is λ_k for $1 \leq k \leq l$. To do this, let $\hat{P}_k := \hat{\mathsf{P}}(\mathsf{w}_{\mathsf{c}}(T^{(k)}))$ for all $1 \leq k \leq l$. The shape of \hat{P}_1 is (λ_1) , and thus, the number of cells starting with y_1 in \hat{P}_1 is equal to λ_1 . Let us fix $2 \leq k \leq l$. By (3.6), the set of entries in the first column of \hat{P}_k is $\{y_1, y_2, \ldots, y_k\}$ and \hat{P}_{k-1} is $\{y_1, y_2, \ldots, y_{k-1}\}$. One can easily see that for all $1 \leq i \leq k - 1$, the number of cells in the row starting with y_i in \hat{P}_k is greater than or equal to that of cells in the row starting with y_i in \hat{P}_{k-1} . Combining this with

$$\lambda(\operatorname{sh}(\hat{P}_{k-1})) = (\lambda_1, \dots, \lambda_{k-1}) \text{ and } \lambda(\operatorname{sh}(\hat{P}_k)) = (\lambda_1, \lambda_2, \dots, \lambda_k),$$

we deduce that

- for $1 \le i < k$, the number of cells in the row starting with y_i in \hat{P}_k is equal to that of cells in the row starting with y_i in \hat{P}_{k-1} , and
- the number of cells in the row starting with y_i in \hat{P}_k is λ_k .

Consequently, for $1 \leq k \leq l$, the number of cells in the row starting with y_k in $\hat{P}_l = \hat{\mathsf{P}}(\mathsf{w}_{\mathsf{c}}(T))$ is equal to that of cells in the row starting with y_k in \hat{P}_k .

Example 3.16. Let $\sigma = 52783146 \in \mathfrak{S}_8$. Then,

$$\mathsf{P}(\sigma) = \begin{array}{c|c} 5 \\ \hline 2 & 7 & 8 \\ \hline 1 & 3 & 4 & 6 \end{array}$$

This shows that $\operatorname{sh}(\mathsf{P}(\sigma)) = (4, 3, 1)$. Furthermore, $\underline{\mathsf{IES}}_1(\sigma) = \{2\}$, $\underline{\mathsf{IES}}_2(\sigma) = \{2, 5\}$, and $\underline{\mathsf{IES}}_3(\sigma) = \{1, 2, 5\}$. Under the notation in Theorem 3.15(2), this computation shows that $i_1 = 3, i_2 = 1$, and $i_3 = 2$. By Theorem 3.15(2), we deduce that $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = (1, 4, 3)$.

Indeed,

$$\hat{\mathsf{P}}(\sigma) = \begin{array}{c|c} 5 & 7 & 8 \\ 2 & 3 & 4 & 6 \\ \hline 1 & & \\ \end{array}.$$

4. Distinguished filtrations of \mathcal{V}_{α} and X_{α}

We begin by presenting the background of the problem under consideration. Let M be a finitely generated $H_n(0)$ -module and let

$$\operatorname{ch}([M]) = \sum_{\alpha \models n} c_{\alpha} F_{\alpha} \quad (c_{\alpha} \in \mathbb{Z}_{\geq 0}).$$

Given a composition series

$$0 =: M_0 \subsetneq M_1 \subsetneq M_2 \subsetneq \cdots \subsetneq M_l := M$$

of M, the following hold:

- For each *i*, $ch([M_i/M_{i-1}]) = F_{\alpha}$ for some $\alpha \models n$ with $c_{\alpha} > 0$.
- For each $\alpha \models n$, the number of composition factors M_i/M_{i-1} (with $1 \le i \le l$) satisfying $ch([M_i/M_{i-1}]) = F_{\alpha}$ is equal to c_{α} .

Now, suppose that ch([M]) expands positively in \mathcal{B} , where $\mathcal{B} = \{\mathcal{B}_{\alpha} \mid \alpha \in \mathcal{I}\}$ is a linearly independent subset of $QSym_n$ such that \mathcal{B}_{α} is *F*-positive for all $\alpha \in \mathcal{I}$, with \mathcal{I} being an index set. One may naturally ask whether a similar phenomenon occurs in this context. To address this question, Kim, Lee, and Oh introduced the notion of distinguished filtrations of an $H_n(0)$ -module.

Definition 4.1. ([22, Definition 6.5]) Let $\mathcal{B} = \{\mathcal{B}_{\alpha} \mid \alpha \in \mathcal{I}\}$ be a linearly independent subset of QSym_n with the property that \mathcal{B}_{α} is *F*-positive for all $\alpha \in \mathcal{I}$, where \mathcal{I} is an index set. Given an $H_n(0)$ -module M, a distinguished filtration of M with respect to \mathcal{B} is an $H_n(0)$ -submodule series of M

$$0 =: M_0 \subsetneq M_1 \subsetneq M_2 \subsetneq \cdots \subsetneq M_l := M$$

such that for all $1 \leq k \leq l$, $ch([M_k/M_{k-1}]) = \mathcal{B}_{\alpha}$ for some $\alpha \in \mathcal{I}$.

As seen in [22, Example 6.6], a distinguished filtration of M with respect to \mathcal{B} may not exist even if ch([M]) expands positively in \mathcal{B} . This is because the category $H_n(0)$ -mod is neither semisimple nor representation-finite when n > 3 ([13, 14]).

Let us review the results presented in [22, Section 6.2]. The authors address the problem in the case where M is the weak Bruhat interval module B(I) associated with a dual plactic-closed left weak Bruhat interval I and \mathcal{B} is the Schur basis $\{s_{\lambda} \mid \lambda \vdash n\}$. It is well known that ch([B(I)]) is a skew Schur function. Under this assumption, it was proved in [22, Theorem 6.7] that B(I) admits a distinguished filtration with respect to the Schur basis. In the proof, the combinatorics related to the Robinson-Schensted-Knuth algorithm plays a crucial role.

Now, consider the basis $S := \{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ of the quasisymmetric Schur functions and the basis $\hat{S} := \{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ of the Young quasisymmetric Schur functions, rather than the Schur basis. It is known that every Schur function s_{λ} expands positively in both Sand \hat{S} . Consequently, one might anticipate that the modules examined in [22] possess a distinguished filtration with respect to these bases. However, as seen in Appendix A, this expectation does not hold in general.

In this paper, the focus is on the cases where $M = \mathcal{V}_{\alpha}$ or X_{α} and $\mathcal{B} = \hat{\mathcal{S}}$ for the following reasons:

- According to [1, Theorem 1.1], $ch([\mathcal{V}_{\alpha}])$ expands positively in $\hat{\mathcal{S}}$, even though it is not a symmetric function.
- According to [25, Theorem 4.10], $ch([X_{\alpha}])$ expands positively in \hat{S} when α is a composition of n obtained by shuffling a partition and (1^k) for some $k \ge 0$.
- These cases can be handled in a combinatorial manner by virtue of Algorithm 3.3.
- By [12, Theorem 3.5], X_{α} is a quotient module of \mathcal{V}_{α} , revealing an important relationship between these two structures.

4.1. Distinguished filtrations of \mathcal{V}_{α} . We introduce a new relation on \mathfrak{S}_n derived from Algorithm 3.3.

Definition 4.2. Define a relation \simeq_M on \mathfrak{S}_n by

$$\sigma \simeq_M \rho$$
 if $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = \operatorname{sh}(\hat{\mathsf{P}}(\rho))$ and $\hat{\mathsf{Q}}(\sigma) = \hat{\mathsf{Q}}(\rho)$.

It is straightforward to verify that \simeq_M is an equivalence relation on \mathfrak{S}_n . Moreover, it follows from (3.4) and (3.5) that

$$\sigma \simeq_M \rho$$
 if and only if $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = \operatorname{sh}(\hat{\mathsf{P}}(\rho))$ and $\mathsf{Q}(\sigma) = \mathsf{Q}(\rho)$.

Recall that the dual Knuth equivalence relation $\stackrel{K^*}{\cong}$ on \mathfrak{S}_n is defined by

$$\sigma \stackrel{K^*}{\cong} \rho$$
 if $\sigma^{-1} \stackrel{K}{\cong} \rho^{-1}$, equivalently, $\mathsf{Q}(\sigma) = \mathsf{Q}(\rho)$.

This indicates that \simeq_M is a refinement of the dual Knuth equivalence relation \cong . To elaborate further, let $\sigma_0 \in \mathfrak{S}_n$ and let λ denote the shape of $\mathsf{P}(\sigma_0)$. The dual Knuth equivalence class C containing σ_0 can be expressed as the union of equivalence classes under \simeq_M as follows:

$$C = \bigsqcup_{\substack{\beta \models n \\ \lambda(\beta) = \lambda}} \{ \sigma \in \mathfrak{S}_n \mid \operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = \beta \text{ and } \hat{\mathsf{Q}}(\sigma) = \hat{\mathsf{Q}}(\sigma_0) \}.$$

Proposition 3.7 implies that if a subset S of \mathfrak{S}_n is closed under \simeq_M , then the quasisymmetric function

$$\sum_{\sigma \in S} F_{\operatorname{comp}(\operatorname{Des}_L(\sigma))}$$

expands positively in \hat{S} . In particular, when S is an equivalence class under \simeq_M , this function is equal to $\hat{\mathscr{S}}_{\beta}$, where β is the shape of $\hat{\mathsf{P}}(\sigma)$ for any $\sigma \in S$.

Let $\alpha \models n$. Recall that $\mathcal{V}_{\alpha} \cong \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}),\mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha}))$ (see Theorem 2.8). We will show that $[\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}),\mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L} \cdot w_{0}$ is closed under \simeq_{M} . To do so, we first review the results of Allen, Hallam, and Mason [1]. Therein, they proved that the dual immaculate quasisymmetric function $\mathfrak{S}^{*}_{\alpha}$ expands positively in $\hat{\mathcal{S}}$. To be precise, they first map a word $w = w_{1}w_{2}\ldots w_{n}$ to a pair $(\hat{\mathsf{P}}(w), \hat{Q}(w))$ of fillings of the same shape. Here, $\hat{Q}(w)$ is obtained by filling the new cell created during the insertion procedure $w_{k} \to \hat{\mathsf{P}}(w[1:k-1])$ with k, for each $1 \leq k \leq n$. They then showed that for $\sigma \in \mathfrak{S}_{n}$, $\hat{Q}(\sigma)$ is a dual immaculate recording tableau (DIRT) with a row strip shape α^{r} if and only if $\sigma \in [\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L} \cdot w_{0}$ (for undefined terms such as dual immaculate recording tableaus and row strip shapes, see [1, Definitions 3.9 and 3.4]). Using this property, they established the bijection

$$\underline{\text{AHM}} : \text{SIT}(\alpha) \to \bigsqcup_{\beta \models n} \text{SYCT}(\beta) \times \text{DIRT}(\beta, \alpha^{r})$$

$$\mathscr{T} \mapsto (\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_{0}), \hat{Q}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_{0})).$$

$$(4.1)$$

Second, we observe that for $\sigma, \rho \in \mathfrak{S}_n$,

if
$$\hat{Q}(\sigma) = \hat{Q}(\rho)$$
, then $\hat{\mathsf{Q}}(\sigma) = \hat{\mathsf{Q}}(\rho)$. (4.2)

Indeed, this can be derived by combining the following facts for a word $w = w_1 w_2 \dots w_n$:

- $\hat{Q}(w)$ records the row and column indices of the new cell created during the insertion procedure $w_k \to \hat{\mathsf{P}}(w[1:k-1])$, for all $1 \le k \le n$,
- $\hat{\mathsf{Q}}(w)$ records only the column index of the new cell created during the insertion procedure $w_k \to \hat{\mathsf{P}}(w[1:k-1])$, for all $1 \le k \le n$.

The following lemma is derived by combining (4.1) and (4.2) with the C-linear isomorphism in Theorem 2.8(1).

Lemma 4.3. For a composition α of n, $[\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L} \cdot w_{0}$ is closed under \simeq_{M} .

Remark 4.4. (1) Lemma 4.3 does not imply that $[\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L}$ is closed under \simeq_{M} . More generally, even if a subset $S \subseteq \mathfrak{S}_{n}$ is closed under \simeq_{M} , it does not necessarily follow that the sets $S \cdot w_{0}, w_{0} \cdot S$, and $w_{0} \cdot S \cdot w_{0}$ are also closed under \simeq_{M} . For instance, consider $S = \{3124, 4123\} \subseteq \mathfrak{S}_{4}$; in this case, none of $S \cdot w_{0}, w_{0} \cdot S$, and $w_{0} \cdot S \cdot w_{0}$ is closed under \simeq_{M} . For instance, consider \simeq_{M} . On the other hand, if S is closed under the dual Knuth equivalence relation $\stackrel{K^{*}}{\cong}$, then the sets $S \cdot w_{0}, w_{0} \cdot S$, and $w_{0} \cdot S \cdot w_{0}$ remain closed under $\stackrel{K^{*}}{\cong}$. This property is an intriguing distinction between \simeq_{M} and $\stackrel{K^{*}}{\cong}$. (2) For $\sigma, \rho \in \mathfrak{S}_n$, the equality $\hat{\mathsf{Q}}(\sigma) = \hat{\mathsf{Q}}(\rho)$ does not necessarily imply that $\hat{Q}(\sigma) = \hat{Q}(\rho)$. For instance, let $\sigma = 621543$ and $\rho = 531426$. Then

$$\hat{\mathsf{Q}}(\sigma) = \hat{\mathsf{Q}}(\rho) = \begin{bmatrix} 6 \\ 3 & 4 \\ 2 & 5 \\ 1 \end{bmatrix}, \text{ but } \hat{Q}(\sigma) = \begin{bmatrix} 3 \\ 6 \\ 2 & 4 \\ 1 & 5 \end{bmatrix} \neq \hat{Q}(\rho) = \begin{bmatrix} 6 \\ 3 \\ 2 & 4 \\ 1 & 5 \end{bmatrix}.$$

For any $\mathscr{T} \in \operatorname{SIT}(\alpha)$, the set of entries in the first column of $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0)$ is equal to the set of entries in the first column of \mathscr{T} . Combined with Theorem 3.15, this observation directly leads to the following lemma.

Lemma 4.5. Let α be a composition of n and $\mathscr{T} \in SIT(\alpha)$.

- (1) The length of the shape of $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0)$ is equal to $\ell(\alpha)$.
- (2) For $1 \le k \le \ell(\alpha)$,

$$\underline{\mathsf{IES}}_k(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0) \subseteq \{\mathscr{T}(i,1) \mid 1 \le i \le \ell(\alpha)\}.$$

In particular, the equality holds when $k = \ell(\alpha)$.

Let \leq_{lex} denote the lexicographic order on the set of all compositions of n. The following lemma is essential in constructing our distinguished filtrations.

Lemma 4.6. Let α be a composition of n and let $T, S \in SIT(\alpha)$. If $w_r(T) \stackrel{K^*}{\cong} w_r(S)$ and $w_r(T) \preceq_L w_r(S)$, then

$$\operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(T)w_0)) \ge_{\operatorname{lex}} \operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(S)w_0)).$$

Proof. We may assume that $w_r(S) = s_i w_r(T)$ for some $1 \le i \le n-1$. Then *i* is strictly below i + 1 in *T* and one of the following conditions holds:

- Neither i nor i + 1 are not in the first column of T, or
- i + 1 is in the first column of T, while i is not.

For simplicity, let $\lambda := \operatorname{sh}(\mathsf{P}(\mathsf{w}_{\mathsf{r}}(T)w_0))$ and $l := \ell(\lambda)$. Let $1 \leq k \leq l$. Note that i + 1 is to the left of i in $\mathsf{w}_{\mathsf{r}}(T)w_0$ when written in one-line notation. From this, it follows that $s_i \cdot u \in \operatorname{Inc}_k(\mathsf{w}_{\mathsf{r}}(S)w_0)$ for any $u \in \operatorname{Inc}_k(\mathsf{w}_{\mathsf{r}}(T)w_0)$. Here, $s_i \cdot u$ denotes the k-tuple of sequences obtained from u by swapping i and i + 1. Therefore,

$$s_i \cdot \underline{\mathsf{IES}}_k(\mathsf{w}_\mathsf{r}(T)w_0) \leq_{\text{set}} \underline{\mathsf{IES}}_k(\mathsf{w}_\mathsf{r}(S)w_0).$$
(4.3)

By Lemma 4.5, there exist subsets $I_k, J_k \subseteq [l]$ with $|I_k| = |J_k| = k$ such that

$$\underline{\mathsf{IES}}_k(\mathsf{w}_\mathsf{r}(T)w_0) = \{T(r,1) \mid r \in I_k\} \text{ and } \underline{\mathsf{IES}}_k(\mathsf{w}_\mathsf{r}(S)w_0) = \{S(r,1) \mid r \in J_k\}.$$

Combining these equalities with (4.3) yields that

$$I_k \leq_{\text{set}} J_k. \tag{4.4}$$

S.-Y. LEE AND Y.-T. OH

Let $\alpha := \operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(T)w_0))$ and $\beta := \operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(S)w_0))$. We claim that $\alpha \geq_{\operatorname{lex}} \beta$. Suppose to the contrary that $\alpha <_{\operatorname{lex}} \beta$. Then, there exists a unique index $1 \leq t_0 \leq l$ such that $\alpha_t = \beta_t$ for all $1 \leq t < t_0$ and $\alpha_{t_0} < \beta_{t_0}$. Thus, we can select an index $1 \leq k_0 < l$ such that $\lambda_{k_0} = \beta_{t_0}$ and $\lambda_{k_0} > \lambda_{k_0+1}$. Since $\lambda(\alpha) = \lambda(\beta) = \lambda$, by Theorem 3.15, we have $I_{k_0} = \{1 \leq r \leq l \mid \alpha_r \geq \lambda_{k_0}\}$ and $J_{k_0} = \{1 \leq r \leq l \mid \beta_r \geq \lambda_{k_0}\}$. Consequently,

$$I_{k_0} = A \sqcup B$$
 and $J_{k_0} = A' \sqcup \{t_0\} \sqcup B'$,

for some $A, A' \subseteq [1, t_0 - 1]$ and $B, B' \subseteq [t_0 + 1, l]$. Note that A = A' since $\alpha_t = \beta_t$ for all $1 \leq t < t_0$. Furthermore, since $|I_{k_0}| = |J_{k_0}|$, it follows that $I_{k_0} >_{\text{set}} J_{k_0}$. This contradicts the inequality in (4.4). Therefore, our assertion follows.

Recall that in the proof of [22, Theorem 6.7], the following order relation plays a key role: for $\sigma, \rho \in \mathfrak{S}_n$ with $\sigma \preceq_L \rho$, it holds that

$$\mathbf{Q}(\sigma) = \mathbf{Q}(\rho) \quad \text{or} \quad \operatorname{sh}(\mathbf{Q}(\sigma)) \triangleright \operatorname{sh}(\mathbf{Q}(\rho)), \tag{4.5}$$

where \triangleright denotes the dominance order on the set of partitions of n. This result can be derived by applying Taşkin's result [33, Proposition 3.2.5] to the weak order¹ on SYT_n given in [33, Definition 3.1.3]. This relation also plays a crucial role in the proof of the subsequent theorem.

Theorem 4.7. For a composition α of n, \mathcal{V}_{α} has a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models n\}.$

Proof. By Theorem 2.8(1), the assertion is equivalent to stating that $\mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}),\mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha}))$ has a distinguished filtration with respect to $\{\mathscr{S}_{\beta} \mid \beta \models n\}$. For simplicity, set $I := [\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}),\mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L}$. Let $\mathcal{L} := \{\hat{\mathsf{Q}}(\sigma w_{0}) \mid \sigma \in I\}$ and choose an arbitrary total order $<_{\mathcal{L}}$ on \mathcal{L} such that

$$T <_{\mathcal{L}} S$$
 if $\lambda(\operatorname{sh}(T)) \triangleright \lambda(\operatorname{sh}(S))$.

For $Q \in \mathcal{L}$, set

$$A_Q := \{ \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) \mid \sigma \in I \text{ and } \hat{\mathsf{Q}}(\sigma w_0) = Q \}$$

Let

$$R := \bigsqcup_{Q \in \mathcal{L}} \{Q\} \times A_Q,$$

and define a total order \ll on R by

 $(T,\beta) \ll (S,\gamma)$ if either $T <_{\mathcal{L}} S$ or $(T = S \text{ and } \beta <_{\text{lex}} \gamma)$.

Enumerate the elements in R in increasing order with respect to the total order \ll so that

$$R = \{ (Q_1, \gamma_1) \ll (Q_2, \gamma_2) \ll \dots \ll (Q_l, \gamma_l) \}.$$
(4.6)

¹This order was originally defined in [27, 2.5.1], where it is called the *induced Duflo order*.

Now, for each $1 \leq k \leq l$, define

$$B_k := \{ \sigma \in I \mid \hat{\mathsf{Q}}(\sigma w_0) = Q_i \text{ and } \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) = \gamma_i \text{ for some } 1 \le i \le k \},$$
(4.7)

with $B_0 = \emptyset$. In particular, $B_l = I$. We prove the assertion in two steps:

Step 1. For each $1 \leq k \leq l$, we claim that $M_k := \mathbb{C}B_k$ is an $H_n(0)$ -submodule of $\mathsf{B}(I)$. To prove this, it suffices to verify that

$$\pi_i \cdot \sigma \subseteq B_k \cup \{0\}$$
 for all $\sigma \in B_k$ and $1 \le i \le n-1$.

Let $\sigma \in B_k$ and $1 \leq i \leq n-1$. If $\pi_i \cdot \sigma$ is σ or 0, we are done. Suppose that $\pi_i \cdot \sigma = s_i \sigma$, that is, $s_i \sigma \in I$ and $\ell(s_i \sigma) = \ell(\sigma) + 1$. Let s (respectively, t) be the smallest positive integer $1 \leq k \leq l$ such that $\sigma w_0 \in B_k$ (respectively, $s_i \sigma w_0 \in B_k$). To prove the claim, it suffices to show that $s \geq t$.

- Suppose that the rearrangements of γ_s and γ_t are the same. This assumption is equivalent to saying that the shapes of $\mathsf{P}(\sigma w_0)$ and $\mathsf{P}(s_i \sigma w_0)$ are the same. Note $\sigma w_0 \succeq_L s_i \sigma w_0$. Therefore, by (4.5), we deduce that $\mathsf{Q}(\sigma w_0) = \mathsf{Q}(s_i \sigma w_0)$, which is equivalent that $\hat{\mathsf{Q}}(\sigma w_0) = \hat{\mathsf{Q}}(s_i \sigma w_0)$. Also, by Lemma 4.6, we have $\gamma_s \geq_{\text{lex}} \gamma_t$. Therefore, $s \geq t$.
- Suppose that the rearrangements of γ_s and γ_t are different. This assumption is equivalent to saying that the shapes of $\mathsf{P}(\sigma w_0)$ and $\mathsf{P}(s_i \sigma w_0)$ are different. Note that $\sigma w_0 \succeq_L s_i \sigma w_0$. Therefore, by (4.5), we deduce that

$$\lambda(\operatorname{sh}(Q_s)) = \operatorname{sh}(\mathsf{Q}(\sigma w_0)) \triangleleft \operatorname{sh}(\mathsf{Q}(s_i \sigma w_0)) = \lambda(\operatorname{sh}(Q_t)).$$

As a result, we have $Q_s >_{\mathcal{L}} Q_t$, which implies that s > t.

Step 2. For each $1 \leq k \leq l$, we claim that $ch([M_k/M_{k-1}]) = \hat{\mathscr{S}}_{\gamma_k}$. Since M_{k-1} and M_k are submodules of B(I), it follows that

$$\operatorname{ch}([M_k/M_{k-1}]) = \operatorname{ch}([M_k]) - \operatorname{ch}([M_{k-1}]) = \sum_{\sigma \in B_k \setminus B_{k-1}} F_{\operatorname{comp}(\operatorname{Des}_L(\sigma))^c}$$

(see (2.2)). On the other hand, in view of Lemma 4.3, we see that the restriction

$$\underline{\operatorname{AHM}} \circ (\Theta_V)^{-1}|_{B_k} \colon B_k \to \bigsqcup_{1 \le i \le k} \operatorname{SYCT}(\gamma_i) \times \{Q_i\}, \quad \sigma \mapsto (\hat{\mathsf{P}}(\sigma w_0), \hat{\mathsf{Q}}(\sigma w_0))$$

gives a bijection and

AHM
$$\circ (\Theta_V)^{-1}(B_k \setminus B_{k-1}) = \text{SYCT}(\gamma_k) \times \{Q_k\}$$
 (4.8)

(for the definition of Θ_V , see Theorem 2.8(1)). Consequently, we have

$$\operatorname{ch}([M_k/M_{k-1}]) = \sum_{\sigma \in B_k \setminus B_{k-1}} F_{\operatorname{comp}(\operatorname{Des}_L(\sigma w_0))} = \hat{\mathscr{P}}_{\gamma_k}$$

where the second is derived from (4.8) and Proposition 3.7.

By Step 1 and Step 2, we conclude that

$$0 \subsetneq M_1 \subsetneq M_2 \subsetneq \cdots \subsetneq M_l = \mathsf{B}(I)$$

forms a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models n\}$.

Example 4.8. Let $\alpha = (2, 2, 2)$. Then, $\mathcal{V}_{\alpha} \cong \mathsf{B}(I)$, where $I := [214365, 615243]_L$. Note that $\{\hat{\mathsf{Q}}(\gamma w_0) \mid \gamma \in I\}$ is given by

$$\left\{Q_1 := \begin{array}{c|c} 6\\5\\5\\1&2&3&4 \end{array}, Q_2 := \begin{array}{c|c} 5&6\\3\\3\\1&2&4 \end{array}, Q_3 := \begin{array}{c|c} 5\\5\\3&4&6\\1&2 \end{array}, Q_4 := \begin{array}{c|c} 5&6\\3&4\\1&2 \end{array}\right\}.$$

Following the method presented in the proof of Theorem 4.7, we will construct a distinguished filtration of B(I) with respect to $\{\hat{\mathscr{S}}_{\alpha} \mid \alpha \models 6\}$. For $1 \le i \le 4$, let

$$A_i := \{ \operatorname{sh}(\widehat{\mathsf{P}}(\sigma w_0)) \mid \sigma \in I \text{ with } \widehat{\mathsf{Q}}(\sigma w_0) = Q_i \}.$$

We see that

 $A_1 = \{(1,1,4)\}, \ A_2 = \{(1,2,3), (2,1,3)\}, \ A_3 = \{(1,2,3), (1,3,2)\}, \ A_4 = \{(2,2,2)\}.$ For $1 \le i \le 4$ and $\alpha \in A_i$, let

$$B'_{i;\alpha} := \{ \sigma \in [214365, 615243]_L \mid \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) = \alpha \text{ and } \hat{\mathsf{Q}}(\sigma w_0) = Q_i \}$$

and

$$B_{i;\alpha} := \left(\bigsqcup_{\substack{1 \le t < i \\ \beta \in A_t}} B'_{t;\beta}\right) \bigsqcup \left(\bigsqcup_{\substack{\beta \in A_i \\ \text{with } \beta \le \text{lex } \alpha}} B'_{i;\beta}\right) \ .$$

Then,

$$0 = \mathbb{C}B_{1;(1,1,4)} \subsetneq \mathbb{C}B_{2;(3,2,1)} \subsetneq \mathbb{C}B_{2;(2,3,1)} \subsetneq \mathbb{C}B_{3;(3,2,1)} \subsetneq \mathbb{C}B_{3;(2,1,3)} \subsetneq \mathbb{C}B_{4;(2,2,2)} = \mathsf{B}(I)$$

is a distinguished filtration of $\mathsf{B}(I)$ with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models 6\}$.

4.2. Distinguished filtrations of X_{α} . In this subsection, α is assumed to be a composition of *n* that is a shuffle of a partition and (1^k) for some $k \ge 0$. Under this assumption, Marcum and Niese [25] showed that restricting the domain of the bijection <u>AHM</u> given in (4.1) to SET(α) induces the bijection

$$\underline{\operatorname{AHM}}|_{\operatorname{SET}(\alpha)} : \operatorname{SET}(\alpha) \to \bigsqcup_{\beta \models n} \operatorname{SYCT}(\beta) \times \operatorname{DIRT}^*(\beta, \alpha^{\mathrm{r}}),$$

where DIRT^{*}($\beta, \alpha^{\rm r}$) is the set of DIRTs in DIRT($\beta, \alpha^{\rm r}$) that have no *exceptions* (for the precise definition of DIRT^{*}($\beta, \alpha^{\rm r}$), see [25, Definition 4.7]). Using this, they [25, Theorem 3.5] proved that \mathcal{E}_{α} expands positively in $\hat{\mathcal{S}}$.

Combined with the C-linear isomorphism in Theorem 2.8(2), the bijection $\underline{AHM}|_{SET(\alpha)}$ and (4.2) yield the following lemma.



FIGURE 4.1. The $H_6(0)$ -action on the basis $[214365, 615243]_L$ for $B(214365, 615243) \cong \mathcal{V}_{(2,2,2)}$ and the sets $B'_{i;\alpha}$'s

Lemma 4.9. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then $[\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathsf{T}'_{\alpha})]_L \cdot w_0$ is closed under \simeq_M .

Moreover, since $\text{SET}(\alpha) \subseteq \text{SIT}(\alpha)$, Lemma 4.5 and Lemma 4.6 remain valid. Combining these lemmas with Lemma 4.9, we can derive the following theorem using the same approach as in the proof of Theorem 4.7. It should be noted that, in this case, R in (4.6) can be described more simply. Specifically, it consists only of γ_i 's rather than (Q_i, γ_i) 's.

Theorem 4.10. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then X_{α} has a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models n\}$.

Proof. By Theorem 2.8(2), the assertion is equivalent to stating that $\mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathsf{T}'_{\alpha}))$ has a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models n\}$. For simplicity, set $I := [\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathsf{T}'_{\alpha})]_{L}$ and $\lambda := \lambda(\alpha)$.

First, we show that

$$\hat{\mathsf{Q}}(\sigma w_0) = \mathsf{T}_{\alpha^{\mathrm{r}}} \text{ for all } \sigma \in I.$$

$$(4.9)$$

For this purpose, we begin by observing that

$$\operatorname{sh}(\mathsf{P}(\sigma w_0)) = \lambda \text{ for all } \sigma \in I.$$
 (4.10)

Let $\sigma \in I$. Then $\sigma = w_r(T)$ for some $T \in SET(\alpha)$. One can see that for $1 \le k \le n$,

$$\mathfrak{i}_k(\sigma w_0) = \lambda_1 + \dots + \lambda_k$$

Indeed, the inequality \geq follows from the fact that every row of T increases from left to right, while the opposite inequality \leq follows from the fact that every increasing subsequence of σw_0 contains at most one entry in each column of T. Combining this equality with Theorem 3.9, we see that $\operatorname{sh}(\mathsf{P}(\sigma w_0)) = \lambda$, as required. Since $I \cdot w_0$ is a left weak Bruhat interval, by (4.10) and (4.5), we have

$$\mathsf{Q}(\sigma w_0) = \mathsf{Q}(\mathsf{w}_\mathsf{r}(T)w_0) \text{ for any } \sigma \in I.$$

Now, due to (3.4) and (3.5), (4.9) follows from $Q(w_r(T_\alpha)w_0) = T_{\alpha^r}$.

Second, we define B_k for each $1 \leq k \leq l$. Let $R = {\operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) \mid \sigma \in I}$. Enumerate the elements in R in increasing order according to the lexicographic order $\leq_{\operatorname{lex}}$ so that

$$R = \{\gamma_1 <_{\text{lex}} \gamma_2 <_{\text{lex}} \cdots <_{\text{lex}} \gamma_l\}.$$

Now, for each $1 \le k \le l$, define

$$B_k := \{ \sigma \in I \mid \operatorname{sh}(\widehat{\mathsf{P}}(\sigma w_0)) = \gamma_i \text{ for some } 1 \le i \le k \},\$$

with $B_0 = \emptyset$. In particular, $B_l = I$. In view of (4.9), we see that

$$B_k = \{ \sigma \in I \mid \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) = \gamma_i \text{ and } \hat{\mathsf{Q}}(\sigma w_0) = \mathsf{T}_{\alpha^{\mathrm{r}}} \text{ for some } 1 \le i \le k \}.$$

Finally, following the same manner as the proof of Theorem 4.7, we can see that for $1 \le k \le l$, $M_k := \mathbb{C}B_k$ is an $H_n(0)$ -submodule of $\mathsf{B}(I)$ and that

 $0 \subsetneq M_1 \subsetneq M_2 \subsetneq \cdots \subsetneq M_l = \mathsf{B}(I)$

forms a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models n\}$.

Example 4.11. Let $\alpha = (3, 1, 2)$. Then, $I := [w_r(T_\alpha), w_r(T'_\alpha)]_L = [321465, 641253]_L$. And, for every $\sigma \in I$, we have

$$\hat{\mathsf{Q}}(\sigma w_0) = \begin{array}{c|c} 4 & 5 & 6 \\ \hline 3 \\ \hline 1 & 2 \end{array}$$

Let

 $R := \{ \gamma_1 := (1, 2, 3) <_{\text{lex}} \gamma_2 := (1, 3, 2) <_{\text{lex}} \gamma_3 := (2, 1, 3) <_{\text{lex}} \gamma_4 := (3, 1, 2) \}.$

Then, by Theorem 4.10,

$$0 \subsetneq \mathbb{C}B_1 \subsetneq \mathbb{C}B_2 \subsetneq \mathbb{C}B_3 \subsetneq \mathbb{C}B_4 = \mathsf{B}(I)$$

is the desired distinguished filtration of $\mathsf{B}(I)$ with respect to $\{\hat{\mathscr{S}}_{\beta} \mid \beta \models 6\}$.



FIGURE 4.2. The $H_6(0)$ -action on the basis $[215436, 641254]_L$ for $B(215436, 641254) \cong X_{(3,1,2)}$ and the sets $B_i \setminus B_{i-1}$'s

5. INDECOMPOSABLE 0-HECKE MODULES FOR YOUNG QUASISYMMETRIC SCHUR FUNCTIONS AND QUASISYMMETRIC SCHUR FUNCTIONS

Given an important F-positive quasisymmetric function f, constructing an indecomposable $H_n(0)$ -module whose image under the quasisymmetric characteristic is f is a significant open problem. However, this problem remains unsolved for \mathscr{S}_{α} and $\hat{\mathscr{S}}_{\alpha}$. For example, $\operatorname{ch}([\mathbf{S}_{\alpha}]) = \mathscr{S}_{\alpha}$, but \mathbf{S}_{α} is not, in general, indecomposable (see Subsection 2.4.1). The objective of this section is to construct and investigate an indecomposable $H_n(0)$ module \mathbf{Y}_{α} such that $\operatorname{ch}([\mathbf{Y}_{\alpha}]) = \hat{\mathscr{S}}_{\alpha}$. By taking the ϕ -twist of this module, we obtain an indecomposable $H_n(0)$ -module whose image under the quasisymmetric characteristic is \mathscr{S}_{α} .

5.1. Indecomposable $H_n(0)$ -modules \mathbf{Y}_{α} and $\phi[\mathbf{Y}_{\alpha}]$. Let $\alpha \models n$ and let

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_{l-1} \subsetneq M_l = \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})) \ (\cong \mathcal{V}_{\alpha})$$

be a distinguished filtration with respect to $\{\hat{\mathscr{S}}_{\alpha} \mid \alpha \models n\}$, constructed in the proof of Theorem 4.7. A remarkable property of this filtration is that the submodule M_{l-1} is uniquely determined, independent of the choice of filtration. Let us explain how this property is derived. Since $\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha})$ is a generator of $\mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}),\mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha}))$, the minimal index $1 \leq i \leq l$ such that $\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}) \in B_i$ is equal to l (for the definition of B_i , see (4.7)). Combining this with the fact that

$$\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha})w_0) = \mathscr{T}_{\alpha} \text{ and } \hat{\mathsf{Q}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha})w_0) = \mathscr{T}_{\alpha^{\mathrm{r}}},$$

we have

$$B_l \setminus B_{l-1} = \{ \sigma \in [\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_L \mid \operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\sigma w_0))) = \alpha \text{ and } \hat{\mathsf{Q}}(\mathsf{w}_{\mathsf{r}}(\sigma w_0)) = \mathscr{T}_{\alpha^{\mathrm{r}}} \}.$$

This shows that the submodule M_{l-1} is given by the \mathbb{C} -span of

$$\{\sigma \in [\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})]_{L} \mid \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_{0})) \neq \alpha \text{ or } \hat{\mathsf{Q}}(\sigma w_{0}) \neq \mathscr{T}_{\alpha^{\mathrm{r}}}\}$$
(5.1)

and it depends only on α , not on the choice of filtration.

Now, define the $H_n(0)$ -module \mathbf{Y}_{α} by

$$\mathbf{Y}_{\alpha} := M_l / M_{l-1}.$$

We have a surjective $H_n(0)$ -module homomorphism

$$\delta: \mathcal{V}_{\alpha} \xrightarrow{\Theta_{V}} \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})) \xrightarrow{\mathrm{pr}} \mathbf{Y}_{\alpha},$$

where pr : $\mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathscr{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\mathscr{T}'_{\alpha})) \to \mathbf{Y}_{\alpha}$ is the natural projection. By (5.1), \mathbf{Y}_{α} can be viewed as the $H_n(0)$ -module whose underlying space is the C-span of

$$\mathcal{K}_{\alpha} := \{ \sigma \in \mathfrak{S}_n \mid \operatorname{sh}(\hat{\mathsf{P}}(\sigma w_0)) = \alpha \text{ and } \hat{\mathsf{Q}}(\sigma w_0) = \mathscr{T}_{\alpha^{\mathrm{r}}} \},$$
(5.2)

equipped with the $H_n(0)$ -action defined as follows: for $\sigma \in \mathcal{K}_{\alpha}$ and $1 \leq i \leq n-1$,

$$\pi_i \cdot \sigma = \begin{cases} \sigma & \text{if } i \in \text{Des}_L(\sigma), \\ s_i \sigma & \text{if } i \notin \text{Des}_L(\sigma) \text{ and } s_i \sigma \in \mathcal{K}_\alpha, \\ 0 & \text{otherwise.} \end{cases}$$

Since $|\text{DIRT}(\alpha, \alpha^r)| = 1$ by [1, Theorem 1.1], the basis \mathcal{K}_{α} can be rewritten as

$$\{\mathsf{w}_{\mathsf{r}}(\mathscr{T}) \mid \mathscr{T} \in \operatorname{SIT}(\alpha) \text{ and } \operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0)) = \alpha\}.$$
(5.3)

Recall that every automorphism of $H_n(0)$ induces an equivalence on the category of finitely generated $H_n(0)$ -modules. Let $\phi : H_n(0) \to H_n(0)$ be the automorphism defined by $\pi_i \mapsto \pi_{n-i}$. For an $H_n(0)$ -module M, let $\phi[M]$ denote the ϕ -twist of M. For more details, see [21, Section 3.4].

Theorem 5.1. Let α be a composition of n.

- (1) \mathbf{Y}_{α} is an indecomposable $H_n(0)$ -module whose image under the quasisymmetric characteristic is $\hat{\mathscr{S}}_{\alpha}$.
- (2) The ϕ -twist $\phi[\mathbf{Y}_{\alpha}]$ of \mathbf{Y}_{α} is an indecomposable $H_n(0)$ -module whose image under the quasisymmetric characteristic is \mathscr{S}_{α} .

Proof. (1) Note that $\mathcal{K}_{\alpha} \cdot w_0$ is an equivalence class under \simeq_M containing $w_r(\mathscr{T}_{\alpha})w_0$ and $ch([\mathbf{Y}_{\alpha}])$ is given by

$$\sum_{\sigma \in \mathcal{K}_{\alpha}} F_{\operatorname{comp}(\operatorname{Des}_{L}(\sigma))^{c}} \left(= \sum_{\sigma \in \mathcal{K}_{\alpha} \cdot w_{0}} F_{\operatorname{comp}(\operatorname{Des}_{L}(\sigma))} \right).$$

This shows that $\operatorname{ch}([\mathbf{Y}_{\alpha}]) = \hat{\mathscr{S}}_{\alpha}$.

Next, we show that \mathbf{Y}_{α} is indecomposable. It was shown in [12, Theorem 3.2] that there exists a surjective $H_n(0)$ -module homomorphism $\Phi : \overline{\mathbf{P}}_{\alpha^c} \to \mathcal{V}_{\alpha}$. Here, $\overline{\mathbf{P}}_{\alpha^c}$ denotes the projective indecomposable $H_n(0)$ -module indexed by α^c as defined in [12, (2.2)]. Then, we have a surjective $H_n(0)$ -module homomorphism $\delta \circ \Phi : \overline{\mathbf{P}}_{\alpha^c} \to \mathbf{Y}_{\alpha}$. The desired result follows from the general fact that every quotient of a projective indecomposable $H_n(0)$ -module is indecomposable.

(2) Note that the automorphism $\boldsymbol{\phi}$ induces an equivalence on the category of finitely generated $H_n(0)$ -modules. Hence, $\boldsymbol{\phi}[\mathbf{Y}_{\alpha}]$ is indecomposable since \mathbf{Y}_{α} is indecomposable by (1). Furthermore, we have $\operatorname{ch}([\boldsymbol{\phi}[\mathbf{Y}_{\alpha}]]) = \mathscr{S}_{\alpha}$ since $\boldsymbol{\phi}[\mathbf{F}_{\beta}] \cong \mathbf{F}_{\beta^{\mathrm{r}}}$ (for instance, see [21, Table 2]).

5.2. A surjection series containing \mathbf{Y}_{α} . Let $\alpha \models n$. It was shown in [12, Corollary 4.6] that there exists a series of surjective $H_n(0)$ -module homomorphisms given by

$$\overline{\mathbf{P}}_{\alpha^{c}} \xrightarrow{\Phi} \mathcal{V}_{\alpha} \xrightarrow{\Gamma} X_{\alpha} \xrightarrow{\tilde{\eta}} \hat{\mathbf{S}}_{\alpha,C},$$

where

- Φ is the surjective $H_n(0)$ -module homomorphism described in [12, (3.2)],
- Γ is the surjective $H_n(0)$ -module homomorphism such that for $\mathscr{T} \in SIT(\alpha)$,

$$\Gamma(\mathscr{T}) = \begin{cases} \mathscr{T} & \text{if } \mathscr{T} \in \text{SET}(\alpha), \\ 0 & \text{otherwise} \end{cases}$$
(5.4)

as given in [12, (3.4)],

• $\tilde{\eta}$ is the surjective $H_n(0)$ -homomorphism such that for $T \in \text{SET}(\alpha)$,

$$\tilde{\eta}(T) = \begin{cases} T & \text{if } T \in \text{SYCT}(\alpha, C), \\ 0 & \text{otherwise,} \end{cases}$$
(5.5)

as detailed in [12, Section 3.3 and Section 4.2].

Remark 5.2. (1) In [12], the map $\tilde{\eta}$ is defined using a different formulation. Nevertheless, it can be readily verified that this definition is equivalent to the one given in (5.5).

(2) Using (5.5), one can easily verify that the essential epimorphism $\eta : \mathbf{P}_{\alpha^c} \twoheadrightarrow \mathbf{S}_{\alpha,C}$ in [12, Theorem 5.3] satisfies

$$\eta = \tilde{\eta} \circ \Gamma \circ \Phi.$$

In this subsection, we show that there is a surjective $H_n(0)$ -module homomorphism $\Upsilon: \mathbf{Y}_{\alpha} \to \hat{\mathbf{S}}_{\alpha,C}$ such that the diagram



commutes. In the case where α is a shuffle of a partition and (1^k) for some $k \ge 0$, we also show that there exists a surjective $H_n(0)$ -module homomorphism $\tilde{\delta} : X_{\alpha} \to \mathbf{Y}_{\alpha}$ such that the diagram



commutes.

Proposition 5.3. For a composition α of n, there is a unique $H_n(0)$ -module homomorphism $\Upsilon : \mathbf{Y}_{\alpha} \to \hat{\mathbf{S}}_{\alpha,C}$ such that $\Upsilon \circ \delta = \tilde{\eta} \circ \Gamma$.

Proof. To prove the assertion, it suffices to show that $\ker(\tilde{\eta} \circ \Gamma) \supseteq \ker(\delta)$. To this end, we first describe $\ker(\tilde{\eta} \circ \Gamma)$ and $\ker(\delta)$ explicitly. By (5.4) and (5.5), $\ker(\tilde{\eta} \circ \Gamma)$ is given by

$$\ker(\tilde{\eta} \circ \Gamma) = \mathbb{C}\{\mathscr{T} \in \operatorname{SIT}(\alpha) \mid \mathscr{T} \notin \operatorname{SYCT}(\alpha; C)\}.$$

On the other hand, by the definition of \mathbf{Y}_{α} , we have $\ker(\delta) = \Theta_V^{-1}(M_{l-1})$. Combining this equality with (5.3), we obtain

$$\ker(\delta) = \mathbb{C}\{\mathscr{T} \in \operatorname{SIT}(\alpha) \mid \operatorname{sh}(\mathsf{P}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0)) \neq \alpha\}.$$

To establish the desired inclusion, we will use mathematical induction on $n = |\alpha|$. Specifically, we aim to show that if $\tau \in \text{SYCT}(\alpha; C)$, then $\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(\tau)w_0) = \tau$. If $\ell(\alpha) = 1$, there is nothing to prove. So, assume that $l := \ell(\alpha) > 1$. Let $\sigma := \mathsf{w}_{\mathsf{r}}(\tau)w_0$ and $\sigma' := \sigma[1:n-1]$. Let τ' be the filling obtained from τ by removing the cell containing $\tau(1, \alpha_1) (= \sigma(n))$. Then, τ' is a Young composition tableau filled with distinct entries and its shape is

$$\beta = \begin{cases} (\alpha_2, \dots, \alpha_l) & \text{if } \alpha_1 = 1, \\ (\alpha_1 - 1, \alpha_2, \dots, \alpha_l) & \text{if } \alpha_1 > 1. \end{cases}$$

By the induction hypothesis, we have $\hat{\mathsf{P}}(\sigma') = \tau'$. Thus, it remains to show that

$$\tau(1, \alpha_1)$$
 is placed in the cell $(1, \alpha_1)$ in $\mathsf{P}(\sigma)$. (5.6)

If $\alpha_1 = 1$, then (5.6) follows immediately since $\tau(1, 1) = 1$. From now on, suppose that $\alpha_1 > 1$. Then (5.6) follows by showing that for $(i, j) \in cd(\beta)$,

if
$$i \geq 2$$
 and $j \geq \alpha_1 - 1$, then $\tau(i, j) > \tau(1, \alpha_1)$.

In the case where $j \ge \alpha_1$, since $\tau \in \text{SYCT}(\alpha; C)$, it follows that $\tau(i, j) \ge \tau(i, \alpha_1) > \tau(1, \alpha_1)$. Now, consider the case where $j = \alpha_1 - 1$. Suppose, for the sake of contradiction, that $\tau(i, \alpha_1 - 1) < \tau(1, \alpha_1)$. By Young triple rule, this inequality implies that $(i, \alpha_1) \in \mathsf{cd}(\alpha)$ and $\tau(i, \alpha_1) < \tau(1, \alpha_1)$. However, this contradicts the assumption that $\tau \in \text{SYCT}(\alpha; C)$.

Remark 5.4. In [34, Section 7], Tewari and van Willigenburg introduced the notion of simple compositions to characterize when \mathbf{S}_{α} is indecomposable. A composition α is called simple if, for any $1 \leq i < j \leq \ell(\alpha)$ with $\alpha_i \geq \alpha_j \geq 2$, there exists an integer k such that i < k < j and $\alpha_k = \alpha_j - 1$. Since $\hat{\mathbf{S}}_{\alpha} = \boldsymbol{\phi}[\mathbf{S}_{\alpha^r}]$ by [12, (4.2)], [34, Theorem 7.6] says that $\hat{\mathbf{S}}_{\alpha}$ is indecomposable, equivalently, $\hat{\mathbf{S}}_{\alpha} = \hat{\mathbf{S}}_{\alpha,C}$ if and only if α^r is simple. Combining this with the equality $ch([\mathbf{Y}_{\alpha}]) = \hat{\mathscr{S}}_{\alpha} = ch([\hat{\mathbf{S}}_{\alpha}])$, it follows from Proposition 5.3 that $\Upsilon : \mathbf{Y}_{\alpha} \to \hat{\mathbf{S}}_{\alpha,C}$ is an isomorphism if and only if α^r is a simple composition.

Next, we introduce the second main result of this subsection.

Proposition 5.5. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then there exists a unique $H_n(0)$ -module homomorphism $\tilde{\delta} : X_{\alpha} \to \mathbf{Y}_{\alpha}$ such that $\delta = \tilde{\delta} \circ \Gamma$.

Proof. As in the proof of Proposition 5.3, it suffices to show that $\ker(\delta) \supseteq \ker(\Gamma)$. Note that

$$\ker(\delta) = \mathbb{C}\{\mathscr{T} \in \operatorname{SIT}(\alpha) \mid \mathsf{w}_{\mathsf{r}}(\mathscr{T}) \notin \mathcal{K}_{\alpha}\} \text{ and} \\ \ker(\Gamma) = \mathbb{C}\{\mathscr{T} \in \operatorname{SIT}(\alpha) \mid \mathscr{T} \notin \operatorname{SET}(\alpha)\}$$

(for the definition of \mathcal{K}_{α} , see (5.4)). Note that $\mathcal{K}_{\alpha} \cdot w_0$ is an equivalence class under \simeq_M and it contains $w_r(\mathscr{T}_{\alpha})w_0$. Combining this with Lemma 4.9, we have

$$\mathcal{K}_{\alpha} \subseteq \{ \mathsf{w}_{\mathsf{r}}(T) \mid T \in \operatorname{SET}(\alpha) \}.$$

Therefore, if $\mathscr{T} \in \operatorname{SIT}(\alpha) \setminus \operatorname{SET}(\alpha)$, then $\mathsf{w}_{\mathsf{r}}(\mathscr{T}) \notin \ker(\delta)$, as required. \Box

5.3. Weak Bruhat interval module structure for \mathbf{Y}_{α} . Unless otherwise specified, in this subsection, α refers to a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$. The purpose of this subsection is to prove that \mathcal{K}_{α} consists of the reading words of SE-decreasing standard extended tableaux and that \mathcal{K}_{α} forms a left weak Bruhat interval. Consequently, \mathbf{Y}_{α} is endowed with the structure of a weak Bruhat interval module. **Definition 5.6.** Let α be a composition of n. We say that a filling T of $cd(\alpha)$ is southeast decreasing (simply, SE-decreasing) if for all $1 \le i < j \le \ell(\alpha)$ with $\alpha_i > \alpha_j$,

$$T(i, k+1) < T(j, k)$$
 for all $1 \le k \le \alpha_j$.

Lemma 5.7. Let α be a composition of n and $\mathscr{T} \in SIT(\alpha)$. If $sh(\mathsf{P}(\mathsf{w}_{\mathsf{r}}(\mathscr{T})w_0)) = \alpha$, then \mathscr{T} is SE-decreasing.

Proof. For simplicity, let $\sigma := \mathsf{w}_{\mathsf{r}}(\mathscr{T})\mathsf{w}_0, \lambda := \lambda(\alpha)$, and $l := \ell(\alpha)$. Assume, for the sake of contradiction, that there exist $1 \le i < j \le l$ with $\alpha_i > \alpha_j$ such that $\mathscr{T}(i, t+1) > \mathscr{T}(j, t)$ for some $1 \le t \le \alpha_j$. Let

$$R := \{ 1 \le r \le l \mid \alpha_r \ge \alpha_i \}, \quad k := |R|,$$

and let $u = (u^{(r)})_{r \in R}$ denote the k-tuple of increasing subsequences of σ defined as

$$u^{(r)} := \begin{cases} \mathscr{T}(j,1) \ \mathscr{T}(j,2) \ \dots \ \mathscr{T}(j,t) \ \mathscr{T}(i,t+1) \ \dots \ \mathscr{T}(i,\alpha_i) & \text{if } r = i, \\ \mathscr{T}(r,1) \ \mathscr{T}(r,2) \ \dots \ \mathscr{T}(r,\alpha_r) & \text{otherwise.} \end{cases}$$

Since $j \notin R$, the length of **u** is $\sum_{r \in R} \alpha_r$ (= $\sum_{1 \leq r \leq k} \lambda_r$). Combining this with $\operatorname{sh}(\mathsf{P}(\sigma)) = \lambda$, we derive $u \in \operatorname{Inc}_k(\sigma)$ from Theorem 3.9. On the other hand, by Theorem 3.15 and Lemma 4.5, the assumption $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = \alpha$ implies that

$$\underline{\mathsf{IES}}_k(\sigma) = \{\mathscr{T}(r,1) \mid r \in R\}.$$

Since $\mathscr{T}(i,1) < \mathscr{T}(j,1)$ with $i \in R$ and $j \notin R$, it follows that $\underline{\mathsf{IES}}_k(\sigma) <_{\text{set}} \mathsf{IES}(u)$, leading to a contradiction.

Let

$$\operatorname{SET}_{\operatorname{sed}}(\alpha) := \{ T \in \operatorname{SET}(\alpha) \mid T \text{ is SE-decreasing} \}.$$

The following theorem presents the first main result of this subsection.

Theorem 5.8. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \ge 0$, then

$$\mathcal{K}_{\alpha} = \{ \mathsf{w}_{\mathsf{r}}(T) \mid T \in \operatorname{SET}_{\operatorname{sed}}(\alpha) \}.$$

Proof. By Lemma 4.9 and Lemma 5.7, we have $\mathcal{K}_{\alpha} \subseteq \{\mathsf{w}_{\mathsf{r}}(T) \mid T \in \operatorname{SET}_{\operatorname{sed}}(\alpha)\}$. Thus, it remains to prove that $\mathcal{K}_{\alpha} \supseteq \{\mathsf{w}_{\mathsf{r}}(T) \mid T \in \operatorname{SET}_{\operatorname{sed}}(\alpha)\}$. By (5.3), this reduces to showing that $\operatorname{sh}(\hat{\mathsf{P}}(\mathsf{w}_{\mathsf{r}}(T)w_0) = \alpha$ for all $T \in \operatorname{SET}_{\operatorname{sed}}(\alpha)$.

Let $T \in \text{SET}_{\text{sed}}(\alpha)$. For simplicity, let $\sigma := \mathsf{w}_{\mathsf{r}}(T)w_0$, $\lambda := \lambda(\alpha)$, and $l := \ell(\lambda)$. Note that every increasing subsequence of σ contains at most one element in each column of T. More precisely, every increasing subsequence of length k can be expressed as

$$T(i_1, j_1)T(i_2, j_2)\dots T(i_k, j_k),$$
 (5.7)

where $l \ge i_1 \ge i_2 \ge \cdots \ge i_k \ge 1$ and $1 \le j_1 < j_2 < \cdots < j_k \le l$. From this, it follows that

$$\mathfrak{i}_k(\sigma) = \lambda_1 + \lambda_2 + \dots + \lambda_k$$

for every $1 \le k \le l$. Therefore, by Theorem 3.9, $\mathsf{P}(\sigma)$ has shape λ . Combining this with Lemma 4.5 and Theorem 3.15, we deduce that $\operatorname{sh}(\hat{\mathsf{P}}(\sigma)) = \alpha$ if and only if

$$\underline{\mathsf{IES}}_k(\sigma) \subseteq \{T(i,1) \mid 1 \le i \le l \text{ and } \alpha_i \ge \lambda_k\} \text{ for all } 1 \le k \le l.$$
(5.8)

Let us fix $1 \leq k \leq l$ and $u = (u^{(i)})_{1 \leq i \leq k} \in \text{Inc}_k(\sigma)$. To prove (5.8), we have only to see that

$$\mathsf{IES}(u) \subseteq \{T(i,1) \mid 1 \le i \le l \text{ with } \alpha_i \ge \lambda_k\}.$$

Case 1: $\lambda_1 = \lambda_k$. Let $1 \le i \le k$. Since $i_1(\sigma) = \lambda_1$, the length of $u^{(i)}$ is λ_1 . By (5.7), we can write $u^{(i)}$ as

$$u^{(i)} = T(i_1, 1)T(i_2, 2)\dots T(i_{\lambda_1}, \lambda_1),$$

where $l \geq i_1 \geq i_2 \geq \cdots \geq i_{\lambda_1} \geq 1$. It suffices to show that $\alpha_{i_1} = \lambda_1$. Suppose, for the sake of contradiction, that $\alpha_{i_1} \neq \lambda_1$, i.e., $\alpha_{i_1} < \lambda_1$. Since $\alpha_{i_{\lambda_1}} = \lambda_1$, we can choose the smallest $1 \leq t \leq \lambda_1$ such that $\alpha_{i_t} = \lambda_1$. Then, $1 < t \leq l$. By the choice of t, $i_{t-1} > i_t$ and $\alpha_{i_{t-1}} < \alpha_{i_t} = \lambda_1$. Since T is SE-decreasing, this implies that $T(i_{t-1}, t-1) > T(i_t, t)$, which contradicts that $u^{(i)}$ is an increasing subsequence.

Case 2: $\lambda_1 > \lambda_k$. Let

$$X := \{ T(i,j) \mid 1 \le i \le l \text{ with } \alpha_i > \lambda_k \text{ and } \lambda_k < j \le \alpha_i \}.$$

Let T' be the subfilling obtained from T by removing all elements in X and σ' the word obtained from σ by removing all elements in X. Note that T' is an extended tableau of shape $\beta = (\beta_1, \ldots, \beta_l)$, where

$$\beta_i = \begin{cases} \lambda_k & \text{if } \alpha_i > \lambda_k, \\ \alpha_i & \text{otherwise} \end{cases}$$

(for the definition of extended tableaux, see [2, Definition 6.16]). Define $u' = (u'^{(i)})_{1 \le i \le k}$ as the k-tuple of increasing subsequences of σ' , where each $u'^{(i)}$ is obtained from $u^{(i)}$ by removing all entries in X. In view of (5.7) and Theorem 3.9, we have $\mathsf{IES}(u) \cap X = \emptyset$, and therefore $\mathsf{IES}(u') = \mathsf{IES}(u)$. Thus, to prove the assertion, it suffices to show that *Case 1* can be applied to σ' and u'. Observe that

- β is a composition that is a shuffle of a partition and $(1^{k'})$ for some $k' \ge 0$,
- T' is SE-decreasing, and
- σ' is the word obtained by reversing the entries of $w_r(T')$.

Therefore, we have only to show that $u' \in \text{Inc}_k(\sigma')$. By the construction, it immediately follows that \mathbf{u}' is a k-increasing subsequence of σ' and

$$\ell(\mathbf{u}') \geq \lambda_1 + \lambda_2 + \dots + \lambda_k - |X| = k\lambda_k.$$

Also, since the shape of $\mathsf{P}(\sigma')$ is $(\underbrace{\lambda_k, \ldots, \lambda_k}_{k \text{ times}}, \lambda_{k+1}, \ldots, \lambda_l)$, Theorem 3.9 says that $\ell(\mathbf{u}') \leq k$

k times

 $k\lambda_k$, as required.



FIGURE 5.1

Next, we show that \mathcal{K}_{α} forms a left weak Bruhat interval. We begin by introducing sequences of cells in $cd(\alpha)$. These sequences are constructed using the following procedure.

Procedure 5.9. Let $\alpha = (\alpha_1, \ldots, \alpha_l)$ be a composition of n that is a shuffle of a partition and (1^k) for some $k \ge 0$. Let i be the smallest index $1 \le t \le l$ such that α_t is the largest part of α . Set the sequences $c^{(1)}$ and $c^{(2)}$ as $c^{(1)} = ((i, \alpha_i))$ and $c^{(2)} = \emptyset$.

- **P1.** If $\alpha_i \geq 3$, proceed to **P2**. Otherwise, proceed to **P3**.
- **P2.** If $\alpha_j < \alpha_i 1$ for all $i < j \leq l$, terminate the procedure. Otherwise, choose the smallest $i < j \leq l$ such that $\alpha_j \geq \alpha_i 1$; in this case, $\alpha_t = 1$ for all i < t < j. Update $c^{(1)}$ to be the concatenation of $((j, \alpha_j))$ and $c^{(1)}$, and update i to j. Then, proceed to **P1**.
- **P3.** Choose the largest $i \leq j \leq l$ such that $\alpha_t = \alpha_i$ for all $i \leq t \leq j$. Update $c^{(1)}$ to be the concatenation of $((j, \alpha_j), \ldots, (i+2, \alpha_{i+2}), (i+1, \alpha_{i+1}))$ and $c^{(1)}$. Let $\beta := (\alpha_{j+1}, \ldots, \alpha_l)$.
 - **P3a.** If $\beta = \emptyset$, terminate the procedure. Otherwise, proceed to **P3b**.
 - **P3b.** Choose the smallest $1 \leq t \leq \ell(\beta)$ such that $(\beta_t, \ldots, \beta_{\ell(\beta)})$ forms a partition. Update $c^{(2)}$ to be the concatenation of $c^{(2)}$ and $((\ell(\beta), \beta_{\ell(\beta)}), \ldots, (t, \beta_t))$ and update β to be $(\beta_1, \ldots, \beta_{t-1}, \beta_t - 1, \ldots, \beta_{\ell(\beta)} - 1)$. Then, proceed to **P3a**.

We denote the two resulting sequences $c^{(1)}$ and $c^{(2)}$ obtained by applying Procedure 5.9 to α as $seq_1(\alpha)$ and $seq_2(\alpha)$, respectively. For instance, if $\alpha = (1, 4, 1, 4, 3, 2, 2, 2, 1, 2, 1, 1, 2, 1)$, we have

$$\begin{aligned} & \mathsf{seq}_1(\alpha) = ((8,2), (7,2), (6,2), (5,3), (4,4), (2,4)) & \text{and} \\ & \mathsf{seq}_2(\alpha) = ((14,1), (13,2), (13,1), (12,1), (11,1), (10,2), (10,1), (9,1)) \end{aligned}$$

See FIGURE 5.1, where the cells in $seq_1(\alpha)$ are highlighted in red and the cells in $seq_2(\alpha)$ are highlighted in blue.

Let r_0 be the positive integer such that (r_0, α_{r_0}) is the initial cell of the sequence $\operatorname{seq}_1(\alpha)$. By the assumption on α and the construction of $\operatorname{seq}_1(\alpha)$ and $\operatorname{seq}_2(\alpha)$, it follows that for $(i,j) \in \operatorname{cd}(\alpha),$

$$(i, j) \in seq_1(\alpha)$$
 if and only if $1 \le i \le r_0$ with $\alpha_i \ge \alpha_{r_0}$ and $j = \alpha_i$, (5.9)

 $(i, j) \in \operatorname{seq}_2(\alpha)$ if and only if $\alpha_{r_0} = 2$ and $i > r_0$. (5.10)

Now, define D_{α} as the diagram obtained from $cd(\alpha)$ by removing all cells in the sequences $seq_1(\alpha)$ and $seq_2(\alpha)$.

Lemma 5.10. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then D_{α} is a composition diagram. Moreover, letting $\tilde{\alpha} := \operatorname{sh}(D_{\alpha})$, we have

- (1) $\widetilde{\alpha}$ is a shuffle of a partition and $(1^{k'})$ for some $k' \geq 0$, and
- (2) for all $1 \leq i < j \leq \ell(\widetilde{\alpha})$, if $\alpha_i > \alpha_j$, then $\widetilde{\alpha}_i > \widetilde{\alpha}_j$.

Proof. Let $l := \ell(\alpha)$ and (r_0, α_{r_0}) be the initial cell of the sequence $\operatorname{seq}_1(\alpha)$. For each $1 \leq i \leq l$, let β_i be the number of cells in the *i*th row of D_{α} . Then,

$$\beta_i = \begin{cases} \alpha_i - 1 & \text{if } (i, \alpha_i) \in \mathsf{seq}_1(\alpha), \\ 0 & \text{if } (i, \alpha_i) \in \mathsf{seq}_2(\alpha), \\ \alpha_i & \text{otherwise.} \end{cases}$$

We prove our assertion by considering the following three cases for α_{r_0} .

Case 1: $\alpha_{r_0} = 1$. In this case, by the construction of $\operatorname{seq}_1(\alpha)$, the largest part of α is 1. Therefore, $\alpha = (1^n)$ and $r_0 = n$. As a result, D_{α} is the empty diagram, implying that $\tilde{\alpha} = \emptyset$. Consequently, $\tilde{\alpha}$ trivially satisfies conditions (1) and (2).

Case 2: $\alpha_{r_0} = 2$. Combining (5.9) and (5.10) yields that for $1 \leq i \leq l$,

$$\beta_i = \begin{cases} \alpha_i - 1 & \text{if } 1 \leq i \leq r_0 \text{ and } \alpha_i \geq 2, \\ 1 & \text{if } 1 \leq i \leq r_0 \text{ and } \alpha_i = 1, \\ 0 & \text{if } r_0 < i \leq l. \end{cases}$$

This shows that D_{α} is a composition diagram and $\tilde{\alpha} = (\beta_1, \beta_2, \dots, \beta_{r_0})$. Furthermore, $\tilde{\alpha}$ satisfies conditions (1) and (2), as can be deduced from the assumption on α and the observation that $\{1 \leq i \leq r_0 \mid \alpha_i = 2\} = [t, r_0]$ for some $1 \leq t \leq r_0$.

Case 3: $\alpha_{r_0} \geq 3$. Combining (5.9) and (5.10) yields that for $1 \leq i \leq l$,

$$\beta_i = \begin{cases} \alpha_i - 1 & \text{if } 1 \le i \le r_0 \text{ and } \alpha_i \ge 2, \\ 1 & \text{if } 1 \le i \le r_0 \text{ and } \alpha_i = 1, \\ \alpha_i & \text{if } r_0 < i \le l. \end{cases}$$

This shows that D_{α} is a composition diagram and that $\tilde{\alpha} = (\beta_1, \beta_2, \dots, \beta_l)$. Furthermore, $\tilde{\alpha}$ satisfies conditions (1) and (2), as can be deduced from the assumption on α and the observation that $\alpha_i < \alpha_{r_0} - 1$ for all $r_0 < i \leq l$.

Recall that the shape of D_{α} is denoted by $\widetilde{\alpha}$ (see Lemma 5.10). We have $cd(\alpha) = cd(\widetilde{\alpha}) \sqcup \{C \mid C \in seq_i(\alpha) \text{ for some } i = 1, 2\}$. Now, define the filling τ'_{α} of $cd(\alpha)$ by

$$\tau'_{\alpha}(C) = \begin{cases} \tau'_{\widetilde{\alpha}}(C) & \text{if } C \in \mathsf{cd}(\widetilde{\alpha}), \\ n+1-k & \text{if } C \text{ is the } k\text{th entry of } \mathsf{seq}(\alpha), \end{cases}$$

where $seq(\alpha)$ is the sequence obtained by concatenating $seq_2(\alpha)$ and $seq_1(\alpha)$.

Lemma 5.11. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \geq 0$, then $\tau'_{\alpha} \in \text{SET}_{sed}(\alpha)$.

Proof. We will prove the assertion using mathematical induction on n. If n = 1, then the assertion holds trivially. Assume n > 1. If $\tilde{\alpha} = \emptyset$, then $\alpha = (1^n)$, and again, the assertion holds trivially. Now, suppose $\tilde{\alpha} \neq \emptyset$. Since $1 \leq |\tilde{\alpha}| < n$, by Lemma 5.10, the induction hypothesis ensures that $\tau'_{\tilde{\alpha}} \in \text{SET}_{\text{sed}}(\tilde{\alpha})$. From this containment one can derive that τ'_{α} is a standard extended tableau. It remains to show that τ'_{α} is SE-decreasing.

Let (r_0, α_{r_0}) be the initial cell of $\operatorname{seq}_1(\alpha)$. Since $\widetilde{\alpha} \neq \emptyset$, we have $\alpha_{r_0} \geq 2$ and $r_0 \leq \ell(\widetilde{\alpha})$. Let $1 \leq i < j \leq l$ with $\alpha_i > \alpha_j$ and $1 \leq k \leq \alpha_j$. We prove our assertion by considering the following three cases.

Case 1: $j > r_0$. In this case, it follows from the definition of τ'_{α} that $\tau'_{\alpha}(j,k) > \tau'_{\alpha}(i,k+1)$. Case 2: $j \leq r_0$ with $\alpha_j > 1$ and $k = \alpha_j$. In this case, we have $\alpha_j \geq \alpha_{r_0}$ by the assumption on α . This implies that $(i, \alpha_i), (j, \alpha_j) \in \mathsf{seq}_1(\alpha)$ by (5.9). Therefore, (j, α_j) appears before (i, α_i) in $\mathsf{seq}_1(\alpha)$, which implies that

$$\tau'_{\alpha}(j,\alpha_j) > \tau'_{\alpha}(i,\alpha_i) \ge \tau'_{\alpha}(i,\alpha_j+1).$$

Case 3: If neither of the above applies, by Lemma 5.10(2), we have $\tilde{\alpha}_i > \tilde{\alpha}_j$. Additionally, it is easy to see that $k \leq \tilde{\alpha}_j$, and thus, $k + 1 \leq \tilde{\alpha}_i$. Combining these observations with $\tau'_{\tilde{\alpha}} \in \text{SET}_{sed}(\tilde{\alpha})$, we derive that

$$\tau'_{\alpha}(j,k) = \tau'_{\widetilde{\alpha}}(j,k) > \tau'_{\widetilde{\alpha}}(i,k+1) = \tau'_{\alpha}(i,k+1).$$

We are now prepared to state the second main result of this subsection.

Theorem 5.12. If α is a composition of n obtained by shuffling a partition and (1^k) for some $k \ge 0$, then

$$\mathcal{K}_{\alpha} = [\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\tau'_{\alpha})]_{L}.$$

Before proceeding with the proof, let us collect the necessary notations. Define the bijection $\iota_{\alpha} : \operatorname{cd}(\alpha) \to [n]$ by

$$\iota_{\alpha}((i,j)) = \sum_{1 \le t \le i} \alpha_t - j + 1.$$

For a filling T of $cd(\alpha)$, define

 $InvCell(T) := \{ ((i_1, j_1), (i_2, j_2)) \in cd(\alpha) \times cd(\alpha) \mid i_1 < i_2 \text{ and } T(i_1, j_1) > T(j_2, j_2) \}.$

If $T \in \text{SET}(\alpha)$, then

$$\operatorname{Inv}_{L}(\mathsf{w}_{\mathsf{r}}(T)) = \{(\iota_{\alpha}(C_{1}), \iota_{\alpha}(C_{2})) \mid (C_{1}, C_{2}) \in \operatorname{InvCell}(T)\} \sqcup A_{\alpha},$$
(5.11)

where

$$A_{\alpha} = \{(\iota_{\alpha}((i,j)), \iota_{\alpha}((i,k))) \mid 1 \le i \le \ell(\alpha) \text{ and } 1 \le k < j \le \alpha_i\}.$$

Proof of Theorem 5.12. For simplicity, let $I := [\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\tau'_{\alpha})]_{L}$ and $l := \ell(\alpha)$.

We first show that $\mathcal{K}_{\alpha} \supseteq I$. Let $\sigma \in I$. Observe that $\mathsf{T}_{\alpha} \in \operatorname{SET}_{\operatorname{sed}}(\alpha)$, and $\tau'_{\alpha} \in \operatorname{SET}_{\operatorname{sed}}(\alpha)$ by Lemma 5.11. By Theorem 5.8, we obtain

$$\operatorname{sh}(\mathbf{\hat{P}}(\mathbf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha})w_{0})) = \alpha = \operatorname{sh}(\mathbf{\hat{P}}(\mathbf{w}_{\mathsf{r}}(\tau_{\alpha}')w_{0})),$$

and thus

$$\operatorname{sh}(\mathsf{P}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha})w_{0})) = \lambda(\alpha) = \operatorname{sh}(\mathsf{P}(\mathsf{w}_{\mathsf{r}}(\tau_{\alpha}')w_{0})).$$

On the other hand, since $w_r(\tau'_{\alpha})w_0 \preceq_L \sigma w_0 \preceq_L w_r(\mathsf{T}_{\alpha})w_0$, it follows from (4.5) that $\operatorname{sh}(\mathsf{P}(\sigma w_0)) = \lambda(\alpha)$. Therefore, by Lemma 4.6, we derive that $\operatorname{sh}(\mathsf{P}(\sigma w_0)) = \alpha$, and thus, $\sigma \in \mathcal{K}_{\alpha}$.

Next, we show that $\mathcal{K}_{\alpha} \subseteq I$. Let $T \in \text{SET}_{\text{sed}}(\alpha)$. Since it has been already shown that $\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}) \preceq_{L} \mathsf{w}_{\mathsf{r}}(T)$, it suffices to show that $\mathsf{w}_{\mathsf{r}}(T) \preceq_{L} \mathsf{w}_{\mathsf{r}}(\tau'_{\alpha})$, that is, $\text{Inv}_{L}(\mathsf{w}_{\mathsf{r}}(T)) \subseteq \text{Inv}_{L}(\mathsf{w}_{\mathsf{r}}(\tau'_{\alpha}))$. By (5.11), this is equivalent to

$$InvCell(T) \subseteq InvCell(\tau'_{\alpha}).$$
(5.12)

Before proving this inclusion, we partition $\operatorname{InvCell}(T)$ into three disjoint subsets. Let (r_0, α_{r_0}) be the cell in the first entry of $\operatorname{seq}_1(\alpha)$. Let $T^{(1)}$ be the subfilling of T consisting of all cells not in $\operatorname{seq}_1(\alpha)$ or $\operatorname{seq}_2(\alpha)$, and let $T^{(2)}$ be the subfilling of T consisting of the cells in $\operatorname{seq}_2(\alpha)$. For later use, we note that the standardizations of $T^{(1)}$ and $T^{(2)}$ are SE-decreasing standard extended tableaux. In particular, when T is equal to τ'_{α} , it is not difficult to see that the standardizations of $T^{(1)}$ and $T^{(2)}$ are τ'_{α} and τ'_{β} , respectively, where $\beta = (\alpha_{r_0+1}, \ldots, \alpha_l)$. We claim that

$$\operatorname{InvCell}(T) = \operatorname{InvCell}(T^{(1)}) \sqcup \operatorname{InvCell}(T^{(2)}) \sqcup X, \qquad (5.13)$$

where

$$X = \{ (C_1, C_2) \in \operatorname{InvCell}(T) \mid C_1 \in \operatorname{seq}_1(\alpha), C_2 \in \operatorname{cd}(\widetilde{\alpha}) \}.$$

The inclusion \supseteq is evident. To show the reverse inclusion \subseteq , we will verify that for $(i_1, j_1), (i_2, j_2) \in cd(\alpha)$ with $i_1 < i_2$, the following conditions are satisfied:

(i) if $(i_2, j_2) \in seq_1(\alpha)$, then $T(i_1, j_1) < T(i_2, j_2)$ and

(ii) if $(i_1, j_1) \notin \text{seq}_2(\alpha)$ and $(i_2, j_2) \in \text{seq}_2(\alpha)$, then $T(i_1, j_1) < T(i_2, j_2)$.

For (i), it suffices to consider the case where $\alpha_{i_1} > 1$. By the assumption on α , we see that $\alpha_{i_1} \ge \alpha_{i_2}$. Since $j_2 = \alpha_{i_2}$ and $T \in \text{SET}_{sed}(\alpha)$, it follows that

$$T(i_1, j_1) \le T(i_1, \alpha_{i_1}) < T(i_2, \alpha_{i_2}) = T(i_2, j_2).$$

S.-Y. LEE AND Y.-T. OH

Now, it remains to show (ii). From the condition $\sec_2(\alpha) \neq \emptyset$, we deduce that $\alpha_{r_0} = 2$ and $\alpha_{r_0+1} = 1$. Since T is SE-decreasing, it follows that $T(r_0, 2) < T(r_0 + 1, 1)$. Moreover, we have

$$T(i_1, j_1) \leq T(r_0, 2)$$
 and $T(r_0 + 1, 1) \leq T(i_2, j_2)$

where the first inequality follows from (i), while the second one follows from $T \in \text{SET}_{sed}(\alpha)$. Combining these three inequalities verifies (ii).

In what follows, we prove the inclusion given in (5.12) using mathematical induction on $n = |\alpha|$. For n = 0, 1, this inclusion trivially holds. Now, assume that n > 1 and the inclusion (5.12) holds for all compositions of size < n obtained by shuffling a partition and (1^k) for some $k \ge 0$. Let α be a composition of n obtained by shuffling a partition and (1^k) for some $k \ge 0$. By Lemma 5.10, $\tilde{\alpha}$ is a composition obtained by shuffling a partition and (1^k) for some $k \ge 0$. Since $|\tilde{\alpha}| < n$ and the standardization of $T^{(1)}$ is an element of SET_{sed} $(\tilde{\alpha})$, by the induction hypothesis, we have

$$\operatorname{InvCell}(T^{(1)}) \subseteq \operatorname{InvCell}(\tau'_{\widetilde{\alpha}}).$$
(5.14)

In a similar manner, viewing τ'_{β} as the subfilling of τ'_{α} consisting of the cells in $\operatorname{seq}_2(\alpha)$, we obtain

$$\operatorname{InvCell}(T^{(2)}) \subseteq \operatorname{InvCell}(\tau'_{\beta}).$$
(5.15)

On the other hand, applying (5.13) to the case where T is equal to τ'_{α} , it follows from the definition of τ'_{α} that

$$\operatorname{InvCell}(\tau'_{\alpha}) = \operatorname{InvCell}(\tau'_{\widetilde{\alpha}}) \sqcup \operatorname{InvCell}(\tau'_{\beta}) \sqcup Y,$$

where

$$Y = \{ ((i_1, j_1), (i_2, j_2)) \mid (i_1, j_1) \in \mathsf{seq}_1(\alpha), \ (i_2, j_2) \in \mathsf{cd}(\widetilde{\alpha}), \text{ and } i_1 < i_2 \}.$$

Finally, by combining (5.14) and (5.15) with the inclusion $X \subseteq Y$, we obtain the desired result.

Theorem 5.12 tells us that \mathbf{Y}_{α} is endowed with the structure of a weak Bruhat interval module:

$$\mathbf{Y}_{\alpha} \cong \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\tau_{\alpha}')).$$

Combining this with [21, Table 1] implies that the ϕ -twist $\phi[\mathbf{Y}_{\alpha}]$ of \mathbf{Y}_{α} also has a weak Bruhat interval module structure:

$$\Phi[\mathbf{Y}_{\alpha}] \cong \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha})^{w_0}, \mathsf{w}_{\mathsf{r}}(\tau_{\alpha}')^{w_0}).$$

Example 5.13. Let $\alpha = (3, 1, 2)$. Then

$$\tau'_{\alpha} = \begin{array}{c} 4 & 6 \\ 3 \\ 1 & 2 & 5 \end{array}$$



FIGURE 5.2. The $H_6(0)$ -action on $\mathcal{K}_{(2,3,1)}$

On the other hand, considering FIGURE 4.2, we have $\mathbf{Y}_{\alpha} = \mathbb{C}B_3/\mathbb{C}B_2$, and thus, we have $\mathbf{Y}_{\alpha} \cong \mathsf{B}(321465, 521364) = \mathsf{B}(\mathsf{w}_{\mathsf{r}}(\mathsf{T}_{\alpha}), \mathsf{w}_{\mathsf{r}}(\tau'_{\alpha})).$

Remark 5.14. For a general composition α , \mathbf{Y}_{α} does not necessarily have a weak Bruhat interval module structure. For instance, let $\alpha = (2, 3, 1)$. In this case, the $H_6(0)$ -action on \mathcal{K}_{α} is illustrated in FIGURE 5.2. We claim that there is no left weak Bruhat interval Iin \mathfrak{S}_6 such that $\mathbf{Y}_{\alpha} \cong \mathsf{B}(I)$. Suppose, for the sake of contradiction, that such an interval $I = [\sigma, \rho]_L$ exists. Let $f : \mathbf{Y}_{\alpha} \to \mathsf{B}(\sigma, \rho)$ be an $H_6(0)$ -module isomorphism. Note that there exists a unique $\gamma \in \mathcal{K}_{\alpha}$ such that $\mathrm{Des}_L(\gamma) \supseteq \{1, 3, 4\}$ and this unique element is 215436. Therefore, there exists a unique $\gamma' \in [\sigma, \rho]_L$ such that $\mathrm{Des}_L(\gamma') \supseteq \{1, 3, 4\}$ and $f(215436) = \gamma'$. Since 215436 is a generator for $\mathbf{Y}_{\alpha}, \gamma'$ must be σ . This contradicts our assumption that I is a left weak Bruhat itnerval.

6. Remarks concerned with future research

(1) As seen in Remark 5.14, for a general composition α , \mathcal{K}_{α} is not necessarily a left weak Bruhat interval. It would be nice to characterize when \mathcal{K}_{α} is a left weak Bruhat interval, more generally, when \mathbf{Y}_{α} has a weak Bruhat interval module structure.

(2) Let I be a left weak Bruhat interval in \mathfrak{S}_n . It was shown in [22, Section 6.2] that if I is a dual plactic-closed, then B(I) admits a distinguished filtration with respect to the Schur basis. A natural next step is to characterize the conditions under which the module B(I) admits a distinguished filtration with respect to $\{\mathscr{S}_\alpha \mid \alpha \models n\}$ or $\{\widehat{\mathscr{S}}_\alpha \mid \alpha \models n\}$ (see Appendix A).

(3) When I is a dual plactic-closed left weak Bruhat interval in \mathfrak{S}_n , let

$$0 = M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_l = \mathsf{B}(I)$$

be a distinguished filtration of B(I) with respect to the Schur basis, as constructed in the proof of [22, Theorem 6.7]. Since $ch([M_i/M_{i-1}])$ is a Schur function, it is natural to ask whether M_i/M_{i-1} admits a distinguished filtration with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ or $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$. The validity of this question has been checked for values of n up to 9 with the aid of the computer program SAGEMATH.

Appendix A. Examples of $H_n(0)$ -modules that have no distinguished filtrations with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ or $\{\widehat{\mathscr{S}}_{\alpha} \mid \alpha \models n\}$.

In this appendix, we present an example to show that when P is a regular Schur labeled poset with the underlying set [n], the associated $H_n(0)$ -module M_P , as defined in [22, Definition 2.8], does not necessarily admit a distinguished filtration with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models n\}$ or $\{\widehat{\mathscr{S}}_{\alpha} \mid \alpha \models n\}$.

Consider the $H_8(0)$ -modules $X_{(5,2,1)}$ and its ϕ -twist $\phi[X_{(5,2,1)}]$. In view of [22, Appendix], one sees that each of these modules is isomorphic to the $H_8(0)$ -module associated with a regular Schur labeled skew shape poset with underlying set [8].

Claim 1. $X_{(5,2,1)}$ has no distinguished filtrations with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models 8\}$. We begin by observing that

$$ch([X_{(5,2,1)}]) = s_{(5,2,1)} = \sum_{\substack{\alpha \models 8\\\lambda(\alpha) = (5,2,1)}} \mathscr{S}_{\alpha}$$

and $X_{(5,2,1)} \cong B(54321768, 87641523)$ as $H_8(0)$ -modules. Suppose on the contrary that B(54321768, 87641523) has a distinguished filtration

$$0 =: M_0 \subsetneq M_1 \subsetneq \cdots \subsetneq M_6 := \mathsf{B}(54321768, 87641523)$$

with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models 8\}$. We note that the sink of B(54321768, 87641523) is isomorphic to $\mathbf{F}_{(1,1,2,4)}$ and thus $[F_{(1,1,2,4)}]ch([M_i]) > 0$ for $1 \leq i \leq 6$. Combining this with the fact that

$$[F_{(1,1,2,4)}]s_{(5,2,1)} = 1$$
 and $[F_{(1,1,2,4)}]\mathscr{S}_{(1,2,5)} = 1$,

we see that $ch([M_1]) = \mathscr{S}_{(1,2,5)}$ and $\dim M_1 = 16$. Given a composition series of M_1

 $0 =: N_0 \subsetneq N_1 \subsetneq \cdots \subsetneq N_{16} := M_1,$

let $\{v_1, v_2, \ldots, v_{16}\}$ be a basis for M_1 such that such that $\{v_1, v_2, \ldots, v_j\}$ is a basis for N_j for $1 \leq j \leq 16$. Since $[F_{(2,2,4)}] \mathscr{S}_{(1,2,5)} = 1$, there exists a unique $1 \leq j \leq 16$ such that

$$N_j/N_{j-1} \cong \mathbf{F}_{(2,2,4)}.$$
 (A.1)

For each $\sigma \in [54321768, 87641523]_L$, observe that $\pi_{w_0(\{1,3,5,6,7\})} \cdot \sigma$ is one of 0, 87421635, or 87621435. This implies that

$$\pi_{w_0(\{1,3,5,6,7\})} v_j \in \mathbb{C}\{87421635, 87621435\}.$$

Let

$$\pi_{w_0(\{1,3,5,6,7\})}v_i = c\,87421635 + d\,87621435$$

By (A.1), it holds that $\pi_{w_0(\{1,3,5,6,7\})}v_j - v_j \in N_{j-1}$ and $\pi_{w_0(\{1,3,5,6,7\})}v_j \neq 0$. It follows that $N_j = \mathbb{C}\{v_1, v_2, \dots, v_{j-1}, c\,87421635 + d\,87621435\}.$

If c = 0 or d = 0, then $L := \mathsf{B}(87621534, 87641523)$ is a submodule of M_1 . However, this cannot occur since

 $[F_{(2,1,5)}]\mathscr{S}_{(1,2,5)} = 0$ and $[F_{(2,1,5)}]ch([L]) = 1.$

Next, suppose that $c, d \neq 0$. Since

$$\pi_2 \cdot (c\,87421635 + d\,87621435) = c\,87431625 + d\,87631425$$

and

$$\pi_3\pi_2 \cdot (c87421635 + d87621435) = c87431625,$$

it follows that $87631425 \in M_1$. Therefore, $L' := \mathsf{B}(87631425, 87641523)$ is a submodule of M_1 . However, this cannot occur since

$$[F_{(1,2,1,4)}]$$
ch $([L']) = 1$ and $[F_{(1,2,1,4)}]\mathscr{S}_{(1,2,5)} = 0.$

For a pictorial description of L and L', along with the $H_8(0)$ -action on B(54321768, 87641523), see FIGURE A.1. Consequently, B(54321768, 87641523)($\cong X_{(5,2,1)}$) does not admit a distinguished filtration with respect to $\{\mathscr{S}_{\alpha} \mid \alpha \models 8\}$.

Claim 2. $\phi[X_{(5,2,1)}]$ has no distinguished filtrations with respect to $\{\hat{\mathscr{S}}_{\alpha} \mid \alpha \models 8\}$.

This follows from *Claim 1* since ϕ induces an equivalence on the category of finitely generated $H_n(0)$ -modules and $\phi[\mathbf{F}_\beta] \cong \mathbf{F}_{\beta^r}$.





Acknowledgments. The first author was supported by NRF grant funded by Basic Science Research Program through NRF funded by the Ministry of Education (No. RS-2023-00271282), NRF grant funded by the Korea government(MSIT) (No. RS-2024-00342349), and the Sogang University Research Grant of 2024(No. 202412001.01). The second author was supported by NRF grant funded by the Korea government(MSIT) (No. RS-2024-00342349).

References

- E. E. Allen, J. Hallam, and S. Mason. Dual immaculate quasisymmetric functions expand positively into Young quasisymmetric Schur functions. J. Combin. Theory Ser. A, 157:70–108, 2018.
- [2] S. Assaf and D. Searles. Kohnert polynomials. *Experiment. Math.*, pages 1–27, 2019.
- [3] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki. A lift of the Schur and Hall-Littlewood bases to non-commutative symmetric functions. *Canad. J. Math.*, 66(3):525–565, 2014.
- [4] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki. Indecomposable modules for the dual immaculate basis of quasi-symmetric functions. *Proc. Amer. Math. Soc.*, 143(3):991–1000, 2015.
- [5] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki. Multiplicative structures of the immaculate basis of non-commutative symmetric functions. J. Combin. Theory Ser. A, 152:10–44, 2017.
- [6] N. Bergeron and H. Li. Algebraic structures on Grothendieck groups of a tower of algebras. J. Algebra, 321(8):2068–2084, 2009.
- [7] N. Bergeron, J. Sánchez-Ortega, and M. Zabrocki. The pieri rule for dual immaculate quasisymmetric functions. Annals of Combinatorics, 20:283–300, 2016.
- [8] C. Bessenrodt, K. Luoto, and S. van Willigenburg. Skew quasisymmetric Schur functions and noncommutative Schur functions. Adv. Math., 226(5):4492–4532, 2011.
- [9] A. Björner and F. Brenti. Combinatorics of Coxeter groups, volume 231 of Graduate Texts in Mathematics. Springer, New York, 2005.
- [10] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. Modules of the 0-Hecke algebra arising from standard permuted composition tableaux. J. Combin. Theory Ser. A, 179:105389, 34, 2021.
- [11] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. Homological properties of 0-Hecke modules for dual immaculate quasisymmetric functions. *Forum Math. Sigma*, 10:e91, 2022.
- [12] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. The projective cover of tableau-cyclic indecomposable H_n(0)-modules. Trans. Amer. Math. Soc., 375(11):7747–7782, 2022.
- [13] B. Deng and G. Yang. Representation type of 0-Hecke algebras. Sci. China Math., 54(3):411–420, 2011.
- [14] G. Duchamp, F. Hivert, and J.-Y. Thibon. Noncommutative symmetric functions. VI. Free quasisymmetric functions and related algebras. *Internat. J. Algebra Comput.*, 12(5):671–717, 2002.
- [15] G. Duchamp, D. Krob, B. Leclerc, and J.-Y. Thibon. Fonctions quasi-symétriques, fonctions symétriques non commutatives et algèbres de Hecke à q = 0. C. R. Acad. Sci. Paris Sér. I Math., 322(2):107–112, 1996.
- [16] W. Fulton. Young tableaux, volume 35 of London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [17] I. M. Gessel. Multipartite P-partitions and inner products of skew Schur functions. In Combinatorics and algebra (Boulder, Colo., 1983), volume 34 of Contemp. Math., pages 289–317. Amer. Math. Soc., Providence, RI, 1984.
- [18] C. Greene. An extension of Schensted's theorem. Adv. Math., 14:254–265, 1974.

- [19] J. Haglund, K. Luoto, S. Mason, and S. van Willigenburg. Quasisymmetric Schur functions. J. Combin. Theory Ser. A, 118(2):463–490, 2011.
- [20] J. Haglund, K. Luoto, S. Mason, and S. van Willigenburg. Refinements of the Littlewood–Richardson rule. Trans. Amer. Math. Soc., 363(3):1665–1686, 2011.
- [21] W.-S. Jung, Y.-H. Kim, S.-Y. Lee, and Y.-T. Oh. Weak Bruhat interval modules of the 0-Hecke algebra. Math. Z., 301(4):3755–3786, 2022.
- [22] Y.-H. Kim, S.-Y. Lee, and Y.-T. Oh. Regular Schur labeled skew shape posets and their 0-hecke modules. Forum Math. Sigma, 12:e110, 2024.
- [23] S. König. The decomposition of 0-Hecke modules associated to quasisymmetric Schur functions. Algebr. Comb., 2(5):735–751, 2019.
- [24] K. Luoto, S. Mykytiuk, and S. van Willigenburg. An introduction to quasisymmetric Schur functions. SpringerBriefs in Mathematics. Springer, New York, 2013.
- [25] C. Marcum and E. Niese. Positive expansions of extended Schur functions in the Young quasisymmetric Schur basis. *Involve*, Vol. 17(2):217–232, 2024.
- [26] S. Mason. A decomposition of Schur functions and an analogue of the Robinson-Schensted-Knuth algorithm. Sém. Lothar. Combin., 57:Art. B57e, 24, 2006/08.
- [27] A. Melnikov. On orbital variety closures in sl_n . I. Induced Duflo order. J. Algebra, 271(1):179–233, 2004.
- [28] P. Norton. 0-Hecke algebras. J. Austral. Math. Soc. Ser. A, 27(3):337–357, 1979.
- [29] B. E. Sagan. The symmetric group representations, combinatorial algorithms, and symmetric functions. Wadsworth & Brooks / Cole mathematics series. Wadsworth, 1991.
- [30] C. Schensted. Longest increasing and decreasing subsequences. Canad. J. Math., 13:179–191, 1961.
- [31] D. Searles. Indecomposable 0-Hecke modules for extended Schur functions. Proc. Amer. Math. Soc., 148(5):1933–1943, 2020.
- [32] R. Stanley. Enumerative combinatorics. Vol. 2, volume 62 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999.
- [33] M. Taskin. Properties of four partial orders on standard Young tableaux. ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)–University of Minnesota.
- [34] V. Tewari and S. van Willigenburg. Modules of the 0-Hecke algebra and quasisymmetric Schur functions. Adv. Math., 285:1025–1065, 2015.

(S.-Y. Lee) Department of Mathematics, Sogang University, Seoul 04107, Republic of Korea

Email address: sylee0814@sogang.ac.kr

(Y.-T. Oh) DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, REPUBLIC OF KOREA

Email address: ytoh@sogang.ac.kr