NON UNIFORM EXPANSION AND ADDITIVE NOISE IMPLY RANDOM HORSESHOES

JEROEN S.W. LAMB^{1,2,3}, GIUSEPPE TENAGLIA*1, AND DMITRY TURAEV¹

ABSTRACT. We propose a notion of random horseshoe and prove density of random horseshoes for non uniformly expanding random dynamical systems with additive noise

1. INTRODUCTION

In the study of time-evolving systems, sensitivity to initial conditions is the hallmark of chaotic behaviour and is often associated with the positivity of the top Lyapunov exponent with respect to an ergodic measure. Indeed, the top Lyapunov exponent is an ergodic average that measures the exponential rate of expansion along the most expanded direction. If it is positive, then we observe that typically nearby orbits eventually diverge.

It is natural to ask whether more refined topological properties can be deduced from the positivity of the top Lyapunov exponent. A celebrated result by Katok [11] establishes that $C^{1+\beta}$ deterministic diffeomorphisms preserving an ergodic measure μ with all non zero Lyapunov exponents have dense Smale horseshoes in the support of μ , generated by transverse intersection of stable and unstable manifolds of hyperbolic periodic points. Thereafter, Katok's result was extended to infinite dimensional settings [15, 16, 13, 17].

In the random context, establishing a correspondence between positivity of the top Lyapunov exponent and hyperbolic horseshoes presents non-trivial difficulties: an immediate challenge is that systems with noise typically do not have periodic orbits, which are an essential ingredient in the construction of Smale horseshoes.

In a previous paper [12], we constructed a random version of the Smale horseshoe for a special classes of so-called "predominantly expanding" RDSs, which have positive Lyapunov exponent.

In this paper, we generalize this result (Theorem 2.2) by showing that for $C^{1+\beta}$ maps with non-degenerated critical points, subject to diffusive noise, if there exists an ergodic stationary measure with all Lyapunov exponents positive, then random horseshees are dense in the support of the measure.

The main novelty of this paper is that we are able to build the same object as in [12] under essentially no assumptions. Indeed, the random horseshoe construction in [12] relies crucially on the following key properties of predominantly expanding RDSs:

 (i) the existence of a large expanding region, which provides control the probabilistic evolution of balls of large enough size;;

²⁰²⁰ Mathematics Subject Classification. 37H20, 37B10, 37D25, 60J05.

Key words and phrases. Chaotic Behaviour, Positive Lyapunov exponents, Symbolic dynamics, Random dynamical systems.

(ii) the noise strength being significantly larger than the size of the contracting region, which implies large deviations estimates for the Lyapunov exponent.

These two properties are used in [12] to establish that any ball is eventually onto and to control the tail of the statistics of this phenomenon.

As generically balls are not eventually onto as in [12], instead here we rely on a modification of Young times, the concept which was introduced in [3] to build a random Young tower for predominantly expanding RDSs. In [3], conditions (i) - (ii) were key to control Young times.

In this paper, we control the statistics of the modified Young times under the sole assumptions of diffusive additive noise and positivity of all Lyapunov exponents with respect to an ergodic measure. In particular

- We leverage ergodicity to control the evolution of large balls;
- We exploit the Nagaev method for Markov chains [18] to prove large deviations for the Lyapunov exponents in the presence of diffusive additive noise.

Another novelty is that we infer the existence of random horseshoes directly from the existence of Young times, leveraging the iid nature of the noise.

To put our result in context, prior to [12], horseshoe constructions for random systems had been established either in the context of non-hyperbolic horseshoes [8, 9, 10], or, in the uniformly hyperbolic case, where horseshoes were built for uniformly hyperbolic RDSs either mixing on fibers [7], or with an equicontinuous or quasi-periodic base [6]). This paper substantially generalizes and strengthens these results in the multi-dimensional setting.

This paper is organized as follows: In Section 2, we recall the basic setup of random dynamical systems, state our hypotheses and our main results. In Section 3 we establish large deviations results for the Lyapunov exponent and for recurrences to the critical set. In Section 4 we study the annealed properties of orbits of large balls. In Section 5, after a review of the main properties of Young times, we combine the results of Sections 3 and 4 to obtain the desired control on the statistics of Young times. In Section 6 we deduce, from the properties of Young times, the density of the random horseshoes.

2. Statement of the result

2.1. Hypotheses and main result. Let X a compact n-dimensional subset of \mathbb{R}^n , and consider a map $f \in \mathcal{C}^0(X, X)$, that is $C^{1+\beta}$ everywhere except in the critical set \mathcal{C} , i.e. the set of points where the derivative of f either does not exists or fails to be invertible. Furthermore, given the probability space $([-\sigma, \sigma]^n, \Sigma, \nu^{\otimes n})$, with $\nu = \frac{\operatorname{Leb}_{|[-\sigma,\sigma]}}{2\sigma}$, consider the product space $\Omega := ([-\sigma, \sigma]^n)^{\otimes \mathbb{N}}$, equipped with the product measure $\mathbb{P} = (\nu^{\otimes n})^{\otimes \mathbb{N}}$ and the product σ -algebra \mathcal{F} . Let

$$\theta: \Omega \to \Omega, \qquad \theta(\{\omega_i\}_{i \ge 0}) = \{\omega_{i+1}\}_{i \ge 0}$$

be the shift map acting on Ω . Then, $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ is an ergodic dynamical system.

We consider skew-product systems of the form

(2.1)
$$\Theta: \Omega \times X \to \Omega \times X, \qquad \Theta(\omega, x) := (\theta(\omega), f_{\omega}(x)),$$

where $\{f_{\omega}\}_{\omega\in\Omega}$ is a family of piecewise $C^{1+\alpha}$ endomorphisms of the form

(2.2)
$$f_{\omega}(x) = f(x) + \omega_0$$

with $f \in C^{1+\beta}(X \setminus \mathcal{C}, X)$. Our first assumption is about the structure of the critical set \mathcal{C} .

 $\mathbf{2}$

Hypothesis (H1). The critical set C is contained in an n-1 dimensional sub-manifold of X, which has at most a finite number of connected components and admits C^1 parametrization with bounded Jacobian on such components. Furthermore, the normal direction $v: C \to \mathbb{S}^{n-1}$ is of class C^1 . Furthermore, there are constants B > 1 and $\beta > 0$ such that

(1) for all $x \in X$ and $v \in \mathbb{R}^n$

(2.3)
$$\frac{1}{B}dist(x,\mathcal{C})^{\beta} \leq \frac{||df(x) \cdot v||}{||v||} \leq Bdist(x,\mathcal{C})^{-\beta}$$

(2) for all $x, y \in X \setminus C$ with $|x - y| \le \frac{dist(x,C)}{2}$

(2.4)
$$\left|\log ||df(x)^{-1}|| - \log ||df^{-1}(y)||\right| \le \frac{B}{dist(x, \mathcal{C})^{\beta}}|x - y|$$

and

(2.5)
$$\left|\log\left|\det(df(x))\right| - \log\left|\det(df(y))\right|\right| \le \frac{B}{dist(x,\mathcal{C})^{\beta}}|x-y|$$

The first part of Hypothesis (H1) is used in the large deviations estimates to compute integrals of the form $\int_{B_{\delta}(\mathcal{C})} h(\operatorname{dist}(x,\mathcal{C})) dx$, with $h \in L^1(dx)$. Indeed, note that

$$B_{\delta}(x) = \bigcup_{x \in \mathcal{C}} \operatorname{graph} \left\{ s \in [-\delta, \delta] \mapsto x + v(x)s \right\}.$$

This follows from the fact that the minimal distance between a point and C is orthogonal to C. Let $t \in [0,1]^{n-1} \mapsto g(t)$ be a parametrization for C, then

(2.6)
$$\int_{B_{\delta}(\mathcal{C})} h(\operatorname{dist}(x,\mathcal{C})) dx = \int_{t \in [0,1]^{n-1}} \int_{s \in [-\delta,\delta]} h(|sv(g(t)|)J(g(t))dtds \lesssim \int_{s \in [-\delta,\delta]} h(s)ds,$$

since the Jacobian is bounded. The second part of (H1) is standard in the theory of hyperbolic times. Given the family $\{f_{\omega}\}_{\omega\in\Omega}$ of circle endomorphisms in (2.2), the random composition f_{ω}^{n}

(2.7)
$$f_{\omega}^{n}(x) = f_{\theta^{n-1}(\omega)}(x) \circ \cdots \circ f_{\omega}(x),$$

is a Markov process with associated transition kernel $\{\mathcal{P}_x(dy) := \mathbb{P}(f_\omega(x) \in dy)\}_{x \in X}$. We recall that μ is a stationary measure for this Markov process if

$$\mu(A) = \int_X \mathcal{P}_x(A) d\mu(x),$$

for all Borel sets A. Furthermore, μ is ergodic if and only if the measure $\mathbb{P} \otimes \mu$ is ergodic for the skew product Θ introduced in (2.1). If μ is ergodic then, by Birkhoff Ergodic Theorem

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=0}^{n-1} T(f^i_{\omega}(x)) = \int_X T(x) d\mu(x) \qquad \mathbb{P} \otimes \mu \text{ a.s.}$$

for all $T \in L^1(\mu)$. For a more detailed background on Markov processes, see e.g. [19, 5]. Given $(\omega, x) \in \Omega \times X$, we can define the following quantity

$$\lambda(\omega, x) := \limsup_{n \to \infty} \frac{1}{n} \log \left| \left| \left(df(f_{\omega}^{i}(x))^{-1} \right| \right|.$$

If μ is an invariant measure for f_{ω}^n , then it follows from the *BV* estimates for μ in [14], Hypothesis (H1) and the parametrization in (2.6) that

$$\int_{B_{\delta}(\mathcal{C})} \log ||df(x)^{-1}|| dx \approx \int_{B_{\delta}(\mathcal{C})} \log \operatorname{dist}(x, \mathcal{C}) dx \lessapprox \delta \log(\delta)$$

i.e. $\log ||df(x)^{-1}|| \in L^1(\mu)$. For an ergodic process, this means that $\lambda(\omega, x)$ is almost surely constant $\mathbb{P} \otimes \mu$ and equal to

$$\lambda := \int_X \log ||df(x)^{-1}||d\mu(x).$$

Note that in dimension one the condition $\lambda < 0$ is equivalent to the positivity of the Lyapunov exponent. In dimension n, the same condition implies that for $\mathbb{P} \otimes \mu$ almost (ω, x) all Lyapunov exponents are positive, i.e. f_{ω}^{n} is uniformly expanding. In this paper, we assume there exists at least one ergodic component for the process f_{ω}^{n} with $\lambda < 0$, as stated in the following assumption.

Hypothesis (H2). The Markov process associated to f_{ω}^n admits an invariant ergodic component $Y \subset X$, so that the process restricted to Y admits unique ergodic measure μ satisfying

$$\lambda = \int_Y \log ||df(x)^{-1}||d\mu(x) < 0$$

Now we give the definition of random horseshoe we will use throughout the paper. This is a slightly different version than the one in [12].

Definition 2.1. Let I_0, I_1 be two balls such that $I_0 \cap I_1 = \emptyset$. Let $\kappa > 1$. Then the pair (I_0, I_1) is a κ -horseshoe if there exists a sequence of variables $\{n_k(\omega)\}_{k\geq 0}$ satisfying almost surely

(2.8)
$$\limsup_{k \to \infty} \frac{n_k(\omega)}{k} = \mathbb{E}[n_0],$$

(2.9)
$$f_{\theta^{n_k(\omega)}(\omega)}^{n_{k+1}(\omega)-n_k(\omega)}(I_i) \supseteq I_0 \cup I_1 \qquad \forall i \in \{0,1\}.$$

Moreover there exists balls $J(k, \omega)_{i,j} \subset I_i$ such that

(2.10)
$$f_{\theta^{n_k(\omega)}(\omega)}^{n_{k+1}(\omega)-n_k(\omega)}(J(k,\omega)_{i,j}) = I_j$$

and

(2.11)
$$\left\| \left(df_{\theta^{n_k+1}(\omega)-n_k(\omega)}^{n_{k+1}(\omega)-n_k(\omega)} |_{J(k,\omega)_{i,j}} \right)^{-1} \right\| < \kappa^{-1} \qquad \forall k \ge 0$$

With this definition, we may consider the set $P(\omega)$, which consists of all points x such that $f_{\omega}^{n_k(\omega)}(x) \in I_0 \cup I_1$, for all $k \ge 0$, i.e.

(2.12)
$$P(\omega) := \{ x \in X \colon f_{\omega}^{n_k(\omega)}(x) \in I_0 \cup I_1 \, \forall k \ge 0 \}.$$

Then, the set $P(\omega)$ is hyperbolic and in one-to-one correspondence with the sequences of zeros and ones. Note that Definition 2.1 is similar to the definition of horseshoe in [8], with the additional hyperbolicity requirement (2.11). Our main result is the following

Theorem 2.2. Let f_{ω}^n the random composition in (2.7) satisfying Hypotheses (H1)-(H2). Then, there exists $\kappa > 0$ such that the set of κ -horseshoes is dense, which means that for every open set A there are balls $I_1, I_2 \Subset A$ such that (I_1, I_2) is a κ -horseshoe.

3. Large Deviation estimates

Given $\delta > 0$ and $\omega \in \Omega$, we define the δ -truncated distance from x to the critical set C as

(3.1)
$$\operatorname{dist}_{\delta}(x, \mathcal{C}) := \begin{cases} 1 & \text{if } \operatorname{dist}(x, \mathcal{C}) > \delta, \\ \operatorname{dist}(x, \mathcal{C}) & \text{otherwise.} \end{cases}$$

Consider the following Birkhoff averages

$$S_n(\omega, x) := \frac{1}{n} \sum_{i=0}^n \log \left| \left| df^{-1}(f^i_\omega(x)) \right| \right|$$
$$Z_n(\omega, x) := \frac{1}{n} \sum_{i=0}^n -\log \operatorname{dist}_{\delta}(f^i_\omega(x), \mathcal{C}).$$

By Birkhoff ergodic theorem

$$S_n(\omega, x) \to \lambda, \qquad Z_n(\omega, x) \to \int_X \log \operatorname{dist}_{\delta}(x, \mathcal{C}) d\mu \approx \delta \log(\delta) \qquad \mathbb{P} \otimes \mu \ a.s.$$

We wish to estimate large deviations result for the convergence of the above limits. In particular, we aim to prove the following Proposition.

Proposition 3.1. For every $\varepsilon > 0$ small enough, there exists C > 1, $\gamma < 1$ and a family of sets $\{A_n\}_{n\geq 0}$ such that, for all $n \geq 0$, $m(A_n) \leq C\gamma^n$ and if $x \notin A_n$

(3.2)
$$\mathbb{P}\{S_n(\omega, x) > -\lambda + \varepsilon, \ Z_n(\delta, \omega, x) > \varepsilon\} < C\gamma^n.$$

The following Lemma gives sufficient conditions for the statement of Proposition 3.1 to hold.

Lemma 3.2. Let

$$\tilde{S}_n(\omega, x) := \sum_{i=1}^n \log \left| \left| df^{-1}(f^i_\omega(x)) \right| \right|$$
$$\tilde{Z}_n(\omega, x) := \sum_{i=1}^n -\log \operatorname{dist}_{\delta}(f^i_\omega(x), \mathcal{C}).$$

Suppose that

(3.3)
$$\mathbb{P}\{S_n(\omega, x) > (-\lambda + \varepsilon)n\} + \mathbb{P}\{Z_n(\delta, \omega, x) > \varepsilon n\} < C\gamma^n \qquad \forall x \in X.$$

Then, the statement of Proposition 3.1 holds.

Proof. Suppose (3.3) holds. Note that

$$\{x \in X : -\log\left(\operatorname{dist}_{\delta}(x, \mathcal{C})\right) \ge n\varepsilon\} \subset \{x \in X : \operatorname{dist}_{\delta}(x, \mathcal{C}) \le e^{\varepsilon n}\} \subset A_n := B_{e^{-\varepsilon n}}(\mathcal{C})$$

Hence, if $x \notin A_n$ and $m \ge n$

$$\mathbb{P}\left\{Z_m(\omega, x) > 2m\varepsilon\right\} = \mathbb{P}\left\{-\log\left(\operatorname{dist}_{\delta}(x, \mathcal{C}) + \tilde{Z}_m(\omega, x) > 2m\varepsilon\right\}$$
$$\leq \mathbb{P}\left\{\tilde{Z}_m(\omega, x) \ge m\varepsilon\right\} \le \gamma^m.$$

Using equation (2.3) in Hypothesis (H1), we can argue analogously for $\tilde{S}_n(\omega, x)$ and conclude the proof.

Because of the above Lemma, it remains to prove equation equation (3.3), i.e it remains to establish large deviation estimates in annealed sense both for \tilde{S}_n and \tilde{Z}_n . In order to do it, we exploit an adaptation of Nagaev method as outlined in [18, 4]. Since the proofs for \tilde{S}_n and \tilde{Z}_n are similar, we will outline the details for \tilde{S}_n and leave the rest to the reader.

Let $g(x) := \log |f^{-1}(x)|$. Combining (2.3) and (2.6), it is easy to see that

• g is bounded uniformly outside of $B_{\delta}(\mathcal{C})$ and

(3.4)
$$\int_{B_{\delta}(\mathcal{C})} e^{\theta g(x)} \lesssim \delta^{1-\beta\theta},$$

• the following set inclusion holds

$$(3.5)\qquad \{|g|>\alpha\}\subset B_{Ce^{-\alpha}}(\mathcal{C})$$

for all α large enough,

• Because of (2.3), we also have that

(3.6)
$$\int_{B_{\delta}(\mathcal{C}} g(x) dx \lessapprox \delta \log(\delta)$$

For $\theta > 0$, consider the family of operators $\{\mathcal{P}_{\theta}\}_{\theta > 0}$ defined as

$$P_{\theta} \colon \mathcal{C}^{0}(X) \to \mathcal{C}^{0}$$
$$h \to x \to \mathbb{E}\left[(e^{\theta g}h) \circ f_{\omega}(x) \right].$$

First we show it is well defined in $\mathcal{C}^0(X)$. Indeed

$$||P_{\theta}(h)||_{\mathcal{C}^{0}} \le ||h||_{\mathcal{C}^{0}} \frac{|e^{\theta g}|_{L^{1}}}{2\sigma},$$

and also

$$\begin{aligned} |P_{\theta}(h)(x) - P_{\theta}(h)(y)| &\lesssim ||h||_{C^{0}} \int_{B_{\sigma}(f(x))\Delta B_{\sigma}(f(y))} e^{\theta g(z)} dz \\ &\lesssim \sqrt{\left(\int_{X} e^{2\theta g}\right)} \sqrt{\operatorname{Leb}\left(B_{\sigma}(f(x))\Delta B_{\sigma}(f(y))\right)} \end{aligned}$$

where the integral $\int_X e^{2\theta g}$ is bounded because of (3.4). Furthermore, note that

(3.7)
$$\mathbb{P}\left\{\tilde{S}_{n}(\omega, x) > (\lambda + \varepsilon)n\right\} \leq e^{-(\lambda + \varepsilon)\theta n} \mathcal{P}_{\theta}^{n}(1)(x).$$

The following Lemma is used to find the asymptotic growth of $||\mathcal{P}_{\theta}^{n}(1)||_{\mathcal{C}^{0}}$ as $n \to \infty$.

Lemma 3.3. There exists $\theta_0 > 0$ such that, for all $\theta < \theta_0$, the operator \mathcal{P}_{θ} can be written as

$$\mathcal{P}_{\theta} = \lambda_{\theta} \Pi_{\theta} + Q_{\theta},$$

where Π_{θ} is the projection into the simple eigenvalues associated to the spectral radius λ_{θ} , $Q_{\theta} = Id - \Pi_{\theta}$ and

 $||Q^n|| \le Cr^n,$

for some r < 1.

Proof. Note that

$$\begin{aligned} |\mathcal{P}_{\theta}(h)(x) - \mathcal{P}_{0}(h)(x)| &\leq \frac{||h||_{\mathcal{C}^{0}}}{4\sigma^{2}} \int_{B_{\sigma(f(x))}} \left(e^{\theta g(y)} - 1 \right) dy \\ &\lesssim ||h||_{\mathcal{C}^{0}} \left(\int_{|g| < \theta^{-\frac{1}{3}}} \left(e^{\theta g(y)} - 1 \right) dy + \int_{|g| > \theta^{-\frac{1}{3}}} \left(e^{\theta g(y)} - 1 \right) dy \right) \end{aligned}$$

If $|g| < \theta^{-\frac{1}{3}}$, Then the Taylor series for $e^{\theta g}$ is well defined and we have

$$e^{\theta g(y)} - 1 = \theta g(y) + O(\theta^{\frac{4}{3}}),$$

which implies that the first term is $O(\theta^{\frac{2}{3}})$ as $\theta \to 0$. For the second term note that, because of equation (3.5)

$$\left\{|g| > \theta^{-\frac{1}{3}}\right\} \subset B_{Ce^{-\theta^{-\frac{1}{3}}}}\left(\mathcal{C}\right) := \mathcal{N}_{\theta}.$$

As a result

$$\int_{|g|>\theta^{-\frac{1}{3}}} \left(e^{\theta g(y)} - 1\right) dy \le \int_{\mathcal{N}_{\theta}} e^{\theta g} dx + C e^{-\theta^{-\frac{1}{3}}}$$
$$\lesssim C e^{-\theta^{-\frac{1}{3}}} + C e^{-\theta^{-\frac{1}{3}}(1-\theta)} \lesssim e^{-\frac{1}{2}\theta^{-\frac{1}{3}}}.$$

Combining all estimates, we obtain

$$(3.8) \qquad \qquad ||\mathcal{P}_{\theta} - \mathcal{P}_{0}||_{\mathcal{C}^{0}} = O(\theta^{\frac{2}{3}}).$$

The statement of Lemma 3.3 follows from (3.8) and [4, Proposition 2.3].

By Lemma 3.3 and equation (3.7), one has the Large deviations formula:

(3.9)
$$\sup_{x \in X} \mathbb{P}\left\{\tilde{S}_n(\omega, x) > (\lambda + \varepsilon) n\right\} \le C e^{-(\lambda + \varepsilon)\theta n} \lambda_{\theta}^n,$$

for θ small enough. All it remains to prove is to find an approximation for λ_{θ} as $\theta \to 0$. The following Lemma gives such approximation and concludes the proof of the Proposition.

Proposition 3.4. As $\theta \to 0$, one has

(3.10)
$$\lambda_{\theta} = 1 + \theta \lambda + O(\theta^{\frac{4}{3}}) \approx e^{\theta \lambda}.$$

Proof. By [18, Lemma 1.3]

(3.11)
$$\lambda_{\theta} = \int_{X} e^{\theta g(x)} d\mu(x) + O(\theta^{\frac{4}{3}}).$$

We need to take care of the integral term above. Write

$$\int_X e^{\theta g(x)} d\mu(x) = \int_{|g| < \theta^{-\frac{1}{3}}} e^{\theta g(x)} d\mu(x) + \int_{|g| > \theta^{-\frac{1}{3}}} e^{\theta g(x)} d\mu(x) := G + B$$

First we estimate the bad term B:

$$B = \int_{|g| > \theta^{-\frac{1}{3}}} e^{\theta g(x)} d\mu(x)$$
$$\lesssim \int_{\mathcal{N}_{\theta}} e^{\theta g(x)} dx$$

$$\approx e^{-\frac{1}{2}\theta^{\frac{1}{3}}}$$

Now we consider the good part G. Since $|g| < \theta^{-\frac{1}{3}}$, we can write $e^{\theta g}$ using Taylor expansion, and obtain that

$$G = \int_{|g| < \theta^{-\frac{1}{3}}} 1 + \theta g(x) d\mu(x) + O(\theta^{\frac{4}{3}})$$
$$= O(\theta^{\frac{4}{3}}) + 1 + \theta \lambda - \int_{|g| > \theta^{-\frac{1}{3}}} (1 + \theta g) dx$$
$$\lessapprox O(\theta^{\frac{4}{3}}) + 1 - \theta \lambda - e^{-\theta^{-\frac{1}{3}}},$$

where the last inequality follows from the fact that

$$\int_{|g|<\theta^{-\frac{1}{3}}}gd\mu(x)\approx e^{-\theta^{-\frac{1}{3}}}$$

To conclude, we have that

$$\Lambda_{\theta} = 1 + \theta \lambda + O(\theta^{\frac{4}{3}}) \approx e^{\theta \lambda} \qquad \theta \to 0$$

which concludes the proof.

4. ANNEALED PROPERTIES OF ORBITS OF LARGE BALLS

In this section, we study the annealed properties of orbits of balls of "large scale", which will be crucial to establish existence of Young times in section 4. Given an open set A, and $J \subset A$, we consider the family of events:

$$E_J(I,N,\iota) := \left\{ \exists i \le N \colon \exists \text{ an ball } \tilde{I} \subset I \colon f^i_{\omega}(\tilde{I}) \text{ contains } J, \inf_{j \le i} \inf_{x \in \tilde{I}} d\left(f^j_{\omega}(x), \mathcal{C}\right) > \iota \right\},$$

where I is an ball in Y, $N \in \mathbb{N}$, and $\iota > 0$. Our main result of this section is the following Proposition.

Proposition 4.1. For every open set $A \subset Y$, and any $\varepsilon > 0$ there exists a reference set J, $\iota, \rho > 0$ and $N \in \mathbb{N}$ for which

$$\inf_{|I|\subset Y: |I|\geq\varepsilon} \mathbb{P}\left\{E_J(I,N,\iota)\right\} > \rho.$$

Proof. The above statement is a consequence of the following Lemma.

Lemma 4.2. For every open set A and $\varepsilon > 0$, there exists one point $(\omega, x) \in \Omega \times Y$, $N \in \mathbb{N}$ and $\delta_0, \delta_1 > 0$ such that, for all balls I of size $\geq \varepsilon$, there exists a j such that $y := f_{\omega}^j(x) \in I$ with $B_{\delta_0}(y) \in I$, $f_{\theta^j(\omega)}^{N-j}(B_{\delta_0}(y)) \in A$ and contains $B_{\delta_1}(f_{\omega}^N(x))$. Furthermore,

$$\inf_{j \le N, i \le N-j} d\left(f^i_{\theta^j(\omega)}(B_{\delta_0}(f^j_{\omega}(x))), \mathcal{C}\right) > 0.$$

Indeed, suppose the above Lemma holds. Given $A \subset Y$ and $\varepsilon > 0$, by Proposition 4.2 there exists one point $(\omega, x) \in \Omega \times X, \delta_0 > 0$ and $N \in \mathbb{N}$ such that the quantity

$$l := \inf_{j \le N, \, i \le N-j} d\left(f^i_{\theta^j(\omega)}(B_{\delta_0}(f^j_{\omega}(x))), \mathcal{C}\right) > 0$$

is positive. Consider the set

$$S_{\gamma} := \{ w \in \Omega \colon |\omega_j - w_j| \le \gamma \ \forall j \le N \}.$$

Note that $\mathbb{P}\{S_{\gamma}\} > \gamma^{N+1}$. Choose γ small enough so that for all $w \in S_{\gamma}$, the following conditions hold:

$$\inf_{j\geq 0, i\leq N-j} \inf_{z\in B_{\delta_0}(f^j_w(x))} d\left(f^i_{\theta^j(w)}(z), \mathcal{C}\right) > l/4,$$

$$f_{\theta^j(w)}^{N-j}(B_{\delta_0}(f_w^j(x))) \supset B_{\frac{\delta_1}{20}}(f_w^N(x)) \supseteq B_{\frac{\delta_1}{100}}(f_\omega^N(x)), \qquad \forall j \ge 0,$$

The first condition can be satisfied using continuity, whilst the second follows by Lemma 4.2 and the fact that the Holder regularity of f implies

$$\sup_{j\geq 0, w\in S_{\gamma}} \mathcal{H}\left(f^{N-j}_{\theta^{j}(\omega)}(B_{\delta_{0}}(f^{j}_{\omega}(x))), f^{N-j}_{\theta^{j}(w)}(B_{\delta_{0}}(f^{j}_{w}(x)))\right) \to 0 \ as \ \gamma \to 0,$$

where $\mathcal{H}(\cdot, \cdot)$ denotes the Haussdorf distance. Choosing then

$$J := B_{\frac{\delta_1}{100}}(f^N_{\omega}(x))$$

with $\delta_1, N, \varepsilon_0$ and δ_0 as in Lemma 4.2 and the event S_{γ} , we prove Proposition 4.1. It remains to prove Lemma 4.2

Proof. Fix an open set $A \subset Y$ and $\varepsilon > 0$. Let $\delta := \frac{\varepsilon}{4}$, and let $\{A_i^{\delta}\}_{i \in I}$ be a partition of X into balls such that

$$|A_i^{\delta}| \le \delta \qquad \forall i \in I.$$

Let $J \subset A$ be a reference set of size η and let Δ_c be the ball at the center of Δ of size $\eta/4$. By Birkhoff ergodic theorem, for $\mathbb{P} \times \mu$ almost surely, there exists a time

$$N = N(\delta, \eta, \omega, x)$$

such that the orbit $\{f^i_{\omega}(x)\}_{i\leq N}$ satisfies:

• for all $i \in I$ there exists j such that

$$f^j_{\omega}(x) \in \operatorname{Int}\left(A^{\delta}_i\right)$$

- $f_{\omega}^{N}(x) \in \operatorname{Int}(\Delta_{c}).$ $\{f_{\omega}^{i}(x)\}_{i \leq N} \cap \mathcal{C} = \emptyset.$

Now we fix one (ω, x) and we obtain $N = N(\delta, \eta)$. Take I_c the ball at the center of I of size 2 δ . Then exists j such that $y = f_{\omega}^{j}(x) \in I_{c}$. Let

$$l = \inf_{i \le N} d\left(f_{\omega}^{i}(x), \mathcal{C}\right) > 0,$$

and also

•

$$M := \sup_{x \in X} |f'(x)|$$
$$m := \inf_{x \in X \setminus B_{l/4}(\mathcal{C})} |f'(x)|$$

For r > 0, let $B_r(y)$ be a ball of radius r around $y = f_{\omega}^j(x)$, Then, one can choose $r_0 > 0$ such that, for all $r \leq r_0$.

$$\inf_{j} \inf_{i \leq N-j} d\left(f_{\theta^{j}(\omega)}^{i}(B_{r}(f_{\omega}^{j}(x)), \mathcal{C}\right) > \frac{l}{2},$$

and

$$f^{N-j}_{\theta^j(\omega)}(B_r(f^j_\omega(x))) \Subset \Delta$$

For such r, $f_{\theta^j(\omega)}^{N-j}(y)$ is a ball around $f_{\omega}^N(x) \in \Delta_c$ of size R with

$$rm^n \leq R \leq rM^N.$$

Now choosing $\delta_0 = r$, with $r < r_0$ small enough such that $r_0 + \delta < \varepsilon$ and $\delta_1 = \frac{m^N r}{2}$, we conclude the proof of Lemma 4.2.

5. EXISTENCE OF YOUNG TIMES

In this section we define the Young times. Before, we recall the definition of hyperbolic times. The following definition is adapted from [1, Section 6.1.1]. We start fixing a real number 0 < b < 1/2.

Definition 5.1. Given $\sigma \in (0,1)$ and r > 0, we say that n is a (σ^2, r) -hyperbolic time for $(\omega, x) \in \Omega \times \mathbb{S}^1$ if for every $0 \le k < n$,

$$\prod_{i=k}^{n-1} \left| \left| df^{-1}(f^i_{\omega}(x)) \right| \right| \le \sigma^{2(n-k)}, \text{ and } \operatorname{dist}_r(f^{n-k}_{\omega}(x), \mathcal{C}) \ge \sigma^{b(n-k)},$$

For $x \in X$, let B(x, r) be the open ball around x of radius r. The following Proposition is standard in the context of hyperbolic times and was first proved in [2, Proposition 4.9]. Here we state it as in [12, Proposition 3.2]. Essentially, it shows that, whenever n is a hyperbolic time for (ω, x) , f_{ω}^n uniformly expands a sufficiently small neighbourhood of x to a neighbourhood of $f_{\omega}^n(x)$ of a fixed radius δ_1 .

Proposition 5.2. Given $\sigma, r > 0$, there exist $\tilde{\delta} = \tilde{\delta}(\sigma, r)$ and $C = C(\sigma, r) > 0$, such that for every $(\omega, x) \in \Omega \times \mathbb{S}^1$ for which $n \in \mathbb{N}$ is a (σ^2, r) -hyperbolic time and for every $0 < \delta < \tilde{\delta}$ the following items hold:

- (i) There exists an open neighbourhood $V_n^{\delta}(\omega, x)$ of x, such that f_{ω}^n maps $V_n^{\delta}(\omega, x)$ diffeomorphically to $B_{\delta}(f_{\omega}^n(x))$.
- (ii) for every $z, y \in V_n^{\delta}(x)$ $a. |f_{\omega}^{n-k}(z) - f_{\omega}^{n-k}(y)| \leq \sigma^k |f_{\omega}^n(z) - f_{\omega}^n(y)|$, for every $1 \leq k \leq n$; $b. \log \frac{|\mathrm{d}f_{\omega}^n(z)|}{|\mathrm{d}f_{\omega}^n(y)|} \leq C|f_{\omega}^n(z) - f_{\omega}^n(y)|$.

Fix an open set $A \subset Y$. Then, applying Proposition 4.1 with $\varepsilon = \delta_1$, we infer the existence of $J \subset Y$, $N \in \mathbb{N}$, $\iota, \rho > 0$ such that the event

(5.1)
$$\mathcal{E}(I) := E_J(I, N, \iota) = \left\{ \exists i \le N \colon \exists \tilde{I} \subset I \colon f^i_{\omega}(\tilde{I}) \text{ contains } J, \inf_{j \le i} \inf_{x \in \tilde{I}} d\left(f^j_{\omega}(x), \mathcal{C}\right) > \iota \right\}$$

satisfies

$$\mathbb{P}(\mathcal{E}(I)) > \rho \qquad \forall I \subset Y \colon |I| \ge \delta_1.$$

10

Now consider the family $\{\tau_i : \Omega \times \mathbb{S}^1 \to \mathbb{N} \cup \{\infty\}\}_{i \in \mathbb{N}}$, of N-sparse hyperbolic times, where

$$\tau_i(\omega, x) := \begin{cases} \min\{n; \ n \text{ is a } (\sigma^2, r) \text{ hyperbolic time for } (\omega, x) \} &, \ if \ i = 1, \\ \min\{n; \ n > N + \tau_{i-1}(\omega, x) \text{ is a } (\sigma^2, r) \text{ hyperbolic time for } (\omega, x) \} &, \ if \ i > 1. \end{cases}$$

Definition 5.3. Let $\{\tau_i\}_{i=1}^{\infty}$ be the family of *N*-sparse (σ^2, r) -hyperbolic times. For every $n \in \mathbb{N}$, consider the set

(5.2)
$$Y_n(\omega, x) := \left\{ i \in \{1, \dots, n\} \cap \{\tau_i(\omega, x)\}_{i=1}^{\infty}; \ \theta^i \omega \in \mathcal{E}\left(B_{\delta_1}\left(f_{\omega}^i(x)\right)\right) \right\}.$$

We say that i is a (σ^2, r) -Young time for (ω, x) if there exists $n \in \mathbb{N}$, such that $i \in Y_n(\omega, x)$.

The importance of Young times is due to the fact that if n is a Young time, then f_{ω}^{n} uniformly expands a small neighbourhood of x right into J, keeping uniform distortion bounds, as stated in the following proposition (see [3, Lemma 4.1])

Proposition 5.4. Suppose *n* is a (σ^2, r) -Young time for (ω, x) . Then, there exists $C \ge 1, \delta_1 > 0$ such that there exists an interval $\tilde{I} \subset B\left(x, C\delta_1\kappa_1^{-\frac{n}{2}}\right)$ containing *x* such that $f_{\omega}^n(\tilde{I}) = J$, where *J* is the fixed reference set in (5.1). Furthermore

$$|| (df^n_{\omega}(x))^{-1} || < \sigma^n \qquad \forall x \in \tilde{I},$$

and

$$\left. \frac{df_{\omega}^n(x)}{df_{\omega}^n(y)} \right| \le C \qquad \forall x, y \in \tilde{I}.$$

The following Proposition gives tail estimates for the event that the set $|Y_n(\omega, x)|$ is not dense in $\{1, \ldots, n\}$.

Proposition 5.5. Let f_{ω}^n be a RDS satisfying Hypotheses (H1)-(H2). Then, for all open sets A there exists $\theta_1 > 0$ such that, for all $n \ge 0$ there exists a set E_n with $m(E_n) \le Ce^{-\gamma n}$ such that, if $x \notin E_n$ and m > n

(5.3)
$$\mathbb{P}\left\{|Y_m(\omega, x)| \le \theta_1 m\right\} \le C e^{-\gamma m}$$

Proof. Let $S_n(\omega, x) := \{1, \ldots, n\} \cap \{\tau_i(\omega, x)\}_{i \ge 1}$. By [3, Proposition 3.5] and Proposition 3.1, there exists u > 0 there exists a set $E_n \subset Y$ with $m(E_n) \le Ce^{-\gamma n}$, such that if $x \notin E_n$ and $m \ge n$

$$\mathbb{P}\left\{|S_m(\omega, x)| \le um\right\} \le Ce^{-\gamma m}.$$

As a result, if $x \notin E_n$, $m \ge n$ and $\lambda > 0$

$$\begin{split} \mathbb{P}\left\{|Y_m(\omega, x)| \le \lambda m\right\} \le \mathbb{P}\left\{|Y_m(\omega, x)| \le \lambda m, |S_m(\omega, x)| \le um\right\} + \mathbb{P}\left\{|Y_m(\omega, x)| \le \lambda m, |S_m(\omega, x)| \ge um\right\} \\ \le \mathbb{P}\left\{|Y_m(\omega, x)| \le \lambda m, |S_m(\omega, x)| \ge um\right\} + Ce^{-\gamma m}. \end{split}$$

The result now follows following the lines of [3, Theorem 3.7].

The definition of Young times can be extended from points to balls.

Definition 5.6. We say that n is a young time for (ω, I) if n is an (ω, x) -Young time for x at distance less or equal than |I|/2 from the center of I and $n > \frac{\log(\frac{2\delta}{|I|})}{\log(\lambda)}$.

Consider the following stopping time

$$m(\omega, I) := \min\{n \ge 0 : n \text{ is an } (\omega, I) \text{-young time}\}.$$

In order to construct the horseshoes, we need to estimate the tail $a_k := \mathbb{P}\{m(\omega, I) > k\}$. This task is carried out in the following Proposition.

Proposition 5.7. Consider the event

(5.4)
$$E_n(\omega) := |\{x \in \mathbb{S}^1 \colon |Y_n(\omega, x)| < \theta_1 n\}|$$

There exists a random variable $n(\omega)$ such that, for all $k > \frac{\log(\frac{2C}{|C|})}{\gamma}$

$$\mathbb{P}\left(n_0 > k\right) \le C e^{-\gamma k}$$

and for any $n \ge n_0(\omega)$

$$Leb(E_n(\omega)) < \frac{|I|}{2}.$$

In particular, if $n > \frac{\log\left(\frac{2C}{|I|}\right)}{\gamma}$

$$\mathbb{P}\left\{E_n(\omega) > |I|/2\right\} \le \mathbb{P}\left(n_0 > n\right) \le Ce^{-\gamma n}$$

and (5.5)

$$\mathbb{E}m(\omega, I) \approx \log(|I|^{-1}).$$

Proof. First note that

$$m \otimes \mathbb{P}\{(\omega, x) \colon |Y_n(\omega, x)| < \theta_1 n\} \le m(E_n) + Ce^{-\gamma n}$$

Then, the first part of Proposition 5.7 follows from the lines of [12, Proposition 4.3]. Now let
$$k_0 := -A \log(B|I|) > \max\left\{\frac{\log\left(\frac{2C}{|I|}\right)}{\gamma}, \frac{1 + \frac{\log\left(\frac{2K}{|I|}\right)}{\log(\lambda)}}{\theta}\right\}.$$
 If $k > k_0$, then $\{m(\omega, I) > k\} \le \{E_k(\omega) > |I|/2\} \subset \{n_0 > k\}.$

Indeed, if $E_k(\omega) < |I|/2$ and $k > k_0$, then

$$\{E_k(\omega) < |I|/2\} \subset \{\exists \theta k - \log(1/|I|) \ (\omega, I) \text{-Young times}\}.$$

As a result

(5.6)
$$\mathbb{E}[m(\omega)] = \sum_{k \le k_0} 1 + C \sum_{k \ge k_0} e^{-\gamma k} \approx \log(1/|I|) + C$$

6. Existence of the horseshoe

6.1. **Preliminary work.** Before diving into the construction of horseshoes, we prove a few genera Propositions that will be used later. The first Proposition ensures that the Birkhoff averages of IID random variables with infinite average pointwise explode almost surely.

Proposition 6.1. Let $\{X_n\}_{n\geq 0}$ be a sequence of non negative iid random variables such that $E[X_0] = \infty$. Then

$$\liminf_{n \to \infty} \frac{X_0 + \dots + X_{n-1}}{n} = \infty.$$

Proof. Let

$$Y_n^k := \min\{X_n, k\} \le X_n,$$

Note that, for all $k \ge 0$

$$\frac{X_0 + \dots + X_{n-1}}{n} \ge \frac{Y_0^k + \dots + Y_{n-1}^k}{n},$$

so that

$$\liminf_{n \to \infty} \frac{X_0 + \dots + X_{n-1}}{n} \ge \liminf_{n \to \infty} \frac{Y_0^k + \dots + Y_{n-1}^k}{n} = \mathbb{E}\left[Y_0^k\right],$$

where the latter follows from Cental Limit Theorem for iid random variables. We claim

$$\lim_{k \to \infty} \mathbb{E}\left[Y_0^k\right] = \infty$$

Indeed, $Y_0^k \ge 0$ for all $k, Y_0^{k+1} \ge Y_0^k$ and they converge pointwise to X_0 . The claim follows by Monotone convergence theorem.

Now suppose we have a sequence $\{n_j\}_{j\geq 0}$. We want to know if there is any kind of relationship between the asymptotic behaviour of the sequence $\left\{\frac{n_j}{j}\right\}_{j\geq 0}$ and the sequence

$$\left\{\frac{|\{1,\ldots,j\}\cap\{n_k\}_{k\geq 0}|}{j}\right\}_{j\geq 0}$$

where the *j*-th element is the proportion of elements of $\{n_j\}_{j\geq 0}$ in the set $\{1,\ldots,j\}$. This task is carried out in the following two propositions.

Proposition 6.2. Let $\{n_j\}_{j\geq 0}$ be an increasing sequence. Then

(6.1)
$$\limsup_{n \to \infty} \frac{n_k}{k} \le \alpha \implies \liminf_{n \to \infty} \frac{|\{i : n_i \le n\}|}{n} \ge \frac{1}{3\alpha}$$

and

(6.2)
$$\liminf_{n \to \infty} \frac{|\{i \colon n_i \le n\}|}{n} \ge \beta \implies \limsup_{n \to \infty} \frac{n_k}{k} \le \frac{3}{\beta}.$$

Proof. First we prove (6.1). Suppose that

$$\limsup_{n \to \infty} \frac{n_k}{k} \le \alpha.$$

Then, for $k > k_0$

$$n_k < 2\alpha k$$

Then, for any $k \ge 0$ such that $\lfloor k/2\alpha \rfloor > k_0$. Then

$$\begin{split} |\{i: n_i < k\}| &= |\{i: n_i \le 2\alpha \left(\frac{k}{2\alpha}\right)\}|\\ &\geq |\{i: n_i \le 2\alpha \left(\left\lfloor\frac{k}{2\alpha}\right\rfloor\right)\}|\\ &\geq \left\lfloor\frac{k}{2\alpha}\right\rfloor\\ &\geq \frac{k}{3\alpha} \end{split}$$

since , by assumption $n_j \leq j$ for all $j > k_0$ and the sequence n_j is increasing. Now we prove (6.2). Suppose that

$$\liminf_{n \to \infty} \frac{|\{i \colon n_i \le n\}|}{n} \ge \beta$$

. Then, for $k > k_0$

$$|\{i: n_i \le k\}| \ge \frac{\beta}{2}k.$$

Then, for all k such that $\frac{2}{\beta}k > k_0$

$$|\{i: n_i \le \left\lfloor \frac{2}{\beta}k \right\rfloor + 1\}| \ge \frac{\beta}{2} \left(\left\lfloor \frac{2}{\beta}k \right\rfloor + 1 \right) \ge \beta/2 \left((2/\beta)k - 1 + 1 \right) > k$$

As a result, for k large enough

$$n_k \le (3/\beta) \, k.$$

This concludes

We also need the following Proposition that establishes an even stronger link between the two asymptotic fr

Proposition 6.3. Let $\{n_k\}_{k\geq 0}$ be an increasing sequence. Suppose that

$$\limsup_{n \to \infty} |\{i \colon n_i \le k\}|/k > \beta.$$

Then

$$\liminf_{n \to \infty} \frac{n_k}{k} < \frac{2}{\beta}$$

Proof. Suppose the first item holds. Then there exists an increasing sequence $s_r \to \infty$ as $r \to \infty$ such that

$$\lim_{r \to \infty} \frac{|i: n_i \le s_r|}{s_r} > \beta.$$

Then for $r > r_0$

$$|i: n_i \le s_r| \ge \beta s_r.$$

Consider the sequence

$$q_r := \left\lfloor \frac{\beta}{2} s_r \right\rfloor + 1.$$

Then, for $r > r_0$

$$n_{q_r} \le s_r \le \frac{2}{\beta} q_r$$

As a consequence

$$\limsup_{r \to \infty} \frac{n_{q_r}}{q_r} < \frac{2}{\beta},$$

which concludes the proof.

6.2. Horseshoe construction. Now we are going to leverage (5.6), Proposition 6.2 and 6.3. First, Let J be the reference ball and consider M large which will be defined later. Let I_1, \ldots, I_M a sequence of balls size $\approx \frac{|J|}{M}$. For any of these balls we can consider $m(I_i)_0 = 0$, $m^i(I_i)_1$ the first (ω, I_i) -Young time and

$$m(I_i)_{k+1} = m(I_i)_1(\theta^{m(I_i)_k}(\omega)).$$

Since the sequence $\{m(I_i)_k\}_{k\geq 0}$ has independent and identically distributed differences, we have that

$$\lim_{k \to \infty} \frac{m(I_i)_k}{k} = \mathbb{E}m(I_i)_1 < C + \log\left(\frac{M}{|J|}\right)$$

Now we consider the family

(6.3)
$$H_i^n(\omega) := \{m(I_i)_k(\omega)\}_{k \ge 0} \cap \{1, \dots, n\}.$$

Note that

$$|H_i^j(\omega)| = |\{k \ge 1 \colon m(I_i)_k(\omega) \le n\}|.$$

Hence, by Proposition 6.2

(6.4)
$$\liminf_{n \to \infty} \frac{|H_i^n|}{n} > \frac{1}{2\left(C + \log\left(\frac{M}{|J|}\right)\right)} := V_M$$

We choose M such that $MV_M > 1$. By the Bonferroni inequality, for every n, we have

(6.5)
$$n \ge \left| \bigcup_{j=1}^{M} H_{j}^{n} \right| \ge \sum_{j=1}^{M} |H_{j}^{n}| - \sum_{1 \le j < i \le n} |H_{j}^{n} \cap H_{i}^{n}|.$$

Using the above and equation (6.4). there exists $n_0(\omega)$ such that for all $n \ge n_0(\omega)$

$$\sum_{1 \le j < i \le n} \left| H_j^n \cap H_i^n \right| \ge \sum_{j=1}^M |H_j^n| - n$$
$$\ge (MV_M - 1)n.$$

Then, for every $n > n_0(\omega)$, there exists indices $i(n, \omega)$ and $j(n, \omega)$ such that

(6.6)
$$\left| H_{j(n,\omega)}^n \cap H_{i(n,\omega)}^n \right| > Z_M n,$$

with $Z_M = \frac{V_M}{M} - 1/M^2$. Since there are at most M^2 pairs of $(i, j) \in \{1, \dots, M\}^2$, then there exist with probability one indices $i(\omega)$ and $j(\omega)$ such that

(6.7)
$$\left| H_{j(\omega)}^n \cap H_{i(\omega)}^n \right| > Z_M n$$
 for infinitely many n s,

or equivalently

(6.8)
$$\limsup_{n \to \infty} \frac{\left| H_{j(\omega)}^n \cap H_{i(\omega)}^n \right|}{n} > Z_M$$

Note that we can write

$$\bigcup_{n\geq 0} \ H^n_{j(\omega)}\cap H^n_{i(\omega)} = \left\{n_k^{i(\omega),j(\omega)}(\omega)\right\}_{k\geq 0}.$$

By (6.8) and Proposition 6.3,

(6.9)
$$\liminf_{k \to \infty} \frac{n_k^{i(\omega), j(\omega)}(\omega)}{k} < \frac{2}{Z_M}$$

Now consider the event

$$E := \left\{ \exists i, j \colon \liminf_{k \to \infty} \frac{n_k^{i,j}(\omega)}{k} < \frac{2}{Z_M} \right\},$$

we have $\mathbb{P}(E) = 1$. Note that

$$E = \bigcup_{i,j} E_{i,j},$$

where

$$E_{i,j} := \left\{ \liminf_{k \to \infty} \frac{n_k^{i,j}(\omega)}{k} < \frac{2}{Z_M} \right\}.$$

Hence, there exists i, j such that

$$\mathbb{P}(E_{i,j}) > 0.$$

Now we consider the event

(6.10)
$$m^{i,j}(\omega)_1 := \min\{n \ge 0 : n = m(I_i)_u = m(I_j)_v(\omega) \text{ for some } u, v \in \mathbb{N}\}$$

i.e. $m^{i,j}(\omega)_1$ is the first time both I_i and I_j cover J. Note that $m^{i,j}(\omega)_1$ is also the smallest element of $H^i(\omega) \cap H^j(\omega)$. One then can define inductively the stopping times $\{m^{i,j}(\omega)_k\}_{k \leq k_0}$, where $k_0 \in \mathbb{N} \cup \{+\infty\}$ is such that $m^{i,j}(\omega)_{k_0} = \infty$.

One can also prove by induction that

(6.11)
$$m^{i,j}(\omega)_k = n_k^{i,j}(\omega) \qquad \forall k \ge 0, \ \omega \in E_{i,j}.$$

In particular

(6.12)
$$\liminf_{k \to \infty} \frac{m^{i,j}(\omega)_k}{k} < \frac{2}{Z_M} \qquad \forall \omega \in E_{i,j}$$

By proposition 6.1 $E[m_1^{i,j}]$ is finite. By Law of large number, then

(6.13)
$$\lim_{k \to \infty} \frac{m_k^{i,j}}{k} = \mathbb{E}m^{i,j},$$

with probability one.

References

- J. F. Alves. Nonuniformly Hyperbolic Attractors-Geometric and Probabilistic Aspects. Springer Monographs in Mathematics. Springer, Cham, [2020] ©2020.
- Jose F. Alves and Helder Vilarinho. Strong stochastic stability for non-uniformly expanding maps. Ergodic Theory and Dynamical Systems, 33(3):647–692, 2013.
- [3] Matheus M Castro and Giuseppe Tenaglia. Random young towers and quenched decay of correlations for predominantly expanding multimodal circle maps. arXiv preprint arXiv:2303.16345, 2023.
- [4] Sébastien Gouëzel. Limit theorems in dynamical systems using the spectral method. Hyperbolic dynamics, fluctuations and large deviations, 89:161–193, 2015.
- [5] Martin Hairer. Convergence of markov processes. Lecture notes, https://www.hairer.org/notes/Convergence.pdf, 2010.

- [6] Wen Huang and Zeng Lian. Horseshoes for Anosov systems on fibers driven by an equicontinuous system. Acta Mathematica Sinica, 38(1):281–290, 2022.
- [7] Wen Huang, Zeng Lian, and Kening Lu. Ergodic theory of random anosov systems mixing on fibers. arXiv preprint arXiv:1612.08394, 2016.
- [8] Wen Huang and Kening Lu. Entropy, chaos, and weak horseshoe for infinite-dimensional random dynamical systems. Communications on Pure and Applied Mathematics, 70(10):1987–2036, 2017.
- Wen Huang and Jianhua Zhang. Full-horseshoes for the galerkin truncations of 2d navier-stokes equation with degenerate stochastic forcing. arXiv preprint arXiv:2303.05027, 2023.
- [10] Wen Huang and Jianhua Zhang. Observable full-horseshoes for lagrangian flows advected by stochastic 2d navier-stokes equations. arXiv preprint arXiv:2311.05193, 2023.
- [11] Anatole Katok. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. Publications Mathématiques de l'IHÉS, 51:137–173, 1980.
- [12] Jeroen SW Lamb, Giuseppe Tenaglia, and Dmitry Turaev. Horsehoes for a class of nonuniformly expanding random dynamical systems on the circle. arXiv preprint arXiv:2304.03685, 2023.
- [13] Zeng Lian and Xiao Ma. Existence of periodic orbits and horseshoes for mappings in a separable banach space. Journal of Differential Equations, 269(12):11694–11738, 2020.
- [14] Zeng Lian and Mikko Stenlund. Positive Lyapunov exponent by a random perturbation. Dynamical Systems, 27:239 – 252, 2012.
- [15] Zeng Lian and Lai-Sang Young. Lyapunov exponents, periodic orbits and horseshoes for mappings of hilbert spaces. Annales Henri Poincaré, 12(6):1081–1108, September 2011.
- [16] Zeng Lian and Lai-Sang Young. Lyapunov exponents, periodic orbits, and horseshoes for semiflows on hilbert spaces. *Journal of the American Mathematical Society*, 25:637–665, 2012.
- [17] Xiao Ma. Existence of periodic orbits and horseshoes for semiflows on a separable banach space. Calculus of Variations and Partial Differential Equations, 61(6):217, October 2022.
- [18] S. V. Nagaev. Some limit theorems for stationary markov chains. Theory of Probability & Its Applications, 2(4):378–406, 1957.
- [19] James R. Norris. Markov chains, volume 2 of Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 1998. Reprint of 1997 original.

 $^2\mathrm{INternational}$ Research Center for Neurointelligence, The University of Tokyo, Tokyo 113-0033, Japan

³CENTRE FOR APPLIED MATHEMATICS AND BIOINFORMATICS, DEPARTMENT OF MATHEMATICS AND NATURAL SCIENCES, GULF UNIVERSITY FOR SCIENCE AND TECHNOLOGY, HALWALLY 32093, KUWAIT

Email address: jeroen.lamb@imperial.ac.uk

¹Department of Mathematics, Imperial College London, London SW7 2AZ, UK

Email address: giuseppe.tenaglia20@imperial.ac.uk

Email address: d.turaev@imperial.ac.uk