CONVERGENCE OF DISCONTINUOUS GALERKIN METHODS FOR QUASICONVEX AND RELAXED VARIATIONAL PROBLEMS

G. GREKAS ¹, K. KOUMATOS ², C. MAKRIDAKIS ^{2,3,4}, AND A. VIKELIS ⁵

ABSTRACT. In this work, we establish that discontinuous Galerkin methods are capable of producing reliable approximations for a broad class of nonlinear variational problems. In particular, we demonstrate that these schemes provide essential flexibility by removing inter-element continuity while also guaranteeing convergent approximations in the quasiconvex case. Notably, quasiconvexity is the weakest form of convexity pertinent to elasticity. Furthermore, we show that in the non-convex case discrete minimisers converge to minimisers of the relaxed problem. In this case, the minimisation problem corresponds to the energy defined by the *quasiconvex envelope* of the original energy. Our approach covers all discontinuous Galerkin formulations known to converge for convex energies. This work addresses an open challenge in the vectorial calculus of variations: developing and rigorously justifying numerical schemes capable of reliably approximating nonlinear energy minimization problems with potentially singular solutions, which are frequently encountered in materials science.

1. INTRODUCTION

In this work, we establish that discontinuous Galerkin methods are capable of producing reliable approximations for a broad class of nonlinear variational problems. Our aim is twofold: first, to show that these methods are applicable to variational problems involving *quasiconvex energies*; and second, to establish that, even in the absence of convexity assumptions, the approximations converge to the minimisers of the *relaxed* problem. In this case, the minimisation problem corresponds to the energy defined by the *quasiconvex envelope* of the original energy.

By addressing this, we contribute to a long-standing algorithmic and theoretical challenge in the vectorial calculus of variations: the development and rigorous justification of numerical schemes capable of reliably approximating nonlinear energy minimisation problems with possibly singular solutions, which commonly arise in materials science.

The insightful observation by J.M. Ball in [2]—"minimisers of variational problems may have singularities, but natural numerical schemes may fail

Date: January 22, 2025.

¹Computer, Electrical, Mathematical Sciences and Engineering Division, KAUST.

²Department of Mathematics, University of Sussex.

³Institute of Applied & Computational Mathematics, Foundation for Research & Technology-Hellas.

⁴Department of Mathematics and Applied Mathematics, University of Crete.

⁵Faculty of Civil Engineering-Czech Technical University of Prague.

to detect them"—remains highly relevant today. As discussed in detail in Ball [2] and Carstensen [12], although standard finite element methods provide a natural discrete framework for seeking minimisers of such problems, the inter-element continuity imposed on the discrete spaces appears to lack sufficient flexibility to make these methods the optimal choice. This limitation is particularly evident in the computation of minimisers for variational problems where the *Lavrentiev phenomenon* may occur. As first noted in [3], standard finite element methods can converge to energy values that are strictly greater than the exact energy of the problem, leading to physically irrelevant approximations.

In the finite element literature, there are alternatives to standard finite elements that offer greater flexibility, such as nonconforming finite elements and discontinuous Galerkin methods. The spaces introduced by Crouzeix and Raviart [20], originally developed for the Stokes and Navier-Stokes equations, were employed by Ortner [33] (see also [34] and [1]) to demonstrate that these spaces provide sufficient flexibility to overcome the limitations of standard methods, particularly in addressing the Lavrentiev gap phenomenon.

Discontinuous Galerkin methods, which are completely discontinuous, have become especially popular for nonlinear PDEs. While these methods offer high-order approximation capabilities, and great flexibility, they require specially constructed energy functionals, which are often nontrivial to design and analyse [38], [11], [27]. Γ -convergence for convex energies associated with the energy functional proposed in [38] was established in Buffa and Ortner [11], while the authors of this paper derived Γ -convergence for the functional introduced in [27].

Despite the progress outlined above, allowing possibly noncontinuous discrete approximations comes at a cost: the analytical treatment of energies involving the discrete gradient heavily relies on the convexity of W. Specifically, due to the absence of a true gradient structure in the discrete gradient —i.e., $\operatorname{curl} G_h(u_h) \neq 0$, $G_h(u_h)$ being an appropriate "discrete gradient" the lower semicontinuity of the energy term is guaranteed for convex integrands W, but remains uncertain when W is quasiconvex. This represents a significant technical challenge that hinders further advancements in the design and analysis of approximation methods applicable to elasticity and other variational problems.

In this work, we resolve this issue by showing that discontinuous Galerkin schemes not only provide the necessary flexibility by eliminating inter-element continuity but also ensure convergent approximations in the quasiconvex case. To compensate the lack of differential structure and regain lower semicontinuity, we transition to the limit of reconstructed approximations from Karakashian and Pascal [28] and demonstrate that the associated error terms can be adequately controlled. Furthermore, we show that in the non-convex case discrete minimisers converge to minimisers of the relaxed problem.

Plan of the paper. For $\Omega \subset \mathbb{R}^d$, we consider the minimisation problem associated with the energy

(1.1)
$$\mathcal{E}(u) = \int_{\Omega} W(\nabla u(x)) \, dx$$
, where $u : \Omega \to \mathbb{R}^N$, $u = u_0$, on $\partial \Omega$,

over an admissible set of functions which is a subset of a Sobolev space $W^{1,p}$ and where $W : \mathbb{R}^{N \times d} \to \mathbb{R}$. Detailed assumptions on the corresponding energy minimisation problem are in Section 3. We start with some preliminary material in Section 2 including standard notation on the discontinuous Galerkin spaces $V_h(\Omega)$. Lemma 2.1 is an extension in the L^p setting of the classical result in [28] regarding the error between $u_h \in V_h(\Omega)$ and its continuous reconstruction $w_h \in V_h(\Omega) \cap W^{1,p}(\Omega)$. This is a crucial tool in our analysis.

In Section 3 we introduce our discontinuous Galerkin method [27] and we show that, in the absence of convexity, our discrete energies Γ -converge to the relaxed problem. Similarly with [27], a penalty term ensures the discrete stability, but in this case it is appropriately modified to include also energies which are not convex. Our results cover both discontinuous Galerkin formulations known to converge for convex W. The proof for the method based on discrete gradients [38] is postponed to the Section 3.2. The discrete gradients are defined through standard lifting operators, defined in the same section. Although the use of the lifting operators and the discrete gradients was the main tool in the analysis in the convex case, as mentioned, this is no longer true if we would like to consider quasiconvex integrands. The main obstacle in proving the convergence of DG schemes for quasiconvex energies lies in the fact that the finite element function $\nabla_h y_h$ used to approximate the gradient of an admissible function ∇y is not itself a gradient, i.e. $\operatorname{curl} \nabla_h y_h \neq 0$, and hence in general the lower semicontinuity of \mathcal{E} with respect to weak convergence in $W^{1,p}$ is not true. That is, for a quasiconvex W, the inequality

$$\liminf_{h \to 0} \int_{\Omega} W(\nabla_h y_h) \ge \int_{\Omega} W(\nabla y)$$

does not hold in general when

$$\nabla_h y_h \rightarrow \nabla y$$
, in L^p as $h \rightarrow 0$.

The main results of this paper are Theorems 3.1 and 3.2 where we show that the discrete minimisers of disontinuous Galerkin schemes of [27] and [38] converge to a minimiser of the relaxed problem defined as

$$\mathcal{E}^{rel}(y) = \int_{\Omega} W^{qc}(\nabla y(x)) \, dx$$

where W^{qc} denotes the quasiconvex envelope of W, i.e. the largest quasicnovex function that lies below W. In particular, if W is quasiconvex, then $W = W^{qc}$, and as a consequence we establish convergence of the discrete minimisers to a minimiser of the continuous quasiconvex energy.

In Section 4, we provide detailed numerical evidence using specifically designed examples. We begin with a problem involving polyconvex energy and thoroughly analyze the impact of the new penalty term introduced in this work (see Section 3.1) in comparison to the penalty term employed in [27].

Subsequently, we examine a frame-indifferent two-well strain energy function where, under specific boundary conditions, a phase mixture emerges. This energy leads to increasingly fine microstructures as the mesh size h is reduced, approaching the infimum of the total potential energy. An intriguing aspect of our experiments, consistent with the theoretical results, is the observation that for this problem, minimizing sequences with distinct laminate structures exist. Nevertheless, all such sequences strongly converge in L^2 to the same function. This behavior can be attributed to the diminishing amplitude of gradient oscillations, which become negligible as h decreases.

An additional interesting observation is that, by leveraging Dacorogna's formula for the quasiconvex envelope of a general energy density W, our scheme can approximate $W^{qc}(F)$ pointwise for each F. This suggests that quasiconvex envelope approximations can be constructed using discrete energy minimisation techniques. While the full implementation of this idea lies beyond the scope of the current paper, we present preliminary results for this pointwise approximation in Figures 5, 6, 7. As mentioned, these numerical experiments are for non-quasiconvex, double-well potentials W, which are typical in modeling phase transitions in solids.

Further references. The systematic study of numerical methods for energy minimisation problems arising in elasticity started in the late 80-90's with the works of Ball and Knowles [3], Collins, Kinderlehrer, and Luskin [18], Collins and Luskin [19], Pedregal [36], Carstensen and Plecháč [14], [15], [16], Carstensen and Roubíček [17]. For reviews, open problems and remarks of the effectivity of various numerical approaches, see e.g., [31], [2], [12], [37], [13], [6] and their references. Early approaches emphasising the benefits of potential flexibility offered by nonconforming or discontinuous methods were discussed in [31], [24]. Crouzeix-Raviart nonconforming spaces are used in total variation and other convex minimisation problems in [8], [9]. Discontinuous Galerkin schemes for problems with higher gradients were considered in [10], [26]. To obtain feasible discrete solutions for the relaxed problem using its definition, it is necessary to compute approximations of polyconvex or quasiconvex envelopes. Such approaches have been explored in [30], [23], [5], [7], [29], [32], among others, along with the references therein.

2. Preliminaries.

2.0.1. Finite element spaces and trace operators. For simplicity, let us assume that Ω is a polytope domain, and consider a triangulation T_h of Ω of mesh size h > 0. We seek discontinuous Galerkin approximations to (1.1). To this end, we introduce the space of L^p piecewise polynomial functions

(2.1)
$$V_h^q(\Omega) := \left\{ v \in L^p(\Omega) : v \Big|_K \in \mathbb{P}_q(K), K \in T_h \right\},$$

where $\mathbb{P}_q(K)$ denotes the space of polynomial functions on K of degree $q \in \mathbb{N}$ which is henceforth fixed. Moreover, we adopt the following standard notation: The *trace* of functions in $V_h^q(\Omega)$ belongs to the space

(2.2)
$$T(E_h) := \prod_{e \in E_h} L^p(e),$$

where E_h is the set of mesh edges. The *average* and *jump* operators over $T(E_h)$ are defined by:

(2.3)
$$\{\!\!\{ \cdot \}\!\!\} : T(E_h) \mapsto L^p(E_h) \\ \{\!\!\{ w \}\!\!\} := \begin{cases} \frac{1}{2} (w|_{K_{e^+}} + w|_{K_{e^-}}), & \text{for } e \in E_h^i \\ w, & \text{for } e \in E_h^b \end{cases}$$

(2.4)
$$\begin{split} \llbracket \cdot \rrbracket : T(E_h) &\mapsto L^p(E_h) \\ \llbracket v \rrbracket := v \big|_{K_{e^+}} - v \big|_{K_{e^-}}, \quad \text{for } e \in E_h^i \end{split}$$

We often apply the jump operator on dyadic products $v \otimes n$, for $v \in \mathbb{R}^N$, $n \in \mathbb{R}^d$ and we write

(2.5)
$$\llbracket v \otimes n_e \rrbracket := v \otimes n_{e^+} |_{K_{e^+}} + v \otimes n_{e^-} |_{K_{e^-}}, \quad \text{for } e \in E_h^i$$

where $(v \otimes n)_{ij} = v_i n_j$, K_{e^+} , K_{e^-} are the elements that share the internal edge e with corresponding outward pointing normals n_{e^+} , n_{e^-} , and E_h^i , E_h^b denote respectively the set of internal and boundary edges.

The space $V_h^q(\Omega)$ is equipped with the norm

$$\|v\|_{W^{1,p}(\Omega,T_h)}^p = \|v\|_{L^p(\Omega)}^p + |v|_{W^{1,p}(\Omega,T_h)}^p,$$

where the seminorm $|\cdot|_{W^{1,p}(\Omega,T_h)}$ is defined by

(2.6)
$$|v|_{W^{1,p}(\Omega,T_h)}^p := \sum_{K \in T_h} \int_K |\nabla v|^p + \sum_{e \in E_h^i} \frac{1}{h^{p-1}} \int_e |[v]|^p.$$

We note that in the above expression ∇v denotes the gradient, in the classical sense, of the polynomial function v over the set K. As a function in $SBV(\Omega)$, $v \in V_h^q(\Omega)$, admits a distributional gradient which is a Radon measure and decomposes as

(2.7)
$$Dv = \nabla v \mathcal{L}^d \sqcup \Omega + \llbracket v \rrbracket \otimes n_e \mathcal{H}^{d-1} \sqcup E_h$$

where ∇v denotes the approximate gradient. To avoid confusion, we denote the above approximate gradient by $\nabla_h v$, i.e. we write $\nabla_h : V_h^k \to V_h^{k-1}$ for the operator defined by

$$(\nabla_h v)\big|_K = \nabla(v\big|_K).$$

In particular, it holds that

$$\int_{\Omega} W(\nabla_h u_h) = \sum_{K \in T_h} W(\nabla u_h).$$

2.0.2. Approximation results and Poincaré inequalities in DG spaces. Here we approximate discontinuous functions $u_h \in V_h(\Omega)$ by continuous piecewise functions $w_h \in V_h(\Omega) \cap W^{1,p}(\Omega)$ and prove their error estimates by generalising the case of p = 2 presented in [28, Theorem 2.2]. From this result we derive a Poincaré inequality for functions in $V_h(\Omega)$ while these estimates are constituents for the Γ -convergence result when the energy function Wis quasiconvex. First we generalise the algebraic inequality of [28, Lemma 2.2] **Lemma 2.1.** Given n real numbers $c_1, c_2, ..., c_n$, let $m = \frac{1}{n} \sum_{i=1}^n c_i$. If $r \ge 1$, then

$$\sum_{i=1}^{n} |c_i - m|^r \le C \sum_{i=1}^{n-1} |c_{i+1} - c_i|^r,$$

where C depends only on n and r.

Proof. Here we follow partially the proof of [28, Lemma 2.2]. For $r \ge 1$ one can use Jensen's inequality for an upper bound

$$|c_j - m|^r = \frac{1}{N^r} \left| \sum_{i=1}^n (c_j - c_i) \right|^r \lesssim \sum_{i=1}^n |c_j - c_i|^r.$$

Now, summing over j we deduce that

$$\sum_{j=1}^{n} |c_j - m|^r \lesssim \sum_{j=1}^{n} \sum_{i=j+1}^{n} |c_j - c_i|^r = \sum_{j=1}^{n} \sum_{i=j+1}^{n} \left| \sum_{k=i}^{j-1} (c_{k+1} - c_k) \right|^r$$
$$\lesssim \sum_{j=1}^{n} \sum_{i=j+1}^{n} \sum_{k=i}^{j-1} |c_{k+1} - c_k|^r \lesssim \sum_{j=1}^{n} |c_{j+1} - c_j|^r.$$

Using the above Lemma, we extend the approximation result [28, Theorem 2.2] for the case $p \ge 2$. We omit the details of the proof emphasising more on the points where our case differs from the classical case p = 2.

Lemma 2.2. Let $u_h \in V_h(\Omega)$, then for any multi-index α with $|\alpha| = 0, 1$ there exists $w_h \in V_h(\Omega) \cap W^{1,p}(\Omega)$ (continuous finite element space), such that

(2.8)
$$\sum_{K\in T_h} \|\nabla^{\alpha} u_h - \nabla^{\alpha} w_h\|_{L^p(K)}^p \lesssim \sum_{e\in E_h^i} h_e^{1-p|\alpha|} \int_e |\llbracket u_h]\|^p ds,$$

and for $\Gamma \subset \partial \Omega$:

(2.9)
$$\|\nabla^{\alpha} u_h - \nabla^{\alpha} w_h\|_{L^p(\Gamma)}^p \lesssim \sum_{e \in E_h^i} h_e^{-p|\alpha|} \int_e |\llbracket u_h \rrbracket|^p.$$

Proof. We proceed similarly to the proof of [28] adopting partially their notation. For $K \in T_h$ the set of the Lagrange nodes of K is denoted by $\mathcal{N}_K = \{x_{K,j} : j = 1, ..., m\}$ and $\{\phi_K^{(x_{K,j})} : j = 1, ..., m\}$ the corresponding basis functions. In addition to this, we set $\mathcal{N} := \bigcup_{K \in \mathcal{T}_h} \mathcal{N}_K$. For each $\nu \in \mathcal{N}$, let $\omega_{\nu} := \{K \in T_h : \nu \in K\}$ and $|\omega_{\nu}|$ its cardinality. Now, we write

$$u_h(x) := \sum_{\nu \in \mathcal{N}} \sum_{K \in \omega_\nu} a_K^{(\nu)} \phi_K^{(\nu)}(x)$$

and at every node we define the continuous w_h to be the average of the values of u_h , i.e.

$$w_h = \sum_{\nu \in \mathcal{N}} \beta^{(\nu)} \phi^{(\nu)}, \text{ where } \beta^{(\nu)} = \frac{1}{|\omega_\nu|} \sum_{K \in \omega_\nu} a_K^{(\nu)}, \ \nu \in \mathcal{N},$$

and $\phi^{(\nu)}|_K = \phi_K^{(\nu)}$. After a simple scaling argument we know that $\|\nabla^{\alpha}\phi_K^{(\nu)}\|_{L^p(K)}^p \lesssim h_K^{d-p|\alpha|}$. Hence, employing this upper bound and Jensen's inequality we obtain

$$\begin{split} \sum_{K\in T_h} \int_K |\nabla^{\alpha} u_h - \nabla^{\alpha} w_h|^p dx &= \sum_{K\in T_h} \int_K \left| \sum_{\nu\in\mathcal{N}_K} (\alpha_K^{(\nu)} - \beta^{(\nu)}) \nabla^{\alpha} \phi_K^{(\nu)}(x) \right|^p dx \\ &\lesssim \sum_{K\in T_h} \sum_{\nu\in\mathcal{N}_K} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p \int_K |\nabla^{\alpha} \phi_K^{(\nu)}(x)|^p dx \lesssim \sum_{K\in T_h} \sum_{\nu\in\mathcal{N}_K} h_K^{d-p|\alpha|} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p \\ &= \sum_{\nu\in\mathcal{N}} \sum_{K\in\omega_\nu} h_K^{d-p|\alpha|} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p \lesssim \sum_{\nu\in\mathcal{N}} h_\nu^{d-p|\alpha|} \sum_{K\in\omega_\nu} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p, \end{split}$$

where $h_{\nu} = \max_{K \in \omega_{\nu}} h_K$. Following the same strategy as in [28] for the ordering of ω_{ν} and applying instead Lemma 2.1 we infer that

$$\sum_{K \in T_h} \|\nabla^{\alpha} u_h - \nabla^{\alpha} w_h\|_{L^p(K)}^p \lesssim \sum_{e \in E_h^i} h_e^{d-p|\alpha|} \|[\![u_h]\!]|_{L^{\infty}(e)}^p \lesssim \sum_{e \in E_h^i} h_e^{1-p|\alpha|} \|[\![u_h]\!]|_{L^p(e)}^p,$$

where for the last estimate an inverse inequality has been used. Now for (2.9) first notice that from the scaling argument one obtains $\|\nabla^{\alpha}\phi_{K}^{(\nu)}\|_{L^{p}(e)}^{p} \lesssim h_{e}^{d-1-p|\alpha|}$. Defining the set $K_{e} = \{K \in T_{h} : e \in K\}$ and proceeding as before

$$\sum_{e \in E_h^b \cap \Gamma} \int_e |\nabla^\alpha u_h - \nabla^\alpha w_h|^p dx = \sum_{e \in E_h^b \cap \Gamma} \int_e \left| \sum_{\nu \in \mathcal{N}_{K_e}} (\alpha_{K_e}^{(\nu)} - \beta^{(\nu)}) \nabla^\alpha \phi_{K_e}^{(\nu)}(x) \right|^p ds$$
$$\lesssim \sum_{e \in E_h^b \cap \Gamma} h_e^{d-1-p|\alpha|} \sum_{\nu \in \mathcal{N}_{K_e}} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p \lesssim \sum_{\nu \in \mathcal{N} \cap \Gamma} h_\nu^{d-1-p|\alpha|} \sum_{K \in \omega_\nu} |\alpha_K^{(\nu)} - \beta^{(\nu)}|^p,$$

where $h_{\nu} = \max_{e \in E_{\nu}^{b} \cap \Gamma} h_{e}$. Following the same steps as before

$$\sum_{e \in E_h^b \cap \Gamma} \int_e |\nabla^\alpha u_h - \nabla^\alpha w_h|^p dx \lesssim \sum_{e \in E_h^i} h_e^{d-1-p|\alpha|} |\llbracket u_h \rrbracket|_{L^\infty(e)}^p \lesssim \sum_{e \in E_h^i} h_e^{-p|\alpha|} |\llbracket u_h \rrbracket|_{L^p(e)}^p.$$

From Lemma 2.2 it is straightforward to prove Poincaré type inequalities for DG spaces. Similar results have been presented in [11].

Theorem 2.1 (Poincaré inequality for DG spaces.). Let $u_h \in V_h(\Omega)$ and $\Gamma \subset \partial \Omega$ with $|\Gamma| > 0$, then for all $h \leq 1$ holds

(2.10)
$$\|u_h\|_{L^p(\Omega)} \lesssim |u_h|_{W^{1,p}(\Omega)} + \|u_h\|_{L^p(\Gamma)}.$$

Proof. The proof can be found in e.g. [27].

3. Convergence of Discrete Minimisers

In this section we introduce our discontinuous Galerkin method and we show that, in the absence of convexity, our discrete energies Γ -converge to the relaxed problem. Similarly with [27], a penalty term ensures the discrete stability, but in this case it is slightly modified to include also energies which are not convex.

For $\Omega \subset \mathbb{R}^d$ a (Lipschitz) bounded domain, we consider the minimisation problem associated with the energy

(3.1)
$$\mathcal{E}(u) = \int_{\Omega} W(\nabla u(x)) \, dx$$
, where $u : \Omega \to \mathbb{R}^N$, $u = u_0$, on $\partial \Omega$.

We assume throughout that $W : \mathbb{R}^{N \times d} \to \mathbb{R}$ is of class C^1 and satisfies the following growth, coercivity, and continuity assumptions

(3.2)
$$-1 + |\xi|^p \lesssim W(\xi) \lesssim 1 + |\xi|^p$$

(3.3)
$$|W(\xi_1) - W(\xi_2)| \lesssim \left(1 + |\xi_1|^{p-1} + |\xi_1|^{p-1}\right) |\xi_1 - \xi_2|,$$

for some p > 1. Hence, we naturally pose the problem of minimising (3.1) on the Sobolev space

$$\mathbb{A}(\Omega) = \left\{ u \in W^{1,p}(\Omega)^N : u = u_0 \ \mathcal{H}^{d-1} \text{ a.e. on } \partial\Omega \right\}.$$

Due to the lack of any convexity assumption on W and hence the possible absence of minimisers of (3.1), we introduce the relaxed energy

(3.4)
$$\mathcal{E}^{\rm qc}(u) = \int_{\Omega} W^{qc}(\nabla u(x)) \, dx,$$

where $W^{qc}: \mathbb{R}^{N \times d} \to \mathbb{R}$ is the quasiconvex envelope of W, and it is defined as

$$W^{qc}(\xi) := \sup\{g(\xi) : g \le W \text{ and } g \text{ quasiconvex}\}.$$

It is known that W^{qc} satisfies (3.2) and it is quasiconvex and hence the minimisation problem

$$\min_{u\in\mathbb{A}(\Omega)}\mathcal{E}^{\rm qc}(u)$$

admits minimisers.

Employing the presented ideas in [27] for the discretisation of energy functionals in the DG setting and applying Dirichlet boundary conditions through Nitsche's approach, for $u_h \in V_h^q$ our discrete energy functional has the form (3.5)

$$\mathcal{E}_h(u_h) = \sum_{K \in T_h} \int_K W(\nabla u_h(x)) - \sum_{e \in E_h^i} \int_e S(\{\!\!\{\nabla u_h\}\!\!\}) \cdot [\!\![u_h \otimes n_e]\!\!] + \alpha \operatorname{Pen}(u_h),$$

where α will be chosen large enough and the penalty $Pen(u_h)$ is defined as

(3.6)
$$\operatorname{Pen}(u_h) := \left(1 + \sum_{K \in T_h} \int_K W(\nabla u_h) + \sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p \right)^{\frac{p-1}{p}} \times \left(\sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p \right)^{\frac{1}{p}}.$$

where for the boundary faces we use the notation

(3.7)
$$\llbracket u_h \rrbracket = u_h^- - u_0, \text{ for } e \in E_h^b = E_h \cap \partial \Omega.$$

Recall that u_0 are the boundary conditions encoded in the continuous function space $\mathbb{A}(\Omega)$ of eq. (3).

Remark 3.1. Alternatively, one may use the penalty term

(3.8)
$$\operatorname{Pen}(u_h) := \left(1 + |u_h|_{W^{1,p}(\Omega,T_h)}^p\right)^{\frac{p-1}{p}} \left(\sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p\right)^{\frac{1}{p}},$$

We remark that for the purposes of the proofs we will use the penalty term (3.8) to avoid lengthy expressions. Note that the proofs remain unaffected as, due to the growth and coercivity conditions (3.2),

$$1 + |u_h|_{W^{1,p}(\Omega,T_h)}^p \lesssim 1 + \sum_{K \in T_h} \int_K W(\nabla u_h) + \sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p \lesssim 1 + |u_h|_{W^{1,p}(\Omega,T_h)}^p.$$

3.1. Γ -convergence of the discrete energies. We may now state our main result:

Theorem 3.1. Let $W : \mathbb{R}^{N \times d} \to \mathbb{R}$ be of class C^1 , satisfying assumptions (3.2) and (3.3). Suppose that $u_h \in V_h^q(\Omega)$ is a sequence of minimisers of \mathcal{E}_h , i.e.

$$\mathcal{E}_h(u_h) = \min_{V_h^q(\Omega)} \mathcal{E}_h.$$

Then, there exists $u \in \mathbb{A}(\Omega)$ such that, up to a subsequence,

$$u_h \to u$$
 in $L^p(\Omega)$

and

$$\mathcal{E}^{\mathrm{qc}}(u) = \min_{\mathbb{A}(\Omega)} \mathcal{E}^{\mathrm{qc}}.$$

The proof of Theorem 3.1 is a direct consequence of the fact that \mathcal{E}_h Γ -converges to \mathcal{E}^{qc} with respect to the strong topology of $L^p(\Omega)$. The remainder of this section is devoted to the proof of the Γ -convergence result. We start by proving compactness in an appropriate topology for sequences of bounded energy.

Lemma 3.1. For $u_h \in V_h^q(\Omega)$ it holds that

$$\|u_h\|_{W^{1,p}(\Omega,T_h)}^p \lesssim 1 + \mathcal{E}_h(u_h).$$

In particular, if

$$\sup_h \mathcal{E}_h(u_h) \le C$$

there exists $u \in \mathbb{A}(\Omega)$ such that, up to a subsequence,

$$u_h \to u$$
 in $L^p(\Omega)$.

Proof. For $u_h \in V_h^q(\Omega)$, using assumption (3.3), we estimate the second term in the discrete energy (3.5) as follows:

$$\begin{split} &\int_{\Gamma_{\rm int}} \left| DW(\{\{\nabla_h u_h\}\}) : [\![v_h \otimes n_e]\!] \Big| ds = \int_{\Gamma_{\rm int}} \left| h^{\frac{p-1}{p}} DW(\{\{\nabla_h u_h\}\}) : h^{-\frac{p-1}{p}} [\![v_h \otimes n_e]\!] \right| \\ &\lesssim \left(\int_{\Gamma_{\rm int}} h(1 + |\{\{\nabla_h u_h\}\}|^{p-1})^{\frac{p}{p-1}} ds \right)^{\frac{p-1}{p}} \left(\int_{\Gamma_{\rm int}} h^{1-p} |[\![v_h \otimes n_e]\!]|^p ds \right)^{1/p} \\ &\lesssim \left(\int_{\Gamma_{\rm int}} h(1 + |\{\{\nabla_h u_h\}\}|^p) ds \right)^{\frac{p-1}{p}} \left(\int_{\Gamma_{\rm int}} h^{1-p} |[\![v_h \otimes n_e]\!]|^p ds \right)^{1/p}, \end{split}$$

where for the first term on the RHS we further use an inverse inequality to infer that

$$\begin{split} \int_{\Gamma_{\text{int}}} h(1+|\{\{\nabla_h u_h\}\}|^p) ds &\lesssim \int_{\Gamma_{\text{int}}} h(1+|\nabla_h u_h^+|^p+|\nabla_h u_h^-|^p) ds \\ &\lesssim \sum_{K\in\mathcal{T}_h} \int_{\partial K} h(1+|\nabla_h u_h|^p) ds \\ &\lesssim \sum_{K\in\mathcal{T}_h} \int_K 1+|\nabla_h u_h|^p dx. \end{split}$$

Hence, by Young's inequality, we infer that

(3.9)
$$\begin{aligned} \int_{\Gamma_{\text{int}}} \left| DW(\{\{\nabla_h u_h\}\}) : \llbracket v_h \otimes n_e \rrbracket \right| &\lesssim \delta^{1-p} \sum_{e \in E_h^i} h_e^{1-p} \int_e |\llbracket u_h \rrbracket|^p \\ &+ \delta \int_{\Omega} \left(1 + |\nabla_h u_h|^p \right). \end{aligned}$$

The coercivity assumption on W says that

$$W(\xi) \gtrsim -1 + |\xi|^p.$$

Thus, choosing δ appropriately as a function of p, by (3.9) we find that

$$\mathcal{E}_h(u_h) \gtrsim \int_{\Omega} |\nabla_h u_h|^p - 1 - C(p) \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p + \alpha \operatorname{Pen}(u_h).$$

Next, note that since

(3.10)
$$|u_h|_{W^{1,p}(\Omega,T_h)}^p \ge \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p$$

10

the penalty term in (3.6) satisfies

$$\begin{aligned} \operatorname{Pen}(u_h) &= \left(1 + |u_h|_{W^{1,p}(\Omega,T_h)}^p\right)^{\frac{p-1}{p}} \left(\sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p\right)^{\frac{1}{p}} \\ &\geq \left(1 + \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p\right)^{\frac{p-1}{p}} \left(\sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p\right)^{\frac{1}{p}} \\ &\geq \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p, \end{aligned}$$

where the last inequality follows from the fact that the conjugate exponents (p-1)/p and 1/p add up to 1. Then, provided that $\alpha > C(p)$, we deduce that

$$\mathcal{E}_h(u_h) \gtrsim \int_{\Omega} |\nabla_h u_h|^p - 1 + (\alpha - C(p)) \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p$$

and hence that

$$|u_h|_{W^{1,p}(\Omega,T_h)}^p \lesssim 1 + \mathcal{E}_h(u_h).$$

Furthermore, if $\mathcal{E}_h(u_h) \leq C$, $||u_h||_{L^p(\partial\Omega)}$ is uniformly bounded which implies from the Poincaré inequality (2.10) that $||u_h||_{L^p(\Omega)} \leq C'$. From the compactness of the space $W^{1,p}(\Omega, T_h)$, see [11, Theorem 5.2], we infer that there exists a function $u \in W^{1,p}(\Omega)$ such that

$$u_h \to u$$
 in $L^p(\Omega)$.

To prove $u \in \mathbb{A}(\Omega)$ note that

$$||u - u_0||_{L^p(\partial\Omega)} \le ||u_h - u_0||_{L^p(\partial\Omega)} + ||u - u_h||_{L^p(\partial\Omega)}$$

As $h \to 0$ the first term of the right hand side converge to zero due to the assumption that $\sup_h \mathcal{E}_h(u_h) \leq C$ and $||u - u_h||_{L^p(\partial\Omega)} \to 0$ from the compact embedding in [11, Lemma 8].

We next prove the liminf-inequality, providing a lower bound for the sequence $\mathcal{E}_h(u_h)$.

Lemma 3.2. Suppose that $u_h \in V_h^q(\Omega)$ and $u \in \mathbb{A}(\Omega)$ satisfy $u_h \to u$ in $L^p(\Omega)$. Then,

$$\liminf_{h \to 0} \mathcal{E}_h(u_h) \ge \mathcal{E}^{\rm qc}(u).$$

Proof. Given u_h , u as in the statement, suppose that $\liminf \mathcal{E}_h(u_h) \leq C$ as otherwise there is nothing to prove. By passing to a subsequence, we may thus assume that

$$\sup_h \mathcal{E}_h(u_h) \le C$$

and by Lemma 3.1 we have that $||u_h||_{W^{1,p}(\Omega,T_h)} \leq C$ and that

(3.11)
$$u_h \to u \text{ in } L^p(\Omega),$$

where $u \in W^{1,p}(\Omega)$. Now, from Lemma 2.2, we extract a sequence $w_h \in V_h(\Omega) \cap W^{1,p}(\Omega)$ such that

(3.12)
$$\int_{\Omega} |u_h - w_h|^p \lesssim \sum_{e \in E_h^i} h_e \int_e |\llbracket u_h \rrbracket|^p ds,$$

(3.13)
$$\int_{\Omega} |\nabla_h u_h - \nabla w_h|^p \lesssim \sum_{e \in E_h^i} h_e^{1-p} \int_e |\llbracket u_h \rrbracket|^p ds,$$

and so combined with (3.11) we get that $w_h \to u$ in $L^p(\Omega)$. In addition to this, since

(3.14)
$$\int_{\Omega} |\nabla w_h|^p \lesssim \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p + \int_{\Omega} |\nabla_h u_h|^p \lesssim |u_h|_{W^{1,p}(\Omega,\mathcal{T}_h)}^p,$$

we infer that (∇w_h) is bounded in $L^p(\Omega)$ and hence there exists $G \in L^p(\Omega)$ such that $\nabla w_h \rightharpoonup G$ in $L^p(\Omega)$. However, the strong convergence of w_h in $L^p(\Omega)$ and the distributional continuity of the operator $\nabla : L^p(\Omega) \rightarrow L^p(\Omega)$, give us that $G = \nabla u$ and so

$$\nabla w_h \rightharpoonup \nabla u$$
, in $L^p(\Omega)$.

Note that since W^{qc} is quasiconvex and $W \ge W^{qc}$, it holds that (3.15)

$$\liminf_{h \to 0} \int_{\Omega} W(\nabla w_h) \ge \liminf_{h \to 0} \int_{\Omega} W^{qc}(\nabla w_h) \ge \int_{\Omega} W^{qc}(\nabla u) = \mathcal{E}^{qc}(u).$$

We thus add and subtract the term

$$\int_{\Omega} W(\nabla w_h)$$

in the discrete energy \mathcal{E}_h to find that

$$\mathcal{E}_{h}(u_{h}) = \int_{\Omega} W(\nabla w_{h}) + \int_{\Omega} W(\nabla_{h} u_{h}) - W(\nabla w_{h})$$
$$- \int_{\Gamma_{\text{int}}} DW(\{\{\nabla_{h} u_{h}\}\}) : \llbracket v_{h} \otimes n_{e} \rrbracket + \alpha \operatorname{Pen}(u_{h})$$
$$=: I_{h} + II_{h} + III_{h} + \alpha \operatorname{Pen}(u_{h}).$$

By (3.15) it suffices to show that

(3.16)
$$\liminf_{h \to 0} \left[II_h + III_h + \alpha \operatorname{Pen}(u_h) \right] \ge 0.$$

Regarding the term III_h we argue similarly to Lemma 3.1 to get that

$$|III_h| \lesssim \left(\sum_{K \in \mathcal{T}_h} \int_K 1 + |\nabla_h u_h|^p dx\right)^{1/p'} \left(\int_{\Gamma_{\text{int}}} h^{1-p} |\llbracket v_h \otimes n_e \rrbracket|^p ds\right)^{1/p},$$

which in turns implies that

(3.17)
$$|III_{h}| \leq C \left(1 + |u_{h}|_{W^{1,p}(\Omega,T_{h})}^{p}\right)^{\frac{p-1}{p}} \left(\sum_{e \in E_{h}^{i}} \frac{1}{h_{e}^{p-1}} \int_{e} |\llbracket u_{h}]|^{p}\right)^{\frac{1}{p}} \leq C \operatorname{Pen}(u_{h}).$$

Next, we estimate the term II_h . By Hölder's inequality, (3.13), (3.14) and (3.3) we find that (3.18)

$$\begin{aligned} |II_{h}| \lesssim \int_{\Omega} \left(1 + |\nabla_{h}u_{h}|^{p-1} + |\nabla w_{h}|^{p-1} \right) |\nabla_{h}u_{h} - \nabla w_{h}| \\ \lesssim \left[\int_{\Omega} \left(1 + |\nabla_{h}u_{h}|^{p-1} + |\nabla w_{h}|^{p-1} \right)^{\frac{p}{p-1}} \right]^{\frac{p-1}{p}} \left(\int_{\Omega} |\nabla_{h}u_{h} - \nabla w_{h}|^{p} \right)^{\frac{1}{p}} \\ \lesssim \left[\int_{\Omega} \left(1 + |\nabla_{h}u_{h}|^{p} + |\nabla w_{h}|^{p} \right) \right]^{\frac{p-1}{p}} \left(\sum_{e \in E_{h}^{i}} \frac{1}{h_{e}^{p-1}} \int_{e} |[u_{h}]|^{p} \right)^{\frac{1}{p}} \\ \lesssim \left(1 + |u_{h}|^{p}_{W^{1,p}(\Omega,T_{h})} \right)^{\frac{p-1}{p}} \left(\sum_{e \in E_{h}^{i}} \frac{1}{h_{e}^{p-1}} \int_{e} |[u_{h}]|^{p} \right)^{\frac{1}{p}} \\ \leq \tilde{C} \operatorname{Pen}(u_{h}). \end{aligned}$$

Combining the latter estimate with (3.17), we get that

$$II_h + III_h + \alpha \operatorname{Pen}(u_h) \ge \left(\alpha - C - \tilde{C}\right) \operatorname{Pen}(u_h)$$

and choosing $\alpha > C + \tilde{C}$ we infer (3.16) which concludes the proof.

The remaining ingredient in the proof of Theorem 3.1 is providing a recovery sequence which is given in the following lemma.

Lemma 3.3. For every $u \in \mathbb{A}(\Omega)$ if $q \ge 1$ and h < 1 there exist $u_h \in V_h^q(\Omega)$ such that $u_h \to u$ in $L^p(\Omega)$ and

$$\limsup_{h \to 0} \mathcal{E}_h(u_h) \le \mathcal{E}^{\rm qc}(u).$$

Proof. First, similarly to [27][Lemma 3.6], we show that for all $v \in \mathbb{A}(\Omega)$ there exists a recovery sequence $(u_h) \subset V_h^q$ such that

(3.19)
$$||v - u_h||_{W^{1,p}(\Omega,T_h)}^p + \sum_{e \in E_h^b} \frac{1}{h_e^{p-1}} \int_e |u_h - u_0|^p \to 0, \text{ as } h \to 0.$$

and

(3.20)
$$\lim_{h \to 0} \mathcal{E}_h(u_h) = \mathcal{E}(v).$$

First note that $C^{\infty}(\overline{\Omega})$ is dense in $W^{1,p}(\Omega)$ and one can create a sequence $(u_{\delta}) \subset C^{\infty}(\overline{\Omega})$ with the properties

(3.21)
$$||v - u_{\delta}||_{W^{1,p}(\Omega)} \lesssim \delta \quad \text{and} \quad |u_{\delta}|_{W^{2,p}(\Omega)} \lesssim \frac{1}{\delta} |v|_{W^{1,p}(\Omega)}$$

Choosing the sequence $u_{h,\delta} = I_h^q u_\delta$, $I_h^q : W^{s,p}(\Omega) \to V_h^q(\Omega)$ is the standard nodal interpolation operation, we recall the error estimates for all $q \ge 1$

(3.22)
$$|u_{\delta} - u_{h,\delta}|_{W^{i,p}(K)} \lesssim h_{K}^{2-i} |u_{\delta}|_{W^{2,p}(K)},$$

(3.23)
$$|u_{\delta} - u_{h,\delta}|_{L^{p}(e)} \lesssim h_{K}^{2-\frac{1}{p}} |u_{\delta}|_{W^{2,p}(K)}.$$

In the following, assuming that h < 1, the above error estimates are combined with eq. (3.21)

$$(3.24) \quad \sum_{K \in T_{h}} |u_{\delta} - u_{h,\delta}|_{W^{i,p}(K)}^{p} \lesssim h^{p} |u_{\delta}|_{W^{2,p}(\Omega)}^{p} \lesssim \frac{h^{p}}{\delta^{p}} |v|_{W^{1,p}(\Omega)}^{p},$$

$$(3.25) \quad \sum_{e \in E_{h}} \frac{1}{h_{e}^{p-1}} || \llbracket u_{\delta} - u_{h,\delta} \rrbracket ||_{L^{p}(e)}^{p} \lesssim h^{p} \sum_{e \in E_{h}} |u_{\delta}|_{W^{2,p}(K_{e})}^{p} \lesssim \frac{h^{p}}{\delta^{p}} |v|_{W^{1,p}(\Omega)}^{p},$$

here K_e denotes the set of the elements containing $e, e \in E_h$. For the boundary terms applying Dirichlet boundary conditions through Nitsche's approach, the above estimates together with eq. (3.21) imply

(3.26)
$$\sum_{e \in E_h^b} \frac{1}{h_e^{p-1}} \int_e |u_{h,\delta} - u_0|^p \lesssim \sum_{e \in E_h^b} \frac{1}{h_e^{p-1}} \int_e (|u_{h,\delta} - u_\delta|^p + |u_\delta - v|^p) \\ \lesssim \left(\frac{h^p}{\delta^p} + \frac{\delta^p}{h^{p-1}}\right) ||v||_{W^{1,p}(\Omega)}^p.$$

For $\beta \in (\frac{p-1}{p}, 1)$, choosing $\delta = h^{\beta}$ equations (3.24), (3.25) and (3.26) $\rightarrow 0$ as $h \rightarrow 0$. Here $v \in W^{1,p}(\Omega)$ (u_{δ} is continuous by construction), which means $\llbracket v \rrbracket = 0$ on every $e \in E_h^i$ and as $h \rightarrow 0$

$$(3.27) ||v - u_{h,\delta}||_{W^{1,p}(\Omega,T_h)}^p \le ||v - u_{\delta}||_{W^{1,p}(\Omega)} + ||u_{\delta} - u_{h,\delta}||_{W^{1,p}(\Omega,T_h)}^p \to 0.$$

Therefore, we found a sequence in $V_h^q(\Omega)$ such that eq. (3.19) holds.

Next, for the convergence of the discrete functional, eq. (3.20) and considering again the sequence (w_h) of Lemma 2.2, we note that

(3.28)
$$\nabla w_h \to \nabla v \text{ in } L^p(\Omega).$$

Indeed, recalling that $\llbracket v \rrbracket = 0$, we find that

$$\begin{split} \int_{\Omega} |\nabla w_h - \nabla v|^p &\lesssim \int_{\Omega} |\nabla_h u_h - \nabla v|^p + \int_{\Omega} |\nabla_h u_h - \nabla w_h|^p \\ &\lesssim \int_{\Omega} |\nabla_h u_h - \nabla v|^p + \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h] |^p \\ &= \sum_K \int_K |\nabla_h u_h - \nabla v|^p + \sum_{e \in E_h^i} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h - v] |^p \\ &= |u_h - v|_{W^{1,p}(\Omega,T_h)}^p \to 0. \end{split}$$

As in Lemma 3.2, we can write the discrete energy as

$$\mathcal{E}_{h}(u_{h}) = \int_{\Omega} W(\nabla w_{h}) + \int_{\Omega} W(\nabla_{h}u_{h}) - W(\nabla w_{h})$$
$$- \int_{\Gamma_{\text{int}}} DW(\{\{\nabla_{h}u_{h}\}\}) : \llbracket u_{h} \otimes n_{e} \rrbracket + \alpha \operatorname{Pen}(u_{h})$$
$$=: I_{h} + II_{h} + III_{h} + \alpha \operatorname{Pen}(u_{h}).$$

Vitali's convergence theorem, (3.28), and the growth of W say that

(3.29)
$$I_h = \int_{\Omega} W(\nabla w_h) \to \int_{\Omega} W(\nabla v) = \mathcal{E}(v) \text{ as } h \to 0,$$

and since the boundary terms converge to zero, eq. (3.19), it remains to show that II_h , III_h as well as $Pen(u_h)$ all vanish in the limit $h \to 0$. Using the estimates (3.17) and (3.18), we find that

$$|II_h| + |III_h| \lesssim \operatorname{Pen}(u_h)$$

and it thus suffices to prove that

$$\limsup_{h \to 0} \operatorname{Pen}(u_h) = 0.$$

Indeed, since $|u_h - v|_{W^{1,p}(\Omega,T_h)} \to 0$, it follows that

$$|u_h|_{W^{1,p}(\Omega,T_h)} \le C$$

and hence

$$\operatorname{Pen}(u_{h}) \lesssim \left(\sum_{e \in E_{h}} \frac{1}{h_{e}^{p-1}} \int_{e} |\llbracket u_{h} \rrbracket|^{p}\right)^{\frac{1}{p}} \lesssim \left(\sum_{e \in E_{h}} \frac{1}{h_{e}^{p-1}} \int_{e} |\llbracket u_{h} - v]\|^{p}\right)^{\frac{1}{p}}$$
$$\lesssim \left(|u_{h} - v|_{W^{1,p}(\Omega,T_{h})} + \sum_{e \in E_{h}^{b}} \frac{1}{h_{e}^{p-1}} \int_{e} |\llbracket u_{h} - v]\|^{p}\right)^{\frac{1}{p}} \to 0.$$

This completes the proof of (3.20).

Let now $u \in \mathbb{A}(\Omega)$, then from [21][Theorem 9.1] the exists a sequence $(u_k) \subset \mathbb{A}(\Omega)$ such that

$$u_k \to u$$
, in $L^p(\Omega)$ and $\mathcal{E}_k(u_k) \to \mathcal{E}^{\mathrm{qc}}(u)$,

for $k \to \infty$. Using the first part of our proof, for each $k \in \mathbb{N}$, we can further extract a sequence $u_{k,h} \subset V_h^q$ such that

$$u_{k,h} \to u_k$$
, in $L^p(\Omega)$, and $\mathcal{E}_h(u_{k,h}) \to \mathcal{E}(u_k)$,

for $h \to 0$. Hence, combining the above convergences we have that

$$\lim_{k \to \infty} \lim_{h \to 0} \mathcal{E}_h(u_{k,h}) = \lim_{k \to \infty} \mathcal{E}(u_k) = \mathcal{E}^{\mathrm{qc}}(u),$$

and

$$\lim_{k \to \infty} \lim_{h \to 0} \|u_{k,h} - u\|_{L^p(\Omega)} = 0.$$

The latter implies that we can choose a subsequence (k_h) such that for $u_{k_h,h} \subset V_h^q$ it holds that

$$\lim_{h \to 0} \mathcal{E}_h(u_{k_h,h}) = \mathcal{E}^{\mathrm{qc}}(u), \text{ and,}$$
$$\lim_{h \to 0} \|u_{k_h,h} - u\|_{L^p(\Omega)} = 0,$$

which concludes the proof of our Lemma.

We conclude by demonstrating the proof of Theorem 3.1.

Proof of Theorem 3.1. Suppose that $u_h \in V_h^q$ is given such that

$$\mathcal{E}_h(u_h) = \inf_{V_h^q} \mathcal{E}_h$$

Note that for any $v \in \mathbb{A}(\Omega)$, by Lemma 3.3 we can find a sequence $v_h \in V_h^q$ such that

$$v_h \to v \text{ in } L^p(\Omega)$$

and $\limsup \mathcal{E}_h(v_h) \leq \mathcal{E}^{\mathrm{qc}}(v)$. In particular, up to a subsequence,

$$\mathcal{E}_h(u_h) \leq \mathcal{E}_h(v_h) \leq 1 + \limsup \mathcal{E}_h(v_h) \leq 1 + \mathcal{E}^{\mathrm{qc}}(v) < \infty.$$

Hence, by Lemma 3.1, there exists $u \in \mathbb{A}(\Omega)$ such that

$$u_h \to u$$
 in $L^p(\Omega)$

and, for any $v \in \mathbb{A}(\Omega)$,

$$\mathcal{E}^{\mathrm{qc}}(u) \leq \liminf_{h \to 0} \mathcal{E}_h(u_h) \leq \limsup_{h \to 0} \mathcal{E}_h(v_h) \leq \mathcal{E}^{\mathrm{qc}}(v),$$

where v_h is the recovery sequence for v whose existence is guaranteed by Lemma 3.3. That is,

$$\mathcal{E}^{\rm qc}(u) = \inf_{\mathbb{A}(\Omega)} \mathcal{E}^{\rm qc}$$

completing the proof.

3.2. **DG methods based on discrete gradients.** We start with a definition of discrete gradients in the discontinuous Galerkin setting. To this end, define first the lifting operator $R_h: T(E_h) \to V_h^{q-1}(\Omega)$ as

(3.30)
$$\int_{\Omega} R_h(\varphi) : w_h \, dx = \sum_{e \in E_h^i} \int_e \{\!\!\{w_h\}\!\!\} : [\!\![\varphi \otimes n_e]\!] \, ds, \quad \forall w_h \in V_h^{q-1}(\Omega).$$

This operator leads to the definition of the discrete gradient G_h for $u_h \in V_h^q$ as

(3.31)
$$G_h(u_h) = \nabla_h u_h - R_h(u_h).$$

The method of Ten Eyck and Lew (2006) [38] is based on minimising over the DG space the functional defined on V_h^q :

(3.32)
$$\mathcal{E}_{G,h}[u_h] = \sum_{K \in T_h} \int_K W(\nabla u_h(x) - R_h(u_h)) + \alpha \operatorname{Pen}(u_h)$$
$$= \sum_{K \in T_h} \int_K W(G_h(u_h)) + \alpha \operatorname{Pen}(u_h).$$

Convergence in the case of convex W was derived in [11] and error estimates for smooth solutions were derived in [35]. The evaluation of this functional requires the computation of $R_h(u_h)$, and thus of the "discrete gradient" $G_h(u_h)$ which appears in W. Computationally, the cost of this operation is reasonable, since it involves the solution of local systems. In the convex case, the weak convergence properties of the discrete gradient were enough to establish lim-inf inequalities and conclude the proof of convergence of minimisers. In our setting this is no longer the case. As in the previous section the key idea is still to use Lemma 2.2 and to extract a sequence $w_h \in V_h(\Omega) \cap W^{1,p}(\Omega)$ such that (3.12), (3.13) will hold. Then, (3.11) will imply that $w_h \to u$ in $L^p(\Omega)$. Repeating the same arguments as in Lemma 3.2, we conclude (3.33)

$$\liminf_{h \to 0} \int_{\Omega} W(\nabla w_h) \ge \liminf_{h \to 0} \int_{\Omega} W^{qc}(\nabla w_h) \ge \int_{\Omega} W^{qc}(\nabla u) = \mathcal{E}^{qc}(u).$$

And thus it suffices to show that

(3.34)
$$\liminf_{h \to 0} \left[A_h + \alpha \operatorname{Pen}(u_h) \right] \ge 0.$$

where

$$\mathcal{E}_{\mathcal{G},h}(u_h) = \int_{\Omega} W(\nabla w_h) + \int_{\Omega} W(G_h(u_h)) - W(\nabla w_h) + \alpha \operatorname{Pen}(u_h)$$
$$=: \int_{\Omega} W(\nabla w_h) + A_h + \alpha \operatorname{Pen}(u_h).$$

To estimate A_h we shall use the known bound for the lifting operator, [11, 22],

(3.35)
$$\int_{\Omega} |R_h(v_h)|^p \le C_R \sum_{e \in E_h^i} h_e^{1-p} \int_e |\llbracket v_h \rrbracket|^p, \qquad v_h \in V_h^q(\Omega),$$

where the constant C_R is independent of h. A consequence of this bound, see [27], is that for all $v_h \in V_h^q(\Omega)$ it holds that

(3.36)
$$\|G_h(v_h)\|_{L^p(\Omega)} \le C |v_h|_{W^{1,p}(\Omega,T_h)}.$$

Next, proceeding as in Lemma 3.2, we conclude

$$\begin{aligned} |A_{h}| &\lesssim \int_{\Omega} \left(1 + |G_{h}(u_{h})|^{p-1} + |\nabla w_{h}|^{p-1} \right) |G_{h}(u_{h}) - \nabla w_{h}| \\ &\lesssim \left[\int_{\Omega} \left(1 + |G_{h}(u_{h})|^{p-1} + |\nabla w_{h}|^{p-1} \right)^{\frac{p}{p-1}} \right]^{\frac{p-1}{p}} \left(\int_{\Omega} |G_{h}(u_{h}) - \nabla w_{h}|^{p} \right)^{\frac{1}{p}} \\ &\lesssim \left[\int_{\Omega} \left(1 + |G_{h}(u_{h})|^{p} + |\nabla w_{h}|^{p} \right) \right]^{\frac{p-1}{p}} \left[\left(\int_{\Omega} |\nabla_{h}u_{h} - \nabla w_{h}|^{p} \right)^{\frac{1}{p}} + \left(\int_{\Omega} |R_{h}(v_{h})|^{p} \right)^{\frac{1}{p}} \right] \\ &\lesssim \left(1 + |u_{h}|^{p}_{W^{1,p}(\Omega,T_{h})} \right)^{\frac{p-1}{p}} \left(\sum_{e \in E_{h}^{i}} \frac{1}{h_{e}^{p-1}} \int_{e} |\llbracket u_{h}] |^{p} \right)^{\frac{1}{p}} \\ &\leq \tilde{C} \operatorname{Pen}(u_{h}). \end{aligned}$$

As a result,

$$A_h + \alpha \operatorname{Pen}(u_h) \ge \left(\alpha - \tilde{C}\right) \operatorname{Pen}(u_h)$$

Thus, if $\alpha > \tilde{C}$, (3.34) follows. The lim-sup inequality follows by adopting in similar fashion arguments from the previous section. We therefore obtain the following theorem.

Theorem 3.2. Let $W : \mathbb{R}^{N \times d} \to \mathbb{R}$ be of class C^1 , satisfying assumptions (3.2) and (3.3). Suppose that $u_h \in V_h^q(\Omega)$ is a sequence of minimisers of $\mathcal{E}_{G,h}$, i.e.

$$\mathcal{E}_{\mathrm{G},h}(u_h) = \min_{V_h^q(\Omega)} \mathcal{E}_{\mathrm{G},h}.$$

Then, there exists $u \in \mathbb{A}(\Omega)$ such that, up to a subsequence,

$$u_h \to u$$
 in $L^p(\Omega)$

and

$$\mathcal{E}^{\rm qc}(u) = \min_{\mathbb{A}(\Omega)} \mathcal{E}^{\rm qc}.$$

4. Computational Results

In this section it is illustrated how the proposed numerical scheme approximates minimisers for the total potential energy when polyconvex (and hence quasiconvex) and multi-well strain energy functions are employed. In all computations we use the scheme described in Section 3, see (3.5). For the first case we compare our scheme with a generalisation of the standard L^2 interior penalty method for convex energies, [27], while for the latter case we show how a sequence of discrete minimisers is formed which converges to the minimiser of the relaxed problem.

In the sequel $y : \mathbb{R}^2 \supset \Omega \rightarrow \Omega^*$ will denote plane deformations from Ω (undeformed body) to $\Omega^* \subset \mathbb{R}^2$ (deformed configuration). Under given boundary conditions we seek deformations that minimise the total potential energy

(4.1)
$$\min_{\substack{y \in W^{1,p}(\Omega) \\ y = y_0 \text{ on } \partial\Omega}} \mathcal{E}(y) = \min_{\substack{y \in W^{1,p}(\Omega) \\ y = y_0 \text{ on } \partial\Omega}} \int_{\Omega} W(\nabla y) dx,$$

where W is the strain energy function characterising the material properties.

Examples with a polyconvex strain energy function. We would like to approximate minimisers of eq. (4.1) through the discretised energy (3.5) for the polyconvex strain energy function $W(F) = |\det F|^2$. We compare the proposed penalty of eq. (3.6) with an appropriate penalty for convex energies, [27], given by

(4.2)
$$\operatorname{Pen}(u_h) := \left(1 + |u_h|_{W^{1,p}(\Omega,T_h)}^{p-2}\right) \left(\sum_{e \in E_h} \frac{1}{h_e^{p-1}} \int_e |\llbracket u_h \rrbracket|^p\right)^{\frac{2}{p}},$$

which for p = 2 reduces to the standard L^2 error. In eq. (4.1) we choose $y_0(x) = F_0 x$ where

(4.3)
$$F_0 = \begin{pmatrix} 1 & 0 \\ 0 & 0.9 \end{pmatrix}.$$

corresponding to uniaxial compression. Here y_0 is a homogeneous deformation, therefore polyconvexity implies that

(4.4)
$$\int_{\Omega} W(\nabla y) \ge \int_{\Omega} W(\nabla y_0) \text{ for all } y \in W^{1,4}(\Omega) \text{ with } y = y_0 \text{ on } \partial\Omega,$$

which means that every admissible deformation results in a potential energy no less than that of y_0 . We seek discrete minimisers in $V_h^1(\Omega)$, i.e. piecewise polynomials of first degree. The structure of the described minimisation problem indicates that for big enough stabilisation parameter α in eq. (3.5), the computed minimiser should satisfy inequality (4.4). As it is illustrated in fig. 1 where the blue curve approaches the black dashed line, large values of the stabilisation parameter α are required to approach $\int_{\Omega} W(\nabla y_0)$ when the penalty of eq. (4.2) is used. On the contrary for the penalty of eq. (3.6), even small values of α bring the discrete minimiser close to $W^{1,4}(\Omega)$, i.e. the orange curve is always close to the black dashed line. To examine further this behaviour we plot $1/\det \nabla y$ in the deformed configuration, fig. 2. For both penalties y_h approaches y_0 as α increases but for the term (4.2) large values of α are required while for (3.6) even small values produce a minimiser close to y_0 . These observations are quantified in fig. 3 where we compare the $L^1(\Omega)$ and $W^{1,1}(\Omega)$ errors between y_0 and the computed minimisers y_h varying the penalty parameter α .

Non rank-one convex energies and minimising sequences. We construct a frame indifferent two-well strain energy function where under specific boundary conditions a phase mixture emerges with finer and finer microstructure, approaching the infimum of the total potential energy as we decrease the mesh size h. Mimicking the cubic to orthorhombic phase transitions in crystals we define the energy $W(F) = |C - V^2||C - I|^2$, where $C = F^T F$ is the right Cauchy-Green tensor, I is the identity and

(4.5)
$$V = \begin{pmatrix} \frac{a_o + b_o}{2} & \frac{b_o - a_o}{2} \\ \frac{b_o - a_o}{2} & \frac{a_o + b_o}{2} \end{pmatrix}.$$

Choosing $a_0 = \sqrt{2 - b_0^2}$ and $b_0 = 0.9$, equation $RV - I = d \otimes n$ has two solutions namely $\{R_1, d_1, n_1\}$, $\{R_2, d_2, n_2\}$ such that $n_1 \approx (1, 0)$ and $n_2 \approx$ (0, 1), where $R_i \in SO(2)$, see [4, Proposition 4]. This essentially means, that there exist continuous deformations with discontinuous deformation gradients across the planes defined by n_1 and n_2 . Note that W is zero when the deformation gradient lies at SO(2)V or at SO(2)I. Now we define y_0 in eq.(4.1) such that

(4.6)
$$\nabla y_0 = 0.5I + 0.5R_1V,$$

and we initialise y_h by the interpolation $y_h = I_h^1 y_0$, where $I_h^1 : W^{1,p}(\Omega) \to$ $V_h^1(\Omega)$. Discretising the domain through equal squares with normals n_1 and n_2 , where each rectangle is divided to four triangles, perfect interfaces between discontinuous deformation gradients (phases) can occur due to this triangulation. For a comparison of meshes with or without the above property see [25, Section 4.5]. Minimising the discrete version of eq. (4.1) the total potential energy and $||y_h - y_0||_{L^2(\Omega)}$ decrease for smaller mesh sizes, fig. 4. The energy is reduced through a minimising sequence as it is depicted in fig. 5, describing the deformed state. In some regions perfect interfaces with zero contribution are formed between $\nabla y_h = I$ and $\nabla y_h = R_1 V$ across the direction $n_1 = (1,0)$ and between $\nabla y_h = I$ and $\nabla y_h = R_2 V$ across the direction $n_2 = (0, 1)$. These regions illustrate the minimising sequence in the reference configuration, see fig. 6. Note that R_1V is not rank-one connected to R_2V , $R_1 \neq R_2$. Hence, a transition layer is formed between these two deformations, green color in fig. 6. This is the only non-zero part of the energy. As finer and finer microstructures appear on each side of a green curve $y_h \to y_0$ in $L^{\infty}(\Omega)$ as $h \to 0$, which means a "macroscopic" deformation is formed equal to y_0 and the energy at the transition layer tends to zero as $h \to 0$, see also fig. 7. In this case, the discrete minimiser captures a minimising sequence $\{y_k\} \in W^{1,\infty}(\Omega)$ such that $y_k \stackrel{*}{\rightharpoonup} y_0$ in $W^{1,\infty}(\Omega)$ where $\{\nabla y_k\}$ generates the homogeneous Young measure $\nu_x = 0.5\delta_I + 0.5\delta_V$. Note that y_0 is the minimiser of the relaxed problem.

The above simulations have been performed by choosing $\alpha = 80$ in eq. (3.5). Note that, for the proposed multi-well energy

$$-1 + |F|^8 \lesssim W(F) \lesssim 1 + |F|^8.$$




FIGURE 1. Total potential energy (Vertical axis) for the homogeneous deformation $y_0(x) = F_0 x$ (dashed black line), for the computed minimiser with respect to the stabilisation parameter α employing the penalty of eq. 4.2 (blue squares) and the penalty of eq. 3.6 (orange circles).

We have noticed that when p = 8 roundoff errors affects the discrete minimisation process. Note that the second term of the proposed penalty, eq.(3.6), is getting less important numerically, due to the fact that the power of 1/8appears and small numbers are described by finite precision. One can rewrite the jump terms of eq. (3.5) as

(4.7)
$$jumps = \sum_{e \in E_h} h_e \int_e \left| \frac{\llbracket u_h \rrbracket}{h_e} \right|^p ds,$$

and the penalty term as

(4.8)
$$\alpha \operatorname{Pen}(u_h) = \frac{\alpha}{2} \left(\int_{\Omega} 1 + W(\nabla u_h) dx + jumps \right)^{\frac{p-1}{p}} (2^p jumps)^{\frac{1}{p}}.$$

Under the above modifications we have obtained minimisers that capture minimising sequences of the continuous problem. The above minimising sequence appears for both $y_h \in V_h^q(\Omega) \cap C^0(\Omega)$ and $y_h \in V_h^q(\Omega)$, where both converge to the minimiser of the relaxed problem, since the proved convergence results hold for conforming finite element spaces as well. DG FOR NON-CONVEX ENERGIES




FIGURE 2. Density $(1/\det \nabla y)$ in the deformed configuration under uniaxial compression (10% strain). Employing the penalty of eq. (4.2), the density for the computed minimisers are illustrated: (a) when $\alpha = 20$ and (b) when $\alpha = 160$ (see blue squares of Figs. 3a and 3b at $\alpha = 20, 160$). (c): Computed solution when the proposed penalty of eq.(3.6) is used for $\alpha = 20$. (d) Density of the exact homogeneous minimiser y_0 .




FIGURE 3. Comparing the accuracy of the two proposed stabilisation penalty terms (3.6) (circles) and (4.2) (squares) for the polyconvex $W(F) = |\det F|^2$ strain energy under 10% uniaxial compression. Horizontal axis: values of the penalty parameter α . Vertical axes: $|y_h - y_0|_{L^1(\Omega)}$ and $|y_h - y_0|_{W^{1,1}(\Omega)}$ errors for various mesh resolutions, specifically for 1024 (blue), 2034 (green) and 4096 (orange) triangles, where y_h denote the numerical solutions and y_0 the homogeneous minimiser. In (a) circular error belongs in the range of 10^{-8} and 10^{-9} , while in (b) circular errors are of the order 10^{-6} .




FIGURE 4. Imposing the homogeneous boundary conditions $y_0(x) = (0.5R_1V + 0.5I)x$, minimisers are computed varying the mesh resolution (horizontal axis). Blue curve denotes the total potential energy (scale of left vertical axis) and red dashed curve is the error $||y_h - y_0||_{L^2(\Omega)}$ (scale of right vertical axis) where y_h denote the computed minimiser.




FIGURE 5. The maximum eigenvalue (λ_{max}) in the deformed configuration of the computed minimiser for each triangle. The wells I, V have the eigenvalues (1, 1) and 0.9, 1.09871212respectively. In blue regions $\lambda_{max} = 1$ which indicates that well I is attained up to a rotation. Similarly, red color, $\lambda_{max} = 1.09871212$, corresponds to $R_i V$ up to a rotation, i = 1, 2.




FIGURE 6. The maximum eigenvalue (λ_{max}) of the discrete minimiser for each triangle expressed in the reference configuration. The wells I, V have the eigenvalues (1, 1) and 0.9, 1.09871212 respectively. In blue regions $\lambda_{max} = 1$ which indicates the well I appears up to a rotation. Similarly, in red color $\lambda_{max} = 1.09871212$ correspond to $R_i V$ up to a rotation, where i=1 when the interface between the two phases is normal to $n_1 = (1, 0)$ and i=2 when the interface is normal to $n_2 = (0, 1)$.




FIGURE 7. Displacement of the computed minising sequence varying the mesh resolution. (a): Pointwise values of $|u| = \sqrt{u \cdot u}$, $u : \Omega \to \mathbb{R}^2$ is the displacement vector. For 200, 800, and 32768 triangles the values of |u| along the diagonal of the domain (white line in (a)) are illustrated in (b), (c) and (d) respectively.

References

- A. K. Balci, C. Ortner, and J. Storn. "Crouzeix-Raviart finite element method for non-autonomous variational problems with Lavrentiev gap". Numer. Math. 151.4 (2022), pp. 779–805. URL: https://doi.org/10.1007/s00211-022-01303-1.
- J. M. Ball. "Singularities and computation of minimizers for variational problems". Foundations of computational mathematics (Oxford, 1999). Vol. 284. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 2001, pp. 1–20.
- J. M. Ball and G. Knowles. "A numerical method for detecting singular minimizers". Numer. Math. 51.2 (1987), pp. 181–197. URL: https: //doi.org/10.1007/BF01396748.
- [4] J. M. Ball and R. D. James. "Fine phase mixtures as minimizers of energy". Analysis and Continuum Mechanics. Springer, 1989, pp. 647– 686.
- S. Bartels, C. Carstensen, K. Hackl, and U. Hoppe. "Effective relaxation for microstructure simulations: algorithms and applications". *Comput. Methods Appl. Mech. Engrg.* 193.48-51 (2004), pp. 5143–5175. URL: https://doi.org/10.1016/j.cma.2003.12.065.
- S. Bartels. Numerical methods for nonlinear partial differential equations. Vol. 47. Springer Series in Computational Mathematics. Springer, Cham, 2015, pp. x+393. URL: https://doi.org/10.1007/978-3-319-13797-1.
- S. Bartels. "Reliable and efficient approximation of polyconvex envelopes". SIAM J. Numer. Anal. 43.1 (2005), pp. 363–385. URL: https://doi.org/10.1137/S0036142903428840.
- [8] S. Bartels. "Total variation minimization with finite elements: convergence and iterative solution". SIAM J. Numer. Anal. 50.3 (2012), pp. 1162–1180. URL: https://doi.org/10.1137/11083277X.
- S. Bartels and A. Kaltenbach. "Explicit and efficient error estimation for convex minimization problems". *Math. Comp.* 92.343 (2023), pp. 2247-2279. URL: https://doi.org/10.1090/mcom/3821.
- [10] A. Bonito, R. H. Nochetto, and D. Ntogkas. "DG approach to large bending plate deformations with isometry constraint". *Math. Models Methods Appl. Sci.* 31.1 (2021), pp. 133–175. URL: https://doi.org/ 10.1142/S0218202521500044.

REFERENCES

- [11] A. Buffa and C. Ortner. "Compact embeddings of broken Sobolev spaces and applications". *IMA journal of numerical analysis* 29.4 (2009), pp. 827–855.
- [12] C. Carstensen. "Numerical analysis of microstructure". Theory and numerics of differential equations (Durham, 2000). Universitext. Springer, Berlin, 2001, pp. 59–126.
- C. Carstensen. "Ten remarks on nonconvex minimisation for phase transition simulations". Comput. Methods Appl. Mech. Engrg. 194.2-5 (2005), pp. 169–193. URL: https://doi.org/10.1016/j.cma.2004.05.012.
- [14] C. Carstensen and P. Plecháč. "Adaptive algorithms for scalar nonconvex variational problems". Proceedings of the International Centre for Mathematical Sciences Conference on Grid Adaptation in Computational PDEs: Theory and Applications (Edinburgh, 1996). Vol. 26.
 1-2. 1998, pp. 203–216. URL: https://doi.org/10.1016/S0168-9274(97)00089-5.
- [15] C. Carstensen and P. Plecháč. "Numerical analysis of compatible phase transitions in elastic solids". SIAM J. Numer. Anal. 37.6 (2000), pp. 2061– 2081. URL: https://doi.org/10.1137/S0036142998337697.
- C. Carstensen and P. Plecháč. "Numerical solution of the scalar doublewell problem allowing microstructure". *Math. Comp.* 66.219 (1997), pp. 997–1026. URL: https://doi.org/10.1090/S0025-5718-97-00849-1.
- C. Carstensen and T. Roubíček. "Numerical approximation of Young measures in non-convex variational problems". *Numer. Math.* 84.3 (2000), pp. 395–415. URL: https://doi.org/10.1007/s002110050003.
- [18] C. Collins, D. Kinderlehrer, and M. Luskin. "Numerical approximation of the solution of a variational problem with a double well potential". *SIAM J. Numer. Anal.* 28.2 (1991), pp. 321–332. URL: https://doi. org/10.1137/0728018.
- [19] C. Collins and M. Luskin. "Optimal-order error estimates for the finite element approximation of the solution of a nonconvex variational problem". *Math. Comp.* 57.196 (1991), pp. 621–637. URL: https:// doi.org/10.2307/2938708.
- [20] M. Crouzeix and P.-A. Raviart. "Conforming and nonconforming finite element methods for solving the stationary Stokes equations. I". Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge 7 (1973), pp. 33–75.

REFERENCES

- [21] B. Dacorogna. Direct methods in the calculus of variations. Vol. 78. Springer Science & Business Media, 2007.
- [22] D. Di Pietro and A. Ern. "Discrete functional analysis tools for discontinuous Galerkin methods with application to the incompressible Navier-Stokes equations". *Mathematics of Computation* 79.271 (2010), pp. 1303–1330.
- [23] G. Dolzmann and N. J. Walkington. "Estimates for numerical approximations of rank one convex envelopes". Numer. Math. 85.4 (2000), pp. 647–663. URL: https://doi.org/10.1007/PL00005395.
- [24] M. K. Gobbert and A. Prohl. "A discontinuous finite element method for solving a multiwell problem". SIAM J. Numer. Anal. 37.1 (1999), pp. 246–268. URL: https://doi.org/10.1137/S0036142998333791.
- [25] G. Grekas. "Modelling, Analysis and Computation of Cell-Induced Phase Transitions in Fibrous Biomaterials". PhD thesis. University of Crete, 2019.
- [26] G. Grekas, K. Koumatos, C. Makridakis, and P. Rosakis. "Approximations of energy minimization in cell-induced phase transitions of fibrous biomaterials: Γ-convergence analysis". SIAM J. Numer. Anal. 60.2 (2022), pp. 715–750.
- [27] G. Grekas, K. Koumatos, C. Makridakis, and A. Vikelis. A class of Discontinuous Galerkin methods for nonlinear variational problems. 2023, To appear in Math. Comp. arXiv: 2308.12891 [math.NA]. URL: https://arxiv.org/abs/2308.12891.
- [28] O. A. Karakashian and F. Pascal. "A posteriori error estimates for a discontinuous Galerkin approximation of second-order elliptic problems". SIAM J. Numer. Anal. 41.6 (2003), pp. 2374–2399.
- [29] M. Köhler, T. Neumeier, M. A. Peter, D. Peterseim, and D. Balzani. "Hierarchical rank-one sequence convexification for the relaxation of variational problems with microstructures". *Comput. Methods Appl. Mech. Engrg.* 432 (2024), Paper No. 117321, 20. URL: https://doi. org/10.1016/j.cma.2024.117321.
- [30] M. Kružík. "Numerical approach to double well problems". SIAM J. Numer. Anal. 35.5 (1998), pp. 1833–1849. URL: https://doi.org/ 10.1137/S0036142995286477.
- [31] M. Luskin. "On the computation of crystalline microstructure". Vol. 5. Acta Numer. Cambridge Univ. Press, Cambridge, 1996, pp. 191–257. URL: https://doi.org/10.1017/S0962492900002658.

28

REFERENCES

- [32] T. Neumeier, M. A. Peter, D. Peterseim, and D. Wiedemann. "Computational polyconvexification of isotropic functions". *Multiscale Model. Simul.* 22.4 (2024), pp. 1402–1420. URL: https://doi.org/10.1137/ 23M1589773.
- [33] C. Ortner. "Nonconforming finite-element discretization of convex variational problems". IMA J. Numer. Anal. 31.3 (2011), pp. 847–864. URL: https://doi.org/10.1093/imanum/drq004.
- [34] C. Ortner and D. Praetorius. "On the convergence of adaptive nonconforming finite element methods for a class of convex variational problems". SIAM J. Numer. Anal. 49.1 (2011), pp. 346–367. URL: https: //doi.org/10.1137/090781073.
- [35] C. Ortner and E. Süli. "Discontinuous Galerkin finite element approximation of nonlinear second-order elliptic and hyperbolic systems". *SIAM J. Numer. Anal.* 45.4 (2007), pp. 1370–1397.
- [36] P. Pedregal. "On the numerical analysis of non-convex variational problems". Numer. Math. 74.3 (1996), pp. 325–336. URL: https:// doi.org/10.1007/s002110050219.
- [37] P. Pedregal. Variational methods in nonlinear elasticity. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2000, pp. xii+99. URL: https://doi.org/10.1137/1.9780898719529.
- [38] A. Ten Eyck and A. Lew. "Discontinuous Galerkin methods for nonlinear elasticity". International Journal for Numerical Methods in Engineering 67.9 (2006), pp. 1204–1243.