

# The Laguerre constellation of classical orthogonal Polynomials

R. S. Costas-Santos<sup>1</sup> 

<sup>1</sup> *Department of Quantitative Methods, Universidad Loyola Andalucía, E-41704, Sevilla, Spain*

URL: <http://www.rscosan.com>

E-mail: [rscosa@gmail.com](mailto:rscosa@gmail.com)

Received **January 23, 2025** in final form ?????; Published online ????

doi:10.3842/JOURNAL.2024.\*\*\*

## Abstract

A linear functional  $\mathbf{u}$  is classical if there exist polynomials,  $\phi$  and  $\psi$ , with  $\deg \phi \leq 2$ ,  $\deg \psi = 1$ , such that  $\mathcal{D}(\phi(x)\mathbf{u}) = \psi(x)\mathbf{u}$ , where  $\mathcal{D}$  is a certain differential, or difference, operator. The polynomials orthogonal with respect to the linear functional  $\mathbf{u}$  are called **classical orthogonal polynomials**. In the theory of orthogonal polynomials, a correct characterization of the classical families is of great interest. In this work, on the one hand, we present the Laguerre constellation, which is formed by all the classical families for which  $\deg \phi = 1$ , obtaining for all of them new algebraic identities such as structure formulas, orthogonality properties as well as new Rodrigues formulas; on the other hand, we present a theorem that characterizes the classical families belonging to the Laguerre constellation.

*Key words:* Recurrence relation; Characterization Theorem; Classical orthogonal polynomials; Laguerre constellation;

*2020 Mathematics Subject Classification:* 42C05; 33C45; 33D45

## 1 Introduction

Orthogonal polynomials  $(p_n(x))_n$  associated with a measure on the real line, i.e.,

$$\int_{\mathbb{R}} p_m(x)p_n(x)d\mu(x) = d_n^2 \delta_{mn},$$

satisfy a three-term recurrence equation

$$xp_n(x) = \alpha_n p_{n+1}(x) + \beta_n p_n(x) + \gamma_n p_{n-1}(x), \quad (1.1) \text{ ttrr}$$

where  $p_{-1}(x) = 0$ ,  $p_0(x) = 1$  and, according to the Favard theorem (cf. [6, p. 21]), if  $\gamma_n \neq 0$  for all  $n \in \mathbb{N}$  this recurrence completely characterizes such a polynomial sequence.

Due to the property of orthogonality of such a polynomial sequence, it is well-known the following relation between the recurrence coefficients:

$$\gamma_n = \alpha_{n-1} \frac{d_n^2}{d_{n-1}^2}, \quad n = 1, 2, \dots, \quad (1.2) \text{ gamrel}$$

where  $d_n^2$  is the squared norm of  $p_n$ .

Such polynomial sequence is said to be classical if it is orthogonal with respect to a linear functional  $\mathbf{u} : \mathbb{P} \rightarrow \mathbb{C}$  which fulfills the Pearson-type equation

$$\mathcal{D}(\phi(x)\mathbf{u}) = \psi(x)\mathbf{u}, \quad (1.3) \text{ peq}$$

where  $\phi$  is a polynomial of degree at most 2,  $\psi$  is a polynomial of degree 1, and  $\mathcal{D}$  is the differential operator in the continuous case, the forward ( $\Delta$ ) or backward ( $\nabla$ ) difference operator in the discrete case, and the Hahn operator ( $\mathcal{D}_q$ ) in the  $q$ -discrete case.

**Remark 1.1.** Note that if a linear functional  $\mathbf{u}$  fulfills (1.3) then it also fulfills the Pearson-type equation [9]

$$\mathcal{D}^*(\phi^*(x)\mathbf{u}) = \psi^*(x)\mathbf{u},$$

where

- In the continuous case  $\mathcal{D}^* = \mathcal{D}$ , so  $\phi^*(x) = \phi(x)$  and  $\psi^*(x) = \psi(x)$ .
- In the discrete case  $\phi^*(x) = \phi(x) + \psi(x)$ ,  $\psi^*(x) = \psi(x)$ ,  $\Delta^* = \nabla$ ,  $\nabla^* = \Delta$ .
- In the  $q$ -discrete case,  $\phi^*(x) = \phi(x) + (q-1)x\psi(x)$ ,  $\psi^*(x) = q\psi(x)$ , and  $\mathcal{D}_q^* = \mathcal{D}_{q^{-1}}$ .

?⟨def:1.2⟩?

**Definition 1.2.** The polynomial sequence  $(p_n(x))_n$  belongs to the Laguerre constellation (in short, LC) if it is a classical orthogonal polynomial sequence and  $\deg \phi = 1$  or  $\deg \phi^* = 1$ .

The so-called characterization Theorems play a fundamental role, i.e., the Theorems which collect those properties that completely define and characterize the classical orthogonal polynomials. One of the many ways to characterize a family of continuous classical polynomials (Hermite, Laguerre, Jacobi, and Bessel), which was first posed by R. Askey and proved by W. A. Al-Salam and T. S. Chihara [1] (see also [17]), is the structure relation

$$\phi \mathcal{D} p_n(x) = \tilde{a}_n p_{n+1}(x) + \tilde{b}_n p_n(x) + \tilde{c}_n p_{n-1}(x), \quad (1.4) \quad \boxed{\text{eq:1}}$$

where  $\tilde{c}_n \neq 0$ .

A. G. Garcia et al. proved in [11] that the relation (1.4) also characterizes the discrete classical orthogonal polynomials (Hahn, Krawtchouk, Meixner, and Charlier polynomials) when the forward replaces the derivative difference operator  $\Delta$ .

Later on, J. C. Medem et al. [18] characterized the orthogonal polynomials which belong to the  $q$ -Hahn class by a structure relation obtained from (1.4) replacing the derivative by the  $q$ -difference operator  $\mathcal{D}_q$  (see also [2, 3, 4]). One of the most general characterization theorem for the  $q$ -polynomials in the  $q$ -quadratic lattice was done in [8].

The structure of this work is the following: in Section 2, we introduce some notations and definitions used throughout the paper. In Section 3 we present the main results of the polynomials that belongs to the Laguerre constellation and the algebraic properties supporting the presented results. In Section 4 we present the specific identities about the different families of the Laguerre Constellation. Several Wolfram Mathematica<sup>®</sup> codes used to obtain the different expressions presented in this work have been included at the end of this work.

## 2 Preliminaries

⟨sec:2⟩

In this section, we will give a brief survey of the operational calculus that we will use as well as some basic notations that we will need in the rest of this work.

### 2.1 Basic concepts and results

We adopt the following set notations:  $\mathbb{N}_0 := \{0\} \cup \mathbb{N} = \{0, 1, 2, \dots\}$ , and we use the sets  $\mathbb{Z}$ ,  $\mathbb{R}$ ,  $\mathbb{C}$ , which represents the integers, real numbers, and complex numbers, respectively. Let  $\mathbb{P}$  be the linear space of polynomial functions in  $\mathbb{C}$  (in the following, we will refer to them as polynomials) with complex coefficients and  $\mathbb{P}^*$  be its algebraic dual space, i.e.,  $\mathbb{P}^*$  is the linear space of all linear maps  $\mathbf{u} : \mathbb{P} \rightarrow \mathbb{C}$ . In the following, we will call the elements of  $\mathbb{P}^*$  as functionals. In general, we will represent the action of a linear functional over a polynomial by

$$\langle \mathbf{u}, \pi \rangle, \quad \mathbf{u} \in \mathbb{P}^*, \quad \pi \in \mathbb{P}.$$

Therefore, a functional is completely determined by a sequence of complex numbers  $\langle \mathbf{u}, x^n \rangle = u_n$ ,  $n = 0, 1, \dots$ , the so-called moments of the functional.

**Definition 2.1.** Let  $\mathbf{u} \in \mathbb{P}^*$  be a functional. We say that  $\mathbf{u}$  is a quasi-definite functional if there exists a polynomial sequence  $(p_n)$  which is orthogonal with respect to  $\mathbf{u}$ , i.e.

$$\langle \mathbf{u}, p_m p_n \rangle = k_n \delta_{mn}, \quad k_n \neq 0, \quad n = 0, 1, \dots,$$

where  $\delta_{mn}$  is the Kronecker delta.

In order to obtain our derived identities, we rely on properties of the shifted factorial, or Pochhammer symbol,  $(a)_n$ , and the  $q$ -shifted factorial, or  $q$ -Pochhammer symbol,  $(a; q)_n$ . For any  $n \in \mathbb{N}_0$ ,  $a, q \in \mathbb{C}$ , the shifted factorial is defined as

$$(a)_n = a(a+1) \cdots (a+n-1),$$

the  $q$ -shifted factorial is defined as

$$(a; q)_n = (1-a)(1-aq) \cdots (1-aq^{n-1}).$$

We will also use the common notational product conventions

$$\begin{aligned} (a_1, \dots, a_k)_n &= (a_1)_n \cdots (a_k)_n, \\ (a_1, \dots, a_k; q)_n &= (a_1; q)_n \cdots (a_k; q)_n. \end{aligned}$$

The hypergeometric series  ${}_rF_s$ , where  $s, r \in \mathbb{N}_0$ , is defined by the series [14, (1.4.1)]

$${}_rF_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r)_k}{(b_1, \dots, b_s)_k} \frac{z^k}{k!}, \tag{2.1} \text{?eq:2.1?}$$

and the basic hypergeometric series  ${}_r\phi_s$ , is defined by the series as [14, (1.10.1)]

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r; q)_k}{(q, b_1, \dots, b_s; q)_k} \left( (-1)^k q^{\binom{k}{2}} \right)^{1+s-r} z^k. \tag{2.2} \text{?eq:2.2?}$$

Observe that both, hypergeometric and basic hypergeometric series, are entire functions of  $z$  if  $s+1 > r$ , are convergent for  $|z| < 1$  for  $s+1 = r$ , and divergent if  $s+1 < r$  and  $z \neq 0$ , unless it terminates.

Note that when we refer to a hypergeometric or basic hypergeometric function with *arbitrary argument*  $z$ , we mean that the argument does not necessarily depend on the other parameters, namely the  $a_j$ 's,  $b_j$ 's. However, for the arbitrary argument  $z$ , it very-well may be that the domain of the argument is restricted, such as for  $|z| < 1$ .

The next theorem [6, p. 8] is useful if one works with linear functionals.

**Theorem 2.2.** Let  $\mathbf{u} \in \mathbb{P}^*$  be a linear functional with moments  $(u_n)$ . Then,  $\mathbf{u}$  is quasi-definite if and only if the Hankel determinants  $H_n := \det \left( u_{i+j} \right)_{i,j=0}^n \neq 0$ ,  $n = 0, 1, \dots$

## 2.2 Definition of the operators in $\mathbb{P}$ and $\mathbb{P}^*$

Next, we will define the backward and forward difference operators as well the so called  $q$ -derivative operator, or Hahn operator.

$$\begin{aligned} \Delta f(x) &= f(x+1) - f(x), \\ \nabla f(x) &= f(x) - f(x-1), \\ \mathcal{D}_q f(x) &= \frac{f(qx) - f(x)}{x(q-1)}, \quad q \neq 1, \quad x \neq 0. \end{aligned}$$

Since the polynomial sequences depend on  $n$ ,  $x$  and their parameters and we are going to focus more on the variable  $n$  along this work ( $n$  as a discrete variable) than in the variable  $x$ , we need to consider along the paper as well the operators  $\Delta_n$  and  $\nabla_n$  that are defined analogously as the operators  $\Delta$  and  $\nabla$ , i.e.

$$\nabla_n f(n; x) = f(n, x) - f(n-1, x), \quad \Delta_n f(n; x) = f(n+1, x) - f(n, x).$$

**Definition 2.3.** Let  $\mathbf{u} \in \mathbb{P}^*$  and  $\pi \in \mathbb{P}$ , let  $\Delta\mathbf{u}$ ,  $\nabla\mathbf{u}$  and  $\mathcal{D}_q^*\mathbf{u}$  be the linear functionals defined by

$$\left\langle \frac{d}{dx}\mathbf{u}, \pi \right\rangle = -\langle \mathbf{u}, \pi' \rangle, \quad (2.3) \text{?eq:difdef?}$$

$$\langle \Delta\mathbf{u}, \pi \rangle = -\langle \mathbf{u}, \nabla\pi \rangle, \quad (2.4) \text{?eq:backdef?}$$

$$\langle \nabla\mathbf{u}, \pi \rangle = -\langle \mathbf{u}, \Delta\pi \rangle, \quad (2.5) \text{?eq:forwdef?}$$

$$\langle \mathcal{D}_q^*\mathbf{u}, \pi \rangle = -q\langle \mathbf{u}, \mathcal{D}_q\pi \rangle. \quad (2.6) \text{?eq:qdifdef?}$$

Notice that we use the same notation for the operators on  $\mathbb{P}$  and  $\mathbb{P}^*$ . Whenever it is not specified the linear space where an operator acts, it will be understood that it acts on the polynomial space  $\mathbb{P}$ .

### 3 The Laguerre constellation

(sec:3)

In this section, we are going to present several identities that the polynomials belonging to LC fulfill. First, we are going to show some theoretical aspects and results related to them.

(lem:3.1)

**Lemma 3.1.** Let  $\mathbf{u} \in \mathbb{P}^*$  be a quasi-definite classical functional, let  $(p_n)$  be the polynomial sequence orthogonal with respect to  $\mathbf{u}$ . If  $(p_n)$  belong to the LC then, there exists a numerical sequence  $(\lambda_n)_n$  so that  $(\lambda_n p_n)_n$  fulfills the recurrence relation (1.1) for which  $\gamma_0 = 0$  and  $\alpha_n + \beta_n + \gamma_n = c$  for all  $n \in \mathbb{N}$ , where  $c$  is a root of  $\phi(x)$ , or  $\phi^*(x)$ .

**Proof.** To prove this result it is enough to consider all the families of the Hypergeometric orthogonal polynomials scheme and the basic Hypergeometric orthogonal polynomials scheme (see e.g. [14, 18, 19]) that are of Laguerre-type, i.e.,  $\deg \phi = 1$  or  $\deg \phi^* = 1$ . These families are the Laguerre (L), Charlier (C), Meixner (M), big  $q$ -Laguerre (bqL),  $q$ -Meixner (qM), little  $q$ -Laguerre (lqL),  $q$ -Laguerre (qL),  $q$ -Charlier (qC), Stieltjes-Wigert (SW), and the 0-Laguerre/Bessel (OLB) polynomials

Once we obtain all the families belonging to LC it is enough to verify that the conditions established for each of the families (see Table 1) are satisfied. Observe that the value of  $p_n(c)$ , where  $c$  is a zero of  $\phi$  or  $\phi^*$ , is known (see [15, 20]), and these values are non-zero so one can define  $\lambda_n$  as  $1/p_n(c)$ . ■

**Remark 3.2.** Note that Lemma 3.1 is not a Characterization Theorem since other families of the basic Hypergeometric orthogonal polynomials scheme fulfill the condition about the recurrence coefficients, in short (RC), i.e.  $\alpha_n + \beta_n + \gamma_n$  is a constant, however they do not belong to the LC, for example, the Askey-Wilson polynomials (see [14, (14.1.4)]).

In the next result we write the recurrence relation as a structure-type relation on  $n$ .

(lem:3.3)?

**Lemma 3.3.** Let  $(p_n)_n$  be a polynomial sequence that belongs to the Laguerre constellation. For any  $x \in \mathbb{C}$  the recurrence relation (1.1) can be written as:

$$\phi(x)p_n(x) = \alpha_n \Delta_n p_n(x) - \gamma_n \Delta_n p_{n-1}(x), \quad (3.1) \text{eq:sr1}$$

if  $\deg \phi = 1$ ; and

$$\phi^*(x)p_n(x) = \alpha_n \Delta_n p_n(x) - \gamma_n \Delta_n p_{n-1}(x),$$

if  $\deg \phi^* = 1$ .

**Proof.** Let us assume that  $\deg \phi = 1$ , i.e.,  $\phi(x) = x - c$ , then by Lemma 3.1 we have that the coefficients of the recurrence relation (1.1) fulfill  $\alpha_n + \beta_n + \gamma_n = c$ , then rewriting the recurrence assuming this relation the result holds. The  $\deg \phi^* = 1$  case is analogous. ■

**Remark 3.4.** For the sake of convenience, we are going to replace  $x - c$  by  $\phi(x)$  assuming that  $\deg \phi = 1$ . In the case that  $\deg \phi \neq 1$  and  $\deg \phi^* = 1$ , then one must replace  $\phi$  by  $\phi^*$  in the further results since such identities and results hold similarly.

We write the recurrence relation in the next result as a Sturm-Liouville form difference equation on  $n$ .

(lem:3.4)

**Lemma 3.5.** *Let  $(p_n)_n$  be a polynomial sequence that belongs to the Laguerre constellation. For any  $x \in \mathbb{C}$  the recurrence relation (1.1) can be written as:*

$$\phi(x)p_n(x) = d_n^2 \nabla_n \frac{\gamma_n}{d_n^2} \Delta_n p_n(x), \tag{3.2} \text{?eq:3.1?}$$

$$= d_n^2 \Delta_n \frac{\alpha_n}{d_n^2} \nabla_n p_n(x). \tag{3.3} \text{eq:3.2}$$

**Proof.** Starting from (3.1), using (1.2) and taking into account the definition of  $\nabla_n$  and  $\Delta_n$  the results follow. ■

?<thm:3.1)?

**Theorem 3.6.** *Let  $(p_n)_n$  be a polynomial sequence that belongs to the Laguerre constellation. For any  $x \in \mathbb{C}$ , the sequence  $(\nabla_n p_n(x))$  is orthogonal with respect to  $\alpha_n/d_n^2$ , and the sequence  $(\Delta_n p_n(x))$  is orthogonal with respect to  $\gamma_n/d_n^2$ .*

**Proof.** Let us fix  $x \in \mathbb{C}$  such that  $\omega(x) \neq 0$ . By Korovkin's Theorem (see [16] or [12, Theorem 2.1]) we have the following closure relation:

$$\sum_{n=0}^{\infty} \frac{p_n(x)p_n(y)}{d_n^2} = \frac{1}{\omega(x)} \delta(x-y),$$

where  $\omega$  is the weight function and both the left- and right-hand sides should be treated as distributions. From this expression, using (3.3) and Abel's lemma on partial sums [13, p. 313] for  $y \neq x$  we obtain

$$0 = \sum_{n=0}^{\infty} \frac{\phi(x)p_n(x)p_n(y)}{d_n^2} = \sum_{n=0}^{\infty} \left( \Delta_n \frac{\alpha_n}{d_n^2} \nabla_n p_n(x) \right) p_n(y) = - \sum_{n=0}^{\infty} \frac{\alpha_n}{d_n^2} \nabla_n p_n(x) \nabla_n p_n(y).$$

Moreover,

$$\sum_{n=0}^{\infty} \frac{\alpha_n}{d_n^2} \nabla_n p_n(x) \nabla_n p_n(x) = \frac{\phi(x)}{\omega(x)}.$$

The proof for the sequence  $(\Delta_n p_n(x))$  is analogous. Hence the result follows. ■

**Remark 3.7.** *Since we have the data for all the families which belong to the LC it is a straightforward calculation to check that  $1/d_{-1}^2 = 0$  (since  $\gamma_0 = 0$ ) and*

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{d_n^2} = 0.$$

**Remark 3.8.** *Observe the previous result does not characterize families belonging to LC since, for example, for the Jacobi polynomials the following identity holds:*

$$\nabla_n \left( \frac{P_n^{(\alpha,\beta)}(x)}{P_n^{(\alpha,\beta)}(1)} \right) = \frac{P_n^{(\alpha,\beta)}(x)}{P_n^{(\alpha,\beta)}(1)} - \frac{P_{n-1}^{(\alpha,\beta)}(x)}{P_{n-1}^{(\alpha,\beta)}(1)} = \frac{2n + \alpha + \beta}{2(1+a)} (x-1) \frac{P_{n-1}^{(\alpha+1,\beta)}(x)}{P_{n-1}^{(\alpha+1,\beta)}(1)},$$

and in such a case  $\phi(x) = 1 - x^2$ .

?<thm:3.2)?

**Theorem 3.9** (Characterization Theorem). *Let  $(p_n)_n$  be an orthogonal polynomial sequence with respect to  $\omega$ , such that  $\mathcal{D}(\phi\omega) = \psi\omega$ . For any  $x \in \mathbb{C}$  such that  $\omega(x) \neq 0$ , the following statements are equivalent:*

1.  $(p_n)_n$  belongs to the Laguerre constellation.

2. The polynomial sequence  $(p_n(x))$  fulfills the second order difference equation

$$\phi(x)p_n(x) = \alpha_n \Delta_n p_n(x) - \gamma_n \Delta_n p_{n-1}(x),$$

which is equal to its discrete structure-type relation.

3. The polynomial sequence  $(p_n(x))$  satisfies the Sturm-Liouville difference equations


$$\phi(x)p_n(x) = d_n^2 \nabla_n \frac{\gamma_n}{d_n^2} \Delta_n p_n(x) = d_n^2 \Delta_n \frac{\alpha_n}{d_n^2} \nabla_n p_n(x).$$

**Proof.** 1.  $\leftrightarrow$  2. since the 2rd item is the recurrence relation in such a case. 2.  $\leftrightarrow$  3. is a consequence of Lemma 3.5.  $\blacksquare$

## 4 The families

(sec:4)

Along this section we present several identities related to the different families that belong to LC. Since there are some relation among them (see Figure 1) we will present such identities only for some of the families in order to avoid duplicities. We consider the Laguerre, Meixner, Charlier, big  $q$ -Laguerre, little  $q$ -Laguerre and Stieltjes-Weigert cases.



(fig:1) **Figure 1.** Relations between the families in the Laguerre constellation. The gray lines are the particular cases. The black lines are the limiting cases.

Before presenting the main results let us show the relations between the families we are going to consider with respect to the rest of the families that belong to LC (see [7, p. 20],

[14, remark p. 526]):

$$p_n(x; a, b, 1/q) = \frac{1}{(q/b; q)_n} M_n(xq/a; 1/a, -b; q), \tag{4.1} \{?\}$$

$$p_n(x; q^\alpha | 1/q) = \frac{(q; q)_n}{(q^{\alpha+1}; q)_n} L_n^{(\alpha)}(-x; q), \tag{4.2} \{?\}$$

$$L_n^{(\alpha)}(x; q) = \frac{1}{(q; q)_n} C_n(-x; -q^{-\alpha}; q), \tag{4.3} \{?\}$$

$$S_n(x/a, 1/q) = {}_2\phi_0\left(q^{-n}, 0; -; q, -\frac{x}{a}\right) =: l_n(x; a), \tag{4.4} \{?\}$$

where  $l_n(x; a)$  denote the 0-Laguerre/Bessel polynomials ( $l_n(x; a)$ ) (see [5], [19, p. 244]).

### 4.1 The Laguerre polynomials

The Laguerre polynomials can be written in terms of hypergeometric series as [14, §9.12]

$$L_n^{(\alpha)}(x) = \frac{(\alpha + 1)_n}{n!} {}_1F_1\left(\begin{matrix} -n \\ \alpha + 1 \end{matrix}; x\right).$$

In this case  $\phi(x) = x$ , i.e., this family belongs to the Laguerre constellation where the zero of  $\phi$  is  $c = 0$ , and  $L_n^{(\alpha)}(c) = (\alpha + 1)_n/n!$ . Taking this into account we can state the following result.

(lem:3.5)

**Lemma 4.1.** *For any  $\alpha \in \mathbb{C}$  and any  $n \in \mathbb{N}_0$ , the following identities hold:*

$$\nabla_n L_n^{(\alpha)}(x) = L_n^{(\alpha-1)}(x), \tag{4.5} \boxed{\text{lownL}}$$

$$\alpha L_n^{(\alpha)}(x) - (n + 1)\Delta_n L_n^{(\alpha)}(x) = x L_n^{(\alpha+1)}(x), \tag{4.6} \boxed{\text{?raiupnL?}}$$

$$\alpha L_n^{(\alpha)}(x) - (n + \alpha)\nabla_n L_n^{(\alpha)}(x) = x L_{n-1}^{(\alpha+1)}(x), \tag{4.7} \boxed{\text{raidonL}}$$

$$\nabla_\alpha L_n^{(\alpha)}(x) = L_{n-1}^{(\alpha)}(x), \tag{4.8} \boxed{\text{lowaL}}$$

$$(n + \alpha - x)L_n^{(\alpha)}(x) - (\alpha + n)\nabla_\alpha L_n^{(\alpha)}(x) = (n + 1)L_{n+1}^{(\alpha-1)}(x), \tag{4.9} \boxed{\text{raidoaL}}$$

$$(n + 1 + \alpha - x)L_n^{(\alpha)}(x) - x\Delta_\alpha L_n^{(\alpha)}(x) = (n + 1)L_{n+1}^{(\alpha)}(x), \tag{4.10} \boxed{\text{raiupaL}}$$

where  $\Delta_\alpha f(x, \alpha) = f(x, \alpha + 1) - f(x, \alpha)$ , and  $\nabla_\alpha f(x, \alpha) = \Delta_\alpha f(x, \alpha - 1)$ .

**Proof.** These identities can be checked by identifying the polynomial coefficients on the left and right-hand sides of each identity. Hence, the results follow. ■

A direct consequence of the former result is stated as follows.

(thm:3.3)?

**Theorem 4.2.** *For any  $\alpha \in \mathbb{C}$ , any  $x \in \mathbb{C}$ ,  $x \neq 0$ , and any  $n \in \mathbb{N}_0$ , the polynomial sequence  $(L_n^{(\alpha+k)}(x))_k$  is orthogonal with respect to certain moment functional.*

**Proof.** By combining (4.8), (4.9) and (4.10) we have the following second-order difference equations:

$$nL_n^{(\alpha)}(x) = -(\alpha + n)\nabla_\alpha \Delta_\alpha L_n^{(\alpha)}(x) + (\alpha + n - x)\Delta_\alpha L_n^{(\alpha)}(x) \tag{4.11} \boxed{\text{?eq:sodeL1?}}$$

$$= -x\Delta_\alpha \nabla_\alpha L_n^{(\alpha)}(x) + (\alpha + n - x)\nabla_\alpha L_n^{(\alpha)}(x), \tag{4.12} \boxed{\text{?eq:sodeL2?}}$$

which is connected with the Charlier polynomials case (see [14, (9.14.5)]). Using the theory of discrete Sturm-Liouville and assuming that for any  $n \in \mathbb{N}_0$ , in this case,  $\alpha$  is variable, and the rest of the elements are fixed, the result holds. ■

**Remark 4.3.** *Observe that one can construct, under certain condition on  $\alpha$  and  $x$ , certain integral representation for such moment functional.*

From Lemma 4.1 we can deduce some new identities related to the Laguerre polynomials.

?{thm:3.4}?

**Theorem 4.4.** For any  $\alpha \in \mathbb{C}$ , and any  $n, k \in \mathbb{N}_0$ , the following Rodrigues-type identities hold:

$$L_n^{(\alpha+k)}(x) = \frac{(\alpha+k)_n}{x^k} (\alpha+k)(\alpha+k-1) \nabla_\alpha \cdots (\alpha+1) \alpha \nabla_\alpha \frac{L_{n+k}^{(\alpha)}(x)}{(\alpha)_{n+k}}, \quad (4.13) \text{ ?eq:LalRF?}$$

$$L_n^{(\alpha+k)}(x) = \frac{(-1)^k (\alpha+1)_{n+k}}{n! x^k} \nabla_n^k \frac{(n+k)!}{(\alpha+1)_{n+k}} L_{n+k}^{(\alpha)}(x), \quad (4.14) \text{ ?eq:LnRF?}$$

$$L_n^{(\alpha)}(x) = (-1)^k \frac{(n+1)_\alpha}{x^\alpha} \Delta_\alpha^k \frac{x^\alpha}{(n-k+1)_k} L_{n-k}^{(\alpha)}(x), \quad (4.15) \text{ ?eq:LalRF3?}$$

$$L_n^{(\alpha)}(x) = \Delta_n^k L_{n-k}^{(\alpha+k)}(x), \quad (4.16) \text{ ?eq:LnRF2?}$$

$$L_n^{(\alpha+k)}(x) = \Delta_\alpha^k L_{n+k}^{(\alpha)}(x). \quad (4.17) \text{ ?eq:LalRF2?}$$

**Proof.** The first identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.7) written in the following way:

$$\frac{n!}{(\alpha+k+1)_n} L_n^{(\alpha+k)}(x) = \frac{(\alpha+k)(\alpha+k-1)}{x(n+1)} \nabla_\alpha \frac{(n+1)!}{(\alpha+k)_{n+1}} L_{n+1}^{(\alpha+k-1)}(x).$$

The second identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.9) written in the following way:

$$\frac{n!}{(\alpha+k+1)_n} L_n^{(\alpha+k)}(x) = -\frac{\alpha+k}{x} \nabla_n \frac{(n+1)!}{(\alpha+k)_{n+1}} L_{n+1}^{(\alpha+k-1)}(x).$$

The third identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.10) written in the following way:

$$x^\alpha L_n^{(\alpha)}(x) = -(n+1)_\alpha \Delta_n \frac{x^\alpha}{(n)_\alpha} L_{n-1}^{(\alpha)}(x).$$

The fourth and the fifth relation hold from (4.5) and (4.8). ■

The last result, but not least interesting, concerning the operators associated with the Laguerre polynomials is as follows.

?{pro:1}?

**Proposition 4.5.** The Laguerre polynomials fulfill the following identity:

$$(n+1)L_{n+1}^{(\alpha+1)}(x) = (2+a-x)L_n^{(\alpha)}(x) + (3x-4-2a)(L_n^{(\alpha)}(x))' + (2+a-3x)(L_n^{(\alpha)}(x))'' + x(L_n^{(\alpha)}(x))'''.$$

**Proof.** The identity follows using the identities:

$$L_n^{(\alpha)}(x) - (L_n^{(\alpha)}(x))' = L_n^{(\alpha+1)}(x), \quad (4.18) \text{ {?}}$$

$$x(L_n^{(\alpha)}(x))' - (x-\alpha)L_n^{(\alpha)}(x) = (n+1)L_{n+1}^{(\alpha-1)}(x), \quad (4.19) \text{ {?}}$$

applying the first twice, and later the second one once mapping  $\alpha \mapsto \alpha+2$ . ■

## 4.2 The Charlier polynomials

The Charlier polynomials can be written in terms of hypergeometric series as [14, §9.14]

$$C_n(x; a) = {}_2F_0 \left( \begin{matrix} -n, -x \\ - \\ -\frac{1}{a} \end{matrix} \right).$$

In this case  $\phi(x) = x$  and  $\phi^*(x) = a$ , i.e.,  $c = 0$ , and  $C_n(c; a) = 1$ . Taking this into account we can state the following result.



(lem:3.6)

**Lemma 4.6.** For any  $a \in \mathbb{C}$  and any  $n \in \mathbb{N}_0$ , the following identities hold:

$$a\Delta_n C_n(x; a) = -x C_n(x - 1; a), \tag{4.20} \text{rainC}$$

$$nC_n(x; a) + a(C_n(x; a))' = (n + 1)C_{n-1}(x; a), \tag{4.21} \text{lowDC}$$

$$(a - n)C_n(x; a) + n\nabla_n C_n(x; a) = aC_n(x + 1; a), \tag{4.22} \text{?lowdonC?}$$

$$(a - n - 1)C_n(x; a) + a\Delta_n C_n(x; a) = aC_{n+1}(x + 1; a). \tag{4.23} \text{?lowupnC?}$$

**Proof.** (4.21) is a direct consequence of the identity presented in Remark in [14, p. 249]

$$\frac{(-a)^n}{n!} C_n(x; a) = L_n^{(x-n)}(a).$$

The other identities can be checked by identifying the polynomial coefficients on the left and right-hand sides of each identity. Hence, the results follow ■

From Lemma 4.6 we can deduce some new identities related to the Charlier polynomials.

(thm:3.5)

**Theorem 4.7.** For any  $a \in \mathbb{C}$ , and any  $n, k \in \mathbb{N}_0$ , the following Rodrigues-type identities hold:

$$C_n(x; a) = \frac{(-a)^k}{(x + 1)_k} \Delta_n^k C_n(x + k; a), \tag{4.24} \text{?eq:CnRF?}$$

$$C_n(x; a) = \frac{n!}{a^n} \Delta_n^k \frac{a^{n-k}}{(n - k)!} C_{n-k}(x - k; a). \tag{4.25} \text{?eq:CnRF2?}$$

**Proof.** The first identity is a direct consequence of (4.20); and the second is due the identity:

$$(n + 1)! \Delta_n \frac{a^n}{n!} C_n(x; a) = a^{n+1} C_{n+1}(x + 1; a). \tag{4.26}$$

■

### 4.3 The Meixner polynomials

The Meixner polynomials can be written in terms of hypergeometric series as [14, §9.10]

$$M_n(x; \beta, c) = {}_2F_1 \left( \begin{matrix} -n, -x \\ \beta \end{matrix}; 1 - \frac{1}{c} \right).$$

In this case  $\phi(x) = x$  and  $\phi^*(x) = c(x + \beta)$ , therefore we must consider two cases,  $c_1 = 0$ , and  $c_2 = -\beta$ , for which we have  $M_n(c_1; \beta, c) = 1$ , and  $M_n(c_2; \beta, c) = 1/c^n$ . Taking into account Lemma 3.1 we can state the next result.

?(lem:3.7)?

**Lemma 4.8.** The polynomial sequence  $(M_n(x; \beta, c)/M_n(c_2; \beta, c))_n$  fulfills the recurrence relation

$$(x + \beta)p_n(x) = \alpha_n^M p_{n+1}(x) - (\alpha_n^M + \gamma_n^M)p_n(x) + \gamma_n^M p_{n-1}(x),$$

with initial condition  $p_0(x) = 1$ ,

$$\alpha_n^M = \frac{n + \beta}{c - 1}, \quad \gamma_n^M = \frac{nc}{c - 1},$$

and fulfills the second order difference equation

$$(x + \beta)p_n(x) = d_n^2 \Delta_n \frac{\gamma_n^M}{d_n^2} \nabla_n p_n(x) = d_n^2 \nabla_n \frac{\alpha_n^M}{d_n^2} \Delta_n p_n(x) = \gamma_n^M \nabla_n \Delta_n p_n(x) + (\alpha_n^M - \gamma_n^M) \Delta_n p_n(x).$$

As in the Laguerre polynomial case, the next result follows.

(lem:3.8)

**Lemma 4.9.** For any  $\beta, c \in \mathbb{C}$ ,  $\beta \notin \{0, 2\}$ ,  $c \notin \{0, 1\}$ , and any  $n \in \mathbb{N}_0$ , the following identities hold:

$$\frac{\beta c}{c-1} \Delta_n M_n(x; \beta, c) = x M_n(x-1; \beta+1, c), \quad (4.26) \text{rainM}$$

$$M_n(x; \beta, c) + \frac{c}{c-1} \Delta_n M_n(x; \beta, c) = \frac{x+\beta}{\beta} M_n(x; \beta+1, c), \quad (4.27) \text{?raiupnM?}$$

$$M_n(x; \beta, c) + \frac{1}{c-1} \nabla_n M_n(x; \beta, c) = \frac{x+\beta}{\beta} M_{n-1}(x; \beta+1, c), \quad (4.28) \text{raidonM}$$

$$\frac{c\beta(1-\beta)}{c-1} \nabla_\beta M_n(x; \beta, c) = xn M_{n-1}(x-1; \beta+1, c), \quad (4.29) \text{lowbM}$$

$$\begin{aligned} \frac{\beta(\beta-1)c}{(\beta+n)(c-1)} M_{n+1}(x+1; \beta-1, c) &= \left( x + \beta + \frac{\beta(\beta-1)}{(\beta+n)(c-1)} \right) M_n(x; \beta, c) \\ &\quad + (x+\beta) \Delta_\beta M_n(x; \beta, c), \end{aligned} \quad (4.30) \text{raiupbM}$$

$$\begin{aligned} \frac{(\beta-1)(\beta-2)c}{(\beta-1+n)(c-1)} M_{n+1}(x+1; \beta-2, c) &= \left( x + \beta - 1 + \frac{(\beta-1)(\beta-2)}{(\beta-1+n)(c-1)} \right) M_n(x; \beta, c) \\ &\quad - \frac{(\beta-1)(\beta-2)}{(\beta-1+n)(c-1)} \nabla_\beta M_n(x; \beta, c), \end{aligned} \quad (4.31) \text{raidobM}$$

$$\frac{\partial}{\partial c} c^n (\beta)_n M_n(x; \beta, c) = \frac{n(x+\beta)}{c+n+\beta} c^n (\beta+1)_n M_{n-1}(x; \beta+1, c), \quad (4.32) \text{lowcM}$$

$$\begin{aligned} c(1-\beta) c^n (\beta)_n \beta M_{n+1}(x; \beta-1, c) &= c(1-c) \frac{\partial}{\partial c} c^n (\beta)_n M_n(x; \beta, c) \\ &\quad - ((c-1)x + n - (n+1)c + c\beta) c^n (\beta)_n \beta M_n(x; \beta, c). \end{aligned} \quad (4.33) \text{?raidcM?}$$

**Proof.** These identities can be checked by identifying the polynomial coefficients on the left and right-hand sides of each identity and with the help of Wolfram Mathematica 13<sup>®</sup>. ■

A direct consequence is the fact that this polynomial sequence is orthogonal with respect to the parameter  $\beta$ .

(thm:3.6)?

**Theorem 4.10.** For any  $\beta, c \in \mathbb{C}$ , with  $-\beta \notin \mathbb{N}_0$ , any  $x \in \mathbb{C}$ ,  $x \notin \{0, -\beta\}$ , and any  $n \in \mathbb{N}_0$ , the polynomial sequence  $(M_n(x; \beta+k, c))_k$  is orthogonal with respect to certain moment functional.

**Proof.** By combining (4.29), (4.30) and (4.31) we have the following second order difference equations:

$$nx M_n(x; \beta, c) = \frac{\beta(\beta-1)}{c-1} \nabla_\beta \Delta_\beta M_n(x; \beta, c) - \left( \frac{\beta(\beta-1)}{c-1} + (\beta+x)(\beta+n) \right) \Delta_\beta M_n(x; \beta, c), \quad (4.34) \text{?eq:sodeM?}$$

which is connected with the Continuous Hahn polynomials case (see [14, (9.4.5)]). Using the theory of discrete Sturm-Liouville and assuming that for any  $n \in \mathbb{N}_0$ , in this case,  $\beta$  is variable, and the rest of the elements are fixed, the result holds. ■

**Remark 4.11.** • Note that in [10] the authors extended the orthogonality relations for the Meixner polynomials through the orthogonality relations of the Continuous Hahn Polynomials.

- Observe that one can construct, under certain condition on the parameters,  $n$  and  $x$ , certain integral representation for such moment functional.

From Lemma 4.9 we can deduce some new identities related to the Meixner polynomials.

(thm:3.6)?

**Theorem 4.12.** For any  $\beta, c \in \mathbb{C}$ , and any  $n, k \in \mathbb{N}_0$ , the following Rodrigues-type identities hold:

$$M_n(x; \beta + k, c) = \frac{(\beta)_k c^k}{(c-1)^k (x+1)_k} \nabla_n^k M_{n+k}(x; \beta, c), \tag{4.35} \text{?eq:MnRF?}$$

$$M_n(x; \beta + k, c) = \frac{(\beta)_k}{(1-c)^k (x+\beta)_k (-c)^n} \Delta_n^k (-c)^{n+k} M_{n+k}(x; \beta, c), \tag{4.36} \text{?eq:MnRF2?}$$

$$M_n(x; \beta + k, c) = \frac{c^k (\beta + k - 1)(2 - k - \beta)}{(c-1)^k (x+1)_k (n+1)_k} \nabla_\beta \cdots \beta(1-\beta) \nabla_\beta M_{n+k}(x+k; \beta, c), \tag{4.37} \text{?eq:Mbrf?}$$

$$c^n (\beta + k)_n M_n(x; \beta + k, c) = \frac{(c+n+\beta+k)}{(n+1)_k (x+\beta)_k} \frac{\partial}{\partial c} \cdots (c+n+\beta+k) \frac{\partial}{\partial c} c^{n+k} (\beta)_{n+k} M_{n+k}(x; \beta, c). \tag{4.38} \text{?eq:LalRF2?}$$

**Proof.** The first identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.26). The second identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.28) written in the following way:

$$\Delta_n (-c)^n M_n(x; \beta, c) = \frac{x+\beta}{\beta} (1-c) (-c)^n M_{n-1}(x; \beta+1, c).$$

The third identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.29). The fourth and the fifth relation hold from (4.32). ■

The last result concerning the operators associated with the Meixner polynomials is as follows.

?(pro:2)?

**Proposition 4.13.** The following identity holds:

$$\begin{aligned} & \frac{(\beta c + cx + c - x)}{c} M_n(\beta, c, x) + \frac{(2\beta c^2 + 2c^2 x + 2c^2 - cx - x)}{(c-1)c} \nabla M_n(\beta, c, x) \\ & + \frac{c(-2\beta + 3\beta c + 3cx + 3c - 3x - 2)}{(c-1)^2} \nabla^2 M_n(\beta, c, x) + \frac{c^2(\beta + x + 1)}{(c-1)^2} \nabla^3 M_n(\beta, c, x) \\ & = \frac{(\beta + n)(\beta + n + 1)}{\beta} M_{n+1}(\beta + 1, c, x). \end{aligned}$$

**Proof.** The proof follows after using an algorithm written in Wolfram Mathematica 13<sup>®</sup>. ■

### 4.4 The big $q$ -Laguerre polynomials

The big  $q$ -Laguerre polynomials can be written in terms of basic hypergeometric series as [14, §14.11]

$$p_n(x; a, b; q) = {}_3\phi_2 \left( \begin{matrix} q^{-n}, 0, x \\ aq, bq \end{matrix}; q, q \right).$$

In this case  $\phi(x) = (x - aq)(x - bq)$  and  $\phi^*(x) = abq(1 - x)$ , i.e.,  $\mathbf{c} = 1$ , and  $p_n(\mathbf{c}; a, b; q) = 1$ . Observe these polynomials are symmetric in the parameters  $a$  and  $b$ . Taking this into account we can state the following result.

(lem:4.5)

**Lemma 4.14.** For any  $a, b \in \mathbb{C}$ , and any  $n \in \mathbb{N}_0$ , the following identities hold:

$$\phi(1)q^n \Delta_n p_n(x; a, b; q) = (x-1)p_n(xq; aq, bq; q), \quad (4.39) \text{rainbqL}$$

$$\begin{aligned} (-1+a)(-1+b)q^{n+1} p_{n+1}(x/q; a/q, b/q; q) &= (abq^{n+1} - aq^{n+1} - bq^{n+1} + 1)p_n(x; a, b; q) \\ &\quad + \alpha_n^{\text{bqL}} \Delta_n p_n(x; a, b; q), \end{aligned} \quad (4.40) \text{?raiupnbqL?}$$

$$\begin{aligned} (-1+a)(-1+b)q^{n+1} p_n(x/q; a/q, b/q; q) &= q(abq^n - aq^n - bq^n + 1)p_n(x; a, b; q) \\ &\quad + q^n \gamma_n^{\text{bqL}} \nabla_n p_n(x; a, b; q), \end{aligned} \quad (4.41) \text{?raidonbqL?}$$

$$(a-1)\phi(1) \mathcal{D}_{q,a} p_n(x; a, b; q) = \frac{a(1-q^n)}{q^{n-1}} (x-1)p_{n-1}(xq; aq, bq; q), \quad (4.42) \text{lowabqL}$$

$$\begin{aligned} (a-1) bq^{n+1} (aq-1) p_n(x; a/q, b; q) &= (aq^{n+1} - 1)(x-aq) \mathcal{D}_{q,a} p_n(x; a, b; q) \\ &\quad + ((1-aq)(x - (a-b+ab)q^{n+1}) + (aq^{n+1} - 1)(x-aq)) p_n(x; a, b; q), \end{aligned} \quad (4.43) \text{raiupbbqL}$$

$$\begin{aligned} (a-1)(aq-1)(bq^{n+1} - 1) p_{n+1}(x; a/q, b; q) &= (1-a)b(aq-1)q^{n+1} \mathcal{D}_{1/q,a} p_n(x; a, b; q) \\ &\quad + (aq-1)(x + abq^{n+1} - a - bq^{n+1}) p_n(x; a, b; q) \end{aligned} \quad (4.44) \text{raidobbqL}$$

where  $\alpha_n^{\text{bqL}}$  and  $\gamma_n^{\text{bqL}}$  are the big  $q$ -Laguerre recurrence relation coefficients (see Table 1), and

$$\mathcal{D}_{q,a} f(x, a) = \frac{f(x, qa) - f(x, a)}{a(q-1)}.$$

**Proof.** These identities can be checked by identifying the polynomial coefficients on the left and right-hand sides of each identity and with the help of Wolfram Mathematica 13<sup>®</sup>. ■

A direct consequence is the fact that this polynomial sequence is orthogonal with respect to the parameters  $a$  and  $b$ .

?{thm:4.6}?

**Theorem 4.15.** For any  $a, b \in \mathbb{C}$ , any  $x \in \mathbb{C}$ ,  $x \notin \{1, aq, bq\}$ , and any  $n \in \mathbb{N}_0$ , the polynomial sequences  $(p_n(x; aq^k, b; q))_k$  and  $(p_n(x; a, bq^k; q))_k$  are orthogonal with respect to certain moment functional.

**Proof.** By combining (4.42), (4.43) and (4.44) we have the following second order difference equations:

$$(x-1)a(1-q^n)p_n(x; a, b; q) = \mathbf{t}_n(a) \mathcal{D}_{1/q,a} \mathcal{D}_{q,a} p_n(x; a, b; q) - (\mathbf{t}_n(a) - \mathbf{t}_n^*(a)) \mathcal{D}_{q,a} p_n(x; a, b; q), \quad (4.45) \text{?eq:sodeM1?}$$

$$= \mathbf{t}_n^*(a) \mathcal{D}_{q,a} \mathcal{D}_{1/q,a} p_n(x; a, b; q) - (\mathbf{t}_n(a) - \mathbf{t}_n^*(a)) \mathcal{D}_{1/q,a} p_n(x; a, b; q), \quad (4.46) \text{?eq:sodeM2?}$$

where  $\mathbf{t}_n(a) = (aq^{n+1} - 1)(x - aq)$ , and  $\mathbf{t}_n^*(a) = (a-1)bq^{n+1}(aq-1)$ , which is connected with the big  $q$ -Jacobi polynomials case (see [14, (14.5.5)]). By using the theory of Sturm-Liouville, the result holds. The result also holds for the parameter  $b$  by symmetry. ■

From Lemma 4.14 we can deduce some new identities related to the big  $q$ -Laguerre polynomials.

?{thm:3.6}?

**Theorem 4.16.** For any  $a, b \in \mathbb{C}$ , and any  $n, k \in \mathbb{N}_0$ , the following Rodrigues-type identities hold:

$$p_n(q^k x; aq^k, bq^k; q) = \frac{(aq, bq; q)_k}{(x; q)_k} \nabla_n^k p_{n+k}(x; a, b; q), \quad (4.47) \text{?eq:bqLnRF?}$$

$$p_n(q^k x; aq^k, bq^k; q) = \frac{q^{nk} q^{\binom{k+1}{2}} (bq; q)_k}{(q^{n+1}, x; q)_k} \frac{(aq^{k-1} - 1)(1 - aq^k)}{aq^k} \mathcal{D}_{q,a} \cdots \frac{(a-1)(1-aq)}{aq} \mathcal{D}_{q,a} p_{n+k}(x; a, b; q). \quad (4.48) \text{?eq:bqLaRF?}$$

**Proof.** The identities holds by mathematical induction on  $k$  after a straightforward simplification and using (4.39) and (4.42). ■

The last result concerning the operators associated with the big  $q$ -Laguerre polynomials is as follows.

⟨pro:3⟩

**Proposition 4.17.** *The following identity holds:*

$$\frac{(cq^{n+1} - 1)(cq^{n+2} - 1)(\beta q^{n+1} - 1)(\beta q^{n+2} - 1)}{q^{4n}(cq - 1)(\beta q - 1)} p_{n+1}(q^4 x, aq, bq; q) = \sum_{k=0}^4 p_k(x) \mathcal{D}_q^k p_n(x; a, b; q),$$

where the polynomial coefficients are

$$\begin{aligned} p_0(x) &= \frac{1}{q} ((q^4 x - 1)(a + b + aq + bq + abq) + (a + b)(1 + q)(aq^2 - 1)(bq^2 - 1)), \\ p_1(x) &= \frac{(q-1)}{q^2} ((q^4 x - 1)(ab + a^2 q + 2abq + a^2 bq + b^2 q + ab^2 q + abq^2 + a^2 bq^2 + ab^2 q^2) \\ &\quad + (aq^2 - 1)(bq^2 - 1)(ab + a^2 q + 2abq + b^2 q + abq^2)), \\ p_2(x) &= \frac{ab(q-1)^2}{q^2} ((q^4 x - 1)(a + b + ab + aq + a^2 q + bq + 2abq + b^2 q + abq^2) \\ &\quad + (a + b)(1 + q)(aq^2 - 1)(bq^2 - 1)), \\ p_3(x) &= \frac{a^2 b^2 (q-1)^3}{q^2} ((q^4 x - 1)(1 + a + b + aq + bq) + (aq^2 - 1)(bq^2 - 1)), \\ p_4(x) &= \frac{a^3 b^3 (q-1)^4}{q^2} (q^4 x - 1). \end{aligned}$$

**Proof.** The proof follows after using an algorithm written in Wolfram Mathematica 13<sup>®</sup>. ■

## 4.5 The little $q$ -Laguerre/Wall polynomials

The little  $q$ -Laguerre polynomials can be written in terms of basic hypergeometric series as [14, §14.20]

$$p_n(x; a|q) = {}_2\phi_1 \left( \begin{matrix} q^{-n}, 0 \\ aq \end{matrix}; q, qx \right).$$

In this case  $\phi(x) = (1-x)x$  and  $\phi^*(x) = ax$ , i.e.,  $c = 0$ , and  $p_n(c; a|q) = 1$ . Taking this into account we can state the following result.

⟨lem:4.6⟩

**Lemma 4.18.** *For any  $a \in \mathbb{C}$ , and any  $n \in \mathbb{N}_0$ , the following identities hold:*

$$\frac{q^{1-n} x}{aq-1} p_{n-1}(x; aq|q) = \nabla_n p_n(x; a|q), \tag{4.49} \underline{\text{?lownlql?}}$$

$$(1-a)(x-1)p_n(x/q; a/q|q) = (aq^{n+1} - 1)\Delta_n p_n(x; a|q) - (ax + 1 - a)p_n(x; a|q), \tag{4.50} \underline{\text{?lownuplql?}}$$

$$(1-a)(x-1)p_{n-1}(x/q; a/q|q) = (ax + (q^n - 1)a)\nabla_n p_n(x; a|q) - (ax + 1 - a)p_n(x; a|q), \tag{4.51} \underline{\text{?lowndolql?}}$$

$$\frac{a(1 - q^n)x}{q^{n-1}(1-a)(1-aq)} p_{n-1}(x; aq|q) = \mathcal{D}_{1/q,a} p_n(x; a|q), \tag{4.52} \underline{\text{?lowalql?}}$$

$$\begin{aligned} (a-1)q^n(aq-1)p_n(x; a/q|q) &= (aq^{n+1} - 1)x\mathcal{D}_{1/q,a} p_n(x; a|q) \\ &\quad + (aq(q^n - 1)x + q^n(aq - 1)(a - 1))p_n(x; a|q), \end{aligned} \tag{4.53} \underline{\text{?raiadolql?}}$$

$$\begin{aligned} a(a-1)q^{n-1}(q^n - 1)p_{n-1}(x; a/q|q) &= (a-1)(x + aq^{2n-1} - aq^{n-1})\mathcal{D}_{q,a} p_n(x; a|q) \\ &\quad + (a(q^n - 1)x + (1-a)aq^{n-1}(q^n - 1))p_n(x; a|q). \end{aligned} \tag{4.54} \underline{\text{?raiadolql?}}$$

**Proof.** These identities can be checked by identifying the polynomial coefficients on the left and right-hand sides of each identity and with the help of Wolfram Mathematica 13<sup>®</sup>. ■

A direct consequence is the fact that this polynomial sequence is orthogonal with respect to the parameter  $a$ .

?{thm:4.8}?

**Theorem 4.19.** *For any  $a \in \mathbb{C}$ , any  $x \in \mathbb{C}$ ,  $x \notin \{1, aq\}$ , and any  $n \in \mathbb{N}_0$ , the polynomial sequence  $(p_n(x; aq^k|q))_k$  is orthogonal with respect to certain moment functional.*

**Proof.** By combining (4.42), (4.43) and (4.44) we have the following second-order difference equations:

$$aqx(q^n - 1)p_n(x; a|q) = s_n(a)\mathcal{D}_{1/q,a}\mathcal{D}_{q,a}p_n(x; a|q) - (s_n(a) - s_n^*(a))\mathcal{D}_{q,a}p_n(x; a|q) \quad (4.55) \text{?eq:sodelqL1?}$$

$$= s_n^*(a)\mathcal{D}_{q,a}\mathcal{D}_{1/q,a}p_n(x; a|q) - (s_n(a) - s_n^*(a))\mathcal{D}_{1/q,a}p_n(x; a|q) \quad (4.56) \text{?eq:sodelqL2?}$$

where  $s_n(a) = (a - 1)q^n(aq - 1)$ , and  $s_n^*(a) = (1 - aq^{n+1})x$ , which is connected with the big  $q$ -Laguerre polynomials case (see [14, (14.11.5)]). Using the theory of discrete Sturm-Liouville and assuming that for any  $n \in \mathbb{N}_0$ , in this case,  $a$  is variable, and the rest of the elements are fixed, the result holds.  $\blacksquare$

The last result concerning the operators associated with the little  $q$ -Laguerre polynomials is as follows.

?{pro:4}?

**Proposition 4.20.** *The following identity holds:*

$$\frac{(aq^{n+1} - 1)(aq^{n+2} - 1)}{q^{3n}(q - 1)(aq - 1)}p_{n+1}(q^4x, aq|q) = \sum_{k=0}^4 p_k(x)\mathcal{D}_q^k p_n(x; a|q)$$

where the polynomial coefficients are

$$p_0(x) = \frac{1}{q^3}(q^4x(aq^3 - aq - 1 - q - q^2) + (aq^2 - 1)(aq^3 - 1 - q - q^2)),$$

$$p_1(x) = \frac{(q-1)}{q^5}(aq^9x^2 + q^4x(a^2q^4 + a^2q^3 - 1 - a - q - 2aq - q^2 - aq^2 + aq^3 + aq^4) + (aq^2 - 1)(aq^4 - 1 - q - q^2 + aq^3)),$$

$$p_2(x) = \frac{(q-1)^2}{q^6}(aq^8x^2(1 + q + aq) + q^3x(a^2q^5 + 2a^2q^4 - a - q - 2aq - 2aq^2 + a^2q^3 + aq^4) + (aq^2 - 1)(aq^3 - 1)),$$

$$p_3(x) = \frac{a(q-1)^3x}{q^4}(q^5x(1 + a + aq) + (1 + q)(-1 + aq^3)),$$

$$p_4(x) = a^2(q-1)^4x^2.$$

**Proof.** The proof follows after using an algorithm written in Wolfram Mathematica 13<sup>®</sup>.  $\blacksquare$

## 4.6 The Stieltjes-Wigert polynomials

These polynomials can be written in terms of basic hypergeometric series as [14, §14.21]

$$S_n(x; q) = {}_1\phi_1\left(\begin{matrix} q^{-n} \\ 0 \end{matrix}; q, -q^{n+1}x\right). \quad (4.57) \text{?sw-def?}$$

In this case  $\phi(x) = x^2$  and  $\phi^*(x) = x$ , i.e.,  $c = 0$ , and  $S_n(c; q) = 1$ . Taking this into account we can state the following result.

{lem:4.7}

**Lemma 4.21.** *For any  $n \in \mathbb{N}_0$ , the following identities hold:*

$$\nabla_n S_n(x; q) = -q^n x S_{n-1}(qx; q), \quad (4.58) \text{rainSW}$$

$$S_n(x; q) + q^{-n-1} \Delta_n S_n(x; q) = S_{n+1}(x/q; q), \quad (4.59) \text{lowupnSW}$$

$$S_n(x; q) - (1 - q^{-n}) \nabla_n S_n(x; q) = S_n(x/q; q). \quad (4.60) \text{?lowdownSW?}$$

From Lemma 4.21 we can deduce some a new identities related to the Stieltjes-Wigert polynomials.

?{thm:4.9}?

**Theorem 4.22.** For any  $n, k \in \mathbb{N}_0$ , the following Rodrigues-type identities hold:

$$S_n(x; q) = \frac{1}{(-x)^k} q^{-n} \nabla_n \cdots q^{-n} \nabla_n S_{n+k}(x/q^k; q), \quad (4.61) \text{ ?SWnRF?}$$

$$S_n(x; q) = (-1)^n (q; q)_n q^{-n} \Delta_n \cdots q^{-n+k} \Delta_n \frac{(-1)^n}{(q; q)_{n-k}} S_{n-k}(xq^k; q). \quad (4.62) \text{ ?SWnRF2?}$$

**Proof.** The first identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.58).

The second identity holds by mathematical induction on  $k$  after a straightforward simplification and using (4.59) written in the following way:

$$\frac{(-q)^n}{(q; q)_n} S_n(x; q) = \Delta_n \frac{(-1)^n}{(q; q)_{n-1}} S_{n-1}(xq; q).$$

■

The last result concerning the operators associated with the Stieltjes-Wigert polynomials is as follows.

?{pro:5}?

**Proposition 4.23.** The following identity holds:

$$(q-x)S_n(x; q) + (1-q)x\mathcal{D}_{1/q}S_n(x; q) = qS_{n+1}(x/q^2; q).$$

**Proof.** The proof follows after using an algorithm written in Wolfram Mathematica 13<sup>®</sup>.

■

## A Code in wolfram Mathematica<sup>®</sup> to check the identities

We are going to present different codes we can use to check all the identities presented in this work. Let us start with expression (4.5):

```
1 Table[LaguerreL[n, a, x] - LaguerreL[n-1, a, x] - LaguerreL[n, a-1, x], {n, 1, 6}] // Factor
```

One can check this for all the terms as one desires; using similar expressions one can check (4.9), (4.10) and the rest of the expressions where one uses the operators  $\nabla_n$  or  $\Delta_n$ .

In an analogous way one can check, for instance, (4.29):

```
1 mep[n_, b_, c_, x_] := Hypergeometric2F1[-n, -x, b, 1-1/c]; Table[mep[n, b, c, x] - mep[n, b-1, c, x] - x n(c-1)/(b-b^2)/c mep[n-1, b+1, c, x-1], {n, 1, 6}] // Factor
```

One can check identity used in the proof Theorem 4.7:

```
1 chp[n_, a_, x_] := HypergeometricPFQ[{-n, -x}, {}, -1/a]; Table[a^{n+1}/(n+1)! - chp[n+1, a, x] - a^n/n! chp[n, a, x] - a^{n+1}/(n+1)! chp[n+1, a, x+1], {n, 1, 6}] // Factor
```

We also used different codes to know what kind of expression we expect

```
1 m:=4; Solve[CoefficientList[mep[m+1, b, c, x] - mep[m, b, c, x] - (Ax+B)mep[m, bb, c, x-dd], x] == Table[0, {i, 1, m+2}], {A, B, bb, dd}]
```

obtaining in such a case

```
1 {{A -> (-1+c)/(bc), B->0, bb->1+b, dd->1}}
```

Let me finish this sequence of expressions with the only one challenging related with the big  $q$ -Laguerre polynomial expression, proposition 4.17:

	$\alpha_n$	$\beta_n$	$\gamma_n$	$d_n^2$
L	$-n - \alpha - 1$	$2n + \alpha + 1$	$-n$	$\frac{n! \Gamma^2(\alpha + 1)}{\Gamma(n + \alpha + 1)}$
C	$-a$	$n + a$	$-n$	$\frac{n!}{a^n}$
M	$\frac{c(n + \beta)}{c - 1}$	$-\frac{n + c(n + \beta)}{c - 1}$	$\frac{n}{c - 1}$	$\frac{n! \Gamma(\beta)}{\Gamma(\beta + n) c^n}$
M	$\frac{n + \beta}{c - 1}$	$-\beta - \alpha_n - \gamma_n$	$\frac{nc}{c - 1}$	$\frac{n! \Gamma(\beta) c^n}{\Gamma(\beta + n)}$
bqL	$(1 - aq^{n+1})(1 - bq^{n+1})$	$1 - \alpha_n - \gamma_n$	$abq^{n+1}(q^n - 1)$	$\frac{(q^{-1}; q^{-1})_n q^n}{(a^{-1}q^{-1}, b^{-1}q^{-1}; q^{-1})_n}$
qM	$\frac{(c + q^{n+1})(bq^{n+1} - 1)}{q^{2n+1}}$	$bq - \alpha_n - \gamma_n$	$\frac{cq(q^n - 1)}{q^{2n+1}}$	$\frac{(q; q)_n}{(bq, -c^{-1}q; q)_n q^n}$
lqL	$q^n(aq^{n+1} - 1)$	$-\alpha_n - \gamma_n$	$aq^n(q^n - 1)$	$\frac{(q; q)_n a^n q^n}{(aq; q)_n}$
qL	$\frac{q^{n+1+\alpha} - 1}{q^{2n+1+\alpha}}$	$-\alpha_n - \gamma_n$	$\frac{q(q^n - 1)}{q^{2n+\alpha+1}}$	$\frac{(q; q)_n}{(q^{(\alpha+1)}; q)_n q^n}$
qC	$-\frac{a + q^{n+1}}{q^{2n+1}}$	$-\alpha_n - \gamma_n$	$\frac{aq(q^n - 1)}{q^{2n+1}}$	$\frac{(q; q)_n}{(-a^{-1}q, q)_n q^n}$
OLB	$-aq^{2n+1}$	$-\alpha_n - \gamma_n$	$-aq^n(q^n - 1)$	$\frac{(q, q)_n a^n q^n}{(aq, q)_n}$
SW	$\frac{-1}{q^{2n+1}}$	$-\alpha_n - \gamma_n$	$\frac{q(q^n - 1)}{q^{2n+1}}$	$\frac{(q; q)_n}{q^n}$

**Table 1.** Essential data of the families belonging to the Laguerre constellation.



```

1 dfqu [k_, f_] := If [k==0, f, If [k==1, Return [(f/.{x->q x})-f]/(qx-x)], Return
  [dfqu [k-1, dfqu [1, f]]]];
2 bqLp [n_, a_, b_, x_] := QHypergeometricPFQ [{q^{-n}, 0, x}, {a q, b q}, q, q];
3 AAbqL [k_, a_, b_, x_] := A [k] bqLp [k+1, aq, bq, Bx] - Sum [AAbqL [1, a, b, x] dfqu [1,
  bqLp [k, a, b, x]], {1, 0, k-1}]/dfqu [k, bqLp [k, a, b, x]]//FunctionExpand//
  Factor

```

By using the Solve & coefficientList former algorithm modified to be used for the big  $q$ -Laguerre polynomials one used previously one can obtain the leading term  $A[k]$ , and assuming the expression should be finite is not hard to obtain  $B \rightarrow q^4$ .

## References

- [mr316772] [1] W. A. Al-Salam and T. S. Chihara. Another characterization of the classical orthogonal polynomials. *SIAM J. Math. Anal.*, 3:65–70, 1972. doi:10.1137/0503007.
- [mr2293537] [2] M. Alfaro and R. Álvarez Nodarse. A characterization of the classical orthogonal discrete and  $q$ -polynomials. *J. Comput. Appl. Math.*, 201:48–54, 2007. doi:10.1016/j.cam.2006.01.031.
- [mr2241592] [3] R. Álvarez Nodarse. On characterizations of classical polynomials. *J. Comput. Appl. Math.*, 196:320–337, 2006. doi:10.1016/j.cam.2005.06.046.
- [mr1771452] [4] R. Álvarez Nodarse and J. Arvesú. On the  $q$ -polynomials in the exponential lattice  $x(s) = c_1 q^s + c_3$ . *Integral Transforms Spec. Funct.*, 8:299–324, 1999. doi:10.1080/10652469908819236.
- [MR1850541] [5] R. Álvarez Nodarse and J. C. Medem.  $q$ -classical polynomials and the  $q$ -Askey and Nikiforov-Uvarov tableaux. *J. Comput. Appl. Math.*, 135:197–223, 2001. doi:10.1016/S0377-0427(00)00585-9.
- [mr0481884] [6] T. S. Chihara. *An introduction to orthogonal polynomials*. Mathematics and its Applications, Vol. 13. Gordon and Breach Science Publishers, New York-London-Paris, 1978. Mathematics and its Applications, Vol. 13, Reprinted by Dover Publications.
- [Jacob.tPhD] [7] J. S. Christiansen. Indeterminate moment problems within the askey-scheme. Master’s thesis, University of Copenhagen, Denmark, 2004.
- [MR2653053] [8] R. S. Costas-Santos and F. Marcellán.  $q$ -classical orthogonal polynomials: a general difference calculus approach. *Acta Appl. Math.*, 111(2):107–128, 2010. doi:10.1007/s10440-009-9536-z.
- [mr2741220] [9] R. S. Costas-Santos and J. J. Moreno-Balcázar. The semiclassical Sobolev orthogonal polynomials: a general approach. *J. Approx. Theory*, 163:65–83, 2011. doi:10.1016/j.jat.2010.03.001.
- [MR2494714] [10] R. S. Costas-Santos and J. F. Sánchez-Lara. Extensions of discrete classical orthogonal polynomials beyond the orthogonality. *J. Comput. Appl. Math.*, 225:440–451, 2009. doi:10.1016/j.cam.2008.07.055.
- [mr1340932] [11] A. G. García, F. Marcellán, and L. Salto. A distributional study of discrete classical orthogonal polynomials. In *Proceedings of the Fourth International Symposium on Orthogonal Polynomials and their Applications (Evian-Les-Bains, 1992)*, volume 57, pages 147–162, 1995. doi:10.1016/0377-0427(93)E0241-D.
- [MR4496948] [12] M. E. H. Ismail, R. Zhang, and K. Zhou.  $q$ -fractional Askey-Wilson integrals and related semigroups of operators. *Physica D*, 442:Paper No. 133534, 15, 2022. doi:10.1016/j.physd.2022.133534.
- [ZBMATH2576281] [13] K. Knopp. Theory and application of infinite series. London: Blackie & Sons. XII, 571 p. (1928)., 1928.
- [mr2656096] [14] R. Koekoek, P. A. Lesky, and René F. Swarttouw. *Hypergeometric orthogonal polynomials and their  $q$ -analogues*. Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder. doi:10.1007/978-3-642-05014-5.

- 
- [MR2217787](#) [15] W. Koepf and M. Masjed-Jamei. A generic formula for the values at the boundary points of monic classical orthogonal polynomials. *J. Comput. Appl. Math.*, 191:98–105, 2006. doi:[10.1016/j.cam.2005.04.024](#).
- [MR0150565](#) [16] P. P. Korovkin. *Linear operators and approximation theory*. Russian Monographs and Texts on Advanced Mathematics and Physics, Vol. III. Gordon and Breach Publishers, Inc., New York; Hindustan Publishing Corp. (India), Delhi, 1960. Translated from the Russian ed. (1959).
- [mr1273613](#) [17] F. Marcellán, A. Branquinho, and J. Petronilho. Classical orthogonal polynomials: a functional approach. *Acta Appl. Math.*, 34:283–303, 1994. doi:[10.1007/BF00998681](#).
- [mr1850540](#) [18] J. C. Medem, R. Álvarez Nodarse, and F. Marcellán. On the  $q$ -polynomials: a distributional study. *J. Comput. Appl. Math.*, 135:157–196, 2001. doi:[10.1016/S0377-0427\(00\)00584-7](#).
- [mr1421640](#) [19] A. F. Nikiforov and V. B. Uvarov. Polynomial solutions of hypergeometric type difference equations and their classification. *Integral Transforms Spec. Funct.*, 1:223–249, 1993. doi:[10.1080/10652469308819023](#).
- [mr2324843](#) [20] J. Petronilho. Generic formulas for the values at the singular points of some special monic classical  $H_{q,\omega}$ -orthogonal polynomials. *J. Comput. Appl. Math.*, 205:314–324, 2007. doi:[10.1016/j.cam.2006.05.005](#).