

A family of convolution operators, part one

Zipeng Wang

Abstract

We study a family of convolution operators defined in \mathbb{R}^n . The regarding Fourier multipliers are defined in terms of distributions having singularity on \mathbb{S}^{n-1} . We prove the L^p -boundedness of these operators.

1 Introduction

Let $\alpha \in \mathbb{C}$. As was first investigated by Strichartz [5], Ω^α is a distribution defined in \mathbb{R}^n by analytic continuation from

$$\operatorname{Re}\alpha > \frac{n-1}{2}, \quad \Omega^\alpha(x) = \pi^{\alpha - \frac{n+1}{2}} \Gamma^{-1}\left(\alpha - \frac{n-1}{2}\right) \left(\frac{1}{1 - |x|^2} \right)_+^{\frac{n+1}{2} - \alpha}. \quad (1.1)$$

Equivalently, it can be defined as

$$\begin{aligned} \widehat{\Omega}^\alpha(\xi) &= \left(\frac{1}{|\xi|} \right)^{\frac{n}{2} - [\frac{n+1}{2} - \alpha]} J_{\frac{n}{2} - [\frac{n+1}{2} - \alpha]}(2\pi|\xi|) \\ &= \left(\frac{1}{|\xi|} \right)^{\alpha - \frac{1}{2}} J_{\alpha - \frac{1}{2}}(2\pi|\xi|), \quad \alpha \in \mathbb{C}. \end{aligned} \quad (1.2)$$

This family of distributions arises from the study of the propagation of waves.

- ◊ Throughout, we regard $\mathfrak{B} > 0$ as a generic constant whose value depends on its sub-indices.
- ◊ We always write $c > 0$ for some fixed, large enough constant.

Consider

$$\widehat{\mathbf{P}}^\alpha(\xi) = \widehat{\phi}(\xi) \int_{\mathbb{R}} e^{-2\pi i r} \widehat{\Omega}^\alpha(r\xi) \omega(r) |r|^{2\alpha-1} dr \quad (1.3)$$

where $\widehat{\phi}$ is a radially symmetric, smooth cut-off function supported in the shell: $\frac{1}{3} < |\xi| \leq 3$. On the other hand, $\omega \in C^\infty(\mathbb{R})$ satisfies

$$|\omega(r)| \leq \mathfrak{B} [1 + |r|]^{-1}. \quad (1.4)$$

Moreover, we require $\omega(r) \sim r^{-1}$ in the sense of

$$\omega(r) - r^{-1} = \mathbf{O}(r^{-2}), \quad |r| \rightarrow \infty. \quad (1.5)$$

Define

$$\mathbf{I}^\alpha f(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{f}(\xi) \widehat{\mathbf{P}}^\alpha(\xi) d\xi. \quad (1.6)$$

Theorem One For $\frac{n-1}{n} < \operatorname{Re}\alpha < 1$, we have

$$\|\mathbf{I}^\alpha f\|_{L^p(\mathbb{R}^n)} \leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)}, \quad \frac{n-1}{2n} \leq \frac{1}{p} \leq \frac{n+1}{2n}. \quad (1.7)$$

Remark 1.1 We leave the implication of **Theorem One** to the second part of this project.

The remaining paper is organized as follows. In the next section, we develop 3 lemmas, stated as **Lemma One**, **Lemma Two** and **Lemma Three**. Then, we prove **Theorem One** by using these lemmas within an interpolation argument.

Section 5, 6 and 7 are devoted to the proof of **Lemma One**, **Lemma Two** and **Lemma Three** respectively. We add 2 appendices in the end for certain preliminary estimates.

2 Proof of Theorem One

Let $\widehat{\mathbf{P}}^\alpha$ defined in (1.3)-(1.5) for $\frac{n-1}{n} < \operatorname{Re}\alpha < 1$. We assert

$$\begin{aligned} \widehat{\mathbf{P}}^\alpha(\xi) &= \sum_{j \in \mathbb{Z}} \widehat{\mathbf{P}}_j^\alpha(\xi), \\ \widehat{\mathbf{P}}_j^\alpha(\xi) &= \widehat{\phi}(\xi) \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r \xi} \widehat{\Omega}^\alpha(r\xi) \omega(r) |r|^{2\alpha-1} dr. \end{aligned} \quad (2.1)$$

From (1.2), we find

$$\begin{aligned} \widehat{\Omega}^\alpha(r\xi) |r|^{2\alpha-1} &= \left(\frac{1}{|r\xi|} \right)^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(2\pi|r\xi|) |r|^{2\alpha-1} \\ &= \pi^{\alpha-1} \Gamma(\alpha) \left\{ \int_{-1}^1 e^{2\pi i |\xi| rs} (1-s^2)^{\alpha-1} ds \right\} |r|^{2\alpha-1} \quad \text{by (A.1)} \end{aligned} \quad (2.2)$$

which is locally integrable in r .

Recall $\mathbf{I}^\alpha f$ defined in (1.6). We have

$$\widehat{\mathbf{I}^\alpha f}(\xi) = \widehat{f}(\xi) \widehat{\mathbf{P}}^\alpha(\xi). \quad (2.3)$$

Define

$$\begin{aligned} \widehat{\mathbf{I}_<^\alpha f}(\xi) &= \widehat{f}(\xi) \sum_{j \leq 0} \widehat{\mathbf{P}}_j^\alpha(\xi) \\ &= \widehat{f}(\xi) \widehat{\phi}(\xi) \int_{-1}^1 e^{-2\pi i r \xi} \widehat{\Omega}^\alpha(r\xi) \omega(r) |r|^{2\alpha-1} dr. \end{aligned} \quad (2.4)$$

On the other hand, for $j > 0$, we introduce an ingenious dyadic decomposition implicitly given by Fefferman [1] and later refined by Seeger, Sogge and Stein [2] in their study of Fourier integral operators.

Given $j > 0$, $\{\xi_j^\nu\}_\nu$ is a collection of points equally distributed on \mathbb{S}^{n-1} having a grid length between $2^{-j/2-1}$ and $2^{-j/2}$. Let $\varphi \in C_0^\infty(\mathbb{R})$ such that $\varphi(t) = 1$ for $|t| \leq 1$ and $\varphi(t) = 0$ for $|t| > 2$. Define

$$\varphi_j^\nu(\xi) = \frac{\varphi\left[2^{j/2}\left|\frac{\xi}{|\xi|} - \xi_j^\nu\right|\right]}{\sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \varphi\left[2^{j/2}\left|\frac{\xi}{|\xi|} - \xi_j^\nu\right|\right]} \quad (2.5)$$

whose support is contained in

$$\Gamma_j^\nu = \left\{ \xi \in \mathbb{R}^n : \left| \frac{\xi}{|\xi|} - \xi_j^\nu \right| < 2^{-j/2+1} \right\}. \quad (2.6)$$

We write

$$\widehat{\mathbf{P}}_j^\alpha(\xi) = \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \widehat{\mathbf{P}}_j^{\alpha \nu}(\xi), \quad \widehat{\mathbf{P}}_j^{\alpha \nu}(\xi) = \varphi_j^\nu(\xi) \widehat{\mathbf{P}}_j^\alpha(\xi). \quad (2.7)$$

Let $\xi_j^\mu \in \{\xi_j^\nu\}_\nu$ for some μ . \mathbf{L}_μ is an $(n \times n)$ -orthogonal matrix with $\det \mathbf{L}_\mu = 1$. Moreover,

$$\mathbf{L}_\mu^T \xi_j^\mu = (1, 0)^T \in \mathbb{R} \times \mathbb{R}^{n-1}. \quad (2.8)$$

We define

$$\begin{aligned} \Psi_j^\alpha(x) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x), \\ \Psi_j^{\alpha \mu}(x) &= \varphi\left[2^{-\epsilon j}\left|2^j - (\mathbf{L}_\mu^T x)_1\right|\right] \varphi\left[2^{-\epsilon j}\left|2^j + (\mathbf{L}_\mu^T x)_1\right|\right] \prod_{i=2}^n \varphi\left[2^{-[\frac{1}{2}+\epsilon]j}\left|(\mathbf{L}_\mu^T x)_i\right|\right] \end{aligned} \quad (2.9)$$

where $\epsilon = \epsilon(\operatorname{Re}\alpha) > 0$ is a small, adjustable number.

Let

$$\mathcal{U}_j^{\alpha \mu} = \left\{ \nu: \left| \xi_j^\mu \pm \xi_j^\nu \right| > c 2^{-j/2} 2^{\epsilon j} \right\}, \quad \mathcal{V}_j^{\alpha \mu} = \left\{ \nu: \left| \xi_j^\mu \pm \xi_j^\nu \right| \leq c 2^{-j/2} 2^{\epsilon j} \right\}. \quad (2.10)$$

We consider

$$\begin{aligned} \Pi^\alpha f(x) &= \sum_{j>0} \Pi_j^\alpha f(x), \quad \Pi_j^\alpha f(x) = \int_{\mathbb{R}^n} f(x-u) \mathbf{U}_j^\alpha(u) du, \\ \mathbf{U}_j^\alpha(u) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} \mathbf{P}_j^{\alpha \nu}(u) + \mathbf{P}_j^\alpha(u) [1 - \Psi_j^\alpha(u)] \end{aligned} \quad (2.11)$$

and

$$\begin{aligned} \text{III}^\alpha f(x) &= \sum_{j>0} \text{III}_j^\alpha f(x), \quad \text{III}_j^\alpha f(x) = \int_{\mathbb{R}^n} f(x-u) \mathbf{V}_j^\alpha(u) du, \\ \mathbf{V}_j^\alpha(u) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} \mathbf{P}_j^{\alpha \nu}(u). \end{aligned} \quad (2.12)$$

Remark 2.1 From (2. 4) and (2. 11)-(2. 12), we find

$$\mathbf{I}^\alpha f(x) = \mathbf{I}_<^\alpha f(x) + \mathbf{II}^\alpha f(x) + \mathbf{III}^\alpha f(x). \quad (2. 13)$$

$\# \Omega^\alpha$ is a distribution defined by analytic continuation from

$$\operatorname{\mathbf{Re}}\alpha > \frac{n}{2}, \quad \# \Omega^\alpha(x) = \pi^{\alpha - \frac{n+2}{2}} \Gamma^{-1}\left(\alpha - \frac{n}{2}\right) \left(\frac{1}{1 - |x|^2} \right)_+^{\frac{n+2}{2} - \alpha}. \quad (2. 14)$$

Equivalently, it can be defined by

$$\begin{aligned} \# \widehat{\Omega}^\alpha(\xi) &= \left(\frac{1}{|\xi|} \right)^{\frac{n}{2} - [\frac{n+2}{2} - \alpha]} J_{\frac{n}{2} - [\frac{n+2}{2} - \alpha]}(2\pi|\xi|) \\ &= \left(\frac{1}{|\xi|} \right)^{\alpha-1} J_{\alpha-1}(2\pi|\xi|), \quad \alpha \in \mathbb{C}. \end{aligned} \quad (2. 15)$$

Given $j \in \mathbb{Z}$, we define

$$\begin{aligned} \# \widehat{\mathbf{P}}^{\alpha \beta}(\xi) &= \sum_{j \in \mathbb{Z}} \# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi), \\ \# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) &= \widehat{\phi}(\xi) \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r \# \widehat{\Omega}^\alpha(r\xi)} \omega(r) |r|^{2\beta-1} dr, \quad \alpha, \beta \in \mathbb{C}. \end{aligned} \quad (2. 16)$$

Remark 2.2 Suppose $\operatorname{\mathbf{Re}}\alpha \geq [\frac{2n}{2n-1}] \operatorname{\mathbf{Re}}\beta$ and $\frac{2n-1}{2n} < \operatorname{\mathbf{Re}}\beta < \frac{2n-1}{2n-2}$. We have

$$|\# \widehat{\mathbf{P}}^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{\mathbf{Re}}\alpha \operatorname{\mathbf{Re}}\beta} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\alpha|} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2} - \varepsilon}, \quad |\xi| \neq 1 \quad (2. 17)$$

for some $\varepsilon = \varepsilon(\operatorname{\mathbf{Re}}\alpha, \operatorname{\mathbf{Re}}\beta) > 0$.

Lemma One Suppose $\operatorname{\mathbf{Re}}\alpha \geq [\frac{2n}{2n-1}] \operatorname{\mathbf{Re}}\beta$ and $\frac{2n-1}{2n} < \operatorname{\mathbf{Re}}\beta < \frac{2n-1}{2n-2}$. We have

$$\sum_{j \leq 0} |\# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{\mathbf{Re}}\alpha \operatorname{\mathbf{Re}}\beta} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\alpha|}. \quad (2. 18)$$

Given $j > 0$, we have

$$|\# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{\mathbf{Re}}\alpha \operatorname{\mathbf{Re}}\beta} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\alpha|} 2^{j/2} 2^{-j\varepsilon}, \quad |1 - |\xi|| \leq 2^{-j} \quad (2. 19)$$

and

$$|\# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{\mathbf{Re}}\alpha \operatorname{\mathbf{Re}}\beta} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\alpha|} e^{\mathbf{C}|\operatorname{\mathbf{Im}}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} 2^{-j\varepsilon}, \quad |1 - |\xi|| > 2^{-j} \quad (2. 20)$$

for some $\varepsilon = \varepsilon(\operatorname{\mathbf{Re}}\alpha, \operatorname{\mathbf{Re}}\beta) > 0$.

Let φ_j^ν defined in (2. 5). Write

$$\#\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) = \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \#\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\xi), \quad \#\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\xi) = \varphi_j^\nu(\xi) \#\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi). \quad (2. 21)$$

Similar to (2. 11)-(2. 12), we assert

$$\#\mathbf{U}_j^{\alpha \beta}(u) = \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} \#\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(u) + \#\widehat{\mathbf{P}}_j^{\alpha \beta}(u) [1 - \Psi_j^\alpha(u)] \quad (2. 22)$$

and

$$\#\mathbf{V}_j^{\alpha \beta}(u) = \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} \#\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(u). \quad (2. 23)$$

Lemma Two Suppose $\operatorname{Re}\alpha \geq [\frac{2n}{2n-1}] \operatorname{Re}\beta$ and $\frac{2n-1}{2n} < \operatorname{Re}\beta < \frac{2n-1}{2n-2}$. Given $j > 0$, we have

$$|\#\widehat{\mathbf{U}}_j^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} e^{\mathbf{C}|\operatorname{Im}\beta|} 2^{j/2} 2^{-j\varepsilon} \quad (2. 24)$$

and

$$|\#\widehat{\mathbf{V}}_j^{\alpha \beta}(\xi)| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} e^{\mathbf{C}|\operatorname{Im}\beta|} 2^{-j\varepsilon} \quad (2. 25)$$

for some $\varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0$.

${}^b\Omega^\alpha$ is a distribution defined by analytic continuation from

$$\operatorname{Re}\alpha > 0, \quad {}^b\Omega^\alpha(x) = \pi^{\alpha-1} \Gamma^{-1}(\alpha) \left(\frac{1}{1-|x|^2} \right)_+^{1-\alpha}. \quad (2. 26)$$

Alternatively, it is defined by

$$\begin{aligned} {}^b\widehat{\Omega}^\alpha(\xi) &= \left(\frac{1}{|\xi|} \right)^{\frac{n}{2}-(1-\alpha)} J_{\frac{n}{2}-(1-\alpha)}(2\pi|\xi|) \\ &= \left(\frac{1}{|\xi|} \right)^{\frac{n-1}{2}+\alpha-\frac{1}{2}} J_{\frac{n-1}{2}+\alpha-\frac{1}{2}}(2\pi|\xi|), \quad \alpha \in \mathbb{C}. \end{aligned} \quad (2. 27)$$

Given $j \in \mathbb{Z}$, we define

$$\begin{aligned} {}^b\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) &= \sum_{j \in \mathbb{Z}} {}^b\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi), \\ {}^b\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) &= \widehat{\phi}(\xi) \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r {}^b\widehat{\Omega}^\alpha(r\xi)} \omega(r) |r|^{2\beta-1} dr, \quad \alpha, \beta \in \mathbb{C}. \end{aligned} \quad (2. 28)$$

Let φ_j^ν defined in (2. 5). Write

$$\#{^b\widehat{\mathbf{P}}_j^{\alpha \beta}}(\xi) = \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} {}^b\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\xi), \quad {}^b\widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\xi) = \varphi_j^\nu(\xi) {}^b\widehat{\mathbf{P}}_j^{\alpha \beta}(\xi). \quad (2. 29)$$

Similar to (2. 11)-(2. 12), we assert

$${}^b\mathbf{U}_j^{\alpha \beta}(u) = \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} {}^b\mathbf{P}_j^{\alpha \beta \nu}(u) + {}^b\mathbf{P}_j^{\alpha \beta}(u) [1 - \Psi_j^\alpha(u)] \quad (2. 30)$$

and

$${}^b\mathbf{V}_j^{\alpha \beta}(u) = \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} {}^b\mathbf{P}_j^{\alpha \beta \nu}(u). \quad (2. 31)$$

Lemma Three Suppose $\operatorname{Re}\alpha > 0$ and $0 < \operatorname{Re}\beta < \frac{1}{2}$. We have

$$\int_{\mathbb{R}^n} \left| \sum_{j \leq 0} {}^b\mathbf{P}_j^{\alpha \beta}(x) \right| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|}. \quad (2. 32)$$

Given $j > 0$, we have

$$\int_{\mathbb{R}^n} \left| {}^b\mathbf{U}_j^{\alpha \beta}(x) \right| dx \leq \mathfrak{B}_{N \operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-jN} 2^{-j\varepsilon}, \quad N \geq 0 \quad (2. 33)$$

and

$$\int_{\mathbb{R}^n} \left| {}^b\mathbf{V}_j^{\alpha \beta}(x) \right| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-j\varepsilon} \quad (2. 34)$$

for some $\varepsilon = \varepsilon(\operatorname{Re}\alpha) > 0$.

Remark 2.3 There are $\mathbf{a}_1 > 0, 0 < \mathbf{b}_1 < \frac{1}{2}$ and $\mathbf{a}_2 \geq [\frac{2n}{2n-1}] \mathbf{b}_2, \frac{2n-1}{2n} < \mathbf{b}_2 < \frac{2n-1}{2n-2}$ such that $\operatorname{Re}\alpha = \frac{\mathbf{a}_1}{n} + \mathbf{a}_2 \left(\frac{n-1}{n} \right) = \frac{\mathbf{b}_1}{n} + \mathbf{b}_2 \left(\frac{n-1}{n} \right)$. As $\operatorname{Re}\alpha \rightarrow 1$, we must have $\mathbf{b}_1 \rightarrow \frac{1}{2}, \mathbf{b}_2 \rightarrow \frac{2n-1}{2n-2}$ together with $\mathbf{a}_1 \rightarrow 0, \mathbf{a}_2 \rightarrow \frac{n}{n-1}$.

Denote $\mathbf{a} = (\mathbf{a}_1, \mathbf{a}_2)$ and $\mathbf{b} = (\mathbf{b}_1, \mathbf{b}_2)$ for $\mathbf{a}_i, \mathbf{b}_i \in \mathbb{R}, i = 1, 2$. For $0 \leq \operatorname{Re}z \leq 1$, we define

$$\begin{aligned} \widehat{\Omega}_z^{\alpha \mathbf{a}}(\xi) &= \left(\frac{1}{|\xi|} \right)^{\mathbf{a}_1 z + \mathbf{a}_2(1-z) - \frac{1}{2} + \left(\frac{n-1}{2} \right) z - \frac{1}{2}(1-z) + i\operatorname{Im}\alpha} J_{\mathbf{a}_1 z + \mathbf{a}_2(1-z) - \frac{1}{2} + \left(\frac{n-1}{2} \right) z - \frac{1}{2}(1-z) + i\operatorname{Im}\alpha} (2\pi|\xi|) \\ &= \pi^{\mathbf{a}_1 z + \mathbf{a}_2(1-z) - 1 + \left(\frac{n-1}{2} \right) z - \frac{1}{2}(1-z) + i\operatorname{Im}\alpha} \Gamma^{-1} \left(\mathbf{a}_1 z + \mathbf{a}_2(1-z) + \left(\frac{n-1}{2} \right) z - \frac{1}{2}(1-z) + i\operatorname{Im}\alpha \right) \\ &\quad \int_{-1}^1 e^{2\pi i |\xi| s} (1-s^2)^{\mathbf{a}_1 z + \mathbf{a}_2(1-z) - 1 + \left(\frac{n-1}{2} \right) z - \frac{1}{2}(1-z) + i\operatorname{Im}\alpha} ds \end{aligned} \quad (2. 35)$$

and

$$\widehat{\mathbf{P}}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(\xi) = \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \widehat{\mathbf{P}}_{j z}^{\alpha \mathbf{a} \mathbf{b} \nu}(\xi), \quad (2. 36)$$

$$\widehat{\mathbf{P}}_{j z}^{\alpha \mathbf{a} \mathbf{b} \nu}(\xi) = \widehat{\phi}(\xi) \varphi_j^\nu(\xi) \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r \widehat{\Omega}_z^{\alpha \mathbf{a}}(r\xi)} \omega(r) |r|^{2[\mathbf{b}_1 z + \mathbf{b}_2(1-z)] - 1 + 2\operatorname{Im}\alpha} dr.$$

At $z = \frac{1}{n}$, we find $\widehat{\Omega}_{\frac{1}{n}}^{\alpha \mathbf{a}}(\xi) = \widehat{\Omega}^{\alpha}(\xi)$ and $\widehat{\mathbf{P}}_{j \frac{1}{n}}^{\alpha \mathbf{a} \mathbf{b} \nu}(\xi) = \widehat{\mathbf{P}}_j^{\alpha \nu}(\xi)$.

Consider

$$\mathbf{I}_{< z}^{\alpha \mathbf{a} \mathbf{b}} f(x) = \int_{\mathbb{R}^n} f(x-u) \sum_{j \leq 0} \mathbf{P}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) du. \quad (2.37)$$

On the other hand, recall $\Psi_j^{\alpha \mu}$, Ψ_j^α defined in (2. 9) and $\mathcal{U}_j^{\alpha \mu}$, $\mathcal{V}_j^{\alpha \mu}$ defined in (2. 10). Assert

$$\begin{aligned} \mathbf{II}_z^{\alpha \mathbf{a} \mathbf{b}} f(x) &= \sum_{j>0} \mathbf{II}_{j z}^{\alpha \mathbf{a} \mathbf{b}} f(x), \quad \mathbf{II}_{j z}^{\alpha \mathbf{a} \mathbf{b}} f(x) = \int_{\mathbb{R}^n} f(x-u) \mathbf{U}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) du, \\ \mathbf{U}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{v \in \mathcal{U}_j^{\alpha \mu}} \mathbf{P}_{j z}^{\alpha \mathbf{a} \mathbf{b} v}(u) + \mathbf{P}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) [1 - \Psi_j^\alpha(u)] \end{aligned} \quad (2.38)$$

and

$$\begin{aligned} \mathbf{III}_z^{\alpha \mathbf{a} \mathbf{b}} f(x) &= \sum_{j>0} \mathbf{III}_{j z}^{\alpha \mathbf{a} \mathbf{b}} f(x), \quad \mathbf{III}_{j z}^{\alpha \mathbf{a} \mathbf{b}} f(x) = \int_{\mathbb{R}^n} f(x-u) \mathbf{V}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) du, \\ \mathbf{V}_{j z}^{\alpha \mathbf{a} \mathbf{b}}(u) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{v \in \mathcal{V}_j^{\alpha \mu}} \mathbf{P}_{j z}^{\alpha \mathbf{a} \mathbf{b} v}(u). \end{aligned} \quad (2.39)$$

From (2. 18) in **Lemma One** and (2. 32) in **Lemma Three**, we conclude

$$\begin{aligned} \|\mathbf{I}_{< 0+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^2(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_2 \mathbf{b}_2} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} \|f\|_{L^2(\mathbb{R}^n)}, \\ \|\mathbf{I}_{< 1+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^p(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_1 \mathbf{b}_1} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 \leq p \leq \infty. \end{aligned} \quad (2.40)$$

By using (2. 40) and applying Stein interpolation theorem [4], we obtain (1. 7) for $\mathbf{I}_{<}^{\alpha} f$.

Given $j > 0$, (2. 24)-(2. 25) in **Lemma Two** implies

$$\begin{aligned} \|\mathbf{II}_{j 0+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^2(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_2 \mathbf{b}_2} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} 2^{j/2} 2^{-j\varepsilon} \|f\|_{L^2(\mathbb{R}^n)}, \\ \|\mathbf{III}_{j 0+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^2(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_2 \mathbf{b}_2} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} 2^{-j\varepsilon} \|f\|_{L^2(\mathbb{R}^n)} \end{aligned} \quad (2.41)$$

for some $\varepsilon = \varepsilon(\mathbf{a}_2, \mathbf{b}_2) > 0$.

On the other hand, (2. 33)-(2. 34) in **Lemma Three** implies

$$\begin{aligned} \|\mathbf{II}_{j 1+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^p(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_1 \mathbf{b}_1 N} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} 2^{-jN} 2^{-j\varepsilon} \|f\|_{L^p(\mathbb{R}^n)}, \quad N \geq 0, \quad 1 \leq p \leq \infty \\ \|\mathbf{III}_{j 1+i\text{Im}z}^{\alpha \mathbf{a} \mathbf{b}} f\|_{L^p(\mathbb{R}^n)} &\leq \mathfrak{B}_{\mathbf{a}_1 \mathbf{b}_1} e^{c|\text{Im}\alpha|} e^{c|\text{Im}z|} 2^{-j\varepsilon} \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 \leq p \leq \infty \end{aligned} \quad (2.42)$$

for some $\varepsilon = \varepsilon(\mathbf{a}_1, \mathbf{b}_1) > 0$.

By using (2. 41)-(2. 42) and applying Stein interpolation theorem [4], we obtain

$$\|\mathbf{II}_j^{\alpha} f\|_{L^p(\mathbb{R}^n)} + \|\mathbf{III}_j^{\alpha} f\|_{L^p(\mathbb{R}^n)} \leq \mathfrak{B}_{\mathbf{Re}\alpha} e^{c|\text{Im}\alpha|} 2^{-j\varepsilon} \|f\|_{L^p(\mathbb{R}^n)}, \quad \frac{n-1}{2n} \leq \frac{1}{p} \leq \frac{n+1}{2n} \quad (2.43)$$

for some $\varepsilon = \varepsilon(\mathbf{Re}\alpha) > 0$. By summing over every $j > 0$, we conclude that both $\mathbf{II}^{\alpha} f$ and $\mathbf{III}^{\alpha} f$ satisfy the L^p -norm inequality in (1. 7).

3 Proof of Lemma One

Recall $\sharp\Omega^\alpha$ defined in (2. 14)-(2. 15). In order to prove (2. 18) and (2. 19)-(2. 20), we reformulate $\sharp\widehat{\Omega}^\alpha(\xi)$ into an $(n+1)$ -dimensional framework. Let $(x, s) \in \mathbb{R}^n \times \mathbb{R}$ and $(\xi, \rho) \in \mathbb{R}^n \times \mathbb{R}$.

\mathfrak{Q}^α is a distribution defined by analytic continuation from

$$\operatorname{Re}\alpha > \frac{n}{2}, \quad \mathfrak{Q}^\alpha(x, s) = \pi^{\alpha - \frac{n+2}{2}} \Gamma^{-1}\left(\alpha - \frac{n}{2}\right) \left\{ \frac{1}{1 - |x|^2 - s^2} \right\}_+^{\frac{n+2}{2} - \alpha}. \quad (3. 1)$$

Equivalently, \mathfrak{Q}^α can be defined by

$$\begin{aligned} \widehat{\mathfrak{Q}}^\alpha(\xi, \rho) &= \left\{ \frac{1}{\sqrt{|\xi|^2 + \rho^2}} \right\}_{\frac{n+1}{2} - [\frac{n+2}{2} - \alpha]}^{\frac{n+1}{2} - [\frac{n+2}{2} - \alpha]} J_{\frac{n+1}{2} - [\frac{n+2}{2} - \alpha]}(2\pi \sqrt{|\xi|^2 + \rho^2}) \\ &= \left\{ \frac{1}{\sqrt{|\xi|^2 + \rho^2}} \right\}_{\alpha - \frac{1}{2}}^{\alpha - \frac{1}{2}} J_{\alpha - \frac{1}{2}}(2\pi \sqrt{|\xi|^2 + \rho^2}), \quad \alpha \in \mathbb{C}. \end{aligned} \quad (3. 2)$$

By using the norm estimate in (A. 5), we find

$$\begin{aligned} \int_{\mathbb{R}} |\widehat{\mathfrak{Q}}^\alpha(\xi, \rho)| d\rho &\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \int_{\mathbb{R}} \left(\frac{1}{1 + |\rho|} \right)^{\operatorname{Re}\alpha} d\rho \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|}, \quad \operatorname{Re}\alpha > 1. \end{aligned} \quad (3. 3)$$

Observe that

$$\mathfrak{Q}^\alpha(x, 0) = \sharp\Omega^\alpha(x), \quad \operatorname{Re}\alpha > \frac{n}{2}. \quad (3. 4)$$

We have

$$\begin{aligned} \sharp\widehat{\Omega}^\alpha(\xi) &= \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} \mathfrak{Q}^\alpha(x, 0) dx \\ &= \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} \left\{ \iint_{\mathbb{R}^{n+1}} e^{2\pi i [x \cdot \eta + 0]} \widehat{\mathfrak{Q}}^\alpha(\eta, \rho) d\eta d\rho \right\} dx \\ &= \iint_{\mathbb{R}^n \times \mathbb{R}^n} e^{2\pi i x \cdot (\eta - \xi)} \left\{ \int_{\mathbb{R}} \widehat{\mathfrak{Q}}^\alpha(\eta, \rho) d\rho \right\} dx d\eta \\ &= \int_{\mathbb{R}} \widehat{\mathfrak{Q}}^\alpha(\xi, \rho) d\rho, \quad \operatorname{Re}\alpha > \frac{n}{2}. \end{aligned} \quad (3. 5)$$

Remark 3.1 The equality in (3. 5) remains valid for $\operatorname{Re}\alpha > 1$.

Given $\xi \in \mathbb{R}^n$, $\rho \in \mathbb{R}$, both $\sharp\widehat{\Omega}^\alpha(\xi)$ and $\widehat{\mathfrak{Q}}^\alpha(\xi, \rho)$ are analytic functions of $\alpha \in \mathbb{C}$. Furthermore, (3. 3) implies that $\int_{\mathbb{R}} \widehat{\mathfrak{Q}}^\alpha(\xi, \rho) d\rho$ is analytic whenever $\operatorname{Re}\alpha > 1$. Therefore, by the principal of analytic continuation, we must have (3. 5) hold for every $\operatorname{Re}\alpha > 1$.

Recall $\# \widehat{\mathbf{P}}_j^{\alpha \beta}, j \in \mathbb{Z}$ defined in (2. 16). From (3. 2) and (3. 5), we have

$$\begin{aligned} \# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) &= \widehat{\phi}(\xi) \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi ir} \left\{ \int_{\mathbb{R}} \widehat{\mathfrak{Q}}^\alpha(r\xi, \rho) d\rho \right\} \omega(r) |r|^{2\beta-1} dr \\ &= \widehat{\phi}(\xi) \int_{2^{j-1} \leq |r| < 2^j} \int_{\mathbb{R}} e^{-2\pi ir} \left\{ \frac{1}{\sqrt{r^2|\xi|^2 + \rho^2}} \right\}^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(2\pi \sqrt{r^2|\xi|^2 + \rho^2}) \\ &\quad \omega(r) |r|^{2\beta-1} d\rho dr \end{aligned} \quad (3. 6)$$

where ω satisfies (1. 4)-(1. 5) and $\text{supp } \widehat{\phi} \subset \{\xi \in \mathbb{R}^n : \frac{1}{3} < |\xi| \leq 3\}$.

Let $\operatorname{Re} \alpha \geq \left[\frac{2n}{2n-1} \right] \operatorname{Re} \beta$ and $\frac{2n-1}{2n} < \operatorname{Re} \beta < \frac{2n-1}{2n-2}$. We find

$$\operatorname{Re} \alpha \geq \left[\frac{2n}{2n-1} \right] \operatorname{Re} \beta > 1, \quad 2\operatorname{Re} \beta - \operatorname{Re} \alpha \leq \left[\frac{2n-2}{2n-1} \right] \operatorname{Re} \beta < 1. \quad (3. 7)$$

By using (3. 3) to the first integral in (3. 6), we have

$$\begin{aligned} \left| \sum_{j \leq 0} \# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) \right| &\leq \mathfrak{B}_{\operatorname{Re} \alpha} e^{\mathbf{C} |\operatorname{Im} \alpha|} \int_{|r| < 1} \left\{ \int_{\mathbb{R}} \left(\frac{1}{1+|\rho|} \right)^{\operatorname{Re} \alpha} d\rho \right\} |\omega(r)| |r|^{2\beta-1} dr \\ &\leq \mathfrak{B}_{\operatorname{Re} \alpha} e^{\mathbf{C} |\operatorname{Im} \alpha|}. \end{aligned} \quad (3. 8)$$

Given $j > 0$, we write the second integral in (3. 6) as

$$\begin{aligned} \# \widehat{\mathbf{P}}_j^{\alpha \beta}(\xi) &= \widehat{\phi}(\xi) |\xi|^{-2\beta} \int_{2^{j-1} |\xi| \leq |s| < 2^j |\xi|} \int_{\mathbb{R}} e^{-2\pi is/|\xi|} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(2\pi \sqrt{s^2 + \rho^2}) \\ &\quad \omega(s/|\xi|) |s|^{2\beta-1} d\rho ds. \end{aligned} \quad (3. 9)$$

We assert

$$\# \widehat{\mathbf{Q}}_j^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{-2\pi is/|\xi|} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(2\pi \sqrt{s^2 + \rho^2}) s^{2\beta-2} d\rho ds \quad (3. 10)$$

and

$$\begin{aligned} \# \widehat{\mathbf{R}}_j^{\alpha \beta}(\xi) &= \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{-2\pi is/|\xi|} [1 - e^{-2\pi is/|\xi|}] \\ &\quad \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(2\pi \sqrt{s^2 + \rho^2}) [\omega(s) - s^{-1}] s^{2\beta-1} d\rho ds. \end{aligned} \quad (3. 11)$$

Remark 3.2 Later, it should be clear that the same estimates regarding $\# \widehat{\mathbf{Q}}_j^{\alpha \beta}$ and $\# \widehat{\mathbf{R}}_j^{\alpha \beta}$ hold if $s \in [-2^j, -2^{j-1}]$.

First, the norm estimate in (A. 5) implies

$$\left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\operatorname{Re} \alpha - \frac{1}{2}} \left| J_{\alpha-\frac{1}{2}}(2\pi \sqrt{s^2 + \rho^2}) \right| \leq \mathfrak{B}_{\operatorname{Re} \alpha} e^{\mathbf{C} |\operatorname{Im} \alpha|} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\operatorname{Re} \alpha}. \quad (3. 12)$$

By using (1. 5) and (3. 12), we have

$$\begin{aligned}
\left| \# \widehat{\mathbf{R}}_j^{\alpha \beta}(\xi) \right| &\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\operatorname{Re}\alpha} s^{2\operatorname{Re}\beta - 3} d\rho ds \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{1 + |\rho|} \right\}^{1+\delta} s^{2\operatorname{Re}\beta - \operatorname{Re}\alpha - 2 + \delta} d\rho ds \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1 + \delta]}
\end{aligned} \tag{3. 13}$$

where $\delta = \delta(\operatorname{Re}\alpha) > 0$ is sufficiently small.

Next, we claim

$$\left| \# \widehat{\mathbf{Q}}_j^{\alpha \beta}(\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{j/2} 2^{-j\varepsilon}, \quad |1 - |\xi|| \leq 2^{-j} \tag{3. 14}$$

and

$$\left| \# \widehat{\mathbf{Q}}_j^{\alpha \beta}(\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} e^{\mathbf{C}|\operatorname{Im}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} 2^{-j\varepsilon}, \quad |1 - |\xi|| > 2^{-j} \tag{3. 15}$$

for some $\varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0$.

Note that (3. 14)-(3. 15) together with (3. 13) and **Remark 3.2** imply (2. 19)-(2. 20).

By using the asymptotic expansion of Bessel functions given at (A. 2), we write

$$\# \widehat{\mathbf{Q}}_j^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{-2\pi i s / |\xi|} [\mathfrak{S}^\alpha(s, \rho) + \mathcal{E}^\alpha(s, \rho)] s^{2\beta - 2} d\rho ds \tag{3. 16}$$

where

$$\mathfrak{S}^\alpha(s, \rho) = \frac{1}{\pi} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha \cos \left[2\pi \sqrt{s^2 + \rho^2} - \frac{\pi}{2} \alpha \right] \tag{3. 17}$$

and

$$|\mathcal{E}^\alpha(s, \rho)| \leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \begin{cases} \sqrt{s^2 + \rho^2}^{-\operatorname{Re}\alpha}, & \sqrt{s^2 + \rho^2} \leq 1, \\ \sqrt{s^2 + \rho^2}^{-\operatorname{Re}\alpha-1}, & \sqrt{s^2 + \rho^2} > 1. \end{cases} \tag{3. 18}$$

By using (3. 18), we find

$$\begin{aligned}
\left| \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{-2\pi i s / |\xi|} \mathcal{E}^\alpha(s, \rho) s^{2\beta - 2} d\rho ds \right| &\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\operatorname{Re}\alpha+1} s^{2\operatorname{Re}\beta - 2} d\rho ds \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{1 + |\rho|} \right\}^{1+\delta} s^{2\operatorname{Re}\beta - \operatorname{Re}\alpha - 2 + \delta} d\rho ds \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1 + \delta]}.
\end{aligned} \tag{3. 19}$$

Because of Euler's formulae, we replace the cosine function inside (3. 17) with $e^{2\pi i \sqrt{s^2 + \rho^2}}$ multiplied by $\mathfrak{B}_{\text{Re}\alpha} e^{\mathfrak{c}|\text{Im}\alpha|}$. Consider

$$\#\widehat{\mathbf{A}}_j^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} d\rho ds. \quad (3. 20)$$

We aim to show that $\#\widehat{\mathbf{A}}_j^{\alpha \beta}(\xi)$ satisfies the norm inequalities in (3. 14)-(3. 15). Our estimates remain valid by switching $e^{2\pi i \sqrt{s^2 + \rho^2}}$ to $e^{-2\pi i \sqrt{s^2 + \rho^2}}$.

Let $\varphi \in C_c^\infty(\mathbb{R})$ such that $\varphi(t) = 1$ for $|t| \leq 1$ and $\varphi(t) = 0$ for $|t| > 2$. We write

$$\#\widehat{\mathbf{A}}_j^{\alpha \beta}(\xi) = \#\widehat{\mathbf{F}}_j^{\alpha \beta}(\xi) + \#\widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) \quad (3. 21)$$

for which

$$\#\widehat{\mathbf{F}}_j^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} \varphi(\rho) d\rho ds \quad (3. 22)$$

and

$$\#\widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} [1 - \varphi(\rho)] d\rho ds. \quad (3. 23)$$

Observe that

$$\begin{aligned} |\#\widehat{\mathbf{F}}_j^{\alpha \beta}(\xi)| &\leq \int_{2^{j-1}}^{2^j} \int_{0 < |\rho| \leq 2} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\text{Re}\alpha} s^{2\text{Re}\beta-2} d\rho ds \\ &\leq 2^j [2\text{Re}\beta - \text{Re}\alpha - 1]. \end{aligned} \quad (3. 24)$$

Given $\mathbf{t} > 0$, define

$$\#\widehat{\mathbf{G}}_{j \mathbf{t}}^{\alpha \beta}(\xi) = \int_{2^{j-1}}^{2^j} \int_{|\rho| > \mathbf{t}s} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} [1 - \varphi(\rho)] d\rho ds. \quad (3. 25)$$

By integration by parts w.r.t ρ , we have

$$\begin{aligned} \#\widehat{\mathbf{G}}_{j \mathbf{t}}^{\alpha \beta}(\xi) &= \frac{1}{2\pi i} \int_{2^{j-1}}^{2^j} \int_{|\rho| > \mathbf{t}s} \frac{2\pi i \rho}{\sqrt{s^2 + \rho^2}} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \frac{1}{\rho} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} [1 - \varphi(\rho)] s^{2\beta-2} d\rho ds \\ &= -\frac{1}{2\pi i} \int_{2^{j-1}}^{2^j} \int_{|\rho| > \mathbf{t}s} e^{2\pi i [\sqrt{s^2 + \rho^2} - s/|\xi|]} \partial_\rho \left\{ \frac{1}{\rho} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} [1 - \varphi(\rho)] \right\} s^{2\beta-2} d\rho ds \\ &\quad + \frac{1}{\pi i} \mathbf{t}^{-1} (\sqrt{1 + \mathbf{t}^2})^{1-\alpha} \int_{2^{j-1}}^{2^j} e^{2\pi i [s \sqrt{1 + \mathbf{t}^2} - s/|\xi|]} [1 - \varphi(\mathbf{t}s)] s^{2\beta-\alpha-2} ds \\ &\doteq \text{main term} + \text{boundary term}. \end{aligned} \quad (3. 26)$$

For every $\mathbf{t} > 0$, we find

$$\begin{aligned}
|\text{boundary term}| &\leq \mathfrak{B} \mathbf{t}^{-1} \left(\sqrt{1 + \mathbf{t}^2} \right)^{1 - \text{Re}\alpha} \int_{2^{j-1}}^{2^j} s^{2\text{Re}\beta - \text{Re}\alpha - 2} ds \\
&\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} \mathbf{t}^{-1} \left(\sqrt{1 + \mathbf{t}^2} \right)^{1 - \text{Re}\alpha} 2^{j[2\text{Re}\beta - \text{Re}\alpha - 1]} \\
&\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} 2^{j[2\text{Re}\beta - \text{Re}\alpha - 1]} \begin{cases} 1, & \mathbf{t} \geq 1, \\ \mathbf{t}^{-1}, & 0 < \mathbf{t} < 1. \end{cases}
\end{aligned} \tag{3. 27}$$

Return to the **main term** in (3. 26). A direct computation shows

$$\partial_\rho \left\{ \frac{1}{\rho} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} [1 - \varphi(\rho)] \right\} = - \frac{1}{\rho^2} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} [1 - \varphi(\rho)] \tag{1}$$

$$+ (1 - \alpha) \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha+1} [1 - \varphi(\rho)] \tag{2} \tag{3. 28}$$

$$+ \frac{1}{\rho} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} \partial_\rho [1 - \varphi(\rho)]. \tag{3}$$

Regarding (1) in (3. 28), we have

$$\begin{aligned}
&\left| \int_{2^{j-1}}^{2^j} \int_{|\rho| > \mathbf{t}s} e^{2\pi i \left[\sqrt{s^2 + \rho^2} - s/|\xi| \right]} \frac{1}{\rho^2} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\alpha-1} [1 - \varphi(\rho)] s^{2\beta - 2} d\rho ds \right| \\
&\leq \int_{2^{j-1}}^{2^j} \int_{|\rho| > 1, |\rho| > \mathbf{t}s} \frac{\sqrt{s^2 + \rho^2}}{|\rho|} \frac{1}{|\rho|} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\text{Re}\alpha} s^{2\text{Re}\beta - 2} d\rho ds \\
&\leq \mathfrak{B} \mathbf{t}^{-1} \sqrt{1 + \mathbf{t}^2} \int_{2^{j-1}}^{2^j} \int_{|\rho| > 1} \frac{1}{|\rho|} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\text{Re}\alpha} s^{2\text{Re}\beta - 2} d\rho ds \\
&\leq \mathfrak{B} \mathbf{t}^{-1} \sqrt{1 + \mathbf{t}^2} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{1 + |\rho|} \right\} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\text{Re}\alpha - \delta + \delta} s^{2\text{Re}\beta - 2} d\rho ds \\
&\leq \mathfrak{B} \mathbf{t}^{-1} \sqrt{1 + \mathbf{t}^2} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{1 + |\rho|} \right\}^{1+\delta} s^{2\text{Re}\beta - \text{Re}\alpha - 2 + \delta} d\rho ds \\
&\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} 2^{j[2\text{Re}\beta - \text{Re}\alpha - 1 + \delta]} \begin{cases} 1, & \mathbf{t} \geq 1, \\ \mathbf{t}^{-1}, & 0 < \mathbf{t} < 1. \end{cases}
\end{aligned} \tag{3. 29}$$

Regarding (2) in (3. 28), we have

$$\begin{aligned}
& \left| (1-\alpha) \int_{2^{j-1}}^{2^j} \int_{|\rho|>ts} e^{2\pi i [\sqrt{s^2+\rho^2}-s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^{\alpha+1} [1-\varphi(\rho)] s^{2\beta-2} d\rho ds \right| \\
& \leq |1-\alpha| \int_{2^{j-1}}^{2^j} \int_{|\rho|>1, |\rho|>ts} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^{\text{Re}\alpha+1} s^{2\text{Re}\beta-2} d\rho ds \\
& \leq \mathfrak{B}_{\text{Re}\alpha} e^{c|\text{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^{\text{Re}\alpha-\delta+1+\delta} s^{2\text{Re}\beta-2} d\rho ds \\
& \leq \mathfrak{B}_{\text{Re}\alpha} e^{c|\text{Im}\alpha|} \int_{2^{j-1}}^{2^j} \int_{\mathbb{R}} \left\{ \frac{1}{1+|\rho|} \right\}^{1+\delta} s^{2\text{Re}\beta-\text{Re}\alpha-2+\delta} d\rho ds \\
& \leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} 2^{j[2\text{Re}\beta-\text{Re}\alpha-1+\delta]}.
\end{aligned} \tag{3. 30}$$

Remark 3.3 $\partial_\rho [1-\varphi(\rho)] = 0$ for $|\rho| < 1$ and $|\rho| > 2$.

Regarding (3) in (3. 28), we have

$$\begin{aligned}
& \left| \int_{2^{j-1}}^{2^j} \int_{|\rho|>ts} e^{2\pi i [\sqrt{s^2+\rho^2}-s/|\xi|]} \frac{1}{\rho} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^{\alpha-1} \partial_\rho [1-\varphi(\rho)] s^{2\beta-2} d\rho ds \right| \\
& \leq t^{-1} \int_{2^{j-1}}^{2^j} \int_{1 \leq |\rho| \leq 2} \frac{1}{s} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^{\text{Re}\alpha-1} s^{2\text{Re}\beta-2} d\rho ds \\
& \leq \mathfrak{B} t^{-1} \int_{2^{j-1}}^{2^j} s^{2\text{Re}\beta-\text{Re}\alpha-2} ds \\
& \leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} t^{-1} 2^{j[2\text{Re}\beta-\text{Re}\alpha-1]}.
\end{aligned} \tag{3. 31}$$

From (3. 25) to (3. 31), we conclude

$$\left| \widehat{\mathbf{G}}_j^\alpha \mathbf{G}_t^\beta (\xi) \right| \leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} t^{-1} 2^{-j\varepsilon} \tag{3. 32}$$

for some $\varepsilon = \varepsilon(\text{Re}\alpha, \text{Re}\beta) > 0$.

In particular, at $t = 2^{-j/2}$, we find

$$\begin{aligned}
& \left| \int_{2^{j-1}}^{2^j} \int_{|\rho|>2^{-j/2}s} e^{2\pi i [\sqrt{s^2+\rho^2}-s/|\xi|]} \left\{ \frac{1}{\sqrt{s^2+\rho^2}} \right\}^\alpha [1-\varphi(\rho)] s^{2\beta-2} d\rho ds \right| \\
& \leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} 2^{j/2} 2^{-j\varepsilon}.
\end{aligned} \tag{3. 33}$$

On the other hand, we have

$$\begin{aligned}
& \left| \int_{2^{j-1}}^{2^j} \int_{|\rho| \leq 2^{-j/2}s} e^{2\pi i [\sqrt{s^2 + \rho^2} - s / |\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha [1 - \varphi(\rho)] s^{2\beta-2} d\rho ds \right| \\
& \leq \int_{2^{j-1}}^{2^j} \int_{|\rho| \leq 2^{-j/2}s} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^{\operatorname{Re}\alpha} s^{2\operatorname{Re}\beta-2} d\rho ds \\
& \leq \mathfrak{B} 2^{-j/2} \int_{2^{j-1}}^{2^j} s^{2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1} ds \\
& \leq \mathfrak{B}_{\operatorname{Re}\alpha, \operatorname{Re}\beta} 2^{j/2} 2^j [2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1].
\end{aligned} \tag{3. 34}$$

By putting together (3. 33) and (3. 34), we obtain

$$\left| \# \widehat{\mathbf{G}}_j^{\alpha, \beta}(\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha, \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} 2^{j/2} 2^{-j\varepsilon}, \quad \varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0. \tag{3. 35}$$

Let $|\xi| < 1$. Define

$$t = \frac{\sqrt{3|\xi|^{-2} - 2|\xi|^{-1} - 1}}{|\xi|^{-1} + 1} = \frac{\sqrt{3|\xi|^{-1} + 1}}{|\xi|^{-1} + 1} \sqrt{|\xi|^{-1} - 1}. \tag{3. 36}$$

The norm estimate in (3. 32) gives

$$\left| \# \widehat{\mathbf{G}}_j^{\alpha, \beta}(t\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha, \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} 2^{-j\varepsilon}, \quad \varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0. \tag{3. 37}$$

Denote $s = R \cos \vartheta$ and $\rho = R \sin \vartheta$. Take into account $\cos \vartheta = \frac{s}{\sqrt{s^2 + \rho^2}}$. A direct computation shows

$$|\rho| \leq st \implies |\xi|^{-1} \cos \vartheta - 1 \geq \frac{1}{2} [|\xi|^{-1} - 1]. \tag{3. 38}$$

Recall $\frac{1}{3} < |\xi| < 1$. Define θ implicitly by

$$\cos \theta = \frac{|\xi|^{-1} + 1}{2|\xi|^{-1}}, \quad \theta \in \left(0, \arccos\left(\frac{2}{3}\right)\right). \tag{3. 39}$$

As an necessity, $\vartheta \in [-\theta, \theta]$ if $\cos \vartheta$ satisfies the inequality in (3. 38). We consider

$$\begin{aligned}
\# \widehat{\mathbf{B}}_j^{\alpha, \beta}(t\xi) &= \int_{2^{j-1}}^{2^j} \int_{|\rho| \leq ts} e^{2\pi i [\sqrt{s^2 + \rho^2} - s / |\xi|]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} [1 - \varphi(\rho)] d\rho ds \\
&= \int_{-\theta}^{\theta} \left\{ \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 1} [1 - \varphi(R \sin \vartheta)] dR \right\} \cos^{2\beta-2} \vartheta d\vartheta.
\end{aligned} \tag{3. 40}$$

By integration by parts *w.r.t* R , we find

$$\begin{aligned}
& \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 1} [1 - \varphi(R \sin \vartheta)] dR \\
&= \frac{1}{2\pi i} \frac{1}{1 - |\xi|^{-1} \cos \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 1} [1 - \varphi(R \sin \vartheta)] \Big|_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} \\
&\quad - \frac{1}{2\pi i} [2\beta - \alpha - 1] \frac{1}{1 - |\xi|^{-1} \cos \vartheta} \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 2} [1 - \varphi(R \sin \vartheta)] dR \\
&\quad - \frac{1}{2\pi i} \frac{1}{1 - |\xi|^{-1} \cos \vartheta} \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 1} \partial_R [1 - \varphi(R \sin \vartheta)] dR. \tag{3. 41}
\end{aligned}$$

Note that $\partial_R [1 - \varphi(R \sin \vartheta)] = -\varphi'(R \sin \vartheta) \sin \vartheta$ where $\varphi'(R \sin \vartheta) = 0$ for $R < |\sin \vartheta|^{-1}$ and $R > 2|\sin \vartheta|^{-1}$. For $\vartheta \in [-\theta, \theta]$, (3. 38) implies $|1 - |\xi|^{-1} \cos \vartheta| \geq \frac{1}{2} ||\xi|^{-1} - 1|$. We have

$$\begin{aligned}
& \left| \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R [1 - \cos \vartheta / |\xi|]} R^{2\beta - \alpha - 1} [1 - \varphi(R \sin \vartheta)] dR \right| \\
&\leq \mathfrak{B} \frac{1}{|1 - |\xi|^{-1}|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \\
&\quad + \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} \frac{1}{|1 - |\xi|^{-1}|} \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} R^{2\operatorname{Re}\beta - \operatorname{Re}\alpha - 2} dR \tag{3. 42} \\
&\quad + \mathfrak{B} \frac{1}{|1 - |\xi|^{-1}|} \int_{|\sin \vartheta|^{-1} \leq R \leq 2|\sin \vartheta|^{-1}} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} |\sin \vartheta| dR \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} \frac{1}{|1 - |\xi|^{-1}|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]}.
\end{aligned}$$

Remark 3.4 Suppose $|\xi|^{-1} = 1 + 2^{-\ell}$ for $\ell > 0$. From (3. 39), we find $\cos \theta \approx 1 - 2^{-\ell}$ suggesting $\theta \approx 2^{-\ell/2}$.

This can be easily seen by writing out the Taylor expansion of $\cos \theta$.

From (3. 40), (3. 42) and **Remark 3.4**, we find

$$\begin{aligned}
\left| \widehat{\mathbf{B}}_j^\alpha \mathbf{t}^\beta (\xi) \right| &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \int_{[-\theta, \theta]} \frac{1}{|1 - |\xi|^{-1}|} d\vartheta \\
&\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}}. \tag{3. 43}
\end{aligned}$$

Recall (3. 23) and (3. 25). We have

$$\widehat{\mathbf{G}}_j^\alpha \mathbf{t}^\beta (\xi) = \widehat{\mathbf{G}}_j^\alpha \mathbf{t}^\beta (\xi) + \widehat{\mathbf{B}}_j^\alpha \mathbf{t}^\beta (\xi), \quad \mathbf{t} > 0. \tag{3. 44}$$

By putting together (3. 37) and (3. 43), we obtain

$$\left| \# \widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} 2^{-j\varepsilon} \quad (3. 45)$$

for some $\varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0$.

Let $|\xi| > 1$. We consider

$$\# \widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) = \# \widehat{\mathbf{G}}_{j1}^{\alpha \beta}(\xi) + \# \widehat{\mathbf{B}}_{j1}^{\alpha \beta}(\xi). \quad (3. 46)$$

Clearly, (3. 32) implies

$$\left| \# \widehat{\mathbf{G}}_{j1}^{\alpha \beta}(\xi) \right| \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} 2^{-j\varepsilon}, \quad \varepsilon = \varepsilon(\operatorname{Re}\alpha, \operatorname{Re}\beta) > 0. \quad (3. 47)$$

On the other hand, we have

$$\begin{aligned} \# \widehat{\mathbf{B}}_{j1}^{\alpha \beta}(\xi) &= \int_{2^{j-1}}^{2^j} \int_{|\rho| \leq s} e^{2\pi i \left[\sqrt{s^2 + \rho^2} - s/|\xi| \right]} \left\{ \frac{1}{\sqrt{s^2 + \rho^2}} \right\}^\alpha s^{2\beta-2} [1 - \varphi(\rho)] d\rho ds \\ &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \left\{ \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R \left[1 - \cos \vartheta / |\xi| \right]} R^{2\beta-\alpha-1} [1 - \varphi(R \sin \vartheta)] dR \right\} \cos^{2\beta-2} \vartheta d\vartheta. \end{aligned} \quad (3. 48)$$

By carrying out the same estimates in (3. 41)-(3. 42), we find

$$\begin{aligned} &\left| \int_{2^{j-1} \cos^{-1} \vartheta}^{2^j \cos^{-1} \vartheta} e^{2\pi i R \left[1 - \cos \vartheta / |\xi| \right]} R^{2\beta-\alpha-1} [1 - \varphi(R \sin \vartheta)] dR \right| \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \left| 1 - \frac{\cos \vartheta}{|\xi|} \right|^{-1}, \quad \vartheta \in [-\frac{\pi}{4}, \frac{\pi}{4}]. \end{aligned} \quad (3. 49)$$

Remark 3.5 Suppose $|\xi|^{-1} = 1 - 2^{-\ell}$ for $\ell > 0$. We have $\cos \vartheta = 1 - 2^{-\ell+k}$ implying $\vartheta \approx 2^{-\ell/2+k/2}$ for $0 \leq k < \ell$.

We have

$$\begin{aligned} \left| \# \widehat{\mathbf{B}}_{j1}^{\alpha \beta}(\xi) \right| &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \left| 1 - \frac{\cos \vartheta}{|\xi|} \right|^{-1} d\vartheta \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} \sum_{k=0}^{\infty} 2^{\ell-k} 2^{-\ell/2+k/2} \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]} 2^{\ell/2} \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|} e^{c|\operatorname{Im}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} 2^{j[2\operatorname{Re}\beta - \operatorname{Re}\alpha - 1]}. \end{aligned} \quad (3. 50)$$

By putting together (3. 47) and (3. 50), we obtain (3. 45) again.

Upon to this point, we finish the proof of **Lemma One**. Next, we conclude **Remark 2.3**.

Suppose $2^{-\ell} \leq |1 - |\xi|| < 2^{-\ell+1}$, $\ell > 0$. We have

$$\begin{aligned} \sum_{0 < j \leq \ell} \left| \widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) \right| &\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} \sum_{0 < j \leq \ell} 2^{j/2} 2^{-j\varepsilon} \quad \text{by (3. 35)} \\ &\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2} - \varepsilon} \end{aligned} \quad (3. 51)$$

and

$$\begin{aligned} \sum_{j > \ell} \left| \widehat{\mathbf{G}}_j^{\alpha \beta}(\xi) \right| &\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} e^{c|\text{Im}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2}} \sum_{j > \ell} 2^{-j\varepsilon} \quad \text{by (3. 45)} \\ &\leq \mathfrak{B}_{\text{Re}\alpha \text{ Re}\beta} e^{c|\text{Im}\alpha|} e^{c|\text{Im}\beta|} \left| \frac{1}{1 - |\xi|} \right|^{\frac{1}{2} - \varepsilon}. \end{aligned} \quad (3. 52)$$

Together with the previous estimates in (3. 8)-(3. 24), we obtain (2. 17).

4 Proof of Lemma Two

Recall $\widehat{\mathbf{P}}_j^{\alpha \beta \nu}$ and $\# \mathbf{U}_j^{\alpha \beta}, \# \mathbf{V}_j^{\alpha \beta}, j > 0$ defined in (2. 21) and (2. 22)-(2. 23). We consider

$$\begin{aligned} \widehat{\mathbf{V}}_j^{\alpha \beta}(\xi) &= \int_{\mathbb{R}^n} e^{-2\pi i u \cdot \xi} \left\{ \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(u) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} \# \mathbf{P}_j^{\alpha \beta \nu}(u) \right\} du \\ &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} \int_{\mathbb{R}^n} \widehat{\Psi}_j^{\alpha \mu}(\xi - \zeta) \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\zeta) d\zeta \end{aligned} \quad (4. 1)$$

where $\mathcal{V}_j^{\alpha \mu}$ is defined in (2. 10).

Let $\zeta = \mathbf{L}_\mu \eta$ inside (4. 1). \mathbf{L}_μ is the orthogonal matrix defined in (2. 8) with $\det \mathbf{L}_\mu = 1$. Moreover, $\mathbf{L}_\mu^T \xi_j^\mu = (1, 0)^T \in \mathbb{R} \times \mathbb{R}^{n-1}$.

Recall Γ_j^ν defined in (2. 6). We find $\# \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) \neq 0$ if $\mathbf{L}_\mu \eta \in \Gamma_j^\nu \cap \{\eta \in \mathbb{R}^n : \frac{1}{3} < |\eta| \leq 3\}$. Define

$$\mathcal{S}_\ell = \left\{ \eta \in \mathbb{R}^n : 2^{-\ell} \leq |1 - |\eta|| < 2^{-\ell+1} \right\}, \quad \ell = 0, 1, \dots \quad (4. 2)$$

For each μ fixed and $\nu \in \mathcal{V}_j^{\alpha \mu}$, write

$$\begin{aligned} \int_{\mathbb{R}^n} \widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) d\eta &= \\ \sum_{\ell=0}^{\infty} \int_{\mathcal{S}_\ell} \widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) d\eta. \end{aligned} \quad (4. 3)$$

We find

$$\text{vol} \left\{ \mathcal{S}_\ell \cap \text{supp} \sum_{\nu \in V_j^{\alpha \mu}} \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu} \right\} \leq \mathfrak{B} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} 2^{-\ell}. \quad (4.4)$$

Note that $\# \widehat{\mathbf{P}}_j^{\alpha \beta \nu} = \varphi_j^\nu \# \widehat{\mathbf{P}}_j^{\alpha \beta}$ where φ_j^ν is defined in (2.5). Suppose $2^{-\ell} \leq |1 - |\eta|| < 2^{-\ell+1}$.

Lemma One implies

$$\begin{aligned} \left| \sum_{\nu \in V_j^{\alpha \mu}} \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu} (\mathbf{L}_\mu \eta) \right| &\leq \mathfrak{B}_{\text{Re}\alpha} \text{Re}\beta e^{c|\text{Im}\alpha|} 2^{j/2} 2^{-j\varepsilon}, \quad j \leq \ell; \\ \left| \sum_{\nu \in V_j^{\alpha \mu}} \# \widehat{\mathbf{P}}_j^{\alpha \beta \nu} (\mathbf{L}_\mu \eta) \right| &\leq \mathfrak{B}_{\text{Re}\alpha} \text{Re}\beta e^{c|\text{Im}\alpha|} e^{c|\text{Im}\beta|} 2^{\ell/2} 2^{-j\varepsilon}, \quad j > \ell \end{aligned} \quad (4.5)$$

for some $\varepsilon = \varepsilon(\text{Re}\alpha, \text{Re}\beta) > 0$.

Recall $\Psi_j^{\alpha \mu}$ defined in (2.9). We have

$$\widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) = \int_{\mathbb{R}^n} e^{-2\pi i [\mathbf{L}_\mu^T \xi - \eta] \cdot u} \varphi \left[2^{-\epsilon j} |2^j - |u_1|| \right] \prod_{i=2}^n \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} |u_i| \right] du \quad (4.6)$$

where $\epsilon = \epsilon(\text{Re}\alpha) > 0$ is small and adjustable.

Remark 4.1 The volume of the support of $\left\{ \varphi \left[2^{-\epsilon j} |2^j - |u_1|| \right] \prod_{i=2}^n \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} |u_i| \right] \right\}$ is bounded by $\mathfrak{B} 2^{j(\frac{n-1}{2})} 2^{jne}$.

Suppose that there is an $i \in \{2, \dots, n\}$ such that $|(\mathbf{L}_\mu^T \xi - \eta)_i| \gtrsim 2^{-j/2}$. By integration by parts w.r.t u_i inside (4.6), we find

$$\begin{aligned} \left| \widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) \right| &\leq \mathfrak{B}_N 2^{j(\frac{n-1}{2})} 2^{jne} 2^{j(\frac{1}{2})N} 2^{-j[\frac{1}{2}+\epsilon]N} \\ &\leq \mathfrak{B}_{\text{Re}\alpha} \end{aligned} \quad (4.7)$$

by taking sufficiently large depending on $\epsilon = \epsilon(\text{Re}\alpha) > 0$.

Remark 4.2 Given $\xi \in \mathbb{R}^n$, we have $\xi \in \Gamma_j^{\bar{\mu}}$ defined in (2.6) for some $\bar{\mu}$. Assert

$$\Gamma_{j-2}^{\bar{\mu}} = \left\{ \xi \in \mathbb{R}^n : \left| \frac{\xi}{|\xi|} - \xi_j^{\bar{\mu}} \right| < 2^{-j/2+2} \right\}.$$

There are at most a constant multiple of $2^{\epsilon(n-1)j}$ other μ 's such that

$$\Gamma_{j-2}^{\bar{\mu}} \cap \bigcup_{\nu \in V_j^{\alpha \mu}} \Gamma_j^\nu \neq \emptyset. \quad (4.8)$$

We say $\mu \in G_1$ if it belongs to the $\mathfrak{B}2^{2\epsilon(n-1)j}$ many's in **Remark 6.2**. Otherwise, we say $\mu \in G_2$.

Remark 4.3 Suppose $\mu \in G_2$. We necessarily have $\left|(\mathbf{L}_\mu^T \xi - \eta)_i\right| \gtrsim 2^{-j/2}$ for some $i \in \{2, \dots, n\}$ whenever $\mathbf{L}_\mu \eta \in \bigcup_{\nu \in \mathcal{V}_j^\alpha} \Gamma_j^\nu$.

From (4. 1) to (4. 3), we have

$$\begin{aligned}
\left| \widehat{\mathbf{V}}_j^{\alpha \beta}(\xi) \right| &\leq \sum_{\mu \in G_2} \sum_{\ell=0}^{\infty} \left| \int_{\mathcal{S}_\ell} \widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) \sum_{\nu \in \mathcal{V}_j^\alpha} \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) d\eta \right| \\
&+ \sum_{\mu \in G_1} \sum_{\ell=0}^{\infty} \left| \int_{\mathcal{S}_\ell} \widehat{\Psi}_j^{\alpha \mu}(\xi - \mathbf{L}_\mu \eta) \sum_{\nu \in \mathcal{V}_j^\alpha} \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) d\eta \right| \\
&\leq \sum_{\mu \in G_2} \mathfrak{B}_{\mathbf{Re}\alpha} \sum_{\ell=0}^{\infty} \int_{\mathcal{S}_\ell} \left| \sum_{\nu \in \mathcal{V}_j^\alpha} \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) \right| d\eta \quad \text{by Remark 4.3 and (4. 7)} \\
&+ \sum_{\mu \in G_1} \mathfrak{B}2^{j(\frac{n-1}{2})} 2^{jn\epsilon} \sum_{\ell=0}^{\infty} \int_{\mathcal{S}_\ell} \left| \sum_{\nu \in \mathcal{V}_j^\alpha} \widehat{\mathbf{P}}_j^{\alpha \beta \nu}(\mathbf{L}_\mu \eta) \right| d\eta \quad \text{by Remark 4.1} \\
&\leq \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} \sum_{\mu \in G_2} \sum_{\ell=0}^{\infty} 2^{\ell/2} 2^{-j\epsilon} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} 2^{-\ell} \\
&+ \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} \sum_{\mu \in G_1} 2^{j(\frac{n-1}{2})} 2^{jn\epsilon} \sum_{\ell=0}^{\infty} 2^{\ell/2} 2^{-j\epsilon} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} 2^{-\ell} \\
&\leq \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} \sum_{\mu \in G_2} 2^{-j\epsilon} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} \\
&+ \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} \sum_{\mu \in G_1} 2^{j(\frac{n-1}{2})} 2^{jn\epsilon} 2^{-j\epsilon} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} \\
&\leq \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} 2^{j(\frac{n-1}{2})} 2^{-j\epsilon} 2^{-j(\frac{n-1}{2})} 2^{\epsilon(n-1)j} \\
&+ \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} 2^{\epsilon(n-1)j} 2^{-j\epsilon} 2^{jn\epsilon} 2^{\epsilon(n-1)j} \\
&\leq \mathfrak{B}_{\mathbf{Re}\alpha} \mathbf{Re}\beta e^{\mathbf{c}|\mathbf{Im}\alpha|} e^{\mathbf{c}|\mathbf{Im}\beta|} 2^{-j\epsilon/2} \tag{4. 9}
\end{aligned}$$

provided that $\epsilon = \epsilon(\mathbf{Re}\alpha) > 0$ is chosen sufficiently small.

Observe that (4. 9) implies (2. 25). On the other hand, we have $\# \mathbf{U}_j^{\alpha \beta}(x) + \# \mathbf{V}_j^{\alpha \beta}(x) = \# \mathbf{P}_j^{\alpha \beta}(x)$. From **Lemma One** and (2. 25), we conclude (2. 24).

5 Proof of Lemma Three

Recall ${}^b\Omega^\alpha$ defined by analytic continuation from (2. 26). Alternatively, its Fourier transform is given in (2. 27).

Let ${}^b\widehat{\mathbf{P}}_j^{\alpha \beta}$ defined in (2. 28) for $\operatorname{Re}\alpha > 0$ and $0 < \operatorname{Re}\beta < \frac{1}{2}$. We have

$${}^b\mathbf{P}_j^{\alpha \beta}(x) = \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r} \left\{ \int_{\mathbb{R}^n} \phi(x-u) {}^b\Omega_{|r|}^\alpha(u) du \right\} \omega(r) |r|^{2\beta-1} dr \quad (5. 1)$$

where ${}^b\Omega_r^\alpha(u) = r^{-n} {}^b\Omega^\alpha\left(\frac{u}{r}\right)$, $r > 0$.

Note that $\widehat{\phi} \in C_o^\infty(\mathbb{R}^n)$ where $\operatorname{supp} \widehat{\phi} \subset \{\xi \in \mathbb{R}^n : \frac{1}{3} < |\xi| \leq 3\}$. Moreover, $|\omega(r)| \leq \mathfrak{B}[1+|r|]^{-1}$. We find

$$\begin{aligned} \int_{\mathbb{R}^n} \left| \sum_{j \leq 0} {}^b\mathbf{P}_j^{\alpha \beta}(x) \right| dx &= \int_{\mathbb{R}^n} \left| \int_{|r|<1} e^{-2\pi i r} \left\{ \int_{\mathbb{R}^n} \phi(x-u) {}^b\Omega_{|r|}^\alpha(u) du \right\} \omega(r) |r|^{2\beta-1} dr \right| dx \\ &\leq \int_{|r|<1} \left\{ \int_{\mathbb{R}^n} \|\phi\|_{L^1(\mathbb{R}^n)} \left| {}^b\Omega_{|r|}^\alpha(u) \right| du \right\} |\omega(r)| |r|^{2\beta-1} dr \\ &\quad \text{by Minkowski integral inequality} \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} \int_{|r|<1} \left\{ \int_{|u|<1} \left(\frac{1}{1-|u|^2} \right)^{1-\operatorname{Re}\alpha} du \right\} |\omega(r)| |r|^{2\operatorname{Re}\beta-1} dr \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{c|\operatorname{Im}\alpha|}. \end{aligned} \quad (5. 2)$$

Recall φ_j^ν defined in (2. 5). Given $j > 0$, we have

$$\begin{aligned} {}^b\mathbf{P}_j^{\alpha \beta}(x) &= \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} {}^b\mathbf{P}_j^{\alpha \beta \nu}(x), \\ {}^b\mathbf{P}_j^{\alpha \beta \nu}(x) &= \int_{2^{j-1} \leq |r| < 2^j} e^{-2\pi i r} \left\{ \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \varphi_j^\nu(\xi) {}^b\widehat{\Omega}^\alpha(r\xi) d\xi \right\} \omega(r) |r|^{2\beta-1} dr. \end{aligned} \quad (5. 3)$$

Remark 5.1 By symmetry, it is suffice to assert $2^{j-1} \leq r < 2^j$.

Define

$$\begin{aligned} {}^b\mathbf{P}_r^\alpha(x) &= \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) {}^b\widehat{\Omega}^\alpha(r\xi) d\xi \\ &= \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \left(\frac{1}{r|\xi|} \right)^{\frac{n-1}{2} + \alpha - \frac{1}{2}} J_{\frac{n-1}{2} + \alpha - \frac{1}{2}}(2\pi r|\xi|) d\xi, \quad \text{by (2. 27)} \end{aligned} \quad (5. 4)$$

and

$${}^b\mathbf{P}_r^\alpha \varphi_j^\nu(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \varphi_j^\nu(\xi) \widehat{\phi}(\xi) {}^b\widehat{\Omega}^\alpha(r\xi) d\xi. \quad (5. 5)$$

By using the asymptotic expansion of Bessel functions in (A. 2)-(A. 3), we write

$${}^b\mathbf{P}_r^\alpha(x) = {}^b\mathbf{Q}_r^\alpha(x) + \sum_{k=1}^M {}^b\mathbf{Q}_{r,k}^\alpha(x) + {}^b\mathbf{R}_{r,M}^\alpha(x) \quad (5. 6)$$

for some $M \geq 1$ sufficiently large.

First, we have a principal term

$${}^b\mathbf{Q}_r^\alpha(x) = \frac{1}{\pi} \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \left(\frac{1}{r|\xi|} \right)^{\frac{n-1}{2}+\alpha} \cos \left[2\pi r |\xi| - \left(\frac{n}{2} + \alpha - 1 \right) \frac{\pi}{2} - \frac{\pi}{4} \right] d\xi. \quad (5. 7)$$

Next, there are finitely many

$$\begin{aligned} {}^b\mathbf{Q}_{r,k}^\alpha(x) &= \\ (2\pi)^{-2k} \frac{\mathbf{a}_k}{\pi} \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \left(\frac{1}{r|\xi|} \right)^{\frac{n-1}{2}+\alpha+2k} \cos \left[2\pi r |\xi| - \left(\frac{n}{2} + \alpha - 1 \right) \frac{\pi}{2} - \frac{\pi}{4} \right] d\xi \\ + (2\pi)^{-2k+1} \frac{\mathbf{b}_k}{\pi} \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \left(\frac{1}{r|\xi|} \right)^{\frac{n-1}{2}+\alpha+2k-1} \sin \left[2\pi r |\xi| - \left(\frac{n}{2} + \alpha - 1 \right) \frac{\pi}{2} - \frac{\pi}{4} \right] d\xi \\ k &= 1, 2, \dots, M \end{aligned} \quad (5. 8)$$

where $\mathbf{a}_k, \mathbf{b}_k$ are constants given in (A. 2).

Finally, there is a remainder term

$${}^b\mathbf{R}_{r,M}^\alpha(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\phi}(\xi) \mathbf{H}^\alpha(2\pi r |\xi|) d\xi \quad (5. 9)$$

of which

$$\partial_\xi^\gamma \mathbf{H}^\alpha(2\pi r |\xi|) = \mathbf{O}(r^{-2M-\frac{1}{2}}), \quad r \rightarrow \infty \quad (5. 10)$$

for every multi-index γ . The implied constant in (5. 10) is bounded by $\mathfrak{B}_{\text{Re}\alpha} \gamma e^{c|\text{Im}\alpha|}$.

By integration by parts *w.r.t* ξ inside (5. 9), we find

$$|{}^b\mathbf{R}_{r,M}^\alpha(x)| \leq \mathfrak{B}_{\text{Re}\alpha} e^{c|\text{Im}\alpha|} \left(\frac{1}{1+|x|} \right)^{n+1} r^{-2M-\frac{1}{2}}. \quad (5. 11)$$

Furthermore, we write

$${}^b\mathbf{P}_{r,j}^{\alpha,\nu}(x) = {}^b\mathbf{Q}_{r,j}^{\alpha,\nu}(x) + \sum_{k=1}^M {}^b\mathbf{Q}_{r,j,k}^{\alpha,\nu}(x) + {}^b\mathbf{R}_{r,j,M}^{\alpha,\nu}(x) \quad (5. 12)$$

of which ${}^b\mathbf{Q}_{r,j}^{\alpha,\nu}$ and $\mathbf{Q}_{r,j,k}^{\alpha,\nu}, {}^b\mathbf{R}_{r,j,N}^{\alpha,\nu}$ are defined as (5. 7) and (5. 8)-(5. 9) with $\widehat{\phi}$ replaced by $\widehat{\phi}\varphi_j^\nu$.

Recall ${}^b\mathbf{U}_j^{\alpha,\beta}$ and ${}^b\mathbf{V}_j^{\alpha,\beta}$ defined in (2. 30) and (2. 31).

We consider

$$\begin{aligned} {}^b\mathbf{U}_{r,j}^\alpha(x) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x) \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} {}^b\mathbf{Q}_{r,j}^{\alpha \nu}(x) + {}^b\mathbf{Q}_r^\alpha(x) [1 - \Psi_j^\alpha(x)], \\ {}^b\mathbf{U}_{r,j,k}^\alpha(x) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x) \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} {}^b\mathbf{Q}_{r,j,k}^{\alpha \nu}(x) + {}^b\mathbf{Q}_r^\alpha(x) [1 - \Psi_j^\alpha(x)], \quad k = 1, 2, \dots, M \end{aligned} \quad (5.13)$$

and

$$\begin{aligned} {}^b\mathbf{V}_{r,j}^\alpha(x) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} {}^b\mathbf{Q}_{r,j}^{\alpha \nu}(x), \\ {}^b\mathbf{V}_{r,j,k}^\alpha(x) &= \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x) \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} {}^b\mathbf{Q}_{r,j,k}^{\alpha \nu}(x), \quad k = 1, 2, \dots, M \end{aligned} \quad (5.14)$$

where $\Psi_j^{\alpha \mu}$, Ψ_j^α are defined in (2.9) and $\mathcal{U}_j^{\alpha \mu}$, $\mathcal{V}_j^{\alpha \mu}$ are defined in (2.10).

We claim

$$\int_{\mathbb{R}^n} |{}^b\mathbf{U}_{r,j}^\alpha(x)| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-jN} 2^{-j\varepsilon}, \quad N \geq 0 \quad (5.15)$$

and

$$\int_{\mathbb{R}^n} |{}^b\mathbf{V}_{r,j}^\alpha(x)| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-j\varepsilon} \quad (5.16)$$

for some $\varepsilon = \varepsilon(\operatorname{Re}\alpha) > 0$.

Remark 5.2 An analogue to the estimates regarding (5.15)-(5.16) gives

$$\int_{\mathbb{R}^n} |{}^b\mathbf{U}_{r,j,k}^\alpha(x)| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-j[N+2k-1]} 2^{-j\varepsilon}, \quad N \geq 0 \quad (5.17)$$

and

$$\int_{\mathbb{R}^n} |{}^b\mathbf{V}_{r,j,k}^\alpha(x)| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha \operatorname{Re}\beta} e^{\mathbf{C}|\operatorname{Im}\alpha|} 2^{-j(2k-1)} 2^{-j\varepsilon} \quad (5.18)$$

for every $k = 1, 2, \dots, N$.

Recall **Remark 5.1** and ${}^b\mathbf{P}_j^{\alpha \beta}$, ${}^b\mathbf{P}_j^{\alpha \beta \nu}$ given at (5.3). The above estimates in (5.11)-(5.18) together imply (2.33)-(2.34).

By using Euler's formulae, we replace the cosine function inside (5.7) with $\exp\{\pm 2\pi i r |\xi|\}$ multiplied by $\mathfrak{B}_{\operatorname{Re}\alpha} e^{\mathbf{C}|\operatorname{Im}\alpha|}$.

Note that $2^{j-1} \leq r < 2^j$. We consider

$${}^b\mathbf{A}_j^\alpha(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi \pm 2^j |\xi|} \widehat{\phi}(\xi) \left(\frac{1}{2^j |\xi|} \right)^{\frac{n-1}{2} + \alpha} d\xi \quad (5.19)$$

and

$${}^b\mathbf{A}_j^{\alpha \nu}(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi \pm 2^j |\xi|} \widehat{\phi}(\xi) \varphi_j^\nu(\xi) \left(\frac{1}{2^j |\xi|} \right)^{\frac{n-1}{2} + \alpha} d\xi. \quad (5.20)$$

Case One We aim to show

$$\int_{\mathbb{R}^n} \left| {}^b A_j^\alpha(x) \right| \left[1 - \Psi_j^\alpha(x) \right] dx \leq \mathfrak{B}_{\mathbf{Re}\alpha} e^{c|\text{Im}\alpha|} 2^{-jN} 2^{-j\varepsilon} \quad (5.21)$$

and

$$\sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n} \Psi_j^\alpha{}^\mu(x) \sum_{\nu \in \mathcal{U}_j^\alpha{}^\mu} \left| {}^b A_j^\alpha{}^\nu(x) \right| dx \leq \mathfrak{B}_{\mathbf{Re}\alpha} e^{c|\text{Im}\alpha|} 2^{-jN} 2^{-j\varepsilon} \quad (5.22)$$

for every $N \geq 0$. From (5.21)-(5.22), we conclude (5.15).

By changing $\xi \rightarrow 2^{-j}\xi$ inside (5.19), we write

$${}^b A_j^\alpha(x) = 2^{-jn} \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}x) \cdot \xi \pm |\xi|]} \widehat{\phi}(2^{-j}\xi) \left(\frac{1}{|\xi|} \right)^{\frac{n-1}{2} + \alpha} d\xi. \quad (5.23)$$

Moreover,

$$\begin{aligned} {}^b A_j^\alpha(x) &= \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} {}^b A_j^\alpha{}^\nu(x), \\ {}^b A_j^\alpha{}^\nu(x) &= 2^{-jn} \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}x) \cdot \xi \pm |\xi|]} \widehat{\phi}(2^{-j}\xi) \varphi_j^\nu(\xi) \left(\frac{1}{|\xi|} \right)^{\frac{n-1}{2} + \alpha} d\xi. \end{aligned} \quad (5.24)$$

Given ν fixed, \mathbf{L}_ν is the $n \times n$ -orthogonal matrix with $\det \mathbf{L}_\nu = 1$ defined as (2.8). Let $\xi = \mathbf{L}_\nu \eta$ and $\eta = (\eta_1, \eta')^T \in \mathbb{R} \times \mathbb{R}^{n-1}$. Denote $\eta_j^\nu = (1, 0)^T \in \mathbb{R} \times \mathbb{R}^{n-1}$. We have

$$\xi_j^\nu = \mathbf{L}_\nu \eta_j^\nu. \quad (5.25)$$

Recall that $\widehat{\phi} \in C_o^\infty(\mathbb{R}^n)$ is radially symmetric and $\text{supp} \widehat{\phi} \subset \{\xi \in \mathbb{R}^n : \frac{1}{3} < |\xi| \leq 3\}$. Let $x = \mathbf{L}_\nu u$. Rewrite (5.24) as

$${}^b A_j^\alpha{}^\nu(\mathbf{L}_\nu u) = 2^{-jn} \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}u) \cdot \eta \pm |\eta|]} \widehat{\phi}(2^{-j}\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} d\eta. \quad (5.26)$$

Consider

$$\begin{aligned} {}^b A_j^\alpha{}^\nu(\mathbf{L}_\nu u) &= 2^{-jn} \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}u) \cdot \eta \pm |\eta_1|]} \Theta_j^\alpha{}^\nu(\eta) d\eta, \\ \Theta_j^\alpha{}^\nu(\eta) &= e^{\pm 2\pi i [|\eta| - |\eta_1|]} \widehat{\phi}(2^{-j}\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha}. \end{aligned} \quad (5.27)$$

Recall $\text{supp} \varphi_j^\nu \subset \Gamma_j^\nu$ defined in (2.6).

Remark 5.3 Whenever $\widehat{\phi}(2^{-j}\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \neq 0$, we have $|\eta_1| \sim 2^j$ and $|\eta'| \lesssim 2^{j/2}$. Hence

$$\text{vol} \left\{ \text{supp} \Theta_j^\alpha{}^\nu \right\} \leq \mathfrak{B} 2^j 2^{j(\frac{n-1}{2})}. \quad (5.28)$$

Define a differential operator

$$\mathbf{D}f = Id + 2^{2j}\partial_{\eta_1}^2 f + 2^j \Delta_{\eta'} f. \quad (5.29)$$

We claim

$$|\mathbf{D}^N \Theta_j^\alpha \nu(\eta)| \leq \mathfrak{B}_{N \operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]}, \quad N \geq 0. \quad (5.30)$$

1. Denote $\chi(\eta) = \pm [| \eta | - |\eta_1|]$. We show

$$|\partial_\eta^\gamma \chi(\eta)| \leq \mathfrak{B}_\mu 2^{-j\gamma_1} 2^{-(j/2)[|\gamma| - \gamma_1]}, \quad \mathbf{L}_\nu \eta \in \Gamma_j^\nu \cap \{2^{j-2} \leq |\eta| \leq 2^{j+2}\} \quad (5.31)$$

for every multi-index γ . To conclude (5.31), we follow the lines in 4.5, chapter IX of Stein [3].

Note that $\chi(\eta)$ is homogeneous of degree 1 in η : $\chi(s\eta) = s\chi(\eta), s > 0$. We have

$$|\partial_\eta^\gamma \chi(\eta)| \leq \mathfrak{B}_\gamma |\eta|^{1-|\gamma|} \quad (5.32)$$

for every multi-index γ .

Indeed, $\partial_\eta^\gamma \chi(s\eta) = s^{|\gamma|} \partial_\xi^\gamma \chi(\xi)$ for $\xi = s\eta$. On the other hand, $\partial_\eta^\gamma \chi(s\eta) = \partial_\eta^\gamma [s\chi(\eta)] = s\partial_\eta^\gamma \chi(\eta)$. Choose $s = |\eta|^{-1}$. We find $\partial_\eta^\gamma \chi(\eta) = |\eta|^{1-|\gamma|} \partial_\xi^\gamma \chi(\xi)$ where $|\xi| = 1$.

Suppose $|\gamma| - \gamma_1 \geq 2$. From **Remark 5.3** and (5.32), we find

$$\begin{aligned} |\partial_\eta^\gamma \chi(\eta)| &\leq \mathfrak{B}_\gamma 2^{-j(|\gamma|-1)} \leq \mathfrak{B}_\gamma 2^{-j\gamma_1} 2^{-j[|\gamma| - \gamma_1 - 1]} \\ &\leq \mathfrak{B}_\gamma 2^{-j\gamma_1} 2^{-(j/2)[|\gamma| - \gamma_1]}. \end{aligned} \quad (5.33)$$

Suppose $|\gamma| - \gamma_1 = 0$ or 1. Denote $\eta^1 = (\eta_1, 0)^T \in \mathbb{R} \times \mathbb{R}^{n-1}$. A direct computation shows

$$(\nabla_\eta \chi)(\eta^1) = 0, \quad (\partial_{\eta_1}^N \nabla_{\eta'} \chi)(\eta^1) = (\nabla_{\eta'} \partial_{\eta_1}^N \chi)(\eta^1) = 0, \quad N \geq 0. \quad (5.34)$$

By writing out the Taylor expansion of $\partial_{\eta_1}^N \chi(\eta)$ and $\nabla_{\eta'} \partial_{\eta_1}^N \chi(\eta)$ in the η' -subspace and using $(\nabla_{\eta'} \partial_{\eta_1}^N \chi)(\eta^1) = 0$ from (5.34), we have

$$\partial_{\eta_1}^N \chi(\eta) = \mathbf{O}(|\eta'|^2 |\eta|^{-N-1}), \quad \nabla_{\eta'} \partial_{\eta_1}^N \chi(x, \eta) = \mathbf{O}(|\eta'| |\eta|^{-N-1}) \quad (5.35)$$

for every $N \geq 0$.

From **Remark 5.3** and (5.35), we find

$$|\partial_{\eta_1}^N \chi(\eta)| \leq \mathfrak{B}_N 2^{-jN}, \quad |\nabla_{\eta'} \partial_{\eta_1}^N \chi(\eta)| \leq \mathfrak{B}_N 2^{-jN} 2^{-j/2}, \quad N \geq 0. \quad (5.36)$$

2. We clearly have

$$\left| \partial_\eta^\gamma \left\{ \widehat{\phi} \left(2^{-j}\eta \right) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} \right\} \right| \leq \mathfrak{B}_\gamma \mathfrak{B}_{\operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} 2^{-j|\gamma|} \quad (5.37)$$

for every multi-index γ .

3. Recall φ_j^ν defined in (2. 5). Note that $\varphi_j^\nu(\mathbf{L}_\nu \eta)$ is homogeneous of degree 0 in η . Moreover, $\partial_\eta \varphi_j^\nu(\mathbf{L}_\nu \eta) = 0$ whenever $\left| \frac{\eta}{|\eta|} - \eta_j^\nu \right| \leq 2^{-j/2}$. A direct computation shows

$$\left| \partial_\eta^\gamma [\varphi_j^\nu(\mathbf{L}_\nu \eta)] \right| \leq \mathfrak{B}_\gamma 2^{|\gamma|j/2} |\eta|^{-|\gamma|} \quad (5. 38)$$

for every multi-index γ .

Geometrically, for every $\mathbf{L}_\nu \eta \in \Gamma_j^\nu$, the angle between η and η^1 is bounded by $\arcsin(2^{-j/2+1})$. Denote $\rho = |\eta|$. We find

$$\frac{\partial}{\partial \eta_1} = \left(\frac{\partial \rho}{\partial \eta_1} \right) \frac{\partial}{\partial \rho} + \mathbf{O}(2^{-j/2}) \cdot \nabla_{\eta'}, \quad \mathbf{L}_\nu \eta \in \Gamma_j^\nu. \quad (5. 39)$$

Because $\varphi_j^\nu(\mathbf{L}_\nu \eta)$ is homogeneous of degree 0 in η , we have

$$\partial_\rho [\varphi_j^\nu(\mathbf{L}_\nu \eta)] = 0. \quad (5. 40)$$

From (5. 39)-(5. 40), we obtain

$$\left| \partial_{\eta_1}^N [\varphi_j^\nu(\mathbf{L}_\nu \eta)] \right| \leq \mathfrak{B}_N |\eta|^{-N}, \quad N \geq 0. \quad (5. 41)$$

By putting together the above estimates in 1-3, we obtain (5. 30).

Recall ${}^b \mathbf{A}_j^{\alpha \nu}(\mathbf{L}_\nu u)$ from (5. 27). An N -fold integration by parts w.r.t \mathbf{D} shows

$${}^b \mathbf{A}_j^{\alpha \nu}(\mathbf{L}_\nu u) = (2\pi i)^{-2N} \left[1 + [u_1 \pm 2^j]^2 + 2^j \sum_{i=2}^n \left(\frac{u_i}{2^j} \right)^2 \right]^{-N} 2^{-jn} \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}u) \cdot \eta \pm |\eta_1|]} \mathbf{D}^N \Theta_j^{\alpha \nu}(\eta) d\eta. \quad (5. 42)$$

By using (5. 28) and (5. 30), we have

$$\left| \int_{\mathbb{R}^n} e^{2\pi i [(2^{-j}u) \cdot \eta \pm |\eta_1|]} \mathbf{D}^N \Theta_j^{\alpha \nu}(\eta) d\eta \right| \leq \mathfrak{B}_N \mathbf{Re} \alpha e^{c|\mathbf{Im} \alpha|} 2^{-j[\frac{n-1}{2} + \mathbf{Re} \alpha]} 2^j 2^{j(\frac{n-1}{2})}. \quad (5. 43)$$

Recall $\Psi_j^{\alpha \mu}$, Ψ_j^α defined in (2. 9). Define

$$\widetilde{\Psi}_j^{\alpha \mu}(x) = \begin{cases} 1, & \Psi_j^{\alpha \mu}(x) = 1, \\ 0, & \Psi_j^{\alpha \mu}(x) \neq 1. \end{cases} \quad (5. 44)$$

Let

$$R_j^\epsilon = \left\{ u \in \mathbb{R}^n : 2^j - 2^{\epsilon j} \leq |u_1| \leq 2^j + 2^{\epsilon j}, |u_i| \leq 2^{(\frac{1}{2}+\epsilon)j}, i = 2, \dots, n \right\} \quad (5. 45)$$

where $\epsilon = \epsilon(\mathbf{Re} \alpha) > 0$ is the small, adjustable number in (2. 9).

Observe that $\Psi_j^\alpha \mu(\mathbf{L}_\mu u) = 1$ if $u \in R_j^\epsilon$. We have

$$\begin{aligned}
& \int_{\mathbb{R}^n} \left| {}^\flat \mathbf{A}_j^\alpha(x) \right| \left[1 - \Psi_j^\alpha(x) \right] dx \\
& \leq \int_{\mathbb{R}^n} \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \left| {}^\flat \mathbf{A}_j^{\alpha \nu}(x) \right| \left[1 - \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \Psi_j^{\alpha \mu}(x) \right] dx \\
& \leq \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n \setminus \bigcup_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \text{supp } \tilde{\Psi}_j^{\alpha \mu}} \left| {}^\flat \mathbf{A}_j^{\alpha \nu}(x) \right| dx \\
& \leq \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n \setminus \text{supp } \tilde{\Psi}_j^{\alpha \nu}} \left| {}^\flat \mathbf{A}_j^{\alpha \nu}(x) \right| dx \\
& \leq \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n \setminus R_j^\epsilon} \left| {}^\flat \mathbf{A}_j^{\alpha \nu}(x) \right| dx.
\end{aligned} \tag{5. 46}$$

By using the estimates in (5. 42)-(5. 43), we find

$$\begin{aligned}
& \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} \int_{\mathbb{R}^n \setminus R_j^\epsilon} \left| {}^\flat \mathbf{A}_j^{\alpha \nu}(x) \right| dx \\
& \leq \mathfrak{B}_{N \operatorname{Re} \alpha} e^{\mathfrak{c} |\operatorname{Im} \alpha|} \sum_{\nu: \xi_j^\nu \in \mathbb{S}^{n-1}} 2^{-jn} 2^{-j[\frac{n-1}{2} + \operatorname{Re} \alpha]} 2^j 2^{j(\frac{n-1}{2})} \\
& \quad \int_{\mathbb{R}^n \setminus R_j^\epsilon} \left[1 + [u_1 \pm 2^j]^2 + 2^j \sum_{i=2}^n \left(\frac{u_i}{2^j} \right)^2 \right]^{-N} du \\
& \leq \mathfrak{B}_{N \operatorname{Re} \alpha} e^{\mathfrak{c} |\operatorname{Im} \alpha|} 2^{j(\frac{n-1}{2})} 2^{-jn} 2^{-j[\frac{n-1}{2} + \operatorname{Re} \alpha]} 2^j 2^{j(\frac{n-1}{2})} \int_{\mathbb{R}^n \setminus R_j^\epsilon} \left[1 + [u_1 \pm 2^j]^2 + \sum_{i=2}^n \left(\frac{u_i}{2^{j/2}} \right)^2 \right]^{-N} du \\
& = \mathfrak{B}_{N \operatorname{Re} \alpha} e^{\mathfrak{c} |\operatorname{Im} \alpha|} 2^{-j[\frac{n-1}{2} + \operatorname{Re} \alpha]} \int_{\mathbb{R}^n \setminus R_j^\epsilon} \left[1 + [u_1 \pm 2^j]^2 + \sum_{i=2}^n \left(\frac{u_i}{2^{j/2}} \right)^2 \right]^{-N} du \\
& = \mathfrak{B}_{N \operatorname{Re} \alpha} e^{\mathfrak{c} |\operatorname{Im} \alpha|} 2^{-j[\frac{n-1}{2} + \operatorname{Re} \alpha]} 2^{j(\frac{n-1}{2})} \\
& \quad \int_{\mathbb{R}^n \setminus \{2^{j-2^{\epsilon j}} \leq |u_1| \leq 2^j + 2^{\epsilon j}, |u_i| \leq 2^{\epsilon j}\}} \left[1 + [u_1 \pm 2^j]^2 + \sum_{i=2}^n u_i^2 \right]^{-N} du \\
& \leq \mathfrak{B}_{N \operatorname{Re} \alpha} e^{\mathfrak{c} |\operatorname{Im} \alpha|} 2^{-j \operatorname{Re} \alpha} 2^{-j\epsilon[N-(n+1)]}
\end{aligned} \tag{5. 47}$$

for every $N \geq 0$.

Remark 5.4 Suppose $v \in \mathcal{U}_j^{\alpha \mu} = \left\{ v : \left| \xi_j^\mu \pm \xi_j^\nu \right| > c2^{-j/2} \right\}$. Let $x = \mathbf{L}_v u \in \text{supp} \Psi_j^{\alpha \mu}$. There is at least one $i \in \{2, \dots, n\}$ such that $|u_i| \gtrsim 2^{j[\frac{1}{2}+\epsilon]}$.

Consider $x \in \text{supp} \Psi_j^{\alpha \mu}$ for j sufficiently large. By definition of $\Psi_j^{\alpha \mu}$ in (2. 9), we must have $|x| \approx 2^j$ and $\prod_{i=2}^n \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} \left| (\mathbf{L}_\mu^T x)_i \right| \right] \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} \left| (\mathbf{L}_\nu^T x)_i \right| \right] = 0$ whenever $v \in \mathcal{U}_j^{\alpha \mu}$. Suppose $\left| (\mathbf{L}_\nu^T x)_i \right| \leq 2^{j[\frac{1}{2}+\epsilon]}$ for every $i = 2, \dots, n$. We find $\prod_{i=2}^n \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} \left| (\mathbf{L}_\nu^T x)_i \right| \right] = 1$. This is a contradiction because $\prod_{i=2}^n \varphi \left[2^{-[\frac{1}{2}+\epsilon]j} \left| (\mathbf{L}_\mu^T x)_i \right| \right] \neq 0$ for x inside the support of $\Psi_j^{\alpha \mu}$.

The differential inequality in (5. 30) shows $\left| \partial_{\eta_\nu}^N \Theta_j^{\alpha \nu}(\eta) \right| \leq \mathfrak{B}_{N \operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} 2^{-jN/2}$ for every $N \geq 0$. By integration by parts w.r.t η_ν , we find

$$\begin{aligned} \left| {}^b \mathbf{A}_r^{\alpha \nu}(\mathbf{L}_\nu u) \right| &\leq \mathfrak{B}_N 2^{-jn} \left(2^{-j} u_i \right)^{-N} \left| \int_{\mathbb{R}^n} e^{2\pi i \left[(2^{-j} u) \cdot \eta \pm |\eta| \right]} \partial_{\eta_\nu}^N \Theta_j^{\alpha \nu}(\eta) d\eta \right| \\ &\leq \mathfrak{B}_N \operatorname{Re}\alpha e^{c|\operatorname{Im}\alpha|} 2^{-jn} 2^{jN/2} 2^{-jeN} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} 2^{-jN/2} 2^j 2^{j(\frac{n-1}{2})} \\ &= \mathfrak{B}_{\operatorname{Re}\alpha N} e^{c|\operatorname{Im}\alpha|} 2^{-jeN} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} 2^{-j(\frac{n-1}{2})}, \quad N \geq 0. \end{aligned} \quad (5. 48)$$

As defined in (2. 9), the volume of each $\text{supp} \Psi_j^{\alpha \mu}$ is bounded by $\mathfrak{B} 2^{j(\frac{n-1}{2})} 2^{jn\epsilon}$. We have

$$\begin{aligned} &\sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \sum_{\nu \in \mathcal{U}_j^{\alpha \mu}} \int_{\mathbb{R}^n} \Psi_j^{\alpha \mu}(\mathbf{L}_\nu u) \left| {}^b \mathbf{A}_r^{\alpha \nu}(\mathbf{L}_\nu u) \right| du \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha N} e^{c|\operatorname{Im}\alpha|} 2^{j(\frac{n-1}{2})} 2^{j(\frac{n-1}{2})} 2^{j(\frac{n-1}{2})} 2^{jn\epsilon} 2^{-jeN} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} 2^{-j(\frac{n-1}{2})} \quad \text{by (5. 48)} \\ &\leq \mathfrak{B}_{\operatorname{Re}\alpha N} e^{c|\operatorname{Im}\alpha|} 2^{j(\frac{n-1}{2})} 2^{jn\epsilon} 2^{-j\operatorname{Re}\alpha} 2^{-jeN}, \quad N \geq 0. \end{aligned} \quad (5. 49)$$

From (5. 47) and (5. 49), we obtain (5. 21)-(5. 22).

Case Two Let ${}^b \mathbf{A}_j^{\alpha \nu}$ defined in (5. 20). We aim to show

$$\sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \sum_{\nu \in \mathcal{V}_j^{\alpha \mu}} \int_{\mathbb{R}^n} \Psi_j^{\alpha \mu}(x) \left| {}^b \mathbf{A}_j^{\alpha \nu}(x) \right| dx \leq \mathfrak{B}_{\operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|} 2^{-je}, \quad \varepsilon = \varepsilon(\operatorname{Re}\alpha) > 0. \quad (5. 50)$$

This further implies (5. 16).

Remark 5.5 There are at most $\mathfrak{B} 2^{\epsilon(n-1)j}$ many $\nu \in \mathcal{V}_j^{\alpha \mu} = \left\{ \nu : \left| \xi_j^\mu \pm \xi_j^\nu \right| \leq c2^{-j/2} \right\}$.

Given μ and ν fixed, we have

$$\begin{aligned} \int_{\mathbb{R}^n} \Psi_j^{\alpha \mu}(x) \left| {}^b \mathbf{A}_j^{\alpha \nu}(x) \right| dx = \\ 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} \int_{\mathbb{R}^n} \Psi_j^{\alpha \mu}(x) \left| \int_{\mathbb{R}^n} e^{2\pi i [x \cdot \xi \pm 2^j |\xi|]} \widehat{\phi}(\xi) \varphi_j^\nu(\xi) \left(\frac{1}{|\xi|} \right)^{\frac{n-1}{2} + \alpha} d\xi \right| dx. \end{aligned} \quad (5.51)$$

Let $\xi = \mathbf{L}_\nu \eta$ as (5.25) and

$${}^b \mathbf{B}_j^{\alpha \nu}(x) = \int_{\mathbb{R}^n} e^{2\pi i [\mathbf{L}_\nu^T x \cdot \eta \pm 2^j |\eta|]} \widehat{\phi}(\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} d\eta. \quad (5.52)$$

Define

$$\Psi_j^{\alpha \mu \nu}(x) = \Psi_j^{\alpha \mu}(A_\nu x) \quad (5.53)$$

where A_ν is another $n \times n$ -orthogonal matrix with $\det A_\nu = 1$. We require

$$\mathbf{L}_\nu^T A_\nu x = (x_1, 0)^T \in \mathbb{R} \times \mathbb{R}^{n-1}. \quad (5.54)$$

In particular, $\operatorname{vol} \{ \operatorname{supp} \Psi_j^{\alpha \mu} \} = \operatorname{vol} \{ \operatorname{supp} \Psi_j^{\alpha \mu \nu} \} \leq \mathfrak{V} 2^{j(\frac{n-1}{2})} 2^{j n \epsilon}$.

Observe that

$$\begin{aligned} & \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \sum_{\nu \in \mathcal{V}_j^\alpha} \int_{\mathbb{R}^n} \Psi_j^{\alpha \mu}(x) \left| {}^b \mathbf{A}_j^{\alpha \nu}(x) \right| dx \\ & \leq \sum_{\mu: \xi_j^\mu \in \mathbb{S}^{n-1}} \sum_{\nu \in \mathcal{V}_j^\alpha} 2^{-j[\frac{n-1}{2} + \operatorname{Re}\alpha]} \int_{\operatorname{supp} \Psi_j^{\alpha \mu \nu}} \left| {}^b \mathbf{B}_j^{\alpha \nu}(A_\nu x) \right| dx. \end{aligned} \quad (5.55)$$

Because $\epsilon = \epsilon(\operatorname{Re}\alpha) > 0$ can be chosen sufficiently small, it is suffice to show

$$\begin{aligned} {}^b \mathbf{B}_j^{\alpha \nu}(A_\nu x) &= \int_{\mathbb{R}^n} e^{2\pi i [x_1 \eta_1 \pm 2^j |\eta|]} \widehat{\phi}(\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} d\eta \\ &= \mathbf{O}\left(2^{-j(\frac{n-1}{2})}\right), \quad j \rightarrow \infty \end{aligned} \quad (5.56)$$

where the implied constant is bounded by $\mathfrak{V}_{\operatorname{Re}\alpha} e^{c|\operatorname{Im}\alpha|}$.

Let

$$\Phi(\eta_1, \eta') = \pm [|\eta| - |\eta_1|]. \quad (5.57)$$

We consider

$$\begin{aligned} & \int_{\mathbb{R}^n} e^{2\pi i [x_1 \eta_1 \pm 2^j |\eta|]} \widehat{\phi}(\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} d\eta = \\ & e^{2\pi i [x_1 \eta_1 \pm 2^j |\eta_1|]} \int_{\mathbb{R}} \left\{ \int_{\mathbb{R}^{n-1}} e^{2\pi i 2^j \Phi(\eta_1, \eta')} \widehat{\phi}(\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2} + \alpha} d\eta' \right\} d\eta_1. \end{aligned} \quad (5.58)$$

Remark 5.6 For $\widehat{\phi}(\eta)\varphi_j^v(\mathbf{L}_v\eta) \neq 0$, we necessarily have $|\eta_1| \sim 1$ and $|\eta'| \lesssim 2^{-j/2}$.

A direct computation shows

$$\det \left[\frac{\partial^2 \Phi}{\partial \eta' \partial \eta'} \right] (\eta_1, \eta') = (-1)^{n-1} \frac{\eta_1^2}{[|\eta'|^2 + \eta_1^2]^{1+\frac{n-1}{2}}}. \quad (5.59)$$

Here, $\left[\frac{\partial^2 \Phi}{\partial \eta' \partial \eta'} \right]$ denotes the $(n-1) \times (n-1)$ Hessian matrix of Φ . See appendix B in the end.

Observe that $\Phi(\eta_1, 0) = 0$ and $\nabla_{\eta'} \Phi(\eta_1, 0) = 0$ whenever $|\eta_1| > 0$. Moreover, $\Phi(\eta_1, \eta')$ is non-degenerate at $\eta' = 0$. Next, we follow the lines in p. 346-347, chapter VIII of Stein [3]. Write

$$\begin{aligned} \Phi(\eta_1, \eta') &= \int_0^1 \frac{d}{dt} [\Phi(\eta_1, t\eta')] dt \\ &= \eta' \cdot \nabla_{\eta'} \Phi(\eta_1, \eta') - \int_0^1 t \frac{d^2}{dt^2} [\Phi(\eta_1, t\eta')] dt \\ &= \int_0^1 (1-t) \frac{d^2}{dt^2} [\Phi(\eta_1, t\eta')] dt. \end{aligned} \quad (5.60)$$

From (5.60), we find

$$\Phi(\eta_1, \eta') = \sum_{i,k=2}^n \xi_i \xi_k \dot{\Phi}_{ik}(\eta_1, \eta'), \quad (5.61)$$

$$\dot{\Phi}_{ik}(\eta_1, \eta') = \int_0^1 (1-t) \partial_{\eta_i} \partial_{\eta_k} \Phi(\eta_1, t\eta') dt.$$

In particular, $\dot{\Phi}_{ik}(\eta_1, 0) = \frac{1}{2} \left[\frac{\partial^2 \Phi}{\partial \eta_i \partial \eta_k} \right] (\eta_1, 0)$.

Consider $\Phi(\eta_1, \eta') = (\eta')^T \mathbf{A}(\eta_1, \eta') \eta'$ where

$$\mathbf{A}(\eta_1, \eta') = \begin{bmatrix} \dot{\Phi}_{22}(\eta_1, \eta') & \dots & \dot{\Phi}_{2n}(\eta_1, \eta') \\ \vdots & \ddots & \vdots \\ \dot{\Phi}_{n2}(\eta_1, \eta') & \dots & \dot{\Phi}_{nn}(\eta_1, \eta') \end{bmatrix}. \quad (5.62)$$

Denote $\lambda_i = \lambda_i(\eta_1, 0)$, $i = 2, \dots, n$ to be the $n-1$ real eigenvalues of $\mathbf{A}(\eta_1, 0)$.

Remark 5.7 We assume $\dot{\Phi}_{ii}(\eta_1, 0) = \lambda_i$ for every $i = 2, \dots, n$ and $\dot{\Phi}_{ik}(\eta_1, 0) = 0$ if $i \neq k$.

Indeed, write

$$\mathbf{A}(\eta_1, 0) = Q \begin{bmatrix} \lambda_2 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} Q^T \quad (5.63)$$

for which Q is an orthogonal matrix with $\det Q = 1$. By applying a linear transformation $\eta' \rightarrow Q\eta'$, we can verify **Remark 5.7**.

Note that $|\eta'| \lesssim 2^{-j/2}$ whenever $\varphi_j^\nu(\mathbf{L}_\nu \eta) \neq 0$. For j sufficiently large, we have $|\dot{\Phi}_{ii}(\eta_1, \eta')| > \frac{1}{2}$, $i = 2, \dots, n$ and $|\dot{\Phi}_{ik}(\eta_1, \eta')| \ll \frac{1}{2}$ if $i \neq k$. Now, define

$$\dot{\zeta}_2 = [\pm \dot{\Phi}_{22}(\eta_1, \eta')]^{\frac{1}{2}} \left[\eta_2 \pm \sum_{i=3}^n \frac{\eta_i \dot{\Phi}_{i2}(\eta_1, \eta')}{\dot{\Phi}_{22}(\eta_1, \eta')} \right] \quad (5.64)$$

where the \pm sign depends on the value of $\lambda_2 = \dot{\Phi}_{22}(\eta_1, 0)$.

From direct computation, we find

$$\Phi(\eta_1, \eta') = \pm \dot{\zeta}_2^2 + \sum_{i=3}^n \sum_{k=3}^n \eta_i \eta_k \ddot{\Phi}_{ik}(\eta_1, \eta') \quad (5.65)$$

where

$$\ddot{\Phi}_{ii}(\eta_1, \eta') = \dot{\Phi}_{ii}(\eta_1, \eta') \pm \frac{\dot{\Phi}_{2i}(\eta_1, \eta') \dot{\Phi}_{i2}(\eta_1, \eta')}{|\dot{\Phi}_{22}(\eta_1, \eta')|} \quad (5.66)$$

and

$$\ddot{\Phi}_{ik}(\eta_1, \eta') = \dot{\Phi}_{ik}(\eta_1, \eta') \pm 2 \frac{\dot{\Phi}_{2i}(\eta_1, \eta') \dot{\Phi}_{2k}(\eta_1, \eta')}{|\dot{\Phi}_{22}(\eta_1, \eta')|}, \quad i \neq k. \quad (5.67)$$

Observe that $\ddot{\Phi}_{ii}(\eta_1, 0) = \dot{\Phi}_{ii}(\eta_1, 0) = \lambda_i$ for $i = 3, \dots, n$ and $\ddot{\Phi}_{ik}(\eta_1, 0) = \dot{\Phi}_{ik}(\eta_1, 0) = 0$ if $i \neq k$. We repeat this process until

$$\Phi(\eta_1, \eta') = \sum_{i=2}^n \pm \dot{\zeta}_i^2. \quad (5.68)$$

Consequently, we find

$$\begin{aligned} \frac{\partial \dot{\zeta}_i}{\partial \eta_i}(\eta_1, 0) &= [\pm \lambda_i]^{\frac{1}{2}}, \quad i = 2, \dots, n \\ \frac{\partial \dot{\zeta}_i}{\partial \eta_k}(\eta_1, 0) &= 0, \quad i \neq k. \end{aligned} \quad (5.69)$$

Lastly, we define $\dot{\zeta}_i = |\lambda_i|^{\frac{1}{2}} \zeta_i$, $i = 2, \dots, n$. Denote $\zeta' = (\zeta_2, \dots, \zeta_n)^T \in \mathbb{R}^{n-1}$.

Recall **Remark 5.6**.

Remark 5.8 From the above construction, we find $|\zeta'| \lesssim 2^{-j/2}$.

For j sufficiently large, the Inverse Function Theorem guarantees

$$\begin{aligned} &\int_{\mathbb{R}^{n-1}} e^{2\pi i 2^j \Phi(\eta_1, \eta')} \widehat{\phi}(\eta) \varphi_j^\nu(\mathbf{L}_\nu \eta) \left(\frac{1}{|\eta|} \right)^{\frac{n-1}{2}+\alpha} d\eta' = \\ &\int_{\mathbb{R}^{n-1}} e^{2\pi i 2^j [\lambda_2 \zeta_2^2 + \dots + \lambda_n \zeta_n^2]} \widehat{\phi} \left[\sqrt{\eta_1^2 + |\eta'(\zeta')|^2} \right] \varphi_j^\nu [\mathbf{L}_\nu(\eta_1, \eta'(\zeta'))^T] \\ &\left\{ \frac{1}{\sqrt{\eta_1^2 + |\eta'(\zeta')|^2}} \right\}^{\frac{n-1}{2}+\alpha} \mathfrak{J}(\zeta') d\zeta' \end{aligned} \quad (5.70)$$

where $\eta'(\zeta')$ is continuously differentiable and

$$\Im(\zeta') = \det \begin{bmatrix} \frac{\partial \eta_2}{\partial \zeta_2}(\zeta') & \dots & \frac{\partial \eta_n}{\partial \zeta_2}(\zeta') \\ \vdots & \ddots & \vdots \\ \frac{\partial \eta_2}{\partial \zeta_n}(\zeta') & \dots & \frac{\partial \eta_n}{\partial \zeta_n}(\zeta') \end{bmatrix}. \quad (5.71)$$

Let $\lambda \in \mathbb{R}$ and $\zeta, y \in \mathbb{R}$. We have

$$\int_{-\infty}^{\infty} e^{2\pi i 2^j \lambda \zeta^2} e^{-2\pi i \zeta y} d\zeta = (-2i 2^j \lambda)^{-\frac{1}{2}} e^{-\pi y^2 / (-2i 2^j \lambda)}. \quad (5.72)$$

By changing $\zeta = z/\sqrt{-2i 2^j \lambda}$ in the left hand side of (5.72), the integrand $e^{-\pi z^2} e^{-2\pi i z y / \sqrt{-2i 2^j \lambda}}$ is analytic whose norm decays to zero as $|z| \rightarrow \infty$ inside the region: $|\text{Im} z| < |\text{Re} z|$. By applying Cauchy's theorem, we find

$$\begin{aligned} \int_{-\infty}^{\infty} e^{2\pi i 2^j \lambda \zeta^2} e^{-2\pi i \zeta y} d\zeta &= (-2i 2^j \lambda)^{-\frac{1}{2}} \int_{-\infty}^{\infty} e^{-\pi z^2} e^{-2\pi i z y / \sqrt{-2i 2^j \lambda}} dz \\ &= (-2i 2^j \lambda)^{-\frac{1}{2}} e^{-\pi y^2 / (-2i 2^j \lambda)} \end{aligned} \quad (5.73)$$

where the second equality holds because

$$e^{-\pi z^2} = \int_{-\infty}^{\infty} e^{-\pi s^2} e^{-2\pi i s z} ds, \quad z \in \mathbb{C}. \quad (5.74)$$

By using the identity in (5.72) and applying Plancherel theorem for $(n-1)$ -times, we have (5.70) equal to

$$\begin{aligned} &\prod_{i=2}^n \left(\frac{i}{2^{j+1} \lambda_i} \right)^{\frac{1}{2}} \int_{\mathbb{R}^{n-1}} e^{-\pi [y_2^2/\lambda_2 + \dots + y_n^2/\lambda_n] \left(\frac{i}{2^{j+1}} \right)} \Psi(y') dy', \\ \Psi(y') &= \int_{\mathbb{R}^{n-1}} e^{-2\pi i y' \cdot \zeta'} \widehat{\phi} \left[\sqrt{\eta_1^2 + |\eta'(\zeta')|^2} \right] \varphi_j^\nu \left[\mathbf{L}_\nu(\eta_1, \eta'(\zeta'))^T \right] \\ &\quad \left\{ \frac{1}{\sqrt{\eta_1^2 + |\eta'(\zeta')|^2}} \right\}^{\frac{n-1}{2} + \alpha} \Im(\zeta') d\zeta'. \end{aligned} \quad (5.75)$$

By changing dilations $\zeta' \rightarrow 2^{-j/2} \zeta'$ and $y' \rightarrow 2^{j/2} y'$, we rewrite (5.75) as

$$2^{-j(\frac{n-1}{2})} \prod_{i=2}^n \left(\frac{i}{2\lambda_i} \right)^{\frac{1}{2}} \int_{\mathbb{R}^{n-1}} e^{-\pi [y_2^2/\lambda_2 + \dots + y_n^2/\lambda_n] (\frac{i}{2})} \Psi(2^{j/2} y') dy',$$

$$\begin{aligned} \Psi(2^{j/2} y') &= \int_{\mathbb{R}^{n-1}} e^{-2\pi i y' \cdot \zeta'} \widehat{\phi} \left[\sqrt{\eta_1^2 + |\eta'(2^{-j/2} \zeta')|^2} \right] \varphi_j^\nu \left[\mathbf{L}_\nu(\eta_1, \eta'(2^{-j/2} \zeta'))^T \right] \\ &\quad \left\{ \frac{1}{\sqrt{\eta_1^2 + |\eta'(2^{-j/2} \zeta')|^2}} \right\}^{\frac{n-1}{2} + \alpha} \Im(2^{-j/2} \zeta') d\zeta'. \end{aligned} \quad (5.76)$$

Recall φ_j^ν defined in (2. 5) satisfying the differential inequality in (5. 38). A direct computation shows

$$\begin{aligned} & \left| \partial_{\zeta'}^\gamma \left\{ \widehat{\phi} \left[\sqrt{\eta_1^2 + |\eta'(2^{-j/2}\zeta')|^2} \right] \varphi_j^\nu \left[\mathbf{L}_\nu(\eta_1, \eta'(2^{-j/2}\zeta'))^T \right] \left\{ \frac{1}{\sqrt{\xi_1^2 + |\eta'(2^{-j/2}\zeta')|^2}} \right\}^{\frac{n-1}{2}+\alpha} \Im(2^{-j/2}\zeta') \right\} \right| \\ & \leq \mathfrak{B}_\gamma \operatorname{Re} \alpha e^{c|\operatorname{Im} \alpha|} \end{aligned} \quad (5. 77)$$

for every multi-index γ . By using (5. 77) and integration by parts, we find $\Psi(2^{j/2}\gamma')$ in (5. 76) to be an integrable function in \mathbb{R}^{n-1} .

Lastly, (5. 59) implies $\prod_{i=2}^n \lambda_i = \det \left[\frac{\partial^2 \Phi}{\partial \eta' \partial \eta'} \right](\eta_1, 0) = (-1)^{n-1} [\eta_1^2]^{-\frac{n-1}{2}} \approx \pm 1$.

A Some estimates regarding Bessel functions

Recall a number of well known results for Bessel functions.

- For $a > -\frac{1}{2}, b \in \mathbb{R}$ and $\rho > 0$, a Bessel function J_{a+ib} has an integral formula

$$J_{a+ib}(\rho) = \frac{(\rho/2)^{a+ib}}{\pi^{\frac{1}{2}} \Gamma(a + \frac{1}{2} + ib)} \int_{-1}^1 e^{i\rho s} (1-s^2)^{a-\frac{1}{2}+ib} ds. \quad (\text{A. 1})$$

- For every $a > -\frac{1}{2}, b \in \mathbb{R}$ and $\rho > 0$, we have

$$\begin{aligned} J_{a+ib}(\rho) & \sim \left(\frac{\pi \rho}{2} \right)^{-\frac{1}{2}} \cos \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \\ & + \left(\frac{\pi \rho}{2} \right)^{-\frac{1}{2}} \sum_{k=1}^{\infty} \cos \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \mathbf{a}_k \rho^{-2k} + \sin \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \mathbf{b}_k \rho^{-2k+1}, \\ \mathbf{a}_k & = (-1)^k [a+ib, 2k] 2^{-2k}, \quad \mathbf{b}_k = (-1)^{k+1} [a+ib, 2k-1] 2^{-2k+1}, \\ [a+ib, m] & = \frac{\Gamma(\frac{1}{2} + a + ib + m)}{m! \Gamma(\frac{1}{2} + a + ib - m)}, \quad m = 0, 1, 2, \dots \end{aligned} \quad (\text{A. 2})$$

in the sense of that

$$\begin{aligned} & \left(\frac{d}{d\rho} \right)^\ell \left[J_{a+ib}(\rho) - \left(\frac{\pi \rho}{2} \right)^{-\frac{1}{2}} \cos \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \right. \\ & \left. - \left(\frac{\pi \rho}{2} \right)^{-\frac{1}{2}} \sum_{k=1}^N \cos \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \mathbf{a}_k \rho^{-2k} + \sin \left[\rho - (a+ib) \frac{\pi}{2} - \frac{\pi}{4} \right] \mathbf{b}_k \rho^{-2k+1} \right] \quad (\text{A. 3}) \\ & = \mathbf{O}(\rho^{-2N-\frac{1}{2}}) \quad N \geq 0, \quad \ell \geq 0, \quad \rho \rightarrow \infty \end{aligned}$$

where the implied constant is bounded by $\mathfrak{B}_a e^{\mathfrak{C}|b|}$.

- For every $a, b \in \mathbb{R}$ and $\rho > 0$,

$$J_{a-1+ib}(\rho) = 2 \left[\frac{a+ib}{\rho} \right] J_{a+ib}(\rho) - J_{a+1+ib}(\rho). \quad (\text{A. 4})$$

By using (A. 1) and (A. 2) together with (A. 4), we find the norm estimate in below.

- For every $a, b \in \mathbb{R}$ and $\rho > 0$,

$$\left| \frac{1}{\rho^{a+ib}} J_{a+ib}(\rho) \right| \leq \mathfrak{B}_a \left(\frac{1}{1+\rho} \right)^{\frac{1}{2}+a} e^{\mathfrak{C}|b|}. \quad (\text{A. 5})$$

More discussion of Bessel functions can be found in the book of Watson [6].

Let $z \in \mathbb{C}$. Ω^z is a distribution defined by analytic continuation from

$$\operatorname{Re} z < 1, \quad \Omega^z(x) = \pi^{-z} \Gamma^{-1}(1-z) \left(\frac{1}{1-|x|^2} \right)_+^z. \quad (\text{A. 6})$$

- Ω^z can be equivalently defined by

$$\widehat{\Omega}^z(\xi) = \left(\frac{1}{|\xi|} \right)^{\frac{n}{2}-z} J_{\frac{n}{2}-z}(2\pi|\xi|), \quad z \in \mathbb{C}. \quad (\text{A. 7})$$

From (A. 6), we have

$$\begin{aligned} \widehat{\Omega}^z(\xi) &= \pi^{-z} \Gamma^{-1}(1-z) \int_{|x|<1} e^{-2\pi i x \cdot \xi} \left(\frac{1}{1-|x|^2} \right)^z dx \\ &= \pi^{-z} \Gamma^{-1}(1-z) \int_0^\pi \left\{ \int_0^1 e^{-2\pi i |\xi|r \cos \vartheta} (1-r^2)^{-z} r^{n-1} dr \right\} \omega_{n-2} \sin^{n-2} \vartheta d\vartheta \\ &= \omega_{n-2} \pi^{-z} \Gamma^{-1}(1-z) \int_{-1}^1 \left\{ \int_0^1 e^{2\pi i |\xi|rs} (1-r^2)^{-z} r^{n-1} dr \right\} (1-s^2)^{\frac{n-3}{2}} ds \quad (-s = \cos \vartheta) \\ &= \omega_{n-2} \pi^{-z} \Gamma^{-1}(1-z) \int_0^1 \left\{ \int_{-1}^1 \cos(2\pi|\xi|rs) (1-s^2)^{\frac{n-3}{2}} ds \right\} (1-r^2)^{-z} r^{n-1} dr \\ &= \omega_{n-2} \pi^{-z} \Gamma^{-1}(1-z) \int_0^1 \left\{ \int_{-1}^1 \cos(2\pi|\xi|rs) (1-s^2)^{\frac{n-3}{2}} ds \right\} (1-r^2)^{-z} r^{n-1} dr \end{aligned} \quad (\text{A. 8})$$

where

$$\omega_{n-2} = 2\pi^{\frac{n-1}{2}} \Gamma^{-1}\left(\frac{n-1}{2}\right). \quad (\text{A. 9})$$

Recall the Beta function identity:

$$\frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)} = \int_0^1 r^{z-1} (1-r)^{w-1} dr, \quad \operatorname{Re} z > 0, \quad \operatorname{Re} w > 0. \quad (\text{A. 10})$$

By writing out the Taylor expansion of the cosine function inside (A. 8), we find

$$\begin{aligned}
& \omega_{n-2} \pi^{-z} \Gamma^{-1} (1-z) \int_0^1 \left\{ \int_{-1}^1 \cos(2\pi|\xi|rs) (1-s^2)^{\frac{n-3}{2}} ds \right\} (1-r^2)^{-z} r^{n-1} dr \\
&= \omega_{n-2} \pi^{-z} \Gamma^{-1} (1-z) \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \left\{ \int_{-1}^1 s^{2k} (1-s^2)^{\frac{n-3}{2}} ds \right\} \left\{ \int_0^1 r^{2k+n-1} (1-r^2)^{-z} dr \right\} \\
&= \frac{1}{2} \omega_{n-2} \pi^{-z} \Gamma^{-1} (1-z) \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \left\{ \int_0^1 t^{k+\frac{1}{2}-1} (1-t)^{\frac{n-1}{2}-1} dt \right\} \left\{ \int_0^1 \rho^{k+\frac{n}{2}-1} (1-\rho)^{1-z-1} d\rho \right\} \quad (t=s^2, \rho=r^2) \\
&= \frac{1}{2} \omega_{n-2} \pi^{-z} \Gamma^{-1} (1-z) \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \frac{\Gamma(k+\frac{1}{2}) \Gamma(\frac{n-1}{2})}{\Gamma(k+\frac{n}{2})} \frac{\Gamma(k+\frac{n}{2}) \Gamma(1-z)}{\Gamma(k+\frac{n}{2}+1-z)} \quad \text{by (A. 10)} \\
&= \pi^{\frac{n-1}{2}-z} \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \frac{\Gamma(k+\frac{1}{2})}{\Gamma(k+\frac{n}{2}+1-z)}, \quad \text{by (A. 9).}
\end{aligned} \tag{A. 11}$$

On the other hand, we have

$$\begin{aligned}
\left(\frac{1}{|\xi|} \right)^{\frac{n}{2}-z} J_{\frac{n}{2}-z}(2\pi|\xi|) &= \pi^{\frac{n-1}{2}-z} \Gamma^{-1}\left(\frac{n+1}{2}-z\right) \int_{-1}^1 e^{2\pi i |\xi| s} (1-s^2)^{\frac{n-1}{2}-z} ds \quad \text{by (A. 1)} \\
&= \pi^{\frac{n-1}{2}-z} \Gamma^{-1}\left(\frac{n+1}{2}-z\right) \int_{-1}^1 \cos(2\pi|\xi|s) (1-s^2)^{\frac{n-1}{2}-z} ds \\
&= \pi^{\frac{n-1}{2}-z} \Gamma^{-1}\left(\frac{n+1}{2}-z\right) \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \int_{-1}^1 s^{2k} (1-s^2)^{\frac{n-1}{2}-z} ds \\
&= \pi^{\frac{n-1}{2}-z} \Gamma^{-1}\left(\frac{n+1}{2}-z\right) \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \int_0^1 \rho^{k+\frac{1}{2}-1} (1-\rho)^{\frac{n+1}{2}-z-1} d\rho \quad (\rho=s^2) \\
&= \pi^{\frac{n-1}{2}-z} \sum_{k=0}^{\infty} (-1)^k \frac{(2\pi|\xi|)^{2k}}{(2k)!} \frac{\Gamma(k+\frac{1}{2})}{\Gamma(k+\frac{n}{2}+1-z)} \quad \text{by (A. 10).}
\end{aligned} \tag{A. 12}$$

B Determinant of the Hessian matrix

Let $\xi = (\xi_1, \xi') \in \mathbb{R} \times \mathbb{R}^{n-1}$. Consider $\Phi(\xi_1, \xi') = \sqrt{|\xi'|^2 + \xi_1^2} - |\xi_1|$. We find

$$\partial_{\xi_i} \Phi(\xi_1, \xi') = \frac{\xi_i}{\sqrt{|\xi'|^2 + \xi_1^2}}, \quad i = 2, \dots, n \quad (\text{B. 1})$$

and

$$\begin{aligned} \partial_{\xi_i}^2 \Phi(\xi_1, \xi') &= \frac{1}{\sqrt{|\xi'|^2 + \xi_1^2}} - \frac{\xi_i^2}{\left[|\xi'|^2 + \xi_1^2\right]^{\frac{3}{2}}}, & \partial_{\xi_i} \partial_{\xi_k} \Phi(\xi_1, \xi') &= - \frac{\xi_i \xi_k}{\left[|\xi'|^2 + \xi_1^2\right]^{\frac{3}{2}}}, \\ && i, k &= 2, \dots, n. \end{aligned} \quad (\text{B. 2})$$

By using (B. 2), we write

$$\begin{aligned} \det \left[\frac{\partial^2 \Phi}{\partial \xi' \partial \xi'} \right] (\xi_1, \xi') &= \left[|\xi'|^2 + \xi_1^2 \right]^{-\frac{3}{2}(n-1)} \det \begin{bmatrix} |\xi'|^2 + \xi_1^2 - \xi_2^2 & \cdots & -\xi_2 \xi_n \\ \vdots & \ddots & \vdots \\ -\xi_n \xi_2 & \cdots & |\xi'|^2 + \xi_1^2 - \xi_n^2 \end{bmatrix} \\ &\doteq \left[|\xi'|^2 + \xi_1^2 \right]^{-\frac{3}{2}(n-1)} \det \mathbf{A}(\xi_1, \xi'). \end{aligned} \quad (\text{B. 3})$$

The determinant of $\mathbf{A}(\xi_1, \xi')$ can be computed as follows.

$$\begin{aligned} \det \mathbf{A}(\xi_1, \xi') &= \\ \det \begin{bmatrix} 1 & \xi_2 & \cdots & \xi_n \\ 0 & |\xi'|^2 + \xi_1^2 - \xi_2^2 & \cdots & -\xi_2 \xi_n \\ \vdots & \vdots & \ddots & \vdots \\ 0 & -\xi_n \xi_2 & \cdots & |\xi'|^2 + \xi_1^2 - \xi_n^2 \end{bmatrix} &= \det \begin{bmatrix} 1 & \xi_2 & \cdots & \xi_n \\ \xi_2 & |\xi'|^2 + \xi_1^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \xi_n & 0 & \cdots & |\xi'|^2 + \xi_1^2 \end{bmatrix} \\ &= \det \begin{bmatrix} 1 - \frac{\xi_2^2}{|\xi'|^2 + \xi_1^2} - \cdots - \frac{\xi_n^2}{|\xi'|^2 + \xi_1^2} & \xi_2 & \cdots & \xi_n \\ 0 & |\xi'|^2 + \xi_1^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & |\xi'|^2 + \xi_1^2 \end{bmatrix}. \end{aligned} \quad (\text{B. 4})$$

From (B. 3)-(B. 4), we have

$$\begin{aligned} \det \left[\frac{\partial^2 \Phi}{\partial \xi' \partial \xi'} \right] (\xi_1, \xi') &= \left[|\xi'|^2 + \xi_1^2 \right]^{-\frac{3}{2}(n-1)} \left[1 - \frac{\xi_2^2}{|\xi'|^2 + \xi_1^2} - \cdots - \frac{\xi_n^2}{|\xi'|^2 + \xi_1^2} \right] \left[|\xi'|^2 + \xi_1^2 \right]^{n-1} \\ &= \frac{\xi_1^2}{\left[|\xi'|^2 + \xi_1^2 \right]^{1+\frac{n-1}{2}}}. \end{aligned} \quad (\text{B. 5})$$

References

- [1] C. Fefferman, *A note on spherical summation multipliers*, Israel J. Math **15**: 44-52, 1973.
- [2] A. Seeger, C. D. Sogge and E. M. Stein, *Regularity Properties of Fourier Integral Operators*, Annals of Mathematics, **134**: 231-251, 1991.
- [3] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton University Press, 1993.
- [4] E. M. Stein, *Interpolation of Linear Operators*, Transactions of the A. M. S **83**: no.2, 482-492, 1956.
- [5] R. Strichartz, *Convolutions with kernels having singularity on a sphere*, Transaction of the American Mathematical Society **148**: 461-471, 1970.
- [6] G. N. Watson, *Theory of Bessel Functions*, Cambridge University Press, Cambridge, 1944.

Department of Mathematics, Westlake University
wangzipeng@westlake.edu.cn