TENSOR PRODUCT SURFACES AND QUADRATIC SYZYGIES

MATTHEW WEAVER

ABSTRACT. For $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b))$ a four-dimensional vector space, a basis $\{p_0, p_1, p_2, p_3\}$ of U defines a rational map $\phi_U : \mathbb{P}^1 \times \mathbb{P}^1 \longrightarrow \mathbb{P}^3$. The tensor product surface associated to U is the closed image X_U of the map ϕ_U . These surfaces arise within the field of geometric modelling, in which case it is particularly desirable to obtain the implicit equation of X_U . In this paper, we study X_U via the syzygies of the associated bigraded ideal $I_U = (p_0, p_1, p_2, p_3)$ when U is free of basepoints, i.e. ϕ_U is regular. Expanding upon work of Duarte and Schenck [13] for such ideals with a linear syzygy, we address the case that I_U has a quadratic syzygy.

1. INTRODUCTION

A classical problem within algebraic geometry is to find the implicit equations of the image or graph of a rational map between projective spaces. This so-called *implicitization problem* has been studied to great length by algebraic geometers and commutative algebraists alike. More recently, the problem has gained interest within the geometric modeling community for its applications to computer-aided geometric design (CAGD), see e.g. [12, 20, 21, 22]. In this context, knowledge of the implicit equations of a curve or surface allows for more efficient computation and geometric rendering. For instance, determining whether a point lies on a surface is trivial given an implicit form, but requires solving a possibly large polynomial system of equations given a parametric form.

In this context, two situations often considered are rational maps $\mathbb{P}^2 \dashrightarrow \mathbb{P}^3$ and $\mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow \mathbb{P}^3$. Surfaces defined as the image of the first map are *triangular surfaces*, whereas surfaces defined as the image of the second are called *tensor product surfaces*. Implicitization of these surfaces has been studied extensively using a variety of techniques, such as Gröbner bases, resultants, and syzygies. Whereas these first two methods are classical and straightforward to implement, they are typically computationally intensive and slow. In contrast, syzygy techniques, and methods borrowed from the study of *Rees algebras*, are much more effective, see e.g. [1, 2, 3, 4]. One such tool in this realm is the *approximation complex* \mathcal{Z} introduced by Herzog, Simis, and Vasconcelos [17, 18], which has proven particularly useful for implicitization purposes [1, 3, 7]. We refer the reader to [8] for a wonderful overview of syzygy methods used for implicitization.

In this paper, we study tensor product surfaces and adopt the following setting. Let R = k[s, t, u, v] be a polynomial ring over an algebraically closed field k, bigraded by setting bideg s, t = (1, 0) and bideg u, v = (0, 1). We note that the global sections $H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b))$ correspond to the bigraded components $R_{a,b}$ of R. Let $U \subseteq R_{a,b}$ denote a subspace with basis $\{p_0, p_1, p_2, p_3\}$, such that p_0, p_1, p_2, p_3 have no common zeros on $\mathbb{P}^1 \times \mathbb{P}^1$, i.e. U is basepoint free. With this, consider the regular map

$$\phi_U: \mathbb{P}^1 \times \mathbb{P}^1 \longrightarrow \mathbb{P}^3$$

defined by $\{p_0, p_1, p_2, p_3\}$ and let $X_U = \phi_U(\mathbb{P}^1 \times \mathbb{P}^1) \subseteq \mathbb{P}^3$. Writing $I_U = (p_0, p_1, p_2, p_3)$, the assumption that U is free of basepoints is equivalent to $\sqrt{I_U} = (s, t) \cap (u, v)$.

From [2], it is well known that the implicit equation of X_U can be determined from the approximation complex \mathcal{Z} on a generating set of I_U (see Section 2 for details). As this complex relates to the module of syzygies of I_U , knowledge of a free resolution of I_U is particularly fruitful in this direction. In [19], tensor product surfaces X_U for $U \subset R_{2,1}$ are studied through a free resolution of I_U , of which the possible shapes are determined. There it is noted that the existence of a *linear* syzygy, in bidegree (0, 1) or (1, 0), on I_U yields certain constraints. This idea is further developed in [13], where this phenomenon is shown to extend beyond generation in bidegree (2, 1). Furthermore, in [13] it is shown how the implicit equation of X_U can be determined from the presence of a linear syzygy, without the need of a full resolution of I_U . In this direction, we show in the present article that a similar result can be achieved if I_U has a *quadratic* syzygy. The key aspect in the arguments of [13] is that a syzygy in bidegree (0, 1), or (1, 0), allows one to construct a specific generating set of I_U . This, in turn, allows for more detailed study of the remaining syzygies, with respect to this generation. In particular, one may produce a subset of syzygies which determines the bigraded strand $\mathcal{Z}_{2a-1,b-1}$ of the approximation complex, and hence the implicit equation of X_U , following [1].

In the case that I_U has a syzygy in either bidegree (0, 2) or (2, 0), we show that a similar phenomenon occurs, within a handful of cases. The main innovation presented here is the construction of a particular subspace $V \subseteq U$, determined by the quadratic syzygy (see Section 3 for details), with dim V serving as the invariant for each case. The main results of this article, Theorems 4.3 and 5.4, are summarized as follows.

Theorem. Assume that $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b))$ is basepoint free, with $b \ge 3$. Let I_U denote the ideal of U, and assume that I_U has a minimal first syzygy Q in bidegree (0, 2), and no linear syzygy. Write V to denote the subspace of U associated to the syzygy Q.

- (i) If dim V = 2, then I_U has two syzygies S_1, S_2 of bidegree (a, b 2) such that $\langle Q, S_1, S_2 \rangle$ determines the first differential d_1 of $\mathcal{Z}_{2a-1,b-1}$.
- (ii) If dim V = 3, then I_U has one syzygy S_1 of bidegree (a, b 2) and two syzygies S_2, S_3 of bidegree (a, b 1) such that $\langle Q, S_1, S_2, S_3 \rangle$ determines the first differential d_1 of $\mathcal{Z}_{2a-1,b-1}$.

Moreover, following [1], the determinant of a $2ab \times 2ab$ matrix representation of d_1 is a power of the implicit equation F of X_U .

By symmetry, a similar result holds if I_U has a syzygy in bidegree (2,0) and $a \ge 3$. The key aspect of the result above is the formulation of the subspace V, from a suitable basis of which the additional syzygies are constructed. Furthermore, the description of these syzygies is formulaic, and so this process may be easily implemented into a computer algebra system, such as Macaulay2 [15]. In particular, the methods presented here yield more efficient computation of the implicit equation of X_U , in this setting.

To illustrate this, we consider the following, which will be our running example for the case that $\dim V = 2$.

Example 1.1. Suppose that

$$U = \{s^2 u^3 + t^2 u^2 v, s^2 u v^2 + t^2 v^3, s^2 v^3, t^2 u^3\} \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(2,3))$$

and let I_U denote the ideal associated to U. A computation shows that I_U has a first syzygy in degree (0, 2). Moreover, upon constructing the subspace $V \subseteq U$, further computations show that dim V = 2 and also that I_U has seven minimal first syzygies in bidegrees

$$(0,2), (2,1), (2,1), (0,4), (2,3), (4,2), (6,1).$$

Following Theorem 4.3, the syzygies of bidegree (0,2), (2,1), and (2,1) are the columns of the matrix

$$M = \begin{bmatrix} v^2 & 0 & t^2 u \\ -u^2 & s^2 v & 0 \\ 0 & -s^2 u - t^2 v & 0 \\ 0 & 0 & -s^2 u - t^2 v \end{bmatrix}$$
(1.1)

and moreover, these syzygies determine the first differential of $\mathcal{Z}_{(2a-1,b-1)} = \mathcal{Z}_{3,2}$. Hence by Theorem 2.5, these syzygies yield the implicit equation of X_U .

We continue this example in Section 5, verifying the construction of the two syzygies in (1.1) of bidegree (2, 1) from the syzygy in bidegree (0, 2). Additionally, we show precisely how a matrix representation of the differential d_1 is obtained. Following Theorem 2.5, the determinant of the resulting 12×12 matrix is then a power of the implicit equation of X_U .

We briefly describe how this paper is organized. In Section 2, we provide the preliminary material required for the scope of this article. We review the construction of the approximation complex of [17, 18] and recall the techniques of [1], using this complex for the purpose of multigraded implicitization. In Section 3, we introduce the main setting of this article, and show that a syzygy in bidegree (0, 2) implies constraints on the generation of I_U . The subspace $V \subseteq U$ is introduced, and we consider two cases based on its dimension. In Section 4, we address the first case where dim V = 2, and produce a pair of additional syzygies that determine the implicit equation of X_U . In Section 5, we consider the case when dim V = 3 and produce a set of three additional syzygies which determine the implicit equation of X_U in this setting. We conclude the paper in Section 6 with some further observations and questions related to the results presented here.

2. Preliminaries

Here we briefly describe the preliminary material necessary for this paper. We begin by introducing the approximation complex \mathcal{Z} [17, 18] associated to the ideal I_U . Moreover, we then recall the applications of this complex to the implicitization of tensor product surfaces, as outlined in [1]. We refer the reader to [7] for a nice overview of the techniques presented here.

2.1. Approximation Complex. We recall the construction of the *approximation complex* Z, introduced by Herzog, Simis, and Vasconcelos [17, 18]. This complex may be defined more generally, but for our purposes we consider the following setting. Let

$$I = (f_0, \dots, f_n) \subseteq R = k[x_0, \dots, x_d]$$

be a homogeneous ideal of R, and consider the Koszul complex $\mathcal{K}(\underline{f})$ on the sequence $\underline{f} = f_0, \ldots, f_n$, with differentials d_i^f . Moreover, for new indeterminates $\underline{T} = T_0, \ldots, T_n$, let $S = k[T_0, \ldots, T_n]$ and consider the Koszul complex $\mathcal{K}(\underline{T})$ with differential d_i^S . We construct a hybrid complex from this data.

Definition 2.1. Writing $Z_i = \ker d_i^f$ to denote the *i*th Koszul cycle, the approximation complex \mathcal{Z} is the complex

$$\mathcal{Z} : \cdots \to \mathcal{Z}_{i+1} \xrightarrow{d_{i+1}} \mathcal{Z}_i \xrightarrow{d_i} \mathcal{Z}_{i-1} \xrightarrow{d_{i-1}} \cdots$$

where $\mathcal{Z}_i = S \otimes_k Z_i$ and $d_i = d_i^S$.

A direct computation shows that $d_i^f d_{i+1}^S + d_{i+1}^S d_i^f = 0$. Hence for any $g \in Z_i$, we see that $d_i^f d_{i+1}^S(g) = -d_{i+1}^S d_i^f(g) = 0$, and so $d_i(g) \in Z_{i-1}$ and these maps are well defined. Moreover, the fact that Z is a complex follows as its differentials are inherited from $\mathcal{K}(\underline{T})$. Much like the Koszul complex, the approximation complex Z depends only on the ideal I, and not the choice of generating set.

Whereas the higher homology of \mathcal{Z} is often obscure, we note that the zeroth homology is familiar. Notice that $Z_0 = R$ and $Z_1 = \operatorname{syz}(I)$, the module of syzygies on I. It follows that the first differential d_1 of \mathcal{Z} maps $d_1 : S \otimes_k \operatorname{syz}(I) \to S \otimes_k R$ by

$$(a_0, \dots, a_n) \longmapsto a_0 T_0 + \dots + a_n T_n.$$

$$(2.1)$$

With this, we see that $H_0(\mathcal{Z}) \cong \mathcal{S}(I)$, where $\mathcal{S}(I)$ is the symmetric algebra of I.

Remark 2.2. With the description in (2.1), one may determine the image of d_1 from a free resolution, or even a free presentation, of I. Indeed, if $R^m \xrightarrow{\varphi} R^{n+1} \to I \to 0$ is such a presentation, with syzygy matrix φ , then im $d_1 = (\ell_1, \ldots, \ell_m) \subset R[T_0, \ldots, T_n] \cong S \otimes_k R$, where $[\ell_1 \ldots \ell_m] = [T_0 \ldots T_n] \cdot \varphi$.

With this, we note that it suffices to understand the syzygy matrix φ to determine the first differential d_1 of \mathcal{Z} . We make use of this observation throughout this article.

2.2. Multigraded Implicitization. We now recall the applications of the approximation complex Z to the implicitization of tensor product surfaces. This complex has been used in multiple instances for implicitization purposes [2, 3, 7], but for conciseness, we refer to the tools developed in [1].

Notice that $R \otimes_k S$ is naturally bigraded, hence we may take graded strands of \mathcal{Z} . Indeed, for ν a fixed degree within the grading of R, the complex \mathcal{Z}_{ν} is

$$\mathcal{Z}_{\nu} : \longrightarrow S \otimes_k (Z_i)_{\nu} \xrightarrow{d_i} S \otimes_k (Z_{i-1})_{\nu} \xrightarrow{d_{i-1}} \cdots$$

Moreover, we note that if R is bigraded, as in the proceeding sections, one may also take a *bigraded* strand \mathcal{Z}_{ν} consisting of S-modules as well.

Lemma 2.3 ([1, 7.3]). Let $U = \{p_0, p_1, p_2, p_3\} \subset R_{a,b}$ and let $\phi_U : \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow \mathbb{P}^3$ denote the rational map defined by U. Assume that either ϕ_U has no basepoints or ϕ_U has finitely many basepoints that are locally a complete intersection. Let $\nu = (2a - 1, b - 1)$ (equivalently $\nu = (a - 1, 2b - 1)$) and let $\Delta_{\nu} = \det \mathcal{Z}_{\nu}$. Then

$$\deg(\Delta_{\nu}) = 2ab - \dim(H_2)_{4a-1,3b-1}.$$

Moreover, the differential $d_1: (\mathcal{Z}_1)_{\nu} \to (\mathcal{Z}_0)_{\nu}$ has a matrix representation which is square of size $2ab \times 2ab$ if and only if $(H_2)_{4a-1,3b-1} = 0$.

Here $(H_2)_{4a-1,3b-1}$ denotes the second homology module of the bigraded strand $\mathcal{Z}_{4a-1,3b-1}$. The notion of the determinant of a complex is a general formulation, but we refer the reader to [7] for a concise description aimed towards applications similar to those presented here. We will be most interested in the case that U is free of basepoints, in which case $(H_2)_{4a-1,3b-1}$ vanishes and the determinant of \mathcal{Z}_{ν} is simply the determinant of a square matrix representation of d_1 .

Lemma 2.4 ([1, 7.4]). With the conditions of Lemma 2.3, suppose that the basepoints of U (if any) have multiplicity e_x . One has

$$\deg(\phi_U)\deg(F) = 2ab - \sum e_x$$

where $F \in S = k[T_0, T_1, T_2, T_3]$ is the implicit equation of X_U .

Combining Lemmas 2.3 and 2.4, we obtain our primary tool to determine the implicit equation F of X_U .

Theorem 2.5 ([1, 7.5]). With the assumptions of Lemma 2.3, we have that $\Delta_{\nu} = F^{\deg \phi_U}$. In particular, from Lemma 2.4 we have

$$\deg \Delta_{\nu} = \deg(F^{\deg \phi_U}) = (\deg F)(\deg \phi_U) = 2ab - \sum e_x.$$

Hence by Lemma 2.3, it follows that $\dim(H_2)_{4a-1,3b-1} = \sum e_x$.

Thus in the absence of basepoints, the differential $d_1 : (\mathcal{Z}_1)_{\nu} \to (\mathcal{Z}_0)_{\nu}$ is a square $2ab \times 2ab$ matrix. Additionally, its determinant is a power of the implicit equation F of X_U , and this power is precisely deg ϕ_U .

3. QUADRATIC SYZYGIES

We now introduce the setting for the duration of the paper. Whereas many of the conventions have been stated in the introduction, we briefly restate them here for clarity. Our primary setting is the following.

Setting 3.1. Let R = k[s, t, u, v] with bideg s, t = (1, 0) and bideg u, v = (0, 1). Let $U \subseteq R_{a,b}$ be a subspace with basis $\{p_0, p_1, p_2, p_3\}$ and let $I_U = (p_0, p_1, p_2, p_3) \subseteq R$. Assume that U is basepoint free and write $\phi_U : \mathbb{P}^1 \times \mathbb{P}^1 \longrightarrow \mathbb{P}^3$ to denote the regular map defined by U, with image X_U . Assume that $b \ge 3$ and that I_U has a first syzygy Q of bidegree (0, 2). Moreover, assume that I_U has no linear first syzygy.

We may safely assume that I_U has no linear syzygy in bidegree (0, 1) or (1, 0), as this case has already been examined in [13]. Moreover, we note that by symmetry, the case that I_U has a syzygy in bidegree (2, 0)is also addressed if $a \ge 3$.

To begin our initial treatment in the setting above, we apply a technique from the proof of [19, 4.8], and introduce similar constructions. As I_U has a first syzygy of bidegree (0, 2), there exist coefficients $a_i, b_i, c_i \in k$ such that

$$\sum_{i=0}^{3} (a_i u^2 + b_i uv + c_i v^2) p_i = 0.$$
(3.1)

Rearranging, we have

$$0 = \sum_{i=0}^{3} (a_i u^2 + b_i uv + c_i v^2) p_i = (\sum_{i=0}^{3} a_i p_i) u^2 + (\sum_{i=0}^{3} b_i p_i) uv + (\sum_{i=0}^{3} c_i p_i) v^2.$$
(3.2)

Writing $f_0 = \sum_{i=0}^3 a_i p_i$, $f_1 = \sum_{i=0}^3 b_i p_i$, and $f_2 = \sum_{i=0}^3 c_i p_i$, we note that $[f_0, f_1, f_2]$ is a syzygy on $[u^2, uv, v^2]$. As a free resolution of the ideal (u^2, uv, v^2) is well known, by say the Hilbert-Burch theorem [14, 20.15], it follows that

$$\begin{bmatrix} f_0 \\ f_1 \\ f_2 \end{bmatrix} = \alpha \begin{bmatrix} v \\ -u \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 0 \\ v \\ -u \end{bmatrix} = \begin{bmatrix} \alpha v \\ \beta v - \alpha u \\ -\beta u \end{bmatrix}$$
(3.3)

for some $\alpha, \beta \in R_{a,b-1}$.

Remark 3.2. Notice that, as I_U has no linear syzygy, both $f_0 \neq 0$ and $f_2 \neq 0$. Indeed, as $\{p_0, p_1, p_2, p_3\}$ is linearly independent, if $f_0 = 0$ we see that $a_0 = a_1 = a_2 = a_3 = 0$. Hence from (3.1) we have

$$0 = \sum_{i=0}^{3} (b_i uv + c_i v^2) p_i = v \sum_{i=0}^{3} (b_i u + c_i v) p_i,$$

and so $\sum_{i=0}^{3} (b_i u + c_i v) p_i = 0$, which contradicts the assumption that I_U has no linear syzygy. A similar argument shows that $f_2 \neq 0$. In particular, following (3.3), we see that both α and β are nonzero.

We note that, whereas both f_0 and f_2 are nonzero, there is no such restriction on the vanishing of f_1 . In the case that $f_1 = 0$, (3.2) and Proposition 3.7 will show that I_U has a reduced Koszul syzygy in bidegree (0,2), which may be taken as $Q = [v^2, -u^2, 0, 0]$; see Example 1.1.

With this, we will eventually consider two cases: when the set $\{f_0, f_1, f_2\}$ is k-linearly independent and when it is not. To this end, let $V = \text{Span}_k\{f_0, f_1, f_2\}$ denote the subspace of U spanned by $\{f_0, f_1, f_2\}$.

Proposition 3.3. With V the subspace spanned by $\{f_0, f_1, f_2\}$, we have that $2 \leq \dim V \leq 3$.

Proof. The last inequality is clear, hence we need only verify the first. By Remark 3.2, we see that $V \neq 0$ and so it suffices to show that dim $V \neq 1$. Recall from Remark 3.2 that $f_0 \neq 0$, hence if dim V = 1 then $f_1 = d_1 f_0$ and $f_2 = d_2 f_0$ for some $d_1, d_2 \in k$. Thus (3.2) shows that $(u^2 + d_1 uv + d_2 v^2) f_0 = 0$, however this is a contradiction as R is a domain.

Remark 3.4. We note that the subspace V is inspired by a similar construction in [19, 4.8], within the study of tensor product surfaces of bidegree (2, 1). However, the formulation of such a subspace associated to a syzygy holds quite generally, a topic which we explore in Section 6. In particular, adapting this method to the setting of a *linear* syzygy in [13], the proof of [13, 2.1] shows the resulting subspace V has dim V = 2 in this setting. As a consequence, the statement of Theorem 4.3 reads very similarly to [13, 2.2].

With Proposition 3.3, we may consider two cases, namely when dim V = 2 and when dim V = 3. We note that this figure dictates the largest size of a subset of $\{f_0, f_1, f_2\}$ that may be taken as part of a minimal generating set of I_U . With this, we proceed as in [13] and establish particular generating sets of I_U , in each case. Additional syzygies, based upon this generation, are then determined in the proceeding sections.

Remark 3.5. We note that the dimension of V is easily computed from the data of (3.2). Indeed, write φ for the 4 × 3 coefficient matrix

$$\varphi = \begin{vmatrix} a_0 & b_0 & c_0 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$
(3.4)

and note that $[f_0, f_1, f_2] = [p_0, p_1, p_2, p_3] \cdot \varphi$. As $\{p_0, p_1, p_2, p_3\}$ is a basis of U, and hence linearly independent, it follows that dim $V = \operatorname{rank} \varphi$, the latter of which is a linear algebra computation.

Additionally, we note that dim V is precisely the largest size of a subset of $\{f_0, f_1, f_2\}$ that may be taken as part of a minimal generating set of I_U . Clearly one has $(f_0, f_1, f_2) \subseteq I_U = (p_0, p_1, p_2, p_3)$ and, as every polynomial involved has the same bidegree, the size of this subset is exactly the largest-sized nonzero, and hence invertible, minor of φ .

3.1. Case 1: dim V = 2. With the conditions of Setting 3.1, we consider the first case of Proposition 3.3 and assume that dim V = 2. With this, we aim to produce a particular generating set of I_U , which will be used to determine additional syzygies of I_U in Section 4.

With the assumption that dim V = 2, clearly the set $\{f_0, f_1, f_2\}$ is linearly dependent. Hence there is an equation of dependence

$$d_0 f_0 + d_1 f_1 + d_2 f_2 = 0 (3.5)$$

for $d_0, d_1, d_2 \in k$, with at least one nonzero. With this, we consider the (non-exclusive) subcases depending on the non-vanishing of the coefficients.

(i) Suppose that $d_2 \neq 0$. After rescaling d_2 to 1, rearranging, and relabelling in (3.5), we may write

$$f_2 = d_0 f_0 + d_1 f_1 \tag{3.6}$$

for $d_0, d_1 \in k$. With this, we see that $\{f_0, f_1\}$ is then a basis for V. We also note that, as $f_2 \neq 0$ by Remark 3.2, it follows that at least one of d_0, d_1 is nonzero.

Recall from (3.2) that $[f_0, f_1, f_2]$ is a syzygy on the ideal (u^2, uv, v^2) . With this and (3.6) we have

$$f_0u^2 + f_1uv + (d_0f_0 + d_1f_1)v^2 = 0,$$

hence

$$f_0(u^2 + d_0v^2) + f_1(uv + d_1v^2) = 0.$$
(3.7)

With this, we observe that $u^2 + d_0v^2$, $uv + d_1v^2$ is an *R*-regular sequence. Indeed, clearly the two are not unit multiples of each other, and so $ht(u^2 + d_0v^2, uv + d_1v^2) = 1$ only if these polynomials have a common linear factor in k[u, v]. However, factoring it from (3.7), it would then follow that I_U has a linear syzygy, which contradicts the assumptions of Setting 3.1.

From (3.7) it follows that $[f_0, f_1]$ is a syzygy on the ideal $(u^2 + d_0v^2, uv + d_1v^2)$. Since this ideal is generated by a regular sequence, its syzygy module is spanned by a single Koszul syzygy. Hence

$$\begin{cases} f_0 = h(uv + d_1v^2) \\ f_1 = -h(u^2 + d_0v^2) \end{cases}$$

for some $h \in R_{a,b-2}$. Lastly, we note that since $u^2 + d_0v^2$ and $uv + d_1v^2$ share no common factor, and $uv + d_1v^2 = v(u + d_1v)$, we have that $u + d_1v \nmid u^2 + d_0v^2$. With this, we see that $d_1^2 + d_0 \neq 0$, recalling that d_0 and d_1 cannot simultaneously vanish by (3.6) and Remark 3.2.

(ii) Alternatively, suppose that $d_1 \neq 0$. After rescaling d_1 to 1, rearranging, and relabelling in (3.5) accordingly, we may write

$$f_1 = d_0 f_0 + d_2 f_2 \tag{3.8}$$

for $d_0, d_2 \in k$. With this, we note that $\{f_0, f_2\}$ is then a basis for V. With this and (3.2), it follows that

$$f_0u^2 + (d_0f_0 + d_2f_2)uv + f_2v^2 = 0,$$

hence

$$f_0(u^2 + d_0uv) + f_2(v^2 + d_2uv) = 0. (3.9)$$

A similar argument as before shows that $u^2 + d_0 uv, v^2 + d_2 uv$ is a regular sequence. Noting from (3.9) that $[f_0, f_2]$ is a syzygy on this sequence, it follows that

$$\begin{cases} f_0 = h(v^2 + d_2 uv) \\ f_2 = -h(u^2 + d_0 uv) \end{cases}$$

for some $h \in R_{a,b-2}$. Moreover, since $u^2 + d_0 uv, v^2 + d_2 uv$ form a regular sequence, they have no common factor. As $v^2 + d_2 uv = v(v + d_2 u)$ and $u^2 + d_0 uv = u(u + d_0 v)$, we see that $v + d_2 u$ and $u + d_0 v$ are not unit multiples of each other, hence $d_0 d_2 - 1 \neq 0$.

Remark 3.6. We purposely omit the third case that $d_0 \neq 0$ in (3.5), as it is superfluous. Indeed, if $d_0 \neq 0$, then at least one of d_1, d_2 is nonzero as well, as $f_0 \neq 0$ by Remark 3.2. Hence this setting belongs to at least one of the cases above. We also note that instead, one could consider the two cases that $d_1 \neq 0$ and $d_2 \neq 0$. However, repeating as before, this latter case follows identically to the first case above by symmetry in the monomial sequence $\{u^2, uv, v^2\}$.

We now address the generation of I_U in the case dim V = 2. In order to make notation consistent, we adjust the indices of the coefficients in the discussion above.

Proposition 3.7. With the assumptions of Setting 3.1, if dim V = 2, then after possibly reindexing, we have $I_U = (hg_0, hg_1, p_2, p_3)$ for some $h \in R_{a,b-2}$ and $g_0, g_1 \in R_{0,2}$ where either

$$\begin{cases} g_0 = uv + d_0 v^2 \\ g_1 = u^2 + d_1 v^2 \\ with \ d_0^2 + d_1 \neq 0 \end{cases} \quad or \quad \begin{cases} g_0 = v^2 + d_0 uv \\ g_1 = u^2 + d_1 uv \\ with \ d_0 d_1 - 1 \neq 0 \end{cases}$$

for some $d_0, d_1 \in k$.

Proof. This follows from Remark 3.5 and the previous discussion, after adjusting the indices involved. \Box

Remark 3.8. Although the case that dim V = 2 splits into two subcases, one may easily determine the coefficients d_0 and d_1 , and which case of Proposition 3.7 one has. Indeed, recall that $\{p_0, p_1, p_2, p_3\}$ is linearly independent, and one has the matrix equation in Remark 3.5. Thus finding the coefficients in (3.5) corresponds to finding a basis for the kernel of the coefficient matrix φ , which is an exercise in linear algebra.

With the generating set of I_U established in Proposition 3.7, it will be much easier to describe other syzygies of I_U . In particular, in Section 4 we will produce a set of additional syzygies which will inevitably determine the complex of Lemma 2.3. First however, we consider the generation of I_U when dim V = 3.

3.2. Case 2: dim V = 3. With the conditions of Setting 3.1, we now consider the second case of Proposition 3.3 and proceed under the assumption that dim V = 3, i.e. $\{f_0, f_1, f_2\}$ is a basis of V. As $\{f_0, f_1, f_2\}$ is linearly independent, these polynomials may be taken as minimal generators of I_U , following Remark 3.5.

Proposition 3.9. With the assumptions of Setting 3.1, if dim V = 3, then after possibly reindexing, we have $I_U = (\alpha u, \beta v - \alpha u, -\beta u, p_3)$ for some $\alpha, \beta \in R_{a,b-1}$.

Proof. This follows from (3.3) and Remark 3.5.

With suitable generating sets of I_U established under the assumptions of Setting 3.1, within both cases of Proposition 3.3, we may proceed in the following sections with a deeper study of the syzygies of I_U . In particular, we determine a subset of syzygies that is sufficient to determine the bigraded strand $Z_{2a-1,b-1}$ in Lemma 2.3, and hence determine the implicit equation of X_U .

4. Syzygies in the case $\dim V = 2$

With the conditions of Setting 3.1, we proceed under the assumption that dim V = 2 for the duration of this section. Recall from Proposition 3.7 that I_U may be generated as $I_U = (hg_0, hg_1, p_2, p_3)$, for particular $g_0, g_1 \in R_{0,2}$, and some $h \in R_{a,b-2}$. By selecting this generating set, we note that the syzygy Q of bidegree (0, 2) in Setting 3.1 is the reduced Koszul syzygy

$$Q = \begin{bmatrix} g_1 \\ -g_0 \\ 0 \\ 0 \end{bmatrix}.$$
 (4.1)

With this specific choice of generating set for I_U , we aim to produce a set of additional syzygies which, with Q, will determine the complex $\mathbb{Z}_{2a-1,b-1}$ in Lemma 2.3. Before we are able to describe these syzygies, we must verify the containment of ideals $I_U \subseteq (g_0, g_1)$, which follows from the proceeding lemma.

Lemma 4.1. We have the containment of ideals $(u, v)^3 \subseteq (g_0, g_1)$.

Proof. We write each cubic monomial of k[u, v] in terms of g_0 and g_1 , in both cases of Proposition 3.7.

(i) In the first case where $g_0 = uv + d_0v^2$ and $g_1 = u^2 + d_1v^2$ with $d_0^2 + d_1 \neq 0$, we have the following equations, which are easily verified.

$$u^{3} = \left(\frac{-d_{0}d_{1}}{d_{0}^{2}+d_{1}}u - \frac{d_{1}^{2}}{d_{0}^{2}+d_{1}}v\right)g_{0} + \left(u + \frac{d_{0}d_{1}}{d_{0}^{2}+d_{1}}v\right)g_{1},$$

$$u^{2}v = \left(\frac{d_{1}}{d_{0}^{2}+d_{1}}u - \frac{d_{0}d_{1}}{d_{0}^{2}+d_{1}}v\right)g_{0} + \left(\frac{d_{0}}{d_{0}^{2}+d_{1}}v\right)g_{1},$$

$$uv^{2} = \left(\frac{d_{0}}{d_{0}^{2}+d_{1}}u + \frac{d_{1}}{d_{0}^{2}+d_{1}}v\right)g_{0} - \left(\frac{d_{0}}{d_{0}^{2}+d_{1}}v\right)g_{1},$$

$$v^{3} = \left(\frac{-1}{d_{0}^{2}+d_{1}}u + \frac{d_{0}}{d_{0}^{2}+d_{1}}v\right)g_{0} + \left(\frac{1}{d_{0}^{2}+d_{1}}v\right)g_{1}.$$
(4.2)

(ii) Similarly, if $g_0 = v^2 + d_0 uv$ and $g_1 = u^2 + d_1 uv$ with $d_0 d_1 - 1 \neq 0$, we have the following.

$$u^{3} = \left(\frac{-d_{1}^{2}}{d_{0}d_{1}-1}u\right)g_{0} + \left(u + \frac{d_{1}}{d_{0}d_{1}-1}v\right)g_{1},$$

$$u^{2}v = \left(\frac{d_{1}}{d_{0}d_{1}-1}u\right)g_{0} - \left(\frac{1}{d_{0}d_{1}-1}v\right)g_{1},$$

$$uv^{2} = \left(\frac{-1}{d_{0}d_{1}-1}u\right)g_{0} + \left(\frac{d_{0}}{d_{0}d_{1}-1}v\right)g_{1},$$

$$v^{3} = \left(\frac{d_{0}}{d_{0}d_{1}-1}u + v\right)g_{0} - \left(\frac{d_{0}^{2}}{d_{0}d_{1}-1}v\right)g_{1}.$$
(4.3)

Hence we have $(u, v)^3 \subseteq (g_0, g_1)$, in each case of Proposition 3.7.

Remark 4.2. From Lemma 4.1, we have that $I_U \subseteq (g_0, g_1)$, as I_U is generated in bidegree (a, b) with $b \ge 3$. Hence one has $V(g_0, g_1) \subseteq V(I_U)$, and we note that this is not a contradiction to the assumption that U is basepoint free. Indeed, recall that g_0, g_1 is a regular sequence in k[u, v], hence it follows that $V(g_0, g_1) = \emptyset$ in \mathbb{P}^1 . Alternatively, one may also see that $V(g_0)$ and $V(g_1)$ have no intersection from Lemma 5.1.

We now produce an additional pair of syzygies that, along with Q, determine the implicit equation of X_U . Following the discussion in Remark 3.4, we note that the statement and proof below are remarkably similar to those of [13, 2.2]. We revisit this observation in Section 6, by making a more general conjecture.

Theorem 4.3. With the conditions of Setting 3.1, assume that dim V = 2. The ideal I_U has two syzygies S_1 , S_2 of bidegree (a, b - 2) such that dim $\langle Q, S_1, S_2 \rangle_{2a-1,b-1} = 2ab$.

Proof. We proceed in a manner similar to the proof of [13, 2.2]. Recall from Proposition 3.7 that we may take $p_0 = hg_0$ and $p_1 = hg_1$. Moreover, as $(p_0, p_1, p_2, p_3) = I_U \subseteq (g_0, g_1)$ by Lemma 4.1, we may write

$$\begin{cases} p_2 = q_0 g_0 + q_1 g_1 \\ p_3 = r_0 g_0 + r_1 g_1 \end{cases}$$
(4.4)

for some $q_0, q_1, r_0, r_1 \in R_{a,b-2}$. With this, note that $q_0p_0 + q_1p_1 - hp_2 = 0$, and also $r_0p_0 + r_1p_1 - hp_3 = 0$. Hence both

$$S_1 = \begin{bmatrix} q_0 \\ q_1 \\ -h \\ 0 \end{bmatrix} \quad \text{and} \quad S_2 = \begin{bmatrix} r_0 \\ r_1 \\ 0 \\ -h \end{bmatrix}$$

are syzygies of I_U . Thus the syzygy module of I_U contains the span of the columns of

$$M = \begin{bmatrix} g_1 & q_0 & r_0 \\ -g_0 & q_1 & r_1 \\ 0 & -h & 0 \\ 0 & 0 & -h \end{bmatrix}.$$
(4.5)

Deleting the first row of M yields an upper triangular matrix, hence the columns of M span a free R-module. The claim will follow once it has been shown that $M_{2a-1,b-1}$ consists of 2ab linearly independent columns, the independence following from this previous observation.

We note that the number of columns contributed to $M_{2a-1,b-1}$ and also a matrix representation of d_1 in $\mathcal{Z}_{2a-1,b-1}$ by each syzygy agree. Moreover, this is a matter of counting monomials in certain bidegrees. Indeed, as the syzygy Q has bidegree (0, 2), it yields

$$h^{0}(\mathcal{O}_{\mathbb{P}^{1}\times\mathbb{P}^{1}}(2a-1,b-3)) = 2a(b-2)$$
(4.6)

columns of d_1 . Similarly, as both S_1 and S_2 are syzygies of bidegree (a, b-2), they each give rise to

$$h^{0}(\mathcal{O}_{\mathbb{P}^{1} \times \mathbb{P}^{1}}(a-1,1)) = 2a$$
(4.7)

columns of d_1 . Moreover, the fact that these columns are linearly independent follows as $\{Q, S_1, S_2\}$ spans a free *R*-module.

Corollary 4.4. With the assumptions of Theorem 4.3, the first differential d_1 of the bigraded strand $\mathcal{Z}_{2a-1,b-1}$ of the approximation complex \mathcal{Z} is determined by the syzygies $\{Q, S_1, S_2\}$.

Proof. This follows from Theorem 4.3, Lemma 2.3, and Theorem 2.5.

We conclude this section with an example, using the syzygies of I_U constructed in Theorem 4.3 to determine the implicit equation of X_U . We note that all of the necessary tools are in Section 2.

Example 4.5. We continue and finish Example 1.1 from the introduction, noting that the columns of (1.1) are precisely the syzygies constructed in the proof of Theorem 4.3. With this, the bidegree (2a - 1, b - 1) = (3, 2) component of (1.1) is generated by the image of

where $h = s^2 u + t^2 v$. The fact that each syzygy contributes four columns follows from (4.6) and (4.7). Multiplying by $[T_0, T_1, T_2, T_3]$ and contracting against the monomials of R in bidegree (3, 2) shows that a matrix representation of d_1 in the bigraded strand $\mathcal{Z}_{3,2}$ is the 12×12 matrix

$$d_1 = \begin{bmatrix} -T_1 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & T_1 & -T_2 & 0 & 0 & 0 & -T_3 & 0 & 0 \\ T_0 & 0 & 0 & 0 & 0 & T_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -T_1 & 0 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & T_1 & -T_2 & 0 & 0 & 0 & -T_3 \\ 0 & T_0 & 0 & 0 & 0 & 0 & 0 & T_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -T_1 & 0 & 0 & 0 & 0 & 0 & T_0 & 0 & 0 & 0 \\ 0 & 0 & -T_1 & 0 & 0 & 0 & 0 & 0 & -T_3 & T_0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & T_0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -T_2 & 0 & 0 & 0 & -T_3 & T_0 \end{bmatrix}$$

with determinant

$$(T_0^3 T_1^3 - T_0^4 T_2^2 - 2 T_0^2 T_1^2 T_2 T_3 - T_1^4 T_3^2)^2.$$

By Lemma 2.3 and Theorem 2.5, the implicit equation of X_U is $F = T_0^3 T_1^3 - T_0^4 T_2^2 - 2 T_0^2 T_1^2 T_2 T_3 - T_1^4 T_3^2$ and the degree of the rational map defined by U is deg $\phi_U = 2$.

Remark 4.6. Following the procedure above, one needs only to determine the syzygies $\{Q, S_1, S_2\}$ to determine the implicit equation of X_U . Following their construction in Theorem 4.3, recall from Remark 3.8 that the polynomials g_0 , g_1 , and h of Proposition 3.7 may be found easily from the coefficient matrix (3.4). Additionally, one may write p_2 and p_3 in terms of the monomial basis $\{u^3, u^2v, uv^2, v^3\}$, and then use the equations of Lemma 4.1 to find suitable q_0, q_1, r_0, r_1 in (4.4). In particular, this method is significantly less computationally intensive than computing the entire syzygy module of I_U to produce $\mathcal{Z}_{2a-1,b-1}$.

5. Syzygies in the case dim V = 3

We now consider the second case of Proposition 3.3 and proceed under the conditions of Setting 3.1, with the assumption that dim V = 3. As before, we aim to produce a subset of syzygies of I_U that determines the first differential of $\mathbb{Z}_{2a-1,b-1}$, and hence the implicit equation of X_U , following Theorem 2.5.

With the assumption that dim V = 3, recall from Proposition 3.9 that the ideal I_U may be generated as $I_U = (\alpha u, \beta v - \alpha u, -\beta u, p_3)$ for some $\alpha, \beta \in R_{a,b-2}$. By selecting this generating set, the syzygy Q of bidegree (0, 2) in Setting 3.1 is then

$$Q = \begin{bmatrix} u^2 \\ uv \\ v^2 \\ 0 \end{bmatrix}$$
(5.1)

following (3.2).

Similar to the approach of the previous section, we begin our treatment by examining the syzygies of the subideal $(\alpha v, \beta v - \alpha u, -\beta u)$ of I_U , generated by the basis elements of V (3.3). First however, we provide a short lemma, which is particularly useful for tensor product surfaces free of basepoints.

Lemma 5.1 ([16, V.1.4.3]). Let $f \in R_{a,b}$ and $g \in R_{c,d}$ such that gcd(f,g) = 1. The curves V(f) and V(g) in $\mathbb{P}^1 \times \mathbb{P}^1$ meet at ad + bc points.

With this, we investigate the syzygy module of the subideal $J = (\alpha v, \beta v - \alpha u, -\beta u)$ of I_U .

Proposition 5.2. The ideal $J = (\alpha v, \beta v - \alpha u, -\beta u)$ is a perfect *R*-ideal of height 2.

Proof. Since $\alpha, \beta \in R_{a,b-1}$ and $b \ge 3$, we have that $\alpha, \beta \in (u, v)^2 = (u, v^2) \cap (u^2, v)$. Hence we may write

$$\begin{cases} \alpha = q_0 u + q_1 v^2 \\ \beta = r_0 u^2 + r_1 v \\ 9 \end{cases}$$
(5.2)

for some $q_0, r_1 \in R_{a,b-2}$ and $q_1, r_0 \in R_{a,b-3}$. With this, notice that the generators of J are precisely the signed 2×2 minors of

$$\begin{bmatrix} u^2 & -q_1u + r_1 \\ uv & -r_0u - q_1v \\ v^2 & q_0 - r_0v \end{bmatrix}$$
(5.3)

and so the claim will follow from the Hilbert-Burch theorem [14, 20.15], once it has been shown that $ht J \geq 2$.

By Remark 3.2 we have that $J \neq 0$, hence it suffices to show that $ht J \neq 1$. Suppose, to the contrary, that ht J = 1. Thus f_0, f_1, f_2 have a non-unit common factor, and write h to denote the greatest common factor. Notice that $I_U \subseteq (h, p_3)$, hence h and p_3 have no common factor, as $ht I_U = 2$ since $\sqrt{I_U} = (s, t) \cap (u, v)$. However, as h is a non-unit, we may write bideg h = (c, d) for either $c \ge 1$ or $d \ge 1$ and, from the containment of ideals, we have $V(I_U) \supseteq V(h, p_3)$. Thus by Lemma 5.1 it follows that $V(I_U)$ contains ad + bc > 0 points, which contradicts the assumption that I_U is basepoint free.

Corollary 5.3. The polynomials α, β form an *R*-regular sequence.

Proof. This follows from Proposition 5.2, noting that
$$J \subseteq (\alpha, \beta)$$
.

We may now state the main result of this section. Similar to Theorem 4.3, we produce a set of additional syzygies which, with Q in (5.1), are enough to determine the implicit equation of X_U .

Theorem 5.4. With the assumptions of Setting 3.1, assume that dim V = 3. The ideal I_U has a syzygy S_1 of bidegree (a, b-2) and two syzygies S_2 , S_3 of bidegree (a, b-1) such that dim $\langle Q, S_1, S_2, S_3 \rangle_{2a-1,b-1} = 2ab$.

Proof. Recall from Proposition 3.9 that we may take $p_0 = \alpha v$, $p_1 = \beta v - \alpha u$, and $p_2 = -\beta u$. With this, we begin with the syzygy in bidegree (a, b - 2). By Proposition 5.2 and the Hilbert-Burch theorem [14, 20.15], the matrix (5.3) is precisely the syzygy matrix of $J = (\alpha v, \beta v - \alpha u, -\beta u)$. Hence, we may extend its columns to syzygies on I_U . Doing so yields Q in (5.1) and also

$$S_1 = \begin{bmatrix} -q_1 u + r_1 \\ -r_0 u - q_1 v \\ q_0 - r_0 v \\ 0 \end{bmatrix}$$

as a syzygy of I_U in bidegree (a, b - 2).

For the remaining two syzygies, we must involve the last generator of I_U . As bideg $p_3 = (a, b)$ and $b \ge 3$, we note that $p_3 \in (u^2, v^2)$. Thus we may write

$$p_3 = m_0 u^2 + m_1 v^2 \tag{5.4}$$

for some $m_0, m_1 \in R_{a,b-2}$. With this, notice that $-m_1vp_0 + m_0up_1 + m_0vp_2 + \alpha p_3 = 0$ and also $m_1up_0 + m_1vp_1 - m_0up_2 - \beta p_3 = 0$. Hence

$$S_2 = \begin{bmatrix} -m_1 v \\ m_0 u \\ m_0 v \\ \alpha \end{bmatrix} \quad \text{and} \quad S_3 = \begin{bmatrix} m_1 u \\ m_1 v \\ -m_0 u \\ -\beta \end{bmatrix}$$

are syzygies of I_U in bidegree (a, b - 2).

Consider the matrix

$$M = \begin{bmatrix} u^2 & -q_1u + r_1 & -m_1v & m_1u \\ uv & -r_0u - q_1v & m_0u & m_1v \\ v^2 & q_0 - r_0v & m_0v & -m_0u \\ 0 & 0 & \alpha & -\beta \end{bmatrix}$$
(5.5)

with columns $\{Q, S_1, S_2, S_3\}$. We first show that the columns of $M_{2a-1,b-1}$ are linearly independent, i.e. $M_{2a-1,b-1}$ is injective. We note that M itself is not injective, and we claim that its kernel is spanned by

$$N = \begin{bmatrix} m_0(q_1u - r_1) + m_1(r_0v - q_0) \\ p_3 \\ \beta \\ \alpha \end{bmatrix}$$

To verify this, consider the sequence of bigraded R-modules

A direct computation shows this is a complex. Moreover, this complex is exact by the Buchsbaum-Eisenbud acyclicity criterion [6, Cor. 1], with the required conditions following from Proposition 5.2 and Corollary 5.3. As (5.6) is a bigraded free complex, we may consider the bigraded strand in bidegree (2a - 1, b - 1). By degree considerations it then follows that $N_{2a-1,b-1} = 0$, hence $M_{2a-1,b-1}$ is indeed injective.

Now that the columns of $M_{2a-1,b-1}$ have been shown to be linearly independent, we need only count them to verify the assertion. As previously noted, this is also the number of columns contributed to a matrix representation of d_1 , by the syzygies above. The syzygy Q has bidegree (0, 2), hence it yields

$$h^{0}(\mathcal{O}_{\mathbb{P}^{1}\times\mathbb{P}^{1}}(2a-1,b-3)) = 2a(b-2)$$

columns of d_1 . Similarly, the syzygy S_1 in bidegree (a, b - 2) yields

$$h^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a-1,1)) = 2a$$

columns. Lastly, the syzygies S_2 and S_3 in bidegree (a, b - 1) each give rise to

$$h^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a-1,0)) = a$$

columns of d_1 , and the claim follows.

Corollary 5.5. With the assumptions of Theorem 5.4, the first differential d_1 of the bigraded strand $Z_{2a-1,b-1}$ of the approximation complex Z is determined by the syzygies $\{Q, S_1, S_2, S_3\}$.

Proof. This follows from Theorem 5.4, Lemma 2.3, and Theorem 2.5.

Remark 5.6. Once the syzygies $\{Q, S_1, S_2, S_3\}$ have been constructed, the procedure to determine a matrix representation of d_1 in $\mathcal{Z}_{2a-1,b-1}$, and obtain the implicit equation of X_U , is the same as the process in Example 4.5. Following Theorem 5.4, we note that the entries of these syzygies can be found easily from the decompositions (3.3), (5.2), and (5.4). As a consequence, finding the implicit equation with this procedure is computationally simple, compared to computing the full syzygy module of I_U in order to produce $\mathcal{Z}_{2a-1,b-1}$.

6. Further observations and questions

In this final section, we address some additional questions and possibilities for future directions, related to the results presented here. As the primary technique of this article is the construction of the subspace Vassociated to a given syzygy, one natural question is how to extend this method to more general settings. A further question is whether V, or rather its dimension, is always a sufficient invariant, as it was here.

The first question is easily answered. Suppose one has $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b))$ with basis $\{p_0, p_1, p_2, p_3\}$, as before. If I_U has a syzygy C with entries in R = k[s, t, u, v] of a given bidegree, say (c, d), then one may repeat the process described in Section 3. Letting $\{m_0, \ldots, m_n\}$ denote the monomials in $R_{c,d}$, one has

$$\sum_{j=0}^{3} \left(\sum_{i=0}^{n} a_{ij} m_i\right) p_j = 0$$

for some $a_{ij} \in k$. This may be rearranged as

$$0 = \sum_{j=0}^{3} \left(\sum_{i=0}^{n} a_{ij} m_i \right) p_j = \sum_{i=0}^{n} \left(\sum_{j=0}^{3} a_{ij} p_j \right) m_i = \sum_{i=0}^{n} f_i m_i,$$

by letting $f_i = \sum_{j=0}^3 a_{ij}p_j$, for $0 \le i \le n$. Then it is clear that $[f_0, \ldots, f_n]$ is a syzygy of the monomial ideal $J = (m_0, \ldots, m_n)$. Hence one proceeds by examining syz(J), noting that a free resolution (non-minimal in general) of J is readily available from Taylor's resolution [14, Ex. 17.11]. In particular, the study of syz(J) is very approachable in this setting.

Moreover, one may write V to denote the subspace $V = \text{Span}_k\{f_0, \ldots, f_n\}$ of U. An argument similar to the proof of Proposition 3.3 shows that $2 \leq \dim V \leq 4$. Moreover, similar to Remark 3.5, dim V agrees with the rank of the coefficient matrix $A = (a_{ij})$, and this is the maximum number of the f_i which may be taken as part of a minimal generating set of I_U .

Whereas the formulation of such a subspace V associated to a syzygy is quite general, we note that its dimension is rarely a sufficient invariant, as it was in Sections 4 and 5, even for syzygies of low total degree. We illustrate this next in a setting where this fails to be the case. However, we then consider a setting where it is believed that dim V is the correct invariant, in an effort to extend the work presented here and in [13].

6.1. Syzygies of bidegree (1, 1). Recall that the assumptions of Setting 3.1 may be modified to address tensor product surfaces with a syzygy in bidegree (2, 0). Thus the natural question, which has not yet been answered, is how to proceed in the remaining case of a quadratic syzygy, namely in bidegree (1, 1). We may proceed in the manner above, however we will see that the behavior of the remaining syzygies is more erratic.

If the ideal $I_U \subseteq R = k[s, t, u, v]$ has such a syzygy in bidegree (1, 1), we may write

$$\sum_{i=0}^{3} (a_i s u + b_i s v + c_i t u + d_i t v) p_i = 0.$$

for some $a_i, b_i, c_i, d_i \in k$. Rewriting this, we have

$$0 = \sum_{i=0}^{3} (a_i su + b_i sv + c_i tu + d_i tv) p_i = (\sum_{i=0}^{3} a_i p_i) su + (\sum_{i=0}^{3} b_i p_i) sv + (\sum_{i=0}^{3} c_i p_i) tu + (\sum_{i=0}^{3} d_i p_i) tv.$$
(6.1)

Writing $f_0 = \sum_{i=0}^3 a_i p_i$, $f_1 = \sum_{i=0}^3 b_i p_i$, $f_2 = \sum_{i=0}^3 c_i p_i$, and $f_3 = \sum_{i=0}^3 d_i p_i$, we see that $[f_0, f_1, f_2, f_3]$ is a syzygy on (su, sv, tu, tv). A resolution of this ideal is easily computed, from which one has

$$\begin{bmatrix} f_0 \\ f_1 \\ f_2 \\ f_3 \end{bmatrix} = \alpha \begin{bmatrix} v \\ -u \\ 0 \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 0 \\ 0 \\ v \\ -u \end{bmatrix} + \gamma \begin{bmatrix} t \\ 0 \\ -s \\ 0 \end{bmatrix} + \delta \begin{bmatrix} 0 \\ t \\ 0 \\ -s \end{bmatrix} = \begin{bmatrix} \alpha v + \gamma t \\ -\alpha u + \delta t \\ \beta v - \gamma s \\ -\beta u - \delta s \end{bmatrix}$$

for $\alpha, \beta \in R_{a,b-1}$ and $\gamma, \delta \in R_{a-1,b}$.

Writing $V = \text{Span}_k \{f_0, f_1, f_2, f_3\}$, we show that dim V alone is inadequate to dictate the behavior of the remaining syzygies of I_U in this setting. We consider the following two examples, with dim V = 4 in both, and so I_U may be generated as $I_U = (f_0, f_1, f_2, f_3)$.

Example 6.1. Consider $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(3,3))$ with basis $\{p_0, p_1, p_2, p_3\}$ where

$$p_{0} = -2t^{3}u^{2}v - s^{2}tv^{3} - t^{3}v^{3}$$

$$p_{1} = -s^{2}tu^{3} + t^{3}u^{3}$$

$$p_{2} = -s^{2}tuv^{2} + s^{3}v^{3} + st^{2}v^{3} - t^{3}v^{3}$$

$$p_{3} = s^{3}u^{3} + st^{2}u^{3} + s^{2}tu^{2}v + t^{3}uv^{2}$$

A computation through Macaulay2 [15] shows that U is basepoint free and I_U has a syzygy in bidegree (1,1). Formulating the subspace V as above, one computes the rank of the coefficient matrix and sees that dim V = 4. The ideal I_U has 10 minimal first syzygies in bidegree

(1,1), (0,5), (2,3), (2,3), (3,2), (3,2), (5,2), (5,2), (6,1), (6,1).

In particular, the only syzygies which can contribute to a matrix representation of d_1 in $\mathcal{Z}_{5,2}$ are the five in bidegrees (1, 1), (3, 2), and (5, 2). Further computations show that this is the case, and these five syzygies do determine d_1 , and hence the implicit equation, in a manner similar to the process outlined in Example 4.5.

Example 6.2. Consider $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(3,3))$ with basis $\{p_0, p_1, p_2, p_3\}$ where

$$\begin{array}{rcl} p_0 = & -t^3 u^3 - s^3 u^2 v - st^2 u^2 v - st^2 u v^2 \\ p_1 = & s^3 u^3 + st^2 u^3 - st^2 u v^2 - t^3 v^3 \\ p_2 = & st^2 u^3 - s^2 t u^2 v + s^2 t u v^2 - s^3 v^3 \\ p_3 = & s^2 t u^3 + s^3 u v^2 + s^2 t u v^2 + st^2 v^3. \end{array}$$

Similar computations show that U is basepoint free, I_U has a syzygy in bidegree (1, 1), and also dim V = 4. The ideal I_U has 10 minimal first syzygies in bidegree

(1,1), (1,9), (1,5), (2,5), (2,3), (2,3), (3,2), (3,2), (5,1), (7,1).

Moreover, the only syzygies which can contribute to d_1 in $\mathcal{Z}_{5,2}$ are the four in bidegrees (1,1), (3,2), and (5,1). As before, a further computation shows that these syzygies do determine d_1 in this example.

Remark 6.3. In each of the previous examples, we see that $U \subseteq H^0(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(3,3))$ is basepoint free, I_U has a syzygy in bidegree (1,1), and dim V = 4. However, in the examples above, the differential d_1 is derived from a different number of syzygies in differing bidegrees. Hence one cannot achieve a result analogous to Theorem 4.3 or Theorem 5.4 in this setting, based on dim V alone.

6.2. Syzygies of bidegree (0, n). From the behavior in Examples 6.1 and 6.2, it is apparent that the results presented in Sections 4 and 5 are not a consequence of the low total degree of the initial syzygy Q in bidegree (0, 2). Rather, their success is likely due to the fact that Q consists of homogeneous entries in the subring k[u, v]. In this setting, the ideal of entries of Q is much simpler, as are its syzygies. With this, we briefly discuss the case that I_U has a syzygy of bidegree (0, n).

Similar to Setting 3.1, we may assume that n is minimal, i.e. I_U has no syzygy in bidegree (0, m) for m < n. Moreover, we also assume that $U \subseteq R_{a,b}$ is basepoint free with $b \ge n + 1$. Following the approach of Section 3, we have

$$\sum_{j=0}^{3} \left(\sum_{i=0}^{n} a_{ij} u^{n-i} v^{i} \right) p_{j} = 0$$

for some $a_{ij} \in k$. Once more, we rearrange this as

$$0 = \sum_{j=0}^{3} \left(\sum_{i=0}^{n} a_{ij} u^{n-i} v^{i}\right) p_{j} = \sum_{i=0}^{n} \left(\sum_{j=0}^{3} a_{ij} p_{j}\right) u^{n-i} v^{i} = \sum_{i=0}^{n} f_{i} u^{n-i} v^{i}$$

where $f_i = \sum_{j=0}^3 a_{ij}p_j$. Thus $[f_0, f_1, \ldots, f_n]$ is a syzygy of the ideal $J = (u^n, u^{n-1}v, \ldots, v^n)$. As a resolution of this ideal is easily produced for any n, say by the Hilbert-Burch theorem [14, 20.15], one notes that syz(J) is free and spanned by the columns of the $(n + 1) \times n$ matrix

$$\begin{bmatrix} v & & & \\ -u & v & & \\ & -u & \ddots & \\ & & \ddots & v \\ & & & -u \end{bmatrix}, \text{ and so } \begin{bmatrix} f_0 \\ f_1 \\ \vdots \\ f_{n-1} \\ f_n \end{bmatrix} = \begin{bmatrix} \alpha_1 v \\ \alpha_2 v - \alpha_1 u \\ \vdots \\ \alpha_n v - \alpha_{n-1} u \\ -\alpha_n u \end{bmatrix}$$

for some $\alpha_1, \ldots, \alpha_n \in R_{a,b-1}$. Write $V = \text{Span}_k\{f_0, \ldots, f_n\}$ and note that $2 \leq \dim V \leq 4$ following the argument of Proposition 3.3. Moreover, notice that $\dim V$ agrees with the rank of the coefficient matrix $A = (a_{ij})$, as in Remark 3.5.

It is suspected, with support from experimentation through Macaulay2 [15], that dim V is the correct invariant to distinguish between the behavior of the syzygies required to determine $\mathcal{Z}_{2a-1,b-1}$ in this setting. If correct, there are three cases to consider, based on the dimension of V. However, each case is likely fraught with subcases, similar to Section 4, perhaps making the approach presented here impractical. Nevertheless, the author intends to study this setting in a future paper.

Before we conclude this article, we present a conjecture within this setting in the case that $\dim V = 2$, based on observations made here.

Conjecture 6.4. Suppose that $U \subseteq R_{a,b}$ is basepoint free with $b \ge n+1$ and I_U has a syzygy C in bidegree (0, n), and no syzygy in k[u, v] of smaller degree. If dim V = 2, then I_U has two syzygies S_1, S_2 of bidegree (a, b-n) such that the first differential d_1 of $\mathcal{Z}_{2a-1,b-1}$, and hence the implicit equation of X_U , is determined by $\{C, S_1, S_2\}$.

We note that this conjecture is true in the case n = 1 by [13, 2.2] following Remark 3.4, as well as the case that n = 2 by Theorem 4.3. If confirmed, it is curious if there are similar results for the cases dim V = 3 and dim V = 4, when I_U has a syzygy from the subring k[u, v].

Acknowledgements

The use of Macaulay2 [15] was helpful in the preparation of this article, offering numerous examples to support the results presented here, and verifying the direct computations and equations in Sections 4 and 5.

References

- [1] N. Botbol, The implicit equation of a multigraded hypersurface, J. Algebra 348 (2011), 381–401.
- [2] N. Botbol, A. Dickenstein, M. Dohm, Matrix representations for toric parametrizations, Comput. Aided Geom. Design 26 (2009), 757–771.
- [3] L. Busé, M. Chardin, Implicitizing rational hypersurfaces using approximation complexes, J. Symbolic Computation 40 (2005), 1150–1168.
- [4] L. Busé, J.-P. Jouanolou, On the closed image of a rational map and the implicitization problem, J. Algebra 265 (2003), 312–357.
- [5] W. Bruns and J. Herzog, Cohen-Macaulay rings, Cambridge Studies in Advanced Mathematics 39, Cambridge University Press, Cambridge, 1993.
- [6] D.A. Buchsbaum and D. Eisenbud, What makes a complex exact?, J. Algebra 25 (1973), 259–268.
- [7] M. Chardin, Implicitization using approximation complexes, in "Algebraic geometry and geometric modeling", Math. Vis., Springer, Berlin (2006), 23–35.
- [8] D. Cox, Curves, surfaces, and syzygies, Contemp. Math. 334 (2003), 131-150.
- [9] D. Cox, Equations of parametric curves and surfaces via syzygies, Contemp. Math. 286 (2001), 1-20.
- [10] D. Cox, The moving curve ideal and the Rees algebra, Theoret. Comput. Sci. 392 (2008), 23-36.
- [11] D. Cox, J. W. Hoffman, and H. Wang, Syzygies and the Rees algebra, J. Pure Appl. Algebra 212 (2008), 1787–1796.
- [12] D. Cox, R. Goldman, M. Zhang, On the validity of implicitization by moving quadrics for rational surfaces with no basepoints, J. Symbolic Computation 29 (2000), 419–440.
- [13] E. Duarte and H. Schenck, Tensor product surfaces and linear syzygies, Proc. Amer. Math. Soc. 144 (2016), 65-72.
- [14] D. Eisenbud, Commutative algebra: with a view toward algebraic geometry, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.
- [15] D. R. Grayson and M. E. Stillman, Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/
- [16] R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics 52, Springer-Verlag, New York-Heidelberg, 1977.
- [17] J. Herzog, A. Simis, and W. V. Vasconcelos, Approximation complexes of blowing-up rings, J. Algebra 74 (1982), 466–493.
- [18] J. Herzog, A. Simis, and W. V. Vasconcelos, Approximation complexes of blowing-up rings. II, J. Algebra 82 (1983), 53–83.
- [19] H. Schenck, A. Seceleanu, and J. Validashti, Syzygies and singularities of tensor product surfaces of bidegree (2, 1), Math. Comp. 83 (2014), 1337–1372.
- [20] T. W. Sederberg and F. Chen, Implicitization using moving curves and surfaces, Proceedings of SIGGRAPH (1995), 301–308.
- [21] T. W. Sederberg, R. N. Goldman, and H. Du, Implicitizing rational curves by the method of moving algebraic curves, J. Symb. Comput. 23 (1997), 153–175.
- [22] T. W. Sederberg, T. Saito, D. Qi, and K. S. Klimaszewksi, Curve implicitization using moving lines, Comput. Aided Geom. Des. 11 (1994), 687–706.

School of Mathematical and Statistical Sciences, Arizona State University, Wexler Hall, Tempe AZ 85281 *Email address:* matthew.j.weaver@asu.edu