

From the Schaar and Lösch-Schoblick integrals to representations of the Glaisher-Kinkelin constant

Jean-Christophe PAIN^{1,2,*}

¹CEA, DAM, DIF, F-91297 Arpajon, France

²Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
F-91680 Bruyères-le-Châtel, France

January 24, 2025

Abstract

In this article, we present two integral representations of the logarithm of the Glaisher-Kinkelin constant, relying on two different integral formulations of the so-called Binet function $\mu(x)$. The first one is attributed to Schaar (and also often referred to as “the second Binet formula”), and the second one is due to Lösch and Schoblick. It seems that the two new expressions (formulas (28) and (33) of the present article) of the Glaisher-Kinkelin constant, can not be easily deduced from know ones.

1 Introduction

The Glaisher-Kinkelin constant A [1–3] plays a major role in different fields of mathematics and theoretical physics. For instance, it appears in the Gaudin model of quantum spin chains [4]. A is also related to prime numbers, through [5]

$$\prod_{p \text{ prime}} p^{\frac{1}{p^2-1}} = \frac{A^{12}}{2\pi e^\gamma}, \quad (1)$$

where γ is the usual Euler-Mascheroni constant. The Glaisher-Kinkelin constant can be expressed as

$$A = \lim_{n \rightarrow \infty} \frac{(2\pi)^{\frac{n}{2}} n^{\frac{n^2}{2} - \frac{1}{12}} e^{-\frac{3n^2}{4} + \frac{1}{12}}}{G(n+1)}, \quad (2)$$

involving in the denominator the Barnes G -function:

$$G(n) = \prod_{k=1}^{n-2} k!. \quad (3)$$

*jean-christophe.pain@cea.fr

One has for instance

$$G(1/2) = \frac{2^{1/24} e^{1/8}}{A^{3/2} \pi^{1/4}} \quad (4)$$

or also

$$G(1/4) = \frac{1}{2^{9/16} A^{9/8} \pi^{3/16} \varpi^{3/8}} \exp\left(\frac{3}{32} - \frac{\beta(2)}{4\pi}\right), \quad (5)$$

where β is the Dirichlet beta function¹:

$$\beta(s) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s}, \quad (6)$$

and

$$\varpi = \frac{\Gamma(1/4)^2}{2\sqrt{2\pi}} \quad (7)$$

denotes the transcendental lemniscate constant (ratio of the perimeter of Bernoulli's lemniscate to its diameter).

The logarithm of the Glaisher-Kinkelin constant A can be expressed in different manners, such as [2]:

$$\int_0^{1/2} \log[\Gamma(x+1)] dx = -\frac{1}{2} - \frac{7}{24} \log 2 + \frac{1}{4} \log \pi + \frac{3}{2} \log A, \quad (8)$$

where Γ represents the usual Gamma function, or also [2, 6]:

$$\log A = \frac{1}{12} - 2 \int_0^{\infty} \frac{x \log x}{e^{2\pi x} - 1} dx. \quad (9)$$

Other (but few) integral representations are actually available in the literature (see for instance the work of Choi and Nash [7]. More recently, we obtained [8]:

$$\log A = \frac{1}{9} \log 2 + \frac{1}{24} + \frac{1}{3} \int_0^{\infty} \frac{(1 - e^{-t/2}) [t \coth(t/2) - 2]}{t^3} dt, \quad (10)$$

$$\log A = \frac{1}{3} + \frac{7}{36} \log 2 - \frac{1}{6} \log \pi + \frac{1}{12} \int_0^{\infty} \frac{e^{-t} [(8 - 3t) e^t - 8 e^{t/2} - t]}{t^2(e^t - 1)} dt, \quad (11)$$

as well as [9]:

$$\begin{aligned} \log A = & \frac{1}{3} + \frac{7}{36} \log 2 - \frac{\log \pi}{6} \\ & + \frac{2}{3} \int_0^{\infty} \left[\frac{e^{-t}}{8} - \frac{1}{(1+t)^{3/2} \log^2(1+t)} - \frac{1}{2} \frac{(\log(1+t) - 2)}{(1+t) \log^2(1+t)} \right] \frac{dt}{t}, \end{aligned} \quad (12)$$

and

$$\log A = \frac{\log 2}{36} + \frac{1}{3} \int_0^{\infty} \left[\frac{\tanh\left(\frac{t}{4}\right)}{t} - \frac{e^{-t}}{4} \right] dt. \quad (13)$$

¹ $\beta(2)$ is also the Catalan constant.

Using two different improper integral representations of the so-called Binet function [10]:

$$\mu(x) = \log [\Gamma(x + 1)] - \left(x + \frac{1}{2}\right) \log x + x - \log(\sqrt{2\pi}), \quad (14)$$

the first one due to Schaar (see Sec. 2), and the second one to Lösch and Schoblik (see Sec. 3)), we derive two integral representations of the logarithm of the Glaisher-Kinkelin constant $\log A$, which, to our knowledge, were not published elsewhere.

2 Integral representation deduced from the Schaar formula

The following important result is due to Schaar [10–12] (a proof is given in Appendix A):

$$\mu(x) = 2x \int_0^\infty \frac{\arctan(t)}{e^{2\pi xt} - 1} dt. \quad (15)$$

Making the change of variable $u = 2\pi xt$, one gets

$$\mu(x) = \frac{1}{\pi} \int_0^\infty \frac{\arctan\left(\frac{u}{2\pi x}\right)}{e^u - 1} du. \quad (16)$$

We have, integrating Eq. (14) from 0 to 1/2:

$$\int_0^{1/2} \mu(x) dx = \int_0^{1/2} \log [\Gamma(x + 1)] dx + \int_0^{1/2} \left\{ \left(x + \frac{1}{2}\right) \log x - x + \log(\sqrt{2\pi}) \right\} dx. \quad (17)$$

It is easy to prove that

$$\int_0^{1/2} \left\{ \left(x + \frac{1}{2}\right) \log x - x + \log(\sqrt{2\pi}) \right\} dx = \frac{1}{16} (-7 - 2 \log 2 + 4 \log \pi). \quad (18)$$

Using the Fubini-Tonelli theorem, one can write

$$\int_0^{1/2} \mu(x) dx = \frac{1}{\pi} \int_0^{1/2} \int_0^\infty \frac{\arctan\left(\frac{u}{2\pi x}\right)}{e^u - 1} du dx = \frac{1}{\pi} \int_0^\infty \int_0^{1/2} \frac{\arctan\left(\frac{u}{2\pi x}\right)}{e^u - 1} dx du \quad (19)$$

or

$$\int_0^{1/2} \mu(x) dx = \frac{1}{\pi} \int_0^\infty \frac{1}{e^u - 1} \int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx du. \quad (20)$$

One has to calculate the integral

$$\int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx. \quad (21)$$

We have

$$\arctan(x) + \arctan\left(\frac{1}{x}\right) = \frac{\pi}{2} \quad (22)$$

and thus

$$\int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx = \int_0^{1/2} \left[\frac{\pi}{2} - \arctan\left(\frac{2\pi x}{u}\right)\right] dx = \frac{\pi}{4} - \frac{u}{2\pi} \int_0^{\frac{\pi}{u}} \arctan(v) dv. \quad (23)$$

Integrating the last integral by parts gives

$$\int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx = \frac{\pi}{4} - \frac{u}{2\pi} (v \arctan(v)) \Big|_0^{\frac{\pi}{u}} + \frac{u}{2\pi} \int_0^{\frac{\pi}{u}} \frac{v}{v^2+1} dv \quad (24)$$

and thus

$$\int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx = \frac{\pi}{4} - \frac{1}{2} \arctan\left(\frac{\pi}{u}\right) + \frac{u}{4\pi} \int_0^{\frac{\pi}{u}} \log(1+v^2) dv. \quad (25)$$

The latter result enables us to write

$$\int_0^{1/2} \arctan\left(\frac{u}{2\pi x}\right) dx = \frac{\pi}{4} - \frac{1}{2} \arctan\left(\frac{\pi}{u}\right) + \frac{u}{4\pi} \log\left(1 + \frac{\pi^2}{u^2}\right). \quad (26)$$

We thus have

$$\begin{aligned} \int_0^{1/2} \log[\Gamma(x+1)] dx &= \frac{1}{\pi} \int_0^\infty \frac{1}{e^u-1} \left[\frac{\pi}{4} - \frac{1}{2} \arctan\left(\frac{\pi}{u}\right) + \frac{u}{4\pi} \log\left(1 + \frac{\pi^2}{u^2}\right) \right] du \\ &\quad + \frac{1}{16} (-7 - 2 \log 2 + 4 \log \pi). \end{aligned} \quad (27)$$

Inserting this expression in Eq. (8) yields

$$\log A = \frac{\log 2}{9} + \frac{1}{24} + \frac{2}{3\pi} \int_0^\infty \frac{1}{e^t-1} \left[\frac{\pi}{4} - \frac{1}{2} \arctan\left(\frac{\pi}{t}\right) + \frac{t}{4\pi} \log\left(1 + \frac{\pi^2}{t^2}\right) \right] dt, \quad (28)$$

which is the first main result of the present work.

3 Integral representation resulting from a representation due to Lösch and Schoblik

Lösch and Schoblik proposed the following integral representation of the Binet function [13,14]:

$$\mu(x) = -\frac{x}{\pi} \int_0^\infty \frac{\log(1 - e^{-2\pi t})}{t^2 + x^2} dt. \quad (29)$$

Using the Fubini theorem, one can write

$$\begin{aligned} \int_0^{1/2} \mu(x) dx &= -\frac{1}{\pi} \int_0^{1/2} \int_0^\infty \frac{x \log(1 - e^{-2\pi t})}{t^2 + x^2} dt dx \\ &= -\frac{1}{2\pi} \int_0^\infty \log(1 - e^{-2\pi t}) \int_0^{1/2} \frac{2x}{t^2 + x^2} dx dt, \end{aligned} \quad (30)$$

where

$$\int_0^{1/2} \frac{2x}{t^2 + x^2} dx = \log \left(1 + \frac{1}{4t^2} \right). \quad (31)$$

We thus have

$$\begin{aligned} \int_0^{1/2} \log [\Gamma(x + 1)] dx &= -\frac{1}{2\pi} \int_0^\infty \log(1 - e^{-2\pi t}) \log \left(1 + \frac{1}{4t^2} \right) dt \\ &\quad + \frac{1}{16}(-7 - 2 \log 2 + 4 \log \pi), \end{aligned} \quad (32)$$

which, inserted into Eq. (8) again, yields

$$\log A = \frac{1}{24} + \frac{\log 2}{9} - \frac{1}{3\pi} \int_0^\infty \log(1 - e^{-2\pi t}) \log \left(1 + \frac{1}{4t^2} \right) dt, \quad (33)$$

which is the second main result of the present work.

4 Conclusion

In this article, we proposed two integral representations of the logarithm of the Glaisher-Kinkelin constant. Both are based on a definite integral representation involving $\log [\Gamma(x + 1)]$, which can itself be expressed as an improper integral, either using a representation originally proposed by Schaar and Binet or utilizing a formula by Löscher and Schoblik. These two new expressions may prove useful in exploring new properties of the Glaisher-Kinkelin constant. Other relations should be obtained from integral representations of the derivative of the Binet function (see Appendix B).

References

- [1] H. Kinkelin, *Über eine mit der Gammafunktion verwandte Transcendente und deren Anwendung auf die Integralrechnung*, J. reine angew. Math. **57**, 122-158 (1860).
- [2] J. W. L. Glaisher, *On the product $1^1 \cdot 2^2 \cdot 3^3 \dots n^n$* , Messenger Math. **7**, 43-47 (1878).
- [3] S. R. Finch, *Mathematical Constants* (Cambridge Univ. Press, New York, 2003).
- [4] M. Gaudin, *Diagonalisation d'une classe d'hamiltoniens de spin*, J. Physique **37**, 1087-1098 (1976).
- [5] R. A. Van Gorder, *Glaisher-type products over the primes*, Int. J. Number Theory **8**, 543-550 (2012).
- [6] G. Almkvist, *Asymptotic formulas and generalized Dedekind sums*, Experim. Math. **7**, 343-359 (1998).
- [7] J. Choi and C. Nash, *Integral representations of the Kinkelin's constant A*, Math. Japon. **45**, 223-230 (1997).

- [8] J.-C. Pain, *Two integral representations for the logarithm of the Glaisher-Kinkelin constant*, 2024, <https://arxiv.org/abs/2405.05264>
- [9] J.-C. Pain, *Representations of the Glaisher-Kinkelin constant deduced from integrals due to Féaux and Kummer*, 2024, <https://arxiv.org/abs/2410.22338>
- [10] R. Campbell, *Les intégrales Eulériennes et leurs applications* (Dunod, 1966).
- [11] E. T. Whittaker and G. N. Watson, *A course in modern analysis*, 4th ed. (Cambridge, England: Cambridge University Press, 1990).
- [12] T. H. Gronwall, *The Gamma function in the integral calculus*, *Annals Math.* **20**, 35-124 (1918).
- [13] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher transcendental functions*, Vol. 1, (New York: Krieger, 1981), pp. 20-21.
- [14] F. Lössch and F. Schöblich, *Die Fakultät (Gammafunktion) und verwandte Funktionen mit besonderer Berücksichtigung ihrer Anwendungen* (Teubner, Leipzig, 1952).
- [15] D. F. Connon, *Some applications of the Stieltjes constants*, 2009, arxiv.org/pdf/0901.2083
- [16] T. M. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Mathematics (Springer-Verlag, New York-Heidelberg, 1976).
- [17] M. W. Coffey, *The Stieltjes constants, their relation to the η_j coefficients, and representation of the Hurwitz zeta function*, 2007, [arXiv:math-ph/0612086](https://arxiv.org/abs/math-ph/0612086)
- [18] B. C. Berndt, *Ramanujan's Notebooks*, Parts I-III (Springer-Verlag, 1985-1991).
- [19] Z. Sasvari, *An elementary proof of Binet's formula for the Gamma function*, *Amer. Math. Mon.* **106**, 156-158 (1999).

A Derivation of the Schaar-Binet formula

The Stieltjes constants are the coefficients of the Laurent expansion of the Riemann Zeta function $\zeta(x)$ around $x = 1$ [15]:

$$\zeta(x) = \frac{1}{x-1} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \gamma_n (x-1)^n. \quad (34)$$

In the case of the Hurwitz Zeta function $\zeta(x, z)$ [16], we have the generalized Stieltjes constants:

$$\zeta(x, z) = \sum_{n=0}^{\infty} \frac{1}{(n+z)^x} = \frac{1}{x-1} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \gamma_n(z) (x-1)^n. \quad (35)$$

Let us differentiate $\zeta(x, z)$ with respect to variable x and set $x = 0$. Using the Lerch identity

$$\log \Gamma(z) = \zeta'(0, z) + \frac{1}{2} \log(2\pi), \quad (36)$$

one gets

$$\frac{1}{2} \log(2\pi) - \log \Gamma(z) = 1 + \sum_{n=0}^{\infty} \frac{\gamma_{n+1}(z)}{n!}. \quad (37)$$

Now, using the Coffey expression of the generalized Stieltjes constants [17]:

$$\gamma_n(z) = \frac{1}{2z} \log^n z - \frac{1}{n+1} \log^{n+1} z - 2\Re \int_0^{\infty} \frac{i(z+iv) \log^n(z-iv)}{(z^2+v^2)(e^{2\pi v}-1)} dv, \quad (38)$$

and inserting expression (38) into Eq. (37) yields

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\gamma_{n+1}(z)}{n!} &= \frac{1}{2z} \sum_{n=0}^{\infty} \frac{1}{n!} \log^{n+1} z - \sum_{n=0}^{\infty} \frac{1}{n!} \frac{1}{(n+2)} \log^{n+2} z \\ &\quad - 2\Re \int_0^{\infty} \frac{i(z+iv) \sum_{n=0}^{\infty} \frac{1}{n!} \log^{n+1}(z-iv)}{(z^2+v^2)(e^{2\pi v}-1)} dv. \end{aligned} \quad (39)$$

Then, using a Taylor expansion series, one gets

$$\sum_{n=0}^{\infty} \frac{1}{n!} \log^{n+1} z = z \log z. \quad (40)$$

One has also

$$\sum_{n=0}^{\infty} \frac{y^{n+2}}{(n+2)!} = e^y - 1 - y, \quad (41)$$

or equivalently

$$\sum_{n=0}^{\infty} \frac{1}{n!} \frac{y^n}{(n+2)} = \frac{e^y(y-1)-1}{y^2}, \quad (42)$$

and thus

$$\sum_{n=0}^{\infty} \frac{1}{n!} \frac{\log^{n+2} z}{(n+2)} = z(\log z - 1) + 1. \quad (43)$$

Since

$$\sum_{n=0}^{\infty} \frac{1}{n!} \log^{n+1}(z-iv) = (z-iv) \log(z-iv), \quad (44)$$

one gets

$$\sum_{n=0}^{\infty} \frac{\gamma_{n+1}(z)}{n!} = \frac{\log z}{2} - z(\log z - 1) - 1 - 2 \int_0^{\infty} \frac{\arctan(\frac{x}{z})}{e^{2\pi x} - 1} dx, \quad (45)$$

yielding

$$\mu(x) = 2x \int_0^{\infty} \frac{\arctan(t)}{e^{2\pi xt} - 1} dt. \quad (46)$$

This expression was also obtained by Ramanujan [18] for $x > 0$.

B Expressions for the function $\mu(x)$ and its derivative

In addition to expressions (15) and (29), other formulas are known for the Binet function $\mu(x)$, such as the following one, attributed to Binet as well [19]:

$$\mu(x) = \int_0^\infty e^{-xt} \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{dt}{t}. \quad (47)$$

Some formulas are available also for the function $\nu(z)$ defined as

$$\nu(x) = \frac{d\mu(x)}{dx} + \frac{1}{2x}. \quad (48)$$

The two following ones are due to Poisson:

$$\nu(x) = \int_0^\infty e^{-xt} \left(\frac{1}{e^t - 1} - \frac{1}{t} \right) dt, \quad (49)$$

and

$$\nu(x) = \frac{1}{2x} + 2 \int_0^\infty \frac{t}{(1+t^2)(1-e^{2\pi xt})} dt. \quad (50)$$