HIGHER DIMENSIONAL TETER RINGS

TONY J. PUTHENPURAKAL

ABSTRACT. Let (A, \mathfrak{m}) be a complete Cohen-Macaulay local ring. Assume A is not Gorenstein. We say A is a Teter ring if there exists a complete Gorenstein ring (B, \mathfrak{n}) with dim $B = \dim A$ and a surjective map $B \to A$ with e(B) - e(A) = 1 (here e(A) denotes multiplicity of A). We give an intrinsic characterization of Teter rings which are domains. We say a Teter ring is a strongly Teter ring if $G(B) = \bigoplus_{i \ge 0} \mathfrak{n}^i/\mathfrak{n}^{i+1}$ is also a Gorenstein ring. We give an intrinsic characterizations of strongly Teter rings which are domains. If k is algebraically closed field of characteristic zero and R is a standard graded Cohen-Macaulay k-algebra of finite representation type (and not Gorenstein) then we show that $\widehat{R}_{\mathfrak{M}}$ is a Teter ring (here \mathfrak{M} is the maximal homogeneous ideal of R).

1. INTRODUCTION

Let (A, \mathfrak{m}) be an Artinian local ring. Note there exists an Artinian Gorenstein local ring B with a surjective map $B \to A$. Let $\ell(A)$ denote length of A. The Gorenstein co-length of A is defined as

 $g(A) = \min\{\ell(B) - \ell(A) \mid B \text{ is Gorenstein Artin local ring mapping onto } A\}.$

We note that g(A) = 0 if and only if A is Gorenstein. Observe that g(A) = 1 if and only if A is not Gorenstein and $A = B/\operatorname{soc}(B)$ for a Gorenstein Artin local ring B. W. Teter gives an intrinsic characterization for such rings in his paper [9]. In their paper [5], C. Huneke and A. Vraciu refer to these rings as Teter's rings.

We generalize the definition of Teter rings to higher dimensions. Let (A, \mathfrak{m}) be a local Cohen-Macaulay local ring of dimension d. We assume A is a quotient of a regular local ring T. Let e(A) denote the multiplicity of A. We note that there exists a Gorenstein local ring B mapping onto A with dim $B = \dim A$. Let U(A)denote the set of all Gorenstein local rings B which are quotients of T, mapping onto A with dim $B = \dim A$. Set

$$g(A) = \min\{e(B) - e(A) \mid B \in U(A) \}.$$

We note that g(A) = 0 if and only if A is a Gorenstein local ring. If g(A) = 1 then we call A to be a *Teter* ring. If $B \in U(A)$ such that e(B) - e(A) = 1 then we call B to be a *Teter Gorenstein approximation* of A. Note we are *not* asserting that B is uniquely determined by A.

In general giving an intrinsic characterization of Teter's ring seems to be difficult. However for domains we have

Date: January 24, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary 13H10; Secondary 13A30.

Key words and phrases. Gorenstein rings, Finite Cohen–Macaulay type, canonical module, associated graded rings and modules.

Theorem 1.1. Let (A, \mathfrak{m}) be a Cohen-Macaulay local domain. Assume A is a quotient of a regular local ring T. Assume A is not Gorenstein. Let ω_A denote the canonical module of A. The following assertions are equivalent:

(i) A is a Teter ring.

(ii) There exists a proper ideal J of A with $\omega_A \cong J$ and $\operatorname{codim} A/J \leq 1$.

Furthermore if A is a Teter-ring then the Teter Gorenstein approximation of A is unique (up to isomorphism).

Also if (R, \mathfrak{m}) is a local Noetherian ring then $\operatorname{codim} R = \mu(\mathfrak{m}) - \dim R$ (here $\mu(\mathfrak{m})$ denotes the minimal number of generators of \mathfrak{m}).

If (R, \mathfrak{n}) is a Noetherian local ring then $G(R) = \bigoplus_{i \ge 0} \mathfrak{n}^i/\mathfrak{n}^{i+1}$ is the associated graded ring of R. We say A is strongly Teter if A has a Teter Gorenstein approximation (B, \mathfrak{n}) with G(B) Gorenstein. Being strongly Gorenstein is a very restricted property. We show

Theorem 1.2. Let (A, \mathfrak{m}) be a Cohen-Macaulay local ring. Assume A is a quotient of a regular local ring T. If A is strongly Teter then G(A) is Cohen-Macaulay.

Theorem 1.2 can be used to give examples of rings which are Teter but not strongly Teter, see 5.7. We give an intrinsic characterization of Cohen-Macaulay local domains which are strongly Teter.

Theorem 1.3. Let (A, \mathfrak{m}) be a Cohen-Macaulay local domain. Assume A is a quotient of a regular local ring T. Assume A is not Gorenstein. The following assertions are equivalent:

- (i) A is a strongly Teter ring.
- (ii) A satisfies the following properties.
 - (a) G(A) is Cohen-Macaulay.
 - (b) There exists a proper ideal J of A with $\omega_A \cong J$ and $\operatorname{codim} A/J \leq 1$.
 - (c) Consider the filtration $\mathcal{F} = \{J_n = J \cap \mathfrak{m}^n\}_{n \ge 0}$ on J. Then $G_{\mathcal{F}}(J)$ is isomorphic to the graded canonical module of G(A) (up to a shift).

In [5], the authors stated that rings of utmost countable representation type should be close to Gorenstein rings. We prove this for homogeneous rings of finite representation type. Recall that a graded algebra over a field has finite representation type if it has up to isomorphisms and shifts in the grading, only a finite number of indecomposable maximal Cohen-Macaulay modules.

Theorem 1.4. Let k be an algebraically closed field of characteristic zero. Let R be a standard Cohen-Macaulay k-algebra with maximal homogeneous ideal \mathfrak{M} . Assume R is NOT Gorenstein. Further assume R is of finite Cohen-Macaulay type. Let $A = \widehat{R}_{\mathfrak{M}}$ be its completion with respect to \mathfrak{M} . Then A is a Teter ring.

We now describe in brief the contents of this paper. In section two we prove Theorem 1.1. In section three we give proof of Theorem 1.2. In section four we give a proof of Theorem 1.3. Finally in section five we give examples and prove Theorem 1.4.

2. Proof of Theorem 1.1

In this section we give a proof of Theorem 1.1. We first prove


Proof of $(i) \implies (ii)$. Let dim A = d. As A is a Teter ring there exists a d-dimensional Gorenstein local ring B mapping onto A with e(B) - e(A) = 1. We have an exact sequence

$$0 \to M \xrightarrow{g} B \xrightarrow{f} A \to 0$$
, where $e(M) = 1$.

We note that M is a MCM B-module with e(M) = 1. Dualizing the above exact sequence we obtain an exact sequence

$$0 \to \omega_A \xrightarrow{f^{\vee}} B \xrightarrow{g^{\vee}} M^{\vee} \to 0.$$

We note that M^{\vee} is a Cohen-Macaulay ring of dimension d. As $e(M^{\vee}) = e(M) = 1$, it follows that M^{\vee} is a regular local ring. Denote it by Q. Consider the map $\omega_A \xrightarrow{f \circ f^{\vee}} A$ and let K be the kernel of this map. Set $V = \omega_A/K$ and let $i' \colon V \to A$ be the induced injection. We have the following commutative diagram:



We first claim that either N or Q'' is zero. Suppose if possible both of them are non-zero. We note that as associate primes of ω_A have dimension d it follows that dim N = d. Similarly dim Q'' = d. It follows that the modified multiplicity e(N), e(Q'') are positive. As dimensions of N, M, Q'' is d we get by [1, 4.7.7]; e(N) + e(Q'') = e(M) = 1. This is a contradiction

Suppose if possible Q'' = 0. Then Q' = Q. We note that $Q = M^{\vee} \neq 0$. We note that Q is a *d*-dimensional quotient of A which is not A (as A is NOT regular). This is a contradiction as A is a domain. So N = 0. Thus $V = \omega_A$. We note that M = Q' is a MCM Q-module with multiplicity one. So $Q'' \cong Q$. Thus Q' = Q/(a) for some a. If Q' = 0 then note that $\omega_A \cong A$. This forces A to be Gorenstein, a contradiction. So Q' is non-zero and note that codim $Q' \leq 1$. The result follows.

We have to prove a slightly more general version of the assertion (ii) \implies (i) of Theorem 1.1.

Theorem 2.1. Let (A, \mathfrak{m}) be a Cohen-Macaulay local ring of dimension $d \geq 1$. Assume A is not Gorenstein. Assume A is a quotient of a regular local ring T. Suppose the canonical module ω_A of A is isomorphic to a proper ideal J of A such that codim $A/J \leq 1$. Then A is a Teter ring.

2.2. Before proving the above result we make the following remarks:

TONY J. PUTHENPURAKAL

- (1) The ring A/J is a hypersurface ring (or regular local). We note that it suffices to show A/J is Cohen-Macaulay. To prove this note that depth $A/J \ge d-1$. As $e(J) = e(\omega) = e(A)$, it follows from [1, 4.7.7] that dim $A/J \le d-1$. The assertion follows.
- (2) Note that $\omega_A \cong J$ has a rank as A/J is a torsion A-module. It follows that A is generically Gorenstein, see [1, 3.3.18].
- (3) A/J = Q/(a) for some regular local ring Q of dimension d. This follows from the fact that A/J is a local ring which has utmost hypersurface singularity is a quotient of a regular local ring T.

We now give

4

Proof of Theorem 2.1. By 2.2 there exists a complete regular local ring Q of dimension d such that A/J = Q/(a) for some non-zero a. Let B be the fiber-product of $A \to Q/(a) = A/J$ and $Q \to Q/(a)$. It is routine to verify that the induced homomorphisms $p: B \to Q$ and $q: B \to A$ are surjective. Also notice B is local. We have the following commutative diagram

$$0 \longrightarrow \omega_A \xrightarrow{i} B \xrightarrow{p} Q \longrightarrow 0$$
$$\downarrow^f \qquad \downarrow^q \qquad \downarrow^{\pi'} Q \xrightarrow{\pi'} 0$$
$$0 \longrightarrow \omega_A \xrightarrow{i'} A \xrightarrow{\pi} Q/(a) \longrightarrow 0$$

We note that f is an isomorphism, [6, Chapter III, 5.4]. As A, Q are quotient of T, so B is also a quotient of T. As we have an exact sequence $0 \to \omega_A \to B \xrightarrow{p} Q \to 0$ is exact it follows that B is Cohen-Macaulay of dimension d. We may by using prime avoidance construct a sequence $\mathbf{x} = x_1, \ldots, x_d$ which is an $B \oplus \omega_A \oplus Q \oplus A$ regular sequence, such that images of \mathbf{x} generate the maximal ideal of Q. Going mod \mathbf{x} we obtain an exact sequence

$$0 \to \omega_A / \mathbf{x} \omega A \to B / \mathbf{x} B \xrightarrow{\overline{p}} k \to 0.$$


If $t \in \operatorname{soc}(B/\mathbf{x}B)$ then $\overline{p(t)} = 0$. It follows that $\operatorname{soc}(B/\mathbf{x}B) \cong \operatorname{soc}(\omega_A/\mathbf{x}\omega_A) \cong k$ as $\omega_A/\mathbf{x}\omega_A$ is the canonical module of $A/\mathbf{x}A$. Thus $B/\mathbf{x}B$ is a Gorenstein ring and hence B is a Gorenstein ring. We note that $\ker(B \xrightarrow{q} A) \cong \ker(Q \to Q/(a)) = Q$. This we have an exact sequence $0 \to M \to B \to A \to 0$ where $M \cong Q$. So e(M) = 1. Thus e(B) = e(A) + 1. It follows that A is a Teter ring. \Box

We now give

Proof of Theorem 1.1 continued. We already have proved (i) \implies (ii). The assertion (ii) \implies (i) follows from Theorem 2.1. We now have to prove uniqueness of the Teter Gorenstein approximation of A. Set $C = A \times_{Q/(a)} Q$ the fiber-product and let $j: C \to A$ and $l: C \to Q$ be the natural maps. By the proof of (ii) \implies (i) we get that C is a Teter Gorenstein approximation of A. We prove $B \cong C$. By our proof of (i) \implies (ii) we obtain a commutative diagram

$$0 \longrightarrow \omega_A \xrightarrow{i} B \xrightarrow{p} Q \longrightarrow 0$$
$$\downarrow^g \qquad \qquad \downarrow^q \qquad \qquad \downarrow^{\pi'} 0 \longrightarrow \omega_A \xrightarrow{i'} A \xrightarrow{\pi} Q/(a) \longrightarrow 0$$

where g is an isomorphism and p, q are surjective. By the universal property of fiber products we have a ring map $\psi: B \to C$ such that $p = l \circ \psi$ and $q = j \circ \psi$. So we have a commutative diagram



We note that $g = f \circ u$ where $f : \omega_A \to \omega_A$ is the isomorphism induced by the fiber-product diagram. As g is an isomorphism it follows that u is an isomorphism. So ψ is an isomorphism. The result follows.

3. Proof of Theorem 1.2

In this section we give

Proof of Theorem 1.2. Let (B, \mathfrak{n}) be a Teter Gorenstein approximation of A. So we have an exact sequence $0 \to M \to B \to A \to 1$ where M is a MCM B-module of multiplicity one. We note that we may assume that the residue field of B is infinite (otherwise we may consider $B' = B[X]_{\mathfrak{n}B[X]}$ and $A' = A \otimes_B B'$).

We induct on dimension d of A. When d = 0 we have nothing to prove.

The case when d = 1 is the most important case. Suppose if possible G(A) is not Cohen-Macaulay.

Consider the filtration $\mathcal{F} = \{\mathfrak{n}^n \cap M\}_{n \geq 0}$ on M. We note that \mathcal{F} is an \mathfrak{n} -stable filtration on M. We have an exact sequence of G(B)-modules

$$0 \to G_{\mathcal{F}}(M) \to G(B) \to G(A) \to 0.$$

It follows that $G_{\mathcal{F}}(M)$ is Cohen-Macaulay. Let x be $B \oplus A$ -superficial and Msuperficial with respect to \mathcal{F} . We note that $e(M) = \ell(M/xM) = 1$. So $xM = \mathfrak{m}M$ and $\mu(M) = 1$. We shift \mathcal{F} to get a new filtration $\mathcal{G} = \{\mathcal{G}_n\}_{n\geq 0}$ of M such that $\mathcal{G}_0 = M$ and $\mathcal{G}_1 \neq 1$. We note that $G_{\mathcal{G}}(M)$ is Cohen-Macaulay, say its h-polynomial is $a_0 + a_1 z + \cdots + a_s z^s$. Then as $G_{\mathcal{G}}(M)$ is Cohen-Macaulay we get $a_i \geq 0$. Also $a_0 \neq 0$ by construction. As $e(G_{\mathcal{G}}(M)) = 1$ it follows that $a_0 = 1$ and $a_i = 0$ for i > 0. Thus $e_1(G_{\mathcal{G}}(M)) = 0$. It follows that $G_{\mathcal{G}}(M) \cong k[Y]$. It follows that $G_{\mathcal{F}}(M) = k[Y](-a)$. Let α be the generator of M. Say image of α in B is ξ . By going mod a $B \oplus A$ and M-superficial element with respect to \mathcal{F} we obtain an exact sequence $0 \to k \to \overline{B} \to \overline{A} \to 0$. It follows that $\overline{\alpha}$ maps to δ , a generator of soc (\overline{B}) . Say $\operatorname{soc}(\overline{B}) = \mathfrak{n}^r$. Thus $\xi = \delta + \theta x$ where $\theta \in B$. So $\operatorname{ord}_n(\xi) \leq r$. So $a \leq r$.

If J is a non-zero homogeneous ideal of G(B) set

 $i(J) = \min\{\deg t \mid t \in J \text{ where } t \text{ is non-zero and homogeneous } \}.$

We note that if $J \subseteq J'$ then $i(J') \leq i(J)$. We note that $i(G_{\mathcal{F}}(M)) = a$.

Consider a primary decomposition of $G_{\mathcal{F}}(M)$ as an ideal in G(B). As G(A) is not Cohen-Macaulay we have

$$G_{\mathcal{F}}(M) = Q_1 \cap Q_2 \cap \dots \cap Q_s \cap L,$$

where L is \mathfrak{M} -primary (here \mathfrak{M} is the unique homogeneous ideal of G(B)) and Q_i is P_i -primary where P_i is some minimal prime of G(B). Set $Q = Q_1 \cap Q_2 \cap \cdots \cap Q_s$.

Then Q is a Cohen-Macaulay ideal in G(B) and $Q \supseteq G_{\mathcal{F}}(M)$. We have an exact sequence

$$0 \to G(B)/G_{\mathcal{F}}(M) \to G(B)/Q \oplus G(B)/L \to G(B)/(L+Q) \to 0.$$

As L is a zero-dimensional ideal of G(B) we obtain e(G(B)/Q) = e(G(A)) = e(B) - 1. It follows that e(Q) (as a G(B)-module) is one. We have an exact sequence of MCM G(B)-modules $0 \to Q \to G(B) \to G(B)/Q \to 0$. Going mod a regular element x^* of degree one we obtain a sequence $0 \to Q/x^*Q \to G(\overline{B}) \to G(B)/(Q + x^*G(B)) \to 0$. As e(Q) = 1 it follows that $Q/x^*Q = k[-c]$ for some c. We note that the degree of socle of $G(\overline{B})$ is also $\operatorname{ord}_{\overline{B}}(\operatorname{soc}(\overline{B})) = r$. So c = r. Thus Q = k[Y](-r). Thus i(Q) = r. As $Q \supseteq G_{\mathcal{F}}(M)$ we obtain $r = i(Q) \le i(G_{\mathcal{G}}(M)) = a \le r$. So $i(G_{\mathcal{F}}(M)) = r$. Thus $G_{\mathcal{F}}(M) = k[Y](-r)$.

We now consider the exact sequence $0 \to G_{\mathcal{F}}(M) \to G(B) \to G(A) \to 0$. Going mod x^* where x is $B \oplus A$ and M-superficial with respect to \mathcal{F} we obtain a sequence

$$0 \to (0 \colon {}_{G(A)}x^*) \to k[-r] \xrightarrow{f} G(\overline{B}) \to G(A)/x^*G(A) \to 0.$$

We note that $f([\alpha]) = [\delta] \neq 0$. Thus f is injective and hence $(0: _{G(A)}x^*) = 0$. So G(A) is Cohen-Macaulay.

Now assume dim $A = d \geq 2$. Let $0 \to M \to B \to A \to 0$. Let $\mathbf{x} = x_1, \ldots, x_{d-1}$ be a $A \oplus B \oplus M$ -superficial sequence. So we have an exact sequence $0 \to \overline{M} \to \overline{B} \to \overline{A} \to 0$. We note $e(\overline{M}) = 1$ and $G(\overline{B})$ is Gorenstein. So by above argument $G(\overline{A})$ is Cohen-Macaulay. By Sally descent, see [4, 2.2], G(A) is Cohen-Macaulay. \Box

4. Proof of Theorem 1.3


In this sections we give

Proof of Theorem 1.3. We may assume that the residue field of T is infinite. Otherwise consider the extension $T \to T[X]_{\mathfrak{n}T[X]}$. The hypotheses and conclusion of the theorem are invariant under this transformation.

Let $d = \dim A \ge 1$. As A is a Teter and a domain, by Theorem 1.1 we have there exists a proper ideal J of A with $J \cong \omega_A$ and $\operatorname{codim} A/J \le 1$. Furthermore A/J = Q/(a) where Q is a regular local ring of dimension d and $a \ne 0$. By Theorem 1.1 we get $B = A \times_{Q/(a)} Q$ is the unique (up to isomorphism) Teter Gorenstein approximation of A.

By 1.2 we may assume that G(A) is Cohen-Macaulay. We note that by the exact sequence $0 \to J \to A \to Q/(a) \to 0$ we obtain an exact sequence $0 \to G_{\mathcal{F}}(J) \to G(A) \to G(Q)/(a^*) \to 0$. So $G_{\mathcal{F}}(J)$ is Cohen-Macaulay.

Suppose $d \geq 2$. Then we choose $\mathbf{x} = x_1, \ldots, x_{d-1}$ which is $Q \oplus B \oplus A$ and ω_A superficial with respect to \mathcal{F} . Set $\overline{(-)} = (-) \otimes_T T/(\mathbf{x})$. Then note G(B) is Gorenstein if and only if $G(\overline{B})$ is Gorenstein. Furthermore $G_{\mathcal{F}}(\omega)$ is isomorphic to the canonical module of G(A) (up to shift) if and only if $G_{\overline{\mathcal{F}}}(\overline{\omega})$ is isomorphic to the canonical module of $G(\overline{A})$ (up to shift). We also get $\overline{B} \cong \overline{A} \times_{Q/(a,\mathbf{x})} \overline{Q}$. However note \overline{A} need not be a domain. But this is not pertinent to our result (the only requirement is that \overline{B} is an appropriate fiber-product). Thus we may assume that d = 1. Consider the fiber-product exact diagram




We note g is an isomorphism.

Let \mathfrak{n} be the maximal ideal of B. As Q is a DVR we have $\mathfrak{n}Q = (x)$. We note j' is multiplication by x^r for some $r \ge 1$. On Q (on the left hand side) we have two filtration's. First set $\mathcal{H} = \{j^{-1}(\mathfrak{n}^n)\}_{n\ge 0}$ and $\mathcal{H}' = \{(j')^{-1}((x^n))\}_{n\ge 0}$. Furthermore we have a natural injective map $\mathcal{H}_n \to \mathcal{H}'_n$ for all $n \ge 0$. It is evident that $\mathcal{H}' = \{(x)^{n-r}\}_{n\ge 0}$ (here $(x)^l = Q$ if $l \le 0$). Note that $\mathfrak{n} = \mathfrak{m} \times_{(x)/(x^r)} (x)$. Let $n \ge r$. We note that $(0, x^n) \in \mathfrak{n}^n$. So by a diagram chase we obtain $\mathcal{H}_n \supseteq (x^{n-r})$ As $\mathcal{H}_n \subseteq \mathcal{H}'_n = (x^{n-r})$. It follows that $\mathcal{H}_n = (x^{n-r})$ for all $n \ge r$. For $0 \le n \le r-1$ we note that $\mathcal{H}_n \supseteq \mathcal{H}_r = Q$. So $\mathcal{H}_n = Q$ for $n \le r$. It follows that $\mathcal{H} \cong \mathcal{H}'$.

We note that $G_{\mathcal{H}}(Q) = k[X](-r)$. We also have an exact sequence $0 \to G_{\mathcal{H}}(Q) \to G(B) \to G(A) \to 0$. As G(A) is Cohen-Macaulay we obtain that G(B) is Cohen-Macaulay.

We note that we have an exact sequence $0 \to \omega_A \to B \xrightarrow{p} Q \to 0$. Set $\mathcal{F}' = \{\omega_A \cap \mathfrak{n}^n\}_{n \geq 0}$. We also have an exact sequence $0 \to J \to A \to Q/(a) \to 0$, with $J \cong \omega_A$ and the filtration $\mathcal{F} = \{\mathfrak{m}^n \cap J\}_{n \geq 0}$. We had that $G_{\mathcal{F}}(J)$ is Cohen-Macaulay. We have a commutative diagram



We had earlier shown that $\mathcal{H}' \cong \mathcal{H}$. So G(g) is an isomorphism. It follows by Snake Lemma that ψ is an isomorphism. Thus we have an exact sequence $0 \to G_{\mathcal{F}}(J) \to G(B) \to G(Q) \to 1$. By going x^* which is G(B)-regular and of degree one we obtain an exact sequence

$$0 \to G_{\mathcal{F}}(J)/x^*G_{\mathcal{F}}(J) \to G(\overline{B}) \to k \to 0.$$

It follows that $\operatorname{soc}(G_{\mathcal{F}}(J)/x^*G_{\mathcal{F}}(J)) = \operatorname{soc}(G(\overline{B}))$. We note that G(B) is Gorenstein if and only if $\operatorname{soc} G(\overline{B}) = k$. We also have $G_{\mathcal{F}}(J)$ isomorphic to the canonical module of G(A) up to shift if and only if $\operatorname{soc}(G_{\mathcal{F}}(J)/x^*G_{\mathcal{F}}(J)) = k$. The result follows

5. Examples and proof of Theorem 1.4

In this section we give several examples which illustrate our results. We also give a proof of Theorem 1.4. Eisenbud and Herzog gave a complete list of homogeneous Cohen-Macaulay rings over an algebraically closed field k of characteristic zero which are of finite representation type. Our proof of Theorem 1.4 is by a case by case analysis of the examples of Eisenbud and Herzog.

We first show

Proposition 5.1. Let $R = \bigoplus_{n\geq 0} R_n$ be a standard algebra over an infinite field $k = R_0$. We assume dim R = 1 and that R is generically Gorenstein. Assume R is Cohen-Macaulay with minimal multiplicity and that R is not Gorenstein. Let \mathfrak{M} be the maximal homogeneous ideal of R. Then $A = \widehat{R}_{\mathfrak{M}}$ is a strong Teter ring.

Proof. We note that $h_R(z) = 1 + hz$. We also have that R is a level ring. So all generators of ω have the same degree. We have a homogeneous inclusion $J = \omega_R(r) \to R$ with J a proper ideal. Choose $x \in R_1$ which is R-regular. We note that $R_n = xR_{n-1}$ for $n \ge 2$. If all generators of J are of degree ≥ 2 then J = xJ'for some ideal J' of R. We note that $J' \cong J[+1] \cong \omega_R(r+1)$. Thus we can assume that there exists an ideal K generated in degree one with $K \cong \omega_R(s)$. We have $h_K(z) = hz + z^2$. So $h_{R/K}(z) = 1 + z$. Also dim R/K = 0. It follows that R/K is a graded hypersurface ring. After completing we obtain that A is a Teter ring by Theorem 2.1.

We note that G(A) = A and $G(\overline{R}/\overline{K}) = R/K$. Thus the exact sequence $0 \to \omega_A \to A \to \widehat{R/K} \to 0$ induces an exact sequence $0 \to G_{\mathcal{F}}(\omega_A) \to R \to R/K \to 0$. So $G_{\mathcal{F}}(\omega_A) \cong K$ is isomorphic to the canonical module of G(A) = R up to shift. It follows from 1.3 that A is a strong Teter ring.

Example 5.2. Let R = k[X, Y, Z]/(XY, XZ, YZ). Then R is of minimal multiplicity. Also it is not difficult to prove that R is reduced. So by Theorem 5.1 we get that $A = \widehat{R_{(X,Y,Z)}} = k[[X, Y, Z]]/(XY, XZ, YZ)$ is a strongly Teter ring.

Example 5.3. Let k be a field. Set S = k[X, Y] and let $R = S^{\langle m \rangle}$ for some $m \geq 3$. Then $A = \hat{R}$ is strongly Teter.

Remark 5.4. We note that $k[X,Y]^{<2>} = k[X^2, XY, Y^2]$ is a hypersurface ring and so Gorenstein.

We now give

Proof of Example 5.3. We note that $\omega_S = XYS \subseteq S$. By [3, 3.1.3] we get that $\omega_R = \omega_S^{<m>}$. We note that $\omega_R \subseteq R$. We note that $\dim_k R_1 = m+1$. Also note that $(\omega_R)_1 = (\omega_S)_m = (X,Y)^{(m-2)}XY$. So $\dim_k(\omega_R)_1 = m-1$. Thus the maximal ideal of R/ω_R is generated by two elements and it is a Cohen-Macaulay ring of dimension one. So R/ω_R is a hypersurface ring. After completing A is a Teter ring by Theorem 1.1. By an argument similar to proof of Proposition 5.1 we get that A is strongly Teter.

5.5. Let H be an additive subsemigroup of $\mathbb{N} = \{0, 1, 2, \dots\}$ with $0 \in H$ such that $\mathbb{N} \setminus H$ is finite. Let R = K[H] be the numerical semigroup ring of H. Set $A = \hat{R} = k[[H]]$. We give a few examples of semigroup rings whose completions are

Teter. The crucial observation is that

$$\omega_R = \sum_{\alpha \in \mathbb{N} \setminus H} Rt^{-\alpha}, \quad \text{see } [3, 2.1.9].$$

Example 5.6. Consider $R = k[t^3, t^4, t^5]$. Then $A = \hat{R} = k[[t^3, t^4, t^5]]$ is a Teter ring.

Proof: We note that $H = \{0, n, \text{where } n \geq 3\}$. So $\omega_R = Rt^{-1} + Rt^{-2}$. Explicitly

$$\omega_R = kt^{-2} + kt^{-1} + \sum_{n \ge 1} kt^n.$$

We note that $J = t^6 \omega_R \subseteq R$ and note that $t^4, t^5 \in J$. Thus the maximal ideal of R/J is generated by one element and dim R/J = 0. Hence it is a hypersurface ring. So A is a Teter ring by Theorem 1.1.

Example 5.7. Consider $R = k[t^4, t^5, t^{11}]$. Then $A = \hat{R} = k[[t^4, t^5, t^{11}]]$ is a Teter ring. However A is *not* strongly Teter.

Proof: We note that $H = \{0, 4, 5, n, \text{where } n \ge 8\}$. So

$$\omega_R = Rt^{-1} + Rt^{-2} + Rt^{-3} + Rt^{-6} + Rt^{-7}.$$

Explicitly

$$\omega_R = kt^{-7} + kt^{-6} + kt^{-3} + kt^{-2} + kt^{-1} + \sum_{n \ge 1} kt^n.$$

We note that $J = t^{11}\omega_R \subseteq R$ and note that $t^4, t^5 \in J$. Thus the maximal ideal of R/J is generated by one element and dim R/J = 0. Hence it is a hypersurface ring. So A is a Teter ring by Theorem 1.1. By [7, p. 20], G(A) is NOT Cohen-Macaulay. So by 1.2 we get that A is NOT strongly Teter.

5.8. Integer matrix rings: Let $A = {}^{t}(a_1, \ldots, a_n)$ be an integer matrix and assume that the g.c.d of $\{a_1, \ldots, a_n\}$ is 1. Define a grading of $S = k[x_1, \ldots, x_n]$ by defining deg $x_i = a_i$. We have $S = \bigoplus_{n \in \mathbb{Z}} S_n$. Set $S_0 = R(A)$. We have that each S_n is a R(A)-module. Let H is the sub-semigroup of \mathbb{N}^n consisting of $\alpha = (\alpha_1, \ldots, \alpha_n)$ with $\sum_{i=1}^n a_i \alpha_i = 0$ then R(A) is the semigroup ring $k[x^{\alpha} \mid \alpha \in H]$ where x^{α} denotes $x^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}$. Note R(A) is naturally an \mathbb{N} -graded ring. Set $\widehat{R(A)} = k[[x^{\alpha} \mid \alpha \in H]]$. We note that R(A) and so $\widehat{R(A)}$ are Cohen-Macaulay rings, see [10, 16.1].

Example 5.9. Consider the integer matrix ring $A = {}^{t}(2, -1, -1, -1)$. Then $\widehat{R}(A)$ is a Teter ring.

Proof. By [10, 16.10] we have

$$R(A) = k[x_1x_2^2, x_1x_2x_3, x_1x_3^2, x_1x_3x_4, x_1x_4^2, x_1x_2x_4].$$

We note that dim R(A) = 3. We also have $S_1 = (x_1x_2, x_1x_3, x_1x_4)R(A)$ is isomorphic to the canonical module of R(A) (up-to shift). We have $x_2S_1 \subseteq R(A)$ and note that

$$x_2S_1 = (x_1x_2^2, x_1x_2x_3, x_1x_2x_4).$$

So the graded maximal ideal of $R(A)/x_2S_1$ is generated by 3 elements. As $\dim R(A)/x_2S_1$ is two and it is Cohen-Macaulay it follows that $R(A)/x_2S_1$ is a hypersurface ring. Taking completions it follows that $\widehat{R(A)}$ is a Teter ring by Theorem 1.1.

Example 5.10. Consider the integer matrix ring $C = {}^{t}(2, 1, -1, -1)$. Then $\widehat{R}(\widehat{C})$ is a Teter ring.

Proof. By [10, 16.12] we have

 $R(C) = k[x_1x_3^2, x_1x_3x_4, x_1x_4^2, x_2x_3, x_2x_4].$

We note that dim R(C) = 3. We also have $S_{-1} = (x_3, x_4)R(C)$ is isomorphic to the canonical module of R(A) (up-to shift). We have $x_1x_3S_{-1} \subseteq R(A)$ and note that

$$x_1 x_3 S_{-1} = (x_1 x_3^2, x_1 x_3 x_4).$$

So the graded maximal ideal of $D = R(A)/x_1x_3S_{-1}$ is generated by 3 elements. As dim D is two and it is Cohen-Macaulay it follows that D is a hypersurface ring. Taking completions it follows that $\widehat{R(C)}$ is a Teter ring by Theorem 1.1.

5.11. Let k be an algebraically closed field of characteristic zero. Let R be a standard graded Cohen-Macaulay k-algebra of dimension d. Assume R is *not* Gorenstein and is of finite representation type. Then by [2]; R is one of the following rings :

d = 1. Then R is isomorphic to k[X, Y, Z]/(XY, XZ, YZ).

d = 2. Then R is isomorphic to $k[X, Y]^{<m>}$ for some $m \ge 3$.

d = 3. Then R is isomorphic to either $U = k[X, Y, Z]^{<2>}$ or R is isomorphic to $V = k[X_1, X_2, X_3, X_4, X_5]/(X_1^2 - X_2X_3, X_2X_5 - X_1X_4, X_1X_5 - X_3X_4)$. We now give

Proof of Theorem 1.4. We prove case by case. Set $d = \dim R$. When d = 1 we note that R is isomorphic to k[X, Y, Z]/(XY, XZ, YZ). So by 5.2 we get $\widehat{R_{\mathfrak{M}}}$ is a Teter ring. When d = 2 we have R is isomorphic to $k[X, Y]^{\leq m \geq}$ for some $m \geq 3$. So by 5.3 we get that $\widehat{R_{\mathfrak{M}}}$ is a Teter ring. When d = 3 we note that R(t(2, -1, -1, -1)) is isomorphic to the completion of U, see [10, 16.10]. So by 5.9 we get that $\widehat{U_{\mathfrak{M}}}$ is Teter. We also have R(t(2, 1, -1, -1)) is isomorphic to the completion of V, see [10, 16.12]. So by 5.10 we get that $\widehat{V_{\mathfrak{M}}}$ is Teter. The result follows.

Remark 5.12. Let k be a field of characteristic zero. Consider $A = k[[t^3, t^4, t^5]]$. In 5.6 we proved that A is Teter. Note A bi-rationally dominates $k[[t^3, t^4]]$ (which is E_6). So A has finite representation type. Thus there exists examples of completions of non-homogeneous rings of finite representation type which are Teter.

Remark 5.13. We note that all the examples in 5.11 have minimal multiplicity. Consider $A = k[[t^4, t^5, t^{11}]]$ which is a Teter ring, see 5.7. We note A does not have minimal multiplicity.

Example 5.14. If (A, \mathfrak{m}) is a Teter ring then it is easily verified that both $A[X_1, \ldots, X_n]_{\mathfrak{n}}$ (where $\mathfrak{n} = (\mathfrak{m}, X_1, \ldots, X_n)$) and $A[[X_1, \ldots, X_n]]$ are Teter rings. Applying this result to our examples creates bountiful examples of Teter rings which are domains.

It is perhaps pertinent to give an example of a local domain which is not Teter. To enable this we prove:

Proposition 5.15. Let (A, \mathfrak{m}) be a Cohen-Macaulay local ring of dimension d. If A is Teter then type $(A) = \mu(\mathfrak{m}) - d$.

Remark 5.16. In the above proposition type(A) denotes the Cohen-Macaulay type of A, i.e., the $k = A/\mathfrak{m}$ dimension of $\operatorname{Ext}_{A}^{d}(k, A)$.

TETER RINGS

We now give

Proof of Proposition 5.15. We may assume that the residue field of A is infinite. Let B be a Gorenstein Teter approximation of A. We have an exact sequence $0 \to M \to B \to A \to 0$; where e(M) = 1. We go mod \mathbf{x} an $M \oplus B \oplus A$ -superficial sequence. So we have $0 \to k \to \overline{B} \to \overline{A} \to 0$. So \overline{A} is Teter. We note that $\operatorname{type}(A) = \operatorname{type}(\overline{A})$ and $\mu(\overline{\mathfrak{m}}) = \mu(\mathfrak{m}) - d$. Thus we can assume A is an Artin Teter ring.

Dualizing the sequence $0 \to k \to B \to A \to 0$ we obtain an exact sequence $0 \to \omega_A \to B \to k \to 0$. So we obtain a surjective map $\omega_A \to \mathfrak{m}$ with kernel k. Thus we have exact sequence $0 \to k \to \omega_A \to \mathfrak{m} \to 0$. We apply the functor $\operatorname{Hom}_A(k, -)$ and noting that ω_A is the canonical module of A we obtain $\operatorname{Hom}_A(k, \mathfrak{m}) \cong \operatorname{Ext}_A^1(k, k)$. Thus $\operatorname{type}(A) = \mu(\mathfrak{m})$.

Example 5.17. Consider the following example from [8, p. 395]. Let

 $A = k[[t^5, t^6, t^7, t^9]]$. We have $\mu(\mathfrak{m}) = 4$. But type(A) = 2. So A is not Teter by 5.15.

References

- W. Bruns and J. Herzog, Cohen-Macaulay Rings, revised edition, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, 1998.
- [2] D. Eisenbud and J. Herzog, The classification of homogeneous Cohen-Macaulay rings of finite representation type, Math. Ann. 280 (1988), no. 2, 347–352.
- [3] S. Goto and K. Watanabe, On graded rings. I, J. Math. Soc. Japan 30 (1978), no. 2, 179–213.
- [4] S. Huckaba and T. Marley, Hilbert coefficients and the depths of associated graded rings, J. London Math. Soc. (2) 56 (1997), no. 1, 64–76.
- [5] C. Huneke and A. Vraciu, *Rings that are almost Gorenstein*, Pacific J. Math. 225 (2006), no. 1, 85–102.
- [6] E. Kunz, Introduction to commutative algebra and algebraic geometry, Birkhäuser Boston, Inc., Boston, MA, 1985.
- [7] J. D. Sally, On the associated graded ring of a local Cohen-Macaulay ring, J. Math. Kyoto Univ. 17 (1977), no. 1, 19–21.
- [8] _____, Cohen-Macaulay local rings of embedding dimension e + d 2, J. Algebra 83 (1983), no. 2, 393–408.
- W. Teter, Rings which are a factor of a Gorenstein ring by its socle, Invent. Math. 23 (1974), 153-162.
- [10] Y. Yoshino, Cohen-Macaulay modules over Cohen-Macaulay rings, London Mathematical Society Lecture Note Series, 146. Cambridge University Press, Cambridge, 1990.

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA *Email address*: tputhen@gmail.com