

UNIQUE CONTINUATION FOR LOCALLY UNIFORMLY DISTRIBUTED MEASURES

MAX ENGELSTEIN AND IGNASI GUILLÉN-MOLA

ABSTRACT. In this note we show that the support of a locally k -uniform measure in \mathbb{R}^{n+1} satisfies a kind of unique continuation property. As a consequence, we show that locally uniformly distributed measures satisfy a weaker unique continuation property. This continues work of Kirchheim and Preiss in [KP02] and David, Kenig and Toro in [DKT01] and lends additional evidence to the conjecture proposed by Kowalski and Preiss in [KP87] that the support of a locally n -uniform measure in \mathbb{R}^{n+1} is contained in the zero set of a quadratic polynomial.

1. INTRODUCTION

In this note we consider locally uniform measures, first introduced by Kowalski and Preiss in [KP87] and later studied by David, Kenig and Toro in [DKT01]. For $0 \leq k \leq n + 1$ integer, a Radon measure μ with $\text{supp } \mu \subset \mathbb{R}^{n+1}$ is **locally k -uniform** if

$$(1.1) \quad \mu(B(x, r)) = w_k r^k \text{ whenever } x \in \text{supp } \mu \text{ and } 0 < r \leq 1,$$

where $w_k = \mathcal{H}^k(B^k(0, 1))$ is the measure of the unit ball $B^k(0, 1)$ in \mathbb{R}^k . When $\text{supp } \mu$ is a smooth manifold and $\mu = \mathcal{H}^k|_{\text{supp } \mu}$ it is a natural question in differential geometry to ask what the condition (1.1) tell us about the manifold $\text{supp } \mu$, see e.g., [KP89, Oss75].

For general μ , the condition (1.1) and its global cousin (i.e., (1.1) without the restriction that $r \leq 1$) are extremely well studied, as they represent “end point cases” in the study of measure densities, see e.g., [Pre87, Mar64]. These measure density theorems have also been applied to understand rectifiability in measure spaces (see [DL08, Mer22]) and higher notions of smoothness, see [DKT01, PTT09, Tol15].

The condition (1.1), is extremely rigid and constrains not just the measure, but also its support. For example any two locally uniform measures (in fact locally uniformly distributed, see (1.2)) with the same support must be scalar multiples of one another, see for instance [Mat95, Theorem

Date: January 24, 2025.

2020 Mathematics Subject Classification. 28A75, 28C15, 58C35. Secondary: 28A78, 49Q15.

Key words and phrases. Locally uniform measure, Locally uniformly distributed measure, Unique continuation.

Thanks: M.E. would like to thank Carlos Kenig for introducing him to Conjecture 1 many years ago. I.G.M. would like to thank the first author, Max Engelstein, for his hospitality and mentorship during I.G.M.’s stay at the University of Minnesota (UMN). This work began when I.G.M. was visiting M.E. at the UMN, during Spring 2024. Both authors would like to thank the university for its hospitality. M.E. was partially supported by NSF DMS CAREER 2143719. I.G.M. was supported by Generalitat de Catalunya’s Agency for Management of University and Research Grants (AGAUR) (2021 FI_B 00637 and 2023 FI-3 00151), and also partially supported by 2021-SGR-00071 (AGAUR, Catalonia) and by the Spanish State Research Agency (AEI) projects PID2020-114167GB-I00 and PID2021-125021NAI00. I.G.M. thanks CERCA Programme/Generalitat de Catalunya for institutional support.

3.4]¹. Furthermore, by Marstrand’s theorem, [Mar64], there are no locally k -uniform measures for non-integer k .

For globally uniform measures ((1.1) without the restriction $r \leq 1$), even more is known. It was proven by Kowalski and Preiss in [KP87] that globally k -uniform measures in \mathbb{R}^{n+1} must be of the form $\mathcal{H}^k|_{\Sigma}$ where Σ is a quadratic variety. When $k = n$ (the co-dimension 1 case) the admissible quadratic varieties were fully characterized, with Σ given by (after a translation and rotation)

$$\begin{aligned}\Sigma &= \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_{n+1} = 0\}, \text{ or} \\ \Sigma &= \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_4^2 = x_1^2 + x_2^2 + x_3^2\} \text{ if } n \geq 3.\end{aligned}$$

In higher co-dimension classifying globally 0-uniform and globally 1-uniform measures is straightforward and it follows from work in [KP87] that any connected component of the support of a globally 2-uniform measure must be a sphere or a plane². However, the characterization of the support of globally k -uniform measures when $2 < k < n$ remains an open problem and even constructing interesting examples is extremely difficult (see [Nim22]).

However, given their success in classifying globally n -uniform measures in \mathbb{R}^{n+1} it was natural for Kowalski and Preiss to make the following conjecture about the support of locally n -uniform measures in \mathbb{R}^{n+1} .

Conjecture 1 (See [KP87, Conjecture 5.14]). *Let μ be a locally n -uniform measure in \mathbb{R}^{n+1} . Then each connected component of the support of μ is contained in the zero set of a quadratic polynomial.*

This conjecture remains completely open for $n \geq 3$ with little progress (apart from work of [DKT01] which we will discuss below). In this note, provide some evidence towards Conjecture 1, in any codimension. In particular, we prove that the support of a locally k -uniform measure in \mathbb{R}^{n+1} must satisfy the following “strong unique continuation property” in terms of the mean curvature vector:

Theorem 1.1 (Strong Unique Continuation). *Let μ be a locally k -uniform measure in \mathbb{R}^{n+1} . Assume there is a point $z_0 \in \text{supp } \mu$ such that the mean curvature vector of $\text{supp } \mu$ vanishes at z_0 . Then the connected component of $\text{supp } \mu$ containing z_0 must be a k -plane.*

We prove this theorem in Section 3, where we also recall the definition of the mean curvature vector and establish a seemingly novel and key identity, Lemma 3.1. The assumption that the mean curvature vector vanishes (and is, in particular, well-defined) requires $\text{supp } \mu$ be of class C^2 in a neighborhood of $z_0 \in \text{supp } \mu$. For $k < n$, all that is known (to the best of the authors’ knowledge) is that $\text{supp } \mu$ is a k -dimensional $C^{1,\beta}$ -manifold ($0 < \beta < 1$) away from a closed set \mathcal{S} such that $\mathcal{H}^k(\mathcal{S}) = 0$, see [PTT09, Theorem 1.9]. For $k = n$, work of [DKT01, Theorem 1.10] by David, Kenig and Toro, combined with [PTT09, Corollary 1.11] by Preiss, Tolsa and Toro, shows $\text{supp } \mu = \mathcal{R} \cup \mathcal{S}$, where $\dim \mathcal{S} \leq n - 3$ (discrete if $n = 3$) and around every $z_0 \in \mathcal{R}$ there exists a $r_0 > 0$ such that $B(z_0, r_0) \cap \text{supp } \mu$ is the graph of a C^∞ function. Thus in co-dimension 1 asking that the mean curvature vector exists is not particularly restrictive, while in higher co-dimensions it may in fact place additional constraints on μ (though it seems more likely that a similar regularity theory is true in the higher co-dimension case).

¹The theorem is written for uniformly distributed measures but the proof only uses the local structure of the measures.

²In unpublished work, it seems that Kowalski showed that the support of any locally 2-uniform measure must be a union of spheres and planes in \mathbb{R}^3 , see also [KP89, Proposition 3.1].

Observe that we cannot say much more if we ask that $\mu = \mathcal{H}^k|_{\text{supp } \mu}$, as a standard density argument³ shows that in any ball, B , such that $\text{supp } \mu \cap B$ is sufficiently smooth, we have $\mu|_B = \mathcal{H}^k|_{\text{supp } \mu \cap B}$. That is to say (1.1) (and the work of [PTT09]) immediately implies that $\mu = \mathcal{H}^k|_{\text{supp } \mu}$. Furthermore, in the setting of Theorem 1.1 above, we can conclude that $\mu|_P = \mathcal{H}^k|_P$, where P is the k -plane coinciding with the connected component of $\text{supp } \mu$ containing z_0 .

By the curvature restrictions on the support of (locally smooth) locally n -uniform measures in \mathbb{R}^{n+1} , see [KP87, Corollary 3.1], Theorem 1.1 still holds if the mean curvature is replaced by scalar curvature in co-dimension 1.

Theorem 1.1 provides some evidence towards Conjecture 1 (in any co-dimension), since if $\Sigma = \{Q = 0\}$ for some quadratic polynomial Q , and the second fundamental form of Σ vanishes at a point, then Σ must be a plane. We should mention that we know of only one other result regarding the mean curvature of the support of a locally n -uniform measure in \mathbb{R}^{n+1} : [KP87, Theorem 3.3] shows, under the much stronger assumption that $\text{supp } \mu$ is minimal (i.e., the mean curvature vanishes everywhere) that $\text{supp } \mu$ must be a union of planes.

As a simple corollary, Theorem 1.1 implies a “weak” unique continuation type property for the support of a broader class of measures, which we call locally uniformly distributed. A Radon measure μ with $\text{supp } \mu \subset \mathbb{R}^{n+1}$ is **locally uniformly distributed** if

$$(1.2) \quad 0 < \mu(B(x, r)) = \mu(B(y, r)) < \infty \text{ whenever } x, y \in \text{supp } \mu \text{ and } 0 < r \leq 1.$$

Every locally uniform measure is locally uniformly distributed but the opposite is not always the case. Indeed, $\mathcal{H}^3|_{\mathbb{S}^3}$ is locally uniformly distributed but a straightforward computation shows that it is not locally 3-uniform in \mathbb{R}^4 (one can notice that \mathbb{S}^3 does not satisfy the curvature conditions of [KP87, Corollary 3.1]). In contrast $\mathcal{H}^2|_{\mathbb{S}^2}$ is locally 2-uniform in \mathbb{R}^3 (and locally uniformly distributed). Locally uniformly distributed measures naturally arise as the conical part of globally uniform measures ((1.1) without the restriction $r \leq 1$), and so understanding their structure is an important step towards producing more examples of globally uniform measures, see e.g., [Nim22].

Very little is known even about globally uniformly distributed measures ((1.2) without the restriction that $r \leq 1$). Kirchheim and Preiss showed in [KP02] that globally uniformly distributed measures are supported on analytic varieties. Given the few known examples, it is natural to conjecture that the support of a locally uniformly distributed measure must also be an analytic variety. Our next theorem supports this conjecture.

Theorem 1.2 (Weak Unique Continuation). *Let μ be a locally uniformly distributed measure in \mathbb{R}^{n+1} . Assume there is an open ball $B(z_0, r_0)$ (with $z_0 \in \text{supp } \mu$ and $r_0 > 0$) and a k -plane P such that $B(z_0, r_0) \cap \text{supp } \mu = B(z_0, r_0) \cap P$. Then there is a unique connected component of $\text{supp } \mu$ which intersects non-trivially with $B(z_0, r_0)$ and this connected component is the k -plane P .*

This follows from the strong unique continuation Theorem 1.1 for locally k -uniform measures in \mathbb{R}^{n+1} , and its proof is given in Section 2.

2. PROOF OF THEOREM 1.2: WEAK UNIQUE CONTINUATION PROPERTY

Before proving Theorem 1.1, we show how the weak unique continuation property in Theorem 1.2 follows from our Theorem 1.1.

³See for instance the proof of [KP87, Theorem 4.5], or the proof of Theorem 1.2, more precisely (2.2), in Section 2.

Proof of Theorem 1.2. We mainly follow the first part of [KP02, Proof of Theorem 1.4(ii)]. The Lebesgue differentiation theorem implies that the limit

$$K(y) := \lim_{r \rightarrow 0} \frac{\mu(B(y, r))}{(\mu + \mathcal{H}^k|_{\text{supp } \mu})(B(y, r))}$$

exists at $(\mu + \mathcal{H}^k|_{\text{supp } \mu})$ -a.e. $y \in \text{supp } \mu$, and that

$$(2.1) \quad \mu(E) = \int_E K(y) d(\mu + \mathcal{H}^k|_{\text{supp } \mu})(y) \text{ for any } E \subset B(z_0, r_0),$$

since μ is (clearly) absolutely continuous with respect to $\mu + \mathcal{H}^k|_{\text{supp } \mu}$, i.e., $\mu \ll (\mu + \mathcal{H}^k|_{\text{supp } \mu})$. Note also that the lower bound in (1.2) implies $K(y) > 0$ for $(\mu + \mathcal{H}^k|_{\text{supp } \mu})$ -a.e. $y \in \text{supp } \mu$.

Fixed a (any) point $p \in \text{supp } \mu$, we define

$$g_\mu(r) := \mu(B(p, r)) \text{ for } 0 < r \leq 1,$$

see (1.2). By the flatness assumption $B(z_0, r_0) \cap \text{supp } \mu = B(z_0, r_0) \cap P$ with k -plane P , for $y \in B(z_0, r_0) \cap \text{supp } \mu$ we have

$$K(y) = \lim_{r \rightarrow 0} \frac{g_\mu(r)}{g_\mu(r) + w_k r^k} \leq 1,$$

which is independent of $y \in B(z_0, r_0) \cap \text{supp } \mu$. Since $K(y)$ exists $(\mu + \mathcal{H}^k|_{\text{supp } \mu})$ -a.e., we conclude that the limit $K(y)$ exists for all $y \in B(z_0, r_0) \cap \text{supp } \mu$. Furthermore, since $K(y) > 0$ for $(\mu + \mathcal{H}^k|_{\text{supp } \mu})$ -a.e. $y \in \text{supp } \mu$, it follows that $0 < K(y) \leq 1$ for all $y \in B(z_0, r_0) \cap \text{supp } \mu$.

We define (before checking it makes sense)

$$c := \lim_{r \rightarrow 0} \frac{g_\mu(r)}{w_k r^k}.$$

We have that this limit exists by the existence of the limit $K(y)$ for all $y \in B(z_0, r_0) \cap \text{supp } \mu$ and since

$$\frac{1}{c} = \lim_{r \rightarrow 0} \frac{w_k r^k}{g_\mu(r)} = \lim_{r \rightarrow 0} \frac{w_k r^k + g_\mu(r)}{g_\mu(r)} - 1 = \frac{1}{K(y)} - 1.$$

Moreover, the fact that $K(y) > 0$ implies that $c > 0$. Furthermore if $c = \infty$ then $K(y) = 1$ for $y \in B(z_0, r_0) \cap \text{supp } \mu$ and therefore (2.1) would give $\mathcal{H}^k|_{\text{supp } \mu}(E) = 0$ for any $E \subset B(z_0, r_0) \cap \text{supp } \mu$, but this cannot happen since $B(z_0, r_0) \cap \text{supp } \mu = B(z_0, r_0) \cap P$.

Since $c \in (0, \infty)$, we can write $K(y) = \frac{c}{c+1} \in (0, 1)$ for all $y \in B(z_0, r_0) \cap \text{supp } \mu$, and therefore by (2.1) we conclude that

$$(2.2) \quad \mu(E) = c \mathcal{H}^k|_{\text{supp } \mu}(E) \text{ for any } E \subset B(z_0, r_0).$$

The equality (2.2) in particular implies $g_\mu(r) = \mu(B(x, r)) = c w_k r^k$ for all $x \in B(z_0, r_0/2) \cap \text{supp } \mu$ and $0 < r \leq r_0/4$. However, by the locally uniformly distributed condition (1.2), the same holds for any $x \in \text{supp } \mu$. That is,

$$g_\mu(r) = \mu(B(x, r)) = c w_k r^k \text{ for all } x \in \text{supp } \mu \text{ and } 0 < r \leq r_0/4.$$

Thus, the measure

$$\tilde{\mu}(\cdot) := \frac{1}{c} \frac{1}{\left(\frac{r_0}{4}\right)^k} g\left(\frac{r_0}{4} \cdot\right)$$

satisfies the local uniform condition (1.1), is flat in the open ball $B(4z_0/r_0, 4) = B(z_0, r_0)/(r_0/4)$, and $\text{supp } \tilde{\mu}$ is a dilation of $\text{supp } \mu$. The result follows by Theorem 1.1 applied to the locally k -uniform measure $\tilde{\mu}$. \square

3. PROOF OF THEOREM 1.1: STRONG UNIQUE CONTINUATION PROPERTY

In this section we prove the strong continuation-type property in Theorem 1.1 for locally k -uniform measures in \mathbb{R}^{n+1} that is, if the mean curvature vector of the support vanishes at a point, then the whole connected component containing the point must be a k -plane.

Let us recall that the mean curvature vector of a k -dimensional C^2 -manifold in \mathbb{R}^{n+1} is the trace of the (vector-valued) second fundamental form, divided by k . Let $\mathcal{M} \cap B(z_0, r_0)$, with $z_0 \in \mathcal{M}$ and $r_0 > 0$, be a C^2 -manifold of dimension k . If $\{\hat{n}_j(z)\}_{j=k+1}^{n+1}$ is an orthonormal basis of the normal space $N_{\mathcal{M}}(z)$ at $z \in \mathcal{M} \cap B(z_0, r_0)$ then, intuitively, the j -component of the mean curvature vector is a measure of ‘‘flatness’’ in that it infinitesimally captures how much the measure of \mathcal{M} changes under perturbations in the direction of $\hat{n}_j(z)$.

For our purposes it is enough to work in coordinates, where the definition becomes more concrete. Assume that $\mathcal{M} \cap B(z_0, r_0)$ can be described by a C^2 function $\varphi = (\varphi_{k+1}, \dots, \varphi_{n+1}) : \mathbb{R}^k \rightarrow \mathbb{R}^{n+1-k}$ such that $z = (z', \varphi(z'))$ for $z \in \mathcal{M} \cap B(z_0, r_0)$, with $z' \in \mathbb{R}^k$. Note that, in this case, the tangent space $T_{\mathcal{M}}(z)$ is

$$T_{\mathcal{M}}(z) := \text{span} \left(\{t_j\}_{j=1}^k \right), \text{ with } t_j(z) := \frac{e_j^k + \partial_{x_j} \varphi(z')}{\sqrt{1 + |\partial_{x_j} \varphi(z')|^2}},$$

and $\{e_j^k\}_{j=1}^k$ is the standard orthonormal basis of \mathbb{R}^k . As above, $\{\hat{n}_j(z)\}_{j=k+1}^{n+1}$ is an orthonormal basis of the normal space $N_{\mathcal{M}}(z)$. Then the vector-valued second fundamental form $\mathbb{I}_{\mathcal{M}}(\cdot, \cdot)(z) : T_{\mathcal{M}}(z) \times T_{\mathcal{M}}(z) \rightarrow N_{\mathcal{M}}(z)$ is given by

$$\mathbb{I}_{\mathcal{M}}(u, v)(z) := \Pi_{N_{\mathcal{M}}(z)}(u \cdot \nabla v) = \sum_{j=k+1}^{n+1} \langle u \cdot \nabla v, \hat{n}_j(z) \rangle \hat{n}_j(z).$$

(Where the inner product is the Euclidean inner product).

The mean curvature vector is then the trace so we have that

$$H_{\mathcal{M}}(z) = \frac{1}{k} \sum_{i,j=1}^k g^{ij}(z) \sum_{\ell=k+1}^{n+1} \langle t_j(z) \cdot \nabla t_i(z), \hat{n}_\ell(z) \rangle \hat{n}_\ell(z),$$

where $g^{ij}(z)$ is the inverse of the first fundamental form, given by $g_{ij}(z) = \langle t_i(z), t_j(z) \rangle$.

In generality, this expression is extremely complicated, but it simplifies greatly if we assume that $T_{\mathcal{M}}(0) = \mathbb{R}^k \times \{0\}^{n+1-k}$ (that is, $\partial_{x_i} \varphi(0) = 0$, for every $i = 1, \dots, k$). In this setting we can take $t_j(0) = e_j$, $j = 1, \dots, k$, and $\hat{n}_j(0) = e_j$, $j = k+1, \dots, n+1$ (here $\{e_i\}_{i=1}^{n+1}$ is the standard orthonormal basis of \mathbb{R}^{n+1}). It follows that $g^{ij}(0) = \delta_{ij}$ (i.e., is 1 when $i = j$ and 0 otherwise). Thus our formula simplifies to

$$H_{\mathcal{M}}(0) = \frac{1}{k} \sum_{i=1}^k \sum_{\ell=k+1}^{n+1} \langle e_i \cdot \nabla t_i(0), e_\ell \rangle e_\ell.$$

We recall that $\nabla t_i(0)$ is a matrix and we need only compute its (ℓ, i) -entry. This is a straightforward computation that gives

$$(\nabla t_i(0))_{\ell, i} = \partial_{x_i} \left(\frac{\partial_{x_i} \varphi_\ell}{\sqrt{1 + |\partial_{x_i} \varphi|^2}} \right) (0) = \partial_{x_i}^2 \varphi_\ell(0),$$

where we have used in the second equality that $\partial_{x_i} \varphi(0) = 0$ for all $i = 1 \dots, k$. Thus we have the final form

$$(3.1) \quad H_{\mathcal{M}}(0) = \frac{1}{k} \sum_{\ell=k+1}^{n+1} \Delta \varphi_\ell(0) e_\ell = \frac{1}{k} (0, \Delta \varphi(0)), \text{ if } T_{\mathcal{M}}(0) = \mathbb{R}^k \times \{0\}^{n+1-k}.$$

Before we begin our proofs, let us introduce some preliminary notation. We define the vector $b_{z,r}$ and its nonnormalized version $\tilde{b}_{z,r}$ as

$$(3.2a) \quad b_{z,r} := \frac{k+2}{2w_k r^{k+2}} \int_{B(z,r)} (r^2 - |y-z|^2) (y-z) d\mu(y),$$

$$(3.2b) \quad \tilde{b}_{z,r} := \frac{2w_k r^{k+2}}{k+2} b_{z,r} = \int_{B(z,r)} (r^2 - |y-z|^2) (y-z) d\mu(y),$$

and the quadratic form $Q_{z,r}$ and its normalized version $\tilde{Q}_{z,r}$ on \mathbb{R}^{n+1} as

$$(3.3a) \quad Q_{z,r}(x) := \frac{k+2}{w_k r^{k+2}} \int_{B(z,r)} \langle x, y-z \rangle^2 d\mu(y),$$

$$(3.3b) \quad \tilde{Q}_{z,r}(x) := \frac{w_k r^{k+2}}{k+2} Q_{z,r} = \int_{B(z,r)} \langle x, y-z \rangle^2 d\mu(y),$$

all first introduced in [KP87, (4.3) and (4.4)], see also [DKT01, (7.2) and (7.3)]. Throughout this section, $b_{z,r}^j$ and $\tilde{b}_{z,r}^j$ denote the j -component ($1 \leq j \leq n+1$) of the vectors $b_{z,r}$ and $\tilde{b}_{z,r}$ respectively.

Here we present the key equality in the proof of our main result Theorem 1.1. This relates the orthogonal (see (3.6)) vector $\tilde{b}_{0,r}$ and the mean curvature (given in local coordinates by $\Delta \varphi(0)$, see (3.1)), to the integral of “height” of the support squared. This identity is the main novel part of our work.

Lemma 3.1 (Key equality). *Let μ be a locally k -uniform measure in \mathbb{R}^{n+1} and $0 < r < 1/2$. Assume $0 \in \text{supp } \mu$ and that $\text{supp } \mu \cap B(0, r_0)$ (for some $r_0 > 0$) is given by the graph of a C^2 function $\varphi = (\varphi_{k+1}, \dots, \varphi_{n+1}) : \mathbb{R}^k \rightarrow \mathbb{R}^{n+1-k}$ with $|\nabla \varphi_j(0)| = 0$ for all $j = k+1, \dots, n+1$. Then there holds*

$$\frac{1}{2} \sum_{j=k+1}^{n+1} \tilde{b}_{0,r}^j \Delta \varphi_j(0) = \sum_{j=k+1}^{n+1} \int_{B(0,r)} y_j^2 d\mu(y).$$

We prove this in Section 3.1. We are now ready to turn to the proof of Theorem 1.1.

Proof of Theorem 1.1. Since we are C^2 -smooth in a neighborhood of z_0 , we can work in coordinates: after a harmless rotation and translation, let $\text{supp } \mu$ in a neighborhood of $0 \in \text{supp } \mu$ be given by $(x', \varphi(x'))$, $x' \in \mathbb{R}^n$, with $\varphi(x) = (\varphi_{k+1}(x'), \dots, \varphi_{n+1}(x'))$ satisfying $\varphi(0) = 0$ and $|\nabla \varphi_j(0)| = 0$ for all $j = k+1, \dots, n+1$. As we computed in (3.1) above, this implies that

$$H_{\text{supp } \mu}(0) = \frac{1}{k} (0, \Delta \varphi(0)).$$

Fixed $0 < r < 1/2$, by Lemma 3.1 we have

$$\frac{1}{2} \sum_{j=k+1}^{n+1} \tilde{b}_{0,r}^j \Delta \varphi_j(0) = \sum_{j=k+1}^{n+1} \int_{B(0,r)} y_j^2 d\mu(y).$$

Since $(0, \Delta \varphi(0)) = 0$ (in particular $\Delta \varphi_j(0) = 0$ for all $j = k+1, \dots, n+1$), then the integral in the right-hand side is zero, implying that $y_j = 0$ for μ -a.e. $y \in B(0, r)$ for all $j = k+1, \dots, n+1$. But this is only possible if $B(0, r) \cap \text{supp } \mu = B(0, r) \cap P$ for the k -plane $P := \mathbb{R}^k \times \{0\}^{n+1-k}$. Since this holds for every $0 < r < 1/2$, we conclude that $\text{supp } \mu \cap B(0, 1/2) = P \cap B(0, 1/2)$.

To show that $B(z_0, 1/2) \cap \text{supp } \mu = B(z_0, 1/2) \cap P$ for any z_0 in the same connected component of $\text{supp } \mu$, consider a chain of balls of radius $1/2$ centered at points in $\text{supp } \mu$ such that the center of each new ball is in the previous ball with 0 as the center of the first ball and z_0 as the center of the last ball. The result follows by applying this argument to each ball in the chain in sequence. \square

3.1. Proof of Key equality: Lemma 3.1. Recall the definition of $b_{z,r}$ and $Q_{z,r}$ from (3.2a) and (3.3a). Let us state two fundamental properties satisfied by a locally k -uniform measure μ in \mathbb{R}^{n+1} . The first is the elementary identity

$$(3.4) \quad \int_{B(z,r)} |y - z|^2 d\mu(y) = \frac{k w_k}{k+2} r^{k+2} \text{ for all } z \in \text{supp } \mu \text{ and } 0 < r < 1,$$

for the detailed straightforward computation (for $k = n$) see [KP87, (4.5)] for instance. Secondly, proved in [DKT01, (10.3)],⁴ for $z \in \text{supp } \mu$ and $0 < r < 1/2$ there holds

$$(3.5) \quad |\langle 2b_{z,r}, x - z \rangle + Q_{z,r}(x) - |x|^2| \leq C \frac{|x - z|^3}{r} \text{ for all } x \in \text{supp } \mu \cap B(z, r/2).$$

Since $Q_{z,r}(x) \leq \frac{k+2}{w_k} |x|^2$, this in particular implies that for $|x - z|$ small enough there holds

$$|\langle 2b_{z,r}, x - z \rangle| \leq C' |x - z|^2.$$

From this latter inequality we conclude that, if $\text{supp } \mu \cap B(z, r_0)$ (for some $r_0 > 0$) is at least of class C^1 , then either $b_{z,r} = 0$ or

$$(3.6) \quad b_{z,r} \text{ is orthogonal to the tangent } k\text{-plane of } \text{supp } \mu \text{ at } z \in \text{supp } \mu.$$

Having recalled these facts we can now establish the following algebraic relation, which gets us most of the way to Lemma 3.1.

Lemma 3.2. *Under the assumptions of Lemma 3.1, there holds*

$$(3.7) \quad \sum_{j=k+1}^{n+1} b_{0,r}^j \langle \nabla^2 \varphi_j(0) e, e \rangle = 1 - Q_{0,r}(e) \text{ for all } e \in \mathbb{S}^{k-1} \times \{0\}^{n+1-k}.$$

Proof of Lemma 3.2. During the proof, by the C^2 local representation of $\text{supp } \mu$ we write $x = (x', \varphi(x'))$ with $x' \in \mathbb{R}^k$ and $\varphi(x') = (\varphi_{k+1}(x'), \dots, \varphi_{n+1}(x')) \in \mathbb{R}^{n+1-k}$. As we are assuming that $0 \in \text{supp } \mu$ and $|\nabla \varphi_j(0)| = 0$ for all $j = k+1, \dots, n+1$, by Taylor's theorem we then have

$$\varphi_j(x') = \frac{\langle \nabla^2 \varphi_j(0) x', x' \rangle}{2} + o(|x'|^2) \text{ for all } j = k+1, \dots, n+1.$$

(We use little o notation, so that $o(s)$ is a quantity that when divided by s goes to zero as $s \downarrow 0$).

⁴The codimension one version is proved in [DKT01, (10.3)], but its proof does not depend on the codimension. For a statement in any codimension, see [PTT09, Proposition 2.3 (2.12)].

With $z = 0$, (3.5) reads as

$$(3.8) \quad |\langle 2b_{0,r}, x \rangle + Q_{0,r}(x) - |x|^2| \leq C \frac{|x|^3}{r} \text{ for all } x \in \text{supp } \mu \cap B(0, r/2).$$

First, from (3.6) we have that $b_{0,r}$ is orthogonal to the tangent k -plane of $\text{supp } \mu$ at 0, or it may be zero. That is, in local coordinates, this means $b_{0,r} \in (\mathbb{R}^k \times \{0\}^{n+1-k})^\perp$, since $|\nabla \varphi_j(0)| = 0$ for all $j = k+1, \dots, n+1$. This implies $b_{0,r}^j = 0$ for all $j = 1, \dots, k$, and therefore,

$$\langle 2b_{0,r}, x \rangle = 2 \sum_{j=k+1}^{n+1} b_{0,r}^j \varphi_j(x') = \sum_{j=k+1}^{n+1} b_{0,r}^j \langle \nabla^2 \varphi_j(0) x', x' \rangle + o(|x'|^2).$$

Second,

$$\begin{aligned} Q_{0,r}(x) &= \frac{k+2}{w_k r^{k+2}} \int_{B(0,r)} \langle x, y \rangle^2 d\mu(y), \\ &= \frac{k+2}{w_k r^{k+2}} \int_{B(0,r)} \left(\langle x', y' \rangle + \sum_{j=k+1}^{n+1} \varphi_j(x') y_j \right)^2 d\mu(y), \\ &= \frac{k+2}{w_k r^{k+2}} \int_{B(0,r)} \langle x', y' \rangle^2 d\mu(y) + o(|x'|^2). \end{aligned}$$

Finally, simply $|x|^2 = |x'|^2 + o(|x'|^3)$. Also from this, we have that $|x| \approx |x'|$ for $x \in \text{supp } \mu \cap B(0, r/2)$ with small enough modulus.

From (3.8) and these estimates, there exists a $\theta \in (0, 1)$ (depending on $\|\varphi\|_{C^2}$) such that for all $x \in \text{supp } \mu \cap B(0, r/2)$ with $|x| \leq \theta r$ we get

$$(3.9) \quad \left| \sum_{j=k+1}^{n+1} b_{0,r}^j \langle \nabla^2 \varphi_j(0) x', x' \rangle + \frac{k+2}{w_k r^{k+2}} \int_{B(0,r)} \langle x', y' \rangle^2 d\mu(y) - |x'|^2 \right| \leq \frac{1}{r} o(|x'|^2).$$

Note that we can simply write the middle term as $Q_{0,r}((x', 0))$.

Given $e \in \mathbb{S}^{k-1} \times \{0\}^{n+1-k}$, let $\theta r/2 > \varepsilon > 0$ and $x' = \varepsilon e$ in (3.9). Dividing both sides of (3.9) by $|x'|^2 = \varepsilon^2$ and letting $\varepsilon \downarrow 0$, we get (3.7). \square

We now turn to the proof of Lemma 3.1.

Proof of Lemma 3.1. Multiplying the equality (3.7) in Lemma 3.2 by $w_k r^{k+2}/(k+2)$ we have

$$(3.10) \quad \frac{1}{2} \sum_{j=k+1}^{n+1} \tilde{b}_{0,r}^j \langle \nabla^2 \varphi_j(0) e, e \rangle = \frac{w_k}{k+2} r^{k+2} - \tilde{Q}_{0,r}(e) \text{ for all } e \in \mathbb{S}^{k-1} \times \{0\}^{n+1-k}.$$

For $1 \leq i \leq k$, taking $e = e_i \in \mathbb{S}^{k-1} \times \{0\}^{n+1-k}$ in (3.10), recall $\{e_i\}_{i=1}^{n+1}$ is the standard orthonormal basis of \mathbb{R}^{n+1} , we have

$$(3.11) \quad \frac{1}{2} \sum_{j=k+1}^{n+1} \tilde{b}_{0,r}^j \partial_{x_i, x_i} \varphi_j(0) = \frac{w_k}{k+2} r^{k+2} - \int_{B(0,r)} y_i^2 d\mu(y).$$

Adding up (3.11) for $1 \leq i \leq k$ we get

$$\frac{1}{2} \sum_{j=k+1}^{n+1} \tilde{b}_{0,r}^j \Delta \varphi_j(0) = \frac{k w_k}{k+2} r^{k+2} - \sum_{i=1}^k \int_{B(0,r)} y_i^2 d\mu(y).$$

The lemma follows from the above after applying the identity (3.4). \square

REFERENCES

- [DKT01] Guy David, Carlos Kenig, and Tatiana Toro. Asymptotically optimally doubling measures and Reifenberg flat sets with vanishing constant. *Comm. Pure Appl. Math.*, 54(4):385–449, 2001. (Cited on pages 1, 2, 6, and 7)
- [DL08] Camillo De Lellis. *Rectifiable sets, densities and tangent measures*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008. (Cited on page 1)
- [KP87] Oldřich Kowalski and David Preiss. Besicovitch-type properties of measures and submanifolds. *J. Reine Angew. Math.*, 379:115–151, 1987. (Cited on pages 1, 2, 3, 6, and 7)
- [KP89] Leon Karp and Mark Pinsky. Volume of a small extrinsic ball in a submanifold. *Bull. London Math. Soc.*, 21(1):87–92, 1989. (Cited on pages 1 and 2)
- [KP02] Bernd Kirchheim and David Preiss. Uniformly distributed measures in Euclidean spaces. *Math. Scand.*, 90(1):152–160, 2002. (Cited on pages 1, 3, and 4)
- [Mar64] John M. Marstrand. The (φ, s) regular subsets of n -space. *Trans. Amer. Math. Soc.*, 113:369–392, 1964. (Cited on pages 1 and 2)
- [Mat95] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability. (Cited on page 2)
- [Mer22] Andrea Merlo. Geometry of 1-codimensional measures in Heisenberg groups. *Invent. Math.*, 227(1):27–148, 2022. (Cited on page 1)
- [Nim22] A. Dali Nimer. Conical 3-uniform measures: a family of new examples and characterizations. *J. Differential Geom.*, 121(1):57–99, 2022. (Cited on pages 2 and 3)
- [Oss75] Robert Osserman. Isoperimetric and related inequalities. In *Differential geometry (Proc. Sympos. Pure Math., Vol. XXVII, Part 1, Stanford Univ., Stanford, Calif., 1973)*, Proc. Sympos. Pure Math., Vol. XXVII, Part 1, pages 207–215. Amer. Math. Soc., Providence, RI, 1975. (Cited on page 1)
- [Pre87] David Preiss. Geometry of measures in \mathbf{R}^n : distribution, rectifiability, and densities. *Ann. of Math. (2)*, 125(3):537–643, 1987. (Cited on page 1)
- [PTT09] D. Preiss, X. Tolsa, and T. Toro. On the smoothness of Hölder doubling measures. *Calc. Var. Partial Differential Equations*, 35(3):339–363, 2009. (Cited on pages 1, 2, 3, and 7)
- [Tol15] Xavier Tolsa. Uniform measures and uniform rectifiability. *J. Lond. Math. Soc. (2)*, 92(1):1–18, 2015. (Cited on page 1)

MAX ENGELSTEIN, SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN, 55455, USA.
Email address: `mengelst@umn.edu`

IGNASI GUILLÉN-MOLA, DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA.
Email address: `ignasi.guillen@uab.cat`