Can one-loop corrections to the one-gluon exchange potential adequately describe the charmed meson spectrum?

A. Capelo-Astudillo,^{1, *} T. Aguilar,¹ M. Conde-Correa,¹ A. Duenas-Vidal,^{1, †} P. G. Ortega,^{2, 3, ‡} and J. Segovia^{4, §}

¹Departamento de Física, Escuela Politécnica Nacional, Quito 170143, Ecuador

²Departamento de Física Fundamental, Universidad de Salamanca, E-37008 Salamanca, Spain

³Instituto Universitario de Física Fundamental y Matemáticas (IUFFyM),

⁴Departamento de Sistemas Físicos, Químicos y Naturales,

Universidad Pablo de Olavide, E-41013 Sevilla, Spain

(Dated: January 27, 2025)

We investigate the charmed meson spectrum using a constituent quark model (CQM) with oneloop corrections applied to the one-gluon exchange (OGE) potential. The study aims to understand if these one-loop corrections sufficiently account for the charmed meson spectrum observed experimentally, without invoking exotic configurations like tetraquarks. Within this model, charmed mesons' masses are computed, comparing theoretical predictions to experimental data. The results suggest that the model, with one-loop OGE corrections, generally reproduces mass splittings and level ordering observed for charmed mesons. Discrepancies, particularly in *P*-wave states, are addressed by incorporating higher-order interaction terms. The findings emphasize that while the traditional quark model is limited in fully describing charmed mesons, enhanced potential terms may bridge the gap with experimental observations. The study contributes a framework for predicting excited charmed meson states for future experimental validation.

I. INTRODUCTION

A simple analysis about the properties of mesons containing a single heavy quark, Q = c or b, can be carried out in the limit of $m_Q \to \infty$. In such a regime, the heavy quark acts as a static color source for the rest of the heavy-light meson, *i.e.* its spin s_Q is decoupled from the total angular momentum of the light antiquark, j_q , and they are separately conserved. As a result, heavy-light mesons are grouped into doublets, each associated with a specific value of j_q and parity. The members of each doublet differ from the orientation of s_Q with respect to j_q and they are degenerate in the heavy quark symmetry (HQS) limit [1], whose mass degeneracy is broken at order $1/m_Q$.

For $Q\bar{q}$ states, and following HQS, one can write $\vec{j}_q = \vec{s}_q + \vec{\ell}$, where s_q is the light antiquark spin and ℓ is its orbital angular momentum relative to the static heavy quark. Therefore, the lowest-lying $Q\bar{q}$ mesons correspond to $\ell = 0$ with $j_q^P = \frac{1}{2}^-$. This doublet comprises two S-wave states with spin-parity $J^P = (0^-, 1^-)$, where $\vec{J} = \vec{j}_q + \vec{s}_Q$. For $\ell = 1$, it could be either $j_q^P = \frac{1}{2}^+$ or $j_q^P = \frac{3}{2}^+$, and thus the two corresponding doublets are $J^P = (0^+, 1^+)$ and $J^P = (1^+, 2^+)$, respectively. The mesons with $\ell = 2$ are collected either in the $j_q^P = \frac{3}{2}^-$ doublet, consisting of states with $J^P = (1^-, 2^-)$, or in the $j_q^P = \frac{5}{2}^-$ with $J^P = (2^-, 3^-)$; and so forth and so on.

If we now focus on the spectrum of charmed mesons, $(c\bar{n})$ -states with n = u or d quark, it contains a number of long known and well established states collected in the Review of Particle Physics (RPP) of Particle Data Group (PDG) [2]. We find the lowest-lying S-wave states with quantum numbers $J^P = 0^-$ and 1^- , denoted as D and D^* mesons. The P-wave ground states with spin-parity quantum numbers 0^+ ($D_0^*(2300)$), 1^+ ($D_1(2420)$ and $D_1(2430)$) and 2^+ ($D_2^*(2460)$) are also given in Ref. [2]. In addition, the PDG lists as wellestablished state, a highly-excited charmed meson, with spin-parity $J^P = 3^-$, denoted as $D_3^*(2750)$. It was observed as a resonant substructure in the $B^0 \to \overline{D}^0 \pi^+ \pi^$ and $B^- \to D^+ \pi^- \pi^-$ decays analyzed with the Dalitz plot technique [3, 4].

Over the past 15 years or so, several new signals in the charmed meson sector have been observed. The now named $D_0(2550)$, $D_1^*(2600)$, $D_2(2740)$ and $D_3^*(2750)$ were observed for the first time by the BaBar collaboration in 2010 [5], and were confirmed by the LHCb experiment with slightly different masses in 2013 [6]. Furthermore, the LHCb collaboration reported in Refs. [6] two new higher *D*-meson excitations, $D_J^*(3000)$ and $D_J(3000)$, with natural and unnatural parities,¹ respectively, collectively named by the PDG as $D(3000)^0$. In 2015, again the LHCb collaboration observed a new state $D_1^*(2760)$ with spin-parity quantum numbers $J^P = 1^$ in the $D^+\pi^-$ channel by analyzing the $B^- \to D^+K^-\pi^$ decay [7]. Finally, there have been observed two more

Universidad de Salamanca, E-37008 Salamanca, Spain

^{*} andre.capelo@epn.edu.ec

[†] alvaro.duenas@epn.edu.ec

[‡] pgortega@usal.es

[§] jsegovia@upo.es

¹ Natural parity means that the bosonic meson field behaves under reflection as +1 for even spin and -1 for odd spin; note then that, for heavy-light mesons, the superindex "*" is used for those having natural parity.

states which are not collected in the RPP of PDG. The first one is the named $D^*(2640)^{\pm}$ seen in Z decays by Abreu *et al.* [8] but missing in the searches performed in Refs. [9, 10], thus requiring confirmation. The second was observed in 2016 by the LHCb collaboration in the $D^+\pi^$ channel when analyzing the $B^- \to D^+\pi^-\pi^-$ decay [4]; they assigned to this signal the name $D_2^*(3000)$ with spinparity $J^P = 2^+$ because its resonance parameters were inconsistent with the previously observed $D(3000)^0$ [6].

Theoretical predictions of the spectrum of charmed mesons dates from the early days of phenomenological quark models [11–14]. In the recent years, many theoretical studies have been carried out within different theoretical approaches such as lattice-regularized QCD [15–19], unitarized coupled-channels T-matrix analyses [20–22], heavy meson effective theory [23, 24], Regge-based phenomenology [25, 26] and phenomenological quark models [27–36]. This is mainly because two reasons; the first one is the recent experimental measurements in the subject which provide sustained progress in the field as well as the breadth and depth necessary for a vibrant theoretical research environment. The second is related mostly with the fact that $D_0^*(2300)$ and $D_1(2420)$ charmed mesons, which belong to the doublet $j_q^P = \frac{1}{2}^+$ predicted by HQS, have surprisingly light masses, compared with naive quark model expectations, and are located below $D\pi$ and $D^*\pi$ thresholds, respectively. This implies that these states are narrow. These facts have stimulated a fruitful line of research, suggesting that their structure is much richer than what one might guess assuming the $q\bar{q}$ picture [37–39].

The quark model has been notably successful in describing the heavy quark-antiquark system since the early days of charmonium studies (see, for example, Refs. [40– 47]). Predictions from this framework on the properties of heavy quarkonium, including those related to decays and interactions, have proven highly valuable for guiding experimental searches. Additionally, the quark model's adaptability makes it well-suited for exploratory research. Thus, the theoretical results presented here are based on a constituent quark model (CQM), initially proposed in Ref. [48], and recently applied to conventional mesons containing heavy quarks, capturing a broad range of physical observables related to spectra [49–52], strong decays [53–55], hadronic transitions [56–58], and both electromagnetic and weak reactions [59–61]. To improve the accuracy of mass splittings, we adopt the approach in Ref. [62] and incorporate one-loop corrections to the One-Gluon Exchange (OGE) potential as derived by Gupta and Radford [63]. These corrections include a spin-dependent term that impacts only mesons with quarks of different flavors. Our primary objective is to determine whether the entire spectrum of experimentally observed charmed mesons can be described within the quark-antiquark model alone, without needing to invoke more exotic configurations.

The manuscript is organized as follows. After this introduction, the theoretical framework is briefly presented in Sec II. Section III is mainly devoted to the analysis and discussion of our theoretical results. Finally, we summarize and draw some conclusions in Sec. IV.

II. THEORETICAL FORMALISM

Dynamical chiral symmetry breaking in Quantum Chromodynamics (QCD) is responsible, among other phenomena, of generating constituent quark masses and Goldstone-boson exchanges between light quarks. This together with one-gluon exchange and confining interactions consist on the main pieces of our constituent quark model [48, 64].

Under chiral transformations, the following Lagrangian

$$\mathcal{L} = \bar{\psi}(i \partial \!\!\!/ - M(q^2) U^{\gamma_5}) \psi, \qquad (1)$$

is invariant [65]. In Eq. (1), $M(q^2)$ is the dynamical momentum-dependent constituent quark mass and $U^{\gamma_5} = e^{i\lambda_a \phi^a \gamma_5 / f_{\pi}}$, with $\phi = \{\vec{\pi}, K, \eta_8\}$, is the matrix of Goldstone-boson fields that can be expanded as

$$U^{\gamma_5} = 1 + \frac{i}{f_{\pi}} \gamma^5 \lambda^a \phi^a - \frac{1}{2f_{\pi}^2} \phi^a \phi^a + \dots$$
 (2)

One can guess that the first term of the expansion provides the constituent quark mass, the second gives rise to one-boson exchange interactions between quarks and the main contribution of the third term comes from the twopion exchange which is simulated in our case by means of a scalar-meson exchange interaction. Note, however, that in the presence of heavy quarks chiral symmetry is explicitly broken and Goldstone-boson exchanges do not appear. However, it constrains the model parameters through the light-meson phenomenology [66].

At higher energy scales, the CQM incorporates QCD perturbative effects by taking into account one-gluon fluctuations around the instanton vacuum through the vertex Lagrangian

$$\mathcal{L}_{qqg} = i\sqrt{4\pi\alpha_s}\,\bar{\psi}\gamma_\mu G^\mu_c \lambda^c \psi\,,\tag{3}$$

with λ^c the SU(3) color matrices and G_c^{μ} the gluon field. The α_s is a scale-dependent effective strong coupling constant that allows a comprehensive description of light, strange and heavy meson spectra [48, 64]:

$$\alpha_s(\mu) = \frac{\alpha_0}{\ln\left(\frac{\mu_{ij}^2 + \mu_0^2}{\Lambda_0^2}\right)},\tag{4}$$

in which μ_{ij} is the reduced mass of the meson's constituent $q\bar{q}$ pair and α_0 , μ_0 and Λ_0 are parameters of the model.

The potential derived from Eq. (3) contains central, tensor, and spin-orbit contributions given by

$$\begin{split} V_{\text{OGE}}^{\text{C}}(\vec{r}_{ij}) &= \frac{1}{4} \alpha_s(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}} - \frac{1}{6m_i m_j} (\vec{\sigma}_i \cdot \vec{\sigma}_j) \frac{e^{-r_{ij}/r_0(\mu)}}{r_{ij} r_0^2(\mu)} \right], \\ V_{\text{OGE}}^{\text{T}}(\vec{r}_{ij}) &= -\frac{1}{16} \frac{\alpha_s}{m_i m_j} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}} \left(\frac{1}{r_{ij}^2} + \frac{1}{3r_g^2(\mu)} + \frac{1}{r_{ij} r_g(\mu)} \right) \right] S_{ij}, \\ V_{\text{OGE}}^{\text{SO}}(\vec{r}_{ij}) &= -\frac{1}{16} \frac{\alpha_s}{m_i^2 m_j^2} (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) \left[\frac{1}{r_{ij}^3} - \frac{e^{-r_{ij}/r_g(\mu)}}{r_{ij}^3} \left(1 + \frac{r_{ij}}{r_g(\mu)} \right) \right] \times \\ &\times \left[((m_i + m_j)^2 + 2m_i m_j) (\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2) (\vec{S}_- \cdot \vec{L}) \right], \end{split}$$
(5)

where $S_{ij} = 3(\vec{\sigma}_i \cdot \hat{r}_{ij})(\vec{\sigma}_j \cdot \hat{r}_{ij}) - \vec{\sigma}_i \cdot \vec{\sigma}_j$ is the quark tensor operator and $\vec{S}_{\pm} = \frac{1}{2}(\vec{\sigma}_i \pm \vec{\sigma}_j)$ are the symmetric and antisymmetric spin-orbit operators. Besides, $r_0(\mu) = \hat{r}_0 \frac{\mu_{nn}}{\mu_{ij}}$ and $r_g(\mu) = \hat{r}_g \frac{\mu_{nn}}{\mu_{ij}}$ are regulators which depend on μ_{ij} , the reduced mass of the meson's constituent $q\bar{q}$

pair. The contact term of the central potential has been regularized as $\delta(\vec{r}_{ij})\approx(1/4\pi r_0^2)\cdot e^{-r_{ij}/r_0}/r_{ij}$

To improve the description of charmed mesons, we follow the proposal of Ref. [62] and include one-loop corrections to the OGE potential as derived by Gupta *et al.* [63]. As in the case of V_{OGE} , $V_{\text{OGE}}^{1-\text{loop}}$ contains central, tensor and spin-orbit contributions, given by [67]

$$\begin{split} V_{\text{OGE}}^{1-\text{loop,C}}(\vec{r}_{ij}) &= 0, \\ V_{\text{OGE}}^{1-\text{loop,T}}(\vec{r}_{ij}) &= \frac{C_F}{4\pi} \frac{\alpha_s^2}{m_i m_j} \frac{1}{r^3} S_{ij} \left[\frac{b_0}{2} \left(\ln(\mu r_{ij}) + \gamma_E - \frac{4}{3} \right) + \frac{5}{12} b_0 - \frac{2}{3} C_A \\ &\quad + \frac{1}{2} \left(C_A + 2C_F - 2C_A \left(\ln(\sqrt{m_i m_j} r_{ij}) + \gamma_E - \frac{4}{3} \right) \right) \right], \\ V_{\text{OGE}}^{1-\text{loop,SO}}(\vec{r}_{ij}) &= \frac{C_F}{4\pi} \frac{\alpha_s^2}{m_i^2 m_j^2} \frac{1}{r^3} \times \\ \times \left\{ (\vec{S}_+ \cdot \vec{L}) \left[\left((m_i + m_j)^2 + 2m_i m_j \right) \left(C_F + C_A - C_A \left(\ln(\sqrt{m_i m_j} r_{ij}) + \gamma_E \right) \right) \\ &\quad + 4m_i m_j \left(\frac{b_0}{2} \left(\ln(\mu r_{ij}) + \gamma_E \right) - \frac{1}{12} b_0 - \frac{1}{2} C_F - \frac{7}{6} C_A + \frac{C_A}{2} \left(\ln(\sqrt{m_i m_j} r_{ij}) + \gamma_E \right) \right) \\ &\quad + \frac{1}{2} (m_j^2 - m_i^2) C_A \ln\left(\frac{m_j}{m_i}\right) \right] \\ + (\vec{S}_- \cdot \vec{L}) \left[(m_j^2 - m_i^2) \left(C_F + C_A - C_A \left(\ln(\sqrt{m_i m_j} r_{ij}) + \gamma_E \right) \right) \\ &\quad + \frac{1}{2} (m_i + m_j)^2 C_A \ln\left(\frac{m_j}{m_i}\right) \right] \right], \end{split}$$

where $C_F = 4/3$, $C_A = 3$, $b_0 = 9$, $\gamma_E = 0.5772$ and the scale $\mu \approx 1 \text{ GeV}$.

Finally, an important non-perturbative term of our CQM is color confining interaction between quarks and antiquarks to ensure colorless hadrons. The potential used here is linearly-rising for short interquark distances, but acquires a plateau at large distances to mimic the effect of sea quarks, which induces the breakdown of the color binding string [68]. Its explicit expression is

$$V_{\text{CON}}^{\text{C}}(\vec{r}_{ij}) = \left[-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta\right] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c),$$

$$V_{\text{CON}}^{\text{SO}}(\vec{r}_{ij}) = -\left(\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c\right) \frac{a_c \mu_c e^{-\mu_c r_{ij}}}{4m_i^2 m_j^2 r_{ij}} \left[((m_i^2 + m_j^2)(1 - 2a_s) + 4m_i m_j(1 - a_s))(\vec{S}_+ \cdot \vec{L}) + (m_j^2 - m_i^2)(1 - 2a_s)(\vec{S}_- \cdot \vec{L}) \right],$$
(7)

Γ

where the model parameters are a_c , Δ , μ_c and a_s , being the last one the mixture between scalar and vector Lorentz structures of the confinement.

Among the different methods to solve the Schrödinger equation in order to find the quark-antiquark bound states, we use the Gaussian Expansion Method [69] because it provides enough accuracy and makes the subsequent evaluation of matrix elements easier. This procedure provides the radial wave function solution of the Schrödinger equation as an expansion in terms of basis functions

$$R_{\alpha}(r) = \sum_{n=1}^{n_{max}} c_n^{\alpha} \phi_{nl}^G(r), \qquad (8)$$

where α refers to the channel quantum numbers. Following Ref. [69], we employ Gaussian trial functions with ranges in geometric progression. This enables the optimization of ranges employing a small number of free parameters. Moreover, the geometric progression is dense at short distances, so that it allows the description of the dynamics mediated by short range potentials. The fast damping of the gaussian tail is not a problem, since we can choose the maximal range much longer than the hadronic size. The coefficients, c_n^{α} , and the eigenvalue, E, are determined from the Rayleigh-Ritz variational principle

$$\sum_{n=1}^{n_{max}} \left[\left(T^{\alpha}_{n'n} - EN^{\alpha}_{n'n} \right) c^{\alpha}_{n} + \sum_{\alpha'} V^{\alpha\alpha'}_{n'n} c^{\alpha'}_{n} = 0 \right], \quad (9)$$

where $T^{\alpha}_{n'n}$, $N^{\alpha}_{n'n}$ and $V^{\alpha\alpha'}_{n'n}$ are the matrix elements of the kinetic energy, the normalization and the potential, respectively. The matrices $T^{\alpha}_{n'n}$ and $N^{\alpha}_{n'n}$ are diagonal whereas the mixing between different channels is given by $V^{\alpha\alpha'}_{n'n}$.

III. RESULTS

Model parameters relevant for this analysis are shown in Table I. As we have mention in the introduction, all model parameters were constrained based on prior investigations into hadron phenomenology (see, for instance, Refs. [48, 50, 64]). Then, from this perspective, we present a parameter-free model-dependent prediction of charmed mesons. Finally, the parameters were fixed to describe a set of hadron observables within a certain

| Quark masses | $m_n ({\rm MeV})$ | 313 |
|--------------|------------------------------|---------|
| · | m_c (MeV) | 1763 |
| OGE | $lpha_0$ | 2.118 |
| | $\Lambda_0 ~({\rm fm}^{-1})$ | 0.113 |
| | $\mu_0 ~({\rm MeV})$ | 36.976 |
| | $\hat{r}_0 ~(\mathrm{fm})$ | 0.181 |
| | \hat{r}_g (fm) | 0.259 |
| Confinement | $a_c \; (MeV)$ | 507.4 |
| | $\mu_c \; ({\rm fm}^{-1})$ | 0.576 |
| | $\Delta (MeV)$ | 184.432 |
| | a_s | 0.81 |

range of agreement and thus a theoretical uncertainty is associated with this model adjustment. To assess this, the results presented in this manuscript show a theoretical uncertainty of ± 25 MeV in the meson's mass.

In Table II, we show the charmed meson masses, in MeV, from constituent quark model (CQM) and experiment [2, 6, 8]. We show, for CQM's energy levels, the quark-antiquark value taking into account the one-gluon exchange potential $\mathcal{O}(\alpha_s)$ and including its one-loop correction $\mathcal{O}(\alpha_s^2)$. For experiment, we distinguish between well established states ([2]) and those levels which still need confirmation and so have been omitted from the summary table ([2]*).

Two charmed mesons with quantum numbers $J^P = 0^$ have been experimentally observed, D and $D_0(2550)$. The first one is the ground level of charmed mesons and it is well established in the RPP of PDG [2]. The second is still omitted from the summary table because even though two experiments observed this state its mass is slightly different. Our theoretical predictions are in reasonable agreement with experiment, confirming that there should be a first excitation of the D-meson at around 2550 MeV; note that the experimental masses measured until now go from 2518 to 2580 MeV for this state. Another important feature to highlight is that $\mathcal{O}(\alpha_s^2)$ OGE corrections are zero in this J^P -channel and thus our naïve quark model must predict correctly these two states from the original global fit of hadron phenomenology.

There are four *P*-wave states measured experimentally and denoted in the RPP of PDG as $D_0^*(2300)$, $D_1(2420)$, $D_1(2430)$, $D_2^*(2460)$. Two of them, $D_0^*(2300)$

TABLE II. Charmed meson masses, in MeV, from constituent quark model (CQM) and experiment [2, 6, 8]. We show, for CQM's energy levels, the quark-antiquark value taking into account the one-gluon exchange potential $\mathcal{O}(\alpha_s)$ and including its one-loop correction $\mathcal{O}(\alpha_s^2)$. For experiment, we distinguish between well established states ([2]) and those levels which still need confirmation and so have been omitted from the summary table ([2]*).

| Meson | J^P | n | The. $\mathcal{O}(\alpha_s)$ | The. $\mathcal{O}(\alpha_s^2)$ | Exp. | Ref. |
|-------------------|---------|----------|------------------------------|--------------------------------|----------------------|-----------|
| D | 0^{-} | 1 | 1876 | 1876 | 1867.95 ± 0.27 | [2] |
| | | 2 | 2595 | 2595 | 2549 ± 19 | $[2]^{*}$ |
| | | 3 | 3154 | 3154 | | |
| D_0^* | 0^+ | 1 | 2416 | 2346 | 2343 ± 10 | [2] |
| | | 2 | 3033 | 2925 | | |
| | | 3 | 3366 | 3292 | | |
| D^* | 1^{-} | 1 | 2010 | 2008 | 2009.12 ± 0.04 | [2] |
| | | 2 | 2656 | 2654 | 2627 ± 10 | $[2]^{*}$ |
| | | 3 | 2835 | 2805 | $2781 \pm 18 \pm 13$ | $[2]^*$ |
| D_1 | 1^+ | 1 | 2433 | 2449 | 2422.1 ± 0.8 | [2] |
| | | 2 | 2495 | 2435 | 2412 ± 9 | [2] |
| | | 3 | 3008 | 3033 | | |
| D_2 | 2^{-} | 1 | 2758 | 2768 | 2747 ± 6 | [2]* |
| | | 2 | 2973 | 2961 | | |
| | | 3 | 3227 | 3234 | | |
| D_2^* | 2^+ | 1 | 2465 | 2476 | 2461.1 ± 0.7 | [2] |
| | | 2 | 3037 | 3059 | | |
| | | 3 | 3217 | 3204 | | |
| D_3^* | 3^{-} | 1 | 2767 | 2788 | 2763.1 ± 3.2 | [2] |
| 0 | | 2 | 3249 | 3260 | | |
| | | 3 | 3429 | 3422 | | |
| D_3 | 3^+ | 1 | 2887 | 2891 | | |
| | | 2 | 3222 | 3217 | | |
| | | 3 | 3408 | 3411 | | |
| $D^*(2640)^{\pm}$ | $?^{?}$ | | | | $2637\pm2\pm6$ | [8] |
| $D(3000)^{0}$ | ?' | | | | $3214 \pm 29 \pm 49$ | [6] |

and $D_1(2430)$, belong to the $j_q^P = \frac{1}{2}^+$ doublet of HQS symmetry and the other two, $D_1(2420)$ and $D_2^*(2460)$, pertain to the $j_q^P = \frac{3}{2}^+$ as shown in Table VI of Ref. [61]. This explains why the $D_1(2420)$ has a slightly higher mass than the $D_1(2430)$ despite their PDG's notation. As one can see in Table II, our theoretical results reproduce correctly the level ordering and they are also in global agreement with the experimental reported masses once the one-loop OGE corrections are incorporated. The addition of the $\mathcal{O}(\alpha_s^2)$ OGE corrections was proposed by Lakhina *et al.* in [62] motivated by the fact that in the one-loop computation there is a spin-dependent term which affects only to mesons with different flavor quarks and it is not negligible for *P*-wave states where theory and experiment find their most significant differences. We demonstrate herein that *naïve* quark models cannot reproduce *P*-wave charmed meson spectrum but, instead of resorting to more complicated solutions such as exotic hadron structures, one should investigate the possibility of having missed potential terms that may be relevant for this sector.

The partner of the *D*-meson which belongs to the $j_q^P = \frac{1}{2}^-$ doublet in heavy quark spin symmetry is the D^* meson. As one can see in Table II, there are three candidates: D^* , $D_1^*(2600)$ and $D_1^*(2760)$; the first one is well established in PDG the other two are omitted from the summary table since they need confirmation. The 1-loop OGE corrections are small to moderate in this channel producing mass shifts from 2 to 20 MeV; however, these corrections are in the correct direction when trying to get agreement between theory and experiment. One may state that our results for $J^P = 1^-$ channel agrees well with the experimental masses reported until now, despite of being a bit larger.

The RPP of PDG reports two more charmed mesons with well established spin-parity quantum numbers, the $D_2(2740)$ and $D_3^*(2750)$ mesons. The first one is omitted from the summary table whereas the second is a well established charmed meson. Theoretically, both are dominant *D*-wave states whose masses are close to the experimental measurements; therefore, we may confirm that the experimental assignment is plausible. When incorporating the $\mathcal{O}(\alpha_s^2)$ OGE corrections, the theoretical masses of these states grow moderately, moving away from the experimental figures, but the change is not dramatic.

Focusing now on the two states whose quantum numbers have not been assigned (see last part of Table II). The $D^*(2640)^{\pm}$ seems to have a mass similar to the expected one for the first excitation of the D^* meson. In fact, there is no other possible case attending to the mass only. The $D(3000)^0$, whose mass is actually $(3214 \pm 29 \pm 49)$ MeV, could be fitted as the first excitation of either D_3^* or D_3 , but could be also assigned as the second excitation of either D_2 or D_2^* mesons.

In summary, the *naïve* constituent quark model is not able to reproduce the spectrum of charmed mesons. However, there are higher-order terms of the gluon exchange potential that seem to be very significant in those channels of charmed mesons where there is a larger discrepancy between theory and experiment. As can be guess from the discussion so far, and see in Fig. 1, when these higher-order interaction terms are included in the model, the spectrum of charmed mesons is described reasonably well. This does not mean that more complex structures such as tetraquark degrees of freedom in the meson's wave function cannot play a role but before resorting to such exotic solutions one should explore simpler refinements. Finally, Fig. 1 also show schematically our theoretical predictions for higher excited states of charmed mesons which should serve as a template in order to find them in future high energy nuclear and particle experiments.



FIG. 1. Charmed meson spectrum from constituent quark model (CQM) and experiment [2]. We show, for CQM's energy levels, the quark-antiquark value taking into account the one-gluon exchange potential $\mathcal{O}(\alpha_s)$ and including its one-loop correction $\mathcal{O}(\alpha_s^2)$. For experiment, we distinguish between well established states (purple bands) and those levels which still need confirmation and thus they have been omitted from the summary table (pink bands). The vertical extension of the experimental band is given by the experimental uncertainty.

IV. SUMMARY

We have evaluated the effectiveness of one-loop corrections to the one-gluon exchange potential in describing the spectrum of charmed mesons within a wellestablished constituent quark model. By incorporating these corrections, particularly spin-dependent terms that mainly affect P-wave states of mesons with different flavor quarks, the model aims to bridge gaps between theoretical predictions and experimental measurements across the charmed meson spectrum. The study investigates both well-established and recently observed states listed by the Particle Data Group.

The model successfully reproduces many S-wave state masses, including the ground state charmed mesons Dand D^* , as well as excited P and D wave states. Notably, P-wave states initially posed significant discrepancies with naive quark model predictions. However, incorporating one-loop corrections to the OGE potential reduced these differences, aligning theoretical predictions more closely with observed values. This adjustment suggests that refinements of the *naïve* constituent quark model can be effectively reproduce the charmed meson spectrum without resorting to exotic configurations, such as tetraquarks or meson-meson molecules.

Overall, this enhanced CQM provides a refined framework for describing the charmed meson spectrum, offering insight into both the nature of charmed mesons and the dynamics governing their mass structure. The results set a foundation for predicting higher-excited charmed states, potentially guiding future experimental searches and broadening the understanding of charmed meson interactions.

ACKNOWLEDGMENTS

Work partially financed by the Escuela Politécnica Nacional under projects PIS-22-04 and PIM 23-01; EU Horizon 2020 research and innovation program, STRONG-2020 project, under grant agreement no. 824093; Ministerio Español de Ciencia e Innovación under grant nos. PID2022-141910NB-I00 and PID2022-140440NB-C22; Junta de Andalucía under contract no. PAIDI FQM-370 and and PCI+D+i under the title: "Tecnologías avanzadas para la exploración del universo y sus componentes" (Code AST22-0001).

- N. Isgur and M. B. Wise, Spectroscopy with heavy quark symmetry, Phys. Rev. Lett. 66, 1130 (1991).
- [2] S. Navas *et al.* (Particle Data Group), Review of particle physics, Phys. Rev. D **110**, 030001 (2024).
- [3] R. Aaij *et al.* (LHCb), Dalitz plot analysis of $B^0 \rightarrow \overline{D}^0 \pi^+ \pi^-$ decays, Phys. Rev. D **92**, 032002 (2015), arXiv:1505.01710 [hep-ex].
- [4] R. Aaij *et al.* (LHCb), Amplitude analysis of $B^- \rightarrow D^+\pi^-\pi^-$ decays, Phys. Rev. D **94**, 072001 (2016), arXiv:1608.01289 [hep-ex].
- [5] P. del Amo Sanchez *et al.* (BaBar), Observation of new resonances decaying to $D\pi$ and $D^*\pi$ in inclusive e^+e^- collisions near $\sqrt{s} = 10.58$ GeV, Phys. Rev. D 82, 111101 (2010), arXiv:1009.2076 [hep-ex].
- [6] R. Aaij *et al.* (LHCb), Study of D_J meson decays to $D^+\pi^-$, $D^0\pi^+$ and $D^{*+}\pi^-$ final states in pp collision, JHEP **09**, 145, arXiv:1307.4556 [hep-ex].
- [7] R. Aaij *et al.* (LHCb), First observation and amplitude analysis of the $B^- \rightarrow D^+ K^- \pi^-$ decay, Phys. Rev. D **91**, 092002 (2015), [Erratum: Phys.Rev.D **93**, 119901 (2016)], arXiv:1503.02995 [hep-ex].
- [8] P. Abreu *et al.* (DELPHI), First evidence for a charm radial excitation, D*-prime, Phys. Lett. B 426, 231 (1998).
- [9] G. Abbiendi *et al.* (OPAL), A Search for a narrow radial excitation of the D*+- meson, Eur. Phys. J. C 20, 445 (2001), arXiv:hep-ex/0101045.
- [10] S. Chekanov *et al.* (ZEUS), Production of excited charm and charm-strange mesons at HERA, Eur. Phys. J. C 60, 25 (2009), arXiv:0807.1290 [hep-ex].
- [11] S. Godfrey and N. Isgur, Mesons in a Relativized Quark Model with Chromodynamics, Phys. Rev. D 32, 189 (1985).
- [12] J. Zeng, J. W. Van Orden, and W. Roberts, Heavy mesons in a relativistic model, Phys. Rev. D 52, 5229 (1995), arXiv:hep-ph/9412269.
- [13] S. N. Gupta and J. M. Johnson, Quantum chromodynamic potential model for light heavy quarkonia and the heavy quark effective theory, Phys. Rev. D 51, 168 (1995), arXiv:hep-ph/9409432.
- [14] T. A. Lahde, C. J. Nyfalt, and D. O. Riska, Spectra and M1 decay widths of heavy light mesons, Nucl. Phys. A 674, 141 (2000), arXiv:hep-ph/9908485.
- [15] D. Mohler and R. M. Woloshyn, D and D_s meson spectroscopy, Phys. Rev. D 84, 054505 (2011), arXiv:1103.5506 [hep-lat].
- [16] G. Moir, M. Peardon, S. M. Ryan, C. E. Thomas, and L. Liu, Excited spectroscopy of charmed mesons from lattice QCD, JHEP 05, 021, arXiv:1301.7670 [hep-ph].
- [17] M. Kalinowski and M. Wagner, Masses of D mesons, D_s mesons and charmonium states from twisted mass lattice QCD, Phys. Rev. D 92, 094508 (2015), arXiv:1509.02396 [hep-lat].
- [18] K. Cichy, M. Kalinowski, and M. Wagner, Continuum limit of the *D* meson, D_s meson and charmonium spectrum from $N_f = 2+1+1$ twisted mass lattice QCD, Phys. Rev. D **94**, 094503 (2016), arXiv:1603.06467 [hep-lat].
- [19] L. Gayer, N. Lang, S. M. Ryan, D. Tims, C. E. Thomas, and D. J. Wilson (Hadron Spectrum), Isospin-1/2 $D\pi$ scattering and the lightest D_0^* resonance from lattice QCD, JHEP 07, 123, arXiv:2102.04973 [hep-lat].
- [20] M. Albaladejo, P. Fernandez-Soler, F.-K. Guo, and

J. Nieves, Two-pole structure of the $D_0^*(2400)$, Phys. Lett. B **767**, 465 (2017), arXiv:1610.06727 [hep-ph].

- [21] M.-L. Du, M. Albaladejo, P. Fernández-Soler, F.-K. Guo, C. Hanhart, U.-G. Meißner, J. Nieves, and D.-L. Yao, Towards a new paradigm for heavy-light meson spectroscopy, Phys. Rev. D 98, 094018 (2018), arXiv:1712.07957 [hep-ph].
- [22] M.-L. Du, F.-K. Guo, C. Hanhart, B. Kubis, and U.-G. Meißner, Where is the lightest charmed scalar meson?, Phys. Rev. Lett. **126**, 192001 (2021), arXiv:2012.04599 [hep-ph].
- [23] Z.-G. Wang, Analysis of strong decays of the charmed mesons D(2550), D(2600), D(2750) and D(2760), Phys. Rev. D 83, 014009 (2011), arXiv:1009.3605 [hep-ph].
- [24] Z.-G. Wang, Analysis of strong decays of the charmed mesons $D_J(2580), D_J^*(2650), D_J(2740), D_J^*(2760), D_J(3000), D_J^*(3000),$ Phys. Rev. D 88, 114003 (2013), arXiv:1308.0533 [hepph].
- [25] D.-M. Li, B. Ma, and Y.-H. Liu, Understanding masses of c anti-s states in Regge phenomenology, Eur. Phys. J. C 51, 359 (2007), arXiv:hep-ph/0703278.
- [26] J.-K. Chen, Regge trajectories for the mesons consisting of different quarks, Eur. Phys. J. C 78, 648 (2018).
- [27] M. Di Pierro and E. Eichten, Excited Heavy Light Systems and Hadronic Transitions, Phys. Rev. D 64, 114004 (2001), arXiv:hep-ph/0104208.
- [28] S. Godfrey, Properties of the charmed P-wave mesons, Phys. Rev. D 72, 054029 (2005), arXiv:hep-ph/0508078.
- [29] F. E. Close and E. S. Swanson, Dynamics and decay of heavy-light hadrons, Phys. Rev. D 72, 094004 (2005), arXiv:hep-ph/0505206.
- [30] F. E. Close, C. E. Thomas, O. Lakhina, and E. S. Swanson, Canonical interpretation of the D(sJ)(2860) and D(sJ)(2690), Phys. Lett. B 647, 159 (2007), arXiv:hepph/0608139.
- [31] B. Zhang, X. Liu, W.-Z. Deng, and S.-L. Zhu, $D_{sJ}(2860)$ and $D_{sJ}(2715)$, Eur. Phys. J. C **50**, 617 (2007), arXiv:hep-ph/0609013.
- [32] W. Wei, X. Liu, and S.-L. Zhu, D wave heavy mesons, Phys. Rev. D 75, 014013 (2007), arXiv:hep-ph/0612066.
- [33] D. Ebert, R. N. Faustov, and V. O. Galkin, Heavy-light meson spectroscopy and Regge trajectories in the relativistic quark model, Eur. Phys. J. C 66, 197 (2010), arXiv:0910.5612 [hep-ph].
- [34] Q.-T. Song, D.-Y. Chen, X. Liu, and T. Matsuki, Higher radial and orbital excitations in the charmed meson family, Phys. Rev. D 92, 074011 (2015), arXiv:1503.05728 [hep-ph].
- [35] J. Ferretti and E. Santopinto, Open-flavor strong decays of open-charm and open-bottom mesons in the ${}^{3}P_{0}$ model, Phys. Rev. D **97**, 114020 (2018), arXiv:1506.04415 [hep-ph].
- [36] J.-B. Liu and C.-D. Lu, Spectra of heavy–light mesons in a relativistic model, Eur. Phys. J. C 77, 312 (2017), arXiv:1605.05550 [hep-ph].
- [37] T. Barnes, F. E. Close, and H. J. Lipkin, Implications of a DK molecule at 2.32-GeV, Phys. Rev. D 68, 054006 (2003), arXiv:hep-ph/0305025.
- [38] E. van Beveren and G. Rupp, Observed $D_s(2317)$ and tentative D(2100-2300) as the charmed cousins of the

light scalar nonet, Phys. Rev. Lett. **91**, 012003 (2003),

- arXiv:hep-ph/0305035.
 [39] J. Vijande, F. Fernandez, and A. Valcarce, Open-charm meson spectroscopy, Phys. Rev. D 73, 034002 (2006), [Erratum: Phys.Rev.D 74, 059903 (2006)], arXiv:hep-ph/0601143.
- [40] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T.-M. Yan, Charmonium: The Model, Phys. Rev. D 17, 3090 (1978), [Erratum: Phys.Rev.D 21, 313 (1980)].
- [41] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T.-M. Yan, Charmonium: Comparison with Experiment, Phys. Rev. D 21, 203 (1980).
- [42] S. N. Gupta, S. F. Radford, and W. W. Repko, Quarkonium Spectra and Quantum Chromodynamics, Phys. Rev. D 26, 3305 (1982).
- [43] S. N. Gupta, S. F. Radford, and W. W. Repko, Quantum Chromodynamic Potential Model for Light and Heavy Quarkonia, Phys. Rev. D 28, 1716 (1983).
- [44] S. N. Gupta, S. F. Radford, and W. W. Repko, b anti-b SPECTROSCOPY, Phys. Rev. D 30, 2424 (1984).
- [45] S. N. Gupta, S. F. Radford, and W. W. Repko, Semirelativistic Potential Model for Charmonium, Phys. Rev. D 31, 160 (1985).
- [46] W. Kwong, P. B. Mackenzie, R. Rosenfeld, and J. L. Rosner, Quarkonium Annihilation Rates, Phys. Rev. D 37, 3210 (1988).
- [47] W. Kwong and J. L. Rosner, D Wave Quarkonium Levels of the Υ Family, Phys. Rev. D 38, 279 (1988).
- [48] J. Vijande, F. Fernandez, and A. Valcarce, Constituent quark model study of the meson spectra, J. Phys. G 31, 481 (2005), arXiv:hep-ph/0411299.
- [49] J. Segovia, A. M. Yasser, D. R. Entem, and F. Fernandez, JPC=1– hidden charm resonances, Phys. Rev. D 78, 114033 (2008).
- [50] J. Segovia, P. G. Ortega, D. R. Entem, and F. Fernández, Bottomonium spectrum revisited, Phys. Rev. D 93, 074027 (2016), arXiv:1601.05093 [hep-ph].
- [51] J. Segovia, D. R. Entem, and F. Fernandez, Charmedstrange Meson Spectrum: Old and New Problems, Phys. Rev. D 91, 094020 (2015), arXiv:1502.03827 [hep-ph].
- [52] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernandez, Spectroscopy of B_c mesons and the possibility of finding exotic B_c-like structures, Eur. Phys. J. C 80, 223 (2020), arXiv:2001.08093 [hep-ph].
- [53] J. Segovia, D. R. Entem, and F. Fernández, Scaling of the ³P₀ Strength in Heavy Meson Strong Decays, Phys. Lett. B **715**, 322 (2012), arXiv:1205.2215 [hep-ph].
- [54] J. Segovia, A. M. Yasser, D. R. Entem, and F. Fernandez, Ds-1 (2536) + decays and the properties of Pwave charmed strange mesons, Phys. Rev. D 80, 054017 (2009).
- [55] J. Segovia, D. R. Entem, and F. Fernandez, Strong char-

monium decays in a microscopic model, Nucl. Phys. A **915**, 125 (2013), arXiv:1301.2592 [hep-ph].

- [56] J. Segovia, D. R. Entem, and F. Fernández, Puzzles in hadronic transitions of heavy quarkonium with two pion emission, Phys. Rev. D 91, 014002 (2015), arXiv:1409.7079 [hep-ph].
- [57] J. Segovia, F. Fernandez, and D. R. Entem, The Role of Spin-Flipping Terms in Hadronic Transitions of $\Upsilon(4S)$, Few Body Syst. **57**, 275 (2016), arXiv:1507.01607 [hep-ph].
- [58] B. Martín-González, P. G. Ortega, D. R. Entem, F. Fernández, and J. Segovia, Toward the discovery of novel Bc states: Radiative and hadronic transitions, Phys. Rev. D 106, 054009 (2022), arXiv:2205.05950 [hepph].
- [59] J. Segovia, D. R. Entem, and F. Fernandez, Charmonium resonances in e+ e- exclusive reactions around the psi(4415) region, Phys. Rev. D 83, 114018 (2011).
- [60] J. Segovia, C. Albertus, D. R. Entem, F. Fernandez, E. Hernandez, and M. A. Perez-Garcia, Semileptonic B and B_s decays into orbitally excited charmed mesons, Phys. Rev. D 84, 094029 (2011), arXiv:1107.4248 [hepph].
- [61] J. Segovia, C. Albertus, E. Hernandez, F. Fernandez, and D. R. Entem, Nonleptonic $B \rightarrow D^{(*)}D_{sJ}^{(*)}$ decays and the nature of the orbitally excited charmed-strange mesons, Phys. Rev. D 86, 014010 (2012), arXiv:1203.4362 [hep-ph].
- [62] O. Lakhina and E. S. Swanson, A Canonical Ds(2317)?, Phys. Lett. B 650, 159 (2007), arXiv:hep-ph/0608011.
- [63] S. N. Gupta and S. F. Radford, Quark Quark and Quark - Anti-quark Potentials, Phys. Rev. D 24, 2309 (1981).
- [64] J. Segovia, D. R. Entem, F. Fernandez, and E. Hernandez, Constituent quark model description of charmonium phenomenology, Int. J. Mod. Phys. E 22, 1330026 (2013), arXiv:1309.6926 [hep-ph].
- [65] D. Diakonov, Instantons at work, Prog. Part. Nucl. Phys. 51, 173 (2003), arXiv:hep-ph/0212026.
- [66] J. Segovia, D. R. Entem, and F. Fernandez, Is chiral symmetry restored in the excited meson spectrum?, Phys. Lett. B 662, 33 (2008).
- [67] P. G. Ortega, J. Segovia, D. R. Entem, and F. Fernandez, Molecular components in P-wave charmedstrange mesons, Phys. Rev. D 94, 074037 (2016), arXiv:1603.07000 [hep-ph].
- [68] G. S. Bali, H. Neff, T. Duessel, T. Lippert, and K. Schilling (SESAM), Observation of string breaking in QCD, Phys. Rev. D 71, 114513 (2005), arXiv:heplat/0505012.
- [69] E. Hiyama, Y. Kino, and M. Kamimura, Gaussian expansion method for few-body systems, Prog. Part. Nucl. Phys. 51, 223 (2003).