

# PAIR CORRELATION OF ZEROS OF THE RIEMANN ZETA FUNCTION I: PROPORTIONS OF SIMPLE ZEROS AND CRITICAL ZEROS

SIEGFRED ALAN C. BALUYOT, DANIEL ALAN GOLDSTON, ADE IRMA SURIAJAYA,  
AND CAROLINE L. TURNAGE-BUTTERBAUGH

*Dedicated to Hugh Montgomery whose insight continues to enlighten and surprise*

ABSTRACT. Assuming the Riemann Hypothesis (RH), Montgomery proved a theorem in 1973 concerning the pair correlation of zeros of the Riemann zeta-function and applied this to prove that at least  $2/3$  of the zeros are simple. In earlier work we showed how to remove RH from Montgomery's theorem and, in turn, obtain results on simple zeros assuming conditions on the zeros that are weaker than RH. Here we assume a more general condition, namely that all the zeros  $\rho = \beta + i\gamma$  with  $T < \gamma \leq 2T$  are in a narrow vertical box centered on the critical line with width  $\frac{b}{\log T}$ . For simplicity, now assume that  $b \rightarrow 0$  as  $T \rightarrow \infty$ . Following Montgomery's method, we prove the generalization of Montgomery's result that at least  $2/3$  of zeros are simple, and also the new result that at least  $2/3$  of the zeros are on the critical line. We also use the pair correlation method to prove that at least  $1/3$  of the zeros are both simple and on the critical line, a result already known unconditionally. Our work thus shows that the pair correlation method can be used to prove results not only on the vertical distribution of zeros but also on their horizontal distribution.

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let  $s = \sigma + it$  with  $\sigma, t \in \mathbb{R}$ , and let  $\rho = \beta + i\gamma$  denote a nontrivial zero of the Riemann zeta-function  $\zeta(s)$  so that  $\beta > 0$  and  $\gamma \in \mathbb{R}$ . All such zeros lie within the *critical strip*  $\{s \in \mathbb{C} \mid 0 < \sigma < 1\}$  and are symmetric with respect to the *critical line*  $\{s \in \mathbb{C} \mid \sigma = 1/2\}$  and also the real axis. The yet unproved Riemann Hypothesis (RH) asserts that  $\beta = 1/2$  for all  $\rho$ .

The best unconditional results for zeros on the critical line are due to Pratt, Robles, Zaharescu, and Zeindler [PRZZ20] who proved that more than 41.7% of the zeros are on the critical line, and also that more than 40.7% of the zeros are on the critical line and simple. Assuming RH, Montgomery [Mon73] obtained  $2/3 = 66.\bar{6}\%$  of the zeros are simple, and soon after, Montgomery and Taylor [Mon75] improved this and showed more than 67.25% of the zeros are simple. Recently Chirre, Gonçalves, and de Laat [CGdL20] obtained by this method 67.9%. By a mollifier method Conrey, Ghosh, and Gonek [CGG98] showed on RH and an additional hypothesis that at least  $19/27 = 70.3\bar{7}0\bar{3}\%$  of the zeros are simple, and later Bui and Heath-Brown [BHB13] showed that this result holds on RH alone.

In the recent paper [BGSTB24], we partially remove the assumption of the Riemann Hypothesis in the pair correlation method, and with weaker assumptions on zeros of  $\zeta(s)$  obtain results on simple zeros. In this paper we show that the pair correlation method for finding simple zeros can be applied to find an equal proportion of zeros on the critical line. The reason behind this is so simple that, except for technicalities, we will give the proof in the next paragraph. The key is that zeros off the critical line come in pairs  $\rho$  and  $1 - \bar{\rho}$ , which are symmetric with respect to the critical line. The main technicality is that our method requires that all the zeros must be in a narrow vertical box centered on the critical line.

---

*Date:* January 27, 2025.

*2010 Mathematics Subject Classification.* 11M06, 11M26.

*Key words and phrases.* Riemann zeta-function, zeros, pair correlation, simple zeros.

Assuming RH, Montgomery applied the pair correlation method to prove that there are simple zeros by using the multiplicity weighted diagonal terms from the correlation of each zero  $\rho$  with itself. This sum is bounded by a kernel weighted sum over the differences of all the pairs of zeros. With a Fejér kernel this bound is  $\frac{4}{3}$  of the number of zeros, and from this bound one easily obtains that at least  $\frac{2}{3}$  of the zeros are simple. As we make explicit below, when RH is not assumed one also obtains *symmetric* diagonal terms from the correlations of each zero  $\rho$  with its symmetric pair  $1 - \bar{\rho}$ , which only occur for zeros off the critical line with  $\beta \neq 1/2$ . The same  $\frac{4}{3}$  of the number of zeros now bounds the sum of the diagonal terms plus the symmetric diagonal terms when we assume all the zeros are in a thin vertical box. Since the diagonal terms trivially have the number of zeros as a lower bound, the number of symmetric diagonal terms can be at most  $\frac{1}{3}$  of the number of zeros. Thus, at least  $\frac{2}{3}$  of the zeros must be on the critical line.

For  $b > 0$ , let  $B_b$  be the thin vertical box centered on the critical line defined by

$$(1.1) \quad B_b := \left\{ s = \sigma + it \mid \frac{1}{2} - \frac{b}{2 \log T} < \sigma < \frac{1}{2} + \frac{b}{2 \log T}, T < t \leq 2T \right\},$$

and let  $N(B_b)$  denote the number of zeros of  $\zeta(s)$  in  $B_b$ . Let  $N_1(B_b)$  count the number of zeros in  $B_b$  that are simple, and let  $N_0(B_b)$  count the number of zeros in  $B_b$  that are on the critical line. Finally, let  $N_{1,0}(B_b)$  denote the number of zeros in  $B_b$  that are simple and on the critical line. Note that in each of these settings, we count zeros with multiplicity.

**Theorem 1.** *Assume that, for all sufficiently large  $T$ , all the zeros  $\rho = \beta + i\gamma$  of  $\zeta(s)$  with  $T < \gamma \leq 2T$  are in  $B_b$ . Then we have, where  $b \rightarrow 0$  as  $T \rightarrow \infty$ ,*

$$(1.2) \quad N_1(B_b) = \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \rho \text{ simple}}} 1 \geq \left( \frac{2}{3} + o(1) \right) \sum_{\substack{\rho \\ T < \gamma \leq 2T}} 1 = \left( \frac{2}{3} + o(1) \right) N(B_b),$$

$$(1.3) \quad N_0(B_b) = \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta = \frac{1}{2}}} 1 \geq \left( \frac{2}{3} + o(1) \right) \sum_{\substack{\rho \\ T < \gamma \leq 2T}} 1 = \left( \frac{2}{3} + o(1) \right) N(B_b),$$

and

$$(1.4) \quad N_{1,0}(B_b) = \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta = \frac{1}{2} \text{ and } \rho \text{ simple}}} 1 \geq \left( \frac{1}{3} + o(1) \right) \sum_{\substack{\rho \\ T < \gamma \leq 2T}} 1 = \left( \frac{1}{3} + o(1) \right) N(B_b).$$

*Remark 1.* The result in (1.4) is known unconditionally and was first proved independently by Heath-Brown [HB79] and Selberg (unpublished), see [Tit86, Sections 10.28 and 10.29].

Using a more complicated kernel based on the kernel used in [Mon75], we can improve the numerical results obtained in Theorem 1. We state the results in the following theorem for two choices of fixed  $b$ , not simply as a statement of proportion, but rather utilizing the asymptotic formula for the number of zeros  $\rho$  with  $T < \gamma \leq 2T$ , which is given in (2.6) below.

**Theorem 2.** *Assume that, for all sufficiently large  $T$ , all the zeros  $\rho = \beta + i\gamma$  of  $\zeta(s)$  with  $T < \gamma \leq 2T$  are in  $B_b$ . Then as  $T \rightarrow \infty$ , we have*

$$(1.5) \quad \begin{aligned} N_1(B_{0.3185}) &\geq \left(0.66666908 + o(1)\right) \frac{T}{2\pi} \log T, \\ N_0(B_{0.3185}) &\geq \left(0.66666908 + o(1)\right) \frac{T}{2\pi} \log T, \\ N_{1,0}(B_{0.3185}) &\geq \left(0.33333816 + o(1)\right) \frac{T}{2\pi} \log T, \end{aligned}$$

and

$$(1.6) \quad \begin{aligned} N_1(B_{0.001}) &\geq \left(0.67250064 + o(1)\right) \frac{T}{2\pi} \log T, \\ N_0(B_{0.001}) &\geq \left(0.67250064 + o(1)\right) \frac{T}{2\pi} \log T, \\ N_{1,0}(B_{0.001}) &\geq \left(0.34500129 + o(1)\right) \frac{T}{2\pi} \log T. \end{aligned}$$

We prove Theorem 2 in the last section of this paper, and there we also include Table 1 which consists of similar results obtained for small values of  $b$ . The table also exhibits the fact that the results deteriorate as  $b$  increases. (The method ultimately fails when  $b \geq 4.2$  for (1.2) and (1.3) and when  $b \geq 2$  in (1.4).) It may seem counterintuitive that increasing the width  $b$  of the box  $B_b$  degrades the results, but the method requires positivity supplied by a kernel that is evaluated on the differences  $\rho - \rho'$  of pairs of zeros in  $B_b$ . The kernel puts less weight on pairs  $\rho = \rho'$  as  $b$  increases, which in turn decreases the lower bounds obtained in Theorem 2.

*Remark 2.* In our first paper [BGSTB24], we used the pair correlation method to obtain results on simple zeros and wrote “The method of proof neither requires nor provides any information on whether any of these zeros are or are not on the critical line where  $\beta = 1/2$ .” The results of Theorem 1 show that this statement is incorrect and that the pair correlation method can be used to prove results not only on the vertical distribution of zeros but also on their horizontal distribution.

## 2. UNCONDITIONAL MONTGOMERY THEOREM

In [Mon73], Montgomery introduced the subject of pair correlation of zeros of the zeta-function. He studied, for  $x > 0$  and  $T \geq 3$ ,

$$(2.1) \quad F(x, T) := \sum_{\substack{\rho, \rho' \\ 0 < \gamma, \gamma' \leq T}} x^{i(\gamma - \gamma')} w(\gamma - \gamma'), \quad \text{where} \quad w(u) := \frac{4}{4 + u^2}.$$

Here the zeros are counted with multiplicity. While the definition of  $F(x, T)$  does not depend on RH, the asymptotic formula Montgomery obtained for  $F(x, T)$  when  $1 \leq x \leq T$  does depend on RH. This dependence, however, is only because RH allowed the real part of the zeros  $\beta$  to be removed out of  $F(x, T)$ . In [BGSTB24] we retained this dependence to obtain an unconditional form of Montgomery’s theorem. To state our result, let

$$(2.2) \quad \mathcal{F}(x, T) := \sum_{\substack{\rho, \rho' \\ T < \gamma, \gamma' \leq 2T}} x^{\rho - \rho'} W(\rho - \rho'), \quad \text{where} \quad W(u) := \frac{4}{4 - u^2}.$$

**Montgomery Theorem (MT).** *For  $x \geq 1$  and  $T \geq 3$ , we have  $\mathcal{F}(x, T) \geq 0$ ,  $\mathcal{F}(x, T) = \mathcal{F}(1/x, T)$ , and*

$$(2.3) \quad \mathcal{F}(x, T) = \frac{T}{2\pi x^2} \log^2 T \left(1 + O\left(\frac{1}{\sqrt{\log T}}\right)\right) + \frac{T}{2\pi} \log x + O(T\sqrt{\log T}),$$

uniformly for  $1 \leq x \leq T$ .

The statement above has been modified from its original formulation in [BGSTB24], with two changes. First, note that the sum  $\mathcal{F}(x, T)$  above is taken over zeros satisfying  $T < \gamma, \gamma' \leq 2T$ , where as in [BGSTB24] the sums is taken over zeros satisfying  $0 < \gamma, \gamma' \leq T$ . Secondly, and more importantly, the error terms appearing above have been corrected from those in the original theorem statement.\* The result stated in [BGSTB24], in contrast to (2.3) above, was

$$\mathcal{F}(x, T) = \left( \frac{T}{2\pi x^2} \log^2 T + \frac{T}{2\pi} \log x \right) \left( 1 + O\left( \frac{1}{\sqrt{\log T}} \right) \right).$$

Note that if, for example,  $x = c \log T$ , then

$$\mathcal{F}(x, T) = \frac{T}{2\pi} \log \log T + \frac{T}{2\pi} \left( \frac{1}{c^2} + \log c \right) + O\left( T \frac{\log \log T}{\sqrt{\log T}} \right),$$

and the factor of  $c$  on the  $T$  term is incorrect. In the corrected version above the error term  $O(T\sqrt{\log T})$  which now holds over the whole range  $1 \leq x \leq T$  absorbs all these terms in this range. We thank Ramūnas Garunkštis and Julija Paliulionytė for pointing out the mistake and for suggesting an argument to correct the issue. At the beginning of the next section, and before we prove Theorem 1, we explain how to correct the proof of [BGSTB24].

Before we prove Theorem MT, we record the following standard results. The Riemann von Mangoldt formula for  $N(T)$  states that

$$(2.4) \quad N(T) := \sum_{0 < \gamma \leq T} 1 = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + O(\log T).$$

From (2.4), we immediately have

$$(2.5) \quad N(T) = \frac{T}{2\pi} \log T + O(T) \quad \text{and} \quad N(T+1) - N(T) \ll \log T.$$

Thus, for example, we see that the sum on the right-hand side of (1.2), (1.3), and (1.4) is

$$(2.6) \quad \sum_{T < \substack{\rho \\ \gamma' \leq 2T}} 1 = N(2T) - N(T) = \frac{T}{2\pi} \log T + O(T).$$

Using the second estimate in (2.5), or directly as in [Dav00, Lemma, Ch. 15], we have

$$(2.7) \quad \sum_{\rho} \frac{1}{1 + (t - \gamma)^2} \ll \log(|t| + 2).$$

Using this estimate, we obtain for  $w(\gamma - \gamma')$  in (2.1),

$$(2.8) \quad \sum_{T < \substack{\rho, \rho' \\ \gamma, \gamma' \leq 2T}} w(\gamma - \gamma') \ll \sum_{0 < \gamma \leq 2T} \sum_{\gamma'} \frac{1}{1 + (\gamma - \gamma')^2} \ll \sum_{0 < \gamma \leq 2T} \log \gamma \leq N(2T) \log(2T) \ll T \log^2 T.$$

---

\*All the applications of Theorem 1 in [BGSTB24], such as Lemma 5 and Lemma 7, remain correct and analogous to those applications obtained using (2.3). The source of the error came from applying Lemma 8 of [GM87], which has the same issue but likewise no effect on any consequential results in that paper. We note that Montgomery and Vaughan (to appear) have obtained on RH a more refined version of Theorem 1 which deals correctly with the lower order terms which we absorb into a larger error term here.

### 3. PROOF OF **MT**

In this section we first correct the proof in [BGSTB24, Theorem 1] to obtain Theorem 1 when  $1 \leq \gamma, \gamma' \leq T$ . Next, using this result, we prove Theorem 1 for  $\mathcal{F}(x, T)$  when  $T < \gamma, \gamma' \leq 2T$ . (This could also be done by modifying each step in the first proof.)

*Proof of Theorem 1 in [BGSTB24].* Let  $I$  be an interval and

$$(3.1) \quad \mathcal{F}_I(x) := \sum_{\substack{\rho, \rho' \\ \gamma, \gamma' \in I}} x^{\rho - \rho'} W(\rho - \rho').$$

Thus we see from (2.2) that  $\mathcal{F}(x, T) = \mathcal{F}_{(T, 2T]}(x)$ . Then by (2.17), (2.18) of [BGSTB24] we have, for  $1 \leq x \leq T$ ,

$$(3.2) \quad \mathcal{F}_{(0, T]}(x) + O(T^{1/2}) + O(x) = \frac{1}{2\pi} R(x, T),$$

where by [BGSTB24, Lemma 1] and following the argument in [Mon73],

$$R(x, T) = \int_0^T |A_1(x, t) + A_2(x, t) + A_3(x, t)|^2 dt,$$

where

$$A_1(x, t) = \frac{\log(|t| + 2)}{x}, \quad A_2(x, t) = - \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^{\frac{1}{2} + it}} \min \left\{ \frac{n}{x}, \frac{x}{n} \right\},$$

$$A_3(x, t) \ll \frac{1}{x} + \frac{x^{\frac{1}{2}}}{(1 + t^2)} + \frac{x^{-\frac{5}{2}}}{|t| + 2}.$$

Now, for  $1 \leq x \leq T$ ,

$$M_1 := \int_0^T |A_1(x, t)|^2 dt = \frac{T}{x^2} (\log^2 T + O(\log T)),$$

and by [GM87, Lemma 7] or [Gol81, Lemma B],

$$M_2 := \int_0^T |A_2(x, t)|^2 dt = T \log x + O(T \sqrt{\log T}),$$

and

$$M_3 := \int_0^T |A_3(x, t)|^2 dt \ll \frac{T}{x^2} + x.$$

Using Cauchy-Schwarz, and an argument of Ramūnas Garunkštis and Julija Paliulionytė [GP24],

$$\begin{aligned} R(x, T) &= \int_0^T |A_1(x, t) + A_2(x, t)|^2 dt + O(\sqrt{M_1 M_3}) + O(\sqrt{M_2 M_3}) + O(M_3) \\ &= M_1 + M_2 + O\left(\left| \int_0^T A_1(x, t) \overline{A_2(x, t)} dt \right|\right) + O(T \sqrt{\log T}) + O\left(\frac{T}{x^2} \log T\right). \end{aligned}$$

Since for  $n \geq 2$

$$\int_0^T n^{it} \log(t + 2) dt \ll \frac{\log T}{\log n},$$

we have

$$\int_0^T A_1(x, t) \overline{A_2(x, t)} dt \ll \frac{\log T}{x} \sum_{n=2}^{\infty} \frac{\Lambda(n)}{n^{\frac{1}{2}} \log n} \min \left\{ \frac{n}{x}, \frac{x}{n} \right\} \ll \frac{\log T}{\sqrt{x} \log 2x} \ll \log T.$$

Thus, by (3.2), we have for  $1 \leq x \leq T$ ,

$$(3.3) \quad \begin{aligned} \mathcal{F}_{(0,T]}(x) &= \frac{T}{2\pi x^2}(\log^2 T + O(\log T)) + \frac{T}{2\pi} \log x + O(T\sqrt{\log T}) \\ &= \frac{T \log^2 T}{2\pi x^2} \left(1 + O\left(\frac{1}{\sqrt{\log T}}\right)\right) + \frac{T}{2\pi} \log x + O(T\sqrt{\log T}), \end{aligned}$$

where we have degraded the first error term insignificantly to simplify applications to sums over differences of zeros. Thus we have obtained **MT** for  $\mathcal{F}_{(0,T]}(x)$ .  $\square$

We now turn to proving **MT** for  $\mathcal{F}_{(T,2T]}(x)$ .

**Lemma 1.** *We have, for  $x > 0$  and  $T \geq 3$ ,*

$$(3.4) \quad \mathcal{F}_{(T,2T]}(x, T) = \frac{2}{\pi} \int_{-\infty}^{\infty} \left| \sum_{\substack{\rho \\ T < \gamma \leq 2T}} \frac{x^{\rho-1/2}}{1 - (\rho - (1/2 + it))^2} \right|^2 dt$$

and

$$(3.5) \quad F_{(T,2T]}(x, T) = \frac{2}{\pi} \int_{-\infty}^{\infty} \left| \sum_{\substack{\rho \\ T < \gamma \leq 2T}} \frac{x^{i\gamma}}{1 + (t - \gamma)^2} \right|^2 dt.$$

This was proved unconditionally as Lemma 3 in [BGSTB24] with the sum over terms with  $0 < \gamma \leq T$ . That proof works line by line for both (3.4) and (3.5).

*Proof of (2.3).* From (3.4) of Lemma 1 we see immediately that  $\mathcal{F}_{(T,2T]}(x, T)$  is real and non-negative, and since we can interchange  $\rho$  and  $\rho'$  in (2.2), we have  $\mathcal{F}_{(T,2T]}(1/x, T) = \mathcal{F}_{(T,2T]}(x, T)$ . Thus,  $\mathcal{F}(T^\alpha, T)$  is real, non-negative, and even. From (2.2) we have

$$\begin{aligned} \mathcal{F}_{(T,2T]}(x, T) &= \sum_{\substack{\rho, \rho' \\ \gamma, \gamma' \in (T, 2T]}} x^{\rho - \rho'} W(\rho - \rho') \\ &= \mathcal{F}_{(0,2T]}(x) - \mathcal{F}_{(0,T]}(x) + O\left( \sum_{\substack{\rho, \rho' \\ 0 < \gamma \leq T \\ T < \gamma' \leq 2T}} x^{|\beta - \beta'|} |W(\rho - \rho')| \right) \\ &= \frac{T \log^2 T}{2\pi x^2} \left(1 + O\left(\frac{1}{\sqrt{\log T}}\right)\right) + \frac{T}{2\pi} \log x \\ &\quad + O(T\sqrt{\log T}) + O\left( \sum_{\substack{\rho, \rho' \\ 0 < \gamma \leq T \\ T < \gamma' \leq 2T}} x^{|\beta - \beta'|} |W(\rho - \rho')| \right), \end{aligned}$$

where we have applied (3.3) for both  $\mathcal{F}_{(0,2T]}(x)$  and  $\mathcal{F}_{(0,T]}(x)$ . Note that zeros off the critical line in the upper half plane occur in pairs  $\rho = \beta + i\gamma$  and  $1 - \bar{\rho} = 1 - \beta + i\gamma$  with  $\beta \neq 1/2$ . Thus, letting

$$(3.6) \quad \Theta(t) := \max\{\beta : \rho = \beta + i\gamma, 0 < \gamma \leq t\},$$

we have

$$\max_{0 < \gamma, \gamma' \leq T} |\beta - \beta'| = 2(\max_{0 < \gamma \leq T} \beta) - 1 = 2\Theta(T) - 1.$$

Hence

$$E_1 := \sum_{\substack{\rho, \rho' \\ 0 < \gamma \leq T \\ T < \gamma' \leq 2T}} x^{|\beta - \beta'|} |W(\rho - \rho')| \ll x^{2\Theta(2T)-1} \sum_{\substack{\rho, \rho' \\ 0 < \gamma \leq T \\ T < \gamma' \leq 2T}} |W(\rho - \rho')|.$$

Using (2.5) and the bound

$$(3.7) \quad |W(\rho - \rho')| \ll w(\gamma - \gamma'),$$

which is (5.5) of [BGSTB24], we see that

$$\begin{aligned} E_1 &\ll x^{2\Theta(2T)-1} \sum_{0 < \gamma \leq T} \left( \sum_{T < \gamma' \leq 2T} w(\gamma - \gamma') \right) \\ &\ll x^{2\Theta(2T)-1} \sum_{0 < \gamma \leq T} \frac{\log T}{(T - \gamma) + 1} \\ &\ll x^{2\Theta(2T)-1} \log^3 T. \end{aligned}$$

Korobov [Kor58b, Kor58a] and Vinogradov [Vin58] (independently) proved that there is a zero-free region for points  $\sigma + it$  in the region in the complex plane determined by  $\sigma > 1 - \eta(t) \geq \Theta(t)$  with

$$\eta(t) = \frac{c}{(\log t)^{2/3} (\log \log t)^{1/3}}, \quad \text{for } t \geq 3,$$

and some constant  $c > 0$ . (It is well-known that there are no zeros in the region  $\sigma \geq 0$  and  $0 \leq t \leq 3$ .) Hence  $E_1 \ll x^{1-2\eta(2T)} \log^3 T$ . Thus, for  $T^{1/2} \leq x \leq T$ ,

$$E_1 \ll x^{1-2\eta(2T)} \log^3 T \ll x \exp\left(-c \frac{\log x}{(\log x)^{2/3} (\log \log x)^{1/3}}\right) \log^3 x \ll x,$$

while, for  $1 \leq x \leq T^{1/2}$ ,

$$E_1 \ll x^{1-2\eta(2T)} \log^3 T \ll T^{1/2}$$

since this error term is increasing in  $x$  and thus we may take  $x = T^{1/2}$  in the previous bound. Since for  $1 \leq x \leq T$ , we have  $E_1 \ll T^{1/2} + x \ll T \ll T\sqrt{\log T}$ , we conclude

$$\mathcal{F}_{(T, 2T]}(x, T) = \frac{T \log^2 T}{2\pi x^2} \left(1 + O\left(\frac{1}{\sqrt{\log T}}\right)\right) + \frac{T}{2\pi} \log x + O(T\sqrt{\log T}).$$

□

#### 4. EVALUATION OF A SUM OVER DIFFERENCES OF ZEROS BY MONTGOMERY'S THEOREM

We can use Montgomery's theorem to evaluate many sums over the differences  $\rho - \rho'$ . If RH is assumed then  $\rho - \rho' = i(\gamma - \gamma')$  and we can use the usual real Fourier transform. Thus letting  $r$  be in  $L^1$ , then  $\widehat{r}$ , the Fourier transform of  $r$ , is well defined and if it is also in  $L^1$  we have

$$(4.1) \quad r(\alpha) = \int_{-\infty}^{\infty} \widehat{r}(t) e(\alpha t) dt, \quad \widehat{r}(t) := \int_{-\infty}^{\infty} r(\alpha) e(-t\alpha) d\alpha, \quad \text{where } e(u) := e^{2\pi i u}.$$

To prove (1.2) on RH, Montgomery [Mon73] used the Fejér kernel

$$(4.2) \quad j_F(\alpha) = \max\{1 - |\alpha|, 0\}, \quad \widehat{j}_F(w) = \left(\frac{\sin \pi t}{\pi t}\right)^2.$$

For improved numerical results, we will also use the Montgomery-Taylor kernel [Mon75, CG93]

$$(4.3) \quad j_M(\alpha) = \frac{1}{1 - \cos \sqrt{2}} \left( \frac{1}{2\sqrt{2}} \sin(\sqrt{2}j_F(\alpha)) + \frac{1}{2}j_F(\alpha) \cos(\sqrt{2}\alpha) \right),$$

with  $j_F(\alpha)$  from (4.2), and

$$(4.4) \quad \widehat{j}_M(w) = \frac{1}{1 - \cos \sqrt{2}} \left( \frac{\sin(\frac{1}{2}(\sqrt{2} - 2\pi w))}{\sqrt{2} - 2\pi w} + \frac{\sin(\frac{1}{2}(\sqrt{2} + 2\pi w))}{\sqrt{2} + 2\pi w} \right)^2.$$

In the formulas that follow we will take the kernel  $j(\alpha)$  to be either  $j = j_F$  or  $j = j_M$ .

Next, without assuming RH we use the complex Fourier transform, with  $z \in \mathbb{C}$ , and define

$$(4.5) \quad \widehat{K}_b(t) := \frac{j(2\pi t)}{\cosh(2\pi bt)},$$

and thus

$$(4.6) \quad K_b(z) = \int_{-\infty}^{\infty} \widehat{K}_b(t) e^{zt} dt = \int_{-\frac{1}{2\pi}}^{\frac{1}{2\pi}} \frac{j(2\pi t)}{\cosh(2\pi bt)} e^{2\pi izt} dt = \frac{1}{2\pi} \int_{-1}^1 \frac{j(\alpha)}{\cosh(b\alpha)} e^{iz\alpha} d\alpha.$$

Note that  $K_b(z)$  is an analytic function for all  $z$  since the integral defining it is over a finite interval [Tit39, 2.83]. Also, note that if  $b = 0$  then  $\widehat{K}_0(t) = j(2\pi t)$ . We call  $K_b(z)$  a Tsang kernel [Tsa93].

We now take  $z = -i(\rho - \rho') \log T$  in (4.6) and have

$$K_b(-i(\rho - \rho') \log T) = \int_{-\infty}^{\infty} \frac{j(2\pi t)}{\cosh(2\pi bt)} T^{2\pi(\rho - \rho')t} dt = \frac{1}{2\pi} \int_{-1}^1 \frac{j(\alpha)}{\cosh(b\alpha)} T^{\alpha(\rho - \rho')} d\alpha.$$

Multiplying by  $W(\rho - \rho')$  and summing over pairs of zeros  $\rho, \rho'$  with  $T < \gamma, \gamma' \leq 2T$ , we obtain by the definition (2.2) that

$$(4.7) \quad \sum_{\substack{\rho, \rho' \\ T < \gamma, \gamma' \leq 2T}} K_b(-i(\rho - \rho') \log T) W(\rho - \rho') = \frac{1}{2\pi} \int_{-1}^1 \frac{j(\alpha)}{\cosh(b\alpha)} \mathcal{F}(T^\alpha, T) d\alpha.$$

By Montgomery's Theorem (2.3) we see that  $\mathcal{F}(T^\alpha, T)$  is an even function of  $\alpha$ , and the right-hand side of (4.7) is equal to

$$\left( \frac{1}{\pi} \int_0^1 \left( \frac{j(\alpha)}{\cosh(b\alpha)} T^{-2\alpha} \log T \left( 1 + O\left( \frac{1}{\sqrt{\log T}} \right) \right) + \frac{\alpha j(\alpha)}{\cosh(b\alpha)} \right) d\alpha \right) \frac{T}{2\pi} \log T + O(T\sqrt{\log T}).$$

Since

$$\begin{aligned} \frac{1}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} T^{-2\alpha} \log T d\alpha &= \frac{1}{\pi} \left( \int_0^{\frac{\log \log T}{\log T}} + \int_{\frac{\log \log T}{\log T}}^1 \right) \frac{j(\alpha)}{\cosh(b\alpha)} T^{-2\alpha} \log T d\alpha \\ &= \left( j(0) + O\left( \frac{\log \log T}{\log T} \right) \right) \frac{1}{\pi} \int_0^{\frac{\log \log T}{\log T}} T^{-2\alpha} \log T d\alpha + O\left( \frac{1}{\log T} \right) = \frac{1}{2\pi} j(0) + O\left( \frac{\log \log T}{\log T} \right), \end{aligned}$$

we conclude that

$$(4.8) \quad \sum_{\substack{\rho, \rho' \\ T < \gamma, \gamma' \leq 2T}} K_b(-i(\rho - \rho') \log T) W(\rho - \rho') = \frac{1}{2\pi} \left( j(0) + 2 \int_0^1 \frac{\alpha j(\alpha)}{\cosh(b\alpha)} d\alpha + O\left( \frac{1}{\sqrt{\log T}} \right) \right) \frac{T}{2\pi} \log T.$$



## 5. THE TSANG KERNEL

To obtain the main properties of the Tsang kernel, we need the following Fourier Transform pair.

**Lemma 2.** *Let  $t, x$  be real variables, and take real numbers  $y$  and  $b$  with  $|y| < b$ . Then we have the Fourier Transform pair*

$$(5.1) \quad h_{y,b}(t) = \frac{\cosh(2\pi yt)}{\cosh(2\pi bt)} \quad \text{and} \quad \widehat{h}_{y,b}(x) = \frac{1}{b} \left( \frac{\cos \frac{\pi y}{2b} \cosh \frac{\pi x}{2b}}{\cos \frac{\pi y}{b} + \cosh \frac{\pi x}{b}} \right) > 0.$$

*Proof of Lemma 2.* By [SS03, Example 3, p. 81], a residue calculation gives

$$\int_{-\infty}^{\infty} \frac{e(-xw)}{\cosh \pi x} dx = \frac{1}{\cosh \pi w},$$

so that  $1/\cosh \pi x$  is its own Fourier transform. Letting  $x = 2bt$  and  $w = z/2b$ , where  $z$  is real, then we have  $xw = zt$ , and taking the real part of both sides of the above equation, we have

$$(5.2) \quad \int_{-\infty}^{\infty} \frac{\cos(2\pi zt)}{\cosh(2\pi bt)} dt = \frac{1}{2b \cosh \frac{\pi z}{2b}}.$$

While this was derived with  $z$  real, if we now take  $z = x + iy$  with  $|y| < b$  we see the integral is absolutely convergent, and therefore each side of this equation is well defined and equals the same analytic function in the horizontal strip  $|y| < b$  in the complex plane.

Recalling the formulas

$\cos z = \cos(x + iy) = \cos x \cosh y - i \sin x \sinh y$ ,  $\cosh z = \cosh(x + iy) = \cosh x \cos y + i \sinh x \sin y$ , we have  $\operatorname{Re}(\cos z) = \cos x \cosh y$ ,  $\operatorname{Re}(\cosh z) = \cosh x \cos y$ , and

$$\operatorname{Re} \left( \frac{1}{\cosh z} \right) = \frac{\operatorname{Re}(\cosh \bar{z})}{|\cosh z|^2} = \frac{\cosh x \cos y}{\cos^2 y + \sinh^2 x} = \frac{\cos y \cosh x}{\frac{1 + \cos 2y}{2} + \frac{\cosh 2x - 1}{2}} = \frac{2 \cos y \cosh x}{\cos 2y + \cosh 2x}.$$

Then, by the above formulas and using that  $h_{y,b}(t)$  is even in the second line below, we have

$$\begin{aligned} \widehat{h}_{y,b}(x) &= \int_{-\infty}^{\infty} e(-xt) h_{y,b}(t) dt \\ &= \int_{-\infty}^{\infty} \frac{\cos(2\pi xt) \cosh(2\pi yt)}{\cosh(2\pi bt)} dt \\ &= \operatorname{Re} \int_{-\infty}^{\infty} \frac{\cos(2\pi zt)}{\cosh(2\pi bt)} dt \\ &= \operatorname{Re} \frac{1}{2b \cosh \frac{\pi z}{2b}} \\ &= \frac{1}{b} \left( \frac{\cos \frac{\pi y}{2b} \cosh \frac{\pi x}{2b}}{\cos \frac{\pi y}{b} + \cosh \frac{\pi x}{b}} \right) > 0. \end{aligned}$$

□

*Remark 3.* Tsang [Tsa93] uses  $h_y(t)$  which is the case  $b = 1$  of the formula above and obtains  $\widehat{h}_y(x)$  from [EMOT54, p. 31, (12)], but this is for half of the Fourier transform formula above after a change of variable. Thus Tsang's  $\widehat{h}_y(x)$  needs to be doubled, but this has no effect on anything in his paper since only the property  $\widehat{h}_y(x) > 0$  is used there.

We next use this lemma to obtain the following lemma, which Tsang [Tsa93, Lemma 1] proved with  $b = 1$ .

**Lemma 3** (K.-M. Tsang). *For fixed  $0 < b \ll 1$ , the kernel  $K_b(z)$  defined by (4.5) and (4.6) is an even entire function for  $z = x + iy$ ,  $x, y \in \mathbb{R}$ , and we have*

$$(5.3) \quad \operatorname{Re} K_b(x + iy) > 0 \quad \text{for all } x \text{ and } |y| < b.$$

Also, for  $z \in \mathbb{C}$ , we have

$$(5.4) \quad K_b(z) \ll \frac{e^{|\operatorname{Im}(z)|}}{1 + |z|^2}.$$

*Proof of Lemma 3.* By (4.6), we have

$$(5.5) \quad K_b(z) = \int_{-\frac{1}{2\pi}}^{\frac{1}{2\pi}} \frac{j(2\pi t)}{\cosh(2\pi bt)} e^{2\pi izt} dt = 2 \int_0^{\frac{1}{2\pi}} \frac{j(2\pi t)}{\cosh(2\pi bt)} \cos(2\pi zt) dt.$$

Letting  $z = x + iy$  and using  $\operatorname{Re}(\cos(u + iv)) = \cos u \cosh v$  for real  $u$  and  $v$ , we have

$$(5.6) \quad \operatorname{Re} K_b(x + iy) = 2 \int_0^{\frac{1}{2\pi}} \frac{j(2\pi t) \cosh(2\pi yt)}{\cosh(2\pi bt)} \cos(2\pi xt) dt = 2 \int_0^{\frac{1}{2\pi}} j(2\pi t) h_{y,b}(t) \cos(2\pi xt) dt.$$

Hence, by Lemma 2, for  $|y| < b$ ,

$$(5.7) \quad \operatorname{Re} K_b(x + iy) = (j(2\pi t) h_{y,b}(t))^\wedge(x) = \widehat{j(2\pi t)} * \widehat{h_{y,b}(t)}(x).$$

Since  $\widehat{j} \geq 0$  by (4.2) and (4.4) and  $\widehat{h_{y,b}} > 0$  by (5.1), we see the convolution is positive and (5.3) follows. We now prove (5.4). By (5.5) with the change of variable  $\alpha = 2\pi t$ , we have

$$(5.8) \quad K_b(z) = \frac{1}{\pi} \int_0^1 J_b(\alpha) \cos(z\alpha) d\alpha, \quad J_b(\alpha) := \frac{j(\alpha)}{\cosh(b\alpha)}.$$

If  $|z| \leq 1$ , then  $|K_b(z)| \ll \int_0^1 j(\alpha) d\alpha \ll 1$ , which proves the estimate in this range. If  $|z| > 1$ , then by integration by parts

$$\begin{aligned} K_b(z) &= \frac{\sin z\alpha}{\pi z} J_b(\alpha) \Big|_0^1 - \int_0^1 \frac{\sin z\alpha}{\pi z} J_b'(\alpha) d\alpha \\ &= \frac{\cos z\alpha}{\pi z^2} J_b'(\alpha) \Big|_0^1 - \int_0^1 \frac{\cos z\alpha}{\pi z^2} J_b''(\alpha) d\alpha \ll \frac{e^{|\operatorname{Im}(z)|}}{|z|^2}. \end{aligned}$$

□

Returning to  $K_b$  in (4.8), we see from (5.3) of Lemma 3 that

$$(5.9) \quad \operatorname{Re} K_b(-i(\rho - \rho') \log T) = \operatorname{Re} K_b((\gamma - \gamma') \log T - i(\beta - \beta') \log T) > 0 \quad \text{if } |\beta - \beta'| < \frac{b}{\log T}.$$

We want the real part of every term of the sum in (4.8) to be positive (or at least non-negative), and the assumption in Theorem 1 that  $|\beta - \frac{1}{2}| < \frac{b}{2 \log T}$  for all zeros with  $T < \gamma \leq 2T$  clearly implies that for any two of these zeros  $\rho$  and  $\rho'$ , we have  $|\beta - \beta'| < \frac{b}{\log T}$ .

**Lemma 4.** *Assume that, for all sufficiently large  $T$ , all zeros  $\rho$  of the Riemann zeta-function with  $T < \gamma \leq 2T$  are in  $B_b$  for some  $0 < b \ll 1$ . Then*

$$(5.10) \quad \sum_{\rho, \rho' \in B_b} \operatorname{Re} K_b(-i(\rho - \rho') \log T) = \frac{1}{2\pi} \left( j(0) + 2 \int_0^1 \frac{\alpha j(\alpha)}{\cosh(b\alpha)} d\alpha + O\left(\frac{1}{\sqrt{\log T}}\right) \right) \frac{T}{2\pi} \log T,$$

where every term in the sum above is positive.

*Proof of Lemma 4.* Since  $W(\rho - \rho')$  is complex-valued we need to remove this weight when  $|\beta - \beta'| < \frac{b}{\log T}$  in (4.8) before applying (5.9). Since  $W(0) = 1$  and  $W(z) - 1 = \frac{z^2}{4 - z^2}$ , we have by (5.4),

$$(5.11) \quad \begin{aligned} \sum_{\rho, \rho' \in B_b} K_b(-i(\rho - \rho') \log T)(W(\rho - \rho') - 1) &\ll \sum_{\rho, \rho' \in B_b} \frac{T^{|\beta - \beta'|}}{1 + |\rho - \rho'|^2 \log^2 T} \frac{|\rho - \rho'|^2}{|4 - (\rho - \rho')^2|} \\ &\leq \frac{1}{\log^2 T} \sum_{T < \gamma, \gamma' \leq 2T} \frac{T^{b/\log T}}{|4 - (\rho - \rho')^2|} \ll \frac{e^b}{\log^2 T} \sum_{T < \gamma, \gamma' \leq 2T} w(\gamma - \gamma') \ll e^b T \ll T, \end{aligned}$$

where we used (2.8) and (3.7) in the last line.

To complete the proof, our assumption in Lemma 4 allows us to insert the condition  $|\beta - \beta'| < \frac{b}{\log T}$  into the sum in (4.8), which then allows us to remove  $W(\rho - \rho')$  from the terms in the sum with an error  $\ll T \ll T\sqrt{\log T}$ . Taking the real parts proves (5.10).  $\square$

## 6. PROOF OF THEOREM 1

*Proof of Theorem 1.* In Lemma 4, taking the “diagonal” terms with  $\rho = \rho'$  and using (5.8), we obtain the sum

$$\sum_{\substack{\rho \\ T < \gamma \leq 2T}} m_\rho \operatorname{Re} K_b(0) = \sum_{\substack{\rho \\ T < \gamma \leq 2T}} m_\rho \left( \frac{1}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} d\alpha \right),$$

where  $m_\rho$  is the multiplicity of the zero  $\rho$ .<sup>§</sup> Similarly, the sum of the “symmetric diagonal terms,” which are the terms that have  $\rho' = 1 - \bar{\rho}$  and  $\beta' = 1 - \beta \neq 1/2$ , is

$$\sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \operatorname{Re} K_b(-i(2\beta - 1) \log T) = \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \left( \frac{1}{2\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} \left( T^{(2\beta-1)\alpha} + T^{-(2\beta-1)\alpha} \right) d\alpha \right).$$

Since all the terms in the sum in Lemma 4 are positive, these diagonal and symmetric diagonal terms provide a lower bound for the sum, and we obtain

$$(6.1) \quad \begin{aligned} \sum_{\substack{\rho \\ T < \gamma \leq 2T}} m_\rho \left( \frac{1}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} d\alpha \right) + \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \left( \frac{1}{2\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} \left( T^{(2\beta-1)\alpha} + T^{-(2\beta-1)\alpha} \right) d\alpha \right) \\ \leq \frac{1}{2\pi} \left( j(0) + 2 \int_0^1 \frac{\alpha j(\alpha)}{\cosh(b\alpha)} d\alpha + o(1) \right) \frac{T}{2\pi} \log T. \end{aligned}$$

We now take  $j(\alpha) = j_F(\alpha) = 1 - \alpha$  for  $0 \leq \alpha \leq 1$ . Under the assumption in Theorem 1 that  $b \rightarrow 0$  as  $T \rightarrow \infty$ , we have  $\cosh(b\alpha) \rightarrow 1$  uniformly for  $0 \leq \alpha \leq 1$ . Similarly, since  $e^{-b\alpha} \leq T^{\pm(2\beta-1)\alpha} \leq e^{b\alpha}$ , we have  $T^{\pm(2\beta-1)\alpha} \rightarrow 1$  uniformly for  $0 \leq \alpha \leq 1$  when  $b \rightarrow 0$  as  $T \rightarrow \infty$ . Thus, we have

$$\sum_{\substack{\rho \\ T < \gamma \leq 2T}} m_\rho \left( \frac{1}{2\pi} + o(1) \right) + \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \left( \frac{1}{2\pi} + o(1) \right) \leq \frac{1}{2\pi} \left( 1 + 2 \int_0^1 \alpha(1 - \alpha) d\alpha + o(1) \right) \frac{T}{2\pi} \log T,$$

<sup>§</sup>Note that  $\rho = \rho'$  is counted with multiplicity in the sum and also weighted by its multiplicity; for example if  $m_\rho = 2$  then  $\rho = \rho'$  occurs in 4 ways, but is only counted twice in the sum if not weighted by  $m_\rho$ .

and we conclude that

$$(6.2) \quad \sum_{T < \gamma \leq 2T} m_\rho + \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \leq \left( \frac{4}{3} + o(1) \right) \frac{T}{2\pi} \log T.$$

To prove (1.2) we use Montgomery's argument. By (6.2),

$$\sum_{T < \gamma \leq 2T} m_\rho \leq \left( \frac{4}{3} + o(1) \right) \frac{T}{2\pi} \log T,$$

and, on recalling (2.6), we have

$$(6.3) \quad \sum_{\substack{T < \gamma \leq 2T \\ \rho \text{ simple}}} 1 \geq \sum_{T < \gamma \leq 2T} (2 - m_\rho) \geq \left( 2 - \frac{4}{3} + o(1) \right) \frac{T}{2\pi} \log T = \left( \frac{2}{3} + o(1) \right) \frac{T}{2\pi} \log T.$$

To prove (1.3), since

$$\sum_{T < \gamma \leq 2T} m_\rho \geq \sum_{T < \gamma \leq 2T} 1 = (1 + o(1)) \frac{T}{2\pi} \log T,$$

we have by (6.2)

$$(6.4) \quad \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \leq \left( \frac{4}{3} + o(1) \right) \frac{T}{2\pi} \log T - \sum_{T < \gamma \leq 2T} m_\rho \leq \left( \frac{1}{3} + o(1) \right) \frac{T}{2\pi} \log T.$$

Thus

$$\sum_{\substack{T < \gamma \leq 2T \\ \beta = \frac{1}{2}}} 1 = \sum_{T < \gamma \leq 2T} 1 - \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} 1 \geq (1 + o(1)) \frac{T}{2\pi} \log T - \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \geq \left( \frac{2}{3} + o(1) \right) \frac{T}{2\pi} \log T.$$

To prove (1.4), we have by (6.3) and (6.4),

$$\begin{aligned} \sum_{\substack{T < \gamma \leq 2T \\ \beta = \frac{1}{2} \text{ and } \rho \text{ simple}}} 1 &= \sum_{\substack{T < \gamma \leq 2T \\ \rho \text{ simple}}} 1 - \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2} \text{ and } \rho \text{ simple}}} 1 \geq \left( \frac{2}{3} + o(1) \right) \frac{T}{2\pi} \log T - \sum_{\substack{T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \\ &\geq \left( \frac{1}{3} + o(1) \right) \frac{T}{2\pi} \log T. \end{aligned}$$

□

## 7. PROOF OF THEOREM 2

In Theorem 2 we no longer assume  $b \rightarrow 0$  as  $T \rightarrow \infty$ . Then, in the second sum in (6.1), we group each term with  $\beta > \frac{1}{2}$  with its symmetric diagonal term that has  $\text{Re}(\rho) = 1 - \beta$ , so that this sum is equal to

$$\sum_{\substack{T < \gamma \leq 2T \\ \beta > \frac{1}{2}}} m_\rho \left( \frac{2}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} \cosh((2\beta - 1)\alpha \log T) d\alpha \right).$$

Since  $\cosh(x) \geq \cosh(0) = 1$  for any real number  $x$ , we conclude that this sum is

$$\geq \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta > \frac{1}{2}}} m_\rho \left( \frac{2}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} d\alpha \right) = \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \left( \frac{1}{\pi} \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} d\alpha \right).$$

We thus conclude that in place of (6.2) we have

$$(7.1) \quad \sum_{\substack{\rho \\ T < \gamma \leq 2T}} m_\rho + \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho \leq (\mathcal{C}_b(j) + o(1)) \frac{T}{2\pi} \log T,$$

where

$$\mathcal{C}_b(j) := \frac{j(0) + 2 \int_0^1 \frac{\alpha j(\alpha)}{\cosh(b\alpha)} d\alpha}{2 \int_0^1 \frac{j(\alpha)}{\cosh(b\alpha)} d\alpha}.$$

Exactly as in the previous section, we have

$$(7.2) \quad \begin{aligned} N_1(B_b) &\geq (2 - \mathcal{C}_b(j) + o(1)) \frac{T}{2\pi} \log T, & N_0(B_b) &\geq (2 - \mathcal{C}_b(j) + o(1)) \frac{T}{2\pi} \log T, \\ \sum_{\substack{\rho \\ T < \gamma \leq 2T \\ \beta \neq \frac{1}{2}}} m_\rho &\leq (\mathcal{C}_b(j) - 1 + o(1)) \frac{T}{2\pi} \log T, & N_{1,0}(B_b) &\geq (3 - 2\mathcal{C}_b(j) + o(1)) \frac{T}{2\pi} \log T. \end{aligned}$$

We now take  $j(\alpha) = j_M(\alpha)$  and by Mathematica it is easy to verify Theorem 2 and compute Table 1 and Table 2.

#### ACKNOWLEDGEMENT AND FUNDING

The authors thank the American Institute of Mathematics for its hospitality and for providing a pleasant research environment where the authors met and started this research. The first author was supported by NSF DMS-1854398 FRG. The third author was supported by JSPS KAKENHI Grant Number 22K13895 and Inamori Research Grant 2024. The fourth author is partially supported by NSF CAREER DMS-2239681.

#### REFERENCES

- [BGSTB24] Siegfried Alan C. Baluyot, Daniel Alan Goldston, Ade Irma Suriajaya, and Caroline L. Turnage-Butterbaugh. An unconditional Montgomery theorem for pair correlation of zeros of the Riemann zeta-function. *Acta Arith.*, 214:357–376, 2024.
- [BHB13] H. M. Bui and D. R. Heath-Brown. On simple zeros of the Riemann zeta-function. *Bull. Lond. Math. Soc.*, 45(5):953–961, 2013.
- [CG93] A. Y. Cheer and D. A. Goldston. Simple zeros of the Riemann zeta-function. *Proc. Amer. Math. Soc.*, 118(2):365–372, 1993.
- [CGdL20] Andrés Chirre, Felipe Gonçalves, and David de Laat. Pair correlation estimates for the zeros of the zeta function via semidefinite programming. *Adv. Math.*, 361:106926, 22, 2020.
- [CGG98] J. B. Conrey, A. Ghosh, and S. M. Gonek. Simple zeros of the Riemann zeta-function. *Proc. London Math. Soc. (3)*, 76(3):497–522, 1998.
- [Dav00] Harold Davenport. *Multiplicative number theory*, volume 74 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2000. Revised and with a preface by Hugh L. Montgomery.
- [EMOT54] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Tables of integral transforms. Vol. I*. McGraw-Hill Book Co., Inc., New York-Toronto-London, 1954. Based, in part, on notes left by Harry Bateman.

$b$	$N_1(B_b)$ or $N_0(B_b)$	$N_{1,0}(B_b)$
0.001	.67250	.34500
0.2	.67019	.34038
0.4	.66333	.32666
0.6	.65208	.30416
0.8	.63670	.27339
1	.61748	.23496
1.2	.59475	.18951
1.4	.56884	.13768
1.6	.54003	.08007
1.8	.50862	.01724
2	.47485	
2.2	.43894	
2.4	.40109	
2.6	.36149	
2.8	.32027	
3	.27760	
3.2	.23357	
3.4	.18832	
3.6	.14194	
3.8	.09451	
4	.04612	
4.187	.00007	

TABLE 1. Lower bounds obtained using  $j_M$  for  $N_1(B_b)$  or  $N_0(B_b)$  and  $N_{1,0}(B_b)$  under the assumption of Theorem 1.

- [GM87] Daniel A. Goldston and Hugh L. Montgomery. Pair correlation of zeros and primes in short intervals. In *Analytic number theory and Diophantine problems (Stillwater, OK, 1984)*, volume 70 of *Progr. Math.*, pages 183–203. Birkhäuser Boston, Boston, MA, 1987.
- [Gol81] Daniel Alan Goldston. *LARGE DIFFERENCES BETWEEN CONSECUTIVE PRIME NUMBERS*. ProQuest LLC, Ann Arbor, MI, 1981. Thesis (Ph.D.)–University of California, Berkeley.
- [GP24] R. Garunkštis and J. Paliulionytė, 2024. personal communication.
- [HB79] D. R. Heath-Brown. Simple zeros of the riemann zeta-function on the critical line. *Bull. London Math. Soc.*, 11:17–18, 1979.
- [Kor58a] N. M. Korobov. Estimates of trigonometric sums and their applications. *Uspehi Mat. Nauk*, 13(4(82)):185–192, 1958.
- [Kor58b] N. M. Korobov. Weyl’s estimates of sums and the distribution of primes. *Dokl. Akad. Nauk SSSR*, 123:28–31, 1958.
- [Mon73] H. L. Montgomery. The pair correlation of zeros of the zeta function. In *Analytic number theory (Proc. Sympos. Pure Math., Vol. XXIV, St. Louis Univ., St. Louis, Mo., 1972)*, volume Vol. XXIV of *Proc. Sympos. Pure Math.*, pages 181–193. Amer. Math. Soc., Providence, RI, 1973.
- [Mon75] Hugh L. Montgomery. Distribution of the zeros of the Riemann zeta function. In *Proceedings of the International Congress of Mathematicians (Vancouver, B.C., 1974)*, Vol. 1, pages 379–381. Canad. Math. Congr., Montreal, QC, 1975.
- [PRZZ20] Kyle Pratt, Nicolas Robles, Alexandru Zaharescu, and Dirk Zeindler. More than five-twelfths of the zeros of  $\zeta$  are on the critical line. *Res. Math. Sci.*, 7(2):Paper No. 2, 74, 2020.
- [SS03] Elias M. Stein and Rami Shakarchi. *Complex analysis*, volume 2 of *Princeton Lectures in Analysis*. Princeton University Press, Princeton, NJ, 2003.
- [Tit39] E. C. Titchmarsh. *The theory of functions*. Oxford University Press, Oxford, second edition, 1939.

$b$	$N_1(B_b) _{j_F}$	$N_1(B_b) _{j_M}$
0.001	.66666	.67250
0.2	.66422	.67019
0.4	.65697	.66333
0.6	.64509	.65208
0.8	.62886	.63670
1	.60861	.61748
1.2	.58468	.59475
1.4	.55743	.56884
1.6	.52719	.54003
1.8	.49424	.50862
2	.45887	.47485
2.2	.42130	.43894
2.4	.38176	.40109
2.6	.34043	.36149
2.8	.29747	.32027
3	.25304	.27760
3.2	.20727	.23357
3.4	.16026	.18832
3.6	.11214	.14194
3.8	.06298	.09451
4	.01288	.04612
4.0508	.00022	.03368
4.187	0	.00007

TABLE 2. Comparison of using  $j_F$  and  $j_M$  for  $N_1(B_b)$  (or  $N_0(B_b)$ ).

- [Tit86] E. C. Titchmarsh. *The theory of the Riemann zeta-function*. The Clarendon Press, Oxford University Press, New York, second edition, 1986. Edited and with a preface by D. R. Heath-Brown.
- [Tsa93] Kai Man Tsang. The large values of the Riemann zeta-function. *Mathematika*, 40(2):203–214, 1993.
- [Vin58] I. M. Vinogradov. A new estimate of the function  $\zeta(1+it)$ . *Izv. Akad. Nauk SSSR Ser. Mat.*, 22:161–164, 1958.

MATHEMATICS DEPARTMENT, EAST CAROLINA UNIVERSITY, GREENVILLE, NC 27858  
*Email address:* baluyots24@ecu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, SAN JOSE STATE UNIVERSITY  
*Email address:* daniel.goldston@sjsu.edu

FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY  
*Email address:* adeirmasuriajaya@math.kyushu-u.ac.jp

CARLETON COLLEGE  
*Email address:* cturnageb@carleton.edu