arXiv:2501.15159v1 [gr-qc] 25 Jan 2025 arXiv:2501.15159v1 [gr-qc] 25 Jan 2025

Reproducing Λ CDM-like Solutions in $f(Q)$ Gravity: A Comprehensive Study Across All Connection Branches

Saikat Chakraborty \mathbb{D} [,](https://orcid.org/0000-0002-6097-454X) ^{1, 2, *} Jibitesh Dutta \mathbb{D} , ^{3, 4, [†](#page-0-1)} Daniele

Gregoris \mathbb{D} [,](https://orcid.org/0000-0002-0448-3447)^{5,[‡](#page-0-2)} Khamphee Karwan,^{1, [§](#page-0-3)} and Wompherdeiki Khyllep \mathbb{D}^{6} \mathbb{D}^{6} \mathbb{D}^{6} ,

 1 ¹The Institute for Fundamental Study "The Tah Poe Academia Institute",

Naresuan University, Phitsanulok 65000, Thailand

² Center for Space Research, North-West University, Potchefstroom 2520, South Africa

³Mathematics Division, Department of Basic Sciences and Social Sciences,

North Eastern Hill University, Shillong, Meghalaya 793022, India

⁴ Inter University Centre for Astronomy and Astrophysics, Pune 411 007, India

⁵ School of Science, Jiangsu University of Science and Technology, Zhenjiang 212100, China

⁶Department of Mathematics, St. Anthony's College, Shillong, Meghalaya 793001, India

Given the remarkable success of the ΛCDM model in fitting various cosmological observations, a pertinent question in assessing the phenomenological viability of modified gravity theories is whether they can reproduce an exactly ΛCDM-like cosmic background evolution. In this paper, we address this question in the context of $f(Q)$ gravity, where Q denotes the nonmetricity scalar. It is known that there are three possible symmetric teleparallel connection branches that respect the cosmological principles of spatial homogeneity, isotropy, and global spatial flatness. By enforcing a ΛCDM-like background evolution via the cosmographic condition $j(z) = 1$, where j is the jerk parameter, we reconstruct the Λ CDM-mimicking $f(Q)$ theory for each of the three possible connection branches. For the first connection branch, also known as the "coincident gauge" in cosmology, we recover the previously known result that a theory of the form $f(Q) = -2\Lambda + \alpha Q + \beta \sqrt{-Q}$ can exactly reproduce a ΛCDM-like cosmic evolution. Furthermore, we establish that the stability of the ΛCDM-like cosmic solution within this reconstructed $f(Q)$, as well as the robustness of the reconstructed $f(Q)$ form with respect to small errors in the astrophysical measurements of the jerk parameter. For the second connection branch, we analytically reconstruct the Λ CDM-mimicking $f(Q)$ to be of the form $f(Q) = -2\Lambda + \alpha Q - \beta Q^2$. For the third connection branch, we could decouple the evolution equation for the dynamical connection function, which enabled us to perform a numerical reconstruction. Our analysis proves that, at least at the background level, it is possible to obtain Λ CDM-mimicking $f(Q)$ models for all the three possible connection branches. These models effectively reduce to (STE)GR in the past while possessing a positive effective gravitational coupling throughout.

I. INTRODUCTION

General Relativity (GR), based on the curvature of spacetime, has been fundamental in understanding the physics of gravitational interaction and the dynamics of the universe's large-scale structure. Its achievements encapsulated within the ΛCDM cosmological model include the successful description of the cosmic microwave background (CMB), baryonic acoustic oscillations (BAO), and the accelerating expansion of the universe [\[1,](#page-28-0) [2\]](#page-28-1). Still, specific unresolved theoretical problems such as the cosmological constant problem [\[3\]](#page-28-2) and observational tensions such as the Hubble tension [\[4\]](#page-28-3) provide reasons to study alternatives to GR.

The geometric foundation of GR is, however, part of a broader "Trinity of Gravity" that also includes two equivalent formulations of GR: the teleparallel equivalent of GR (TEGR) and the symmetric teleparallel equivalent of GR $(STEGR)$ [\[5\]](#page-28-4). In the TEGR framework, gravity is described through the torsion of the spacetime [\[6\]](#page-28-5), while in STEGR, it is interpreted in terms of nonmetricity of the spacetime [\[7\]](#page-28-6). These alternative formulations of non-Riemannian geometry provide fertile grounds for developing modified gravity theories based on arbitrary scalar functions of their respective scalar quantities, such as $f(T)$ or $f(Q)$, where T and Q represent the torsion and nonmetricity scalars, respectively [\[8–](#page-28-7)[10\]](#page-28-8).

There has been tremendous interest in $f(Q)$ gravity in recent years due to its exceptional geometric interpretation without using curvature or torsion. However, we point out that recently there has been a discussion about potential presence of ghost and strong coupling issues in $f(Q)$ gravity [\[11\]](#page-28-9), and whether such ghost degrees of freedom can actually propagate [\[12\]](#page-29-0). In fact, similar issues also showed up in the study of $f(T)$ gravity, where certain degrees of

[∗] [saikat.ch@nu.ac.th;](mailto:saikat.ch@nu.ac.th) 48674087@nwu.ac.za

[†] jibitesh@nehu.ac.in

[‡] danielegregoris@libero.it

[§] [khampheek@nu.ac.th;](mailto:khampheek@nu.ac.th) Corresponding author

[¶] sjwomkhyllep@gmail.com

freedom lose their kinetic terms on the FLRW background [\[13\]](#page-29-1). It was pointed out in [\[14\]](#page-29-2) that a direct coupling of the matter field to the connection or a nonminimal coupling between the matter and the geometry sector might alter the situation leading to a healthy theory. A class of ghost free theories in symmetric teleparallel geometry was recently investigated in [\[15\]](#page-29-3). Strong coupling and ghost issues, as well as the question of the actual number of propagating degrees of freedom underscore the need for perturbative stability analysis and Hamiltonian analysis in $f(Q)$ theory. Investigating these issues are important from the theoretical point of view for establishing the viability of modifying gravity in the symmetric teleparallel framework. Indeed, this is an active area of research currently.

Although we are aware of the above theoretical issues, this is not the main focus of our investigation here. Rather, our focus is to examine the phenomenological viability of modifying gravity in the symmetric teleparallel framework, along a line of similar works [\[16](#page-29-4)[–20\]](#page-29-5). The framework of symmetric teleparallelism gives promising insights into cosmological dynamics. $f(Q)$ gravity has proven to be compatible with the observational data at the level of the background and of the perturbations, such as CMB, supernovae, BAO, redshift-space distortions, growth measurements [\[21–](#page-29-6)[23\]](#page-29-7). In addition, it does not have problems with the constraints imposed by Big Bang Nucleosynthesis, which means it could successfully compete with other modified gravity theories [\[24\]](#page-29-8).

The ΛCDM model based on Einstein's GR remains a very good fit with available cosmological datasets at least according to Planck 20[1](#page-1-0)8 $[25]$ ¹. In the context of modified gravity theories, from a phenomenological point of view, an important question is whether a modified gravity theory can give rise to exactly the same kind of background evolution as the GR-based ΛCDM model. Such modified gravity theories will then be indistinguishable from the ΛCDM model at the background level, and typically one needs to go to the perturbation level to find any distinctive signature. This particular question has been addressed in the context of $f(R)$ gravity [\[28–](#page-29-10)[30\]](#page-29-11), $F(R, G)$ and modified Gauss-Bonnet gravity [\[31\]](#page-29-12) and coincident gauge $f(Q)$ gravity (i.e. considering only the trivial connection branch where $Q = -6H^2$) [\[17,](#page-29-13) [32\]](#page-29-14). However, in general there are three different symmetric teleparallel connection branches respecting spatial flatness, homogeneity and isotropy [\[33\]](#page-29-15), which we denote later as Γ_1 , Γ_2 , Γ_3 , with Γ_2 , Γ_3 commonly being referred to in the literature as the so-called non-coincident gauges $[34]^2$ $[34]^2$. In this paper, we reconstruct the $f(Q)$ gravity that reproduces exactly a ΛCDM-like background evolution for each of the three connection branches, recovering also the earlier reconstructed $f(Q)$ for Γ_1 .

There has been some work on the cosmological reconstruction of $f(Q)$ gravity, primarily within the trivial connection branch Γ_1 , i.e., the so-called coincident gauge. For instance, [\[35\]](#page-29-17) numerically reconstructed $f(Q)$ starting from a cosmographic series with Pad´e approximations, while [\[36\]](#page-29-18) presented reconstruction algorithms for Bianchi-I and FLRW cosmology. In [\[17\]](#page-29-13), a designer approach was adopted to reconstruct $f(Q)$ corresponding to various dark energy equation of state parametrizations, and [\[37\]](#page-29-19) explored cosmological reconstruction of both $f(Q)$ gravity and mimetic $f(Q)$ gravity. Similarly, [\[32\]](#page-29-14) reconstructed $f(Q)$ gravity that reproduces a Λ CDM-like background evolution.

To our knowledge, the paper [\[38\]](#page-29-20) is the first study to attempt reconstruction beyond the coincident gauge. Using the OHD dataset and assuming vanishing hypermomentum, the authors numerically reconstructed $f(Q)$ for the connection branch Γ_2 [\[38,](#page-29-20) Eq.30]. In the general case with nonvanishing hypermomentum, they numerically reconstructed the dynamical connection function for Γ_2 and Γ_3 by assuming particular forms of $f(Q)$.

In this paper, we perform the cosmological reconstruction of $f(Q)$ starting from the Λ CDM-like cosmic evolution for all three connection branches. For the connection branch Γ_1 , we recover the results earlier obtained in [\[17,](#page-29-13) [32\]](#page-29-14). For the connection branch Γ2, we could perform an analytical reconstruction and obtain a functional form for the Λ CDM-mimicking $f(Q)$. For the connection branch Γ_3 , which we found to be the most complicated, we could obtain a decoupled evolution equation for the dynamical connection function, which ultimately enabled us for a numerical reconstruction of the Λ CDM-mimicking $f(Q)$.

The paper is organized as follows: In sections [II,](#page-2-0) [III,](#page-4-0) we give some brief and self-contained exposures to the cosmographic condition corresponding to the Λ CDM-like cosmic evolution and $f(Q)$ gravity respectively. In section [IV,](#page-5-0) we introduce the three connection branches of the FLRW cosmology in the symmetric teleparallel framework and write down the corresponding cosmological field equations in section [V.](#page-6-0) The sections [VI,](#page-7-0) [VII,](#page-12-0) [VIII](#page-17-0) are dedicated to the reconstruction of the ΛCDM-mimicking $f(Q)$ for the connection branches Γ_1 , Γ_2 and Γ_3 respectively. In each case, we identify the free parameters of the reconstructed $f(Q)$ model and possible bounds for them, as much as possible. We also show the evolution of the effective gravitational coupling in each case. For the case of Γ_1 , we also investigate the stability of the Λ CDM-like cosmological solution in the reconstructed $f(Q)$ in the subsection VID and the robustness of the reconstructed $f(Q)$ against astrophysical errors of the measurement of the jerk parameter in section [VI E.](#page-10-0) Finally, we conclude in section [IX](#page-23-0) summarizing the key takeaways from our paper and adding some relevant discussions. Additionally, in appendix [A,](#page-26-0) we comment on how one can go ahead to investigate the stability of

¹ The recent DESI data [\[26\]](#page-29-21) apparently favours an evolving dark energy, although there has been arguments that their conclusion is highly driven by the choice of the priors [\[27\]](#page-29-22).

² There is a slight misnomer here. For any particular connection branch Γ_i , one can find a coordinate system in which the connection can be trivialized [\[33\]](#page-29-15). This particular coordinate system is the coincident gauge of the connection branch in question. It happens that the coincident gauge for the connection branch Γ_1 is the usual Cartesian coordinate system.

the ΛCDM-like cosmological solution in the reconstructed $f(Q)$ for the case of Γ_2 and add some comments about the dynamical connection function for Γ_2 in appendix [B.](#page-27-0)

Throughout the paper, we use the metric signature $(-, +, +, +)$ and the geometrized unit system $8\pi G = c = 1, G$ being Newton's gravitational constant and c is the speed of light in vacuum.

II. HUBBLE RATE FROM COSMOGRAPHIC RECONSTRUCTION

Cosmography is a model-independent approach to derive the universe's expansion history from astrophysical datasets, characterized by a hierarchy of parameters involving time derivatives of the scale factor $a(t)$. The first two terms of this hierarchy are the Hubble rate and deceleration function, respectively:

$$
H := \frac{\dot{a}}{a}, \qquad q := -\frac{\ddot{a}a}{\dot{a}^2} = -1 - \frac{\dot{H}}{H^2}, \qquad (1)
$$

while the third one is the so-called jerk parameter [\[39\]](#page-29-23):

$$
j = \frac{\dddot{a}}{aH^3} = 1 + 3\frac{\dot{H}}{H^2} + \frac{\ddot{H}}{H^3}.
$$
 (2)

The cosmographic parameters characterize the 'kinematics' of the universe, capturing information from the time derivatives of its scale factor. However, they do not provide an understanding of its dynamics. A relationship of the cosmographic parameters to the matter abundance parameters can be found only if a specific 'model', as characterized by the Friedmann and Raychaudhuri equations , is specified [\[40\]](#page-29-24). In other words, while the cosmographic parameters are completely kinematic, the particular relationship between the matter abundance parameter to the cosmographic parameters depends on the 'dynamics'. For example, in the ΛCDM model, the matter abundance parameter is related to the cosmographic deceleration parameter as

$$
\Omega_{m0} = \frac{2}{3}(1+q_0),\tag{3}
$$

where the subscript '0' implies present day values. For alternative late time models, this relationship will change, as we will see later on in the text for $f(Q)$ gravity. It is worth mentioning that a different algebraic combinations of the time derivatives could also be considered as in statefinder diagnostic approach [\[41\]](#page-29-25).

The evolution of the cosmographic functions with respect to the redshift can be inferred in a model-independent (or non-parametric) manner directly from relevant datasets. Deviations from the a ΛCDMlike evolution being constrained to be no more than 10 (per cent) [\[42,](#page-29-26) Fig.11]-[\[43,](#page-29-27) Fig.10]-[\[44,](#page-29-28) Fig.4]-[\[45\]](#page-29-29). The spatially flat ΛCDM model exhibits a very special cosmographic property of a constant jerk parameter $j(z) = 1$ all along its evolution. A stout theory should not be fine-tuned to just a very specific numerical value of the jerk, but rather be applicable to all its astrophysical values within an interval of observational uncertainty. We shall therefore set $j = 1 + \varepsilon = const.$ for "small" ε ; the latter should nevertheless be regarded as the previously mentioned largest possible uncertainty in the observational estimate of the jerk function so that the evolution provided by different redshift parametrization would all fall within this interval. This choice has been shown not to trouble the clustering properties of dark matter accounting for a consistent structure formation cosmic epoch [\[46\]](#page-29-30), but sill allowing for small departures from a ΛCDM-like evolution which may come in handy in taming current observational tensions [\[47\]](#page-29-31). By integrating (2) , we obtain [\[48,](#page-29-32) Eq.(8)]:

$$
H^{2}(z) = H_{0}^{2}[c_{1}(1+z)^{k_{1}} + (1-c_{1})(1+z)^{k_{2}}], \qquad k_{1,2} = \frac{3 \pm \sqrt{1+8j}}{2}, \tag{4}
$$

with c_1 , for now, an arbitrary mathematical constant of integration to be determined through appropriate boundary conditions. By computing the deceleration function

$$
q(z) = -1 + \frac{H_0^2[c_1k_1(1+z)^{k_1} + (1-c_1)k_2(1+z)^{k_2}]}{H^2(z)} = -1 + \frac{k_1H^2(z) + (1-c_1)(k_2 - k_1)H_0^2(1+z)^{k_2}}{2H^2(z)},
$$
(5)

we can actually relate this mathematical integration constant to the present day value of the deceleration parameter and the jerk parameter

$$
c_1 = 1 + \frac{k_1 - 2(1 + q_0)}{k_2 - k_1},
$$
\n⁽⁶⁾

where $k_{1,2}$ have been given in Eq.[\(4\)](#page-2-2). From the above equation, it is seen that the constant c_1 is completely kinematical in nature. It is determined in general by the present day values of the deceleration and jerk parameters, both of which can be estimated from astrophysical datasets without referring to any particular model.

To reconstruct a modified gravity theory in the context of the late-time universe, one can proceed two ways. Firstly, one can reconstruct starting from a particular cosmic evolution $a(t)$ or $H(t)$ [\[28,](#page-29-10) [29,](#page-29-33) [32,](#page-29-14) [49–](#page-30-0)[51\]](#page-30-1). One can also start by specifying a cosmographic condition [\[52\]](#page-30-2), since a particular cosmic evolution can also be specified by a particular cosmographic condition [\[53\]](#page-30-3) (e.g. the condition $j(z) = 1$ corresponds to a Λ CDM-like evolution). On the contrary, one can also start the reconstruction starting from a given form of the dark energy equation of state parameter (e.g. $w_{\text{DE}}(z) = -1$). The later is the so-called designer approach [\[16,](#page-29-4) [17,](#page-29-13) [30,](#page-29-11) [54–](#page-30-4)[56\]](#page-30-5). The approach adopted in this paper is the first one. In particular, we make use of the cosmographic condition $j(z) = 1$. We prefer the former approach because it is more grounded on available astrophysical datasets e.g. the luminosity distance-redshift data, which gives us bounds on the cosmographic parameters.

The Hubble rate [\(4\)](#page-2-2) smoothly reduces to the following ΛCDM-like behavior in the limit $\varepsilon \to 0$:

$$
H2(z) = H02h2(z) = H02[c1(1 + z)3 + (1 - c1)],
$$
\n(7)

with $0 < c_1 < 1$. We define a ACDM-like evolution to be a cosmic evolution which admits well-defined asymptotic past and asymptotic future epochs at which $H(z) \propto (1+z)^3$ and $H(z) \to const.$, respectively [\[40\]](#page-29-24). It is important at this point to clarify the distinction between the ΛCDM-like evolution and the ΛCDM 'model' itself. Of course, the ΛCDM model based on Einstein's GR does give rise to the ΛCDM-like evolution [\(7\)](#page-3-0), but it is possible for other models, in particular modified gravity based models to give rise to a ΛCDM-like evolution as well. See, for example, Ref.[\[57\]](#page-30-6) for $f(R)$ gravity or Ref.[\[32\]](#page-29-14) for coincident gauge $f(Q)$ gravity reproducing ΛCDM-like evolution. How the mathematical integration constant c_1 is related to Ω_{m0} and $\Omega_{\text{DE}0}$ is model dependent. For the particular Λ CDM 'model', one has $c_1 = \Omega_{m0}$, and hence the mathematical constant c_1 assumes the physical meaning of the present-day abundance of dark matter. However, in general, and one would obtain $c_1 = c_1(\Omega_{m0}, \Omega_{\text{DE0}})$, and consequently its physical interpretation will change. The particular function $c_1(\Omega_{m0}, \Omega_{\text{DE0}})$, and consequently its physical interpretation, is, therefore, model dependent^{[3](#page-3-1)} While the parameter c_1 is a purely kinematical and model-independent parameter (Eq.[\(6\)](#page-2-3)), the available estimates of the present-day abundance of dark matter Ω_{m0} are model-dependent. The latter constitute possible sources to the Hubble tension [\[59\]](#page-30-7).

It can be checked that for a ΛCDM-like evolution [\(4\)](#page-2-2) the deceleration parameter is

$$
q = -1 - \frac{\dot{H}}{H^2} = \frac{1}{2} - \frac{3}{2} \left(\frac{1 - c_1}{h^2} \right),
$$
\n(8)

where we have used the relation $\frac{d}{dt} \equiv -H(1+z)\frac{d}{dz}$. For Λ CDM-like evolution we have

$$
\frac{dq}{Hdt} = (2q - 1)(q + 1),\tag{9}
$$

and consequently q decreases monotonically from $q = \frac{1}{2}$ to $q = -1$. It can be calculated that c_1 is related to q_0 as

$$
c_1 = \frac{2}{3}(1+q_0),\tag{10}
$$

whose relationship is consistent with [\(6\)](#page-2-3) for $j = 1$. Indeed, in the ΛCDM-like paradigm, Cosmology is a search for two numbers, H_0 and q_0 , [\[60\]](#page-30-8). Therefore, we can express the Λ CDM-like cosmic evolution as

$$
H^{2}(z) = H_{0}^{2}h^{2}(z) = H_{0}^{2}\left[\frac{2}{3}(1+q_{0})(1+z)^{3} + \frac{1}{3}(1-2q_{0})\right],
$$
\n(11)

such that it guarantees $H(z = 0) = H_0, q(z = 0) = q_0$.

Lastly, we propose a definition for the "almost" ΛCDM-like evolution. We define it such that all its cosmographic quantities and relations between them are "almost" like that in the ΛCDM-like evolution modulo regular terms with respect to a small parameter ε . Whether the same is also true for the reconstructed theory function $f(Q)$ or the dynamical connection function $\gamma(z)$ and the non-metricity variable $Q(z)$, is, however, not a trivial question due to the non-linearity of the field equations and demands explicitly investigation. In a clearer manner, the question can be posed as: does small deviation in cosmological kinematics also imply small deviation in cosmological dynamics?

³ For some discussion about this important conceptual point, the reader is referred to the discussion around Eq.(8) of Ref.[\[58\]](#page-30-9) and [\[40,](#page-29-24) Sec.2].

III. BASICS OF $f(Q)$ GRAVITY THEORY

In metric-affine theories, the affine connection can be decomposed into three distinct components as follows:

$$
\Gamma^{\lambda}_{\mu\nu} = \mathring{\Gamma}^{\lambda}_{\mu\nu} + K^{\lambda}_{\mu\nu} + L^{\lambda}_{\mu\nu},\tag{12}
$$

where $\mathring{\Gamma}^{\lambda}_{\mu\nu}$ is the Levi-Civita connection of the metric $g_{\mu\nu}$

$$
\mathring{\Gamma}^{\lambda}_{\mu\nu} = \frac{1}{2} g^{\lambda\beta} \left(\partial_{\mu} g_{\beta\nu} + \partial_{\nu} g_{\beta\mu} - \partial_{\beta} g_{\mu\nu} \right) , \qquad (13)
$$

 $K^{\lambda}{}_{\mu\nu}$ is the contorsion tensor

$$
K^{\lambda}{}_{\mu\nu} = \frac{1}{2}g^{\lambda\beta} \left(-T_{\mu\beta\nu} - T_{\nu\beta\mu} + T_{\beta\mu\nu} \right) , \qquad (14)
$$

and $L^{\lambda}{}_{\mu\nu}$ is the disformation tensor

$$
L^{\lambda}{}_{\mu\nu} = \frac{1}{2} g^{\lambda\beta} \left(-Q_{\mu\beta\nu} - Q_{\nu\beta\mu} + Q_{\beta\mu\nu} \right). \tag{15}
$$

It may be noted that the contorsion is constructed from the torsion tensor $T^{\lambda}{}_{\mu\nu}$ which is due to the anti-symmetric nature of the connection i.e.

$$
T^{\lambda}{}_{\mu\nu} = \Gamma^{\lambda}{}_{\mu\nu} - \Gamma^{\lambda}{}_{\nu\mu} \,. \tag{16}
$$

On the other hand, the disformation tensor is constructed from nonmetricity tensors as

$$
Q_{\rho\mu\nu} \equiv \nabla_{\rho} g_{\mu\nu} = \partial_{\rho} g_{\mu\nu} - \Gamma^{\beta}_{\mu\rho} g_{\beta\nu} - \Gamma^{\beta}_{\nu\rho} g_{\mu\beta}.
$$
\n(17)

One may also defined curvature tensor of the connection as

$$
R^{\sigma}{}_{\rho\mu\nu} = \partial_{\mu}\Gamma^{\sigma}{}_{\nu\rho} - \partial_{\nu}\Gamma^{\sigma}{}_{\mu\rho} + \Gamma^{\sigma}{}_{\mu\lambda}\Gamma^{\lambda}{}_{\nu\rho} - \Gamma^{\sigma}{}_{\mu\lambda}\Gamma^{\lambda}{}_{\nu\rho} \,. \tag{18}
$$

In analogy to the Ricci scalar obtained in GR due to the Levi-Civita connection, in nonmetrcity theory, one can obtain the non-metricity scalar Q due to the affine connection $\Gamma_{\mu\nu}^{\lambda}$ which is defined as the trace of the non-metricity tensor and given by [\[7\]](#page-28-6)

$$
Q = Q_{\alpha\mu\nu} P^{\alpha\mu\nu} \,, \tag{19}
$$

where $P^{\alpha}_{\ \mu\nu}$ the non-metricity conjugate given by

$$
P^{\alpha}_{\ \mu\nu} = -\frac{1}{2}L^{\alpha}_{\ \mu\nu} + \frac{1}{4}\left(Q^{\alpha} - \overline{Q}^{\alpha}\right)g_{\mu\nu} - \frac{1}{4}\delta^{\alpha}_{(\mu}Q_{\nu)}\,,\tag{20}
$$

with $Q_{\alpha} = Q_{\alpha\mu\nu}g^{\mu\nu}, \overline{Q}_{\alpha} = Q_{\mu\nu\alpha}g^{\mu\nu}$ as traces of the non-metricity tensor.

In this work, we consider an affine connection with vanishing curvature and torsion but a non-vanishing nonmetricity. Under such connection, a generic non-metricity $f(Q)$ theory was formulated whose action is given by [\[33,](#page-29-15) [61,](#page-30-10) [62\]](#page-30-11)

$$
S = \int d^4x \sqrt{-g} \left[\frac{1}{2} f(Q) + \mathcal{L}_m \right], \qquad (21)
$$

where \mathcal{L}_m is the matter Lagrangian density, g is the determinant of the metric $g_{\mu\nu}$, and $f(Q)$ is an arbitrary function of the invariant non-metricity scalar Q. The particular case $f(Q) = Q$ differs from Einstein-Hilbert action by only a boundary term. Therefore, at the level of the field equation, they are equivalent formulations [\[5\]](#page-28-4). Hence, the theory given by $f(Q) = Q$ is called the Symmetric Teleparallel Equivalent of GR (henceforth STEGR in short).

Since the action [\(21\)](#page-4-1) is constructed from the metric and the connection, so there are two equations of motion. One of the equations of motion is due to variation with respect to the metric given by

$$
f_{Q}G_{\mu\nu} + \frac{1}{2}g_{\mu\nu}(f_{Q}Q - f) + 2f_{QQ}\nabla_{\alpha}QP^{\alpha}{}_{\mu\nu} = T_{\mu\nu}\,,\tag{22}
$$

6

where $G_{\mu\nu}$ is the Einstein tensor and $T_{\mu\nu}$ is the energy-momentum tensor of matter. Here, $f_Q = \partial f/\partial Q$, $f_{QQ} =$ $\partial^2 f/\partial Q^2.$

Another equation of motion due to variation with respect to the connection is

$$
\nabla_{\mu}\nabla_{\nu}\left(\sqrt{-g}f_{Q}P^{\mu\nu}_{\sigma}\right) = 0. \tag{23}
$$

Since, the matter Lagrangian is not coupled to the connection, the energy momentum tensor is separately conserved i.e. $\nabla_{\mu}T^{\mu\nu} = 0$, here covariant derivative with respect to the Levi-Civita connection. On performing the Levi-Civita covariant derivative of [\(22\)](#page-4-2) and using Bianchi identity, one can obtain the connection equation [\(23\)](#page-5-1) [\[63\]](#page-30-12). Thus, the two equations of motion are related to each other.

Let us mention an important physical viability condition here which will be used extensively later on to put constraints on the free parameters of the reconstructed $f(Q)$. The effective gravitational constant $\frac{1}{f_Q}$ dictates how the geometry responds to matter energy-momentum in $f(Q)$ gravity. The situation $f_Q < 0$ may be interpreted as anti-gravity. The borderline case $f_Q = 0$ implies infinite gravitational "constant" and corresponds to a singularity. Hence, physically viable models should obey the requirement $f_Q > 0$ [\[64\]](#page-30-13).

IV. FLRW COSMOLOGY IN $f(Q)$ THEORY

In this work, we consider the most general forms of the metric and connection in the context of symmetric telleparallel gravity, which are compatible with the cosmological principle on homogeneity and isotropy. The homogeneity and isotropy nature is determined by the invariance of the metric and the connection with respect to the spatial rotational and translational transformations. Mathematically, the condition is equivalent to the vanishing Lie derivatives of the connection with respect to the Killing vectors corresponding to spatial rotations and translation [\[33\]](#page-29-15).

A general form of spatially flat metric with homogeneous and isotropic condition is the FLRW metric given by

$$
ds^{2} = -dt^{2} + a(t)^{2} (dr^{2} + r^{2} d\theta^{2} + r^{2} \sin^{2} \theta d\phi^{2}).
$$
\n(24)

The non-zero components of a general non-vanishing torsion, affine connection respecting homogeneity and isotropy are [\[33\]](#page-29-15)

$$
\Gamma_{tt}^{t} = C_1, \quad \Gamma_{rr}^{t} = C_2, \quad \Gamma_{\theta\theta}^{t} = C_2 r^2, \quad \Gamma_{\phi\phi}^{t} = C_2 r^2 \sin^2 \theta,
$$

\n
$$
\Gamma_{tr}^{r} = C_3, \Gamma_{\theta\theta}^{r} = -r, \quad \Gamma_{\phi\phi}^{r} = -r \sin^2 \theta,
$$

\n
$$
\Gamma_{t\theta}^{\theta} = C_3, \Gamma_{\theta r}^{\theta} = \frac{1}{r}, \Gamma_{\phi\phi}^{\theta} = -\cos \theta \sin \theta,
$$

\n
$$
\Gamma_{t\phi}^{\phi} = C_3, \Gamma_{\phi r}^{\phi} = \frac{1}{r}, \Gamma_{\theta\phi}^{\phi} = -\cot \theta,
$$

where C_1, C_2, C_3 are purely temporal functions. Imposing a curvature free condition, the above affine connection has three possible branches depending on the values of C_1, C_2, C_3 .

The first type of connection Γ_1 corresponds to:

$$
(C_1, C_2, C_3) = (\gamma, 0, 0).
$$

The second type of connection Γ_2 corresponds to

$$
(C_1, C_2, C_3) = \left(\gamma + \frac{\dot{\gamma}}{\gamma}, 0, \gamma\right),
$$

and the third type of connection Γ_3 corresponds to

$$
(C_1, C_2, C_3) = \left(-\frac{\dot{\gamma}}{\gamma}, \gamma, 0\right).
$$

For details, the reader is referred to the paper [\[33\]](#page-29-15).

V. COSMOLOGICAL FIELD EQUATIONS

To derive the cosmological field equations for the different branches of connections in $f(Q)$ gravity, we follow the conventions laid out in [\[34\]](#page-29-16). Specifically, we adopt the expression $Q = -6H^2$ for Γ_1 . In what follows, we write the field equations for each set of connections:

• Branch 1 (Γ_1) :

For the first branch, the nonmetricity scalar Q is related to the Hubble parameter H as:

$$
Q = -6H^2,\tag{25}
$$

leading to the Friedmann and Raychaudhuri equations:

$$
3H^2 f_Q + \frac{1}{2} \left(f - Q f_Q \right) = \rho_m \,, \tag{26a}
$$

$$
-2\frac{d}{dt}\left(Hf_Q\right) - 3H^2f_Q - \frac{1}{2}\left(f - Qf_Q\right) = 0.
$$
\n(26b)

• Branch 2 (Γ_2) :

In this branch, the scalar Q acquires additional terms from the time-dependent function $\gamma(t)$:

$$
Q = -6H^2 + 9\gamma H + 3\dot{\gamma},\tag{27}
$$

modifying the Friedmann and Raychaudhuri equations as:

$$
3H^2 f_Q + \frac{1}{2} \left(f - Q f_Q \right) + \frac{3\gamma \dot{Q} f_{QQ}}{2} = \rho_m \,, \tag{28a}
$$

$$
-2\frac{d}{dt}\left(Hf_Q\right) - 3H^2f_Q - \frac{1}{2}\left(f - Qf_Q\right) + \frac{3\gamma\dot{Q}f_{QQ}}{2} = 0\,,\tag{28b}
$$

with an additional equation governing the evolution of Q :

$$
\dot{Q}^2 f_{QQQ} + (\ddot{Q} + 3H\dot{Q}) f_{QQ} = 0.
$$
\n(29)

• Branch 3 (Γ_3) :

For the third branch, Q depends on both the time-dependent function $\gamma(t)$ and the scale factor $a(t)$:

$$
Q = -6H^2 + \frac{3\gamma}{a^2}H + \frac{3\dot{\gamma}}{a^2},\tag{30}
$$

with the corresponding Friedmann and Raychaudhuri equations:

$$
3H^2 f_Q + \frac{1}{2} \left(f - Q f_Q \right) - \frac{3\gamma \dot{Q} f_{QQ}}{2a^2} = \rho_m \,, \tag{31a}
$$

$$
-2\frac{d}{dt}\left(Hf_Q\right) - 3H^2f_Q - \frac{1}{2}\left(f - Qf_Q\right) + \frac{\gamma \dot{Q}f_{QQ}}{2a^2} = 0\,,\tag{31b}
$$

Here, the equation for Q evolution is given by:

$$
\dot{Q}^2 f_{QQQ} + \left[\ddot{Q} + \left(H + \frac{2\dot{\gamma}}{\gamma} \right) \dot{Q} \right] f_{QQ} = 0. \tag{32}
$$

On the redefinition $\gamma \to a^2 \gamma^4$ $\gamma \to a^2 \gamma^4$, the equations become

$$
Q = -6H^2 + 9\gamma H + 3\dot{\gamma},\tag{33a}
$$

$$
3H^2 f_Q + \frac{1}{2} \left(f - Q f_Q \right) - \frac{3\gamma \dot{Q} f_{QQ}}{2} = \rho_m \,, \tag{33b}
$$

$$
-2\frac{d}{dt}\left(Hf_Q\right) - 3H^2f_Q - \frac{1}{2}\left(f - Qf_Q\right) + \frac{\gamma \dot{Q}f_{QQ}}{2} = 0\,,\tag{33c}
$$

$$
\dot{Q}^2 f_{QQQ} + \left[\ddot{Q} + \left(5H + \frac{2\dot{\gamma}}{\gamma} \right) \dot{Q} \right] f_{QQ} = 0. \tag{33d}
$$

The energy conservation for pressureless matter reads as $\dot{\rho} = -3H\rho$ and it can be separately integrated into

$$
\rho_m = \rho_{m0} \left(\frac{a(t)}{a_0} \right)^3 \rho_{m0} (1+z)^3.
$$
\n(34)

That the energy conservation equation remains the same as in GR is due to the fact that particles still follow geodesics governed by the Levi-Civita connection [\[65,](#page-30-14) p.3].

VI. RECONSTRUCTING ΛCDM-MIMICKING $f(Q)$ WITH Γ_1

For the connection Γ_1 , it is possible to formulate a general reconstruction method for $f(Q)$ gravity. By combining Eqs. $(26a)$ and $(26b)$, we obtain

$$
2\frac{d}{dt}\left(Hf_Q\right) = -\rho_m\,,\tag{35}
$$

which can be expressed in terms of the redshift z

$$
\frac{d}{dz}(f_Q H) = \frac{\rho_{m0}(1+z)^2}{2H(z)}\,. \tag{36}
$$

Using the chain rule and noting that for Γ_1 , $Q = -6H^2$ we can express f_Q as

$$
f_Q = -\frac{f'(z)}{12H(z)H'(z)},
$$
\n(37)

where a ' denotes derivative with respect to z. If the background evolution $H(z)$ is known, Eq.[\(36\)](#page-7-2) can now be integrated to obtain $f_Q(z)$. With the relation [\(37\)](#page-7-3), we can now integrate Eq.[\(36\)](#page-7-2) to obtain

$$
f'(z) = -18H_0^2 \Omega_{m0} H'(z) \int \frac{(1+z)^2}{H(z)} dz + A_1 H'(z), \qquad (38)
$$

where A_1 is a constant of integration and we have used $\Omega_{m0} = \frac{\rho_{m0}}{3H_0^2}$. A second integration yields

$$
f(z) = -18H_0^2 \Omega_{m0} \int \left(H'(z) \int \frac{(1+z)^2}{H(z)} dz \right) dz + A_1 H(z) + A_2 , \qquad (39)
$$

where A_2 is another constant of integration.

As an example, let us now investigate the case of a ΛCDM-like evolution [\(7\)](#page-3-0). Then we have

$$
f(z) = -\frac{12\Omega_{m0}}{c_1} \int H'(z)H(z)dz + A_1H(z) + A_2 = -\frac{6\Omega_{m0}}{c_1}H^2(z) + A_1H(z) + A_2.
$$
 (40)

Redefining $\frac{A_1}{\sqrt{6}} \to \beta$, $A_2 \to -2\Lambda$, the reconstructed $f(Q)$ takes the form

$$
f(Q) = -2\Lambda + \frac{\Omega_{m0}}{c_1}Q + \beta\sqrt{-Q}.
$$
\n(41)

Note that the reconstruction method does not impose any constraints on the sign of the coefficient β or the the cosmological constant term Λ.

⁴ Essentially what we do is to redefine $\gamma \to \tilde{\gamma} = \frac{\gamma}{a^2}$; since the dynamical connection function $\gamma(t)$ is a free function of time, we have this freedom of redefinition.

A. Identifying the free parameters in the reconstructed $f(Q)$ (Γ₁)

The way the reconstructed $f(Q)$ is expressed in Eq.[\(41\)](#page-7-4) may give one an impression that the reconstructed $f(Q)$ is characterized by four independent parameters $\Omega_{m0}, c_1, \beta, \Lambda$. However, this is not the case. In this subsection we identify the free parameters of the reconstructed $f(Q)$ [\(41\)](#page-7-4).

For the reconstructed theory [\(41\)](#page-7-4), writing the Friedmann equation [\(26a\)](#page-6-1) at $z = 0$

$$
\frac{3H_0^2\Omega_{m0}}{c_1} = \rho_{m0} + \Lambda \,, \quad \Rightarrow \quad \frac{\Omega_{m0}}{c_1} = \Omega_{m0} + \frac{\Lambda}{3H_0^2} \,, \quad \Rightarrow \quad \Omega_{m0} = \frac{2\Lambda}{3H_0^2} \left(\frac{1+q_0}{1-2q_0}\right) \,. \tag{42}
$$

 5 where at the last step we used the relation [\(10\)](#page-3-2). One can get the same equation from the Raychaudhuri equation [\(26b\)](#page-6-1) as well, which is expected, since in the presence of a nondynamical connection like Γ_1 , the Friedmann and the Raychaudhuri equations are not independent, given the conservation of the energy-momentum tensor. For a particular cosmological trajectory specified by a given value of c_1 (which delivers the observed value q_0), the above equation determines the value of the matter abundance parameter Ω_{m0} in terms of the model parameter Λ . In other words, the model parameter Λ , though not affecting the effective gravitational constant $\kappa_{\text{eff}} = \frac{1}{f_Q}$, actually determines the value of the present day matter abundance Ω_{m0} . Alternatively, one can demand that the class of reconstructed $f(Q)$ [\(41\)](#page-7-4) to deliver any preferred value of the present day matter abundance Ω_{m0} by suitably choosing the value of the model parameter Λ.

Therefore the reconstructed $f(Q)$ [\(41\)](#page-7-4) is characterized by only two free model parameters, namely β and Λ , which are related respectively to the effective gravitational coupling and the present day value of the matter abundance. Taking into account the relation [\(10\)](#page-3-2), the reconstructed $f(Q)$ that reproduces the ΛCDM-like cosmic evolution [\(11\)](#page-3-3) can ultimately be expressed as

$$
f(Q) = -2\Lambda + \left(\frac{\Lambda/H_0^2}{1 - 2q_0}\right)Q + \beta\sqrt{-Q}.
$$
 (43)

For the theory [\(43\)](#page-8-1), the Friedmann equation is

$$
3H^2 \left(\frac{\Lambda/H_0^2}{1 - 2q_0} \right) = \rho_{m0} (1 + z)^3 + \Lambda \,. \tag{44}
$$

If we demand that at high redshift, GR acts as a cosmological past attractor for the theory, then the relation $3H^2 \approx \rho_m$ must hold at high redshift, which demands setting the value of the model parameter Λ as $\Lambda = H_0^2(1 - 2q_0)$. In that case we have the following identification

$$
\{\Omega_{m0}, \Omega_{\Lambda 0}\} = \left\{\frac{2}{3}(1+q_0), \frac{1}{3}(1-2q_0)\right\},\tag{45}
$$

where we have defined $\Omega_{\Lambda 0} \equiv \frac{\Lambda}{3H_0^2}$. For this particular case, the model parameter Λ as well as the present day matter abundance Ω_{m0} is completely determined in terms of a single quantity, namely the cosmographic parameter q_0 , whose value can be estimated in a model-independent manner from cosmographic data sets. β is the only free parameter in the reconstructed theory that could not be determined in terms of q_0 .

Note that, setting $\Lambda = H_0^2(1 - 2q_0)$ is not necessary a-priori as long as obtaining a particular Λ CDM-like solution giving rise to the required value of q_0 (see [\(11\)](#page-3-3)) is concerned. The requirement $\Lambda = H_0^2(1-2q_0)$ comes from demanding that GR acts as a past cosmological attractor. If we do, however, set $\Lambda = H_0^2(1 - 2q_0)$, then the reconstructed $f(Q)$ [\(43\)](#page-8-1) simplifies to

$$
f(Q) = -2H_0^2(1 - 2q_0) + Q + \beta\sqrt{-Q}.
$$
\n(46)

The above $f(Q)$ admits the Λ CDM solution [\(11\)](#page-3-3) with the present day value of the deceleration parameter q_0 , as well as admits GR (more precisely, STEGR) as the cosmological past attractor.

One might wonder if successive derivatives of the Raychaudhuri equation might give rise to a new constraint that help us determine β. But this is not the case. Since Raychaudhuri equation is identically satisfied by the reconstructed $f(Q)$ [\(43\)](#page-8-1) for all z, all it's successive derivatives are satisfied as well. β is truly a free parameter.

⁵ Even though the Friedmann equation at $z = 0$ [\(42\)](#page-8-2) looks very close to that in GR, in general it is different from GR. This is because in general one cannot naively set $\Omega_{m0} = c_1$, as neither of them are the free parameters as we have shown in this section.

B. Constraints on the free model parameters of the reconstructed $f(Q)$ (Γ₁)

Now that we have identified the free model parameters in the reconstructed $f(Q)$ [\(43\)](#page-8-1), namely β and Λ, in this subsection we try to find possible constraints they need to satisfy.

• Eq.[\(42\)](#page-8-2) pins down the sign of the model parameter Λ . Since $q_0 \approx -0.55$ and $0 < \Omega_{m0} < 1$,

$$
0 < \frac{\Lambda}{H_0^2} < \frac{3}{2} \left(\frac{1 - 2q_0}{1 + q_0} \right) \,. \tag{47}
$$

• One must have $f_Q > 0$, otherwise the effective gravitational constant would be negative [\[64\]](#page-30-13). Substituting the explicit form of $f(Q)$ gives the condition

$$
\frac{\Omega_{m0}}{c_1} > \frac{\beta}{2\sqrt{6}H}, \qquad \text{or} \qquad \frac{\beta}{H_0} < 2\sqrt{6} \frac{\Lambda/H_0^2}{1 - 2q_0} \, h(z) \, \forall z \,. \tag{48}
$$

where in the last step we have used Eqs.[\(42\)](#page-8-2) and [\(10\)](#page-3-2). This, however, does not pin down the sign of β .

• One has the following two conditions coming from the Friedmann and the Raychaudhuri equation [\(26a\)](#page-6-1) and [\(26b\)](#page-6-1)

$$
f - 2Qf_Q = 2\rho_m > 0, \qquad f_Q + 2Qf_{QQ} = \frac{\rho_m}{2H^2(1+q)} > 0.
$$
\n(49)

However, taking into account [\(10\)](#page-3-2) and [\(42\)](#page-8-2), one can check that these equations are satisfied. Therefore, they do not put any additional constraint on the free model parameters.

C. Effective gravitational coupling in the reconstructed $f(Q)$ (Γ₁)

Lets us consider the $f(Q)$ in Eq.[\(46\)](#page-8-3). The effective gravitational constant is $\kappa_{\text{eff}} = \frac{1}{f_Q}$ [6](#page-9-1) with

$$
f_Q(z) = 1 - \frac{\beta}{2\sqrt{-Q}} = 1 - \frac{\beta/H_0}{2\sqrt{6}h(z)}.
$$
\n(50)

Now, suppose we consider the $f(Q)$ theory that reproduces the Λ CDM-like cosmology with $q_0 = -0.55$, which corresponds to $c_1 = 0.3$. Because we have chosen the $f(Q)$ form from Eq.[\(46\)](#page-8-3), this also implies from [\(45\)](#page-8-4) that ${\Omega_{m0}, \Omega_{\Lambda0}} = {0.3, 0.7}$. Amazingly, we have arrived at the same value of Ω_{m0} and $\Omega_{\Lambda0}$ even without the underlying theory being GR. To respect the physical viability condition (48) for all values of z, one needs to enforce the following condition on the model parameter β

$$
\frac{\beta}{H_0} < 2\sqrt{6} \, h_{\text{min}} = 2\sqrt{6} \times 0.7 \approx 3.43 \,. \tag{51}
$$

For various values of the model parameter β (taken from [\[17\]](#page-29-13)), all of which satisfy the above mentioned condition, we show in Fig[.1](#page-10-1) the evolution of the effective gravitational constant, along with that for the actual General Relativistic Λ CDM model. It is possible to constrain the quantity $\kappa_{\text{eff}} - 1$ at $z = 0$ using astrophysical datasets, which is expected to put some constraints on the model parameter β .

D. Stability of a solution in the reconstructed $f(Q)$ for Γ_1

In general, a solution $H(t)$ is characterized by a phase trajectory in the phase space, whereas fixed points characterize the asymptotic states of the solutions that are interpreted as cosmological epochs. For example, in the ΛCDM phase space, ΛCDM solutions are trajectories, whereas matter and Λ-dominated phases are fixed points (see [\[66,](#page-30-15) Fig.2]). In

$$
^6
$$
 $\kappa_{\text{eff}}=8\pi G_{\text{eff}}=\frac{8\pi G}{f_Q}.$ Since we are working in the geometrized unit system, $\kappa_{\text{eff}}=\frac{1}{f_Q}.$

FIG. 1: Evolution of $\kappa_{\rm eff}$ for the 1-parameter family of reconstructed $f(Q)$ [\(46\)](#page-8-3) that reproduces a ACDM-like cosmic evolution [\(11\)](#page-3-3) for the connection $\Gamma_1.$ The evolution shows (STE)GR as a cosmological past attractor.

this section, following the idea of [\[67,](#page-30-16) [68\]](#page-30-17), we analyze the stability of the ΛCDM cosmological solution as a whole against small homogeneous and isotropic perturbations.

Eq.[\(26b\)](#page-6-1) can be written as

$$
2\dot{H}f_Q - 24\dot{H}H^2f_{QQ} + 6H^2f_Q + \frac{1}{2}f = 0.
$$
\n(52)

For a given $f(Q)$, this is a differential equation in $H(t)$. Consider any given solution of this equation $H(t)$, and take a homogeneous and isotropic perturbation around it: $H(t) \to H(t) + \delta H(t)$. Substitute it back to the above equation. Keeping terms of only linear order we get (keep in mind that $Q = -6H^2 \Rightarrow \delta Q = -12H\delta H$)

$$
\delta H + \lambda(Q)\delta H = 0. \tag{53}
$$

where

$$
\lambda(Q) = 3H - 36H\dot{H}\frac{f_{QQ} - 4H^2 f_{QQQ}}{f_Q - 12H^2 f_{QQ}} = 3H + 36(1+q)H^3 \frac{f_{QQ} + \frac{2}{3}Qf_{QQQ}}{f_Q + 2Qf_{QQ}}.
$$
\n(54)

 $\lambda(Q)$ determines the stability of the given solution $H(t)$. For stability of this solution with respect to small homogeneous and isotropic perturbations, one needs $\lambda(Q) > 0$.

Consider, now, the theory given by Eq. (43) . For this theory, we have

$$
\lambda(Q) = 3H \,,\tag{55}
$$

which is always positive for an expanding cosmology. This implies that within the theory (43) , any expanding cosmological solution, ΛCDM-like solution included, is always stable[7](#page-10-2) . If we start with a particular trajectory and move to a neighboring trajectory in its vicinity by a small homogeneous and isotropic perturbation at any point during its course of evolution, the new trajectory will show the same qualitative behavior (notably, the same past and future attractor) as the original trajectory we started with.

E. Robustness of the reconstructed $f(Q)$ against slight deviation from the condition $j = 1$

Building upon our discussion from Sect[.II,](#page-2-0) we will now demonstrate explicitly the robustness of our reconstructed $f(Q)$ in the framework of Γ_1 , by investigating the a slightly more general scenario $j = 1 + \varepsilon = const.$ for "small" ε . This will be done by checking explicitly that there exists a smooth matching between the non-metricity function Q and

⁷ Even though we reconstructed the theory [\(43\)](#page-8-1) starting from the ΛCDM-like evolution, the theory can in principle admit as a solution not only the ΛCDM-like evolution but many other different kinds of evolution.

the reconstructed $f(Q)$ for the "almost Λ CDM" case to their exactly Λ CDM-mimicking counterpart^{[8](#page-11-0)}. The smoothness of the non-metricity directly follows from that of the Hubble rate via [\(25\)](#page-6-2), and it reads as

$$
Q = -6H^2 = -6H_{\Lambda \text{CDM}}^2 + \frac{2\varepsilon[(z^3 + 3z^2 + 3z + 2)c_1 - 1]}{3}\ln(1+z) + O(\varepsilon^2). \tag{56}
$$

Next, for assessing the Raychaudhuri equation [\(26b\)](#page-6-1), we recall the deceleration function [\(5\)](#page-2-4) for the cosmic rate reconstructed from a general constant jerk, where we should also use [\(34\)](#page-7-5), and re-write the redshift by using the Friedmann equation [\(26a\)](#page-6-1) with the conservation of dark matter, that is

$$
(1+z) = \left(\frac{\rho_m}{\rho_{m0}}\right)^{1/3} = \left(\frac{f - 2Qf_Q}{2\rho_{m0}}\right)^{1/3}.
$$
\n(57)

Then, from the Raychaudhuri equation

$$
2H^2(1+q)f_Q - 24(1+q)H^4f_{QQ} - \frac{f}{2} - Qf_Q = 0 \quad \Rightarrow \quad \frac{Q}{3}(1+q)(f_Q + 2Qf_{QQ}) + \frac{f}{2} - Qf_Q = 0,
$$
 (58)

we can obtain the following closed-form differential equation for the reconstruction of the theory $f(Q)$:

$$
\left[\frac{k_1Q}{6} + (1 - c_1)(k_1 - k_2)H_0^2 \left(\frac{f - 2Qf_Q}{2\rho_{m0}}\right)^{k_2/3}\right](f_Q + 2Qf_{QQ}) + \frac{f}{2} - Qf_Q = 0.
$$
\n(59)

This equation can be recast in terms of the deviation ε from the exact Λ CDM-like jerk parameter

$$
\left[\frac{(3+\sqrt{9+8\varepsilon})Q}{12} + (1-c_1)\sqrt{9+8\varepsilon}H_0^2 \left(\frac{f-2Qf_Q}{2\rho_{m0}}\right)^{\frac{1}{2}-\frac{\sqrt{9+8\varepsilon}}{6}}\right](f_Q+2Qf_{QQ}) + \frac{f}{2} - Qf_Q = 0.
$$
 (60)

The non-linear nature of the above equation and its dependence on the value of the jerk parameter $j = 1 + \varepsilon$ should be appreciated. It allows for an explicit confirmation that it consistently reproduces the case $\varepsilon = 0$ with

$$
f(Q) = -2\Lambda + \frac{\Omega_{m0}}{c_1}Q + \beta\sqrt{-Q},\qquad(61)
$$

which is basically what we obtained in Eq.[\(41\)](#page-7-4). By plugging the latter into the reconstruction differential equation (60) which holds for a general constant j, we obtain

$$
\sqrt{9+8\varepsilon}\,\Omega_{m0}\left(\frac{1-c_1}{c_1}\right)\left(\frac{h^2}{c_1}-\frac{\Omega_{\Lambda 0}}{\Omega_{m0}}\right)^{\frac{1}{2}-\frac{\sqrt{9+8\varepsilon}}{6}}-\frac{3}{2}\frac{\Omega_{m0}}{c_1}h^2\left(\frac{\sqrt{9+8\varepsilon}}{3}-1\right)-3\Omega_{\Lambda 0}=0.\tag{62}
$$

For $|\varepsilon| \ll 1$, linearizing the left hand side of the above equation with respect to ε , we arrive at the following condition

$$
\frac{2\Omega_{m0}}{3c_1} \left[(1 - c_1) \left(2 - \ln \left| \frac{h^2}{c_1} - \frac{\Omega_{\Lambda 0}}{\Omega_{m0}} \right| \right) - h^2 \right] \varepsilon + O(\varepsilon^2) \approx 0. \tag{63}
$$

Taking into consideration the expression [\(7\)](#page-3-0), the relations [\(10\)](#page-3-2) and [\(42\)](#page-8-2), the above condition can be expressed as

$$
\left[\frac{1}{3}(1-2q_0)(1-\ln(1+z)) - \frac{2}{3}(1+q_0)(1+z)^3\right] \varepsilon + O(\varepsilon^2) \approx 0. \tag{64}
$$

For the term linear in ε , the quantity inside the square bracket is regular for $-1 < z < \infty$. Therefore the term as a whole remains small, of the order $\mathcal{O}(\varepsilon)$. Hence, the reconstruction solution [\(41\)](#page-7-4) for $j = 1$ "almost solves" the reconstruction differential equation for a constant $j = 1 + \varepsilon$ for any redshift $-1 < z < \infty$. Hence, we conclude that our reconstructed LCDM-like $f(Q)$ [\(41\)](#page-7-4) holds within small observational error in the astrophysical estimate of j, and that "almost Λ CDM" can be achieved by "almost the same $f(Q)$ ".

Note that, we could not have inferred the same if the quantity inside the square bracket was divergent at any redshift within the range $-1 < z < \infty$. Also, one can consider more accurate variation in the astrophysical estimate of the jerk parameter e.g. $j(z) = 1 + \varepsilon(z)$ (e.g. see [\[43\]](#page-29-27)) or $j(z) = 1 + \varepsilon q(z)$, which also characterize an "almost ΛCDM" evolution. Such deviations are not considered here.

⁸ For the case of Γ_1 , the dynamical connection function is always set to zero as a geometrical property while constructing the $f(Q)$ theory: this applies regardless of the cosmic history and consequently a variation on the astrophysical value of the jerk does not bear any consequence on it. For the case of Γ_2 which we will consider in the next section, the smooth matching of the dynamical connection function will also have to be taken into consideration, allowing for potential discrimination between the two scenarios.

VII. RECONSTRUCTING ΛCDM-MIMICKING $f(Q)$ WITH Γ_2

For Γ_2 , we were not able to formulate a generic reconstruction method unlike the Γ_1 case. Fortunately, we could perform the analytical reconstruction for the Λ CDM-like background evolution. Eq.[\(29\)](#page-6-3) can be written as

$$
0 = \frac{d}{dt} \left(\dot{Q} f_{QQ} \right) + 3H (\dot{Q} f_{QQ}) = \frac{d}{dt} \left(f_Q \right) + 3H \dot{f}_Q
$$

=
$$
\frac{d^2 f_Q}{dt^2} + 3H \frac{df_Q}{dt} = 0.
$$
 (65)

Eq.[\(65\)](#page-12-1) can be integrated for any generic cosmic history as $[38, Eq.(17)]$

$$
\frac{df_Q}{dt} = Ca^{-3}, \quad \frac{df_Q}{dz} = -C\frac{(1+z)^2}{H}, \quad f_Q(z) = -C\int \frac{(1+z)^2dz}{H(z)} + D\,,\tag{66}
$$

where C is a dimensionful integration constant and D is a dimensionless integration constant. From the second equality above, we see that the sign of the integration constant C determines how the effective gravitational coupling $\kappa_{\text{eff}} = \frac{1}{f_Q}$ evolves with time in an expanding universe. A positive (negative) C implies that the effective gravitational coupling decreases (increases) with time^{[9](#page-12-2)}.

Now, we try to reconstruct the Λ CDM-mimicking $f(Q)$ for Γ_2 . Therefore, in whatever follows in this section, H actually refers to the Λ CDM-like $H(z)$ as given in [\(7\)](#page-3-0), unless stated otherwise explicitly^{[10](#page-12-3)}. Then we get

$$
f_Q(z) = D - \frac{2c}{3c_1} \sqrt{c_1(z+1)^3 + (1-c_1)} = D - \frac{2c}{3c_1} h(z),\tag{67}
$$

where we have defined $c \equiv \frac{C}{H_0}$, which is now dimensionless. Taking the sum of [\(28a\)](#page-6-4) and [\(28b\)](#page-6-4):

$$
3\gamma \dot{Q} f_{QQ} - \rho_m - 2\frac{d}{dt}(f_Q H) = 0, \qquad (68)
$$

that can be re-written as

$$
3\gamma \dot{f}_Q - \rho_m - 2\frac{d}{dt}(f_Q H) = 0.
$$
\n(69)

Now use the conservation for dust and the result for the reconstruction from (67) :

$$
\left(\frac{4}{3}h - \frac{\gamma}{H_0} - \frac{Dc_1}{c}\right)\frac{2c}{c_1}\frac{\dot{h}}{H_0} - 3\Omega_{m0}(1+z)^3 = 0.
$$
\n(70)

For the time derivative of the Hubble function we use

$$
\frac{\dot{h}}{H_0} = (-h)(1+z)\frac{dh/dz}{H_0} = -\frac{3c_1(1+z)^3}{2}.
$$
\n(71)

Consequently, we reconstruct

$$
\frac{\gamma}{H_0} = \frac{4}{3}h + \frac{\Omega_{m0}}{c} - \frac{Dc_1}{c} \,. \tag{72}
$$

Therefore

$$
\frac{\dot{\gamma}}{H_0^2} = -2c_1(1+z)^3 = 2(1-c_1-h^2). \tag{73}
$$

Inserting the above equations into $Eq.(29)$ $Eq.(29)$, we get

$$
\frac{Q}{H_0^2} = -6h^2 + 9\frac{\gamma}{H_0}h + 3\frac{\dot{\gamma}}{H_0^2}
$$

= -6h^2 + 9h\left(\frac{4}{3}h + \frac{\Omega_{m0}}{c} - D\frac{c_1}{c}\right) + 6(1 - c_1 - h^2)
= 6(1 - c_1) + \frac{9h}{c}(\Omega_{m0} - c_1D). (74)

 $9 \ C = 0$ gives a constant effective gravitational coupling, which can be reduced to GR with a constant rescaling of the metric.

¹⁰ When there is a scope of confusion, we will specify the Λ CDM-like H as H_{Λ CDM.

Thus, both the reconstructed Λ CDM-like dynamical connection function and non-metricity for Γ_2 are linear in the Hubble parameter and monotonically increasing with respect to the redshift. Having in mind the previously established negativity of the constant c, we also understand that they are both allowed to vanish at some cosmic epoch as well as possibly switch their sign. For the nonmetricity, this constitutes a sharp difference than its behavior for Γ_1 , which is negative all along the comic history by construction [\(25\)](#page-6-2), and may provide a viable astrophysical route for discriminating among these scenarios. Next, using Eq.[\(67\)](#page-12-4) we get

$$
\frac{df}{dQ} = D + \frac{2c^2}{27c_1\left(\Omega_{m0} - c_1D\right)} \left[6(1 - c_1) - \frac{Q}{H_0^2}\right],\tag{75}
$$

which allows us to ultimately reconstruct the Λ CDM-mimicking $f(Q)$ theory for the connection Γ_2

$$
f(Q) = -2\Lambda + \left(D + \frac{4}{9} \frac{c^2 (1 - c_1)}{c_1 (\Omega_{m0} - c_1 D)}\right) Q - \frac{c^2}{27 c_1 (\Omega_{m0} - c_1 D) H_0^2} Q^2.
$$
 (76)

Screening mechanisms are likely to kill the effects of the cosmological constant Λ on galactic length scales, in which context therefore our reconstructed theory is of quadratic nature, constituting a realistic and healthy model [\[69,](#page-30-18) $Eq.(13)$.

A. Identifying the free parameters in the reconstructed $f(Q)$ (Γ₂)

The way the reconstructed $f(Q)$ is expressed in Eq.[\(76\)](#page-13-0) may give an impression that the reconstructed $f(Q)$ is characterized by five independent parameters $\Omega_{m0}, c_1, c, D, \Lambda$ (H_0 is not a "free" parameter as it is obtained from data in a model-independent way). However this is not the case. In this subsection we identify the free parameters of the reconstructed $f(Q)$ [\(76\)](#page-13-0).

Eqs.[\(28a\)](#page-6-4) and [\(28b\)](#page-6-4) can be combined such that γ is eliminated. In the resulting equation, using the reconstructed form of various quantities from Eqs.[\(67\)](#page-12-4), [\(72\)](#page-12-5), [\(73\)](#page-12-6), [\(74\)](#page-12-7), [\(75\)](#page-13-1), as well as the ΛCDM-like evolution [\(7\)](#page-3-0), after some steps of straightforward manipulations, we arrive at the following expression, which is, surprisingly, independent of the redshift z:

$$
4c^{2}(1-c_{1})^{2} + 3c_{1}^{2}D(-3(1-c_{1})D + 2\lambda) - 6c_{1}\lambda\Omega_{m0} + 9(1-c_{1})(\Omega_{m0})^{2} = 0,
$$
\n(77)

where $\lambda \equiv \Lambda/H_0^2$. The same equation can be obtained by evaluating the Friedmann equation [\(28a\)](#page-6-4) at $z = 0$. The above equation can be solved for Ω_{m0}

$$
\Omega_{m0} = \frac{2}{3}\lambda \left(\frac{1+q_0}{1-2q_0}\right) \pm \frac{2}{3}\sqrt{(1+q_0)^2 \left(\frac{\lambda}{1-2q_0} - D\right)^2 - \frac{c^2}{3}(1-2q_0)},\tag{78}
$$

where at the last step we have used the relation [\(10\)](#page-3-2). For a particular cosmological trajectory specified by a given value of c_1 (which delivers the observed value q_0), Eq.[\(78\)](#page-13-2) determines the value of the matter abundance parameter Ω_{m0} in terms of the model parameters Λ, c, D . The model parameter Λ , although not affecting the effective gravitational constant $\kappa_{\text{eff}} = \frac{1}{f_Q}$, contributes towards determining the value of the present day matter abundance Ω_{m0} via Eq.[\(78\)](#page-13-2). Alternatively, one can demand that the class of reconstructed $f(Q)$ [\(76\)](#page-13-0) to deliver any preferred value of the present day matter abundance Ω_{m0} by suitably choosing the values of the free model parameters Λ , c, D that satisfy Eq.[\(77\)](#page-13-3). Therefore the reconstructed $f(Q)$ [\(76\)](#page-13-0) is characterized by three free model parameters, namely c, D and Λ. Taking into account the relation [\(10\)](#page-3-2), the reconstructed $f(Q)$ that reproduces the ΛCDM-like cosmic evolution [\(11\)](#page-3-3) can ultimately be expressed as

$$
f(Q) = -2\Lambda + \left[D + \frac{1}{3} \left(\frac{1 - 2q_0}{1 + q_0} \right) \left(\frac{2c^2}{3\Omega_{m0} - 2(1 + q_0)D} \right) \right] Q - \left[\frac{c^2}{6(1 + q_0)(3\Omega_{m0} - 2(1 + q_0)D)H_0^2} \right] Q^2, \tag{79}
$$

with Ω_{m0} given by Eq.[\(78\)](#page-13-2). This clearly shows that the reconstructed $f(Q)$ is a 3-parameter model; the free parameters being Λ, c, D .

The parameter D is actually related to the redshift z_* at which the reconstructed $f(Q)$ coincides with STEGR. Supposing that $f_Q(z_*) = 1$ at some redshift z_* , we get from Eq.[\(67\)](#page-12-4)

$$
D = 1 - \frac{2|c|}{3c_1}h_*,\tag{80}
$$

where $h_* = h(z_*)$. Therefore, instead of D, z_* (or equivalently h_*) can also be used as a free parameter of the reconstructed $f(Q)$ model.

One may wonder, and justifiably so, what is the meaning of the two "branches" of Ω_{m0} given by the ' \pm ' sign in Eq. [\(78\)](#page-13-2). Since Λ , c, D are free parameters, it is possible for both branches to produce a physically consistent value of $\Omega_{m0} \in (0,1)$ for some range of values of the parameters Λ, c, D . This particular question demands some explanation. Applying Eq. (80) to Eq. (72) , we get

$$
\frac{\gamma}{H_0} = \frac{4h}{3} + \frac{2h_*}{3} + \frac{\Omega_{m0} - c_1}{c} \,. \tag{81}
$$

Eq.[\(81\)](#page-14-0) shows that the different value of Ω_{m0} leads to different constant term in the expression for γ/H_0 . Inserting D from Eq. (80) into Eq. (74) , we get

$$
\frac{Q}{H_0^2} = 6(1 - c_1 - 6hh_*) - \frac{9h(\Omega_{m0} - c_1)}{|c|} = 6(1 - c_1) - h\left(\frac{9(\Omega_{m0} - c_1)}{|c|} + 6h_*\right),\tag{82}
$$

which shows that Q always varies with h linearly, but with different rates for different values of Ω_{m0} . In other words, different value of Ω_{m0} , can lead to the same kinematic Λ CDM evolution, while different quantitative evolution of $Q(z)^{11}$ $Q(z)^{11}$ $Q(z)^{11}$

Since Λ , c, D are free parameters of the reconstructed $f(Q)$ [\(79\)](#page-13-5), we have some freedom to choose them. Let us consider the parameter choice

$$
\lambda = 1 - 2q_0, \qquad c^2 = \frac{3(1 - D)^2 (1 + q_0)^2}{1 - 2q_0}.
$$
\n(83)

It can be verified by straightforward substitution in Eq.[\(78\)](#page-13-2) that the above parameter choice yields a unique value for Ω_{m0} ; $\Omega_{m0} = \frac{2}{3}(1+q_0) = c_1$. The particular choice of the model parameters in Eq.[\(83\)](#page-14-2) gives us back the same identification as in Eq.[\(45\)](#page-8-4)

$$
\{\Omega_{m0}, \Omega_{\Lambda 0}\} = \left\{\frac{2}{3}(1+q_0), \frac{1}{3}(1-2q_0)\right\}.
$$
 (84)

Moreover, it can be checked that the choice [\(83\)](#page-14-2) also make the coefficient of the linear term in the reconstructed $f(Q)$ [\(79\)](#page-13-5) unity, so that the reconstructed $f(Q)$ is of the form $f(Q) = -2\Lambda + Q + F(Q)$ with $F(Q) \propto -Q^2$. Indeed, in many works found in the literature $f(Q)$ is indeed taken in the form $f(Q) = Q + F(Q)$ [\[16–](#page-29-4)[18,](#page-29-34) [38,](#page-29-20) [70\]](#page-30-19), to make it explicit that at some limit the $f(Q)$ matches with the STEGR with the correct Newtonian gravitational constant ^{[12](#page-14-3)}. For the parameter choice (83) , the reconstructed $f(Q)$ [\(79\)](#page-13-5) simplifies to

$$
f(Q) = -2H_0^2(1 - 2q_0) + Q - \frac{1}{4H_0^2} \left(\frac{1 - D}{1 - 2q_0}\right) Q^2.
$$
 (85)

The above $f(Q)$ is a 1-parameter $f(Q)$ model, parametrized by the single model parameter D, that admits the Λ CDM solution [\(11\)](#page-3-3) with the present day value of the deceleration parameter q_0 for the connection branch Γ_2 .

B. Constraints on the model parameters of the reconstructed $f(Q)$ (Γ₂)

Now that we have identified the free model parameters in the reconstructed $f(Q)$ [\(79\)](#page-13-5), namely Λ, c, D , in this subsection we find the range of values of the parameters for which f_Q can be unity at some particular redshift $z = z_* > 0$ and f_Q is always positive. The requirement that $f_Q(z_*) = 1$ at some redshift $z_* > 0$ is a "desirable" physical demand ensuring that the reconstructed theory coincides with STEGR at some point in the past, although not strictly necessary if just reconstructing a ΛCDM-like background evolution is concerned. Beyond the cutoff redshift z∗, there is no point of considering a modified gravity anymore. In other words, we say that the effect of gravity modification becomes apparent only after the cutoff redshift $z = z_*$. The condition $f_Q(z) > 0$ ensures the positivity of effective gravitational coupling.

¹¹ Different values of Ω_{m0} implies different rate of evolution for $Q(z)$, which may lead to different values of Ω_m at high redshift.

¹² Again, we remind, we are using a geometrized unit system with $\kappa = 8\pi G = 1$

• Firstly, we note from Eq. $(67)^{13}$ $(67)^{13}$ $(67)^{13}$ $(67)^{13}$ that if $C > 0$, then f_Q definitely becomes negative at some redshift irrespective of the value of D. To ensure that $f_Q > 0$ for all redshift, we demand that $C < 0$, so that Eq.[\(67\)](#page-12-4) can be written as

$$
f_Q = \frac{2|c|}{3c_1}h + D.
$$
\n(86)

where we remind that $c \equiv C/H_0$. Since $h(z)$ is a monotonically increasing function of z for a Λ CDM-like evolution $(Eq.(11)), h_* = h(z_*) > h(z = 0) = 1$ $(Eq.(11)), h_* = h(z_*) > h(z = 0) = 1$ $(Eq.(11)), h_* = h(z_*) > h(z = 0) = 1$. This gives an upper bound on D

$$
D < 1 - \frac{2|c|}{3c_1} \,. \tag{87}
$$

• Inserting D from Eq.[\(80\)](#page-13-4) into Eq.[\(86\)](#page-15-1), we get

$$
f_Q = 1 + \frac{2|c|}{3c_1}(h - h_*)\,. \tag{88}
$$

It follows from Eq.[\(88\)](#page-15-2) that $f_Q = 0$ when

$$
h = h_* - \frac{3c_1}{2|c|} \,. \tag{89}
$$

Hence, f_Q will never vanish for $h > 0$ if h in the left–hand–side of the above equation is negative or zero, yielding the condition

$$
\frac{3c_1}{2|c|} \ge h_* \,. \tag{90}
$$

The above condition, in conjunction with Eq. (80) , implies

$$
D \ge 0,\tag{91}
$$

which is a lower bound on D.

• Combining [\(87\)](#page-15-3) and [\(91\)](#page-15-4), we get the following bound on the model parameters

$$
c < 0, \qquad 0 \le D < 1 - \frac{2|c|}{3c_1} < 1. \tag{92}
$$

If one trades D for z_* to be the free parameter, then one can instead write from Eq.[\(90\)](#page-15-5) the following bounds on the model parameters c, z_*

$$
-\frac{3c_1}{2h_*} = -\frac{3c_1}{2h(z_*)} \le c < 0. \tag{93}
$$

Note that, $f_Q(z) > 0$ can be directly verified from the positivity of D, or by substituting [\(90\)](#page-15-5) into [\(88\)](#page-15-2). The bound [\(92\)](#page-15-6) or [\(93\)](#page-15-7) ensure both $f_Q(z) > 0$ for all z and coincidence with STEGR at some z_* in the past.

• As of now no bound has been given on the model parameter Λ . Note that the relation [\(78\)](#page-13-2) is of the form $\Omega_{m0} = \Omega_{m0}(\Lambda, c, D)^{14}$ $\Omega_{m0} = \Omega_{m0}(\Lambda, c, D)^{14}$ $\Omega_{m0} = \Omega_{m0}(\Lambda, c, D)^{14}$. The condition for a real valued Ω_{m0} puts the following constraint on Λ .

$$
\left| \frac{\Lambda}{H_0^2 (1 - 2q_0)} - D \right| \ge \frac{|c|}{1 + q_0} \sqrt{\frac{1 - 2q_0}{3}} = \frac{2|c|}{3c_1} \sqrt{1 - c_1} \,. \tag{94}
$$

The condition $0 < \Omega_{m0} < 1$ puts a further bound on Λ . However, its form is complicated and we deem in not necessary to write down the explicit inequality here. The point we make here is that, given a choice of values of the parameters c and D according to (92) , the choice of the free parameter Λ is not completely arbitrary. It is to be chosen such that one gets a real valued Ω_{m0} from Eq.[\(78\)](#page-13-2) and that $0 < \Omega_{m0} < 1$.

¹³ To play with f_Q , we have made use of the simpler expression [\(67\)](#page-12-4): $f_Q = D - \frac{2c}{3c_1}h$. We could have also determined f_Q from the reconstructed $f(Q)$ [\(79\)](#page-13-5), which gives $f_Q = D + \frac{4}{9} \frac{c^2 (1 - c_1)}{c_1 (\Omega_{m0} - c_1 D)} - \frac{2c^2}{27c_1 (\Omega_{m0})}$ $\frac{2c^2}{27c_1(\Omega_{m0}-c_1D)}\frac{Q}{H_0^2}$. Using the expression of Q along a Λ CDM trajectory (74) , we see that the two expressions for f_Q are in fact equivalent.

¹⁴ H_0 and q_0 are not a model parameters as they are obtained from cosmographic datasets in a model-independent manner.

• Let us see what the above bounds imply for the simpler 1-parameter reconstructed $f(Q)$ [\(85\)](#page-14-4). Firstly, for this choice one has $\Omega_{m0} = \frac{2}{3}(1+q_0)$ with $q_0 \approx -0.55$; see Eq.[\(84\)](#page-14-5). So, the condition $0 < \Omega_{m0} < 1$ is automatically satisfied.

Substituting D from Eq.[\(80\)](#page-13-4) into the expression for c in Eq.[\(83\)](#page-14-2)

$$
1 = \frac{3}{1 - 2q_0} h_*^2 = \frac{1}{1 - c_1} h_*^2.
$$
\n(95)

Since $c_1 < 1$, $h_* = h(z_*) > 1$ for any $z_* > 0$, meaning the right hand side of the above equation is always > 1. In that case, clearly the above equality can never be satisfied. This suggests that the value of c from Eq.[\(83\)](#page-14-2) is not compatible with the value of D in Eq.[\(80\)](#page-13-4). Physically, it means that at no point in the past the reconstructed f(Q) [\(85\)](#page-14-4) coincides with STEGR. However, the above equation can be satisfied if $h_* = h(z_*) < 1$, which can be achieved when $z_* < 0$. In the latter case the reconstructed $f(Q)$ [\(85\)](#page-14-4) may coincide with STEGR at some point in the future.

C. Effective gravitational coupling in the reconstructed $f(Q)$ (Γ₂)

Let us first consider the 1-parameter reconstructed $f(Q)$ in Eq.[\(85\)](#page-14-4). The effective gravitational constant is $\kappa_{\text{eff}} = \frac{1}{f_Q}$ with

$$
f_Q = 1 - \frac{1}{2} \left(\frac{1 - D}{1 - 2q_0} \right) \frac{Q}{H_0^2} \,. \tag{96}
$$

with $\frac{Q}{H_0^2}$ given from Eq.[\(74\)](#page-12-7). Taking into account the relations [\(10\)](#page-3-2), [\(83\)](#page-14-2) and [\(84\)](#page-14-5), as well as the bounds in [\(92\)](#page-15-6), we see that $f(Q)$ can be written as

$$
f_Q(z) = D + \sqrt{\frac{3}{1 - 2q_0}} \left(1 - D\right) h(z) \,. \tag{97}
$$

Since $0 \leq D < 1$ [\(92\)](#page-15-6), the reconstructed $f(Q)$ asymptote to STEGR at $z \to -1$ irrespective of whatever is the value of D, implying that (STE)GR acts not as a past attractor but as a future attractor. This conclusion is in line with what we observed at the last item of the previous subsection.

Suppose we consider one such $f(Q)$ theory that reproduces the ΛCDM-like cosmology with $q_0 = -0.55$, which corresponds to $c_1 = 0.3$. Because we have chosen the $f(Q)$ form from Eq.[\(85\)](#page-14-4), this also implies from [\(84\)](#page-14-5) that ${\Omega_{m0},\Omega_{\Lambda0}} = {0.3,0.7}$. For four different values of D, the evolution of the effective gravitational constant is shown in Fig[.2\(a\),](#page-16-0) along with that for the actual General Relativistic Λ CDM model.

FIG. 2: (a) Evolution of $\kappa_{\rm eff}$ for the 1-parameter family of reconstructed $f(Q)$ [\(85\)](#page-14-4) that reproduces a Λ CDM-like cosmic evolution [\(11\)](#page-3-3) for the connection Γ_2 . The evolution shows STEGR as a cosmological future attractor. (b) Evolution of $\kappa_{\rm eff}$ for the reconstructed $f(Q)$ [\(79\)](#page-13-5) that reproduces a Λ CDM-like cosmic evolution [\(11\)](#page-3-3) for the connection Γ_2 . $z_* = 5$ is chosen.

We expect a matter dominated phase to be there at some high redshift. If the reconstructed $f(Q)$ does not reduce to STEGR around that redshift, there might be very different signatures coming from the analysis of matter perturbations. This may potentially render the connection branch Γ_2 ultimately non-viable, even though it reproduces a Λ CDM-like evolution with correct values of q_0 , Ω_{m0} .

Let us instead consider the generic reconstructed $f(Q)$ [\(79\)](#page-13-5), which is parametrized by three parameters Λ, c, D . In general f_Q depends only on c and D

$$
f_Q(z) = D - \frac{2c}{3c_1}h(z).
$$
\n(98)

If one trades D for z_* to be the free parameter, then one can write $((88))$ $((88))$ $((88))$

$$
f_Q(z) = 1 - \frac{c}{1 + q_0} [h(z) - h(z_*)].
$$
\n(99)

We can choose any value of z_* that we want, but must choose a value of c satisfying the bound in [\(93\)](#page-15-7) to ensure that the reconstructed $f(Q)$ coincides with STEGR at some high redshift, while also satisfying $f_Q(z) > 0$ for all z. For $z_* = 5$ and different values of c, the evolution of the effective gravitational coupling is shown in Fig[.2\(b\).](#page-16-1)

VIII. RECONSTRUCTING ΛCDM-MIMICKING $f(Q)$ WITH Γ_3

To reconstruct the Λ CDM-mimicking $f(Q)$ for the case of Γ_3 , we had to ultimately resort to numerical analysis. Eq.[\(33d\)](#page-7-6) can be written as

$$
\frac{d^2 f_Q}{dt^2} + \left(5H + 2\frac{\dot{\gamma}}{\gamma}\right) \frac{df_Q}{dt} = 0,\tag{100}
$$

which can be integrated to give

$$
\frac{df_Q}{dt} = \frac{C}{a^5 \gamma^2}, \quad \frac{df_Q}{dz} = -\frac{C(1+z)^4}{\gamma^2 H}.
$$
\n(101)

Unlike the case of Γ_2 , for the case of Γ_3 the time-derivative of f_Q depends on γ , so that it can be integrated only if the evolution of γ is known. This brings in a lot of complications. The evolution equation for γ can be obtain by firstly combining Eqs. $(33b)$ and $(33c)$ such that f is eliminated:

$$
(\gamma + 2H)\dot{f}_Q + 2\dot{H}f_Q + \rho_m = 0.
$$
\n(102)

Using Eq. (101) , Eq. (102) can be written as

$$
\frac{C(1+z)^5(\gamma+2H)}{\gamma^2} + 2\dot{H}f_Q + \rho_m = 0.
$$
\n(103)

The expression for f_Q can be obtained by solving Eq.[\(103\)](#page-17-3) as

$$
f_Q = -\frac{C(1+z)^5(\gamma + 2H) + \gamma^2 \rho_m}{2\gamma^2 H} \,. \tag{104}
$$

Differentiating Eq.[\(103\)](#page-17-3) with respect to time, and then inserting the expression for f_Q from Eq.[\(104\)](#page-17-4) into the resulting equation, we get

$$
(1+z)^5 \gamma^3 \left(\ddot{H} + 3H\dot{H}\right)\rho_m + \gamma C \left(10\dot{H}H^2 + \dot{H}\left(\dot{\gamma} - 4\dot{H}\right) + 2H\ddot{H}\right) + \gamma^2 C \left(\ddot{H} + 5H\dot{H}\right) + 4\dot{\gamma}CH\dot{H} = 0. \tag{105}
$$

We see that we have been able to obtain an evolution equation for $\gamma(z)$ that is decoupled from $Q(z)$. Since $\rho_m(z) = \rho_{m0}(1+z)^3$, the above equation can be used to solve for $\gamma(z)$ once an $H(z)$ is supplied. Once we know the solution $\gamma(z)$, Eqs.[\(33a\)](#page-7-6) and [\(104\)](#page-17-4) can be used to solve for $Q(z)$ and $f_Q(z)$ respectively. One can, in principle, invert the expression of $Q(z)$ to solve for $z(Q)$, substitute it back into the expression of $f_Q(z)$ and get the reconstructed $f_Q(Q) = f_Q(z(Q))$, which can then be integrated to obtain $f(Q)$. Up to now, the relations we have derived are true for any generic cosmic history $H(z)$. Lets us now attempt the above exercise for a Λ CDM background. Therefore, in

the rest of this section, unless stated otherwise explicitly, H , q etc corresponds to that of a Λ CDM-like evolution as specified in Eq. $(11)^{15}$ $(11)^{15}$ $(11)^{15}$ $(11)^{15}$.

At this point we impose the condition that the Universe undergoes a ΛCDM-like evolution [\(7\)](#page-3-0), which corresponds to j = 1. From the definition of the jerk parameter [\(2\)](#page-2-1), we see that j = 1 implies $\ddot{H} = -3H\dot{H}$. Substituting this, $Eq.(105)$ $Eq.(105)$ simplifies to

$$
\gamma \left(\dot{\gamma} + 4H^2 - 4\dot{H}\right) + 2\gamma^2 H + 4\dot{\gamma}H = 0.
$$
\n(106)

Surprisingly, explicit dependence on redshift and the matter density is washed away once we fix the background evolution to be ΛCDM-like. Also, for a ΛCDM-like evolution, we can write

$$
\dot{H} = -\frac{3}{2}c_1H_0^2(1+z)^3 = -\frac{3}{2}[H^2 - (1-c_1)H_0^2],\tag{107}
$$

so that Eq. (106) becomes

$$
\dot{\gamma} = -\frac{2H\gamma^2 + (10H^2 - 6(1 - c_1)H_0^2)\gamma}{\gamma + 4H}.
$$
\n(108)

Substituting Eq. (108) into Eq. $(33a)$, we get

$$
Q = \frac{3\left(6\gamma(1 - c_1)H_0^2 - 8H^3 + \gamma^2 H\right)}{\gamma + 4H}.
$$
\n(109)

Eq.[\(108\)](#page-18-2), or equivalently Eq.[\(106\)](#page-18-1) can be solved, at least numerically, for $\gamma(z)$ given the ΛCDM-like $H(z)$ from [\(11\)](#page-3-3). Using the solution $\gamma(z)$ one can compute numerically $Q(z)$ and $f_O(z)$ from Eqs. [\(109\)](#page-18-3) and [\(104\)](#page-17-4). One can then, in principle, invert $Q = Q(z)$ to obtain $z = z(Q)$ and substitute it back to the expression of $f_Q(z)$ to obtain $f_Q(Q) = f_Q(z(Q))$. Integrating that will give the Λ CDM-mimicking $f(Q)$ for Γ_3 .

A. The system of equations for numerical reconstruction and the initial conditions

For the purpose of numerical reconstruction, in proves to be useful to write the equations in terms of the following dimensionless quantities

$$
x \equiv \frac{\gamma}{H}, \qquad \tilde{Q} \equiv \frac{Q}{H^2}, \qquad c \equiv \frac{C}{H_0^3},^{16} \qquad N \equiv \ln a \,. \tag{110}
$$

In terms of the above variables, Eqs. (104) , (106) and (109) can be written as

$$
f_Q = \frac{c(x+2)(1+z)^5 + 3hx^2(1+z)^3\Omega_{m0}}{3hx^2(c_1+h^2-1)},
$$
\n(111)

$$
\frac{dx}{dN} = \frac{x[(-1+q)x-4]}{x+4},
$$
\n(112)

$$
\tilde{Q} = \frac{3\left[(2-4q)x + x^2 - 8 \right]}{x+4}.
$$
\n(113)

Next, we need to specify the initial conditions, which we set by considering the reasonable assumption that the physics of gravity do not deviate much from GR during the matter dominated epoch, which occurs at high redshift; any modification of gravity becomes apparent at late time to drive cosmic acceleration. Then one can write

$$
\Omega_{m0}(1+z)^3 \simeq h^2(z) \simeq c_1(1+z)^3. \tag{114}
$$

In the above, the first equality comes from the fact that during the matter dominated epoch the physics of gravity is approximately GR so that the approximate equality $3H^2 \approx \rho_m$ holds. The second equality comes from the fact

¹⁵ Again, if there is a scope of confusion, we will specify the dynamical quantities corresponding to the ΛCDM-like evolution with a subscript ΛCDM.

¹⁶ Readers are advised to take note of the difference in the definition of the dimensionless parameter c between the cases Γ_2 and Γ_3 .

that it occurs at high redshift. This gives rise to the identification $\Omega_{m0} = c_1$. The cosmic evolution during a matter dominated epoch in GR is characterized by the cosmographic condition $q = 1/2$, or equivalently $H = -3H^2/2$. Let us assume that the reconstructed $f(Q)$ exactly coincides with STEGR at a redshift $z = z_* > 0$ during this epoch, so that $f_Q(z_*) = 1$. Setting $f_Q = 1$, Eq.[\(111\)](#page-18-5) gives

$$
x \simeq -2 \,, \quad \text{at} \quad z \simeq z_* \,. \tag{115}
$$

Inserting $x = -2$ into Eq.[\(113\)](#page-18-5), we get

$$
\tilde{Q} \simeq -6 \,, \quad \text{at} \quad z \simeq z_* \,. \tag{116}
$$

Eqs. [\(115\)](#page-19-0) and [\(116\)](#page-19-1) will be used to define the initial conditions during matter dominated epoch in the following analysis.

B. Identifying the free parameters in the reconstructed $f(Q)$ (Γ₃)

Up to this point, we already have two model parameters Ω_{m0} and C in our analysis. Since we have to resort to numerical integration ultimately, we have to set the initial conditions at some redshift value z∗, which acts as another free model parameter^{[17](#page-19-2)}. Since our approach involves numerically finding $Q(z)$ and $f_Q(z)$, inverting $Q(z)$ to find $z(Q)$ and substituting it back to the expression of $f_Q(z)$ to obtain $f_Q(Q)$, we need to ultimately perform an integration to obtain $f(Q)$. This integration will produce the cosmological constant term -2Λ , which is another model parameter. So, there appears to be a total of four model parameters $\Lambda, c, z_*, \Omega_{m0}$. However, not all of them are independent. One can combine Eqs.[\(33b\)](#page-7-6) and [\(33c\)](#page-7-6) such that γ is eliminated, yielding

$$
(1+z)^{-5}\gamma^2 \left[f_Q \left(6H^2 + 6\dot{H} - Q \right) + f + \rho_m \right] + 6CH = 0. \tag{117}
$$

In terms of the variable $x \equiv \frac{\gamma}{H}$, the above equation can be written as

$$
f = -\frac{6c(1+z)^5}{hx^2} + f_Qh^2(6q+\tilde{Q}) - 3(1+z)^3\Omega_{m0}.
$$
\n(118)

Eq.[\(118\)](#page-19-3), evaluated at $z = 0$, puts a constraint between the model parameters Λ , c , z_*, Ω_{m0} , which can be used to determine Ω_{m0} in terms of of the other three. Therefore, ultimately, the reconstructed $f(Q)$ for the case of Γ₃ is in general parametrized by three model parameters just like in the case for Γ_2 . The free model parameters are Λ, c, z_* .

C. Approximate analytic reconstruction during the matter dominated epoch

It is difficult to analytically solve the system of equations (104) , (112) , (113) and obtain an analytic function $f(Q)$ valid for all z , precisely because of which we will go for a numerical reconstruction. However, before moving on to that, let us consider the solutions around the vicinity of the redshift $z = z_*$, while allowing for some deviations from GR (i.e. c_1 is not necessarily equal to Ω_{m0}). It happens that we can reconstruct an approximate analytical $f(Q)$ during this epoch. The redshift $z = z_*$ is typically considered in a high enough redshift epoch so that one can safely approximate

$$
h^2 \approx c_1 (1+z)^3 \implies q = \frac{1}{2} \,. \tag{119}
$$

Setting $q = 1/2$, Eq. [\(112\)](#page-18-5) becomes

$$
\frac{dx}{dN} = -\frac{x(x+8)}{2x+8} \,. \tag{120}
$$

This differential equation can be solved analytically, and the solutions are

$$
x(a) = -4 \pm \sqrt{16 + \frac{c_3}{a}}.
$$
\n(121)

¹⁷ The situation is parallel to the case of Γ_2 , where we have mentioned that z_* can also be used instead of D as the independent model parameter.

21

The constant of integration c_3 can be determined by setting $x(z_*) = -2$ according to Eq.[\(115\)](#page-19-0). Hence, we get

$$
x(z) = -4 + 2\sqrt{4 - 3\frac{1+z}{1+z_*}}\,,\tag{122}
$$

where the plus sign in Eq.[\(121\)](#page-19-4) is chosen to match with Eq.[\(115\)](#page-19-0). It can be seen that this solution becomes imaginary for $z > \frac{1}{3}(1+4z_*)$. At $z = \frac{1}{3}(1+4z_*)$, we have $x = -4$ and consequently $dx/dN \to \infty$ according to Eq.[\(120\)](#page-19-5). For $q = 1/2$, Eq. [\(113\)](#page-18-5) becomes

$$
\tilde{Q} = \frac{3\left(x^2 - 8\right)}{x + 4},\tag{123}
$$

which shows that $Q \to \infty$ at $x = -4$. This blowing up of dx/dN and Q is confirmed in numerical integration later on. Hence, the evolution of the system has a singularity during matter dominated epoch at the redshift $z = \frac{1}{3}(1 + 4z_*)$. Hence, an inherent assumption in our subsequent reconstruction attempt is that the matter dominated epoch starts after the redshift value $z < \frac{1}{3}(1 + 4z_*)^{18}$ $z < \frac{1}{3}(1 + 4z_*)^{18}$ $z < \frac{1}{3}(1 + 4z_*)^{18}$.

According to Eq.[\(111\)](#page-18-5), it is possible for f_Q to vanish for some value of x and some values of the parameters c and Ω_{m0} . To find a range of parameter such that f_Q never vanish within the interval $x > -4$, first we compute the value of x that makes f_Q vanish:

$$
x_0 = -\frac{4}{y} \left(1 \pm \sqrt{1 - y} \right) , \qquad y \equiv \frac{24c_1 h}{c(1 + z)^2} \simeq \frac{24c_1^{3/2}}{c\sqrt{1 + z}} . \tag{124}
$$

The last equality in the definition of y comes from Eq.[\(119\)](#page-19-6). To ensure that f_Q never vanish for real x, one must demand $y > 1$, which places the following bounds on c

$$
c < \frac{24c_1^{3/2}}{\sqrt{1+z}} \ \forall z \in \left(0, \frac{1}{3}(1+4z_*)\right) \quad \Rightarrow c < \frac{12\sqrt{3}c_1^{3/2}}{\sqrt{1+z_*}} \,. \tag{125}
$$

To avoid a singularity of x, we analyze the solution near $z = z_*$. Expanding Eq.[\(122\)](#page-20-1) around $z = z_*$, we get

$$
x \simeq 1 - \frac{3(1+z)}{1+z_*} \,. \tag{126}
$$

Inserting Eq.[\(122\)](#page-20-1) into Eq.[\(123\)](#page-20-2) and expanding the resulting expression around $z = z_*$, we get

$$
Q \simeq 3H_*^2 \left(1 - \frac{3(1+z)}{1+z_*}\right),\tag{127}
$$

where $H_* = H(z_*)$. The expression for f_Q can be obtained by substituting Eq.[\(122\)](#page-20-1) into Eq.[\(111\)](#page-18-5) yielding f_Q around $z = z_*$ as

$$
f_Q \simeq 1 - \frac{C\left(z - z_*\right)\left(1 + z_*\right)^4}{4\left(H_*\right)^3},\tag{128}
$$

Using Eqs. (128) and (127) , we can write

$$
f_Q \simeq 1 + \frac{C\left(1+z_*\right)^5 \left(6\left(H_*\right)^2 + Q\right)}{36\left(H_*\right)^5},\tag{129}
$$

which can be integrated as

$$
f(Q) \simeq -2\Lambda + Q + \frac{CQ\left(1+z_*\right)^5 \left(12\left(H_*\right)^2 + Q\right)}{72\left(H_*\right)^5},\tag{130}
$$

¹⁸ In fact, this singularity can possibly be avoided in a more realistic scenario involving radiation, provided the transition from radiation to matter domination occurs at a redshift $z < \frac{1}{3}(1 + 4z_*)$.

where Λ is the constant of integration. Inserting the above equations for x, Q and f_Q into Eq.[\(117\)](#page-19-7) and evaluating at $z = z_*$, we get

$$
\left(C - \frac{2\Lambda H_*}{(1+z_*)^5}\right) \left[2(1+z_*) + \ln(1+z_*)\right] \simeq 0\,,\tag{131}
$$

which can be satisfied when

$$
C = \frac{2\Lambda H_*}{\left(1 + z_*\right)^5}, \quad \Rightarrow \quad \lambda \equiv \frac{\Lambda}{H_0^2} = \frac{12h_*}{y_*},\tag{132}
$$

where $y_* = y(z_*)$. This relation suggests that Λ relates to Ω_{m0} through C and z_* .

For the case of Γ_2 , we had an integration constant D arising in the expression of f_Q , which we could later relate to z_* . Also for the case of Γ_3 we can have a similar integration constant D. To show this, we insert x from Eq.[\(126\)](#page-20-5) into $Eq.(101)$ $Eq.(101)$, and perform an integration:

$$
f_Q = -\frac{C\left(1+z_*\right)^3 \left(7\left(1+z_*\right)^2 + 2(2(1+z)-9)\left(1+z_*\right) + 7\right)}{16\left(H_*\right)^3} + D\,,\tag{133}
$$

where D is the integration constant. Demanding that $f_Q = 1$ at $z = z_*$, we get

$$
D = 1 + \frac{C\left(11\left(1+z_*\right)^2 - 18\left(1+z_*\right) + 7\right)\left(1+z_*\right)^3}{16H_*^3} \,. \tag{134}
$$

For the case of Γ_2 , where we could carry out the analytic reconstruction for the ΛCDM-mimicking $f(Q)$, we could express γ and Q in terms of H (Eqs.[\(72\)](#page-12-5) and [\(74\)](#page-12-7)). We can do the same here for the approximate reconstructed $f(Q)$ for Γ_3 in the vicinity of the matter-dominated epoch. For the sake of completeness we give the respective expressions below. Since the matter dominated epoch occurs at a high redshift, $h^2 \approx c_1(1+z)^3$ holds. From [\(122\)](#page-20-1) and [\(123\)](#page-20-2), straightforward substitution yields

$$
\frac{\gamma}{H_0} = -4h \left(1 - \sqrt{1 - \frac{3}{4} \left(\frac{h}{h_*} \right)^{2/3}} \right) , \qquad \frac{Q}{H_0^2} = 2h^2 \left[\left(1 - \frac{3}{4} \left(\frac{h}{h_*} \right)^{2/3} \right)^{-1/2} + 2 \left(1 - \frac{3}{4} \left(\frac{h}{h_*} \right)^{2/3} \right)^{1/2} - 4 \right].
$$
\n(135)

D. Numerical reconstruction

Let us now do the full numerical reconstruction, setting $\Omega_{m0} = c_1 = 0.3$. We choose two values of z_* for comparison: $z_* = 20$ and $z_* = 100$. For the purpose of the numerical integration z is limited within the range $(0, \frac{1}{3}(1 + 4z_*))$ to avoid the singularity at $z = \frac{1}{3}(1 + 4z_*)$. The initial conditions are set according to Eqs.[\(115\)](#page-19-0) and [\(116\)](#page-19-1). The plots of $x = \frac{\gamma}{H}$ and $\tilde{Q} = \frac{Q}{H^2}$ vs z are shown in Figs[.3\(a\)](#page-22-0) and [3\(b\).](#page-22-1) It follows from the plots that \tilde{Q} grows rapidly and changes sign when $x \rightarrow -4$.

FIG. 3: Plots (a) and (b) shows the evolution of $x \equiv \frac{\gamma}{H}$ and $\tilde{Q} = \frac{Q}{H^2}$ as functions of z as obtained by solving the equations [\(112\)](#page-18-5) and [\(113\)](#page-18-5) for the parameter value $c = 0.1$. The blue and the orange plots correspond to $z_* = 20$ and $z_* = 100$. The initial conditions are chosen according to Eqs.[\(115\)](#page-19-0) and [\(116\)](#page-19-1).

To show that f_Q can vanish during cosmic evolution leading to a divergence of κ_{eff} if c is not properly chosen, we numerically compute f_Q for two choices of the parameter c: $c = 0.1$ and $c = 1$. For our particular choice of the parameter values $\Omega_{m0} = c_1 = 0.3$ and for both the cases $z_* = 20$ and $z_* = 100$, it can be checked that $c = 0.1$ satisfies the bound in Eq.[\(125\)](#page-20-6), while $c = 1$ doesn't. Accordingly, the plots in Fig[.4\(a\)](#page-22-2) show that f_Q can change sign from positive to be negative when $z > z_*$ for the case of $c = 1$. We show the plots for κ_{eff} from z_* to the present epoch in Fig[.4\(b\).](#page-22-3)

FIG. 4: The plots for the f_Q and $\kappa_{\rm eff} = \frac{1}{f_Q}$ as obtained from Eq.[\(111\)](#page-18-5). To show that the model parameter c should be properly chosen, we have considered two values of c, c = 0.1 and c = 1 for each value of z_* . The blue and the green plots represent the cases $z_* = 20$. The orange and the red plots represent the cases $z_* = 100$. The blue and the orange plots represent the cases $c = 0.1$. The green and the red lines represent the cases $c = 1$.
For the latter case we can see that f_Q changes sign at some high r

Finally, the plots for f as functions of the redshift and Q are shown in Figs[.5](#page-23-1) and [6,](#page-23-2) respectively. For these plots, we just consider the value $c = 0.1$, which ensures the positivity of the effective gravitational coupling within the redshift range considered.

FIG. 5: The plots of f as a function of the redshift z for $c = 0.1$ within the range $z \in (0, z_*)$. The left panel shows the case of $z_* = 20$, while the right panel shows the case of $z_* = 100$.

FIG. 6: The plots of f as a function of Q for $c = 0.1$. The left panel shows the case of $z_* = 20$, while the right panel shows the case of $z_* = 100$.

IX. SUMMARY AND DISCUSSIONS

It is well known that there are three symmetric teleparallel connection branches that respect homogeneity and isotropy [\[33\]](#page-29-15). Most cosmological studies in $f(Q)$ gravity have focused on the trivial connection branch Γ_1 , where Cartesian coordinates serve as the coincidence gauge. Only in recent years have alternative connection branches garnered significant attention. In this work, we explore the possibility of reproducing an exact ΛCDM-like background evolution [\(11\)](#page-3-3) within the framework of $f(Q)$ gravity, considering each of the three connection branches individually. While this question has been previously addressed for the trivial branch Γ_1 in [\[32\]](#page-29-14)—with which our results are consistent—we extend the analysis to the two nontrivial branches. Furthermore, we carefully identify the free parameters of the reconstructed $f(Q)$ for each case and examine the possible constraints among these parameters. The key takeaways from our reconstruction analysis are as follows:

• For the trivial connection Γ_1 , the general $f(Q)$ theory that matches Λ CDM evolution is given by the following 2-parameter family [\(43\)](#page-8-1):

$$
f(Q) = -2\Lambda + \left(\frac{\Lambda/H_0^2}{1 - 2q_0}\right)Q + \beta\sqrt{-Q},
$$
\n(136)

with β and the cosmological constant term Λ acting as the two free parameters. The physical viability condition, $f_Q > 0$, imposes the following bounds on the free parameters:

$$
0 < \frac{\Lambda}{H_0^2} < \frac{3}{2} \left(\frac{1 - 2q_0}{1 + q_0} \right) \,, \qquad \frac{\beta}{H_0} < 2\sqrt{6} \, \frac{\Lambda / H_0^2}{1 - 2q_0} \, h(z) \, \forall \, z \,. \tag{137}
$$

Demanding that STEGR acts as a cosmological past attractor simplifies the above to a 1-parameter family :

$$
f(Q) = -2H_0^2(1 - 2q_0) + Q + \beta\sqrt{-Q}, \qquad \frac{\beta}{H_0} < 2\sqrt{6}h(z) \ \forall z.
$$
 (138)

For the connection Γ_2 , the general $f(Q)$ theory matching Λ CDM evolution is given by the following 3-parameter family [\(79\)](#page-13-5):

$$
f(Q) = -2\Lambda + \left[D + \frac{1}{3} \left(\frac{1 - 2q_0}{1 + q_0} \right) \left(\frac{2c^2}{3\Omega_{m0} - 2(1 + q_0)D} \right) \right] Q - \left[\frac{c^2}{6(1 + q_0)(3\Omega_{m0} - 2(1 + q_0)D)H_0^2} \right] Q^2,
$$
(139)

with c, D and the cosmological constant term Λ acting as the three free parameters, and Ω_{m0} given in terms of the free parameters as [\(78\)](#page-13-2). One can trade D for z_* as a free parameter ([\(80\)](#page-13-4)), z_* being the redshift at which the reconstructed $f(Q)$ coincides with STEGR. The physical viability requirement $f_Q > 0$ for all z and the physically desirable requirement that the reconstructed $f(Q)$ coincides with STEGR at some point in the past (i.e. $z_* > 0$) is ensured by the following bounds on the free parameters :

$$
-\frac{3c_1}{2h(z_*)} < c < 0, \qquad 0 < D < 1 - \frac{2|c|}{3c_1} < 1. \tag{140}
$$

With suitable parameter choices [\(83\)](#page-14-2) it is possible to obtain a simpler 1-parameter family [\(85\)](#page-14-4):

$$
f(Q) = -2H_0^2(1 - 2q_0) + Q - \frac{1}{4H_0^2} \left(\frac{1 - D}{1 - 2q_0}\right) Q^2.
$$
 (141)

However, this simpler family has the flaw that STEGR acts not as a cosmological past attractor, but rather as a cosmological future attractor $(f_Q(z \to -1) = 1)$, as shown in section VIIC. Such a theory will significantly deviate from STEGR during the matter dominated epoch in the past, leading to a signature in the structure formation scenario very different than observed.

• For the connection Γ_3 , we were able to decouple the evolution equation for the dynamical connection for a generic cosmic history (Eq.[\(105\)](#page-17-5)) as well as for the ΛCDM-like evolution in particular (Eq.[\(108\)](#page-18-2)). This enabled us to perform a subsequent numerical reconstruction. The result of the numerical reconstruction is shown in Fig[.6.](#page-23-2) The reconstructed $f(Q)$ is a 3-parameter family in general, as discussed in section [VIII B,](#page-19-8) parametrized by c, z_* and the cosmological constant term Λ . An approximate analytic form of the reconstructed $f(Q)$ during the matter dominated epoch is given by (130) :

$$
f(Q) \simeq -2\Lambda + Q + \frac{cQ\left(1+z_*\right)^5 \left(12h_*^2 + Q/H_0^2\right)}{72h_*^5}, \qquad h_* = h(z_*)\,. \tag{142}
$$

By appropriately choosing the free parameters, it is possible to ensure the positivity of f_Q within the range $z \in (0, z_*)$ (see Fig[.4\(a\)\)](#page-22-2).

To alleviate the cosmological constant problem [\[3,](#page-28-2) [71\]](#page-30-20), lately there have been proposals where the cosmological constant term in the gravitational Lagrangian shows up as an integration constant, decoupled from the microphysics of vacuum energy [\[72,](#page-30-21) [73\]](#page-30-22). Notably, the cosmological constant term also appears as an integration constant during the reconstruction of Λ CDM-mimicking $f(Q)$ gravity for all the three possible connection branches. In fact, one encounters a similar situation while reconstructing the Λ CDM-mimicking $f(R)$ gravity [\[28\]](#page-29-10).

In the context of modified gravity theories, a very pertinent question is whether one can achieve an exactly ΛCDM-like evolution with a vanishing cosmological constant term. If feasible, then one can attribute the observed late time cosmic evolution solely to additional *dynamical* degrees of freedom coming the theory, and the cosmological constant issue simply never arises. Our results are as follows:

• For the trivial connection Γ_1 , this is not possible, as Eq. [\(42\)](#page-8-2) indicates that $\Lambda = 0$ implies $\Omega_{m0} = 0$.

• For the connection Γ_2 , it is possible to set $\Lambda = 0$, provided the remaining model parameters c and D satisfy the following bounds:

$$
|c| < \frac{3}{2} \left(\frac{c_1}{1 + \sqrt{1 - c_1}} \right), \qquad \frac{2|c|}{3c_1} \sqrt{1 - c_1} \le D < \frac{2|c|}{3c_1} \sqrt{1 - c_1} \sqrt{1 + \frac{3}{2c^2 (1 - c_1)}}. \tag{143}
$$

Here, the first bound is obtained by combining the inequalities [\(92\)](#page-15-6) and [\(94\)](#page-15-9) setting $\Lambda = 0$. The second bound is obtained by combining the inequalities [\(94\)](#page-15-9) and the requirement $0 < \Omega_{m0} < 1$, using the expression of Ω_{m0} from [\(78\)](#page-13-2) and setting $\Lambda = 0$.

• For the connection Γ_3 , we were unable to derive an analytic form for the reconstructed $f(Q)$ that reproduces the full Λ CDM-like cosmic history. However, for the approximate reconstructed $f(Q)$ [\(130\)](#page-20-7), valid during the matter-dominated epoch, we see that setting $\Lambda = 0$ results in $C = 0$ (Eq. [\(131\)](#page-21-0)), which reduces the reconstructed $f(Q)$ [\(130\)](#page-20-7) to STEGR, $f(Q) \simeq Q$.

Conclusively, we can say that only the nontrivial connection branch Γ_2 allows us to achieve an exactly Λ CDM-like cosmic evolution without actually a cosmological constant term, thereby addressing the cosmological constant problem. It is worth noting that we have specified the ΛCDM-like cosmic evolution as (Eq.[\(7\)](#page-3-0))

$$
h_{\Lambda \text{CDM}}(z) = \sqrt{[c_1(1+z)^3 + (1-c_1)]}, \qquad c_1 = \frac{2}{3}(1+q_0), \qquad (144)
$$

instead of

$$
h_{\Lambda \text{CDM}}(z) = \sqrt{\left[\Omega_{m0}(1+z)^3 + (1-\Omega_{m0})\right]}.
$$
\n(145)

While the identification $\Omega_{m0} = c_1 = \frac{2}{3}(1+q_0)$ holds for the actual General Relativistic ΛCDM model, it does not generally apply to alternative late-time models. In particular, for our reconstructed $f(Q)$ models, Ω_{m0} is expressed in terms of the other free model parameters and the present-day cosmographic parameters as follows:

• For Γ_1 (see [\(42\)](#page-8-2)):

$$
\Omega_{m0} = \frac{2\Lambda}{3H_0^2} \left(\frac{1+q_0}{1-2q_0}\right) \,. \tag{146}
$$

• For Γ_2 (see [\(78\)](#page-13-2)):

$$
\Omega_{m0} = \frac{2}{3} \lambda \left(\frac{1+q_0}{1-2q_0} \right) \pm \frac{2}{3} \sqrt{(1+q_0)^2 \left(\frac{\lambda}{1-2q_0} - D \right)^2 - \frac{c^2}{3} (1-2q_0)}.
$$
\n(147)

• For Γ_3 , Ω_{m0} is determined by solving the following equation(see [\(118\)](#page-19-3)):

$$
f_0 = -\frac{6c}{x_0^2} + f_{Q0}(6q_0 + \tilde{Q}_0) - 3\Omega_{m0},
$$
\n(148)

the subscript '0' denoting present day values.

However, the converse is not always true. For example:

- For the case of Γ_1 , demanding that STEGR acts as a cosmological past attractor for the reconstructed $f(Q)$ ultimately leads us to the condition $\Omega_{m0} = c_1$, even though the theory starts deviating from STEGR at lower redshifts; see section [VI A.](#page-8-5)
- For the case of Γ_2 , employing the parameter choice of Eq.[\(83\)](#page-14-2), we arrive at the 1-parameter reconstructed $f(Q)$ [\(85\)](#page-14-4), which also produces $\Omega_{m0} = c_1$, while the gravity theory is still different from STEGR.

By combining CC+BAO+Pantheon+CMB data, and employing a Gaussian process, one can obtain a modelindependent estimate for q_0 typically as [\[43,](#page-29-27) Table 5] $q_0 = -0.647 \pm 0.069$. Similarly, combining CC+Type 1a Supernovae+BAO+Redshift Space Distorsion data and employing a Gaussian process one can obtain a modelindependent estimate for Ω_{m0} typically as 0.224 ± 0.066 [\[74\]](#page-30-23). The quantity $|c_1 - \Omega_{m0}|/\Omega_{m0} \approx 5\%$ gives an estimate of the difference from the General Relativistic ΛCDM model according to a combination of different datasets. The

difference between c_1 and Ω_{m0} holds the potential of possible amelioration of the Hubble tension [\[59\]](#page-30-7). Our analysis reveals that the connection branch Γ_2 exhibits greater phenomenological potential than Γ_1 , as it can maintain coincidence with STEGR at high redshifts without requiring $\Omega_{m0} = c_1$.

Our work paves the way for reconstructing $f(Q)$ theories from cosmological data for all three possible branches of symmetric teleparallel homogeneous and isotropic connections. By assuming no coupling between the matter Lagrangian and the connection, we simplify the equations by setting the hypermomentum to zero. However, in general, there is no inherent physical motivation to neglect hypermomentum within the teleparallel framework. We hope this work serves as a foundation for future efforts to reconstruct $f(Q)$ models, either analytically or numerically, in scenarios that include hypermomentum.

Our work paves the way for reconstructing $f(Q)$ theories starting from a cosmology for all the three possible branches of the symmetric teleparallel homogeneous and isotropic connections. We have not assumed any coupling between the matter Lagrangian and the connection, leading to a vanishing hypermomentum, considerably simplifying the equations. However, in general, though, there is no inherent physical motivation to ignore hypermomentum in a teleparallel framework. We hope this work serves as a foundation for future efforts to reconstruct $f(Q)$ models, either analytically or numerically, in scenarios that include hypermomentum.

Finally, we emphasize that proving the existence of ΛCDM-mimicking modified gravity theories is not sufficient. It is crucial to explore ways to distinguish ΛCDM-mimicking models from the actual ΛCDM model. While the two scenarios may yield indistinguishable background evolution, differences will arise in the perturbation evolution. Recently, a cosmographic approach was used to analyze perturbations in ΛCDM-mimicking f(R) models, identifying distinctive signatures such as dispersion in the growth-index parameter [\[75\]](#page-30-24). A similar analysis should be extended to Λ CDM-mimicking $f(Q)$ models across all connection branches.

Open questions remain, such as whether an analytic reconstruction for Γ_3 is possible and how the inclusion of hypermomentum might change the results. We leave them for future work.

ACKNOWLEDGMENTS

SC acknowledges funding support from the NSRF via the Program Management Unit for Human Resources and Institutional Development, Research and Innovation [grant number B13F670063]. JD acknowledges the support of IUCAA, Pune (India) through the visiting associateship program. DG is a member of the GNFM working group of Italian INDAM, and also acknowledges financial support from the start-up funding plan of Jiangsu University of Science and Technology.

Appendix A: On the stability of the ΛCDM-like cosmic solution $j(z) = 1$ in the reconstructed $f(Q)$ (Γ₂)

In this section, we comment on the stability of the ΛCDM cosmological solution against small homogeneous and isotropic perturbations for the reconstructed $f(Q)$ for the case of Γ_2 . To this goal, we first take a time derivative of the Raychaudhuri equation [\(28b\)](#page-6-4). After quite a few steps of calculations we arrive at the following equation

$$
2(\ddot{H} + 9H\dot{H} + 9H^3)f_Q + 2(2\dot{H} + 3H^2)f_{QQ}\dot{Q} + 3(f - Qf_Q)H - f_{QQ}Q\dot{Q} = 0.
$$
 (A1)

In deriving the above equation we have made use of the equation [\(29\)](#page-6-3). Consider now a given solution $H(t)$ of the above equation, and take a homogeneous and isotropic perturbation around it: $H(t) \to H(t) + \delta H(t)$. We substitute it back in the above equation as well as in Eq.[\(29\)](#page-6-3). Keeping terms of only linear order we get

$$
2f_Q\delta \ddot{H} + (18Hf_Q + 4f_{QQ}\dot{Q})\delta H + [18(1-q)H^2f_Q + 12f_{QQ}H\dot{Q} + 3(f - Qf_Q)]\delta H
$$

= $-f_{QQ}[2(1-2q)H^2 - Q]\delta Q - [2(j-6q+2)H^3f_{QQ} + 2(1-2q)H^2\dot{Q}f_{QQQ} - (\dot{Q} + 3HQ)f_{QQ} - Q\dot{Q}f_{QQQ}]\delta Q,$
(A2)

$$
f_{QQ}\ddot{\delta Q} + (2\dot{Q}f_{QQQ} + 3Hf_{QQ})\dot{\delta Q} + [\dot{Q}^2f_{QQQQ} + (\ddot{Q} + 3H\dot{Q})f_{QQQ}]\delta Q = -3\dot{Q}f_{QQ}\delta H. \tag{A3}
$$

For the particular 1-parameter reconstructed $f(Q)$ [\(85\)](#page-14-4) that reproduces the ΛCDM-like background evolution [\(11\)](#page-3-3), the above equations become

$$
\mathcal{A}(z)\delta\ddot{H} + \mathcal{B}(z)\delta\dot{H} + \mathcal{C}(z)\delta H = \mathcal{D}(z)\delta Q + \mathcal{E}(z)\delta Q, \tag{A4}
$$

$$
\ddot{\delta Q} + 3H\dot{\delta Q} = -3\dot{Q}\delta H\,,\tag{A5}
$$

where

$$
\mathcal{A}(z) = 2\left(D - \frac{2c}{3c_1}h_{\Lambda \text{CDM}}(z)\right),\tag{A6a}
$$

$$
\mathcal{B}(z) = \left[18h_{\Lambda\text{CDM}}(z)D - \frac{4c}{3c_1}(7 - 2q_{\Lambda\text{CDM}}(z))h_{\Lambda\text{CDM}}^2(z)\right]H_0,\tag{A6b}
$$

$$
\mathcal{C}(z) = 18(1 - q_{\Lambda \text{CDM}}(z))H_{\Lambda \text{CDM}}^2(z) \left(1 - \frac{Q_{\Lambda \text{CDM}}(z)}{6(1 - c_1)H_0^2}\right) - \frac{2}{(1 - c_1)H_0^2}H\dot{Q}_{\Lambda \text{CDM}}(z) + 3\left(\Lambda + \frac{1}{12(1 - c_1)H_0^2}Q_{\Lambda \text{CDM}}^2\right),\tag{A6c}
$$

$$
\mathcal{D}(z) = \frac{1}{6(1 - c_1)H_0^2} [2(1 - 2q_{\Lambda \text{CDM}}(z))H_{\Lambda \text{CDM}}^2(z) - Q_{\Lambda \text{CDM}}(z)], \tag{A6d}
$$

$$
\mathcal{E}(z) = \frac{1}{6(1 - c_1)H_0^2} [2(3 - 6q_{\Lambda \text{CDM}}(z))H_{\Lambda \text{CDM}}^3(z) - (\dot{Q}_{\Lambda \text{CDM}}(z) + 3H_{\Lambda \text{CDM}}(z)Q_{\Lambda \text{CDM}}(z))]. \tag{A6e}
$$

In the above, the subscript 'ΛCDM' implies that the respective dynamical quantities are calculated for ΛCDM-like background evolution [\(11\)](#page-3-3), $c_1 = \frac{2}{3}(1+q_0)$ according to Eq.[\(10\)](#page-3-2) and c is given by the equation [\(83\)](#page-14-2).

We will not execute the full analysis here due to it's computational complexity, but we will chalk out the schematics of how one can proceed with it. Taking a time derivative of Eq.[\(37\)](#page-7-3) and utilizing Eq.[\(A5\)](#page-26-1), one can get the following equation

$$
\mathcal{A}\ddot{\delta}\ddot{H} + (\dot{\mathcal{A}} + \mathcal{B})\dot{\delta}\dot{H} + (\dot{\mathcal{B}} + \mathcal{C})\dot{\delta}H + (\mathcal{C} + 3\mathcal{D}\dot{Q})\delta H = (\dot{\mathcal{D}} + \mathcal{E} - 3\mathcal{D}H)\dot{\delta}Q + \dot{\mathcal{E}}\delta Q. \tag{A7}
$$

Eqs. (37) and $(A7)$ can be solved to obtain

$$
\delta Q = \delta Q(\delta H, \delta H, \delta \ddot{H}, \delta \dddot{H}), \qquad \dot{\delta Q} = \dot{\delta Q}(\delta H, \delta H, \delta \dddot{H}, \delta \dddot{H}). \tag{A8}
$$

In particular, one can take another derivative of the expression for δQ to obtain

$$
\ddot{\delta Q} = \ddot{\delta Q} (\delta H, \delta H, \delta H, \dot{\delta H}, \ddot{\delta H}). \tag{A9}
$$

One can then substitute the expressions for δQ and δQ thus obtained to Eq.[\(A5\)](#page-26-1) to obtain a 4-th order homogeneous ordinary differential equation for δH

$$
(\ldots)\ddot{\delta}\dddot{H} + (\ldots)\ddot{\delta}\dddot{H} + (\ldots)\delta\dot{H} + (\ldots)\delta H = 0, \qquad (A10)
$$

with the coefficients depending on background related dynamical quantities. The signs of the roots of the characteristic polynomial of the above differential equation gives the stability of the ΛCDM cosmological solution with respect to small homogeneous isotropic perturbations within the reconstructed $f(Q)$ [\(85\)](#page-14-4). In particular, for the solution to be stable, all the roots must be negative.

Appendix B: Some comments regarding the dynamical connection function (Γ_2)

For any generic cosmic history $H(z)$, in the context of the connection Γ_2 , the inverse of the effective gravitational coupling relates to the redshift via the Hubble rate as [\(66\)](#page-12-8),

$$
f_Q(z) = -cJ(z) + D\,,\tag{B1}
$$

where we have defined

$$
\mathcal{I}(z) = \int \frac{(1+z)^2 dz}{h(z)}.
$$
 (B2)

Qualitatively, we can conclude from [\(66\)](#page-12-8) that $f_Q(z)$ is a monotonic function of z for any cosmic history of the universe (whether monotonically increasing or decreasing depends on the sign of the parameter c). The dynamical connection function for any generic cosmic history $H(z)$ can be expressed as [\[38,](#page-29-20) Eq.(29)]

$$
\gamma(z) = \frac{2h(z)H'(z)\left[c\int \frac{(1+z)^2dz}{h(z)} - D\right]}{3c(1+z)^2} + \frac{2H(z)}{3} + \frac{\Omega_{m0}H_0}{c} = -\frac{2h(z)H'(z)f_Q(z)}{3c(1+z)^2} + \frac{2H(z)}{3} + \frac{\Omega_{m0}H_0}{c},\tag{B3}
$$

or

$$
\frac{\gamma}{H_0} = \frac{2c_1(1+q)h^2(z)[c\Im(z)-D]}{3c(h_{\Lambda \text{CDM}}^2(z)-1+c_1)} + \frac{2h(z)}{3} + \frac{\Omega_{m0}}{c} = -\frac{2c_1(1+q(z))h^2(z)f_Q(z)}{3c(h_{\Lambda \text{CDM}}^2(z)-1+c_1)} + \frac{2h(z)}{3} + \frac{\Omega_{m0}}{c}.
$$
 (B4)

For the Hubble rate corresponding to a generic constant jerk parameter throughout the entire comic history (Eq.[\(4\)](#page-2-2)), the relevant integral $\mathfrak{I}(z)$ cannot be computed in a closed-form. We can nevertheless get a taste of the robustness of the reconstructed Λ CDM-mimicking $f(Q)$ within the range of astrophysical uncertainty for the observed jerk function by noticing that

$$
H(z) = H_{\Lambda \text{CDM}}(z) + \frac{H_0}{3} \left(h_{\Lambda \text{CDM}}(z) + \frac{2(c_1 - 1)}{h_{\Lambda \text{CDM}}(z)} \right) \ln(1 + z) \varepsilon + O(\varepsilon^2) ,\tag{B5}
$$

and consequently

$$
\mathcal{I}(z) := \int \frac{(1+z)^2 dz}{h(z)} = \int \frac{(1+z)^2 dz}{h_{\Lambda\text{CDM}}(z)} - \frac{\varepsilon}{3} \int (1+z)^2 \left(h_{\Lambda\text{CDM}}(z) + \frac{2(c_1-1)}{h_{\Lambda\text{CDM}}(z)} \right) \ln(1+z) dz + O(\varepsilon^2),
$$

\n
$$
= \frac{2h_{\Lambda\text{CDM}}(z)}{3c_1} + \frac{2\varepsilon}{243c_1} \left[30(1-c_1)^{3/2} \arctan\left(\frac{h_{\Lambda\text{CDM}}(z)}{\sqrt{1-c_1}}\right) - \left(2[15(1-c_1) - h_{\Lambda\text{CDM}}^2(z)] + 9[6(1-c_1) - h_{\Lambda\text{CDM}}^2(z)] \ln\left(\frac{h_{\Lambda\text{CDM}}^2(z) - 1 + c_1}{c_1}\right)^{1/3} \right) h_{\Lambda\text{CDM}}(z) + O(\varepsilon^2).
$$
\n(B6)

The deceleration parameter can be calculate similarly from Eq.[\(5\)](#page-2-4).

The case of the dynamical connection is therefore richer than that of the effective coupling, potentially providing a route for discriminating between different non-metrical models. In fact, its increasing/decreasing properties appear to be sensitive in a non-trivial manner to the algebraic sign of ε because the former addendum is the product of the three functions $-2/(1+z)^3$ (negative and monotonically increasing), $f_Q(z)/c$ (negative and monotonically decreasing, at least in the limit $D \to 0$), and $(1+q(z))h^2(z) \equiv c_1k_1(1+z)^{k_1} + (1-c_1)k_2(1+z)^{k_2}$ (monotonically increasing with its algebraic sign depending on the value of k_2), and also the properties of the second addendum depend on the sign of the parameter k_2 . Our result $(B4)$ constitutes the starting point for discriminating between different underlying models since the reconstructed dynamical functions from astrophysical data indeed exhibit different qualitative behaviors [\[38\]](#page-29-20). Furthermore, the dynamical connection in turns determines the Riemann curvature entering the equation of geodesic deviations [\[76\]](#page-30-25), potentially bearing consequences on the propagation of gravitational wave signals [\[77\]](#page-30-26): identifying the signature of the deperature from a ΛCDM-mimiking model, and specifically towards larger or smaller values of the jerk function, on their spectra is left for future investigations.

[2] S. Perlmutter et al. Measurements of Ω and Λ from 42 High Redshift Supernovae. Astrophys. J., 517:565–586, 1999.

- [7] Jose Beltrán Jiménez, Lavinia Heisenberg, and Tomi Koivisto. Coincident General Relativity. Phys. Rev. D, 98(4):044048, 2018.
- [8] Christian G. Boehmer and Erik Jensko. Modified gravity: A unified approach to metric-affine models. J. Math. Phys., 64(8):082505, 2023.
- [9] Christian G. Boehmer and Erik Jensko. Modified gravity: A unified approach. Phys. Rev. D, 104(2):024010, 2021.
- [10] Christian G. Boehmer, Erik Jensko, and Ruth Lazkoz. Cosmological dynamical systems in modified gravity. Eur. Phys. J. C, 82(6):500, 2022.
- [11] Débora Aguiar Gomes, Jose Beltrán Jiménez, Alejandro Jiménez Cano, and Tomi S. Koivisto. Pathological Character of Modifications to Coincident General Relativity: Cosmological Strong Coupling and Ghosts in f(Q) Theories. Phys. Rev. Lett., 132(14):141401, 2024.

^[1] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. Astron. J., 116:1009–1038, 1998.

^[3] Steven Weinberg. The Cosmological Constant Problem. Rev. Mod. Phys., 61:1–23, 1989.

^[4] L. Verde, T. Treu, and A. G. Riess. Tensions between the Early and the Late Universe. Nature Astron., 3:891, 2019.

^[5] Jose Beltrán Jiménez, Lavinia Heisenberg, and Tomi S. Koivisto. The Geometrical Trinity of Gravity. Universe, 5(7):173, 2019.

^[6] R. Aldrovandi and J. G. Pereira. Teleparallelism: A New Way to Think the Gravitational Interaction. Ciencia Hoje, 55:32, 2015.

- [12] Kun Hu, Makishi Yamakoshi, Taishi Katsuragawa, Shin'ichi Nojiri, and Taotao Qiu. Nonpropagating ghost in covariant f(Q) gravity. Phys. Rev. D, 108(12):124030, 2023.
- [13] Jose Beltrán Jiménez, Alexey Golovnev, Tomi Koivisto, and Hardi Veermäe. Minkowski space in $f(T)$ gravity. Phys. Rev. D, 103(2):024054, 2021.
- [14] Lavinia Heisenberg, Manuel Hohmann, and Simon Kuhn. Cosmological teleparallel perturbations. JCAP, 03:063, 2024.
- [15] Antonio G. Bello-Morales, Jose Beltrán Jiménez, Alejandro Jiménez Cano, Tomi S. Koivisto, and Antonio L. Maroto. A class of ghost-free theories in symmetric teleparallel geometry. JHEP, 12:146, 2024.
- [16] Luís Atayde and Noemi Frusciante. Can $f(Q)$ gravity challenge Λ CDM? Phys. Rev. D, 104(6):064052, 2021.
- [17] Inês S. Albuquerque and Noemi Frusciante. A designer approach to $f(Q)$ gravity and cosmological implications. Phys. Dark Univ., 35:100980, 2022.
- [18] Shambel Sahlu, Alvaro de la Cruz-Dombriz, and Amare Abebe. Structure growth in $f(Q)$ cosmology. $arXiv:2405.07361$ $\sqrt{qr-qc}$, 5 2024.
- [19] Tiago B. Gonçalves, Luís Atayde, and Noemi Frusciante. Cosmological study of a symmetric teleparallel gravity model. Phys. Rev. D, 109(8):084003, 2024.
- [20] Avik De and Tee-How Loo. On the viability of $f(Q)$ gravity models. Class. Quant. Grav., $40(11):115007$, 2023.
- [21] Ruth Lazkoz, Francisco S. N. Lobo, María Ortiz-Baños, and Vincenzo Salzano. Observational constraints of $f(Q)$ gravity. Phys. Rev. D, 100(10):104027, 2019.
- [22] Ismael Ayuso, Ruth Lazkoz, and Vincenzo Salzano. Observational constraints on cosmological solutions of $f(Q)$ theories. Phys. Rev. D, 103(6):063505, 2021.
- [23] Noemi Frusciante. Signatures of $f(Q)$ -gravity in cosmology. Phys. Rev. D, 103(4):044021, 2021.
- [24] Fotios K. Anagnostopoulos, Viktor Gakis, Emmanuel N. Saridakis, and Spyros Basilakos. New models and big bang nucleosynthesis constraints in $f(Q)$ gravity. Eur. Phys. J. C, 83(1):58, 2023.
- [25] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. Astron. Astrophys., 641:A6, 2020. [Erratum: Astron.Astrophys. 652, C4 (2021)].
- [26] A. G. Adame et al. The Early Data Release of the Dark Energy Spectroscopic Instrument. Astron. J., 168(2):58, 2024.
- [27] Marina Cortês and Andrew R. Liddle. Interpreting DESI's evidence for evolving dark energy. JCAP, 12:007, 2024.
- [28] Peter K. S. Dunsby, Emilo Elizalde, Rituparno Goswami, Sergei Odintsov, and Diego Saez Gomez. On the LCDM Universe in f(R) gravity. Phys. Rev. D, 82:023519, 2010.
- [29] Jian-hua He and Bin Wang. Revisiting $f(R)$ gravity models that reproduce ΛCDM expansion. Phys. Rev. D, 87(2):023508, 2013.
- [30] S. Fay, S. Nesseris, and L. Perivolaropoulos. Can $f(R)$ Modified Gravity Theories Mimic a LCDM Cosmology? Phys. Rev. D, 76:063504, 2007.
- [31] E. Elizalde, R. Myrzakulov, V. V. Obukhov, and D. Saez-Gomez. LambdaCDM epoch reconstruction from F(R,G) and modified Gauss-Bonnet gravities. Class. Quant. Grav., 27:095007, 2010.
- [32] Gaurav N. Gadbail, Sanjay Mandal, and P. K. Sahoo. Reconstruction of ΛCDM universe in f(Q) gravity. Phys. Lett. B, 835:137509, 2022.
- [33] Manuel Hohmann. General covariant symmetric teleparallel cosmology. Phys. Rev. D, 104(12):124077, 2021.
- [34] Andronikos Paliathanasis. The impact of the non-coincidence gauge on the dark energy dynamics in f(Q)-gravity. Gen. Rel. Grav., 55(11):130, 2023.
- [35] Salvatore Capozziello and Rocco D'Agostino. Model-independent reconstruction of f(Q) non-metric gravity. Phys. Lett. B, 832:137229, 2022.
- [36] Fabrizio Esposito, Sante Carloni, Roberto Cianci, and Stefano Vignolo. Reconstructing isotropic and anisotropic f(Q) cosmologies. Phys. Rev. D, 105(8):084061, 2022.
- [37] Shin'ichi Nojiri and S. D. Odintsov. Well-defined f(Q) gravity, reconstruction of FLRW spacetime and unification of inflation with dark energy epoch. *Phys. Dark Univ.*, 45:101538, 2024.
- [38] Yuhang Yang, Xin Ren, Bo Wang, Yi-Fu Cai, and Emmanuel N. Saridakis. Data reconstruction of the dynamical connection function in $f(Q)$ cosmology. Mon. Not. Roy. Astron. Soc., 533(2):2232–2241, 2024.
- [39] E. R. Harrison. Observational tests in cosmology. Nature, 260(5552): 591–592, April 1976.
- [40] Saikat Chakraborty, Daniele Gregoris, and B. Mishra. On the uniqueness of ΛCDM-like evolution for homogeneous and isotropic cosmology in General Relativity. Phys. Lett. B, 842:137962, 2023.
- [41] Varun Sahni, Tarun Deep Saini, Alexei A. Starobinsky, and Ujjaini Alam. Statefinder: A New geometrical diagnostic of dark energy. JETP Lett., 77:201–206, 2003.
- [42] Jose Luis Bernal, Licia Verde, and Adam G. Riess. The trouble with H_0 . JCAP, 10:019, 2016.
- [43] Purba Mukherjee and Narayan Banerjee. Non-parametric reconstruction of the cosmological jerk parameter. Eur. Phys. J. $C, 81(1):36, 2021.$
- [44] Jun-Qian Jiang, Davide Pedrotti, Simony Santos da Costa, and Sunny Vagnozzi. Nonparametric late-time expansion history reconstruction and implications for the Hubble tension in light of recent DESI and type Ia supernovae data. Phys. Rev. D, 110(12):123519, 2024.
- [45] Levon Pogosian, Marco Raveri, Kazuya Koyama, Matteo Martinelli, Alessandra Silvestri, Gong-Bo Zhao, Jian Li, Simone Peirone, and Alex Zucca. Imprints of cosmological tensions in reconstructed gravity. Nature Astron., 6(12):1484–1490, 2022.
- [46] James M. Bardeen. Gauge Invariant Cosmological Perturbations. Phys. Rev. D, 22:1882–1905, 1980.
- [47] Orlando Luongo. Dark energy from a positive jerk parameter. *Mod. Phys. Lett. A*, 28:1350080, 2013.
- [48] Hassan Amirhashchi and Soroush Amirhashchi. Recovering ΛCDM model from a cosmographic study. Gen. Rel. Grav., 52(2):13, 2020.
- [49] María Ortiz-Baños, Mariam Bouhmadi-López, Ruth Lazkoz, and Vincenzo Salzano. ΛCDM suitably embedded in $f(R)$ with a non-minimal coupling to matter. Eur. Phys. J. C, $81(3):237, 2021$.
- [50] Shibendu Gupta Choudhury, Ananda Dasgupta, and Narayan Banerjee. Reconstruction of f(R) gravity models for an accelerated universe using the Raychaudhuri equation. Mon. Not. Roy. Astron. Soc., 485(4):5693–5699, 2019.
- [51] Gaurav N. Gadbail, Avik De, and P. K. Sahoo. Cosmological reconstruction and ΛCDM universe in $f(Q, C)$ gravity. Eur. Phys. J. C, 83(12):1099, 2023.
- [52] Sante Carloni, Rituparno Goswami, and Peter K. S. Dunsby. A new approach to reconstruction methods in f(R) gravity. Class. Quant. Grav., 29:135012, 2012.
- [53] Maciej Dunajski and Gary Gibbons. Cosmic Jerk, Snap and Beyond. Class. Quant. Grav., 25:235012, 2008.
- [54] Yong-Seon Song, Wayne Hu, and Ignacy Sawicki. The Large Scale Structure of f(R) Gravity. Phys. Rev. D, 75:044004, 2007.
- [55] Levon Pogosian and Alessandra Silvestri. The pattern of growth in viable f(R) cosmologies. Phys. Rev. D, 77:023503, 2008. [Erratum: Phys.Rev.D 81, 049901 (2010)].
- Rohin Kumar. A New Class of Cosmologically 'Viable' $f(R)$ Models. $arXiv: 1611.03728$ [qr-qc], 11 2016.
- [57] Saikat Chakraborty, Kelly MacDevette, and Peter Dunsby. A model independent approach to the study of $f(R)$ cosmologies with expansion histories close to Λ CDM. Phys. Rev. D, 103(12):124040, 2021.
- [58] Zhong-Xu Zhai, Ming-Jian Zhang, Zhi-Song Zhang, Xian-Ming Liu, and Tong-Jie Zhang. Reconstruction and constraining of the jerk parameter from OHD and SNe Ia observations. Phys. Lett. B, 727:8–20, 2013.
- [59] Davide Pedrotti, Jun-Qian Jiang, Luis A. Escamilla, Simony Santos da Costa, and Sunny Vagnozzi. Multidimensionality of the Hubble tension: The roles of Ωm and $ωc$. Phys. Rev. D, 111(2):023506, 2025.
- [60] A. R. Sandage. Cosmology: a search for two numbers. Physics Today, 23(2):34–41, January 1970.
- [61] Manuel Hohmann. Metric-affine Geometries With Spherical Symmetry. Symmetry, 12(3):453, 2020.
- [62] Dehao Zhao. Covariant formulation of $f(Q)$ theory. Eur. Phys. J. C, 82(4):303, 2022.
- [63] Laur Järv, Mihkel Rünkla, Margus Saal, and Ott Vilson. Nonmetricity formulation of general relativity and its scalar-tensor extension. Phys. Rev. D, 97(12):124025, 2018.
- $[64]$ Maria-Jose Guzman, Laur Järv, and Laxmipriya Pati. Exploring the stability of $f(Q)$ cosmology near general relativity limit with different connections. Phys. Rev. D, 110(12):124013, 2024.
- [65] Jose Beltrán Jiménez, Lavinia Heisenberg, Tomi Sebastian Koivisto, and Simon Pekar. Cosmology in $f(Q)$ geometry. Phys. Rev. D, 101(10):103507, 2020.
- [66] Christian G. Boehmer and Nyein Chan. Dynamical systems in cosmology. 2017.
- [67] John D. Barrow and A. C. Ottewill. The Stability of General Relativistic Cosmological Theory. J. Phys. A, 16:2757, 1983.
- [68] Alvaro de la Cruz-Dombriz and Diego Saez-Gomez. On the stability of the cosmological solutions in $f(R, G)$ gravity. Class. Quant. Grav., 29:245014, 2012.
- [69] Qingqing Wang, Xin Ren, Yi-Fu Cai, Wentao Luo, and Emmanuel N. Saridakis. Observational Test of f(Q) Gravity with Weak Gravitational Lensing. Astrophys. J., 974(1):7, 2024.
- [70] Shambel Sahlu and Endalkachew Tsegaye. Linear Cosmological perturbations in $f(Q)$ Gravity. arXiv:2206.02517 [gr-qc], 6 2022.
- [71] Y. B. Zeldovich. Cosmological Constant and Elementary Particles. JETP Lett., 6:316, 1967.
- [72] Priidik Gallagher and Tomi Koivisto. The Λ and the CDM as Integration Constants. Symmetry, 13(11):2076, 2021.
- [73] Justin C. Feng and Pisin Chen. Cosmological constant as an integration constant. Eur. Phys. J. C, 84(12):1331, 2024.
- [74] Jaime Ruiz-Zapatero, Carlos García-García, David Alonso, Pedro G. Ferreira, and Richard D. P. Grumitt. Model-independent constraints on Ωm and H(z) from the link between geometry and growth. Mon. Not. Roy. Astron. Soc., 512(2):1967–1984, 2022.
- [75] Kelly MacDevette, Jess Worsley, Peter Dunsby, and Saikat Chakraborty. A model independent approach to the study of structure growth in $f(R)$ gravity. $arXiv:2408.03998$ [gr-qc], 8 2024.
- [76] Jing-Theng Beh, Tee-how Loo, and Avik De. Geodesic deviation equation in f(Q)-gravity. Chin. J. Phys., 77:1551–1560, 2022.
- [77] Salvatore Capozziello, Maurizio Capriolo, and Shin'ichi Nojiri. Gravitational waves in f(Q) non-metric gravity via geodesic deviation. Phys. Lett. B, 850:138510, 2024.