

Remark to a Theorem of van Geemen

Eberhard Freitag

Mathematisches Institut

Im Neuenheimer Feld 288

D69120 Heidelberg

freitag@mathi.uni-heidelberg.de

Riccardo Salvati Manni

Dipartimento di Matematica,

Piazzale Aldo Moro, 2

I-00185 Roma, Italy.

salvati@mat.uniroma1.it

Heidelberg-Roma 2025

Introduction

We consider the Riemann theta series

$$\vartheta[m](\tau, z) = \sum_{p \in \mathbb{Z}^g} \exp\left(\pi i \tau \left[p + \frac{a}{2} \right] + 2 \left(p + \frac{a'}{2} \right) \left(z + \frac{b}{2} \right)\right).$$

Here $m \in \mathbb{Z}^{2g}$ is a column vector that is divided into two columns $a, b \in \mathbb{Z}^g$,

$$m = \begin{pmatrix} a \\ b \end{pmatrix}.$$

The matrix τ is from the Siegel upper half plane \mathcal{H}_g . This means that τ is a symmetric $g \times g$ -matrix such that its imaginary part is positive definite. Finally z is a column in \mathbb{C}^g . Of particular interest are the theta nullwerte

$$\vartheta[m](\tau) := \vartheta[m](\tau, 0).$$

Up to sign they depend only on $m \bmod 2$. In this context the columns $m \in \mathbb{Z}^{2g}$ are called characteristics. A characteristic m is called even (odd) if $a'/b \equiv 0 \pmod{2}$ ($\equiv 1 \pmod{2}$). One knows that a theta nullwert vanishes if and only if m is odd.

Let T_g be the vector space generated by the fourth powers of the theta nullwerte. These are modular forms of weight 2 with respect to the principal congruence subgroup of level two and a joint non-trivial quadratic character. In [Ge], Bert van Geemen's result states

$$\dim(T_g) = (2^g + 1)(2^{g-1} + 1)/3.$$

In [SM2], it has been observed, as consequence of results of Fay [Fa], that all linear relations between the θ_m^4 are consequences of the quartic Riemann relations.

In this note, we want to give a new proof of these results and extend them. Starting from Fay's results about Riemann's relations [Fa], we prove that the representation of $\mathrm{Sp}(g, \mathbb{F}_2)$, induced by a character of the theta group modulo 2, has two irreducible components that result to be isomorphic to the eigenspaces of Fay's matrix M^+ . One component is isomorphic to the space of Riemann's relations, the other is isomorphic to T_g . Then we prove a similar result for vector valued modular forms that are attached to odd characteristics. Here we have to use Fay's matrix M^- .

It is a pleasure to acknowledge the discussions we had with B. van Geemen

1. The symplectic group over the field of two elements

We use the following notations for matrices A . The transposed of A is denoted by A' . Its trace is denoted by $\mathrm{tr}(A)$ and A_0 is the diagonal of A written as column vector. Let A be an $n \times n$ -matrix and B an $n \times m$ -matrix, then

$$A[B] := B'AB.$$

Let $G = G_g = \mathrm{Sp}(g, \mathbb{F}_2)$ be the symplectic group over the field of two elements. We recall the affine action of G on \mathbb{F}^{2g} ,

$$\sigma\{m\} = \sigma'^{-1}m + \begin{pmatrix} (CD')_0 \\ (AB')_0 \end{pmatrix}$$

Here

$$\sigma = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

denotes the decomposition of σ into 4 $g \times g$ -blocks. It is an action in the sense

$$(\sigma\tau)\{m\} = \sigma\{\tau\{m\}\}.$$

In this context the elements of \mathbb{F}_2^{2g} are also called characteristics. A characteristic

$$m = \begin{pmatrix} a \\ b \end{pmatrix}$$

is called even if $a'b = 0$ otherwise odd. We denote the subset of even (odd) characteristics by

$$(\mathbb{F}_2^{2g})_{\mathrm{even}}, \quad (\mathbb{F}_2^{2g})_{\mathrm{odd}}.$$

Their orders are

$$k_g^\pm = 2^{g-1}(2^g \pm 1).$$

It is known the the affine action preserves the parity. Moreover, the group G acts double transitive on the set of all even (odd) characteristics and it acts transitive on the set of all pairs (m, n) where m is even and n odd (see [Ig1], Chapt. V, Prop. 2).

We introduce the stabilizers

$$H(m) = \{\sigma \in G; \quad \sigma\{m\} = m\}.$$

In the case $m = 0$ this is the so-called theta group. We denote it by $H = H(0)$. We will have to use the following function on $G \times \mathbb{F}_2^{2g}$

$$\varepsilon_m(\sigma) = (-1)^{\text{tr}(B'C) + (B'D)[a] + (A'C)[b]}.$$

This function arises in the theta transformation formalism as we will see in the next section. At the moment we take it as it is. It has three important properties.

1.1 Lemma. *The following rules hold.*

LvH

- a) *One has that $\varepsilon_m(\sigma)$ for fixed m is a non-trivial (quadratic) character on $H(m)$. For $h \in H$ one has $\varepsilon_m(h) = \varepsilon_0(h)$.*
- b) *For $h \in H, \sigma \in G$ one has*

$$\varepsilon_m(h)\varepsilon_m(\sigma) = \varepsilon_m(h\sigma).$$

- c) *For $\sigma, \tau \in G$ one has*

$$\varepsilon_m(\tau\sigma) = \varepsilon_m(\sigma)\varepsilon_{\sigma\{m\}}(\tau).$$

Proof. It can be proved directly by means of somewhat tedious calculations. Since it is also a direct consequence of the theta transformation formalism which we treat in the next section, we skip the direct proof and leave it to the interested reader as an exercise. \square

1.2 Lemma. *The following rule holds.*

LvK

$$e(\sigma\{m\}, \sigma\{n\}) = e(m, n)\varepsilon_m(\sigma)e_n(\sigma).$$

Proof. We use the notations

$$\sigma = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad \sigma(m) := \sigma'^{-1}m$$

and

$$m = \begin{pmatrix} a \\ b \end{pmatrix}, \quad n = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad v = \begin{pmatrix} (CD')_0 \\ (AB')_0 \end{pmatrix}$$

In the proof we will use

$$\begin{aligned} (D'B)_0 + D'(AB')_0 + B'(CD')_0 &= 0, \\ (A'C)_0 + C'(AB')_0 + A'(CD')_0 &= 0. \end{aligned}$$

This follows from $\sigma^{-1}\{\sigma\{0\}\} = 0$. Now we get

$$\begin{aligned} e(\sigma\{m\}, \sigma\{n\}) &= e(\sigma(m), \sigma(n))e(\sigma(m), v)e(v, \sigma(n)) \\ &= e(m, n)e(\sigma(m+n), v) \\ &= e(m, n)(-1)^{(CD')'_0(B(a+\alpha)+A(b+\beta))-(AB')'_0(D(a+\alpha)+C(b+\beta))} \\ &= e(m, n)(-1)^{(D'B)'_0(a+\alpha)+(A'C)'_0(b+\beta)} \\ &= e(m, n)\varepsilon_m(\sigma)\varepsilon_n(\sigma) \end{aligned}$$

which proves the Lemma. □

As special case of the Lemma we get a character

$$\varepsilon_H = \varepsilon_0 : H \longrightarrow \{\pm 1\}.$$

We induce the character ε_H on H to a G -representation $\text{Ind}_H^G(\varepsilon)$. So $\text{Ind}_H^G(\varepsilon)$ consists of all functions $f : G \rightarrow \mathbb{C}$ with the property $f(hg) = \varepsilon_H(h)f(g)$. The group G acts by translation from the right on this space. We mention that this representation has been studied by Frame [Fra]. He proved that it decomposes into two irreducible non-isomorphic representations and he determined the dimensions of the two components. We will reprove this result with a completely different method.

We will construct a basis $X(m)$ of $\text{Ind}_H^G(\varepsilon)$. It is parameterized by even characteristics. Notice that the number of even characteristics equals the index of H in G which is the dimension of $\text{Ind}_H^G(\varepsilon)$. So, let m be an even characteristic. We choose $\sigma \in G$ such that $\sigma\{m\} = 0$. Then we define the following function,

$$X(m) : G \longrightarrow \mathbb{C}.$$

It is defined to be 0 outside the coset $H\sigma$, and on this coset it is defined by

$$X(m)(h\sigma) = \varepsilon_H(h)\varepsilon_m(\sigma).$$

It is essential that this is independent on the choice of σ . The proof follows immediately from Lemma .

We associate to each even m a variable e_m and consider the vector space

$$\sum_{m \text{ even}} \mathbb{C}e_m.$$

We use the basis above to define an isomorphism

$$\text{Ind}_H^G(\varepsilon) \xrightarrow{\sim} \sum_{m \text{ even}} \mathbb{C}e_m.$$

Just map $X(m)$ to e_m .

Next we describe an action of G on the right hand side such that this isomorphism is G -equivariant. We have to associate to each $\sigma \in G$ an isomorphism

$$\sigma : \sum_{m \text{ even}} \mathbb{C}e_m \xrightarrow{\sim} \sum_{m \text{ even}} \mathbb{C}e_m.$$

On basis elements it is defined by

$$\sigma(e_m) = \varepsilon_m(\sigma)e_{\sigma\{m\}}.$$

It is easy to check that this defines a representation of G and that the isomorphism

$$\text{Ind}_H^G(\varepsilon) \xrightarrow{\sim} \sum_{m \text{ even}} \mathbb{C}e_m$$

is G -equivariant.

2. An operator of Fay

We treat the space

$$\sum_{m \text{ even}} \mathbb{C}e_m.$$

together with its G -action. We use the notations

$$\langle m, n \rangle = a'\beta + b'\alpha, \quad m = \begin{pmatrix} a \\ b \end{pmatrix}, \quad n = \begin{pmatrix} \alpha \\ \beta \end{pmatrix},$$

and

$$e(m, n) = (-1)^{\langle m, n \rangle}.$$

Notice that $\langle m, n \rangle$ is the symplectic pairing whose invariance group is $G = \text{Sp}(g, \mathbb{F}_2)$. Fay [Fa] defined the following operator

$$M^+ : \sum_{m \text{ even}} \mathbb{C}e_m \longrightarrow \sum_{m \text{ even}} \mathbb{C}e_m, \quad e_m \longmapsto \sum_n e(m, n)e_n.$$

(Actually Fay ordered the even characteristics and introduced M^+ as a matrix, see [Fa], formula (6).)

2.1 Theorem (Fay). *The operator M^+ has two eigen values, namely -2^{g-1} and 2^g . Let* Tfay

$$V^+ \subset \mathbb{C}^{k_g^+}, \quad W^+ \subset \mathbb{C}^{k_g^+}$$

be the eigenspaces. Then we have

$$\mathbb{C}^{k_g^+} = V^+ \oplus W^-.$$

The dimension of the eigenspaces are

$$\dim V^+ = (2^{2g} - 1)/3, \quad \dim W^+ = (2^g + 1)(2^{g-1} + 1)/3.$$

We need the following result.

2.2 Lemma. *The operator M^+ commutes with the action of G .* Lmp

Proof. Use Lemma . □

As a consequence the two eigenspaces are G -invariant. So we see that $\text{Ind}_H^G(\varepsilon)$ admits two invariant subspaces, one isomorphic to V^+ the other to W^+ .

We need the decomposition of G into double cosets with respect to H . This can be found also in [Fra].

2.3 Lemma. *There are two double cosets in $H \backslash G / H$.* Ldc

We present a different proof. Let $g_1, g_2 \in G - H$. We have to show that they are in the same double coset. We consider the two characteristics $m_1 = g_1\{0\}$, $m_2 = g_2\{0\}$. They are both different from zero. The group H acts transitively on the set of non-zero characteristics (since G acts double transitive on the set of even characteristics). Hence there exist $h_1 \in H$ such that $h_1\{m_1\} = m_2$. This implies $h_2 := g_2^{-1}h_1g_1 \in H$. It follows $g_2 = h_1g_1h_2^{-1}$. □

2.4 Lemma. *The representation $\text{Ind}_H^G(\varepsilon)$ is either irreducible or decomposes into two non-isomorphic irreducible components.* Lrd

Proof. The statement is equivalent to

$$\dim \text{Hom}_G(\text{Ind}_H^G(\varepsilon), \text{Ind}_H^G(\varepsilon)) \leq 2.$$

We use Frobenius reciprocity. For a representation of G on a finite dimensional vector space V the evaluation map

$$\text{Hom}_G(V, \text{Ind}_H^G(\varepsilon)) \xrightarrow{\sim} \text{Hom}_H(V, \mathbb{C}(\varepsilon))$$

is an isomorphism. Here $\mathbb{C}(\varepsilon)$ is the one dimensional vector space \mathbb{C} where H acts by means of the character ε . So we have to show

$$\dim \text{Hom}_H(\text{Ind}_H^G(\varepsilon), \mathbb{C}(\varepsilon)) \leq 2.$$

Since both representations are real, they are isomorphic to their duals. Hence it is equivalent to prove

$$\dim \text{Hom}_H(\mathbb{C}(\varepsilon), \text{Ind}_H^G(\varepsilon)) \leq 2.$$

Such a homomorphism is defined by the image f of $1 \in \mathbb{C}$ and f must have the property

$$f(hg) = \varepsilon_H(h)f(g) \quad \text{and} \quad f(gh) = \varepsilon_H(h)f(g).$$

Since G has only two double cosets mod H , the dimension of the space of all f is ≤ 2 . \square

We found two non-zero invariant subspaces of $\text{Ind}_H^G(\varepsilon)$. So Lemma shows that they are irreducible and non-isomorphic. So we have proved the following theorem.

2.5 Theorem (Frame). *The representation $\text{Ind}_H^G(\varepsilon)$ decomposes into two non-isomorphic representations of dimensions* TF

$$(2^{2g} - 1)/3, \quad (2^g + 1)(2^{g-1} + 1)/3.$$

3. Theta series

The symplectic group $\text{Sp}(g, \mathbb{R})$ acts on the Siegel half plane \mathcal{H}_g through

$$\sigma(\tau) = (A\tau + B)(C\tau + D)^{-1}.$$

Remember that the Riemann theta functions are defined by

$$\vartheta[m](\tau, z) = \sum_{p \in \mathbb{Z}^g} \exp\left(\pi i \tau \left[p + \frac{a}{2} \right] + 2\left(p + \frac{a'}{2}\right) \left(z + \frac{b}{2}\right)\right).$$

Here

$$m = \begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{Z}^{2g}$$

is an integral column vector. The theta transformation formula states [Ig1], Chapt. II, Corollary to Theorem 6.)

3.1 Theorem. *We have*

Ttt

$$\vartheta[\sigma\{m\}](\sigma(\tau), (C\tau + D)^{-1}z) = \exp \pi i \{((C\tau + D)^{-1}C)[z]\} \\ v(\sigma, m) \sqrt{\det(C\tau + D)} \vartheta[m](\tau, z)$$

for all $\sigma \in \Gamma_n$. Here $v(\sigma, m)$ denotes a system of complex numbers which is independent of (τ, z) . One has $v(\sigma, m)^8 = 1$.

Corollary. *For the theta nullwerte this means*

$$\vartheta[\sigma\{m\}](\sigma(\tau)) = v(\sigma, m) \sqrt{\det(C\tau + D)} \vartheta[m](\tau)$$

The system of numbers $v(\sigma, m)$ is very complicated, We will need only the 4th powers. It is known that these depend only on $m \bmod 2$ and on the image of σ with respect to the homomorphism $\mathrm{Sp}(g, \mathbb{Z}) \rightarrow \mathrm{Sp}(g, \mathbb{F}_2)$. This homomorphism is surjective. So we can define

$$v(\sigma, m)^4 \quad \text{for } \sigma \in \mathrm{Sp}(g, \mathbb{F}_2), \quad m \in \mathbb{F}_2^{2g}.$$

This function is known explicitly.

3.2 Lemma. *One has*

Lveps

$$v(\sigma, m)^4 = \varepsilon_m(\sigma) = (-1)^{\mathrm{tr}(B'C) + (B'D)[a] + (A'C)[b]}$$

A proof can be found in [Ig2], Lemma 10. □

This was the expression we introduced in Sect. 1. It gets now a natural interpretation. The Lemma is an easy consequences of this interpretation.

Now we consider the linear map

$$\mathrm{Ind}_H^G(\varepsilon) \longrightarrow T_g, \quad X(m) \longmapsto \vartheta[m]^4.$$

The theta transformation formula shows that this map is compatible with the action of G . It is also clear that it is surjective. Its dimension is smaller than that of $\mathrm{Ind}_H^G(\varepsilon)$. To show this, one needs that there exists at least on non-trivial relation between the $\vartheta[m]^4$. The quartic Riemann relations give one. (In the next Theorem we will describe a space of such relations.) Therefore one of the two irreducible components must map isomorphically to T_g . We have to work out which one.

For this we formulate some of the Riemann theta relations in a way that has been described by Fay [Fa]. In Sect. 1 we introduced the operator

$$M^+ : \sum_{m \text{ even}} \mathbb{C} e_m \longrightarrow \sum_{m \text{ even}} \mathbb{C} e_m$$

and described the eigenspaces. The original meaning of this operator is also due to Fay [Fa]. He showed that the Riemann quartic relations imply the following result.

3.3 Theorem (Fay). *Let v be an eigen vector of M^+ with eigenvalue -2^{g-1} (i.e., $v \in V^+$). Then the relation* Tfayz

$$\sum_m v(m)\vartheta[m]^4 = 0$$

holds.

This shows that the irreducible component that maps isomorphically to V^+ goes to zero in T_g . Hence the other component W^+ maps isomorphically to T_g . We obtain the first main result.

3.4 Theorem. *We have* TvG

$$\dim T_g = (2^g + 1)(2^{g-1} + 1)/3.$$

The linear relations between the $\vartheta[m]^4$ are consequences of Riemann's quartic relations.

4. A variant of van Geemen's result

Let $f(\tau, z)$ be a holomorphic function on $\mathcal{H}_g \times \mathbb{C}^g$. Then we can define the holomorphic vector valued function

$$J(f) : \mathcal{H}_g \longrightarrow \mathbb{C}^g, \quad J(f) = \text{grad}_z f(\tau, z)|_{z=0}.$$

We are interested in $J(\vartheta[m])$. This is different from 0 if and only if m is odd.

4.1 Theorem. *For $\sigma \in \Gamma_g$ and odd $m \in \mathbb{Z}^{2g}$ the transformation formula* Tot

$$J(\vartheta[\sigma\{m\}])(\sigma(\tau)) = v(M, \sigma) \sqrt{\det(C\tau + D)} (C\tau + D)(\vartheta[m])(\tau)$$

holds.

This is a consequence of Theorem (see for example [SM1]). □

We consider the fourth symmetric tensor product $\text{Sym}^4(\mathbb{C}^g)$ and define the functions

$$S(\vartheta[m]) : \mathcal{H}_g \rightarrow \text{Sym}^4(\mathbb{C}^g), \quad \tau \longmapsto \text{Sym}^4 J(\vartheta[m])(\tau).$$

The space generated by these k_g^- tensors is called by S_g . A special case of Theorem states

$$S(\vartheta[\sigma\{m\}])(\sigma(\tau)) = \varepsilon_m(\sigma) \det(C\tau + D)^2 \text{Sym}^4(C\tau + D) S(\vartheta[m])(\tau).$$

This shows that the $S(\vartheta[m])$ are vector valued modular forms on the principal congruence subgroup of level two with the character ε_0 and with respect to the $\mathrm{GL}(n, \mathbb{C})$ representation

$$\det(CZ + D)^2 \mathrm{Sym}^4(CZ + D)$$

This representation is irreducible and has highest weight $(6, 2, \dots, 2)$.

Up to the weight this is the same transformation formula as for the $\vartheta[m]^4$ in the even case.

The tensors $S(\vartheta[m])$ satisfy similar linear relations as the $\theta[m]^4$.

Instead of the theta group H we now have to use the stabilizer $K = H(m)$ of an odd characteristic m . To be concrete, we take

$$m' = (1, 0, \dots, 0, 1, 0, \dots, 0).$$

Now we have to consider the non trivial character $\varepsilon_K = \varepsilon(m)$. The induced representation $\mathrm{Ind}_K^G(\varepsilon_K)$ comes into the game. Its dimension equals the number k_g^- of odd characteristics. We will construct a basis $Y(m)$ of $\mathrm{Ind}_K^G(\varepsilon_K)$. It is parameterized by odd characteristics. So, let m be an odd characteristic. We choose $\sigma \in G$ such that $\sigma\{m\} = 0$. Then we define the following function, $Y(m) : G \rightarrow \mathbb{C}$ analogous to $X(m)$. It is defined to be 0 outside the coset $K\sigma$, and on this coset it is defined by

$$X(m)(k\sigma) = \varepsilon_K(k)\varepsilon_m(\sigma).$$

It is also independent on the choice of σ . Again this follows from Lemma .

We associate to each odd m a variable e_m and consider the vector space

$$\sum_{m \text{ odd}} \mathbb{C}e_m.$$

We use the basis above to define an isomorphism

$$\mathrm{Ind}_K^G(\varepsilon) \xrightarrow{\sim} \sum_{m \text{ odd}} \mathbb{C}e_m.$$

Just map $Y(m)$ to e_m .

Besides M^+ Fay [Fa] defined the following operator

$$M^- \sum_{m \text{ odd}} \mathbb{C}e_m \longrightarrow \sum_{m \text{ odd}} \mathbb{C}e_m, \quad e_m \longmapsto \sum_n e(m, n)e_n.$$

4.2 Theorem (Fay). *The operator M^- has two eigen values, namely 2^{g-1} and -2^g . Let*

$$V^- \subset \mathbb{C}^{k_g^+}, \quad W^- \subset \mathbb{C}^{k_g^+}$$

be the eigenspaces. Then we have

$$\mathbb{C}^{k_g^-} = V^- \oplus W^-.$$

The dimension of the eigenspaces are

$$\dim V^- = (2^{2g} - 1)/3, \quad \dim W^- = (2^g - 1)(2^{g-1} - 1)/3.$$

The same proof as in Lemma shows that M^- commutes with G . Hence the two spaces V^- and W^- are G -invariant subspaces.

With the same method, using Fay's matrix M^- instead of M^+ , one can get now a new proof of the following result of Frame.

4.3 Theorem (Frame). *Let $K = H(m)$ be the stabilizer of an odd character-
istic m . Denote by ε_K the distinguished character of K . Then the representa-
tion $\text{Ind}_K^G(\varepsilon_K)$ decomposes into two non-isomorphic irreducibles representation
of degrees* Rfr

$$(2^{2g} - 1)/3, \quad (2^g - 1)(2^{g-1} - 1)/3.$$

As we have seen, the group $\text{Sp}(g, \mathbb{F}_2)$ acts on the space S_g . The linear map

$$\text{Ind}_K^G(\varepsilon_K) \longrightarrow S_g, \quad Y(m) \longmapsto S(\vartheta[m]),$$

is G -linear. It is surjective. Hence it is an isomorphism or one of the two irreducible complements maps isomorphically. We will show that the second case holds. For this we have to make use of Fay's theta relations. A special case of [Fa] (23) says for $z_1 - z_2 = z_3 - z_4$ and $w \in W^-$

$$\sum_{m \text{ odd}} w_m \vartheta[m](\tau, z_1) \vartheta[m](\tau, z_2) \vartheta[m](\tau, z_3) \vartheta[m](\tau, z_4) = 0$$

or

$$\sum_{m \text{ odd}} w_m \vartheta[m](\tau, z_1) \vartheta[m](\tau, z_2) \vartheta[m](\tau, z_3) \vartheta[m](\tau, -z_1 + z_2 + z_3) = 0$$

with three independent variables z_1, z_2, z_3 . We take the gradient with respect to the variable z_3 and take then $z_3 = 0$. Since $\vartheta[m](\tau, 0) = 0$ for odd m , we obtain

$$\sum_{m \text{ odd}} w_m \vartheta[m](\tau, z_1) \vartheta[m](\tau, z_2) \vartheta[m](\tau, -z_1 + z_2) J(\vartheta[m])(\tau) = 0.$$

Taking the z_2 -gradient gives

$$\sum_{m \text{ odd}} w_m \vartheta[m](\tau, z_1) (\vartheta[m](\tau, -z_1) J(\vartheta[m])(\tau)) \otimes J(\vartheta[m])(\tau) = 0.$$

The theta functions $\vartheta[m](\tau, z)$ are invariant under $z \mapsto -z$ up to a sign. Now differentiating twice by z_1 and then setting $z_1 = 0$ gives

$$\sum_{m \text{ odd}} w_m S(\vartheta[m]) = 0.$$

So we get non trivial linear relations between the $S(\theta[m])$. Hence the space S_g must be isomorphic to one of the two irreducible components of $\text{Ind}_K^G(\varepsilon_K)$.

4.4 Theorem. *The dimension of the space spanned by the tensors $S(\vartheta[m])$ has dimension $(2^{2g} - 1)/3$.* Smt

References

- [Fa] Fay, J.F.: *On the Riemann-Jacobi Formula*, Nachr. Akad. Wiss. Göttingen, 1979
- [Fr] Frame, J.S.: *Some Characters of Orthogonal Groups over the field of two Elements*, Lecture Notes in Mathematics **372**, Proceedings of the Second International Conference on the Theory of groups, Edited by M.F. Newman Springer-Verlag Berlin Heidelberg u.a., 1974
- [Ge] van Geemen, B.: *Siegel modular forms vanishing on the moduli space of curves*, Invent. math. **78**, 1984
- [Ig1] Igusa, J.I.: *Theta Functions*, Grundlehren der mathematischen Wissenschaften, Band **194**, Springer-Verlag Berlin Heidelberg u.a., 1972
- [Ig2] Igusa, J.I.: *On Jacobi's Derivative Formula and its Generalizations*, Am. Journ. Math., Vol, **102**, No. 2, 1980
- [SM1] Salvati Manni, R.: *On the dimension of the vector space $\mathbb{C}[\theta_m]_4$* , Nagoya Math. J. Vol. **98**, 1985
- [SM2] Salvati Manni, R.: *On the Not Identically Zero Nullwerte of Jacobians of Theta Functions with Odd Characteristics*, Advances in Math. **47**, 1983