# DEPTH OF POWERS OF EDGE IDEALS OF EDGE-WEIGHTED INTEGRALLY CLOSED CYCLES

GUANGJUN ZHU<sup>∗</sup> , JIAXIN LI, YIJUN CUI AND YI YANG

Abstract. This paper gives some exact formulas for the depth of powers of the edge ideal of an edge-weighted integrally closed cycle.

# 1. INTRODUCTION

Let G be a finite simple graph with the vertex set  $V(G)$  and the edge set  $E(G)$ . We write xy for an edge of G with x and y as endpoints. Suppose  $\mathbf{w}: E(G) \longrightarrow \mathbb{Z}_{>0}$ is an edge-weighted function (weighted function for short) on  $G$ . We write  $G_{\bf w}$  for the pair  $(G, \mathbf{w})$  and call it an *edge-weighted* graph (or just weighted graph) with the underlying graph G. For a weighted graph  $G_w$  with the vertex set  $V(G)$  =  ${x_1, \ldots, x_n}$ , its *edge-weighted ideal* (edge ideal for short), was introduced in [\[20\]](#page-29-0), is a monomial ideal of the polynomial ring  $S = \mathbb{K}[x_1, \ldots, x_n]$  in n variables over a field K, given by

$$
I(G_{\mathbf{w}}) = (x^{\mathbf{w}(xy)}y^{\mathbf{w}(xy)} \mid xy \in E(G_{\mathbf{w}})).
$$

If **w** is the constant function defined by  $\mathbf{w}(e) = 1$  for all  $e \in E(G)$ , then  $I(G_{\mathbf{w}})$  is the classical edge ideal of the underlying graph  $G$  of  $G_{\mathbf{w}}$ , which has been studied extensively in the literature [\[9,](#page-28-0) [11,](#page-29-1) [13,](#page-29-2) [15,](#page-29-3) [18,](#page-29-4) [19,](#page-29-5) [22,](#page-29-6) [25,](#page-29-7) [26\]](#page-29-8).

In [\[1\]](#page-28-1), Brodmann demonstrated that for a proper ideal  $I \subset S$ , depth $(S/I<sup>t</sup>)$  is a constant for  $t \gg 0$ , and this constant is bounded above by  $n-\ell(I)$ , where  $\ell(I)$  is the analytic spread of I. Both Eisenbud and Huneke in [\[7,](#page-28-2) Proposition 3.3] and Herzog and Hibi in [\[13,](#page-29-2) Theorem 1.2] independently showed that the upper bound can be obtained when the associated graded ring is Cohen–Macaulay. Furthermore, Herzog and Hibi proved [\[13\]](#page-29-2) that  $\text{depth}(S/I^t)$  is a non-increasing function if all the powers of I have a linear resolution. They also provided lower bounds for depth $(S/I<sup>t</sup>)$  when all the powers of I have linear quotients, a condition that entails that all powers of I have linear resolutions. In particular, they showed that the edge ideal of a finite graph whose complementary graph is chordal has linear quotients. Morey in [\[18\]](#page-29-4) gives lower bounds on the depths of all powers when I is the edge ideal of a forest. For the general graph, Fouli1 and Morey in [\[9\]](#page-28-0) gave lower bounds for its edge ideal. In contrast, little is known about the depth of the quotient ring  $S/I<sup>t</sup>$  when I is the edge ideal of a weighted graph.

Recently, there has been a surge of interest in characterizing weights for which the edge ideal of a weighted graph is Cohen-Macaulay. For example, Paulsen and

<sup>∗</sup> Corresponding author.

<sup>2020</sup> *Mathematics Subject Classification*. Primary 13B22, 13F20; Secondary 05C99, 05E40. Keywords: Depth, edge-weighted graph, powers of the edge ideal, integrally closed cycle.

Sather-Wagstaff in [\[20\]](#page-29-0) classified Cohen-Macaulay weighted graphs  $G_w$  where the underlying graph  $G$  is a cycle, a tree, or a complete graph. Seyed Fakhari et al. in  $[8]$ continued this study and classified a Cohen-Macaulay weighted graph  $G_w$  when G is a very well-covered graph. Hien in [\[14\]](#page-29-9) classified Cohen-Macaulay weighted graphs  $G_{\mathbf{w}}$  when G has girth at least 5. Recently, the first second authors of this paper and Trung in [\[16\]](#page-29-10) characterized the Cohen-Macaulayness of the second power of the edge ideal of  $G_w$  when the underlying graph G is a very well-covered graph and all ordinary powers of the edge ideal of  $G_w$  when G is a tree with a perfect matching consisting of pendant edges. In [\[24\]](#page-29-11), Wei classified all Cohen-Macaulay weighted chordal graphs from a purely graph-theoretic point of view. Diem et al. in [\[5\]](#page-28-4) gave a complete characterization of sequentially Cohen-Macaulay edge-weighted graphs. The first three authors of this paper and Duan in [\[28\]](#page-29-12) gave some exact formulas for the depth of the powers of the edge ideal of a weighted star graph and also gave some lower bounds on the depth of powers of the integrally closed weighted path.

In this paper, we are interested in the depth of powers of the edge ideal  $I(C_{\bf w}^n)$  of a weighted integrally closed *n*-cycle  $C_{\mathbf{w}}^n$ . The results we obtained are as follows:

**Theorem 1.1.** Let  $C^n_{\mathbf{w}}$  be a non-trivially weighted integrally closed n-cycle. Then *we have*

(1) *if*  $C_w^n$  has only one edge with non-trivial weight, then

$$
\text{depth}(S/I(C_{\mathbf{w}}^n)^t) = \begin{cases} \lceil \frac{n-t}{3} \rceil, & \text{if } 1 \le t < \lceil \frac{n+1}{2} \rceil, \\ 1, & \text{if } t \ge \lceil \frac{n+1}{2} \rceil, \ n \text{ is even,} \\ 0, & \text{if } t \ge \lceil \frac{n+1}{2} \rceil, \ n \text{ is odd.} \end{cases}
$$

(2) if  $C_{\bf w}^n$  has two edges with non-trivial weights, then

$$
depth(S/I(C_{\mathbf{w}}^n)^t) = \max\{\lceil \frac{n-t}{3} \rceil, 1\} \text{ for all } t \ge 1.
$$

(3) if  $C_{\bf w}^n$  has three edges with non-trivial weights, then

$$
\operatorname{depth}(S/I(C_{\mathbf{w}}^n)^t) = 2 \text{ for all } t \ge 1.
$$

The article is organized as follows. In Section [2,](#page-2-0) we provide the definitions and basic facts that will be used throughout this paper. Sections [3](#page-3-0) and [4](#page-6-0) are devoted to prove our main results. By the depth lemma, we provide exact formulas for depth $(S/I(C_{\bf w}^n))$  for any weighted integrally closed *n*-cycle  $C_{\bf w}^n$ , and exact formulas for depth $(S/I(C_{\bf w}^n)^t)$  if  $C_{\bf w}^n$  has three edges with non-trivial weights or  $C_{\bf w}^3$  has two edges with non-trivial weights. If  $C_{\mathbf{w}}^n$  ( $n \geq 4$ ) has two edges with non-trivial weights, we can give the lower bound for  $\text{depth}(S/I^t)$  by using the depth lemma, then, by choosing suitable  $f \notin I^t$  and using the fact that  $\text{depth}(S/I(C_{\bf w}^n)^t) \leq \text{depth}(S/(I^t :$ f)), we prove that  $\text{depth}(S/I^t)$  just reaches this lower bound. If  $C_{\bf w}^n$  has only one edge with non-trivial weight, we can also give exact formulas for depth $(S/I(C_{\bf w}^n)^t)$ using methods from [\[17,](#page-29-13) [27\]](#page-29-14).

# 2. Preliminaries

<span id="page-2-0"></span>In this section, we will provide the definitions and basic facts that will be used throughout this paper. For detailed information, we refer to [\[2\]](#page-28-5) and [\[12\]](#page-29-15).

Let G be a finite simple graph with the vertex set  $V(G)$  and the edge set  $E(G)$ . For any subset A of  $V(G)$ ,  $G[A]$  denotes the *induced subgraph* of G on the set A, i.e., for any  $u, v \in A$ ,  $uv \in E(G[A])$  if and only if  $uv \in E(G)$ . The induced subgraph of G on the set  $V(G)\backslash A$  is denoted by  $G\backslash A$ . In particular, if  $A = \{v\}$  then we will write  $G\backslash v$ instead of  $G \setminus \{v\}$  for simplicity. For a vertex  $v \in V(G)$ , its *neighborhood* is defined as  $N_G(v) := \{u \in V(G) \mid uv \in E(G)\}\$ , and its degree is defined as  $\deg_G(v) = |N_G(v)|$ . A connected graph G is called a *cycle* if  $deg_G(v) = 2$  for all  $v \in V(G)$ . A cycle with n vertices is called an n-cycle, denoted by  $C<sup>n</sup>$ . A connected graph with vertex set  $\{x_1, \ldots, x_n\}$ , is called a *path*, if  $E(G) = \{x_i x_{i+1} \mid 1 \leq i \leq n-1\}$ . Such a path is denoted by  $P^n$ .

Given a weighted graph  $G_{\mathbf{w}}$ , we denote its vertex and edge sets by  $V(G_{\mathbf{w}})$  and  $E(G_{\bf w})$ , respectively. Any concept valid for graphs automatically applies to weighted graphs. For example, the *neighborhood* of a vertex v in a weighted graph  $G_w$  with the underlying graph G is defined as  $N_{G_{\mathbf{w}}}(v) := \{u \in V(G) \mid uv \in E(G)\}\.$  For a subset A of  $V(G)$ , the *induced subgraph* of  $G_w$  on the set A is the graph  $G_w[A]$ with vertex set A, and for any  $u, v \in A$ ,  $uv \in E(G_{\bf w}[A])$  if and only if  $uv \in E(G_{\bf w}),$ and the weight function **w'** satisfies  $\mathbf{w}'(uv) = \mathbf{w}(uv)$ . The induced subgraph of  $G_{\mathbf{w}}$ on the set  $V(G) \setminus A$  is denoted by  $G_w \setminus A$ . In particular, if  $A = \{v\}$  then we will write  $G_{\mathbf{w}} \setminus v$  instead of  $G_{\mathbf{w}} \setminus \{v\}$  for simplicity. A weighted graph is said to be a *non-trivially* weighted graph if there is at least one edge with a weight greater than 1, otherwise, it is said to be a *trivially* weighted graph. An edge  $e \in E(G_{\bf w})$  is said to be *non-trivial* if its weight  $w(e) \geq 2$ . Otherwise, it is said to be *trivial*.

<span id="page-2-1"></span>The following lemmas are often used to compute the depth of a module.

**Lemma 2.1.** ([\[15,](#page-29-3) Lemmas 2.1]) *Let*  $0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$  *be a short exact sequence of finitely generated graded S-modules. Then*

- (1) depth $(N) \geq \min{\{\text{depth}(M), \text{depth}(P)\}}$ *, the equality holds if* depth $(P) \neq$  $depth(M) - 1;$
- (2) depth $(M) \ge \min\{\text{depth}(N), \text{depth}(P)+1\}$ *, the equality holds if* depth $(N) \ne$ depth(P)*;*
- (3) depth $(P) \ge \min{\{\operatorname{depth}(M) 1, \operatorname{depth}(N)\}\}\$ , the equality holds if depth $(M) \ne$  $depth(N)$ .

<span id="page-2-2"></span>**Lemma 2.2.** ([\[15,](#page-29-3) Lemma 2.2]) Let  $S_1 = \mathbb{K}[x_1, \ldots, x_m]$  and  $S_2 = \mathbb{K}[x_{m+1}, \ldots, x_n]$ *be two polynomial rings over a field* K*. Let*  $S = S_1 \otimes_{\mathbb{K}} S_2$  *and*  $I \subset S_1$ *,*  $J \subset S_2$  *be two nonzero homogeneous ideals. Then*

- (1) depth $(S/(I+J)) =$  depth $(S_1/I) +$  depth $(S_2/J)$ ;
- (2)  $\text{depth}(S/JI) = \text{depth}(S_1/I) + \text{depth}(S_2/J) + 1.$

<span id="page-2-3"></span>Lemma 2.3. ([\[21,](#page-29-16) Corollary 1.3], [\[15,](#page-29-3) Lemma 2.1]) *Let* I ⊂ S *be a monomial ideal* and  $f \notin I$  *be a monomial of degree s. Then* 

$$
\text{depth}(S/I) \le \text{depth}(S/(I : f)).
$$

*Furthermore, it follows from Lemma [2.1](#page-2-1) and the following short exact sequence*

<span id="page-3-4"></span>(1) 
$$
0 \longrightarrow \frac{S}{I:f}(-s) \stackrel{f}{\longrightarrow} \frac{S}{I} \longrightarrow \frac{S}{(I,f)} \longrightarrow 0
$$

*that*

$$
\operatorname{depth}(S/I) \ge \min\{\operatorname{depth}(S/(I : f)), \operatorname{depth}(S/(I, f))\}.
$$

<span id="page-3-2"></span>**Lemma 2.4.** ([\[17,](#page-29-13) Theorem 3.6]) Let  $P_{\bf w}^n$  be a trivially weighted path, then for all  $t \geq 1$ ,

$$
depth(S/I(P_w^n)^t) = \max\{\lceil \frac{n-t+1}{3}\rceil, 1\}.
$$

#### 3. The depth of the edge ideal

<span id="page-3-0"></span>Calculating or even estimating the depth of edge ideals of weighted cycles with arbitrary weight functions is a difficult problem. In this section, we will give precise formulas for the depth of the edge ideal of a non-trivially weighted integrally closed n-cycle. First, we recall that the definition of an ideal is integrally closed.

Definition 3.1. ([\[12,](#page-29-15) Definition 1.4.1]) *Let* I *be an ideal in a ring* R*. An element* f ∈ R *is said to be* integral *over* I *if there exists an equation*

$$
f^k + c_1 f^{k-1} + \cdots + c_{k-1} f + c_k = 0
$$
 with  $c_i \in I^i$ .

*The set*  $\overline{I}$  *of elements in* R *which are integral over* I *is the* integral closure *of* I. *If*  $I = \overline{I}$ *, then I is said to be* integrally closed. A weighted graph  $G_{\omega}$  *is said to be* integrally closed *if its edge ideal*  $I(G_\omega)$  *is integrally closed.* 

We will use the following notation for the remainder of this paper. Let  $n$  be a positive integer, the notation [n] denotes by convention the set  $\{1, 2, \ldots, n\}$ . If  $n \geq 3$ , let  $C_{\mathbf{w}}^{n}$  denote a non-trivially weighted integrally closed *n*-cycle with the vertex set  $V(C_{\mathbf{w}}^n) = \{x_1, \ldots, x_n\}$  and the edge set  $E(C_{\mathbf{w}}^n)$ . We also denote  $e_i = x_i x_{i+1} \in E(C_{\mathbf{w}}^n)$ with  $x_{n+1} = x_1$  and  $\mathbf{w}_i = \mathbf{w}(e_i)$  for all  $i \in [n]$ . By symmetry, we can assume that  $\mathbf{w}_1 \geq 2$ ,  $\mathbf{w}_1 \geq \mathbf{w}_3$  and  $\mathbf{w}_3 \geq \mathbf{w}_5$  if  $e_5 \in E(C_\mathbf{w}^n)$ . Let  $P_\mathbf{w}^n$  be an induced subgraph of the  $(n + 1)$ -cycle  $C_{\mathbf{w}}^{n+1}$  on the set  $\{x_1, x_2, ..., x_n\}$ .

According to [\[12,](#page-29-15) Theorem 1.4.6], every trivially weighted graph  $G_w$  is integrally closed. The following lemma gives a complete characterization of a non-trivially weighted graph that is integrally closed.

<span id="page-3-1"></span>**Lemma 3.2.** ([\[6,](#page-28-6) Theorem 3.6] and [\[23,](#page-29-17) Theorem 2.8]) Let  $G_w$  be a non-trivially *weighted graph, then* G<sup>w</sup> *is integrally closed if and only if* G<sup>w</sup> *does not contain any of the following three graphs as its induced subgraphs.*

- $(1)$  *A path*  $P_w^3$  *of length* 2 *where all edges have non-trivial weights.*
- (2) *The disjoint union*  $P_w^2 \sqcup P_w^2$  *of two paths*  $P_w^2$  *of length* 1 *where all edges have non-trivial weights.*
- (3) *A* 3*-cycle*  $C_w^3$  *where all edges have non-trivial weights.*

<span id="page-3-3"></span>From the above lemma, we can derive

**Corollary 3.3.** Let  $C^n_{\bf w}$  be a non-trivially weighted integrally closed n-cycle. If n = 6*, it can have at most three edges with non-trivial weights, otherwise it can have up to two edges with non-trivial weights.*

<span id="page-4-0"></span>The following lemmas provide some formulas for the depth of powers of the edge ideal of a non-trivially integrally closed path.

**Lemma 3.4.** ([\[28,](#page-29-12) Theorems 4.8 and 4.9]) Let  $P_{\mathbf{w}}^n$  be a non-trivially weighted inte*grally closed path.* Let  $\mathbf{w}_i \geq 2$  *and*  $\mathbf{w}_i \geq \mathbf{w}_{i+2}$  *if*  $e_{i+2} \in E(P_{\mathbf{w}}^n)$ *. Then* 

(1) if  $n \leq 3$ , then depth $(S/I(P_{\bf w}^n)) = 1$ ; (2) if  $n = 4$ , then depth $(S/I(P_{\bf w}^n)) = 2 - a$ , where  $a =$  $\int 0, \quad \text{if }$ **w**<sub>2</sub> = 1*,* 1, *otherwise. ;* (3) *if*  $n \ge 5$ *, then* depth $(S/I(P_{\bf w}^n)) = \min\{\lceil \frac{i}{3} \rceil + \lceil \frac{n-i-b}{3} \rceil\}$  $\left\lfloor\frac{i-b}{3}\right\rceil, \left\lceil\frac{i-2}{3}\right\rceil$  $\frac{-2}{3}$  +  $\frac{n-i-2}{3}$  $\frac{1}{3}$ <sup> $\frac{1}{3}$ </sup> + 1 },  $where b =$  $\int 0, \quad \text{if }$ **w**<sub>i+2</sub> = 1*,* 1, *otherwise*. *.*

<span id="page-4-2"></span>**Lemma 3.5.** ([\[28,](#page-29-12) Theorems 4.12, 4.17, 4.18 and Proposition 4.13]) Let  $P_{\bf w}^n$  be a *non-trivially weighted integrally closed path as in Lemma [3.4.](#page-4-0)* Then for all  $t \geq 2$ , *we have the following results:*

(1) If  $n \leq 3$ , then depth $(S/I(P_{\bf w}^n)^t) = 1$ ;

(2) *If*  $n = 4$ *, then* 

$$
\text{depth}(S/I(P_{\mathbf{w}}^n)^t) \ge \begin{cases} 2, & \text{if } \mathbf{w}_1, \mathbf{w}_3 > 1 \text{ and } \mathbf{w}_2 = 1, \\ 1, & \text{otherwise.} \end{cases};
$$

(3) If 
$$
n \geq 5
$$
 and  $i = 1$ , then

$$
\text{depth}(S/I(P_{\mathbf{w}}^{n})^{t}) \geq \begin{cases} \max\{\lceil \frac{n-t+1}{3} \rceil, 1\}, & \text{if } \mathbf{w}_{3} = 1, \\ \max\{\lceil \frac{n-t+1}{3} \rceil, 2\}, & \text{if } \mathbf{w}_{3} > 1. \end{cases};
$$

(4) If  $n \geq 5$  and  $i > 1$ , then

$$
\text{depth}(S/I(P_{\mathbf{w}}^{n})^{t}) \geq \begin{cases} \max\{\lceil \frac{n-t}{3} \rceil, 2\}, & \text{if } \mathbf{w}_{i+2} > 1, \\ \lceil \frac{n-1}{3} \rceil, & \text{if } \mathbf{w}_{i+2} = 1 \text{ and } i \equiv 1 \pmod{3}, n \equiv 2 \pmod{3}, \text{ and } t = 2, \\ \max\{\lceil \frac{n-t}{3} \rceil, 1\}, & \text{otherwise.} \end{cases}
$$

The following lemma provides some exact formulas for the depth of powers of the edge ideal of a trivially weighted cycle.

Lemma 3.6. ( [\[3,](#page-28-7) Proposition 1.3]*,* [\[17,](#page-29-13) Theorem 1.1] *and* [\[22,](#page-29-6) Theorems 4.4 and 5.1]) Let  $C_{\mathbf{w}}^n$  be a trivially weighted n-cycle, then

$$
\text{depth}(S/I(C_{\mathbf{w}}^n)^t) = \begin{cases} \lceil \frac{n-1}{3} \rceil, & \text{if } t = 1, \\ \lceil \frac{n-t+1}{3} \rceil, & \text{if } 2 \le t < \lceil \frac{n+1}{2} \rceil, \\ 1, & \text{if } n \text{ is even and } t \ge \frac{n}{2} + 1, \\ 0, & \text{if } n \text{ is odd and } t \ge \frac{n+1}{2}. \end{cases}
$$

<span id="page-4-1"></span>Next, we will compute the depth of the edge ideal of a non-trivially weighted integrally closed n-cycle.

**Theorem 3.7.** Let  $C_{\bf w}^n$  be a non-trivially weighted integrally closed n-cycle, then

$$
\text{depth}(S/I(C_{\mathbf{w}}^n)) = \lceil \frac{n-1}{3} \rceil.
$$

*Proof.* Let  $I = I(C_{\bf w}^n)$  and  ${\bf w}_1 = \max{\{\bf w}_1,\ldots,\bf w}_n\}$  by symmetry. If  $n = 3$ , then  $\mathbf{w}_2 = 1$  or  $\mathbf{w}_3 = 1$  by Lemma [3.2.](#page-3-1) We can assume  $\mathbf{w}_3 = 1$  by symmetry. It is easy to see that  $(I : x_3) = (x_2^{w_2} x_3^{w_2-1}, x_1)$  and  $(I, x_3) = (x_1^{w_1} x_2^{w_1}, x_3)$ . Thus  $depth(S/(I : x_3)) = depth(S/(I, x_3)) = 1.$  Applying Lemma [2.1](#page-2-1) to the following exact sequence

$$
0 \longrightarrow \frac{S}{I:x_3}(-1) \xrightarrow{.x_3} \frac{S}{I} \longrightarrow \frac{S}{(I,x_3)} \longrightarrow 0,
$$

we can observe that  $\text{depth}(S/I) = 1$ .

In the following, we assume that  $n \geq 4$ . We distinguish between the following two cases:

*Case* 1: If  $C_w^n$  has at most two edges with non-trivial weights, then we can assume by symmetry and Lemma [3.2](#page-3-1) that  $\mathbf{w}_1 \geq \mathbf{w}_3$  and  $\mathbf{w}_i = 1$  for all  $i \neq 1, 3$ . It is easy to see that  $(I, x_2) = I(C_{\mathbf{w}}^n \backslash x_2) + (x_2)$  and  $(I : x_2) = I(P_{\mathbf{w}}^{n-2}) + (x_1^{\mathbf{w}_1} x_2^{\mathbf{w}_1-1}, x_3),$ where  $P_n^{n-2}$  is an induced subgraph of  $C_n^n$  on the set  $\{x_1, \hat{x_2}, \hat{x_3}, x_4, \dots, x_n\}$  and  $\hat{x_2}$ denotes the element  $x_2$  omitted from  $\{x_1, x_2, \ldots, x_n\}$ . Furthermore,  $(I : x_2x_1) =$  $I(P_{\mathbf{w}}^{n-4}) + (x_1^{\mathbf{w}_1-1}x_2^{\mathbf{w}_1-1}, x_3, x_n)$  and  $((I: x_2), x_1) = I(P_{\mathbf{w}}^{n-3}) + (x_1, x_3)$ , where  $P_{\mathbf{w}}^{n-4}$ and  $P_{\mathbf{w}}^{n-3}$  are induced subgraphs of  $C_{\mathbf{w}}^n$  on the sets  $\{x_4, \ldots, x_{n-1}\}$  and  $\{x_4, \ldots, x_n\}$ , respectively. It follows from Lemmas [2.2](#page-2-2) and [2.4](#page-3-2) that depth $(S/(I: x_2 x_1)) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$ ] and depth $(S/((I:x_2),x_1)) = \lceil \frac{n}{3} \rceil$  $\frac{n}{3}$ . We consider the following two subcases:

(i) If  $w_3 = 1$ , then depth $(S/(I, x_2)) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$  by Lemma [2.4.](#page-3-2) It follows that  $\operatorname{depth}(S/I) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$  by applying Lemma [2.1](#page-2-1) to the following short exact sequences

<span id="page-5-0"></span>
$$
(2) \quad 0 \quad \longrightarrow \quad \frac{S}{I:x_2}(-1) \quad \frac{3}{I}\quad \longrightarrow \quad \frac{S}{I} \quad \longrightarrow \quad \frac{S}{(I,x_2)} \quad \longrightarrow \quad 0,
$$
\n
$$
0 \quad \longrightarrow \quad \frac{S}{I:x_2x_1}(-1) \quad \frac{3}{I:x_2} \quad \longrightarrow \quad \frac{S}{I:x_2} \quad \longrightarrow \quad \frac{S}{((I:x_2),x_1)} \quad \longrightarrow \quad 0.
$$

(ii) If  $w_3 \geq 2$ , then it follows from Lemmas [2.2](#page-2-2) and [3.4](#page-4-0) that

$$
depth(S/(I, x_2)) = \begin{cases} \lceil \frac{n-1}{3} \rceil, & \text{if } n = 4, \\ \lceil \frac{n}{3} \rceil, & \text{if } n \ge 5. \end{cases}
$$

Using Lemma [2.1](#page-2-1) to the short exact sequences [\(2\)](#page-5-0), we get depth $(S/I) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$ .

*Case* 2: If  $C_w^n$  has three edges with non-trivial weights, then  $n = 6$  by Corol-lary [3.3.](#page-3-3) In this case, we can assume by symmetry that  $\mathbf{w}_1 \geq \mathbf{w}_3 \geq \mathbf{w}_5 \geq 2$  and  $\mathbf{w}_2 = \mathbf{w}_4 = \mathbf{w}_6 = 1$ . It is easy to see that  $(I : x_2) = I(P_{\mathbf{w}}^4) + (x_1^{\mathbf{w}_1} x_2^{\overline{\mathbf{w}}_1 - 1}, x_3)$ and  $(I, x_2) = I(P_w^5) + (x_2)$ , where  $P_w^4$  and  $P_w^5$  are induced subgraphs of  $C_w^6$  on the sets  $\{x_1, x_4, x_5, x_6\}$  and  $\{x_1, x_3, x_4, x_5, x_6\}$ , respectively. Furthermore,  $(I : x_2x_1) =$  $(x_1^{w_1-1}x_2^{w_1-1}, x_3, x_4x_5, x_6)$  and  $((I : x_2), x_1) = (x_1, x_3, x_4x_5, (x_5x_6)^{w_5})$ . It follows from Lemmas [2.2,](#page-2-2) [2.4](#page-3-2) and [3.4](#page-4-0) that depth $(S/(I, x_2)) = \text{depth}(S/(I : x_2x_1)) =$  $depth(S/(I : x_2), x_1)) = 2$ . Applying Lemma [2.1](#page-2-1) to the above exact sequences  $(2)$ , we get depth $(S/I) = 2$ .

### 4. The depth of powers of the edge ideal

<span id="page-6-0"></span>In this section, we will give precise formulas for the depth of powers of the edge ideal of a non-trivially integrally closed n-cycle. Thus, in the rest of this section, we let  $C_{\mathbf{w}}^{n}$  be a non-trivially weighted integrally closed *n*-cycle.

4.1.  $C_w^n$  has at least two edges with non-trivial weights. In this subsection we assume that  $\mathbf{w}_1 \geq \mathbf{w}_3 \geq 2$ . First, we consider the case where  $C^n_{\mathbf{w}}$  has three edges with non-trivial weights.

**Theorem 4.1.** If  $C^n_{\bf w}$  has three edges with non-trivial weights, then

 $\operatorname{depth}(S/I(C_{\mathbf{w}}^n)^t) = 2$  *for all*  $t \geq 1$ .

*Proof.* Let  $I = I(C_{\mathbf{w}}^n)$ . By Corollary [3.3,](#page-3-3) we get  $n = 6$ . By symmetry we can assume that  $\mathbf{w}_1 \geq \mathbf{w}_3 \geq \mathbf{w}_5 \geq 2$  and  $\mathbf{w}_2 = \mathbf{w}_4 = \mathbf{w}_6 = 1$ . Now we prove the statements by induction on t with the case  $t = 1$  verified in Theorem [3.7.](#page-4-1) Thus, in the following, we can assume that  $t \geq 2$ . By [\[27,](#page-29-14) Lemmas 4.1 and 4.3], we see that  $(I^t, x_1) = I(C_w^6 \setminus x_1)^t + (x_1), (I^t : x_1 x_6) = I^{t-1}, ((I^t : x_1), x_6) = ((I^t, x_6) :$  $(x_1)$ ,  $(I^t, x_6) = I(C_w^6 \setminus x_6)^t + (x_6)$  and  $((I^t, x_6), x_1) = I(P_w^4)^t + (x_1, x_6)$ , where  $P_w^4$ is the induced subgraph of  $C_w^6$  on  $\{x_2, x_3, x_4, x_5\}$ . It follows from the induction and Lemmas [2.2](#page-2-2) and [3.5](#page-4-2) that  $\text{depth}(S/(I^t, x_1)) \geq 2$ ,  $\text{depth}(S/(I^t : x_1 x_6)) = 2$ ,  $\operatorname{depth}(S/(I^t,x_6)) \geq 2$  and  $\operatorname{depth}(S/((I^t,x_6),x_1)) = 1$ . Thus, by Lemma [2.1](#page-2-1) and the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{I^t:x_1}(-1) & \xrightarrow{\cdot x_1} & \frac{S}{I^t} & \longrightarrow & \frac{S}{(I^t,x_1)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{I^t:x_1x_6}(-1) & \xrightarrow{\cdot x_6} & \frac{S}{I^t:x_1} & \longrightarrow & \frac{S}{((I^t:x_1),x_6)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{((I^t:x_1),x_6)}(-1) & \xrightarrow{\cdot x_1} & \frac{S}{(I^t,x_6)} & \longrightarrow & \frac{S}{((I^t,x_6),x_1)} & \longrightarrow & 0, \\
\text{nth}(S/I^t) = 2. & \square & \square & \square\n\end{array}
$$

we get depth $(S/I^t)$ 

Next, we consider the case where  $C_{\mathbf{w}}^n$  has two edges with non-trivial weights. First we consider the case where  $n = 3$ .

**Theorem 4.2.** If  $C_w^3$  has two edges with non-trivial weights, then

 $\operatorname{depth}(S/I(C_{\mathbf{w}}^3)^t) = 1$  *for all*  $t \geq 1$ .

*Proof.* Let  $I = I(C_{\bf w}^3)$ ,  ${\bf w}_1 \ge {\bf w}_3 \ge 2$  and  ${\bf w}_2 = 1$ . We prove the statements by induction on t with the case  $t = 1$  verified in Theorem [3.7.](#page-4-1) Thus, in the following, we can assume that  $t \geq 2$ .

From the proof of [\[25,](#page-29-7) Theorem 4.2], we see that  $(I^t : x_2 x_3) = I^{t-1}$ ,  $((I^t : x_2)$ ,  $x_3) =$  $(x_1^{w_1t}x_2^{w_1t-1}, x_3)$ , and  $(I^t, x_2) = ((x_1x_3)^{w_3t}, x_2)$ . By induction, we have depth $(S/((I^t :$  $(x_2), (x_3)$  = depth $(S/(I^t, x_2))$  = depth $(S/(I^t : x_2x_3)) = 1$ . Using Lemma [2.1](#page-2-1) to the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{I^t:x_2}(-1) & \xrightarrow{·x_2} & \frac{S}{I^t} & \longrightarrow & \frac{S}{(I^t,x_2)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{I^t:x_2x_3}(-1) & \xrightarrow{..x_3} & \frac{S}{I^t:x_2} & \longrightarrow & \frac{S}{((I^t:x_2),x_3)} & \longrightarrow & 0,\n\end{array}
$$

we can observe that  $\text{depth}(S/I^t) = 1$ .

<span id="page-7-0"></span>Next, we consider the case where  $n \geq 4$ .

**Theorem 4.3.** Let  $C_{\bf w}^n$  be an n-cycle with  $n \geq 4$ . If  $C_{\bf w}^n$  has two edges with non*trivial weights, then*

$$
\mathrm{depth}(S/I(C_{\mathbf{w}}^n)^t) \ge \max\{\lceil\frac{n-t}{3}\rceil, 1\} \text{ for all } t \ge 1.
$$

*Proof.* Let  $I = I(C_{\bf w}^n)$ . First, by the assumption and Lemma [3.2,](#page-3-1) we have  ${\bf w}_i = 1$  for all  $i \neq 1, 3$ . We prove the statements by induction on t with the case  $t = 1$  verified in Theorem [3.7.](#page-4-1) Thus, in the following, we can assume that  $t \geq 2$ .

By [\[27,](#page-29-14) Lemmas 4.1 and 4.3], it is easy to see that  $(I^t, x_3) = I(C_{\bf w}^n \backslash x_3)^t + (x_3)$ ,  $(I^t: x_3x_2) = I^{t-1}, ((I^t: x_3), x_2) = ((I^t, x_2): x_3), ((I^t, x_2), x_3) = I(P_{\mathbf{w}}^{n-2})^t + (x_2, x_3),$ where  $P_n^{n-2}$  is an induced subgraph of  $C_n^n$  on the set  $\{x_1, \hat{x_2}, \hat{x_3}, x_4, \dots, x_n\}$  and  $\hat{x_2}$ denotes the element  $x_2$  omitted from  $\{x_1, x_2, \ldots, x_n\}$ . By induction and Lemmas [2.2](#page-2-2) and [3.5,](#page-4-2) we have

$$
\text{depth}(S/(I^t, x_3)) \ge \max\{\lceil \frac{(n-1)-t+1}{3} \rceil, 1\} = \max\{\lceil \frac{n-t}{3} \rceil, 1\},\
$$
\n
$$
\text{depth}(S/((I^t : x_3) : x_2)) \ge \max\{\lceil \frac{n-(t-1)}{3} \rceil, 1\} = \max\{\lceil \frac{n-t+1}{3} \rceil, 1\},\
$$
\n
$$
\text{depth}(S/(I^t, x_2)) \ge \max\{\lceil \frac{(n-1)-t+1}{3} \rceil, 1\} = \max\{\lceil \frac{n-t}{3} \rceil, 1\},\
$$
\n
$$
\text{depth}(S/((I^t, x_2), x_3)) = \ge \max\{\lceil \frac{(n-2)-t+1}{3} \rceil, 1\} = \max\{\lceil \frac{n-t-1}{3} \rceil, 1\}.
$$

Using Lemma [2.1](#page-2-1) to the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{I^t:x_3}(-1) & \stackrel{\cdot x_3}{\longrightarrow} & \frac{S}{I^t} & \longrightarrow & \frac{S}{(I^t,x_3)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{I^t:x_3x_2}(-1) & \stackrel{\cdot x_2}{\longrightarrow} & \frac{S}{I^t:x_3} & \longrightarrow & \frac{S}{((I^t:x_3),x_2)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{((I^t:x_3),x_2)}(-1) & \stackrel{\cdot x_3}{\longrightarrow} & \frac{S}{I^t:x_2} & \longrightarrow & \frac{S}{((I^t,x_2),x_3)} & \longrightarrow & 0, \\
\text{oth}(S/I^t) \geq \{\max\left\lceil \frac{n-t}{3} \right\rceil, 1\}.\n\end{array}
$$

we get depth $(S/I^t) \geq {\max\left\lceil \frac{n-t}{3} \right\rceil}$ 

<span id="page-7-1"></span>For a monomial ideal  $I \subset S$ , let  $\mathcal{G}(I)$  denote the unique minimal set of its monomial generators.

**Theorem 4.4.** If  $C_w^4$  has two edges with non-trivial weights, then

$$
\operatorname{depth}(S/I(C_{\mathbf{w}}^4)^t) \le 1 \text{ for all } t \ge 2.
$$

*Proof.* Let  $I = I(C_{\bf w}^4), \mathbf{w}_1 \ge \mathbf{w}_3 \ge 2$  and  $\mathbf{w}_2 = \mathbf{w}_4 = 1$ . Choose  $f = x_1(x_3x_4)^{(t-1)\mathbf{w}_3}$ , then  $(I^t : f) = (x_1, x_2, x_4)$ . In fact,

$$
x_1 f = x_1^2 (x_3 x_4)^{(t-1)\mathbf{w}_3} = x_3^{\mathbf{w}_3} x_4^{\mathbf{w}_3 - 2} (x_3 x_4)^{(t-2)\mathbf{w}_3} (x_4 x_1)^2 \in I^t,
$$
  
\n
$$
x_2 f = x_1 x_2 (x_3 x_4)^{(t-1)\mathbf{w}_3} = (x_3 x_4)^{\mathbf{w}_3 - 1} (x_2 x_3) (x_3 x_4)^{(t-2)\mathbf{w}_3} (x_4 x_1) \in I^t,
$$
  
\n
$$
x_4 f = x_1 x_4 (x_3 x_4)^{(t-1)\mathbf{w}_3} = (x_3 x_4)^{(t-1)\mathbf{w}_3} (x_4 x_1) \in I^t.
$$

So  $(x_1, x_2, x_4) \subseteq (I^t : f)$ .

On the other hand, let  $u \in \mathcal{G}(I^t : f)$  be a monomial, then we can write  $uf$  as

<span id="page-8-0"></span>(3) 
$$
u(x_1(x_3x_4)^{(t-1)\mathbf{w}_3}) = (x_1x_2)^{k_1\mathbf{w}_1}(x_2x_3)^{k_2}(x_3x_4)^{k_3\mathbf{w}_3}(x_4x_1)^{k_4}h,
$$

where h is a monomial and  $\sum_{n=1}^4$  $i=1$  $k_i = t$  with  $k_i \geq 0$  for any  $i \in [4]$ . If  $k_3 = t$ , then Eq. [\(3\)](#page-8-0) becomes  $u(x_1(x_3x_4)^{(t-1)\mathbf{w}_3}) = (x_3x_4)^{t\mathbf{w}_3}h$ . It follows that  $u \in (x_4) \subseteq (x_1, x_2, x_4)$ . Now, let  $0 \leq k_3 \leq t-1$ . Consider the following two cases:

(1) If  $k_1 \geq 1$  or  $k_2 \geq 1$ , then  $u \in (x_2)$  by the choice of f and Eq. [\(3\)](#page-8-0).

(2) If  $k_1 = k_2 = 0$ , then Eq. [\(3\)](#page-8-0) becomes  $u(x_1(x_3x_4)^{(t-1)w_3}) = (x_3x_4)^{k_3w_3}(x_4x_1)^{k_4}h$ with  $k_3 + k_4 = t$ . Thus  $u \in (x_4) \subseteq (x_1, x_2, x_4)$  if  $k_3 = t - 1$ ; otherwise,  $u \in (x_1) \subseteq$  $(x_1, x_2, x_4)$ . Hence,  $(I^t : f) \subseteq (x_1, x_2, x_4)$ .

Therefore, depth $(S/I(\tilde{C}_{\bf w}^4)^t) \leq$  depth $(S/(I(\tilde{C}_{\bf w}^4)^t : f)) = 1$  by Lemma [2.3.](#page-2-3)

A direct consequence follows from Theorems [3.7,](#page-4-1) [4.3](#page-7-0) and [4.4.](#page-7-1)

**Corollary 4.5.** If  $C_w^4$  has two edges with non-trivial weights, then

$$
\operatorname{depth}(S/I(C_{\mathbf{w}}^4)^t) = 1 \text{ for all } t \ge 1.
$$

In the following, we use the notation  $u_i = (x_i x_{i+1})^{\mathbf{w}_i}$  where  $x_{n+1} = x_1$ , and  $\prod^j$  $\ell = i$  $u_{\ell} = 1$  if  $i > j$ .

<span id="page-8-1"></span>**Theorem 4.6.** Let  $C^n_{\bf w}$  be an n-cycle with  $n \geq 5$ . If  $C^n_{\bf w}$  has two edges with non*trivial weights. Let*  $t \geq 2$  *and choose* 

$$
f = \begin{cases} x_5(x_3x_4)^{\mathbf{w}_3} \prod_{k=5}^{t+2} (x_kx_{k+1}), & \text{if } 2 \le t \le n-3, \\ x_5(x_3x_4)^{(t-n+3)\mathbf{w}_3} \prod_{k=5}^n (x_kx_{k+1}), & \text{if } t \ge n-2. \end{cases}
$$

*Then*

$$
(I(C_{\mathbf{w}}^n) : f) = \begin{cases} (x_2) + \sum_{k=4}^{t+4} (x_k) + I(C_{\mathbf{w}}^n), & \text{if } 2 \le t \le n-3, \\ \sum_{j \in [n] \setminus \{3\}} (x_j), & \text{if } t \ge n-2. \end{cases}
$$

*Proof.* Let  $I = I(C_{\bf w}^n)$  and  ${\bf w}_1 \geq {\bf w}_3 \geq 2$  and  ${\bf w}_i = 1$  for all  $i \neq 1, 3$ . Then it is trivial that  $I(C_{\mathbf{w}}^n) \subseteq (I^t : f)$  by the choice of f, and  $x_2 \in (I^t : f)$  if  $2 \le t \le n-3$ , since  $x_2 f = (x_3 x_4)^{\mathbf{w}_3 - 1} u_2 \prod^{t+2}$  $k=4$  $u_k \in I^t$ . Now we consider the case where  $2 \le t \le n-3$ i

and 
$$
4 \le i \le t + 4
$$
. In this case, if *i* is even, then  $x_i f = u_3(\prod_{\ell=2}^{\frac{i}{2}-1} u_{2\ell+1}^2)(\prod_{\ell=i}^{t+2} u_{\ell}) \in I^t$ .  
Otherwise,  $x_i f = x_3^{w_3} x_4^{w_3-2}(\prod_{\ell=2}^{\frac{i-1}{2}} u_{2\ell}^2)(\prod_{\ell=i}^{t+2} u_{\ell}) \in I^t$ . Therefore,  $(x_2) + \sum_{k=4}^{t+4} (x_k) + I(C_{\mathbf{w}}^n) \subseteq (I^t : f)$ .

On the other hand, let  $u \in \mathcal{G}(I^t : f)$  be a monomial, then uf can be written as

<span id="page-9-0"></span>(4) 
$$
ux_5(x_3x_4)^{w_3}\prod_{k=5}^{t+2} (x_kx_{k+1}) = h \prod_{i=1}^{n} u_i^{k_i}
$$

for some monomial h, where  $\sum_{i=1}^{n} k_i = t$  and each  $k_i \geq 0$ . It is easy to see that if  $i=1$  $k_i \geq 1$  for some  $i \in \{1, 2, t + 3\}$  then  $u \in (x_2) + (x_{t+4})$ , and if  $k_i \geq 1$  for some  $\{t+4,\ldots,n\}$  then  $u \in (x_ix_{i+1}) \subseteq I(C_{\bf w}^n)$ . In the following, we assume that  $k_i = 0$ for each  $i \in \{1,2\} \cup \{t+3, t+4, \ldots, n\}$ . In this case, Eq. [\(4\)](#page-9-0) becomes

<span id="page-9-1"></span>(5) 
$$
ux_5(x_3x_4)^{w_3}\prod_{k=5}^{t+2} (x_kx_{k+1}) = h\prod_{i=3}^{t+2} u_i^{k_i},
$$

where  $\sum_{i=1}^{t+1}$  $i=3$  $k_i = t$ . We distinguish into three subcases: (a) If  $k_3 \geq 2$ , then  $u \in ((x_3x_4)^{w_3}) \subseteq I(C_{\mathbf{w}}^n)$ . (b) If  $k_3 = 1$ , then Eq. [\(5\)](#page-9-1) becomes  $ux_5$  $\prod^{t+2}$  $k=5$  $(x_kx_{k+1}) = h$  $\prod_{i=1}^{t+1}$  $i=4$  $u_i^{k_i}$  with  $\sum_{i=1}^{t+1}$  $i=4$  $k_i = t-1.$ When  $k_4 \geq 1$ ,  $u \in (x_4)$ . Otherwise, Eq. [\(5\)](#page-9-1) becomes  $ux_5$  $\prod_{i=1}^{t+1}$  $(x_kx_{k+1})=h$  $\prod_{i=1}^{t+1}$  $u_i^{k_i}$ 

with 
$$
\sum_{i=5}^{t+2} k_i = t - 1
$$
. Thus  
\n
$$
u \frac{x_5 \prod_{k=5}^{t+2} (x_k x_{k+1})}{\gcd(x_5 \prod_{k=5}^{t+2} (x_k x_{k+1}), \prod_{i=5}^{t+2} u_i^{k_i})} = h \frac{\prod_{i=5}^{t+2} u_i^{k_i}}{\gcd(x_5 \prod_{k=5}^{t+2} (x_k x_{k+1}), \prod_{i=5}^{t+2} u_i^{k_i})}.
$$
\n
$$
t+2
$$

Since the degrees of monomials  $x_5$  $\prod^{t+2}$  $k=5$  $(x_kx_{k+1})$  and  $\prod^{t+2}$  $i=5$  $u_i^{k_i}$  are  $2t - 3$  and  $2t - 4$ , respectively, we can deduce that  $u \in$  $\sum_{i=1}^{t+3}$  $i=5$  $(x_k)$  by comparing the degree of  $x_i$  for any  $i\in\{5,6,\ldots,t+3\}.$ 

(c) If  $k_3 = 0$ , then Eq. [\(5\)](#page-9-1) becomes  $ux_5(x_3x_4)^{w_3} \prod_{k=1}^{t+2}$  $k=5$  $(x_kx_{k+1}) = h$  $\prod_{i=1}^{t+1}$  $i=4$  $u_i^{k_i}$  with  $\sum_{i=1}^{t+1}$  $i=4$  $k_i = t$ . In this case, if  $k_4 \geq 3$ , then  $u \in (x_5)$  by comparing the degree of  $x_5$ ; otherwise,  $x_3^{\mathbf{w}_3} x_4^{\mathbf{w}_3-2}$  | h. It follows that  $ux_4^2x_5$  $\prod^{t+2}$  $k=5$  $(x_kx_{k+1}) = ($  $\prod^{t+2}$  $i=4$  $u_i^{k_i}\big) \frac{h}{x^{\mathbf{w}_3}x^{\mathbf{v}_4}}$  $\frac{h}{x_3^{\mathbf{w}_3}x_4^{\mathbf{w}_3-2}}$ . By similar arguments to (b), we can get that  $u \in$  $\sum_{i=1}^{t+3}$  $i=4$  $(x_k)$ . In breif,  $(I^t : f) \subseteq (x_2) +$  $\sum_{i=1}^{t+4}$  $k=4$  $(x_k) + I(C_{\mathbf{w}}^n).$ 

Now we consider the case where  $t \geq n-2$  and divide into the following four cases:

*Case* 1: If both *n* and *i* are odd, then  $x_i f \in I^t$ , since

$$
x_i f = \begin{cases} u_3^{t-n+3}(\prod_{\ell=2}^{\frac{n-1}{2}} u_{2\ell+1}^2), & \text{if } i = 1, \\ x_3^{\mathbf{w}_3} x_4^{\mathbf{w}_3-2} u_3^{t-n+2}(\prod_{\ell=2}^{\frac{i-1}{2}} u_{2\ell}^2)(\prod_{\ell=i}^n u_\ell), & \text{if } i \neq 1, 3. \end{cases}
$$

*Case* 2: If both *n* and *i* are even, then  $x_i f \in I^t$ , since

$$
x_i f = \begin{cases} (x_3 x_4)^{\mathbf{w}_3 - 1} u_2 u_3^{t - n + 2}(\prod_{\ell = 4}^n u_\ell), & \text{if } i = 2, \\ u_3^{t - n + 3}(\prod_{\ell = 2}^{\frac{i}{2} - 1} u_{2\ell+1}^2)(\prod_{\ell = i}^n u_\ell), & \text{if } i \neq 2. \end{cases}
$$

*Case* 3: If *n* is odd and *i* is even, then  $x_i f \in I^t$ , since

$$
x_i f = \begin{cases} (x_3 x_4)^{\mathbf{w}_3 - 1} u_2 u_3^{t - n + 2}(\prod_{\ell=4}^n u_\ell), & \text{if } i = 2, \\ u_3^{t - n + 3}(\prod_{\ell=2}^{\frac{i}{2} - 1} u_{2\ell}^2)(\prod_{\ell=i}^n u_\ell), & \text{if } i \neq 2. \end{cases}
$$

*Case* 4: If *n* is even and *i* is odd, then  $x_i f \in I^t$ , since

$$
x_i f = \begin{cases} x_3^{\mathbf{w}_3} x_4^{\mathbf{w}_3 - 2} u_3^{t - n + 2}(\prod_{\ell=2}^{\frac{n}{2}} u_{2\ell}^2), & \text{if } i = 1, \\ x_3^{\mathbf{w}_3} x_4^{\mathbf{w}_3 - 2} u_3^{t - n + 2}(\prod_{\ell=2}^{\frac{i-1}{2}} u_{2\ell}^2)(\prod_{\ell=i}^n u_\ell), & \text{if } i \neq 1, 3. \end{cases}
$$

In brief,  $\sum$  $i \in [n] \backslash \{3\}$  $(x_i) \subseteq (I^t : f).$ 

Next, we will prove that  $(I^t : f) \subseteq \sum$  $i \in [n] \backslash \{3\}$  $(x_i)$  by induction on t.

(a) If  $t = n - 2$ , then  $f = x_5(x_3x_4)^{w_3} \prod_{r=1}^{n}$  $k=5$  $(x_k x_{k+1})$ . Let  $u \in \mathcal{G}(I^t : f)$  be a monomial, then we can write  $uf$  as

<span id="page-10-0"></span>(6) 
$$
ux_5(x_3x_4)^{w_3}\prod_{k=5}^n(x_kx_{k+1})=h\prod_{i=1}^n u_i^{k_i}
$$

for some monomial h, where  $\sum_{n=1}^{\infty}$  $i=1$  $k_i = t$  and each  $k_i \geq 0$ .

If  $k_1 \geq 1$  or  $k_2 \geq 1$ , then  $u \in (x_2)$ . In the following, we will assume that  $k_1 = k_2 = 0$ . In this case, Eq. [\(6\)](#page-10-0) becomes

<span id="page-10-1"></span>(7) 
$$
ux_5(x_3x_4)^{w_3}\prod_{k=5}^n(x_kx_{k+1})=h\prod_{i=3}^n u_i^{k_i}
$$

with  $\sum_{n=1}^{\infty}$  $i=3$  $k_i = t$ . We consider the following three subcases: (i) If  $k_3 \geq 2$ , then  $u \in (x_4)$  by comparing the degree of  $x_4$  in Eq. [\(7\)](#page-10-1).

(ii) If  $k_3 = 1$ , then Eq. [\(7\)](#page-10-1) becomes  $ux_5 \prod_{ }^n$  $k=5$  $(x_k x_{k+1}) = h \prod_{k=1}^{n}$  $i=4$  $u_i^{k_i}$  with  $\sum_{i=4}^n$  $k_i = t-1.$ Further, if  $k_4 \geq 1$ , then  $u \in (x_4)$  by comparing the degree of  $x_4$ ; otherwise, Eq. [\(7\)](#page-10-1) becomes  $ux_5 \prod_{n=1}^n$  $k=5$  $(x_k x_{k+1}) = h \prod_{k=1}^{n}$  $i=5$  $u_i^{k_i}$  with  $\sum_{i=5}^n$  $k_i = t - 1$ . Similar arguments as in (1)(b), we can get  $u \in (x_1) + \sum_{n=1}^{\infty}$  $i=5$  $(x_k)$ .

(iii) If  $k_3 = 0$ , then Eq. [\(7\)](#page-10-1) becomes  $ux_5(x_3x_4)^{w_3} \prod_{r=1}^{n}$  $k=5$  $(x_k x_{k+1}) = h \prod_{k=1}^{n}$  $i=4$  $u_i^{k_i}$  with  $\sum_{n=1}^{\infty}$ 

 $i=4$  $k_i = t$ . Furthermore, if  $k_4 \geq 3$ , then  $u \in (x_5)$  by comparing the degree of  $x_5$ ; otherwise, by similar arguments as in (1)(c), we can see that  $u \in (x_1) + \sum_{k=1}^{n} (x_k)$ .

 $k=4$ (b) If  $t \geq n-1$ . Let  $u \in \mathcal{G}(I^t : f)$  be a monomial, then we can write uf as

 $ux_5(x_3x_4)^{(t-n+3)w_3} \prod_{r=1}^n$  $k=5$  $(x_k x_{k+1}) = h \prod^{n}$  $i=1$  $u_i^{k_i}$  for some monomial h, where  $\sum^n$  $i=1$  $k_i = t$ and each  $k_i \geq 0$ . We consider the following two subcases:

(i) If  $k_3 \geq 1$ , then  $u \frac{f}{(x_3x_4)^{w_3}} \in I^{t-1}$ . From induction it follows that  $(I^t : f) \subseteq$  $(I^{t-1}: \frac{f}{(x_3x_4)^{w_3}}) \subseteq \sum_{I \subset I^{(1)}_1}$  $i \in [n] \backslash \{3\}$  $(x_i).$ 

(ii) If  $k_3 = 0$ . We consider two cases: (i) if  $k_1 \geq 1$  or  $k_2 \geq 1$ , then  $u \in (x_2)$  by comparing the degree of  $x_2$ ; (ii) if  $k_1 = k_2 = 0$ , then  $u \in (x_1) + \sum_{n=1}^{\infty}$  $k=4$  $(x_k)$  by similar arguments as in  $(2)(a)(iii)$ . We complete the proof.

**Theorem 4.7.** Let  $C_{\bf w}^n$  be an n-cycle with  $n \geq 4$ . If  $C_{\bf w}^n$  has two edges with non*trivial weights, then* depth $(S/I(C_{\mathbf{w}}^n)^t) = \max\left\{ \left\lceil \frac{n-t}{3} \right\rceil, 1 \right\}$  *for all*  $t \geq 1$ *.* 

*Proof.* By Theorem [4.3,](#page-7-0) depth $(S/I(C_{\bf w}^n)^t) \geq {\max\left\lceil \frac{n-t}{3} \right\rceil}$  $\frac{-t}{3}$ , 1}. On the other hand, by Lemmas [2.3](#page-2-3) and [2.4,](#page-3-2) Theorems [4.4](#page-7-1) and [4.6,](#page-8-1) we can obtain that depth $(S/I(C_{\bf w}^n)^t) \leq$  $\{\max\left\lceil\frac{n-t}{3}\right\rceil\}$  $\frac{-t}{3}$ , 1}.

<span id="page-11-1"></span>4.2.  $C_{\bf w}^n$  has only one edge with non-trivial weights. In this section, we suppose  $\mathbf{w}_1 \geq 2$  and  $\mathbf{w}_i = 1$  for all  $i \neq 1$ . We need the following lemmas.

**Lemma 4.8.** ([\[17,](#page-29-13) Lemma 3.4]) Let  $P^n_{\mathbf{w}}$  be a trivially weighted path and H its sub*graph. Then* depth  $(S/(I(P_w^n)^t + I(H))) \geq \lceil \frac{n-t+1}{3} \rceil$  for any  $t < n$ .

<span id="page-11-0"></span>**Lemma 4.9.** ([\[28,](#page-29-12) Lemma 4.10]) *Let*  $P_{\mathbf{w}}^n$  *be a weighted path and*  $\mathbf{w}_{n-1} = 1$ *. Then for all*  $t \geq 2$ *, we have* 

<span id="page-11-2"></span>(1)  $(I(P_{\mathbf{w}}^{n})^{t} : x_{n-1}x_{n}) = I(P_{\mathbf{w}}^{n})^{t-1};$ (2)  $((I(P_w^n)^t : x_n), x_{n-1}) = (I(P_w^n \backslash x_{n-1})^t, x_{n-1})$ ; (3)  $(I(P_{\mathbf{w}}^{n})^{t}, x_{n}) = (I(P_{\mathbf{w}}^{n} \backslash x_{n})^{t}, x_{n});$ (4)  $(I(P_{\mathbf{w}}^n)^t, x_{n-1}) = (I(P_{\mathbf{w}}^n \backslash x_{n-1})^t, x_{n-1});$ (5)  $((I(P_w^n)^t : x_{n-1}), x_n) = ((I(P_w^n \setminus x_n)^t : x_{n-1}), x_n).$ 12

**Theorem 4.10.** Let  $P_{\bf w}^n$  be a path with only one edge of non-trivial weight. Let  $\mathbf{w}_i \geq 2$ ,  $\mathbf{w}_j = 1$  *for all*  $j \neq i$ , and let  $P_{\mathbf{w}}^{n-i+1}$  be the induced subgraph of  $P_{\mathbf{w}}^n$  on the set  $\{x_i, x_{i+1}, \ldots, x_n\}$  and  $H_{\mathbf{w}}$  be any subgraph of  $P_{\mathbf{w}}^{n-i+1}$ . Then for any  $1 \leq t < \lceil \frac{n+1}{2} \rceil$  $\frac{+1}{2}$ , *we have*

(1) depth 
$$
(S/(I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}}))) \ge \begin{cases} \lceil \frac{n-t+1}{3} \rceil, & \text{if } i = 1, \\ \lceil \frac{n-t}{3} \rceil, & \text{if } i \ge 2. \end{cases}
$$

(2) If  $i \geq 2$  and  $P_{\mathbf{w}}^i$  is the induced subgraph of  $P_{\mathbf{w}}^n$  on  $\{x_1, x_2, \ldots, x_i\}$ , then

depth 
$$
\left(S/(I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}}) + I(P_{\mathbf{w}}^i)\right) \ge \lceil \frac{n-t}{3} \rceil
$$
.

*Proof.* Let  $I = I(P_{\bf w}^n)$ . We first prove the statement (1) by induction on t. If  $t = 1$ , then  $I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}}) = I(P_{\mathbf{w}}^n)$ . By Lemma [3.4,](#page-4-0) we can get

$$
\text{depth}(S/I(P^n_{\mathbf{w}})) \ge \begin{cases} \lceil \frac{n}{3} \rceil, & \text{if $i = 1$,} \\ \lceil \frac{n-1}{3} \rceil, & \text{if $i \ge 2$.} \end{cases}
$$

In the following, we can assume that  $t \geq 2$ . Note that  $\sqrt{I(P_{\bf w}^n)^t + I(H_{\bf w})} = \sqrt{I(P_{\bf w}^n)}$ , we have  $\mathfrak{m} \notin Ass_S(I(P_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}))$  for all i, t and  $H_{\mathbf{w}}$ , where  $Ass_S(I(P_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}))$  $I(H_{\mathbf{w}})$  is the set of associated prime ideals of the S-module  $S/(I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}})).$ So depth $(S/(I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}}))) \geq 1 \geq \lceil \frac{n-t+1}{3} \rceil$  if  $n \leq 4$ .

Now we assume that  $n \geq 5$  and  $A = I(P_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}})$ . First, we consider the case where  $i = 1$  and divide into the following four cases:

*Case* 1: If  $x_{n-2}x_{n-1}, x_{n-1}x_n \notin E(H_{\mathbf{w}})$ , then, by Lemma [4.9,](#page-11-0) we see that

$$
(A, x_{n-1}) = I(P_{\mathbf{w}}^{n} \setminus x_{n-1})^{t} + I(H_{\mathbf{w}}) + (x_{n-1}),
$$
  
\n
$$
(A: x_{n-1}x_{n}) = I(P_{\mathbf{w}}^{n})^{t-1} + I(H_{\mathbf{w}}),
$$
  
\n
$$
((A: x_{n-1}), x_{n}) = (I(P_{\mathbf{w}}^{n} \setminus x_{n})^{t} : x_{n-1}) + I(H_{\mathbf{w}}) + (x_{n}),
$$
  
\n
$$
(((A: x_{n-1}), x_{n}), x_{n-2}) = I(P_{\mathbf{w}}^{n} \setminus \{x_{n-2}, x_{n}\})^{t} + I(H_{\mathbf{w}} \setminus x_{n-2}) + (x_{n-2}, x_{n}),
$$
  
\n
$$
(((A: x_{n-1}), x_{n}) : x_{n-2}) = \begin{cases} I(P_{\mathbf{w}}^{n} \setminus x_{n})^{t-1} + I(H_{\mathbf{w}}) + (x_{n}), \text{if } x_{n-3}x_{n-2} \notin E(H_{\mathbf{w}}), \\ B, \text{ otherwise,} \end{cases}
$$

where  $B = I(P_{\mathbf{w}}^n \setminus \{x_{n-3}, x_n\})^{t-1} + I(H_{\mathbf{w}} \setminus x_{n-3}) + (x_{n-3}, x_n)$ . Therefore, by induction and Lemma [2.2,](#page-2-2) we obtain that

$$
\text{depth}(S/(A, x_{n-1})) \ge \lceil \frac{(n-2)-t+1}{3} \rceil + 1 = \lceil \frac{n-t+2}{3} \rceil,
$$

$$
\text{depth}(S/(A: x_{n-1}x_n)) \ge \lceil \frac{n-(t-1)+1}{3} \rceil = \lceil \frac{n-t+2}{3} \rceil,
$$

$$
\text{depth}(S/(((A: x_{n-1}), x_n), x_{n-2})) \ge \lceil \frac{(n-3)-t+1}{3} \rceil + 1 = \lceil \frac{n-t+1}{3} \rceil.
$$

Furthermore, if  $x_{n-3}x_{n-2} \notin E(H_{\mathbf{w}})$ , then depth $(S/(((A : x_{n-1}), x_n) : x_{n-2})) \ge$  $\lceil \frac{(n-1)-(t-1)+1}{3} \rceil$  $\left\lfloor\frac{(t-1)+1}{3}\right\rceil = \left\lceil\frac{n-t+1}{3}\right\rceil$  $\frac{t+1}{3}$ ; otherwise, we let  $K_{\ell} = I(P_{\mathbf{w}}^n \setminus \{x_{n-3}, x_n\})^{\ell} + I(H_{\mathbf{w}} \setminus x_{n-3}) +$  $(x_{n-3}, x_n)$  for all  $\ell \in [t-1]$ , then  $K_1 = I(P_{\mathbf{w}}^n \setminus \{x_{n-3}, x_n\}) + I(H_{\mathbf{w}} \setminus x_{n-3}) + (x_{n-3}, x_n)$ ,  $(K_{\ell}, x_{n-2}x_{n-1}) = I(P_{\mathbf{w}}^{n} \setminus \{x_{n-3}, x_{n-2}, x_{n-1}, x_n\})^{\ell} + I(H_{\mathbf{w}} \setminus x_{n-3}) + (x_{n-3}, x_{n-2}x_{n-1}, x_n)$ 

for all  $\ell \in [t-1]$ ,  $(K_{\ell}: x_{n-2}x_{n-1}) = K_{\ell-1}$  for all  $2 \leq \ell \leq t-1$ , and  $K_{t-1} = (((A :$  $(x_{n-1}), x_n): x_{n-2}$ . By induction, Lemmas [2.2](#page-2-2) and [3.4,](#page-4-0) we get that depth $(S/K_1) \ge$  $\lceil \frac{n-4}{3} \rceil$  $\frac{-4}{3}$ ] + 1 =  $\frac{n-1}{3}$  $\frac{-1}{3}$  and depth $(S/(K_{\ell}, x_{n-2}x_{n-1})) \geq \lceil \frac{(n-4)-(t-1)+1}{3} \rceil + 1 = \lceil \frac{n-t+1}{3} \rceil$  $\frac{t+1}{3}$  for any  $2 \leq \ell \leq t-1$ . Applying Lemma [2.1](#page-2-1) to the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{K_{t-2}}(-2) & \frac{x_{n-2}x_{n-1}}{K_{t-1}} & \frac{S}{K_{t-1}} & \longrightarrow & \frac{S}{(K_{t-1}, x_{n-2}x_{n-1})} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{K_{t-3}}(-2) & \frac{x_{n-2}x_{n-1}}{K_{t-2}} & \frac{S}{K_{t-2}} & \longrightarrow & \frac{S}{(K_{t-2}, x_{n-2}x_{n-1})} & \longrightarrow & 0, \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
0 & \longrightarrow & \frac{S}{K_1}(-2) & \frac{x_{n-2}x_{n-1}}{K_2} & \frac{S}{K_2} & \longrightarrow & \frac{S}{(K_{2}, x_{n-2}x_{n-1})} & \longrightarrow & 0,\n\end{array}
$$

we deduce depth $(S/K_{t-1}) \geq \lceil \frac{n-t+1}{3} \rceil$ . In both cases, we always have depth $(S/K_{t-1}) \geq$  $\lceil \frac{n-t+1}{3} \rceil$  $\frac{t+1}{3}$ .

Using Lemma [2.1](#page-2-1) to the following short exact sequences

<span id="page-13-0"></span>
$$
\begin{array}{ccccccccc}\n & 0 & \longrightarrow & \frac{S}{A:x_{n-1}}(-1) & \xrightarrow{x_{n-1}} & \frac{S}{A} & \longrightarrow & \frac{S}{(A,x_{n-1})} & \longrightarrow & 0, \\
 & & \frac{S}{A:x_{n-1}x_n}(-1) & \xrightarrow{x_n} & \frac{S}{A:x_{n-1}} & \longrightarrow & \frac{S}{(A:x_{n-1}),x_n} & \longrightarrow & 0, \\
 & & & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{A:x_{n-1}} & \longrightarrow & \frac{S}{(A:x_{n-1}),x_n} & \longrightarrow & 0, \\
 & & & & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{(A:x_{n-1}),x_n} & \longrightarrow & 0, \\
 & & & & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{(A:x_{n-1})x_n} & \longrightarrow & \frac{S}{(A:x_{n-1})x_n} & \longrightarrow & 0, \\
 & & & & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{A:x_{n-1}x_n} & \longrightarrow & \frac{S}{(A:x_{n-1})x_n} & \longrightarrow & \frac{S}{(A:x
$$

$$
0 \longrightarrow \frac{S}{((A:x_{n-1}),x_n):x_{n-2}}(-1) \stackrel{x_{n-2}}{\longrightarrow} \frac{S}{((A:x_{n-1}),x_n)} \longrightarrow \frac{S}{(((A:x_{n-1}),x_n),x_{n-2})} \longrightarrow 0,
$$

we obtain that depth $(S/A) \geq \lceil \frac{n-t+1}{3} \rceil$ .

*Case* 2: If  $x_{n-2}x_{n-1}, x_{n-1}x_n \in E(H_{\mathbf{w}})$ , then by Lemma [4.9,](#page-11-0) we have

$$
(A, x_{n-1}) = I(P_{\mathbf{w}}^n \setminus x_{n-1})^t + I(H_{\mathbf{w}} \setminus x_{n-1}) + (x_{n-1}),
$$
  
\n
$$
(A: x_{n-1}) = I(P_{\mathbf{w}}^n \setminus \{x_{n-2}, x_n\})^t + I(H_{\mathbf{w}} \setminus \{x_{n-2}, x_n\}) + (x_{n-2}, x_n).
$$

By induction and Lemma [2.2,](#page-2-2) we get that

$$
\text{depth}(S/(A, x_{n-1})) \ge \lceil \frac{(n-2)-t+1}{3} \rceil + 1 = \lceil \frac{n-t+2}{3} \rceil,
$$
  

$$
\text{depth}(S/(A: x_{n-1})) \ge \lceil \frac{(n-3)-t+1}{3} \rceil + 1 = \lceil \frac{n-t+1}{3} \rceil.
$$

It follows from Lemma [2.1](#page-2-1) and the following exact sequence

$$
0 \longrightarrow \frac{S}{A:x_{n-1}}(-1) \stackrel{\cdot x_{n-1}}{\longrightarrow} \frac{S}{A} \longrightarrow \frac{S}{(A,x_{n-1})} \longrightarrow 0
$$

that depth $(S/A) \geq \lceil \frac{n-t+1}{3} \rceil$ . *Case* 3: If  $x_{n-2}x_{n-1} \in E(H_{\mathbf{w}})$  and  $x_{n-1}x_n \notin E(H_{\mathbf{w}})$ , then by Lemma [4.9,](#page-11-0) we have  $(A, x_{n-1}) = I(P_{\mathbf{w}}^n \backslash x_{n-1})^t + I(H_{\mathbf{w}} \backslash x_{n-1}) + (x_{n-1}),$  $(A: x_{n-1}x_n) = I(P_{\mathbf{w}}^n \setminus \{x_{n-2}\})^{t-1} + I(H_{\mathbf{w}} \setminus x_{n-2}) + (x_{n-2}),$ 

$$
((A: x_{n-1}), x_n) = (I(P_{\mathbf{w}}^n \setminus \{x_{n-2}, x_n\})^t) + I(H_{\mathbf{w}} \setminus x_{n-2}) + (x_{n-2}, x_n).
$$

Therefore, by induction and Lemma [2.2,](#page-2-2) we can deduce that

$$
\text{depth}(S/(A, x_{n-1})) \ge \lceil \frac{(n-2) - t + 1}{3} \rceil + 1 = \lceil \frac{n - t + 2}{3} \rceil,
$$
  

$$
\text{depth}(S/((A: x_{n-1}), x_n)) \ge \lceil \frac{(n-3) - t + 1}{3} \rceil + 1 = \lceil \frac{n - t + 1}{3} \rceil.
$$

Similarly to the calculation of the depth of  $S/K_{t-1}$  in *Case* 1, we obtain depth $(S/(A :$  $(x_{n-1}x_n)$ ) ≥  $\lceil \frac{n-t+1}{3} \rceil$ . Applying Lemma [2.1](#page-2-1) to the first two exact sequences of the sequences [\(8\)](#page-13-0), we can get depth $(S/A) \geq \lceil \frac{n-t+1}{3} \rceil$ .

*Case* 4: If  $x_{n-2}x_{n-1} \notin E(H_{\mathbf{w}})$  and  $x_{n-1}x_n \in E(H_{\mathbf{w}})$ , then by Lemma [4.9,](#page-11-0) we have

$$
(A, x_{n-1}) = I(P_{\mathbf{w}}^{n} \setminus x_{n-1})^{t} + I(H_{\mathbf{w}} \setminus x_{n-1}) + (x_{n-1}),
$$
  

$$
((A : x_{n-1}), x_{n-2}) = (I(P_{\mathbf{w}}^{n} \setminus \{x_{n-2}, x_{n}\})^{t}) + I(H_{\mathbf{w}} \setminus \{x_{n-2}, x_{n}\}) + (x_{n-2}, x_{n}),
$$
  

$$
(A : x_{n-1}x_{n-2}) = \begin{cases} I(P_{\mathbf{w}}^{n} \setminus x_{n})^{t-1} + I(H_{\mathbf{w}} \setminus x_{n}) + (x_{n}), & \text{if } x_{n-3}x_{n-2} \notin E(H_{\mathbf{w}}), \\ C, & \text{otherwise}, \end{cases}
$$

where  $C = I(P_{\mathbf{w}}^n \setminus \{x_{n-3}, x_n\})^{t-1} + I(H_{\mathbf{w}} \setminus \{x_{n-3}, x_n\}) + (x_{n-3}, x_n).$ Therefore, from induction and Lemma [2.2,](#page-2-2) we can deduce that

$$
depth(S/(A, x_{n-1})) \ge \lceil \frac{(n-2) - t + 1}{3} \rceil + 1 = \lceil \frac{n - t + 2}{3} \rceil
$$

$$
depth(S/((A:x_{n-1}), x_{n-2})) \ge \lceil \frac{(n-3)-t+1}{3} \rceil + 1 = \lceil \frac{n-t+1}{3} \rceil.
$$

⌉,

Similarly to the calculation of the depth of  $S/(((A:x_{n-1}),x_n):x_{n-2})$  in *Case* 1, we can get depth $(S/(A: x_{n-1}x_{n-2})) \geq \lceil \frac{n-t+1}{3} \rceil$ . Again, using Lemma [2.1](#page-2-1) to the first two exact sequences of the sequence [\(8\)](#page-13-0), we can get depth $(S/A) \geq \lceil \frac{n-t+1}{3} \rceil$ .

Now, we consider the case where  $i \geq 2$ . If  $n \leq 5$  and  $t \geq 2$ , or  $i = 3$ ,  $n \leq 6$  and  $t \geq 3$ , then  $\mathfrak{m} \notin Ass_S(I(P_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}))$  for all i, t and  $H_{\mathbf{w}}$ . So depth $(S/(I(P_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}})))$  $I(H_w)) \geq 1 \geq \lceil \frac{n-t}{3} \rceil$ . Therefore, in the following, we consider the four remaining cases: (I)  $i = 2, n \ge 6$  and  $t \ge 2$ ; (II)  $i = 3, n = 6$  and  $t = 2$ ; (III)  $i = 3, n \ge 7$  and  $t > 2$ ; (IV)  $i > 4$ ,  $n > 6$  and  $t > 2$ .

First, if the case (I) or (III) occurs, then, by similar arguments as in the case where  $i = 1$ , we can deduce depth $(S/A) \geq \lceil \frac{n-t}{3} \rceil$ .

If the case (II) occurs, then it remains to show that depth $(S/A) \geq 2$ . If  $x_4x_5 \in$  $E(H_{\mathbf{w}})$ , then depth $(S/A) \geq 2$  by similar arguments as in *Case* 2 and *Case* 3 for  $i = 1$ . Otherwise, there are two cases to consider:

(a) If  $x_5x_6 \in E(H_{\mathbf{w}})$ , then by Lemma [4.9,](#page-11-0) we have

$$
(A, x_5) = I(P_w^6 \setminus x_5)^2 + I(H_w \setminus x_5) + (x_5),
$$
  
\n
$$
(A: x_5x_4) = \begin{cases} I(P_w^6 \setminus x_6) + (x_6), & \text{if } x_3x_4 \notin E(H_w), \\ I(P_w^6 \setminus x_6) + (x_3^{\mathbf{w}} x_4^{\mathbf{w}-1}, x_6), & \text{if } x_3x_4 \in E(H_w), \end{cases}
$$
  
\n
$$
((A: x_5), x_4) = I(P_w^6 \setminus \{x_4, x_6\})^2 + (x_4, x_6).
$$

By the inductive hypothesis and Lemma [2.2,](#page-2-2) we get that depth $(S/((A : x_5), x_4)) \ge$ 2 and depth $(S/(A, x_5)) \geq 2$ . To compute depth of  $S/A$ , we need to compute depth $(S/(A: x_5x_4))$  and distinguish between two subcases: (i) if  $x_3x_4 \notin E(H_{\mathbf{w}})$ , then depth $(S/(A: x_5x_4)) \ge 2$  by Lemma [3.4\(](#page-4-0)3). (ii) If  $x_3x_4 \in E(H_{\mathbf{w}})$ , then  $(A:$  $x_5x_4x_3 = (x_2, (x_3x_4)^{\mathbf{w}-1}, x_4x_5, x_6)$  and  $((A : x_5x_4), x_3) = I(P_{\mathbf{w}}^6 \setminus {x_3, x_6}) + (x_3, x_6)$ by Lemma [4.9.](#page-11-0) It follows from Lemmas [2.2](#page-2-2) and [3.4](#page-4-0) that depth $(S/(A: x_5x_4x_3)) = 2$ and depth $(S/((A: x_5x_4), x_3)) = 2$ . Applying Lemma [2.1](#page-2-1) to the following exact

sequences

$$
(9) \quad 0 \longrightarrow \frac{S}{A:x_5x_4x_3}(-1) \xrightarrow{\cdot x_3} \frac{S}{A:x_5x_4} \longrightarrow \frac{S}{((A:x_5x_4),x_3)} \longrightarrow 0, \n0 \longrightarrow \frac{S}{A:x_5x_4}(-1) \xrightarrow{\cdot x_4} \frac{S}{A:x_5} \longrightarrow \frac{S}{((A:x_5),x_4)} \longrightarrow 0, \n0 \longrightarrow \frac{S}{A:x_5}(-1) \xrightarrow{\cdot x_5} \frac{S}{A} \longrightarrow \frac{S}{((A,x_5),x_4)} \longrightarrow 0,
$$

we deduce that depth $(S/A) \geq 2$ .

(b) If  $x_5x_6 \notin E(H_w)$ , then  $(A : x_5x_6) = I(P_w^6) + I(H_w)$  and  $(A, x_5x_6) = I(P_w^6)^2 + I(H_w)$  $I(H_w) + (x_5x_6)$  by Lemma [4.9.](#page-11-0) It follows from Lemma [3.4](#page-4-0) and the above (a) that  $depth(S/(A: x_5x_6)) \geq 2$  and  $depth(S/(A, x_5x_6)) \geq 2$ . This forces depth $(S/A) \geq 2$ by using Lemma [2.1](#page-2-1) to the following short exact sequence

$$
0 \longrightarrow \frac{S}{A:x_5x_6}(-2) \xrightarrow{\cdot x_5x_6} \frac{S}{A} \longrightarrow \frac{S}{(A,x_5x_6)} \longrightarrow 0.
$$

If the case (IV) occurs, then

$$
(A, x_2) = I(P_{\mathbf{w}}^n \backslash x_2)^t + I(H_{\mathbf{w}}) + (x_2),
$$
  
\n
$$
(A: x_2x_1) = I(P_{\mathbf{w}}^n)^{t-1} + I(H_{\mathbf{w}}),
$$
  
\n
$$
((A: x_2), x_1) = (I(P_{\mathbf{w}}^n \backslash x_1)^t : x_2) + I(H_{\mathbf{w}}) + (x_1),
$$
  
\n
$$
(((A: x_2), x_1) : x_3) = I(P_{\mathbf{w}}^n \backslash x_1)^{t-1} + I(H_{\mathbf{w}}) + (x_1),
$$
  
\n
$$
(((A: x_2), x_1), x_3) = I(P_{\mathbf{w}}^n \backslash \{x_1, x_3\})^t + I(H_{\mathbf{w}}) + (x_1, x_3).
$$

By induction and Lemma [2.2,](#page-2-2) we can deduce that

$$
\text{depth}(S/(A, x_2)) \ge \lceil \frac{(n-2) - t}{3} \rceil + 1 = \lceil \frac{n - t + 1}{3} \rceil,
$$
  

$$
\text{depth}(S/(A : x_2 x_1)) \ge \lceil \frac{n - (t - 1)}{3} \rceil = \lceil \frac{n - t + 1}{3} \rceil,
$$
  

$$
\text{depth}(S/(((A : x_2), x_1) : x_3)) \ge \lceil \frac{(n-1) - (t - 1)}{3} \rceil = \lceil \frac{n - t}{3} \rceil,
$$
  

$$
\text{depth}(S/(((A : x_2), x_1), x_3)) \ge \lceil \frac{(n-3) - t}{3} \rceil + 1 = \lceil \frac{n - t}{3} \rceil.
$$

By Lemma [2.1](#page-2-1) and the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{A:x_2}(-1) & \stackrel{\cdot x_2}{\longrightarrow} & \frac{S}{A} & \longrightarrow & \frac{S}{(A,x_2)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{A:x_2x_1}(-1) & \stackrel{\cdot x_1}{\longrightarrow} & \frac{S}{A:x_2} & \longrightarrow & \frac{S}{((A:x_2),x_1)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{((A:x_2),x_1):x_3}(-1) & \stackrel{\cdot x_3}{\longrightarrow} & \frac{S}{((A:x_2),x_1)} & \longrightarrow & \frac{S}{(((A:x_2),x_1),x_3)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{((A:x_2),x_1):x_3}(-1) & \stackrel{\cdot x_3}{\longrightarrow} & \frac{S}{((A:x_2),x_1)} & \longrightarrow & \frac{S}{(((A:x_2),x_1),x_3)} & \longrightarrow & 0,\n\end{array}
$$

we can get depth $(S/A) \geq \lceil \frac{n-t}{3} \rceil$ .

Final, we will prove the statement  $(2)$  by induction on t, its proof is similar as in the previous case.

If  $t = 1$ , then  $I(P_{\mathbf{w}}^n)^t + I(P_{\mathbf{w}}^i) + I(H_{\mathbf{w}}) = I(P_{\mathbf{w}}^n)$ . So depth $(S/I(P_{\mathbf{w}}^n)) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$  by Lemma [3.4,](#page-4-0) since  $i \geq 2$ .

Next, we assume that  $t \geq 2$ . Since  $\sqrt{I(P_{\mathbf{w}}^n)^t + I(P_{\mathbf{w}}^i) + I(H_{\mathbf{w}})} = \sqrt{I(P_{\mathbf{w}}^n)}$ , m  $\notin$  $Ass_S(I(P_{\mathbf{w}}^{n})^t + I(P_{\mathbf{w}}^{i}) + I(H_{\mathbf{w}}))$  for all i, t and  $H_{\mathbf{w}}$ . So depth $(S/(I(P_{\mathbf{w}}^{n})^t + I(P_{\mathbf{w}}^{i}) + I(H_{\mathbf{w}}^{n})))$  $I(H_{\mathbf{w}})) \geq 1 \geq \lceil \frac{n-t}{3} \rceil$  if  $n \leq 5$ .

In the following, we assume that  $n \geq 6$  and  $D = I(P_{\mathbf{w}}^{n})^t + I(P_{\mathbf{w}}^{i}) + I(H_{\mathbf{w}})$ . (i) If  $i = 2$ , then  $(D : x_1) = I(P_w^n \setminus x_2)^t + I(H_w \setminus x_2) + (x_2)$  and  $(D, x_1) =$  $I(P_{\mathbf{w}}^n \backslash x_1)^t + I(H_{\mathbf{w}}) + (x_1)$  by Lemma [4.9,](#page-11-0) it follows that depth $(S/(D : x_1)) \ge$  $\lceil \frac{(n-2)-t+1}{3} \rceil$  $\frac{(-t+1)}{3}$  + 1 =  $\frac{n-t+2}{3}$  $\frac{t+2}{3}$  and depth $(S/(D, x_1)) \geq \lceil \frac{(n-1)-t+1}{3} \rceil = \lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$  by Lemmas [2.2,](#page-2-2) [4.8](#page-11-1) and the statement (1). By Lemma [2.1](#page-2-1) and the exact sequence for  $s = 1$ 

<span id="page-16-0"></span>(10) 
$$
0 \longrightarrow \frac{S}{D:x_s}(-1) \stackrel{\cdot x_s}{\longrightarrow} \frac{S}{D} \longrightarrow \frac{S}{(D,x_s)} \longrightarrow 0,
$$

we can get depth $(S/D) \geq \lceil \frac{n-t}{3} \rceil$ .

(ii) If  $i \geq 3$ , then  $(D: x_2) = I(P_{\mathbf{w}}^n \setminus \{x_1, x_3\})^t + I(P_{\mathbf{w}}^i \setminus \{x_1, x_3\}) + I(H_{\mathbf{w}} \setminus x_3) + (x_1, x_3)$ and  $(D, x_2) = I(P_{\mathbf{w}}^n \backslash x_2)^t + I(P_{\mathbf{w}}^i \backslash x_2) + I(H_{\mathbf{w}}) + (x_2)$  by Lemma [4.9.](#page-11-0) By induction, Lemmas [2.2,](#page-2-2) [4.8](#page-11-1) and the statement (1), we can obtain that depth $(S/(D : x_2)) \geq$  $\lceil \frac{(n-3)-t}{3} \rceil$  $\frac{3}{3}$  +1 =  $\frac{n-t}{3}$  $\frac{-t}{3}$  and depth $(S/(D, x_2)) \geq \lceil \frac{(n-2)-t}{3} \rceil + 1 = \lceil \frac{n-t+1}{3} \rceil$  $\frac{t+1}{3}$ . Using Lemma [2.1](#page-2-1) to the short exact sequence [\(10\)](#page-16-0) for  $s = 2$ , we can get depth $(S/D) \geq \lceil \frac{n-t}{3} \rceil$ .

<span id="page-16-1"></span>**Lemma 4.11.** ([\[17,](#page-29-13) Lemma 3.10]) Let  $C_{\mathbf{w}}^n$  be a trivially weighted n-cycle and H its *non-empty subgraph. Then*

$$
\text{depth}\left(S/(I(C_{\mathbf{w}}^n)^t + I(H))\right) \ge \lceil \frac{n-t+1}{3} \rceil \text{ for all } t \ge 2.
$$

**Theorem 4.12.** Let n, t be two integers with  $n \geq 4$  and  $1 \leq t < \lceil \frac{n+1}{2} \rceil$  $\frac{+1}{2}$ . Let  $C_{\mathbf{w}}^n$ *be an* n*-cycle with only one edge of non-trivial weight and* H<sup>w</sup> *its subgraph with*  $V(H_{\mathbf{w}}) = \{x_1, \ldots, x_{2t-2}\}$  *and*  $E(H_{\mathbf{w}}) =$  $\begin{bmatrix}t-1\\1\end{bmatrix}$  $j=1$  ${x_{2j-1}}{x_{2j}}$ *. Assume that*  $H'_{\mathbf{w}}$  *is a nonempty subgraph of* Hw*. Then*

$$
\text{depth}(S/(I(C_{\mathbf{w}}^n)^t + I(H_{\mathbf{w}}^{\prime}))) \geq \lceil \frac{n-t}{3} \rceil.
$$

*Proof.* We prove the statement by induction on t. Let  $J = I(C_{\mathbf{w}}^{n})^{t} + I(H_{\mathbf{w}}^{'})$ . If  $t = 1$ , then  $J = I(C_{\mathbf{w}}^n)$ . By Theorem [3.7,](#page-4-1) depth $(S/J) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$ . Therefore, in the following, we can assume that  $t \geq 2$ . There are two cases:

*Case* 1: If  $E(H'_w) = \{x_1x_2\}$ , then  $J = I(C_w^n \backslash e_1)^t + (u_1)$ . In this case, let  $J_1 =$  $(J : x_2x_3)$  and  $J_2 = (J, x_2x_3)$ . First we compute depth $(S/J_1)$ . Note that  $(J_1, x_1)$  $I(C_{\bf w}^n \backslash x_1)^{t-1} + (x_1)$  by Lemma [4.9,](#page-11-0) so depth $(S/(J_1, x_1)) = \lceil \frac{(n-1)-(t-1)+1}{3} \rceil$  $\left\lfloor\frac{(t-1)+1}{3}\right\rceil = \left\lceil \frac{n-t+1}{3}\right\rceil$  $rac{t+1}{3}$ by Lemmas [2.2](#page-2-2) and [2.4.](#page-3-2) Again using Lemma [4.9,](#page-11-0) we can get

$$
(J_1: x_1) = \begin{cases} I(C_{\mathbf{w}}^n \setminus \{x_n, x_1\}) + ((x_1 x_2)^{\mathbf{w}_1 - 1}, x_n), & \text{if } t = 2, \\ (I(C_{\mathbf{w}}^n \setminus e_1)^{t-1} : x_1) + ((x_1 x_2)^{\mathbf{w}_1 - 1}), & \text{if } t \ge 3. \end{cases}
$$

Consequently, if  $t = 2$ , then  $\frac{\text{depth}(S/(J_1 : x_1)) \geq \lceil \frac{n-2}{3} \rceil}{\lceil \frac{n-2}{3} \rceil}$  by Lemmas [2.2,](#page-2-2) [2.4](#page-3-2) and [3.4.](#page-4-0) Applying Lemma [2.1](#page-2-1) to the short exact sequence

<span id="page-16-2"></span>(11) 
$$
0 \longrightarrow \frac{S}{J_1:x_1}(-1) \stackrel{\cdot x_1}{\longrightarrow} \frac{S}{J_1} \longrightarrow \frac{S}{(J_1,x_1)} \longrightarrow 0,
$$

we obtain depth $(S/J_1) \geq \lceil \frac{n-2}{3} \rceil$ .

If  $t \geq 3$ , then  $(J_1 : x_1x_n) = I(C_{\mathbf{w}}^n \setminus e_1)^{t-2} + ((x_1x_2)^{\mathbf{w}_1-1}), ((J_1 : x_1), x_n) =$  $I(C_{\bf w}^n\backslash\{\overline{x}_1,\overline{x}_n\})^{t-1} + ((x_1x_2)^{{\bf w}_1-1},x_n)$  by Lemma [4.9.](#page-11-0) By the induction and Lemmas [2.2,](#page-2-2) [4.8,](#page-11-1) [4.11](#page-16-1) and Theorem [4.10,](#page-11-2) we get that  $\frac{\text{depth}(S/(J_1: x_1x_n)) \geq \lceil \frac{n-(t-2)}{3} \rceil}{1 - \frac{1}{t}}$ 

 $\lceil \frac{n-t+2}{3} \rceil$  $\frac{t+2}{3}$  and depth $(S/((J_1: x_1), x_n)) \geq \lceil \frac{(n-1)-(t-1)+1}{3} \rceil = \lceil \frac{n-t+1}{3} \rceil$  $\frac{t+1}{3}$ . It follows from the exact sequence [\(11\)](#page-16-2) and

$$
0 \longrightarrow \frac{S}{J_1:x_1x_n}(-1) \stackrel{\cdot x_n}{\longrightarrow} \frac{S}{J_1:x_1} \longrightarrow \frac{S}{((J_1:x_1),x_n)} \longrightarrow 0
$$

that depth $(S/(J_1 : x_1)) \geq \lceil \frac{n-t}{3} \rceil$  and depth $(S/J_1) \geq \lceil \frac{n-t}{3} \rceil$ .

Next, we compute depth $(S/J_2)$ . Note that  $(J_2 : x_3x_4) = I(C_\mathbf{w}^n \backslash x_2)^{t-1} + (x_2),$  $(J_2, x_3) = I(C_{\mathbf{w}}^n \setminus \{x_2, x_3\})^t + (u_1, x_3)$  and  $((J_2 : x_3), x_4) = I(C_{\mathbf{w}}^n \setminus \{x_2, x_4\})^t + (x_2, x_4)$ by Lemma [4.9,](#page-11-0) we have that depth $(S/(J_2, x_3)) \geq \lceil \frac{n-t+1}{3} \rceil$ , depth $(S/(J_2 : x_3x_4))$  =  $\lceil \frac{(n-1)-(t-1)+1}{3} \rceil$  $\left\lfloor\frac{(t-1)+1}{3}\right\rfloor = \left\lceil \frac{n-t+1}{3} \right\rceil$  $\frac{t+1}{3}$  and depth $(S/((J_2: x_3), x_4)) = 1 + \lceil \frac{(n-3)-t+1}{3} \rceil$  $\left\lfloor \frac{n-t+1}{3} \right\rfloor = \left\lceil \frac{n-t+1}{3} \right\rceil$  $rac{t+1}{3}$ by Lemmas [2.2,](#page-2-2) [2.4](#page-3-2) and Theorem [4.10.](#page-11-2) Using the exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{J_2:x_3}(-1) & \xrightarrow{3} & \frac{S}{J_2} & \longrightarrow & \frac{S}{(J_2,x_3)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{J_2:x_3x_4}(-1) & \xrightarrow{3} & \frac{S}{J_2:x_3} & \longrightarrow & \frac{S}{((J_2:x_3),x_4)} & \longrightarrow & 0,\n\end{array}
$$

we can get depth $(S/J_2) \geq \lceil \frac{n-t}{3} \rceil$ .

As a result, using the short exact sequence

$$
0 \longrightarrow \frac{S}{J_1}(-2) \stackrel{\cdot x_2 x_3}{\longrightarrow} \frac{S}{J} \longrightarrow \frac{S}{J_2} \longrightarrow 0,
$$

we can obtain depth $(S/J) \geq \lceil \frac{n-t}{3} \rceil$ .

*Case* 2: If  $E(H'_{\mathbf{w}}) \neq \{x_1x_2\}$ , then we let  $m = \max\{i \in [t-2] \mid x_{2i+1}x_{2i+2} \in$  $E(H'_{\mathbf{w}})$  and  $K_j = J + \sum_{j=1}^{j}$  $s = 2m + 2$  $(x_sx_{s+1})$  for  $2m+2 \leq j \leq n$ . Thus  $(J : x_{2m+2}x_{2m+3}) =$  $I(C_{\mathbf{w}}^n \setminus x_{2m+1})^{t-1} + I(H_{\mathbf{w}}^{\prime} \setminus x_{2m+1}) + (x_{2m+1})$  and  $(K_j : x_{j+1}x_{j+2}) = I(C_{\mathbf{w}}^n \setminus x_j)^{t-1} +$  $I(H'_{\mathbf{w}} \backslash x_j) + (x_j)$  for  $2m+2 \leq j \leq n-1$ . By Theorem [4.10,](#page-11-2) we get that depth $(S/(J :$  $(x_{2m+2}x_{2m+3})\geq \lceil \frac{(n-1)-(t-1)}{3}\rceil = \lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$ , depth $(S/(K_j : x_{j+1}x_{j+2})) \geq \lceil \frac{(n-1)-(t-1)}{3} \rceil$  =  $\lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$  for  $2m + 2 \le j \le n - 1$ . It follows from the exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{J:x_{2m+2}x_{2m+3}}(-2) & \frac{x_{2m+2}x_{2m+3}}{J} & \frac{S}{J} & \longrightarrow & \frac{S}{K_{2m+2}} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{K_{2m+2}:x_{2m+3}x_{2m+4}}(-2) & \frac{x_{2m+3}x_{2m+4}}{J} & \frac{S}{K_{2m+2}} & \longrightarrow & \frac{S}{K_{2m+3}} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{K_{2m+3}:x_{2m+4}x_{2m+5}}(-2) & \frac{x_{2m+4}x_{2m+5}}{J} & \frac{S}{K_{2m+3}} & \longrightarrow & \frac{S}{K_{2m+4}} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{K_{n-1}:x_{n}x_{1}}(-2) & \frac{x_{n}x_{1}}{J} & \frac{S}{K_{n-1}} & \longrightarrow & \frac{S}{K_{n}} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{K_{n-1}:x_{n}x_{1}}(-2) & \frac{x_{n}x_{1}}{J} & \frac{S}{K_{n-1}} & \longrightarrow & \frac{S}{K_{n}} & \longrightarrow & 0, \\
\end{array}
$$

that depth $(S/J) \geq \lceil \frac{n-t}{3} \rceil$  if depth $(S/K_n) \geq \lceil \frac{n-t}{3} \rceil$ . It remains to prove that depth $(S/K_n) \geq \lceil \frac{n-t}{3} \rceil$ .

Let  $H^1_{\mathbf{w}}$  be a subgraph of  $C_{\mathbf{w}}^n$  with  $V(H^1_{\mathbf{w}}) = V(H'_{\mathbf{w}}) \cup \{x_{2m+3}, \ldots, x_n, x_1\}, E(H^1_{\mathbf{w}}) =$  $E(H'_{\mathbf{w}})\cup (\bigcup^n$  $i=2m+2$  ${x_i x_{i+1}}$ ,  $p_1 = \max\{j \in [n] \mid x_j x_{j+1} \notin E(H_w^1)\}\$ and  $\mathfrak{P}_1 =$  $I(C_{\mathbf{w}}^{n})^{t} + I(H_{\mathbf{w}}^{1})$ . For  $i \geq 2$ , let  $H_{\mathbf{w}}^{i}$  be a subgraph of  $C_{\mathbf{w}}^{n}$  with  $V(H_{\mathbf{w}}^{i}) = V(H_{\mathbf{w}}^{i-1}) \cup$  ${x_{p_{i-1}}, x_{p_{i-1}+1}}, E(H_w^i) = E(H_w^{i-1}) \cup {x_{p_{i-1}}x_{p_{i-1}+1}}, p_i = \max\{j \in [n] \mid x_jx_{j+1} \notin$  $E(H_w^i)$  and  $\mathfrak{P}_i = I(C_w^n)^t + I(H_w^i)$ . Since  $x_2x_3 \notin E(H_w^{'})$ ,  $p_i \geq 2$ . By the definition of  $p_i$  in above iterative process of  $H_w^i$ , we know that  $p_\ell = 2$  for some  $\ell \geq 2$ .

By Lemma [4.9,](#page-11-0) we obtain that for any  $i \in [\ell-1]$ ,

$$
(\mathfrak{P}_i : x_i x_{i+1}) = \begin{cases} I(C^n_{\mathbf{w}} \setminus x_{i+2})^{t-1} + I(G_{\mathbf{w}} \setminus x_{i+2}) + (x_{i+2}), & \text{if } x_{i-1} x_i \notin E(H^i_{\mathbf{w}}), \\ E, & \text{if } x_{i-1} x_i \in E(H^i_{\mathbf{w}}), \end{cases}
$$

where  $E = I(C_{\mathbf{w}}^n \setminus \{x_{i-1}, x_{i+2}\})^{t-1} + I(G_{\mathbf{w}} \setminus \{x_{i-1}, x_{i+2}\}) + (x_{i-1}, x_{i+2}).$ 

Similarly calculation of depth $(S/(((A : x_{n-1}), x_n) : x_{n-2}))$  as in *Case* 1 of the proof of Theorem [4.10\(](#page-11-2)1), we deduce  $\text{depth}(S/(\mathfrak{P}_i : x_i x_{i+1})) \geq \lceil \frac{n-t}{3} \rceil$  for any  $i \in [\ell-1]$ . To prove that  $\text{depth}(S/K_n) \geq \lceil \frac{n-t}{3} \rceil$ , by the following sequences

<span id="page-18-0"></span>
$$
(12) \quad \begin{array}{ccccccc}\n0 & \longrightarrow & \frac{S}{\mathfrak{P}_{1}:x_{p_{1}}x_{p_{1}+1}}(-2) & \xrightarrow{\cdot x_{p_{1}}x_{p_{1}+1}} & \frac{S}{K_{n}} & \longrightarrow & \frac{S}{\mathfrak{P}_{2}} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{\mathfrak{P}_{2}:x_{p_{2}}x_{p_{2}+1}}(-2) & \xrightarrow{\cdot x_{p_{2}}x_{p_{2}+1}} & \frac{S}{\mathfrak{P}_{2}} & \longrightarrow & \frac{S}{\mathfrak{P}_{3}} & \longrightarrow & 0, \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
0 & \longrightarrow & \frac{S}{\mathfrak{P}_{\ell-1}:x_{p_{\ell-1}}x_{p_{\ell-1}+1}}(-2) & \xrightarrow{\cdot x_{p_{\ell-1}}x_{p_{\ell-1}+1}} & \frac{S}{\mathfrak{P}_{\ell-1}} & \longrightarrow & \frac{S}{\mathfrak{P}_{\ell}} & \longrightarrow & 0,\n\end{array}
$$

we need to compute depth $(S/\mathfrak{P})$ . We consider the following two cases:

(a) If  $x_1x_2 \notin E(H'_{\mathbf{w}})$ , then  $(\mathfrak{P}_\ell : x_2x_3) = I(C_{\mathbf{w}}^n \setminus x_4)^{t-1} + I(H_{\mathbf{w}}^{\ell} \setminus x_4) + (x_4)$  and  $(\mathfrak{P}_{\ell}, x_2x_3) = I(C_{\mathbf{w}}^{n} \backslash e_1) + ((x_1x_2)^{t_{\mathbf{w}_1}})$  by Lemma [4.9.](#page-11-0) Thus depth $(S/(\mathfrak{P}_{\ell}: x_2x_3)) \ge$  $\lceil \frac{(n-1)-(t-1)}{3} \rceil$  $\frac{-(t-1)}{3}$  =  $\lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$ , depth $(S/(\mathfrak{P}_{\ell}, x_2x_3)) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$  by Lemma [3.7](#page-4-1) and Theorem [4.10.](#page-11-2) We can get  $\text{depth}(S/\mathfrak{P}_\ell) \geq \lceil \frac{n-t}{3} \rceil$  by the exact sequence

$$
0 \longrightarrow \frac{S}{\mathfrak{P}_{\ell} : x_2 x_3}(-2) \stackrel{x_2 x_3}{\longrightarrow} \frac{S}{\mathfrak{P}_{\ell}} \longrightarrow \frac{S}{(\mathfrak{P}_{\ell}, x_2 x_3)} \longrightarrow 0.
$$

(b) If  $x_1x_2 \in E(H'_\mathbf{w})$ , then  $\mathfrak{P}_\ell = I(C_\mathbf{w}^n \backslash e_2) + ((x_2x_3)^t)$ . It follows that  $(\mathfrak{P}_\ell, x_2x_3) =$  $I(C_{\bf w}^n), \; (\mathfrak{P}_\ell: x_2x_3x_1) = I(C_{\bf w}^n\setminus\{x_2,x_4,x_n\}) + ((x_1x_2)^{{\bf w}_1-1}, (x_2x_3)^{t-1}) + (x_4, x_n),$  $((\mathfrak{P}_{\ell}:x_2x_3),x_1) = I(C_{\mathbf{w}}^n \setminus \{x_1,x_2,x_4\}) + ((x_2x_3)^{t-1}) + (x_1,x_4)$  by Lemma [4.9.](#page-11-0) Thus  $\text{depth}(S/(\mathfrak{P}_{\ell}, x_2x_3)) = \lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$ , depth $(S/(\mathfrak{P}_{\ell} : x_2x_3x_1)) = \lceil \frac{n-5}{3} \rceil$  $\frac{-5}{3}$  + 1 =  $\frac{n-2}{3}$  $\frac{-2}{3}$ , depth $(S/((\mathfrak{P}_{\ell}: x_2x_3), x_1)) = \lceil \frac{n-4}{3} \rceil$  $\frac{-4}{3}$  + 1 =  $\frac{n-1}{3}$  $\frac{-1}{3}$  by Lemmas [2.2,](#page-2-2) [2.4,](#page-3-2) [3.4](#page-4-0) and Theorem [3.7.](#page-4-1) By Lemma [2.1](#page-2-1) and the following short exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{\mathfrak{P}_{\ell}:x_{2}x_{3}}(-2) & \stackrel{\cdot x_{2}x_{3}}{\longrightarrow} & \frac{S}{\mathfrak{P}_{\ell}} & \longrightarrow & \frac{S}{(\mathfrak{P}_{\ell},x_{2}x_{3})} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{\mathfrak{P}_{\ell}:x_{2}x_{3}x_{1}}(-1) & \stackrel{\cdot x_{1}}{\longrightarrow} & \frac{S}{\mathfrak{P}_{\ell}:x_{2}x_{3}} & \longrightarrow & \frac{S}{((\mathfrak{P}_{\ell}:x_{2}x_{3}),x_{1})} & \longrightarrow & 0,\n\end{array}
$$

we can get depth $(S/\mathfrak{P}_\ell) \geq \lceil \frac{n-t}{3} \rceil$ . Again using the exact sequences [\(12\)](#page-18-0), we obtain depth $(S/K_n) \geq \lceil \frac{n-t}{3} \rceil$  and finish the proof.

<span id="page-18-1"></span>**Theorem 4.13.** Let  $C_w^n$  be an n-cycle with  $n \geq 4$ , and  $\mathbf{w}_1 \geq 2$  and  $\mathbf{w}_i = 1$  for any  $i \neq 1$ . Let  $2 \leq t < \lceil \frac{n+1}{2} \rceil$  $\frac{+1}{2}$  be an integer and  $f = (x_1x_2)^{w_1}$   $\prod_{n=1}^{2t-2}$  $_{k=3}$ xk*, then*  $\operatorname{depth}(S/(I(C_{\mathbf{w}}^n)^t, f)) \geq \lceil \frac{n-t}{2} \rceil$ 3 ⌉.

*Proof.* Let  $f_j = (x_1 x_2)^{w_1} \prod^{2t-2j}$  $_{k=3}$  $x_k$  for each  $j \in [t-2]$ , and let  $H_w$  be any subgraph of  $C_{\mathbf{w}}^n$  whose edges are taken from the set  $\{e_{2t-2j+1}, e_{2t-2j+3}, \ldots, e_{2t-1}\}$ . then  $f_1 = f$ , 19

$$
f_j = x_{2t-2j-1}x_{2t-2j}f_{j+1} \text{ and } I(C_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}) + (f_j) = (I(C_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}) + (f_{j+1})) \cap (I(C_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}}) + (x_{2t-2j-1}x_{2t-2j})). \text{ The desired results follow from exact sequence}
$$
  
\n
$$
0 \longrightarrow \frac{S}{(A, f_j)} \longrightarrow \frac{S}{(A, f_{j+1})} \oplus \frac{S}{(A, x_{2t-1-2j}x_{2t-2j})} \longrightarrow \frac{S}{(A, f_{j+1}, x_{2t-1-2j}x_{2t-2j})} \longrightarrow 0,
$$
  
\nwhere  $A = I(C_{\mathbf{w}}^{n})^t + I(H_{\mathbf{w}})$  for any  $1 \le j \le t - 2$ .

<span id="page-19-2"></span>**Theorem 4.14.** Let t, f and  $C_w^n$  be as in Theorem [4.13.](#page-18-1) Then

$$
(I(C_{\mathbf{w}}^{n})^{t}: f) = I(C_{\mathbf{w}}^{n}) + (x_{n}^{2}) + (\bigcup_{i=0}^{t-2} \{x_{3}x_{2i+3}\}) + (\bigcup_{j=1}^{t-1} \{x_{2j+1}x_{n}\}) + (\bigcup_{p=1}^{t-2} \bigcup_{q=0}^{t-p-2} \{x_{2p}x_{2p+2q+3}\}).
$$

*Proof.* Let  $I = I(C_{\bf w}^n)$ . First we prove that the inclusion relation " $\supseteq$ " holds. It is easy to see that  $(I^t : f) \supseteq I$  and  $(x_n^2) \in (I^t : f)$ , since  $f = (x_1 x_2)^{w_1} \prod_{i=1}^{2t-2}$  $_{k=3}$  $x_k =$ 

 $u_1$  $\prod_{i=1}^{t-1} u_{2\ell+1} \in I^{t-1}$ , and  $x_n f = x_1^{w_1-2} x_2^{w_1} u_n^2$  $\ell = 1$   $\ell = 1$  $\prod_{i=1}^{t-2} u_{2\ell+1} \geq I^t$ . Now we divide into three cases: (i) If  $p \in [t-2]$  and  $q \in [t-p-2] \cup \{0\}$ , then  $x_{2p}x_{2p+2q+3}f =$  $u_1($ p−1<br>∏  $_{\ell=1}$  $u_{2\ell+1})$ (  $\prod^{p+q+1}$  $\ell = p$  $u_{2\ell})$ (  $t-2$ <br>∏  $\ell = p+q+1$  $u_{2\ell+1} \in I^t$ ; (ii) If  $i \in [t-2] \cup \{0\}$ , then  $x_3x_{2i+3}f =$  $x_1^{\mathbf{w}_1} x_2^{\mathbf{w}_1-2} u_2^2$ (  $\prod^{i+1}$  $_{\ell=2}$  $u_{2\ell})$ (  $t=2$ <br>∏  $\ell = i + 1$  $u_{2\ell+1} \in I^t$ ; (iii) If  $j \in [t-1]$ , then  $x_{2j+1}x_nf = (x_1x_2)^{\mathbf{w}_1-1}u_2$ .  $u_n(\prod^j)$  $_{\ell=2}$  $u_{2\ell})$ (  $t-2$ <br>∏  $_{\ell=j}$  $u_{2\ell+1} \in I^t$ . Therefore, the inclusion relation " $\supseteq$ " holds.

Next, we prove the inclusion relation " $\subseteq$ ". Let  $u \in \mathcal{G}(I^n : f)$  be a monomial, then  $uf = h \prod_{n=1}^{n}$  $i=1$  $u_i^{k_i}$  for some monomial h, where  $\sum^n$  $i=1$  $k_i = t$  and each  $k_i \geq 0$ . Thus

<span id="page-19-0"></span>(13) 
$$
u(x_1 x_2)^{w_1} \prod_{k=3}^{2t-2} x_k = h \prod_{i=1}^n u_i^{k_i}.
$$

Therefore, if  $k_i \geq 1$  for some  $2t - 1 \leq i \leq n - 1$ , then  $u \in (x_i x_{i+1}) \subseteq I(C_{\bf w}^n)$ , and if  $k_2 \geq 3$ , then  $u \in (x_3^2)$  by comparing the degree of  $x_3$  in Eq. [\(13\)](#page-19-0). If  $k_i \geq 2$  for some  $i \in \{3, 4, ..., 2t - 2\} \cup \{1, n\}$ , then  $u \in (x_n^2) + I(C_{\bf w}^n)$ . It remain to consider the cases:  $k_i = 0$  for all  $2t - 1 \le i \le n - 1$  and  $k_2 \le 2$  and  $k_i \le 1$  for all  $i \in \{3, 4, \ldots, 2t - 2\} \cup \{1, n\}.$  In this case, Eq. [\(13\)](#page-19-0) becomes

<span id="page-19-1"></span>(14) 
$$
u(x_1x_2)^{\mathbf{w}_1} \prod_{k=3}^{2t-2} x_k = h u_n^{k_n} \prod_{i=1}^{2t-2} u_i^{k_i},
$$

where  $k_n +$  $\sum_{i=1}^{2t-2}$  $i=1$  $k_i = t$ . We distinguish into two subcases:

(a) When 
$$
k_2 = 2
$$
, Eq. (14) becomes  $u(x_1x_2)^{w_1} \prod_{k=3}^{2t-2} x_k = hu_2^2 u_1^{k_1} u_n^{k_n} \prod_{i=3}^{2t-2} u_i^{k_i}$ . Furthermore, if  $k_1 = 1$ , then  $u \prod_{k=3}^{2t-2} x_k = hu_2^2 u_n^{k_n} \prod_{i=3}^{2t-2} u_i^{k_i}$ , which forces  $u \in (x_2x_3)$ . It

is easy to see that  $u \in (x_3x_{2t-1})$  if  $k_{2t-2} = 1$ , and  $u \in (x_3x_n)$  if  $k_n = 1$ . If  $k_1 = k_{2t-2} = k_n = 0$ , then Eq. [\(14\)](#page-19-1) becomes  $u(x_1x_2)^{\mathbf{w}_1} \prod^{2t-2}$  $_{k=3}$  $x_k = h$ <sup>2t−3</sup>  $i=2$  $u_i^{k_i}$ . In this case,  $k_3 = 1$  or  $k_{2i} = k_{2i+1} = 1$  for some  $i \in [t-2] \setminus \{1\}$ . In fact, if  $k_3 = 0$ , and  $k_{2i} = 0$  or  $k_{2i+1} = 0$  for any  $i \in [t-2] \setminus \{1\}$ , then  $\sum_{n=1}^{2t-3}$  $i=2$  $k_i \leq (2t-3)-(t-2)=t-1,$ a contradiction. By comparing the degrees of  $x_3$  and  $x_{2i+1}$  in the above expression, we get that  $u =$  $h \prod_{i=2}^{2t-3} u_i^{k_i}$  $(x_1x_2)^{\mathbf{w}_1} \prod_{k=3}^{2t-2} x_k$ ∈ (  $\begin{bmatrix}t-3\\1\end{bmatrix}$  $i=0$  ${x_3x_{2i+3}}$ .

(b) If  $0 \leq k_2 \leq 1$ , then there are three subcases:

(i) When  $k_n = 1$ , Eq. [\(14\)](#page-19-1) becomes  $u(x_1x_2)^{w_1} \prod^{2t-2}$  $_{k=3}$  $x_k = h u_n$  $\prod^{2t-2}$  $i=1$  $u_i^{k_i}$ . Furthermore, if  $k_1 = 1$  then  $u \in (x_n x_1) \subseteq I(C_{\mathbf{w}}^n)$ , and if  $k_{2t-2} = 1$  then  $u \in (x_{2t-1} x_n) \subseteq$ (  $\begin{bmatrix}t-1\\1\end{bmatrix}$  $j=1$  ${x_{2j+1}x_n}$ . If  $k_1 = k_{2t-2} = 0$ , then Eq. [\(14\)](#page-19-1) becomes  $u(x_1x_2)^{w_1}\prod_{i=1}^{2t-2}$  $_{k=3}$  $x_k =$  $hu_n$ <sup>2t−3</sup>  $i=2$  $u_i^{k_i}$ . In this case,  $k_{2i} = k_{2i+1} = 1$  for some  $i \in [t-2]$ . In fact, if  $k_{2i} = 0$  or  $k_{2i+1} = 0$  for any  $i \in [t-2]$ , then  $\sum_{i=1}^{2t-3}$  $i=2$  $k_i \le (2t-4) - (t-2) = t-2$ , a contradiction. By comparing the degrees of  $x_n$  and  $x_{2i+1}$  in the above expression, we get that  $u =$  $hu_n \prod_{i=2}^{2t-3} u_i^{k_i}$  $(x_1x_2)^{w_1} \prod^{2t-2} x_k$  $k=3$ ∈ (  $\begin{bmatrix} t-2 \\ 1 \end{bmatrix}$  $i=1$  ${x_{2i+1}x_n}.$ 

(ii) When  $k_n = 0$  and  $k_{2t-2} = 1$ , Eq. [\(14\)](#page-19-1) becomes  $u(x_1x_2)^{w_1} \prod^{2t-2}$  $_{k=3}$  $x_k = h u_{2t-2}$ <sup>2t−3</sup>  $i=1$  $u_i^{k_i}.$ By comparing the degrees of  $x_{2t-2}$  and  $x_{2t-1}$ , we can get that  $u \in (x_{2t-2}x_{2t-1}) \subseteq$  $I(C_{\mathbf{w}}^n)$  if  $k_{2t-3} = 1$ . If  $k_{2t-3} = 0$ , then Eq. [\(14\)](#page-19-1) becomes  $u(x_1x_2)^{\mathbf{w}_1} \prod_{i=1}^{2t-2}$  $_{k=3}$  $x_k =$  $hu_{2t-2}$  $\prod^{2t-4}$  $\frac{i=1}{i}$  $u_i^{k_i}$ . By arguments similar to the case (a) above, we deduce  $k_{2i-1} = k_{2i} = 1$ for some  $i \in [t-2]$ . Again comparing the degrees of  $x_{2i}$  and  $x_{2t-1}$  in the above expression, we can get that  $u \in$  $\begin{bmatrix} t-2 \\ 1 \end{bmatrix}$  $\frac{i=1}{i}$  $(x_{2i}x_{2t-1}) \subseteq ($  $\begin{bmatrix} t-2 \\ 1 \end{bmatrix}$  $p=1$  $\bigcup_{i=1}^{t-p-2}$  $q=0$  $\{x_{2p}x_{2p+2q+3}\}.$ (iii) When  $k_n = k_{2t-2} = 0$ , Eq. [\(14\)](#page-19-1) becomes

<span id="page-20-0"></span>(15) 
$$
u(x_1x_2)^{w_1} \prod_{k=3}^{2t-2} x_k = h \prod_{i=1}^{2t-3} u_i^{k_i}.
$$

Likewise,  $k_{2i-1} = k_{2i} = 1$  for some  $i \in [t-2]$ . Let  $a = \min\{i \in [t-2] \mid k_{2i-1} = k_{2i} = 1\}$ 1}. We divide into two subcases:

(a) If  $a = 1$ , then  $k_{2b} = k_{2b+1} = 1$  for some  $b \in [t-2]$ . Otherwise,  $\sum_{n=1}^{2t-3}$  $j=1$  $k_i \leq$  $(2t-3)-(t-2) = t-1$ , a contradiction. By comparing the degrees of  $x_2$  and  $x_{2b+1}$ in Eq. [\(15\)](#page-20-0), we can get  $u \in (x_2x_{2b+1}) \subseteq I(C_{\bf w}^n) + ($  $\begin{bmatrix} t-2 \\ 1 \end{bmatrix}$  $p=1$  $\bigcup_{i=1}^{t-p-1}$  $q=0$  ${x_{2p}x_{2p+2q+3}}$ .

(b) If  $2 \le a \le t-2$ , then  $k_{2i-1} = 0$  or  $k_{2i} = 0$  for any  $i \in [a-1]$ . So  $k_{2b} = k_{2b+1} = 1$ for some  $b \in [t-2] \setminus [a-1]$ . Otherwise,  $\sum_{n=1}^{2t-3}$  $j=1$  $k_i \leq (2t-3)-(a-1)-(t-a-1)=t-1,$ a contradiction. By comparing the degrees of  $x_{2a}$  and  $x_{2b+1}$  in Eq. [\(15\)](#page-20-0), we can get that  $u \in (x_{2a}x_{2b+1}) \subseteq I(C_{\bf w}^n) + ($  $\begin{bmatrix} t-2 \\ 1 \end{bmatrix}$  $p=1$  $\begin{bmatrix} t-p-2 \\ 1 \end{bmatrix}$  $q=0$  ${x_{2p}x_{2p+2q+3}}$ ).

<span id="page-21-1"></span>**Theorem 4.15.** Let t and  $C_w^n$  be as in Theorem [4.13,](#page-18-1) and let

$$
f = \begin{cases} x_3(x_1x_2)^{\mathbf{w}_1}, & \text{if } t = 2, \\ x_3(x_1x_2)^{\mathbf{w}_1} \prod_{k=3}^t (x_kx_{k+1}), & \text{if } 3 \le t < \lceil \frac{n+1}{2} \rceil, \end{cases}
$$

then  $(I(C_{\mathbf{w}}^n)^t : f) = I(C_{\mathbf{w}}^n) +$  $\sum_{i=1}^{t+1}$  $k=2$  $(x_k) + (x_n)$ .

*Proof.* Let  $I = I(C_{\bf w}^n)$ . If  $t = 2$ , then it is trival that  $I \subseteq (I^2 : f)$ . Since  $x_2 f =$  $(x_1x_2)^{\mathbf{w}_1}(x_2x_3) \in I^2$ ,  $x_3f = x_1^{\mathbf{w}_1}x_2^{\mathbf{w}_1-2}(x_2x_3)^2 \in I^2$ ,  $x_4f = (x_1x_2)^{\mathbf{w}_1}(x_3x_4) \in I^2$  and  $x_n f = (x_1 x_2)^{\mathbf{w}_1 - 1} (x_2 x_3)(x_n x_1) \in I^2$ , it follows that  $\sum_{n=1}^4$  $k=2$  $(x_k) + (x_n) \subseteq (I^2 : f).$ 

On the other hand, let  $u \in \mathcal{G}(I^2 : f)$  be a monomial, then  $uf = h \prod_{i=1}^{n}$  $i=1$  $u_i^{k_i}$  for some monomial h, where  $\sum_{n=1}^{\infty}$  $i=1$  $k_i = 2$  and each  $k_i \geq 0$ . Thus

<span id="page-21-0"></span>(16) 
$$
ux_3(x_1x_2)^{\mathbf{w}_1} = h \prod_{i=1}^n u_i^{k_i}.
$$

If  $k_i \geq 1$  for some  $i \geq 3$ , then  $u \in I(C_{\mathbf{w}}^n) + (x_4, x_n)$ . Next, we assume that  $k_i = 0$ for any  $i \geq 3$ . Then Eq. [\(16\)](#page-21-0) becomes  $ux_3(x_1x_2)^{\mathbf{w}_1} = h \prod_{i=1}^{2}$  $i=1$  $u_i^{k_i}$ . It is easy to see that  $u \in (x_2)$  if  $k_1 = 2$  and  $k_2 = 0$ , or  $k_1 = 1$  and  $k_2 = 1$ , and that  $u \in (x_3)$  if  $k_1 = 0$  and  $k_2 = 2.$ 

If  $3 \leq t < \left\lceil \frac{n+1}{2} \right\rceil$  $\frac{+1}{2}$ , then  $(I^t : f) = I(C_{\mathbf{w}}^n) +$  $\sum_{i=1}^{t+1}$  $k=2$  $(x_k) + (x_n)$  as similarly discussed in the proof of Theorem [4.6\(](#page-8-1)1).

**Theorem 4.16.** *Let*  $1 \le t < \frac{n+1}{2}$  $\frac{+1}{2}$  be an integer and  $C_{\mathbf{w}}^n$  as in Theorem [4.13,](#page-18-1) then  $\operatorname{depth}(S/I(C_{\mathbf{w}}^n)^t) = \lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$ .

*Proof.* Let  $I = I(C_{\bf w}^n)$ . If  $t = 1$ , then the desired result follows from Theorem [3.7.](#page-4-1) Now we assume that  $2 \leq t < \lceil \frac{n+1}{2} \rceil$  $\frac{+1}{2}$ . If  $t = 2$ , we choose  $f = x_3(x_1x_2)^{\mathbf{w}_1}$ , otherwise, we choose  $f = x_3(x_1x_2)^{\mathbf{w}_1} \prod_{i=1}^{t}$  $k=3$  $(x_kx_{k+1})$ . Thus, by Lemma [2.3](#page-2-3) and Theorem [4.15,](#page-21-1) depth $(S/I^t) \leq$  depth $(S/(I^t : f)) = 1 + \lceil \frac{(n-1)-(t+2)}{3} \rceil$  $\frac{-(t+2)}{3}$ ] =  $\lceil \frac{n-t}{3} \rceil$  $\frac{-t}{3}$ .

It remains to prove depth $(S/I^t) \geq \lceil \frac{n-t}{3} \rceil$ . Let  $g = (x_1 x_2)^{w_1} \prod_{i=1}^{2t-2}$  $_{k=3}$  $x_k$  and  $\mathfrak{Q} = (I^t :$ g). By the sequence [\(1\)](#page-3-4) in Lemma [2.3](#page-2-3) and Theorem [4.13,](#page-18-1) it suffices to prove that depth $(S/\mathfrak{Q}) \geq \lceil \frac{n-t}{3} \rceil$ . By Lemma [4.14,](#page-19-2) we deduce

$$
(\mathfrak{Q}:x_3) = [I + (x_n^2) + (\bigcup_{i=0}^{t-2} \{x_3x_{2i+3}\}) + (\bigcup_{j=1}^{t-1} \{x_{2j+1}x_n\}) + (\bigcup_{p=1}^{t-2} \bigcup_{q=0}^{t-p-2} \{x_{2p}x_{2p+2q+3}\})]: x_3
$$
  
=  $I(C_{\mathbf{w}}^n \setminus \{x_2, x_4\}) + (\bigcup_{i=0}^{t-2} \{x_{2i+3}\}) + (\bigcup_{p=1}^{t-2} \bigcup_{q=0}^{t-p-2} \{x_{2p}x_{2p+2q+3}\}) + (x_2, x_4, x_n)$   
=  $I(P_{\mathbf{w}}^{n-2t}) + (\bigcup_{i=0}^{t-2} \{x_{2i+3}\}) + (x_2, x_4, x_n)$ 

where  $P_{\mathbf{w}}^{n-2t}$  is an induced subgraph of  $C_{\mathbf{w}}^n$  on the set  $\{x_{2t}, \ldots, x_{n-1}\}$ , and

$$
((\mathfrak{Q}, x_3) : x_n) = [I + (x_3, x_n^2) + (\bigcup_{j=2}^{t-1} \{x_{2j+1}x_n\}) + (\bigcup_{p=1}^{t-2} \bigcup_{q=0}^{t-p-2} \{x_{2p}x_{2p+2q+3}\})] : x_n
$$
  

$$
= I(C_{\mathbf{w}}^n \setminus \{x_1, x_{n-1}\}) + (\bigcup_{j=2}^{t-1} \{x_{2j+1}\}) + (\bigcup_{p=1}^{t-2} \bigcup_{q=0}^{t-p-2} \{x_{2p}x_{2p+2q+3}\})
$$
  

$$
+ (x_1, x_3, x_{n-1}, x_n)
$$
  

$$
= I(P_{\mathbf{w}}^{n-2t-1}) + (\bigcup_{j=1}^{t-1} \{x_{2j+1}\}) + (x_1, x_{n-1}, x_n),
$$

where  $P_{\mathbf{w}}^{n-2t-1}$  is an induced subgraph of  $C_{\mathbf{w}}^n$  on the set  $\{x_{2t}, \ldots, x_{n-2}\}$ . There-fore, by Lemmas [2.2,](#page-2-2) [2.4](#page-3-2) and [3.4,](#page-4-0) we obtain that depth $(S/(\mathfrak{Q}:x_3)) \geq \lceil \frac{n-t}{3} \rceil$  and depth $(S/((\mathfrak{Q},x_3):x_n)) \geq \lceil \frac{n-t}{3} \rceil$ . We consider the following two cases:

(1) If  $t = 2$ , then  $(\mathfrak{Q}, x_3, x_n) = I(C_{\mathbf{w}}^n) + (x_3, x_n)$  by Lemma [4.14.](#page-19-2) Thus, we have depth $(S/(\mathfrak{Q},x_3,x_n)) = \lceil \frac{n-4}{3} \rceil$  $\frac{-4}{3}$ ] + 1 =  $\lceil \frac{n-1}{3} \rceil$  $\frac{-1}{3}$ . Using Lemma [2.1](#page-2-1) to the following exact sequences

<span id="page-22-0"></span>(17) 
$$
\begin{array}{ccccccccc}\n & 0 & \longrightarrow & \frac{S}{\mathfrak{Q}:\mathfrak{x}_3}(-1) & \xrightarrow{\cdot x_3} & \frac{S}{\mathfrak{Q}} & \longrightarrow & \frac{S}{(\mathfrak{Q},\mathfrak{x}_3)} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{(\mathfrak{Q},\mathfrak{x}_3):\mathfrak{x}_n}(-1) & \xrightarrow{\cdot x_n} & \frac{S}{(\mathfrak{Q},\mathfrak{x}_3)} & \longrightarrow & \frac{S}{(\mathfrak{Q},\mathfrak{x}_3,\mathfrak{x}_n)} & \longrightarrow & 0,\n\end{array}
$$

we can get depth $(S/\mathfrak{Q}) \geq \lceil \frac{n-t}{3} \rceil$ .

(2) If 
$$
3 \le t < \lceil \frac{n+1}{2} \rceil
$$
, then let  $M_j = (((\mathfrak{Q}, x_3, x_n) + (\bigcup_{s=j+1}^t \{x_{2s-1}\})) : x_{2j-1})$   
and  $H_j = (\mathfrak{Q}, x_3, x_n) + (\bigcup_{s=j}^t \{x_{2s-1}\})$  for any  $3 \le j \le t$ , where  $(\bigcup_{s=j}^t \{x_{2s-1}\}) = 0$  if

 $j = t$ . By Lemma [4.14,](#page-19-2) we get that  $H_3 = I(P_{\mathbf{w}}^{n-2t}) + \left(\bigcup_{i=1}^{t} I_{i}^{(i)}\right)^T$  $j=3$  ${x_{2j-1}}$ ) +  $(u_1, x_3, x_n)$  and  $M_j = I(P_{\mathbf{w}}^{n-2t}) + (\bigcup_{i=1}^t$  $s=j+1$  $\{x_{2s-1}\}\$  + ( j S−2  $m=1$  $\{x_{2m}\}\right) + (x_3, x_{2j-2}, x_{2j}, x_n)$  for any  $3 \leq j \leq t$ ,

where  $P_{\mathbf{w}}^{n-2t}$  is an induced subgraph of  $C_{\mathbf{w}}^n$  on  $\{x_{2t}, \ldots, x_{n-1}\}$ . Therefore, we can deduce that depth $(S/H_3) \geq \lceil \frac{n-t}{3} \rceil$  and depth $(S/M_j) \geq \lceil \frac{n-t}{3} \rceil$ . By the following exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{M_t}(-1) & \xrightarrow{x_{2t-1}} & \frac{S}{(\mathfrak{Q},x_3,x_n)} & \longrightarrow & \frac{S}{H_t} & \longrightarrow & 0, \\
0 & \longrightarrow & \frac{S}{M_{t-1}}(-1) & \xrightarrow{x_{2t-3}} & \frac{S}{H_t} & \longrightarrow & \frac{S}{H_{t-1}} & \longrightarrow & 0, \\
\vdots & & & \vdots & & \vdots \\
0 & \longrightarrow & \frac{S}{M_3}(-1) & \xrightarrow{x_5} & \frac{S}{M_4} & \longrightarrow & \frac{S}{H_3} & \longrightarrow & 0,\n\end{array}
$$

we obtain depth $(S/(\mathfrak{Q}, x_3, x_n)) \geq \lceil \frac{n-t}{3} \rceil$ . Again using the exact sequence [\(17\)](#page-22-0), we get that  $\text{depth}(S/\mathfrak{Q}) \geq \lceil \frac{n-t}{3} \rceil$ .

**Theorem 4.17.** Let  $C^n_{\bf w}$  be an *n*-cycle, where *n* is odd. Then

$$
depth(S/I(C_{\mathbf{w}}^n)^t) = 0 \text{ for all } t \ge \frac{n+1}{2}.
$$

*Proof.* Let  $I = I(C_{\bf w}^n)$  and  ${\bf m} = (x_1, x_2, \ldots, x_n)$ . It suffices to prove  ${\bf m} \in Ass_S(I^t)$ by Lemma [2.3.](#page-2-3) Choose  $f = (x_1 x_2)^{(t - \frac{n-1}{2})w_1} \prod_{r=1}^{n}$  $j=3$  $x_j$ , then  $f \notin I^t$  and  $(I^t : f) \subseteq \mathfrak{m}$ . On the other hand,  $x_1 f = u_1^{t - \frac{n-1}{2}}$  $\frac{n-1}{2}$  $_{k=1}$  $u_{2k+1} \in I^t$  and  $x_2 f = u_1^{t-\frac{n-1}{2}}$  $\frac{n-1}{2}$  $k=1$  $u_{2k} \in I^t$ . If *i* is

an integer with  $3 \leq i \leq n$ . We distinguish between two cases:

(a) If *i* is even, then 
$$
x_i f = u_1^{t-\frac{n-1}{2}} \left( \prod_{k=1}^{\frac{i}{2}-1} u_{2k+1} \right) \left( \prod_{m=\frac{i}{2}}^{\frac{n-1}{2}} u_{2m} \right) \in I^t
$$
.  
\n(b) If *i* is odd, then  $x_i f = (x_1 x_2)^{\mathbf{w}_1 - 1} u_1^{t-\frac{n-1}{2} - 1} \left( \prod_{k=1}^{\frac{i-1}{2}} u_{2k} \right) \left( \prod_{m=\frac{i-1}{2}}^{\frac{n-1}{2}} u_{2m+1} \right) \in I^t$ .

Therefore,  $(I^t : f) \supseteq \mathfrak{m}$ , which implies that  $(I^t : f) = \mathfrak{m}$  and so  $\mathfrak{m} \in Ass_S(I^t)$ . The desired result follows.

<span id="page-23-0"></span>**Theorem 4.18.** Let  $C_w^n$  be an n-cycle, where n is even, and  $w_1 \geq 2$  and  $w_i = 1$ for any  $i \neq 1$ *. Let*  $t \geq \frac{n}{2} + 1$  *be an integer and*  $f = (x_1 x_2)^{(t-\frac{n}{2})w_1} \prod_{i=1}^n$  $_{k=3}$ xk*, then*

$$
(I(C_{\mathbf{w}}^{n})^{t}: f) = (\bigcup_{k=1}^{\frac{n}{2}-1} \{x_{1}x_{2k}\}) + (\bigcup_{i=2}^{\frac{n}{2}-1} \bigcup_{j=1}^{\frac{n}{2}-1} \{x_{2i+1}x_{2j}\}) + (x_{3}, x_{n}).
$$
  
24

*Proof.* Let  $I = I(C_{\bf w}^n)$ . First note that  $x_3 f = x_1^{{\bf w}_1 - 1} x_2^{{\bf w}_1 - 2} u_1^{t - \frac{n}{2} - 1} u_2^2$  $\frac{\frac{n}{2}}{n}$  $_{\ell=2}$  $u_{2\ell} \in I^t$ and  $x_n f = x_1^{\mathbf{w}_1 - 2} x_2^{\mathbf{w}_1 - 1} u_1^{t - \frac{n}{2} - 1} u_n^2$  $\frac{n}{2}-1$  $_{\ell=1}$  $u_{2\ell} \in I^t$ . Furthermore, if  $k \in [\frac{n}{2} - 1]$ , then  $x_1x_{2k}f = u_1^{t-\frac{n}{2}}$  $k-1$   $\prod$  $\ell = 1$  $u_{2\ell+1}$  $\frac{\frac{n}{2}}{\prod}$  $_{\ell=k}$  $u_{2\ell} \in I^t$ , and if  $i \in \left[\frac{n}{2} - 1\right] \setminus \{1\}$  and  $j \in \left[\frac{n}{2} - 1\right]$ , then  $x_{2i+1}x_{2j}f =$  $\sqrt{ }$  $\int$  $\overline{\mathcal{L}}$  $u_1^{t-\frac{n}{2}}$ j−1<br>∏  $_{\ell=1}$  $u_{2\ell+1} \prod_{}^i$  $\ell = j$  $u_{2\ell}$  $\frac{n}{2}-1$  $\ell = i$  $u_{2\ell+1}$ , if  $i \geq j$ ,  $(x_1x_2)^{{\bf w}_1-1}u_1^{t-\frac{n}{2}-1}\prod^i$  $_{\ell=1}$  $u_{2\ell}$ j−1<br>∏  $\ell = i$  $u_{2\ell+1}$  $\frac{\frac{n}{2}}{\prod}$  $\ell = j$  $u_{2\ell}$ , if  $i < j$ .

This implies that  $x_{2i+1}x_{2j}f \in I^t$ . Therefore, the inclusion relation " $\supseteq$ " holds.

On the other hand, let  $u \in \mathcal{G}(I^n : f)$  be a monomial, then  $uf = h \prod_{i=1}^n$  $i=1$  $u_i^{k_i}$  for some monomial h, where  $\sum_{n=1}^{\infty}$  $i=1$  $k_i = t$  and each  $k_i \geq 0$ . Thus

<span id="page-24-0"></span>(18) 
$$
u(x_1x_2)^{(t-\frac{n}{2})\mathbf{w}_1} \prod_{k=3}^n x_k = h \prod_{i=1}^n u_i^{k_i}.
$$

Now we prove that the inclusion relation " $\subseteq$ " holds by induction on t. If  $t = \frac{n}{2} + 1$ , then Eq. [\(18\)](#page-24-0) becomes

<span id="page-24-1"></span>(19) 
$$
u(x_1 x_2)^{\mathbf{w}_1} \prod_{k=3}^n x_k = h \prod_{i=1}^n u_i^{k_i}.
$$

If  $k_2 \geq 2$  or  $k_n \geq 2$ , then  $u \in (x_3, x_n)$ . If  $k_i \geq 2$  for some  $i \in [n-1] \setminus \{2\}$ , then  $u \in (u_i) \subseteq ($  $\begin{bmatrix} \frac{n}{2} - 1 \\ 1 \end{bmatrix}$  $k=1$  ${x_1x_{2k}}$ ) + (  $\begin{bmatrix} \frac{n}{2} - 1 \\ 1 \end{bmatrix}$  $i=2$  $\begin{bmatrix} \frac{n}{2} - 1 \\ 1 \end{bmatrix}$  $j=1$  ${x_{2i+1}x_{2j}}$ ). In the following, we consider the case where  $0 \leq k_i \leq 1$  for any  $i \in [n]$  and divide into two cases:

(I) If  $k_1 = 1$ , then Eq. [\(19\)](#page-24-1) becomes u  $\prod_{n=1}^n$  $k=3$  $x_k = h \prod^n$  $i=2$  $u_i^{k_i}$  with  $\sum_{i=2}^n$  $k_i = t - 1$ . There are four subcases:

(a) If  $k_2 = k_n = 1$ , then  $u \in (x_1 x_2)$ .

(b) If  $k_2 = 1$  and  $k_n = 0$ , then Eq. [\(19\)](#page-24-1) becomes u  $\prod_{n=1}^{n}$  $k=4$  $x_k = hx_2 \prod^{n-1}$  $i=3$  $u_i^{k_i}$  with  $\sum_{n=1}^{n-1}$  $i=3$  $k_i = t - 2$ . Hence  $k_{2i} = k_{2i+1} = 1$  for some  $i \in [\frac{n}{2} - 1]$ . Indeed, if  $k_{2i} = 0$  or  $k_{2i+1} = 0$  for any  $i \in [\frac{n}{2} - 1]$ , then  $\sum_{n=1}^{n-1}$  $i=3$  $k_i \leq (n-3) - (\frac{n}{2} - 1) = \frac{n}{2} - 2 = t - 3$ , a contradiction. By comparing the degrees of  $x_2$  and  $x_{2i+1}$  in the above expression, we can get  $u \in ($  $\frac{n}{2}-1$  $i=1$  $\{x_2x_{2i+1}\}\subseteq ($  $\frac{n}{2}-1$  $i=2$  $\frac{n}{2}-1$  $j=1$  $\{x_{2i+1}x_{2j}\}\mathcal{)} + (x_3).$ 25

(c) If  $k_2 = 0$  and  $k_n = 1$ , then similar to the discussion of the case (b), we deduce that  $u \in (\bigcup^2)$  $\frac{i=1}{i}$  ${x_1x_{2i}}\) \subseteq (\bigcup^{n-1}$  $i=1$  ${x_1x_{2i}}$  +  $(x_n)$ . (d) If  $k_2 = k_n = 0$ , then Eq. [\(19\)](#page-24-1) becomes

<span id="page-25-0"></span> $u\prod_{i=1}^{n}$  $k=3$  $x_k = h \prod^{n-1}$  $i=3$ (20)  $u \prod x_k = h \prod u_i^{k_i},$ 

where  $\sum_{n=1}^{n-1}$  $i=3$  $k_i = t-1$ . Likewise, we can get that  $k_{2i-1} = k_{2i} = 1$  for some  $2 \le i \le \frac{n}{2} - 1$ . Let  $a_1 = \min\{2 \le i \le \frac{n}{2} - 1 \mid k_{2i-1} = k_{2i} = 1\}$ . We consider two subcases:

(i) If  $a_1 = 2$ , then  $k_{2b_1} = k_{2b_1+1} = 1$  for some  $2 \le b_1 \le \frac{n}{2} - 1$  by arguments similar to case (b) above. By comparing the degrees of  $x_{2a_1}$  and  $x_{2b_1+1}$  in Eq. [\(20\)](#page-25-0), we can get that  $u \in (x_{2a_1}x_{2b_1+1}) \subseteq ($  $\begin{array}{c}\frac{n}{2}-1\\1\end{array}$  $i=2$  $\begin{array}{c}\frac{n}{2}-1\\1\end{array}$  $j=1$  $\{x_{2i+1}x_{2j}\}.$ 

(ii) If  $3 \le a_1 \le \frac{n}{2} - 1$ , then by the definition of  $a_1$ , we have  $k_{2i-1} = 0$  or  $k_{2i} = 0$ for any  $2 \le i \le a_1 - 1$ , and  $k_{2b_2} = k_{2b_2+1} = 1$  for some  $b_2 \in [\frac{n}{2} - 1] \setminus [a_1 - 1]$ . Indeed, if  $k_{2b_2} = 0$  or  $k_{2b_2+1} = 0$ , then  $\sum^{n-1}$  $i=3$  $k_i \leq (n-3) - (a_1 - 2) - (\frac{n}{2} - a_1) = \frac{n}{2} - 1 = t - 2,$ a contradiction. By comparing the degrees of  $x_{2a_1}$  and  $x_{2b_2+1}$  in Eq. [\(20\)](#page-25-0), we can get that  $u \in (x_{2a_1}x_{2b_2+1}) \subseteq ($  $\begin{array}{c}\frac{n}{2}-1\\1\end{array}$  $i=2$  $\begin{array}{c}\n\frac{n}{2} - 1 \\
\end{array}$  $j=1$  $\{x_{2i+1}x_{2j}\}.$ (II) If  $k_1 = 0$ , then Eq. [\(19\)](#page-24-1) becomes

<span id="page-25-1"></span>(21) 
$$
u(x_1 x_2)^{\mathbf{w}_1} \prod_{k=3}^n x_k = h \prod_{i=2}^n u_i^{k_i},
$$

where  $\sum_{i=1}^{n} k_i = t$ . Likewise, we have  $k_{2i-1} = k_{2i} = 1$  for some  $2 \leq i \leq \frac{n}{2}$  $i=2$  $\frac{n}{2}$ . Let  $a_2 = \min\{2 \leq i \leq \frac{n}{2}\}\$  $\frac{n}{2}$  |  $k_{2i-1} = k_{2i} = 1$ }. we divide into two subcases:

(i) If  $a_2 = 2$ , then  $k_{2b_3} = k_{2b_3+1} = 1$  for some  $b_3 \in \left[\frac{n}{2} - 1\right]$  by arguments similar to case (b) above. By comparing the degrees of  $x_{2a_2}$  and  $x_{2b_3+1}$  in Eq. [\(21\)](#page-25-1), we can deduce that  $u \in (x_{2a_2}x_{2b_3+1}) \subseteq ($  $\frac{n}{2}-1$  $i=2$  $\begin{array}{c}\n\frac{n}{2} - 1 \\
\end{array}$  $j=1$  $\{x_{2i+1}x_{2j}\}\mathcal{)} + (x_3).$ 

(ii) If  $3 \leq a_2 \leq \frac{n}{2}$  $\frac{n}{2}$ , then by the definition of  $a_2$ , we have  $k_{2i-1} = 0$  or  $k_{2i} = 0$  for any  $2 \le i \le a_2 - 1$ , and  $k_{2b_4} = k_{2b_4+1} = 1$  for some  $b_4 \in [\frac{n}{2} - 1] \setminus [a_2 - 2]$ . Indeed, if  $k_{2b_4} = 0$  or  $k_{2b_4+1} = 0$ , then  $\sum_{i=2}^{n}$  $k_i \leq (n-1) - (a_2 - 2) - (\frac{n}{2} - a_2 + 1) = \frac{n}{2} = t - 1$ , a contradiction. By comparing the degrees of  $x_{2a_2}$  and  $x_{2b_4+1}$  in Eq. [\(21\)](#page-25-1), we can get that  $u \in (x_{2a_2}x_{2b_4+1}) \subseteq ($  $\frac{n}{2}-1$  $i=2$  $\frac{n}{2}-1$  $j=1$  $\{x_{2i+1}x_{2j}\}\mathcal{)} + (x_n).$ n

Therefore, if  $t = \frac{n}{2} + 1$ , then the desired conclusion holds. In the following, we assume that  $t \geq \frac{n}{2} + 2$ . We consider two cases:

(i) If 
$$
k_1 \ge 1
$$
, then  $(I^t : f) \subseteq (I^{t-1} : \frac{f}{(x_1x_2)^{w_1}}) \subseteq (\bigcup_{k=1}^{\frac{n}{2}-1} \{x_1x_{2k}\})+(\bigcup_{i=2}^{\frac{n}{2}-1} \bigcup_{j=1}^{\frac{n}{2}-1} \{x_{2i+1}x_{2j}\})+$ 

 $(x_3, x_n)$  by Eq. [\(18\)](#page-24-0) and induction.

(ii) If  $k_1 = 0$ , then Eq. [\(18\)](#page-24-0) becomes

<span id="page-26-0"></span>(22) 
$$
u(x_1x_2)^{(t-\frac{n}{2})\mathbf{w}_1} \prod_{k=3}^n x_k = h \prod_{i=2}^n u_i^{k_i},
$$

where  $\sum_{i=1}^{n} k_i = t$ . Likewise, we have  $k_{2i-1} = k_{2i} = 1$  for some  $2 \leq i \leq \frac{n}{2}$  $i=2$  $\frac{n}{2}$ . Let  $a_3 = \min\{2 \le i \le \frac{n}{2}\}$  $\frac{n}{2}$  |  $k_{2i-1} = k_{2i} = 1$ }. We divide into two subcases:

(i) If  $a_3 = 2$ , then  $k_{2b_5} = k_{2b_5+1} = 1$  for some  $b_5 \in \left[\frac{n}{2} - 1\right]$  by arguments similar to the case  $a_2 = 2$  above. By comparing the degrees of  $x_{2a_3}$  and  $x_{2b_5+1}$  in Eq. [\(22\)](#page-26-0), we can deduce that  $u \in (x_{2a_3}x_{2b_5+1}) \subseteq ($  $\begin{array}{c}\frac{n}{2}-1\\1\end{array}$  $i=2$  $\begin{array}{c}\n\frac{n}{2} - 1 \\
\end{array}$  $j=1$  $\{x_{2i+1}x_{2j}\}\mathcal{)} + (x_3).$ 

(ii) If  $3 \leq a_3 \leq \frac{n}{2}$  $\frac{n}{2}$ , then, by arguments similar to the case  $3 \le a_2 \le \frac{n}{2}$  $\frac{n}{2}$  above, we deduce that  $k_{2i-1} = 0$  or  $k_{2i} = 0$  for any  $2 \le i \le a_3 - 1$ , and  $k_{2b_6} = k_{2b_6+1} = 1$  for some  $b_6 \in \left[\frac{n}{2} - 1\right] \setminus [a_3 - 2]$ . By comparing the degrees of  $x_{2a_3}$  and  $x_{2b_6+1}$  in Eq. [\(22\)](#page-26-0), we can get that  $u \in (x_{2a_3}x_{2b_6+1}) \subseteq ($  $\frac{n}{2}-1$  $\frac{n}{2}-1$  ${x_{2i+1}x_{2j}}$  +  $(x_n)$ .

For a monomial  $u = x_1^{a_1} \cdots x_n^{a_n}$ , we set  $\deg_{x_i}(u) = a_i$  and  $\text{supp}(u) = \{x_i \mid a_i > 0\}.$ For a monomial ideal I, we set  $\text{supp}(I) = \overrightarrow{\bigcup}$  $u{\in}\mathcal{G}(I)$  $supp(u)$ .

 $j=1$ 

 $i=2$ 

**Theorem 4.19.** Let t and  $C^n_{\mathbf{w}}$  be as in Theorem [4.18.](#page-23-0) Then depth $(S/I(C^n_{\mathbf{w}})^t) = 1$ .

*Proof.* Let  $I = I(C_{\mathbf{w}}^n)$  and  $f = (x_1x_2)^{(t-\frac{n}{2})\mathbf{w}_1} \prod_{r=1}^n$  $k=3$  $x_k$ . Then by Theorem [4.18,](#page-23-0) we have

$$
(It: f) = (\bigcup_{k=1}^{\frac{n}{2}-1} \{x_1x_{2k}\}) + (\bigcup_{i=2}^{\frac{n}{2}-1} \bigcup_{j=1}^{\frac{n}{2}-1} \{x_{2i+1}x_{2j}\}) + (x_3, x_n)
$$
  
=  $(x_1, \hat{x}_3, x_5, \dots, x_{n-1})(x_2, x_4, \dots, x_{n-2}, \hat{x}_n) + (x_3, x_n).$ 

where  $\hat{x}_3$  denotes the element  $x_3$  omitted from  $\{x_1, x_3, x_5, \ldots, x_{n-1}\}.$  Thus

 $\operatorname{depth}(S/I^t) \leq \operatorname{depth}(S/(I^t : f)) = 1$ 

by Lemmas [2.2](#page-2-2) and [2.3.](#page-2-3)

Next, we will prove that  $\text{depth}(S/I^t) \geq 1$ . Let  $L^{(t)} = \{L_1^{(t)}\}$  $L_1^{(t)}, L_2^{(t)}, \ldots, L_r^{(t)}$ } be a specific totally ordered set of all elements in  $\mathcal{G}(I^t)$  as defined in [\[27,](#page-29-14) Setting 4.5]. For simplicity, we set  $L_i = L_i^{(1)}$  $i^{(1)}$ . Let  $C = \{L_i^{(t)}\}$  $\sum_{i}^{(t)}$  :  $L_1|^{edge}L_i^{(t)}$  $i^{(t)}$  for  $i \in [r]$  be a set of  $L_i^{(t)}$  divided by  $L_1$  as an edge and  $c = |C|$ . For each  $i \in [c]$ , we write  $L_i^{(t)}$  $i^{(t)}$  as  $L_i^{(t)} = L_{i_1}^{a_{i_1}}$  $\frac{a_{i_1}}{i_1}L_{i_2}^{a_{i_2}}$  $\stackrel{a_{i_2}}{_{i_2}}\cdots\stackrel{a_{i_{k_i}}}{L_{i_{k_i}}}$  $i_{i_{k_i}}^{k_i}$  with  $i_1 = 1$  and  $2 \leq i_2 < \cdots < i_{k_i} \leq n$ , where  $\sum_{i=1}^{k_i}$  $\sum_{j=1}^{n_t} a_{i_j} = t$  with  $a_{i_j} > 0$  for all  $j \in [k_i]$ . Assume that  $J_i = (L_{i+1}^{(t)}, \ldots, L_r^{(t)})$  for all  $i \in [c]$ . From the proof of [\[27,](#page-29-14) Theorem 4.11], we obtain  $J_c = I(P_w^n)^t$ , where  $I(P_{\mathbf{w}}^{n}) = (x_2x_3, x_3x_4, \ldots, x_{n-1}x_n, x_nx_1)$ . So

$$
\mathrm{depth}(S/J_c) = \mathrm{depth}(S/I(P_{\mathbf{w}}^n)^t) = \max\{\lceil \frac{n-t+1}{3}\rceil, 1\} \ge 1
$$

by Lemma [2.4.](#page-3-2) Let  $s_i$  be the degree of  $L_i^{(t)}$  $i<sup>i</sup>$  for each  $i \in [c]$ . By the exact sequences

$$
\begin{array}{ccccccccc}\n0 & \longrightarrow & \frac{S}{(J_1:L_1^{(t)})}(-s_1) & \xrightarrow{L_1^{(t)}} & \frac{S}{J_1} & \longrightarrow & \frac{S}{I^t} & \longrightarrow & 0 \\
0 & \longrightarrow & \frac{S}{(J_2:L_2^{(t)})}(-s_2) & \xrightarrow{L_2^{(t)}} & \frac{S}{J_2} & \longrightarrow & \frac{S}{J_1} & \longrightarrow & 0 \\
0 & \longrightarrow & \frac{S}{(J_3:L_3^{(t)})}(-s_3) & \xrightarrow{L_3^{(t)}} & \frac{S}{J_3} & \longrightarrow & \frac{S}{J_2} & \longrightarrow & 0 \\
& & \vdots & & \vdots & & \vdots \\
0 & \longrightarrow & \frac{S}{(J_c:L_c^{(t)})}(-s_c) & \xrightarrow{L_{c-1}^{(t)}} & \frac{S}{J_c} & \longrightarrow & \frac{S}{J_{c-1}} & \longrightarrow & 0,\n\end{array}
$$

we know that  $\text{depth}(S/I^t) \geq 1$  if  $\text{depth}(S/(J_i: L_i^{(t)}))$  $i_i^{(i)}$   $\geq$  2 for any  $i \in [c]$ . It remains to prove that  $\text{depth}(S/(J_i: L_i^{(t)}))$  $i^{(i)}$   $) \geq 2$  for any  $i \in [c]$ .

By [\[27,](#page-29-14) Theorems 4.10], we obtain that for each  $i \in [c]$ ,  $(J_i : L_i^{(t)})$  $\binom{v}{i} = K_i +$  $Q_i$ , where  $K_i = (L_4, L_5, \ldots, L_{n-2}) + \sum_{i=1}^{p_i}$  $\sum_{j=2}^{p_i} (x_{i_j+2}) + (x_3, x_n), Q_i = \sum_{j=0}^{q_i}$  $j=0$  $(x_{n-2j}), p_i =$  $\int k_i - 1$ , if  $i_{k_i} = n$ ,  $k_i$ , otherwise, and  $q_i = \max\{\ell_i : L_{n+1-2s} |^{edge} L_i^{(t)}\}$  $i^{(i)}$  for all  $0 \leq s \leq \ell_i$ .

If  $k_i = 1$ , or  $k_i = 2$  and  $i_{k_i} = n$ , then  $(J_1 : L_1^{(t)})$  $\binom{a}{1} = (L_4, L_5, \ldots, L_{n-2}) + (x_3, x_n).$ Thus depth $(S/(J_1: L_1^{(t)})$  $\binom{[t]}{1}$   $\geq$  2 by Lemma [2.4.](#page-3-2) In the following, we consider the case  $k_i \geq 3$ , or  $k_i = 2$  and  $i_{k_i} < n$ .

Note that if  $i_{k_i} = n$ , then  $p_i = k_i - 1$ , thus  $2 \leq i_{p_i} < i_{k_i} = n$  by the choice of  $k_i$ , and if  $i_{k_i} < n$ , then  $2 \le i_{p_i} = i_{k_i} < n$ . So  $2 \le i_j \le n-1$  for any  $2 \le j \le p_i$ , which forces  $i_j + 2 \in \{4, 5, \ldots, n\} \cup \{n+1\}$ . It follows that  $x_2 \notin \mathcal{G}(K_i)$ . Next we prove that  $x_2 \notin \mathcal{G}(Q_i)$ , which forces  $x_2 \notin \mathcal{G}(J_i : L_i^{(t)})$  $\binom{(\iota)}{i}$ .

Conversely, if  $x_2 \in \mathcal{G}(Q_i)$ , then  $q_i = \frac{n}{2} - 1$  by the expression of  $Q_i$ . By the definition of  $q_i$ ,  $L_{n+1-2s}$  |<sup>edge</sup>  $L_i^{(t)}$  $i_j^{(t)}$  for all  $0 \leq s \leq \frac{n}{2} - 1$ . So we can write  $L_i^{(t)}$  $i^{(t)}$  as  $L_i^{(t)} =$  $h_1$  $\frac{\frac{n}{2}}{n}$  $s=1$  $L_{n+1-2s}$  for some monomial  $h_1 \in \mathcal{G}(I^{t-\frac{n}{2}})$ . This implies  $h_1$  $\frac{\frac{n}{2}}{\prod}$  $_{k=1}$  $L_{2k} \in \mathcal{G}(I^t)$ . By the expression of  $L_i^{(t)}$  $i^{(t)}$ , we know that  $L_i^{(t)} = (x_1 x_2)^{\mathbf{w}_1 - 1} v$  with  $v = h_1$  $\frac{\frac{n}{2}}{\prod}$  $k=1$  $L_{2k}$ , which contradicts with  $L_i^{(t)} \in \mathcal{G}(I^t)$ .

By the definition of  $Q_i$  and  $K_i$ , we have  $(x_1x_2)^{\mathbf{w}_1}, x_2x_3 \notin \mathcal{G}(Q_i)$ ,  $(x_1x_2)^{\mathbf{w}_1}, x_2x_3 \notin$  $\mathcal{G}(K_i)$ , which means that  $x_2x_3, (x_1x_2)^{\mathbf{w}_1} \notin \mathcal{G}(J_i : L_i^{(t)})$  $\binom{t}{i}$ . Thus  $x_2 \notin \text{supp}(J_i : L_i^{(t)})$  $\binom{(\iota)}{i}$ for any  $i \in [c]$ . Therefore, depth $(S/(J_i : L_i^{(t)})$  $\binom{(t)}{i}$   $\geq 1$ . Note that  $(J_i : L_i^{(t)})$  $\binom{u}{i} =$   $(L_4, L_5, \ldots, L_{n-2}) + \sum_{i=1}^{p_i}$  $\sum_{j=2}^{p_i} (x_{i_j+2}) + \sum_{j=0}^{q_i}$  $j=0$  $(x_{n-2j}) + (x_3, x_n)$ , it follows from Lemma [2.4](#page-3-2) that depth $(S/(J_i: L_i^{(t)})$  $\binom{t}{i}$ ) = 1 if and only if  $(J_i: L_i^{(t)})$  $i^{(l)}_i$  =  $(x_1, \hat{x}_2, x_3, \ldots, x_n)$  for any  $i \in [c]$ . Therefore, it is remains to prove  $(J_i: L_i^{(t)})$  $\hat{u}^{(t)}(x) \neq (x_1, \hat{x}_2, x_3, \dots, x_n).$  In fact, if  $(J_i \,:\, L_i^{(t)}$  $\hat{z}^{(t)}$ ) =  $(x_1, \hat{x}_2, x_3, \ldots, x_n)$  for some  $i \in [c]$ , then  $x_{2k-1} \in (J_i : L_i^{(t)})$  $i^{(t)}$ ) for any  $k \in \left[\frac{n}{2}\right]$  $\frac{n}{2}$ . Since for any  $k \in \{1\} \cup \{3, 4, \ldots, \frac{n}{2}\}$  $\{\frac{n}{2}\}, x_{2k-1} \notin (L_4, L_5, \ldots, L_{n-2}) +$  $\sum_{i=1}^{q_i}$  $j=0$  $(x_{n-2j}) + (x_3, x_n)$ , we have  $x_{2k-1} \in \sum_{i=1}^{p_i}$  $\sum_{j=2}^{\infty} (x_{i_j+2})$ . Depending on the range of k, we see that  $\frac{n}{2}-1$  $k=1$  $\{2k+1\} \subseteq \bigcup^{p_i}$  $j=2$  $\{i_j\}$ . Since  $p_i \leq k_i$ ,  $L_{i_j}$  |<sup>edge</sup>  $L_i^{(t)}$  $i^{(i)}$  for all  $2 \leq j \leq p_i$  by the expression  $L_i^{(t)} = L_{i_1}^{a_{i_1}}$  $\frac{a_{i_1}}{i_1}L_{i_2}^{a_{i_2}}$  $\frac{a_{i_2}}{i_2}\cdots L_{i_{k_i}}^{a_{i_{k_i}}}$  $\sum_{i_{k_i}}^{a_{i_{k_i}}}$ . Thus  $L_{2k+1}$  |<sup>edge</sup>  $L_i^{(t)}$  $i^{(t)}$  for all  $k \in \left[\frac{n}{2} - 1\right]$ . Again by the definition of C, we get  $L_1 \mid^{edge} L_i^{(t)}$  $i^{(t)}$ . So  $L_i^{(t)}$ in by the definition of C, we get  $L_1 \mid^{edge} L_i^{(t)}$ . So  $L_i^{(t)}$  can be expressed as  $L_i^{(t)} = \frac{n}{2}$  $h_2$   $\prod^2$  $s=1$  $L_{2s+1}$  for some monomial  $h_2 \in \mathcal{G}(I^{t-\frac{n}{2}})$ . Note that  $L_i^{(t)}$  $i^{(l)}$  can also be written as  $L_i^{(t)} = (x_1 x_2)^{\mathbf{w}_1 - 1} v$  with  $v = h_2$  $\frac{\frac{n}{2}}{n}$  $k=1$  $L_{2k}$ , which contradicts with  $L_i^{(t)} \in \mathcal{G}(I^t)$ . Therefore, depth $(S/(J_i: L_i^{(t)})$  $\binom{t}{i}$   $\geq 2$  and we finish the proof.

### Acknowledgments

This research is supported by the Natural Science Foundation of Jiangsu Province (No. BK20221353) and the National NaturalScience Foundation of China (No. 12471246). The authors are grateful to the software systems [\[4\]](#page-28-8) and [\[10\]](#page-29-18) for providing us with a large number of examples.

### **REFERENCES**

- <span id="page-28-1"></span>[1] M. Brodmann, The asymptotic nature of the analytic spread, Math. Proc. Cambridge Philos Soc., 86 (1979), 35–39.
- <span id="page-28-7"></span><span id="page-28-5"></span>[2] W. Bruns, J. Herzog, *Cohen-Macaulay rings*, Revised Edition, Cambridge University Press, 1998.
- [3] M. Cimpoeas, On the Stanley depth of edge ideals of line and cyclic graphs, *Rom. J. Math. Comput. Sci.*, 5(2015), 70-75.
- <span id="page-28-8"></span>[4] CoCoATeam, CoCoA: a system for doing computations in Commutative Algebra, Avaible at [http://cocoa.dima.unige.it.](http://cocoa.dima.unige.it)
- <span id="page-28-4"></span>[5] L. T. K. Diem, N. C. Minh, and T. Vu, The sequentially Cohen-Macaulay property of edge ideals of edge-weighted graphs, [arXiv:2308.05020.](http://arxiv.org/abs/2308.05020)
- <span id="page-28-6"></span>[6] S. Y. Duan, G. J. Zhu, Y. J. Cui and J. X. Li, Integral closure and normality of edge ideals of some edge-weighted graphs, To appear in *J. Algebra Appl.*,<http://DOI:> 10.1142/S0219498826500921, [arXiv:2308.06016.](http://arxiv.org/abs/2308.06016)
- <span id="page-28-2"></span>[7] D. Eisenbud and C. Huneke, Cohen–Macaulay Rees algebras and their specializations, *J. Algebra*, 81, (1983), 202–224.
- <span id="page-28-3"></span>[8] S. A. Seyed Fakhari, K. Shibata, N. Terai and Siamak Yassemi, Cohen-Macaulay edgeweighted edge ideals of very well-covered graphs, *Comm. Algebra*, 49(10) (2021), 4249-4257.
- <span id="page-28-0"></span>[9] L. Fouli and S. Morey, A lower bound for depths of powers of edge ideals, *J. Algebr Comb.*, 42 (2015), 829-848.
- <span id="page-29-18"></span>[10] D. R. Grayson and M. E. Stillman, *Macaulay2*, a software system for research in algebraic geometry. Available at https://math.uiuc.edu/Macaulay2/.
- <span id="page-29-1"></span>[11] N. T. Hang, H. T. Truong and T. Vu, Depth of powers of edge ideals of Cohen-Macaulay trees, *Comm. Algebra* 52(12) (2024), 5049–5060.
- <span id="page-29-15"></span><span id="page-29-2"></span>[12] J. Herzog and T. Hibi, *Monomial ideals*, New York, NY, USA: Springer-Verlag, 2011.
- <span id="page-29-9"></span>[13] J. Herzog and T. Hibi, The depth of powers of an ideal, *J. Algebra*, 291, (2005), 534–550.
- <span id="page-29-3"></span>[14] T. T. Hien, Cohen-Macaulay edge-weighted graphs of girth 5 or greater, [arXiv:2309.05056.](http://arxiv.org/abs/2309.05056)
- [15] L. T. Hoa and N. D. Tam, On some invariants of a mixed product of ideals, *Arch. Math*, 94 (4) (2010), 327-337.
- <span id="page-29-13"></span><span id="page-29-10"></span>[16] J. X. Li, T. N. Trung and G. J. Zhu, Cohen-Macaulayness of powers of edge ideals of edgeweighted graphs, preprint.
- <span id="page-29-4"></span>[17] N. C. Minh, T. N. Trung and T. Vu, Depth of powers of edge ideals of cycles and starlike trees, To appear in *Rocky MT. J. Math.*, [arXiv:2308.00874.](http://arxiv.org/abs/2308.00874)
- <span id="page-29-5"></span>[18] S. Morey, Depths of powers of the edge ideal of a tree, Comm. Algebra, 38, (2010), 4042-4055.
- [19] S. Morey and R. H. Villarreal, *Edge ideals: algebraic and combinatorial properties*, in Progress in Commutative Algebra, Combinatorics and Homology, Vol. 1 (C. Francisco, L. C. Klingler, S. Sather-Wagstaff, and J. C. Vassilev, Eds.), De Gruyter, Berlin, 2012, 85-126.
- <span id="page-29-0"></span>[20] C. Paulsen and S. Sather-Wagstaff, Edge ideals of weighted graphs, *J. Algebra Appl.*, 12 (2013), 1250223-1-24.
- <span id="page-29-16"></span><span id="page-29-6"></span>[21] A. Rauf, Depth and sdepth of multigraded modules, *Comm. Algebra*, 38, (2010), 773–784.
- <span id="page-29-17"></span>[22] T. N. Trung, Stability of depths of powers of edge ideals, *J. Algebra* 452, (2016), 157–187.
- <span id="page-29-11"></span>[23] T. Vu and G. J. Zhu, Normal edge ideals of edge-weighted graphs, [arXiv:2407.16118.](http://arxiv.org/abs/2407.16118)
- <span id="page-29-7"></span>[24] S. Wei, Cohen-Macaulay weighted chordal graphs, [arXiv:2309.02414.](http://arxiv.org/abs/2309.02414)
- [25] G. J. Zhu, Projective dimension and regularity of the path ideal of the line graph, *J. Algebra Appl.*, 17(4), (2018), 1850068-1-15.
- <span id="page-29-8"></span>[26] G. J. Zhu, Projective dimension and regularity of path ideals of cycles, *J. Algebra Appl.*, 17(10) (2018), 1850188-1-22.
- <span id="page-29-14"></span><span id="page-29-12"></span>[27] G. J. Zhu, Y. J. Cui, J. X. Li and Y. Yang, Regularity of powers of edge ideals of edge-weighted integrally closed cycles, [arXiv:2407.05077.](http://arxiv.org/abs/2407.05077)
- [28] G. J. Zhu, S. Y. Duan, Y. J. Cui and J. X. Li, Edge ideals of some edge-weighted graphs, To appear in *Rocky MT J. Math.*, [arXiv:2401.02111.](http://arxiv.org/abs/2401.02111)

School of Mathematical Sciences, Soochow University, Suzhou, Jiangsu, 215006, P. R. China

*Email address*: zhuguangjun@suda.edu.cn(Corresponding author:Guangjun Zhu), 237546805@qq.com(Yijun Cui), lijiaxinworking@163.com(Jiaxin Li), 3201088194@qq.com(Yi Yang).