SINGULARITIES AND THEIR PROPAGATION IN OPTIMAL TRANSPORT

PIERMARCO CANNARSA, WEI CHENG, TIANQI SHI AND WENXUE WEI

ABSTRACT. In this paper, we investigate the singularities of potential energy functionals $\phi(\cdot)$ associated with semiconcave functions ϕ in the Borel probability measure space and their propagation properties. Our study covers two cases: when ϕ is a semiconcave function and when u is a weak KAM solution of the Hamilton-Jacobi equation H(x, Du(x)) = c[0] on a smooth closed manifold. By applying previous work on Hamilton-Jacobi equations in the Wasserstein space, we prove that the singularities of $u(\cdot)$ will propagate globally when u is a weak KAM solution, and the dynamical cost function C^t is the associated fundamental solution. We also demonstrate the existence of solutions evolving along the cut locus, governed by an irregular Lagrangian semiflow on the cut locus of u.

1. INTRODUCTION

In principle, the formation and evolution of singularities in optimal control, partial differential equations (PDEs), and geometry arise from the crossing and focusing of underlying characteristics or geodesics. Once formed, these singularities will propagate. This phenomenon reflects a certain irreversibility in the evolution of Hamiltonian systems. In a series of papers [2, 33, 3, 26, 27, 66, 79], the authors developed a theory to study the propagation of singularities using the concept of *generalized characteristics* (see also [40] for this concept in the context of hyperbolic conservation laws). Recently, various notions of generalized characteristics have been clarified, and some refined regularity properties have been established in [29], using ideas from abstract gradient flow theory. In this paper, we aim to study the problem of singularities propagation in optimal transport by applying the ideas from [29], along with the concept of singularities of functionals in probability measure space.

Over the past 30 years, the vibrant development of optimal transport theory has been driven by a series of geometric and dynamical perspectives. A pivotal advancement was made through the work of Gangbo and McCann ([55]), and later by McCann ([72]), who introduced the concept of displacement interpolation under a quadratic cost function in Euclidean space. Shortly thereafter, Benamou and Brenier reformulated these findings as an action-minimization problem within the space of measures ([13, 24]). Otto, in a series of pioneering papers ([76, 77, 78]), introduced the notion of gradient flows in Wasserstein space and explicitly formulated geodesic equations in a Riemannian geometric setting. Building on this foundation, Ambrosio, Gigli, and Savaré systematically developed the theory of gradient flows in metric spaces ([9]). At this stage, the close connection between Hamiltonian systems and optimal transport became evident. Bernard and Buffoni constructed displacement interpolation with cost functions determined by Tonelli Lagrangians,

²⁰²⁰ Mathematics Subject Classification. 35D40,49Q22, 37Jxx,37Kxx.

Key words and phrases. optimal transport, potential energy functional, cut locus, propagation of singularities, Hamilton-Jacobi equation.

establishing a clear link between optimal transport and Aubry-Mather theory in Hamiltonian dynamical systems ([15, 16, 17]). Additionally, Ambrosio, Gangbo, and Tudorascu, utilizing the geometry of the 2-Wasserstein space, considered Hamiltonians and flows within its phase space, deriving continuity equations, Euler-Poisson equations, and energy conservation results ([7, 53]). This work further contributed to the development of weak KAM theory in Wasserstein space by Gangbo and Tudorascu ([54, 59]).

With the continued development of calculus theory on Wasserstein and metric spaces, there has been growing interest in studying viscosity solutions to Hamilton-Jacobi equations on these spaces. Gangbo, Nguyen, and Tudorascu were the first to define viscosity solutions for a class of Hamilton-Jacobi equations in Wasserstein space using subdifferentials and superdifferentials, and to investigate their well-posedness ([56, 60]). Additionally, using the concept of metric derivatives, Ambrosio and Feng, Gangbo and Święch, as well as other researchers, have explored the well-posedness of viscosity solutions to Hamilton-Jacobi equations in general metric spaces ([6, 58, 57, 61]). Building on these foundational results, some related dynamical aspects, such as the asymptotic behavior of viscosity solutions and Hamilton-Jacobi equations with cohomological terms, have been examined ([74, 18]). In the areas of calculus of variations and optimal (or mean-field) control, we refer to [34, 20, 19, 21, 11, 64, 63, 39, 49], where some works provide variational representations of viscosity solutions.

The study of the geometric aspects of optimal transport, with an emphasis on the cut locus, has been the subject of numerous previous works. Under the assumption that the cost function is the square of the distance, the existence of the optimal transport map from an absolutely continuous (with respect to Lebesgue measure) measure was obtained in [23] in Euclidean case and [73] in Riemannian case. The first-order properties of the cut locus, such as Lipschitz continuity and (n - 1)-rectifiability, have been investigated in [35, 65, 69, 71]. The authors in [70] study the second-order characterization of the cut locus acting as a barrier to the regularity of optimal transport maps, including the semiconvexity of the injectivity domain. By virtue of the Ma-Trudinger-Wang (MTW) condition, for a Riemannian manifold M with a nonfocal cut locus satisfying the strong MTW condition, the following results hold:

- For any two C^{∞} positive probability densities f and g on M, the optimal transport map from $\mu(dx) = f(x) \operatorname{vol}(dx)$ to $\nu(dy) = g(y) \operatorname{vol}(dy)$ with cost function $c = d^2$ is C^{∞} ;
- There exists a $\kappa > 0$ such that all injectivity domains are κ -uniformly convex.

Furthermore, by employing a slightly modified version of the MTW condition, a more delicate characterization of the injectivity domains and the regularity of optimal transport maps is obtained (see [70, Theorem 1.8]). If (M, g) is a surface, the authors in [50] prove that the injectivity domains of M are semiconvex if M satisfies an appropriate signed curvature condition. In [52], the authors demonstrated that any C^4 -deformation of the round sphere (\mathbb{S}^2, g^{can}) satisfies the transport continuity property. In their subsequent work [51], they further showed that if (M, g) is a C^4 -deformation of the round sphere (\mathbb{S}^n, g^{can}), then all injectivity domains of M are uniformly convex. For the verification of MTW condition on Riemannian surfaces, see also [42].

To address singularities and cut locus of functionals on Wasserstein spaces, we follow the framework in [21]. For a metric space (X, d), we denote by $\mathscr{P}(X)$ the set of all probability Borel measures on X, and by $\mathscr{P}_c(\mathbb{R}^m)$ the subspace of $\mathscr{P}(\mathbb{R}^m)$ consisting of measures with compact

support. For any functional $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$, we introduce $\partial^{\pm} U(\mu)$, the localized Fréchet superdifferential and subdifferential of U at μ (see Definition 3.1). A measure $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ is called a regular point of U if U is locally differentiable at μ , i.e., both $\partial^+ U(\mu)$ and $\partial^- U(\mu)$ are non-empty. A measure $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ is called a singular point of U if U is not locally differentiable at μ . We denote by $\mathscr{S}(U)$ the set of all singular points of U. We also introduce the notions of semiconcavity for a functional $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ (see Definition 3.3).

We concentrate to the potential energy functional (see [9]) induced by a function $\phi : \mathbb{R}^m \to \mathbb{R}$

$$\Phi(\mu) := \int_{\mathbb{R}^m} \phi(x) \, d\mu, \quad \forall \mu \in \mathscr{P}(\mathbb{R}^m).$$

We denote $\Phi(\cdot)$ by $\phi(\cdot)$ for brevity. That is ϕ is real-valued function on \mathbb{R}^m and $\phi(\cdot)$ is also a functional on $\mathscr{P}(\mathbb{R}^m)$ if there is no confusion. If ϕ is a locally semiconcave function on \mathbb{R}^m , its associated potential energy functional $\phi(\cdot)$ has the following properties:

- (1) If ϕ is locally semiconcave, then potential energy functional $\phi(\cdot)$ is locally strongly semiconcave. (Proposition 3.3)
- (2) If ϕ is locally semiconcave and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, then $\alpha \in \partial^+ \phi(\mu)$ if and only if $\alpha(x) \in D^+ \phi(x)$, the superdifferential of ϕ at x, holds true for μ -a.e. $x \in \mathbb{R}^m$. (Theorem 3.1) Moreover, $\partial^+ \phi(\mu) \neq \emptyset$ and $\phi(\cdot)$ is locally differentiable at μ if and only if $\partial^+ \phi(\mu)$ is a singleton, and $\alpha = \partial \phi(\mu)$ if and only if $\alpha(x) = D\phi(x)$ holds for μ -a.e. $x \in \mathbb{R}^m$. (Corollary 3.1)
- (3) If ϕ is locally semiconcave and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, then $\mu \in \mathscr{S}(\phi(\cdot))$ if and only if $\mu(\text{Sing}(\phi)) > 0$. (Theorem 3.2) Therefore, when $\mu \ll \mathscr{L}^m$, the Lebesgue measure on $\mathbb{R}^m, \mu \notin \mathscr{S}(\phi(\cdot))$.

We aim to study the problem of propagation of singularities of the potential energy functional $u(\cdot)$ with *u* a weak KAM solution of the Hamilton-Jacobi equation

$$H(x, Du(x)) = c[0], \qquad x \in M,$$
(1)

where *M* is a smooth closed manifold, $H : T^*M \to \mathbb{R}$ is a Tonelli Hamiltonian and c[0] is the Mañé's critical value.

Let $L: TM \to \mathbb{R}$ be the Tonelli Lagrangian associated with H, and let $A_t(x, y) = \inf_{\eta} \int_0^t L(\eta, \dot{\eta}) ds$ be the fundamental solution, where the infimum is taken over all absolutely continuous curves $\eta: [0, t] \to M$ connecting $x = \eta(0)$ to $y = \eta(t)$. Set $c^t(x, y) = A_t(x, y)$ and recall the dynamical cost functional

$$C^{t}(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{M \times M} c^{t}(x, y) \, d\gamma.$$

Based on the regularity properties of the fundamental solution $A_t(x, y)$, we have the following regularity results for the dynamical cost functional $C^t(\mu, \nu)$ (Theorem 3.3): Suppose t, a, b > 0 with a < b, and let $K \subset \mathbb{R}^m$ be a convex and compact subset. Then, there exists $\tau_1 > 0$ such that:

- (1) $C^{t}(\mu, \cdot)$ and $C^{t}(\cdot, \mu)$ are superlinear on $\mathscr{P}_{c}(\mathbb{R}^{m})$ endowed with the W_{1} metric, where $\mu \in \mathscr{P}_{c}(\mathbb{R}^{m})$;
- (2) $(t, v) \mapsto C^{t}(\mu, v) \in \operatorname{Lip}([a, b] \times \mathscr{P}(K))$, where $\mu \in \mathscr{P}(K)$;
- (3) For $t \in (0, \tau_1]$, there exists $C_K > 0$ depending on K, such that for any $\mu \in \mathscr{P}(K)$, $C^t(\mu, \cdot)$ is locally strongly semiconcave on $\mathscr{P}(K)$ with constant $\frac{C_K}{t}$.

Moreover, if $(t, v) \mapsto C^t(\mu, v)$ is locally differentiable, the dynamical cost functional satisfies the Hamilton-Jacobi inequality

$$D_t C^t(\mu, \nu) + \int_{\mathbb{R}^m} H(y, \partial_\nu C^t(\mu, \nu)) \, d\nu \leq 0.$$

In general, a strict inequality can be obtained in the Hamilton-Jacobi inequality above.

By addressing the Rubinstein-Kantorovitch duality in terms of abstract Lax-Oleinik operators ([37]), we introduce the random Lax-Oleinik operators:

$$P_t^+\phi(\mu) := \sup_{\nu \in \mathscr{P}(\mathbb{R}^m)} \left\{ \phi(\nu) - C^t(\mu, \nu) \right\},$$
$$P_t^-\phi(\mu) := \inf_{\nu \in \mathscr{P}(\mathbb{R}^m)} \left\{ \phi(\nu) + C^t(\nu, \mu) \right\}.$$

We consider the following Hamilton-Jacobi equation:

$$\partial_t U(t,\mu) + \int_{\mathbb{R}^m} H(x,\partial_\mu U(t,\mu)(x)) \, d\mu = 0, \tag{PHJ}_e)$$

where $(t, \mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$. Then, if ϕ is lower semicontinuous and (κ_1, κ_2) -Lipschitz in the large, then $U(t, \mu) := P_t^- \phi(\mu)$ satisfies the following Cauchy problem for equation (PHJ_e) in the viscosity sense (see Definition 4.2):

$$\begin{cases} \partial_t U(t,\mu) + \int_{\mathbb{R}^m} H\left(x,\partial_\mu U(t,\mu)\right) d\mu = 0, \quad (t,\mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m), \\ U(0,\mu) = \phi(\mu), \qquad \qquad \mu \in \mathscr{P}_c(\mathbb{R}^m). \end{cases}$$

Moreover, *u* is a weak KAM solution of (1) if and only if $P_t^- u + c[0]t = u(\cdot)$ for all $t \ge 0$.

Analogous to classic weak KAM theory ([44, 43]), an absolutely continuous curve $\{\mu_s\}_{s\in I} \subset \mathscr{P}_c(\mathbb{R}^m)$ defined on some interval $I \subset \mathbb{R}$ is called a $(u(\cdot), C, c[0])$ -calibrated curve if for any $a, b \in I$, with $a \leq b$, the following condition holds:

$$u(\mu_b) - u(\mu_a) = C^{b-a}(\mu_a, \mu_b) + c[0](b-a).$$

We say that $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ is a cut point of $u(\cdot)$, the potential energy functional induced by the weak KAM solution u, if all $(u(\cdot), C, c[0])$ -calibrated curves ending at μ cannot be extended forward as $(u(\cdot), C, c[0])$ -calibrated curves. The set of all cut points of $u(\cdot)$ is denoted by $\mathscr{C}(u(\cdot))$, which is called the cut locus of the functional $u(\cdot)$ (with respect to the dynamical cost functional C^t).

In general, it holds that $\mathscr{S}(u(\cdot)) \subset \mathscr{C}(u(\cdot))$, and $\mu \in \mathscr{S}(u(\cdot))$ if and only if $\mu(\text{Sing}(u)) > 0$, where Sing (*u*) denotes the set of singularities of *u*.

Building on the work of the first two authors on intrinsic singular characteristics in [25] (see also [26, 31, 27]), we consider the following result: for any $\phi \in \text{Lip}(\mathbb{R}^m) \cap \text{SCL}_{\text{loc}}(\mathbb{R}^m)$ and any $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, there exists $t_{\phi,\mu} > 0$ depending on ϕ and μ , such that the problem

$$\underset{\nu \in \mathscr{P}(\mathbb{R}^m)}{\arg \max} \left\{ \phi(\nu) - C^t(\mu, \nu) \right\}, \quad t \in (0, t_{\phi, \mu}]$$

has a unique solution (singleton). We define the curve $v_{\phi,\mu}(t)$ as follows:

$$v_{\phi,\mu}(t) := \begin{cases} \mu, & t = 0, \\ \arg\max_{\nu \in \mathscr{P}(\mathbb{R}^m)} \left\{ \phi(\nu) - C^t(\mu, \nu) \right\}, & t \in (0, t_{\phi,\mu}] \end{cases}$$

Then, we have the following convergence result:

$$\lim_{t \to 0^+} v_{\phi,\mu}(t) = \mu \quad \text{in the sense of the metric } W_p, \quad \text{for any fixed } p \ge 1.$$
 (Theorem 5.3)

However, the curve $v_{\phi,\mu}(\cdot)$ does not generally possess a semi-group property. To address this, we invoke the notion of a maximal slope curve from abstract gradient flow theory (see, for instance, [9]). Using an EDI (energy dissipation inequality) framework, we derive a minimizing movement for any given pair (ϕ, C^t) (see also [29] for the case in the calculus of variation).

Now, we formulate the following main results of this paper:

- (1) Let $u(\cdot)$ be the potential energy functional induced by a weak KAM solution u of (1). If $\mu \in \mathscr{C}(u(\cdot)), v_{u,\mu}(t)$ and $t_{u,\mu} > 0$ is mentioned above, then $v_{u,\mu}(t) \in \mathscr{S}(u(\cdot))$ for all $t \in (0, t_{u,\mu}]$. (Theorem 5.4)
- (2) Suppose $\phi \in \text{SCL}(\mathbb{T}^m)$, T > 0 and $\mu_0 \in \mathscr{P}(\mathbb{T}^m)$. Then there exists a Lipschitz curve $\mu(\cdot)$: [0, T] $\to \mathscr{P}(\mathbb{T}^m)$, which satisfies the following continuity equation:

$$\begin{cases} \frac{d}{dt}\mu + \operatorname{div}(H_p(\cdot, \mathbf{p}_{\phi}^{\#}(\cdot)) \cdot \mu) = 0, \\ \mu(0) = \mu_0, \end{cases}$$
(2)

where $\mathbf{p}_{\phi}^{\#}(x) := \arg \min\{H(x, p) : p \in D^{+}\phi(x)\}$ for any $x \in \mathbb{T}^{m}$ (Theorem 5.5). Moreover, if $\phi = u$ in (2) is a weak KAM solution of (1),

- $\mu_0(\overline{\operatorname{Sing}(\phi)}) > 0$ implies $[\mu(t)](\overline{\operatorname{Sing}(\phi)}) > 0$ for all $t \in [0, T]$. (Corollary 5.1)
- $\mu_0 \in \mathscr{C}(u(\cdot))$ implies $\mu(t) \in \mathscr{C}(u(\cdot))$ for all $t \in [0, T]$. (Corollary 5.1)
- (3) For such a Lipschitz curve satisfying (2) we have that for any $f \in C^{\infty}(\mathbb{T}^n)$, the map $t \mapsto f(\mu(t))$ is in Lip ([0, *T*]; \mathbb{R}) and its right derivative exists such that

$$\frac{d^+}{dt}f(\mu(t)) = \int_{\mathbb{T}^m} \langle Df(x), H_p(x, \mathbf{p}_{\phi}^{\#}(x)) \rangle \, d\mu(t), \qquad \forall t \in (0, T].$$

(Theorem 5.6)

We emphasize the following novel aspects of this work:

- Dynamics after the Formation of Singularities: In this paper, we study the dynamics that arise after the formation of singularities in optimal transport. In this setting, the forward dynamics is no longer governed by a regular Lagrangian flow, as discussed in [8]. This makes the classical DiPerna-Lions theory (which is typically used in the regular case) inapplicable to our case.
- Challenges from Non-Compactness: Due to the lack of local compactness in the space of probability measures, our problem cannot be treated as a straightforward extension of results from finite-dimensional settings. This lack of compactness introduces significant technical challenges that distinguish our approach from previous treatments of similar problems.

Acknowledgements. Piermarco Cannarsa was partly supported by the PRIN 2022 PNRR Project P20225SP98 "Some Mathematical Approaches to Climate Change and Its Impacts" (funded by the European Community-Next Generation EU), CUP E53D2301791 0001. He was also supported by the INdAM (Istituto Nazionale di Alta Matematica) Group for Mathematical Analysis, Probability and Applications, and by the MUR Excellence Department Project awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C23000330006.Wei Cheng was partly

supported by the National Natural Science Foundation of China (Grant No. 12231010). Tianqi Shi was partly supported by the "New Cornerstone Investigator Program" led by Jinxin Xue. Wei Cheng also thanks Jiahui Hong for helpful discussions, Bangxian Han and Shibin Chen for comments on some literatures.

Μ	smooth Riemannian manifold without boundary (also compact in certain
	contexts)
\mathbb{R}^m and \mathbb{T}^m	<i>m</i> -dimensional Euclidean space and torus
TM/T^*M	tangent/cotangent bundle of M
π	the canonical projection from T^*M to M
SCL(M)	the set of semiconcave functions with linear modulus on M
$\operatorname{SCL}_{\operatorname{loc}}(\mathbb{R}^m)$	the set of locally semiconcave functions with linear modulus on \mathbb{R}^m
$\operatorname{Lip}(\mathbb{R}^m)$	the set of Lipschitz functions on \mathbb{R}^m
AC	absolutely continuous
$\eta = \eta(s)$	the absolutely continuous deterministic curve on M or \mathbb{R}^m
ϕ	a function on M or \mathbb{R}^m (usually denoted as a semiconcave function)
и	the viscosity solution of a stationary type HJ equation on M or \mathbb{R}^m
$D^+\phi(x)$	the superdifferential of a function ϕ
$\mathbf{p}_{\phi}^{\#}(x)$	the minimal-energy element of $D^+\phi(x)$ w.r.t. the Hamiltonian H
$\operatorname{Sing}(\phi)$	the set of non-differentiability points of a function ϕ
$\operatorname{Cut}(u)$	the cut locus of viscosity solution <i>u</i>
$\tau_u(x)$	the cut time function of viscosity solution <i>u</i>
B(x, R)	open ball with radius R
$C^{1,1}(M)$	the space of differentiable functions with Lipschitz differential on M
$\mathrm{UC}\left(M\right)$	the space of uniformly continuous real-valued functions on M
Φ_H^t	the Hamiltonian flow associated to the Hamiltonian H
X, Y	Polish space, i.e., complete and separable metric space
$\mathscr{P}(X)$	the set of all Borel probability measures on X
$\Gamma(\mu, \nu)$	the set of transport plans between Borel probability measures μ and ν
\mathscr{L}^m	Lebesgue measure on <i>m</i> -dimensional manifold
$(\mathscr{P}_p(X), W_p)$	<i>p</i> -Wasserstein space of $X \ (p \ge 1)$
$\Gamma_p(\mu, \nu)$	the set of minimizers of $W_p(\mu, \nu)$
π_i	the canonical projection from a product space to its <i>i</i> -th component
$\pi_{i,j}$	the canonical projection from product space to another product space of
	<i>i</i> -th and <i>j</i> -th component of the former
$\mathscr{P}_{c}(\mathbb{R}^{m})$	the set of all Borel probability measures on \mathbb{R}^m with compact supports
U	the functional on $\mathscr{P}_{c}(\mathbb{R}^{m})$
$\partial^{\pm} U(\mu)$	the localized Fréchet super/sub-differential of U at μ
$\mathscr{S}(U)$	the set of singular points of U
$\phi(\cdot)$	the potential energy functional induced by the function ϕ on M or \mathbb{R}^m
$C^{t}(\mu, \nu)$	the value of dynamical cost functional at μ, ν
v · · ·	

$\Gamma_o^t(\mu, \nu)$	the set of minimizers of $C^t(\mu, \nu)$
$(\Omega,\mathscr{F},\mathbb{P})$	the probability space of sample space Ω
E	the expectation of random variables on $(\Omega, \mathscr{F}, \mathbb{P})$
$\xi = \xi(s) = \xi(s, \omega)$	the absolutely continuous random curve of $(\Omega, \mathscr{F}, \mathbb{P})$ on <i>M</i> or \mathbb{R}^m
$T_u(\mu)$	the cut time function of measure associated to potential energy $u(\cdot)$
$\mathscr{C}(u(\cdot))$	the cut locus of potential energy functional $u(\cdot)$

2. Preliminaries

Let (M, g) be a smooth Riemannian manifold. d_g is the metric induced by g on M. For any $x \in M$, $T_x M$ and $T_x^* M$ are denoted by the tangent and cotangent spaces of M at x, respectively. TM and T^*M are the tangent and cotangent bundle of M, respectively. Let $\pi : T^*M \to M$ be the canonical projection from T^*M to M, defined as $\pi(x, p) := x$, for all $(x, p) \in T^*M$.

2.1. **Some aspects on semiconcave functions.** Some basic relevant notions on semiconcavity are listed below.

- Let Ω be an open convex subset of \mathbb{R}^m . A continuous function $\phi : \Omega \to \mathbb{R}$ is called a *semicon-cave function (of linear modulus) with constant* $C \ge 0$ if for any $x \in \Omega$ there exists $p \in \mathbb{R}^m$ such that

$$\phi(y) \le \phi(x) + \langle p, y - x \rangle + \frac{C}{2} |x - y|^2, \qquad \forall y \in \Omega,$$
(3)

and we denote as $\phi \in SCL(\Omega)$. The set of covectors *p* satisfying (3) is called the *proximal* superdifferential of ϕ at *x* and we denote it by $D^+\phi(x)$.

- Similarly, ϕ is *semiconvex* if $-\phi$ is semiconcave. The set of $D^-\phi(x) = -D^+(-\phi)(x)$ is called *the proximal subdifferential of* ϕ *at x*.
- For any $\phi \in \text{SCL}(\Omega)$, the set $D^+\phi(x)$ is a singleton if and only if ϕ is differentiable at *x*, and $D^+\phi(x) = \{D\phi(x)\}$. A point *x* is called a *singular point* of a semiconcave function ϕ if $D^+\phi(x)$ is not a singleton. We denote by Sing (ϕ) the set of all singular points of ϕ .
- Given any $\phi \in SCL(\Omega)$. We call $p \in D^*\phi(x)$, the set of *reachable gradients*, if there exists a sequence $x_k \to x$ as $k \to \infty$, ϕ is differentiable at each x_k and $p = \lim_{k\to\infty} D\phi(x_k)$. We have $D^*\phi(x) \subset D^+\phi(x)$ and $D^+\phi(x) = \operatorname{co} D^*\phi(x)$.

For more on the semiconcave functions in Euclid spaces, the readers can refer to [32].

Proposition 2.1. A function $\phi : \Omega \to \mathbb{R}$ is a semiconcave function with constant $C \ge 0$ if and only if there exists a family of C^2 -functions $\{\phi_i\}$ with $D^2\phi_i \le CI_m$ such that

$$\phi = \inf \phi_i.$$

Proposition 2.1 is an important and useful characterization of semiconcavity. Let *S* be a compact topological space and $F: S \times \mathbb{R}^m \to \mathbb{R}$ be a continuous function such that

- (a) $F(s, \cdot)$ is of class C^2 for all $s \in S$ and $||F(s, \cdot)||_{C^2}$ is uniformly bounded by some constant C,
- (b) $D_x F(s, x)$ is continuous on $S \times \mathbb{R}^m$,
- (c) $\phi(x) = \inf\{F(s, x) : s \in S\}.$

We call such a function ϕ a marginal function of a family of C^2 -functions.

Proposition 2.2. Let $\phi(x) = \inf\{F(s, x) : s \in S\}$ be a marginal function of the family of C^2 -functions $\{F(s, \cdot)\}_{s \in S}$.

- (1) ϕ is semiconcave with constant C.
- (2) $D^*\phi(x) \subset \{D_xF(s,x) : x \in M(x)\}, where M(x) := \arg\min\{F(s,x) : s \in S\}.$
- (3) For each $x \in \mathbb{R}^m$,

$$D^+\phi(x) = \begin{cases} D_x F(s, x), & M(x) = \{s\} \text{ is a singleton}; \\ \cos\{D_x F(s, x) : s \in M(x)\}, & otherwise. \end{cases}$$

Remark 2.1. A function $\phi : M \to \mathbb{R}$, with M a manifold, is called a semiconcave function, if there exists a family of C^2 -functions $\{\phi_i\}$ such that $\phi = \inf_i \phi_i$, and the Hessians of ϕ_i 's is uniformly bounded above. The readers can refer to [12, 47, 75] for more discussion on the semiconcavity of functions on manifolds, including the equivalence of various definitions of semiconcave functions. Thus, because of the local nature of our main discussion and for convenience, we will work on Euclidean space instead, then get similar conclusions on manifolds.

2.2. Weak KAM theory. Let *M* be a smooth manifold without boundary. A function L = L(x, v): $TM \rightarrow \mathbb{R}$ is a time-independent *Tonelli Lagrangian*, if *L* is of class C^2 and satisfies the following conditions:

- (L1) The function $v \mapsto L(x, v)$ is strictly convex for all $x \in M$.
- (L2) There exist convex, nondecreasing and superlinear functions $\theta_0, \theta_1 : [0, +\infty) \rightarrow [0, +\infty)$ and positive constants c_0, c_1 , such that

$$\theta_0(|v|_x) - c_0 \leq L(x, v) \leq \theta_1(|v|_x) + c_1, \qquad \forall (x, v) \in TM,$$

where $|v|_x = \sqrt{g_x(v, v)}$.

We call H associated with the Lagrangian L a Tonelli Hamiltonian, defined by

$$H(x, p) = \sup_{v \in T_x M} \{ p(v) - L(x, v) \}, \qquad (x, p) \in T^* M.$$

Given any $x, y \in M$, and t > 0, we denote by $\Gamma_{x,y}^t$ the set of absolutely continuous curves $\eta \in AC([0, t], M)$ with $\eta(0) = x$ and $\eta(t) = y$. We call the function $A_t(x, y) : \mathbb{R}^+ \times M \times M \to \mathbb{R}$ the *fundamental solution* of the associated Hamilton-Jacobi equation, defined by

$$A_t(x,y) := \inf_{\eta \in \Gamma_{x,y}^t} \int_0^t L(\eta(s), \dot{\eta}(s)) \, ds, \qquad t > 0, \quad x, y \in M.$$

For any ϕ : $M \to \mathbb{R}$, $x \in M$, and t > 0, we define

$$T_t^- \phi(x) := \inf_{y \in M} \{ \phi(y) + A_t(y, x) \},$$
(4)

$$T_t^+ \phi(x) := \sup_{y \in M} \{ \phi(y) - A_t(x, y) \}.$$
 (5)

The families $\{T_t^-\}_{t>0}$ and $\{T_t^+\}_{t>0}$ of operators are called *negative and positive Lax-Oleinik evolution* respectively.

Theorem 2.1 ([45]). Let $H : T^*M \to \mathbb{R}$ be a Tonelli Hamiltonian, then $u(t, x) := T_t^-\phi(x)$ and $\check{u}(t, x) = T_t^+\phi(x)$ are the unique viscosity solution to Cauchy problem of evolutionary type Hamilton-Jacobi equation

$$\begin{cases} \partial_t u(t, x) + H(x, \partial_x u(t, x)) = 0, \\ u(0, x) = \phi(x). \end{cases}$$
(HJ_e-)

and

$$\begin{cases} \partial_t \breve{u}(t,x) - H(x,\partial_x \breve{u}(t,x)) = 0\\ \breve{u}(0,x) = \phi(x), \end{cases}$$
(HJ_e+)

respectively.

Theorem 2.2 ([43, 48, 28]). Assume that M is compact and $H : T^*M \to \mathbb{R}$ is a Tonelli Hamiltonian, then there exists a unique $c = c[0] \in \mathbb{R}$ such that

$$H(x, Du(x)) = c, \qquad x \in M \tag{HJ}_s$$

admits a viscosity solution, where c[0] is the Mañé critical value. Moreover, the following statements are equivalent:

- (1) *u* is the viscosity solution to (HJ_s) ;
- (2) $u = T_t^- u + c[0]t$ for any $t \ge 0$. We also call u a weak KAM solution of (HJ_s) ;
- (3) $T_t^- \circ T_t^+ u = u$ for any $t \ge 0$.

If $M = \mathbb{R}^m$, then there exists $c[0] \in \mathbb{R}$ such that (HJ_s) admits a viscosity solution $u \in Lip(\mathbb{R}^m) \cap$ SCL (\mathbb{R}^m) satisfying that $T_t^-u + c[0]t = u$ for all $t \ge 0$. If c < c[0], there is no viscosity solution of (HJ_s) .

For more conclusions on the weak KAM theory and viscosity solutions of Hamilton-Jacobi equations in Euclid space \mathbb{R}^m or other non-compact manifold, readers can refer to [48, 46]. We collect some regularity properties of Lax-Oleinik operators as follows.

Proposition 2.3 ([43, Proposition 4.6.6]). Suppose M is compact, $\phi \in C(M)$, $t_0 > 0$ and c[0] = 0.

- (1) $\{T_t^-\}_{t>0}$ and $\{T_t^+\}_{t>0}$ are the semigroups with respect to $t \in (0, +\infty)$;
- (2) $T_t^-, T_t^+ : C(M) \to C(M);$
- (3) $\lim_{t\to 0^+} T_t^- \phi = \lim_{t\to 0^+} T_t^+ \phi = \phi$. Then we define $T_0^- \phi = T_0^+ \phi = \phi$;
- (4) For any t > 0, $T_t^- \phi \in \text{SCL}(M)$ (resp. $-T_t^+ \phi \in \text{SCL}(M)$) and $\{T_t^- \phi\}_{t \ge t_0}$ (resp. $\{-T_t^+ \phi\}_{t \ge t_0}$) are semiconcave uniformly to t;
- (5) $t \mapsto T_t^- \phi$ ($t \mapsto T_t^+ \phi$) is uniformly continuous on $[0, +\infty)$;
- (6) $(t, x) \mapsto T_t^- \phi(x) ((t, x) \mapsto T_t^+ \phi(x))$ is continuous on $[0, +\infty) \times M$, locally Lipschitz on $(0, +\infty) \times M$ and equi-Lipschitz on $[t_0, +\infty) \times M$ with respect to ϕ .

2.3. Cut locus. This subsection is inspired by [28]. For any weak KAM solution *u* of equation (HJ_s) , we define the function $B_u : [0, +\infty) \times M \to \mathbb{R}$ as follows:

$$B_u(t, x) := u(x) - T_t^+ u(x) + c[0]t, \quad (t, x) \in [0, +\infty) \times M.$$

According to the definition of weak KAM solution, it is straightforward to verify that

$$B_{u}(t, x) = T_{t}^{-} \circ T_{t}^{+} u(x) - T_{t}^{+} \circ T_{t}^{-} u(x).$$

We denote by Cut (*u*) the *cut locus of u*, which consists of points $x \in M$ where any calibrated curve ending at *x* cannot be extended further as a calibrated curve. More precisely, let $\tau : M \to \mathbb{R}$ be the *cut time function of u*, defined for any $x \in M$ as

$$\tau_u(x) := \sup\{t \ge 0 : \exists \eta \in C^1([0, t]; M), u(\eta(t)) - u(x) = A_t(x, \eta(t)) + c[0]t\}.$$

Thus $Cut(u) = \{x \in M : \tau_u(x) = 0\}.$

Proposition 2.4. Suppose $\tau_u(x) > 0$ and $t \in (0, \tau_u(x))$. Then there exists a unique (u, L, c[0])-calibrated curve $\eta_x : [0, t] \to M$ with $\eta_x(0) = x$, and we have

$$\eta_x(s) = \pi \circ \Phi_H^s(x, Du(x)) = \pi \circ \Phi_H^{s-t}(\eta_x(t), Du(\eta_x(t))), \ s \in [0, t].$$
(6)

Recall that the (*projected*) Aubry set of a weak KAM solution u of (HJ_s)

$$I(u) := \{x \in M : \exists u \text{-calibrated curve } \eta : [0, +\infty) \to M, \eta(0) = x\}.$$

Proposition 2.5 ([26, 27]). Suppose u is the weak KAM solution of (HJ_s) .

- (1) Given t > 0 and $x \in M$, then $T_t^+u(x) c[0]t = u(x)$ if and only if there exists a u-calibrated curve $\eta : [0, t] \to M$ such that $\eta(0) = x$;
- (2) $\tau_u(x) = \sup\{t \ge 0 : B_u(t, x) = 0\}$. If $t \in [0, \tau_u(x)]$, then $B_u(t, x) = 0$;
- (3) τ_u is upper semicontinuous and Cut (*u*) is a G_{δ} -set;
- (4) $I(u) = \{x \in M : \tau_u(x) = +\infty\}.$

2.4. A priori estimates on fundamental solutions. The following estimates about $A_t(x, y)$ are well-known.

Proposition 2.6 ([30, Section 2.4]). Any least action curve of $A_t(x, y)$ is class of C^2 . Moreover, there exists a monotone nondecreasing, superlinear and convex function of $F : [0, +\infty) \rightarrow$ $[0, +\infty)$, such that for any t, R > 0, when $d(x, y) \leq R$, the minimizer η of $A_t(x, y)$ satisfies

$$|\dot{\eta}(s)| \leq F\left(\frac{R}{t}\right), \quad \forall s \in [0, t].$$

Definition 2.1 (Lipschitz in the large). Let $\kappa_1 \ge 0, \kappa_2 > 0$. We say ϕ is a (κ_1, κ_2) -Lipschitz in the *large* function in metric space (X, d), if for any $x, y \in X$,

$$|\phi(x) - \phi(y)| \le \kappa_1 d(x, y) + \kappa_2.$$

Recall that if $(X, d) = (M, g), \phi \in UC(M)$ if and only if for arbitrary $\varepsilon > 0$, there exists $K_{\varepsilon} > 0$, such that ϕ is $(K_{\varepsilon}, \varepsilon)$ -Lipschitz in the large.

Proposition 2.7 ([30, Lemma 3.1]). Assume that ϕ is lower semicontinuous and (κ_1, κ_2) -Lipschitz in the large. Then for any $x \in M$, there exists $y \in M$, $T_t^-\phi(x) = \phi(y) + A_t(y, x)$. Furthermore, there exists $\lambda_{\phi} > 0$, which only depends on L and $\kappa_1 > 0$ satisfies that for such minimizer $y_{t,x}$,

$$d(y_{t,x}, x) \leq \lambda_{\phi} t + \kappa_2.$$

More specifically, $\lambda_{\phi} := c_0 + \theta_1(0) + \theta_0^*(\kappa_1 + 1) + c_1$, where θ_0^* is the Fenchel-Legendre transformation of θ_0 .

Similarly, If ϕ is upper semicontinuous and (κ_1, κ_2) -Lipschitz in the large, then for any $x \in M$, there exists $z \in M$, $T_t^+\phi(x) = \phi(z) - A_t(x, z)$ and such maximizer $z_{t,x}$ safisfies

$$d(z_{t,x}, x) \leq \lambda_{\phi} t + \kappa_2.$$

Proposition 2.8 ([36, 25]). Assume that K is a compact and convex subset of \mathbb{R}^m and $\lambda > 0$. Then there exist $\tau_1 \ge \tau_2 > 0$ and $C_{\lambda}, C'_{\lambda}, C''_{\lambda} > 0$, such that $x \in K^1$,

(1) Functions $(t, y) \mapsto A_t(x, y)$ and $(t, y) \mapsto A_t(y, x)$ are both semiconcave and semiconvex (thus $C_{loc}^{1,1}$) on

$$S(x,\lambda,\tau_1) := \{(t,y) \in \mathbb{R}^+ \times K : t \in (0,\tau_1], d(y,x) \le \lambda t\}$$

with constant C_{λ}/t and $-C'_{\lambda}/t$ respectively; In this case, for any $(t, y) \in S(x, \lambda_{\phi}, \tau_1)$,

$$D_{y}A_{t}(x, y) = p(t), \ D_{t}A_{t}(x, y) = -H(\eta(t), p(t)), \ D_{x}A_{t}(x, y) = -p(0),$$

where $\eta \in \Gamma_{x,y}^t$ is the unique minimizer of $A_t(x, y)$, and $p(s) := L_v(\eta(s), \dot{\eta}(s))$, $s \in [0, t]$ is the dual arc of η ;

(2) When $t \in (0, \tau_2]$, $A_t(x, \cdot)$ is strictly convex on $B(x, \lambda t)$ with constant C''_{λ}/t .

In particlular, if $\phi \in SCL(K)$ and $\lambda = \lambda_{\phi}$, then we can find $\tau_1(\phi) = \tau_1$, $\tau_2(\phi) = \tau_2$ and $C_{\lambda_{\phi}}, C'_{\lambda_{\phi}}, C''_{\lambda_{\phi}} > 0$ depend on λ_{ϕ} , which make (1) and (2) hold true, as well as $\phi(\cdot) - A_t(x, \cdot)$ is strictly convex on $B(x, \lambda_{\phi}t)$, thus has a unique maximizer.

2.5. Lasry-Lions regularization, Arnaud's theorem and intrinsic singular characteristic. In the case of compact manifold *M*, the following Lasry-Lions regularization type result in the context of weak KAM theory is due to Patrick Bernard.

Proposition 2.9 ([14]). Suppose *M* is compact and $\phi \in SCL(M)$. Then there exists $0 < \tau_3(\phi) \leq \tau_2(\phi)$, such that when $t \in (0, \tau_3(\phi)]$, $T_t^+ \phi \in C^{1,1}(M)$.

Since ϕ is semiconcave, $D^+\phi(x) \neq \emptyset$ holds for all $x \in M$. Now we recall the evolution of the pseudo-graphs

graph
$$(D^+\phi) := \{(x, p) \in T^*M : p \in D^+\phi(x), x \in M\},$$

graph $(DT_t^+\phi) := \{(x, DT_t^+\phi(x)) : x \in M\}, t \in (0, \tau_3(\phi)].$

under the Hamiltonian flow $\{\Phi_H^t\}_{t \in (0, \tau_3(\phi)]}$.

Proposition 2.10 ([10]). *For any* $t \in (0, \tau_3(\phi)]$ *, we have*

$$\Phi_H^t(\operatorname{graph}(DT_t^+\phi)) = \operatorname{graph}(D^+\phi).$$

Suppose *M* is compact, for any fixed $x \in M$, we define curve $\mathbf{y}_x(t) : [0, \tau_3(\phi)] \to M$,

$$\mathbf{y}_{x}(t) := \begin{cases} x, & t = 0, \\ \arg \max\{\phi(y) - A_{t}(x, y) : y \in M\}, & t \in (0, \tau_{3}(\phi)] \end{cases}$$

We call \mathbf{y}_x an *intrinsic singular characteristic* from *x*.

Proposition 2.11 ([29]). Suppose $\phi \in SCL(M)$. Then

$$\mathbf{y}_{x}(t) = \pi \circ \Phi_{H}^{t}(x, DT_{t}^{+}\phi(x)), \ x \in M, t \in (0, \tau_{3}(\phi)].$$

Let $\eta_{t,x} \in \Gamma_{x,\mathbf{v},(t)}^t$ be the unique minimizer of $A_t(x,\mathbf{y}_x(t))$, then

$$\eta_{t,x}(s) = \pi \circ \Phi_{H}^{s}(x, DT_{t}^{+}\phi(x)) = \pi(\eta_{t,x}(s), DT_{t-s}^{+}\phi(\eta_{t,x}(s))), \ \forall s \in [0, t).$$

¹In general, τ_1 and τ_2 depend on x in non-compact case.

In local coordinates, $\eta_{t,x}$ satisfies differential equation

$$\begin{pmatrix} \dot{\eta}_{t,x}(s) = H_p(\eta_{t,x}(s), DT^+_{t-s}\phi(\eta_{t,x}(s))), & s \in [0,t], \\ \eta_{t,x}(0) = x. \end{cases}$$

Moreover, there exists a constant C > 0, which is independent to x, such that

$$\|\mathbf{y}_x - \eta_{t,x}\|_{\infty} \leq Ct.$$

In [25], authors show that if ϕ is the viscosity solution to (HJ_s) , then $x \in Sing(\phi)$ implies $\mathbf{y}_x(t) \in Sing(\phi)$ for $t \in [0, \tau_3(\phi)]$. This is also true for $\phi \in SCL(M)$ only with

$$\underset{p \in D^+\phi(x)}{\arg\min}\{H(x, p)\} \notin D^*\phi(x),$$

where $D^*\phi(x)$ is the set of reachable gradients of ϕ at *x*.

2.6. **Optimal transport.** Suppose (X, d) is a metric space. We denote $\mathscr{P}(X)$ the set of all probability measures on *X*.

2.6.1. *Monge-Kantorovich problem and Kantorovich duality*. For two fixed Polish spaces *X* and *Y*, the modern formulation of Monge's problem has probability measures $\mu \in \mathscr{P}(X)$, $\nu \in \mathscr{P}(Y)$ and a Borel cost function $c(x, y) : X \times Y \to \mathbb{R}$, representing the cost of shipping a unit mass from *x* to *y*. Given these data, the problem is

$$\inf\left\{\int_X c(x, T(x)) \, d\mu : T : X \to Y \text{ is Borel}, T_{\#}\mu = \nu\right\},\tag{M}$$

where *T* is called the transport map from μ to ν . Under certain conditions (for example, [80, Theorem 10.28]), Monge's problem (M) can have a solution. Brenier ([22, 23]) and Knott-Smith ([67]) proved that (M) has a solution when $X = Y = \mathbb{R}^m$, $c(x, y) = \frac{1}{2}|x - y|^2$ and $\mu \ll \mathcal{L}^m$.

Remark 2.2. We remark that the conclusion due to Brenier and Knott-Smith provides us with the existence of a probability space $(\Omega, \mathscr{F}, \mathbb{P})$ such that for arbitrary $\mu \in \mathscr{P}_2(\mathbb{R}^m)$ (see Definition 2.3), we can always find a random variable $T : \Omega \to \mathbb{R}^m$, which satisfies law $(T) = \mu$. Indeed, we choose $\Omega = \mathbb{R}^m$, \mathscr{F} is the set of Lebesgue measurable subsets of \mathbb{R}^m and $\mathbb{P} = \mathscr{L}^m|_{B(0,R)}$, where R > 0 makes $\mathscr{L}^m(B(0,R)) = 1$. If $c(x,y) = \frac{1}{2}|x - y|^2$, for the corresponding Monge's transport problem, there exists an optimal map $T : \mathbb{R}^m \to \mathbb{R}^m$, i.e., law $(T) = \mu$. By Example 10.36 of [80], the argument for the case X = Y = M with M compact manifold and $\mu \in \mathscr{P}(M)$ is similar.

For Polish spaces X and Y, we call $\gamma \in \mathscr{P}(X \times Y)$ the *transport plan* between $\mu \in \mathscr{P}(X)$ and $\nu \in \mathscr{P}(Y)$, if

$$(\pi_X)_{\#}\gamma = \mu, \qquad (\pi_Y)_{\#}\gamma = \nu,$$

where π_X and π_Y are the canonical projection maps from $X \times Y$ to X and Y respectively. We denote $\Gamma(\mu, \nu)$ the set of all transport plans between μ and ν . Obviously, $\mu \times \nu \in \Gamma(\mu, \nu)$.

Kantorovich's problem asks to find

$$C(\mu, \nu) := \inf \left\{ \int_{X \times Y} c(x, y) \, d\gamma : \gamma \in \Gamma(\mu, \nu) \right\}.$$
(K)

The following is one of the most basic results of the theory of optimal transport, namely the Kantorovich-Rubinstein duality theorem.

Theorem 2.3 ([80, 5]). If $c : X \times Y \to \mathbb{R}$ is lower-semicontinuous, then

$$C(\mu,\nu) = \sup_{(\phi,\psi)\in I_c} \left\{ \int_Y \psi \, d\nu - \int_X \phi \, d\mu \right\} = \sup_{(\phi,\psi)\in K_c} \left\{ \int_Y \psi \, d\nu - \int_X \phi \, d\mu \right\},\tag{KR}$$

where

$$I_c := \{(\phi, \psi) : \psi(y) - \phi(x) \le c(x, y) \ \forall \ x \in X, y \in Y\},\$$
$$K_c := \left\{(\phi, \psi) \in I_c : \psi = \inf_{x \in X} \{\phi(x) + c(x, \cdot)\}, \phi = \sup_{y \in Y} \{\psi(y) - c(\cdot, y)\}\right\}.$$

In many literatures, elements in K_c are referred to as Kantorovich admissible pairs.

Definition 2.2 ([37]). For any $\phi : X \to \mathbb{R}$, $\psi : Y \to \mathbb{R}$, the positive and negative type of abstract *Lax-Oleinik operators* are defined as

$$T^+\psi(x) := \sup_{y \in Y} \{\psi(y) - c(x, y)\}, \quad T^-\phi(y) := \inf_{x \in X} \{\phi(x) + c(x, y)\}.$$

Let $\phi : X \to \mathbb{R}$ and $\psi : Y \to \mathbb{R}$. For any $\mu \in \mathscr{P}(X)$ and $\nu \in \mathscr{P}(Y)$, we try to find a function $\phi : X \to \mathbb{R}$ and a function $\psi : Y \to \mathbb{R}$ such that

$$\int_{Y} T^{-}\phi \, d\nu = \inf_{\gamma \in \Gamma(\mu,\nu)} \int_{X \times Y} \phi(x) + c(x,y) \, d\gamma, \tag{K}^{-}$$

$$\int_{X} T^{+} \psi \ d\mu = \sup_{\gamma \in \Gamma(\mu, \nu)} \int_{X \times Y} \psi(y) - c(x, y) \ d\gamma. \tag{K}^{+}$$

In terms of abstract Lax-Oleinik operators, the equivalence of the problems (K^-) and (K^+) and well known Rubinstein-Kantorovich duality can be showed as follows.

Theorem 2.4 ([37]).

- (1) If $\phi : X \to \mathbb{R}$ is a solution of (K⁻), then $(\phi, T^-\phi) \in I_c$ is a solution of (KR). Conversely, if $(\phi, \psi) \in I_c$ is a solution of (KR), then ϕ solves (K⁻).
- (2) If $\psi : Y \to \mathbb{R}$ is a solution of (\mathbf{K}^+) , then $(T^+\psi,\psi) \in I_c$ is a solution of (\mathbf{KR}) . Conversely, if $(\phi,\psi) \in I_c$ is a solution of (\mathbf{KR}) , then ψ solves (\mathbf{K}^+) .

Invoking the equivalence above, it is useful to refine the analysis of the dynamical nature of the associated optimal transport problem, as initiated by Bernard and Buffoni ([15, 16, 17]), which also forms the basis of the main content in this paper.

2.6.2. Wasserstein space.

Definition 2.3 (Wasserstein space). Let *X* be a Polish space. For $p \in [1, +\infty)$, set

$$\mathscr{P}_p(X) := \left\{ \mu \in \mathscr{P}(X) : \int_X d^p(x, x_0) \, d\mu < +\infty \right\}, \quad \exists x_0 \in X,$$

We call

$$W_p(\mu,\nu) := \left\{ \inf_{\gamma \in \Gamma(\mu,\nu)} \int_{X^2} d^p(x,y) \, d\gamma \right\}^{\frac{1}{p}}.$$

For all $p \ge 1$, we denote $\Gamma_p(\mu, \nu)$ is the set of all optimal plans of $W_p(\mu, \nu)$. The set $\mathscr{P}_p(X)$ endowed the *Wasserstein distance* W_p is called *p*-*Wasserstein space*.

Remark 2.3. If X is a Polish space, then so is $\mathscr{P}_p(X)$. If X is compact, $\mathscr{P}_p(X)$ is tight in the sense of Prokhorov. However, when X is only locally compact, $(\mathscr{P}_p(X), W_p)$ may not be. See, for instance, [80, 9].

Given μ -measurable maps $r, s : (X, \mu) \to X$, a very useful inequality giving an estimate from above of the Wasserstein distance is

$$W_p(\boldsymbol{r}_{\#}\boldsymbol{\mu},\boldsymbol{s}_{\#}\boldsymbol{\mu}) \leq \boldsymbol{d}(\boldsymbol{r},\boldsymbol{s})_{L^p(\boldsymbol{\mu},\boldsymbol{X})} = \left\{ \int d^p(x_1,x_2)d\gamma \right\}^{1/p}$$
(7)

where $\gamma = (\mathbf{r}, \mathbf{s})_{\#} \boldsymbol{\mu} \in \Gamma(\mathbf{r}_{\#} \boldsymbol{\mu}, \mathbf{s}_{\#} \boldsymbol{\mu}).$

A *constant speed geodesic* of a metric space (X, d) is a curve $\eta : [0, 1] \to X$, which satisfies that for any $0 \le s \le t \le 1$,

$$d(\eta(s), \eta(t)) = (t - s)d(\eta(0), \eta(1)).$$
(8)

By triangle inequality, if the equality is replaced by " \leq " in (8) for any $0 \leq s \leq t \leq 1$, then η is a constant speed geodesic equivalently.

For $\mu_i \in \mathscr{P}(X_i)$, $i = 1, \dots, N$, we call $\mu \in \Gamma(\mu_1, \mu_2, \dots, \mu_N)$, if $(\pi_i)_{\#}\mu = \mu_i$, where π_i is the canonical projection from $X_1 \times \dots \times X_N$ to X_i . In the case $X_i = X$, a Hilbert space, $i = 1, \dots, N$. and $\mu \in \mathscr{P}(X^N)$, $N \ge 2$, $1 \le i, j, k \le N$ and $\lambda \in [0, 1]$, set

$$\pi_{i \to j}^{\lambda} := (1 - \lambda)\pi_i + \lambda\pi_j : X^N \to X, \tag{9}$$

$$\pi_{i \to j,k}^{\lambda} := (1 - \lambda)\pi_{i,k} + \lambda\pi_{j,k} : X^N \to X^2,$$
(10)

$$\pi_{k,i \to j}^{\lambda} := (1 - \lambda)\pi_{k,i} + \lambda\pi_{k,j} : X^N \to X^2, \tag{11}$$

$$\boldsymbol{\mu}_{i \to j}^{\lambda} := (\pi_{i \to j}^{\lambda})_{\#} \boldsymbol{\mu} \in \mathscr{P}(X), \tag{12}$$

$$\boldsymbol{\mu}_{i \to j,k}^{\lambda} \coloneqq (\pi_{i \to j,k}^{\lambda})_{\#} \boldsymbol{\mu} \in \mathscr{P}(X^2), \tag{13}$$

$$\boldsymbol{\mu}_{k,i\to j}^{\lambda} \coloneqq (\pi_{k,i\to j}^{\lambda}) \# \boldsymbol{\mu} \in \mathscr{P}(X^2), \tag{14}$$

where π_i is the canonical projection from X^N to the *i*-th *X*, and $\pi_{i,j}$ is the canonical projection from X^N to the product space of *i*-th and *j*-th *X*.

The following theorem implies that, when X is a Hilbert space, $(\mathscr{P}_p(X), W_p)$ is a geodesic space, and geodesics are determined by the optimal plans between the two prescribed measures.

Theorem 2.5 ([9, Lemma 7.2.1 & Theorem 7.2.2]). Assume that $(X, \langle \cdot, \cdot \rangle)$ is a Hilbert space, and p > 1. If $\mu_1, \mu_2 \in \mathscr{P}_p(X)$ and $\mu \in \Gamma_p(\mu_1, \mu_2)$, then the curve of $\mathscr{P}_p(X)$, $\eta(t) := \mu_{1 \to 2}^t$ is a constant speed geodesic connecting μ_1 and μ_2 .

Conversely, for each constant speed geodesic $\gamma : [0,1] \rightarrow \mathscr{P}_p(X)$ connecting μ_1 and μ_2 , there exists $\mu \in \Gamma_p(\mu_1, \mu_2)$ such that $\gamma(t) = \mu_{1 \rightarrow 2}^t$, $\Gamma_p(\mu_1, \gamma(t)) = \{\mu_{1,1 \rightarrow 2}^t\}$ and $\Gamma_p(\gamma(t), \mu_2) = \{\mu_{1 \rightarrow 2,2}^t\}$, $t \in (0, 1)$.

Theorem 2.6 ([9, Theorem 7.3.2]). For arbitrary $\mu_1, \mu_2, \nu \in \mathscr{P}_2(X)$ and $\mu_{1\to 2} \in \Gamma_2(\mu_1, \mu_2)$,

$$(1 - \lambda)W_2(\mu_1, \nu) + \lambda W_2(\mu_2, \nu) - W_2(\mu_{1 \to 2}^{\lambda}, \nu) \leq \lambda (1 - \lambda) W_2^2(\mu_1, \mu_2),$$

i.e., $(\mathscr{P}_2(X), W_2)$ is a positively curved space (see [8, Definition 3.19]).

It can be shown that when (X, d) is a positively curved space, so is $(\mathscr{P}_2(X), W_2)$; However, it is worth noting that if (X, d) is a non-positively curved space, $(\mathscr{P}_2(X), W_2)$ is not necessarily non-positively curved ([8, Example 3.21]).

3. Generalized differential and semiconcavity of functionals of $\mathscr{P}_{c}(\mathbb{R}^{m})$

In the section, we assume $M = \mathbb{R}^m$ and we work on the space

$$\mathscr{P}_{c}(\mathbb{R}^{m}) := \{ \mu \in \mathscr{P}(\mathbb{R}^{m}) : \operatorname{supp}(\mu) \text{ is a compact subset of } \mathbb{R}^{m} \}.$$

It is obvious that $\mathscr{P}_c(\mathbb{R}^m) \subset \mathscr{P}_p(\mathbb{R}^m)$ for all $p \ge 1$. If *M* is a compact manifold, $\mathscr{P}_c(M) = \mathscr{P}(M)$. It is worth noting that for any $p \ge 1$, $(\mathscr{P}_c(\mathbb{R}^m), W_p)$ as a metric space, is not locally compact. See, for instance, the example in [9, Remark 7.1.9].

3.1. Generalized differential and semiconcavity. To deal with singularities of the functionals on Wasserstein spaces, we need some notions of \mathscr{P}_2 -subdifferential calculus in the setting of compact supported measures, developed in [9, Section 9.1, 9.2 & 10.3] (see also the seminal work [72]). We follow the setting in [21]. We write a functional $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ to mean the restriction of an extended real-valued functional $U : \mathscr{P}_2(\mathbb{R}^m) \to \mathbb{R} \cup \{\pm \infty\}$ such that U is finite valued on $\mathscr{P}_c(\mathbb{R}^m)$. Given $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ and R > 0, we call $B_R(\mu) := \bigcup_{x \in \text{supp}(\mu)} B(x, R)$ the *R*-fattening of supp (μ).

Definition 3.1 (Localized Fréchet generalized differential). Assume that $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$, $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, we say $\alpha \in L^2(\mathbb{R}^m;\mu)$ belongs to $\partial^+ U(\mu) = \partial^+_{loc} U(\mu)$, the set of *localized Fréchet* superdifferential of U at μ , if for any R > 0 and any $\nu \in \mathscr{P}(B_R(\mu))$,

$$U(\nu) - U(\mu) \leq \sup_{\gamma \in \Gamma_2(\mu,\nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma \right\} + o_R(W_2(\mu,\nu)), \tag{15}$$

where $o_R(W_2(\mu, \nu))$, depending on R, represents the high order infinitesimal of $W_2(\mu, \nu)$.

Similarly, $\beta \in L^2(\mathbb{R}^m; \mu)$ belongs to $\partial^- U(\mu) = \partial^-_{loc} U(\mu)$, the set of *localized Fréchet subdifferential* of U at μ , if for any R > 0 and any $\nu \in \mathscr{P}(B_R(\mu))$,

$$U(\nu) - U(\mu) \ge \inf_{\gamma \in \Gamma_2(\mu, \nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \beta(x), y - x \rangle \, d\gamma \right\} + o_R(W_2(\mu, \nu)). \tag{16}$$

For a time dependent functional $U : \mathbb{R} \times \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$, we say $(q, \alpha) \in \mathbb{R} \times L^2(\mathbb{R}^m; \mu) \in \partial^+ U(t, \mu)$, if for any R > 0 and $a, b \in \mathbb{R}$ satisfying $t \in (a, b)$ and $(s, \nu) \in [a, b] \times \mathscr{P}(B_R(\mu))$,

$$U(s,\nu) - U(t,\mu) \leq \sup_{\gamma \in \Gamma_2(\mu,\nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma \right\} + q(s-t) + o_R(W_2(\mu,\nu) + |t-s|).$$

The definition of $\partial^- U(t, \mu)$ is similar.

Remark 3.1. We regard two elements in $\partial^{\pm} U(\mu)$ to be the same if they coincide up to a μ -negligible set.

Proposition 3.1. Assume functionals $U, V : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ and $U \leq V$. If there exists $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ such that $U(\mu) = V(\mu)$, we have $\partial^+ V(\mu) \subset \partial^+ U(\mu)$ and $\partial^- U(\mu) \subset \partial^- V(\mu)$.

Proof. For any $\alpha \in \partial^+ V(\mu)$, due to (15) and the relation $U \leq V$, we get

$$\begin{split} U(\nu) - U(\mu) &\leq V(\nu) - V(\mu) \\ &\leq \sup_{\gamma \in \Gamma_2(\mu,\nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma \right\} + o_R(W_2(\mu,\nu)). \end{split}$$

This implies that $\alpha \in \partial^+ U(\mu)$. One can show that $\partial^- U(\mu) \subset \partial^- V(\mu)$ in a similar way.

Definition 3.2 (Singularity of functional in $\mathscr{P}_c(\mathbb{R}^m)$). Let $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$.

- (1) μ is called a *regular point* of U if both $\partial^+ U(\mu)$ and $\partial^- U(\mu)$ are non-empty. We also say U is *locally differentiable* at μ .
- (2) μ is called a *singular point* of U if U is not locally differentiable at μ . We denote by $\mathscr{S}(U)$ the set of all singular points of U.

Lemma 3.1. Suppose $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, $f \in C_c^{\infty}(\mathbb{R}^m)$ and R > 0. There exists $\delta_{f,R} > 0$ such that $v_t := (\mathrm{id} + tDf)_{\#} \mu \in \mathscr{P}(B_R(\mu))$ and

$$\Gamma_2(\mu, \nu_t) = \{ [\operatorname{id} \times (\operatorname{id} + tDf)]_{\#} \mu \}, \qquad \forall t \in [0, \delta_{f,R}].$$

Proposition 3.2. Suppose $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ is locally differentiable at μ , then $\partial^+ U(\mu)$ and $\partial^- U(\mu)$ are singleton. In this case, we denote

$$\partial^+ U(\mu) = \partial^- U(\mu) = \{\partial U(\mu)\}$$

We provide proofs of Lemma 3.1 and Proposition 3.2 in the appendix.

Definition 3.3 ([21], Semiconcavity of functionals in $\mathscr{P}_c(\mathbb{R}^m)$). $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ is *locally* (*geodesically*) *semiconcave* with constant C_R , if for any fixed R > 0, there exists $C_R > 0$, such that for all $\mu_1, \mu_2 \in \mathscr{P}(B(0, R))$ and $\gamma \in \Gamma_2(\mu_1, \mu_2)$ and $t \in [0, 1]$,

$$(1-t)U(\mu_1) + tU(\mu_2) - U(\gamma_{1\to 2}^t) \leq C_R t(1-t)W_2^2(\mu_1,\mu_2),$$

where $\gamma_{1\to 2}^t$ is given by (12). We say $U : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ is *locally strongly semiconcave*, if for any $\mu_1, \mu_2 \in \mathscr{P}(B(0, R))$ and $\gamma \in \Gamma(\mu_1, \mu_2)$,

$$(1-t)U(\mu_1) + tU(\mu_2) - U(\gamma_{1\to 2}^t) \le C_R t(1-t)W_{2,\gamma}^2(\mu_1,\mu_2)$$

for each $t \in [0, 1]$, where $\gamma_{1 \to 2}^t$ is defined in (12), $W_{2,\gamma}^2(\mu_1, \mu_2) := \int_{\mathbb{R}^{2m}} d^2(x, y) d\gamma$.

Theorem 2.6 implies 2-Wasserstein metric W_2 is globally semiconcave in $\mathscr{P}_2(\mathbb{R}^m)$.

Remark 3.2. For time-dependent functional $U : \mathbb{R} \times \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$, we only need to restrict to $[a, b] \times B(0, R)$, where $a, b \in \mathbb{R}$ and R > 0. We can similarly define the semiconcavity of functionals of $\mathbb{R} \times \mathscr{P}_c(\mathbb{R}^m)$.

3.2. Potential energy functional.

Definition 3.4 ([9], potential energy). Given a Borel measurable function $\phi : \mathbb{R}^m \to [-\infty, +\infty]$. A functional of $\mathscr{P}(\mathbb{R}^m)$ induced by ϕ

$$\Phi(\mu) := \int_{\mathbb{R}^m} \phi(x) \, d\mu \qquad (\mu \in \mathscr{P}(\mathbb{R}^m))$$

is called *potential energy functional of* ϕ . We denote $\Phi(\cdot)$ by $\phi(\cdot)$ for brevity if there is no confusion.

The following properties on generalized differential and semiconcavity for potential energy functional are useful.

Proposition 3.3. If $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R}^m)$, then the potential energy functional $\phi(\cdot) : \mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$ is locally strongly semiconcave.

Proof. Let R > 0, $\mu_1, \mu_2 \in \mathscr{P}(B(0, R))$ and $\gamma \in \Gamma(\mu_1, \mu_2)$. Then

$$(1-t)\phi(\mu_{1}) + t\phi(\mu_{2}) - \phi(\gamma_{1\to 2}^{t})$$

= $\int_{\mathbb{R}^{2m}} (1-t)\phi(x) + t\phi(y) - \phi((1-t)x + ty)d\gamma$
 $\leq \int_{\mathbb{R}^{2m}} C_{R}t(1-t)|x-y|^{2} d\mu = C_{R}t(1-t)W_{2,\gamma}^{2}(\mu_{1},\mu_{2}).$

Thus, $\phi(\cdot)$ is locally strongly semiconcave by Definition 3.3.

The following proposition is firstly claimed in [7], which discussed analogous conclusions on convex analysis in $\mathscr{P}_2(\mathbb{R}^m)$. We afford a proof in the appendix.

Proposition 3.4. Suppose $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R}^m)$ and $\alpha \in L^2(\mathbb{R}^m; \mu)$. Then for the potential energy functional $\phi(\cdot)$, the following statements are equivalent:

- (1) $\alpha \in \partial^+ \phi(\mu)$;
- (2) For arbitrary R > 0 and $v \in \mathscr{P}(B_R(\mu))$,

$$\phi(\nu) - \phi(\mu) \leq \inf_{\gamma \in \Gamma_2(\mu,\nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma \right\} + o_R(W_2(\mu,\nu)); \tag{17}$$

(3) For any R > 0 and $v \in \mathscr{P}(B(0, R))$, there exists $C_R > 0$, such that

$$\phi(\nu) - \phi(\mu) \leq \inf_{\gamma \in \Gamma_2(\mu,\nu)} \left\{ \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma \right\} + C_R W_2^2(\mu,\nu). \tag{18}$$

Generalized differential of functionals in $\mathscr{P}_p(X)$ was studied in literature [9, 8]. We restate the relevant conclusions in $\mathscr{P}_c(\mathbb{R}^m)$ and give a proof for the sake of rigorousness and completeness.

Theorem 3.1. Suppose $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R}^m)$ and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$. Then $\alpha \in \partial^+ \phi(\mu)$ if and only if $\alpha(x) \in D^+ \phi(x)$ holds true for μ -a.e. $x \in \mathbb{R}^m$.

See the appendix for the proof of Theorem 3.1. For the potential energy functional $\phi(\cdot)$ (i.e., $\Phi(\cdot)$) with ϕ a semiconcave function, Theorem 3.1 clarifies the relation between $\partial^+\phi$ and $D^+\phi$. This is a key point allowing us to study the singularities of potential energy functional $\phi(\cdot)$ by means of the singularities of the semiconcave function ϕ .

Corollary 3.1. Let $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R}^m)$ and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$. Then $\partial^+\phi(\mu) \neq \emptyset$ and $\phi(\cdot)$ is locally differentiable at μ if and only if $\partial^+\phi(\mu)$ is a singleton. Furthermore, $\alpha = \partial\phi(\mu)$ if and only if $\alpha(x) = D\phi(x)$ holds for μ -a.e. $x \in \mathbb{R}^m$.

Proof. For the first assertion, we note that $D^+\phi(x) \neq \emptyset$ for all $x \in \mathbb{R}^m$. If $\phi(\cdot)$ is locally differentiable at μ , by Proposition 3.2, $\partial^+\phi(\mu)$ is a singleton. Conversely, if $\partial^+\phi(\mu)$ is a singleton, we can find a μ -full-measure set E such that for $x \in E$, $D^+\phi(x) = \{\alpha(x)\}$. This implies ϕ is differentiable at $x, D^-\phi(x) = \{\alpha(x)\}$ and $\alpha \in \partial^-\phi(\mu)$. Then Proposition 3.2 shows that $\phi(\cdot)$ is localized differentiable at μ . The rest of proof is obvious.

Theorem 3.2. Let $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R}^m)$. Then $\mu \in \mathscr{S}(\phi(\cdot))$, i.e., μ is a singular point of $\phi(\cdot)$: $\mathscr{P}_c(\mathbb{R}^m) \to \mathbb{R}$, if and only if $\mu(\text{Sing}(\phi)) > 0$, where $\text{Sing}(\phi)$ denotes the set of singularities of ϕ . Consequently, $\mu \notin \mathscr{S}(\phi(\cdot))$ if $\mu \ll \mathscr{L}^m$, meaning that μ is absolutely continuous with respect to the Lebesgue measure \mathscr{L}^m .

Proof. If $\mu(\text{Sing}(\phi)) = 0$, there exists E with $\mu(E) = 1$ such that ϕ is differentiable at x for all $x \in E$. In other words, $D\phi \in \partial^+\phi(\mu) \cap \partial^-\phi(\mu)$ and $\phi(\cdot)$ is locally differentiable at μ . If $\mu(\text{Sing}(\phi)) > 0$, then one can find $\alpha_1, \alpha_2 \in \partial^+\phi(\mu)$ which are distinct by removing a μ -null set. It yields μ is the singular point of $\phi(\cdot)$.

Finally, recall that Sing (ϕ) is an (m - 1)-countably rectifiable set (see [32, Corollary 4.1.13]) and $\mathscr{L}^m(\text{Sing}(\phi)) = 0$. If $\mu \ll \mathscr{L}^m$, then $\mu(\text{Sing}(\phi)) = 0$ and this implies μ is a regular point of $\phi(\cdot)$.

We need also consider the time-dependent potential energy functionals. Suppose $\phi : \mathbb{R} \times \mathbb{R}^m \to [-\infty, +\infty]$ is Borel measurable. The potential energy functional of $\mathscr{P}_c(\mathbb{R}^m)$ induced by ϕ is

$$\Phi(t,\mu) := \int_{\mathbb{R}^m} \phi(t,x) \, d\mu = \int_{\mathbb{R}^{m+1}} \phi(r,x) \, d(\delta_t \times \mu), \quad \forall (t,\mu) \in \mathbb{R} \times \mathscr{P}_c(\mathbb{R}^m)$$

Again, we write $\Phi(t, \mu) = \phi(t, \mu)$ for the sake of brevity if without confusion.

Remark 3.3. Assume that $\phi \in \text{SCL}_{\text{loc}}(\mathbb{R} \times \mathbb{R}^m)$ and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$.

- (1) According to Definition 3.1 and Theorem 3.1, $(q, \alpha) \in \partial^+ \phi(t, \mu)$ if and only if $(q, \alpha(x)) \in D^+ \phi(t, x)$ for μ -a.e. $x \in \mathbb{R}^m$;
- (2) Similar to Corollary 3.1, $\partial^+ \phi(t,\mu) \neq \emptyset$, and $\phi(\cdot, \cdot)$ is locally differentiable at (t,μ) if and only if $\partial^+ \phi(t,\mu)$ is a singleton, and $(q,\alpha) = \partial \phi(t,\mu)$ if and only if $(q,\alpha(x)) = (D_t\phi(t,x), D_x\phi(t,x))$ for μ -a.e. $x \in \mathbb{R}^m$.

3.3. Dynamical cost functionals.

Definition 3.5 (Dynamical cost functional). The dynamical cost functional of $c^t(\cdot, \cdot)$ is defined as

$$C^{t}(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \int_{M \times M} c^{t}(x, y) \, d\gamma$$

We denote by $\Gamma_{\alpha}^{t}(\mu, \nu)$ the set of minimizers of $C^{t}(\mu, \nu)$.

Now we consider the case that *L* is a Tonelli Lagrangian and the time-dependent cost function $c^t(x, y) = A_t(x, y)$ is the fundamental solution with respect to *L*. In this context, a connection from optimal transportation to Mather theory and weak KAM theory is firstly studied by Bernard and Buffoni in [16].

Definition 3.6 (Random curves). For a given probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and T > 0, An absolutely continuous *random curve* on [0, T] refers to a measurable map $\xi : [0, T] \times \Omega \to M$, which satisfies for any fixed $\omega \in \Omega$, sample path $\xi(\cdot, \omega)$ is absolutely continuous curve on M. Moreover, $\xi = \xi(s)$ can also be treated as a random process for $s \in [0, T]$.

For the setting on work space $\mathscr{P}_c(\mathbb{R}^m)$, We select the probability measure space $(\Omega, \mathscr{F}, \mathbb{P})$ for the random curves according to Remark 2.2. In the subsequent discussion, we usually omit the component $\omega \in \Omega$ of the random curves and denote the expectation of a random variable (or vector) on $(\Omega, \mathscr{F}, \mathbb{P})$ by \mathbb{E} for simplicity when no confusion arises.

Definition 3.7 (Dynamical coupling). We say an absolutely continuous random curve ξ is a *dynamical coupling* of μ and ν for t > 0, if law ($\xi(0)$) = μ and law ($\xi(t)$) = ν . The set of all dynamical couplings of μ and ν for t > 0 is denoted by $L_{u,v}^t$.

Proposition 3.5 ([16, 80]). For any $v_1, v_2 \in \mathscr{P}(\mathbb{R}^m)$ such that $C^t(v_1, v_2)$ is finite,

$$C^{t}(\nu_{1},\nu_{2}) = \min_{\xi \in L^{t}_{\nu_{1},\nu_{2}}} \mathbb{E}\left(\int_{0}^{t} L(\xi(s),\dot{\xi}(s)) \, ds\right).$$

The minimizer of the last term is called dynamical optimal coupling of v_1 and v_2 for t > 0. $\mu_s := \text{law}(\xi_*(s)) (s \in [0, t])$ as the law of a dynamical optimal coupling ξ_* is called a displacement interpolation of $C^t(v_1, v_2)$. In this case, we have

- ξ_* is a random solution of the Euler-Lagrange equation;
- the path $\{\mu_s\}_{s \in [0,t]}$ is a minimizing curve for the action functional defined on $\mathscr{P}(\mathbb{R}^m)$ by

$$\mathbb{A}^{t}(\{\mu_{s}\}_{s\in[0,t]}) := \min_{\xi} \mathbb{E}\left(\int_{0}^{t} L(\xi(s),\dot{\xi}(s)) \, ds\right),$$

where the last minimum is over all random curves ξ such that law $(\xi(s)) = \mu_s$ for $s \in [0, t]$.

For more discussions on dynamical optimal coupling and displacement interpolation, readers can refer to Chapter 7 in [80].

Corollary 3.2. Suppose $\mu, \nu \in \mathscr{P}_c(\mathbb{R}^m)$ and t > 0.

- (1) There exists a compact set $K_{\mu,\nu,t} \subset \mathbb{R}^m$ depending on μ , ν and t, such that for any displacement interpolation $\{\mu_s\}_{s\in[0,t]}$ of $C^t(\mu,\nu)$, $\operatorname{supp}(\mu_s) \subset K_{\mu,\nu,t}$ for all $s \in [0,t]$. Thus, $\{\mu_s\}_{s\in[0,t]} \subset \mathscr{P}_c(\mathbb{R}^m)$;
- (2) $\{\mu_s\}_{s \in [0,t]} \in \text{Lip}([0,t]; \mathscr{P}_c(\mathbb{R}^m)).$

Proof. Every displacement interpolation $\{\mu_s\}_{s\in[0,t]}$ of $C^t(\mu, \nu)$ is the law of some random actionminimizing curve ξ at times s. Invoking Proposition 2.6, we have that $\{\xi(\cdot, \omega)\}_{\omega\in\Omega}$ is equi-Lipschitz, and for each $\omega \in \Omega$, Lip $(\xi(\cdot, \omega)) \leq F(|\xi(0, \omega) - \xi(t, \omega)|/t)$, where F is a non-decreasing function given in Proposition 2.6. Since $\mu = \text{law}(\xi(0))$ and $\nu = \text{law}(\xi(t))$, the equi-Lipschitz constant Lip (ξ) of $\{\xi(\cdot, \omega)\}_{\omega\in\Omega}$ is determined by μ, ν and t > 0. Due to the compactness of supp (μ) and supp (ν) , supp $(\mu_s) \subset B_{\text{Lip}(\xi)t}(\mu) =: K_{\mu,\nu,t}$ for all $s \in [0, t]$.

Now we turn to show the Lipschitz property of $\{\mu_s\}_{s \in [0,t]}$. Since $\mu_s = \text{law}(\xi(s))$ for $s \in [0, t]$, we have

$$W_2(\mu_s, \mu_{s'}) \leq \mathbb{E}(d^2(\xi(s), \xi(s')))^{1/2} \leq \operatorname{Lip}(\xi)|s-s'|, \quad \forall s, s' \in [0, t].$$

It means $\{\mu_s\}_{s\in[0,t]}$ is a Lipschitz curve in $\mathscr{P}_c(\mathbb{R}^m) \subset \mathscr{P}_2(\mathbb{R}^m)$.

The following essentially known results are the technical support of the study of dynamical cost functionals.

Proposition 3.6 ([5, Theorem 2.4], disintegration of measures). Let X, Y be polish spaces, $\mu \in \mathscr{P}(X)$ and $f: X \to Y$ be a Borel function. Then there exists a family $\{\mu_y\}_{y \in Y} \subset \mathscr{P}(X)$ such that

(1) $y \mapsto \mu_y$ is Borel, i.e. $y \mapsto \mu_y(A)$ is Borel for any Borel set $A \subset X$;

(2) $\mu = \int_{Y} \mu_{y} d(f_{\#}\mu)$, *i.e. for any Borel subset* $A \subset X$, $\mu(A) = \int_{Y} \mu_{y}(A) d(f_{\#}\mu)$;

(3) μ_y is concentrated on $f^{-1}(y)$ for $(f_{\#}\mu)$ -a.e. $y \in Y$.

Any other family $\{\mu'_{v}\}_{v \in Y} \subset \mathscr{P}(X)$ with these properties satisfies $\mu'_{v} = \mu_{v}$ for $(f_{\#}\mu)$ -a.e. $v \in Y$.

Particularly, let $X = X_1 \times X_2$, $Y = X_i$ and $f = \pi_i$, i = 1, 2. For $\mu \in \mathscr{P}(X_1 \times X_2)$, we say $\mu = \int_{X_1} \mu_{x_1} d(\pi_1) \# \mu$ and $\mu = \int_{X_2} \mu_{x_2} d(\pi_2) \# \mu$ are the disintegrations of μ with respect to $(\pi_1) \# \mu$ and $(\pi_2) \# \mu$ (or X_1 and X_2), respectively.

Definition 3.8 (Gluing measure). Let $\mu_i \in \mathscr{P}(X_i)$ for i = 1, 2, 3. Set $\gamma \in \Gamma(\mu_1, \mu_2)$ and $\gamma' \in \Gamma(\mu_1, \mu_3)$. Then we denote

$$\gamma * \gamma' := \int_{X_1} \gamma_{x_1} \times \gamma'_{x_1} d\mu_1 \in \Gamma(\mu_1, \mu_2, \mu_3) \subset \mathscr{P}(X_1 \times X_2 \times X_3)$$

as the gluing measure of γ and γ' (with respect to μ_1), where $\gamma = \int_{X_1} \gamma_{x_1} d\mu_1$ and $\gamma' = \int_{X_1} \gamma'_{x_1} d\mu_1$ are the disintegration of γ and γ' with respect to μ_1 respectively.

Theorem 3.3. Suppose t > 0, $K \subset \mathbb{R}^m$ is a convex and compact subset.

- (1) $C^{t}(\mu, \cdot)$ and $C^{t}(\cdot, \mu)$ are superlinear on $\mathscr{P}_{c}(\mathbb{R}^{m})$ endowed with W_{1} metric, where $\mu \in \mathscr{P}_{c}(\mathbb{R}^{m})$;
- (2) $(t, v) \mapsto C^{t}(\mu, v) \in \operatorname{Lip}([a, b] \times \mathscr{P}(K)), where \mu \in \mathscr{P}(K);$
- (3) for $t \in (0, 1)$, there exists $C_K > 0$ depends on K, such that for any $\mu \in \mathscr{P}(K)$, $C^t(\mu, \cdot)$ is locally strongly semiconcave on $\mathscr{P}(K)$ with constant $\frac{C_K}{t}$;

See the appendix for the proof of Theorem 3.3.

Theorem 3.4 (Generalized superdifferential of $C^t(\mu, \nu)$). Define $\mu, \nu \in \mathscr{P}_c(\mathbb{R}^m)$, t > 0 and $\gamma \in \Gamma_o^t(\mu, \nu)$, as well as $\gamma = \int_{\mathbb{R}^m} \gamma_x d\mu(x) = \int_{\mathbb{R}^m} \gamma_y d\nu(y)$ are the disintegrations of γ with respect to μ and ν respectively. Moreover, we choose $\eta_{x,y} \in \Gamma_{x,y}^t$, the minimizer of $A_t(x, y)$, which is measurable with respect to $(x, y) \in \text{supp}(\mu) \times \text{supp}(\nu)^2$. Then

$$\mathbf{p}_{\nu}(y) := \int_{\mathbb{R}^m} p_{x,y}(t) \, d\gamma_y \in \partial_{\nu}^+ C^t(\mu, \nu),$$

$$-\mathbf{p}_{\mu}(x) := \int_{\mathbb{R}^m} -p_{x,y}(0) \, d\gamma_x \in \partial_{\mu}^+ C^t(\mu, \nu),$$

$$-\int_{\mathbb{R}^{2m}} H(y, p_{x,y}(t)) \, d\gamma \in D_t^+ C^t(\mu, \nu),$$

where $p_{x,y}(s) := L_v(\eta_{x,y}(s), \dot{\eta}_{x,y}(s)), s \in [0, t].$

For the proof of Theorem 3.4, see the appendix.

Corollary 3.3. If $(t, v) \mapsto C^t(\mu, v)$ is locally differentiable, we have

$$D_t C^t(\mu, \nu) + \int_{\mathbb{R}^m} H(y, \partial_\nu C^t(\mu, \nu)) \, d\nu \leq 0.$$

The strict inequality holds in general.

Proof. If $(t, v) \mapsto C^{t}(\mu, v)$ is localized differentiable, according to Theorem 3.4,

$$D_t C^t(\mu, \nu) = -\int_{\mathbb{R}^{2m}} H(y, p_{x,y}(t)) \, d\gamma, \qquad \partial_\nu C^t(\mu, \nu) = \int_{\mathbb{R}^m} p_{x,y}(t) \, d\gamma_y.$$

Due to the fact that $H(x, \cdot)$ is strictly convex, Jensen's inequality implies

$$D_t C^t(\mu, \nu) + \int_{\mathbb{R}^m} H(y, \partial_\nu C^t(\mu, \nu)) \, d\nu$$

= $-\int_{\mathbb{R}^{2m}} H(y, p_{x,y}(t)) \, d\gamma + \int_{\mathbb{R}^m} H\left(y, \int_{\mathbb{R}^m} p_{x,y}(t) \, d\gamma_y\right) \, d\nu$
 $\leqslant -\int_{\mathbb{R}^{2m}} H(y, p_{x,y}(t)) \, d\gamma + \int_{\mathbb{R}^m} \int_{\mathbb{R}^m} H\left(y, p_{x,y}(t)\right) \, d\gamma_y d\nu = 0.$

²This can be guaranteed by standard measurable selection theorem (see [38]).

In general, the inequality above is strict if μ and ν are not Dirac measures.

Theorem 3.5. Given $\mu_1 \in \mathscr{P}_c(\mathbb{R}^m)$. Suppose $\lambda, R > 0$ with $\operatorname{supp}(\mu_1) \subset B(0, R)$ and let $t \in (0, t_{\lambda,R})$, where $t_{\lambda,R}$ is determined by Proposition 2.8. Then for all $\mu_2, \mu_3 \in \mathscr{P}(B(0, R))$ those satisfy the following condition: one can find $\gamma_{1,i} \in \Gamma_o^t(\mu_1, \mu_i)$, i = 2, 3 such that $|x_1 - x_i| \leq \lambda t$ holds true for $\gamma_{1,i}$ -a.e. (x_1, x_i) , we have

$$(1-s)C^{t}(\mu_{1},\mu_{2})+sC^{t}(\mu_{1},\mu_{3})-C^{t}(\mu_{1},\gamma_{2\to3}^{s}) \geq \frac{C_{\lambda,R}}{t}s(1-s)W_{2,\gamma}^{2}(\mu_{2},\mu_{3}),$$

where $\gamma := \gamma_{1,2} * \gamma_{1,3} \in \Gamma(\mu_1, \mu_2, \mu_3)$ and $W^2_{2,\gamma}(\mu_2, \mu_3) := \int_{\mathbb{R}^{3m}} |x_2 - x_3|^2 d\gamma$. In this case, we call $C^t(\mu_1, \cdot)$ locally generalized semiconvex with constant $\frac{C_{\lambda,R}}{t}$.

Proof. Taking $\mu_2, \mu_3 \in \mathscr{P}(B(0, R))$ satisfying the condition in the theorem above, then for all $s \in [0, 1], \gamma_{1, 2 \to 3}^s \in \Gamma(\mu_1, \gamma_{2 \to 3}^s)$ and

$$\begin{split} &(1-s)C^{t}(\mu_{1},\mu_{2})+sC^{t}(\mu_{1},\mu_{3})-C^{t}(\mu_{1},\gamma_{2\to3}^{s})\\ &\geqslant \int_{\mathbb{R}^{2m}}(1-s)A_{t}(x,y)\,d\gamma_{1,2}+\int_{\mathbb{R}^{2m}}sA_{t}(x,y)\,d\gamma_{1,3}-\int_{\mathbb{R}^{2m}}A_{t}(x,y)\,d\gamma_{1,2\to3}^{s}\\ &\geqslant \int_{\mathbb{R}^{3m}}(1-s)A_{t}(x_{1},x_{2})+sA_{t}(x_{1},x_{3})-A_{t}(x_{1},(1-s)x_{2}+sx_{3})\,d\gamma\\ &\geqslant \int_{\mathbb{R}^{3m}}\frac{C_{\lambda,R}}{t}s(1-s)|x_{2}-x_{3}|^{2}\,d\gamma\\ &=\frac{C_{\lambda,R}}{t}s(1-s)W_{2,\gamma}^{2}(\mu_{2},\mu_{3}). \end{split}$$

Here we used the semiconvexity result of fundamental solutions in Proposition 2.8.

4. RANDOM LAX-OLEINIK OPERATORS AND VISCOSITY SOLUTIONS

In this section, we define Lax-Oleinik operators in $\mathscr{P}(\mathbb{R}^m)$ and investigate its properties in $\mathscr{P}_c(\mathbb{R}^m)$. It can be seen that they inherit many characteristics of the classical Lax-Oleinik operators, so it can be proved that they are viscosity solutions of the corresponding Hamilton-Jacobi equation in $\mathscr{P}_c(\mathbb{R}^m)$. We highlight the approach of using the Hopf-Lax semigroup, as developed in previous works [68, 62], within the context of optimal transport.

4.1. Definitions and properties on Random Lax-Oleinik operators.

Definition 4.1 (Random Lax-Oleinik operators). For any Borel measurable function $\phi : \mathbb{R}^m \to [-\infty, +\infty], t > 0$ and $\mu \in \mathscr{P}(\mathbb{R}^m)$, we define

$$P_t^+\phi(\mu) := \sup_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{\phi(\nu) - C^t(\mu, \nu)\},\tag{19}$$

$$P_t^-\phi(\mu) := \inf_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{\phi(\nu) + C^t(\nu, \mu)\}.$$
(20)

 P_t^+ and P_t^- are called *positive* and *negative type Lax-Oleinik operators* respectively.

The following theorem and corollary show that $\mathscr{P}_c(\mathbb{R}^m)$ is a suitable working space for the operator P_t^{\pm} . Before proceeding, we introduce a lemma necessary for later discussion.

Lemma 4.1 ([37, Lemma 4.3]). Assume that X and Y are two Polish spaces and $\gamma \in \mathscr{P}(X \times Y)$.

(1) $\pi_X(\operatorname{supp}(\gamma)) \subset \operatorname{supp}((\pi_X)_{\#}\gamma);$

(2) $\pi_X(\operatorname{supp}(\gamma)) = \operatorname{supp}((\pi_X)_{\#}\gamma)$ if Y is compact.

Theorem 4.1. Suppose that t > 0, ϕ is lower semicontinuous on \mathbb{R}^m and (κ_1, κ_2) -Lipschitz in the large. Then the followings hold.

(1) For any $v \in \mathscr{P}_c(\mathbb{R}^m)$, there exists a minimizer $\mu = \mu(t, v) \in \mathscr{P}_c(\mathbb{R}^m)$ such that

$$P_t^-\phi(\nu) = \phi(\mu) + C^t(\mu, \nu).$$

- (2) $P_t^-\phi(v) = T_t^-\phi(v)$ is finite for all $v \in \mathscr{P}_c(\mathbb{R}^m)$. That is $P_t^-\phi$ is the potential energy functional induced by $T_t^-\phi$ on $\mathscr{P}_c(\mathbb{R}^m)$.
- (3) For $v \in \mathscr{P}_c(\mathbb{R}^m)$, each minimizer $\mu(t, v)$ is contained in $\mathscr{P}_c(\mathbb{R}^m)$ and makes (K⁻) hold true.
- (4) For any minimizer $\mu(t, v)$, there exist constants $C_1, C_2 > 0$ such that

$$W_1(\mu(t,\nu),\nu) \leq C_1 t + C_2,$$

where $C_1 = c_0 + \theta_0^*(\kappa_1 + 1) + \theta_1(0) + c_1$ and $C_2 = \kappa_2$, with θ_0^* the convex conjugate function of θ_0 .

Proof. Firstly, to prove part (1), we begin by noting that there exists a probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and a random variable $y_v : \Omega \to \mathbb{R}^m$ such that the law of y_v is v, by Remark 2.2. Leveraging the a priori estimates on the minimizer of $T_t^-\phi$ (refer to Lemma 2.7), we can express the marginal function as:

$$T_t^-\phi(y_\nu(\omega)) = \min_{x \in B(y_\nu(\omega), \lambda_\phi t + \kappa_2)} \{\phi(x) + A_t(x, y_\nu(\omega))\}, \quad \forall \omega \in \Omega,$$

where $\lambda_{\phi} > 0$ depends only on *L* and κ_1 . It's straightforward to verify that the map $\omega \mapsto B(y_{\nu}(\omega), \lambda_{\phi}t + \kappa_2)$ is a measurable and compact set-valued map. By invoking Theorem 18.19 from [4], we can admit a measurable selection $x_{t,\nu}$

$$\omega \mapsto x_{t,\nu}(\omega) \in \underset{x \in B(y_{\nu}(\omega), \lambda_{\phi}t + \kappa_{2})}{\operatorname{arg\,min}} \{\phi(x) + A_{t}(x, y_{\nu}(\omega))\} \subset B(y_{\nu}(\omega), \lambda_{\phi}t + \kappa_{2}).$$

Consequently, defining $\mu(t, \nu) := \text{law}(x_{t,\nu})$, we derive the inequality:

$$\int_{\mathbb{R}^m} T_t^- \phi(y) \, dv = \mathbb{E}(T_t^- \phi(y_v))$$

$$= \mathbb{E}(\phi(x_{t,v}) + A_t(x_{t,v}, y_v))$$

$$\geq \phi(\mu(t, v)) + C^t(\mu(t, v), v) \geq P_t^- \phi(v).$$
(21)

On the other hand, since $T_t^-\phi(y) \leq \phi(x) + A_t(x, y)$ for all $x, y \in \mathbb{R}^m$, then given $v \in \mathscr{P}_c(\mathbb{R}^m)$, for arbitrary $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ and $\gamma \in \Gamma(\mu, v)$, we have

$$\int_{\mathbb{R}^m} T_t^- \phi(y) \, d\nu = \int_{\mathbb{R}^{2m}} T_t^- \phi(y) \, d\gamma \leq \int_{\mathbb{R}^{2m}} \phi(x) + A_t(x, y) \, d\gamma$$

This implies that

$$T_t^-\phi(\nu) = \int_{\mathbb{R}^m} T_t^-\phi(y) \, d\nu \leq \inf_{\mu \in \mathscr{P}(\mathbb{R}^m)} \inf_{\gamma \in \Gamma(\mu,\nu)} \int_{\mathbb{R}^{2m}} \phi(x) + A_t(x,y) \, d\gamma = P_t^-\phi(\nu). \tag{22}$$

Combining (21) and (22) we find that

$$P_{t}^{-}\phi(\nu) = \phi(\mu(t,\nu)) + C^{t}(\mu(t,\nu),\nu)$$

22

Additionally, we claim that $\mu(t, v) \in \mathscr{P}_c(\mathbb{R}^m)$. Indeed, by Corollary 5.22 from [80] and the existence of random curves, we can find a dynamical optimal coupling $\xi \in L^t_{\mu,v}$ such that

$$T_t^-\phi(y_v) = \phi(x_{t,v}) + \int_0^t L(\xi(s), \dot{\xi}(s)) \, ds$$

Furthermore, by Proposition 2.6, for any $\omega \in \Omega$, $\xi(\cdot, \omega) \in \text{Lip}([0, t]; \mathbb{R}^m)$ with a Lipschitz constant dependent only on *t*, *L*, and (κ_1, κ_2) . Therefore, setting the uniform Lipschitz constant as $\text{Lip}(\xi)$, we have that

$$\sup (\mu(t, \nu)) = \sup (\operatorname{law} (\xi(0)))$$
$$\subset \bigcup_{\omega \in \Omega} B(\xi(t, \omega), \operatorname{Lip} (\xi)t) = \bigcup_{y \in \operatorname{supp} (\nu)} B(y, \operatorname{Lip} (\xi)t),$$

Conclusion (2) follows directly from (21) and (22).

Now, we turn to prove part (3). Let $\mu \in \arg \min_{\mathscr{P}(\mathbb{R}^m)} \{\phi(\cdot) + C^t(\cdot, \nu)\}$ and let $\gamma \in \Gamma_o^t(\mu, \nu)$. From part (2), it follows that

$$\int_{\mathbb{R}^m} T_t^- \phi(y) \, dv = P_t^- \phi(v) = \phi(\mu) + C^t(\mu, v) = \int_{\mathbb{R}^{2m}} \phi(x) + A_t(x, y) \, d\gamma$$

We also have that

$$T_t^-\phi(y) = \phi(x) + A_t(x, y), \quad \gamma - \text{a.e.} (x, y) \in \mathbb{R}^{2m}.$$
 (23)

This implies that the minimizer μ satisfies equation (K⁻). Furthermore, we can deduce that

$$T_t^-\phi(y) = \phi(x) + A_t(x, y), \quad \forall (x, y) \in \operatorname{supp}(\gamma).$$
(24)

In fact, by combining the lower semicontinuity of ϕ with the properties of $T_t^-\phi$, we can define the function

$$F_{\phi}(x, y) := \phi(x) + A_t(x, y) - T_t^- \phi(y)$$

which is a non-negative and lower semicontinuous function on \mathbb{R}^{2m} . If there were some $(x_0, y_0) \in$ supp (γ) such that $F_{\phi}(x_0, y_0) > 0$, then we could find some $R_0 > 0$ such that $F_{\phi}(x, y)$ is positive on the ball $B((x_0, y_0), R_0)$ with $\gamma(B((x_0, y_0), R_0)) > 0$, which would contradict equation (23). According to part (1) of Lemma 4.1, we know that

$$\pi_2(\operatorname{supp}(\gamma)) \subset \operatorname{supp}((\pi_2)_{\#}\gamma) = \operatorname{supp}(\nu), \tag{25}$$

For any fixed $y \in \mathbb{R}^m$ satisfying equation (23), by Lemma 2.7, when *x* satisfies the same equation, *x* must be in $B(y, \lambda_{\phi}t + \kappa_2)$. Due to the compact projection, we have:

$$\operatorname{supp}(\gamma) \subset \bigcup_{y \in \operatorname{supp}(\nu)} \{(z, y) : z \in B(y, \lambda_{\phi}t + \kappa_2)\}.$$

Additionally, from the compact projection, there exists a compact set $K \subset \mathbb{R}^m$ such that supp $(\gamma) \subset \mathbb{R}^m \times K$. Following from part (2) of Lemma 4.1, we get:

$$\sup p(\mu) = \sup ((\pi_1)_{\#}\gamma)$$

= $\pi_1(\operatorname{supp}(\gamma))$
 $\subset \pi_1\left(\bigcup_{y\in\operatorname{supp}(\nu)} \{(z, y) : z \in B(y, \lambda_{\phi}t + \kappa_2)\}\right) = \bigcup_{y\in\operatorname{supp}(\nu)} B(y, \lambda_{\phi}t + \kappa_2),$ (26)

which means that supp (μ) is compact and every minimizer of $P_t^-\phi(\nu)$ is contained in $\mathscr{P}_c(\mathbb{R}^m)$.

Finally, we aim to prove part (4). Since μ is the minimizer, we have $\phi(\mu) + C^t(\mu, \nu) \leq \phi(\nu) + C^t(\nu, \nu)$. Moreover, by Theorem 3.3 and the superlinearity of C^t , for any $k \in \mathbb{R}$:

$$C^{t}(\mu, \nu) \ge kW_{1}(\mu, \nu) - (c_{0} + \theta_{0}^{*}(k))t.$$

We also have:

$$kW_1(\mu,\nu) - (c_0 + \theta_0^*(k))t - C^t(\nu,\nu) \le C^t(\mu,\nu) - C^t(\nu,\nu) \le \phi(\nu) - \phi(\mu).$$
(27)

Suppose $\gamma \in \Gamma_1(\mu, \nu)$. Because ϕ is (κ_1, κ_2) -Lipschitz in the large, (27) yields:

$$\begin{split} kW_1(\mu,\nu) - (c_0 + \theta_0^*(k))t - C^t(\nu,\nu) &\leq \int_{\mathbb{R}^{2m}} \phi(y) - \phi(x) \, d\gamma \\ &\leq \int_{\mathbb{R}^{2m}} \kappa_1 |x - y| + \kappa_2 \, d\gamma \\ &= \kappa_1 W_1(\mu,\nu) + \kappa_2. \end{split}$$

Taking $k = \kappa_1 + 1$, we obtain from property (L2) of *L*:

$$\begin{split} W_1(\mu, \nu) &\leq (c_0 + \theta_0^*(\kappa_1 + 1))t + C^t(\nu, \nu) + \kappa_2 \\ &\leq (c_0 + \theta_0^*(\kappa_1 + 1))t + \int_{\mathbb{R}^m} A_t(y, y) \, d\nu + \kappa_2 \\ &\leq (c_0 + \theta_0^*(\kappa_1 + 1))t + \int_{\mathbb{R}^m} \int_0^t L(y, 0) \, ds \, d\nu + \kappa_2 \\ &\leq (c_0 + \theta_0^*(\kappa_1 + 1) + \theta_1(0) + c_1)t + \kappa_2. \end{split}$$

This completes the proof.

By similar reasoning, we can also derive the corresponding conclusions for the positive random Lax-Oleinik operator P_t^+ .

Corollary 4.1. Suppose t > 0, ϕ is upper semicontinuous on \mathbb{R}^m and (κ_1, κ_2) -Lipschitz in the large. *Then the followings hold.*

(1) For any $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, there exists a maximizer $\nu = \nu(t,\mu) \in \mathscr{P}_c(\mathbb{R}^m)$ such that

$$P_t^+\phi(\mu) = \phi(\nu) - C^t(\mu, \nu).$$

- (2) $P_t^+\phi(\mu) = T_t^+\phi(\mu)$ is finite for all $\mu \in \mathscr{P}_c(\mathbb{R}^m)$. That is $P_t^+\phi$ is the potential energy functional induced by $T_t^+\phi$ on $\mathscr{P}_c(\mathbb{R}^m)$.
- (3) For $\mu \in \mathscr{P}_{c}(\mathbb{R}^{m})$, each maximizer $v(t,\mu)$ is contained in $\mathscr{P}_{c}(\mathbb{R}^{m})$ and makes (K⁺) hold true.
- (4) For any maximizer $v(t, \mu)$, there exist constants $C_1, C_2 > 0$ such that

$$W_1(v(t,\mu),\mu) \le C_1 t + C_2,$$
 (28)

where $C_1 = c_0 + \theta_0^*(\kappa_1 + 1) + \theta_1(0) + c_1$ and $C_2 = \kappa_2$.

Proposition 4.1. Let $\phi \in UC(\mathbb{R}^m)$, $t_0 > 0$ and c[0] = 0. Random Lax-Oleinik operators have the following properties:

- (1) $\{P_t^-\}_{t>0}$ and $\{P_t^+\}_{t>0}$ are semigroups with respect to $t \in (0, +\infty)$;
- (2) $\lim_{t\to 0^+} P_t^- \phi = \lim_{t\to 0^+} P_t^+ \phi = \phi$, thus we can define $P_0^- \phi = P_0^+ \phi = \phi$;
- (3) for arbitrary $t, s \ge 0$, $P_t^- \circ P_s^+ \phi = T_t^- \circ T_s^+ \phi(\cdot)$; $P_t^+ \circ P_s^- \phi = T_t^- \circ T_s^- \phi(\cdot)$;

- (4) for any t > 0, $P_t^-\phi$ (resp. $-P_t^+\phi$) is locally (resp. strongly) semiconcave and $\{P_t^-\phi\}_{t \ge t_0}$ (resp. $\{-P_t^+\phi\}_{t \ge t_0}$) uniformly localized (resp. strongly) semiconcave;
- (5) $t \mapsto P_t^- \phi$ ($t \mapsto P_t^+ \phi$) uniformly continuous on $[0, +\infty)$;
- (6) suppose $K \subset \mathbb{R}^m$ is compact set. $(t,\mu) \mapsto P_t^-\phi(\mu) ((t,x) \mapsto P_t^+\phi(x))$ is continuous on $[0,+\infty) \times \mathscr{P}(K)$, locally Lipschitz on $(0,+\infty) \times \mathscr{P}(K)$ and equi-Lipschitz on $[t_0,+\infty) \times \mathscr{P}(K)$ with respect to ϕ .

Proof. It should be noted that the operators P_t^{\pm} represent the potential energies of the functionals induced by T_t^{\pm} on the space of compactly supported probability measures $\mathscr{P}_c(\mathbb{R}^m)$, as established in Theorem 4.1 and Corollary 4.1. Consequently, by drawing an analogy with the results presented in Proposition 2.3, we can readily deduce the aforementioned conclusions.

Proposition 4.2. The following statements are equivalent.

(1) ϕ is the subsolution to (HJ_s);

(2) $\phi(v) - \phi(\mu) \leq C^t(\mu, v) + c[0]t$ for all t > 0 and $\mu, v \in \mathscr{P}_c(\mathbb{R}^m)$;

- (3) For any $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, $[0, +\infty) \ni t \mapsto P_t^-\phi(\mu) + c[0]t$ is non-decreasing;
- (4) For any $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, $[0, +\infty) \ni t \mapsto P_t^+\phi(\mu) c[0]t$ is non-increasing.

Proof. According to [43], ϕ is a subsolution to equation (HJ_s) if and only if for any $x, y \in \mathbb{R}^m$ and t > 0, the inequality $\phi(y) - \phi(x) \leq A_t(x, y) + c[0]t$ holds. Then, for any fixed $\mu, \nu \in \mathscr{P}_c(\mathbb{R}^m)$ and $\gamma \in \Gamma_o(\mu, \nu)$, integrating both sides of this inequality over (x, y) with respect to γ yields equation (2). Conversely, if equation (2) holds, setting $\mu = \delta_x$ and $\nu = \delta_y$ implies that inequality (1) is satisfied. The equivalence among equations (2), (3), and (4) can be derived from Definition 4.1.

Building on Theorem 4.1 and Corollary 4.1, the following statement is a direct corollary of [28, Lemma 3.1].

Proposition 4.3. Suppose $\phi \in UC(\mathbb{R}^m)$ and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$.

- (1) For any $t \ge 0$, $P_t^- \circ P_t^+ \phi \ge \phi(\cdot)$ and $P_t^- \circ P_t^+ \phi(\mu) = \phi(\mu)$ if and only if there exists $v \in \mathscr{P}_c(\mathbb{R}^m)$ such that $P_t^+ \phi(v) = \phi(\mu) C^t(v, \mu)$.
- (2) For any $t \ge 0$, $P_t^+ \circ P_t^- \phi \le \phi(\cdot)$ and $P_t^+ \circ P_t^- \phi(\mu) = \phi(\mu)$ if and only if there exists $v \in \mathscr{P}_c(\mathbb{R}^m)$ such that $P_t^- \phi(v) = \phi(\mu) + C^t(\mu, v)$;
- (3) For any $t \ge 0$, $P_t^+ \circ P_t^- \circ P_t^+ \phi = P_t^+ \phi$ and $P_t^- \circ P_t^+ \circ P_t^- \phi = P_t^- \phi$.

4.2. Hamilton-Jacobi equations in $\mathscr{P}_{c}(\mathbb{R}^{m})$. Let $\phi \in C(\mathbb{R}^{m})$ be a function (κ_{1}, κ_{2}) -Lipschitz in the large. According to Theorem 2.1, as the value function of the Bolza problem, $u(t, x) := T_{t}^{-}\phi(x)$ is the viscosity solution to equation (HJ_{e}^{-}) with the initial condition ϕ . A natural question arises: whether $U(t, \mu) := P_{t}^{-}\phi(\mu)$ will be the viscosity solution to some corresponding form of the Hamilton-Jacobi equations.

Assume that $H \in C(\mathbb{R}^m \times \mathbb{R}^m)$ is a Tonelli Hamiltonian. We consider the following Hamilton-Jacobi equation:

$$\partial_t U(t,\mu) + \int_{\mathbb{R}^m} H(x,\partial_\mu U(t,\mu)(x)) \, d\mu = 0, \tag{PHJ}_e)$$

where $(t, \mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$.

Definition 4.2 (Viscosity solution of (PHJ_e)). We call the functional $U : \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$ a viscosity subsolution of (PHJ_e) , if for every $(t, \mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$,

$$q + \int_{\mathbb{R}^m} H(x, \alpha(x)) \, d\mu \le 0, \qquad \forall (q, \alpha) \in \partial^+ U(t, \mu);$$
⁽²⁹⁾

Similarly, $U : \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$ is called a *viscosity supersolution* of (PHJ_e) , if for every $(t, \mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m)$,

$$p + \int_{\mathbb{R}^m} H(x, \beta(x)) \, d\mu \ge 0, \quad \forall (p, \beta) \in \partial^- U(t, \mu).$$
(30)

If U is both a subsolution and supersolution to (PHJ_e) , we say U is a viscosity solution of (PHJ_e) .

In a more general and abstract frame, Gangbo, Nguyen and Tudorascu introduced concept of Hamilton-Jacobi equations in Wasserstein spaces in [56], which is further studied in [60] later by Gangbo and Tudorascu. Although, (PHJ_e) can be regarded as a special case of their work, it is closely related to main contents of this paper.

Theorem 4.2. Let ϕ be lower semicontinuous and (κ_1, κ_2) -Lipschitz in the large. Then $U(t, \mu) := P_t^- \phi(\mu)$ is a viscosity solution of the following Cauchy problem of (PHJ_e)

$$\begin{aligned} \partial_t U(t,\mu) + \int_{\mathbb{R}^m} H\left(x, \partial_\mu U(t,\mu)\right) d\mu &= 0, \quad (t,\mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m), \\ U(0,\mu) &= \phi(\mu), \qquad \qquad \mu \in \mathscr{P}_c(\mathbb{R}^m). \end{aligned}$$

Corollary 3.3 indicates that, unlike the fundamental solution $A_t(x, y)$, the dynamical cost functional $C^t(\mu, \nu)$ can only possibly be a (strict) viscosity subsolution in general. This is because it can be regarded as a convex combination of $A_t(x, y)$ based on measures μ and ν . However, under the action of the random Lax-Oleinik operators, which corresponds to the convolution with the potential energy functional, it becomes a viscosity solution.

Proof. Following Theorem 4.1, we can express $U(t, \mu)$ as follows:

$$U(t,\mu) := P_t^- \phi(\mu) = \int_{\mathbb{R}^m} T_t^- \phi(x) \, d\mu = \int_{\mathbb{R}^+ \times \mathbb{R}^m} u(r,x) \, d(\delta_t \times \mu).$$

It should be noted that for any fixed R > 0, $u(t, x) := T_t^- \phi(x)$ is a semiconcave function on the domain $(0, +\infty) \times B(0, R)$. When $\mu \in \mathscr{P}(B(0, R))$, according to Remark 3.3, $(q, \alpha) \in \partial^+ U(t, \mu)$ if and only if $(q, \alpha(x)) \in D^+ u(t, x)$ for almost every $x \in \mathbb{R}^m$ with respect to the measure μ . Furthermore, since u(t, x) is the viscosity solution to equation (HJ_e^-) , we have

$$q + H(x, \alpha(x)) \leq 0, \quad \mu - \text{a.e. } x \in \mathbb{R}^m.$$

By integrating over μ , we derive equation (29). Given the arbitrariness of $(q, \alpha) \in \partial^+ U(t, \mu)$, it follows that $U(t, \mu)$ is a viscosity subsolution to equation (PHJ_e).

If $(p,\beta) \in \partial^- U(t,\mu)$, then according to Remark 3.3, since $\partial^+ U(t,\mu) \cap \partial^- U(t,\mu) \neq \emptyset$, *u* is differentiable on a set of full measure with respect to $\mu \times \delta_t$, and we have

$$D^+u(t, x) \cap D^-u(t, x) = \{(p, \beta(x))\}, \quad \mu - a.e. \ x \in \mathbb{R}^m.$$

Given that u(t, x) is the viscosity supersolution, it follows that

$$p + H(x,\beta(x)) \ge 0, \quad \mu - \text{a.e. } x \in \mathbb{R}^m.$$

By integrating over μ , we obtain equation (30), which implies that $U(t,\mu)$ is the viscosity supersolution to (PHJ_e) . Consequently, $U(t,\mu)$ is the viscosity solution to (PHJ_e) . Furthermore, by Proposition 4.1, $U(0,\cdot) = \phi(\cdot)$, thus U is the solution to the Cauchy problem for (PHJ_e) .

Remark 4.1. If ϕ is upper semicontinuous and (κ_1, κ_2) -Lipschitz in the large, $V(t, \mu) := P_t^+ \phi(\mu)$ is the viscosity solution to the following Cauchy problem

$$\begin{cases} \partial_t V(t,\mu) - \int_{\mathbb{R}^m} H\left(x,\partial_\mu V(t,\mu)\right) d\mu = 0, & (t,\mu) \in \mathbb{R}^+ \times \mathscr{P}_c(\mathbb{R}^m), \\ V(0,\mu) = \phi(\mu), & \mu \in \mathscr{P}_c(\mathbb{R}^m). \end{cases}$$

Proposition 4.4. The following statements are equivalent.

- (1) *u* is a weak KAM solution (viscosity solution) of (HJ_s) on \mathbb{T}^m ;
- (2) For any $t \ge 0$, $T_t^- u + c[0]t = u$;
- (3) For any $t \ge 0$, $P_t^- u + c[0]t = u(\cdot)$;
- (4) For any $t \ge 0$, $P_t^- \circ P_t^+ u = u(\cdot)$.

In this case, $u(\cdot)$ is the viscosity solution of the corresponding stationary type of Hamilton-Jacobi equation

$$\int_{\mathbb{T}^m} H(x, \partial_{\mu} u(\mu)(x)) \, d\mu = c[0], \quad \mu \in \mathscr{P}(\mathbb{T}^m).$$
(PHJ_s)

Proof. The equivalence of conditions (1) and (2) is discussed in [28]. If condition (2) holds, for any $\mu \in \mathscr{P}(\mathbb{T}^m)$, we have

$$P_t^- u(\mu) + c[0]t = \int_{\mathbb{T}^m} T_t^- u + c[0]t \, d\mu = \int_{\mathbb{T}^m} u \, d\mu = u(\mu),$$

which implies condition (3). On the other hand, let $t \ge 0$ and $\mu \in \mathscr{P}(\mathbb{T}^m)$, then $P_t^- u(\mu) + c[0]t = u(\mu)$. By choosing $\mu = \delta_x$, we get $T_t^- u(x) + c[0]t = u(x)$. The arbitrariness of $x \in \mathbb{T}^m$ confirms that condition (2) is satisfied. Due to Theorem 2.2 and condition (4) of Proposition 4.1, condition (4) is equivalent to conditions (1)-(3).

For the final part of this proposition, we observe that $P_t^- u(\mu) + c[0]t = u(\mu)$ and since $P_t^- u(\mu)$ is the viscosity solution to equation (PHJ_e), it follows that $u(\cdot)$ is the viscosity solution to equation (PHJ_s).

A simple fact is: if *u* satisfies $u = T_t^- u + c[0]t$ for some $t \ge 0$, then $u = T_t^- \circ T_t^+ u$ for such *t*. As a direct consequence of this fact, we have the following corollary.

Corollary 4.2. Assume that $u \in \text{Lip}(\mathbb{R}^m) \cap \text{SCL}(\mathbb{R}^m)$ is a viscosity solution of (HJ_s) . Then for such u, we can also get $T_t^- \circ T_t^+ u = u$ for arbitrary $t \ge 0$. As a direct consequence,

$$P_t^- u + c[0]t = u(\cdot), \ P_t^- \circ P_t^+ u = u(\cdot), \ \forall t \ge 0.$$

5. Cut locus and singularity propagation

In Section 3.2, we identified an equivalent statement regarding the singular points of potential energy functionals induced by semiconcave functions. Subsequently, we introduce the concept of the cut locus of potential energy functionals induced by viscosity solutions to Hamilton-Jacobi equations in $\mathscr{P}_c(\mathbb{R}^m)$, drawing inspiration from the definition of the cut locus of viscosity solutions. We then investigate the propagation of its singularities in the context of mass transport.

5.1. Cut locus and cut time function. In this subsection, we let $H : \mathbb{R}^{2m} \to \mathbb{R}$ be a Tonelli Hamiltonian and $u \in \text{Lip}(\mathbb{R}^m) \cap \text{SCL}(\mathbb{R}^m)$ be a viscosity solution of (HJ_s)

$$H(x, Du(x)) = c[0].$$

Theorem 2.2 provides the existence of such solutions.

Definition 5.1 (Calibrated curve). An absolutely continuous curve³ $\{\mu_s\}_{s \in I} \subset \mathscr{P}_c(\mathbb{R}^m)$ defined on some interval $I \subset \mathbb{R}$ is called a $(u(\cdot), C, c[0])$ -calibrated curve, if for any $a, b \in I, a \leq b$, we have

$$u(\mu_b) - u(\mu_a) = C^{b-a}(\mu_a, \mu_b) + c[0](b-a).$$

Remark 5.1. Suppose *u* is a viscosity solution. For $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, there is a $(u(\cdot), C, c[0])$ -calibrated curve ending at μ . This is a direct consequence of the following fact mentioned in [43]: for each $x \in \mathbb{R}^m$, there exists a (u, L, c[0])-calibrated curve $\eta \in C^2((-\infty, 0], \mathbb{R}^m)$ with $\eta(0) = x$ in \mathbb{R}^m . In other words, for any $a \le b \le 0$,

$$u(\eta(b)) - u(\eta(a)) = \int_a^b L(\eta(s), \dot{\eta}(s)) \, ds.$$

Define $\mu = \text{law}(x(\cdot))$. Then there exists a random curve $\xi = \xi(t) = \xi(t, \omega)$ such that $\xi(0, \omega) = x(\omega)$ and $\xi(\cdot, \omega) \in C^2((-\infty, 0], \mathbb{R}^m)$ is a calibrated curve for each $\omega \in \Omega$. Suppose $\mu_s := \text{law}(\xi(s))$, then $\{\mu_s\}_{s \in (-\infty, 0]}$ is we need.

Proposition 5.1. Given a $(u(\cdot), C, c[0])$ -calibrated curve $\{\mu_s\}_{s \in [a,b]} \subset \mathscr{P}_c(\mathbb{R}^m)$.

- (1) $\{\mu_s\}_{s \in [a,b]}$ is a displacement interpolation of $C^{b-a}(\mu_a,\mu_b)$;
- (2) for any subinterval $I' \subset [a, b]$, $\{\mu_s\}_{s \in I'}$ is also a $(u(\cdot), C, c[0])$ -calibrated curve.

Proof. According to Definition 5.1, for any $a \le t_1 < t_2 < t_3 \le b$, we have that

$$u(\mu_{t_2}) - u(\mu_{t_1}) = C^{t_2 - t_1}(\mu_{t_1}, \mu_{t_2}) + c[0](t_2 - t_1),$$

$$u(\mu_{t_3}) - u(\mu_{t_2}) = C^{t_3 - t_2}(\mu_{t_2}, \mu_{t_3}) + c[0](t_3 - t_2),$$

$$u(\mu_{t_3}) - u(\mu_{t_1}) = C^{t_3 - t_1}(\mu_{t_1}, \mu_{t_3}) + c[0](t_3 - t_1),$$

then $C^{t_3-t_1}(\mu_{t_1},\mu_{t_3}) = C^{t_2-t_1}(\mu_{t_1},\mu_{t_2}) + C^{t_3-t_2}(\mu_{t_2},\mu_{t_3})$. Due to the arbitrariness of t_1, t_2 and t_3 , $\{\mu_s\}_{s\in[a,b]}$ is a displacement interpolation of $C^{b-a}(\mu_a,\mu_b)$. The rest of proposition is an immediate consequence of Definition 5.1.

Definition 5.2 (Cut point). We call $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ a *cut point* of $u(\cdot)$, the potential energy functional induced by viscosity solution u, if all $(u(\cdot), C, c[0])$ -calibrated curves ending at μ are unable to extend forward still as $(u(\cdot), C, c[0])$ -calibrated curves. The set of all cut points of $u(\cdot)$ is denoted by $\mathscr{C}(u(\cdot))$, which is called the *cut locus* of functional $u(\cdot)$.

Lemma 5.1. $u(\cdot)$ is locally differentiable on the interior of $(u(\cdot), C, c[0])$ -calibrated curve. Furthermore, $\mathscr{S}(u(\cdot)) \subset \mathscr{C}(u(\cdot))$.

Proof. Let I = [a, b] and $\{\mu_s\}_{s \in I} \subset \mathscr{P}_c(\mathbb{R}^m)$ be a $(u(\cdot), C, c[0])$ -calibrated curve, where $-\infty \leq a < b < +\infty$. Then for any $c, d \in [a, b]$ with $c \leq d$, we have

$$u(\mu_d) - u(\mu_c) = C^{d-c}(\mu_c, \mu_d) + c[0](d-c).$$
(31)

³For the definition on absolutely continuous curves of metric space, readers can see [9, Definition 1.1.1].

According to Corollary 3.1, $\partial^+ u \neq \emptyset$ on $\mathscr{P}_c(\mathbb{R}^m)$. Thus, we only need to prove $\partial^- u(\mu_s) \neq \emptyset$ for all $s \in (a, b)$.

Note that *u* is obviously a subsolution, which means

$$u(y) \le u(x) + A_t(x, y) + c[0]t, \quad \forall x, y \in \mathbb{R}^m, \ t > 0.$$

Integrating the inequality above by $\gamma \in \Gamma_{\rho}^{b-s}(\mu, \mu_b)$ for any fixed $\mu \in \mathscr{P}_{c}(\mathbb{R}^m)$, we obtain

$$u(\mu_b) - u(\mu) \le C^{b-s}(\mu, \mu_b) + c[0](b-s).$$
(32)

We set d = b and c = s in (31), then

$$u(\mu_b) - u(\mu_s) = C^{b-s}(\mu_s, \mu_b) + c[0](b-s).$$
(33)

By combining equations (32) and (33), we derive

$$U(\mu) := u(\mu_s) - u(\mu) \le C^{b-s}(\mu, \mu_b) - C^{b-s}(\mu_s, \mu_b) =: V(\mu),$$

with equality when $\mu = \mu_s$. Using Proposition 3.1, we find that $\partial^+ V(\mu_s) \subset \partial^+ U(\mu_s)$, which implies

$$\partial^+ C^{b-s}(\mu,\mu_b)\Big|_{\mu=\mu_s} \subset \partial^+(-u(\mu_s)) = -\partial^- u(\mu_s).$$

According to Theorem 3.4, $-\mathbf{p}_{\mu_s} \in \partial^+ C^{b-s}(\mu_s, \mu_b) \subset -\partial^- u(\mu_s)$, hence $\partial^- u(\mu_s) \neq \emptyset$. Since $s \in (a, b)$ is arbitrary, $u(\cdot)$ is locally differentiable on the interior of the $(u(\cdot), C, c[0])$ -calibrated curve. Consequently, if $u(\cdot)$ is singular at μ , μ is not in the interior of the $(u(\cdot), C, c[0])$ -calibrated curve, indicating that the $(u(\cdot), C, c[0])$ -calibrated curve ending at μ cannot be extended further forward; that is, μ is the cut point of $u(\cdot)$.

Inspired by the concept of cut time function, we introduce the *cut time function of measure* with respect to the potential energy induced by viscosity solution *u*:

$$T_{u}(\mu) := \sup\{t \ge 0 : \exists v(\cdot) \in \operatorname{AC}([0, t]; \mathscr{P}_{c}(\mathbb{R}^{m})),$$

such that $u(v(t)) = u(\mu) + C^{t}(\mu, v(t)) + c[0]t\}.$
(34)

Then $\mu \in \mathscr{C}(u(\cdot))$ if and only if $T_u(\mu) = 0$.

Theorem 5.1. Assume that u is the viscosity solution to (HJ_s) on \mathbb{R}^m and $\mu \in \mathscr{P}_c(\mathbb{R}^m)$. Then

$$T_{u}(\mu) = \sup\{t \ge 0 : P_{t}^{-} \circ P_{t}^{+}u(\mu) = P_{t}^{+} \circ P_{t}^{-}u(\mu)\}$$
(35)

$$= \sup\{t \ge 0 : \mu(\{x \in \mathbb{R}^m : B_u(t, x) = 0\}) = 1\}$$
(36)

$$= \inf\{\tau_u(x) : x \in \operatorname{supp}(\mu)\}.$$
(37)

Proof. We first define the set on the right-hand side of equation (34) as $S(u,\mu)$. Note that u is a viscosity solution. Corollary 4.2 states that $P_t^- \circ P_t^+ u = u$ and $u = P_t^- u + c[0]t$ for any $t \ge 0$. Thus, t satisfies $P_t^- \circ P_t^+ u(\mu) = P_t^+ \circ P_t^- u(\mu)$ if and only if

$$0 = P_t^- \circ P_t^+ u(\mu) - P_t^+ \circ P_t^- u(\mu) = u(\mu) - P_t^+ u(\mu) + c[0]t.$$
(38)

According to the definition of P_t^+ , the Lipschitz property of displacement interpolation (Corollary 3.2) and Corollary 4.1, there exists a function $v(\cdot) \in AC([0, t]; \mathscr{P}_c(\mathbb{R}^m))$ such that

$$0 = u(\mu) - u(v(t)) + C^{t}(\mu, v(t)) + c[0]t,$$

which implies that equation (38) is equivalent to $t \in S(u, \mu)$, i.e., equality (35) holds.

Recall that the equality $P_t^- \circ P_t^+ u(\mu) - P_t^+ \circ P_t^- u(\mu) = 0$ can be expressed as

$$\int_{\mathbb{R}^m} T_t^- \circ T_t^+ u(x) - T_t^+ \circ T_t^- u(x) \, d\mu = 0.$$
(39)

According to [28, Lemma 3.1], for any $t \ge 0$, we have $T_t^- \circ T_t^+ u \ge u \ge T_t^+ \circ T_t^- u$. This implies that equation (39) holds if and only if for almost every $x \in \mathbb{R}^m$ with respect to μ , $B_u(t, x) = 0$. In other words, equality (36) is satisfied.

We now proceed to the proof of equation (37). When $t_0 < T_u(\mu)$, we have

$$\mu(\{x \in \mathbb{R}^m : B_u(t_0, x) = 0\}) = 1.$$

We assert that for any $x_0 \in \text{supp}(\mu)$, $B_u(t_0, x_0) = 0$. Indeed, if $B_u(t_0, x_0) > 0$, the lower semicontinuity of $B_u(t_0, \cdot)$ implies that there exists $R_0 > 0$ such that $B_u(t_0, x) > 0$ for all $x \in B(x_0, R_0)$. Since $x_0 \in \text{supp}(\mu)$, we have $\mu(B(x_0, R_0)) > 0$, which contradicts $\mu(\{x \in \mathbb{R}^m : B_u(t_0, x) = 0\}) = 1$. This assertion implies that $t_0 \leq \inf\{\tau_u(x) : x \in \text{supp}(\mu)\}$. By the arbitrariness of t_0 , we conclude that $T_u(\mu) \leq \inf\{\tau_u(x) : x \in \text{supp}(\mu)\}$.

On the other hand, if $T_u(\mu) < \inf\{\tau_u(x) : x \in \text{supp }(\mu)\}$, there exists some $t \in \mathbb{R}$ such that

$$T_u(\mu) < t < \inf\{\tau_u(x) : x \in \operatorname{supp}(\mu)\}.$$
(40)

For all $x \in \text{supp}(\mu)$, we have $t < \tau_u(x)$, which implies $B_u(t, x) = 0$. Therefore,

$$\operatorname{supp}(\mu) \subset \{x \in \mathbb{R}^m : B_u(t, x) = 0\},\$$

and it follows that

$$1 = \mu(\operatorname{supp}(\mu)) \leq \mu(\{x \in \mathbb{R}^m : B_u(t, x) = 0\}),$$

i.e., $T_u(\mu) \ge t$, but this contradicts equation (40). In conclusion, we have $T_u(\mu) = \inf\{\tau_u(x) : x \in \sup (\mu)\}$.

Following the concept of the Aubry set as discussed in Section 2.3, the Aubry set of the potential energy function $u(\cdot)$ is defined as the collection of all measures μ for which there exists a $(u(\cdot), C, c[0])$ -calibrated curve $v(\cdot) : [0, +\infty) \to \mathscr{P}(\mathbb{T}^m)$ such that $v(0) = \mu$. Consequently, a measure μ is in the Aubry set of $u(\cdot)$ if and only if $T_u(\mu) = +\infty$.

Corollary 5.1. Under the assumptions in Theorem 5.1,

- (1) *if* supp $(\mu) \cap \text{Cut}(u) \neq \emptyset$, then $\mu \in \mathscr{C}(u(\cdot))$;
- (2) if τ_u is continuous, then supp $(\mu) \cap Cut(u) \neq \emptyset$ if and only if $\mu \in \mathscr{C}(u(\cdot))$;
- (3) μ is in the Aubry set of $u(\cdot)$ if and only if supp $(\mu) \subset I(u)$;
- (4) if $\mu = \text{law}(Z)$ is the distribution of an \mathbb{R}^m -valued random variable $Z : \Omega \to \mathbb{R}^m$, then $\mu \in \mathscr{C}(u(\cdot))$ if and only if, for any $\varepsilon > 0$,

$$\mathbb{P}\left(\{\omega \in \Omega : \tau_u(Z(\omega)) < \varepsilon\}\right) > 0.$$

Proof. If the support of μ intersects the cut locus of u, i.e., $\operatorname{supp}(\mu) \cap \operatorname{Cut}(u) \neq \emptyset$, then there exists an $x \in \operatorname{supp}(\mu)$ such that $\tau_u(x) = 0$. This implies that $T_u(\mu) = 0$ by equation (37), and thus $\mu \in \mathscr{C}(u(\cdot))$. For the proof of part (2), we only need to consider the "if" direction. Since μ is a cut point of $u(\cdot)$, equation (37) gives us

$$T_u(\mu) = \inf\{\tau_u(x) : x \in \text{supp}(\mu)\} = 0.$$

Together with the continuity of τ_u , there exists an $x_0 \in \text{supp}(\mu)$ such that $\tau_u(x_0) = 0$, which means $x_0 \in \text{supp}(\mu) \cap \text{Cut}(u)$.

By using T_u , μ is in the Aubry set of $u(\cdot)$ if and only if $T_u(\mu) = +\infty$. Then (3) is a direct consequence.

We now turn to prove (4). Given any fixed $\varepsilon > 0$, we first note that $\tau_u(Z(\omega)) < \varepsilon$ if and only if $B_u(\varepsilon, Z(\omega)) > 0$ for $\omega \in \Omega$. Due to this fact, we have

$$\mathbb{P}(\{\omega : B_u(\varepsilon, Z(\omega)) > 0\}) = \mathbb{P}(\{\omega : \tau_u(Z(\omega)) < \varepsilon\}) > 0.$$

Therefore, we can deduce the following:

$$\mu(\{x \in \mathbb{R}^m : B_u(\varepsilon, x) = 0\}) = 1 - \mu(\{x \in \mathbb{R}^m : B_u(\varepsilon, x) > 0\})$$

$$\leq 1 - \mu(\{x \in \mathbb{R}^m : \exists \omega \in \Omega, Z(\omega) = x, B_u(\varepsilon, x) > 0\})$$

$$= 1 - \mathbb{P}(\{\omega \in \Omega : \tau_u(Z(\omega)) < \varepsilon\}) < 1.$$

This implies $T_u(\mu) \leq \varepsilon$ by (36). Since ε is arbitrary, we conclude that $T_u(\mu) = 0$. On the other hand, if there exists some $\varepsilon_0 > 0$ such that $\mathbb{P}(\{\omega \in \Omega : \tau_u(Z(\omega)) < \varepsilon_0\}) = 0$, i.e., $B_u(\varepsilon_0, Z(\omega)) = 0$ for \mathbb{P} -a.e. $\omega \in \Omega$, then we have

$$\mu(\{x \in \mathbb{R}^m : B_u(\varepsilon_0, x) = 0\}) \ge \mathbb{P}(\{\omega \in \Omega : B_u(\varepsilon_0, Z(\omega)) = 0\}) = 1,$$

which means $T_u(\mu) \ge \varepsilon_0 > 0$. This contradicts the assumption that $\mu \in \mathscr{C}(u(\cdot))$. This completes the proof.

Lemma 5.2. Suppose X, Y are Polish spaces, $\mu \in \mathscr{P}(X)$ and $f : X \to Y$ be a homeomorphism. Then supp $(f_{\#}\mu) = f(\text{supp }(\mu))$.

Proof. We denote the ball with radius $R \ge 0$ and center $x \in X$ (resp. $y \in Y$) in X (resp. Y) as $B_X(x, R)$ (resp. $B_Y(y, R)$). Given $y \in f(\text{supp }(\mu))$, there exists a unique $x \in \text{supp }(\mu)$ such that y = f(x). Moreover, for any R > 0, we can choose some r > 0 such that $f^{-1}(B_Y(y, R))$, as a neighborhood of x, contains $B_X(x, r)$. In this case,

$$f_{\#}\mu(B_Y(y,R)) = \mu(f^{-1}(B_Y(y,R))) \ge \mu(B_X(x,r)) > 0,$$

which implies that $y \in \text{supp}(f_{\#}\mu)$.

Conversely, let $y \in \text{supp}(f_{\#}\mu)$. There exists a unique $x \in X$ such that y = f(x). We need to verify that $x \in \text{supp}(\mu)$. For any R > 0, there exists some r > 0 such that $f(B_X(x, R)) \supset B_Y(y, r)$, which means $B_X(f^{-1}(y), R) \supset f^{-1}(B_Y(y, r))$. This implies

$$\mu(B_X(x,R)) = \mu(B_X(f^{-1}(y),R)) \ge \mu(f^{-1}(B_Y(y,r))) = f_{\#}\mu(B_Y(y,r)) > 0.$$

By the arbitrariness of R > 0, it follows that $x \in \text{supp}(\mu)$ and $y = f(x) \in f(\text{supp}(\mu))$.

Theorem 5.2. Given a viscosity solution u of (HJ_s) on \mathbb{R}^m . Suppose $\mu \in \mathscr{P}_c(\mathbb{R}^m)$ and $T_u(\mu) > 0$. Let $\mu_t := [\pi \circ \Phi_H^t(\cdot, Du(\cdot))]_{\#}\mu$ for $t \in [0, T_u(\mu)]$.

(1) $\{\mu_t\}_{t \in [0,T_u(\mu)]}$ is a $(u(\cdot), C, c[0])$ -calibrated curve;

- (2) $\mu = [\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))]_{\#}\mu_t \text{ for } t \in [0, T_u(\mu));$
- (3) If $t \in [0, T_u(\mu))$ and $\mu \ll \mathscr{L}^m$, then $\mu_t \ll \mathscr{L}^m$.

П

In this case, the expression $\pi \circ \Phi_{H}^{t}(\cdot, Du(\cdot)) : \mathbb{R}^{m} \to \mathbb{R}$ should be interpreted as follows: there exists a measurable set A with $\mu(A) = 1$ such that u is differentiable at each $x \in A$ and $\pi \circ \Phi_{H}^{t}(\cdot, Du(\cdot))$ is defined on A. One can extend the map $\pi \circ \Phi_{H}^{t}(\cdot, Du(\cdot))$ to \mathbb{R}^{m} as a map $\varphi_{t}(\cdot)$ such that $\varphi_{t}|_{\mathbb{R}^{m}\setminus A} = 0$. Since the set-valued map $x \rightsquigarrow D^{+}u$ is upper semicontinuous, $\pi \circ \Phi_{H}^{t}(\cdot, Du(\cdot)) \in C(A)$ and $\varphi_{t}(\cdot)$ is measurable. Therefore, $[\varphi_{t}]_{\#}\mu$ is well-defined. Similarly, $[\pi \circ \Phi_{H}^{-t}(\cdot, Du(\cdot))]_{\#}\mu_{t}$ is also well-defined.

Proof. Assume $T_u(\mu) > 0$. According to Theorem 5.1, for any $x \in \text{supp}(\mu)$, we have $\tau_u(x) \ge T_u(\mu) > 0$. By Proposition 2.4, for each $x \in \text{supp}(\mu)$, there exists a unique calibrated curve starting from x:

$$\varphi_t(x) = \pi \circ \Phi_H^t(x, Du(x)), \quad t \in [0, T_u(\mu)], \quad x \in \operatorname{supp}(\mu).$$

For any fixed $t \in [0, T_u(\mu)]$, the map

$$\varphi_t(\cdot) : \operatorname{supp}(\mu) \to \mathbb{R}^m$$

 $x \mapsto \pi \circ \Phi^t_H(x, Du(x))$

is well-defined on supp (μ). Moreover, since u is of class $C^{1,1}$ on the interior of the (u, L, c[0])calibrated curve, $\varphi_t(\cdot) \in \text{Lip}(\text{supp}(\mu); \mathbb{R}^m)$. Then, for any $0 \leq s < t \leq T_u(\mu)$, combining with Proposition 4.2, we obtain

$$C^{t-s}(\mu_s,\mu_t) + c[0](t-s) \ge u(\mu_t) - u(\mu_s)$$

=
$$\int_{\mathbb{T}^m} u(\varphi_t(x)) - u(\varphi_s(x)) \, d\mu$$

=
$$\int_{\mathbb{T}^m} A_{t-s}(\varphi_s(x),\varphi_t(x)) \, d\mu + c[0](t-s)$$

$$\ge C^{t-s}(\mu_s,\mu_t) + c[0](t-s),$$

which implies that $\{\mu_t\}_{t \in [0, T_u(\mu)]}$ is a $(u(\cdot), C, c[0])$ -calibrated curve.

Furthermore, by invoking Proposition 2.4, for any fixed $t \in [0, T_u(\mu))$, we have

$$\pi \circ \Phi_H^{-t}(\pi \circ \Phi_H^t(x, Du(x)), Du(\pi \circ \Phi_H^t(x, Du(x)))) = \pi \circ \Phi_H^{-t}(\varphi_t(x), Du(\varphi_t(x))) = x.$$

The map $\varphi_t(\cdot)$ is continuous and injective, and thus a bi-Lipschitz homeomorphism from supp (μ) onto its image for each $t \in [0, T_u(\mu))$, by the compactness of supp (μ) . That is, $\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))$ is the Lipschitz inverse of $\varphi_t(\cdot)$ on supp (μ_t) due to Lemma 5.2. It follows that

$$[\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))]_{\#}\mu_t = [\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))]_{\#}[\pi \circ \Phi_H^t(\cdot, Du(\cdot))]_{\#}\mu = \mu, \quad t \in [0, T_u(\mu)).$$

This proves part (2). Note that $\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))$ is Lipschitz and maps Lebesgue zero measure sets into Lebesgue zero measure sets. Consequently, if $\mu = [\pi \circ \Phi_H^{-t}(\cdot, Du(\cdot))]_{\#}\mu_t \ll \mathscr{L}^m$, then $\mu_t \ll \mathscr{L}^m$ for $t \in [0, T_u(\mu))$. This completes the proof of part (3).

5.2. Propagation of singularities.

Theorem 5.3. Assume that $\phi \in \text{Lip}(\mathbb{R}^m) \cap \text{SCL}_{\text{loc}}(\mathbb{R}^m)$. Then for any $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, there exists $t_{\phi,\mu} > 0$, depending on ϕ and μ , such that

$$\underset{\nu \in \mathscr{P}(\mathbb{R}^m)}{\arg \max\{\phi(\nu) - C^{t}(\mu, \nu)\}},$$

is a singleton for all $t \in (0, t_{\phi,\mu}]$. Define

$$\nu_{\phi,\mu}(t) := \begin{cases} \mu, & t = 0, \\ \arg\max_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{\phi(\cdot) - C^t(\mu, \cdot)\}, & t \in (0, t_{\phi,\mu}], \end{cases}$$

then $\lim_{t\to 0^+} v_{\phi,\mu}(t) = \mu$ in the sense of metric W_p for any fixed $p \ge 1$.

Remark 5.2. When considering the case of a compact manifold, for example, $M = \mathbb{T}^m$, the time $t_{\phi} = t_{\phi,\mu}$ mentioned in Theorem 5.3 can be independent of μ , being determined only by ϕ .

Proof. First, Theorem 4.1 asserts that if $\mu \in \mathscr{P}_c(\mathbb{R}^m)$,

$$\emptyset \neq \underset{\nu \in \mathscr{P}(\mathbb{R}^m)}{\operatorname{arg max}} \{ \phi(\nu) - C^t(\mu, \nu) \} \subset \mathscr{P}_c(\mathbb{R}^m).$$

Then, for any $\nu \in \arg \max_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{ \phi(\nu) - C^t(\mu, \nu) \}$ and $\gamma \in \Gamma_o^t(\mu, \nu)$,

$$P_t^+\phi(\mu) = \int_{\mathbb{R}^m} T_t^+\phi(x) \, d\mu = \int_{\mathbb{R}^{2m}} T_t^+\phi(x) \, d\gamma = \int_{\mathbb{R}^{2m}} \phi(y) - A_t(x,y) \, d\gamma.$$

Moreover, for arbitrary $x, y \in \mathbb{R}^m$, $T_t^+\phi(x) \ge \phi(y) - A_t(x, y)$. Thus, $T_t^+\phi(x) = \phi(y) - A_t(x, y)$ holds for γ -a.e. $(x, y) \in \mathbb{R}^{2m}$. According to Lemma 2.7, $|x - y| \le \lambda_{\phi} t$ for γ -a.e. $(x, y) \in \mathbb{R}^{2m}$. In addition, there exists R > 0 such that supp $(\mu) \subset B(0, R)$ (determined by μ), and we can find some $t_R > 0$ such that when $t \in (0, t_R)$,

$$\operatorname{supp}(\nu) \subset B_{\lambda_{\phi}t}(\mu) \subset B(0, R).$$

We denote $C_R > 0$ as the semiconcave constant of ϕ on B(0, R). Let $t \in (0, t_R)$ and assume that $v_1, v_2 \in \mathscr{P}_c(\mathbb{R}^m)$ satisfy

$$v_1, v_2 \in \underset{v \in \mathscr{P}(\mathbb{R}^m)}{\operatorname{arg max}} \{ \phi(v) - C^t(\mu, v) \}.$$

Then, $v_1, v_2 \in \mathscr{P}(B(0, R))$. We have two optimal plans $\gamma_i \in \Gamma_o^t(\mu, v_i)$, such that $|x - y_i| \leq \lambda_{\phi} t$ for γ_i -a.e. $(x, y_i) \in \mathbb{R}^{2m}$, i = 1, 2. For $s \in [0, 1]$, we define

$$\gamma_{x,1\to2}^s := ((1-s)\pi_{x,y_1} + s\pi_{x,y_2})_{\#}\gamma,$$

$$\gamma_{1\to2}^s := ((1-s)\pi_{y_1} + s\pi_{y_2})_{\#}\gamma,$$

where we reset $\gamma \in \Gamma(\mu, \nu_1, \nu_2)$ and $(\pi_{x,y_1})_{\#}\gamma = \gamma_1$, $(\pi_{x,y_2})_{\#}\gamma = \gamma_2$. In this case, $\gamma_{1\to2}^s$ is the generalized geodesic from ν_1 to ν_2 with μ as the base point, and $\gamma_{x,1\to2}^s \in \Gamma(\mu, \gamma_{1\to2}^s)$. Furthermore, Theorem 3.5 implies the existence of $t_{\lambda_{\phi},R} > 0$ and $C_{\phi,\mu} > 0$ such that when $t \in (0, t_{\lambda_{\phi},R}]$,

$$(1-s)C^{t}(\mu,\nu_{1}) + sC^{t}(\mu,\nu_{2}) - C^{t}(\mu,\gamma_{1\to2}^{s}) \ge \frac{C_{\phi,\mu}}{t}s(1-s)W_{2,\gamma}^{2}(\nu_{1},\nu_{2}).$$
(41)

On the other hand, ϕ is semiconcave on B(0, R) with constant C_R . Invoking Proposition 3.3, we obtain

$$(1-s)\phi(\nu_1) + s\phi(\nu_2) - \phi(\gamma_{1\to 2}^s) \le C_R s(1-s) W_{2,\gamma}^2(\nu_1,\nu_2).$$
(42)

We set $t_{\phi,\mu} := \min \{ t_{\lambda_{\phi},R}, \frac{C_{\phi,\mu}}{C_R} \}$. Combining this with equations (41) and (42), when $t \in (0, t_{\phi,\mu}]$,

$$(1-s)[\phi(v_1) - C^t(\mu, v_1)] + s[\phi(v_2) - C^t(\mu, v_2)] - [\phi(\gamma_{1 \to 2}^s) - C^t(\mu, \gamma_{1 \to 2}^s)] \\ \leq \left(C_R - \frac{C_{\phi,\mu}}{t}\right) s(1-s)W_{2,\gamma}^2(v_1, v_2) < 0.$$

This implies that the function $s \mapsto \phi(\gamma_{1 \to 2}^s) - C^t(\mu, \gamma_{1 \to 2}^s)$ is strictly concave on [0, 1], and thus has a unique maximizer. Therefore, $v_1 = v_2$, which means that $\arg \max_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{\phi(\nu) - C^t(\mu, \nu)\}$ is a singleton.

For $\phi \in \text{Lip}(\mathbb{R}^m)$, we have $(\kappa_1, \kappa_2) = (\text{Lip}(\phi), 0)$. Equation (28) states

$$\lim_{t \to 0^+} W_1(\mu, \nu_{\phi,\mu}(t)) = \lim_{t \to 0^+} \lambda_{\phi} t = 0$$

Note that by Theorem 4.1, the support of $v_{\phi,\mu}(t)$ is contained in the union of balls $\bigcup_{x \in \text{supp }(\mu)} B(x, \lambda_{\phi} t)$. For all $t \in [0, t_{\phi,\mu}]$, the supports supp $(v_{\phi,\mu}(t))$ are contained within the same compact subset K, which implies that the W_p -distances for all $p \ge 1$ are equivalent on K. Consequently, for any fixed W_p -metric, we have that $\lim_{t\to 0^+} v_{\phi,\mu}(t) = \mu$. This completes the proof. \Box

Remark 5.3. When considering a compact manifold, such as \mathbb{T}^m , the singularity curve $v_{\phi,\mu}(t)$ mentioned in Theorem 5.4 below can be more precisely characterized. In fact, by the application of Propositions 2.10 and 2.11, we have

$$\nu_{\phi,\mu}(t) = [\pi \circ \Phi_H^t(\cdot, DT_t^+\phi(\cdot))]_{\#}\mu, \quad t \in (0, t_{\phi,\mu}].$$

Lemma 5.3. Suppose $t \ge 0$. Given $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, a locally differentiable point of $u(\cdot)$, we define

$$\psi_s(x) := \begin{cases} \pi \circ \Phi_H^{s-t}(x, Du(x)), & x \in A, \\ 0, & x \in \mathbb{R}^m \setminus A \end{cases}$$

where A is a measurable set with $\mu(A) = 1$ such that u is differentiable at each $x \in A$. Let $\{\mu_s\}_{s \in [0,t]}$ be a subset of $\mathscr{P}_c(\mathbb{R}^m)$, defined as $\mu_s = (\psi_s)_{\#}\mu$. Then for any $s \in [0, t]$,

$$u(\mu_t) = u(\mu_s) + C^{t-s}(\mu_s, \mu_t) + c[0](t-s).$$
(43)

Thus, $\{\mu_s\}_{s \in [0,t]}$ is a $(u(\cdot), C, c[0])$ -calibrated curve.

Proof. For any $t \ge 0$, we first note that $\mu_t = \mu$ and

$$u(\mu_t) - u(\mu_0) \leq C^t(\mu_0, \mu_t) + c[0]t$$

Then, there exists a random curve ξ such that law $(\xi(s)) = \mu_s$ for $s \in [0, t]$. This allows us to analyze the problem by considering individual points. For a differentiable point *x* of *u*, there exists a unique (u, L, c[0])-calibrated curve $\eta_x \in C^2((-\infty, t]; \mathbb{R}^m)$ satisfying $\eta_x(t) = x$. For all t > 0, η_x is the minimizer of the following Bolza problem:

$$\min_{\eta \in AC([0,t];\mathbb{R}^m)} \left\{ u(\eta(0)) + \int_0^t L(\eta(s), \dot{\eta}(s)) \, ds : \eta(t) = x \right\}.$$

Simultaneously, for $s \in [0, t]$,

$$u(\eta_x(t)) - u(\eta_x(s)) = A_{t-s}(\eta_x(s), \eta_x(t)) + c[0](t-s).$$
(44)

Therefore, $p_x(t) := L_v(\eta_x(t), \dot{\eta}_x(t)) = Du(x)$ and η_x solves the Euler-Lagrange equation. By invoking the well-posedness of the Cauchy problem for ODEs, we claim that $\eta_s(x) = \psi_s(x)$ holds for any $s \in [0, t]$ and μ -almost every $x \in \mathbb{R}^m$. Then, $(\psi_s)_{\#\mu_t} = \text{law}(\xi(s))$ is the displacement interpolation of $C^t(\mu_0, \mu_t)$ and $\mu_s = (\psi_s)_{\#\mu_t}$. Finally, integrating (44) by μ_t , we obtain (43).

Theorem 5.4. Assume that $\mu \in \mathcal{C}(u(\cdot))$. Let $v_{u,\mu}(t)$ and $t_{u,\mu} > 0$ be as mentioned in Theorem 5.3, where $u(\cdot)$ is the potential energy functional induced by the viscosity solution u of equation (HJ_s). Then, $v_{u,\mu}(t) \in \mathcal{S}(u(\cdot))$ for all $t \in (0, t_{u,\mu}]$.

Proof. We prove this argument by contradiction: for any fixed $t \in (0, t_{u,\mu}]$, we assume that $u(\cdot)$ is locally differentiable at $v_{u,\mu}(t)$. By Corollary 4.1, there exists $\gamma_t \in \Gamma_o^t(\mu, v_{u,\mu}(t))$ such that

$$P_t^+ u(\mu) = \int_{\mathbb{R}^m} T_t^+ u(x) \, d\mu = \int_{\mathbb{R}^{2m}} [u(y) - A_t(x, y)] \, d\gamma_t$$

Thus, we have for γ_t -almost every $(x, y) \in \mathbb{R}^{2m}$,

$$u(x) - A_t(x, y) = T_t^+ u(x).$$

Moreover, because $u(\cdot)$ is locally differentiable at $v_{u,\mu}(t)$ by our assumption, u is differentiable at $v_{u,\mu}(t)$ -almost every $y \in \mathbb{R}^m$. Fermat's rule implies

$$0 \in D_{y}^{+}(u(y) - A_{t}(x, y)) = Du(y) - D_{y}^{-}A_{t}(x, y), \quad \gamma_{t}\text{-almost every } (x, y) \in \mathbb{R}^{2m}.$$

Therefore, $D_y A_t(x, y) = Du(y)$ for γ_t -almost every $(x, y) \in \mathbb{R}^{2m}$. According to Lemma 5.3, $x = \psi_t(y), (\psi_t)_{\#}(v_{u,\mu}(t)) = \mu$ and

$$[0,t] \ni s \mapsto (\psi_s)_{\#} v_{u,\mu}(t)$$

is a $(u(\cdot), C, c[0])$ -calibrated curve starting from μ . That is, a $(u(\cdot), C, c[0])$ -calibrated curve ending with μ can be extended further (for the existence of such a curve, see Remark 5.1). This conclusion contradicts the assumption that $\mu \in \mathcal{C}(u(\cdot))$.

Remark 5.4. Although when $t > t_{u,\mu}$, the set $\arg \max_{\nu \in \mathscr{P}(\mathbb{R}^m)} \{\phi(\nu) - C^t(\mu, \nu)\}$ is not necessarily a singleton, we can still establish that

$$\underset{v \in \mathscr{P}(\mathbb{R}^m)}{\arg \max\{u(v) - C^{t}(\mu, v)\}} \subset \mathscr{S}(u(\cdot)).$$

Remark 5.5. Suppose *u* is a viscosity solution of (HJ_s) on \mathbb{R}^m and $\mu \ll \mathscr{L}^m$.

(1) Let $0 < T_u(\mu) \leq +\infty$. For $t \in [0, T_u(\mu))$, we have

$$\mu_t := [\pi \circ \Phi_H^t(\cdot, Du(\cdot))]_{\#} \mu = \underset{\nu \in \mathscr{P}(\mathbb{R}^m)}{\arg \max\{u(\nu) - C^t(\mu, \nu)\}} \ll \mathscr{L}^m.$$

In fact, this case is consistent with the DiPerna-Lions theory ([41]). More precisely, μ_t is driven by the flow $\mathbf{X}_1(t, x) := \pi \circ \Phi_H^t(x, Du(x))$ with the vector field

$$\mathbf{b}_1(t, x) = H_p(x, Du(x)),$$

where it is defined. Since *u* is of class $C^{1,1}$ on the interior of the (u, L, c[0])-calibrated curve, **b**₁ satisfies certain regularity conditions.

- (a) $\text{Lip}(\mathbf{b}_1(t, \cdot)) \in L^1([0, t]);$
- (b) $\mathbf{b}_1 \in L^1([0, t]; \operatorname{Lip}(\mathbb{R}^m; \mathbb{R}^m));$
- (c) div($\mathbf{b}_1(t, \cdot)$) $\in L^1([0, t]; L^{\infty}(\mathbb{R}^m));$

(d)
$$\frac{|\mathbf{b}_1|}{1+|\mathbf{y}|} \in L^1([0,t]; L^{\infty}(\mathbb{R}^m))$$

According to the DiPerna-Lions theory, there exists a unique regular Lagrangian flow associated with the vector field \mathbf{b}_1 .

(2) Given $T_u(\mu) = 0$, it follows that $\mu \in \mathscr{C}(u(\cdot))$ and there exists a positive time $t_{u,\mu}$ determined by Theorem 5.3. For $t \in (0, t_{u,\mu}]$,

$$v_{u,\mu}(t) = \underset{v \in \mathscr{P}(\mathbb{R}^m)}{\arg \max} \{ u(v) - C^t(\mu, v) \} \not\ll \mathscr{L}^m$$

are the singular points of $u(\cdot)$. Specifically, for the case of $\mathscr{P}(\mathbb{T}^m)$, due to Arnaud's theorem, $v_{u,\mu}$ is driven by the local Lipschitz flow on $t \in (0, t_u] \times \mathbb{T}^m$

$$\mathbf{X}_2(t, x) := \pi \circ \Phi_H^t(x, DT_t^+ u(x)),$$

which can be extended to $[0, \infty) \times \mathbb{T}^m$, for instance, as a t_u -periodic vector field $\mathbf{b}_2(t, x)$. Note that Lip $(DT_t^+ u) \leq \frac{C}{t}$ (see [29]). Therefore, \mathbf{b}_2 does not generally satisfy conditions (a)-(d) when the time step $t_u \to 0^+$, implying that the DiPerna-Lions theory is not applicable directly in this case.

In other words, the transportation of an absolutely continuous measure (with respect to the Lebesgue measure) along the characteristics before the cut time can ensure absolute continuity. However, after the cut time, it loses this absolute continuity and becomes a singular measure.

5.3. Variational construction of strict generalized measure singular characteristics. In the last subsection, we constructed a Lipschitz curve $v_{u,\mu}(\cdot)$ on $[0, t_u)$ (Remark 5.2), which can be extended to $[0, +\infty)$ and propagates singularities globally, provided the initial data is singular. However, this construction depends critically on the time step t_u . To obtain a semiflow evolution (at least for dynamical cost induced by mechanical Lagrangian) on the cut locus, we need to let $t_u \rightarrow 0^+$. This idea is inspired by the concept of *minimizing movements*, which is widely used in the theory of abstract gradient flows.

Lemma 5.4 ([29]). Suppose *M* is a Riemannian manifold, $\phi \in \text{SCL}_{\text{loc}}(M)$ and $\eta : \mathbb{R} \to M$ is a locally absolutely continuous curve, then

$$\frac{d}{dt}\phi(\eta(t)) = p(\dot{\eta}(t)), \ a.e. \ t \in \mathbb{R}, \ \forall p \in D^+\phi(\eta(t)).$$

Theorem 5.5. Suppose $\phi \in \text{SCL}(\mathbb{T}^m)$, T > 0, and $\mu_0 \in \mathscr{P}(\mathbb{T}^m)$. Then there exists a Lipschitz curve $\mu(\cdot) : [0, T] \to \mathscr{P}(\mathbb{T}^m)$ satisfying the following continuity equation:

$$\begin{cases} \frac{d}{dt}\mu + \operatorname{div}(H_p(\cdot, \mathbf{p}_{\phi}^{\#}(\cdot)) \cdot \mu) = 0, \\ \mu(0) = \mu_0, \end{cases}$$
(45)

where $\mathbf{p}_{\phi}^{\#}(x) := \arg\min\{H(x, p) : p \in D^{+}\phi(x)\}$ for any $x \in \mathbb{T}^{m}$.

Proof. Given T > 0 and $\mu_0 \in \mathscr{P}(\mathbb{T}^m)$, let

$$\Delta : 0 = t_0 < t_1 < \dots < t_{N-1} < t_N = T$$

be a partition of the interval [0, T] with $||\Delta|| < t_{\phi}$, where t_{ϕ} is as determined in Remark 5.2. For $N \in \mathbb{N}$ and $k = 0, 1, \dots, N - 1$, we define μ_{k+1} inductively as

$$\mu_{k+1} := \arg \max\{\phi(\nu) - C^{t_{k+1}-t_k}(\mu_k, \nu) : \nu \in \mathscr{P}(\mathbb{T}^m)\}.$$

More precisely,

$$P_{t_{k+1}-t_k}^+\phi(\mu_k) = \phi(\mu_{k+1}) - C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}), \quad k = 0, 1, \cdots, N-1.$$

In this case, by Corollary 4.1, we have the following sequence of equalities:

$$\begin{split} \phi(\mu_{k+1}) - \phi(\mu_k) &= P_{t_{k+1}-t_k}^+ \phi(\mu_k) - \phi(\mu_k) + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}) \\ &= \int_{\mathbb{T}^m} [T_{t_{k+1}-t_k}^+ \phi(x) - \phi(x)] \, d\mu_k + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}) \\ &= \int_{\mathbb{T}^m} \int_{t_k}^{t_{k+1}} \frac{d}{ds} T_{s-t_k}^+ \phi(x) \, ds \, d\mu_k + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}) \\ &= \int_{\mathbb{T}^m} \int_{t_k}^{t_{k+1}} H(x, DT_{s-t_k}^+ \phi(x)) \, ds \, d\mu_k + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}) \\ &= \int_{t_k}^{t_{k+1}} \int_{\mathbb{T}^m} H(x, DT_{s-t_k}^+ \phi(x)) \, d\mu_k \, ds + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}). \end{split}$$
(46)

Summing this equation from k = 0 to k = N - 1, we obtain

$$\phi(\mu_N) - \phi(\mu_0) = \sum_{k=0}^{N-1} \left[\int_{t_k}^{t_{k+1}} \int_{\mathbb{T}^m} H(x, DT^+_{s-t_k}\phi(x)) \, d\mu_k \, ds + C^{t_{k+1}-t_k}(\mu_k, \mu_{k+1}) \right]. \tag{47}$$

It is noteworthy that for each k, Remark 5.3 gives us

$$\mu_{k+1} = [\pi \circ \Phi_H^{t_{k+1}-t_k}(\cdot, DT_{t_{k+1}-t_k}^+\phi(\cdot))]_{\#}\mu_k.$$

In this framework, we define

$$\mu_{\Delta}(s) := [\pi \circ \Phi_{H}^{s-t_{k}}(\cdot, DT_{t_{k+1}-t_{k}}^{+}\phi(\cdot))]_{\#}\mu_{k}, \quad s \in [t_{k}, t_{k+1})$$
(48)

for $k = 0, 1, \dots, N-1$ with $\mu_{\Delta}(0) := \mu_0$. Then $\{\mu_{\Delta}(s)\}_{s \in [t_k, t_{k+1}]}$ is a displacement interpolation of $C^{t_k, t_{k+1}}(\mu_k, \mu_{k+1})$ by Proposition 2.11. Moreover, the family $\{\mu_{\Delta} : \|\Delta\| \ll 1\}$ is equi-Lipschitz (with a constant depending only on *L* and ϕ) in the sense of W_1 metric (and W_p metric for all $p \ge 1$). If we set $\tau_{\Delta}(\cdot) : [0, T] \rightarrow [0, T]$ as $\tau_{\Delta}(s) = t_k$ for $s \in [t_k, t_{k+1})$, $k = 0, 1, \dots, N-1$, and $\tau_{\Delta}(T) = T$, then (47) can be rewritten as

$$\phi(\mu_N) - \phi(\mu_0) = \int_0^T \int_{\mathbb{T}^m} H(x, DT^+_{s-\tau_\Delta(s)}\phi(x)) \, d\mu_\Delta(\tau_\Delta(s)) \, ds + \sum_{k=0}^{N-1} \mathbb{A}^{t_{k+1}-t_k}(\mu_\Delta)$$

$$= \int_0^T \int_{\mathbb{T}^m} H(x, DT^+_{s-\tau_\Delta(s)}\phi(x)) \, d\mu_\Delta(\tau_\Delta(s)) \, ds + \mathbb{A}^T(\mu_\Delta).$$
(49)

Now, consider a sequence of partitions $\{\Delta_n\}_{n\geq 1}$ such that $\lim_{n\to\infty} ||\Delta_n|| = 0$. By invoking the Arzelà-Ascoli theorem ([5, Theorem 10.3 & 10.4]), we assume that $\mu_{\Delta_n} \to \mu \in \operatorname{Lip}([0, T]; \mathscr{P}(\mathbb{T}^m))$ uniformly. Also, since $\tau_{\Delta_n}(t) \to t$ as $||\Delta_n|| \to 0$, $\mu_{\Delta_n}(\tau_{\Delta_n}(t)) \stackrel{*}{\to} \mu(t)$ for any $t \in [0, T]$. Furthermore, from ([29, Proposition 4.2]), we recall that for any $x \in \mathbb{T}^m$,

$$\lim_{t \to 0^+} DT_t^+ \phi(x) = \arg\min_{p \in D^+ \phi(x)} \{H(x, p)\} =: \mathbf{p}_{\phi}^{\#}(x).$$
(50)

Equation (49) and (50), along with Fatou's lemma, imply that

$$\phi(\mu(T)) - \phi(\mu(0)) = \lim_{n \to \infty} \int_0^T \int_{\mathbb{T}^m} H(x, DT^+_{s - \tau_{\Delta_n}(s)}\phi(x)) \, d\mu_{\Delta_n}(\tau_{\Delta_n}(s)) \, ds + \mathbb{A}^T(\mu_{\Delta_n})$$

$$\geq \int_0^T \int_{\mathbb{T}^m} H(x, \mathbf{p}^{\#}_{\phi}(x)) \, d\mu(s) \, ds + \mathbb{A}^T(\mu)$$

$$= \int_0^T \int_{\mathbb{T}^m} H(x, \mathbf{p}^{\#}_{\phi}(x)) \, d\mu(s) \, ds + \min_{\xi} \mathbb{E}\left(\int_0^T L(\xi(s), \dot{\xi}(s)) \, ds\right)$$
(51)

where the last minimum is over all absolutely continuous random curves $\xi : [0, T] \times \Omega \to M$ such that $\mu(s) = \text{law}(\xi(s))$ for $s \in [0, T]$. We choose one of such minimizers and denote it by ξ_* . Now we can obtain from (51) that

$$\mathbb{E}(\phi(\xi(T)) - \phi(\xi(0))) \ge \mathbb{E}\left(\int_0^T [H(\xi_*(s), \mathbf{p}_{\phi}^{\#}(\xi_*(s))) + L(\xi_*(s), \dot{\xi}_*(s))] \, ds\right)$$
(52)

by Fubini's theorem. Due to Lemma 5.4 and Fenchel's inequality, (52) is actually an equality. Furthermore, $\mu(t)$ is the law at time *t* of a random solution ξ_* of the following equation:

$$\dot{\mathbf{x}}(t) = H_p(\mathbf{x}(t), \mathbf{p}_{\phi}^{\#}(\mathbf{x}(t))).$$

In other words, ξ_* satisfies

$$\dot{\xi}_*(t,\omega) = H_p(\xi_*(t,\omega), \mathbf{p}_\phi^{\#}(\xi_*(t,\omega))), \quad \mathcal{L}^1\text{-a.e. } t \in [0,T], \quad \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

Consequently, $\mu(\cdot) : [0, T] \to \mathscr{P}(\mathbb{T}^m)$ satisfies (45). This completes the proof.

Corollary 5.2. Assume that $\phi = u$ in (45) is a viscosity solution to (HJ_s) , , and $\mu(\cdot) : [0,T] \to \mathbb{T}^m$ is the solution of (45).

(1) If $\mu_0(\overline{\operatorname{Sing}(u)}) > 0$, then $[\mu(t)](\overline{\operatorname{Sing}(u)}) > 0$ for all $t \in [0, T]$. (2) If $\mu_0 \in \mathscr{C}(u(\cdot))$, then $\mu(t) \in \mathscr{C}(u(\cdot))$ for all $t \in [0, T]$.

Proof. For the proof of (1), recall Theorem 1.2 in [1]. If $\xi(0, \omega) \in \overline{\text{Sing}(u)}$ for a fixed $\omega \in \Omega$, then $\xi_*(t, \omega) \in \overline{\text{Sing}(u)}$ for all $t \in [0, T]$, where ξ_* is the minimizer as in the proof of Theorem 5.5. Thus,

$$\{\omega \in \Omega : \xi_*(0,\omega) \in \overline{\operatorname{Sing}(u)}\} \subset \{\omega \in \Omega : \xi_*(t,\omega) \in \overline{\operatorname{Sing}(u)}\}, \quad \forall t \in [0,T].$$

Since $\mu_0(\overline{\text{Sing}(u)}) > 0$, for any $t \in [0, T]$, we have

$$\begin{split} & [\mu(t)](\overline{\operatorname{Sing}\,(u)}) = \mathbb{P}(\{\omega \in \Omega : \xi_*(t,\omega) \in \overline{\operatorname{Sing}\,(u)}\}) \\ & \geq \mathbb{P}(\{\omega \in \Omega : \xi_*(0,\omega) \in \overline{\operatorname{Sing}\,(u)}\}) \\ & = \mu_0(\overline{\operatorname{Sing}\,(u)}) > 0. \end{split}$$

Now we turn to the proof of (2). Since $\mu(\cdot) : [0, T] \to \mathbb{T}^m$ is the solution of (45), we know that $\mu(t)$ is a law at time *t* of a random solution ξ_* of the equation

$$\dot{\xi}_*(t,\omega) = H_p(\xi_*(t,\omega), \mathbf{p}_{\phi}^{\#}(\xi_*(t,\omega))) \in \operatorname{co} H_p(\xi_*(t,\omega), D^+\phi(\xi_*(t,\omega)))$$

for a.e. $t \in [0, T]$ and \mathbb{P} -a.e. $\omega \in \Omega$. Since $\mu_0 \in \mathscr{C}(\phi(\cdot))$, Corollary 5.1 implies that

$$\mathbb{P}(\{\omega \in \Omega : \tau_u(\xi(0,\omega)) < \varepsilon\}) > 0, \quad \forall \varepsilon > 0.$$

Moreover, according to [29, Theorem 5.7], $\tau_u(\xi_*(t,\omega)) \leq \tau_u(\xi_*(0,\omega))$ for all $t \in [0, T]$ and \mathbb{P} -a.e. $\omega \in \Omega$. Hence, for any $\varepsilon > 0$,

$$\mathbb{P}(\{\omega \in \Omega : \tau_u(\xi_*(t,\omega)) < \varepsilon\}) \ge \mathbb{P}(\{\omega \in \Omega : \tau_u(\xi_*(0,\omega)) < \varepsilon\}) > 0$$

This shows that $\mu(t) \in \mathscr{C}(u(\cdot))$ by Corollary 5.1.

Remark 5.6. Given $\phi \in \text{SCL}(\mathbb{T}^m)$, we define⁴

$$\overline{\mathscr{S}}(\phi(\cdot)) := \left\{ \mu \in \mathscr{P}(\mathbb{T}^m) : \mu(\overline{\operatorname{Sing}\,(\phi)}) > 0 \right\}.$$

(1) $\overline{\mathscr{S}}(\phi(\cdot))$ is an F_{σ} -set in the sense of weak-* topology. In fact, we can consider the sets

$$A_k := \left\{ \mu \in \mathscr{P}(\mathbb{T}^m) : \mu(\overline{\operatorname{Sing}\,(\phi)}) \ge \frac{1}{k} \right\}, \quad k \in \mathbb{N}.$$

By the Portmanteau theorem, for any sequence $\{\mu_n\}_{n \in \mathbb{N}} \subset A_k$ with $\mu_n \stackrel{*}{\rightharpoonup} \mu$,

$$\mu_n(\overline{\operatorname{Sing}\,(\phi)}) \ge \frac{1}{k} \Rightarrow \ \mu(\overline{\operatorname{Sing}\,(\phi)}) \ge \limsup_{n \to \infty} \mu_n(\overline{\operatorname{Sing}\,(\phi)}) \ge \frac{1}{k},$$

i.e., $\mu \in A_k$. Therefore, each A_k is a closed subset, and $\overline{\mathscr{S}}(\phi(\cdot)) = \bigcup_{k=1}^{\infty} A_k$ is an F_{σ} -set.

(2) Suppose *u* is a viscosity solution of (HJ_s) . Then $Sing(u) \subset Cut(u) \subset \overline{Sing(u)}$ (see [29]). However, $\mathscr{C}(u(\cdot))$ and $\overline{\mathscr{S}}(u(\cdot))$ are not mutually contained in general, and we have $\mathscr{S}(u(\cdot)) \subsetneq \mathscr{C}(u(\cdot)) \cap \overline{\mathscr{S}}(u(\cdot))$.

Lemma 5.5 ([29]). Given T > 0. Let M be a closed manifold, $\phi \in SCL(M)$, and $H \in C^2(T^*M)$ be a Tonelli Hamiltonian. If $\eta : [0, T] \to M$ is a strict singular characteristic, i.e., η satisfies the equality in the following variational inequality

$$\phi(\eta(t_2)) - \phi(\eta(t_1)) \leq \int_{t_1}^{t_2} L(\eta(s), \dot{\eta}(s)) + H(\eta(s), \mathbf{p}_{\phi}^{\#}(\eta(s))) \, ds, \quad \forall t_1, t_2 \in [0, T], \ t_1 < t_2,$$

or equivalently,

$$\phi(\eta(T)) - \phi(\eta(0)) \leq \int_0^T L(\eta(s), \dot{\eta}(s)) + H(\eta(s), \mathbf{p}_{\phi}^{\#}(\eta(s))) \, ds,$$

then

$$\dot{\eta}^+(t) = H_p(\eta(t), \mathbf{p}^{\#}_{\phi}(\eta(t))), \quad \forall t \in [0, T).$$

Theorem 5.6. Under the same assumptions as in Theorem 5.5, the Lipschitz curve mentioned in Theorem 5.5 that satisfies equation (45) also has the following property: for any $f \in C^{\infty}(\mathbb{T}^m)$, the map $t \mapsto f(\mu(t))$ is in Lip $([0, T]; \mathbb{R})$ and its right derivative

$$\frac{d^+}{dt}f(\mu(t)) = \int_{\mathbb{T}^m} \langle Df(x), H_p(x, \mathbf{p}_{\phi}^{\#}(x)) \rangle \, d\mu(t)$$

exists for all $t \in [0, T)$ *.*

⁴This definition can be extended similarly to the case of Euclidean space \mathbb{R}^{m} .

Proof. By Lemma 5.5, we can deduce from the proof of Theorem 5.5 that

$$\dot{\xi}_*^+(t,\omega) = H_p(\xi_*(t,\omega), \mathbf{p}_{\phi}^{\#}(\xi_*(t,\omega))), \quad \forall t \in [0,T), \ \mathbb{P} - \text{a.e.} \ \omega \in \Omega.$$

Now, for an arbitrary fixed $f \in C^{\infty}(\mathbb{T}^m)$, consider the map $F(t) := f(\mu(t))$ with $t \in [0, T]$. According to Proposition 16.3 in [5] and Theorem 5.5, $F \in \text{Lip}([0, T]; \mathbb{R})$. Moreover,

$$\begin{split} \lim_{\tau \downarrow t} \frac{F(\tau) - F(t)}{\tau - t} &= \lim_{\tau \downarrow t} \frac{1}{\tau - t} \left\{ \int_{\mathbb{T}^m} f(x) d[\mu(\tau)] - \int_{\mathbb{T}^m} f(x) d[\mu(t)] \right\} \\ &= \lim_{\tau \downarrow t} \frac{1}{\tau - t} \mathbb{E} \left(f(\xi_*(\tau)) - f(\xi_*(t)) \right) \\ &= \mathbb{E} \left(\lim_{\tau \downarrow t} \frac{f(\xi_*(\tau)) - f(\xi_*(t))}{\tau - t} \right) \\ &= \mathbb{E} \left(\langle Df(\xi_*(t)), \dot{\xi}_*^+(t) \rangle \right) \\ &= \mathbb{E} \left(\langle Df(\xi_*(t)), H_p(\xi_*(t), \mathbf{p}_{\phi}^{\#}(\xi_*(t))) \rangle \right) \\ &= \int_{\mathbb{T}^m} \langle Df(x), H_p(x, \mathbf{p}_{\phi}^{\#}(x)) \rangle d[\mu(t)]. \end{split}$$

That is, $\dot{F}^+(t) = \int_{\mathbb{T}^m} \langle Df(x), H_p(x, \mathbf{p}_{\phi}^{\#}(x)) \rangle d[\mu(t)]$ for any $t \in [0, T)$. Since the argument above holds for any $f \in C^{\infty}(\mathbb{T}^m)$, we have completed the proof.

Appendix A. Proofs of some statements

Proof of Proposition 2.4. According to the definition of the cut time function, for $x \in M$ with $\tau_u(x) > 0$, there exists a curve $\eta_x : [0, \tau_u(x)] \to M$ such that

$$u(\eta_x(\tau_u(x))) - u(x) = A_{\tau_u(x)}(x, \eta_x(\tau_u(x))) + c[0]\tau_u(x)$$

This implies that

$$T^{+}_{\tau_{u}(x)}u(x) - c[0]\tau_{u}(x) \ge u(\eta_{x}(\tau_{u}(x))) - A_{\tau_{u}(x)}(x,\eta_{x}(\tau_{u}(x))) - c[0]\tau_{u}(x) = u(x)$$
$$\ge T^{+}_{\tau_{u}(x)}u(x) - c[0]\tau_{u}(x).$$

Thus, we have

$$T^{+}_{\tau_{u}(x)}u(x) - c[0]\tau_{u}(x) = u(x)$$

and

$$DT^+_{\tau_u(x)}u(x) = Du(x).$$

The term $A_{\tau_u(x)}(x, \eta_x(\tau_u(x)))$ has a unique minimizer, which we denote by η_x . Therefore, η_x is a (u, L, c[0])-calibrated curve starting from x. Additionally, η_x is also a (u, L, c[0])-calibrated curve on any subinterval [0, t] of $[0, \tau_u(x)]$.

Furthermore, we have

$$T^{-}_{\tau_{u}(x)}u(\eta_{x}(\tau_{u}(x))) + c[0]\tau_{u}(x) = u(\eta_{x}(\tau_{u}(x)))$$
$$= u(x) + A_{\tau_{u}(x)}(x,\eta_{x}(\tau_{u}(x))) + c[0]\tau_{u}(x),$$

which implies that $x \in \arg \min\{u(\cdot) + A_{\tau_u(x)}(\cdot, \eta(\tau_u(x)))\}$. By Fermat's rule,

$$Du(x) = D_x A_{\tau_u(x)}(x, \eta_x(\tau_u(x))) = L_v(\eta_x(0), \dot{\eta}_x(0)).$$

Thus, the first equality in (6) is satisfied. Note that u is differentiable at the interior points of calibrated curves. Consequently, for $t \in (0, \tau_u(x))$,

$$T_t^- u(\eta_x(t)) + c[0]t = u(\eta_x(t)) = u(x) + A_t(x, \eta_x(t)) + c[0]t$$

and

$$DT_{t}^{-}u(\eta_{x}(t)) = Du(\eta_{x}(t)) = D_{y}A_{t}(x,\eta_{x}(t)) = L_{v}(\eta_{x}(t),\dot{\eta}_{x}(t))$$

which confirms that the second equality in (6) holds true.

Proof of Lemma 3.1. Without loss of generality we suppose $f \neq 0$. Observe that there exists $\delta_f > 0$ depending on f such that $I_m + tD^2 f$ is positively definite for $t \in [0, \delta_f]$, i.e., id + tDf is the gradient of smooth convex function $\psi_t := \frac{1}{2} |\cdot|^2 + tf(\cdot)$. It is easy to check that

$$\Gamma := \{ (x, D\psi_t(x)) : x \in \mathbb{R}^m \}$$

is a *c*-cyclically monotone closed graph with $c(x, y) = \frac{1}{2}|x - y|^2$. Then for $t \in [0, \delta_f]$, the map id + tDf is optimal and

$$[\mathrm{id} \times (\mathrm{id} + tDf)]_{\#} \mu \in \Gamma_2(\mu, \nu_t).$$

By inequality (7) for any $s, t \in [0, \delta_f]$,

$$W_{2}(\nu_{t},\nu_{s}) = W_{2}((\mathrm{id} + tDf)_{\#}\mu, (\mathrm{id} + sDf)_{\#}\mu)$$
$$\leq \left\{ \int_{\mathbb{R}^{m}} (t-s)^{2} |Df|^{2} d\mu \right\}^{\frac{1}{2}} \leq |t-s|\delta_{f}^{-1}W_{2}(\mu,\nu_{\delta_{f}}).$$

it follows {(id + tDf)# μ }_{0 $\leq t \leq \delta_f$} is a constant speed geodesic in $\mathscr{P}_2(\mathbb{R}^m)$. Invoking Theorem 2.5, $\Gamma_2(\mu, \nu_t)$ contains a unique transport plan for $t \in [0, \delta_f]$. That is

$$\Gamma_2(\mu, \nu_t) = \{ [\operatorname{id} \times (\operatorname{id} + tDf)]_{\#} \mu \}, \qquad \forall t \in [0, \delta_f].$$

Since μ is compactly supported and $f \in C_c^{\infty}(\mathbb{R}^m)$, we claim that for any R > 0, there exists a $\delta_{f,R} > 0$ such that

$$(\mathrm{id} + tDf)(\mathrm{supp}\,(\mu)) \subset B_R(\mu), \qquad t \in [0, \delta_{f,R}]. \tag{53}$$

Indeed, we take $\delta_{f,R} = \min\{\delta_f, R/||f||_{C^1}\}$. Thus

$$|(\mathrm{id} + tDf)(x) - x| = |tDf(x)| \le t||f||_{C^1} \le R, \qquad \forall x \in \mathbb{R}^m$$

and (53) follows. By (53) we obtain

$$\nu_t(\mathbb{R}^m \setminus B_R(\mu)) = \mu\left(\left[\operatorname{id} + tDf\right]^{-1}(\mathbb{R}^m \setminus B_R(\mu))\right) \leq \mu(\mathbb{R}^m \setminus \operatorname{supp}(\mu)) = 0.$$

This implies that $v_t \in \mathscr{P}(B_R(\mu)) \subset \mathscr{P}_c(\mathbb{R}^m)$.

Proof of Proposition 3.2. Given $\alpha \in \partial^+ U(\mu)$ and $\beta \in \partial^- U(\mu)$. Then for any R > 0, $\nu \in \mathscr{P}(B_R(\mu))$, both (15) and (16) hold true. Invoking Lemma 3.1, for any $f \in C_c^{\infty}(\mathbb{R}^m)$ and R > 0, $\nu_t = (\mathrm{id} + tDf)_{\#}\mu \in \mathscr{P}(B_R(\mu))$ and

$$\Gamma_2(\mu, \nu_t) = \{ [\operatorname{id} \times (\operatorname{id} + tDf)]_{\#} \mu \},\$$

for any $t \in [0, \delta_{f,R}]$. Thus

$$\int_{\mathbb{R}^m} \left\langle \beta(x), tDf \right\rangle d\mu + o_R\left(t ||Df||_{L^2(\mu)}\right) \leq \int_{\mathbb{R}^m} \left\langle \alpha(x), tDf \right\rangle d\mu + o_R\left(t ||Df||_{L^2(\mu)}\right)$$

and it follows

$$\int_{\mathbb{R}^m} \left\langle \alpha(x) - \beta(x), Df \right\rangle d\mu \ge 0.$$

Therefore, $\alpha = \beta$, μ -a.e. since $f \in C_c^{\infty}(\mathbb{R}^m)$ is arbitrary.

Proof of Proposition 3.4. It is easy to check that (3) implies (2) and (2) implies (1). Thus, it is sufficient to prove (1) implies (3).

Given R > 0, we assume $\nu \in \mathscr{P}(B(0, R))$ and let $\gamma \in \Gamma_2(\mu, \nu)$. We define $\gamma_{1\to 2}^t$ for γ by (12). Theorem 2.5 implies $\Gamma_2(\mu, \gamma_{1\to 2}^t) = \{\gamma_{1,1\to 2}^t\}$ (defined in (14)) and $\gamma_{1\to 2}^t$ is the constant speed geodesic. Thus $W_2(\mu, \gamma_{1,1\to 2}^t) = tW_2(\mu, \nu)$ for all $t \in [0, 1]$. Now suppose $\alpha \in \partial^+ \phi(\mu)$. We obtain

$$\begin{split} \phi(\gamma_{1\to2}^{t}) - \phi(\mu) &\leq \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma_{1,1\to2}^{t} + o_R(W_2(\mu, \gamma_{1\to2}^{t})) \\ &= \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma_{1,1\to2}^{t} + o_R(tW_2(\mu, \nu)) \\ &= \int_{\mathbb{R}^{2m}} \langle \alpha(x), (1 - t)x + ty - x \rangle \, d\gamma + o_R(t) \\ &= \int_{\mathbb{R}^{2m}} \langle \alpha(x), t(y - x) \rangle \, d\gamma + o_R(t). \end{split}$$

Since $\phi(\cdot)$ is locally semiconcave, there exists $C_R > 0$ such that

$$(1-t)\phi(\mu) + t\phi(\nu) - \phi(\gamma_{1\to 2}^t) \le C_R t(1-t) + W_2^2(\mu,\nu).$$

It follows

$$\phi(\nu) - \phi(\mu) - C_R(1-t)W_2^2(\mu,\nu) \le \frac{\phi(\gamma_{1\to 2}^t) - \phi(\mu)}{t},$$

and

$$\phi(\nu) - \phi(\mu) - C_R W_2^2(\mu, \nu) \leq \limsup_{t \to 0^+} \frac{\phi(\gamma_{1 \to 2}^t) - \phi(\mu)}{t} \leq \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma.$$

Assertion (3) holds by taking the infimum over all $\gamma \in \Gamma_2(\mu, \nu)$ on the righthand side of the inequality above.

Proof of Theorem 3.1. Since $\mu \in \mathscr{P}_c(\mathbb{R}^m)$, there exists R > 0, such that supp $(\mu) \subset B(0, R)$. By local semiconcavity of ϕ , there exists $C_R > 0$ such that for any $x, y \in B(0, R)$ and any $t \in [0, 1]$,

$$(1-t)\phi(x) + t\phi(y) - \phi(tx + (1-t)y) \le C_R t(1-t)|x-y|^2.$$

We begin with necessity and we will prove by contradiction. Suppose there exists $A \subset B(0, R)$ with $\mu(A) > 0$ such that $\alpha(x) \notin D^+\phi(x)$ for all $x \in A$. We define a set-valued map

$$G: A \rightrightarrows \overline{B(0,R)}$$
$$x \mapsto G(x) := \underset{y \in \overline{B(0,R)}}{\operatorname{arg\,max}} \left\{ \phi(y) - \phi(x) - \langle \alpha(x), y - x \rangle - C_R |y - x|^2 \right\}.$$

42

By a standard measurable selection theorem (see, for example [4, Theorem 18.19]), there is a Borel measurable selection $g : A \to \overline{B(0,R)}$ of *G*, i.e. $x \in A$, $g(x) \in G(x)$. Due to our assumption, $\alpha(x) \notin D^+\phi(x)$ and g(x) is positive on *A*. The map $h : \mathbb{R}^m \to \mathbb{R}^m$,

$$h(x) = \begin{cases} x, & x \notin A \\ g(x), & x \in A \end{cases}$$

is measurable. Set $v := h_{\#}\mu$. Since $h(\overline{B(0,R)}) \subset \overline{B(0,R)}$, we have

$$h_{\#}\mu(\mathbb{R}^m \setminus \overline{B(0,R)}) = \mu(h^{-1}[\mathbb{R}^m \setminus \overline{B(0,R)}]) = \mu(\mathbb{R}^m \setminus [h^{-1}(\overline{B(0,R)}]) \leq \mu(\mathbb{R}^m \setminus \overline{B(0,R)}) = 0.$$

That is $v \in \mathscr{P}(\overline{B(0,R)})$. By Proposition 3.4 we conclude that if $\alpha \in \partial^+ \phi(\mu)$ then

$$\phi(\nu) - \phi(\mu) - \int_{\mathbb{R}^m} \langle \alpha(x), y - x \rangle \, d(\mathrm{id} \times h)_{\#} \mu \leqslant C_R W_{2,(\mathrm{id} \times h)_{\#} \mu}^2(\mu, \nu).$$
(54)

However, the definition of *h* and the fact that g > 0 on *A* imply

$$\begin{split} \phi(v) &- \phi(\mu) - \int_{\mathbb{R}^m} \langle \alpha(x), y - x \rangle \, d(\mathrm{id} \times h)_{\#\mu} \\ &= \int_{\mathbb{R}^m} \phi(h(x)) - \phi(x) - \langle \alpha(x), h(x) - x \rangle \, d\mu \\ &= \int_A \phi(g(x)) - \phi(x) - \langle \alpha(x), g(x) - x \rangle \, d\mu \\ &> \int_A C_R |g(x) - x|^2 \, d\mu = \int_{\mathbb{R}^m} C_R |h(x) - x|^2 \, d\mu \\ &= C_R W_{2,(\mathrm{id} \times h)_{\#\mu}}^2(\mu, \nu). \end{split}$$

This leads to a contradiction to (54).

Now we turn to prove the sufficiency part. Suppose $\phi \in \operatorname{Lip}_{\operatorname{loc}}(\mathbb{R}^m)$, $\alpha(x) \in D^+\phi(x)$ for μ -a.e. *x*. Then $\alpha \in L^{\infty}(\mathbb{R}^m;\mu)$ and

$$\phi(y) - \phi(x) \leq \langle \alpha(x), y - x \rangle + C_R |y - x|^2,$$

for every $y \in B(0, R)$ and μ -a.e. $x \in \mathbb{R}^m$. Thus, for any $v \in \mathscr{P}(B(0, R))$ and $\gamma \in \Gamma_2(\mu, v)$, integrating both sides of the inequality above, we get

$$\begin{split} \phi(\nu) - \phi(\mu) &\leq \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle + C_R |y - x|^2 \, d\gamma \\ &= \int_{\mathbb{R}^{2m}} \langle \alpha(x), y - x \rangle \, d\gamma + C_R W_2^2(\mu, \nu). \end{split}$$

Taking the infimum over all $\gamma \in \Gamma_2(\mu, \nu)$, we have (18). This implies $\alpha \in \partial^+ \phi(\mu)$.

Lemma A.1 ([9, Lemma 5.3.2]). Let X_i (i = 1, 2, 3) be Polish spaces, $\gamma^{1,2} \in \mathscr{P}(X_1 \times X_2)$ and $\gamma^{1,3} \in \mathscr{P}(X_1 \times X_3)$ with $(\pi_1)_{\#}\gamma^{1,2} = (\pi_1)_{\#}\gamma^{1,3} = \mu^1$. Then there exists $\mu \in \mathscr{P}(X_1 \times X_2 \times X_3)$, which satisfies

$$(\pi_{1,2})_{\#}\mu = \gamma^{1,2}, \qquad (\pi_{1,3})_{\#}\mu = \gamma^{1,3}.$$
 (55)

Moreover, if $\gamma^{1,2} = \int \gamma_{x_1}^{1,2} d\mu^1$, $\gamma^{1,3} = \int \gamma_{x_1}^{1,3} d\mu^1$ and $\mu = \int \mu_{x_1} d\mu^1$ are the disintegrations of $\gamma^{1,2}$, $\gamma^{1,3}$ and μ with respect to μ^1 respectively, then (55) is equivalent to

$$\mu_{x_1} \in \Gamma(\gamma_{x_1}^{1,2}, \gamma_{x_1}^{1,3}) \subset \mathscr{P}(X_2 \times X_3) \qquad for \, \mu^1 - a.e. \, x_1 \in X_1.$$

Lemma A.2. Suppose $\mu_i \in \mathscr{P}(\mathbb{R}^m)$ with i = 1, 2, 3 and $\gamma \in \Gamma(\mu_2, \mu_3)$. For any fixed $\lambda \in [0, 1]$ and t > 0, let $\mu_\lambda \in \Gamma_o^t(\mu_1, \gamma_{2\to 3}^{\lambda})$. Then there exists $\gamma_\lambda \in \Gamma(\mu_1, \gamma)$ such that $(\gamma_\lambda)_{1,2\to 3}^{\lambda} = \mu_\lambda$.

Proof. The following is inspired by the proof of [9, Proposition 7.3.1]. Observe that maps

$$\Sigma_{\lambda} : \mathbb{R}^{2m} \to \mathbb{R}^{2m}$$

(x₂, x₃) \mapsto (x₂, (1 - λ)x₂ + λ x₃) and
$$\Lambda_{\lambda} : \mathbb{R}^{3m} \to \mathbb{R}^{3m}$$

(x₁, x₂, x₃) \mapsto (x₁, x₂, (1 - λ)x₂ + λ x₃)

are homeomorphisms for any fixed $\lambda \in (0, 1]$. Thus, $\gamma_{\lambda} \in \Gamma(\mu_1, \gamma)$ satisfies $(\gamma_{\lambda})_{1,2\to 3}^{\lambda} = \mu_{\lambda}$ if and only if for $\nu_{\lambda} := (\Lambda_{\lambda})_{\#} \gamma_{\lambda}$,

$$(\pi_{1,3})_{\#}\nu_{\lambda} = \mu_{\lambda}, \qquad (\pi_{2,3})_{\#}\nu_{\lambda} = (\Sigma_{\lambda})_{\#}\gamma. \tag{56}$$

Then Lemma A.1 implies the existence of v_{λ} and this ensures (56) hold true. This guarantees the existence of γ_{λ} since Λ_{λ} is invertible.

For the case of $\lambda = 0$, we only need to check that the existence of $\gamma_0 \in \Gamma(\mu_1, \gamma)$ satisfying $(\pi_{1,2})_{\#}\gamma_0 = \nu$ for each $\nu \in \Gamma_o^t(\mu_1, \mu_2)$. This can be guaranteed by Lemma A.1 as well.

Proof of Theorem 3.3. Note that for any $v \in \mathbb{R}^m$, $L(x, v) \ge \theta_0(|v|) - c_0$, where θ_0 is convex, increasing and superlinear function in \mathbb{R}^+ . We assume that $\xi \in L^t_{\mu,v}$ is the dynamical optimal coupling of $C^t(\mu, v)$, then by Jensen's inequality and the monotonicity of θ_0 , for $\mu, v \in \mathscr{P}_c(\mathbb{R}^m)$,

$$C^{t}(\mu,\nu) = \mathbb{E}\left(\int_{0}^{t} L(\xi(s),\dot{\xi}(s)) \, ds\right)$$

$$\geq \mathbb{E}\left(\int_{0}^{t} \theta_{0}(|\dot{\xi}(s)|) - c_{0} \, ds\right)$$

$$\geq t\mathbb{E}\left(\theta_{0}\left(\left|\frac{1}{t}\int_{0}^{t}\dot{\xi}(s) \, ds\right|\right) - c_{0}\right)$$

$$= t\mathbb{E}\left(\theta_{0}\left(\left|\frac{\xi(t) - \xi(0)}{t}\right|\right) - c_{0}\right)$$

$$\geq t\theta_{0}\left(\mathbb{E}\left(\left|\frac{\xi(t) - \xi(0)}{t}\right|\right)\right) - c_{0}t$$

$$\geq t\theta_{0}\left(\frac{1}{t}W_{1}(\mu,\nu)\right) - c_{0}t.$$

Thus, for any $k \in \mathbb{R}$,

$$C^{t}(\mu,\nu) \ge t\theta_{0}\left(\frac{1}{t}W_{1}(\mu,\nu)\right) - c_{0}t \ge kW_{1}(\mu,\nu) - (c_{0} + \theta_{0}^{*}(k))t$$

where θ_0^* is Fenchel-Legendre duality of θ_0 , which implies $C^t(\mu, \cdot)$ is superlinear on $\mathscr{P}_1(\mathbb{R}^m)$. The proof of superlinearity of $C^t(\cdot, \mu)$ on $\mathscr{P}_c(\mathbb{R}^m)$ is similar.

Given $\mu, v_1, v_2 \in \mathscr{P}(K)$ and $t_1, t_2 \in [a, b]$, we assume $\gamma_2 \in \Gamma_o^{t_2}(\mu, v_2)$ and $\gamma_{1,2} \in \Gamma_1(v_1, v_2)$. Lemma A.1 provides $\gamma_2 * \gamma_{1,2} \in \Gamma(v_1, v_2, \mu) \in \mathscr{P}(\mathbb{R}^{3m})$ with $(\pi_{3,1})_{\#}(\gamma_2 * \gamma_{1,2}) \in \Gamma(\mu, v_1)$ and $(\pi_{3,2})_{\#}(\gamma_2 * \gamma_{1,2}) = \gamma_2$. Therefore,

$$\begin{split} &C^{t_1}(\mu, \nu_1) - C^{t_2}(\mu, \nu_2) \\ &\leqslant \int_{\mathbb{R}^{2m}} A_{t_1}(x, y_1) \, d(\pi_{3,1})_{\#}(\gamma_2 * \gamma_{1,2}) - \int_{\mathbb{R}^{2m}} A_{t_2}(x, y_2) \, d\gamma_2 \\ &= \int_{\mathbb{R}^{3m}} A_{t_1}(x, y_1) - A_{t_2}(x, y_2) \, d(\gamma_2 * \gamma_{1,2}) \\ &\leqslant \int_{\mathbb{R}^{3m}} l_{K,a,b}(d(y_1, y_2) + |t_1 - t_2|) \, d(\gamma_2 * \gamma_{1,2}) \\ &= l_{K,a,b}(W_1(\nu_1, \nu_2) + |t_1 - t_2|). \end{split}$$

where $l_{K,a,b}$ denotes the Lipschitz constant of function $(t, y) \mapsto A_t(x, y)$ depending on K, a and b.

Suppose $\mu_i \in \mathscr{P}(K)$, $i = 1, 2, 3, \gamma \in \Gamma(\mu_2, \mu_3)$ and $t \in (0, \tau_1]$ fixed. By Lemma A.2 we have that for any $\gamma \in \Gamma(\mu_2, \mu_3)$ and $\mu_\lambda \in \Gamma_o^t(\mu_1, \gamma_{2\to 3}^{\lambda})$, there exists $\gamma_\lambda \in \Gamma(\mu_1, \gamma)$ satisfying $(\gamma_\lambda)_{1,2\to 3}^{\lambda} = \mu_\lambda$. We also have $(\pi_{1,2})_{\#}\gamma_\lambda \in \Gamma(\mu_1, \mu_2)$ and $(\pi_{1,3})_{\#}\gamma_\lambda \in \Gamma(\mu_1, \mu_3)$. In addition, for fixed $t \in (0, \tau_1]$, there exists some $\lambda_K > 0$ such that $K \subset B(x, \lambda_K t)$ for all $x \in K$. Now we denote $C_K := C_{\lambda_K}$ in Proposition 2.8, then

$$(1 - \lambda)C^{t}(\mu_{1}, \mu_{2}) + \lambda C^{t}(\mu_{1}, \mu_{3}) - C^{t}(\mu_{1}, \gamma_{2 \to 3}^{\lambda})$$

$$= (1 - \lambda) \inf_{\mu \in \Gamma(\mu_{1}, \mu_{2})} \int_{\mathbb{R}^{2m}} A_{t}(x, y) d\mu + \lambda \inf_{\mu \in \Gamma(\mu_{1}, \mu_{3})} \int_{\mathbb{R}^{2m}} A_{t}(x, y) d\mu$$

$$= \inf_{\mu \in \Gamma(\mu_{1}, \gamma_{2 \to 3}^{\lambda})} \int_{\mathbb{R}^{2m}} A_{t}(x, y) d(\pi_{1,2})_{\#} \gamma_{\lambda} + \int_{\mathbb{R}^{2m}} \lambda A_{t}(x, y) d(\pi_{1,3})_{\#} \gamma_{\lambda} - \int_{\mathbb{R}^{2m}} A_{t}(x, y) d\mu_{\lambda}$$

$$= \int_{\mathbb{R}^{3m}} \left[(1 - \lambda)A_{t}(x_{1}, x_{2}) + \lambda A_{t}(x_{1}, x_{3}) - A_{t}(x_{1}, (1 - \lambda)x_{2} + \lambda x_{3}) \right] d\gamma_{\lambda}$$

$$\leq \lambda (1 - \lambda) \int_{\mathbb{R}^{3m}} \frac{C_{K}}{t} |x_{2} - x_{3}|^{2} d\gamma_{\lambda}$$

$$= \lambda (1 - \lambda) \int_{\mathbb{R}^{2m}} \frac{C_{K}}{t} |x_{2} - x_{3}|^{2} d\gamma = \frac{C_{K}}{t} \lambda (1 - \lambda) W_{2,\gamma}^{2}(\mu_{2}, \mu_{3}).$$
(58)

Consequently, (58) implies $C^t(\mu, \cdot)$ is locally strongly semiconcave on $\mathscr{P}(K)$ with constant C_K/t .

Proof of Theorem 3.4. For $v, v' \in \mathscr{P}_c(\mathbb{R}^m)$, $\gamma' \in \Gamma_2(v, v')$ and $\gamma \in \Gamma_o^t(\mu, v)$, we know $(\pi_2)_{\#}\gamma = (\pi_2)_{\#}\gamma' = v$. Following from Lemma A.1, $\gamma * \gamma' \in \Gamma(\mu, v, v')$ with $(\pi_{2,3})_{\#}(\gamma * \gamma') = \gamma'$ and

 $(\pi_{1,2})_{\#}(\gamma * \gamma') = \gamma$. Simultaneously, $(\pi_{1,3})_{\#}(\gamma * \gamma') \in \Gamma(\mu, \nu')$. Here we can get

$$C^{t}(\mu, \nu') - C^{t}(\mu, \nu)$$

$$= \inf_{\mu \in \Gamma(\mu, \nu')} \int_{\mathbb{R}^{2m}} A_{t}(x, y) \, d\mu - \inf_{\mu \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^{2m}} A_{t}(x, y) \, d\mu$$

$$\leq \int_{\mathbb{R}^{2m}} A_{t}(x, y) \, d(\pi_{1,3})_{\#}(\gamma * \gamma') - \int_{\mathbb{R}^{2m}} A_{t}(x, y) \, d(\pi_{1,2})_{\#}(\gamma * \gamma')$$

$$= \int_{\mathbb{R}^{3m}} A_{t}(x_{1}, x_{3}) - A_{t}(x_{1}, x_{2}) \, d(\gamma * \gamma')$$

$$\leq \int_{\mathbb{R}^{3m}} \left(\langle L_{\nu}(\eta_{x_{1}, x_{2}}(t), \dot{\eta}_{x_{1}, x_{2}}(t)), x_{3} - x_{2} \rangle + \frac{C_{\operatorname{supp}(\gamma * \gamma')}}{2t} |x_{3} - x_{2}|^{2} \right) d(\gamma * \gamma'),$$

where $C_{\text{supp}(\gamma*\gamma')}$ is a constant depending on supp $(\gamma*\gamma')$. Recall the construction of $\gamma*\gamma'$ mentioned in the proof of Lemma A.1, $\gamma*\gamma' = \int_{\mathbb{R}^m} \gamma_{x_2} \times \gamma'_{x_2} d\nu(x_2)$, where $\gamma = \int_{\mathbb{R}^m} \gamma_{x_2} d\nu(x_2)$ and $\gamma' = \int_{\mathbb{R}^m} \gamma'_{x_2} d\nu(x_2)$. It follows

$$C^{t}(\mu,\nu') - C^{t}(\mu,\nu)$$

$$\leq \int_{\mathbb{R}^{3m}} \left(\langle p_{x_{1},x_{2}}(t), x_{3} - x_{2} \rangle + \frac{C_{\text{supp}(\gamma*\gamma')}}{2t} |x_{3} - x_{2}|^{2} \right) d(\gamma*\gamma')$$

$$= \int_{\mathbb{R}^{m}} \int_{\mathbb{R}^{2m}} \left(\langle p_{x_{1},x_{2}}(t), x_{3} - x_{2} \rangle + \frac{C_{\text{supp}(\gamma*\gamma')}}{2t} |x_{3} - x_{2}|^{2} \right) d(\gamma_{x_{2}} \times \gamma'_{x_{2}}) d\nu$$

$$= \int_{\mathbb{R}^{m}} \int_{\mathbb{R}^{m}} \left(\langle \int_{\mathbb{R}^{m}} p_{x_{1},x_{2}}(t) d\gamma_{x_{2}}, x_{3} - x_{2} \rangle + \frac{C_{\text{supp}(\gamma*\gamma')}}{2t} |x_{3} - x_{2}|^{2} \right) d\gamma'_{x_{2}} d\nu$$

$$= \int_{\mathbb{R}^{2m}} \langle \mathbf{p}_{\nu}(x_{2}), x_{3} - x_{2} \rangle d\gamma' + \frac{C_{\text{supp}(\gamma*\gamma')}}{2t} |x_{3} - x_{2}|^{2} d\gamma'$$

$$= \int_{\mathbb{R}^{2m}} \langle \mathbf{p}_{\nu}(x_{2}), x_{3} - x_{2} \rangle d\gamma' + \frac{C_{\text{supp}(\gamma*\gamma')}}{2t} W_{2}^{2}(\nu,\nu'),$$

where $p_{x,y}(t) := L_{\nu}(\eta_{x,y}(t), \dot{\eta}_{x,y}(t))$. Since our choice of $\gamma' \in \Gamma_2(\mu, \nu)$ is arbitrary, Definition 3.1 shows that $\mathbf{p}_{\nu}(x_2) \in \partial_{\nu}^+ C^t(\mu, \nu)$. $-\mathbf{p}_{\mu}(x_1) \in \partial_{\mu}^+ C^t(\mu, \nu)$ can be obtained by a similar argument. For the rest of the proof, we assume that $\gamma \in \Gamma_o^t(\mu, \nu)$ and $t' \in (t - \varepsilon, t + \varepsilon)$ with $\varepsilon > 0$,

$$C^{t'}(\mu,\nu) - C^{t}(\mu,\nu) \leq \int_{\mathbb{R}^{2m}} A_{t'}(x,y) \, d\gamma - \int_{\mathbb{R}^{2m}} A_{t}(x,y) \, d\gamma$$
$$\leq -(t'-t) \int_{\mathbb{R}^{2m}} H(\eta_{x,y}(t), p_{x,y}(t)) \, d\gamma + \frac{C_{K,t,\varepsilon}}{2} |t'-t|^2$$

where $K = \operatorname{supp} \mu \times \operatorname{supp} \nu$ is a compact subset. This estimate completes the proof.

References

- [1] Paolo Albano. Propagation of singularities for solutions of Hamilton-Jacobi equations. J. Math. Anal. Appl., 411(2):684-687, 2014.
- [2] Paolo Albano and Piermarco Cannarsa. Propagation of singularities for solutions of nonlinear first order partial differential equations. Arch. Ration. Mech. Anal., 162(1):1–23, 2002.
- [3] Paolo Albano, Piermarco Cannarsa, Khai Tien Nguyen, and Carlo Sinestrari. Singular gradient flow of the distance function and homotopy equivalence. *Math. Ann.*, 356(1):23–43, 2013.
- [4] Charalambos D. Aliprantis and Kim C. Border. *Infinite dimensional analysis: A hitchhiker's guide*. Springer, Berlin, third edition, 2006.

- [5] Luigi Ambrosio, Elia Brué, and Daniele Semola. *Lectures on optimal transport*, volume 130 of *Unitext*. Springer, 2021.
- [6] Luigi Ambrosio and Jin Feng. On a class of first order Hamilton-Jacobi equations in metric spaces. J. Differential Equations, 256(7):2194–2245, 2014.
- [7] Luigi Ambrosio and Wilfred Gangbo. Hamiltonian ODEs in the Wasserstein space of probability measures. *Comm. Pure Appl. Math.*, 61(1):18–53, 2008.
- [8] Luigi Ambrosio and Nicola Gigli. A user's guide to optimal transport. In *Modelling and optimisation of flows on networks*, volume 2062 of *Lecture Notes in Math.*, pages 1–155. Springer, Heidelberg, 2013.
- [9] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. Gradient flows in metric spaces and in the space of probability measures. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, second edition, 2008.
- [10] M.-C. Arnaud. Pseudographs and the Lax-Oleinik semi-group: a geometric and dynamical interpretation. Nonlinearity, 24(1):71–78, 2011.
- [11] Zeinab Badreddine and Hélène Frankowska. Solutions to Hamilton-Jacobi equation on a Wasserstein space. Calc. Var. Partial Differential Equations, 61(1):Paper No. 9, 41, 2022.
- [12] Victor Bangert. Analytische Eigenschaften konvexer Funktionen auf Riemannschen Mannigfaltigkeiten. J. Reine Angew. Math., 307(308):309–324, 1979.
- [13] Jean-David Benamou and Yann Brenier. A numerical method for the optimal time-continuous mass transport problem and related problems. In *Monge Ampère equation: applications to geometry and optimization (Deerfield Beach, FL, 1997)*, volume 226 of *Contemp. Math.*, pages 1–11. Amer. Math. Soc., Providence, RI, 1999.
- [14] Patrick Bernard. Existence of C^{1,1} critical sub-solutions of the Hamilton-Jacobi equation on compact manifolds. Ann. Sci. École Norm. Sup. (4), 40(3):445–452, 2007.
- [15] Patrick Bernard and Boris Buffoni. The Monge problem for supercritical Mañé potentials on compact manifolds. Adv. Math., 207(2):691–706, 2006.
- [16] Patrick Bernard and Boris Buffoni. Optimal mass transportation and Mather theory. J. Eur. Math. Soc. (JEMS), 9(1):85–121, 2007.
- [17] Patrick Bernard and Boris Buffoni. Weak KAM pairs and Monge-Kantorovich duality. In Asymptotic analysis and singularities—elliptic and parabolic PDEs and related problems, volume 47 of Adv. Stud. Pure Math., pages 397–420. Math. Soc. Japan, Tokyo, 2007.
- [18] Ugo Bessi. Hamilton-Jacobi in metric spaces with a homological term. Comm. Partial Differential Equations, 45(7):776–819, 2020.
- [19] Benoît Bonnet and Hélène Frankowska. Differential inclusions in Wasserstein spaces: the Cauchy-Lipschitz framework. J. Differential Equations, 271:594–637, 2021.
- [20] Benoît Bonnet and Hélène Frankowska. Necessary optimality conditions for optimal control problems in Wasserstein spaces. Appl. Math. Optim., 84(suppl. 2):S1281–S1330, 2021.
- [21] Benoît Bonnet and Hélène Frankowska. Semiconcavity and sensitivity analysis in mean-field optimal control and applications. J. Math. Pures Appl. (9), 157:282–345, 2022.
- [22] Yann Brenier. Décomposition polaire et réarrangement monotone des champs de vecteurs. C. R. Acad. Sci. Paris Sér. I Math., 305(19):805–808, 1987.
- [23] Yann Brenier. Polar factorization and monotone rearrangement of vector-valued functions. *Comm. Pure Appl. Math.*, 44(4):375–417, 1991.
- [24] Yann Brenier. Extended Monge-Kantorovich theory. In Optimal transportation and applications (Martina Franca, 2001), volume 1813 of Lecture Notes in Math., pages 91–121. Springer, Berlin, 2003.
- [25] Piermarco Cannarsa and Wei Cheng. Generalized characteristics and Lax-Oleinik operators: global theory. Calc. Var. Partial Differential Equations, 56(5):Art. 125, 2017.
- [26] Piermarco Cannarsa, Wei Cheng, and Albert Fathi. On the topology of the set of singularities of a solution to the Hamilton-Jacobi equation. C. R. Math. Acad. Sci. Paris, 355(2):176–180, 2017.
- [27] Piermarco Cannarsa, Wei Cheng, and Albert Fathi. Singularities of solutions of time dependent Hamilton-Jacobi equations. Applications to Riemannian geometry. *Publ. Math. Inst. Hautes Études Sci.*, 133(1):327–366, 2021.
- [28] Piermarco Cannarsa, Wei Cheng, and Jiahui Hong. Topological and control theoretic properties of Hamilton– Jacobi equations via Lax-Oleinik commutators. *Nonlinear Anal. Real World Appl.*, 84:Paper No. 104282, 2025.

- [29] Piermarco Cannarsa, Wei Cheng, Jiahui Hong, and Kaizhi Wang. Variational construction of generalized characteristics and maximal slope curve. preprint, arXiv:2409.00961, 2024.
- [30] Piermarco Cannarsa, Wei Cheng, Liang Jin, Kaizhi Wang, and Jun Yan. Herglotz' variational principle and Lax-Oleinik evolution. J. Math. Pures Appl. (9), 141:99–136, 2020.
- [31] Piermarco Cannarsa, Wei Cheng, Marco Mazzola, and Kaizhi Wang. Global generalized characteristics for the Dirichlet problem for Hamilton-Jacobi equations at a supercritical energy level. SIAM J. Math. Anal., 51(5):4213– 4244, 2019.
- [32] Piermarco Cannarsa and Carlo Sinestrari. Semiconcave functions, Hamilton-Jacobi equations, and optimal control, volume 58 of Progress in Nonlinear Differential Equations and their Applications. Birkhäuser Boston, Inc., Boston, MA, 2004.
- [33] Piermarco Cannarsa and Yifeng Yu. Singular dynamics for semiconcave functions. J. Eur. Math. Soc. (JEMS), 11(5):999–1024, 2009.
- [34] Pierre Cardaliaguet and Panagiotis E. Souganidis. Regularity of the value function and quantitative propagation of chaos for mean field control problems. *NoDEA Nonlinear Differential Equations Appl.*, 30(2):Paper No. 25, 37, 2023.
- [35] Marco Castelpietra and Ludovic Rifford. Regularity properties of the distance functions to conjugate and cut loci for viscosity solutions of Hamilton-Jacobi equations and applications in Riemannian geometry. ESAIM Control Optim. Calc. Var., 16(3):695–718, 2010.
- [36] Cui Chen, Wei Cheng, and Qi Zhang. Lasry-Lions approximations for discounted Hamilton-Jacobi equations. J. Differential Equations, 265(2):719–732, 2018.
- [37] Wei Cheng, Jiahui Hong, and Tianqi Shi. Optimal transport in the frame of abstract lax-oleinik operator revisited,. preprint, arXiv:2402.04159, 2024.
- [38] Francis Clarke. Functional analysis, calculus of variations and optimal control, volume 264 of Graduate Texts in Mathematics. Springer, London, 2013.
- [39] Andrea Cosso, Fausto Gozzi, Idris Kharroubi, Huyên Pham, and Mauro Rosestolato. Master Bellman equation in the Wasserstein space: uniqueness of viscosity solutions. *Trans. Amer. Math. Soc.*, 377(1):31–83, 2024.
- [40] Constantine M. Dafermos. Hyperbolic conservation laws in continuum physics, volume 325 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, second edition, 2005.
- [41] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
- [42] Shi-Zhong Du and Qi-Rui Li. Positivity of Ma-Trudinger-Wang curvature on Riemannian surfaces. *Calc. Var. Partial Differential Equations*, 51(3-4):495–523, 2014.
- [43] Albert Fathi. Weak KAM theorem in Lagrangian dynamics. Cambridge University Press, Cambridge (to appear).
- [44] Albert Fathi. Théorème KAM faible et théorie de Mather sur les systèmes lagrangiens. C. R. Acad. Sci. Paris Sér. I Math., 324(9):1043–1046, 1997.
- [45] Albert Fathi. Weak KAM from a PDE point of view: viscosity solutions of the Hamilton-Jacobi equation and Aubry set. Proc. Roy. Soc. Edinburgh Sect. A, 142(6):1193–1236, 2012.
- [46] Albert Fathi. Viscosity solutions of the Hamilton-Jacobi equation on a noncompact manifold. In *Hamiltonian systems: dynamics, analysis, applications*, volume 72 of *Math. Sci. Res. Inst. Publ.*, pages 111–172. Cambridge Univ. Press, Cambridge, 2024.
- [47] Albert Fathi and Alessio Figalli. Optimal transportation on non-compact manifolds. *Israel J. Math.*, 175:1–59, 2010.
- [48] Albert Fathi and Ezequiel Maderna. Weak KAM theorem on non compact manifolds. NoDEA Nonlinear Differential Equations Appl., 14(1-2):1–27, 2007.
- [49] Jin Feng and Andrzej Święch. Optimal control for a mixed flow of Hamiltonian and gradient type in space of probability measures. *Trans. Amer. Math. Soc.*, 365(8):3987–4039, 2013. With an appendix by Atanas Stefanov.
- [50] A. Figalli, L. Rifford, and C. Villani. Tangent cut loci on surfaces. Differential Geom. Appl., 29(2):154–159, 2011.
- [51] A. Figalli, L. Rifford, and C. Villani. Nearly round spheres look convex. Amer. J. Math., 134(1):109–139, 2012.
- [52] Alessio Figalli and Ludovic Rifford. Continuity of optimal transport maps and convexity of injectivity domains on small deformations of S². Comm. Pure Appl. Math., 62(12):1670–1706, 2009.

- [53] W. Gangbo, T. Nguyen, and A. Tudorascu. Euler-Poisson systems as action-minimizing paths in the Wasserstein space. Arch. Ration. Mech. Anal., 192(3):419–452, 2009.
- [54] W. Gangbo and A. Tudorascu. Lagrangian dynamics on an infinite-dimensional torus; a weak KAM theorem. *Adv. Math.*, 224(1):260–292, 2010.
- [55] Wilfrid Gangbo and Robert J. McCann. The geometry of optimal transportation. Acta Math., 177(2):113–161, 1996.
- [56] Wilfrid Gangbo, Truyen Nguyen, and Adrian Tudorascu. Hamilton-Jacobi equations in the Wasserstein space. *Methods Appl. Anal.*, 15(2):155–183, 2008.
- [57] Wilfrid Gangbo and Andrzej Święch. Existence of a solution to an equation arising from the theory of mean field games. J. Differential Equations, 259(11):6573–6643, 2015.
- [58] Wilfrid Gangbo and Andrzej Święch. Metric viscosity solutions of Hamilton-Jacobi equations depending on local slopes. *Calc. Var. Partial Differential Equations*, 54(1):1183–1218, 2015.
- [59] Wilfrid Gangbo and Adrian Tudorascu. A weak KAM theorem; from finite to infinite dimension. In *Optimal transportation, geometry and functional inequalities*, volume 11 of *CRM Series*, pages 45–72. Ed. Norm., Pisa, 2010.
- [60] Wilfrid Gangbo and Adrian Tudorascu. On differentiability in the Wasserstein space and well-posedness for Hamilton-Jacobi equations. J. Math. Pures Appl. (9), 125:119–174, 2019.
- [61] Yoshikazu Giga, Nao Hamamuki, and Atsushi Nakayasu. Eikonal equations in metric spaces. Trans. Amer. Math. Soc., 367(1):49–66, 2015.
- [62] Nicola Gigli, Kazumasa Kuwada, and Shin-Ichi Ohta. Heat flow on Alexandrov spaces. Comm. Pure Appl. Math., 66(3):307–331, 2013.
- [63] Ryan Hynd and Hwa Kil Kim. Infinite horizon value functions in the Wasserstein spaces. J. Differential Equations, 258(6):1933–1966, 2015.
- [64] Ryan Hynd and Hwa Kil Kim. Value functions in the Wasserstein spaces: finite time horizons. J. Funct. Anal., 269(4):968–997, 2015.
- [65] Jin-ichi Itoh and Minoru Tanaka. The Lipschitz continuity of the distance function to the cut locus. Trans. Amer. Math. Soc., 353(1):21–40, 2001.
- [66] Konstantin Khanin and Andrei Sobolevski. On dynamics of Lagrangian trajectories for Hamilton-Jacobi equations. Arch. Ration. Mech. Anal., 219(2):861–885, 2016.
- [67] M. Knott and C. S. Smith. On the optimal mapping of distributions. J. Optim. Theory Appl., 43(1):39–49, 1984.
- [68] Kazumasa Kuwada. Duality on gradient estimates and Wasserstein controls. J. Funct. Anal., 258(11):3758–3774, 2010.
- [69] Y. Li and L. Nirenberg. The distance function to the boundary, Finsler geometry, and the singular set of viscosity solutions of some Hamilton-Jacobi equations. *Comm. Pure Appl. Math.*, 58(1):85–146, 2005.
- [70] Grégoire Loeper and Cédric Villani. Regularity of optimal transport in curved geometry: the nonfocal case. Duke Math. J., 151(3):431–485, 2010.
- [71] Carlo Mantegazza and Andrea Carlo Mennucci. Hamilton-Jacobi equations and distance functions on Riemannian manifolds. Appl. Math. Optim., 47(1):1–25, 2003.
- [72] Robert J. McCann. A convexity principle for interacting gases. Adv. Math., 128(1):153–179, 1997.
- [73] Robert J. McCann. Polar factorization of maps on Riemannian manifolds. *Geom. Funct. Anal.*, 11(3):589–608, 2001.
- [74] Atsushi Nakayasu and Tokinaga Namba. Stability properties and large time behavior of viscosity solutions of Hamilton-Jacobi equations on metric spaces. *Nonlinearity*, 31(11):5147–5161, 2018.
- [75] Shin-ichi Ohta. Uniform convexity and smoothness, and their applications in Finsler geometry. *Math. Ann.*, 343(3):669–699, 2009.
- [76] Felix Otto. Dynamics of labyrinthine pattern formation in magnetic fluids: a mean-field theory. Arch. Rational Mech. Anal., 141(1):63–103, 1998.
- [77] Felix Otto. Evolution of microstructure in unstable porous media flow: a relaxational approach. Comm. Pure Appl. Math., 52(7):873–915, 1999.
- [78] Felix Otto. The geometry of dissipative evolution equations: the porous medium equation. Comm. Partial Differential Equations, 26(1-2):101–174, 2001.

PIERMARCO CANNARSA, WEI CHENG, TIANQI SHI AND WENXUE WEI

- [79] Thomas Strömberg. Propagation of singularities along broken characteristics. Nonlinear Anal., 85:93–109, 2013.
- [80] Cédric Villani. *Optimal transport: old and new*, volume 338 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 2009.

(Piermarco Cannarsa) Dipartimento di Matematica, Università di Roma "Tor Vergata", Via della Ricerca Scientifica 1, 00133 Roma, Italy

Email address: cannarsa@mat.uniroma2.it

(Wei Cheng) School of Mathematics, Nanjing University, Nanjing 210093, China *Email address*: chengwei@nju.edu.cn

(Tianqi Shi) Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China *Email address*: tqshi.math@gmail.com

(Wenxue Wei) School of Mathematics, Nanjing University, Nanjing 210093, China *Email address:* wwx3708@gmail.com