# THE ČECH HOMOTOPY GROUPS OF A SHRINKING WEDGE OF SPHERES

JEREMY BRAZAS

ABSTRACT. We compute the Čech homotopy groups of the *m*-dimensional infinite earring space  $\mathbb{E}_m$ , i.e. a shrinking wedge of *m*-spheres. In particular, for all  $n, m \ge 2$ , we prove that  $\check{\pi}_n(\mathbb{E}_m)$  is isomorphic to a direct sum of countable powers of homotopy groups of spheres:  $\bigoplus_{1 \le j \le \frac{n-1}{m-1}} (\pi_n(S^{mj-j+1}))^{\mathbb{N}}$ . Equipped with this isomorphism and infinite-sum algebra, we also construct new elements of  $\pi_n(\mathbb{E}_m)$  with a view toward characterizing the image of the canonical homomorphism  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$ . We prove that  $\Psi_n$  is a split epimorphism when  $n \le 2m-1$  and we identify a candidate for the image of  $\Psi_n$  when n > 2m - 1.

## 1. INTRODUCTION

In 1962, Barratt and Milnor [1] proved that the rational homology groups of a shrinking wedge of *m*-spheres  $(m \ge 2)$ , which we refer to as the *m*-dimensional earring space  $\mathbb{E}_m$ , are uncountable in arbitrarily high dimensions. In 2000, Eda and Kawamura proved that  $\mathbb{E}_m$  is (m-1)-connected and locally (m-1)-connected and that  $\pi_m(\mathbb{E}_m)$  is canonically isomorphic to the Baer-Specker group  $\mathbb{Z}^{\mathbb{N}}$ . Built into the proof of Eda-Kawamura's Theorem is the striking fact that higher homotopy groups are "infinitely commutative" in a strong sense.

It remains an open problem to give a full description of the groups  $\pi_n(\mathbb{E}_m)$ , n > m in the spirit of the Hilton-Milnor Theorem. In this paper we give a complete description of the Čech homotopy groups  $\check{\pi}_n(\mathbb{E}_m)$  in terms of the homotopy groups of spheres for all  $n, m \ge 2$ . We build off of the computation of  $\check{\pi}_{m+1}(\mathbb{E}_m), m \ge 2$  established by K. Kawamura in [7]. Our first main result is the following.

**Theorem 1.1.** For all  $m, n \ge 2$ , there is a isomorphism

$$\check{\pi}_n(\mathbb{E}_m) \cong \bigoplus_{1 \le j \le \frac{n-1}{m-1}} \left( \pi_n(S^{mj-j+1}) \right)^{\mathbb{N}}.$$

The isomorphism in Theorem 1.1 is not canonical but depends only an enumeration of the union of coherently chosen Hall bases  $H_k$  on finite sets  $A_k = \{a_1, \ldots, a_k\}$ . The upper bound  $\frac{n-1}{m-1}$  on the index j simply indicates when the homotopy groups of spheres might be non-trivial (if  $j > \frac{n-1}{m-1}$ , then n < mj-j+1 and  $\pi_n(S^{mj-j+1}) = 0$ ). Low-dimensional cases of special interest are highlighted in Example 3.14.

Date: January 28, 2025.

<sup>2010</sup> Mathematics Subject Classification. Primary 55Q07,55Q52; Secondary 55Q20,55Q15,08A65.

Key words and phrases. Čech homotopy group, shape homotopy group, Hilton-Milnor Theorem, homotopy groups of spheres, n-dimensional infinite earring.

When n > m, the group  $\pi_n(\mathbb{E}_m)$  is potentially more complicated than  $\check{\pi}_n(\mathbb{E}_m)$ . However, there is a canonical homomorphism  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$  relating the two. Using the isomorphism in Theorem 1.1 as a book-keeping device, we construct and distinguish new elements of  $\pi_n(\mathbb{E}_m)$  in an effort to characterize the image of  $\Psi_n$ . We do not attempt to address whether or not  $\Psi_n$  is injective. Our second theorem establishes the surjectivity of  $\Psi_n$  in a certain range of dimensions.

**Theorem 1.2.** If  $m, n \ge 2$  and  $n \le 2m - 1$ , then the canonical homomorphism  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$  is a split epimorphism.

Theorem 1.2 is proved in two cases. The stable range (n < 2m - 1) is proved in Corollary 4.3 where  $\check{\pi}_n(\mathbb{E}_m) \cong \pi_n(S^m)^{\mathbb{N}}$ . The case n = 2m - 1 is proved in Lemma 4.4 where  $\check{\pi}_{2m-1}(\mathbb{E}_m) \cong \pi_{2m-1}(S^m)^{\mathbb{N}} \oplus \mathbb{Z}^{\mathbb{N}}$ . The image of the splitting homomorphism  $\check{\pi}_{2m-1}(\mathbb{E}_m) \to \pi_{2m-1}(\mathbb{E}_m)$  constructed in Lemma 4.4 meets the kernel of the homomorphism induced by the canonical embedding  $\sigma : \mathbb{E}_m \to \prod_{j=1}^{\infty} S^m$  precisely in the subgroup  $\mathcal{W}_{2n-1}(\mathbb{E}_n)$  that is computed in [2]. Moreover, the author has suggested in [2, Conjecture 1.3] that  $\mathcal{W}_{2n-1}(\mathbb{E}_n)$  is equal to  $\ker(\sigma_{\#})$ . It follows from the standard splitting of  $\sigma_{\#}$  reviewed in Section 2.3 that [2, Conjecture 1.3] is equivalent to conjecturing that  $\Psi_{2m-1} : \pi_{2m-1}(\mathbb{E}_m) \to \check{\pi}_{2m-1}(\mathbb{E}_m)$  is an isomorphism for all  $m \ge 2$ .

Although the surjectivity of  $\Psi_n$  remains unclear when n > 2m - 1, we construct a candidate for  $\text{Im}(\Psi_n)$  in this range with our third theorem.

**Theorem 1.3.** If n > 2m - 1, then  $\pi_n(\mathbb{E}_m)$  contains a subgroup isomorphic to

$$\prod_{\mathbb{N}} \pi_n(S^m) \oplus \bigoplus_{2 \le j \le \frac{n-1}{m-1}} \left( \prod_{\mathbb{N}} \bigoplus_{\mathbb{N}} \pi_n(S^{mj-j+1}) \right)$$

on which the canonical homomorphism  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$  is injective.

The author is grateful to Kazuhiro Kawamura for helpful conversations regarding the content of this paper.

## 2. Preliminaries

All topological spaces in this paper are assumed to be Hausdorff. Throughout, I will denote the closed unit interval,  $S^n$  will denote the unit *n*-sphere with basepoint  $(1, 0, 0, \ldots, 0)$ . We only consider higher homotopy groups in this paper and so we generally assume  $n \ge 2$ . For a based topological space  $(X, x_0)$ , we write  $\Omega^n(X, x_0)$  to denote the space of relative maps  $(I^n, \partial I^n) \to (X, x_0)$  and  $\pi_n(X, x_0) = \{[f] \mid f \in \Omega^n(X, x_0)\}$  to denote the *n*-th homotopy group. When the basepoint is clear from context, we may simplify this notation to  $\Omega^n(X)$  and  $\pi_n(X)$ . Fixing a based homeomorphism  $I^n/\partial I^n \cong S^n$ , we identify  $\Omega^n(X)$  with the space of based maps  $S^n \to X$ . For  $f \in \Omega^n(X)$ , let -f denote the reverse map given by  $-f(t_1, t_2, \ldots, t_n) = f(1 - t_1, t_2, \ldots, t_n)$ . We write  $f_1 + f_2 + \cdots + f_m$  for the usual first-coordinate concatenation of maps  $f_i \in \Omega^n(X)$ . If m is a positive integer, then mf denotes the m-fold sum  $f + f + \cdots + f$ , 0f denotes the constant map  $c \in \Omega^n(X)$ , and if m is a negative integer, then mf denotes the (-m)-fold sum  $(-f) + (-f) + \cdots + (-f)$ .

2.1. Whitehead Products. We use commutator bracket notation to denote Whitehead products both on the level of maps and homotopy classes. For an ordered pair of natural numbers  $(p,q) \in \mathbb{N}^2$ , we view  $S^p \vee S^q$  naturally as a subspace of  $S^p \times S^q$ . Let  $a_{p,q} : S^{p+q-1} \to S^p \vee S^q$  denote an attaching map of the unique p + q-cell in  $S^p \times S^q$ . If  $f : S^p \to X$  and  $g : S^q \to X$  are based maps, the composition  $[f,g] = (f \vee g) \circ a_{p,q} : S^{p+q-1} \to X$  is the Whitehead product of the pair of maps (f,g). This operation on maps descends to a well-defined bracket operation  $[-, -] : \pi_p(X) \times \pi_q(X) \to \pi_{p+q-1}(X)$  on homotopy classes, which we also refer to as the Whitehead product.

**Remark 2.1.** Whitehead products satisfy the following standard relations [10]. Let  $p, q, r \ge 2$ . For  $\alpha, \alpha' \in \pi_p(X), \beta, \beta' \in \pi_q(X)$ , and  $\gamma \in \pi_r(X)$ , we have:

- (identity) [a, 0] = 0 and [0, b] = 0,
- (bilinearity)  $[\alpha + \alpha', \beta] = [\alpha, \beta] + [\alpha', \beta]$  and  $[\alpha, \beta + \beta'] = [\alpha, \beta] + [\alpha, \beta']$ ,
- (graded symmetry)  $[\alpha, \beta] = (-1)^{pq} [\beta, \alpha],$
- (graded Jacobi identity)

$$(-1)^{pr}[[\alpha,\beta],\gamma] + (-1)^{pq}[[\beta,\gamma],\alpha] + (-1)^{rq}[[\gamma,\alpha],\beta] = 0.$$

We have need to apply some of the standard relations infinitely many times in a single line of algebraic reasoning. To do this it is necessary to know that whenever representing maps are "small" we can apply these relations using "small" homotopies. The following Proposition is an observation made in [2].

**Proposition 2.2.** Suppose X is a based space,  $p, q \ge 2$ ,  $f, f' \in \Omega^p(X)$ , and  $g, g' \in \Omega^q(X)$ .

- (1) If H is a homotopy from f to f' and G is a homotopy from g to g', then  $[f,g] \simeq [f',g']$  by a homotopy with image equal to  $\operatorname{Im}(H) \cup \operatorname{Im}(G)$ ,
- (2) If f or g is constant, then [f, g] is null-homotopic in  $\text{Im}(f) \cup \text{Im}(g)$ ,
- (3) We have  $[f + f', g] \simeq [f, g] + [f', g]$  by a homotopy with image  $\operatorname{Im}(f) \cup \operatorname{Im}(f') \cup \operatorname{Im}(g)$  and  $[f, g + g'] \simeq [f, g] + [f, g']$  by a homotopy with image  $\operatorname{Im}(f) \cup \operatorname{Im}(g) \cup \operatorname{Im}(g')$ .

A based map  $f : X \to Y$  induces a homomorphism  $f_{\#} : \pi_n(X) \to \pi_n(Y)$  and [f] will denote the based homotopy class of f. Recall that ordinary composition  $f \circ g$  of maps descends to composition  $[f] \circ [g]$  of homotopy classes in the homotopy category.

**Remark 2.3.** If  $h: X \to Y$ ,  $f: S^p \to X$ , and  $g: S^q \to X$  are based maps, then we have equality  $h \circ [f,g] = [h \circ f, h \circ g]$  on the level of continuous maps. For homotopy classes a = [f] and b = [g], we have  $h_{\#}([a,b]) = [h_{\#}(a), h_{\#}(b)]$ .

## 2.2. Infinite sums.

**Definition 2.4.** Let  $(X, x_0)$  be a based space. We say that a sequence  $\{f_j\}_{j \in \mathbb{N}}$  of based maps  $f_j : W_j \to X$  converges to  $x_0$  if for every neighborhood U of  $x_0$ , we have  $\operatorname{Im}(f_j) \subseteq U$  for all but finitely many  $j \in \mathbb{N}$ . The *infinite sum* of a sequence  $\{f_j\}_{j \in \mathbb{N}}$  of based maps  $f_j : S^n \to X$  that converges to  $x_0$  is the map  $\sum_{j=1}^{\infty} f_j \in \Omega^n(X, x_0)$  defined as  $f_j$  on  $\left[\frac{j-1}{j}, \frac{j}{j+1}\right] \times I^{n-1}$  for all  $j \in \mathbb{N}$  and which maps  $\{1\} \times I^{n-1}$  to  $x_0$ .

Infinite sums of maps  $S^n \to X$  are infinitely commutative up to homotopy [3] and thus satisfy a variety of identities. We refer to [2] for the proofs of the next four Propositions.

**Proposition 2.5.** [2] If  $\{f_j\}_{j\in\mathbb{N}}$ ,  $\{g_j\}_{j\in\mathbb{N}}$  are sequences of maps  $S^n \to X$  the converge to  $x_0 \in X$  and  $a, b \in \mathbb{Z}$ , then  $\sum_{j=1}^{\infty} af_j + bg_j \simeq a \sum_{j=1}^{\infty} f_j + b \sum_{j=1}^{\infty} g_j$  by a homotopy with image in  $\bigcup_{j\in\mathbb{N}} \operatorname{Im}(f_j) \cup \operatorname{Im}(g_j)$ .

**Definition 2.6.** Let  $\{f_j\}_{j\in\mathbb{N}}$  and  $\{g_j\}_{j\in\mathbb{N}}$  be sequences of maps  $S^n \to X$  that converge to  $x_0 \in X$ . We say that  $\{f_j\}_{j\in\mathbb{N}}$  and  $\{g_j\}_{j\in\mathbb{N}}$  are sequentially homotopic and we write  $\{f_j\}_{j\in\mathbb{N}} \simeq \{g_j\}_{j\in\mathbb{N}}$  if there exists a sequence  $\{H_j\}_{j\in\mathbb{N}}$  where  $H_j : S^n \times I \to X$  is a based homotopy from  $f_j$  to  $g_j$  (for each  $j \in \mathbb{N}$ ) such that  $\{H_j\}_{j\in\mathbb{N}}$  converges to  $x_0$ .

**Proposition 2.7.** [2] If  $\{H_j\}_{j\in\mathbb{N}}$  is a sequential homotopy between sequences  $\{f_j\}_{j\in\mathbb{N}}$ and  $\{g_j\}_{j\in\mathbb{N}}$  of maps  $f_j, g_j \in \Omega^n(X)$  that converge to  $x_0$ , then  $\sum_{j=1}^{\infty} f_j \simeq \sum_{j=1}^{\infty} g_j$ by a homotopy with image  $\bigcup_{j\in\mathbb{N}} \operatorname{Im}(H_j)$ .

We also require the following relations for combinations of infinite sums and Whitehead products.

**Proposition 2.8.** [2] Let  $(X, x_0)$  be a based space and  $p, q \ge 2$ . Suppose  $\{f_j\}_{j\in\mathbb{N}}$ ,  $\{g_j\}_{j\in\mathbb{N}}$  are sequences of maps  $S^p \to X$  that converge to  $x_0$  and  $\{h_j\}_{j\in\mathbb{N}}, \{k_j\}_{j\in\mathbb{N}}$  are sequences of maps  $S^q \to X$  that converge to  $x_0$ . Then  $\{[f_j, h_j]\}_{j\in\mathbb{N}}$  and  $\{[g_j, k_j]\}_{j\in\mathbb{N}}$  are sequences of maps  $S^{p+q-1} \to X$  that converge to  $x_0$ . Moreover, if  $\{f_j\}_{j\in\mathbb{N}} \simeq \{g_j\}_{j\in\mathbb{N}}$  and  $\{h_j\}_{j\in\mathbb{N}} \simeq \{k_j\}_{j\in\mathbb{N}}$ , then  $\{[f_j, h_j]\}_{j\in\mathbb{N}} \simeq \{[g_j, k_j]\}_{j\in\mathbb{N}}$  and  $\sum_{j=1}^{\infty} [f_j, h_j] \simeq \sum_{j=1}^{\infty} [g_j, k_j]$ .

**Proposition 2.9.** [2] Let  $(X, x_0)$  be a based space and  $p, q \ge 2$ . Suppose  $\{f_j\}_{j \in \mathbb{N}}$ ,  $\{g_j\}_{j \in \mathbb{N}}$  are sequences of maps  $S^p \to X$  that converge to  $x_0$  and  $\{h_j\}_{j \in \mathbb{N}}$ ,  $\{k_j\}_{j \in \mathbb{N}}$  are sequences of maps  $S^q \to X$  that converge to  $x_0$ . Then

- (1)  $\sum_{j=1}^{\infty} [f_j + g_j, h_j] \simeq \sum_{j=1}^{\infty} [f_j, h_j] + \sum_{j=1}^{\infty} [g_j, h_j]$  by a homotopy with image in  $\bigcup_{j \in \mathbb{N}} \operatorname{Im}(f_j) \cup \operatorname{Im}(g_j) \cup \operatorname{Im}(h_j),$
- (2)  $\sum_{j=1}^{\infty} [f_j, h_j + k_j] \simeq \sum_{j=1}^{\infty} [f_j, h_j] + \sum_{j=1}^{\infty} [f_j, k_j] \text{ by a homotopy with image}$  $in \bigcup_{i \in \mathbb{N}} \operatorname{Im}(f_j) \cup \operatorname{Im}(h_j) \cup \operatorname{Im}(k_j).$

## 2.3. Shrinking wedges and their Čech homotopy groups.

**Definition 2.10.** The *shrinking wedge* of countable set  $\{X_j\}_{j\in J}$  of based spaces is the space  $\bigvee_{j\in J} X_j$  whose underlying set is the usual one-point union  $\bigvee_{j\in J} X_j$  with canonical basepoint  $b_0$ . A set U is open in  $\bigvee_{j\in J} X_j$  if

- $U \cap X_j$  is open in  $X_j$  for all  $j \in J$ ,
- and whenever  $b_0 \in U$ , we have  $X_j \subseteq U$  for all but finitely many  $j \in J$ .

We refer to the special case  $\mathbb{E}_m = \bigvee_{i \in \mathbb{N}} S^m$  as the *m*-dimensional earring space.

We note two other useful ways to construct a shrinking wedge  $X = \bigvee_{j \in J} X_j$ . First, X canonically embeds in the direct product  $\prod_{j \in J} X_j$ . We let  $\sigma : X \to \prod_{j \in J} X_j$  denote the embedding and often identify X with its image under  $\sigma$  so that  $\sigma$  is an inclusion map. Second, is construction of  $\bigvee_{j \in J} X_j$  as an inverse limit. Let  $X_{\leq k} = \bigvee_{j=1}^k X_j$  be the wedge of the first k-many spaces and  $b_{k+1,k} : X_{\leq k+1} \to X_{\leq k}$  denote the map that collapses  $X_{k+1}$  to the wedgepoint. The projection maps  $b_k : X \to X_{\leq k}$  collapsing the sub-shrinking wedge  $X_{\geq k+1} = \bigvee_{j>k} X_j$  to the basepoint induce a homeomorphism  $X \to \varprojlim_k (X_{\leq k}, b_{k+1,k})$ . **Theorem 2.11.** [7] Let  $X = \bigvee_{j \in \mathbb{N}} X_j$  be the shrinking wedge of a sequence of path-connected, based spaces  $\{X_j\}_{j \in \mathbb{N}}$ . For all  $n \ge 2$ , the long exact sequence of the pair  $(\prod_{i \in \mathbb{N}} X_j, X)$  gives short exact sequences that split naturally:

$$0 \longrightarrow \pi_{n+1}(\prod_{j \in \mathbb{N}} X_j, X) \xrightarrow{\partial} \pi_n(X) \xrightarrow{\sigma_{\#}} \pi_n(\prod_{j \in \mathbb{N}} X_j) \longrightarrow 0.$$

Moreover, the splitting homomorphism  $\zeta : \pi_n(\prod_{j \in \mathbb{N}} X_j) \to \pi_n(X)$  is given by the infinite sum operation  $\zeta([(f_j)_{j \in \mathbb{N}}]) = \left[\sum_{j=1}^{\infty} f_j\right]$  where  $(f_j)_{j \in \mathbb{N}}$  denotes the map  $S^n \to \prod_{j \in \mathbb{N}} X_j$ , which is  $f_j$  in the j-th coordinate.

Theorem 2.11 implies that  $\pi_n(\bigvee_{i\in\mathbb{N}}X_i)$  is isomorphic to:

$$\pi_{n+1}\left(\prod_{j\in\mathbb{N}}X_j,X\right)\oplus\pi_n\left(\prod_{j\in\mathbb{N}}X_j\right)\cong\pi_{n+1}\left(\prod_{j\in\mathbb{N}}X_j,X\right)\oplus\prod_{j\in\mathbb{N}}\pi_n(X_j)$$

Hence, to characterize the group  $\pi_n(X)$ ,  $n \ge 2$  in terms of the homotopy groups of  $X_j$ , it suffices to characterize the relative homotopy group  $\pi_{n+1}(\prod_{j\in\mathbb{N}} X_j, X)$ . A common tactic for distinguishing elements of such groups comes from shape theory [8].

While the Cech homotopy groups (sometimes called the shape homotopy groups) of any based space is well-defined, we have a simple description in the following case: if  $X = \bigvee_{j \in \mathbb{N}}$  is a shrinking wedge of path-connected, compact, based spaces, then the inverse limit group  $\check{\pi}_n(X) = \varprojlim_k(\pi_n(X_{\leq k}), b_{k+1,k\#})$  is the *n*-th Čech homotopy group of X (this definition is justified by [8, Ch I, §5.4, Theorem 13]). The maps  $b_k : X \to X_{\leq k}$  induced a canonical homomorphism  $\Psi_n : \pi_n(X) \to \check{\pi}_n(X)$ ,  $\Psi_n([f]) = ([b_k \circ f])_{k \in \mathbb{N}}$ , which we call the *n*-th Čech homotopy map.

**Lemma 2.12.** For any sequence of based, path-connected, compact Hausdorff spaces  $\{X_j\}_{j \in \mathbb{N}}$  and  $n \ge 2$ , there is an exact sequence

$$0 \longrightarrow \check{\pi}_{n+1} \left( \prod_{j \in \mathbb{N}} X_j, X \right) \xrightarrow{\check{\partial}} \check{\pi}_n(X) \xrightarrow{\check{\sigma}} \pi_n(\prod_{j \in \mathbb{N}} X_j) \longrightarrow 0$$

that splits naturally.

*Proof.* We utilize the notation above for finite wedge  $X_{\leq k} = \bigvee_{j=1}^{k} X_j$  and bonding maps  $b_{k+1,k}$ . Let  $\sigma_k : X_{\leq k} \to \prod_{j=1}^{k} X_j$  be the canonical embedding and let  $s_k : \pi_n(\prod_{j=1}^{k} X_j) \to \pi_n(X_{\leq k})$  be the canonical section to the induced homomorphism  $\sigma_{k\#} : \pi_n(X_{\leq k}) \to \pi_n(\prod_{j=1}^{k} X_j)$ . We have an inverse sequence of short exact sequences with bonding maps appearing vertically in the following diagram:

$$0 \longrightarrow \pi_{n+1}(\prod_{j=1}^{k+1} X_j, X_{\leq k+1}) \xrightarrow{\sigma_{k+1}} \pi_n(X_{\leq k+1}) \xrightarrow{\sigma_{k+1\#}} \pi_n(\prod_{j=1}^{k+1} X_j) \longrightarrow 0$$

$$\downarrow^{p_{k+1,k\#}} \qquad \qquad \qquad \downarrow^{b_{k+1,k\#}} \qquad \qquad \downarrow^{p_{k+1,k\#}} \qquad \qquad \downarrow^{p_{k+1,k\#}}$$

$$0 \longrightarrow \pi_{n+1}(\prod_{j=1}^k X_j, X_{\leq k}) \xrightarrow{\partial_k} \pi_n(X_{\leq k}) \xrightarrow{\sigma_{k\#}} \pi_n(\prod_{j=1}^k X_j) \longrightarrow 0$$

In the above diagram,  $p_{k+1,k}$  is the projection map and  $\partial_k$  is the usual boundary map. The vertical morphism of exact sequences admits a canonical section induced by the based inclusion maps  $\prod_{j=1}^{k} X_j \to \prod_{j=1}^{k+1} X_j$ . Hence, the inverse sequence satisfies the Mittag-Leffler condition and, in the limit, we obtain an exact sequence.

Set  $\check{\partial} = \varprojlim_k \partial$  and  $\check{\sigma} = \varprojlim_k \sigma_{k\#}$ . Since the space  $X_j$  are compact Hausdorff, the pairs  $(\prod_{j=1}^k X_j, X_{\leq k}), k \in \mathbb{N}$  form an  $\mathbf{HPol}_*^2$ -expansion for the pair  $(\prod_{j\in\mathbb{N}} X_j, X)$ . Hence, we may identify  $\check{\pi}_{n+1} (\prod_{j\in\mathbb{N}} X_j, X)$  with  $\varprojlim_k \pi_{n+1} (\prod_{j=1}^k X_j, X_{\leq k})$  [8, Ch I, §5.4, Theorem 13]. Under this identification, the inverse limit of exact sequences is the desired sequence in the statement of the lemma. Since the canonical sections  $s_k$  of the maps  $\sigma_{k\#}$  are natural,  $\varprojlim_k s_k$  is a section of  $\check{\sigma}$ .

Lemma 2.12 implies that  $\check{\pi}_n\left(\bigvee_{j\in\mathbb{N}}X_j\right)$  splits as

$$\check{\pi}_n\left(\bigvee_{j\in\mathbb{N}}X_j\right)\cong\check{\pi}_{n+1}\left(\prod_{j\in\mathbb{N}}X_j,\bigvee_{j\in\mathbb{N}}X_j\right)\oplus\pi_n\left(\prod_{j\in\mathbb{N}}X_j\right).$$

We refer to the image of the splitting map  $\pi_n\left(\prod_{j\in\mathbb{N}}X_j\right) \to \check{\pi}_n\left(\bigvee_{j\in\mathbb{N}}X_j\right)$  as the weight-1 summand of  $\check{\pi}_n\left(\bigvee_{j\in\mathbb{N}}X_j\right)$ . This terminology will be justified later on. We also obtain the following morphism of exact sequences induced by the projection maps  $b_k$ .

If we identify  $\pi_n(\bigvee_{j\in\mathbb{N}} X_j)$  and  $\check{\pi}_n(\bigvee_{j\in\mathbb{N}} X_j)$  with their splittings, we see that  $\Psi_n$  is the identity on  $\pi_n(\prod_{j\in\mathbb{N}} X_j)$ .

## 3. Applying the Hilton-Milnor Theorem

For the remainder of the paper, fix monotone sequence of integers  $1 \leq r_1 \leq r_2 \leq r_3 \leq \cdots$  and set  $X = \bigvee_{i \in \mathbb{N}} S^{r_i+1}$  with wedgepoint  $x_0$ . Later, we will restrict to the case where  $r_i = m-1$  for all i so that  $X = \mathbb{E}_m$ . As before, let  $X_{\leq k} = \bigvee_{i=1}^k S^{r_i+1}$  be the finite wedge of the first k-spheres so that  $X = \lim_{k \to \infty} (X_{\leq k}, b_{k+1,k})$  with bonding maps  $b_{k+1,k} : X_{\leq k+1} \to X_{\leq k}$  and projection maps and  $b_k : X \to X_{\leq k}$ . We also write  $X_{\geq k} = \bigvee_{i=k}^{\infty} S^{r_i+1}$  for the sub-shrinking wedge that excludes the first (k-1) wedge-summands. Our first aim is to describe the n-th Čech homotopy group  $\check{\pi}_n(X) = \lim_{k \to \infty} (\pi_n(X_{\leq k}), b_{k+1,k\#}).$ 

A non-trivial theorem of Eda-Kawamura (see [3]) establishes some cases using higher connectivity. Since each  $S^{r_i+1}$  is an  $r_i$ -connected CW-complex, we have  $\pi_n(X) = \check{\pi}_n(X) = 0$  for  $1 \leq n \leq r_1$ . When  $n = r_1 + 1$ , we have that  $\check{\pi}_n(X, x_0) \cong \prod_{i=1}^{\infty} \pi_n(S^{r_i+1})$  and that  $\Psi_n$  is an isomorphism. Hence,

$$\pi_{r_1+1}(X) \cong \check{\pi}_{r_1+1}(X) \cong \prod_{r_j=r_1} \pi_{r_1+1}(S^{r_j+1}) \cong \prod_{r_j=r_1} \mathbb{Z},$$

which is either finitely generated free abelian or the Baer-Specker group. Also it is shown in [7] that  $\check{\pi}_{n+1}(\mathbb{E}_n)$  is isomorphic to  $\mathbb{Z}^{\mathbb{N}}$  when n = 2 and  $\mathbb{Z}_2^{\mathbb{N}}$  when  $n \ge 3$ .

To extend the argument used in [7], we recall the notion of a Hall set, which has import in both group theory and Lie theory. In particular, a Hall set on a non-empty finite set  $A_k = \{a_1, a_2, \ldots, a_k\}$  provides a convenient basis for a free Lie algebra on  $A_k$  [4]. Following Hilton's approach in [6], we consider such a Hall set on  $A_k = \{a_1, a_2, \dots, a_k\}$  where  $a_i = [id_{S^{r_i+1}}]$  is the canonical generator of  $\pi_{r_i+1}(S^{r_i+1}).$ 

**Definition 3.1.** A Hall set on a non-empty finite set  $A_k = \{a_1, a_2, \ldots, a_k\}$  is a linearly ordered set  $H_k$  of nested formal commutators determined by the following inductive collection and ordering process [5]: the set  $H_k(1)$  of Hall words of weight 1 is the set  $A_k$  ordered as  $a_1 < a_2 < \cdots < a_k$ . Suppose the set  $H_k(i)$  of Hall words of weight i has been defined for i < m and that  $\bigcup_{i < m} H_k(i)$  has been linearly ordered and so that if i < j,  $x \in H_k(i)$ , and  $y \in H_k(j)$ , then x < y. With this inductive hypothesis, we define a Hall word of weight m to be a formal commutator [x, y]such that

- (1)  $x \in H_k(i)$  and  $y \in H_k(j)$  where i + j = m,
- (2) x < y in  $\bigcup_{i < m} H_k(i)$ , (3) if y = [a, b] for  $a, b \in \bigcup_{i < m} H_k(i)$ , then  $a \leq x$ .

We let  $H_k(m)$  be the set of all Hall words of weight m with an arbitrary linear ordering. Now  $H_k = \bigcup_{i=1}^{\infty} H_k(i)$  is given a linear ordering that extends the linear ordering of each  $\bigcup_{i < m} H_k(i)$  so that a Hall word of smaller weight is always less than another Hall word of greater weight.

For each  $i \in \{1, 2, ..., k\}$  and Hall word  $w \in H_k$ , we let  $\nu_i(w)$  denote the number of times the letter  $a_i$  appears in w as a word (when the brackets are removed). The weight of a Hall word  $w \in H_k$  is the same as the length  $L(w) = \sum_{i=1}^k \nu_i(w)$  of w as a word.

**Example 3.2.** When k = 2, the first elements of the Hall set  $H_2$  are ordered as follows (where an arbitrary ordering is chosen for words at a fixed weight):  $a_1, a_2$ ,  $[a_1, a_2], [a_1, [a_1, a_2]], [a_2, [a_1, a_2]], [a_1, [a_1, [a_1, a_2]]], [a_2, [a_1, [a_1, a_2]]], [a_2, [a_2, [a_1, a_2]]], [a_2, [a_1, a_2]]], [a_3, [a_3, [a_3, a_3]]], [a_3, [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]], [a_3, [a_3, a_3]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]]], [a_3, [a_3, a_3]]]$ 

The number of Hall words  $w \in H_k$  of weight j is the value of necklace polynomial  $M_k(j) = \frac{1}{j} \sum_{d|j} \mu(d) k^{j/d}$  where  $\mu$  is the Möbius function. Next, we show that since  $A_2 \subseteq A_3 \subseteq A_4 \subseteq \cdots$ , we can choose Hall sets  $H_k$  on  $A_k$  in a coherent way.

**Proposition 3.3.** There exists a sequence of linearly ordered sets  $H_2 \subseteq H_3 \subseteq$  $H_4 \subseteq \cdots$  where  $H_k$  is a Hall set on  $A_k$  and  $H_{k+1} \setminus H_k$  consists of all Hall words involving the letter  $a_{k+1}$ .

*Proof.* Since  $H_k(1) = A_k$ , the given ordering on the sets  $A_k$  ensures that  $H_k(1)$  is an initial suborder of  $H_{k+1}(1)$  and that  $H_{k+1}(1)\setminus H_k(1) = \{a_{k+1}\}$ . Suppose that, for all i < m, we have constructed the sets  $H_k(i), k \in \mathbb{N}$  so that  $H_k(i)$  is an initial suborder of  $H_{k+1}(i)$  and that  $H_{k+1}(i) \setminus H_k(i)$  consists of all elements of  $H_{k+1}(i)$ involving the letter  $a_{k+1}$ . This induction hypothesis and the Hall set construction described above ensure that  $H_2(m) \subseteq H_3(m) \subseteq H_4(m) \subseteq \cdots$  as subsets. It follows that  $H_{k+1}(m) \setminus H_k(m)$  is the set of elements of  $H_{k+1}(m)$  that involve the letter  $a_{k+1}$ . We choose the ordering of  $H_2(m)$  arbitrarily and recursively choose the ordering on  $H_{k+1}(m)$ ,  $k \ge 2$  so that  $H_k(m)$  is an initial suborder of  $H_{k+1}(m)$ . This completes the induction and we obtain Hall sets  $H_k$  on  $A_k$  such that for all  $k, i \in \mathbb{N}$ ,  $H_k(i) \subseteq H_{k+1}(i)$  where  $H_{k+1}(i) \setminus H_k(i)$  is the set of Hall words in  $H_{k+1}$  of weight *i* that involve the letter  $a_{k+1}$ . Thus  $H_k \subseteq H_{k+1}$  for all  $k \in \mathbb{N}$  and if  $w \in H_{k+1} \setminus H_k$ has weight i, then  $w \in H_{k+1}(i) \setminus H_k(i)$ . 

Since we have chosen  $a_i$  to be an element of  $\pi_{r_i+1}(S^{r_i+1})$ , a Hall word  $w \in H_k$  may be identified with the corresponding iterated Whitehead product, which lies in some homotopy group of  $X_{\leq k}$ . To identify the dimension of this homotopy group, we borrow the following definition from [6].

**Definition 3.4.** The *height* of a Hall word  $w \in H_k$  is  $h(w) = \sum_{i=1}^k r_i \nu_i(w)$ .

**Remark 3.5.** For a Hall word  $w \in H_k$ , the height h(w) is one less than the dimension of the homotopy group in which w is an element, that is  $w \in \pi_{h(w)+1}(X_{\leq k})$ . For example when k = 3, we have  $a_1 \in \pi_{r_1+1}(S^{r_1+1})$ ,  $a_1 \in \pi_{r_2+1}(S^{r_2+1})$ , and  $a_3 \in \pi_{r_3+1}(S^{r_3+1})$ . Then

- (1)  $w = [a_1, a_2] \in \pi_{r_1+r_2+1}(X_3)$  has height  $h(w) = r_1 + r_2$ ,
- (2)  $w = [a_1, [a_1, a_2]] \in \pi_{2r_1+r_2+1}(X_3)$  has height  $h(w) = 2r_1 + r_2$ ,
- (3)  $w = [a_1, [a_2, a_3]] \in \pi_{r_1+r_2+r_3+1}(X_3)$  has height  $h(w) = r_1 + r_2 + r_3$ .

Because a Hall word is given by nesting Whitehead products in a binary fashion, Relation (1) of Remark 2.1 makes the following proposition evident.

**Proposition 3.6.** Let  $k \ge 1$  and  $w \in H_k$  be a Hall word regarded as a nested Whitehead product in  $\pi_{h(w)+1}(X_{\le k})$ . If we replace any letter of w with 0, then the resulting element of  $\pi_{h(w)+1}(X_{\le k})$  is trivial.

Using Proposition 3.3, we fix Hall sets  $H_k$  on  $A_k$  so that  $H_1 \subseteq H_2 \subseteq H_3 \subseteq \cdots$ . The following is Hilton's initial version of the well-known Hilton-Milnor Theorem. We choose to use Hilton's original formulation in terms of homotopy groups of spheres and Whitehead products since our focus is on shrinking wedges of spheres.

**Theorem 3.7** (Hilton [6]). For given  $k \ge 1$  and  $n > r_1$ , there is an isomorphism  $\phi_{n,k} : \bigoplus_{w \in H_k} \pi_n(S^{h(w)+1}) \to \pi_n(X_{\le k})$  determined by  $\phi_{n,k}(f) = w \circ f$  for each  $f \in \pi_n(S^{h(w)+1})$ .

**Remark 3.8.** Although the direct sum  $\bigoplus_{w \in H_k} \pi_n(S^{h(w)+1})$  in Hilton's Theorem is indexed by the infinite set  $H_k$ , only finitely many summands are non-trivial since n is fixed. Indeed, the set  $H_{n,k} = \{w \in H_k \mid h(w) + 1 \leq n\}$  is finite and the domain of  $\phi_{n,k}$  can be more efficiently expressed as the finite direct sum  $\bigoplus_{w \in H_n,k} \pi_n(S^{h(w)+1})$ .

**Lemma 3.9.** Fix  $k \ge 1$ ,  $n > r_1$ , and  $w \in H_{n,k+1}$ . If  $w \in H_{n,k}$ , then the homomorphism

$$b_{k+1,k\#} \circ \phi_{n,k+1} : \bigoplus_{w \in H_{n,k+1}} \pi_n(S^{h(w)+1}) \to \pi_n(X_{\leq k})$$

maps the summand  $\pi_n(S^{h(w)+1})$  isomorphically onto  $\phi_{n,k}(\pi_n(S^{h(w)+1}))$  and if  $w \in H_{n,k+1} \setminus H_{n,k}$ , then  $b_{k+1,k\#} \circ \phi_{n,k+1}$  trivializes  $\pi_n(S^{h(w)+1})$ .

Proof. If  $f \in \pi_n(S^{h(w)+1})$ , then  $b_{k+1,k\#} \circ \phi_{n,k+1}(f) = [b_{k+1,k}] \circ w \circ f$ . Note that  $[b_{k+1,k}] \circ w$  is still a Whitehead product with the same nested structure but where each appearance of the letter  $a_{k+1}$  is trivialized. Hence, if  $a_{k+1}$  appears in w, then  $[b_{k+1,k}] \circ w = 0$  by Proposition 3.6. It follows that if  $w \in H_{n,k+1} \setminus H_{n,k}$  then  $[b_{k+1,k}] \circ w \circ f = 0$ . On the other hand, if  $w \in H_{n,k}$ , then  $a_{k+1}$  does not appear as a letter in w, which implies that  $[b_{k+1,k}] \circ w = w$  and thus  $b_{k+1,k\#} \circ \phi_{n,k+1}(f) = w \circ f = \phi_{n,k}(f)$ .

Define  $H_{\infty} = \bigcup_{k=1}^{\infty} H_k$  and  $H_{n,\infty} = \bigcup_{k=1}^{\infty} H_{n,k} = \{w \in H_{\infty} \mid h(w) + 1 \leq n\}$  and note that both of these sets are infinite.

**Theorem 3.10.** If  $n > r_1$ , then there is a canonical isomorphism

$$\phi_{n,\infty}: \prod_{w \in H_{n,\infty}} \pi_n(S^{h(w)+1}) \to \check{\pi}_n(X)$$

given by  $\phi_{n,\infty}((f_w)_{w \in H_{n,\infty}}) = \left(\sum_{w \in H_{n,k}} w \circ f_w\right)_{k \in \mathbb{N}}$ .

Proof. For each  $k \ge 1$ , let  $p_{k+1,k} : \bigoplus_{w \in H_{n,k+1}} \pi_n(S^{h(w)+1}) \to \bigoplus_{w \in H_{n,k}} \pi_n(S^{h(w)+1})$ be the unique homomorphism, which trivializes  $\pi_n(S^{h(w)+1})$  if  $w \in H_{n,k+1} \setminus H_{n,k}$  and is the identity on  $\pi_n(S^{h(w)+1})$  if  $w \in H_{n,k}$ . By Lemma 3.9, the following diagram commutes where the horizontal maps are isomorphisms from Hilton's Theorem.

Because the maps  $p_{k+1,k}$  are projection maps of a finite direct product, the inverse limit of the left vertical morphisms is the direct product  $\prod_{w \in H_{n,\infty}} \pi_n(S^{h(w)+1})$ . Hence, taking the corresponding inverse limit of the isomorphisms  $\phi_{n,k}$  as  $k \to \infty$ yields the desired isomorphism.

**Remark 3.11.** Since  $\pi_n(S^{h(w)+1}) = 0$  whenever  $w \in H_{\infty} \setminus H_{n,\infty}$ , we can also express the isomorphism  $\phi_{n,\infty}$  as a map

$$\phi_{n,\infty}:\prod_{w\in H_\infty}\pi_n(S^{h(w)+1})\to\check{\pi}_n(X)$$

with the same formula as that given in Theorem 3.10.

**Remark 3.12.** When  $w = a_i \in H_{\infty}$  has weight L(w) = 1, we have  $h(w) = r_i$ and thus w corresponds to a factor  $\pi_n(S^{r_i+1})$  in the direct product of Theorem 3.10. Collecting all such factors together into a single summand, we have  $\check{\pi}_n(X) \cong \prod_{L(w)>1} \pi_n(S^{h(w)+1}) \oplus \prod_{i=1}^{\infty} \pi_n(S^{r_i+1})$  where  $\prod_{i=1}^{\infty} \pi_n(S^{r_i+1})$  is the previouslydefined weight-1 summand of  $\check{\pi}_n(X)$ . Hence,

$$\check{\pi}_{n+1}\left(\prod_{i=1}^{k} S^{r_i+1}, \bigvee_{i=1}^{k} S^{r_i+1}\right) \cong \prod_{L(w)>1} \pi_n(S^{h(w)+1}).$$

We are particularly interested in the case where  $r_i = m - 1 \ge 1$  so that  $\bigvee_i S^{r_i+1} = \mathbb{E}_m$ . To analyze this case more carefully, we partition  $H_{\infty}$  as follows: Let  $H_{\infty}(j) = \bigcup_{k=1}^{\infty} H_k(j)$  be the Hall words of weight j, e.g.  $H_{\infty}(1) = \{a_1, a_2, a_3, \cdots\}$  and  $H_{\infty}(2) = \{[a_i, a_j] \mid i < j\}$ . Note that for each  $j \ge 1$ ,  $H_{\infty}(j)$  is countably infinite.

Next, we prove our first main result.

Proof of Theorem 1.1. By Theorem 3.10,  $\check{\pi}_n(\mathbb{E}_m)$  is isomorphic to the direct product  $\prod_{w \in H_{\infty}} \pi_n(S^{h(w)+1})$ . We prove the current theorem by rearranging the terms in this product. In this case, we have  $r_i = m - 1$  for all *i*. Thus each Hall word  $w \in H_k$  has height  $h(w) = \sum_{i=1}^k (m-1)\nu_i(k) = (m-1)L(w)$  where L(w) is the

weight/length of w. Recall that we have defined  $H_{\infty}(j) = \{w \in H_{\infty} \mid L(w) = j\}$  to be the Hall words of weight j. Thus

$$\check{\pi}_n(\mathbb{E}_m) \cong \prod_{w \in H_\infty} \pi_n(S^{(m-1)L(w)+1}) \cong \prod_{j=1}^\infty \prod_{w \in H_\infty(j)} \pi_n(S^{(m-1)j+1}).$$

As  $j \to \infty$ , the groups  $\pi_n(S^{(m-1)j+1})$  have the potential to be non-trivial only when  $n \ge (m-1)j+1$  or, equivalently, when  $j \le \frac{n-1}{m-1}$ . Hence,  $\prod_{w \in H_\infty(j)} \pi_n(S^{(m-1)j+1}) = 0$  when  $j > \frac{n-1}{m-1}$ . We have:

$$\prod_{j=1}^{\infty} \prod_{w \in H_{\infty}(j)} \pi_n(S^{(m-1)j+1}) \cong \bigoplus_{1 \leq j \leq \frac{n-1}{m-1}} \prod_{w \in H_{\infty}(j)} \pi_n(S^{(m-1)j+1})$$
$$\cong \bigoplus_{1 \leq j \leq \frac{n-1}{m-1}} \left(\pi_n(S^{(m-1)j+1})\right)^{\mathbb{N}}$$

where the last isomorphism holds since m is fixed and  $H_{\infty}(j)$  is countably infinite.

We will often express  $\check{\pi}_n(\mathbb{E}_m)$  as the direct sum  $\bigoplus_{1 \leq j \leq \frac{n-1}{m-1}} \mathcal{W}_j$  where  $\mathcal{W}_j$  is the image of  $\prod_{w \in H_{\infty}(j)} \pi_n(S^{(m-1)j+1})$  under the isomorphism  $\phi_{n,\infty}$ . We refer to  $\mathcal{W}_j$  as the weight-j summand of  $\check{\pi}_n(\mathbb{E}_m)$ . Since  $\mathcal{W}_1 = \prod_{w \in H_{\infty}(1)} \pi_n(S^m)$  corresponds precisely to the "weight-1 summand" of  $\check{\pi}_n(\mathbb{E}_m)$  as defined in Section 2.3, we have the following.

Corollary 3.13. If  $n > m \ge 2$ , then

$$\check{\pi}_{n+1}\left(\prod_{j\in\mathbb{N}}S^m,\mathbb{E}_m\right)\cong\bigoplus_{2\leqslant j\leqslant\frac{n-1}{m-1}}\left(\pi_n(S^{mj-j+1})\right)^{\mathbb{N}}.$$

**Example 3.14.** Some special cases worth noting are the following.

(1) When n = 3 and m = 2, we have a summand for both  $j = 1, 2 \leq \frac{3-1}{2-1}$ . In this case,

$$\check{\pi}_3(\mathbb{E}_2) \cong (\pi_3(S^2))^{\mathbb{N}} \oplus (\pi_3(S^3))^{\mathbb{N}} \cong \mathbb{Z}^{\mathbb{N}} \oplus \mathbb{Z}^{\mathbb{N}}$$

Although this is abstractly isomorphic to  $\mathbb{Z}^{\mathbb{N}}$ , this decomposition shows that the first summand of  $\mathbb{Z}^{\mathbb{N}} \oplus \mathbb{Z}^{\mathbb{N}}$  is the weight-1 summand of  $\check{\pi}_3(\mathbb{E}_2)$ .

(2) For the groups  $\check{\pi}_{n+1}(\mathbb{E}_n)$ ,  $n \ge 3$ , we only have a summand for  $j = 1 \le \frac{n}{n-1}$ . In this case, we enter the stable range and have

$$\check{\pi}_{n+1}(\mathbb{E}_n) \cong (\pi_{n+1}(S^n))^{\mathbb{N}} \cong \mathbb{Z}_2^{\mathbb{N}},$$

the entirety of which is the weight-1 summand. This agrees with the computation in [7].

(3) For n = 4 and m = 2, we have a summand for  $j = 1, 2, 3 \leq \frac{4-1}{2-1}$ . In this case, we have

$$\check{\pi}_4(\mathbb{E}_2) \cong (\pi_4(S^2))^{\mathbb{N}} \oplus (\pi_4(S^3))^{\mathbb{N}} \oplus (\pi_4(S^4))^{\mathbb{N}} \cong \mathbb{Z}_2^{\mathbb{N}} \oplus \mathbb{Z}_2^{\mathbb{N}} \oplus \mathbb{Z}^{\mathbb{N}}$$

where  $\mathcal{W}_1 \cong (\pi_4(S^2))^{\mathbb{N}} \cong \mathbb{Z}_2^{\mathbb{N}}$  is the weight-1 summand. More generally, when  $n \ge 4$ , we have

$$\check{\pi}_n(\mathbb{E}_2) \cong \pi_n(S^2)^{\mathbb{N}} \oplus \pi_n(S^3)^{\mathbb{N}} \oplus \dots \oplus \pi_n(S^n)^{\mathbb{N}}$$

where  $\mathcal{W}_{i} = \pi_{n}(S^{j+1})^{\mathbb{N}}$  corresponds to the weight-*j* summand.

**Corollary 3.15.** For fixed  $n \ge 0$ , the Čech homotopy groups  $\check{\pi}_{m+n}(\mathbb{E}_m)$  stabilize when  $m \ge n+2$ .

*Proof.* When  $m \ge n+2$ , we have  $\frac{m+n-1}{m-1} < 2$  and thus Theorem 1.1 only gives a single summand corresponding to j = 1. In particular,  $\check{\pi}_{m+n}(\mathbb{E}_m) \cong (\pi_{m+n}(S^m))^{\mathbb{N}}$ . Since the groups  $\pi_{m+n}(S^m)$  stabilize when  $m \ge n+2$ , the result follows.  $\Box$ 

4. More on the image of  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$ 

To identify new elements of the image of the canonical homomorphism  $\Psi_n$ :  $\pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$ , we use the isomorphism  $\phi_{n,\infty} : \bigoplus_{1 \leq j \leq \frac{n-1}{m-1}} (\pi_n(S^{mj-j+1}))^{\mathbb{N}} \to \check{\pi}_n(\mathbb{E}_m)$  as a book-keeping device.

**Remark 4.1.** The set  $H_k(j)$  of all Hall words of weight j and involving letters  $a_1, \ldots, a_k$  provides a convenient indexing set for the elements of the weight-j summand  $\mathcal{W}_j = \phi_{n,\infty}((\pi_n(S^{mj-j+1}))^{\mathbb{N}})$ . According to the formula for  $\phi_{n,\infty}$  in Theorem 3.10, a generic element of  $\mathcal{W}_j$  has the form  $(\sum_{w \in H_k(j)} w \circ f_w)_{k \in \mathbb{N}}$  where  $f_w \in \pi_n(S^{mj-j+1})$  and  $f_w$  may only be non-zero for finitely many  $w \in H_{n,k} \cap H_k(j)$ .

**Proposition 4.2.** For any  $m, n \ge 2$ ,  $W_1$  is a direct summand of  $Im(\Psi_n)$ .

*Proof.* Recall from the end of Section 2, that we have  $\check{\sigma} \circ \Psi_n = \sigma_{\#}$  where  $\operatorname{Im}(\sigma_{\#}) = \mathcal{W}_1$  and  $\sigma_{\#}$  splits. The claim follows.

**Corollary 4.3.** If n < 2m-1, then  $\Psi_n : \pi_n(\mathbb{E}_m) \to \check{\pi}_n(\mathbb{E}_m)$  is a split epimorphism. *Proof.* Since  $\frac{n-1}{m-1} < 2$ , Theorem 1.1 implies that  $\check{\pi}_n(\mathbb{E}_m) = \mathcal{W}_1$ . The splitting of  $\Psi_n$  follows from Proposition 4.2.

Next we consider the special case n = 2m - 1 at the edge of the unstable range. In this situation, we will need to apply infinite sums and Whitehead products described in Section 2.

**Lemma 4.4.** For  $m \ge 2$ , the canonical homomorphism  $\Psi_{2m-1} : \pi_{2m-1}(\mathbb{E}_m) \to \pi_{2m-1}(\mathbb{E}_m)$  is a split epimorphism.

Proof. First, we apply Theorem 1.1 in the case n = 2m - 1. Since  $\frac{2m-2}{m-1} < 3$ , we have  $\check{\pi}_{2m-1}(\mathbb{E}_m) = \mathcal{W}_1 \oplus \mathcal{W}_2$ . A splitting homomorphism  $\mathcal{W}_1 \to \pi_n(\mathbb{E}_m)$  of  $\Psi_n$  is established in Proposition 4.2 and so it suffices to produce a splitting homomorphism  $\mathcal{W}_2 \to \pi_n(\mathbb{E}_m)$ . Recall from Remark 4.1, that a generic element of  $\mathcal{W}_2$  has the form  $\alpha = \left(\sum_{w \in H_k(2)} w \circ f_w\right)_{k \in \mathbb{N}}$  where  $f_w \in \pi_{2m-1}(S^{2m-1})$ . Since  $H_k(2) = \{[a_i, a_j] \mid 1 \leq i < j \leq k\}$ , write  $f_{i,j}$  for  $f_w$  when  $w = [a_i, a_j]$ . Let  $\gamma = [id_{S^{2m-1}}]$  be the generator of  $\pi_{2m-1}(S^{2m-1})$  and write  $f_{i,j} = \epsilon_{i,j}\gamma$  for integer  $\epsilon_{i,j} \in \mathbb{Z}$ . Since  $w \circ -$  is a homomorphism, we have  $w \circ f_{i,j} = [a_i, a_j] \circ \epsilon_{i,j}\gamma = \epsilon_{i,j}([a_i, a_j] \circ \gamma) = \epsilon_{i,j}[a_i, a_j] = [a_i, \epsilon_{i,j}a_j]$ . Altogether, we can express  $\alpha$  more conveniently as

$$\alpha = \left(\sum_{i=1}^{k-1} \sum_{j=i+1}^{k} [a_i, \epsilon_{i,j} a_j]\right)_{k \in \mathbb{N}}$$

where the sums in the sequence are directly extended as k increases.

Let  $\ell_j : S^m \to \mathbb{E}_m$  be the inclusion of the *j*-th sphere so that  $a_j = [\ell_j]$  and define an infinite sum map  $F_\alpha : S^{2m-1} \to \mathbb{E}_m$  by

$$F_{\alpha} = \sum_{i=1}^{\infty} \left[ \ell_i, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_j \right]$$

This sum is continuous since the sequence  $\left\{ \left[ \ell_i, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_j \right] \right\}_{i \in \mathbb{N}}$  converges to the wedgepoint of  $\mathbb{E}_m$ . Fix  $k \in \mathbb{N}$  and let  $b_k : \mathbb{E}_m \to X_{\leq k}$  denote the projection map to the union of the first k-spheres. For given  $i \in \mathbb{N}$ , note that  $b_k \circ \left[ \ell_i, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_j \right]$  is constant if i > k. If  $i \leq k$ , then

$$b_k \circ \left[ \ell_i, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_j \right] = \left[ b_k \circ \ell_i, b_k \circ \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_j \right]$$
$$= \left[ b_k \circ \ell_i, \sum_{j=i+1}^{\infty} b_k \circ \epsilon_{i,j} \ell_j \right]$$

and we consider two cases. If i = k, this is equal to  $[\ell_i, c]$  where c is constant. Such a map is null-homotopic within the *i*-th wedge-summand of  $\mathbb{E}_m$ . Lastly, if i < k, this is homotopic to  $\left[\ell_i, \sum_{j=i+1}^k \epsilon_{i,j} \ell_j\right]$ . All three of these cases are realized by homotopies having image in  $X_{\geq i}$ . Thus, Proposition 2.7 gives

$$b_k \circ F_\alpha \simeq \sum_{i=1}^{k-1} \left[ \ell_i, \sum_{j=i+1}^k \epsilon_{i,j} \ell_j \right].$$

On homotopy classes, we have

$$b_{k\#}([F_{\alpha}]) = \sum_{i=1}^{k-1} \left[ a_i, \sum_{j=i+1}^k \epsilon_{i,j} a_j \right] = \sum_{i=1}^{k-1} \sum_{j=i+1}^k \left[ a_i, \epsilon_{i,j} a_j \right]$$

where the second equality follows from bilinearity of Whitehead products. Since k was arbitrary, we conclude that  $\Psi_n([F_\alpha]) = (b_{k\#}([F_\alpha]))_{k \in \mathbb{N}} = \alpha$ .

For the splitting, it suffices to check that the mapping  $\alpha \mapsto [F_{\alpha}]$  defines a homomorphism  $\mathcal{W}_2 \to \pi_{2m-1}(\mathbb{E}_m)$ . Since our representation of  $\alpha$  is uniquely determined by the integers  $\epsilon_{i,j}, j > i$ , this function is well-defined. We show that  $[F_{\alpha+\beta}] = [F_{\alpha}] + [F_{\beta}]$ . Express  $\beta \in \mathcal{W}_2$  as  $\beta = \left(\sum_{i=1}^{k-1} \sum_{j=i+1}^k [a_i, \delta_{i,j}a_j]\right)_{k\in\mathbb{N}}$  for integers  $\delta_{i,j}, j > i$  and note that  $\alpha + \beta = \left(\sum_{i=1}^{k-1} \sum_{j=i+1}^k [a_i, (\epsilon_{i,j} + \delta_{i,j})a_j]\right)_{k\in\mathbb{N}}$ . Recalling the formula for  $F_{\alpha}$ ,  $F_{\beta}$ , and  $F_{\alpha+\beta}$ , and applying the infinite-sum homotopies from Section 2, we have the following sequence:

$$F_{\alpha} + F_{\beta} = \sum_{i=1}^{\infty} \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_{j} \right] + \sum_{i=1}^{\infty} \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \delta_{i,j} \ell_{j} \right]$$

$$\simeq \sum_{i=1}^{\infty} \left( \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_{j} \right] + \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \delta_{i,j} \ell_{j} \right] \right)$$

$$\simeq \sum_{i=1}^{\infty} \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_{j} + \sum_{j=i+1}^{\infty} \delta_{i,j} \ell_{j} \right]$$

$$\simeq \sum_{i=1}^{\infty} \left[ \ell_{i}, \sum_{j=i+1}^{\infty} \epsilon_{i,j} \ell_{j} + \delta_{i,j} \ell_{j} \right]$$

$$\simeq \sum_{i=1}^{\infty} \left[ \ell_{i}, \sum_{j=i+1}^{\infty} (\epsilon_{i,j} + \delta_{i,j}) \ell_{j} \right].$$

$$= F_{\alpha+\beta}$$

In the unstable range n > 2m - 1, the same argument for surjectivity does not work and it does not appear that Hilton-Milnor invariants overcome the obstacle. Fix  $0 \neq \beta \in \pi_n(S^{2m-1})$ . Note that the example element

$$\left(\left[a_1,\sum_{i=2}^k a_i\right]\circ\beta\right)_{k\in\mathbb{N}} = \left(\left(\sum_{i=2}^k [a_1,a_i]\right)\circ\beta\right)_{k\in\mathbb{N}}$$

of  $\mathcal{W}_2$  is the image of  $\left[\ell_1, \sum_{j=2}^{\infty} \ell_j\right] \circ \beta \in \pi_n(\mathbb{E}_m)$  under  $\Psi_n$ . In contrast, it is not clear if  $\left(\sum_{i=2}^k ([a_1, a_i] \circ \beta)\right)_{k \in \mathbb{N}} \in \mathcal{W}_2$  lies in the image of  $\Psi_n$ . Indeed, one would like to form an infinite sum  $\sum_{j=2}^{\infty} [\ell_1, \ell_j]$  and apply Hilton-Milnor invariants of  $\beta$  in the finite-approximation levels but this infinite sum map is not continuous. The lack of obvious continuous representatives may be more analogous to the situation in dimension n = m = 1 where the homomorphism  $\Psi_1 : \pi_1(\mathbb{E}_1) \to \check{\pi}_1(\mathbb{E}_1)$  fails to be surjective. Here, the analogous sequence of products of commutators

$$(1, [a_1, a_2], [a_1, a_2][a_1, a_3], [a_1, a_2][a_1, a_3][a_1, a_4], \dots) \in \check{\pi}_1(\mathbb{E}_1)$$

has no continuous representative in  $\pi_1(\mathbb{E}_1)$ . That surjectivity holds in the critical case n = 2m - 1 is then quite striking since it is the only situation where one can easily represent such growing sequences of commutators/Whitehead products. While this leaves us with an incomplete picture of the image of  $\Psi_n$  (for n > 2m - 1), we identify a natural candidate.

For  $i \in \mathbb{N}$ , let  $M_i$  be the collection of Hall words  $w \in H_{\infty}$  such that

(1)  $L(w) \ge 2$ ,

(2) i is the smallest index for which  $a_i$  appears as a letter in w.

Let  $\mathcal{G}_n(m) = \prod_{i \in \mathbb{N}} \bigoplus_{w \in M_i} \pi_n(S^{(m-1)L(w)+1})$  and note that  $\mathcal{G}_n(m)$  is naturally a subgroup of  $\prod_{w \in H_\infty} \pi_n(S^{(m-1)L(w)+1})$ . We represent an element of  $\mathcal{G}_n(m)$  as a tuple  $(\alpha_{i,w})_{i,w}$  for  $\alpha_{i,w} \in \pi_n(S^{(m-1)L(w)+1})$  indexed over pairs (i,w) with  $i \in \mathbb{N}$  and  $w \in M_i$ .

**Proposition 4.5.** There is a canonical monomorphism  $\Theta : \mathcal{G}_n(m) \to \pi_n(\mathbb{E}_m)$ . Such that  $\Psi_n \circ \Theta(\mathcal{G}_n(m)) = \phi_{n,\infty}(\mathcal{G}_n(m))$ .

Proof. Define  $\Theta : \mathcal{G}_n(m) \to \pi_n(\mathbb{E}_m)$  as follows. For Hall word w, let  $\ell_w : S^{(m-1)L(w)+1} \to \mathbb{E}_m$  be a the canonical map representing w, e.g. given by replacing each appearance of  $a_i$  with  $\ell_i$ . Given  $\alpha = (\alpha_{i,w})_{i,w} \in \mathcal{G}_n(m)$ , the set  $\operatorname{supp}_i(\alpha) = \{w \in M_i \mid f_{i,w} \neq 0\}$  is finite for each i (by definition of  $\mathcal{G}_n(m)$ ). If  $w \in \operatorname{supp}_i(\alpha)$ , let  $f_{i,w} : S^n \to S^{(m-1)L(w)+1}$  be a map representing  $\alpha_{i,w}$ . Define

$$\Theta(\alpha) = \left\lfloor \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_i(\alpha)} \ell_w \circ f_{i,w} \right\rfloor.$$

To see that  $\Theta$  is well-defined, let  $f'_{i,w}$  be another family of representatives for  $\alpha_{i,w}$ . For given pair (i,w), we have  $f_{i,w} \simeq f'_{i,w}$  and thus  $\sum_{w \in \text{supp}_i(\alpha)} \ell_w \circ f_{i,w} \simeq \sum_{w \in \text{supp}_i(\alpha)} \ell_w \circ f_{i,w}$  in  $X_{\ge i}$ . This forms a sequential homotopy indexed over i and thus  $\sum_{i=1}^{\infty} \sum_{w \in \text{supp}_i(\alpha)} \ell_w \circ f_{i,w} \simeq \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_i(\alpha)} \ell_w \circ f_{i,w}$  by Proposition 2.7. To check that  $\Theta$  is a homomorphism, let  $\beta = (\beta_{i,w})_{i,w} \in \mathcal{G}_n(m)$  with represent-

To check that  $\Theta$  is a homomorphism, let  $\beta = (\beta_{i,w})_{i,w} \in \mathcal{G}_n(m)$  with representing maps  $[g_{i,w}] = \beta_{i,w}$ . Note that  $\operatorname{supp}_i(\alpha + \beta) \subseteq \operatorname{supp}_i(\alpha) \cup \operatorname{supp}_i(\beta)$  where  $F_i = (\operatorname{supp}_i(\alpha) \cup \operatorname{supp}_i(\beta)) \setminus \operatorname{supp}_i(\alpha + \beta)$  is the finite (possibly empty) set of  $w \in M_i$  such that  $\alpha_{i,w}$  and  $\beta_{i,w}$  are non-trivial and  $\alpha_{i,w} = -\beta_{i,w}$ . Note that  $\sum_{i=1}^{\infty} \sum_{w \in F_i} \ell_w \circ (f_{i,w} + g_{i,w})$  is null-homotopic. Thus applying linearity of infinite sums to  $A = [\sum_{i=1}^{\infty} \sum_{w \in F_i} (\ell_w \circ f_{i,w}) + \sum_{i=1}^{\infty} \sum_{w \in F_i} (\ell_w \circ g_{i,w})]$ , we have A = 0. Then applications of Propositions 2.5 and 2.7 give that  $\Theta(\alpha + \beta)$  is equal to:

$$\begin{bmatrix} \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_{i}(\alpha+\beta)} \ell_{w} \circ (f_{i,w} + g_{i,w}) \end{bmatrix}$$

$$= \begin{bmatrix} \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_{i}(\alpha+\beta)} \ell_{w} \circ f_{i,w} + \sum_{w \in \text{supp}_{i}(\alpha+\beta)} \ell_{w} \circ g_{i,w} \end{bmatrix} + A$$

$$= \begin{bmatrix} \sum_{i=1}^{\infty} \left( \sum_{w \in \text{supp}_{i}(\alpha+\beta) \cup F_{i}} \ell_{w} \circ f_{i,w} + \sum_{w \in \text{supp}_{i}(\alpha+\beta) \cup F_{i}} \ell_{w} \circ g_{i,w} \right) \end{bmatrix}$$

$$= \begin{bmatrix} \sum_{i=1}^{\infty} \left( \sum_{w \in \text{supp}_{i}(\alpha)} \ell_{w} \circ f_{i,w} + \sum_{w \in \text{supp}_{i}(\beta)} \ell_{w} \circ g_{i,w} \right) \end{bmatrix}$$

$$= \begin{bmatrix} \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_{i}(\alpha)} \ell_{w} \circ f_{i,w} \end{bmatrix} + \begin{bmatrix} \sum_{i=1}^{\infty} \sum_{w \in \text{supp}_{i}(\beta)} \ell_{w} \circ g_{i,w} \end{bmatrix}$$

$$= \Theta(\alpha) + \Theta(\beta)$$

Proof of Theorem 1.3. The first summand  $\prod_{\mathbb{N}} \pi_n(S^m)$  comes from the splitting of the weight-1 summand  $\mathcal{W}_1 \to \pi_n(\mathbb{E}_m)$  in Theorem 2.12. The second summand we construct as follows. Proposition 4.5 and the bijectivity of  $\phi_{n,\infty}$  imply that  $H = \Theta(\mathcal{G}_n(m))$  is a subgroup of  $\pi_n(\mathbb{E}_m)$  for which the restriction  $\Psi_n|_H : H \to \mathcal{G}_n(m)$  is an isomorphism. To show that H splits in a summand distinct from the weight-1 summand, we must show that  $H \leq \ker(\sigma_{\#}) \leq \pi_n(\mathbb{E}_m)$  where  $\sigma : X \to \prod_{i \in \mathbb{N}} S^m$  is the inclusion. Let  $\alpha = (\alpha_{i,w}) \in \mathcal{G}_n(m)$  with  $\alpha_{i,w} = [f_{i,w}]$ . If  $p_j : \mathbb{E}_m \to X_j$  is the retraction to the *j*-th sphere, then  $p_j \circ \ell_w \circ f_{i,w}$  is null-homotopic for all  $i, j \in \mathbb{N}$  and  $w \in M_i$  (since  $\ell_w$  is an iterated Whitehead product map determined by a Hall word w, which involves at least two distinct letters). Thus  $[\ell_w \circ f_{i,w}] \in \ker(\sigma_{\#})$  for all pairs (i,w) with  $i \in \mathbb{N}$ ,  $w \in M_i$ . It is shown in [2] that  $\ker(\sigma_{\#})$  is "closed under infinite sums" and it follows directly that  $\Theta(\alpha) = \left[\sum_{i=1}^{\infty} \sum_{w \in \operatorname{supp}_i(\alpha)} \ell_w \circ f_{i,w}\right] \in \ker(\sigma_{\#})$ . Hence,  $H \leq \ker(\sigma_{\#})$ .

Finally, it suffices to find an expression for  $\mathcal{G}_n(m)$  that matches the expression in the statement of the theorem. For all  $j \ge 2$ , let  $M_i(j) = \{w \in M_i \mid L(w) = j\}$  be the collection of Hall words of weight j with i being the smallest index for which  $a_i$  appears in w and note that  $M_i(j)$  is countably infinite. Since  $\{M_i(j)\}_{j\ge 2}$  is a partition of  $M_i$  and  $\pi_n(S^{(m-1)j+1})$  can only be non-trivial if  $2 \le j \le \frac{n-1}{m-1}$ , we have

$$\begin{aligned} \mathcal{G}_n(m) &= \prod_{i \in \mathbb{N}} \bigoplus_{w \in M_i} \pi_n(S^{(m-1)L(w)+1}) \\ &\cong \prod_{i \in \mathbb{N}} \left( \bigoplus_{2 \leq j \leq \frac{n-1}{m-1}} \bigoplus_{w \in M_i(j)} \pi_n(S^{(m-1)j+1}) \right) \\ &\cong \bigoplus_{2 \leq j \leq \frac{n-1}{m-1}} \left( \prod_{i \in \mathbb{N}} \bigoplus_{w \in M_i(j)} \pi_n(S^{(m-1)j+1}) \right) \end{aligned}$$

where the last isomorphism is justified by the fact that finite direct sums commute with arbitrary direct products. By enumerating  $M_i(j)$  and replacing this indexing set with  $\mathbb{N}$ , for each *i*, we match the form appearing in the statement of the theorem.

### References

- M.G. Barratt, J. Milnor, An Example of Anomalous Singular Homology, Proc. Amer. Math. Soc. 13 (1962), no. 2, 293-297.
- J. Brazas, Identities for Whitehead products and infinite sums, To Appear in Topology. Appl. 2025. arXiv:2408.10430.
- K. Eda, K. Kawamura, Homotopy and homology groups of the n-dimensional Hawaiian earring, Fundamenta Mathematicae 165 (2000) 17–28.
- [4] M. Hall, A basis for free Lie rings and higher commutators in free groups, Proc. Amer. Math. Soc. 1 (1950), no. 5, 575-581.
- [5] P. Hall, A contribution to the theory of groups of prime-power order, Proc. London Math. Soc. 36 (1934) 29-95.
- [6] P.J. Hilton, On the Homotopy Groups of the Union of Spheres, J. London Math. Soc., Volume s1-30 (1955), no. 2, 154-172.
- [7] K. Kawamura, Low dimensional homotopy groups of suspensions of the Hawaiian earring, Colloq. Math. 96 (2003), no. 1, 27-39.
- [8] S. Mardešić, J. Segal, Shape theory, North-Holland Publishing Company, 1982.
- [9] Serre, Jean-Pierre, Homologie singuliére des espaces fibrés. Applications. Annals of Mathematics, Second Series 54 (1951), no. 3, 425-505.
- [10] J.H.C. Whitehead, On adding relations to homotopy groups, Annals of Mathematics 42 (1941), no. 2, 409-428.
- [11] G.W. Whitehead, On products in homotopy groups, Annals of Mathematics 47 (1946), no. 3, 460-475.
- [12] G.W. Whitehead, *Elements of Homotopy Theory*, Springer Verlag, Graduate Texts in Mathematics. 1978.

West Chester University, Department of Mathematics, West Chester, PA 19383, USA

 $Email \ address: \ \texttt{jbrazas@wcupa.edu}$ 

16