

# On Virial Expansion in Hard Sphere Model

Kiyoharu Kawana,<sup>\*</sup>

<sup>†</sup> *School of Physics, Korean Institute for Advanced Study, Seoul 02455, Korea*

January 28, 2025

## Abstract

Virial expansion is a traditional approach in statistical mechanics that expresses thermodynamic quantities, such as pressure  $p$ , as power series of density or chemical potential. Its radius of convergence can serve as a potential indicator of phase transition. In this study, we investigate the virial expansion of the hard-sphere model, using the known dimensionless virial coefficients  $\tilde{B}_k$  ( $k = 1, 2, \dots$ ) up to the 12th order. We find that it is well fitted by  $\tilde{B}_k = 1.28 \times k^{1.90}$ , corresponding to the analytic continuation of the virial expansion of the pressure as  $\sim \text{Li}_{-1.90}(\eta)$ , where  $\eta$  is the packing fraction and  $\text{Li}_s(x)$  is the polylogarithm function. This implies the absence of singular behavior in the physical parameter space  $\eta \leq \eta_{\text{max}} \approx 0.74$  and no indication of phase transition in the virial expansion approach. In addition, we calculate the cluster-integral coefficients  $\{b_l\}_{l=1}^{\infty}$  and observe that their asymptotic behavior resembles the results obtained in the large dimension limit ( $D \rightarrow \infty$ ), suggesting that  $D = 3$  might be already regarded as large dimension. However, the existence of phase transition in the hard-sphere model has been confirmed by numerous simulations, which clearly indicates that a naive extrapolation of the virial series can lead to unphysical results.

---

<sup>\*</sup>E-mail: kkiyoharu@kias.re.kr

# 1 Introduction

Statistical mechanics of classical fluid systems has a long and rich history. One of the traditional approaches is the virial expansion [1, 2, 3, 4] which represents thermodynamic quantities as power series of density or chemical potential around the ideal gas. However, the convergence of the virial series is often slow, and it can diverge at a finite density. Such a divergence can be interpreted as a potential indication of a phase transition although its occurrence is not necessarily guaranteed.

One of the simplest yet non-trivial models in classical fluid systems is the hard-sphere model. Despite the absence of attractive forces, various numerical simulations [5, 6, 7, 8, 9] have consistently demonstrated that this system exhibits a phase transition within physical density region  $\eta < \eta_{\max} \approx 0.74$ , where  $\eta$  is the packing fraction. This intriguing result naturally motivates us to investigate the asymptotic behaviour of the virial series and its radius of convergence. Indeed, there have already been a number of studies [10, 11, 12, 13, 14, 15], employing various approximations and resummation techniques.

In this paper, we present an updated analysis of the asymptotic behavior of the virial expansion in the hard-sphere model, using the known virial coefficients up to the 12th order [16]. We find that these virial coefficients are well fitted by Eq. (48) and the estimated radius of convergence is  $\eta_c = 1$ , suggesting no phase transition within the physical density region  $\eta < \eta_{\max}$ . Furthermore, the estimated virial coefficients correspond to the analytic continuation of the pressure as Eq. (49), and  $\eta = \eta_c = 1$  is found to be a branch point of the polylogarithm function. We also calculate the cluster-integral coefficients  $b_l$  and observe that their asymptotic behavior resembles the results obtained in the large dimension limit ( $D \rightarrow \infty$ ), suggesting that  $D = 3$  might be already regarded as large  $D$ . However, we should emphasize that it is still possible for the virial coefficients to become negative for  $k > 12$ , which then can lead to a finite radius of convergence below  $\eta_{\max} \approx 0.74$ .

This paper is organized as follows. In Section 2, we provide a brief review the statistical mechanics of classical fluid system. We emphase that the existence of thermodynamic limit is guaranteed for a short-range potential that satisfies a couple of reasonable conditions. In Section 3, we discuss the applicability of virial expansion and present one qualitative way of estimating the radius of convergence in the large  $l$  limit of the cluster integral  $\{b_l(V)\}_{l=1}^{\infty}$ . In Section 4, we investigate the virial expansion in the hard-sphere model. For comparison, we first review  $D = 1$  and  $\infty$  limits, where the model is exactly solvable. We then examine the  $D = 3$  case, estimating the asymptotic behavior of the virial series using the known virial coefficients up to the 12th order. Conclusion is given in Section 5.

## 2 Classical fluid system and thermodynamic limit

We consider thermodynamic equilibrium of classical particles interacting with a Hamiltonian in the  $D$ -dimensional Euclidean space:

$$H_N = \sum_{i=1}^N \frac{p_i^2}{2m} + U_N(\{x_i\}_{i=1}^N), \quad U_N(\{x_i\}_{i=1}^N) = \sum_{i<j}^N v(x_i, x_j), \quad (1)$$

where  $x_i = (x_i^1, x_i^2, \dots, x_i^D)$ ,  $p_i^2 = \sum_{M=1}^D (p_i^M)^2$ , and  $v(x_i, x_j)$  is a general two-body potential. If we impose the translation invariance, the pair potential becomes a mere function of relative coordinate as  $v(x_1, x_2) = v(x_1 - x_2)$ , and we focus on this case in the following.

The grand-canonical partition function is defined by

$$\begin{aligned} \Xi[T, \mu, V] &= \sum_{N=0}^{\infty} \frac{1}{N!} \left( \prod_{i=1}^N \int_V d^D x_i \int_V d^D p_i \right) e^{-\beta H_N + \beta \mu N} \\ &= \sum_{N=0}^{\infty} \frac{z^N}{N!} \left( \prod_{i=1}^N \int_V d^D x_i \right) e^{-\beta \sum_{i<j} v(x_i - x_j)}, \end{aligned} \quad (2)$$

where  $\beta = 1/T$  is the inverse temperature,  $\mu$  is the chemical potential, and  $z = e^{\beta \mu} (2\pi m T)^{D/2}$  is the fugacity. Here,  $\sum_{i<j}$  means the summation over all different pairs  $(x_i, x_j)$  among  $N$  particles. The integrand in Eq. (2) can be written as

$$\begin{aligned} W_N(\{x_i\}_{i=1}^N) &:= e^{-\beta \sum_{i<j} v(x_i, x_j)} = \prod_{i<j} (1 + f_{ij}) \\ &= 1 + \sum_{i<j} f_{ij} + \sum_{i<j} \sum_{\substack{k<l \\ (k,l) \neq (i,j)}} f_{ij} f_{kl} + \dots, \end{aligned} \quad (3)$$

where  $f_{ij} := e^{-\beta v(x_i - x_j)} - 1$  is the Mayer's  $f$  function. Then, the cluster expansion allows us to perform the summation of  $N$  in Eq. (2) as [1, 2]

$$\log \Xi[T, \mu, V] = V \sum_{l=1}^{\infty} b_l(V) z^l, \quad (4)$$

where

$$b_k(V) := \frac{1}{k! V} \left( \prod_{i=1}^k d^D x_i \right) C_k(\{x_i\}_{i=1}^k) \quad (5)$$

is the cluster integral. Here,  $C_k(\{x_i\}_{i=1}^k)$  denotes the contribution in  $W_k(\{x_i\}_{i=1}^k)$  such that all the  $k$ -particles are connected each other by the Mayer's function, i.e. connected diagrams. In addition, the density is similarly expressed as

$$\frac{\langle N \rangle}{V} = \frac{\partial}{\partial(\beta \mu)} \frac{\log \Xi[T, \mu, V]}{V} = \sum_{l=1}^{\infty} l b_l(V) z^l. \quad (6)$$

The existence of the thermodynamic limit

$$\beta p(T, \mu) := \lim_{V \rightarrow \infty} \frac{\log \Xi[T, \mu, V]}{V} \quad (7)$$

is a nontrivial problem for a generic  $N$ -body potential; however, in classical fluid systems, it is rigorously proven for short-range  $N$ -body potentials that satisfy several reasonable conditions [17]. Since this is a crucial result, we briefly overview it below.

We first define the mutual potential energy by

$$M_{N_1 N_2}(\{x_i\}_{i=1}^{N_1}; \{x'_j\}_{j=1}^{N_2}) := U_{N_1+N_2}(\{x_i\}_{i=1}^{N_1}, \{x'_j\}_{j=1}^{N_2}) - U_{N_1}(\{x_i\}_{i=1}^{N_1}) - U_{N_2}(\{x'_j\}_{j=1}^{N_2}) . \quad (8)$$

Then, we say that  $U_N(\{x_i\}_{i=1}^N)$  is *tempered* if there exist  $\exists \lambda > D$  and  $\exists R_0 > 0$ ,  $\exists A \geq 0$  such that

$$M_{N_1 N_2}(\{x_i\}_{i=1}^{N_1}; \{x'_j\}_{j=1}^{N_2}) \leq AN_1 N_2 e^{-\lambda r} \quad (9)$$

whenever  $|x_i - x'_j| \geq r \geq R_0$  for all  $i = 1, 2, \dots, N_1$  and  $j = 1, 2, \dots, N_2$ . Roughly speaking, this temperedness implies the suppression of the positive part of  $U_N$  for the long-distance  $r \rightarrow \infty$ . For example, in the pair potential case (1), this implies (with  $N_1 = N_2 = 1$ )

$$v(x) \leq A|x|^{-\lambda} \quad \text{for } |x| \geq R_0 , \quad (10)$$

which implies that positive (repulsive) part of  $v(x)$  should damp faster than  $|x|^{-\lambda}$  ( $< |x|^{-D}$ ) for  $|x| \rightarrow \infty$ .

In addition, we say that  $U_N(\{x_i\}_{i=1}^N)$  is *stable* if there exists  $\exists B \geq 0$  such that

$$U_N(\{x_i\}_{i=1}^N) \geq -NB \quad (11)$$

for all  $N = 1, 2, \dots$  and  $\forall (x_1, \dots, x_N) \in (\mathbb{R}^D)^N$ . The importance of this stability is that it implies the convergence of the grand-canonical partition function as

$$\begin{aligned} \Xi[T, \mu, V] &= 1 + \sum_{N=1}^{\infty} \frac{z^N}{N!} \left( \prod_{i=1}^N \int_V d^D x_i \right) e^{-\beta U_N(\{x_i\}_{i=1}^N)} \\ &\leq 1 + \sum_{N=1}^{\infty} \frac{z^N}{N!} \left( \prod_{i=1}^N \int_V d^D x_i \right) e^{-N\beta B} \\ &= \exp(Vze^{\beta B}) \end{aligned} \quad (12)$$

for  $\forall z \geq 0$  and  $\forall \beta \geq 0$ . Here, we list typical pair potentials that satisfy temperedness and stability [17]:

- **Hard sphere potential**

$$v_{\text{HS}}(x) = \begin{cases} \infty & \text{for } |x| < 2a \\ 0 & \text{for } |x| \geq 2a \end{cases} , \quad (13)$$

where  $a > 0$  corresponds to the radius of molecule.

• **Pair potential with hard core**

More generally, we can consider

$$v(x) = v_{\text{HS}}(x) + \delta v(x) , \quad (14)$$

where the positive part of  $\delta v(x)$  decreases faster than  $|x|^{-D}$  for  $|x| \rightarrow \infty$  and satisfies

$$\sum_{i=2}^N \delta v(x_j - x_1) \geq -B \quad (15)$$

for all  $N$  and  $\forall (x_1, x_2, \dots, x_N)$  such that  $|x_i - x_1| \geq 2a$  for  $\forall i \neq 1$ .

• **Lennard-Jones potential**

$$v(x) = v_0 \left[ \left( \frac{\sigma}{r} \right)^p - \left( \frac{\sigma}{r} \right)^q \right] \quad (16)$$

with  $p \geq D$  and  $q \geq D + 1$ .

Now the existence theorem of the thermodynamic limit (7) is given as follows.

*Theorem.* The thermodynamic limit (7) exists for a stable and tempered potential  $U_N(\{x_i\}_{i=1}^N)$ .  $\beta p(\mu, T)$  is a convex function of  $\mu$  and  $T$ . In particular,  $p(\mu, T)$  is a continuous function of  $(\mu, T)$  and an increasing function of  $\mu$ .

In the next section, we will argue that this rigorous result clearly highlights the applicability of cluster/virial expansion exclusively to a single unique phase.

### 3 Cluster/Virial expansion and its applicability

We have seen that the grand-canonical partition function is given by Eq. (4) based on cluster expansion. The existence of thermodynamic limit implies that the pressure and density can be written as

$$\beta p(T, \mu) = \sum_{l=1}^{\infty} b_l(\infty) z^l , \quad n(T, \mu) = \frac{\partial(\beta p)}{\partial \log z} = \sum_{l=1}^{\infty} l b_l(\infty) z^l . \quad (17)$$

However, the convergence of these series is not necessarily guaranteed for  $\forall z \geq 0$ . If the radius of convergence is finite and given by  $|z| = z_c < \infty$ , it may be possible to construct an analytic continuation  $p_{\text{ana}}(z)$  of the virial series, whose domain contains some interval of the positive real axis  $z \geq z_c$ , as illustrated in Fig. 1. Then, there are typically three possibilities:




Figure 1: Virial expansion and its analytic continuation. We may find an analytic continuation  $p_{\text{ana}}(z)$  of virial series, but it cannot represent the true pressure  $p(z)$  beyond  $z = z_c$  when a system exhibits a phase transition at this point.

- (a)  $z = z_c$  is a pole or essential singularity of  $p_{\text{ana}}(z)$ .
- (b)  $z = z_c$  is a regular point of  $p_{\text{ana}}(z)$ .
- (c)  $z = z_c$  is a branch point of  $p_{\text{ana}}(z)$ .

In case (a),  $p_{\text{ana}}(z)$  cannot represent the true pressure for  $z \geq z_c$ , as the singular behavior at  $z = z_c$  is inconsistent with the theorem. Thus, it can only describe the equilibrium of classical fluid for  $z < z_c$ , i.e. gas phase. On the other hand, the case (b) can be consistent with the theorem, and  $p_{\text{ana}}(z)$  can represent the true pressure even for  $z \geq z_c$ . In this case, there is no singularity at  $z = z_c$  and the system stays in an unique phase throughout the domain of  $p_{\text{ana}}(z)$ . The case (c) is subtle and may also be consistent with the theorem if  $\lim_{z \rightarrow z_c} p_{\text{ana}}(z) < \infty$  and  $p_{\text{ana}}(z)$  is real and positive for  $z \geq z_c$  in a certain branch.

In any cases, the cluster/virial expansion (17) can only describe a single phase and cannot account for phase transition [18, 19].

Nevertheless, the estimation of radius of convergence offers a potential indication of phase transition or critical point. One qualitative way to estimate the convergence radius is the steepest decent method in the  $l \rightarrow \infty$  limit, based on the irreducible cluster integrals. In general,  $b_k(V)$  can be expressed as

$$l^2 b_l(V) = \sum_{\{n_k\}} \prod_{k=1}^{l-1} \frac{(l \beta_k(V))^{n_k}}{n_k!}, \quad (18)$$

where  $\sum_{\{n_k\}}$  is the summation over all combinations of  $(l - 1)$  positive integers  $\{n_k\}_{i=1}^{l-1}$  satisfying  $\sum_{k=1}^{l-1} k n_k = l - 1$ , and  $\beta_k(V)$  denotes the irreducible contribution in  $b_k(V)$ . By

the residue theorem, Eq. (18) is also expressed as

$$l^2 b_l(V) = \frac{1}{2\pi i} \oint_C d\xi \frac{e^{l \sum_{k=1}^{\infty} \beta_k(V) \xi^k}}{\xi^l}, \quad (19)$$

where  $C$  is an anticlockwise contour around  $\xi = 0$  in the complex  $\xi$  plane. We denote the integrand as

$$e^{l \sum_{k=1}^{\infty} \beta_k(V) \xi^k} / \xi^l = e^{lF(\xi)}, \quad (20)$$

where

$$F(\xi) = \sum_{k=1}^{\infty} \beta_k(V) \xi^k - \log \xi. \quad (21)$$

One can check that a saddle point exists at the point  $\xi_0 \in \mathbb{R}$  satisfying<sup>1</sup>

$$\left. \frac{dF(\xi)}{d\xi} \right|_{\xi=\xi_0} = 0 \quad \leftrightarrow \quad \sum_{k=1}^{\infty} k \beta_k(V) \xi_0^k = 1, \quad (22)$$

and we can modify the contour  $C$  so that it passes this saddle point. Then,  $F(\xi)$  is expanded around the saddle point as

$$F(\xi) = F(\xi_0) + \frac{F''(\xi_0)}{2} (\xi - \xi_0)^2 + \mathcal{O}((\xi - \xi_0)^3), \quad (23)$$

where

$$F''(\xi_0) = \sum_{k=2}^{\infty} \beta_k(V) k(k-1) \xi_0^{k-2} + \frac{1}{\xi_0^2} = \frac{1}{\xi_0^2} \sum_{k=2}^{\infty} k^2 \beta_k(V) \xi_0^k. \quad (24)$$

which can be positive or negative depending on  $\beta_k(V)$ . When it is negative, this already indicates the instability of system at the density  $n = \xi_0$  because the series  $\sum_{k=1}^{\infty} k^2 \beta_k(V) n^k$  corresponds to the second-order derivative of the grand-canonical potential with respect to the density  $n$ . (See Eq. (29) below.) Thus, it suffices to consider the positive case, and the direction of steepest decent is given by

$$\xi - \xi_0 = ix, \quad x \in \mathbb{R}, \quad (25)$$

and the saddle point approximation of Eq. (19) is

$$l^2 b_l(V) \simeq \frac{1}{\sqrt{2\pi l F''(\xi_0)}} e^{lF(\xi_0)} \quad \therefore \quad b_l(V) \simeq \frac{c}{l^{5/2}} b_0^l, \quad (26)$$

---

<sup>1</sup>This statement is always true when  $\beta_k(V) \geq 0$  for  $\forall k$  while it is not when  $\beta_k(V)$  can take negative values.

where

$$b_0 = e^{F(\xi_0)} = \frac{1}{\xi_0} e^{\sum_{k=1}^{\infty} \beta_k(V) \xi_0^k}, \quad c = \frac{1}{\sqrt{2\pi F''(\xi_0)}}, \quad (27)$$

and  $\xi_0$  is determined by Eq. (22). Now, we can estimate the radius of convergence  $z_c$  of the series (4) as

$$z_c = \left( \lim_{l \rightarrow \infty} b_l(V)^{1/l} \right)^{-1} = b_0^{-1} = \xi_0 e^{-\sum_{k=1}^{\infty} \beta_k(V) \xi_0^k}. \quad (28)$$

Namely, the radius of convergence is determined by the irreducible cluster integrals  $\{\beta_k(V)\}_{k=1}^{\infty}$  in the large  $l$  limit. Note that these irreducible cluster integrals also allow us to express the pressure as a power series of density as [1, 18]

$$\sum_{l=1}^{\infty} b_l(V) z^l = n \left( 1 - \sum_{k=1}^{\infty} \frac{k}{k+1} \beta_k(V) n^k \right), \quad n = z e^{-\sum_{k=1}^{\infty} \beta_k(V) z^k}, \quad (29)$$

which is the popular form of the Virial expansion.

## 4 Hard Sphere Model

We study the asymptotic behavior of the virial expansion of the hard-sphere model (13). Note that, the cluster/virial coefficients do not depend on temperature  $T$  because the Mayer's  $f$  function behaves as

$$f(x) = e^{-\beta v(x)} - 1 = \begin{cases} -1 & |x| < 2a \\ 0 & |x| > 2a \end{cases} \quad (30)$$

in the hard-sphere model.

### 4.1 $D = 1$ and $\infty$

The hard-sphere model is exactly solvable for  $D = 1$  and  $\infty$ .

#### $D = 1$

When  $D = 1$ , the canonical partition function can be evaluated as (with  $x_{N+1} = L$ )

$$\begin{aligned} \frac{Z_N(L)}{N!} &= \left( \prod_{i=1}^N \int_0^{x_{i+1}} dx_i \right) \prod_{i=1}^{N-1} e^{-\beta \sum_{i=1}^{N-1} v(x_{i+1} - x_i)} \\ &= \frac{1}{N!} \left( \int_0^{L-2Na} dx e^{-\beta v(x)} \right)^N \\ &= \frac{1}{N!} (L - 2Na - 2a)^N. \end{aligned} \quad (31)$$



By using the Stirling's formula  $N! \sim \sqrt{2\pi N}(N/e)^N$ , we obtain

$$Z_N(L) \sim \sqrt{2\pi(N-1)} \left( \frac{L}{N} - 2a \right)^N . \quad (32)$$

Then, the pressure is given by

$$\beta pa = \lim_{N \rightarrow \infty} a \frac{\partial}{\partial L} \log Z_N(L) = \frac{\eta}{1-\eta} , \quad \eta = \frac{2Na}{L} . \quad (33)$$

In this case, the pressure diverges at  $\eta = \eta_{\max} = 1$ , and there is no phase transition in the physical density region  $\eta < 1$ .

$D = \infty$

For  $D \rightarrow \infty$ , the evaluation of the cluster integral  $b_l(V)$  is extremely simplified because only the connected tree-diagrams survive [20]. Namely, it is

$$\lim_{D \rightarrow \infty} b_l(V) = \frac{l^{l-2}}{l!} (-v)^{l-1} , \quad (34)$$

where  $l^{l-2}$  is the number of connected tree diagrams with  $l$  distinguished particles (Cayler's theorem), and

$$-v = \lim_{D \rightarrow \infty} \int d^D x f(x) = \lim_{D \rightarrow \infty} V_D(a) , \quad V_D(a) = \frac{\pi^{D/2} a^D}{\Gamma(\frac{D}{2} + 1)} . \quad (35)$$

Here,  $V_D(a)$  is the volume of  $D$ -dimensional ball. We assume that this limit is finite.

Now, the pressure and density are given in cluster expansion as

$$\beta pv = v \sum_{l=1}^{\infty} b_l(V) z^l = \sum_{l=1}^{\infty} (-1)^{l+1} \frac{l^{l-2}}{l!} (vz)^l , \quad (36)$$

$$nv = v \sum_{l=1}^{\infty} l b_l(V) z^l = \sum_{l=1}^{\infty} (-1)^{l+1} \frac{l^{l-1}}{l!} (vz)^l , \quad (37)$$

and both series have the same radius of convergence

$$vz_c = \left( \lim_{l \rightarrow \infty} \left( \frac{l^l}{l!} \right)^{\frac{1}{l}} \right)^{-1} = e^{-1} . \quad (38)$$

In this case, we can find approximate analytic continuations by using the Stirling's formula as

$$\beta pv \approx -\frac{1}{\sqrt{2\pi}} \sum_{l=1}^{\infty} \frac{1}{l^{\frac{5}{2}}} (-evz)^l = -\frac{1}{\sqrt{2\pi}} \text{Li}_{\frac{5}{2}}(-evz) , \quad (39)$$

$$nv \approx -\frac{1}{\sqrt{2\pi}} \sum_{l=1}^{\infty} \frac{1}{l^{\frac{3}{2}}} (-evz)^l = -\frac{1}{\sqrt{2\pi}} \text{Li}_{\frac{3}{2}}(-evz) , \quad (40)$$

where  $\text{Li}_s(x)$  is the polylogarithm function defined by analytic continuation of

$$\text{Li}_s(x) = \sum_{k=1}^{\infty} \frac{x^k}{k^s}, \quad (41)$$

which is holomorphic except for a branch cut  $x \geq 1$  in the real axis. This implies that Eqs. (39)(40) provide approximate pressure and density of the system in the entire physical parameter space  $z \geq 0$  in the large  $D$  limit. Apparently, there is no phase transition.

The equation of state is much simpler compared to the above expansions in terms of  $z$ . In the large  $D$  limit, all the irreducible cluster integrals  $\beta_k(V)$  for  $k \geq 2$  are negligible because they contain loops. Thus, only  $\beta_1(V)$  survives, and we have

$$\beta p v = \eta + \frac{1}{2}\eta^2, \quad \eta = v n, \quad (42)$$

which is nothing but the mean-field result.

## 4.2 $D = 3$

Now let us consider  $D = 3$ . In this case, only the first few virial coefficients are known. The virial expansion in the right-hand side in Eq. (29) is often written as

$$\beta p = \sum_{k=1}^{\infty} B_k n^k, \quad B_k = -\frac{k-1}{k} \beta_{k-1}, \quad B_1 = 1. \quad (43)$$

In the hard-sphere model, it is conventional to express this expansion in terms of dimensionless density as

$$\beta p V_3(a) = \sum_{k=1}^{\infty} \tilde{B}_k(T) \eta^k, \quad (44)$$

where

$$\tilde{B}_k(T) := \frac{B_k(T)}{V_3(a)^{k-1}}, \quad \eta := V_3(a) n, \quad (45)$$

Here,  $\eta$  is known as the packing fraction and its maximum value in  $D = 3$  is  $\eta_{\max} \approx 0.74$ .

These virial coefficients are analytically calculated up to the fourth order [21, 22, 23]:

$$\tilde{B}_2 = 4, \quad \tilde{B}_3 = 10, \quad \tilde{B}_4 = \frac{2707\pi + 438\sqrt{2} - 4131 \arccos(1/3)}{70\pi} = 18.3647684. \quad (46)$$

The higher-order virial coefficients are numerically calculated up to the 12th order [12, 16]:

$$\begin{aligned} \tilde{B}_5 &= 28.2245(3), & \tilde{B}_6 &= 39.81514(93), & \tilde{B}_7 &= 53.3444(37), \\ \tilde{B}_8 &= 68.537(18), & \tilde{B}_9 &= 85.812(85), & \tilde{B}_{10} &= 105.77(39), \\ \tilde{B}_{11} &= 127.9(42), & \tilde{B}_{12} &= 113(29). \end{aligned} \quad (47)$$




Figure 2: Upper Left (Right): Virial coefficients (cluster-integral coefficients) of the hard sphere model in  $D = 3$ .

Lower Left: Pressure as a function of the packing fraction: blue for the virial expansion up to 12th order, orange for the predicted infinite power series, dashed green for its analytical continuation (49), dashed magenta (purple) for the Borel-Padé resummation with [9/3] ([6, 6]).

Lower Right: Plot of the ratio  $\tilde{B}_k/\tilde{B}_{k-1}$ .

One can check that these coefficients are well fitted by the following function:

$$\tilde{B}_k^{\text{fit}} = 1.28 \times k^{1.90} . \quad (48)$$

In the upper left panel in Fig. 2, we plot this function by blue, while the known virial coefficients are shown by red. Equation (48) implies that the radius of convergence of virial expansion (44) is  $\eta_c = 1$ , similar to the  $D = 1$  case. Besides, Eq. (48) also predicts an approximate analytic continuation of the pressure as

$$\beta p V_3(a) \approx 1.28 \times \text{Li}_{-1.90}(\eta) , \quad (49)$$

which has a branch cut in the unphysical density region  $\eta \geq 1$ . In the lower-left panel in Fig. 2, we plot  $\beta p V_3(a)$  for various cases.

By using the relation (18), we can also calculate the cluster integrals  $\tilde{b}_l := b_l/V_3(a)^{l-1}$  and find that it is well fitted by the following alternating series:

$$\tilde{b}_l^{\text{fit}} = (-1)^{l+1} 0.0754 \times (b_0)^l l^{-1.98}, \quad \tilde{b}_0 = 14.8. \quad (50)$$

They are also plotted in the upper-right panel in Fig. 2. It is noteworthy that the  $l$ -dependence is similar to the large  $D$  results (36)(37), suggesting that  $D = 3$  might be already regarded as large  $D$ . In the present case, approximate analytic continuations of pressure and density as functions of  $\tilde{z} := zV_3(a)$  are given by

$$\beta p V_3(a) \approx -0.0754 \times \text{Li}_{1.98}(-\tilde{b}_0 \tilde{z}), \quad (51)$$

$$\eta \approx -0.0754 \times \text{Li}_{0.98}(-\tilde{b}_0 \tilde{z}), \quad (52)$$

which should be compared to the large  $D$  limit (36)(37).

Before concluding, an additional comment is necessary. In the above analysis, it was simply assumed that the fitted virial coefficients (48) are all positive for  $k \geq 13$ . However, as pointed out in the literatures [12, 24], we cannot rule out a possibility that these coefficients may become negative. At present, a slight discrepancy between the known  $\tilde{B}_{12}$  and Eq. (48) might indicate a potential oscillation behavior of the virial coefficients. For illustration, we plot the ratio  $\tilde{B}_k/\tilde{B}_{k-1}$  in the lower-right panel in Fig. 2. More information on the higher-order virial coefficients is essential to improve the analysis.

## 5 Conclusion

We have studied the asymptotic behavior of the virial series of the hard-sphere model. We have found that the known virial coefficients are well fitted by Eq. (48), leading to the approximate analytic continuation (49). These results imply that there is no indications of phase transition all the way up to the close-packing density  $\eta_{\text{max}} \approx 0.74$  in the virial expansion approach. However, this result apparently contradicts the well established fact of the Alder phase transition. Additional information on the higher-order virial coefficients is necessary to improve the analysis.

## Acknowledgements

We would like to thank Takeru Yokota for the fruitful comments. This work is supported by KIAS Individual Grants, Grant No. 090901.

## References

- [1] J. Mayer and M. Mayer, *Statistical mechanics*, A Wiley-Interscience publication, Wiley, 1977.

- [2] R. Feynman, *Statistical mechanics: A set of lectures*, Advanced Books Classics, Avalon Publishing, 1998.
- [3] J. Hansen and I. McDonald, *Theory of simple liquids*, Academic Press, 1990.
- [4] A. J. Masters, *Virial expansions*, Journal of Physics: Condensed Matter **20** (2008), no. 28, 283102.
- [5] B. J. Alder and T. E. Wainwright, *Phase transition for a hard sphere system*, Journal of Chemical Physics (U.S.) **Vol: 27** (1957).
- [6] B. J. Alder and T. E. Wainwright, *Studies in molecular dynamics. i. general method*, Journal of Chemical Physics **31** (1959), 459–466.
- [7] B. J. Alder and T. E. Wainwright, *Phase transition in elastic disks*, Phys. Rev. **127** (1962), 359–361.
- [8] M. Isobe and W. Krauth, *Hard-sphere melting and crystallization with event-chain monte carlo*, The Journal of Chemical Physics **143** (2015), no. 8, 084509.
- [9] E. Schwegler, B. M. Rubenstein, and S. B. Libby, *Advances in the computational sciences*, WORLD SCIENTIFIC, 2017.
- [10] A. Baram and M. Luban, *Divergence of the virial series for hard discs and hard spheres at closest packing*, Journal of Physics C: Solid State Physics **12** (1979), no. 17, L659.
- [11] I. C. Sanchez, *Virial coefficients and close-packing of hard spheres and disks*, The Journal of Chemical Physics **101** (1994), no. 8, 7003–7006, [https://pubs.aip.org/aip/jcp/article-pdf/101/8/7003/19115215/7003\\_1\\_online.pdf](https://pubs.aip.org/aip/jcp/article-pdf/101/8/7003/19115215/7003_1_online.pdf).
- [12] N. Clisby and B. M. McCoy, *Ninth and tenth order virial coefficients for hard spheres in  $d$  dimensions*, Journal of Statistical Physics **122** (2005), no. 1, 15–57.
- [13] J. Tian, H. Jiang, Y. Gui, and A. Mulero, *Equation of state for hard-sphere fluids offering accurate virial coefficients*, Phys. Chem. Chem. Phys. **11** (2009), 11213–11218.
- [14] I. C. Sanchez and J. S. Lee, *On the asymptotic properties of a hard sphere fluid*, The Journal of Physical Chemistry B **113** (2009), no. 47, 15572–15580, <https://doi.org/10.1021/jp901041b>, PMID: 19921954.
- [15] Oncak, Malijevsky, Kolafa, and Labik, *On extrapolation of virial coefficients of hard spheres*, Condensed Matter Physics **15** (2012), no. 2, 23004.
- [16] R. J. Wheatley, *Calculation of high-order virial coefficients with applications to hard and soft spheres*, Phys. Rev. Lett. **110** (2013), 200601.
- [17] D. Ruelle, *Statistical mechanics: Rigorous results*, G - Reference, Information and Interdisciplinary Subjects Series, World Scientific, 1999.

- [18] J. E. Mayer and S. F. Harrison, *Statistical mechanics of condensing systems. iii*, The Journal of Chemical Physics **6** (1938), no. 2, 87–100, [https://pubs.aip.org/aip/jcp/article-pdf/6/2/87/18790092/87\\_1\\_online.pdf](https://pubs.aip.org/aip/jcp/article-pdf/6/2/87/18790092/87_1_online.pdf).
- [19] M. V. Ushcats, *Equation of state beyond the radius of convergence of the virial expansion*, Phys. Rev. Lett. **109** (2012), 040601.
- [20] D. Wyler, N. Rivier, and H. L. Frisch, *Hard-sphere fluid in infinite dimensions*, Phys. Rev. A **36** (1987), 2422–2431.
- [21] L. Boltzmann, Versl. Gewone Vergadering Afk. Natuurk. Nederlandse Akad. Wtensch. (1899), 7:484.
- [22] B. R. A. Nijboer and L. Van Hove, *Radial distribution function of a gas of hard spheres and the superposition approximation*, Phys. Rev. **85** (1952), 777–783.
- [23] I. Lyberg, *The fourth virial coefficient of a fluid of hard spheres in odd dimensions*, Journal of Statistical Physics **119** (2005), no. 3–4, 747–764.
- [24] C. Domb, M. Green, and J. Lebowitz, *Phase transitions and critical phenomena*, Phase Transitions and Critical Phenomena, no. V. 13, Academic Press, 1989.