

# GEOMETRY OF THE UNIT BALL OF $\mathcal{L}(X, Y^*)$

T. S. S. R. K. RAO<sup>1</sup>, SUSMITA SEAL<sup>2</sup>

ABSTRACT. In this paper we study the geometry of the unit ball of the space of operators  $\mathcal{L}(X, Y^*)$ , by considering the projective tensor product  $X \hat{\otimes}_\pi Y$  as a predual. We prove that if an elementary tensor (rank one operator) of the form  $x_0^* \otimes y_0^*$  in the unit sphere  $S_{\mathcal{L}(X, Y^*)}$  is a  $w^*$ -strongly extreme point of unit ball, then  $x_0^*$  is  $w^*$ -strongly extreme point of unit ball of  $X^*$  and  $y_0^*$  is  $w^*$ -strongly extreme point of the unit ball of  $Y^*$ . We show that similar conclusion holds if the rank one operator is a point of weak\*-weak continuity for the identity mapping on the unit sphere of  $\mathcal{L}(X, Y^*)$ . We also study extremal phenomenon in the unit ball of  $\mathcal{L}(X, Y^*)^*$ . We show that if a point  $z \in S_{\mathcal{L}(X, Y^*)^*}$  is a  $w^*$ -strongly extreme point of the unit ball, then  $z = x \otimes y$  for some  $w^*$ -strongly extreme points  $x \in S_X$  and  $y \in S_Y$ , provided the space of compact operators,  $\mathcal{K}(X, Y^*)$  is separating for  $X \hat{\otimes}_\pi Y$ .

## 1. INTRODUCTION

Let  $X$  be a Banach space. We denote the closed unit ball of a Banach space  $X$  by  $B_X$ , the unit sphere by  $S_X$ . We first recall some geometric notations that we will be studying in this paper. See [10] and [5]. Let  $C$  be a bounded subset of  $X$ ,  $x^* \in X^*$  and  $\alpha > 0$ . Then the set

$$S(C, x^*, \alpha) = \{x \in C : x^*(x) > \sup x^*(C) - \alpha\}$$

is called the slice determined by  $x^*$  and  $\alpha$ . Without loss of generality, for any slice  $S(C, x^*, \alpha)$ , we can consider the determining functional  $x^* \in S_{X^*}$ . Indeed, for  $x^* \neq 0$  we have  $S(C, x^*, \alpha) = S(C, \frac{x^*}{\|x^*\|}, \frac{\alpha}{\|x^*\|})$  and for  $x^* = 0$  we have  $S(C, x^*, \alpha) = C = S(C, x_0^*, \beta)$  where  $x_0^* \in S_{X^*}$  and  $\beta$  is sufficiently large. Analogously one can define  $w^*$ -slices in  $X^*$  by considering the determining functional from the predual space (instead of dual space).

---

2020 *Mathematics Subject Classification*. Primary 46A22, 46B10, 46B25; Secondary 46B20, 46B22. **January 28, 2025.**

*Key words and phrases*. Projective tensor product, spaces of operators, geometry of Banach spaces, weak\*-weak point of continuity.

Geometric notions of denting points and weak\* denting points (see [10]) can be described in terms of slices of the unit ball.  $x^* \in S_{X^*}$  is a weak\*-denting point if and only if weak\*-slices containing  $x^*$  form a base for the norm neighbourhood system of  $x^*$ . We refer to the monographs by D. Bourgin ([1]), J. Diestel and J. J. Uhl (see [10]) for basic definitions from the geometry of the unit ball of a Banach space, for vector-valued integration theory and tensor product spaces. The monograph [14] by H. E. Lacey is a standard reference for isometric theory of Banach spaces and [11], [8] for general theory of Banach spaces.

Consider the identity mapping,  $i : (S_{X^*}, weak^*) \rightarrow (S_{X^*}, weak)$ . A point of continuity (if it exists) is called a  $w^*$ - $w$  PC. In the case the range space is equipped with the norm topology, we call the point of continuity as  $w^*$ -norm PC. It is known that a point in the dual unit ball is  $w^*$ -denting if and only if it is  $w^*$ -norm PC and an extreme point, see [16]. Thus an interesting new direction is to consider the notions of points of continuity alone. Here we study the stability of the continuity of elementary tensor in the dual of a projective tensor product space and compare its relation to the continuity of the component vectors in the respective unit spheres.

We recall from [10, Chapter VIII] that the algebraic tensor product of  $X$  and  $Y$  is denoted by  $X \otimes Y$  and it is the space spanned by all elements  $x \otimes y$  where  $x \in X$  and  $y \in Y$ . Given the tensor product  $X \otimes Y$ , the projective norm  $\|\cdot\|_\pi$  on  $X \otimes Y$  is determined by

$$\|u\|_\pi = \inf \left\{ \sum_{i=1}^n \|x_i\| \|y_i\| : u = \sum_{i=1}^n x_i \otimes y_i \right\}$$

The pair  $(X \otimes Y, \|\cdot\|_\pi)$  is sometimes denoted by  $X \otimes_\pi Y$  and its completion is called projective tensor product, denoted by  $X \hat{\otimes}_\pi Y$ . We have from [10, Chapter VIII, Corollary 2] that the space  $\mathcal{L}(X, Y^*)$  is linearly isometric to the dual of  $X \hat{\otimes}_\pi Y$ . Here we always consider the weak\*-topology on  $\mathcal{L}(X, Y^*)$  coming from the projective tensor product of  $X, Y$ .

We start by recalling a classical theorem where  $\mathcal{L}(X, Y^*)$  has a unique predual and the weak\*-topology is uniquely determined. If  $Y$  is such that  $Y^* = C(\Omega)$  for a compact set  $\Omega$ , then it follows that  $\Omega$  is a hyperstonean space [14, Theorem 11, page 96] and for a positive measure  $\mu$ ,  $Y = L^1(\mu)$ .

Now by [10, Chapter VIII, Proposition 11] we see that the space of Bochner integrable function,  $L^1(\mu, X)$  is a predual of  $\mathcal{L}(X, Y^*)$ . When  $X$  is reflexive, that  $L^1(\mu, X)$  is the unique predual of  $\mathcal{L}(X, C(\Omega))$  follows from the results in [6] and [17]. Hence in this case the weak\*-topology on  $\mathcal{L}(X, C(\Omega))$  is uniquely determined.

We now briefly recall some literature on extremal structures preserved by projective tensor products. It is easy to see that  $B_{X \hat{\otimes}_\pi Y} = \overline{\text{co}}(B_X \otimes B_Y)$  (co denotes the convex hull and the closure is taken in the norm topology). However, it is still an open problem, whether every extreme point of  $B_{X \hat{\otimes}_\pi Y}$  must be of the form  $x \otimes y$ , for  $x \in B_X$  and  $y \in B_Y$ . Let  $\mathcal{K}_{w^*}(X^*, Y^*)^*$  denotes the space of all  $w^*$ - $w$  continuous compact operators. It is known from [7] and [20] that, the extreme points of the unit ball,  $\text{ext}(B_{\mathcal{K}_{w^*}(X^*, Y^*)^*}) = \text{ext}(B_{X^*}) \otimes \text{ext}(B_{Y^*})$ , are preserved under this operation. As a consequence, if  $X^*$  or  $Y^*$  has RNP and  $X^*$  or  $Y^*$  has approximation property (see [10, Chapter VIII]) then  $\text{ext}(B_{X^* \hat{\otimes}_\pi Y^*}) = \text{ext}(B_{X^*}) \otimes \text{ext}(B_{Y^*})$ . The situation is better for the stronger notions of denting points and strongly exposed points (see [10] for the definitions of these notions). Very recently, in [15], the notions of preserved extreme point and weak-strongly exposed point in projective tensor products were studied and under some additional assumptions such extreme points turn out to be of the type  $x \otimes y$ , where the component vectors have the same extremal property. This is also partly the theme of this paper.

Some of the classical results include the work of Ruess and Stegall [19], who proved that  $\text{strex}(B_{X \hat{\otimes}_\pi Y}) = \text{strex}(B_X) \otimes \text{strex}(B_Y)$ , where  $\text{strex}(A)$  denotes set of all strongly exposed points of  $A$ . Also for denting points, D. Werner [23] proved an analogous result in more general context. He showed  $\text{dent}(\overline{\text{co}}(C \otimes D)) = \text{dent}(C) \otimes \text{dent}(D)$ , where  $C \subset X$  and  $D \subset Y$  are absolutely convex, bounded and closed subsets and  $\text{dent}(A)$  denotes set of all denting points of  $A$ . As a consequence of Werner's result, it is true that if  $x_0 \in S_X$  and  $y_0 \in S_Y$ , then  $x_0 \otimes y_0$  is  $w^*$ -denting point of  $B_{(X \hat{\otimes}_\pi Y)^{**}}$  if and only if  $x_0$  is  $w^*$ -denting point of  $B_{X^{**}}$  and  $y_0$  is  $w^*$ -denting point of  $B_{Y^{**}}$ .

Motivated by the above results in the case of projective tensors, in this note we also study the notion of  $w^*$ -strongly extreme point, which can be seen as amalgam of two notions namely,  $w^*$ - $w$  PC and extreme.

**Definition 1.0.1.** Let  $X$  be Banach space. A point  $x_0^* \in S_{X^*}$  is called

- (1)  $w^*$ -strongly extreme point of  $B_{X^*}$  if the family of  $w^*$ -slices containing  $x_0^*$  forms a local base for the weak topology of  $X^*$  at  $x_0^*$ .
- (2) point of weak\*-weak continuity ( $w^*$ - $w$  PC) of  $B_{X^*}$  if the identity map from  $(B_{X^*}, w^*)$  to  $(B_{X^*}, w)$  is continuous at  $x_0^*$ .

**Theorem 1.0.2.** [2] A point  $x_0^* \in S_{X^*}$  is  $w^*$ - $w$  PC and extreme point of  $B_{X^*}$  if and only if it is  $w^*$ -strongly extreme point of  $B_{X^*}$ .

In Section 2, we show that if  $x^* \otimes y^*$  is a  $w^*$ - $w$  PC of  $\mathcal{L}(X, Y^*)$  for some  $x^* \in S_{X^*}$  and  $y^* \in S_{Y^*}$ , then the component functionals have the same property. Let  $x, y$  be unit vectors and consider  $x \otimes y$  as an element of  $\mathcal{L}(X, Y^*)^*$  under the canonical embedding. We show in Section 3, that if  $x \otimes y$  is a  $w^*$ - $w$  PC of the identity mapping on the unit sphere, then the same is true of the components  $x, y$  when they are considered as vectors in the corresponding biduals. As an application, we show that if a point  $z \in S_{\mathcal{L}(X, Y^*)^*}$  is a  $w^*$ -strongly extreme point of the unit ball, then  $z = x \otimes y$  for some  $w^*$ -strongly extreme points  $x \in S_X$  and  $y \in S_Y$ , provided the space of compact operators,  $\mathcal{K}(X, Y^*)$  is separating for  $X \hat{\otimes}_\pi Y$ . In Section 4, similar conclusion is obtained for points of weak-norm continuity, improving on the result of Werner [23]. We also give examples of spaces where no compact operator is a point of  $w^*$ -norm PC in  $S_{\mathcal{L}(X, Y^*)}$ . In Section 5, we show that the weak density of  $w^*$ - $w$  PC of  $B_{X^*}$  in  $S_{X^*}$  does not imply that this is the case for any closed subspace  $Y$  of  $X$ . However, it is known that if all points of  $S_{X^*}$  are  $w^*$ - $w$  PC, then each closed subspace  $Y$  of  $X$  inherits the same (see [5]).

However, a main question in this domain remain open.

**Question 1.0.3.** If  $S_{\mathcal{L}(X, Y^*)}$  has a  $w^*$ - $w$  PC, then does the ball always have a rank one operator as a  $w^*$ - $w$  PC ?

## 2. STABILITY IN $B_{\mathcal{L}(X, Y^*)}$

Our first result shows that if an operator of rank one is a  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)}$ , then the same is the case for component vectors.

**Theorem 2.0.1.** Let  $x_0^* \in S_{X^*}$  and  $y_0^* \in S_{Y^*}$ . Also let  $x_0^* \otimes y_0^* \in S_{\mathcal{L}(X, Y^*)}$  be  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)}$ . Then  $x_0^*$  is  $w^*$ - $w$  PC of  $B_{X^*}$  and  $y_0^*$  is  $w^*$ - $w$  PC of  $B_{Y^*}$ .

*Proof.* Consider two weakly open sets

$$\bigcap_{i=1}^n S(B_{X^*}, x_i^{**}, \alpha_i) \text{ and } \bigcap_{j=1}^m S(B_{Y^*}, y_j^{**}, \beta_j)$$

containing  $x_0^*$  and  $y_0^*$  respectively. Choose  $0 < \varepsilon < \min\{x_i^{**}(x_0^*)y_j^{**}(y_0^*) : 1 \leq i \leq n, 1 \leq j \leq m\}$  such that for each  $1 \leq i \leq n$  and  $1 \leq j \leq m$

$$x_i^{**}(x_0^*) - \frac{\varepsilon}{y_j^{**}(y_0^*)} > 1 - \alpha_i \quad \text{and} \quad y_j^{**}(y_0^*) - \frac{\varepsilon}{x_i^{**}(x_0^*)} > 1 - \beta_j.$$

Also let  $\gamma_{ij} := 1 - x_i^{**}(x_0^*)y_j^{**}(y_0^*) + \varepsilon$ . Consider weakly open set

$$W := \bigcap_{i=1}^n \bigcap_{j=1}^m S(B_{\mathcal{L}(X, Y^*)}, x_i^{**} \otimes y_j^{**}, \gamma_{ij}).$$

Observe that  $x_0^* \otimes y_0^* \in W$ . Then there exists a  $w^*$ -open set  $\bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)}, \tau_k, \delta_k)$  such that  $x_0^* \otimes y_0^* \in \bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)}, \tau_k, \delta_k) \subset W$ , where  $\tau_k \in S_{\mathcal{L}(X, Y^*)} = S_{(X \hat{\otimes}_\pi Y)}$  for all  $k = 1, \dots, p$ . Without loss of generality,

let  $\tau_k = \sum_{l=1}^{r(k)} \lambda_{k,l}(x_{k,l} \otimes y_{k,l})$  with  $\lambda_{k,l} > 0$  and  $\sum_{l=1}^{r(k)} \lambda_{k,l} = 1$ . Then

$$\begin{aligned} (x_0^* \otimes y_0^*) \left( \sum_{l=1}^{r(k)} \lambda_{k,l}(x_{k,l} \otimes y_{k,l}) \right) &> 1 - \delta_k \quad \forall 1 \leq k \leq p \\ \Rightarrow \sum_{l=1}^{r(k)} (\lambda_{k,l} y_0^*(y_{k,l})) x_0^*(x_{k,l}) &> 1 - \delta_k \quad \forall 1 \leq k \leq p. \end{aligned}$$

Hence  $\sum_{l=1}^{r(k)} \eta_{k,l} x_0^*(x_{k,l}) > 1 - \delta_k$ , where  $\eta_{k,l} := \lambda_{k,l} y_0^*(y_{k,l})$  for all  $k = 1, \dots, p$ .

Thus

$$x_0^* \in \bigcap_{k=1}^p S(B_{X^*}, \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l}, \left\| \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l} \right\| - 1 + \delta_k).$$

Let  $x^* \in \bigcap_{k=1}^p S(B_{X^*}, \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l}, \left\| \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l} \right\| - 1 + \delta_k)$ . Then

$$x^* \otimes y_0^* \in \bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)}, \tau_k, \delta_k) \subset W.$$

Thus  $x_i^{**}(x^*) > \frac{1 - \gamma_{i,j}}{y_j^{**}(y_0^*)} > 1 - \alpha_i$  for all  $i = 1, \dots, n$ . Hence

$$x_0^* \in \bigcap_{k=1}^p S(B_{X^*}, \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l}, \left\| \sum_{l=1}^{r(k)} \eta_{k,l} x_{k,l} \right\| - 1 + \delta_k) \subset \bigcap_{i=1}^n S(B_{X^*}, x_i^{**}, \alpha_i).$$

Similarly,

$$y_0^* \in \bigcap_{k=1}^p S(B_{Y^*}, \sum_{l=1}^{r(k)} \zeta_{k,l} x_{k,l}, \|\sum_{l=1}^{r(k)} \zeta_{k,l} x_{k,l}\| - 1 + \delta_k) \subset \bigcap_{j=1}^m S(B_{Y^*}, y_j^*, \beta_j),$$

where  $\zeta_{k,l} := \lambda_{k,l} x_0^*(x_{k,l})$  for all  $k = 1, \dots, p$ . Consequently,  $x_0^*$  and  $y_0^*$  are  $w^*$ - $w$  PC of  $B_{X^*}$  and  $B_{Y^*}$  respectively.  $\square$

**Corollary 2.0.2.** *Let  $x_0^* \in S_{X^*}$  and  $y_0^* \in S_{Y^*}$ . Also let  $x_0^* \otimes y_0^* \in S_{\mathcal{L}(X, Y^*)}$  be  $w^*$ -strongly extreme point of  $B_{\mathcal{L}(X, Y^*)}$ . Then  $x_0^*$  is  $w^*$ -strongly extreme point of  $B_{X^*}$  and  $y_0^*$  is  $w^*$ -strongly extreme point of  $B_{Y^*}$ .*

*Proof.* Observe that  $x_0^* \otimes y_0^*$  is extreme point of  $B_{\mathcal{L}(X, Y^*)}$  gives  $x_0^*$  and  $y_0^*$  are extreme point of  $B_{X^*}$  and  $B_{Y^*}$  respectively. Rest follows from Theorem 1.0.2 and 2.0.1.  $\square$

We recall that investigation of  $w^*$ - $w$  PC is well connected with the uniqueness of the Hahn-Banach extensions. See [13, Lemma III.2.14].

**Lemma 2.0.3.** [13, Lemma III.2.14] *Let  $X$  be Banach space. Then,  $x^* \in S(X^*)$  is a point of continuity of  $id : (X_1^*, w^*) \rightarrow (X_1^*, w)$  if and only if  $x^*$  has a unique norm preserving extension to  $X^{**}$ .*

In the following proposition in order to show  $w^*$ - $w$  continuity at an elementary tensor, we consider a different additional conditions on the second component. We recall the abstract notion of a unitary from [4],  $y_0 \in S(Y)$  is said to be a unitary if  $Y^* = \text{span}\{y^* \in B_{Y^*} : y^*(y_0) = 1\}$ . It follows from the results in [4] that any unitary is a strongly extreme point and unitaries remain in as unitaries in the bidual  $Y^{**}$ .

**Proposition 2.0.4.** *Let  $Y$  be a finite dimensional Banach space and suppose  $y_0^* \in S_{Y^*}$  is a unitary. Then for any Banach space  $X$  with  $x_0^* \in S_{X^*}$  is  $w^*$ - $w$  PC, we have  $x_0^* \otimes y_0^* \in S_{\mathcal{L}(X, Y^*)}$  is  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)}$ .*

*Proof.* Let  $\dim(Y) = n$  and  $D(y_0^*) = \{y \in S_Y : y_0^*(y) = 1\}$ . Since  $y_0^*$  is unitary, then there exist  $n$  linearly independent element  $y_1, \dots, y_n \in D(y_0^*)$  such that  $\text{span}\{D(y_0^*)\} = \text{span}\{y_1, \dots, y_n\} = Y$ . Also since  $Y$  is finite dimensional, then  $\mathcal{L}(X, Y^*)^* = X^{**} \hat{\otimes}_\pi Y^{**}$  (see [3]). If possible let  $T_1$  and  $T_2$  be two distinct Hahn-Banach extension of  $x_0^* \otimes y_0^*$ , i.e., for each  $i = 1, 2$ ,

$$T_i : \mathcal{L}(X, Y^*)^* \rightarrow \mathbb{R} \text{ such that } T_i \Big|_{X \hat{\otimes}_\pi Y} = x_0^* \otimes y_0^* \text{ and } \|T_i\| = 1$$

For each  $i = 1, 2$  and  $k = 1, \dots, n$ , we define

$$\eta_{i,k} : X^{**} \rightarrow \mathbb{R} \text{ by } \eta_{i,k}(x^{**}) = T_i(x^{**} \otimes y_k)$$

Observe that

$$\eta_{i,k}(x) = T_i(x \otimes y_k) = (x_0^* \otimes y_0^*)(x \otimes y_k) = x_0^*(x)y_0^*(y_k) = x_0^*(x) \quad \forall x \in X$$

Thus  $\eta_{i,k}|_X = x_0^*$ . Also  $\|\eta_{i,k}\| = 1$ . Indeed,  $\|\eta_{i,k}\| \leq \|T_i\| = 1$ . Choose  $x_n \in S_X$  with  $\lim_{n \rightarrow \infty} x_0^*(x_n) = 1$ . Then  $\|\eta_{i,k}\| \geq \lim_{n \rightarrow \infty} \eta_{i,k}(x_n) = \lim_{n \rightarrow \infty} x_0^*(x_n) = 1$ . Hence  $\eta_{i,k}$  ( $i = 1, 2$  and  $k = 1, \dots, n$ ) are Hahn-Banach extensions of  $x_0^*$ . Since  $x_0^*$  is  $w^*$ - $w$  PC, then

$$\begin{aligned} \eta_{1,k} &= \eta_{2,k} & \forall k &= 1, \dots, n \\ \Rightarrow \eta_{1,k}(x^{**}) &= \eta_{2,k}(x^{**}) & \forall k &= 1, \dots, n, \forall x^{**} \in X^{**} \\ \Rightarrow T_1(x^{**} \otimes y_k) &= T_2(x^{**} \otimes y_k) & \forall k &= 1, \dots, n, \forall x^{**} \in X^{**} \\ \Rightarrow T_1 &= T_2 \end{aligned}$$

This is a contradiction. Hence,  $x_0^* \otimes y_0^*$  has unique Hahn-Banach extension to  $\mathcal{L}(X, Y^*)^*$ . Thus,  $x_0^* \otimes y_0^*$  is  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)}$ .  $\square$

Taking into account the fact that  $(X/Y)^* = Y^\perp$ , following proposition ensures the existence of unitaries in finite dimensions.

**Proposition 2.0.5.** *Let  $X$  be a finite dimensional Banach space. Then for any  $x \in S_X$  either  $x$  is unitary in  $X$  or  $\pi(x)$  is a unitary in  $X/Y$ , where  $\pi : X \rightarrow X/Y$  is the quotient map and  $Y = \{x^* \in B_{X^*} : x^*(x) = 1\}^\perp$ .*

### 3. STABILITY IN HIGHER DUALS OF PROJECTIVE TENSOR PRODUCTS

The following proof uses techniques similar to the ones used during the proof of Theorem 2.0.1. For the sake of completeness we give the details of the proof here.

**Theorem 3.0.1.** *Let  $x_0 \in S_X$ ,  $y_0 \in S_Y$  and  $x_0 \otimes y_0$  be  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)}$ . Then  $x_0$  is  $w^*$ - $w$  PC of  $B_{X^{**}}$  and  $y_0$  is  $w^*$ - $w$  PC of  $B_{Y^{**}}$ .*

*Proof.* Consider two weakly open sets

$$\bigcap_{i=1}^n S(B_{X^{**}}, x_i^{***}, \alpha_i) \text{ and } \bigcap_{j=1}^m S(B_{Y^{**}}, y_j^{***}, \beta_j)$$

containing  $x_0$  and  $y_0$  respectively. Choose  $0 < \varepsilon < \min\{x_i^{***}(x_0)y_j^{***}(y_0) : 1 \leq i \leq n, 1 \leq j \leq m\}$  such that for each  $1 \leq i \leq n$  and  $1 \leq j \leq m$

$$x_i^{***}(x_0) - \frac{\varepsilon}{y_j^{***}(y_0)} > 1 - \alpha_i \quad \text{and} \quad y_j^{***}(y_0) - \frac{\varepsilon}{x_i^{***}(x_0)} > 1 - \beta_j.$$

Also let  $\gamma_{ij} := 1 - x_i^{***}(x_0)y_j^{***}(y_0) + \varepsilon$ . Consider weakly open set

$$W := \bigcap_{i=1}^n \bigcap_{j=1}^m S(B_{\mathcal{L}(X, Y^*)^*}, \psi_{i,j}, \gamma_{ij}),$$

where  $\psi_{i,j}$  is a norm preserving extension of  $x_i^{***} \otimes y_j^{***}$  from  $X^{**} \otimes Y^{**}$  to  $\mathcal{L}(X, Y^*)^*$ . Observe that  $x_0 \otimes y_0 \in W$ . Then there exists a  $w^*$ -open set  $\bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)^*}, T_k, \delta_k)$  such that  $x_0 \otimes y_0 \in \bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)^*}, T_k, \delta_k) \subset W$ , where  $T_k \in S_{\mathcal{L}(X, Y^*)}$  for all  $k = 1, \dots, p$ . Then

$$\begin{aligned} (x_0 \otimes y_0)(T_k) &> 1 - \delta_k \quad \forall 1 \leq k \leq p \\ \Rightarrow (y_0 \circ T_k)(x_0) &> 1 - \delta_k \quad \forall 1 \leq k \leq p. \end{aligned}$$

Thus

$$x_0 \in \bigcap_{k=1}^p S(B_{X^{**}}, y_0 \circ T_k, \|y_0 \circ T_k\| - 1 + \delta_k).$$

Let  $x^{**} \in \bigcap_{k=1}^p S(B_{X^{**}}, y_0 \circ T_k, \|y_0 \circ T_k\| - 1 + \delta_k)$ . Then

$$x^{**} \otimes y_0 \in \bigcap_{k=1}^p S(B_{\mathcal{L}(X, Y^*)^*}, T_k, \delta_k) \subset W.$$

Thus  $x_i^{***}(x^{**}) > \frac{1 - \gamma_{i,j}}{y_j^{***}(y_0)} > 1 - \alpha_i$  for all  $i = 1, \dots, n$ . Hence

$$x_0 \in \bigcap_{k=1}^p S(B_{X^{**}}, y_0 \circ T_k, \|y_0 \circ T_k\| - 1 + \delta_k) \subset \bigcap_{i=1}^n S(B_{X^{**}}, x_i^{***}, \alpha_i).$$

Similarly,

$$y_0 \in \bigcap_{k=1}^p S(B_{Y^{**}}, T_k(x_0), \|T_k(x_0)\| - 1 + \delta_k) \subset \bigcap_{j=1}^m S(B_{Y^{**}}, y_j^{***}, \beta_j).$$

Consequently,  $x_0$  and  $y_0$  are  $w^*$ - $w$  PC of  $B_{X^{**}}$  and  $B_{Y^{**}}$  respectively.  $\square$

The example below shows that  $X$  and  $Y$  can be considered in such a way that no simple tensor in  $S_{\mathcal{L}(X, Y^*)^*}$  is  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)^*}$ .



**Example 3.0.2.** Consider  $L^1([0, 1], X) (= L^1[0, 1] \hat{\otimes}_\pi X)$  with Lebesgue measure on  $[0, 1]$ . Suppose  $f \otimes x$  is a  $w^*$ - $w$  PC of  $L^1([0, 1], X)^{**}$ , where  $f \in S_{L^1([0, 1])}$  and  $x \in S_X$ . Then by Theorem 3.0.1,  $f$  is a  $w^*$ - $w$  PC of  $S_{L^1([0, 1])}^{**}$ . Let  $\Omega$  be the Stone space of  $L^\infty([0, 1])$ . Then  $f \in S_{C(\Omega)^*}$  is a  $w^*$ - $w$  PC. Since  $B_{C(\Omega)^*}$  is the weak\*-closure of finitely supported measures, by the  $w^*$ - $w$  continuity of  $f$ , we get that  $f$  is a norm limit of a sequence of finitely supported measures. Since the Lebesgue measure is non-atomic, we get a contradiction.

However, the example above doesnot imply that  $B_{L^1([0, 1], X)^{**}}$  has no  $w^*$ - $w$  PC. Even when  $X$  is an infinite dimensional reflexive space, we do not know if,  $S_{L^1([0, 1], X)^{**}}$  has any  $w^*$ - $w$  PC's ?

**Corollary 3.0.3.** Let  $x_0 \in S_X$  and  $y_0 \in S_Y$ . Also let  $x_0 \otimes y_0 \in S_{\mathcal{L}(X, Y^*)^*}$  be  $w^*$ -strongly extreme point of  $B_{\mathcal{L}(X, Y^*)^*}$ . Then  $x_0$  is  $w^*$ -strongly extreme point of  $B_{X^{**}}$  and  $y_0$  is  $w^*$ -strongly extreme point of  $B_{Y^{**}}$ .

*Proof.* Since  $x_0 \otimes y_0$  is  $w^*$ - $w$  PC and extreme in  $B_{\mathcal{L}(X, Y^*)^*}$ , then by [9, Remark 2.0.6],  $x_0 \otimes y_0$  is extreme point of  $B_{X \hat{\otimes}_\pi Y}$ . Hence  $x_0$  and  $y_0$  are extreme points of  $B_X$  and  $B_Y$  respectively. Rest follows from Theorem 1.0.2 and 3.0.1.  $\square$

From [9, Remark 2.0.6], it is known that, for any Banach space  $X$ ,  $w^*$ -strest  $(B_{X^{**}}) \subset \text{ext}(B_X)$ , where  $w^*$ -strest  $(C)$  denotes the set of  $w^*$ -strongly extreme points of  $C$ . As a consequence, every  $w^*$ -strongly extreme points of  $B_{X^{**}}$  is preserved extreme point of  $B_X$ .

**Theorem 3.0.4.** [15, Theorem 1.1] Let  $X$  and  $Y$  be Banach spaces with  $\mathcal{K}(X, Y^*)$  is separating for  $X \hat{\otimes}_\pi Y$ . If  $z$  is preserved extreme point of  $B_{X \hat{\otimes}_\pi Y}$ , then  $z = x \otimes y$  where  $x$  and  $y$  are preserved extreme points of  $B_X$  and  $B_Y$  respectively.

Therefore, taking into account Corollary 3.0.3 and Theorem 3.0.4, we have following.

**Corollary 3.0.5.** Let  $X$  and  $Y$  be Banach spaces with  $\mathcal{K}(X, Y^*)$  is separating for  $X \hat{\otimes}_\pi Y$ . If  $z \in S_{\mathcal{L}(X, Y^*)^*}$  is  $w^*$ -strongly extreme point of  $B_{\mathcal{L}(X, Y^*)^*}$ , then  $z = x \otimes y$  for some  $w^*$ -strongly extreme points  $x \in S_X$  and  $y \in S_Y$ . In other words,

$$w^*\text{-strest}(B_{\mathcal{L}(X, Y^*)^*}) \subset w^*\text{-strest}(B_{X^{**}}) \otimes w^*\text{-strest}(B_{Y^{**}}) \subset B_X \otimes B_Y.$$

Following technique similar to the ones given in Proposition 2.0.4, we get the following.

**Proposition 3.0.6.** *Let  $Y$  be a finite dimensional Banach space and  $y_0 \in S_Y$  be unitary element. Then for any Banach space  $X$  with  $x_0 \in S_X$  is  $w^*$ - $w$  PC of  $B_{X^{**}}$ , we have  $x_0 \otimes y_0$  is  $w^*$ - $w$  PC of  $B_{\mathcal{L}(X, Y^*)^*}$ .*

#### 4. STABILITY OF WEAK-NORM PC'S

In this section we consider the weaker version of denting points, by considering only points of weak-norm continuity on  $S_X$ . We first obtain a unit ball analogue of the results from [23]. A connection to the weak\*-topology comes from the fact that  $x_0 \in S_X$  is weak-norm PC in  $B_X$  if and only if  $x_0$  is weak\*-norm PC in  $B_{X^{**}}$ , see [12]. In particular we get that  $x_0$  is a weak\*-norm PC in all higher ordered even duals of  $X$ .

**Proposition 4.0.1.** *Let  $X$  and  $Y$  be Banach spaces. Also let  $x_0 \in S_X$ ,  $y_0 \in S_Y$  and  $x_0 \otimes y_0$  be weak-norm PC in  $B_{X \hat{\otimes}_\pi Y}$ . Then  $x_0$  is weak-norm PC in  $B_X$  and  $y_0$  is weak-norm PC in  $B_Y$ .*

*Proof.* Let  $\varepsilon > 0$ . Since  $x_0 \otimes y_0$  is weak-norm PC in  $B_{X \hat{\otimes}_\pi Y}$ , then there exists a weakly open set

$$W = \{\tau \in B_{(X \hat{\otimes}_\pi Y)} : |T_i(\tau) - T_i(x_0 \otimes y_0)| < \gamma, i = 1, \dots, p\}$$

where  $T_1, \dots, T_p \in (X \hat{\otimes}_\pi Y)^* = \mathcal{L}(X, Y^*)$  and  $x_0 \otimes y_0 \in W \subset B(x_0 \otimes y_0, \varepsilon)$ . Consider weakly open sets

$$W_1 = \{x \in B_X : |(y_0 \circ T_i)(x) - (y_0 \circ T_i)(x_0)| < \gamma, i = 1, \dots, p\}$$

$$W_2 = \{y \in B_Y : |(T_i(x_0))(y) - (T_i(x_0))(y_0)| < \gamma, i = 1, \dots, p\}$$

in  $B_X$  and  $B_Y$  respectively. Then  $x_0 \in W_1$  and  $y_0 \in W_2$ . Moreover,  $W_1 \otimes \{y_0\} \subset W$  and  $\{x_0\} \otimes W_2 \subset W$ . Therefore, we can conclude  $x_0 \in W_1 \subset B(x_0, \varepsilon)$  and  $y_0 \in W_2 \subset B(y_0, \varepsilon)$ . Indeed, let  $x \in W_1$ . Then  $x \otimes y_0 \in W$ . Thus  $\|x - x_0\| = \|(x \otimes y_0) - (x_0 \otimes y_0)\| < \varepsilon$ . Hence,  $W_1 \subset B(x_0, \varepsilon)$  and similarly  $W_2 \subset B(y_0, \varepsilon)$ .  $\square$

**Corollary 4.0.2.** *Let  $X$  and  $Y$  be Banach spaces. Also let  $x_0 \in S_X$ ,  $y_0 \in S_Y$  and  $x_0 \otimes y_0$  be weak\*-norm PC in  $B_{\mathcal{L}(X, Y^*)^*}$ . Then  $x_0$  is weak\*-norm PC in  $B_{X^{**}}$  and  $y_0$  is weak\*-norm PC in  $B_{Y^{**}}$ .*

We recall from [13] that a closed subspace  $J \subset X$  is said to be a  $M$ -ideal, if there is a linear projection  $P : X^* \rightarrow X^*$  such that  $\|x^*\| = \|P(x^*)\| + \|x^* - P(x^*)\|$  for all  $x^* \in X^*$  and  $\ker(P) = J^\perp$ . See [13, Chapter VI] for examples of spaces  $X, Y$ , for which  $\mathcal{K}(X, Y)$  is a  $M$ -ideal in  $\mathcal{L}(X, Y)$ . It is known that  $M$ -ideals do not inherit stronger form of extremal structure from  $S_X$ . Our next theorem is a point of continuity version of this nature.

**Theorem 4.0.3.** *Let  $J \subset X$  be a  $M$ -ideal of infinite codimension. Then  $J$  has no weak-norm PC of  $S_X$ .*

*Proof.* Since  $J$  is a  $M$ -ideal of  $X$ , we have,  $X^{**} = (J^*)^\perp \oplus_\infty J^{\perp\perp}$  ( $\ell^\infty$ -sum) and  $(J^*)^\perp$  is infinite dimensional. Suppose  $x_0 \in S_X$  be a weak-norm PC and  $x_0 \in J \subset J^{\perp\perp}$ . Again by the result from [12] quoted earlier, we get that  $x_0$  is a weak\*-norm PC of  $S_{X^{**}}$ . Now since  $(J^*)^\perp$  is infinite dimensional, there is a net  $\{x_\alpha^{**}\}_{\alpha \in \Delta} \subset S_{(J^*)^\perp}$  such that  $x_\alpha^{**} \rightarrow 0$  in the weak\*-topology. Now  $\|x_0 + x_\alpha^{**}\| = \max\{\|x_0\|, 1\} = 1$  and  $x_0 + x_\alpha^{**} \rightarrow x_0$  in the weak\*-topology but not in the norm. This contradiction shows that  $x_0 \notin J$ .  $\square$

The following corollary is now easy to see. See [13, Chapter VI] to see the examples of spaces that satisfy the assumptions of this corollary. In particular, for  $1 < p < \infty$ ,  $\mathcal{K}(\ell^p)$  is a  $M$ -ideal of infinite codimension in  $\mathcal{L}(\ell^p)$ .

**Corollary 4.0.4.** *Suppose  $X, Y$  are such that  $\mathcal{K}(X, Y^*)$  is a  $M$ -ideal of infinite codimension in  $\mathcal{L}(X, Y^*)$ . Then no compact operator is a point of  $w^*$ -norm PC in  $S_{\mathcal{L}(X, Y^*)}$ .*

We conclude with a version of Theorem 4.0.3 for weak\*-strongly extreme points.

**Proposition 4.0.5.** *Let  $J \subset X^*$  be a proper  $M$ -ideal.  $J$  has no weak\*-strongly extreme points of  $B_{X^*}$ .*

*Proof.* We only indicate the modification in the proof of Theorem 4.0.3. Suppose  $x_0^* \in J$  is weak\*-strongly extreme point of  $B_{X^*}$ . As before,  $X^{***} = (J^*)^\perp \oplus_\infty J^{\perp\perp}$  and the decomposition is non-trivial. Again, we have from Lemma 2.0.3 that  $x_0^* \in J^{\perp\perp}$  is an extreme point of  $B_{X^{***}}$ . This contradicts the fact that an extreme point in an  $\ell^\infty$ -sum must have all the coordinates as unit vectors. Thus  $x_0^* \notin J$ .  $\square$

5. SOME RESULTS ON  $w^*$ - $w$  PC

Let  $X$  be a Banach space. We say that  $X$  has Property (\*) if the set of  $w^*$ - $w$  PC of  $B_{X^*}$  is weakly dense in  $S_{X^*}$ . We recall that if weak\* and weak topologies coincide on  $S_{X^*}$ , by Lemma 2.0.3, we get that, under the canonical embedding of  $X \subset X^{**}$ , elements of  $X^*$  have unique norm-preserving extension. Such spaces are called Hahn-Banach smooth spaces (see [21]) and it was proved in [5] that being Hahn-Banach smooth is a hereditary property. Similarly if one assumes, only norm attaining functionals of  $X^*$  have unique norm-preserving extensions, then by the Bishop-Phepls theorem [11] and Lemma 2.0.3, we see that  $w^*$ - $w$  PC's are norm dense in  $S_{X^*}$ .

**Proposition 5.0.1.** *Property (\*) is not hereditary.*

*Proof.* As noted in Section 1, the set of  $w^*$ -denting points of  $B_{X^*}$  is contained in set of  $w^*$ - $w$  PC of  $B_{X^*}$ . Also every Banach space  $X$  can be isometrically embedded into a Banach space  $Z$  such that the set of  $w^*$ -denting points of  $B_{Z^*}$  is dense in  $S_{Z^*}$  (see [22, Corollary 2.8]). Therefore, it follows that every Banach space  $X$  can be isometrically embedded into a Banach space  $Z$  such that the set of  $w^*$ - $w$  PC of  $B_{Z^*}$  is weakly dense in  $S_{Z^*}$ . Thus if Property (\*) were hereditary, then every Banach space would have Property (\*), which is not true. Hence, Property (\*) is not hereditary.  $\square$

**Remark 5.0.2.** *Unlike some of the investigations here, the main difficulty in dealing with  $w^*$ - $w$  points of continuity in a dual space  $X^*$  is that they need not be preserved in the bidual,  $X^{***}$  (under the canonical embedding), see [9]. On the other hand if  $x^* \in B_{X^*}$  is a weak\*-strongly extreme point, by Lemma 2.0.3 we see that  $x^*$  is at least an extreme point of  $B_{X^{***}}$ . Let  $X$  be a Banach space such that under the canonical embedding in  $X^{**}$  it is a  $M$ -ideal. These are called  $M$ -embedded spaces. See [13, Chapter III, VI] for several examples of  $M$ -embedded spaces from function theory and operator theory. It was shown in [18] that for a non-reflexive  $M$ -embedded space  $X$ , any  $x^* \in S_{X^*}$  continues to be a  $w^*$ - $w$  PC of the unit ball of all higher order odd duals of  $X$ .*

Thus we have the following variation of Proposition 2.0.4. For a non-reflexive space  $X$  and a positive integer  $k > 1$ , we denote by  $X^{(k)}$ , the  $k$ th ordered dual of  $X$ . We again recall that since  $Y$  is finite dimensional,  $\mathcal{L}(X^{(k)}, Y) = (X^{(k)} \hat{\otimes}_{\pi} Y^*)^*$ . The following theorem is easy to see.

**Theorem 5.0.3.** *Let  $X$  be a non-reflexive  $M$ -embedded space and let  $Y$  be a finite dimensional space with a unitary  $y_0^* \in S_{Y^*}$ . For any  $x_0^* \in S(X^*)$ ,  $x_0^* \otimes y_0^*$  is a  $w^*$ - $w$  PC of  $S_{\mathcal{L}(X^{(k)}, Y)}$  for any even integer  $k$ .*

#### ACKNOWLEDGMENTS

The Part of this work was done when the second author was visiting Shiv Nadar Institute. She would like to express her deep gratitude to everyone at Shiv Nadar Institute for the warm hospitality provided during her visit. She would also like to sincerely thank Dr. Priyanka Grover for the opportunity to visit. The research of T. S. S. R. K. Rao is supported by a 3-year project ‘Classification of Banach spaces using differentiability’ funded by the Anusandhan National Research Foundation (ANRF), Core Research Grant, CRG2023-000595.

#### REFERENCES

- [1] Bourgin, R. D. *Geometric properties of convex sets with the Radon-Nikodým property*, Lecture Notes in Mathematics, Springer-Verlag Berlin **993** (1983).
- [2] Bandyopadhyay, P.; Basu, S. *On a new asymptotic norming property*, Indag. Mathem. **10** (1999), no. 1, 15–23.
- [3] Botelho, F.; Fleming, R. J. *The dual of the space of bounded operators on a Banach space*, Concr. Oper. **8** (2021), 48–59.
- [4] Bandyopadhyay, P.; Jarosz, K.; Rao, T. S. S. R. K. *Unitaries in Banach spaces*, Illinois J. Math. **48** (2004), no. 1, 339–351.
- [5] Basu, S.; Rao, T. S. S. R. K. *Some stability results for asymptotic norming properties in Banach spaces*, Colloq. Math. **75** (1998), no. 2, 271–284.
- [6] Cambern, M.; Greim, P. *Spaces of continuous vector functions as duals*, Canad. Math. Bull. **31** (1988), no. 1, 70–78.
- [7] Collins, H. S.; Ruess, W. *Weak compactness in spaces of compact operators and of vector-valued functions*, Pac. J. Math. **106** (1983), no. 1, 45–71.
- [8] Diestel, J. *Sequences and series in Banach spaces*, Springer-Verlag, Berlin **92** (1984).
- [9] Daptari, S.; Rao, T. S. S. R. K. *A study of weak\*-weak point of continuity in the unit ball of dual spaces*, Preprint (2024).
- [10] Diestel, J.; Uhl, J. J. *Vector measures*, Mathematical Surveys, Amer. Math. Soc., Providence, RI **15** (1977).
- [11] Holmes, R. B. *Geometric Functional analysis and its applications*, Springer-Verlag, Berlin **24** (1975).
- [12] Hu, Z.; Lin, B. L. *RNP and CPCP in Lebesgue-Bochner function spaces*, Illinois J. Math. **37** (1993) 329–347.

- [13] Harmand, P.; Werner, D.; Werner, W. *M-ideals in Banach spaces and Banach algebras*, Lecture Notes in Mathematics, Springer-Verlag, Berlin **1547** (1993).
- [14] Lacey, H. E. *The isometric theory of classical Banach spaces*, Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin-Heidelberg-New York **208** (1974).
- [15] García-Lirola, L. C.; Grelier, G.; Martínez-Cervantes, G.; Zoca, A. R. *Extremal structure of projective tensor products*, Results Math **78** (2023).
- [16] Lin, B. L. ; Lin, P. K.; Troyanski, S. *Characterizations of denting points*, Proc. Amer. Math. Soc. **102** (1988), no. 3, 526–528.
- [17] Rao, T. S. S. R. K.  $\mathcal{L}(X, C(K))$  as a dual space, Proc. Amer. Math. Soc. **110** (1990), no. 3, 727–729.
- [18] Rao, T. S. S. R. K. *How many unique norm preserving extensions are needed to determine the structure of a subspace?*, preprint (2024).
- [19] Ruess, W. M.; Stegall, C. P. *Exposed and denting points in duals of operator spaces*, Isr. J. Math. **53** (1986), no. 2, 163–190.
- [20] Ruess, W. M.; Stegall, C. P. *Extreme points in duals of operator spaces*, Math. Ann. **261** (1982), no. 4, 535–546.
- [21] Sullivan, F. *Geometric properties determined by higher duals of a Banach space*, Illinois J. Math. **21** (1977), 315–331.
- [22] Sevilla, M. J.; Moreno, J. P. *Renorming Banach spaces with the Mazur Intersection Property*, J. Funct. Anal. **144** (1997), 486–504.
- [23] Werner, D. *Denting points in tensor products of Banach spaces*, Proc. Amer. Math. Soc. **101** (1987), no. 1, 122–126.

<sup>1</sup> DEPARTMENT OF MATHEMATICS, SHIV NADAR INSTITUTION OF EMINENCE. GAUTAM BUDDHA NAGAR-201314, INDIA

*Email address:* [srin@fulbrightmail.org](mailto:srin@fulbrightmail.org)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, RAMAKRISHNA MISSION VIVEKANANDA EDUCATIONAL AND RESEARCH INSTITUTE, BELUR MATH, HOWRAH 711202, WEST BENGAL, INDIA

*Email address:* [susmitaseal1996@gmail.com](mailto:susmitaseal1996@gmail.com)