Recent Developments in Stochastic Inflation

R. P. Woodard^{\dagger}

Department of Physics, University of Florida, Gainesville, FL 32611, UNITED STATES

ABSTRACT

This article is dedicated to the memory of Alexei Starobinsky. I begin with some recollections of him and then review the generalization of his wonderful stochastic formalism from scalar potential models to theories which interact with fermions and photons, and finally to theories with derivative interactions such as nonlinear sigma models and gravity. This entails effective potentials generated by the usual field-dependent masses, as well as by field-dependent field strengths, and by field-dependent Hubble parameters. I also discuss secular loop corrections which cannot be captured by stochastic techniques.

PACS numbers: 04.50.Kd, 95.35.+d, 98.62.-g

[†] e-mail: woodard@phys.ufl.edu

1 Introduction

Alexei Starobinsky was a genius whose contributions to cosmology include one of the earliest models of primordial inflation [1] which is still completely viable [2]. He also made the remarkable suggestion that quantum fluctuations from the epoch of primordial inflation might be observable today [3] and can even be used to constrain models of inflation [4].

Of particular interest to me is the stochastic technique Alexei devised to explain the secular growth one finds in the correlation functions of some quantum field theories on inflationary backgrounds [5]. With Yokoyama, he even developed a way of summing up the leading secular corrections to all orders [6]. I believe this technique will eventually be recognized as equal in significance to the development of the renormalization group by the late Ken Wilson for flat space quantum field theory and statistical mechanics [7,8]. Wilson won the 1982 Nobel Prize for this work and I feel that Alexei's contributions would have been similarly honored, had he only lived.

This article consists of 7 sections. In the next one I share some personal recollections of Alexei from the two decades during which I was privileged to have known him. In section 3 I review the large secular loop corrections which occur in scalar potential models on inflationary backgrounds, and I prove that Alexei's stochastic formalism recovers the leading logarithms produced at each order in perturbation theory. Section 4 explains how to handle charged scalars and those with Yukawa couplings. In section 5 I discuss derivative interactions involving scalars and gravity. Theories with such derivative interactions can experience another sort of leading logarithm loop correction which cannot be described stochastically but is instead amenable to treatment by a variant of the renormalization group. This is described in section 6. My conclusions comprise section 7.

2 Remembering Alexei Starobinsky

American elites are infamous for their view that the only interesting parts of my country are portions of the coasts, extending roughly from Boston to Washington in the East, and from Seattle to Los Angeles in the West. Many dismiss the rest of the nation as "fly-over country" which they only see while traveling between the parts they consider civilized. I'm told that Russian elites take a similar view of Moscow and Saint Petersburg. Alexei was not like that. Although he had many friends and collaborators outside of Russia, and could surely have emigrated had he wished, he stood by his country during hard times, and supported physics throughout its vast expanse. Indeed, he contracted the illness which took his life while attending an event in Kazan. And I first met Alexei in 2002, at a conference in the Siberian city of Tomsk.

At that time of our first meeting I had been applying the Schwinger-Keldysh formalism [9–13], with dimensional regularization on de Sitter background, to work out loop corrections in renormalizable theories of massless, minimally coupled scalars which interact with themselves [14], or with electromagnetism [15,16]. It had long been known that the coincidence limit of the massless, minimally coupled scalar propagator on de Sitter shows secular growth [4,17,18]. This observation had eventually led to a full-blown proof that there are no normalizable de Sitter invariant states in this theory [19]. What I was showing is that the secular growth persists when realistic interactions are added.

I confess to having felt a bit persecuted at the time of the Tomsk meeting because both quantum field theory experts and cosmologists were, for different reasons, dismissing my carefully derived results as nonsense. The point which bothered quantum field theory people is that the secular growth I found breaks de Sitter invariance. They were convinced that de Sitter invariance should prove as powerful an organizing principle for QFT on de Sitter background as Poincaré invariance does on flat space background. They therefore suspected that my de Sitter breaking effects must derive from a poorly chosen vacuum, or from having posed unphysical questions [20], or from outright mistakes. On the other hand, some cosmologists altogether rejected the application of QFT on inflationary backgrounds. They argued that QFT expectation values amounted to taking unphysical averages over portions of the wave function which had de-cohered and could not be simultaneously accessible to any real observer. Therefore, they concluded that my results were as unphysical as averaging over the alive-dead cat superposition in the famous paradox of Schrödinger [21].

I spoke on my work and ended by asking, with some asperity, "What is *wrong* with this?" Alexei raised his hand and said that he didn't think anything was wrong with my results, and that the secular growth they contained could be predicted by his stochastic formalism. I was dumbfounded (pleasantly) by a reaction so different from the ones I usually encountered. But I was also suspicious that there could be a trivial way of capturing the most important parts of dimensionally regulated and fully renormalized QFT calculations which required weeks of hard work. However, the possibility was too exciting to ignore, and I agreed to check Alexei's predictions against some three loop results I had not published. I screwed up the check at first, but Alexei patiently explained how to do it correctly, at which point I became convinced that his formalism must be right. Alexei was used to the skepticism of QFT experts that an ultraviolet finite, stochastic random variable could reproduce important features of an ultraviolet divergent, quantum field theory, and he must of sensed in me a willing tool to clarify for them a truth which he could see directly. In any case, he asked me to devise a proof from quantum field theory.

Alexei's friends and collaborators will confirm that he sometimes posed difficult problems! That particular one took me and Nick Tsamis three years to solve. When we finally found a proof [22] it was my great pleasure to present it to Alexei during a conference at the Lebedev Institute. Since then Alexei and I have overlapped at many sites: Dubna, Moscow, Paris, Ann Arbor, Utrecht, Cargese, Tallinn, as well as innumerable virtual conferences. I treasured these opportunities to learn from him, and I was seldom disappointed.

Like many brilliant scientists, Alexei was sometimes impatient with lesser men. It was my good fortune that he was always willing to talk with me. We twice began working together on projects, but never brought anything to completion. It saddens me that the opportunity to do so has been forever foreclosed by his untimely passing.

Let me close this section by commenting that, for someone so renowned, Alexei was quite shy. My wife, Shun-Pei, who is also a shy person, picked up on this immediately when she first met him, during the fall of 2006. She had recently discovered the first large logarithm induced by a loop of gravitons in a dimensionally regulated and fully renormalized computation [23, 24], and was attending a 3-month advanced school in Paris while still a graduate student. She and Alexei used to chat quietly in the corner during coffee breaks, while more gregarious and less accomplished physicists held forth loudly in the center. Years later, as a full professor in her own right, Shun-Pei organized the Taiwan portion of a Russia-Taiwan funding proposal with Alexei and Andrei Barvinsky. It was a good proposal and I'm sure it would have won support, however, the entire program was peremptorily terminated due to the worsening of relations that followed the Russian invasion of Ukraine, which Alexei opposed.

3 Why Starobinsky's Formalism Works

One can characterize the geometry of cosmology by its scale factor a(t), Hubble parameter H(t) and first slow roll parameter $\epsilon(t)$,

$$ds^{2} = -dt^{2} + a^{2}(t)d\vec{x} \cdot d\vec{x} \quad , \qquad H(t) \equiv \frac{\dot{a}}{a} \quad , \qquad \epsilon(t) \equiv -\frac{\dot{H}}{H^{2}} \,. \tag{1}$$

Spacetime expansion means that the first derivative of a(t) is positive, which corresponds to H(t) > 0. Inflation means that the second derivative is also positive, which corresponds to $0 \le \epsilon(t) < 1$. We know that this can occur because it's taking place now [25,26]. The simplest interpretation of cosmological data [27] is that it also took place during the very early universe [28], and Alexei's model [1] offers one of the best explanations for what caused it.

For light fields which are not conformally invariant the accelerated expansion of inflation can rip virtual, long wavelength quanta out of the vacuum so that they become real particles. This occurs maximally for massless, minimally coupled scalars, and for gravitons whose linearized equation of motion is the same [29]. On de Sitter background $(a(t) = e^{Ht}$ with H constant) the occupation number for a single wave vector \vec{k} experiences staggering growth,

$$N(t,k) = \left[\frac{a(t)H}{2k}\right]^2.$$
 (2)

This is what caused the (still unresolved) tensor power spectrum [3] and its scalar cousin [30], which has been measured to three significant figures.

The exponentially increasing occupation numbers (2) for super-horizon modes is what makes the coincident scalar propagator grow [4, 17, 18]. The thing which got me in so much trouble was asserting that this growth must persist when interactions are present. The simplest example is a scalar potential model,

$$\mathcal{L} = -\frac{1}{2}\partial_{\mu}\phi\partial_{\nu}\phi g^{\mu\nu}\sqrt{-g} - V(\phi)\sqrt{-g} .$$
(3)

Few would argue that the expectation value of the stress tensor is unphysical. For any reasonable vacuum on the cosmological background (1) it must take the perfect fluid form,

$$\langle \Omega | T_{\mu\nu}(t, \vec{x}) | \Omega \rangle = [\rho(t) + p(t)] \delta^0_{\ \mu} \delta^0_{\ \nu} + p(t) g_{\mu\nu} .$$
 (4)

For the case of $V(\phi) = \frac{1}{4!}\lambda\phi^4$ on de Sitter background the 1-loop result can be absorbed into a renormalization of the cosmological constant but the 2-loop

results (suitably renormalized and with appropriate 1-loop state corrections to absorb exponentially falling terms) show secular growth [14, 31, 32],

$$\rho(t) = \frac{\lambda H^4}{2^7 \pi^4} \times \ln^2[a(t)] + O(\lambda^2) , \qquad (5)$$

$$p(t) = \frac{\lambda H^4}{2^7 \pi^4} \times \left\{ -\ln^2[a(t)] - \frac{2}{3}\ln[a(t)] \right\} + O(\lambda^2) .$$
(6)

It is time for a crucial digression on notation. The $\lambda \ln^2[a(t)]$ corrections in (5-6) are *leading logarithm*; the $\lambda \ln[a(t)]$ contribution to (6) is a *sub-leading logarithm*. For order λ^N corrections in a quartic potential the leading logarithm contribution is $\ln^{2N}[a(t)]$, and any term with fewer factors of $\ln[a(t)]$ is sub-leading. For a general monomial potential $V \sim \lambda \phi^K$ the leading logarithm contribution at order λ^{2N} is $\ln^{K \cdot N}[a(t)]$ [22].

Vakif Onemli and I derived (5-6) the hard way, by explicitly computing the various diagrams in dimensional regularization, then adding counterterms and taking the unregulated limit. It took weeks. Starobinsky devised a way of trivially getting the leading logarithm result, not just for the stress tensor of a quartic potential but for any correlator in any scalar potential model (3), and quite a bit before us [5]. The technique is to replace the exact Heisenberg equation for the ultraviolet divergent quantum field $\phi(t, \vec{x})$,

$$-\frac{1}{\sqrt{-g}}\partial_{\mu}[\sqrt{-g}\,g^{\mu\nu}\partial_{\nu}\phi] = -V'(\phi)\;,\tag{7}$$

with a Langevin equation for the ultraviolet finite stochastic field $\varphi(t, \vec{x})$,

$$3H[\dot{\varphi} - \dot{\varphi}_0] = -V'(\varphi) . \tag{8}$$

The stochastic jitter $\dot{\varphi}_0$ is the time derivative of the infrared truncated, free field mode sum,

$$\varphi_0(t,\vec{x}) = \int_H^{Ha(t)} \frac{d^3k}{(2\pi)^3} \frac{He^{i\vec{k}\cdot\vec{x}}}{\sqrt{2k^3}} \left\{ \alpha(\vec{k}) + \alpha^{\dagger}(-\vec{k}) \right\} , \quad [\alpha(\vec{k}), \alpha^{\dagger}(\vec{p})] = (2\pi)^3 \delta^3(\vec{k} - \vec{p}) .$$
(9)

There are two ways to use Starobinsky's Langevin equation (8). The first is to solve it perturbatively for $\varphi(t, \vec{x})$ as an expansion in powers of $\varphi_0(t, \vec{x})$. For the quartic potential this gives,

$$\varphi(t, \vec{x}) = \varphi_0(t, \vec{x}) - \frac{\lambda}{18H} \int_0^t dt' \,\varphi_0^3(t', \vec{x}) + \frac{\lambda^2}{108H^2} \int_0^t dt' \,\varphi_0^2(t', \vec{x}) \int_0^{t'} dt'' \,\varphi_0^3(t'', \vec{x}) + O(\lambda^3) \,. \tag{10}$$

Substituting in the stress tensor and taking the expectation value gives the leading logarithm expansion for the energy density (5) [22],

$$\rho(t) \longrightarrow \frac{\lambda H^4}{2^7 \pi^4} \ln^2[a(t)] \times \left\{ 1 - \frac{\lambda}{12\pi^2} \ln^2[a(t)] + \frac{53\lambda^2}{2^4 3^4 5\pi^4} \ln^4[a(t)] + O\left(\lambda^3 \ln^6[a(t)]\right) \right\}. \tag{11}$$

Note that I have carried this to *fourth order* in the loop expansion.

The second way of using Starobinsky's Langevin equation (8) is to derive a Fokker-Planck equation for the time-dependent probability density $\rho(t, \varphi)$ [6],

$$\dot{\varrho}(t,\varphi) = \frac{\partial}{\partial\varphi} \left[\frac{V'(\varphi)}{3H} \varrho(t,\varphi) \right] + \frac{H3}{8\pi^2} \frac{\partial^2 \varrho(t,\varphi)}{\partial\varphi^2} .$$
(12)

If the potential is bounded below, and steep enough, the system will approach an equilibrium when the inward pressure of the classical force balances the outward pressure of inflationary particle production. In that case the left hand side of equation (12) vanishes and we can solve for the late time probability density,

$$\varrho(t,\varphi) \longrightarrow N \exp\left[-\frac{8\pi^2 V(\varphi)}{3H^4}\right].$$
(13)

This allows one to sum the series of leading logarithms. For example, the late time limit of the energy density for the quartic potential is,

$$\rho(t) \longrightarrow \frac{3H^4}{8\pi^2} \frac{\Gamma(\frac{3}{4})}{\Gamma(\frac{1}{4})} . \tag{14}$$

The physical interpretation is that inflationary particle production pushes the scalar up its quartic potential until an equilibrium is reached with the downward classical force.

Before giving the proof of Starobinsky's formalism let me comment on how very strange it seems from the perspective of quantum field theory. The original scalar potential model (3) describes a quantum field which obeys a second order equation and whose commutator $[\phi(t, \vec{x}), \phi(t', \vec{x}')]$ is nonzero for timelike and lightlike separations. The correlators of this quantum field are typically ultraviolet divergent and require both regularization and renormalization. In contrast, the infrared truncated mode sum (9) for $\varphi_0(t, \vec{x})$ is a stochastic random variable which commutes with itself everywhere, $[\varphi_0(t, \vec{x}), \varphi_0(t', \vec{x}')] = 0$. Because the mode sum is cut off at k = Ha(t), its correlators are ultraviolet finite. These same two properties are inherited by the full field $\varphi(t, \vec{x})$, which obeys the first order equation (8). Yet correlators of φ agree with those of ϕ at leading logarithm order. You can see why quantum field theory experts were suspicious of Alexei's formalism! The first step in proving Starobinsky's formalism is to act the retarded Green's function on the dimensionally regulated Heisenberg field equation (7) to derive the Yang-Feldman equation [22],

$$\phi(x) = \phi_0(x) - \int d^D x' \sqrt{-g(x')} \, i\theta(t-t') [\phi_0(x), \phi(x')] V'\Big(\phi(x')\Big) \,. \tag{15}$$

The free field $\phi_0(x)$ is,

$$\phi_0(t,\vec{x}) = \int_{H} \frac{d^{D-1}k}{(2\pi)^{D-1}} \left\{ \alpha(\vec{k})u(t,k)e^{i\vec{k}\cdot\vec{x}} + \alpha^{\dagger}(\vec{k})u^*(t,k)e^{-i\vec{k}\cdot\vec{x}} \right\}, \quad (16)$$

where α^{\dagger} and α are canonically normalized creation and annihilation operators and the free field mode function in Bunch-Davies vacuum [33–35] is,

$$u(t,k) \equiv i\sqrt{\frac{\pi}{4Ha^{D-1}}} H^{(1)}_{\frac{D-1}{2}}\left(\frac{k}{aH}\right).$$
(17)

Note that I have expressed the retarded Green's function as $i\theta(t - t')$ times the commutator of two free fields. Note also that the infrared limit of the mode function is,

$$k \ll Ha \qquad \Longrightarrow \qquad u(t,k) = \frac{\Gamma(\frac{D-1}{2})}{\sqrt{4\pi H}} \left(\frac{2H}{k}\right)^{\frac{D-1}{2}} \left\{ 1 + O\left(\frac{k^2}{a^2 H^2}\right) \right\}. \tag{18}$$

This is why the free field mode sum (16) has been infrared cut off at k = H to exclude modes which are in correlated Bunch-Davies vacuum at the beginning of inflation [17].

The Yang-Feldman equation (15) is exact; iterating it would produce the usual interaction picture expansion of the full field ϕ in powers of the free field ϕ_0 . Factors of $\ln[a(t)] = Ht$ in correlators derive from two sources:

- 1. The integral of $u(t, k)u^*(t', k)$ from the infrared cutoff at k = H to the point $k = H \times \min[a(t), a(t')]$ at which the infrared limit (18) breaks down; and
- 2. Vertex integrations over the retarded Green's function.

The first source is associated with a pair of free fields which combine to give a propagator; the second source is also associated with the pair of free fields in the commutator of the retarded Green's function. Now recall that reaching leading logarithm order requires that each extra pair of vertices must contribute one factor of $\ln[a]$ for each pair of fields. In the interaction picture this translates to each pair of free fields (including those in the retarded Green's function) contributing a factor of $\ln[a]$. Because these factors derive from the two sources given above, correlators will be unaffected, at leading logarithms order, if we cut off the mode sum in (16) at k = Ha(t) (at which point we can take D = 4), and also retain only the term (18) in the small k/aH expansion of the mode function.

Because the retarded Green's function involves a commutator, we must carry the expansion for these terms out to include the first imaginary part,

$$D = 4 \qquad \Longrightarrow \qquad u(t,k) = \frac{H}{\sqrt{2k^3}} \left\{ 1 + \frac{1}{2} \left(\frac{k}{aH}\right)^2 + \frac{i}{3} \left(\frac{k}{aH}\right)^3 + \dots \right\}.$$
(19)

Hence the retarded Green's function becomes,

$$G(x;x') = i\theta(t-t') \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot(\vec{x}-\vec{x}')} \left\{ u(t,k)u^*(t',k) - u^*(t,k)u(t',k) \right\}, \quad (20)$$

$$\longrightarrow \frac{\theta(t-t')}{3H} \times \left[\frac{1}{a^3(t')} - \frac{1}{a^3(t)}\right] \times \delta^3(\vec{x} - \vec{x}') \ . \tag{21}$$

Of course this allows us to perform the spatial integrations in the Yang-Feldman equation (15). A final simplification becomes apparent after multiplying (21) by the measure factor $\sqrt{-g(x')} = a^3(t') = e^{3Ht'}$ in expression (15). It follows that the factor of $1/a^3(t)$ in (21) can be neglected,

$$\int_0^t dt' a^3(t') \times \left[\frac{1}{a^3(t')} - \frac{1}{a^3(t)}\right] = t - \frac{1}{3H} \left[1 - e^{-3Ht}\right].$$
(22)

The various truncations produce a completely different field $\varphi(t, \vec{x})$ whose correlators nonetheless agree with those of $\phi(t, \vec{x})$ at leading logarithm order. The truncated Yang-Feldman equation obeyed by this field is,

$$\varphi(t,\vec{x}) = \varphi_0(t,\vec{x}) - \frac{1}{3H} \int_0^t dt' V\left(\varphi(t',\vec{x})\right) \,. \tag{23}$$

Here $\varphi_0(t, \vec{x})$ is precisely the infrared truncated free field expansion given in (9). Taking the time derivative of (23), and making some trivial rearrangements, gives Starobinsky's Langevin equation (8). Note that we have not only established its validity, we have clarified its precise relation to quantum field theory: *Starobinsky's formalism reproduces the leading logarithms of scalar potential models* (3).

4 Passive Fields

It is important to distinguish between *Active Fields*, which experience inflationary particle production (2), and *Passive Fields*, which do not. The prime examples of active fields are massless, minimally coupled scalars and gravitons; examples of passive fields include conformally coupled scalars, fermions and photons. We have seen that loops of active fields induce secular growth such as (5-6). Passive fields do not induces such factors, but they can mediate them, and they can modify the way that active fields interact.

Passive fields typically contribute nonzero constants to correlators. For example, one of several 1-loop electromagnetic contributions to the dimensionally regulated graviton self-energy on de Sitter background involves the coincidence limit of two field strengths [36],

$$\langle \Omega | F_{\mu\nu}(x) F_{\rho\sigma}(x) | \Omega \rangle = \frac{H^D}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(D-1)}{\Gamma(\frac{D}{2}+1)} \times \left[g_{\mu\rho} g_{\nu\sigma} - g_{\mu\sigma} g_{\nu\rho} \right].$$
(24)

Such contributions come from the full range of modes, from ultraviolet to infrared. They also involve the full passive field mode function, not just its infrared truncation. This is why no stochastic formulation of a passive field can be correct. For example, stochastically truncating electromagnetism would render the field strength an ultraviolet finite, Hermitian operator whose square must be positive. That is obviously inconsistent with the exact result (24) for the case of $\mu = \rho$ and $\nu = \sigma$.

The right thing to do with passive fields is integrate them out in the presence of a constant active field background. For passive fields coupled to an active scalar field setting this scalar to a constant induces a mass for the passive field, and integrating out the passive field results in a Coleman-Weinberg potential [37]. At this stage the active field has been reduced to a scalar potential model on which Starobinsky's formalism can be applied. Two theories for which this procedure has been applied are a real scalar which is Yukawa-coupled to a fermion [38], and a complex scalar which is coupled to electromagnetism [39].

The Lagrangian for Yukawa theory is,

$$\mathcal{L} = -\frac{1}{2}\partial_{\mu}\phi\partial_{\nu}\phi g^{\mu\nu}\sqrt{-g} - \frac{1}{2}\delta\xi\phi^{2}R\sqrt{-g} - V(\phi)\sqrt{-g} - f\phi\overline{\psi}\psi\sqrt{-g} + \overline{\psi}e^{\mu}_{\ a}\gamma^{a}\left(i\partial_{\mu} - \frac{1}{2}A_{\mu bc}J^{bc}\right)\psi\sqrt{-g} .$$
(25)

The usual notation applies: γ^a denotes the gamma matrices, the vierbein is

 $e^{\mu}_{a}(x)$, the spin connection is $A_{\mu bc}(x)$, and the spin generator is J^{bc} ,

$$g^{\mu\nu} = e^{\mu}_{\ a} e^{\nu}_{\ b} \eta^{ab} \quad , \quad A_{\mu bc} \equiv e^{\nu}_{\ b} (e_{\nu c,\mu} - \Gamma^{\rho}_{\ \mu\nu} e_{\rho c}) \quad , \quad J^{bc} \equiv \frac{i}{4} [\gamma^b, \gamma^c] \; .$$
 (26)

For constant ϕ the last term on the first line of (25) can be recognized as a fermion mass of $m = f\phi$. The massive fermion propagator on de Sitter $i[_iS_j](x;x')$ was derived by Candelas and Raine [40]. Integrating out the fermion out of the scalar field equation (with suitable renormalizations) gives,

$$\partial_{\mu} (\sqrt{-g} g^{\mu\nu} \partial_{\nu} \phi) - \delta \xi \phi R \sqrt{-g} - V'(\phi) \sqrt{-g} - f \overline{\psi} \psi \sqrt{-g} \longrightarrow \partial_{\mu} (\sqrt{-g} g^{\mu\nu} \partial_{\nu} \phi) - [\delta \xi \phi R - V'(\phi)] \sqrt{-g} - f i [_{i}S_{i}](x; x) \sqrt{-g} , \quad (27) \longrightarrow \partial_{\mu} (\sqrt{-g} g^{\mu\nu} \partial_{\nu} \phi) - [\alpha H^{2} \phi + \beta \phi^{3}] \sqrt{-g} + \frac{f H^{3}}{4\pi^{2}} \frac{f \phi}{H} (1 + \frac{f^{2} \phi^{2}}{H^{2}}) \Big\{ \psi (1 + \frac{i f \phi}{H}) + \psi (1 - \frac{i f \phi}{H}) \Big\} \sqrt{-g} . \quad (28)$$

Here $\psi(x) \equiv \frac{d}{dx} \ln[\Gamma(x)]$ is the digamma function and the constants $\alpha \sim f^2$ and $\beta \sim f^4$ are arbitrary.

The Lagrangian of scalar quantum electrodynamics (SQED) is,

$$\mathcal{L} = -(D_{\mu}\phi)^* D_{\nu}\phi g^{\mu\nu}\sqrt{-g} - \delta\xi\phi^*\phi R\sqrt{-g} - V(\phi^*\phi)\sqrt{-g} - \frac{1}{4}F_{\mu\nu}F_{\rho\sigma}g^{\mu\rho}g^{\nu\sigma}\sqrt{-g} .$$
(29)

Here $D_{\mu} \equiv \partial_{\mu} + ieA_{\mu}$ is the covariant derivative operator. Setting the scalar equal to a constant endows the photon with a mass $m^2 = e^2 \phi^* \phi$. The Lorentz gauge massive photon propagator on de Sitter background $i[_{\mu}\Delta_{\nu}](x;x')$ was derived in [41]. Using it to integrate the photon out of the scalar field equation (with suitable renormalization) gives ,¹

$$D_{\mu}(\sqrt{-g} g^{\mu\nu} D_{\nu} \phi) - \delta \xi \phi R \sqrt{-g} - \phi V'(\phi^{*} \phi) \sqrt{-g} \longrightarrow \partial_{\mu}(\sqrt{-g} g^{\mu\nu} \partial_{\nu} \phi) - \phi [\delta \xi R - V'] \sqrt{-g} - e^{2} \phi i [_{\mu} \Delta_{\nu}](x; x) g^{\mu\nu} \sqrt{-g} , \quad (30) \longrightarrow \partial_{\mu}(\sqrt{-g} g^{\mu\nu} \partial_{\nu} \phi) - \phi [\overline{\alpha} + \overline{\beta} \phi^{*} \phi] \sqrt{-g} - \frac{3e^{2} H^{2} \phi}{8\pi^{2}} (1 + \frac{e^{2} \phi^{*} \phi}{H^{2}}) \times \left\{ \psi \left(\frac{3}{2} + \frac{1}{2} [1 - \frac{8e^{2} \phi^{*} \phi}{H^{2}}]^{\frac{1}{2}} \right) + \psi \left(\frac{3}{2} - \frac{1}{2} [1 - \frac{8e^{2} \phi^{*} \phi}{H^{2}}]^{\frac{1}{2}} \right) \right\} \sqrt{-g} . \quad (31)$$

The constants $\overline{\alpha} \sim e^2$ and $\overline{\beta} \sim e^4$ are again arbitrary.

The stochastic formulations of the scalar potential models (28) and (31) agree with explicit dimensionally regulated and fully renormalized computations at 1-loop and 2-loop orders [38, 39, 43, 44]. "Leading logarithm" in

¹The result was first obtained by Allen as a mode sum [42].

these two theories means that each additional factor of f^2 in Yukawa, or of e^2 in SQED, brings an extra factor of $\ln[a(t)]$. Note that ultraviolet regularization and renormalization are required to derive the Coleman-Weinberg potentials, however, the resulting scalar potential model is afterwards UV finite at leading logarithm order. The need for renormalization is reflected in the arbitrary finite parts associated with the constants α and β for (28), and $\overline{\alpha}$ and $\overline{\beta}$ for (31).

Finally, it is worth commenting on signs. The Yukawa effective potential (28) is negative, so this system never reaches a stable equilibrium; the universe instead decays in a Big Rip singularity [38]. In contrast, the effective potential for SQED is bounded below, which means that the system approaches a stable equilibrium. However, it turns out that the vacuum energy is slightly reduced [39], for essentially the same reason that a slab of dielectric is pulled into a charged, parallel plate capacitor. With the quartic potential result (14) this means that scalar models span the range of possibilities:

- The system may or may not approach a stable equilibrium; and
- The vacuum energy may be increased or decreased.

5 Differentiated Active Fields

Because the secular growth factors of $\ln[a(t)]$ derive from logarithmic integrations, the presence of a single derivative suppresses them. The status of differentiated active fields is very similar to that of passive fields: although they induce no secular loop corrections, they make important contributions of order one which come as much from the ultraviolet as from the infrared, and involve the full mode function. Hence it is also wrong to infrared truncate differentiated active fields. For example, in dimensional regularization the coincidence limit of the doubly differentiated, massless, minimally coupled scalar propagator is [14, 31],

$$\lim_{x' \to x} \partial_{\mu} \partial'_{\nu} i \Delta(x; x') = \langle \Omega | \partial_{\mu} \phi_0(x) \partial_{\nu} \phi(x) | \Omega \rangle = -\frac{H^D}{2(4\pi)^{\frac{D}{2}}} \frac{\Gamma(D)}{\Gamma(\frac{D}{2}+1)} \times g_{\mu\nu}(x) .$$
(32)

Were we to replace $\phi_0(x)$ by its infrared truncation $\varphi_0(x)$ the expectation value would be positive for $\mu = \nu$ because it is the square of a finite, Hermitian operator. However, the exact result (32) is *negative* for spatial $\mu = \nu$.

Rather than stochastically simplifying differentiated active fields, the correct procedure is to integrate them out of the field equations in the presence of a constant active field background. One then applies Starobinsky's formalism. This is also the same as for passive fields. The key difference is that imposing a constant active field background does not induce a mass but rather a field strength — for nonlinear sigma models — or a modification in the background geometry — for differentiated actives coupled to gravity.

Another important point is that none of these theories is renormalizable. However, all theories can be renormalized by subtracting BPHZ (Bogoliubov and Parasiuk [45], Hepp [46] and Zimmermann [47,48]) counterterms, orderby-order in the loop expansion. When this is done, the leading secular growth factors are uniquely determined by low energy effective field theory in the sense of Donoghue [49–51].

5.1 Nonlinear Sigma Models

Quantum gravity is *hard*. The weeks it requires to perform a 2-loop scalar computation on de Sitter background [14, 31, 52] become months to perform a 1-loop graviton computation [53–55]. The fundamental $h\partial h\partial h$ interaction of quantum gravity also occurs in nonlinear sigma models such as,

$$\mathcal{L}_1 = -\frac{1}{2} (1 + \frac{\lambda}{2} \phi)^2 \partial_\mu \phi \partial_\nu \phi g^{\mu\nu} \sqrt{-g} , \qquad (33)$$

$$\mathcal{L}_2 = -\frac{1}{2}\partial_\mu A \partial_\nu A g^{\mu\nu} \sqrt{-g} - \frac{1}{2}(1 + \frac{\lambda}{2}A)^2 \partial_\mu B \partial_\nu B g^{\mu\nu} \sqrt{-g} .$$
(34)

For this reason nonlinear sigma models have long been employed as a simple venue to sort out the complexities of derivative interactions, without the tensor indices, the gauge issue and the vast proliferation of interactions which make quantum gravity so difficult [22, 56–59].

The key to integrating out differentiated fields in (33-34) is the observation that working in a constant ϕ or A background just changes the field strengths of the ϕ and B propagators from that of the massless, minimally coupled scalar $i\Delta(x; x')$,

$$\langle \Omega | T[\phi(x)\phi(x')] | \Omega \rangle_{\phi_0} = \frac{i\Delta(x;x')}{(1+\frac{\lambda}{2}\phi_0)^2} \quad , \quad \langle \Omega | T[B(x)B(x')] | \Omega \rangle_{A_0} = \frac{i\Delta(x;x')}{(1+\frac{\lambda}{2}A_0)^2} \; .$$

$$(35)$$

Integrating out the $\partial B \partial B$ term (for constant A_0) from the A field equation (and taking D = 4) gives,

$$\partial_{\mu} \left[\sqrt{-g} \, g^{\mu\nu} \partial_{\nu} A \right] - \frac{\lambda}{2} (1 + \frac{\lambda}{2} A) \partial_{\mu} B \partial_{\nu} B g^{\mu\nu} \sqrt{-g} \longrightarrow \partial_{\mu} \left[\sqrt{-g} \, g^{\mu\nu} \partial_{\nu} A \right] - \frac{\lambda}{2} (1 + \frac{\lambda}{2} A_0) \times -\frac{\frac{3H^4}{8\pi^2} \sqrt{-g}}{(1 + \frac{\lambda}{2} A_0)^2} \,. \tag{36}$$

Applying Starobinsky's formalism to this scalar potential model gives a Langevin equation for the stochastic field $\mathcal{A}(t, \vec{x})$,

$$3H(\dot{\mathcal{A}} - \dot{\mathcal{A}}_0) = \frac{3\lambda H^4}{16\pi^2} \frac{1}{1 + \frac{\lambda}{2}\mathcal{A}} \,. \tag{37}$$

The effective potential for A induced in (36) is $V(A) = -\frac{3H^4}{8\pi^2} \ln |1 + \frac{\lambda}{2}A|$. This is unbounded below so no static limit is approached. However, the time evolution of the background is sedate. It consists of a "classical" contribution, obtained by ignoring the jitter \mathcal{A}_0 , plus a series in powers of \mathcal{A}_0 ,

$$\mathcal{A}(t,\vec{x}) = -\frac{2}{\lambda} + \frac{2}{\lambda} \left[1 + \frac{\lambda^2 H^2 \ln[a(t)]}{16\pi^2} \right]^{\frac{1}{2}} + \mathcal{A}_0(t,\vec{x}) - \frac{\lambda^2 H^3}{32\pi^2} \int_0^t dt' \mathcal{A}_0(t',\vec{x}) + \frac{\lambda^3 H^3}{64\pi^2} \int_0^t dt' \mathcal{A}_0^2(t',\vec{x}) + O(\lambda^4) \ . \ (38)$$

Correlators of \mathcal{A} agree, at leading logarithm order, with dimensionally regulated and fully BPHZ renormalized 1-loop and 2-loop computations of those of \mathcal{A} , for example [59,60],

$$\langle \Omega | \mathcal{A}(t, \vec{x}) | \Omega \rangle = \frac{\lambda H^2 \ln[a(t)]}{16\pi^2} + \frac{\lambda^3 H^4 \ln^2[a(t)]}{2^{10}\pi^4} + O(\lambda^5) .$$
(39)

Note that the basic time dependence of (39) derives from the classical part on the first line of (38), and is understandable in terms of the field rolling down its potential. The stochastic jitter accelerates the roll, essentially because it is easier to fluctuate down a potential than up.

The stochastic formulation of the single field model (33) is crucial for gravity because, in both cases, it is the *same* field whose derivatives are integrated out and which experiences stochastic fluctuations. It is straightforward to integrate out differentiated scalars from the ϕ field equation,

$$(1 + \frac{\lambda}{2}\phi)\partial_{\mu}\left[(1 + \frac{\lambda}{2}\phi)\sqrt{-g}\,g^{\mu\nu}\partial_{\nu}\phi\right] \longrightarrow (1 + \frac{\lambda}{2}\phi_{0})\partial_{\mu}\left[\frac{\lambda}{4}\sqrt{-g}\,g^{\mu\nu}\partial_{\nu}\langle\Omega|\phi^{2}|\Omega\rangle_{\phi_{0}}\right], \quad (40)$$

$$\rightarrow \left(1 + \frac{\lambda}{2}\phi_0\right) \times -\frac{3\lambda H^4}{16\pi^2} \frac{\sqrt{-g}}{(1 + \frac{\lambda}{2}\phi_0)^2} \ . \tag{41}$$

This corresponds to another logarithmic potential $V(\phi) = \frac{3H^4}{8\pi^2} \ln |1 + \frac{\lambda}{2}\phi|$. The key point is that we add it to the stochastic simplification of the same derivative terms from which it emerged,

$$3H(1+\frac{\lambda}{2}\varphi)^{2}[\dot{\varphi}-\dot{\varphi}_{0}] = -\frac{3\lambda H^{4}}{16\pi^{2}}\frac{1}{1+\frac{\lambda}{2}\varphi}.$$
(42)

As with the double field model (38), the solution of this Langevin equation consists of a classical part plus a series in powers of φ_0 ,

$$\varphi(t,\vec{x}) = -\frac{2}{\lambda} + \frac{2}{\lambda} \left[1 - \frac{\lambda^2 H^2 \ln[a(t)]}{8\pi^2} \right]^{\frac{1}{4}} + \varphi_0(t,\vec{x}) + \frac{3\lambda^2 H^3}{32\pi^2} \int_0^t dt' \varphi_0(t',\vec{x}) - \frac{3\lambda^3 H^3}{32\pi^2} \int_0^t dt' \varphi_0^2(t',\vec{x}) + O(\lambda^4).$$
(43)

Also like the double field model (39), correlators of φ agree, at leading logarithm order, with explicit 1-loop and 2-loop computations performed using dimensional regularization and BPHZ renormalization [59],

$$\langle \Omega | \varphi(t, \vec{x}) | \Omega \rangle = -\frac{\lambda H^2 \ln[a(t)]}{16\pi^2} - \frac{15\lambda^3 H^4 \ln^2[a(t)]}{2^{10}\pi^4} + O(\lambda^5) .$$
 (44)

5.2 Scalar Corrections to Gravity

Most matter fields are not active, but the massless, minimally coupled scalar is an exception,

$$\mathcal{L}_{\rm MMC} = -\frac{1}{2} \partial_{\mu} \phi \partial_{\nu} \phi g^{\mu\nu} \sqrt{-g} .$$
 (45)

To integrate differentiated scalar out of the gravitational field equation,

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + (\frac{D-2}{2})\Lambda g_{\mu\nu} = 8\pi G \left\{ \partial_{\mu}\phi\partial_{\nu}\phi - \frac{1}{2}g_{\mu\nu}g^{\rho\sigma}\partial_{\rho}\phi\partial_{\sigma}\phi \right\}, \quad (46)$$

we must define what it means to be in a "constant graviton background". If the temporal coordinate is changed from co-moving time t to conformal time η , such that $d\eta = dt/a(t)$, the background geometry of cosmology (1) takes the form $a^2\eta_{\mu\nu}$. We define the graviton field $h_{\mu\nu}(x)$ by conformally transforming the full metric,

$$g_{\mu\nu} \equiv a^2 \widetilde{g}_{\mu\nu} \equiv a^2 (\eta_{\mu\nu} + \kappa h_{\mu\nu}) \qquad , \qquad \kappa^2 \equiv 16\pi G \;. \tag{47}$$

The transverse-traceless and purely spatial components of $h_{\mu\nu}$ obey the same equation of motion as that of the massless, minimally coupled scalar [29]. Consequently, it is $h_{\mu\nu}$ which experiences inflationary particle production (2), and $h_{\mu\nu}$ which should be held constant when we integrate out differentiated scalars from equation (46).

Constant $h_{\mu\nu}$ means that $\tilde{g}_{\mu\nu}$ is also constant. It turns out that setting $g_{\mu\nu} = a^2 \tilde{g}_{\mu\nu}$, with constant $\tilde{g}_{\mu\nu}$ amounts to being in de Sitter with a different Hubble parameter [61, 62],

$$\partial_{\rho} \widetilde{g}_{\mu\nu} = 0 \qquad \Longrightarrow \qquad H^2 \longrightarrow -\widetilde{g}^{00} H^2 \equiv \widetilde{H}^2 .$$
 (48)

This makes it simple to integrate out the scalars from equation (46),

$$\frac{\kappa^2}{2} \left\{ \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} g^{\rho\sigma} \partial_\rho \phi \partial_\sigma \phi \right\} \longrightarrow \frac{\kappa^2}{2} \left(\delta^\rho_{\ \mu} \delta^\sigma_{\ \nu} - \frac{1}{2} g_{\mu\nu} g^{\rho\sigma} \right) \times \langle \Omega | \partial_\rho \phi \partial_\sigma \phi | \Omega \rangle_h , (49)$$
$$\longrightarrow \frac{\kappa^2}{2} \left(\delta^\rho_{\ \mu} \delta^\sigma_{\ \nu} - \frac{1}{2} g_{\mu\nu} g^{\rho\sigma} \right) \times -\frac{3\tilde{H}^4}{32\pi^2} g_{\rho\sigma} = \frac{3\kappa^2 \tilde{H}^4}{64\pi^2} g_{\mu\nu} . (50)$$

Expression (50) corresponds to a finite renormalization of the cosmological constant [63],

$$\delta\Lambda = -\frac{3\kappa^2 \tilde{H}^4}{64\pi^2} \,. \tag{51}$$

The need for this renormalization to make the graviton self-energy conserved had been discovered previously through explicit computations [55, 64].

5.3 Pure Gravity

There is no more postponing the complexities of gravity. The vast proliferation of interactions is apparent from expressing the invariant Lagrangian in terms of the graviton field $h_{\mu\nu}$ and the conformally transformed metric $\tilde{g}_{\mu\nu} \equiv \eta_{\mu\nu} + \kappa h_{\mu\nu}$ [65],

$$\mathcal{L}_{\rm inv} = a^{D-2} \sqrt{-\widetilde{g}} \, \widetilde{g}^{\alpha\beta} \widetilde{g}^{\rho\sigma} \widetilde{g}^{\mu\nu} \left\{ \frac{1}{2} h_{\alpha\rho,\mu} h_{\nu\sigma,\beta} - \frac{1}{2} h_{\alpha\beta,\rho} h_{\sigma\mu,\nu} + \frac{1}{4} h_{\alpha\beta,\rho} h_{\mu\nu,\sigma} - \frac{1}{4} h_{\alpha\rho,\mu} h_{\beta\sigma,\nu} \right\} + (\frac{D-2}{2}) a^{D-1} H \sqrt{-\widetilde{g}} \, \widetilde{g}^{\rho\sigma} \widetilde{g}^{\mu\nu} h_{\rho\sigma,\mu} h_{\nu0} \, . \tag{52}$$

The simplest gauge is based on adding the gauge fixing term [65, 66],

$$\mathcal{L}_{\rm GF} = -\frac{1}{2}a^{D-2}\eta^{\mu\nu}F_{\mu}F_{\nu} \ , \ F_{\mu} = \eta^{\rho\sigma}[h_{\mu\rho,\sigma} - \frac{1}{2}h_{\rho\sigma,\mu} + (D-2)aHh_{\mu\rho}\delta^{0}{}_{\sigma}] \ . \ (53)$$

The associated ghost Lagrangian is,

$$\mathcal{L}_{\rm gh} = -a^{D-2} \eta^{\mu\nu} \overline{c}_{\mu} \delta F_{\nu} , \qquad (54)$$

where δF_{ν} represents the infinitesimal transformation $x^{\mu} \to x^{\mu} + \kappa \epsilon^{\mu}$ of F_{ν} with the transformation parameter replaced by the ghost field.

The gauge (53-54) results in ghost and graviton propagators whose tensor structure consists of constants formed from $\eta_{\mu\nu}$ and $\delta^0_{\ \mu}$,

$$i[_{\mu\nu}\Delta_{\rho\sigma}](x;x') = \sum_{I=A,B,C} [_{\mu\nu}T^{I}_{\rho\sigma}] \times i\Delta_{I}(x;x') , \qquad (55)$$

$$i[_{\mu}\Delta_{\rho}](x;x') = \overline{\eta}_{\mu\nu} \times i\Delta_A(x;x') - \delta^0_{\ \mu}\delta^0_{\ \nu} \times i\Delta_B(x;x') .$$
(56)

Here and henceforth the purely spatial Minkowski metric is $\overline{\eta}_{\mu\nu} \equiv \eta_{\mu\nu} + \delta^0_{\ \mu} \delta^0_{\nu}$, and the other tensor factors are,

$$[_{\mu\nu}T^{A}_{\rho\sigma}] = 2\overline{\eta}_{\mu(\rho}\overline{\eta}_{\sigma)\nu} - \frac{2}{D-3}\overline{\eta}_{\mu\nu}\overline{\eta}_{\rho\sigma} \qquad , \qquad [_{\mu\nu}T^{B}_{\rho\sigma}] = -4\delta^{0}_{\ (\mu}\overline{\eta}_{\nu)(\rho}\delta^{0}_{\ \sigma)} \ , \tag{57}$$

$$[_{\mu\nu}T^{C}_{\rho\sigma}] = \frac{2}{(D-2)(D-3)}[(D-3)\delta^{0}_{\ \mu}\delta^{0}_{\ \nu} + \overline{\eta}_{\mu\nu}][(D-3)\delta^{0}_{\ \rho}\delta^{0}_{\ \sigma} + \overline{\eta}_{\rho\sigma}] .$$
(58)

Parenthesized indices are symmetrized. The three propagators $i\Delta_I(x; x')$ are those of a minimally coupled scalar with masses,

$$m_A^2 = 0$$
 , $m_B^2 = (D-2)H^2$, $m_C^2 = 2(D-3)H^2$. (59)

Because graviton propagators possess indices, working in a constant $\tilde{g}_{\mu\nu}$ background requires more than just replacing all the factors of H^2 by \tilde{H}^2 , as per equation (48), in the scalar propagators $i\Delta_I(x;x')$. One must also modify the gauge fixing function to [67],

$$\widetilde{\mathcal{L}}_{\rm GF} = -\frac{1}{2}a^{D-2}\sqrt{-\widetilde{g}}\,\widetilde{g}^{\mu\nu}\widetilde{F}_{\mu}\widetilde{F}_{\nu} \quad , \quad \widetilde{F}_{\mu} = \widetilde{g}^{\rho\sigma}[h_{\mu\rho,\sigma} - \frac{1}{2}h_{\rho\sigma,\mu} + (D-2)aHh_{\mu\rho}\delta^{0}_{\sigma}] \quad .$$

$$\tag{60}$$

The ghost Lagrangian (54) suffers a similar change [67],

$$\widetilde{\mathcal{L}}_{\rm gh} = -a^{D-2}\sqrt{-\widetilde{g}}\,\widetilde{g}^{\mu\nu}\overline{c}_{\mu}\delta\widetilde{F}_{\nu} \ . \tag{61}$$

The tensor structures of the ghost and graviton propagators (55-58) are constructed from the Minkowski metric $\eta_{\mu\nu}$ and the timelike 4-vector δ^0_{μ} . Generalizing to a constant $\tilde{g}_{\mu\nu}$ background amounts to stating what becomes of these two tensors. The most straightforward is $\eta_{\mu\nu}$: it generalizes to $\tilde{g}_{\mu\nu}$. Explaining what happens to $\delta^0_{\ \mu}$ is facilitated by the ADM decomposition [68],

$$\widetilde{g}_{\mu\nu}dx^{\mu}dx^{\nu} = -N^2d\eta^2 + \gamma_{ij}(dx^i - N^i d\eta)(dx^j - N^j d\eta) .$$
(62)

This suggests the generalization [67],

$$\delta^0_{\ \mu} \longrightarrow u_\mu \equiv -N \delta^0_{\ \mu} , \qquad (63)$$

$$\overline{\eta}_{\mu\nu} \longrightarrow \overline{\gamma}_{\mu\nu} \equiv \widetilde{g}_{\mu\nu} + u_{\mu}u_{\nu} .$$
(64)

So the graviton and ghost propagators in a constant $\tilde{g}_{\mu\nu}$ background are,

$$i[_{\mu\nu}\widetilde{\Delta}_{\rho\sigma}](x;x') = \sum_{I=A,B,C} [_{\mu\nu}\widetilde{T}^{I}_{\rho\sigma}] \times i\widetilde{\Delta}_{I}(x;x') , \qquad (65)$$

$$i[_{\mu}\widetilde{\Delta}_{\rho}](x;x') = \overline{\gamma}_{\mu\nu} \times i\widetilde{\Delta}_{A}(x;x') - u_{\mu}u_{\nu} \times i\widetilde{\Delta}_{B}(x;x') .$$
(66)

The scalar propagators $i\widetilde{\Delta}_I(x;x')$ are obtained by making the replacement (48) in $i\Delta_I(x;x')$, and the tensor factors $[_{\mu\nu}\widetilde{T}^I_{\rho\sigma}]$ are obtained by making the replacements (63-64) in (57-58).

The sum of the invariant Lagrangian (52) and the new gauge fixing term (60) can be grouped into six terms [67],

$$\mathcal{L}_{1+2+3} = a^{D-2} \sqrt{-\widetilde{g}} \, \widetilde{g}^{\alpha\beta} [-\frac{1}{4} \widetilde{g}^{\gamma\rho} \widetilde{g}^{\delta\sigma} + \frac{1}{8} \widetilde{g}^{\gamma\delta} \widetilde{g}^{\rho\sigma}] h_{\gamma\delta,\alpha} h_{\rho\sigma,\beta} + (\frac{D-2}{2}) a^D \sqrt{-\widetilde{g}} \, \widetilde{g}^{\gamma\alpha} \widetilde{g}^{\beta\rho} \widetilde{g}^{\sigma\delta} h_{\alpha\beta} h_{\rho\sigma} \widetilde{H}^2 u_{\gamma} u_{\delta} \,, \quad (67)$$

$$\mathcal{L}_{4+5} = \sqrt{-\widetilde{g}} \, \widetilde{g}^{\alpha\beta} \widetilde{g}^{\gamma\delta} \widetilde{g}^{\rho\sigma} \partial_{\beta} \Big[-\frac{1}{2} \partial_{\sigma} (a^{D-2} h_{\gamma\rho} h_{\delta\alpha}) + a^{D-2} h_{\gamma\rho} h_{\delta\alpha,\sigma} \Big] \,, \tag{68}$$

$$\mathcal{L}_{6} = \left(\frac{D-2}{2}\right) \kappa H a^{D-1} \sqrt{-\widetilde{g}} \, \widetilde{g}^{\alpha\beta} \widetilde{g}^{\gamma\delta} \widetilde{g}^{\rho\sigma} h_{\rho\sigma,\gamma} h_{\delta\alpha} h_{\beta0} \, . \tag{69}$$

Completing the stochastic simplification of pure gravity entails three steps:

- 1. Find the contribution to the equation of motion from (61) and (67-69);
- 2. Integrate out the ghost fields and the differentiated graviton fields from each contribution; and
- 3. Derive the Langevin kinetic operator from each contribution.

The final step involves distinguishing between the spatial components of the graviton field, which are active, and the other components, which are passive, constrained fields whose dynamics is driven by the active components [69],

$$\kappa h_{\mu\nu} = A_{\mu\nu} + 2u_{(\mu}B_{\nu)} + [u_{\mu}u_{\nu} + \overline{\gamma}_{\mu\nu}]C \qquad , \qquad u^{\rho}A_{\rho\sigma} = 0 = u^{\rho}B_{\rho} \ . \tag{70}$$

Note that the active components $A_{\mu\nu}$ have a stochastic jitter $a_{\mu\nu}$ whereas the passive, constrained fields B_{μ} and C do not.

The first two steps have been implemented [67], and the third is in progress [69]. The complete result for (67) is,

$$\frac{\kappa a^{-4}}{\sqrt{-\tilde{g}}} \times \frac{\delta S_{1+2+3}}{\delta h_{\mu\nu}} \longrightarrow \frac{\kappa^2 \tilde{H}^4}{8\pi^2} [13\tilde{g}^{\mu\nu} + 6u^{\mu}u^{\nu}] - [\tilde{g}^{\mu\rho}\tilde{g}^{\nu\sigma} - \frac{1}{2}\tilde{g}^{\mu\nu}\tilde{g}^{\rho\sigma}]a^{-1}\tilde{H}u^{\alpha}\partial_{\alpha}(A_{\rho\sigma} - a_{\rho\sigma})
+ u^{(\mu}\tilde{g}^{\nu)\rho}\tilde{D}_B B_{\rho} + u^{\mu}u^{\nu}\tilde{D}_C C + \tilde{H}^2 \Big\{ \frac{1}{2}\tilde{g}^{\mu\nu}(B^2 - C^2)
+ 2u^{(\mu}A^{\nu)\rho}B_{\rho} - B^{\mu}B^{\nu} - 2u^{(\mu}B^{\nu)}C + u^{\mu}u^{\nu}(2B^2 - 3C^2) \Big\}.$$
(71)

Although the two operators agree for D = 4, it is useful to retain dimensional regularization in order to see the difference between \widetilde{D}_B and \widetilde{D}_C ,

$$\widetilde{D}_B = \partial_\alpha [a^{D-2}\sqrt{-\widetilde{g}}\,\widetilde{g}^{\alpha\beta}\partial_\beta] - (D-2)\widetilde{H}^2 a^D\sqrt{-\widetilde{g}}\,, \tag{72}$$

$$\widetilde{D}_C = \partial_\alpha [a^{D-2} \sqrt{-\widetilde{g}} \, \widetilde{g}^{\alpha\beta} \partial_\beta] - 2(D-3) \widetilde{H}^2 a^D \sqrt{-\widetilde{g}} \,.$$
(73)

In D = 4 both operators degenerate to the conformal d'Alembertian.

Whereas nonlinear sigma models and scalar corrections to gravity had seen extensive dimensionally regulated and fully renormalized computations prior to their stochastic realizations, the stochastic formulation of pure gravity is far ahead of the explicit calculations with which it might be compared. The 1PI (one-particle-irreducible) 1-graviton function has been computed in the original gauge (53) [70], but it must be re-done in the new gauge (60) to check the stochastic equation. Quantum gravity presumably requires the same sort of finite renormalization of the cosmological constant that was found for scalar corrections to gravity [64]. Also in the old gauge (53), the 1PI 2-point function (the "graviton self-energy") was computed, away from coincidence and without using dimensional regularization or renormalizing [53]. This needs to re-done in the new gauge (60) using dimensional regularization and BPHZ renormalization. A technique was devised for extending the old calculation to a fully renormalized result [71]. This extension was then employed to compute 1-loop corrections to the graviton mode function [72] and to the gravitational response to a point mass [73]. Both of these things need to re-done with a exact computation in the new gauge.

6 The Other Source of Large Logarithms

The first large logarithm from a loop of gravitons was discovered 20 years ago [23, 24]. It was soon shown not to follow from a naive realization of the stochastic formalism [74]. In the meantime a number of similar effects were found from graviton loop corrections to matter [75–81] and in graviton corrections to gravity [71–73]. The same kinds of derivative interactions in matter corrections to gravity also induce large logarithms [36, 55, 82, 83]. Explaining these effects, and devising techniques to resum them, has been a long and confusing struggle for two reasons:

- 1. The correct stochastic formulation involves integrating differentiated active fields out, which entails new sorts of effective potentials based on background-dependent field strengths and Hubble parameters; and
- 2. There is another way large logarithms can be induced which cannot be captured stochastically.

Section 5 concerned the first problem, this section concerns the second.

The second way of inducing large logarithms is the incomplete cancellation between primitive divergences and counterterms. When using dimensional regularization it turns out that the factors of a^{D-2} from vertices are canceled by inverse factors from propagators. For example, the $i\Delta_I(x; x')$ propagators which appear in the graviton and ghost propagators (55-56) have the same leading singularity,

$$i\Delta_I(x;x') = \frac{1}{4\pi^{\frac{D}{2}}} \frac{1}{[aa'(x-x')^2]^{\frac{D}{2}-1}} + \dots ,$$
 (74)

where $(x - x')^2 \equiv (x - x')^{\mu} (x - x')^{\nu} \eta_{\mu\nu}$ is the conformal coordinate interval. Higher terms in the propagator can give rise to additional, integer powers of (aa'), but the *D*-dependence will always be $(aa')^{-\frac{D}{2}}$. On the other hand, counterterms inherit a factor of a^D from the $\sqrt{-g}$ measure. It follows that there is an incomplete cancellation between primitive divergences and counterterms,

$$\frac{(2H)^{D-4}}{D-4} - \frac{(a\mu)^{D-4}}{D-4} = -\ln(\frac{a\mu}{2H}) + O(D-4) .$$
(75)

Nonlinear sigma models exhibit both stochastic logarithms such as (39) and (44), and also renormalization-induced logarithms (74). The large logarithms of most graviton loop corrections to matter fields [75–78,81], and all known matter loop corrections to gravity [36,54,55,82,83], are induced by renormalization. The case of pure gravity [71–73] has yet to be decided.

Renormalization-induced logarithms occur even for passive fields which do not experience inflationary particle production (2), so they cannot be explained stochastically. On the other hand, the close association in expression (75) between the dimensional regularization scale μ and the scale factor a(t)makes these logarithms amenable to a modified renormalization group treatment. I will explain the technique in the context of matter loop corrections to gravity [36, 54, 55, 82, 83] for which a unified treatment is possible.

What has been done is to compute matter loop contributions to the graviton self-energy $-i[^{\mu\nu}\Sigma^{\rho\sigma}](x;x')$ on de Sitter background [36, 54, 55, 82, 83]. This can be used to quantum-correct the linearized Einstein equation,

$$\mathcal{L}^{\mu\nu\rho\sigma}\kappa h_{\rho\sigma} - \int d^4x' \, [^{\mu\nu}\Sigma^{\rho\sigma}](x;x')\kappa h_{\rho\sigma}(x') = \frac{\kappa^2}{2}T^{\mu\nu}(x) \,. \tag{76}$$

Here $\mathcal{L}^{\mu\nu\sigma\rho}$ is the Lichnerowicz operator on de Sitter background and $T^{\mu\nu}$ is the "graviton stress tensor," defined as minus the variation of the matter action with respect to $h_{\mu\nu}$. With $T^{\mu\nu} = 0$, one is solving for 1-loop corrections

to gravitational radiation. There are no corrections in flat space background, but a loop of massless, minimally coupled scalars gives rise to a secular enhancement of the Weyl tensor [55],

$$C_{0i0j}(t,\vec{x}) = C_{0i0j}^{\text{tree}}(t,\vec{x}) \left\{ 1 - \frac{3\kappa^2 H^2}{160\pi^2} \ln[a(t)] + \dots \right\}.$$
 (77)

Setting $T^{\mu\nu}(t, \vec{x}) = -M\delta^{\mu}_{0}\delta^{\nu}_{0}a(t)\delta^{3}(\vec{x})$ gives the response to a point mass, which can be parameterized in terms of two scalar potentials,

$$ds^{2} = -[1 - 2\Psi(t, r)]dt^{2} + a^{2}(t)[1 - 2\Phi(t, r)]d\vec{x} \cdot d\vec{x} .$$
(78)

For a loop of massless, minimally coupled scalars one finds [55,84],

$$\Psi(t,r) = \frac{GM}{a(t)r} \left\{ 1 + \frac{\kappa^2}{320\pi^2 a^2(t)r^2} - \frac{3\kappa^2 H^2}{160\pi^2} \ln[a(t)] + \dots \right\},$$
(79)

$$\Psi(t,r) + \Phi(t,r) = \frac{GM}{a(t)r} \left\{ 0 + \frac{\kappa^2}{240\pi^2 a^2(t)r^2} + \frac{3\kappa^2 H^2}{160\pi^2} + \dots \right\}.$$
 (80)

The fractional κ^2/a^2r^2 correction in (79) is the de Sitter descendant of an old flat space effect [85], whereas the $\kappa^2 H^2$ corrections are new. Corrections to the Weyl tensor and to the two potentials have also been computed for conformally invariant matter theories. For a massless, conformally coupled scalar the results are [83],

$$C_{0i0j}(t,\vec{x}) = C_{0i0j}^{\text{tree}}(t,\vec{x}) \left\{ 1 + \frac{\kappa^2 H^2}{480\pi^2} \ln[a(t)] + \dots \right\},$$
(81)

$$\Psi(t,r) = \frac{GM}{a(t)r} \left\{ 1 + \frac{\kappa^2}{720\pi^2 a^2(t)r^2} + \frac{\kappa^2 H^2}{480\pi^2} \ln[a(t)] + \dots \right\},$$
(82)

$$\Psi(t,r) + \Phi(t,r) = \frac{GM}{a(t)r} \left\{ 0 + \frac{\kappa^2}{1440\pi^2 a^2(t)r^2} - \frac{\kappa^2 H^2}{480\pi^2} + \dots \right\}.$$
(83)

As with its minimally coupled cousin, the κ^2/a^2r^2 corrections to (82-83) are well known from flat space background [86]. All (new and old) of the 1-loop corrections from massless, Dirac fermions are a factor of 6 times those in (81-83) [83,87]. For a loop of photons the factor is 12 [36,82,88,89].

The Lagrangian of general relativity with a cosmological constant Λ is,

$$\mathcal{L}_{\rm GR} = \frac{[R - (D-2)\Lambda]\sqrt{-g}}{16\pi G} \,. \tag{84}$$

On the other hand, all single matter loop corrections to gravity require the same two counterterms [90],

$$\Delta \mathcal{L} = c_1 R^2 \sqrt{-g} + c_2 C^{\alpha\beta\gamma\delta} C_{\alpha\beta\gamma\delta} \sqrt{-g} .$$
(85)

The first problem in applying the renormalization group to general relativity (84) is that the counterterms (85) do not seem to represent renormalizations of the classical action. However, one can easily decompose the Eddington (R^2) counterterm into a higher derivative part and two lower derivative parts which can be viewed as part of the classical action [63],

$$R^{2} = [R - D\Lambda + D\Lambda]^{2} = (R - D\Lambda)^{2} + 2D\Lambda[R - (D - 2)\Lambda] + (D - 4)D\Lambda^{2}.$$
 (86)

The final term in (86) could be regarded as renormalizing the cosmological constant, but the factor of (D-4) makes it irrelevant. In contrast, the middle term of (86) could be regarded as a renormalization of the graviton field strength.

Making a similar decomposition of the Weyl $(C^{\alpha\beta\gamma\delta}C_{\alpha\beta\gamma\delta})$ counterterm is more challenging. We first express it in terms of the Gauss-Bonnet scalar G,

$$C^{\mu\nu\rho\sigma}C_{\mu\nu\rho\sigma} = R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma} - \frac{4}{D-2}R^{\mu\nu}R_{\mu\nu} + \frac{2R^2}{(D-1)(D-2)}, \qquad (87)$$

$$= G + 4\left(\frac{D-3}{D-2}\right)R^{\mu\nu}R_{\mu\nu} - \frac{D(D-3)}{(D-1)(D-2)}R^2 .$$
(88)

The Gauss-Bonnet scalar is,

$$G \equiv R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma} - 4R^{\mu\nu}R_{\mu\nu} + R^2 .$$
(89)

It is significant by virtue of being a total derivative in D = 4 dimensions, which means it cannot affect 1-loop divergences. We now decompose the square of the Ricci tensor similar to (86) into a higher derivative term and two lower derivatives,

$$R^{\mu\nu}R_{\mu\nu} = (R^{\mu\nu} - \Lambda g^{\mu\nu})(R_{\mu\nu} - \Lambda g_{\mu\nu}) + 2\Lambda[R - (D - 2)\Lambda] + (D - 4)\Lambda^2 .$$
(90)

Substituting (86) and (90) in equation (88) gives,

$$C^{\mu\nu\rho\sigma}C_{\mu\nu\rho\sigma} = 4(\frac{D-3}{D-2})(R^{\mu\nu} - \Lambda g^{\mu\nu})(R_{\mu\nu} - \Lambda g_{\mu\nu}) - \frac{D(D-3)}{(D-1)(D-2)}(R - D\Lambda)^{2} - \frac{2(D-2)(D-3)}{D-1}\Lambda[R - (D-2)\Lambda)] - \frac{(D-2)(D-3)(D-4)}{D-1}\Lambda^{2} + G.$$
 (91)

Expressions (86) and (91) imply that the counterterms (85) can be viewed as a graviton field strength renormalization, which gives rise to a gamma function,

$$\delta Z = 2 \Big[D(D-1)c_1 - (D-2)(D-3)c_2 \Big] \kappa^2 H^2 \qquad \Longrightarrow \qquad \gamma = \frac{\partial \ln(1+\delta Z)}{\partial \ln(\mu^2)} . \tag{92}$$

The linearized Weyl tensor and the Newtonian potential can be viewed as 2-point Green's functions, for which the Callan-Symanzik equation implies,

$$\left[\frac{\partial}{\partial \ln(\mu)} + 2\gamma\right] G^{(2)} = 0 .$$
(93)

Relation (75) shows that factors of $\ln(\mu)$ always come in the form $\ln(\mu a)$, so it should be valid to replace the derivative with respect to $\ln(\mu)$ in expression (93) by the derivative with respect to $\ln(a)$. At this point one sees that the it is possible to predict the large temporal logarithms from renormalization.

The results are impressive. For a loop of massless, minimally coupled scalars the constants c_1 and c_2 in (85) are [54, 55],

$$c_1 = \frac{\mu^{D-4}\Gamma(\frac{D}{2})}{2^8\pi^{\frac{D}{2}}} \frac{(D-2)}{(D-1)^2(D-3)(D-4)} , \qquad (94)$$

$$c_2 = \frac{\mu^{D-4}\Gamma(\frac{D}{2})}{2^8\pi^{\frac{D}{2}}} \frac{2}{(D+1)(D-1)^2(D-3)^2(D-4)} .$$
(95)

Substituting these results in expression (92) give the gamma function,

$$\gamma_{\rm MMCS} = \frac{3\kappa^2 H^2}{320\pi^2} \ . \tag{96}$$

Using γ_{MMCS} in the Callan-Symanzik equation (93) not only explains the leading logarithms (77-79), but also permits a full resummation,

$$C_{0i0j}(t, \vec{x}) \longrightarrow C_{0i0j}^{\text{tree}}(t, \vec{x}) \times \left[a(t)\right]^{-\frac{3\kappa^2 H^2}{160\pi^2}},$$
 (97)

$$\Psi(t,r) \longrightarrow \frac{GM}{a(r)r} \times \left[a(t)rH\right]^{-\frac{3\kappa^2H^2}{160\pi^2}}.$$
(98)

For massless, conformally coupled scalars one has $c_1 = 0$ and c_2 equal to [83],

$$c_2 = \frac{\mu^{D-4}\Gamma(\frac{D}{2})}{2^8\pi^{\frac{D}{2}}} \frac{2}{(D+1)(D-1)(D-3)^2(D-4)} \qquad \Longrightarrow \qquad \gamma_{\rm MCCS} = -\frac{\kappa^2 H^2}{960\pi^2} \,. \tag{99}$$

This not only explains the leading logarithms at 1-loop (81-82), and again implies the results,

$$C_{0i0j}(t,\vec{x}) \longrightarrow C_{0i0j}^{\text{tree}}(t,\vec{x}) \times \left[a(t)\right]^{\frac{\kappa^2 H^2}{480\pi^2}}, \qquad (100)$$

$$\Psi(t,r) \longrightarrow \frac{GM}{a(r)r} \times \left[a(t)rH\right]^{\frac{\kappa^2 H^2}{480\pi^2}}.$$
(101)

A loop of Dirac fermions has $\gamma_{\text{Dirac}} = 6 \times \gamma_{\text{MMCS}}$ [83] and a loop of photons gives $\gamma_{\text{EM}} = 12 \times \gamma_{\text{MMCS}}$ [82].

7 Conclusions

Alexei Starobinsky possessed a towering intellect which allowed him to see deeply hidden truths. One of these is that a simple stochastic formalism can describe many of the secular loop corrections one finds in quantum field theory on inflationary backgrounds. This is quite surprising from the context of quantum field theory because the fields which obey Starobinsky's first order Langevin equation are ultraviolet finite, commuting variables, quite unlike the original quantum fields which obey second order equations, harbor ultraviolet divergences, and do not commute on or within the light-cone.

The message of this article is that Starobinsky's stochastic formalism is neither wrong, as many QFT experts believed, nor does it supplant quantum field theory, as some cosmologists believed. What it represents is the leading secular corrections in quantum field theories of undifferentiated "active" fields which experience inflationary particle production (2). It does not directly apply to "passive" fields, which do not experience inflationary particle production, or to differentiated active fields. Those systems make nonsecular, but still nonzero, contributions which come as much from the ultraviolet as from the infrared, and from the full mode function. The correct way to deal with these systems is by integrating out the problematic passive or differentiated active fields in the presence of a constant active field background. For passive fields this typically results in a standard effective potential induced by a field-dependent mass. For differentiated active fields one encounters new sorts of effective potentials induced by field-dependent field strengths and/or field-dependent Hubble parameters.

Finally, it is best avoid dogmatic adherence to statistical mechanics, and to check against explicit computations whenever possible. Although Starobinsky's formalism does have a statistical mechanical interpretation, its derivation rests on quantum field theory, and what it does is to predict a certain class of secular corrections. One must renormalize in the presence of either passive fields or differentiated active fields. And differentiated active fields produce new types of secular corrections from renormalization (75), which cannot be represented by any stochastic formalism. This second source of secular corrections was only discovered by explicit 1-loop and 2-loop computations. It is possible that further surprises await us, which Alexei would have loved.

Acknowledgements

It is a pleasure to acknowledge collaboration, conversation and correspondence on these subjects with A. Foraci, D. Glavan, S. P. Miao, T. Prokopec, N. C. Tsamis and B. Yesilyurt. This work was partially supported by NSF grant PHY-2207514 and by the Institute for Fundamental Theory at the University of Florida.

References

- A. A. Starobinsky, Phys. Lett. B 91, 99-102 (1980) doi:10.1016/0370-2693(80)90670-X
- [2] M. Tristram, A. J. Banday, K. M. Górski, R. Keskitalo, C. R. Lawrence, K. J. Andersen, R. B. Barreiro, J. Borrill, L. P. L. Colombo and H. K. Eriksen, et al. Phys. Rev. D 105, no.8, 083524 (2022) doi:10.1103/PhysRevD.105.083524 [arXiv:2112.07961 [astro-ph.CO]].
- [3] A. A. Starobinsky, JETP Lett. **30**, 682-685 (1979)
- [4] A. A. Starobinsky, Phys. Lett. B 117, 175-178 (1982) doi:10.1016/0370-2693(82)90541-X
- [5] A. A. Starobinsky, Lect. Notes Phys. 246, 107-126 (1986) doi:10.1007/3-540-16452-9_6
- [6] A. A. Starobinsky and J. Yokoyama, Phys. Rev. D 50, 6357-6368 (1994) doi:10.1103/PhysRevD.50.6357 [arXiv:astro-ph/9407016 [astro-ph]].
- [7] K. G. Wilson, Phys. Rev. B 4, 3174-3183 (1971) doi:10.1103/PhysRevB.4.3174
- [8] K. G. Wilson, Phys. Rev. B 4, 3184-3205 (1971) doi:10.1103/PhysRevB.4.3184
- [9] J. S. Schwinger, J. Math. Phys. 2, 407-432 (1961) doi:10.1063/1.1703727
- [10] K. T. Mahanthappa, Phys. Rev. **126**, 329-340 (1962) doi:10.1103/PhysRev.126.329
- [11] P. M. Bakshi and K. T. Mahanthappa, J. Math. Phys. 4, 1-11 (1963) doi:10.1063/1.1703883

- [12] P. M. Bakshi and K. T. Mahanthappa, J. Math. Phys. 4, 12-16 (1963) doi:10.1063/1.1703879
- [13] L. V. Keldysh, Zh. Eksp. Teor. Fiz. 47, 1515-1527 (1964) doi:10.1142/9789811279461_0007
- [14] V. K. Onemli and R. P. Woodard, Class. Quant. Grav. 19, 4607 (2002) doi:10.1088/0264-9381/19/17/311 [arXiv:gr-qc/0204065 [gr-qc]].
- [15] T. Prokopec, O. Tornkvist and R. P. Woodard, Phys. Rev. Lett. 89, 101301 (2002) doi:10.1103/PhysRevLett.89.101301
 [arXiv:astro-ph/0205331 [astro-ph]].
- T. Prokopec, O. Tornkvist and R. P. Woodard, Annals Phys. 303, 251-274 (2003) doi:10.1016/S0003-4916(03)00004-6 [arXiv:gr-qc/0205130 [gr-qc]].
- [17] A. Vilenkin and L. H. Ford, Phys. Rev. D 26, 1231 (1982) doi:10.1103/PhysRevD.26.1231
- [18] A. D. Linde, Phys. Lett. B 116, 335-339 (1982) doi:10.1016/0370-2693(82)90293-3
- [19] B. Allen and A. Folacci, Phys. Rev. D 35, 3771 (1987) doi:10.1103/PhysRevD.35.3771
- [20] D. N. Page and X. Wu, JCAP 11, 051 (2012) doi:10.1088/1475-7516/2012/11/051 [arXiv:1204.4462 [hep-th]].
- [21] L. R. Abramo and R. P. Woodard, Phys. Rev. D 65, 063515 (2002) doi:10.1103/PhysRevD.65.063515 [arXiv:astro-ph/0109272 [astro-ph]].
- [22] N. C. Tsamis and R. P. Woodard, Nucl. Phys. B 724, 295-328 (2005) doi:10.1016/j.nuclphysb.2005.06.031 [arXiv:gr-qc/0505115 [gr-qc]].
- [23] S. P. Miao and R. P. Woodard, Class. Quant. Grav. 23, 1721-1762 (2006) doi:10.1088/0264-9381/23/5/016 [arXiv:gr-qc/0511140 [gr-qc]].
- [24] S. P. Miao and R. P. Woodard, Phys. Rev. D 74, 024021 (2006) doi:10.1103/PhysRevD.74.024021 [arXiv:gr-qc/0603135 [gr-qc]].

- [25] T. M. C. Abbott *et al.* [DES], Phys. Rev. D 105, no.2, 023520 (2022) doi:10.1103/PhysRevD.105.023520 [arXiv:2105.13549 [astro-ph.CO]].
- [26] M. Kamionkowski and A. G. Riess, Ann. Rev. Nucl. Part. Sci. 73, 153-180 (2023) doi:10.1146/annurev-nucl-111422-024107 [arXiv:2211.04492 [astro-ph.CO]].
- [27] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910
 [arXiv:1807.06209 [astro-ph.CO]].
- [28] G. Geshnizjani, W. H. Kinney and A. Moradinezhad Dizgah, JCAP 11, 049 (2011) doi:10.1088/1475-7516/2011/11/049 [arXiv:1107.1241 [astroph.CO]].
- [29] E. Lifshitz, J. Phys. (USSR) 10, no.2, 116 (1946) doi:10.1007/s10714-016-2165-8
- [30] V. F. Mukhanov and G. V. Chibisov, JETP Lett. **33**, 532-535 (1981)
- [31] V. K. Onemli and R. P. Woodard, Phys. Rev. D 70, 107301 (2004) doi:10.1103/PhysRevD.70.107301 [arXiv:gr-qc/0406098 [gr-qc]].
- [32] E. O. Kahya, V. K. Onemli and R. P. Woodard, Phys. Rev. D 81, 023508 (2010) doi:10.1103/PhysRevD.81.023508 [arXiv:0904.4811 [gr-qc]].
- [33] N. A. Chernikov and E. A. Tagirov, Ann. Inst. H. Poincare A Phys. Theor. 9, 109 (1968)
- [34] C. Schomblond and P. Spindel, Ann. Inst. H. Poincare Phys. Theor. 25, 67-78 (1976)
- [35] T. S. Bunch and P. C. W. Davies, Proc. Roy. Soc. Lond. A 360, 117-134 (1978) doi:10.1098/rspa.1978.0060
- [36] C. L. Wang and R. P. Woodard, Phys. Rev. D 92, 084008 (2015) doi:10.1103/PhysRevD.92.084008 [arXiv:1508.01564 [gr-qc]].
- [37] S. R. Coleman and E. J. Weinberg, Phys. Rev. D 7, 1888-1910 (1973) doi:10.1103/PhysRevD.7.1888

- [38] S. P. Miao and R. P. Woodard, Phys. Rev. D 74, 044019 (2006) doi:10.1103/PhysRevD.74.044019 [arXiv:gr-qc/0602110 [gr-qc]].
- [39] T. Prokopec, N. C. Tsamis and R. P. Woodard, Annals Phys. 323, 1324-1360 (2008) doi:10.1016/j.aop.2007.08.008 [arXiv:0707.0847 [gr-qc]].
- [40] P. Candelas and D. J. Raine, Phys. Rev. D 12, 965-974 (1975) doi:10.1103/PhysRevD.12.965
- [41] N. C. Tsamis and R. P. Woodard, J. Math. Phys. 48, 052306 (2007) doi:10.1063/1.2738361 [arXiv:gr-qc/0608069 [gr-qc]].
- [42] B. Allen, Nucl. Phys. B 226, 228-252 (1983) doi:10.1016/0550-3213(83)90470-4
- [43] T. Prokopec, N. C. Tsamis and R. P. Woodard, Class. Quant. Grav. 24, 201-230 (2007) doi:10.1088/0264-9381/24/1/011 [arXiv:gr-qc/0607094 [gr-qc]].
- [44] T. Prokopec, N. C. Tsamis and R. P. Woodard, Phys. Rev. D 78, 043523 (2008) doi:10.1103/PhysRevD.78.043523 [arXiv:0802.3673 [gr-qc]].
- [45] N. N. Bogoliubov and O. S. Parasiuk, Acta Math. 97, 227-266 (1957) doi:10.1007/BF02392399
- [46] K. Hepp, Commun. Math. Phys. 2, 301-326 (1966) doi:10.1007/BF01773358
- [47] W. Zimmermann, Commun. Math. Phys. 11, 1-8 (1968) doi:10.1007/BF01654298
- [48] W. Zimmermann, Commun. Math. Phys. 15, 208-234 (1969) doi:10.1007/BF01645676
- [49] J. F. Donoghue, Phys. Rev. Lett. 72, 2996-2999 (1994) doi:10.1103/PhysRevLett.72.2996 [arXiv:gr-qc/9310024 [gr-qc]].
- [50] J. F. Donoghue, Phys. Rev. D 50, 3874-3888 (1994) doi:10.1103/PhysRevD.50.3874 [arXiv:gr-qc/9405057 [gr-qc]].
- [51] J. Donoghue, Scholarpedia **12**, no.4, 32997 (2017) doi:10.4249/scholarpedia.32997

- [52] T. Brunier, V. K. Onemli and R. P. Woodard, Class. Quant. Grav. 22, 59-84 (2005) doi:10.1088/0264-9381/22/1/005 [arXiv:gr-qc/0408080 [grqc]].
- [53] N. C. Tsamis and R. P. Woodard, Phys. Rev. D 54, 2621-2639 (1996) doi:10.1103/PhysRevD.54.2621 [arXiv:hep-ph/9602317 [hep-ph]].
- [54] S. Park and R. P. Woodard, Phys. Rev. D 83, 084049 (2011) doi:10.1103/PhysRevD.83.084049 [arXiv:1101.5804 [gr-qc]].
- [55] S. P. Miao, N. C. Tsamis and R. P. Woodard, JHEP 07, 099 (2024) doi:10.1007/JHEP07(2024)099 [arXiv:2405.00116 [gr-qc]].
- [56] H. Kitamoto and Y. Kitazawa, Phys. Rev. D 83, 104043 (2011) doi:10.1103/PhysRevD.83.104043 [arXiv:1012.5930 [hep-th]].
- [57] H. Kitamoto and Y. Kitazawa, Phys. Rev. D 85, 044062 (2012) doi:10.1103/PhysRevD.85.044062 [arXiv:1109.4892 [hep-th]].
- [58] H. Kitamoto, Phys. Rev. D 100, no.2, 025020 (2019) doi:10.1103/PhysRevD.100.025020 [arXiv:1811.01830 [hep-th]].
- [59] S. P. Miao, N. C. Tsamis and R. P. Woodard, JHEP 03, 069 (2022) doi:10.1007/JHEP03(2022)069 [arXiv:2110.08715 [gr-qc]].
- [60] R. P. Woodard and B. Yesilyurt, JHEP 06, 206 (2023) doi:10.1007/JHEP06(2023)206 [arXiv:2302.11528 [gr-qc]].
- [61] S. Basu and R. P. Woodard, Class. Quant. Grav. 33, no.20, 205007 (2016) doi:10.1088/0264-9381/33/20/205007 [arXiv:1606.02417 [gr-qc]].
- [62] S. Basu, N. C. Tsamis and R. P. Woodard, JHEP 07, 037 (2017) doi:10.1007/JHEP07(2017)037 [arXiv:1612.07406 [gr-qc]].
- [63] S. P. Miao, N. C. Tsamis and R. P. Woodard, Class. Quant. Grav. 41, no.21, 215007 (2024) doi:10.1088/1361-6382/ad7dc8 [arXiv:2405.01024 [gr-qc]].
- [64] N. C. Tsamis, R. P. Woodard and B. Yesilyurt, Phys. Lett. B 849, 138472 (2024) doi:10.1016/j.physletb.2024.138472 [arXiv:2312.15913 [gr-qc]].

- [65] N. C. Tsamis and R. P. Woodard, Commun. Math. Phys. 162, 217-248 (1994) doi:10.1007/BF02102015
- [66] R. P. Woodard, [arXiv:gr-qc/0408002 [gr-qc]].
- [67] S. P. Miao, N. C. Tsamis and R. P. Woodard, [arXiv:2409.12003 [gr-qc]].
- [68] R. L. Arnowitt, S. Deser and C. W. Misner, Gen. Rel. Grav. 40, 1997-2027 (2008) doi:10.1007/s10714-008-0661-1 [arXiv:gr-qc/0405109 [gr-qc]].
- [69] R. P. Woodard, [arXiv:2501.05077 [gr-qc]].
- [70] N. C. Tsamis and R. P. Woodard, Annals Phys. **321**, 875-893 (2006) doi:10.1016/j.aop.2005.08.004 [arXiv:gr-qc/0506056 [gr-qc]].
- [71] L. Tan, N. C. Tsamis and R. P. Woodard, Class. Quant. Grav. 38, no.14, 145024 (2021) doi:10.1088/1361-6382/ac0233 [arXiv:2103.08547 [gr-qc]].
- [72] L. Tan, N. C. Tsamis and R. P. Woodard, Phil. Trans. Roy. Soc. Lond. A 380, 0187 (2021) doi:10.1098/rsta.2021.0187 [arXiv:2107.13905 [gr-qc]].
- [73] L. Tan, N. C. Tsamis and R. P. Woodard, Universe 8, no.7, 376 (2022) doi:10.3390/universe8070376 [arXiv:2206.11467 [gr-qc]].
- [74] S. P. Miao and R. P. Woodard, Class. Quant. Grav. 25, 145009 (2008) doi:10.1088/0264-9381/25/14/145009 [arXiv:0803.2377 [gr-qc]].
- [75] E. O. Kahya and R. P. Woodard, Phys. Rev. D 76, 124005 (2007) doi:10.1103/PhysRevD.76.124005 [arXiv:0709.0536 [gr-qc]].
- [76] K. E. Leonard and R. P. Woodard, Class. Quant. Grav. 31, 015010 (2014) doi:10.1088/0264-9381/31/1/015010 [arXiv:1304.7265 [gr-qc]].
- [77] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, Class. Quant. Grav. **31**, 175002 (2014) doi:10.1088/0264-9381/31/17/175002
 [arXiv:1308.3453 [gr-qc]].
- [78] C. L. Wang and R. P. Woodard, Phys. Rev. D 91, no.12, 124054 (2015) doi:10.1103/PhysRevD.91.124054 [arXiv:1408.1448 [gr-qc]].

- [79] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, Phys. Rev. D 101, no.10, 106016 (2020) doi:10.1103/PhysRevD.101.106016 [arXiv:2003.02549 [gr-qc]].
- [80] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, Phys. Rev. D 103, no.10, 105022 (2021) doi:10.1103/PhysRevD.103.105022 [arXiv:2007.10395 [gr-qc]].
- [81] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, JHEP 03, 088 (2022) doi:10.1007/JHEP03(2022)088 [arXiv:2112.00959 [gr-qc]].
- [82] A. J. Foraci and R. P. Woodard, [arXiv:2412.11022 [gr-qc]].
- [83] A. J. Foraci and R. P. Woodard, [arXiv:2501.01972 [gr-qc]].
- [84] S. Park, T. Prokopec and R. P. Woodard, JHEP 01, 074 (2016) doi:10.1007/JHEP01(2016)074 [arXiv:1510.03352 [gr-qc]].
- [85] A. F. Radkowski, Ann. Phys. 56, no. 2, 319-354 (1970, doi.org/10.1016/0003-4916(70)90021-7.
- [86] D. M. Capper, Nuovo Cim. A 25, 29 (1975) doi:10.1007/BF02735608
- [87] D. M. Capper and M. J. Duff, Nucl. Phys. B 82, 147-154 (1974) doi:10.1016/0550-3213(74)90582-3
- [88] M. J. Duff and J. T. Liu, Phys. Rev. Lett. 85, 2052-2055 (2000) doi:10.1088/0264-9381/18/16/310 [arXiv:hep-th/0003237 [hep-th]].
- [89] D. M. Capper, M. J. Duff and L. Halpern, Phys. Rev. D 10, 461-467 (1974) doi:10.1103/PhysRevD.10.461
- [90] G. 't Hooft and M. J. G. Veltman, Ann. Inst. H. Poincare A Phys. Theor. 20, 69-94 (1974)