

p -SUMMING WEIGHTED HOLOMORPHIC MAPPINGS

M. G. CABRERA-PADILLA, A. JIMÉNEZ-VARGAS, AND A. KETEN ÇOPUR

ABSTRACT. Given an open subset U of a complex Banach space E , a weight v on U , and a complex Banach space F , let $\mathcal{H}_v^\infty(U, F)$ denote the Banach space of all weighted holomorphic mappings $f: U \rightarrow F$, under the weighted supremum norm $\|f\|_v := \sup \{v(x)\|f(x)\| : x \in U\}$. In this paper, we introduce and study the class $\Pi_p^{\mathcal{H}_v^\infty}(U, F)$ of p -summing weighted holomorphic mappings. We prove that it is an injective Banach ideal of weighted holomorphic mappings which is not generated by composition. Variants for weighted holomorphic mappings of Pietsch Domination Theorem, Pietsch Factorization Theorem and Maurey Extrapolation Theorem are presented. We also identify the spaces of p -summing weighted holomorphic mappings from U into F^* under the norm $\pi_p^{\mathcal{H}_v^\infty}$ with the duals of F -valued \mathcal{H}_v^∞ -molecules on U under a suitable version $d_p^{\mathcal{H}_v^\infty}$ of the Chevet–Saphar tensor norms.

INTRODUCTION

The concept of p -summing operators between Banach spaces, introduced by Grothendieck [13] for $p = 1$ and by Pietsch [21] for any $p > 0$, has played an important role in the theory of Banach spaces (see the monographs [9, 10, 22]).

Its influence has motivated the development of the theory of p -summability for different classes of mappings in addition to continuous linear operators. This is the case of multilinear operators and polynomials studied by Achour and Mezrag [1], Angulo-López and Fernández-Unzueta [2] and Dimant [11] (see also the survey [19] by Pellegrino, Rueda and Sánchez-Pérez about the p -summability in this context); Lipschitz mappings addressed by Farmer and Johnson [12] and Saadi [23]; and holomorphic mappings dealt by Matos [15] and Pellegrino [18].

Let E and F be complex Banach spaces and let U be an open subset of E . Let $\mathcal{H}(U, F)$ be the space of all holomorphic mappings from U into F . A weight v on U is a (strictly) positive continuous function. The space of weighted holomorphic mappings, denoted by $\mathcal{H}_v^\infty(U, F)$, is the Banach space of all mappings $f \in \mathcal{H}(U, F)$ such that

$$\|f\|_v := \sup \{v(x)\|f(x)\| : x \in U\} < \infty,$$

under the weighted supremum norm $\|\cdot\|_v$. In particular, $\mathcal{H}_v^\infty(U) := \mathcal{H}_v^\infty(U, \mathbb{C})$. We refer to the survey by Bonet [4] and the references therein for a complete information about such spaces.

We will develop in this paper a theory of p -summability for weighted holomorphic mappings. Thus, we introduce the notion of p -summing weighted holomorphic mappings and show that it is appropriate to establish variants for the weighted holomorphic setting of some known results of the theory of p -summing operators.

Date: January 28, 2025.

2020 Mathematics Subject Classification. 46B20, 46E40, 46E50, 46T25.

Key words and phrases. Weighted holomorphic mappings, p -summing operators, Chevet–Saphar norms.

The first two authors were partially supported by Junta de Andalucía grant FQM194, and by Ministerio de Ciencia e Innovación grant PID2021-122126NB-C31 funded by MCIN/AEI/ 10.13039/501100011033 and by “ERDF A way of making Europe”.

Thus, Section 1 shows that p -summing weighted holomorphic mappings enjoy an injective Banach ideal property and satisfy Pietsch Domination Theorem, Pietsch Factorization Theorem, Maurey Extrapolation Theorem, and Inclusion and Coincidence Theorem. We also show that every p -summing weighted holomorphic mapping from U to F has a relatively weakly compact v -range in F that becomes relatively compact whenever F is reflexive.

Section 2 contains a tensorial approach to this new class of weighted holomorphic mappings. In this way, we introduce a suitable tensor product space equipped with a Chevet–Saphar type tensor norm in such a way that its dual is isometrically isomorphic to the space of p -summing weighted holomorphic mappings.

We will need to fix some notation and recall some facts on weighted holomorphic maps.

Notation. As usual, $\mathcal{L}(E, F)$ denotes the Banach space of all bounded linear operators from E into F , equipped with the operator canonical norm. In particular, $\mathcal{L}(E, \mathbb{C})$ is denoted by E^* . For $x \in E$ and $x^* \in E^*$, we will sometimes write $\langle x^*, x \rangle$ instead of $x^*(x)$. B_E stands for the closed unit ball of E . We will write $(E, \|\cdot\|_E) \leq (F, \|\cdot\|_F)$ to indicate that $E \subseteq F$ and $\|x\|_F \leq \|x\|_E$ for all $x \in E$. For any $1 < p < \infty$, let $p^* = p/(p-1)$ denote the conjugate index of p .

Preliminaries. Following [3, 5, 6], $\mathcal{G}_v^\infty(U)$ is the space of all linear functionals on $\mathcal{H}_v^\infty(U)$ whose restriction to $B_{\mathcal{H}_v^\infty(U)}$ is continuous for the compact-open topology.

Theorem 0.1. [3, 5, 6] *Let U be an open set of a complex Banach space E and let v be a weight on U .*

- (i) $\mathcal{G}_v^\infty(U)$ is a closed subspace of $\mathcal{H}_v^\infty(U)^*$, and the mapping $J_v: \mathcal{H}_v^\infty(U) \rightarrow \mathcal{G}_v^\infty(U)^*$, given by $J_v(g)(\phi) = \phi(g)$ for $\phi \in \mathcal{G}_v^\infty(U)$ and $g \in \mathcal{H}_v^\infty(U)$, is an isometric isomorphism.
- (ii) For each $x \in U$, the functional $\delta_x: \mathcal{H}_v^\infty(U) \rightarrow \mathbb{C}$, defined by $\delta_x(f) = f(x)$ for $f \in \mathcal{H}_v^\infty(U)$, is in $\mathcal{G}_v^\infty(U)$, and there exists $g_x \in B_{\mathcal{H}_v^\infty(U)}$ such that $g_x(x) = \|\delta_x\| := \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} |g(x)|$.
- (iii) The mapping $\Delta_v: U \rightarrow \mathcal{G}_v^\infty(U)$ given by $\Delta_v(x) = \delta_x$ is in $\mathcal{H}_v^\infty(U, \mathcal{G}_v^\infty(U))$ with $\|\Delta_v\|_v \leq 1$.
- (iv) For every complex Banach space F and every mapping $f \in \mathcal{H}_v^\infty(U, F)$, there exists a unique operator $T_f \in \mathcal{L}(\mathcal{G}_v^\infty(U), F)$ such that $T_f \circ \Delta_v = f$. Furthermore, $\|T_f\| = \|f\|_v$. \square

1. p -SUMMING WEIGHTED HOLOMORPHIC MAPPINGS

From now on, unless otherwise stated, we will suppose that $p \in [1, \infty]$ and E, F and G will denote complex Banach spaces, U will be an open subset of E , and v will be a weight on U .

We begin introducing the following variant for weighted holomorphic mappings of the known concept of p -summing linear operators between Banach spaces.

Definition 1.1. Given $1 \leq p \leq \infty$, a mapping $f \in \mathcal{H}(U, F)$ is said to be p -summing weighted holomorphic if there is a constant $C \geq 0$ such that

$$\left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} \leq C \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} \quad \text{if } 1 \leq p < \infty,$$

$$\max_{1 \leq i \leq n} |\lambda_i| v(x_i) \|f(x_i)\| \leq C \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\max_{1 \leq i \leq n} |\lambda_i| v(x_i) |g(x_i)| \right) \quad \text{if } p = \infty,$$

for any $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. We denote by $\pi_p^{\mathcal{H}_v^\infty}(f)$ the infimum of all constants C satisfying the inequality above, and by $\Pi_p^{\mathcal{H}_v^\infty}(U, F)$ the set of all p -summing weighted holomorphic mappings from U into F .

We now show that every p -summing weighted holomorphic mapping turns out to be a weighted holomorphic mapping and this justifies the terminology used in Definition 1.1.

Remark 1.2. $\Pi_p^{\mathcal{H}_v^\infty}(U, F) \subseteq \mathcal{H}_v^\infty(U, F)$ with $\|f\|_v \leq \pi_p^{\mathcal{H}_v^\infty}(f)$ for all $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$. Indeed, for such a function f , note that

$$v(x)\|f(x)\| = (v(x)^p\|f(x)\|^p)^{\frac{1}{p}} \leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} (v(x)^p|g(x)|^p)^{\frac{1}{p}} \leq \pi_p^{\mathcal{H}_v^\infty}(f)$$

for all $x \in U$.

Our main purpose in this paper is to study the spaces of Banach-valued p -summing weighted holomorphic mappings. We will go presenting their main properties in different subsections.

1.1. Injective Banach ideal property. We first prove that the set of p -summing weighted holomorphic mappings enjoys an injective Banach ideal structure.

Let us recall from [7] that a Banach ideal of weighted holomorphic mappings (in short, a Banach weighted holomorphic ideal) is an assignment $[\mathcal{I}^{\mathcal{H}_v^\infty}, \|\cdot\|_{\mathcal{I}^{\mathcal{H}_v^\infty}}]$ which associates with every pair (U, F) , where U is an open subset of a complex Banach space E and F is a complex Banach space, a set $\mathcal{I}^{\mathcal{H}_v^\infty}(U, F) \subseteq \mathcal{H}_v^\infty(U, F)$ and a function $\|\cdot\|_{\mathcal{I}^{\mathcal{H}_v^\infty}} : \mathcal{I}^{\mathcal{H}_v^\infty}(U, F) \rightarrow \mathbb{R}_0^+$ satisfying the properties:

- (P1) $(\mathcal{I}^{\mathcal{H}_v^\infty}(U, F), \|\cdot\|_{\mathcal{I}^{\mathcal{H}_v^\infty}})$ is a Banach space with $\|f\|_{\mathcal{I}^{\mathcal{H}_v^\infty}} \geq \|f\|_v$ for all $f \in \mathcal{I}^{\mathcal{H}_v^\infty}(U, F)$.
- (P2) For any $h \in \mathcal{H}_v^\infty(U)$ and $y \in F$, the map $h \cdot y : x \mapsto h(x)y$ from U to F is in $\mathcal{I}^{\mathcal{H}_v^\infty}(U, F)$ with $\|h \cdot y\|_{\mathcal{I}^{\mathcal{H}_v^\infty}} = \|h\|_v\|y\|$.
- (P3) The ideal property: if V is an open subset of E such that $V \subseteq U$, $h \in \mathcal{H}(V, U)$ with $c_v(h) := \sup_{x \in V} (v(x)/v(h(x))) < \infty$, $f \in \mathcal{I}^{\mathcal{H}_v^\infty}(U, F)$ and $T \in \mathcal{L}(F, G)$, where G is a complex Banach space, then $T \circ f \circ h \in \mathcal{I}^{\mathcal{H}_v^\infty}(V, G)$ with $\|T \circ f \circ h\|_{\mathcal{I}^{\mathcal{H}_v^\infty}} \leq \|T\|\|f\|_{\mathcal{I}^{\mathcal{H}_v^\infty}}c_v(h)$.

A Banach weighted holomorphic ideal $[\mathcal{I}^{\mathcal{H}_v^\infty}, \|\cdot\|_{\mathcal{I}^{\mathcal{H}_v^\infty}}]$ is called:

- (I) Injective if for any mapping $f \in \mathcal{H}_v^\infty(U, F)$, any complex Banach space G and any isometric linear embedding $\iota : F \rightarrow G$, we have that $f \in \mathcal{I}^{\mathcal{H}_v^\infty}(U, F)$ with $\|f\|_{\mathcal{I}^{\mathcal{H}_v^\infty}} = \|\iota \circ f\|_{\mathcal{I}^{\mathcal{H}_v^\infty}}$ whenever $\iota \circ f \in \mathcal{I}^{\mathcal{H}_v^\infty}(U, G)$.

Proposition 1.3. $[\Pi_p^{\mathcal{H}_v^\infty}, \pi_p^{\mathcal{H}_v^\infty}]$ is an injective Banach weighted holomorphic ideal.

Proof. We will prove the case $1 \leq p < \infty$, and for $p = \infty$ a similar proof works.

(P1) Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. If $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ and $\pi_p^{\mathcal{H}_v^\infty}(f) = 0$, then $\|f\|_v = 0$ by Remark 1.2, and so $f = 0$. Given $f_1, f_2 \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$, one has

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|(f_1 + f_2)(x_i)\|^p \right)^{\frac{1}{p}} &\leq \left(\sum_{i=1}^n (|\lambda_i| v(x_i) \|f_1(x_i)\| + |\lambda_i| v(x_i) \|f_2(x_i)\|)^p \right)^{\frac{1}{p}} \\ &\leq \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f_1(x_i)\|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f_2(x_i)\|^p \right)^{\frac{1}{p}} \\ &\leq (\pi_p^{\mathcal{H}_v^\infty}(f_1) + \pi_p^{\mathcal{H}_v^\infty}(f_2)) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

and thus $f_1 + f_2 \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(f_1 + f_2) \leq \pi_p^{\mathcal{H}_v^\infty}(f_1) + \pi_p^{\mathcal{H}_v^\infty}(f_2)$.

Let $\lambda \in \mathbb{C}$ and $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$. An easy calculation yields

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|(\lambda f)(x_i)\|^p \right)^{\frac{1}{p}} &= |\lambda| \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} \\ &\leq |\lambda| \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

and so $\lambda f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(\lambda f) \leq |\lambda| \pi_p^{\mathcal{H}_v^\infty}(f)$. Hence $\pi_p^{\mathcal{H}_v^\infty}(\lambda f) = 0 = |\lambda| \pi_p^{\mathcal{H}_v^\infty}(f)$ if $\lambda = 0$. For $\lambda \neq 0$, it is clear that $\pi_p^{\mathcal{H}_v^\infty}(f) = \pi_p^{\mathcal{H}_v^\infty}(\lambda^{-1}(\lambda f)) \leq |\lambda|^{-1} \pi_p^{\mathcal{H}_v^\infty}(\lambda f)$, hence $|\lambda| \pi_p^{\mathcal{H}_v^\infty}(f) \leq \pi_p^{\mathcal{H}_v^\infty}(\lambda f)$, and, consequently, $\pi_p^{\mathcal{H}_v^\infty}(\lambda f) = |\lambda| \pi_p^{\mathcal{H}_v^\infty}(f)$. In this way, $(\Pi_p^{\mathcal{H}_v^\infty}(U, F), \pi_p^{\mathcal{H}_v^\infty})$ is a normed space.

To show that it is a Banach space, let $(f_n)_{n \geq 1}$ be a sequence in $\Pi_p^{\mathcal{H}_v^\infty}(U, F)$ such that $\sum_n \pi_p^{\mathcal{H}_v^\infty}(f_n)$ converges. Since $\|f_n\|_v \leq \pi_p^{\mathcal{H}_v^\infty}(f_n)$ for all $n \in \mathbb{N}$ and $(\mathcal{H}_v^\infty(U, F), \|\cdot\|_v)$ is a Banach space, then $\sum_n f_n$ converges in $(\mathcal{H}_v^\infty(U, F), \|\cdot\|_v)$ to a function $f \in \mathcal{H}_v^\infty(U, F)$. Given $m \in \mathbb{N}$, $x_1, \dots, x_m \in U$ and $\lambda_1, \dots, \lambda_m \in \mathbb{C}$, we have

$$\begin{aligned} \left(\sum_{k=1}^m |\lambda_k|^p v(x_k)^p \left\| \sum_{i=1}^n f_i(x_k) \right\|^p \right)^{\frac{1}{p}} &\leq \pi_p^{\mathcal{H}_v^\infty} \left(\sum_{i=1}^n f_i \right) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{k=1}^m |\lambda_k|^p v(x_k)^p |g(x_k)|^p \right)^{\frac{1}{p}} \\ &\leq \sum_{i=1}^n \pi_p^{\mathcal{H}_v^\infty}(f_i) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{k=1}^m |\lambda_k|^p v(x_k)^p |g(x_k)|^p \right)^{\frac{1}{p}} \end{aligned}$$

for all $n \in \mathbb{N}$. Taking limits with $n \rightarrow \infty$, we deduce

$$\left(\sum_{k=1}^m |\lambda_k|^p v(x_k)^p \left\| \sum_{i=1}^{\infty} f_i(x_k) \right\|^p \right)^{\frac{1}{p}} \leq \sum_{i=1}^{\infty} \pi_p^{\mathcal{H}_v^\infty}(f_i) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{k=1}^m |\lambda_k|^p v(x_k)^p |g(x_k)|^p \right)^{\frac{1}{p}},$$

using that, for any $x \in U$, one has

$$\left\| \sum_{i=1}^n v(x) f_i(x) - v(x) f(x) \right\| \leq \left\| \sum_{i=1}^n f_i - f \right\|_v$$

for all $n \in \mathbb{N}$. Hence $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(f) \leq \sum_{n=1}^{\infty} \pi_p^{\mathcal{H}_v^\infty}(f_n)$. Furthermore,

$$\pi_p^{\mathcal{H}_v^\infty} \left(f - \sum_{i=1}^n f_i \right) = \pi_p^{\mathcal{H}_v^\infty} \left(\sum_{i=n+1}^{\infty} f_i \right) \leq \sum_{i=n+1}^{\infty} \pi_p^{\mathcal{H}_v^\infty}(f_i)$$

for all $n \in \mathbb{N}$, and so f is the $\pi_p^{\mathcal{H}_v^\infty}$ -limit of the series $\sum_n f_n$. This proves the completeness of $\pi_p^{\mathcal{H}_v^\infty}$.

(P2) Let $h \in \mathcal{H}_v^\infty(U)$ and $y \in F$. Note that $h \cdot y \in \mathcal{H}_v^\infty(U, F)$ with $\|h \cdot y\|_v = \|h\|_v \|y\|$. We can suppose $h \neq 0$ and obtain

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|(h \cdot y)(x_i)\|^p \right)^{\frac{1}{p}} &= \|y\| \|h\|_v \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \left| \left(\frac{h}{\|h\|_v} \right)(x_i) \right|^p \right)^{\frac{1}{p}} \\ &\leq \|y\| \|h\|_v \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Thus $h \cdot y \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(h \cdot y) \leq \|h\|_v \|y\|$. Conversely, note that $\|h\|_v \|y\| \leq \pi_p^{\mathcal{H}_v^\infty}(h \cdot y)$ by Remark 1.2.

(P3) Let V be an open subset of E such that $V \subseteq U$, $h \in \mathcal{H}(V, U)$ with $c_v(h) < \infty$, $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ and $T \in \mathcal{L}(F, G)$. Observe that $T \circ f \circ h \in \mathcal{H}_v^\infty(V, G)$. For any $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in V$, we obtain

$$\begin{aligned} & \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|(T \circ f \circ h)(x_i)\|^p \right)^{\frac{1}{p}} \leq \|T\| \left(\sum_{i=1}^n |\lambda_i|^p \frac{v(x_i)^p}{v(h(x_i))^p} v(h(x_i))^p \|f(h(x_i))\|^p \right)^{\frac{1}{p}} \\ & \leq \|T\| \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p \frac{v(x_i)^p}{v(h(x_i))^p} v(h(x_i))^p |g(h(x_i))|^p \right)^{\frac{1}{p}} \\ & = \|T\| \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |(g \circ h)(x_i)|^p \right)^{\frac{1}{p}} \\ & \leq \|T\| \pi_p^{\mathcal{H}_v^\infty}(f) c_v(h) \sup_{0 \neq g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|g\|_v^p \left| \left(\frac{g \circ h}{c_v(h)\|g\|_v} \right)(x_i) \right|^p \right)^{\frac{1}{p}} \\ & \leq \|T\| \pi_p^{\mathcal{H}_v^\infty}(f) c_v(h) \sup_{g_0 \in B_{\mathcal{H}_v^\infty}(V)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g_0(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

since $g \circ h \in \mathcal{H}_v^\infty(V)$ with $\|g \circ h\|_v \leq c_v(h)\|g\|_v$. So, $T \circ f \circ h \in \Pi_p^{\mathcal{H}_v^\infty}(V, G)$ and $\pi_p^{\mathcal{H}_v^\infty}(T \circ f \circ h) \leq \|T\| \pi_p^{\mathcal{H}_v^\infty}(f) c_v(h)$.

(I) Let $f \in \mathcal{H}_v^\infty(U, F)$ and $\iota: F \rightarrow G$ be an into linear isometry such that $\iota \circ f \in \Pi_p^{\mathcal{H}_v^\infty}(U, G)$. Given $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$, we have

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|(\iota \circ f)(x_i)\|^p \right)^{\frac{1}{p}} \\ &\leq \pi_p^{\mathcal{H}_v^\infty}(\iota \circ f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

and so $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(f) \leq \pi_p^{\mathcal{H}_v^\infty}(\iota \circ f)$. The opposite inequality follows from (P3). \square

Let us recall (see [7]) that given an operator ideal \mathcal{I} , a mapping $f \in \mathcal{H}_v^\infty(U, F)$ belongs to the composition ideal $\mathcal{I} \circ \mathcal{H}_v^\infty$, and we write $f \in \mathcal{I} \circ \mathcal{H}_v^\infty(U, F)$, if there are a complex Banach space G , an operator $T \in \mathcal{I}(G, F)$ and a mapping $g \in \mathcal{H}_v^\infty(U, G)$ such that $f = T \circ g$.

Let us show that the weighted holomorphic ideal $\Pi_p^{\mathcal{H}_v^\infty}$ is not of this type. We thank Professor Elhadj Dahia for suggesting this result to us.

Proposition 1.4. *For $1 \leq p < \infty$, the Banach ideal of p -summing weighted holomorphic mappings $\Pi_p^{\mathcal{H}_v^\infty}$ and the corresponding composition ideal $\Pi_p \circ \mathcal{H}_v^\infty$ do not coincide.*

Proof. Let $1 \leq p < \infty$. Clearly, $\Delta_v \in \Pi_p^{\mathcal{H}_v^\infty}(\mathbb{C}, \mathcal{G}_v^\infty(\mathbb{C}))$ with $\pi_p^{\mathcal{H}_v^\infty}(\Delta_v) \leq 1$. On the other hand, the equality $\text{id}_{\mathcal{G}_v^\infty(\mathbb{C})} \circ \Delta_v = \Delta_v$ and the uniqueness of the linearization shows that $T_{\Delta_v} = \text{id}_{\mathcal{G}_v^\infty(\mathbb{C})}$ (the identity map on the infinite-dimensional Banach space $\mathcal{G}_v^\infty(\mathbb{C})$). Hence $T_{\Delta_v} \notin \Pi_p(\mathcal{G}_v^\infty(\mathbb{C}), \mathcal{G}_v^\infty(\mathbb{C}))$ by Weak Dvoretzky–Rogers Theorem [10, 2.18]. Finally, [7, Theorem 2.7] asserts that $\Delta_v \notin \Pi_p \circ \mathcal{H}_v^\infty(\mathbb{C}, \mathcal{G}_v^\infty(\mathbb{C}))$. \square

1.2. Pietsch domination. We can characterize p -summing weighted holomorphic mappings in terms of the known property of Pietsch domination (see [21, Theorem 2]).

In light of Theorem 0.1, $H_v^\infty(U)$ is a dual Banach space and, consequently, this space can be endowed with the weak* topology. Let $\mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ be the set of all Borel regular probability measures μ on the compact set $(B_{\mathcal{H}_v^\infty(U)}, w^*)$.

Theorem 1.5. *Let $1 \leq p < \infty$ and $f \in \mathcal{H}_v^\infty(U, F)$. The following statements are equivalent:*

- (i) f is p -summing weighted holomorphic.
- (ii) There is a constant $C \geq 0$ and a measure $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ such that

$$\|f(x)\| \leq C \left(\int_{B_{\mathcal{H}_v^\infty(U)}} |g(x)|^p d\mu(g) \right)^{\frac{1}{p}} \quad (x \in U).$$

In this case, $\pi_p^{\mathcal{H}_v^\infty}(f)$ is the infimum of all constants $C \geq 0$ satisfying the inequality and, in fact, this infimum is attained.

Proof. (i) \Rightarrow (ii): In order to apply an unified abstract version of Pietsch Domination Theorem stated in [20], consider the functions $S : \mathcal{H}_v^\infty(U, F) \times U \times \mathbb{C} \rightarrow [0, \infty[$ and $R : B_{\mathcal{H}_v^\infty(U)} \times U \times \mathbb{C} \rightarrow [0, \infty[$ defined, respectively, by

$$\begin{aligned} S(h, x, \lambda) &= |\lambda| v(x) \|h(x)\|, \\ R(g, x, \lambda) &= |\lambda| v(x) |g(x)|. \end{aligned}$$

Note first that for any $x \in U$ and $\lambda \in \mathbb{C}$, the function $R_{x,\lambda} : B_{\mathcal{H}_v^\infty(U)} \rightarrow [0, \infty[$, given by

$$R_{x,\lambda}(g) = R(g, x, \lambda),$$

is continuous. For every $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$, we have

$$\begin{aligned} \left(\sum_{i=1}^n S(f, x_i, \lambda_i)^p \right)^{\frac{1}{p}} &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} \\ &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} \\ &= \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n R(g, x_i, \lambda_i)^p \right)^{\frac{1}{p}}, \end{aligned}$$

and therefore f is RS -abstract p -summing. Hence, by applying [20, Theorem 3.1], there is a measure $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ such that

$$S(f, x, \lambda) \leq \pi_p^{\mathcal{H}_v^\infty}(f) \left(\int_{B_{\mathcal{H}_v^\infty(U)}} R(g, x, \lambda)^p d\mu(g) \right)^{\frac{1}{p}}$$

for all $x \in U$ and $\lambda \in \mathbb{C}$, and therefore

$$\|f(x)\| \leq \pi_p^{\mathcal{H}_v^\infty}(f) \left(\int_{B_{\mathcal{H}_v^\infty(U)}} |g(x)|^p d\mu(g) \right)^{\frac{1}{p}}$$

for all $x \in U$.

(ii) \Rightarrow (i): Given $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$, we have

$$\begin{aligned} \sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p &\leq C^p \sum_{i=1}^n \int_{B_{\mathcal{H}_v^\infty(U)}} |\lambda_i|^p v(x_i)^p |g(x_i)|^p d\mu(g) \\ &= C^p \int_{B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right) d\mu(g) \\ &\leq C^p \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right). \end{aligned}$$

Hence $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_p^{\mathcal{H}_v^\infty}(f) \leq C$, and thus $\pi_p^{\mathcal{H}_v^\infty}(f) \leq \inf\{C \geq 0 \text{ satisfying (ii)}\}$. \square

1.3. Pietsch factorization. We will now characterize the class of all p -summing weighted holomorphic mappings through the so-called Pietsch factorization (see [21, Theorem 3], also [10, Theorem 2.13]).

Towards this end, we will use the following mappings. Given $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ and $1 \leq p < \infty$, $I_{\infty,p}: L_\infty(\mu) \rightarrow L_p(\mu)$ and $j_\infty: C(B_{\mathcal{H}_v^\infty(U)}) \rightarrow L_\infty(\mu)$ denote the formal inclusion operators. Consider also the mapping $\iota_U: U \rightarrow C(B_{\mathcal{H}_v^\infty(U)})$ defined by

$$\iota_U(x)(g) = g(x) \quad (x \in U, g \in B_{\mathcal{H}_v^\infty(U)}),$$

and the isometric linear embedding $\kappa_F: F \rightarrow \ell_\infty(B_{F^*})$ given by

$$\langle \kappa_F(y), y^* \rangle = y^*(y) \quad (y^* \in B_{F^*}, y \in F).$$

We now construct the following useful mapping.

Lemma 1.6. *Let $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$. Then $j_\infty \circ \iota_U \in \Pi_p^{\mathcal{H}_v^\infty}(U, L_\infty(\mu))$ with $\pi_p^{\mathcal{H}_v^\infty}(j_\infty \circ \iota_U) \leq 1$ for any $1 \leq p < \infty$.*

Proof. Clearly, $j_\infty \circ \iota_U \in \mathcal{H}(U, L_\infty(\mu))$. Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. By Theorem 0.1, for each $i = 1, \dots, n$, there exists $g_{x_i} \in B_{\mathcal{H}_v^\infty(U)}$ such that $|g_{x_i}(x_i)| = \|\iota_U(x_i)\|_\infty := \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} |g(x_i)|$. Given $1 \leq p < \infty$, one has

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|j_\infty(\iota_U(x_i))\|_{L_\infty(\mu)}^p \right)^{\frac{1}{p}} &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|\iota_U(x_i)\|_\infty^p \right)^{\frac{1}{p}} \\ &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g_{x_i}(x_i)|^p \right)^{\frac{1}{p}} \\ &\leq \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}. \end{aligned}$$

\square

We are now ready to state the announced result.

Theorem 1.7. *Let $1 \leq p < \infty$ and $f \in \mathcal{H}_v^\infty(U, F)$. The following assertions are equivalent:*

(i) *f is p -summing weighted holomorphic.*

(ii) There exist a measure $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty}(U))$, an operator $T \in \mathcal{L}(L_p(\mu), \ell_\infty(B_{F^*}))$ and a mapping $h \in \Pi_p^{\mathcal{H}_v^\infty}(U, L_\infty(\mu))$ such that the following diagram commutes:

$$\begin{array}{ccc} L_\infty(\mu) & \xrightarrow{I_{\infty,p}} & L_p(\mu) \\ h \uparrow & & \downarrow T \\ U & \xrightarrow{f} F \xrightarrow{\kappa_F} & \ell_\infty(B_{F^*}) \end{array}$$

In this case, $\pi_p^{\mathcal{H}_v^\infty}(f) = \inf \{ \|T\| \pi_p^{\mathcal{H}_v^\infty}(h) \}$, where the infimum is taken over all factorizations of $\kappa_F \circ f$ as in (ii), and this infimum is attained.

Proof. (i) \Rightarrow (ii): If $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$, Theorem 1.5 yields a measure $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty}(U))$ such that

$$\|f(x)\| \leq \pi_p^{\mathcal{H}_v^\infty}(f) \left(\int_{B_{\mathcal{H}_v^\infty}(U)} |g(x)|^p d\mu(g) \right)^{\frac{1}{p}} \quad (x \in U).$$

By Lemma 1.6, the mapping $h := j_\infty \circ \iota_U$ is in $\Pi_p^{\mathcal{H}_v^\infty}(U, L_\infty(\mu))$ with $\pi_p^{\mathcal{H}_v^\infty}(h) \leq 1$. Take the space $S_p := \overline{\text{lin}}(I_{\infty,p}(h(U))) \subseteq L_p(\mu)$ and the operator $T_0 \in \mathcal{L}(S_p, \ell_\infty(B_{F^*}))$ defined by

$$T_0(I_{\infty,p}(h(x))) = \kappa_F(f(x)) \quad (x \in U).$$

Using Theorem 0.1 and denoting $I_n = \{i \in \{1, \dots, n\} : \alpha_i \neq 0\}$, note that $\|T_0\| \leq \pi_p^{\mathcal{H}_v^\infty}(f)$ since

$$\begin{aligned}
 \left\| T_0 \left(\sum_{i=1}^n \alpha_i I_{\infty,p}(h(x_i)) \right) \right\|_{\infty} &= \left\| \sum_{i=1}^n \alpha_i T_0(I_{\infty,p}(h(x_i))) \right\|_{\infty} = \left\| \sum_{i=1}^n \alpha_i \kappa_F(f(x_i)) \right\|_{\infty} \\
 &\leq \sum_{i=1}^n |\alpha_i| \|\kappa_F(f(x_i))\|_{\infty} = \sum_{i=1}^n |\alpha_i| \|f(x_i)\| \\
 &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i=1}^n |\alpha_i| \left(\int_{B_{\mathcal{H}_v^\infty(U)}} |g(x_i)|^p d\mu(g) \right)^{\frac{1}{p}} \\
 &= \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i=1}^n |\alpha_i| \left(\int_{B_{\mathcal{H}_v^\infty(U)}} |J_v(g)(\delta_{x_i})|^p d\mu(g) \right)^{\frac{1}{p}} \\
 &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i=1}^n |\alpha_i| \left(\int_{B_{\mathcal{H}_v^\infty(U)}} \|g\|_v^p \|\delta_{x_i}\|^p d\mu(g) \right)^{\frac{1}{p}} \\
 &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i=1}^n |\alpha_i| |g_{x_i}(x_i)| = \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i=1}^n |\alpha_i| g_{x_i}(x_i) \\
 &= \pi_p^{\mathcal{H}_v^\infty}(f) \sum_{i \in I_n} \alpha_i \left(\frac{\bar{\alpha}_i}{|\alpha_i|} g_{x_i} \right)(x_i) = \pi_p^{\mathcal{H}_v^\infty}(f) \left| \sum_{i \in I_n} \alpha_i \left(\frac{\bar{\alpha}_i}{|\alpha_i|} g_{x_i} \right)(x_i) \right| \\
 &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left| \sum_{i=1}^n \alpha_i g(x_i) \right| = \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left| \sum_{i=1}^n \alpha_i \iota_U(x_i)(g) \right| \\
 &= \pi_p^{\mathcal{H}_v^\infty}(f) \left\| \sum_{i=1}^n \alpha_i \iota_U(x_i) \right\|_{\infty} = \pi_p^{\mathcal{H}_v^\infty}(f) \left\| \sum_{i=1}^n \alpha_i h(x_i) \right\|_{L_\infty(\mu)} \\
 &= \pi_p^{\mathcal{H}_v^\infty}(f) \left\| \sum_{i=1}^n \alpha_i I_{\infty,p}(h(x_i)) \right\|_{L_p(\mu)}
 \end{aligned}$$

for any $n \in \mathbb{N}$, $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. Since the Banach space $\ell_\infty(B_{F^*})$ is injective (see [10, p. 45]), there is a $T \in \mathcal{L}(L_p(\mu), \ell_\infty(B_{F^*}))$ such that $T|_{S_p} = T_0$ with $\|T\| = \|T_0\|$. Consequently, $\kappa_F \circ f = T \circ I_{\infty,p} \circ h$ with $\|T\| \pi_p^{\mathcal{H}_v^\infty}(h) \leq \pi_p^{\mathcal{H}_v^\infty}(f)$.

(ii) \Rightarrow (i): We can write $\kappa_F \circ f = T \circ I_{\infty,p} \circ h$ as in (ii). Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. We have

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|\kappa_F(f(x_i))\|_{\infty}^p \right)^{\frac{1}{p}} \\ &= \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|T(I_{\infty,p}(h(x_i)))\|_{\infty}^p \right)^{\frac{1}{p}} \\ &\leq \|T\| \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|I_{\infty,p}(h(x_i))\|_{L_p(\mu)}^p \right)^{\frac{1}{p}} \\ &= \|T\| \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|h(x_i)\|_{L_{\infty}(\mu)}^p \right)^{\frac{1}{p}} \\ &\leq \|T\| \pi_p^{\mathcal{H}_v^{\infty}}(h) \sup_{g \in B_{\mathcal{H}_v^{\infty}}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Hence $f \in \Pi_p^{\mathcal{H}_v^{\infty}}(U, F)$ with $\pi_p^{\mathcal{H}_v^{\infty}}(f) \leq \|T\| \pi_p^{\mathcal{H}_v^{\infty}}(h)$. Taking the infimum over all factorizations of $\kappa_F \circ f$ as in (ii), it follows that $\pi_p^{\mathcal{H}_v^{\infty}}(f) \leq \inf \{ \|T\| \pi_p^{\mathcal{H}_v^{\infty}}(h) \}$. \square

1.4. Inclusion and coincidence. The next result establishes some inclusion and coincidence relations between classes of p -summing weighted holomorphic mappings in terms of p , and also of these spaces with other distinguished spaces of weighted holomorphic mappings.

Proposition 1.8. *If $1 \leq p < q \leq \infty$, then*

$$(\Pi_p^{\mathcal{H}_v^{\infty}}(U, F), \pi_p^{\mathcal{H}_v^{\infty}}) \leq (\Pi_q^{\mathcal{H}_v^{\infty}}(U, F), \pi_q^{\mathcal{H}_v^{\infty}}) \leq (\Pi_{\infty}^{\mathcal{H}_v^{\infty}}(U, F), \pi_{\infty}^{\mathcal{H}_v^{\infty}}) = (\mathcal{H}_v^{\infty}(U, F), \|\cdot\|_v).$$

Proof. Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. We first prove the coincidence. Let $f \in \Pi_{\infty}^{\mathcal{H}_v^{\infty}}(U, F)$. For all $x \in U$, we have

$$v(x) \|f(x)\| \leq \pi_{\infty}^{\mathcal{H}_v^{\infty}}(f) \sup_{g \in B_{\mathcal{H}_v^{\infty}}(U)} v(x) |g(x)| \leq \pi_{\infty}^{\mathcal{H}_v^{\infty}}(f).$$

Hence $f \in \mathcal{H}_v^{\infty}(U, F)$ with $\|f\|_v \leq \pi_{\infty}^{\mathcal{H}_v^{\infty}}(f)$. Conversely, given $f \in \mathcal{H}_v^{\infty}(U, F)$, Theorem 0.1 provides

$$\begin{aligned} |\lambda_i| v(x_i) \|f(x_i)\| &= |\lambda_i| v(x_i) \|T_f(\delta_{x_i})\| \leq \|T_f\| |\lambda_i| v(x_i) \|\delta_{x_i}\| \\ &= \|f\|_v |\lambda_i| v(x_i) \sup_{g \in B_{\mathcal{H}_v^{\infty}}(U)} |g(x_i)| = \|f\|_v \sup_{g \in B_{\mathcal{H}_v^{\infty}}(U)} |\lambda_i| v(x_i) |g(x_i)| \end{aligned}$$

for all $i \in \{1, \dots, n\}$, hence

$$\max_{1 \leq i \leq n} |\lambda_i| v(x_i) \|f(x_i)\| \leq \|f\|_v \sup_{g \in B_{\mathcal{H}_v^{\infty}}(U)} \left(\max_{1 \leq i \leq n} |\lambda_i| v(x_i) |g(x_i)| \right),$$

and so $f \in \Pi_{\infty}^{\mathcal{H}_v^{\infty}}(U, F)$ with $\pi_{\infty}^{\mathcal{H}_v^{\infty}}(f) \leq \|f\|_v$.

We now prove the first inequality. Let $f \in \Pi_p^{\mathcal{H}_v^{\infty}}(U, F)$. Assume first $q < \infty$. Taking

$$\beta_i = |\lambda_i|^{\frac{q}{p}} v(x_i)^{\frac{q}{p}-1} \|f(x_i)\|^{\frac{q}{p}-1} \quad (i = 1, \dots, n),$$

we have

$$\begin{aligned} \left(\sum_{i=1}^n |\lambda_i|^q v(x_i)^q \|f(x_i)\|^q \right)^{\frac{1}{p}} &= \left(\sum_{i=1}^n |\beta_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} \\ &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\beta_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Since $q/p > 1$ and $(q/p)^* = q/(q-p)$, Hölder inequality yields

$$\begin{aligned} \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\beta_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} &= \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n (|\lambda_i| v(x_i) \|f(x_i)\|)^{q-p} (|\lambda_i| v(x_i) |g(x_i)|)^p \right)^{\frac{1}{p}} \\ &\leq \left(\sum_{i=1}^n |\lambda_i|^q v(x_i)^q \|f(x_i)\|^q \right)^{\frac{1}{p} - \frac{1}{q}} \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^q v(x_i)^q |g(x_i)|^q \right)^{\frac{1}{q}}, \end{aligned}$$

and thus we obtain

$$\left(\sum_{i=1}^n |\lambda_i|^q v(x_i)^q \|f(x_i)\|^q \right)^{\frac{1}{q}} \leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^q v(x_i)^q |g(x_i)|^q \right)^{\frac{1}{q}}.$$

This shows that $f \in \Pi_q^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_q^{\mathcal{H}_v^\infty}(f) \leq \pi_p^{\mathcal{H}_v^\infty}(f)$ if $q < \infty$. For $q = \infty$, the same follows from the coincidence and Proposition 1.3. \square

We now present some inclusions between spaces of p -summing weighted holomorphic mappings and some spaces of weighted holomorphic mappings $f: U \rightarrow F$ whose v -range, that is, the set $(vf)(U) := \{v(x)f(x) : x \in U\}$, satisfies some type of compactness (see, for example, [7, 14]).

For $\mathcal{I} = \mathcal{K}, \mathcal{W}$, we will denote by $\mathcal{I}(E, F)$ the linear space of all compact (weakly compact) linear operators from E to F , respectively. We refer to the monograph [21] for a complete study of the ideal structure of such operators.

For $\mathcal{I} = \mathcal{K}, \mathcal{W}$, $\mathcal{H}_{v\mathcal{I}}(U, F)$ represents the linear space of all weighted holomorphic mappings $f: U \rightarrow F$ such that $(vf)(U)$ is relatively compact (relatively weakly compact) in F , respectively. It is known that if $f \in \mathcal{H}_v^\infty(U, F)$, then $f \in \mathcal{H}_{v\mathcal{I}}(U, F)$ if and only if $T_f \in \mathcal{I}(\mathcal{G}_v^\infty(U), F)$ (see, for example, [14, Proposition 5.1] for $\mathcal{I} = \mathcal{K}$, and the same proof works for the case $\mathcal{I} = \mathcal{W}$).

Proposition 1.9. *Let $1 \leq p < \infty$ and let F be a complex Banach space.*

- (i) *Every p -summing weighted holomorphic mapping from U to F has a relatively weakly compact v -range.*
- (ii) *If F is reflexive, then every p -summing weighted holomorphic mapping from U to F has a relatively compact v -range.*

Proof. (i) If $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$, then Theorem 1.7 provides a measure $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty}(U))$, an operator $T \in \mathcal{L}(L_p(\mu), \ell_\infty(B_{F^*}))$ and a map $h \in \Pi_p^{\mathcal{H}_v^\infty}(U, L_\infty(\mu))$ such that $\kappa_F \circ f = T \circ I_{\infty,p} \circ h$. Assume $p > 1$. Since $L_p(\mu)$ is reflexive, then $T \circ I_{\infty,p} \in \mathcal{W}(L_\infty(\mu), \ell_\infty(B_{F^*}))$ by Davis–Figiel–Johnson–Pełczyński Theorem [8]. Since $\kappa_F \circ T_f = T \circ I_{\infty,p} \circ T_h$ by Theorem 0.1 and \mathcal{W} is an injective operator ideal, it follows that $T_f \in \mathcal{W}(\mathcal{G}_v^\infty(U), F)$ and this means that $f \in \mathcal{H}_{v\mathcal{W}}(U, F)$ as required. For $p = 1$, the result follows from Proposition 1.8 and from what was proved above.

(ii) If $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F)$, then $f \in \mathcal{H}_{v\mathcal{W}}(U, F)$ by (i), that is, $T_f \in \mathcal{W}(\mathcal{G}_v^\infty(U), F)$. Since F is reflexive, it follows that $T_f \in \mathcal{K}(\mathcal{G}_v^\infty(U), F)$ and thus $f \in \mathcal{H}_{v\mathcal{K}}(U, F)$. \square

1.5. Maurey extrapolation. We now use Pietsch domination of p -summing weighted holomorphic mappings to give a weighted holomorphic version of Maurey Extrapolation Theorem [16]. The proof of the next result is based on that of [10, Theorem 3.17].

Theorem 1.10. *Let $1 < p < q < \infty$ and assume that $\Pi_q^{\mathcal{H}_v^\infty}(U, \ell_q) = \Pi_p^{\mathcal{H}_v^\infty}(U, \ell_q)$. Then $\Pi_q^{\mathcal{H}_v^\infty}(U, F) = \Pi_1^{\mathcal{H}_v^\infty}(U, F)$ for every complex Banach space F .*

Proof. For each $\mu \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$, we can take $h_\mu := j_\infty \circ \iota_U \in \Pi_p^{\mathcal{H}_v^\infty}(U, L_\infty(\mu))$ with $\pi_p^{\mathcal{H}_v^\infty}(h_\mu) \leq 1$ by Lemma 1.6. Then $I_{\infty,q} \circ h_\mu \in \Pi_q^{\mathcal{H}_v^\infty}(U, L_q(\mu))$ with $\pi_q^{\mathcal{H}_v^\infty}(I_{\infty,q} \circ h_\mu) \leq 1$ by Propositions 1.3 and 1.8.

Since $\Pi_q^{\mathcal{H}_v^\infty}(U, \ell_q) = \Pi_p^{\mathcal{H}_v^\infty}(U, \ell_q)$ and $\pi_q^{\mathcal{H}_v^\infty} \leq \pi_p^{\mathcal{H}_v^\infty}$ on $\Pi_p^{\mathcal{H}_v^\infty}(U, \ell_q)$ by Proposition 1.8, the Closed Graph Theorem yields a constant $C \geq 0$ such that $\pi_p^{\mathcal{H}_v^\infty}(f) \leq C\pi_q^{\mathcal{H}_v^\infty}(f)$ for all $f \in \Pi_q^{\mathcal{H}_v^\infty}(U, \ell_q)$. Since $L_q(\mu)$ is an $\mathcal{L}_{q,\lambda}$ -space for each $\lambda > 1$, we can assure that given $n \in \mathbb{N}$ and $x_1, \dots, x_n \in U$, the subspace

$$G_0 = \text{lin} \left(\{I_{\infty,q}(h_\mu(x_1)), \dots, I_{\infty,q}(h_\mu(x_n))\} \right) \subseteq L_q(\mu)$$

embeds λ -isomorphically into ℓ_q , that is, G_0 is contained in a subspace $G \subseteq L_q(\mu)$ for which there exists an isomorphism $T: G \rightarrow \ell_q$ with $\|T\| \|T^{-1}\| < \lambda$.

Since $T \circ I_{\infty,q} \circ h_\mu \in \Pi_q^{\mathcal{H}_v^\infty}(U, \ell_q) = \Pi_p^{\mathcal{H}_v^\infty}(U, \ell_q)$, we have

$$\begin{aligned} & \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \left\| (I_{\infty,q} \circ h_\mu)(x_i) \right\|_{L_q(\mu)}^p \right)^{\frac{1}{p}} \leq \|T^{-1}\| \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \left\| T(I_{\infty,q}(h_\mu(x_i))) \right\|_{\ell_q}^p \right)^{\frac{1}{p}} \\ & \leq \|T^{-1}\| C \pi_q^{\mathcal{H}_v^\infty}(T \circ I_{\infty,q} \circ h_\mu) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} \\ & \leq C \|T^{-1}\| \|T\| \pi_q^{\mathcal{H}_v^\infty}(I_{\infty,q} \circ h_\mu) \sup_{g \in B_{\mathcal{H}_v^\infty(U)}} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

therefore $\pi_p^{\mathcal{H}_v^\infty}(I_{\infty,q} \circ h_\mu) \leq C\lambda$ for all $\lambda > 1$, and thus $\pi_p^{\mathcal{H}_v^\infty}(I_{\infty,q} \circ h_\mu) \leq C$. Now, by Theorem 1.5, there exists a measure $\widehat{\mu} \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ such that

$$\left\| (I_{\infty,q} \circ h_\mu)(x) \right\|_{L_q(\mu)} \leq C \left(\int_{B_{\mathcal{H}_v^\infty(U)}} |g(x)|^p d\widehat{\mu}(g) \right)^{\frac{1}{p}} = C \left\| (I_{\infty,q} \circ h_{\widehat{\mu}})(x) \right\|_{L_p(\widehat{\mu})}$$

for all $x \in U$. In the last equality, we have used that

$$(I_{\infty,q} \circ h_{\widehat{\mu}})(x)(g) = I_{\infty,q}(h_{\widehat{\mu}}(x))(g) = h_{\widehat{\mu}}(x)(g) = j_\infty(\iota_U(x))(g) = \iota_U(x)(g) = g(x)$$

for all $x \in U$ and $g \in B_{\mathcal{H}_v^\infty(U)}$.

Take a complex Banach space F and let $f \in \Pi_q^{\mathcal{H}_v^\infty}(U, F)$. In view of Proposition 1.8, we only must show that $f \in \Pi_1^{\mathcal{H}_v^\infty}(U, F)$. Theorem 1.5 provides again a measure $\mu_0 \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ such that

$$\|f(x)\| \leq \pi_q^{\mathcal{H}_v^\infty}(f) \left\| (I_{\infty,q} \circ h_{\mu_0})(x) \right\|_{L_q(\mu_0)}$$

for all $x \in U$. We claim that there is a constant $c \geq 0$ and a measure $\lambda_0 \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ such that

$$\left\| (I_{\infty,q} \circ h_{\mu_0})(x) \right\|_{L_q(\mu_0)} \leq c \left\| (I_{\infty,q} \circ h_{\lambda_0})(x) \right\|_{L_1(\lambda_0)}$$

for all $x \in U$. Indeed, define $\lambda_0 = \sum_{n=0}^{\infty} (1/2^{n+1}) \mu_n \in \mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$, where $(\mu_n)_{n \geq 1}$ is the sequence in $\mathcal{P}(B_{\mathcal{H}_v^\infty(U)})$ given by $\mu_{n+1} = \widehat{\mu}_n$ for all $n \in \mathbb{N}_0$, where the measure $\widehat{\mu}_n$ is defined using Theorem 1.5.

As $1 < p < q$, then $\theta := (q-p)/p(q-1) \in (0, 1)$ and $1/p = \theta + (1-\theta)/q$, and applying Littlewood's Inequality we have

$$\begin{aligned} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_p(\mu_n)} &= \left(\int_{B_{\mathcal{H}_V^\infty(U)}} |(I_{\infty,q} \circ h_{\mu_n})(x)(g)|^p d\mu_n(g) \right)^{\frac{1}{p}} \\ &\leq \left(\int_{B_{\mathcal{H}_V^\infty(U)}} |(I_{\infty,q} \circ h_{\mu_n})(x)(g)| d\mu_n(g) \right)^\theta \left(\int_{B_{\mathcal{H}_V^\infty(U)}} |(I_{\infty,q} \circ h_{\mu_n})(x)(g)|^q d\mu_n(g) \right)^{\frac{1-\theta}{q}} \\ &= \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_1(\mu_n)}^\theta \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)}^{1-\theta} \end{aligned}$$

for each $n \in \mathbb{N}_0$ and all $x \in U$. Using Hölder's Inequality and the inequality

$$\begin{aligned} &\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_q(\mu_{n+1})} \\ &\leq \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_q(\mu_{n+1})} + \|(I_{\infty,q} \circ h_{\mu_0})(x)\|_{L_q(\mu_0)} \\ &= \sum_{n=-1}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_q(\mu_{n+1})} \\ &= \sum_{n=0}^{\infty} \frac{1}{2^n} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)} \\ &= 2 \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)}, \end{aligned}$$

we now obtain

$$\begin{aligned} &\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)} \\ &\leq C \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_p(\mu_{n+1})} \\ &\leq C \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_1(\mu_{n+1})}^\theta \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_q(\mu_{n+1})}^{1-\theta} \\ &\leq C \left(\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_1(\mu_{n+1})} \right)^\theta \left(\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_q(\mu_{n+1})} \right)^{1-\theta} \\ &\leq C \left(\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_1(\mu_{n+1})} \right)^\theta \left(2 \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)} \right)^{1-\theta} \end{aligned}$$

for all $x \in U$, and we deduce that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_q(\mu_n)} &\leq C^{\frac{1}{\theta}} 2^{\frac{1-\theta}{\theta}} \left(\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_{n+1}})(x)\|_{L_1(\mu_{n+1})} \right) \\ &\leq C^{\frac{1}{\theta}} 2^{\frac{1-\theta}{\theta}} 2 \left(\sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \|(I_{\infty,q} \circ h_{\mu_n})(x)\|_{L_1(\mu_n)} \right) \\ &= (2C)^{\frac{1}{\theta}} \|(I_{\infty,q} \circ h_{\lambda_0})(x)\|_{L_1(\lambda_0)} \end{aligned}$$

for all $x \in U$. From above, we deduce that

$$\frac{1}{2} \|(I_{\infty,q} \circ h_{\mu_0})(x)\|_{L_q(\mu_0)} \leq (2C)^{\frac{1}{\theta}} \|(I_{\infty,q} \circ h_{\lambda_0})(x)\|_{L_1(\lambda_0)}$$

for all $x \in U$, and this proves our claim taking $c = 2(2C)^{\frac{1}{\theta}}$. Therefore we can write

$$\|f(x)\| \leq c\pi_q^{\mathcal{H}_v^\infty}(f) \|(I_{\infty,q} \circ h_{\lambda_0})(x)\|_{L_1(\lambda_0)} = c\pi_q^{\mathcal{H}_v^\infty}(f) \int_{B_{\mathcal{H}_v^\infty(U)}} |g(x)| \, d\lambda_0(g)$$

for all $x \in U$. Hence $f \in \Pi_1^{\mathcal{H}_v^\infty}(U, F)$ with $\pi_1^{\mathcal{H}_v^\infty}(f) \leq c\pi_q^{\mathcal{H}_v^\infty}(f)$ by Theorem 1.5. \square

2. BANACH-VALUED \mathcal{H}_v^∞ -MOLECULES

Our main objective in this section is to study the duality of the spaces of p -summing weighted holomorphic mappings. With this purpose, we will first establish some results and introduce some concepts influenced by the theory on tensor product (see the monograph [22]).

Proposition 2.1. *Let $x \in U$ and $y \in F$. Then the function $v(x)\delta_x \otimes y: \mathcal{H}_v^\infty(U, F^*) \rightarrow \mathbb{C}$ defined by*

$$(v(x)\delta_x \otimes y)(f) = \langle v(x)f(x), y \rangle \quad (f \in \mathcal{H}_v^\infty(U, F^*)),$$

belongs to $\mathcal{H}_v^\infty(U, F^)^*$ and $\|v(x)\delta_x \otimes y\| = v(x) \|\delta_x\| \|y\|$.*

Proof. Clearly, $v(x)\delta_x \otimes y$ is linear. For any $f \in \mathcal{H}_v^\infty(U, F^*)$, we have

$$\begin{aligned} |(v(x)\delta_x \otimes y)(f)| &= |\langle v(x)f(x), y \rangle| \leq v(x) \|f(x)\| \|y\| = v(x) \|T_f(\delta_x)\| \|y\| \\ &\leq v(x) \|T_f\| \|\delta_x\| \|y\| = \|f\|_v v(x) \|\delta_x\| \|y\|, \end{aligned}$$

and thus $v(x)\delta_x \otimes y \in \mathcal{H}_v^\infty(U, F^*)^*$ with $\|v(x)\delta_x \otimes y\| \leq v(x) \|\delta_x\| \|y\|$. For the converse inequality, take $y^* \in B_{F^*}$ such that $|\langle y^*, y \rangle| = \|y\|$ and consider $g_x \cdot y^* \in \mathcal{H}_v^\infty(U, F^*)$. Since $\|g_x \cdot y^*\|_v = \|g_x\|_v \|y^*\| \leq 1$, we deduce that

$$\begin{aligned} \|v(x)\delta_x \otimes y\| &\geq |(v(x)\delta_x \otimes y)(g_x \cdot y^*)| = v(x) |\langle (g_x \cdot y^*)(x), y \rangle| \\ &= v(x) |\langle g_x(x)y^*, y \rangle| = v(x) |g_x(x)| |\langle y^*, y \rangle| = v(x) \|\delta_x\| \|y\|. \end{aligned}$$

\square

The elements of the following tensor product space could be referred to as F -valued \mathcal{H}_v^∞ -molecules on U .

Definition 2.2. Let E and F be complex Banach spaces, let U be an open subset of E and let v be a weight on U . Define the linear space

$$\text{lin}((v\Delta_v)(U)) \otimes F := \text{lin} \{v(x)\delta_x \otimes y : x \in U, y \in F\} \subseteq \mathcal{H}_v^\infty(U, F^*)^*.$$

Each element $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$ can be expressed in a form (not necessarily unique):

$$\gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i,$$

where $n \in \mathbb{N}$, $\lambda_i \in \mathbb{C}$, $x_i \in U$ and $y_i \in F$ for $i = 1, \dots, n$; and its action as a functional on a mapping $f \in \mathcal{H}_v^\infty(U, F^*)$ comes given by

$$\gamma(f) = \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle.$$

2.1. Projective norm. Given two linear spaces E and F , the tensor product space $E \otimes F$, equipped with a norm α , is usually denoted by $E \otimes_\alpha F$, and the completion of $E \otimes_\alpha F$ by $E \widehat{\otimes}_\alpha F$. Consider the projective norm π on $u \in E \otimes F$, defined by

$$\pi(u) = \inf \left\{ \sum_{i=1}^n \|x_i\| \|y_i\| : n \in \mathbb{N}, x_1, \dots, x_n \in E, y_1, \dots, y_n \in F, u = \sum_{i=1}^n x_i \otimes y_i \right\},$$

where the infimum is taken over all such representations of u .

We now show that the projective norm and the operator canonical norm coincide on the space of F -valued \mathcal{H}_v^∞ -molecules on U .

Proposition 2.3. *Let $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$. Then $\|\gamma\| = \pi(\gamma)$, where*

$$\|\gamma\| = \sup \{ |\gamma(f)| : f \in \mathcal{H}_v^\infty(U, F^*), \|f\|_v \leq 1 \}$$

and

$$\pi(\gamma) = \inf \left\{ \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| : \gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \right\}.$$

Proof. Let $\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i$ be a representation of γ . Since $\gamma \in \mathcal{H}_v^\infty(U, F^*)^*$ and

$$\begin{aligned} |\gamma(f)| &= \left| \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle \right| \leq \sum_{i=1}^n |\lambda_i| v(x_i) \|f(x_i)\| \|y_i\| \\ &= \sum_{i=1}^n |\lambda_i| v(x_i) \|T_f(\delta_{x_i})\| \|y_i\| \leq \|f\|_v \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| \end{aligned}$$

for all $f \in \mathcal{H}_v^\infty(U, F^*)$, we have that $\|\gamma\| \leq \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\|$. Since this holds for each representation of γ as above, it is deduced that $\|\gamma\| \leq \pi(\gamma)$.

To prove that $\pi(\gamma) \leq \|\gamma\|$, suppose by contradiction that $\|\gamma\| < \pi(\gamma)$. Note that $\gamma \neq 0$ and consider the set $B = \{\mu \in \text{lin}((v\Delta_v)(U)) \otimes F : \pi(\mu) \leq \|\gamma\|\}$. Clearly, B is a closed convex subset of $\text{lin}((v\Delta_v)(U)) \otimes_\pi F$. Applying the Hahn–Banach Separation Theorem to B and $\{\gamma\}$, we can take a functional $\eta \in (\text{lin}((v\Delta_v)(U)) \otimes_\pi F)^*$ with $\|\eta\| = 1$ such that

$$\|\gamma\| = \sup\{\text{Re}(\eta(\mu)) : \mu \in B\} < \text{Re}(\eta(\gamma)).$$

Define $f_\eta : U \rightarrow F^*$ by

$$\langle f_\eta(x), y \rangle = \eta(\delta_x \otimes y) \quad (x \in U, y \in F).$$

We now show that f_η is holomorphic. By [17, Exercise 8.D], it suffices to prove that for each $y \in F$, the function $f_{\eta,y} : U \rightarrow \mathbb{C}$, defined by

$$f_{\eta,y}(x) = \eta(\delta_x \otimes y) \quad (x \in U),$$

is holomorphic. For it, let $a \in U$ and since $\Delta_v: U \rightarrow \text{lin}((v\Delta_v)(U))$ is holomorphic, there exists $D\Delta_v(a) \in \mathcal{L}(E, \text{lin}((v\Delta_v)(U)))$ such that

$$\lim_{x \rightarrow a} \frac{\delta_x - \delta_a - D\Delta_v(a)(x - a)}{\|x - a\|} = 0.$$

Define the function $T(a): E \rightarrow \mathbb{C}$ by

$$T(a)(x) = \eta(D\Delta_v(a)(x) \otimes y) \quad (x \in E).$$

Clearly, $T(a)$ is linear and

$$\begin{aligned} |T(a)(x)| &= |\eta(D\Delta_v(a)(x) \otimes y)| \leq \|\eta\| \pi(D\Delta_v(a)(x) \otimes y) \\ &\leq \|D\Delta_v(a)(x)\| \|y\| \leq \|D\Delta_v(a)\| \|x\| \|y\| \end{aligned}$$

for all $x \in E$, hence $T(a) \in E^*$. Since

$$\begin{aligned} f_{\eta,y}(x) - f_{\eta,y}(a) - T(a)(x - a) &= \eta(\delta_x \otimes y) - \eta(\delta_a \otimes y) - \eta(D\Delta_v(a)(x - a) \otimes y) \\ &= \eta((\delta_x - \delta_a - D\Delta_v(a)(x - a)) \otimes y) \end{aligned}$$

for all $x \in U$, it follows that

$$\lim_{x \rightarrow a} \frac{f_{\eta,y}(x) - f_{\eta,y}(a) - T(a)(x - a)}{\|x - a\|} = \lim_{x \rightarrow a} \eta \left(\frac{\delta_x - \delta_a - D\Delta_v(a)(x - a)}{\|x - a\|} \otimes y \right) = 0.$$

Thus $f_{\eta,y}$ is holomorphic at a with $Df_{\eta,y}(a) = T(a)$, as required.

Given $x \in U$, we have

$$v(x)|\langle f_{\eta}(x), y \rangle| = v(x) |\eta(\delta_x \otimes y)| \leq v(x) \|\eta\| \pi(\delta_x \otimes y) \leq v(x) \|\delta_x\| \|y\| \leq \|y\|$$

for all $y \in F$, and so $v(x) \|f_{\eta}(x)\| \leq 1$. Therefore $f_{\eta} \in \mathcal{H}_v^{\infty}(U, F^*)$ and $\|f_{\eta}\|_v \leq 1$. Furthermore, $\mu(f_{\eta}) = \eta(\mu)$ for all $\mu \in \text{lin}((v\Delta_v)(U)) \otimes F$. Then $\|\gamma\| \geq |\gamma(f_{\eta})| \geq \text{Re}(\gamma(f_{\eta})) = \text{Re}(\eta(\gamma))$, and we arrive at a contradiction. \square

2.2. p -Chevet–Saphar \mathcal{H}_v^{∞} -norms. The p -Chevet–Saphar norms d_p on the tensor product of two Banach spaces $E \otimes F$ are well known (see, for example, [22, Section 6.2]).

Our study of the duality of the spaces of p -summing weighted holomorphic mappings requires the introduction of the following \mathcal{H}_v^{∞} -variants of such norms.

Definition 2.4. Let E and F be complex Banach spaces, let U be an open subset of E , let v be a weight on U and let $1 \leq p \leq \infty$. For $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$, define:

$$\begin{aligned} d_1^{\mathcal{H}_v^{\infty}}(\gamma) &= \inf \left\{ \left(\sup_{g \in B_{\mathcal{H}_v^{\infty}(U)}} \left(\max_{1 \leq i \leq n} |\lambda_i| v(x_i) |g(x_i)| \right) \right) \left(\sum_{i=1}^n \|y_i\| \right) \right\}, \\ d_p^{\mathcal{H}_v^{\infty}}(\gamma) &= \inf \left\{ \left(\sup_{g \in B_{\mathcal{H}_v^{\infty}(U)}} \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}} \right\} \quad (1 < p < \infty), \\ d_{\infty}^{\mathcal{H}_v^{\infty}}(\gamma) &= \inf \left\{ \left(\sup_{g \in B_{\mathcal{H}_v^{\infty}(U)}} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \right) \right) \left(\max_{1 \leq i \leq n} \|y_i\| \right) \right\}, \end{aligned}$$

where the infimum is taken over all such representations of γ as $\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i$.

Motivated by the analogue concept for tensor product spaces, we introduce the following.

Definition 2.5. A norm α on $\text{lin}((v\Delta_v)(U)) \otimes F$ is said to be a reasonable \mathcal{H}_v^{∞} -crossnorm if it enjoys the following properties:

- (i) $\alpha(v(x)\delta_x \otimes y) = v(x) \|\delta_x\| \|y\|$ for all $x \in U$ and $y \in F$,

- (ii) For every $g \in \mathcal{H}_v^\infty(U)$ and $y^* \in F^*$, the linear functional $g \otimes y^* : \text{lin}((v\Delta_v)(U)) \otimes F \rightarrow \mathbb{C}$ defined by $(g \otimes y^*)(v(x)\delta_x \otimes y) = v(x)g(x)y^*(y)$ is bounded on $\text{lin}((v\Delta_v)(U)) \otimes_\alpha F$ with $\|g \otimes y^*\| \leq \|g\|_v \|y^*\|$.

Theorem 2.6. $d_p^{\mathcal{H}_v^\infty}$ is a reasonable \mathcal{H}_v^∞ -crossnorm on $\text{lin}((v\Delta_v)(U)) \otimes F$ for any $1 \leq p \leq \infty$.

Proof. We will only prove it for $1 < p < \infty$. The other cases follow similarly.

Let $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$ and let $\sum_{i=1}^n \lambda_i v(x_i)\delta_{x_i} \otimes y_i$ be a representation of γ . Clearly, $d_p^{\mathcal{H}_v^\infty}(\gamma) \geq 0$. Given $\lambda \in \mathbb{C}$, since $\sum_{i=1}^n (\lambda \lambda_i) v(x_i)\delta_{x_i} \otimes y_i$ is a representation of $\lambda\gamma$, we have

$$\begin{aligned} d_p^{\mathcal{H}_v^\infty}(\lambda\gamma) &\leq \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda \lambda_i|^{p^*} v(x_i)^{p^*} |g(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}} \\ &= |\lambda| \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}}. \end{aligned}$$

If $\lambda = 0$, we obtain $d_p^{\mathcal{H}_v^\infty}(\lambda\gamma) = 0 = |\lambda| d_p^{\mathcal{H}_v^\infty}(\gamma)$. For $\lambda \neq 0$, since the preceding inequality holds for every representation of γ , we deduce that $d_p^{\mathcal{H}_v^\infty}(\lambda\gamma) \leq |\lambda| d_p^{\mathcal{H}_v^\infty}(\gamma)$. For the converse inequality, note that $d_p^{\mathcal{H}_v^\infty}(\gamma) = d_p^{\mathcal{H}_v^\infty}(\lambda^{-1}(\lambda\gamma)) \leq |\lambda^{-1}| d_p^{\mathcal{H}_v^\infty}(\lambda\gamma)$ by using the proved inequality, thus $|\lambda| d_p^{\mathcal{H}_v^\infty}(\gamma) \leq d_p^{\mathcal{H}_v^\infty}(\lambda\gamma)$ and hence $d_p^{\mathcal{H}_v^\infty}(\lambda\gamma) = |\lambda| d_p^{\mathcal{H}_v^\infty}(\gamma)$.

We now prove the triangular inequality of $d_p^{\mathcal{H}_v^\infty}$. Let $\gamma_1, \gamma_2 \in \text{lin}((v\Delta_v)(U)) \otimes F$ and let $\varepsilon > 0$. If $\gamma_1 = 0$ or $\gamma_2 = 0$, there is nothing to prove. Assume $\gamma_1 \neq 0 \neq \gamma_2$. We can choose representations

$$\gamma_1 = \sum_{i=1}^n \lambda_{1,i} v(x_{1,i})\delta_{x_{1,i}} \otimes y_{1,i}, \quad \gamma_2 = \sum_{i=1}^m \lambda_{2,i} v(x_{2,i})\delta_{x_{2,i}} \otimes y_{2,i},$$

so that

$$\left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_{1,i}|^{p^*} v(x_{1,i})^{p^*} |g(x_{1,i})|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^n \|y_{1,i}\|^p \right)^{\frac{1}{p}} \leq d_p^{\mathcal{H}_v^\infty}(\gamma_1) + \varepsilon$$

and

$$\left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^m |\lambda_{2,i}|^{p^*} v(x_{2,i})^{p^*} |g(x_{2,i})|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^m \|y_{2,i}\|^p \right)^{\frac{1}{p}} \leq d_p^{\mathcal{H}_v^\infty}(\gamma_2) + \varepsilon.$$

Fix arbitrary $r, s \in \mathbb{R}^+$ and define

$$\begin{aligned} \lambda_{3,i} v(x_{3,i})\delta_{x_{3,i}} &= \begin{cases} r^{-1} \lambda_{1,i} v(x_{1,i})\delta_{x_{1,i}} & \text{if } i = 1, \dots, n, \\ s^{-1} \lambda_{2,i-n} v(x_{2,i-n})\delta_{x_{2,i-n}} & \text{if } i = n+1, \dots, n+m, \end{cases} \\ y_{3,i} &= \begin{cases} r y_{1,i} & \text{if } i = 1, \dots, n, \\ s y_{2,i-n} & \text{if } i = n+1, \dots, n+m. \end{cases} \end{aligned}$$

It is clear that $\gamma_1 + \gamma_2 = \sum_{i=1}^{n+m} \lambda_{3,i} v(x_{3,i})\delta_{x_{3,i}} \otimes y_{3,i}$ and thus we have

$$d_p^{\mathcal{H}_v^\infty}(\gamma_1 + \gamma_2) \leq \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^{n+m} |\lambda_{3,i}|^{p^*} v(x_{3,i})^{p^*} |g(x_{3,i})|^{p^*} \right)^{\frac{1}{p^*}} \right) \left(\sum_{i=1}^{n+m} \|y_{3,i}\|^p \right)^{\frac{1}{p}}.$$

An easy verification gives

$$\begin{aligned} & \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^{n+m} |\lambda_{3,i}|^{p^*} v(x_{3,i})^{p^*} |g(x_{3,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} \\ & \leq \left(r^{-1} \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_{1,i}|^{p^*} v(x_{1,i})^{p^*} |g(x_{1,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} + \left(s^{-1} \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^m |\lambda_{2,i}|^{p^*} v(x_{2,i})^{p^*} |g(x_{2,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} \end{aligned}$$

and

$$\sum_{i=1}^{n+m} \|y_{3,i}\|^p = r^p \sum_{i=1}^n \|y_{1,i}\|^p + s^p \sum_{i=1}^m \|y_{2,i}\|^p.$$

Using Young's Inequality, it follows that

$$\begin{aligned} d_p^{\mathcal{H}_v^\infty}(\gamma_1 + \gamma_2) & \leq \frac{1}{p^*} \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^{n+m} |\lambda_{3,i}|^{p^*} v(x_{3,i})^{p^*} |g(x_{3,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} + \frac{1}{p} \sum_{i=1}^{n+m} \|y_{3,i}\|^p \\ & \leq \frac{r^{-p^*}}{p^*} \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_{1,i}|^{p^*} v(x_{1,i})^{p^*} |g(x_{1,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} + \frac{r^p}{p} \sum_{i=1}^n \|y_{1,i}\|^p \\ & \quad + \frac{s^{-p^*}}{p^*} \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^m |\lambda_{2,i}|^{p^*} v(x_{2,i})^{p^*} |g(x_{2,i})|^{p^*} \right)^{\frac{1}{p^*}} \right)^{p^*} + \frac{s^p}{p} \sum_{i=1}^m \|y_{2,i}\|^p. \end{aligned}$$

Since r, s were arbitrary in \mathbb{R}^+ , taking above

$$\begin{aligned} r & = (d_p^{\mathcal{H}_v^\infty}(\gamma_1) + \varepsilon)^{-\frac{1}{p^*}} \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_{1,i}|^{p^*} v(x_{1,i})^{p^*} |g(x_{1,i})|^{p^*} \right)^{\frac{1}{p^*}} \right), \\ s & = (d_p^{\mathcal{H}_v^\infty}(\gamma_2) + \varepsilon)^{-\frac{1}{p^*}} \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^m |\lambda_{2,i}|^{p^*} v(x_{2,i})^{p^*} |g(x_{2,i})|^{p^*} \right)^{\frac{1}{p^*}} \right), \end{aligned}$$

we obtain that $d_p^{\mathcal{H}_v^\infty}(\gamma_1 + \gamma_2) \leq d_p^{\mathcal{H}_v^\infty}(\gamma_1) + d_p^{\mathcal{H}_v^\infty}(\gamma_2) + 2\varepsilon$, and thus $d_p^{\mathcal{H}_v^\infty}(\gamma_1 + \gamma_2) \leq d_p^{\mathcal{H}_v^\infty}(\gamma_1) + d_p^{\mathcal{H}_v^\infty}(\gamma_2)$ by the arbitrariness of ε . Hence $d_p^{\mathcal{H}_v^\infty}$ is a seminorm. To prove that it is a norm, note first that

$$\begin{aligned} \pi(\gamma) & \leq \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| \\ & = \sum_{i=1}^n |\lambda_i| v(x_i) |g_{x_i}(x_i)| \|y_i\| \\ & \leq \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g_{x_i}(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}} \\ & \leq \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}}, \end{aligned}$$

by applying Hölder's Inequality, and therefore $\pi(\gamma) \leq d_p^{\mathcal{H}_v^\infty}(\gamma)$ by taking the infimum over all representations of γ . Now, if $d_p^{\mathcal{H}_v^\infty}(\gamma) = 0$, then $\pi(\gamma) = 0$ and thus $\gamma = 0$ by Proposition 2.3.

Finally, we show that $d_p^{\mathcal{H}_v^\infty}$ is a reasonable \mathcal{H}_v^∞ -crossnorm on $\text{lin}((v\Delta_v)(U)) \otimes F$. First, given $x \in U$ and $y \in F$, we have

$$v(x) \|\delta_x\| \|y\| = \|v(x)\delta_x \otimes y\| = \pi(v(x)\delta_x \otimes y) \leq d_p^{\mathcal{H}_v^\infty}(v(x)\delta_x \otimes y)$$

by Propositions 2.1 and 2.3 and by proved above, and, conversely, one has

$$d_p^{\mathcal{H}_v^\infty}(v(x)\delta_x \otimes y) \leq \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} v(x) |g(x)| \right) \|y\| = v(x) \|\delta_x\| \|y\|.$$

Second, given $g \in \mathcal{H}_v^\infty(U)$ and $y^* \in F^*$,

$$\begin{aligned} |(g \otimes y^*)(\gamma)| &= \left| \sum_{i=1}^n \lambda_i (g \otimes y^*)(v(x_i)\delta_{x_i} \otimes y_i) \right| = \left| \sum_{i=1}^n \lambda_i v(x_i) g(x_i) y^*(y_i) \right| \\ &\leq \sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| |y^*(y_i)| = \sum_{i=1}^n |\lambda_i| v(x_i) |J_v(g)(\delta_{x_i})| |y^*(y_i)| \\ &\leq \|g\|_v \|y^*\| \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| = \|g\|_v \|y^*\| \sum_{i=1}^n |\lambda_i| v(x_i) |g_{x_i}(x_i)| \|y_i\| \\ &\leq \|g\|_v \|y^*\| \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g_{x_i}(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}} \\ &\leq \|g\|_v \|y^*\| \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^{p^*} v(x_i)^{p^*} |g(x_i)|^{p^*} \right)^{\frac{1}{p^*}} \left(\sum_{i=1}^n \|y_i\|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Taking infimum over all the representations of γ , we deduce that $|(g \otimes y^*)(\gamma)| \leq \|g\|_v \|y^*\| d_p^{\mathcal{H}_v^\infty}(\gamma)$. Hence $g \otimes y^* \in (\text{lin}((v\Delta_v)(U)) \otimes_{d_p^{\mathcal{H}_v^\infty}} F)^*$ with $\|g \otimes y^*\| \leq \|g\|_v \|y^*\|$. \square

We now compute $d_p^{\mathcal{H}_v^\infty}$ with a simpler formula for $p = 1$ and $p = \infty$. In fact, the 1-Chevet–Saphar \mathcal{H}_v^∞ -norm coincides with the projective norm.

Proposition 2.7. *For $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$, we have*

$$d_1^{\mathcal{H}_v^\infty}(\gamma) = \inf \left\{ \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| \right\}$$

and

$$d_\infty^{\mathcal{H}_v^\infty}(\gamma) = \inf \left\{ \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \|y_i\| \right) \right\},$$

where the infimum is taken over all such representations of γ as $\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i$.

Proof. Let $\gamma \in \text{lin}((v\Delta_v)(U)) \otimes F$ and let $\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i$ be a representation of γ . We have

$$\begin{aligned} \pi(\gamma) &\leq \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\| = \sum_{i=1}^n |\lambda_i| v(x_i) \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} |g(x_i)| \right) \|y_i\| \\ &\leq \sum_{i=1}^n \max_{1 \leq i \leq n} \left(|\lambda_i| v(x_i) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} |g(x_i)| \right) \|y_i\| = \left(\max_{1 \leq i \leq n} \left(|\lambda_i| v(x_i) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} |g(x_i)| \right) \right) \left(\sum_{i=1}^n \|y_i\| \right) \\ &= \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\max_{1 \leq i \leq n} |\lambda_i| v(x_i) |g(x_i)| \right) \right) \left(\sum_{i=1}^n \|y_i\| \right) \end{aligned}$$

and therefore $\pi(\gamma) \leq d_1^{\mathcal{H}_v^\infty}(\gamma)$. Conversely, since $d_1^{\mathcal{H}_v^\infty}$ is a crossnorm, we have

$$d_1^{\mathcal{H}_v^\infty}(\gamma) \leq \sum_{i=1}^n |\lambda_i| v(x_i) d_1^{\mathcal{H}_v^\infty}(\delta_{x_i} \otimes y_i) = \sum_{i=1}^n |\lambda_i| v(x_i) \|\delta_{x_i}\| \|y_i\|,$$

and thus $d_1^{\mathcal{H}_v^\infty}(\gamma) \leq \pi(\gamma)$.

On the other hand, we have

$$\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \|y_i\| \right) \leq \left(\max_{1 \leq i \leq n} \|y_i\| \right) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \right),$$

and taking the infimum over all representations of γ gives

$$\inf \left\{ \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \|y_i\| \right) : \gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \right\} \leq d_\infty^{\mathcal{H}_v^\infty}(\gamma).$$

Conversely, we can assume without loss of generality that $y_i \neq 0$ for all $i \in \{1, \dots, n\}$ and since $\gamma = \sum_{i=1}^n \lambda_i v(x_i) \|y_i\| \delta_{x_i} \otimes (y_i / \|y_i\|)$, we obtain

$$d_\infty^{\mathcal{H}_v^\infty}(\gamma) \leq \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) \|y_i\| |g(x_i)| \right),$$

and taking the infimum over all representations of γ , we conclude that

$$d_\infty^{\mathcal{H}_v^\infty}(\gamma) \leq \inf \left\{ \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i| v(x_i) |g(x_i)| \|y_i\| \right) : \gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \right\}.$$

□

2.3. Duality. Given $p \in [1, \infty]$, we will show that the dual of the space $\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_p^{\mathcal{H}_v^\infty}} F$ can be canonically identified as the space of p -summing weighted holomorphic mappings from U to F^* .

Theorem 2.8. *For $1 \leq p \leq \infty$, the space $(\Pi_p^{\mathcal{H}_v^\infty}(U, F^*), \pi_p^{\mathcal{H}_v^\infty})$ is isometrically isomorphic to $(\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_p^{\mathcal{H}_v^\infty}} F)^*$, via the mapping $\Lambda: \Pi_p^{\mathcal{H}_v^\infty}(U, F^*) \rightarrow (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_p^{\mathcal{H}_v^\infty}} F)^*$ defined by*

$$\Lambda(f)(\gamma) = \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle$$

for $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$ and $\gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \in \text{lin}((v\Delta_v)(U)) \otimes F$. Furthermore, its inverse comes given by

$$\langle \Lambda^{-1}(\varphi)(x), y \rangle = \varphi(\delta_x \otimes y)$$

for $\varphi \in (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F)^*$, $x \in U$ and $y \in F$.

Proof. We prove it for $1 < p < \infty$. The cases $p = 1$ and $p = \infty$ can be proved similarly.

Let $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$ and let $\Lambda_0(f): \text{lin}((v\Delta_v)(U)) \otimes F \rightarrow \mathbb{C}$ be the linear functional given by

$$\Lambda_0(f)(\gamma) = \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle$$

for $\gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \in \text{lin}((v\Delta_v)(U)) \otimes F$. Note that $\Lambda_0(f) \in (\text{lin}((v\Delta_v)(U)) \otimes_{d_{p^*}^{\mathcal{H}_v^\infty}} F)^*$ with $\|\Lambda_0(f)\| \leq \pi_p^{\mathcal{H}_v^\infty}(f)$ since

$$\begin{aligned} |\Lambda_0(f)(\gamma)| &= \left| \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle \right| \leq \sum_{i=1}^n |\lambda_i| v(x_i) \|f(x_i)\| \|y_i\| \\ &\leq \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f(x_i)\|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n \|y_i\|^{p^*} \right)^{\frac{1}{p^*}} \\ &\leq \pi_p^{\mathcal{H}_v^\infty}(f) \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n \|y_i\|^{p^*} \right)^{\frac{1}{p^*}}, \end{aligned}$$

and taking infimum over all the representations of γ , we deduce that $|\Lambda_0(f)(\gamma)| \leq \pi_p^{\mathcal{H}_v^\infty}(f) d_{p^*}^{\mathcal{H}_v^\infty}(\gamma)$. Since γ was arbitrary, then $\Lambda_0(f)$ is continuous on $\text{lin}((v\Delta_v)(U)) \otimes_{d_{p^*}^{\mathcal{H}_v^\infty}} F$ with $\|\Lambda_0(f)\| \leq \pi_p^{\mathcal{H}_v^\infty}(f)$.

Hence there is a unique continuous function $\Lambda(f): \text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F \rightarrow \mathbb{C}$ that extends $\Lambda_0(f)$.

Further, $\Lambda(f)$ is linear and $\|\Lambda(f)\| = \|\Lambda_0(f)\|$. Let $\Lambda: \Pi_p^{\mathcal{H}_v^\infty}(U, F^*) \rightarrow (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F)^*$ be the mapping so defined.

Clearly, $\Lambda_0: \Pi_p^{\mathcal{H}_v^\infty}(U, F^*) \rightarrow (\text{lin}((v\Delta_v)(U)) \otimes_{d_{p^*}^{\mathcal{H}_v^\infty}} F)^*$ is linear. To show its injectivity, if $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$ and $\Lambda_0(f) = 0$, then $v(x) \langle f(x), y \rangle = \Lambda_0(f)(v(x) \delta_x \otimes y) = 0$ for all $x \in U$ and $y \in F$, hence $f(x) = 0$ for all $x \in U$ and thus $f = 0$. Then Λ is also linear and injective. Indeed, let $\phi \in \text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F$ and let $\{\gamma_n\}$ be a sequence in $\text{lin}((v\Delta_v)(U)) \otimes_{d_{p^*}^{\mathcal{H}_v^\infty}} F$ such that $d_{p^*}^{\mathcal{H}_v^\infty}(\gamma_n - \phi) \rightarrow 0$ when $n \rightarrow \infty$. Given $a, b \in \mathbb{C}$ and $f, g \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$, an easy calculation shows that

$$\Lambda(af + bg)(\gamma_n) = \Lambda_0(af + bg)(\gamma_n) = (a\Lambda_0(f) + b\Lambda_0(g))(\gamma_n) = (a\Lambda(f) + b\Lambda(g))(\gamma_n)$$

for all $n \in \mathbb{N}$, and taking limits with $n \rightarrow \infty$, we have that $\Lambda(af + bg)(\phi) = (a\Lambda(f) + b\Lambda(g))(\phi)$. Hence Λ is linear. For the injectivity of Λ , note that if $f \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$ and $\Lambda(f) = 0$, then $\Lambda_0(f) = 0$ which implies that $f = 0$ by the injectivity of Λ_0 .

To prove that Λ is a surjective isometry, let $\varphi \in (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F)^*$ and define $f_\varphi: U \rightarrow F^*$ by

$$\langle f_\varphi(x), y \rangle = \varphi(\delta_x \otimes y) \quad (x \in U, y \in F).$$

As in the proof of Proposition 2.3, it is similarly proved that $f_\varphi \in \mathcal{H}_v^\infty(U, F^*)$ with $\|f_\varphi\|_v \leq \|\varphi\|$.

We now prove that $f_\varphi \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$. Fix $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and $x_1, \dots, x_n \in U$. Let $\varepsilon > 0$. For each $i \in \{1, \dots, n\}$, there exists $y_i \in F$ with $\|y_i\| \leq 1 + \varepsilon$ such that $\langle f_\varphi(x_i), y_i \rangle = \|f_\varphi(x_i)\|$. Clearly, the function $T: \mathbb{C}^n \rightarrow \mathbb{C}$, defined by

$$T(t_1, \dots, t_n) = \sum_{i=1}^n t_i \lambda_i v(x_i) \|f_\varphi(x_i)\|, \quad \forall (t_1, \dots, t_n) \in \mathbb{C}^n,$$

is linear and continuous on $(\mathbb{C}^n, \|\cdot\|_{p^*})$ with $\|T\| = \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f_\varphi(x_i)\|^p\right)^{\frac{1}{p}}$. For any $(t_1, \dots, t_n) \in \mathbb{C}^n$ with $\|(t_1, \dots, t_n)\|_{p^*} \leq 1$, we have

$$\begin{aligned} |T(t_1, \dots, t_n)| &= \left| \varphi \left(\sum_{i=1}^n t_i \lambda_i v(x_i) \delta_{x_i} \otimes y_i \right) \right| \leq \|\varphi\| d_{p^*}^{\mathcal{H}_v^\infty} \left(\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes t_i y_i \right) \\ &\leq \|\varphi\| \left(\sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}} \right) \left(\sum_{i=1}^n \|t_i y_i\|^{p^*} \right)^{\frac{1}{p^*}} \\ &\leq (1 + \varepsilon) \|\varphi\| \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}}, \end{aligned}$$

therefore

$$\left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f_\varphi(x_i)\|^p \right)^{\frac{1}{p}} \leq (1 + \varepsilon) \|\varphi\| \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}},$$

and since ε was arbitrary, we have

$$\left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p \|f_\varphi(x_i)\|^p \right)^{\frac{1}{p}} \leq \|\varphi\| \sup_{g \in B_{\mathcal{H}_v^\infty}(U)} \left(\sum_{i=1}^n |\lambda_i|^p v(x_i)^p |g(x_i)|^p \right)^{\frac{1}{p}},$$

and we conclude that $f_\varphi \in \Pi_p^{\mathcal{H}_v^\infty}(U, F^*)$ with $\pi_p^{\mathcal{H}_v^\infty}(f_\varphi) \leq \|\varphi\|$.

Finally, for any $\gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \in \text{lin}((v\Delta_v)(U)) \otimes F$, we get

$$\Lambda(f_\varphi)(\gamma) = \sum_{i=1}^n \lambda_i v(x_i) \langle f_\varphi(x_i), y_i \rangle = \sum_{i=1}^n \lambda_i v(x_i) \varphi(\delta_{x_i} \otimes y_i) = \varphi \left(\sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \right) = \varphi(\gamma).$$

Hence $\Lambda(f_\varphi) = \varphi$ on a dense subspace of $\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{d_{p^*}^{\mathcal{H}_v^\infty}} F$ and we conclude that $\Lambda(f_\varphi) = \varphi$.

This also justifies the last statement of the theorem. Moreover, $\pi_p^{\mathcal{H}_v^\infty}(f_\varphi) \leq \|\varphi\| = \|\Lambda(f_\varphi)\|$. \square

In light of Theorem 2.8 and taking into account Propositions 1.8, 2.3 and 2.7, we can identify the space $\mathcal{H}_v^\infty(U, F^*)$ with the dual space of $\text{lin}((v\Delta_v)(U)) \widehat{\otimes} F \subseteq \mathcal{H}_v^\infty(U, F^*)^*$.

Corollary 2.9. *The space $(\mathcal{H}_v^\infty(U, F^*), \|\cdot\|_v)$ is isometrically isomorphic to $(\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{\|\cdot\|} F)^*$, via the mapping $\Lambda: \mathcal{H}_v^\infty(U, F^*) \rightarrow (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{\|\cdot\|} F)^*$ given by*

$$\Lambda(f)(\gamma) = \sum_{i=1}^n \lambda_i v(x_i) \langle f(x_i), y_i \rangle$$

for $f \in \mathcal{H}_v^\infty(U, F^*)$ and $\gamma = \sum_{i=1}^n \lambda_i v(x_i) \delta_{x_i} \otimes y_i \in \text{lin}((v\Delta_v)(U)) \otimes F$, with inverse

$$\langle \Lambda^{-1}(\varphi)(x), y \rangle = \varphi(\delta_x \otimes y)$$

for $\varphi \in (\text{lin}((v\Delta_v)(U)) \widehat{\otimes}_{\|\cdot\|} F)^*$, $x \in U$ and $y \in F$. \square

REFERENCES

- [1] D. Achour and L. Mezrag, On the Cohen strongly p -summing multilinear operators, *J. Math. Anal. Appl.* **327** (2007), no. 1, 550–563.
- [2] J. C. Angulo-López and M. Fernández-Unzueta, Lipschitz p -summing multilinear operators, *J. Funct. Anal.* **279** (2020), no. 4, 108572, 20 pp.
- [3] K. D. Bierstedt and W. H. Summers, Biduals of weighted Banach spaces of analytic functions, *J. Austral. Math. Soc. Ser. A* **54** (1993), no. 1, 70–79.
- [4] J. Bonet, Weighted Banach spaces of analytic functions with sup-norms and operators between them: a survey, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* (2022) 116:184.
- [5] J. Bonet, P. Domanski and M. Lindström, Weakly Compact Composition Operators on Weighted Vector-Valued Banach Spaces of Analytic Mappings, *Ann. Acad. Sci. Fenn. Ser. A I. Math.* **26** (2001), 233–248.
- [6] J. Bonet and M. Friz, Weakly Compact Composition Operators on Locally Convex Spaces, *Math. Nachr.* **245** (2002), 26–44.
- [7] M. G. Cabrera-Padilla, A. Jiménez-Vargas and A. Keten Çopur, Weighted holomorphic mappings associated with p -compact type sets, <https://arxiv.org/submit/5814632>, to appear in *Bull. Malays. Math. Sci. Soc.*
- [8] W. J. Davis, T. Figiel, W. B. Johnson and A. Pełczyński, Factoring weakly compact operators, *J. Funct. Anal.* **17** (1974), 311–327.
- [9] A. Defant and K. Floret, Tensor norms and operator ideals. North-Holland Mathematics Studies, vol. 176, North-Holland Publishing Co., Amsterdam, 1993.
- [10] J. Diestel, H. Jarchow and A. Tonge, Absolutely summing operators, Cambridge Studies in Advanced Mathematics, vol. 43, Cambridge University Press, Cambridge, 1995.
- [11] V. Dimant, Strongly p -summing multilinear mappings, *J. Math. Anal. Appl.* **278** (2003) 182–193.
- [12] J. D. Farmer and W. B. Johnson, Lipschitz p -summing operators, *Proc. Amer. Math. Soc.* **137** (2009), no. 9, 2989–2995.
- [13] A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, *Memoirs American Mathematical Society* 16, Providence, Rhode Island 1955.
- [14] M. Gupta and D. Baweja, Weighted spaces of holomorphic functions on Banach spaces and the approximation property, *Extracta Math.* **31** (2016), no. 2, 123–144.
- [15] M. C. Matos, Absolutely summing holomorphic mappings, *An. Acad. Brasil. Cienc.* **68** (1996) 1–13.
- [16] B. Maurey, Théorèmes de factorisation pour les opérateurs linéaires à valeurs dans les espaces L_p , *Soc. Math. France, Asterisque* 11, Paris, 1974.
- [17] J. Mujica, *Complex Analysis in Banach spaces*, Dover Publications, 2010.
- [18] D. Pellegrino, Strongly almost summing holomorphic mappings, *J. Math. Anal. Appl.* **287** (2003), no. 1, 244–252.
- [19] D. Pellegrino, P. Rueda and E. A. Sánchez-Pérez, Surveying the spirit of absolute summability on multilinear operators and homogeneous polynomials, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **110** (2016), no. 1, 285–302.
- [20] D. Pellegrino and J. Santos, A general Pietsch Domination Theorem, *J. Math. Anal. Appl.* **375** (2011), 371–374.
- [21] A. Pietsch, Absolut p -summierende Abbildungen in normierten Räumen, *Studia Math.* **28** (1967), 333–353.
- [22] R. A. Ryan, *Introduction to tensor products of Banach spaces*, Series: Springer Monographs in Mathematics, Springer, 2002.
- [23] K. Saadi, Some properties for Lipschitz strongly p -summing operators, *J. Math. Anal. Appl.* **423** (2015) 1410–1426.

(M. G. Cabrera-Padilla) DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE ALMERÍA, 04120 ALMERÍA, SPAIN.
Email address: m_gador@hotmail.com

(A. Jiménez-Vargas) DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE ALMERÍA, 04120, ALMERÍA, SPAIN.
Email address: ajimenez@ual.es

(A. Keten Çopur) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, FACULTY OF SCIENCES, NECMETTIN ERBAKAN UNIVERSITY, KONYA, 42090, TÜRKIYE.
Email address: aketen@erbakan.edu.tr