# Categories of skew left braces and trifactorised groups

A. Ballester-Bolinches<sup>\*</sup> P. Pérez-Altarriba<sup>\*</sup> V. Pérez-Calabuig<sup>\*</sup>

#### Abstract

The main objective of this paper is to deepen the relationship between skew left braces and trifactorised groups that encodes the information about skew left braces, their structure, their quotients, and their homomorphisms.

Keywords: category, skew left brace, trifactorised group Mathematics Subject Classification (2020): 16T25, 81R50, 20C35, 20C99, 20D40,

# 1 Introduction

A skew left brace  $(B, +, \cdot)$  consists of a set with two binary operations + and  $\cdot$  such that (B, +) and  $(B, \cdot)$  are groups linked by a distributive-like law a(b + c) = ab - a + ac for  $a, b, c \in B$ . As it is usual in this context, the multiplication symbol is omitted, -a denotes the symmetric element of awith respect to +, and a - b denotes a + (-b). When the operations on B are clear from the context, we will refer to a brace as B instead of  $(B, +, \cdot)$ . Skew left braces were introduced in [11] and generalise the *left braces* introduced in [12], in which the first operation is also assumed to be commutative. Skew left braces are interesting algebraic structures, because they represent the exact algebraic framework to study bijective non-degenerate solutions to the Yang-Baxter equation (see [4, 9, 14]). For simplicity, we will use the word *brace* with the meaning of *skew left brace*.

<sup>\*</sup>Departament de Matemàtiques, Universitat de València, Dr. Moliner, 50, 46100 Burjassot, València, Spain; Adolfo.Ballester@uv.es, Ramon.Esteban@uv.es, Pedro.A.Perez@uv.es, Vicent.Perez-Calabuig@uv.es; ORCID 0000-0002-2051-9075, 0000-0002-2321-8139, 0009-0009-7082-9002, 0000-0003-4101-8656

One of the most important tools in the structural study of braces is the analysis of the trifactorised group associated with the action of the multiplicative group on the additive group. Many brace structural properties can be translated into group structural properties of this trifactorised group which is called here *large trifactorised group* (see [2]). Another trifactorised group associated with a brace B appears in a natural way when identifying B with a regular subgroup of the holomorph of the additive group. This group, called here *small trifactorised group*, is extremely useful to construct braces with some prescribed properties (see [3]). Small trifactorised groups associated with particular braces lead to construction of interesting examples of groups that can be used to answer some important questions about products of groups (see [1, Chapter 6]).

The aim of this paper is twofold. Firstly, we study the complete family of trifactorised groups associated with a given brace and secondly, we study braces and their associated trifactorised groups from a categorical point of view. Many of the problems that we discuss in this paper appear in a natural way when considering representations of braces. The following two remarkable results are proved.

- 1. There is a bijection between the isomorphism classes of the trifactorised groups associated with a brace B and the orbits of an action of the automorphism group of B over some normal subgroups of the multiplicative group. This result will be useful for computing the trifactorised groups associated with B (Theorem 4.7).
- 2. All trifactorised groups associated with a brace B are quotients of the large trifactorised group associated with B and have the small trifactorised group associated with B as a quotient (Theorem 6.11).

We also consider two main categories: the category **SKB** whose objects are braces, and whose morphisms are homomorphisms of braces, that are maps between braces that preserve both operations, and the category **3factGrp** whose objects are trifactorised groups satisfying some restrictions, and the morphisms are group homomorphisms which are compatible with the triple factorisation. We show here that these two categories are closely related and have interesting properties. It is worth mentioning that some properties of **SKB** have been studied in [5].

The contents of the paper are organised as follows. We will begin by introducing two categories in Section 2, corresponding to the large and small trifactorised groups. Then we will introduce **3factGrp** in Section 3, where we will show that for every brace we can define trifactorised groups satisfying

certain conditions and that every trifactorised group with these conditions can be understood as a trifactorised group associated with a brace. We also show how to construct brace homomorphisms from trifactorised group morphisms, and give the exact condition for a brace homomorphism to define a trifactorised group morphism. It turns out that the categories of small and large trifactorised groups are subcategories of **3factGrp**. Our next objective, developed in Section 5, is to identify the substructures of a brace B in a trifactorised group associated with B, and study the substructures of trifactorised groups. We will see that there is a bijection between subbraces and these substructures. We also indicate how to use the trifactorised group to compute images of subgroups of the multiplicative group and preimages of subgroups of the additive group under the associated derivation with the help of the trifactorised group. In Section 6, we characterise the normal subgroups of our trifactorised groups giving quotients that can be regarded as trifactorised groups, and show that they are associated with the corresponding brace quotient.

The efficiency of algorithms for finite groups in computer algebra systems like GAP [10], especially in the case of soluble groups and, in, to a lesser extent, in the case of permutation groups, makes us think that encoding braces as trifactorised groups can help us to simplify computations with braces. Trifactorised groups introduced here are, in general, of smaller order than the groups given in [2]. This can be used to save memory. The saving when using the small trifactorised groups is very relevant in the case of trivial braces, where the small trifactorised group coincides with the additive group.

We also would like to mention that trifactorised groups will be in a forthcoming paper to present a definition of representations of braces over a vector space. We will see there that these brace representations are equivalent to some representations of trifactorised groups over a vector space. It turns out that trifactorised groups are a natural tool to reduce some problems on brace representations to problems on group representations.

# 2 Large and small trifactorised groups

Given a brace B, there is an action of the multiplicative group  $C = (B, \cdot)$ on the additive group K = (B, +) called the *lambda action* or the *lambda* map which is defined by  $\lambda(a) = \lambda_a$  for every  $a \in B$ , where  $\lambda_a(b) = -a + ab$ for all  $a, b \in B$ . Sysak [13] observed that the identity map  $\delta: C \longrightarrow K$  is a bijective derivation or 1-cocycle with respect to the lambda action and that the semidirect product G = [K]C with respect to the lambda action possesses a triple factorisation G = KC = KD = CD, where  $D = \{\delta(c)c \mid c \in C\}$  is a subgroup of G, and  $K \cap C = K \cap D = C \cap D = 1$ . We will refer to it as the *large* trifactorised group associated with B, and denoted by L(B) = (G, K, D, C).

It seems natural to consider the category **L3factGrp** whose objects are trifactorised groups of the form (G, K, H, C) where  $K \leq G, H \leq G, C \leq G,$  $G = KH = KC = HC, K \cap H = K \cap C = H \cap C = 1$ , and whose morphisms  $f: (G_1, K_1, H_1, C_1) \longrightarrow (G_2, K_2, H_2, C_2)$  are group homomorphisms  $f: G_1 \longrightarrow G_2$  such that  $f(K_1) \leq K_2, f(H_1) \leq H_1$ , and  $f(C_1) \leq C_2$ . If  $f: B_1 \longrightarrow B_2$  is a brace homomorphism and  $\mathsf{L}(B_1) = (G_1, K_1, H_1, C_1)$  and  $\mathsf{L}(B_2) = (G_2, K_2, H_2, C_2)$  are the large trifactorised groups corresponding to  $B_1$  and  $B_2$ , respectively, then the induced group homomorphisms  $K_1 \longrightarrow K_2$ and  $C_1 \longrightarrow C_2$  induce a group homomorphism  $G_1 \longrightarrow G_2$  and, given an element  $(\delta(c), c) \in H_1$ , where  $\delta: C_1 \longrightarrow K_1$  is the identity map of  $B_1$ , regarded as a bijective derivation associated with the lambda map, we have that  $(f(\delta(c)), f(c)) = (\bar{\delta}(f(c)), f(c)) \in H_2$ , where  $\bar{\delta}: C_2 \longrightarrow K_2$  is the bijective derivation associated with  $B_2$ . If we call this homomorphism  $\mathsf{L}(f)$ , we have that  $\mathsf{L}: \mathbf{SKB} \longrightarrow \mathbf{L3factGrp}$  becomes a functor.

Another natural approach to braces is by means of regular subgroups of the holomorph of the additive group. When this additive group corresponds to a vector space, we have regular subgroups of the affine group, analysed in [6–8].

Let B be a brace and let K = (B, +). By [11, Theorem 4.2],  $H = \{(a, \lambda_a) \mid a \in B\}$  is a regular subgroup of Hol(K) that is isomorphic to  $(B, \cdot)$ . If we consider the subgroup  $S = KH \leq \text{Hol}(K)$ , it follows that S = KH = KE = HE, where  $E = \{(0, \lambda_b) \mid b \in B\}$  and  $C_E(K) = K \cap E = H \cap E = 1$ . We call S(B) = (S, K, H, E) the small trifactorised group associated with B.

As in the case of large trifactorised groups, we can consider the category **S3factGrp** of small trifactorised groups. The objects of this category are small trifactorised groups (G, K, H, E) with G a group,  $K \leq G$ ,  $H \leq G$ ,  $E \leq G$ , G = KH = KE = HE,  $K \cap E = H \cap E = 1$ , and  $C_E(K) = 1$ . As in **L3factGrp**, a morphism  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  of **S3factGrp** is a group homomorphism  $f: G_1 \longrightarrow G_2$  such that  $f(K_1) \leq K_2$ ,  $f(H_1) \leq H_2$ , and  $f(E_1) \leq E_2$ . However, in general, we cannot associate to a brace homomorphism  $f: B_1 \longrightarrow B_2$ , where  $S(B_1) = (G_1, K_1, H_1, E_1)$ ,  $S(B_2) = (G_2, K_2, H_2, E_2)$ , a morphism  $\bar{f}$  of **S3factGrp** such that  $\bar{f}|_{K_1}$  coincides with the group homomorphism  $K_1 \longrightarrow K_2$  induced by f.

**Example 2.1.** Let  $B_1$  be a trivial brace of order 2 and let  $B_2$  be a brace with additive group isomorphic to the alternating group  $A_5$  of degree 5 and in which the multiplication is the operation opposite to +. We have that  $S(B_1) = (G_1, K_1, H_1, E_1)$  with  $G_1 = H_1 = K_1$  cyclic of order 2 and  $E_1 = 1$ , and  $S(B_2) = (G_2, K_2, H_2, E_2)$  with  $K_2 \cong A_5$ ,  $H_2 = \{(t, \alpha_t^{-1}) \mid t \in A_5\} \cong A_5$ ,  $E_2 = \text{Inn}(A_5)$  and  $G_2 = K_2H_2 \cong A_5 \times A_5$ . It is clear that  $f: B_1 \longrightarrow B_2$ given by f(a) = (1, 2)(3, 4), where a is the generator of the additive group of  $B_1$ , defines a brace homomorphism, but there is no group homomorphism  $f: G_1 \longrightarrow G_2$  such that  $f(K_1) \leq K_2$ ,  $f(H_1) \leq H_2$  and  $f(E_1) \leq E_2$  with  $f(a) = (1, 2)(3, 4) \in K_2$ , because  $f(a) \in f(K_1 \cap H_1) \leq K_2 \cap H_2 = 1$ .

# 3 A new category of trifactorised groups

Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Let  $\lambda: C \longrightarrow \operatorname{Aut}(K)$  be the corresponding lambda map and  $\delta: C \longrightarrow K$  the identity map. Consider  $\eta: C \longrightarrow E$  a group epimorphism such that ker  $\eta \leq \ker \lambda$ . Then we can define the group homomorphism  $\overline{\lambda}: E \longrightarrow \operatorname{Aut}(K)$  such that  $\overline{\lambda}(\eta(c)) = \lambda(c)$  for all  $c \in C$ . Hence E acts on K by automorphisms, and so we can consider the corresponding semidirect product G = [K]E.

Consider the map from C to G defined by  $c \mapsto \delta(c)\eta(c)$ , which is injective with image  $H = \{\delta(c)\eta(c) \mid c \in C\}$ . Moreover, given  $c_1, c_2 \in C$ ,

$$\begin{aligned} (\delta(c_1)\eta(c_1))(\delta(c_2)\eta(c_2)) &= \delta(c_1)\lambda_{\eta(c_1)}(\delta(c_2))\eta(c_1)\eta(c_2) \\ &= \delta(c_1)\lambda_{c_1}(\delta(c_2))\eta(c_1)\eta(c_2) \\ &= \delta(c_1c_2)\eta(c_1c_2). \end{aligned}$$

Therefore H is a subgroup of G isomorphic to C.

Every element  $g \in G$  can be uniquely written as a product  $g = \delta(c_1)\eta(c_2)$ for  $c_1, c_2 \in C$ , and

$$G = KE = KH = HE, \quad K \cap E = H \cap E = 1. \tag{1}$$

The map  $\sigma: H \longrightarrow K$  given by  $\sigma(\delta(c)\eta(c)) = \delta(c)$  is bijective and

$$\sigma((\delta(c_1)\eta(c_1))(\delta(c_2)\eta(c_2))) = \sigma(\delta(c_1)\lambda_{c_1}(\delta(c_2))\eta(c_1c_2))$$
  
=  $\delta(c_1)\lambda_{c_1}(\delta(c_2))$   
=  $\sigma(\delta(c_1)\eta(c_1))\overline{\lambda}_{\eta(c_1)}(\sigma(\delta(c_2)\eta(c_2)))$ 

Therefore,  $\sigma$  is a bijective derivation with respect to the action  $\overline{\lambda} \circ \pi_E|_H \colon H \longrightarrow \operatorname{Aut}(K)$ , where  $\pi_E \colon G \longrightarrow E$  is the natural projection.

**Definition 3.1.** The tuple (G, K, H, E) is called a *(generalised) trifactorised* group associated with  $(B, +, \cdot)$ .

We define the category **3factGrp** of (generalised) trifactorised groups, that has as objects tuples (G, K, H, E) where G is a group, K is a normal

subgroup of G, H and E are subgroups of G, and they satisfy (1), and as morphisms  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  group homomorphisms  $f: G_1 \longrightarrow G_2$  such that  $f(K_1) \leq K_2$ ,  $f(H_1) \leq H_2$ , and  $f(E_1) \leq E_2$ . This category has as subcategories **L3factGrp** and **S3factGrp**. From now on when we talk about a trifactorised group we will mean a tuple in this category.

We have shown above that there are different trifactorised groups associated with a brace. We will prove now that the converse is also true: given a trifactorised group (G, K, H, E), we can associate a brace and a bijective derivation  $\sigma: H \longrightarrow K$ .

**Proposition 3.2.** Let (G, K, H, E) be a trifactorised group. Every element  $g \in G$  can be uniquely written as a product  $g = k_g e_g$ , where  $k_g \in K$  and  $e_g \in E$ . Then:

- 1. H acts on K via the action  $\tilde{\lambda}$  given by  $\tilde{\lambda}(h)(k) = e_h k = e_h k e_h^{-1}$  for  $h \in H, k \in K$ .
- 2. The map  $\sigma: H \longrightarrow K$  given by  $\sigma(h) = k_h$  for  $h \in H$  is a bijective derivation with respect to  $\tilde{\lambda}$ .
- 3. If we define the operation  $\boxdot$  on K as  $k_1 \boxdot k_2 = k_1({}^{e_{k_1}}k_2)$  for  $k_1, k_2 \in K$ , it follows that  $(K, \cdot, \boxdot)$  is a skew left brace, with  $(K, \boxdot) \cong H$  and lambda map  $\tilde{\lambda} \circ \sigma^{-1} \colon (K, \boxdot) \longrightarrow \operatorname{Aut}(K)$ .

Proof. It is easy to check that  $e_{h_1h_2} = e_{h_1}e_{h_2}$  for  $h_1, h_2 \in H$ . This implies that  $\tilde{\lambda} \colon H \longrightarrow \operatorname{Aut}(K)$  is an action of H on K. Given  $k \in K$ , there exists  $e \in E$  and  $h \in H$  such that k = he, hence  $ke^{-1} \in H$ . Let us suppose that there exist  $e_1, e_2 \in E$  such that  $ke_1, ke_2 \in H$ , then  $e_1^{-1}e_2 \in H \cap E = 1$ . Consequently,  $\sigma \colon H \longrightarrow K$  is bijective. Furthermore, if  $h_1 = k_1e_1, h_2 = k_2e_2 \in H$  with  $k_1, k_2 \in K$  and  $e_1, e_2 \in E$ , we have that  $\sigma(h_1h_2) = \sigma(k_1(e_1k_1)e_1e_2) = k_1(e_1k_1) = \sigma(h_1)\tilde{\lambda}(h_1)(\sigma(h_2))$ . That is,  $\sigma$  is a bijective derivation with respect to  $\tilde{\lambda}$ .

By [11, Proposition 1.11], we can define a second operation on K such that K becomes a brace, which is  $k_1 \circ k_2 = \sigma(\sigma^{-1}(k_1)\sigma^{-1}(k_2)) = k_1 \boxdot k_2$ . It is obvious that  $\sigma^{-1} \colon (K, \boxdot) \longrightarrow H$  is a group isomorphism. The lambda map is  $\lambda_{k_1}(k_2) = k_1^{-1}(k_1 \boxdot k_2) = {}^{e_{k_1}}k_2 = \tilde{\lambda}(k_1e_{k_1})(k_2) = (\tilde{\lambda} \circ \sigma^{-1}(k_1))(k_2)$ .

**Definition 3.3.** Given a trifactorised group (G, K, H, E), the brace defined in Proposition 3.2 is called *the associated brace* of (G, K, H, E) and it is denoted by B(G, K, H, E).

**Proposition 3.4.** 1. Every trifactorised group (G, K, H, E) is a trifactorised group associated with the brace B(G, K, H, E).

- 2. Let  $(B, +, \cdot)$  be a brace, and let (G, K, H, E) be a trifactorised group associated with  $(B, +, \cdot)$ , then  $B(G, K, H, E) = (B, +, \cdot)$ .
- Proof. 1. Following the notation of Proposition 3.2 and denoting  $C = (K, \boxdot)$ , we can consider  $\eta = \pi_E|_H \circ \sigma^{-1} \colon C \longrightarrow E$ , where  $\pi_E \colon G \longrightarrow E$  is the natural projection. We have that  $\eta$  is a group epimorphism, because  $\sigma^{-1}$  is a group isomorphism by the argument of the last paragraph of the proof of Proposition 3.4 and  $\pi_E$  is a group epimorphism. Moreover,  $\ker \eta = \sigma(\ker \pi_E \cap H) = \sigma(K \cap H) = K \cap H$ . If  $k \in \ker \eta$ , then  $\lambda(k) = \tilde{\lambda}(k) = \operatorname{id}$ , therefore  $\ker \eta \leq \ker \lambda$ . This epimorphism gives us a trifactorised group associated with  $\mathsf{B}(G, K, H, E)$ , which is exactly (G, K, H, E).
  - 2. Let K and C be the additive group and the multiplicative group of B respectively and let  $\lambda: C \longrightarrow \operatorname{Aut}(K)$  be the lambda map of  $(B, +, \cdot)$ . Write  $\delta: C \longrightarrow K$  the identity map. Consider  $\eta: C \longrightarrow E$  the epimorphism that corresponds to the associated trifactorised group (G, K, H, E). The brace  $\mathsf{B}(G, K, H, E)$ , as a set, is B, and its additive operation is just the additive operation of  $(B, +, \cdot)$ . Furthermore

$$\delta(c_1) \boxdot \delta(c_2) = \delta(c_1)({}^{\eta(c_1)}\delta(c_2)) = \delta(c_1)\lambda_{c_1}(\delta(c_2)) = \delta(c_1c_2).$$

This means that  $(B, \boxdot) = (B, \cdot)$ .

#### 3.1 Morphisms of trifactorised groups

**Definition 3.5.** A morphism  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  between two trifactorised groups is a group homomorphism  $f: G_1 \longrightarrow G_2$  such that  $f(K_1) \leq K_2, f(H_1) \leq H_2$ , and  $f(E_1) \leq E_2$ .

These conditions gives us three group homomorphisms, namely  $f|_{K_1} \colon K_1 \longrightarrow K_2$ ,  $f|_{H_1} \colon H_1 \longrightarrow H_2$ ,  $f|_{E_1} \colon E_1 \longrightarrow E_2$ , which are the restrictions of f as group homomorphism. We prove that  $f|_{K_1}$  and  $f|_{H_1}$  are closely related.

**Lemma 3.6.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a trifactorised group morphism. Let  $\sigma_i: H_i \longrightarrow K_i$  for i = 1, 2 be the corresponding bijective derivations. Then

- 1.  $\sigma_2 \circ f|_{H_1} = f|_{K_1} \circ \sigma_1$ .
- 2. Given  $h_1, h_2 \in H_1$ ,  $f(h_1) = f(h_2)$  if and only if  $f(\sigma_1(h_1)) = f(\sigma_1(h_2))$ .

- *Proof.* 1. We use the notation on Proposition 3.2. Statement (1) follows directly from the fact the f(h) is the product  $f(k_h)f(e_h)$ , where  $f(k_h) \in K_2$ ,  $f(e_h) \in E_2$ , for all  $h \in H$ . Therefore,  $\sigma_2(f(h)) = f(k_h) = f(\sigma_1(h))$ , for all  $h \in H$ .
  - 2. We have that  $f(h_1) = f(h_2)$  if and only if  $\sigma_2(f(h_1)) = \sigma_2(f(h_2))$ , equivalently,  $f(\sigma_1(h_1)) = f(\sigma_1(h_2))$ .

Furthermore, the injectivity, surjectivity and bijectivity of  $f|_{K_1}$ ,  $f|_{H_1}$ ,  $f|_{E_1}$ , and f are also closely related.

**Proposition 3.7.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a morphism, then:

- 1. If f is a monomorphism (resp., epimorphism, isomorphism) then  $f|_{K_1}$ ,  $f|_{H_1}$ , and  $f|_{E_1}$  are monomorphisms (resp., epimorphisms, isomorphisms).
- 2.  $f|_{K_1}$  is a monomorphism (resp., epimorphism, isomorphism) if and only if  $f|_{H_1}$  is a monomorphism (resp., epimorphism, isomorphism).
- 3. If  $f|_{K_1}$  is an epimorphism, then f is a epimorphism.
- 4. If  $f|_{K_1}$  and  $f|_{E_1}$  are monomorphisms, then f is a monomorphism.
- 5. If  $f|_{K_1}$  and  $f|_{E_1}$  are isomorphisms, then f is a isomorphism.
- *Proof.* 1. If f is injective obviously its restrictions are injective, so we only need to prove it for surjectivity and the result for bijectivity will follow. Suppose that f is an epimorphism and let  $k \in K_2$ . There exists  $g \in G_1$  such that f(g) = k. Then  $g = k_1 e_1$  for  $k_1 \in K_1$  and  $e_1 \in E_1$ , hence  $f(k_1)f(e_1) = k$ . Therefore,  $f(e_1) = f(k_1^{-1})k \in K_2 \cap E_2 = 1$ . Consequently,  $f(k_1) = k$  and  $f|_{K_1}$  is an epimorphism. We argue analogously for  $H_1$  and  $E_1$ .
  - 2. This follows from Lemma 3.6.
  - 3. If  $f|_{K_1}$  is an epimorphism, then  $f|_{H_1}$  is an epimorphism too. Since  $G_2 = K_2 H_2$ , the result follows.
  - 4. Let  $g_1, g_2 \in G_1$  such that  $f(g_1) = f(g_2)$ . Since  $G_1 = K_1E_1$  there exist  $k_1, k_2 \in K_1$  and  $e_1, e_2 \in E_1$  such that  $g_1 = k_1e_1$  and  $g_2 = k_2e_2$ . Then  $f(k_2^{-1}k_1) = f(e_2e_1^{-1}) \in K_2 \cap E_2 = 1$ . Hence  $f(k_1) = f(k_2)$  and  $f(e_1) = f(e_2)$ . By hypothesis, we have that  $k_1 = k_2$  and  $e_1 = e_2$ . Thus,  $g_1 = g_2$  and f is a monomorphism.

5. It follows from Statements (3) and (4).

If  $f|_{K_1}$  is a monomorphism, then f is not a monomorphism in general.

**Example 3.8.** Consider  $G = \langle x \rangle \times \langle y \rangle$  with  $x^n = y^n = 1$  for some  $n \in \mathbb{N}$ , then  $(G, \langle x \rangle, \langle xy \rangle, \langle y \rangle)$  and  $(\langle x \rangle, \langle x \rangle, \langle x \rangle, 1)$  are trifactorised groups. The projection  $\pi$  from G to  $\langle x \rangle$  is a morphism of trifactorised groups such that  $\pi|_{\langle xy \rangle}$  and  $\pi|_{\langle xy \rangle}$  are monomorphisms, but  $\pi$  is not a monomorphism.

**Proposition 3.9.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a trifactorised group morphism, then  $\mathsf{B}(f) = f|_{K_1}$  can be viewed as a brace morphism from  $\mathsf{B}(G_1, K_1, H_1, E_1)$  to  $\mathsf{B}(G_2, K_2, H_2, E_2)$ .

*Proof.* This follows directly from Lemma 3.6 because f is compatible with the derivations.

Remark 3.10. If we use the language of categories, Proposition 3.9 shows that B: **3factGrp**  $\longrightarrow$  **SKB** is a functor. If  $\widehat{B}$  is the restriction of B to the category **L3factGrp** of large trifactorised groups, then it is easy to see that  $L \circ \widehat{B}$ : **L3factGrp**  $\longrightarrow$  **L3factGrp** and  $\widehat{B} \circ L$ : **SKB**  $\longrightarrow$  **SKB** are the identity functors. That is, **L3factGrp** and **SKB** are equivalent categories. Proposition 3.7 shows that B preserves monomorphisms, epimorphisms and isomorphisms and reflects epimorphisms. Proposition 3.4 shows that B is essentially surjective. Example 2.1 shows that B is not surjective in morphisms.

We present a sufficient and necessary condition on a brace homomorphism f for the existence of a trifactorised group morphism  $\bar{f}$  such that  $B(\bar{f}) = f$ .

**Proposition 3.11.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a trifactorised group morphism. Let  $\mathsf{B}(G_i, K_i, H_i, E_i) = (K_i, \cdot, \boxdot)$ . Consider  $\eta_i: (K_i, \boxdot) \longrightarrow E_i$  the epimorphism that constructs the trifactorised group  $(G_i, K_i, H_i, E_i)$  for i = 1, 2. Then

$$\mathsf{B}(f)(\ker\eta_1) \le \ker\eta_2. \tag{2}$$

*Proof.* Let  $k \in \ker \eta_1$ , then  $k \in K_1 \cap H_1$ , therefore  $f(k) \in K_2 \cap H_2$ , and  $\eta_2(f(k)) = 1$ , which means that  $\mathsf{B}(f)(k) \in \ker \eta_2$ .

If a brace homomorphism satisfies Equation (2), then it is the image of a trifactorised group morphism from  $(G_1, K_1, H_1, E_1)$  to  $(G_2, K_2, H_2, E_2)$ .

**Proposition 3.12.** For i = 1, 2 consider  $(B_i, +, \cdot)$  a brace with additive group  $K_i$ , multiplicative group  $C_i$ , and lambda map  $\lambda_i \colon C_i \longrightarrow \operatorname{Aut}(K_i)$ . Consider  $(G_i, K_i, H_i, E_i)$  a trifactorised group associated with  $B_i$  with corresponding epimorphism  $\eta_i \colon C_i \longrightarrow E_i$  for i = 1, 2. Let  $f \colon B_1 \longrightarrow B_2$  a brace morphism such that  $f(\ker \eta_1) \leq \ker \eta_2$ , then: 1. The map  $\overline{f}: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  given by

$$\bar{f}(k\eta_1(c)) = f(k)\eta_2(f(c))$$

is a trifactorised group morphism.

- 2.  $\mathsf{B}(\bar{f}) = f$ .
- 3. If f is an epimorphism, then  $\overline{f}$  is an epimorphism.
- 4.  $\bar{f}$  is a monomorphism (resp., isomorphism) if and only if f is a monomorphism (resp., isomorphism), and  $f(\ker \eta_1) = \ker \eta_2$ .

Proof. Since f satisfies Equation (2),  $\bar{f}$  is well defined. Obviously, the restrictions of  $\bar{f}$  to  $K_1$  and  $E_1$  are group homomorphisms,  $\bar{f}(k\eta_1(c)) = \bar{f}(k)\bar{f}(\eta_1(c))$ , and  $\bar{f}(\eta_1(c)k)) = \bar{f}((\lambda_1)_c(k)) = f((\lambda_1)_c(k)) = (\lambda_2)_{f(c)}(f(k)) = {\eta_2(f(c))}f(k) = {\bar{f}(\eta_1(c))}\bar{f}(k)$  for all  $k \in K_1$  and  $c \in C_1$ . This proves that  $\bar{f}$  is a trifactorised group morphism. Moreover,  $\bar{f}$  restricted to  $K_1$  coincides with f, therefore  $\mathsf{B}(\bar{f}) = f$ . Proposition 3.7 shows that if  $\mathsf{B}(\bar{f}) = f$  is an epimorphism, then  $\bar{f}$  is an epimorphism.

Suppose that  $\bar{f}$  is a monomorphism. Then  $\mathsf{B}(\bar{f}) = f$  is a monomorphism by Proposition 3.7. Let  $c \in f^{-1}(\ker \eta_2)$ , then  $\bar{f}(\eta_1(c)) = \eta_2(f(c)) = 1 = \bar{f}(1)$ . Since  $\bar{f}$  is a monomorphism,  $\eta_1(c) = 1$  and  $c \in \ker \eta_2$ . Therefore, ker  $\eta_2 \leq f(\ker \eta_1)$  and the other inclusion is given by hypothesis. Now suppose that f is a monomorphism and  $f(\ker \eta_1) = \ker \eta_2$ . Let  $c \in C_1$  such that  $\eta_2(f(c)) = \bar{f}(\eta_1(c)) = 1$ . We have that  $f(c) \in \ker \eta_2 = f(\ker \eta_1)$ . Since fis a monomorphism, it follows that  $c \in \ker \eta_1$ . This means that  $\bar{f}|_{E_1}$  is a monomorphism. By hypothesis  $\bar{f}|_{K_1} = f$  is a monomorphism. Therefore, by Proposition 3.7  $\bar{f}$  is a monomorphism.  $\Box$ 

As we have seen in Proposition 3.11, given a trifactorised group morphism  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$ , then  $\mathsf{B}(f)$  satisfies the hypothesis of Proposition 3.12, thus  $\mathsf{B}(f)$  induces a trifactorised group morphism  $\overline{\mathsf{B}(f)}: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2).$ 

**Proposition 3.13.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a trifactorised group morphism. Then  $\overline{B}(f) = f$ .

Proof. Let  $B(f): (K_1, \cdot, \boxdot) \longrightarrow (K_2, \cdot, \boxdot)$ . Obviously  $\overline{B(f)}|_{K_1} = f|_{K_1}$ . For i = 1, 2, let  $\eta_i: (K_i, \boxdot) \longrightarrow E_i$  be the epimorphism defining the trifactorised group  $(G_i, K_i, H_i, E_i)$ . Then  $\overline{B(f)}(\eta_1(k)) = \eta_2(B(f)(k)) = \eta_2(f(k))$ . Since  $f(k\eta_1(k)) = f(k)f(\eta_1(k)) \in H_2$ , it follows that  $\eta_2(f(k)) = f(\eta_1(k))$ . Therefore,  $\overline{B(f)}|_{E_1} = f|_{E_1}$ . Hence  $\overline{B(f)} = f$ .

Note that Proposition 3.12 extends the idea of Section 2 of associating to a brace homomorphism a morphism between the large or small trifactorised groups.

**Corollary 3.14.** Let  $f: (B_1, +, \cdot) \longrightarrow (B_2, +, \cdot)$  be a brace homomorphism. Let  $C_i$  the multiplicative group,  $K_i$  the additive group, and  $\lambda_i: C_i \longrightarrow$  $\operatorname{Aut}(K_i)$  the lambda map of  $(B_i+, \cdot)$  for i = 1, 2. Then:

- 1. f extends to a trifactorised group morphism between the large trifactorised groups.
- 2. f extends to a trifactorised group morphism between the small trifactorised groups if and only if  $f(\ker \lambda_1) \leq \ker \lambda_2$ . In particular, brace epimorphisms extend to small trifactorised groups morphisms.

*Proof.* It follows from Proposition 3.12.

# 4 Isomorphism classes of the associated generalised trifactorised groups

All the trifactorised groups associated with a brace  $(B, +, \cdot)$  are constructed using group epimorphisms whose kernel contained in ker  $\lambda$ . It is natural then to think that there is a relationship between two of them.

**Proposition 4.1.** Let  $(B, +, \cdot)$  be a brace with additive group K, multiplicative group C, and lambda map  $\lambda: C \longrightarrow \operatorname{Aut}(K)$ . Consider  $\eta_i: C \longrightarrow E_i$ a group epimorphism such that ker  $\eta_i \leq \ker \lambda$  with  $(G_i, K, H_i, E_i)$  its corresponding trifactorised group associated with  $(B, +, \cdot)$  for i = 1, 2.

- 1. If ker  $\eta_1 \leq \ker \eta_2$ , then there exists a trifactorised group epimorphism  $f: (G_1, K, H_1, E_1) \longrightarrow (G_2, K, H_2, E_2)$  such that  $f|_K$  is an isomorphism.
- 2. If ker  $\eta_1 = \ker \eta_2$ , then  $(G_1, K, H_1, E_1) \cong (G_2, K, H_2, E_2)$ .

Proof. Let  $f: G_1 \longrightarrow G_2$  given by  $f(k\eta_1(c)) = k\eta_2(c)$ . Let  $c_1, c_2 \in C$  such that  $\eta_1(c_1) = \eta_1(c_2)$ , then  $c_1^{-1}c_2 \in \ker \eta_1 \leq \ker \eta_2$ . Hence,  $\eta_2(c_1) = \eta_2(c_2)$  and f is well defined. If we denote by  $\bar{\lambda}_i$  the action of  $E_i$  on K for i = 1, 2, we have that for all  $k_1, k_2 \in K$  and  $c_1, c_2 \in C$ :

$$f((k_1\eta_1(c_1))(k_2\eta_1(c_2))) = f((k_1 + \bar{\lambda}_1(c_1)(k_2))\eta_1(c_1c_2))$$
  
=  $(k_1 + \lambda_{c_1}(k_2))\eta_2(c_1c_2).$   
$$f(k_1\eta_1(c_1)) = f(k_2\eta_1(c_2)) = (k_1\eta_2(c_1))(k_2\eta_2(c_2)) = (k_1 + \bar{\lambda}_2(c_1)(k_2))\eta_2(c_1c_2)$$
  
=  $(k_1 + \lambda_{c_1}(k_2))\eta_2(c_1c_2).$ 

Therefore, f is a group homomorphism. Obviously, f(K) = K,  $f(H_1) = H_2$ , and  $f(E_1) = E_2$ , hence, f is a trifactorised group morphism. By Proposition 3.7, f is surjective. Finally, f restricted to K is injective. Therefore, f restricted to K is an isomorphism.

Moreover, it is easy to see that if ker  $\eta_1 = \ker \eta_2$ , then f is a monomorphism, and therefore, f is an isomorphism.

Remark 4.2. Proposition 4.1 shows that all trifactorised groups associated with  $(B, +, \cdot)$  are isomorphic to a trifactorised group of the form (G, K, H, C/N), where N is a normal subgroup of C contained in ker  $\lambda$ .

**Proposition 4.3.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be an epimorphism of trifactorised groups such that restricted to  $K_1$  is an isomorphism. Then  $B(G_1, K_1, H_1, E_1)$  and  $B(G_2, K_2, H_2, E_2)$  are isomorphic.

*Proof.* This follows directly from the fact that B preserves isomorphisms.  $\Box$ 

Proposition 4.3 shows that if two trifactorised groups are isomorphic, then they are associated with the same brace. This does not imply that ker  $\eta_1 =$ ker  $\eta_2$ , where  $\eta_i$ , i = 1, 2, are the epimorphisms defining the associated trifactorised groups.

**Example 4.4.** Consider  $K_1 = \langle x \rangle$ ,  $K_2 = \langle y \rangle$  where  $x^2 = y^2 = 1$ . Let *B* be the trivial brace of the group  $K = K_1 \times K_2$ . Then  $\lambda \colon K \longrightarrow \operatorname{Aut}(K)$  is trivial, therefore,  $\ker \lambda = K$ , so we can consider whichever epimorphism we want. Let  $\eta_i \colon K_1 \times K_2 \longrightarrow K/K_i$  the natural epimorphism for i = 1, 2. It is obvious that  $\ker \eta_1 = K_1 \neq K_2 = \ker \eta_2$  and the trifactorised groups that they define are  $(G_i, K, H_i, K/K_i)$ , where  $G_i = K_1 \times K_2 \times K/K_i$  for i = 1, 2.

Let  $f: (G_1, K, H_1, K/K_1) \longrightarrow (G_2, K, H_2, K/K_2)$  defined by

$$f(x, 1, K_1) = (1, y, K_2), f(1, y, K_1) = (x, 1, K_1), f(1, 1, yK_1) = (1, 1, xK_2).$$

It is easy to check that f is a trifactorised group isomorphism. Therefore,  $(G_1, K, H_1, K/K_1) \cong (G_2, K, H_2, K/K_2)$ , but ker  $\eta_1 \neq \text{ker } \eta_2$ .

In the particular case of the large and small trifactorised groups the result is true.

**Proposition 4.5.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Let (G, K, H, E) be a trifactorised group associated with  $(B, +, \cdot)$  with corresponding epimorphism  $\eta: C \longrightarrow E$ . Then:

- 1.  $(G, K, H, E) \cong \mathsf{L}(B, +, \cdot)$  if and only if ker  $\eta = 1$ .
- 2.  $(G, K, H, E) \cong \mathsf{S}(B, +, \cdot)$  if and only if ker  $\eta = \ker \lambda$ .

*Proof.* If the kernels are equal, we have seen that the trifactorised groups are isomorphic, so we only need to prove the converse. Let us denote  $\delta: C \longrightarrow K$  the identity map of  $(B, +, \cdot)$ .

If  $(G, K, H, E) \cong S(B, +, \cdot)$ , by Proposition 4.6, there exists a brace automorphism f such that  $f(\ker \lambda) = \ker \eta$ . Since  $\ker \lambda$  is invariant by brace automorphisms, it follows that  $\ker \lambda = \ker \eta$ . The argument is analogous for the large trifactorised group.

**Proposition 4.6.** Let  $(B, +, \cdot)$  be a brace with additive group K, multiplicative group C and identity map  $\delta: C \longrightarrow K$ . If  $(G_i, K, H_i, E_i)$  is a trifactorised group associated with  $(B, +, \cdot)$  with the corresponding epimorphism  $\eta_i: C \longrightarrow E_i$  for i = 1, 2. Then  $(G_1, K, H_1, E_1) \cong (G_2, K, H_2, E_2)$  if and only if there exists  $f \in \operatorname{Aut}(B, +, \cdot)$  such that  $f(\ker \eta_1) = \ker \eta_2$ .

Proof. Let suppose that  $(G_1, K, H_1, E_1) \cong (G_2, K, H_2, E_2)$  and let g be a trifactorised group isomorphism. By Propositions 3.7 and 3.9,  $f = \mathsf{B}(g) \in \operatorname{Aut}(B, +, \cdot)$ . By Proposition 3.11  $f(\ker \eta_1) \leq \ker \eta_2$ , and analogously with  $g^{-1}$  we have that  $\ker \eta_2 \leq f(\ker \eta_1)$ .

Suppose that there exists  $f \in \operatorname{Aut}(B, +, \cdot)$  such that  $f(\ker \eta_1) = \ker \eta_2$ . Then  $\ker \eta_1 = \ker(\eta_2 \circ f)$ . By Proposition 3.12, we can construct a trifactorised group isomorphism  $\overline{f}: (G_1, K, H_1, E_1) \longrightarrow (G_2, K, H_2, E_2)$ .  $\Box$ 

Let  $(B, +, \cdot)$  be a brace with additive group K, multiplicative group C, and lambda map  $\lambda$ . Consider  $f \in \operatorname{Aut}(B, +, \cdot)$  and  $c \in \ker \lambda$ , then  $\lambda_{f(c)}(f(c')) = f(\lambda_c(c')) = f(c)$  for all  $c \in B$ . Therefore,  $f(c) \in \ker \lambda$  and  $f(\ker \lambda) \leq \ker \lambda$ . Arguing similarly with the inverse of f, the other inclusion holds. Hence  $f_C(\ker \lambda) = \ker \lambda$ .

Let  $\Omega = \{N \leq C \mid N \leq \ker \lambda\}$ . We have that  $\operatorname{Aut}(B, +, \cdot)$  acts on  $\Omega$  via  $f \bullet N = f(N)$ . The action is well defined because  $f(N) \leq f(\ker \lambda) = \ker \lambda$ .

**Theorem 4.7.** Let  $(B, +, \cdot)$  a brace with additive group K, multiplicative group C, and lambda map  $\lambda \colon C \longrightarrow \operatorname{Aut}(K)$ . Let  $\Omega = \{N \trianglelefteq C \mid N \le \ker \lambda\}$ . There is a bijection between the isomorphism classes of trifactorised groups associated with  $(B, +, \cdot)$  and the orbits of the action of  $\operatorname{Aut}(B, +, \cdot)$  on  $\Omega$ , which associates to the orbit of  $N \in \Omega$  the trifactorised group associated with  $(B, +, \cdot)$  corresponding to the natural epimorphism  $\eta_N \colon C \longrightarrow C/N$ .

Proof. Let  $N \in \Omega$ . Since ker  $\eta_N = N \leq \ker \lambda$ ,  $\eta_N$  defines a trifactorised group associated with  $(B, +, \cdot)$ . If we consider  $N, M \in \Omega$  such that there exists  $f \in \operatorname{Aut}(B, +, \cdot)$  which f(N) = M, by Proposition 4.6,  $\eta_N$  and  $\eta_M$ define isomorphic trifactorised groups associated with  $(B, +, \cdot)$ . Hence this defines a map from the set of all orbits of the action of  $\operatorname{Aut}(B, +, \cdot)$  on  $\Omega$  to the set of all isomorphism classes of trifactorised groups associated with  $(B, +, \cdot)$ . This map is injective by Proposition 4.6.

If (G, K, H, E) is a trifactorised group associated with  $(B, +, \cdot)$  constructed with the epimorphism  $\eta: C \longrightarrow E$ , by Proposition 4.1, we know that  $\eta_{\ker \eta}$  gives us a trifactorised group isomorphic to (G, K, H, E) associated with  $(B, +, \cdot)$ , therefore, our map is surjective too and hence it is a bijection.  $\Box$ 

# 5 Substructures of braces and generalised trifactorised groups

In this section, we characterise substructures of braces in terms of generalised trifactorised groups. We will identify each of these substructures with the corresponding subgroup L of the additive group, represented by the subgroup K, and the corresponding subgroup  $\sigma^{-1}(L)$  of the multiplicative group, represented by the subgroup H.

It is possible to compute the images of subsets of H in K or the preimages of subsets of K in H by the associated derivation  $\sigma$  by means of the following result (cf. [2, Lemma 3.3]).

**Proposition 5.1.** Consider the trifactorised group (G, K, H, E) and consider  $\sigma: H \longrightarrow K$  the corresponding bijective derivation. Then:

- 1. If  $L \subseteq K$ , then  $\sigma^{-1}(L) = LE \cap H$ .
- 2. If  $S \subseteq H$ , then  $\sigma(S) = SE \cap K$ .
- Proof. 1. Let  $h \in \sigma^{-1}(L)$ . Then  $h = k_h e_h$  with  $\sigma(h) = k_h \in L$  and  $h = k_h e_h \in LE \cap H$ . Conversely, suppose that for some  $l \in L$  and  $e \in E$ , we have that  $h = le \in LE \cap H$ . Since  $l \in K$  and  $e \in E$ ,  $\sigma(h) = k_h = l$  and  $e_h = e$ . Therefore  $\sigma^{-1}(L) = LE \cap H$ .
  - 2. Suppose that  $k \in \sigma(S)$ . Then there exists  $h \in S$  such that  $\sigma(h) = k$ and  $h = ke_h$ . Hence  $k = (ke_h)e_h^{-1} \in SE \cap K$ . Conversely, suppose that  $se \in K$  with  $s \in S$ ,  $e \in E$ . Then  $se = (k_se_s)e \in K$ , which implies that  $e_se = 1$  and so  $\sigma(s) = k_s = se$ . Consequently,  $\sigma(S) = SE \cap K$ .

**Proposition 5.2.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Let (G, K, H, E) be an associated trifactorised group to  $(B, +, \cdot)$ . Consider L a subset of B, then:

1. The following statements are equivalent:

(a)  $(L, +, \cdot)$  is a subbrace of  $(B, +, \cdot)$ .

- (b) L is a subgroup of K and  $LE \cap H$  is a subgroup of H.
- (c) L is a subgroup of K and  $LE \cap H$  is a subgroup of  $N_G(L)$ .
- (d)  $LE \cap LH$  is a subgroup of G.
- 2. The following statements are equivalent:
  - (a) L is a left ideal of  $(B, +, \cdot)$ .
  - (b) L is a subgroup of K and E is a subgroup of  $N_G(L)$ .
  - (c) LE is a subgroup of G.
- 3. The following statements are equivalent:
  - (a) L is a strong left ideal of  $(B, +, \cdot)$ .
  - (b) L is a normal subgroup of G.
- 4. The following statements are equivalent:
  - (a) L is an ideal of  $(B, +, \cdot)$ .
  - (b) L is a normal subgroup of G and  $LE \cap H$  is a normal subgroup of H.
  - (c)  $LE \cap LH$  is a normal subgroup of G.
- *Proof.* 1. 1a if and only if 1b. L is a subbrace if and only if is a subgroup of K and  $LE \cap H = \sigma^{-1}(L)$  is a subgroup of H.

1b implies 1c. Let  $le_l \in LE \cap H$ , where  $l \in L$  and  $e_l \in E$  and consider  $l_1 \in L$ . Then  $(le_l)l_1(le_l)^{-1} = l(e_ll_1e_l^{-1})l^{-1}$ . Since  $le_l, l_1e_{l_1} \in LE \cap H$  we have that  $l(e_ll_1e_l^{-1})e_le_{l_1} = (le_l)(l_1e_{l_1}) \in LE \cap H$ . Therefore,  $l(e_ll_1e_l^{-1}) \in L$ , which means that  $(le_l)l_1(le_l)^{-1} \in L$  because L is a subgroup of K. Therefore,  $LE \cap H$  normalises L.

1c implies 1d. If  $LE \cap H$  is a subgroup of  $N_G(L)$ , then  $L(LE \cap H)$  is a subgroup of G and, by Dedekind's identity,  $LE \cap LH$  is a subgroup of G.

1d implies 1b. Note that

$$LE \cap LH \cap K = L(E \cap K) \cap LH = L \cap LH = L.$$

Hence, L is a subgroup of K. Furthermore,  $LE \cap LH \cap H = LE \cap H$  is a subgroup of H.

2. 2a if and only if 2b. We have that L is a left ideal of B if and only if it is a subgroup of K and is  $\lambda$ -invariant (recall that  $\lambda$  corresponds to the action of E on K). Therefore, L is a left ideal if and only if it is a subgroup of K that is invariant by the conjugation by elements of E, equivalently,  $E \leq N_G(L)$ .

2b implies 2c. Since E normalises L it follows that LE is a subgroup. 2c implies 2b Since LE is a subgroup it follows that  $L = L(K \cap E) = K \cap LE$  is a normal subgroup of LE. Therefore, E normalises L.

3. 3a implies 3b. A strong left ideal is simply a left ideal which is normal in the additive group, therefore, we have that L is a normal subgroup of K and E normalises L. Hence,  $G = KE \leq N_G(L)$ , equivalently, Lis a normal subgroup of G.

3b implies 3a. Since L is a normal subgroup of G we have that LE is a subgroup of G, hence, by 2, it follows that L is a left ideal. Furthermore, L is normal in K, so L is a strong left ideal.

4. 4a if and only if 4b. An ideal is a strong left ideal which is normal in the multiplicative group, therefore, L is an ideal if and only if L is a strong left ideal of B and  $LE \cap H = \sigma^{-1}(L)$  is a normal subgroup of H. By 3, this is equivalent to L being a normal subgroup of G and  $LE \cap H$  being a normal subgroup of H.

4b implies 4c. Since L is a normal subgroup of G we have that  $LE \cap LH = L(LE \cap H)$  is a subgroup of G. Since L is normal in G and  $LE \cap H$  is normal in H it follows that  $LE \cap LH = L(LE \cap H)$  is normalised by H. We want to prove that it is also normalised by E.

Let us prove that  $E \cap LH$  is a normal subgroup of E. First note that  $K(LE \cap H)$  is normal in KH = G. Therefore,  $K(LE \cap H) \cap E$  is a normal subgroup of E. Using Dedekind's identity we obtain that

$$K(LE \cap H) \cap E = KL(LE \cap H) \cap E = K(LE \cap LH) \cap E$$
$$= KL(E \cap LH) \cap E = K(E \cap LH) \cap E$$
$$= (E \cap LH)(K \cap E) = E \cap LH$$

In conclusion,  $E \cap LH$  is a normal subgroup of E. Since L is normal in G and  $E \cap LH$  is normal in E, we have that  $LE \cap LH = L(E \cap LH)$  is normalised by E.

Therefore,  $LE \cap LH$  is normalised by H and E, hence, by HE = G. 4c implies 4b. Since  $LE \cap LH$  is normal in G,  $LE \cap H = LE \cap LH \cap H$ is a normal subgroup of H. Furthermore,  $LE \cap LH \cap K$  is a normal subgroup of G. As we have seen in 1,  $LE \cap LH \cap K = L$ . Hence, L is a normal subgroup of G.

Our next result shows a natural embedding of the trifactorised group associated with the subbrace of a brace B in the trifactorised group associated with B.

**Lemma 5.3.** Consider (G, K, H, E) a trifactorised group. Let  $\sigma \colon H \longrightarrow K$ be the corresponding bijective derivation and let  $\pi_E \colon G \longrightarrow E$  be the natural projection. If  $L \subseteq K$ , then  $\pi_E(\sigma^{-1}(L)) = L^{-1}H \cap E$ .

Proof. Let  $h \in \sigma^{-1}(L)$ , then  $h = k_h e_h$  with  $k_h \in K$  and  $e_h \in H$ . Since  $\sigma(h) = k_h$ , we have that  $k_h \in H$ . Furthermore,  $e_h = \pi_E(h)$ . It follows that  $e_h = k_h^{-1}h \in L^{-1}H \cap E$ . We conclude that  $\sigma^{-1}(L) \subseteq L^{-1}H \cap E$ .

Conversely, let  $e = l^{-1}h \in L^{-1}H \cap E$ . As before, let  $h = k_h e_h$  with  $k_h \in K$ and  $e_h \in E$ . Then  $e = l^{-1}k_h e_h$  and so  $l^{-1}k_h = 1$ , that is,  $k_h = l$ , and  $e = e_h$ . We conclude that  $l = k_h = \sigma(h)$ , in other words,  $h \in \sigma^{-1}(L)$ , and  $\pi_E(h) = e$ . Consequently,  $e \in \pi_E(\sigma^{-1}(L))$ . It follows that  $L^{-1}H \cap E \subseteq \pi_E(\sigma^{-1}(L))$  and so we have the desired equality.

**Proposition 5.4.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Let  $\sigma: H \longrightarrow K$  be the bijective derivation and consider (G, K, H, E) a trifactorised group associated with  $(B, +, \cdot)$  with the corresponding epimorphism  $\eta: C \longrightarrow E$ . If  $L \subseteq K$  such that  $(L, +, \cdot)$  is a subbrace of  $(B, +, \cdot)$  and  $T = L\sigma^{-1}(L)$ , then

$$(T, L, T \cap H, T \cap E) = (LE \cap LH, L, LE \cap H, LH \cap E)$$

is a trifactorised group associated with  $(L, +, \cdot)$ . Furthermore, if  $\delta \colon C \longrightarrow K$ is the identity map and  $D = \delta^{-1}(L)$ , the trifactorised group is constructed by means of the epimorphism  $\eta|_D \colon D \longrightarrow LH \cap E$ .

*Proof.* By Dedekind's identity,  $T = L\sigma^{-1}(L) = L(LE \cap H) = LE \cap LH$ . Then  $T \cap H = LE \cap H$  and  $T \cap E = LH \cap E$ .

By Proposition 5.2,  $LE \cap LH$  is a group and  $LE \cap H$  is a subgroup of G that normalises L. Therefore  $LE \cap LH = L(LE \cap H) \leq N_G(L)$ , thus L is a normal subgroup of  $LE \cap LH$ . Using Dedekind's identity we have that

$$(LH \cap E)(LE \cap H) = LE \cap (LH \cap E)H = LE \cap LH \cap EH = LE \cap LH.$$

The other properties for  $(T, L, T \cap H, T \cap E)$  to be a trifactorised group follow from an application of Dedekind's identity.

We have that  $\eta|_D: D \longrightarrow \eta(D)$  is an epimorphism that defines a trifactorised group  $(\tilde{G}, L, \tilde{H}, \eta(D))$  associated with  $(L, +, \cdot)$ . Consider the map  $\begin{array}{l} \beta \colon C \longrightarrow H \text{ given by } \beta(c) = \delta(c)\eta(c) \text{ for } c \in C. \text{ Then } \delta = \sigma \circ \beta \text{ and} \\ \eta = \pi_E|_H \circ \beta. \text{ Consequently, } \eta(D) = \eta(\delta^{-1}(L)) = \pi_E(\beta(\beta^{-1}(\sigma^{-1}(L)))) = \\ \pi_E(\sigma^{-1}(L)) = L^{-1}H \cap E = LH \cap E \text{ by Lemma 5.3 and the fact that } L \leq K. \\ \text{Hence, } \tilde{G} = L\eta(D) = L(LH \cap E) = T, \text{ and } \tilde{H} = \tilde{H}(H \cap \tilde{E}) = H \cap \tilde{H}\tilde{E} = \\ H \cap T. \end{array}$ 

#### 5.1 Generalised trifactorised subgroups

**Definition 5.5.** Given a trifactorised group (G, K, H, E) we define a *trifac*torised subgroup of (G, K, H, E) as a trifactorised group  $(\tilde{G}, \tilde{K}, \tilde{H}, \tilde{E})$  such that  $\tilde{G} \leq G$ ,  $\tilde{K} \leq K$ ,  $\tilde{H} \leq H$  and  $\tilde{E} \leq E$ . We write  $(\tilde{G}, \tilde{K}, \tilde{H}, \tilde{E}) \leq$ (G, K, H, E).

**Proposition 5.6.** Let (G, K, H, E) be a trifactorised group with bijective derivation  $\sigma: H \longrightarrow K$  and let  $(\tilde{G}, \tilde{K}, \tilde{H}, \tilde{E})$  be a trifactorised subgroup of (G, K, H, E) with bijective derivation  $\tilde{\sigma}: \tilde{H} \longrightarrow \tilde{K}$ . Then:

1.  $\tilde{K} = \tilde{G} \cap K$ ,  $\tilde{H} = \tilde{G} \cap H$  and  $\tilde{E} = \tilde{G} \cap E$ . 2.  $\tilde{\sigma} = \sigma|_{\tilde{H}}$ .

*Proof.* The proofs of all three equalities are analogous, so let us prove it for  $\tilde{K}$ .

$$\tilde{G} \cap K = \tilde{K}\tilde{E} \cap K = \tilde{K}(\tilde{E} \cap K) = \tilde{K}.$$

Given  $h \in \tilde{H}$ , it admits a factorisation  $h = \tilde{k}_h \tilde{e}_h$  with  $\tilde{k}_h \in \tilde{K} \leq K$  and  $\tilde{e}_h \in \tilde{E} \leq E$ . Since this factorisation is unique in G, we have that  $\sigma(h) = \tilde{k}_h = \tilde{\sigma}(h)$ .

Remark 5.7. Proposition 5.6 shows that all trifactorised subgroups are of the form  $(T, T \cap K, T \cap H, T \cap E)$  where T is a subgroup of G, but not every subgroup of G works, for example if T = K, we have  $(K, K, K \cap H, 1)$  which in general is not a trifactorised group because  $\tilde{H}\tilde{E} = K \cap H$ .

**Definition 5.8.** Let (G, K, H, E) be a trifactorised group. A subgroup T of G is a *trifactorised subgroup* if  $(T, T \cap K, T \cap H, T \cap E)$  is a trifactorised subgroup of (G, K, H, E).

Our next result characterises the trifactorised subgroups of G.

**Proposition 5.9.** Let (G, K, H, E) be a trifactorised group and let T be a subgroup of G. The following statements are equivalent:

1. T is a trifactorised subgroup of G.

2.  $T = (T \cap K)E \cap (T \cap K)H.$ 

*Proof.* 1 implies 2. Let  $\sigma: H \longrightarrow K$  the bijective derivation of (G, K, H, E)and  $\tilde{\sigma}: T \cap H \longrightarrow T \cap K$  be the bijective derivation of the trifactorised subgroup T. By Proposition 5.6

$$(T \cap K)E \cap H = \sigma^{-1}(T \cap K) = \tilde{\sigma}^{-1}(T \cap K) = T \cap H.$$

Then, by Dedekind's identity,

$$(T \cap K)E \cap (T \cap K)H = (T \cap K)((T \cap K)E \cap H) = (T \cap K)(T \cap H) = T.$$

2 implies 1. The equalities  $(T \cap K)(T \cap H) = T$  and  $(T \cap K)(T \cap E) = T$  follow directly from Dedekind's identity. Furthermore,

$$(T \cap H)(T \cap E) = T \cap (T \cap H)E$$
  
=  $T \cap ((T \cap K)E \cap (T \cap K)H \cap H)E$   
=  $T \cap ((T \cap K)E \cap H)E$   
=  $T \cap (T \cap K)E \cap HE$   
=  $T \cap (T \cap K)E = T.$ 

Therefore T is a trifactorised subgroup of G.

**Proposition 5.10.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Consider (G, K, H, E) a trifactorised group associated with  $(B, +, \cdot)$ . Then the map  $L \mapsto LH \cap LE$  is a bijection between subbraces of  $(B, +, \cdot)$  and trifactorised subgroups of (G, K, H, E). Furthermore, the restriction of this map to the set all ideals of B defines a bijection between the set of all the ideals of B and the set of all normal trifactorised subgroups of G.

Proof. If L is a subbrace of  $(B, +, \cdot)$ , Proposition 5.4 shows that  $LE \cap LH$ is a trifactorised subgroup of (G, K, H, E). If T is a trifactorised subgroup of G, we have that  $T \cap K$  corresponds to a subbrace of  $(B, +, \cdot)$  because  $(T \cap K)H \cap (T \cap K)E = T \leq G$ . Hence,  $T \mapsto T \cap K$  is a map from the the set of all trifactorised subgroups of (G, K, H, E) to the set of all subbraces of  $(B, +, \cdot)$ . It follows that these two maps are mutually inverses and so they are bijective. The last claim follows from Proposition 5.2 (4).

## 6 Quotients of generalised trifactorised groups

Our main goal in this section is to show how we can make quotients with the substructure introduced in Section 5.1, and how these quotients are related to brace quotients.

Note that not every quotient of a trifactorised group is a trifactorised group.

**Example 6.1.** If (G, K, H, E) is a trifactorised group with  $G \neq K$ , then (G/K, 1, G/K, G/K) is not a trifactorised group because  $G/K \cap G/K \neq 1$ .

Not all the quotients of a trifactorised group are quotients by a normal trifactorised subgroup.

**Example 6.2.** Let (G, K, H, E) be a trifactorised group with E nontrivial and normal in G. Consider T = E. Then

$$KT \cap ET = KE \cap EE = G \cap E = E,$$
$$HT \cap ET = HE \cap EE = G \cap E = E,$$
$$(T \cap K)H \cap (T \cap K)E = (E \cap K)H \cap (E \cap K)E = H \cap E = 1$$

The first two equations prove that (G/T, KT/T, HT/T, ET/T) is a trifactorised group, but T is not a trifactorised subgroup.

The following theorem characterises the normal subgroups of a trifactorised group whose corresponding quotient is again trifactorised.

**Theorem 6.3.** Let (G, K, H, E) be a trifactorised group and T a normal subgroup of G, then the following statements are equivalent:

(3)

1. (G/T, KT/T, HT/T, ET/T) is a trifactorised group.

2. 
$$T = (T \cap K)(T \cap E) = (T \cap H)(T \cap E).$$

3. 
$$T = ((T \cap K)H \cap (T \cap K)E)(T \cap E).$$

*Proof.* 1 implies 2. By hypothesis  $KT \cap ET = HT \cap ET = T$ . Now,

$$(T \cap K)(T \cap E) = (T \cap K)(KT \cap ET \cap E) = (T \cap K)(KT \cap E)$$
$$= KT \cap (T \cap K)E = KT \cap (KT \cap ET \cap K)E$$
$$= KT \cap (K \cap ET)E = KT \cap KE \cap ET$$
$$= KT \cap ET = T.$$

Analogously, we have that  $(T \cap H)(T \cap E) = T$ . 2 implies 3.

$$((T \cap K)H \cap (T \cap K)E)(T \cap E) = (T \cap K)H(T \cap E) \cap (T \cap K)E$$
  
=  $TH \cap (T \cap K)E = (T \cap K)(TH \cap E)$   
=  $(T \cap K)((T \cap E)(T \cap H)H \cap E)$   
=  $(T \cap K)((T \cap E)H \cap E)$   
=  $(T \cap K)(T \cap E)(H \cap E) = T.$ 

3 implies 1.

$$HT = H((T \cap K)H(T \cap E) \cap (T \cap K)E)$$
  
=  $(T \cap K)H(T \cap E) \cap H(T \cap K)E$   
=  $(T \cap K)H(T \cap E) \cap G = (T \cap K)H(T \cap E).$   
$$ET = ((T \cap K)H(T \cap E) \cap (T \cap K)E)E$$
  
=  $(T \cap K)H(T \cap E)E \cap (T \cap K)E$   
=  $G \cap (T \cap K)E = (T \cap K)E.$ 

Then  $HT \cap ET = (T \cap K)H(T \cap E) \cap (T \cap K)E = T$ . Furthermore  $T = T \cap ET = T \cap (T \cap K)E = (T \cap K)(T \cap E)$ . Therefore,

$$KT \cap ET = K(T \cap K)(T \cap E) \cap E(T \cap K)(T \cap E)$$
  
=  $K(T \cap E) \cap E(T \cap K) = (T \cap K)(K(T \cap E) \cap E)$   
=  $(T \cap K)(T \cap E)(K \cap E) = T.$ 

This proves that  $KT/T \cap ET/T = T$  and  $HT/T \cap ET/T = T$ . Therefore (G/T, KT/T, HT/T, ET/T) is a trifactorised group.

**Corollary 6.4.** Let (G, K, H, E) be a trifactorised group, let  $\sigma: H \longrightarrow K$  be its associated bijective derivation, and let T be a normal trifactorised subgroup of (G, K, H, E). Then

is a trifactorised group, with bijective derivation  $\overline{\sigma} \colon HT/T \longrightarrow KT/T$  given by  $\overline{\sigma}(hT) = \sigma(h)T$ .

*Proof.* By definition the equation (3) is satisfied, therefore, the quotient is a trifactorised group.

Given  $g \in G$  its factorisation is  $g = k_g e_g$  with  $k_g \in K$  and  $e_g \in E$ , hence,  $gT = (k_g T)(e_g T)$ . Since the factorisation is unique, we have that  $\overline{\sigma}(hT) = k_h T = \sigma(h)T$ .

Remark 6.5. Given a trifactorised group (G, K, H, E) and a normal subgroup  $T \leq G$ , it follows that (G/T, KT/T, HT/T, ET/T) is a trifactorised group if and only if T is the kernel of some trifactorised group morphism. Therefore, given a trifactorised group morphism  $f: (G_1, K_1, H_1, E_1) \longrightarrow$  $(G_2, K_2, H_2, E_2)$ , it follows that

$$\ker f = ((\ker f \cap K_1)H_1 \cap (\ker f \cap K_1)E_1)(\ker f \cap E_1).$$

**Proposition 6.6.** Let  $(B, +, \cdot)$  be a brace with additive group K, multiplicative group C and identity map  $\delta: C \longrightarrow K$ . Consider (G, K, H, E) a trifactorised group associated with  $(B, +, \cdot)$  with its corresponding epimorphism  $\eta: C \longrightarrow E$ . Let I be an ideal of B and suppose I corresponds to the subgroup  $L \leq K$ . If  $T = LH \cap LE$ , then the trifactorised group (G/T, KT/T, HT/T, ET/T) is associated with the brace  $(B/I, +, \cdot)$  with epimorphism  $\bar{\eta}: \overline{C} \longrightarrow \overline{E}$  given by  $\bar{\eta}(cD) = \eta(c)T$  where  $D = \delta^{-1}(L), \overline{C} = C/D$  and  $\overline{E} = ET/T$ .

*Proof.* Note that the additive group of B/I is K/L = KT/T and the multiplicative group is C/D. The operations of the brace associated to the quotient are:

$$\delta(c_1)T\delta(c_2)T = \delta(c_1)\delta(c_2)T$$
  

$$\delta(c_1)T \boxdot \delta(c_2)T = \overline{\sigma}(\overline{\sigma}^{-1}(\delta(c_1)T)\overline{\sigma}^{-1}(\delta(c_2)T))$$
  

$$= \overline{\sigma}((\sigma^{-1}(\delta(c_1))T)(\sigma^{-1}(\delta(c_2)T)))$$
  

$$= \sigma(\sigma^{-1}(\delta(c_1))\sigma^{-1}(\delta(c_2)))T$$
  

$$= \sigma(\delta(c_1)\delta(c_2)^{\eta(c_1)}\eta(c_1c_2))T = \delta(c_1c_2)T$$

These are the operations of B/I viewed in its additive group KT/T. Therefore, (G/T, KT/T, HT/T, ET/T) is associated with B/I.

The identity map  $\overline{\delta} \colon C/D \longrightarrow KT/T$  of B/I is given by  $\overline{\delta}(cD) = \delta(c)T$ . It follows that  $\overline{\delta}(cD)\eta(c)T = \delta(c)\eta(c)T \in HT/T$ , therefore, the epimorphism that constructs the trifactorised group (G/T, KT/T, HT/T, ET/T) must be  $\overline{\eta} \colon C/D \longrightarrow ET/T$  given by  $\overline{\eta}(cD) = \eta(c)T$ .

For the large trifactorised group associated with  $(B, +, \cdot)$ , we have:

**Proposition 6.7.** Let  $(B, +, \cdot)$  be a brace, denote  $L(B, +, \cdot) = (G, K, H, E)$ and consider I an ideal of B with L its associated subgroup of K. If  $T = LE \cap LH$ , it follows that the trifactorised quotient of G by T is  $L(B/I, +, \cdot)$ .

*Proof.* We already know that (G/T, KT/T, HT/T, ET/T) is a trifactorised group associated with  $(B/I, +, \cdot)$ . To prove that it is the large one, we need to prove that  $KT \cap HT = T$ .

$$KT \cap HT = K(T \cap K)(T \cap H) \cap H(T \cap K)(T \cap H)$$
  
=  $K(T \cap H) \cap H(T \cap K) = (T \cap K)(K(T \cap H) \cap H)$   
=  $(T \cap K)(K \cap H)(T \cap H) = (K \cap H)T = T.$ 

The above proposition is false for the small trifactorised group associated with  $(B, +, \cdot)$ .

**Example 6.8.** Consider the trifactorised group (G, K, H, E) where  $K = \langle x \rangle \cong C_6$ ,  $E = \langle y \rangle \cong C_2$ ,  $G = [K]E \cong D_{12}$  and  $H = \langle x^2, x^3y \rangle \cong S_3$ . Since  $x^y = x^{-1}$  it follows that  $C_E(K) = 1$ , therefore, the trifactorised group is small.

Let  $T = \langle x^2 \rangle = H \cap K$ . Since |G| = 12 and |T| = 3 it follows that |G/T| = 4. In particular, G/T is abelian. Hence,  $|C_{ET/T}(K/T)| = |ET/T| = |E| = 2$ . Thus (G/T, K/T, H/T, ET/T) is not the small trifactorised group.

We bring the paper to a close by showing that every trifactorised group associated to a brace B is a quotient of the large trifactorised group and has the small trifactorised group as a quotient.

**Proposition 6.9.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Consider (G, K, H, E) a trifactorised group associated with B with corresponding epimorphism  $\eta: C \longrightarrow E$ , and T a normal subgroup of G contained in E. Then (G/T, KT/T, HT/T, E/T) is a trifactorised group associated with B with its corresponding epimorphism  $\bar{\eta} = \pi_T \circ \eta$  where  $\pi_T: E \longrightarrow E/T$  is the natural epimorphism.

Proof. Proposition 6.3 shows that (G/T, KT/T, HT/T, E/T) is a trifactorised group. Analogously to 6.6 the operations of the associated brace are  $\delta(c_1)T\delta(c_2)T = \delta(c_1)\delta(c_2)T$  and  $\delta(c_1)T \boxdot \delta(c_2)T = \delta(c_1c_2)T$ . But in this case we have that  $KT/T \cong K$ , therefore, we have that the associated brace is  $(B, +, \cdot)$ .

The elements of our trifactorised group are of the form  $\delta(c_1)\eta(c_2)T$ . Since its bijective derivation is  $\overline{\delta}: C \longrightarrow KT/T$  given by  $\overline{\delta}(c) = \delta(c)T$ , we have that the epimorphism that corresponds to the trifactorised group is  $\overline{\eta}: C \longrightarrow E/T$ given by  $\overline{\eta}(c) = \eta(c)T = (\pi_T \circ \eta)(c)$ .

**Lemma 6.10.** Let  $f: (G_1, K_1, H_1, E_1) \longrightarrow (G_2, K_2, H_2, E_2)$  be a trifactorised group epimorphism such that  $f|_{K_1}$  is an isomorphism. Then ker  $f \leq E$ .

*Proof.* Let  $ke \in \ker f$  with  $k \in K_1$  and  $e \in E_1$ . Then 1 = f(ke) = f(k)f(e). Since  $f(k) \in K_2$  and  $f(e) \in E_2$  we have that f(k) = 1 and f(e) = 1. Hence  $f|_{K_1}$  is an isomorphism, and so, k = 1 and ker  $f \leq E$ .

**Theorem 6.11.** Let  $(B, +, \cdot)$  be a brace with additive group K and multiplicative group C. Consider  $(G_i, K_i, H_i, E_i)$  for i = 1, 2 two trifactorised groups associated with B with corresponding epimorphisms  $\eta_i \colon C \longrightarrow E_i$ . If ker  $\eta_1 \leq \ker \eta_2$ , then  $(G_2, K_2, H_2, E_2)$  is a quotient of  $(G_1, K_1, H_1, E_1)$ . In particular:

- 1. Every associated trifactorised group of B is quotient of  $L(B, +, \cdot)$ .
- 2. Every associated trifactorised group of B has  $S(B, +, \cdot)$  as a quotient.

*Proof.* The first part is a direct application of Proposition 4.1, Lemma 6.10 and Proposition 6.9. The second part follows from the first one and Proposition 4.5.  $\hfill \Box$ 

## 7 Acknowledgements

This work is supported by the grant CIAICO/2023/007 from the Conselleria d'Educació, Universitats i Ocupació, Generalitat Valenciana.

# References

- B. Amberg, S. Franciosi, and F. de Giovanni. *Products of groups*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 1992. Oxford Science Publications.
- [2] A. Ballester-Bolinches and R. Esteban-Romero. Triply factorised groups and the structure of skew left braces. *Commun. Math. Stat.*, 10:353–370, 2022.
- [3] A. Ballester-Bolinches, R. Esteban-Romero, P. Jiménez-Seral, and V. Pérez-Calabuig. A construction of braces whose additive group is a given group of trivial centre. Preprint, 2024.
- [4] R. Baxter. Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. I. Some fundamental eigenvectors. Ann. Physics, 76(1):1–24, 1973.
- [5] D. Bourn, A. Facchini, and M. Pompili. Aspects of the category SKB of skew braces. *Comm. Algebra*, 51(5):2129–2143, 2023.
- [6] A. Caranti, F. Dalla Volta, and M. Sala. Abelian regular subgroups of the affine group and radical rings. *Pub. Mat. Debrecen*, 69(3):297–308, 2006.
- [7] F. Catino, I. Colazzo, and P. Stefanelli. On regular subgroups of the affine group. *Bull. Austral. Math. Soc.*, 91(1):76–85, 2015.
- [8] F. Catino and R. Rizzo. Regular subgroups of the affine group and radical circle algebras. *Bull. Austral. Math. Soc.*, 79(1):103–107, 2009.

- [9] V. G. Drinfeld. On some unsolved problems in quantum group theory. In P. P. Kulish, editor, *Quantum groups. Proceedings of workshops held in the Euler International Mathematical Institute, Leningrad, fall* 1990, volume 1510 of *Lecture Notes in Mathematics*, pages 1–8. Springer-Verlag, Berlin, 1992.
- [10] The GAP Group. GAP Groups, Algorithms, and Programming, Version 4.14.0, 2024. http://www.gap-system.org.
- [11] L. Guarnieri and L. Vendramin. Skew-braces and the Yang-Baxter equation. Math. Comp., 86(307):2519–2534, 2017.
- [12] W. Rump. Braces, radical rings, and the quantum Yang-Baxter equation. J. Algebra, 307:153–170, 2007.
- [13] Ya. P. Sysak. The adjoint group of radical rings and related questions. In M. Bianchi, P. Longobardi, M. Maj, and C. M. Scoppola, editors, *Ischia Group Theory 2010. Proceedings of the Conference, Ischia, Naples, Italy, 14 17 April 2010*, pages 344–365, Singapore, September 2011. World Scientific.
- [14] C. N. Yang. Some exact results for many-body problem in one dimension with repulsive delta-function interaction. *Phys. Rev. Lett*, 19:1312–1315, 1967.