OPTIMAL HÖLDER REGULARITY FOR SOLUTIONS TO SIGNORINI-TYPE OBSTACLE PROBLEMS

KI-AHM LEE¹, SE-CHAN LEE², AND WALDEMAR SCHEFER³

ABSTRACT. We study the existence, uniqueness, and regularity of weak solutions to a class of obstacle problems, where the obstacle condition can be imposed on a subset of the domain. In particular, we establish the optimal Hölder regularity for Signorini-type problems, that is, the obstacle condition is imposed only on a subset of codimension one. For this purpose, we employ capacities, Alt–Caffarelli–Friedman-type and Almgren-type monotonicity formulae, and investigate an associated mixed boundary value problem. Further, we apply this problem to study classical obstacle problems for irregular obstacles.

CONTENTS

1

		-
2.	The general obstacle problem	5
3.	Optimal Hölder regularity for Signorini-type problems under a capacity density condition	12
4.	Almost optimal Hölder regularity in the half-hyperplane obstacle case	18
5.	Optimal Hölder regularity in the half-line obstacle case in two dimensions	24
6.	Application: Obstacles with jump-type discontinuities	31
Ret	ferences	32

1. INTRODUCTION

In this article we are interested in the optimal Hölder regularity of solutions to a class of obstacle problems characterized by the Euler-Lagrange equations

(1.1)
$$\begin{cases} -\Delta u \ge 0 & \text{in } \Omega \\ -\Delta u = 0 & \text{in } \Omega \setminus \{u = \psi\} \\ u \ge \psi & \text{on } F \end{cases}$$

for open domains $\Omega \subset \mathbb{R}^n$, $n \geq 2$, relatively closed subsets $F \subset \Omega$, and obstacles ψ defined on F.

In the case that $F = \Omega$, (1.1) coincides with the classical obstacle problem. In the case that $F = \mathcal{M}$, for a manifold \mathcal{M} of codimension one that separates Ω into two parts, (1.1) coincides with the *thin obstacle* problem, also called Signorini problem. There is a vast literature in these two cases as they relate to various problems in applied sciences as physics, biology, mathematical finance, and in pure mathematics within the study of free boundary problems and variational inequalities. We refer to the comprehensive books [DL76,KS80,Fri82,PSU12] and the survey article [FR22] for motivations and applications.

Obstacle problems for more general subsets F appear naturally in the study of capacitor (condenser) potentials, see [Lan72], as the condenser potential of the pair $(\partial\Omega, F)$ solves (1.1), with respect to the obstacle $\psi = 1$, and vanishes at the boundary. Another motivation comes from stochastic control theory, see [BL74, BL75], which gives rise to obstacle problems with irregular obstacles that are neither continuous nor of finite energy. Our main results show the optimal regularity in the case of Signorini-type problems, that is, F is contained in a manifold \mathcal{M} of codimension one. Moreover, we provide a relation between (1.1) with sufficiently regular obstacles and (classical) obstacle problems with discontinuous obstacles.

Introduction

Date: January 28, 2025.

²⁰²⁰ Mathematics Subject Classification. 35B65, 35J86, 35R35.

Key words and phrases. Obstacle problems; Regularity theory; Monotonicity formula; Mixed boundary value problem.

1.1. Main results. Let $\Omega = B_1(0) \subset \mathbb{R}^n$ be the unit ball centered at the origin, F be a subset of the horizontal hyperplane $\{x_n = 0\}$, and $u \in H^1(\Omega)$ be a weak solution to (1.1). The regularity of the obstacle ψ in (1.1) will be suitably chosen depending on the context of each theorem. Then, our main results can be summarized as follows:

- (i) Theorem 3.1: $u \in C^{1/2}$ if the contact set $\Lambda(u) := \{u = \psi\}$ satisfies a capacity density condition.
- (ii) Theorem 4.8: $u \in C^{1/2-}$ if F equals half of the horizontal hyperplane.
- (iii) Theorem 5.11: $u \in C^{1/2}$ if n = 2 and F equals half of the horizontal line.

Here, $u \in C^{1/2-}$ means that $u \in C^{1/2-\varepsilon}$ for any $\varepsilon \in (0, 1/2)$. Due to a simple example, which goes back to [Sha68], the $C^{1/2}$ -regularity is optimal for solutions to Signorini-type problems. See the discussion provided at the end of Section 2. Let us briefly explain the main difference among the results. The benefit of Theorem 3.1 is that no additional assumption on F is required, although it can be difficult to verify the condition in Theorem 3.1 due to the a priori unknown contact set. However, it is noteworthy that various types of sets satisfy a capacity density condition, see Examples 2.15. Further remarks on the contrary case are also provided at the beginning of Section 3. In Theorem 4.8 and Theorem 5.11 we restrict our attention to the special case when F is half of the horizontal hyperplane. While the approach in Theorem 5.11 we have to limit ourselves to the two dimensional setting. For a detailed explanation see the beginning of Section 5 and Remark 5.4. Within the two dimensional setting, we additionally classify the blowup profiles of u at a free boundary point in Theorem 5.10. In polar coordinates, these are given, up to a multiplicative constant, by

$$u_0(r,\theta) = r^{\kappa} \cos(\kappa\theta) \text{ for some } \kappa \in 2\mathbb{N} \text{ or } \kappa \in \mathbb{N}_0 + \frac{1}{2}.$$

Furthermore, as we discuss in Section 6, we relate the regularity of solutions to (1.1) to the regularity of solutions to classical or thin obstacle problems with discontinuous obstacles. More precisely, in Theorem 6.1 we show that if ψ is an obstacle defined in Ω or on \mathcal{M} that can be separated into two obstacles ψ_1 and ψ_2 defined on a partition F_1 and F_2 of Ω or \mathcal{M} and ψ_1 is strictly larger than ψ_2 in a small neighborhood of $x_0 \in \overline{F_1} \cap \overline{F_2}$ (meaning ψ has a jump-type discontinuity at x_0), then the solution to the obstacle problem related to ψ coincides with the solution to (1.1) with respect to ψ_1 and F_1 in a neighbourhood of x_0 . In combination with the regularity results above, this approach shows the optimal regularity for classical and thin obstacle problems with respect to such jump-type discontinuous obstacles. Note, although the regularity results above discuss the zero obstacle case, we provide generalizations in Theorem 3.9, Theorem 4.10, and Remark 5.12.

Let us mention the literature closely related to our results. The well-posedness of (1.1) in terms of continuous solutions vanishing on the boundary was studied in [LS69] for C^1 -regular obstacles. Higher regularity of the solution was investigated in [Lew68, Kin71] for a special case in two dimensions where F was assumed to be a straight line segment. More precisely, the authors proved Lipschitz regularity for solutions u that vanish at the boundary and the obstacle ψ belongs to $C^{1,\alpha}$, is nonnegative, and vanishes at the endpoints of the line F. One can observe that these structural conditions on u and ψ together with the strong minimum principle imply that the endpoint of F cannot be a free boundary point, unless u is a (trivial) zero solution. Indeed, the setting considered in [Lew68, Kin71] is equivalent to a thin obstacle problem and therefore enables them to achieve higher regularity than $C^{1/2}$, see Lemma 6.6 and the comment afterwards. A crucial difference between this result and ours is the fact that we consider arbitrary boundary values without specific relation between ψ and $u|_{\partial\Omega}$, which allows us only to retrieve $C^{1/2}$ -regularity. Since the $C^{1/2}$ -regularity is optimal, our results Theorem 3.1, Theorem 4.8, and Theorem 5.11 generalize the results in [Lew68, LS69, Kin71] in this regard.

In the articles [FM79, FM82, FM84, Mos86, Mos87], the authors studied classical obstacle problems for irregular obstacles, with varying irregularities. Theorem 6.1 provides a new way to study classical obstacle problems for irregular obstacles having jump-type discontinuities. In particular, the combination with our previous regularity results, give the optimal Hölder regularity of solutions for such cases. Let us also mention that the approach in Section 6 is similar to some results in [Ior83, JOS08], although here the authors required higher regularity of the obstacle sets.

1.2. **Historical background.** The classical and thin obstacle problem are quite well understood. Let us provide a brief overview. The classical obstacle problem originated from [Sta64] while the Signorini problem appeared earlier in [Sig59] and [Fic64]. Since then, various authors contributed to these problems. The first systematic treatment of the well-posedness (in weak sense) was done in [LS67] within the context of variational inequalities, which includes both cases. The optimal regularity of solutions to the classical obstacle problem is $C^{1,1}$, see [Fre72]. For the thin obstacle problem, the optimal regularity is Lipschitz continuity and $C^{1,1/2}$ from both sides up to the obstacle, see [Fre75, Fre77, Ric78, AC04]. In subsequent studies, the regularity and structure of the free boundary $\partial \{u > \psi\}$ were analyzed. In the classical case, the combination of results in [Caf77, KN77, CR77, Caf98, Mon03] implies that the free boundary is smooth up to a set of singular points. The local character of free boundary points was characterized in [Caf98] by blowups, saying: Either a free boundary point x_0 is regular, then its blowup is given by the function $x \mapsto (x_1)^2_+$ (up to translation and rotation), or x_0 is singular, then its blowup is given by the function $x \mapsto x_1^2$ (up to translation and rotation). In the thin case, the problem is more involved. The first result in this direction is contained in [Lew72], which shows in the two-dimensional case that the free boundary is a finite amount of intervals provided that the obstacle is real analytic. Another result in this direction was established in [ACS08], showing that the set of regular points in the free boundary is $C^{1,\alpha}$ -regular. Combined with the bootstrap argument in [KPS15], this further implies smoothness of the set of regular points. In contrast to the classical case, the set of regular points may be empty and the set of non-regular points may be large. See also the survey [FR22]. We note that there are many recent articles related to the finer structure of the free boundary that we do not mention.

In the aforementioned articles on the classical and thin obstacle problems, the usual assumption on the obstacle ψ is to be either zero or to be sufficiently regular in a pointwise sense. On the other hand, motivated by the relation to potential theory and stochastic control theory mentioned above, another line of research was dedicated to the study of (classical) obstacle problems for *irregular obstacles*. One of the first articles in this direction was [LS71], where the authors showed that $C^{1,\alpha}$ and Lipschitz regularity of the obstacle translates into the same regularity of the solution. Such pointwise regularity results were extended to finer moduli of continuity in [CK80] and to Dini-continuity of the gradient in [Ok17]. These results intuitively say "the solution is as regular as the obstacle, up to the optimal regularity". Such results were also achieved in the context of Morrey spaces, [Cho91], and Campanato spaces, [Ele07]. Articles directly influenced by potential theory and stochastic control theory were [FM79, FM82, FM84, Mos86, Mos87]. They investigated pointwise regularity of solutions for *discontinuous* obstacles which are not contained in a Sobolev space. More precisely, they assumed the obstacle to satisfy a *one-sided Hölder condition* or a *unilateral regularity condition of Wiener type* and showed continuity, Hölder continuity, and Wiener-modulus of continuity of solutions.

A similar line of research is focused in Calderón-Zygmund type estimates for classical obstacle problems, which roughly means that Sobolev regularity of the obstacle translates into the same Sobolev regularity of the solution. See, for example, [EH10, BDM11, BCO16]. Further articles in this area generalized the treated operators and associated energy functionals, the underlying function spaces, and domain Ω . See, for example, [Gri21, TNH23, BC24] and the references therein. Many of such results are extended to double obstacle, multi-phase, and parabolic problems.

Let us return to the original problem (1.1) for general subsets F of Ω and try to give a complete overview. The first article related to this problem is [Lew68]. Here, the author studied the existence of continuous solutions and the free boundary in a two-dimensional setting with F being a closed straight line segment. The same problem, for an elliptic operator of divergence type, was treated in [Kin71]. The author proved Lipschitz regularity of solutions if the obstacle is $C^{1,\alpha}(F)$ and vanishes at the endpoints of the line segment F. A similar problem was considered in [Giu71] for higher dimensions. In the follow-up article [LS69] to [Lew68], the authors extended the existence of continuous solutions to the case of arbitrary closed subsets F of positive capacity in any dimension. In [SV72] the authors proved Lipschitz regularity in the case when C is a uniformly convex set and ψ a smooth obstacle on $F = \Omega \setminus C$. The articles [Ior73, Ior82] treat double obstacle problems for obstacles φ and ψ defined on subsets E and F, respectively, and the solution u satisfies $u \leq \varphi$ on E and $u \geq \psi$ on F. The author proved α -Hölder continuity of u for $\alpha = 1 - n/p$ if $\varphi, \psi \in W^{1,p}(\Omega), p > n$, and the sets $\Omega \setminus E$ and $\Omega \setminus F$ are Lipschitz domains. The results were extended in the follow-up paper [Ior83] to include finitely many obstacles φ_i and ψ_i on subsets E_i and F_i . A decade later, in the article [BS93] and the following articles, the authors called (1.1) for subsets $F \subset \Omega$ "inner obstacle problems". In [BS93] the authors showed $C^{1,1}$ -regularity of solutions, under the assumptions that F and ψ are sufficiently regular and satisfy an "egg-shape" condition. In [JOS03, JOS08] the authors improved the results from [Ior82, Ior83] to Lipschitz regularity of solutions under the assumptions that obstacles are smooth and obstacle sets are compact subsets of Ω with smooth boundaries. A similar result appeared in the textbook [KS80, p. 137 - 139]. The article [JOS05] dealt with the convergence of solutions if there is a sequence of obstacle sets F_n and obstacles ψ_n that convergence in an appropriate sense. In [JOS09] the authors studied $W^{2,p}$ -regularity of solutions if Ω and F are "strictly star-shaped" and the obstacle has a concave extension to all of Ω .

1.3. **Strategy of the proofs.** Let us briefly illustrate the main strategies to prove Theorem 3.1, Theorem 4.8, and Theorem 5.11. In the proof of Theorem 3.1 the key tool is the monotonicity formula in Proposition 3.4 for an Alt–Caffarelli–Friedman-type frequency function given by

$$\beta(r) = \beta(r, u) \coloneqq \frac{1}{r} \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx.$$

The major difficulty in our situation is the lack of C^1 -regularity of u, which requires an additional effort to obtain the integral identities involved in the proof of Proposition 3.4. Then, we conclude that the capacity density condition (3.1) for the contact set guarantees the optimal regularity of u with the help of Maz'ya's inequality Lemma 3.6. We refer to [ACF84, CJK02] for the original Alt–Caffarelli–Friedman monotonicity formula and some of its variants. The frequency we use was first introduced in [AC04], but used differently, see Remark 3.3.

The key idea in the proof of the almost optimal regularity result in Theorem 4.8 is to establish a connection between Signorini-type problems and mixed boundary value problems (or mixed BVPs for short). To be precise, if we let Ψ be the solution to the associated mixed BVP, then u can be interpreted as a solution to a classical obstacle problem with obstacle Ψ . Note, in the context of thin obstacle problems, this approach is well known by using Dirichlet boundary value problems instead of mixed BVPs. In this view, we first investigate the almost optimal Hölder regularity of solutions to mixed BVPs by constructing appropriate barrier functions, see Theorem 4.7. Then, we transport this regularity to the solution u of (1.1) with the help of the regularity result for classical obstacle problems in [Caf98]. We expect that this argument is also applicable to the viscosity framework, including fully nonlinear operators in non-divergence form, as it does not rely on a weak formulation or integration by parts formulas.

The optimal $C^{1/2}$ -regularity in Theorem 5.11 follows from blowup arguments. For this purpose, we present a Rellich-type formula in Proposition 5.3 to prove the monotonicity of Almgren's frequency function

$$N(r) = N(r, u) \coloneqq \frac{r \int_{B_r} |\nabla u|^2 \, dx}{\int_{\partial B_r} u^2 \, d\sigma}$$

In fact, Almgren's monotonicity formula allows us to utilize the blowup analysis for u: (i) the rescaling u_r of u converge to a limit u_0 in Proposition 5.8, (ii) u_0 is a global homogeneous solution to a Signorini-type problem in Lemma 5.9, and (iii) u has an optimal regularity coming from the classification of blowups in Theorem 5.10. Again, since we cannot expect C^1 -regularity of solutions on F, there are several delicate difficulties when we follow the standard argument developed in [ACS08] for the thin obstacle problems. Indeed, we would like to point out that the lack of C^1 -regularity can be overcome by capturing a kind of degenerate effect in dimension two. For details of the difficulty of this approach in higher dimensions, see Remark 5.4.

1.4. **Overview.** The paper is organized as follows. In Section 2, we introduce the necessary notations, definitions, and preliminary results for general obstacle problems. Section 3, Section 4, and Section 5 are concerned with the proof of the main results Theorem 3.1, Theorem 4.8, and Theorem 5.11, respectively. More precisely, the proofs in these sections rely on the Alt–Caffarelli–Friedman-type monotonicity formula, the analysis of mixed boundary value problems, and Almgren's monotonicity formula, respectively. Possible generalizations to non-zero obstacles are discussed in Theorem 3.9, Theorem 4.10, and Remark 5.12. Finally, in Section 6, we apply these results to study obstacle problems for obstacles with jump-type discontinuities.

Acknowledgements. The third author thanks Florian Grube for valuable discussions regarding some of the technical parts in the proofs of Section 5. Ki-Ahm Lee is supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP): NRF-2020R1A2C1A01006256. Se-Chan Lee is supported by the KIAS Individual Grant (No. MG099001) at Korea Institute for Advanced Study. Waldemar Schefer is supported by the German Research Foundation (DFG GRK 2235 - 282638148).

2. The general obstacle problem

For any open set $\Omega \subset \mathbb{R}^n$, $n \geq 2$, let $H^1(\Omega)$ denote the usual Sobolev space defined as the subspace of $L^2(\Omega)$ with weak derivatives in $L^2(\Omega)$ and norm

$$\|u\|_{H^{1}(\Omega)} = \left(\|u\|_{L^{2}(\Omega)}^{2} + \|\nabla u\|_{L^{2}(\Omega)}^{2}\right)^{\frac{1}{2}}$$

Let $H_0^1(\Omega)$ denote the closure of the set of smooth compactly supported functions $C_c^{\infty}(\Omega)$ in Ω with respect to the $H^1(\Omega)$ -norm. Then, the *relative* 1-capacity of an open set $U \subset \Omega$ is defined by

$$\operatorname{Cap}_{1}(U;\Omega) \coloneqq \inf \left\{ \left\| u \right\|_{L^{2}(\Omega)}^{2} + \left\| \nabla u \right\|_{L^{2}(\Omega)}^{2} \mid u \in H^{1}(\Omega) \text{ such that } u \geq 1 \text{ a.e. on } U \right\},$$

where we set $\operatorname{Cap}_1(U;\Omega)$ as $+\infty$ if there is no such u. Moreover, for an arbitrary set $A \subset \Omega$ we define

$$\operatorname{Cap}_{1}(A;\Omega) \coloneqq \inf_{\substack{A \subset U \subset \Omega \\ U: \text{ open}}} \operatorname{Cap}_{1}(U;\Omega).$$

From [FOT11, Section 2.1], any $u \in H^1(\Omega)$ has a quasi-continuous representative $\tilde{u} : \Omega \to \mathbb{R}$ with respect to Cap₁, which means that for every $\varepsilon > 0$ there exists an open set $G \subset \Omega$ such that Cap₁($G; \Omega) < \varepsilon$, $\tilde{u} \in C(\Omega \setminus G)$, and $\tilde{u} = u$ almost everywhere. For any set $F \subset \Omega$, $u \in H^1(\Omega)$, and any extended real-valued function $\psi : F \to \mathbb{R} := [-\infty, +\infty]$ we write $u \ge \psi$ quasi-everywhere (q.e.) on F if there exists $N \subset F$ such that Cap₁($N; \Omega$) = 0 and $\tilde{u} \ge \psi$ on $F \setminus N$. In particular, [FOT11, Theorem 2.1.5] implies for arbitrary sets $F \subset \Omega$ that

$$\operatorname{Cap}_1(F;\Omega) = \inf \left\{ \|u\|_{L^2(\Omega)}^2 + \|\nabla u\|_{L^2(\Omega)}^2 \mid u \in H^1(\Omega) \text{ such that } u \ge 1 \text{ q.e. on } F \right\}.$$

In the present paper, we consider the following *obstacle problems*:

Definition 2.1 (General obstacle problems). Let $\Omega \subset \mathbb{R}^n$ be open and $F \subset \Omega$. For boundary data $g \in H^1(\Omega)$ and obstacle $\psi : F \to \overline{\mathbb{R}}$ define

$$\mathcal{K}_{g,\psi,F}(\Omega) \coloneqq \left\{ v \in H^1(\Omega) \mid v - g \in H^1_0(\Omega) \text{ and } v \ge \psi \text{ q.e. on } F \right\}.$$

A function $u: \Omega \to \mathbb{R}$ is called a solution to the $\mathcal{K}_{q,\psi,F}(\Omega)$ -obstacle problem if $u \in \mathcal{K}_{q,\psi,F}(\Omega)$ and

$$\int_{\Omega} |\nabla u|^2 \, dx = \inf_{v \in \mathcal{K}_{g,\psi,F}(\Omega)} \int_{\Omega} |\nabla v|^2 \, dx.$$

We call a solution *u* unique if it is unique in $H^1(\Omega)$. For general $F \subset \mathbb{R}^n$ let $\mathcal{K}_{g,\psi,F}(\Omega)$ denote $\mathcal{K}_{g,\psi,F\cap\Omega}(\Omega)$.

If $F = \Omega$, then the $\mathcal{K}_{g,\psi,\Omega}(\Omega)$ -obstacle problem coincides with the original, often called *classical* or *thick*, obstacle problem. If $F = \mathcal{M}$ for an (n-1)-dimensional manifold $\mathcal{M} \subset \Omega$ that separates Ω into two parts, then the $\mathcal{K}_{g,\psi,\mathcal{M}}(\Omega)$ -obstacle problem coincides with a *thin obstacle problem* or also called *Signorini problem*. In both cases, solutions in the sense of Definition 2.1 are often called *weak* or *variational solutions*.

Let us discuss some cases that are included in Definition 2.1.

(i) Assume that $\operatorname{Cap}_1(F;\Omega) = 0$. Then, every function $v \in H^1(\Omega)$ satisfies $v \ge \psi$ q.e. on F. Thus, $\mathcal{K}_{g,\psi,F}(\Omega) = \{v \in H^1(\Omega) \mid v - g \in H^1_0(\Omega)\}$ and therefore the $\mathcal{K}_{g,\psi,F}(\Omega)$ -obstacle problem coincides with the variational formulation of the Dirichlet boundary value problem

$$\begin{cases} -\Delta u = 0 & \text{in } \Omega \\ u = g & \text{on } \partial \Omega. \end{cases}$$

(*ii*) Assume that $\operatorname{Cap}_1(\Omega^c; \mathbb{R}^n) = 0$. Then, [HKM06, Corollary 2.39, Theorems 2.44 and 2.45] imply that $H^1(\Omega) = H^1_0(\Omega) = H^1(\mathbb{R}^n)$. Thus, in this case the condition $v - g \in H^1_0(\Omega)$ in Definition 2.1 always holds and so we have

$$\mathcal{K}_{g,\psi,F}(\Omega) = \{ v \in H^1(\mathbb{R}^n) \mid v \ge \psi \text{ q.e. on } F \}.$$

(*iii*) Let $\Psi : \Omega \to \mathbb{R}$ be an extension of $\psi : F \to \mathbb{R}$ simply by extending as $-\infty$ on $\Omega \setminus F$. Then, for every $u \in H^1(\Omega)$ we have $u \ge \psi$ q.e. on F if and only if $u \ge \Psi$ q.e. on Ω . Thus, we have

$$\mathcal{K}_{g,\psi,F}(\Omega) = \mathcal{K}_{g,\Psi,\Omega}(\Omega).$$

In this sense, Definition 2.1 coincides with the definition of obstacle problems studied in [BB11] and, up to the generality of considered operators, in [FM84, MZ91].

Remark 2.2. In the case $\Omega = \mathbb{R}^n$, $\operatorname{Cap}_1(\cdot; \mathbb{R}^n)$ is sometimes referred as Sobolev capacity, as for example in [HKM06,BB11]. As we are interested in obstacle problems, we usually assume that $\operatorname{Cap}_1(F;\Omega) > 0$ as in the contrary case the obstacle does not have any influence. Note, this implies that $\operatorname{Cap}_1(F;\mathbb{R}^n) > 0$ due to a simple monotonicity argument. On the other hand, whenever Ω is an H^1 -extension domain, the contrary is also true. Thus, for most problems of interest, one can replace $\operatorname{Cap}_1(\cdot;\Omega)$ by the Sobolev capacity. However, for the sake of possible generalizations, we stick to the relative capacity as it provides a more intrinsic perspective.

Let us also introduce the *relative* 0-capacity $\operatorname{Cap}_0(F;\Omega)$, for open sets Ω and arbitrary subsets $F \subset \Omega$, similar to Cap_1 by setting

$$\operatorname{Cap}_{0}(U;\Omega) \coloneqq \inf \left\{ \|\nabla u\|_{L^{2}(\Omega)} \mid u \in H^{1}_{0}(\Omega) \text{ such that } u \geq 1 \text{ a.e. on } U \right\} \quad \text{for } U \subset \Omega \text{ open.}$$

2.1. Existence and uniqueness. The following result is standard and follows from [LS67]:

Theorem 2.3. Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Suppose that $F \subset \Omega$, $g \in H^1(\Omega)$, and $\psi : F \to \overline{\mathbb{R}}$. If $\mathcal{K}_{q,\psi,F}(\Omega)$ is not empty, then there exists a unique solution to the $\mathcal{K}_{q,\psi,F}(\Omega)$ -obstacle problem.

Before we proceed, let us collect typical criteria for the nonemptiness of $\mathcal{K}_{g,\psi,F}(\Omega)$ and therefore the solvability of the corresponding obstacle problem. Adams' criterion from [Ada82] holds in our setting as follows:

Theorem 2.4 (Adams' criterion, [BB11, Theorem 7.3]). Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Suppose that $F \subset \Omega, g \in H^1(\Omega)$, and $\psi: F \to \overline{\mathbb{R}}$. Then, $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$ if and only if

$$\int_0^\infty t\operatorname{Cap}_0\left(\{x\in F\mid \psi(x)-g(x)>t\};\Omega\right)\,dt<\infty.$$

For classical obstacle problems, there exists an easier criterion in the case that $\psi \in H^1(\Omega)$. It says that $\mathcal{K}_{g,\psi,\Omega}(\Omega) \neq \emptyset$ if and only if $(g - \psi)_+ \in H^1_0(\Omega)$, see [BB11, Proposition 7.4]. We can use this criterion if $\psi: F \to \mathbb{R}$ has an extension $\Psi \in H^1(\Omega)$, that is, $\Psi = \psi$ q.e. on F, in the following way:

Proposition 2.5. Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Suppose that $F \subset \Omega$, $g \in H^1(\Omega)$, and $\psi : F \to \overline{\mathbb{R}}$. If ψ has an extension $\Psi \in H^1(\Omega)$ such that $(\Psi - g)_+ \in H^1_0(\Omega)$, then $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$. If $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$ and there exists some extension $\Psi \in H^1(\Omega)$ of ψ , then there exists an extension $\widetilde{\Psi}$ of ψ satisfying $(\Psi - g)_+ \in H^1_0(\Omega)$.

Proof. First, assume that there exists an extension $\Psi \in H^1(\Omega)$ of ψ such that $(\Psi - g)_+ \in H^1_0(\Omega)$. By [BB11, Proposition 7.4], we know that $\mathcal{K}_{g,\Psi,\Omega}(\Omega) \neq \emptyset$. Since for every $f \in \mathcal{K}_{g,\Psi,\Omega}(\Omega)$ we have $f \geq \Psi = \psi$ q.e. on F, we conclude that $f \in \mathcal{K}_{g,\psi,F}(\Omega)$.

Now assume that there exists $f \in \mathcal{K}_{g,\psi,F}(\Omega)$ and there is an extension $\Psi \in H^1(\Omega)$ of ψ . Then, by setting $\widetilde{\Psi} := \min\{f,\Psi\} \in H^1(\Omega)$, we immediately have $\widetilde{\Psi} = \psi$ q.e. on F. At last, $(\Psi - g)_+ \in H^1_0(\Omega)$ follows from the fact that $(f - g)_+ \in H^1_0(\Omega)$ and $0 \le (\Psi - g)_+ \le (f - g)_+$; see for example [HKM06, Lemma 1.25 (ii)]. \Box

Moreover, since F is allowed to be a proper subset of Ω , Proposition 2.5 has a simple application as only the data close to the boundary matters.

Lemma 2.6. Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Suppose that $F \subset \Omega$ and $g, \psi \in H^1(\Omega)$.

- (i) If dist $(F, \partial \Omega) > 0$, then $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$ for every choice $g, \psi \in H^1(\Omega)$.
- (ii) If dist $(F, \partial \Omega) = 0$, then $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$ if there is a neighbourhood U of $\overline{F} \cap \partial \Omega$ such that $g \geq \psi$ q.e. on U.

Proof. Clearly, if dist $(F, \partial \Omega) > 0$, then there exists a cutoff function $\rho \in C_c^{\infty}(\Omega)$ satisfying $\rho = 1$ on F and $\rho = 0$ outside some neighbourhood U of F such that dist $(U, \partial \Omega) > 0$. For $\tilde{\psi} = \rho \psi + (1 - \rho)g$, we have

$$\widetilde{\psi}|_F = \psi \quad \text{and} \quad \widetilde{\psi}|_{\Omega \setminus U} = g.$$

In particular, we have $(\tilde{\psi} - g)_+ \in H^1_0(\Omega)$ and so Proposition 2.5 implies $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$.

Now, assume we are in the case of (*ii*). Take a cutoff function ρ such that $\rho = 1$ on $\overline{F} \setminus U$ and $\rho = 0$ outside a neighbourhood V of \overline{F} . Here we can choose V such that $\operatorname{dist}(V \setminus U, \partial\Omega) > 0$. Again, by letting $\tilde{\psi} = \rho \psi + (1 - \rho)g$, it turns out that

$$\widetilde{\psi}|_{F \setminus U} = \psi, \quad \widetilde{\psi}|_{F \cap U} \ge \psi, \quad \text{and} \quad \widetilde{\psi}|_{\Omega \setminus V} = g.$$

Moreover, since

 $\widetilde{\psi}-g=\rho(\psi-g)\leq 0\quad \text{on }U\text{ and on a small neighbourhood of }\partial\Omega,$

we obtain $(\tilde{\psi} - g)_+ \in H^1_0(\Omega)$ and so Proposition 2.5 implies $\mathcal{K}_{g,\psi,F}(\Omega) \neq \emptyset$.

2.2. Continuity and harmonicity of solutions. The following can be seen as an extension of some results in [LS69], where the authors derived the same statements for bounded domains and $\psi \in C^1(\overline{\Omega})$. However, note that [LS69] treats more general elliptic operators in divergence form, which our prove can be easily generalized to.

Theorem 2.7. Let $\Omega \subset \mathbb{R}^n$ be open, $F \subset \Omega$ be relatively closed, and $\psi : F \to \overline{\mathbb{R}}$ be upper semi-continuous. Assume that $u \in H^1(\Omega)$ is a solution to the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. Then, the following are true:

- (i) u is continuous in Ω and satisfies $u \ge \psi$ on F.
- (ii) u is superharmonic in Ω .
- (iii) u is harmonic in $\Omega \setminus \Lambda(u)$, where $\Lambda(u) := \{x \in F \mid u(x) = \psi(x)\}$ is the contact set.

The proof of Theorem 2.7 consists of two steps. In Proposition 2.8 we show the superharmonicity of u in Ω and the harmonicity of u in $\Omega \setminus F$. On the other hand, in Proposition 2.9 we prove the continuity of u in Ω and the harmonicity of u in $\Omega \setminus \Lambda(u)$.

Proposition 2.8. Let $\Omega \subset \mathbb{R}^n$ be open. Suppose that $F \subset \Omega$, $\psi : F \to \overline{\mathbb{R}}$, and that $u \in H^1(\Omega)$ is a solution to the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. Then, u is harmonic in $\Omega \setminus F$, that is,

$$\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx = 0 \quad \text{for every } \varphi \in H^1_0(\Omega) \text{ with } \varphi = 0 \text{ q.e. on } F,$$

and superharmonic in Ω , that is,

 $\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx \ge 0 \quad \text{for every nonnegative } \varphi \in H^1_0(\Omega).$

Proof. We use standard techniques from calculus of variations. First, we prove harmonicity. For $\varphi \in H_0^1(\Omega)$ such that $\varphi = 0$ q.e. on F we have $u + t\varphi \in \mathcal{K}_{u,\psi,F}(\Omega)$ for every $t \in \mathbb{R}$. Since u is a minimizer of the Dirichlet integral, we have

$$0 = \frac{d}{dt} \int_{\Omega} |\nabla(u+t\varphi)|^2 \, dx \Big|_{t=0} = \frac{d}{dt} \int_{\Omega} \left(|\nabla u|^2 + 2t\nabla u \cdot \nabla \varphi + t^2 |\nabla \varphi|^2 \right) \, dx \Big|_{t=0} = 2 \int_{\Omega} \nabla u \cdot \nabla \varphi \, dx.$$

Next, assume that $\varphi \in H_0^1(\Omega)$ is nonnegative. Then, $u + t\varphi \in \mathcal{K}_{u,\psi,F}(\Omega)$ for every $t \ge 0$. Since $u + t\varphi = (1-t)u + t(u+\varphi)$, the convexity of the Dirichlet integral on $\mathcal{K}_{u,\psi,F}(\Omega)$ implies

$$0 \le \frac{d}{dt} \int_{\Omega} |\nabla(u + t\varphi)|^2 \, dx \Big|_{t=0} = 2 \int_{\Omega} \nabla u \cdot \nabla \varphi \, dx.$$

The harmonicity implies that u is smooth, therefore continuous, away from F. To show the continuity along F we proceed similar to the proof of [Caf98, Theorem 1].

Proposition 2.9. Let $\Omega \subset \mathbb{R}^n$ be open, $F \subset \Omega$ be relatively closed, and $\psi : F \to \overline{\mathbb{R}}$. Assume that $u \in H^1(\Omega)$ is a solution to the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. If ψ is upper semi-continuous on F, then $u \in C(\Omega)$ and u is harmonic in $\Omega \setminus \Lambda(u)$.

Proof. In this proof let us say a subset $A \subset F$ is *F*-open if it is open with respect to the subset topology of *F*. Similarly, we use *F*-closed and the appropriate counterpart for Ω instead of *F*.

The superharmonicity of u from Proposition 2.8 implies that u is lower semi-continuous on Ω after a possible redefinition on a set of measure zero, see [FRRO22, Lemma 1.17]. Thus, the set $\{u > \psi\} := \{x \in F \mid u(x) > \psi(x)\}$ is F-open, as $u - \psi$ is lower semi-continuous on F.

Let us now show that u is harmonic away from $\{u = \psi\} \coloneqq \{x \in F \mid u(x) = \psi(x)\}$. First, harmonicity in $\{u > \psi\}$ follows from a similar argument as in Proposition 2.8: Since $\{u > \psi\}$ is F-open, $\{u \le \psi\} = F \setminus \{u > \psi\}$ is F-closed and therefore $\Omega \setminus \{u \le \psi\}$ is open as F itself is Ω -closed. For any $\varphi \in C_c^{\infty}(\Omega \setminus \{u \le \psi\})$ we have dist(supp $\varphi, \{u \le \psi\}) > 0$. Thus, there exists some open interval $I_{\varphi} \subset \mathbb{R}$ that contains 0 and $u + t\varphi \in \mathcal{K}_{u,\psi,F}(\Omega)$ for every $t \in I_{\varphi}$. By the variational argument from Proposition 2.8 we conclude that $(\nabla u, \nabla \varphi)_{L^2(\Omega)} = 0$. Hence, u is harmonic in $\Omega \setminus \{u \le \psi\}$. The harmonicity in $\{u < \psi\}$ follows from the observation that Cap₁($\{u < \psi\}; \Omega) = 0$ as $u \in \mathcal{K}_{u,\psi,F}(\Omega)$ and therefore

$$\mathcal{K}_{u,\psi,F}(\Omega) = \mathcal{K}_{u,\psi,F\setminus\{u<\psi\}}(\Omega).$$

This means, u coincides with the solution to the $\mathcal{K}_{u,\psi,F\setminus\{u<\psi\}}(\Omega)$ -obstacle problem, which implies the harmonicity away from $\{u=\psi\}$ by Proposition 2.8.

We now show that u is continuous. Since u is harmonic (thus smooth) in $\Omega \setminus \{u = \psi\}$, we claim that u is continuous on $\{u = \psi\}$ as well. Indeed, let $y_0 \in \{u = \psi\}$ and let us argue by contradiction. More precisely, suppose that there exists a sequence $y_k \to y_0$ such that $u(y_k) \to u(y_0) + \varepsilon_0 = \psi(y_0) + \varepsilon_0$ for some $\varepsilon_0 > 0$, where we used the lower semi-continuity of u. Since ψ is upper semi-continuous, we may assume that $y_k \in \{u > \psi\}$ for $k \in \mathbb{N}$. Otherwise, we can extract a subsequence satisfying this property. Let us denote by z_k the projection of y_k towards $\{u = \psi\} \cap F$, that is, $|y_k - z_k| = \inf_{z \in \{u = \psi\}} |y_k - z_k|$. We note that $z_k \in F$ for large k, since F is Ω -closed. If we set $\delta_k := |y_k - z_k| > 0$, then we clearly have $\delta_k \to 0$, $z_k \to y_0$, and $B_{\delta_k}(y_k) \subset B_{2\delta_k}(z_k)$. Thus, the superharmonicity of u from Proposition 2.8 implies

$$u(z_k) \ge \oint_{B_{2\delta_k}(z_k)} u \, dx = (1 - 2^{-n}) \oint_{B_{2\delta_k}(z_k) \setminus B_{\delta_k}(y_k)} u \, dx + 2^{-n} \oint_{B_{\delta_k}(y_k)} u \, dx \eqqcolon I_1 + I_2.$$

In I_2 we use the harmonicity of u away from $\{u = \psi\}$ to get $I_2 = 2^{-n}u(y_k)$. From the lower semi-continuity there exists an open neighbourhood U of y_0 in Ω such that

(2.1)
$$u(x) \ge u(y_0) - 2^{-n} \varepsilon_0 \text{ for every } x \in U.$$

For large k, we have $B_{2\delta_k}(z_k) \subset U$. Thus, we can use (2.1) in I_1 to get

(2.2)
$$u(z_k) \ge (1 - 2^{-n})u(y_0) - (1 - 2^{-n})2^{-n}\varepsilon_0 + 2^{-n}u(y_k)$$

By assumption we have $\lim_{k\to\infty} u(y_k) = u(y_0) + \varepsilon_0$. If $z_k \in \{u = \psi\}$ we could use $u(z_k) = \psi(z_k)$ together with the upper semi-continuity of ψ to deduce

(2.3)
$$\psi(y_0) \ge \limsup_{k \to \infty} \psi(z_k) = \limsup_{k \to \infty} u(z_k) \ge u(y_0) + 2^{-2n} \varepsilon_0 = \psi(y_0) + 2^{-2n} \varepsilon_0,$$

which leads to a contradiction. However, it is unclear if $z_k \in \{u = \psi\}$ for every k (or some subsequence). But, we know that z_k is contained in the F-closure of $\{u = \psi\}$ for large enough k. In this case, we can use the lower semi-continuity of u and upper semi-continuity of ψ to deduce the following: For x in the F-closure of $\{u = \psi\}$ and any sequence $x_n \in \{u = \psi\}$ converging to x we have

$$\psi(x) \ge \limsup_{n \to \infty} \psi(x_n) = \limsup_{n \to \infty} u(x_n) \ge \liminf_{n \to \infty} u(x_n) \ge u(x).$$

Hence, we have $\psi(z_k) \ge u(z_k)$ that we can use in (2.2) to deduce (2.3).

As a direct consequence of Theorem 2.7, we have shown that any solution $u \in H^1(\Omega)$ to the $\mathcal{K}_{u,\psi,F}(\Omega)$ obstacle problem satisfies the Euler-Lagrange equations

(2.4)
$$\begin{cases} -\Delta u \ge 0 & \text{in } \Omega \\ -\Delta u = 0 & \text{in } \Omega \setminus \Lambda(u) \\ u - \psi \ge 0 & \text{on } F. \end{cases}$$

In fact, these problems are equivalent.

Lemma 2.10. Let $\Omega \subset \mathbb{R}^n$ be open, $F \subset \Omega$ be relatively closed, and $\psi : F \to \overline{\mathbb{R}}$ be upper semi-continuous. Assume that $u \in H^1(\Omega) \cap C(\Omega)$. Then, u satisfies (2.4) if and only if u solves the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. *Proof.* If u solves the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem, then it satisfies (2.4) by Theorem 2.7.

To show the contrary, we first show that solutions to (2.4) are unique in $H^1(\Omega) \cap C(\Omega)$ respective to their boundary values. Assume we have two functions $u_1, u_2 \in H^1(\Omega) \cap C(\Omega)$ satisfying (2.4) and $u_1 - u_2 \in H^1_0(\Omega)$. Let $G := \{u_2 > u_1\}$. Note, G is open due to the continuity of u_1 and u_2 . Moreover, since

$$G \subset \Omega \setminus \Lambda(u_2)$$

we observe that

$$-\Delta u_2 = 0$$
 in G .

Thus, if $G \neq \emptyset$ then $w \coloneqq u_2 - u_1 \in H^1_0(G)$ satisfies in a weak sense:

$$\begin{cases} -\Delta w \le 0 & \text{in } G \\ w = 0 & \text{on } \partial G. \end{cases}$$

Then, the maximum principle yields that $w = u_2 - u_1 \leq 0$ in G, which leads to a contradiction. As the same argument holds if we exchange the roles of u_1 and u_2 in the definition of G, we conclude $u_1 = u_2$.

To finish the proof, let $u \in H^1(\Omega) \cap C(\Omega)$ satisfy (2.4). Then, $u \in \mathcal{K}_{u,\psi,F}(\Omega)$ and Theorem 2.3 gives the existence of a solution $v \in H^1(\Omega)$ to the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. By Theorem 2.7, v satisfies (2.4) and we have $u - v \in H^1_0(\Omega)$. Thus, from the previous line of arguments, u = v and therefore u solves the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem.

2.3. Hölder and Lipschitz regularity. The aim of this section is to discuss preliminary results on the regularity of solutions to the obstacle problem implied by known regularity results for associated Dirichlet boundary value problems. To be precise, we show Hölder regularity of some order in the case that F satisfies a capacity density condition and Lipschitz regularity in the case that F satisfies a uniform exterior ball condition.

Definition 2.11.

(i) We say that a set $E \subset \mathbb{R}^n$ satisfies a *capacity density condition* if there exist constants $c_0 > 0$ and $r_0 > 0$ such that

(2.5)
$$\frac{\operatorname{Cap}_{0}(E \cap B_{r}(x_{0}), B_{2r}(x_{0}))}{\operatorname{Cap}_{0}(B_{r}(x_{0}), B_{2r}(x_{0}))} \ge c_{0}$$

whenever $0 < r < r_0$ and $x_0 \in E$.

(ii) We say that a set $E \subset \mathbb{R}^n$ satisfies a uniform exterior ball condition (with radius r_0) if there exists $r_0 > 0$ satisfying the following condition: For every $x_0 \in \partial E$, there exists a point $y_0 \in E^c$ such that $B_{r_0}(y_0) \subset E^c$ and $x_0 \in \partial B_{r_0}(y_0)$.

While the exterior ball condition is quite intuitive, it is noteworthy that there are also several geometric conditions that imply (2.5). We collect a small list in Examples 2.15. Now, let us prove Hölder and Lipschitz regularity of solutions to obstacle problems when a capacity density condition or an exterior ball condition holds, respectively.

Theorem 2.12. Let $\Omega \subset \mathbb{R}^n$ be open and bounded, $F \subset \Omega$ be relatively closed, and $\psi : F \to \overline{\mathbb{R}}$. Let $u \in H^1(\Omega)$ be a bounded solution to the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem. Assume there exists $\Psi \in H^1(\Omega \setminus F)$ such that

$$\Psi = \psi$$
 q.e. on F and $\Psi \leq u$ on $\partial \Omega$.

(i) If $\Psi \in C^{\alpha}(\partial(\Omega \setminus F))$ for some $\alpha \in (0,1)$ and $(\Omega \setminus F)^c$ satisfies the capacity density condition (2.5), then there exists $\alpha_0 \in (0,\alpha]$ such that u is C^{α_0} -regular in Ω . Moreover, for any compact set $K \subset \Omega$ there is C > 0 such that

$$\|u\|_{C^{\alpha_0}(K)} \le C(\|u\|_{L^{\infty}(\Omega)} + \|\Psi\|_{C^{\alpha}(\partial(\Omega\setminus F))}).$$

(ii) If $\Psi \in \text{Lip}(\partial(\Omega \setminus F))$ and $\Omega \setminus F$ satisfies a uniform exterior ball condition, then u is Lipschitz continuous in Ω . Moreover, for any compact set $K \subset \Omega$ there is C > 0 such that

$$\|u\|_{\operatorname{Lip}(K)} \le C(\|u\|_{L^{\infty}(\Omega)} + \|\Psi\|_{\operatorname{Lip}(\partial(\Omega\setminus F))}).$$

Proof. First, assume that Ψ is Hölder continuous and $(\Omega \setminus F)^c$ satisfies (2.5). Let $v \in H^1(\Omega \setminus F)$ be the unique weak solution to the Dirichlet boundary value problem

$$\begin{cases} -\Delta v = 0 & \text{in } \Omega \setminus F \\ v = \Psi & \text{on } \partial(\Omega \setminus F). \end{cases}$$

Since $\Psi \in C^{\alpha}(\partial(\Omega \setminus F))$, [HKM06, Theorem 6.44] implies the existence of $\alpha_0 \in (0, \alpha]$ such that $v \in C^{\alpha_0}(\overline{\Omega \setminus F})$. Let

$$\overline{v} \coloneqq \begin{cases} v & \text{in } \Omega \setminus F \\ \psi & \text{on } F. \end{cases}$$

As $u \ge \Psi = v$ on $\partial(\Omega \setminus F)$ the comparison principle implies that $u \ge \overline{v} = v$ in $\Omega \setminus F$. Clearly, $u \ge \overline{v} = \psi$ on F. Thus, u solves the $\mathcal{K}_{u,\overline{v},\Omega}(\Omega)$ -obstacle problem. Since it corresponds to a classical obstacle problem, [Caf98, Theorem 2] implies that u is as regular as the obstacle \overline{v} as long as we stay away from $\partial\Omega$. Thus, we conclude that $u \in C^{\alpha_0}(K)$ for any compact subset $K \subset \Omega$ with the claimed bound.

In the case that $\Psi \in \text{Lip}(\partial(\Omega \setminus F))$ and $\Omega \setminus F$ satisfies a uniform exterior ball condition the line of arguments stays the same but we replace [HKM06, Theorem 6.44] by a variant of [Saf08, Lemma 1.2], see Lemma 2.13 below.

Lemma 2.13. Let $\Omega \subset \mathbb{R}^n$ be open and bounded. Assume that Ω satisfies a uniform exterior ball condition. Then, every $u \in H^1(\Omega) \cap C(\overline{\Omega})$ such that $-\Delta u = 0$ and $u|_{\partial\Omega} \in \operatorname{Lip}(\partial\Omega)$ is Lipschitz on $\overline{\Omega}$ and satisfies the estimate

$$\|u\|_{\operatorname{Lip}(\overline{\Omega})} \le C\left(\|u\|_{L^{\infty}(\Omega)} + \|u\|_{\operatorname{Lip}(\partial\Omega)}\right)$$

for a constant C > 0 depending on n, Ω , and r_0 from the uniform exterior ball condition.

Sketch of proof. Use a barrier approach as in [Saf08, Lemma 1.2]: For $x_0 \in \partial \Omega$ consider the functions

$$\varphi_{\pm}(x) \coloneqq u(x_0) \pm c_{\pm} \left(r_0^{-\lambda} - |x - y_0|^{-\lambda} \right),$$

where r_0 is the radius of the uniform exterior ball condition, y_0 is the center of the exterior ball at x_0 , $\lambda \ge n-2$, and c_{\pm} are positive constants chosen large enough such that $\varphi_{-} \le u \le \varphi_{+}$ on $\partial\Omega$. Then, the claimed result follows from the comparison principle.

Remark 2.14.

- (i) The boundedness of solutions to the obstacle problem follows immediately from the boundedness of their boundary data as well as the boundedness of the obstacle (from above) due to the comparison principle, see for example [HKM06, Theorem 3.24] or [BB11, Proposition 7.5]. On the other hand, due to Theorem 2.7 we already know that the solution u of an obstacle problem in Ω with upper semi-continuous obstacle ψ is continuous in Ω , which implies that it is bounded in any compact subset.
- (*ii*) In Theorem 2.12 (*i*), a sufficient condition such that $(\Omega \setminus F)^c$ satisfies a capacity density condition is that both Ω^c and F satisfy a capacity density condition. In Theorem 2.12 (*ii*), $\Omega \setminus F$ satisfies a uniform exterior ball condition if Ω satisfies a uniform exterior ball condition and F satisfies a uniform interior ball condition.
- (*iii*) As the proof of Theorem 2.12 depends only on the regularity of an associated Dirichlet boundary value problem in $\Omega \setminus F$, it is expected that the Lipschitz regularity holds in the case that $\Omega \setminus F$ satisfies a uniform exterior $C^{1,\text{Dini}}$ -condition, see [KK73,KK74]. In particular, the exterior $C^{1,\text{Dini}}$ -condition is necessary for the Lipschitz regularity, see [KK74, Theorem 2]. See also [HLW14] and the references therein.

Examples 2.15. Here, let us collect some examples for sets F that satisfy a uniform exterior ball or capacity density condition.

- (i) It is well known, see for example [AKSZ07, Lemma 2.2], that a bounded domain is a $C^{1,1}$ -domain if and only if it satisfies a uniform (interior and exterior) ball condition.
- (*ii*) A uniform exterior ball condition is satisfied by many domains having bad interior regularity, such as any convex Lipschitz domain or a convex domain with an outward cusp.

- (*iii*) If F contains a cone (or a corkscrew) in a uniform sense at each point $x_0 \in \partial F$, then F satisfies a capacity density condition, see [HKM06, Theorem 6.31] for instance. In particular, any Lipschitz domain satisfies a capacity density condition.
- (iv) A large class of fractals satisfy a capacity density condition. More precisely, any nonempty selfsimilar compact set given as the attractor of an iterated function system of similitudes (see for example [Fal14]) with Hausdorff-dimension $d \in (n - 2, n]$ satisfies a capacity density condition. This includes, ignoring the dimensional constraint, for example Cantor sets, Sierpiński gaskets and carpets, the Vicsek set, and so on. A proof of this fact may be found in [ACM22, Section 3] in the particular case of the middle third Cantor set. However, the proof in the general case is the same.

This list of examples shows, focusing on the obstacle, that whenever F and ψ are "nice" in the sense that F has a nonempty interior and satisfies a uniform interior ball condition and ψ is Lipschitz continuous, we can expect the solutions to the associated obstacle problems to be of at least Lipschitz regularity.

On the other hand, as the capacity density condition is satisfied by a large variety of sets F we can expect some Hölder regularity in most cases of interest. In the upcoming sections we discuss *Signorini-type* problems, that is, F is contained in an (n - 1)-dimensional manifold that separates Ω into two disjoint parts. Our results, in particular Theorem 3.1, suggest that the optimal (local) regularity of solutions to such Signorini-type problems is $C^{1/2}$ under an additional capacity density condition on F.

One might expect better regularity if F is not contained in an (n-1)-dimensional manifold. However, here it seems that a classification of obstacle sets F and their associated optimal regularity is more delicate as it involves the angles of ∂F , similar to the boundary regularity problem for Dirichlet boundary value problems. To aid that claim, let us extend the standard examples for thin-obstacle problems to specific Lipschitz domains F.

Fix n = 2 and consider for every $0 < \alpha \leq 1$ the function $h_{\alpha}(x, y) = -\operatorname{Re}(x + i|y|)^{\alpha}$ or represented in polar coordinates by

$$h_{\alpha}(r,\theta) = -r^{\alpha}\cos(\alpha\theta).$$

It is easy to see, using the Laplacian in polar coordinates, that h_{α} is harmonic in the upper and lower half-plane. Furthermore, the normal derivative of h_{α} on the right side $\{x > 0, y = 0\}$ is zero. Thus, h_{α} is harmonic along the right side, which means

$$-\Delta h_{\alpha} = 0 \quad \text{in } \mathbb{R}^2 \setminus \{ x \le 0, y = 0 \}.$$

Moreover, we can see for r > 0 and $\theta \in [0, \pi]$ that

$$h_{\alpha}(r,\theta) \geq 0$$
 if and only if $\theta \in \left[\frac{\pi}{2\alpha},\pi\right]$

and

$$h_{\alpha}(r,\theta) = 0$$
 if and only if $\theta = \frac{\pi}{2\alpha}$.

Thus, h_{α} can be expected to solve a zero obstacle problem around the origin only if $\alpha \geq 1/2$ as else we have $h_{\alpha} < 0$ on $\{x < 0, y = 0\}$. Hence, we can easily determine when h_{α} , $\alpha \geq 1/2$, solves some zero obstacle problem in $\Omega = B_1(0)$: Let F_{α} be the closed cone at the origin that opens to the left with aperture $\phi = \pi(1 - \frac{1}{2\alpha})$, see Figure 1. Then, h_{α} solves the $\mathcal{K}_{h_{\alpha},0,F_{\alpha}}(B_1(0))$ -obstacle problem.



FIGURE 1. The unit ball with cone F_{α} .

Clearly, h_{α} is C^{α} -regular at the origin. In particular, when $\alpha = 1/2$, the cone $F_{1/2}$ becomes simply the left half-line $\{x \leq 0, y = 0\}$. This simple example limits the optimal regularity in the Signorini-type case to $C^{1/2}$.

For any $\alpha \in (1/2, 1)$, F_{α} is a Lipschitz domain and the regularity at the origin clearly depends on the aperture of the cone. That means, similar to the boundary regularity problem for Dirichlet boundary value problems on Lipschitz domains, the optimal regularity for the obstacle problem in the case that F is a Lipschitz domain depends heavily on the Lipschitz constants of the representing chart as this determines the possible range of angles. In particular, if the aperture of the cone is larger, that is $\phi \ge \pi/2$, then we expect higher regularity than Lipschitz as it is known for the Dirichlet problem. For this, we refer to [MR10] for the weighted L^p -estimates and related embedding results concerning Dirichlet boundary value problems on Lipschitz domains, see also [JK95].

3. Optimal Hölder regularity for Signorini-type problems under a capacity density condition

In this section we consider the unit ball $B_1 := B_1(0) \subset \mathbb{R}^n$, $n \ge 2$, and a relatively closed subset $F \subset B_1$ supported on the horizontal hyperplane $\{x_n = 0\}$, that is, $F \subset B_1 \cap \{x \in B_1 \mid x_n = 0\}$. Let us abbreviate $B'_r := B_r(0) \cap \{x \in B_r(0) \mid x_n = 0\}$. With these special choices of Ω and F, we call the corresponding $\mathcal{K}_{g,\psi,F}(B_1)$ -obstacle problem, for boundary data g and obstacle ψ , a Signorini-type problem.

We now investigate the local regularity of a solution u to the Signorini-type problem for the zero obstacle $\psi = 0$. We consider non-zero obstacles at the end of the section in Theorem 3.9. Let $u \in H^1(B_1)$ be a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. If F coincides with the full horizontal hyperplane, that is, $F = B_1 \cap \{x \in B_1 \mid x_n = 0\}$, then u is a solution to the classical Signorini problem. In this case, as already mentioned, the optimal regularity of u is well known, that is, u is Lipschitz continuous along the hyperplane and $C^{1,1/2}$ from above and below up to the hyperplane. However, in the case that F is a proper subset, the example at the end of Section 2 shows that we can not expect better than $C^{1/2}$ -regularity.

Recall, the contact set is denoted by $\Lambda(u) = \{x \in F \mid u(x) = 0\}$. The main result of this section is:

Theorem 3.1. Let $n \ge 2$ and $F \subset \{x \in B_1 \mid x_n = 0\}$ be relatively closed in B_1 . Assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. If $\Lambda(u)$ satisfies a capacity density condition at $x_0 \in \Lambda(u)$, that is, there exists a constant $c_0 > 0$ and a radius $r_0 > 0$ such that

(3.1)
$$\operatorname{Cap}_0(\Lambda(u) \cap B_r(x_0); B_{2r}(x_0)) \ge r^{n-2}c_0 \quad \text{for every } r \in (0, r_0),$$

then u is $C^{1/2}$ -regular at x_0 .

The remainder of this section is devoted to the proof of Theorem 3.1, but let us first make some comments on Theorem 3.1 and its proof. As the contact set $\Lambda(u)$ is unknown a priori, (3.1) is difficult to verify. However, we want to emphasize the fact that (3.1) is not as restrictive as it may seem. As already mentioned in Examples 2.15 (*iii*) and (*iv*) a uniform capacity density condition, in the sense of Definition 2.11, is satisfied by a large variety of sets including various fractals. Note, (3.1) is truly a local variant of Definition 2.11 as $\operatorname{Cap}_0(B_r(x_0), B_{2r}(x_0)) = cr^{n-2}$ for some constant c independent of r, see for example [HKM06, (2.13)]. Thus, it can be expected that (3.1) holds for a large class of solutions.

Nonetheless, let us discuss some of the contrary cases. First, in a simple case when (3.1) is not satisfied is the following. Assume $u \in H^1(B_1)$ solves the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem such that there exists $\delta > 0$ satisfying $\Lambda(u) \cap B_{\delta}(0) = \{0\}$. Clearly, (3.1) fails in $x_0 = 0$. However, u is harmonic and bounded in $B_{\delta}(0) \setminus \{0\}$ due to Theorem 2.7. Thus, by the removable singularity theorem for harmonic functions, uextends uniquely to a harmonic function in $B_{\delta}(0)$ and is therefore smooth in $x_0 = 0$. A more general version of this removable singularity result, and therefore smoothness of the solutions, also holds if $\Lambda(u) \cap B_{\delta}(x_0)$ is of zero capacity, see [Ser64].

Thus, it remains to study the case when $r^{-(n-2)} \operatorname{Cap}_0(\Lambda(u) \cap B_r(x_0); B_{2r}(x_0))$ is a proper sequence converging to zero. We comment on this case in Remark 3.8.

Remark 3.2. A variant of (3.1) also appears in the articles [ACM22, AC22, AC23] within the context of mixed boundary value problems. See Section 4 and Remark 4.9 for the relation of these articles to Signorini-type problems.

The strategy of the proof of Theorem 3.1 is similar to the proof of [AC04, Theorem 5]. First, in Section 3.1 we introduce an Alt–Caffarelli–Friedman-type frequency function and show its monotonicity in Proposition 3.4. The main difficulty here is the lack of regularity along the hyperplane. The key observation is that the frequency function is absolutely continuous. Thus, any equation involving the derivative of the frequency function holds in an almost everywhere sense. Then, we conclude the proof in Section 3.2 with the help of Maz'ya's inequality, see Lemma 3.6, and the subharmonicity of the positive and negative part of the solution.

Remark 3.3.

(i) The Alt–Caffarelli–Friedman frequency function was first introduced in [ACF84] in the context of a two-phase free boundary problem. There, it is given by

$$J(r,u) = \frac{1}{r^4} \int_{B_r} \frac{|\nabla u_+|^2}{|x|^{n-2}} \, dx \int_{B_r} \frac{|\nabla u_-|^2}{|x|^{n-2}} \, dx,$$

where u denotes the solution of a certain free boundary problem. The monotonicity of J allows the authors to prove regularity of solutions and identify blowup limits.

(ii) In [AC04], see also the lecture notes [Sal12], the authors introduces a variant of the Alt–Caffarelli– Friedman frequency, namely

$$\beta(r,w) = \frac{1}{r} \int_{B_r^+} \frac{|\nabla w|^2}{|x|^{n-2}} \, dx$$

to study thin obstacle problems. One major difference to the Alt–Caffarelli–Friedman frequency is that w represents $\partial_{x_n} u$, where u solves a global thin obstacle problem. Additionally, it is assumed that u is convex along every direction tangential to the hyperplane $\{x_n = 0\}$. The monotonicity of β then implies the $C^{1/2}$ -regularity of w, which implies the optimal $C^{1,1/2}$ -regularity of the solution u in B_1^+ up to the hyperplane.

3.1. Alt–Caffarelli–Friedman-type frequency function.

Proposition 3.4 (Monotonicity of Alt–Caffarelli–Friedman-type frequency function). Let $F \subset B_1 \cap \{x_n = 0\}$ be relatively closed in B_1 and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. The function

$$\beta(r) = \beta(r, u) \coloneqq \frac{1}{r} \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx \quad \text{for } r \in (0, 1).$$

is nonnegative and finite for every $r \in (0,1)$ and $r \mapsto \beta(r)$ is continuous and nondecreasing.

The proof of Proposition 3.4 is similar to [AC04, Lemma 4]. That means, we take the derivative of β with respect to r and prove its nonnegativity using a lower bound on the Rayleigh quotient associated with the spherical Laplacian. As already mentioned, the main difficulty is the lack of regularity along the hyperplane $\{x_n = 0\}$. The key observation is that β is absolutely continuous. Thus, every identity that involves the derivative of β holds in an almost everywhere sense.

An important ingredient in the proof of Proposition 3.4 is the following integration by parts formula:

Lemma 3.5. Let $F \subset B_1 \cap \{x_n = 0\}$ be relatively closed in B_1 and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. Then, for almost every 0 < r < 1 we have

(3.2)
$$\int_{B_r} \nabla u \cdot \nabla \varphi \, dx = \int_{\partial B_r} \varphi \partial_\nu u \, d\sigma \quad \text{for every } \varphi \in H^1(B_1) \text{ such that } \varphi = 0 \text{ q.e. on } \Lambda(u).$$

Here, σ denotes the surface measure.

Before we prove Lemma 3.5 and Proposition 3.4 let us make some simple observations for solutions u to such Signorini-type problems:

- (i) The gradient $\nabla u(x)$ is well defined pointwise for every $x \in B_1 \setminus \Lambda(u)$.
- (ii) For every $r \in (0,1)$ the normal derivative $\partial_{\nu} u(x) = \nabla u(x) \cdot \vec{\nu}(x)$ is well defined for $x \in \partial B_r \setminus B'_1$ where $\vec{\nu}(x)$ denotes the outward normal with respect to B_r . (*iii*) The maps $r \mapsto \int_{\partial B_r} |\nabla u|^2 d\sigma$ and $r \mapsto \int_{\partial B_r} |\partial_{\nu} u|^2 d\sigma$ are in $L^1(0,1)$.
- (iv) For almost every $r \in (0, 1)$ we have $|\nabla u|, \partial_{\nu} u \in L^2(\partial B_r, \sigma)$.

We note that items (i) and (ii) follow from the fact that u is harmonic (and therefore smooth) away from the contact set $\Lambda(u)$. In particular, ∇u and $\partial_{\nu} u$ are well defined pointwise in $B_1 \setminus B'_1$. Item (iii) follows by using $|\partial_{\nu} u|^2 \leq |\nabla u|^2$ and $u \in H^1(B_1)$ and (iv) is implied by (iii).

Proof of Lemma 3.5. Let us assume that u is symmetric with respect to the hyperplane $\{x_n = 0\}$. Since u is harmonic in $B_1 \setminus \Lambda(u)$ it is smooth up to the boundary in

$$B_{r,\varepsilon}^+ \coloneqq \{x = (x', x_n) \in B_r \mid x_n > \varepsilon\} \text{ for } \varepsilon > 0.$$

For any $\varphi \in C_c^{\infty}(B_1 \setminus \Lambda(u))$, integration by parts implies

(3.3)
$$\int_{B_r^+} \nabla u \cdot \nabla \varphi \, dx = \lim_{\varepsilon \to 0} \int_{B_{r,\varepsilon}^+} \nabla u \cdot \nabla \varphi \, dx = \lim_{\varepsilon \to 0} \int_{\partial B_{r,\varepsilon}^+} \varphi \partial_{\nu} u \, d\sigma.$$

We then separate $\partial B_{r,\varepsilon}^+$ into

$$(\partial B_r)_{\varepsilon}^+ = \{ x \in \partial B_r \mid x \in \partial B_{r,\varepsilon}^+ \} \quad \text{and} \quad (B_r')_{\varepsilon}^+ = \{ x = (x',\varepsilon) \mid |x'| < r \}.$$

Let us now show that

(3.4)
$$\lim_{\varepsilon \to 0} \int_{(B'_r)^+_{\varepsilon}} \varphi \partial_{\nu} u \, d\sigma = 0.$$

There exists a smooth open neighbourhood U of $\Lambda(u)$ such that $\varphi = 0$ on U, and thus

$$\int_{(B'_r)^+_{\varepsilon} \cap U} \varphi \partial_{\nu} u \, d\sigma = 0.$$

Since u is harmonic in $B_1 \setminus U$, we have

$$\lim_{\varepsilon \to 0} \int_{(B'_r)^+_{\varepsilon} \setminus U} \varphi \partial_{\nu} u \, d\sigma = \int_{B'_r \setminus U} \varphi \partial_{\nu} u \, d\sigma.$$

The symmetry of u together with the harmonicity implies $\partial_{\nu} u = 0$ on $B'_r \setminus U$. Thus, we arrive at (3.4). Similarly, for $(\partial B_r)^+_{\varepsilon}$ we have

(3.5)
$$\lim_{\varepsilon \to 0} \int_{(\partial B_r)_{\varepsilon}^+} \varphi \partial_{\nu} u \, d\sigma = \lim_{\varepsilon \to 0} \int_{(\partial B_r)_{\varepsilon}^+ \cap U} \varphi \partial_{\nu} u \, d\sigma + \int_{(\partial B_r)_{\varepsilon}^+ \setminus U} \varphi \partial_{\nu} u \, d\sigma = 0 + \int_{(\partial B_r)^+} \varphi \partial_{\nu} u \, d\sigma.$$

Thus, using (3.4) and (3.5) in (3.3) imply

(3.6)
$$\int_{B_r^+} \nabla u \cdot \nabla \varphi \, dx = \int_{(\partial B_r)^+} \varphi \partial_\nu u \, d\sigma \quad \text{for every } \varphi \in C_c^\infty(B_1 \setminus \Lambda(u))$$

By the same arguments, (3.6) is also true when the upper half ball B_r^+ is replaced by the lower half B_r^- . Now, for any $r \in (0,1)$ such that $\partial_{\nu} u \in L^2(\partial B_r)$ we conclude (3.2) for every $\varphi \in H^1(B_1)$ such that $\varphi = 0$ q.e. on $\Lambda(u)$ by approximation with functions in $C_c^{\infty}(B_1 \setminus \Lambda(u))$.

In the case when u is not symmetric the proof is analogous. Let us only comment on the crucial difference. The normal derivative of u at $B'_r \setminus \Lambda(u)$ does not need to vanish. However the sum of normal derivatives $\partial^+_{\nu}u$ and $\partial^-_{\nu}u$ calculated from B^+_r and B^-_r , respectively, vanishes. Thus, (3.4) and (3.6) stay true when considering the sum of integrals with respect to B^+_r and B^-_r .

The proof of the monotonicity of the Alt–Caffarelli–Friedman-type frequency β in Proposition 3.4 consists of the following three steps:

- (i) We show that $\beta \in [0, +\infty)$.
- (*ii*) We show that β is absolutely continuous on (0, 1).
- (*iii*) We show that $\beta' \geq 0$ almost everywhere.

Proof of Proposition 3.4. First, let us show that $\beta(r) < +\infty$ for every 0 < r < 1. For this purpose, choose an approximation $g_{\varepsilon} \in C^{\infty}(\mathbb{R}^n)$ for $g(x) = |x|^{-(n-2)}$ and $\varepsilon > 0$ such that:

- (i) $g_{\varepsilon} = g$ on B_{ε}^c .
- (*ii*) $g_{\varepsilon} \nearrow g$ pointwise.
- (*iii*) $-\Delta g_{\varepsilon} \to c\delta_0$, where δ_0 is the Dirac delta measure at x = 0 and c > 0 is some constant depending only on the dimension.

By the chain rule and integration by parts we have

$$\int_{B_r} |\nabla u|^2 \varphi \, dx = \int_{B_r} \nabla u \cdot \nabla (u\varphi) \, dx + \frac{1}{2} \int_{B_r} u^2 \Delta \varphi \, dx - \frac{1}{2} \int_{\partial B_r} u^2 \partial_\nu \varphi \, d\sigma$$

for every $\varphi \in C^{\infty}(B_1)$ and 0 < r < 1. In particular, for $\varphi = g_{\varepsilon}$ with $\varepsilon < r$ we have

(3.7)
$$\int_{B_r} |\nabla u|^2 g_{\varepsilon} \, dx = \int_{B_r} \nabla u \cdot \nabla (ug_{\varepsilon}) \, dx - \frac{1}{2} \int_{B_r} u^2 (-\Delta g_{\varepsilon}) \, dx + \frac{n-2}{2r^{n-1}} \int_{\partial B_r} u^2 \, d\sigma.$$

Since $ug_{\varepsilon} \in H^1(B_1)$ and $ug_{\varepsilon} = 0$ on $\Lambda(u)$, Lemma 3.5 allows us to rewrite the first term on the right-hand side to

(3.8)
$$\int_{B_r} \nabla u \cdot \nabla (ug_{\varepsilon}) \, dx = \int_{\partial B_r} ug_{\varepsilon} \partial_{\nu} u \, d\sigma = \frac{1}{r^{n-2}} \int_{\partial B_r} u \partial_{\nu} u \, d\sigma \quad \text{for almost every } \varepsilon < r < 1.$$

Taking the limit $\varepsilon \searrow 0$ we conclude from (3.7) and (3.8) that

(3.9)
$$\int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} dx = \frac{1}{r^{n-2}} \int_{\partial B_r} u \partial_\nu u \, d\sigma - \frac{c}{2} u(0)^2 + \frac{n-2}{2r^{n-1}} \int_{\partial B_r} u^2 \, d\sigma \quad \text{for almost every } 0 < r < 1.$$

In particular, there exists a radius r arbitrarily close to 1 such that the right-hand side of (3.9) exists. Since $r \mapsto \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} dx$ is nondecreasing, it is finite for every 0 < r < 1 due to (3.9) and therefore $\beta(r) < +\infty$ for every 0 < r < 1.

Next, let us show that β is absolutely continuous on (0,1) with almost everywhere derivative

(3.10)
$$\beta'(r) = \frac{1}{r^{n-1}} \int_{\partial B_r} |\nabla u|^2 \, d\sigma - \frac{1}{r^2} \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx.$$

It is immediate that $r \mapsto \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx$ is absolutely continuous as

$$s \mapsto \int_{\partial B_s} \frac{|\nabla u|^2}{|x|^{n-2}} \, d\sigma \in L^1\left(0,R\right) \quad \text{for every } 0 < R < 1$$

and

$$\int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx = \int_{B_{r_0}} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx + \int_{r_0}^r \int_{\partial B_s} \frac{|\nabla u|^2}{|x|^{n-2}} \, d\sigma ds \quad \text{for every } 0 < r_0 < r < 1.$$

In particular,

$$\frac{d}{dr} \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, dx = \int_{\partial B_r} \frac{|\nabla u|^2}{|x|^{n-2}} \, d\sigma = \frac{1}{r^{n-2}} \int_{\partial B_r} |\nabla u|^2 \, d\sigma \quad \text{for almost every } 0 < r < 1.$$

Thus, the absolute continuity of β and (3.10) follow.

At last we show $\beta' \geq 0$ almost everywhere, which finishes the proof. Fix some 0 < r < 1 such that (3.9) and (3.10) hold. In the first term in the right-hand side of (3.10) we represent ∇ on ∂B_r by the tangential gradient $\nabla_{\theta} u \coloneqq \nabla u - (\partial_{\nu} u) \vec{\nu}$ for the outer normal vector $\vec{\nu}$ with respect to B_r , which gives:

(3.11)
$$\int_{\partial B_r} |\nabla u|^2 \, d\sigma = \int_{\partial B_r} |\nabla_\theta u|^2 \, d\sigma + \int_{\partial B_r} (\partial_\nu u)^2 \, d\sigma$$

For the second term in (3.10) we use (3.9) and the geometric inequality $2ab \le a^2 + b^2$ to get

$$(3.12) \qquad \frac{1}{r^2} \int_{B_r} \frac{|\nabla u|^2}{|x|^{n-2}} dx \leq \frac{1}{r^n} \int_{\partial B_r} u \partial_\nu u \, d\sigma + \frac{n-2}{2r^{n+1}} \int_{\partial B_r} u^2 \, d\sigma$$
$$\leq \frac{1}{4r^{n+1}} \int_{\partial B_r} u^2 \, d\sigma + \frac{1}{r^{n-1}} \int_{\partial B_r} (\partial_\nu u)^2 \, d\sigma + \frac{n-2}{2r^{n+1}} \int_{\partial B_r} u^2 \, d\sigma$$
$$= \frac{2n-3}{4r^{n+1}} \int_{\partial B_r} u^2 \, d\sigma + \frac{1}{r^{n-1}} \int_{\partial B_r} (\partial_\nu u)^2 \, d\sigma.$$

Thus, (3.11) and (3.12) in (3.10) give

$$\beta'(r) \ge \frac{1}{r^{n-1}} \int_{\partial B_r} |\nabla_\theta u|^2 \, d\sigma - \frac{2n-3}{4r^{n+1}} \int_{\partial B_r} u^2 \, d\sigma = \frac{1}{r^{n+1}} \left(\int_{\partial B_r} u^2 \, d\sigma \right) \cdot \left(\frac{r^2 \int_{\partial B_r} |\nabla_\theta u|^2 \, d\sigma}{\int_{\partial B_r} u^2 \, d\sigma} - \frac{2n-3}{4} \right)$$

After rescaling we have

$$\frac{r^2 \int_{\partial B_r} |\nabla_{\theta} u|^2 \, d\sigma}{\int_{\partial B_r} u^2 \, d\sigma} \ge \inf \left\{ \frac{\int_{\partial B_1} |\nabla_{\theta} w|^2 \, d\sigma}{\int_{\partial B_1} w^2 \, d\sigma} : w \in H^1(\mathbb{S}^{n-1}) \right\}.$$

Thus, the right-hand side becomes the Rayleigh quotient for the Laplace–Beltrami operator Δ_{θ} on the (n-1)-dimensional sphere \mathbb{S}^{n-1} , that is, the right-hand side coincides with the first eigenvalue of Δ_{θ} . From [Shi92, Chapter II, Theorem 4.1] we know that the first eigenvalue is n-1, which implies $\beta'(r) \geq 0$. \Box

3.2. **Proof of Theorem 3.1.** We proceed similar to the proof of [AC04, Theorem 4]. The main difference is to replace the Poincaré inequality by Maz'ya's inequality:

Lemma 3.6 (Maz'ya's inequality). There exists a constant $C_M > 0$, depending only on the dimension, such that for any $f \in H^1(B_r)$, r > 0, we have

(3.13)
$$\int_{B_r} f^2 dx \le \frac{C_M}{\operatorname{Cap}_0(\ker f \cap B_r; B_{2r})} \int_{B_r} |\nabla f|^2 dx.$$

Proof. The proof is analogous to [BB11, Theorem 5.53]. As the dependence on the radius r is important, we include the proof. For $f \in H^1(B_r)$ we may assume that f is nonnegative, else we consider |f|. Denote

$$\bar{f}\coloneqq \left(\oint_{B_r} f^2 \, dx \right)^{1/2}$$

and $\varphi := (1 - f/\bar{f})_+$. Clearly $\varphi \in H^1(B_r)$ with $\varphi \ge 1$ q.e. on ker f. Let $\Phi \in H^1_0(B_{2r})$ be an extension of φ such that there exists a constant $C_M > 0$, independent of r, satisfying

$$\left\|\nabla\Phi\right\|_{L^{2}(B_{2r})}^{2} \leq C_{M} \left\|\nabla\varphi\right\|_{L^{2}(B_{r})}^{2}$$

Such an extension exists, for instance: Choose any linear continuous extension operator $E' : H^1(B_1) \to H^1(B_2)$ and define $E \coloneqq T^{-1}E'T$, where $Tg(x) \coloneqq g(rx)$. We conclude

$$\operatorname{Cap}_{0}(\ker f \cap B_{r}; B_{2r}) \leq \int_{B_{2r}} |\nabla \Phi|^{2} \, dx \leq C_{M} \int_{B_{r}} |\nabla \varphi|^{2} \, dx \leq C_{M} \left(\bar{f}\right)^{-2} \int_{B_{r}} |\nabla f|^{2} \, dx.$$

Lemma 3.7. Let $F \subset B_1 \cap \{x_n = 0\}$ be relatively closed in B_1 and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. Then, there exists a constant C > 0 such that for any $r \in (0,1)$ we have

(3.14)
$$\sup_{B_{r/2}} |u| \le C \frac{r^{(n-1)/2}}{\operatorname{Cap}_0(\Lambda(u) \cap B_r; B_{2r})^{1/2}}$$

Proof. First, it is standard to show that u_{\pm} , the positive and negative part of u, are subharmonic in B_1 . Therefore, we apply the mean value property for subharmonic functions together with Jensen's inequality and Maz'ya's inequality (3.13) to obtain

$$\sup_{B_{r/2}} u_{\pm} \le \left(\int_{B_r} u^2 \, dx \right)^{1/2} \le \frac{C_M^{1/2}}{\operatorname{Cap}_0(\ker u \cap B_r; B_{2r})^{1/2}} \left(\int_{B_r} |\nabla u|^2 \, dx \right)^{1/2}.$$

Using the monotonicity of Cap₀ and the inclusion $\Lambda(u) \subset \ker u$ together with Proposition 3.4 and

$$\int_{B_r} |\nabla u|^2 \, dx \le r^{n-1} \beta(r),$$

$$\sum_{i=1}^{n-1} \beta(1/2)^{1/2}.$$

we conclude (3.14) with constant $C = C_M^{1/2} \beta (1/2)^{1/2}$

Proof of Theorem 3.1. Without loss of generality, we may assume that $x_0 = 0 \in \Lambda(u)$. Then, (3.14) and the capacity density condition (3.1) of $\Lambda(u)$ imply the existence of a constant C > 0 such that for every $x \in B_r$, $r \in (0, r_0)$:

$$|u(0) - u(x)| = |u(x)| \le C|x - 0|^{1/2}.$$

Thus, u is $C^{1/2}$ -regular at x_0 .

Remark 3.8. From the proof of Theorem 3.1, in particular due to (3.14), it is immediate that if one replaces (3.1) in Theorem 3.1 with the condition: there exists $c_0, r_0 > 0$ and $\alpha \in [0, 1)$ such that

$$\operatorname{Cap}_0(\Lambda(u) \cap B_r(x_0); B_{2r}(x_0)) \ge r^{n-2+\alpha} c_0 \quad \text{for every } r \in (0, r_0),$$

then u is $C^{(1-\alpha)/2}$ -regular at x_0 .

3.3. Generalization for non-zero obstacles. One can generalize Theorem 3.1 to include non-zero obstacles as follows:

Theorem 3.9. Let $n \ge 2$ and $F \subset \{x \in B_1 \mid x_n = 0\}$ be relatively closed in B_1 . Assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,\psi,F}(B_1)$ -obstacle problem for an obstacle $\psi : F \to \mathbb{R}$. Moreover, assume there exists an extension $\Psi \in H^1(B_1)$ of ψ satisfying:

- (i) For $x_0 \in \Lambda(u)$ there exists a constant C > 0 and $\alpha \in (0,1)$ such that $\sup_{B_r} |\Psi \Psi(x_0)| \leq Cr^{\alpha}$.
- (ii) $\Delta \Psi$ exists and satisfies

$$\int_{B_1(x_0)} \frac{|\Delta \Psi(x)|}{|x - x_0|^{n-1}} \, dx < +\infty.$$

If $\Lambda(u)$ satisfies a capacity density condition at $x_0 \in \Lambda(u)$, that is, there exists a constant $c_0 > 0$ and a radius $r_0 > 0$ such that

$$\operatorname{Cap}_0(\Lambda(u) \cap B_r(x_0); B_{2r}(x_0)) \ge r^{n-2}c_0 \quad \text{for every } r \in (0, r_0),$$

then u is C^{β} -regular at x_0 for $\beta = \min\{1/2, \alpha\}$.

Sketch of the proof. Without loss of generality, it is enough to consider $x_0 = 0$ and $\psi(0) = 0$. If u solves the homogeneous obstacle problem, then $v \coloneqq u - \Psi$ satisfies the Euler-Lagrange equations

(3.15)
$$\begin{cases} -\Delta v \ge f & \text{in } B_1 \\ -\Delta v = 0 & \text{in } \Omega \setminus \Lambda(v) \\ v \ge 0 & \text{on } F, \end{cases}$$

where $f := \Delta \Psi$ and $\Lambda(v) := \{x \in F \mid v(x) = 0\} = \Lambda(u)$. Now, one proves a variant of the monotonicity formula in Proposition 3.4 similar to [CSS08]. More precisely, one shows the monotonicity of

$$\beta^{f}(r) \coloneqq \beta(r, v) + \int_{0}^{r} \frac{1}{s^{2}} \int_{B_{s}} \frac{fv}{|x|^{n-2}} \, dx \, ds \quad \text{for } r \in (0, 1).$$

Note, for the second term we have

$$\int_0^r \frac{1}{s^2} \int_{B_s} \frac{fv}{|x|^{n-2}} \, dx \, ds \leq \sup_{B_r} |v| \cdot \int_{B_1} \frac{|\Delta \Psi(x)|}{|x|^{n-1}} \, dx < +\infty$$

due to the condition (*ii*). The monotonicity of β^{f} follows similarly as in Proposition 3.4 with the help of a variant of Lemma 3.5:

$$\int_{B_r} \nabla v \cdot \nabla \varphi \, dx = \int_{B_r} f \varphi \, dx + \int_{\partial B_r} \varphi \partial_\nu v \, d\sigma \quad \text{for every } \varphi \in H^1(B_1) \text{ such that } \varphi = 0 \text{ q.e. on } \Lambda(u).$$

To finish the proof, we use the subharmonicity of u_{\pm} as in Lemma 3.7 but incorporate Ψ as follows:

$$\sup_{B_r} u_{\pm} \le \int_{B_r} u_{\pm} \, dx \le \int_{B_r} |v| \, dx + \int_{B_r} |\Psi| \, dx \le \left(\int_{B_r} |v|^2 \, dx \right)^{\frac{1}{2}} + \int_{B_r} |\Psi| \, dx$$

After using Maz'ya's inequality (3.13) on the integral with respect to v, we use the estimate

$$\int_{B_r} |\nabla v|^2 \, dx \le r^{n-1} \beta(r, v) = r^{n-1} \left(\beta^f(r) - \int_0^r \frac{1}{s^2} \int_{B_s} \frac{fv}{|x|^{n-2}} \, dx \, ds \right)$$
$$\le r^{n-1} \left(\beta^f(1/2) + \sup_{B_r} |v| \cdot \int_{B_1} \frac{|\Delta \Psi(x)|}{|x|^{n-1}} \, dx \right),$$

for any $r \in (0, 1/2)$. Hence, we conclude

$$\sup_{B_r} u_{\pm} \le C(r^{1/2} + r^{\alpha}),$$

where the constant depends on c_0 , C_M , $\beta(1/2, u)$, $\sup_{B_{3/4}} |u|$, (i), and (ii).

Remark 3.10. Let us comment on the conditions (i) and (ii) in Theorem 3.9. It is easy to see that both are satisfied if $\Psi \in C^{1,1}(B_1)$. In fact, $\Psi \in W^{2,n+\delta}(B_1)$, for any $\delta > 0$, is sufficient. Note, by Sobolev-Morrey embedding, this implies that $\Psi \in C^{1,\alpha}(B_1)$ for $\alpha = 1 - \frac{n}{n+\delta}$. In particular, in this case Ψ is Lipschitz continuous which gives the optimal $C^{1/2}$ -regularity in Theorem 3.9 as (i) is satisfied for $\alpha = 1$.

Moreover, we expect that (*ii*) implies (*i*) for $\alpha = 1$, thus (*ii*) implies the optimal $C^{1/2}$ -regularity already. The reason is that the integral in (*ii*) coincides with the first order Riesz potential $I_1(|\Delta \Psi| \mathbb{1}_{B_1(x_0)})$ at x_0 , see [AH96]. A simple calculation shows if $w = \Phi * f$, for the fundamental solution Φ of the Laplacian on \mathbb{R}^n , then $|\nabla w| \leq I_1(f)$. Thus, roughly speaking, boundedness of $I_1(f)$ implies Lipschitz continuity of w. However, it is not clear how to utilize this observation in a rigorous manner, especially since the statements in (*i*) and (*ii*) are only pointwise in x_0 .

4. Almost optimal Hölder regularity in the half-hyperplane obstacle case

In this section we are concerned with Signorini-type problems in the unit ball $B_1 \subset \mathbb{R}^n$ and the obstacle region $F = \{(x', x_{n-1}, 0) \mid x' \in \mathbb{R}^{n-2}, x_{n-1} \leq 0\} \cap B_1$ being the left half of the horizontal hyperplane in B_1 . We recall that, in order to investigate regularity of solutions to (classical) Signorini problems, it is useful to take the associated Dirichlet boundary value problems into consideration; see [PSU12,FR16,Sal12,FR22] for instance. Nevertheless, since the obstacle condition is achieved only on a subset F of $\{x_n = 0\}$, it turns out that the behavior of solutions to Signorini-type problems can be described by the associated *mixed boundary value problems* (or mixed BVP for short). Therefore, before we establish the suitable Hölder regularity for solutions of Signorini-type problems, we develop the related mixed BVP theory.

4.1. Well posedness of Mixed BVPs. Let Ω be a Lipschitz domain in \mathbb{R}^n and $\partial\Omega = \Gamma_D \cup \Gamma_N$, where Γ_D and Γ_N denote the Dirichlet boundary part and the Neumann boundary part of $\partial\Omega$, respectively. As Ω is a Lipschitz domain, it is an H^1 -extension domain. Thus, for any function $u \in H^1(\Omega)$ there exists a unique quasi-continuous extension $\tilde{u} : \overline{\Omega} \to \mathbb{R}$ where its boundary values are unique in a quasi-everywhere sense with respect to $\operatorname{Cap}_1(\cdot; \mathbb{R}^n)$. In particular, we say $\Gamma \subset \partial\Omega$ is of *positive capacity* if $\operatorname{Cap}_1(\Gamma; \mathbb{R}^n) > 0$. Hence, statements like " $u = \psi$ q.e. on Γ_D " are well stated in the aforementioned sense.

Then, we are interested in a mixed BVP given by

(4.1)
$$\begin{cases} -\Delta v = f & \text{in } \Omega \\ v = \psi & \text{on } \Gamma_D \\ \partial_{\nu} v = 0 & \text{on } \Gamma_N, \end{cases}$$

where $f \in L^2(\Omega)$, $\psi \in H^1(\Omega)$, and ν is the outward unit normal vector on Γ_N .

Definition 4.1. Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain with $\partial \Omega = \Gamma_D \cup \Gamma_N$, $f \in L^2(\Omega)$, and $\psi \in H^1(\Omega)$. A function $v \in H^1(\Omega)$ is said to be a *weak solution* to the mixed BVP (4.1), if for every $\varphi \in H^1_{\Gamma_D}(\Omega) \coloneqq \{\varphi \in H^1(\Omega) \mid \varphi = 0 \text{ q.e. on } \Gamma_D\}$, we have

$$\int_{\Omega} \nabla v \cdot \nabla \varphi \, dx = (f, \varphi)_{L^2(\Omega)}$$

and $v = \psi$ q.e. on Γ_D .

The well posedness of (4.1) follows from the standard variational argument; see also [BL24, Theorem 8] for a similar result.

Theorem 4.2. Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain with $\partial \Omega = \Gamma_D \cup \Gamma_N$. Moreover, suppose that Γ_D is closed and of positive capacity, $f \in L^2(\Omega)$, and $\psi \in H^1(\Omega)$. Then, there exists a unique weak solution $v \in H^1(\Omega)$ of (4.1). Moreover, there exists a universal constant C > 0 such that

$$\|v\|_{H^1(\Omega)} \le C\left(\|\psi\|_{H^1(\Omega)} + \|f\|_{L^2(\Omega)}\right).$$

Proof. We set the linear functional

$$-\Delta\psi(\varphi) \coloneqq \int_{\Omega} \nabla\psi \cdot \nabla\varphi \, dx \quad \text{for } \varphi \in H^1(\Omega).$$

It follows from the Lax–Milgram theorem, applied to the usual Dirichlet energy on $H^1_{\Gamma_D}(\Omega)$, that there exists a unique $w \in H^1_{\Gamma_D}(\Omega)$ satisfying

$$\int_{\Omega} \nabla w \cdot \nabla \varphi \, dx = \Delta \psi(\varphi) + (f, \varphi)_{L^2(\Omega)} \quad \text{for every } \varphi \in H^1_{\Gamma_D}(\Omega)$$

with the uniform bound

$$||w||_{H^1(\Omega)} \le C \left(||\psi||_{H^1(\Omega)} + ||f||_{L^2(\Omega)} \right).$$

Now our solution can be found by $v \coloneqq w + \psi \in H^1(\Omega)$ with the desired uniform estimate.

We also provide the maximum principle and the comparison principle for weak solutions to (4.1).

Theorem 4.3 (Maximum principle for subsolutions). Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain with $\partial \Omega = \Gamma_D \cup \Gamma_N$. Moreover, suppose that Γ_D is closed and of positive capacity. Let $v \in H^1(\Omega)$ be a weak solution to

$$\begin{cases} -\Delta v \leq 0 & \text{in } \Omega \\ v \leq 0 & \text{in } \Gamma_D \\ \partial_{\nu} v \leq 0 & \text{in } \Gamma_N, \end{cases}$$

that means, for every nonnegative $\varphi \in H^1_{\Gamma_D}(\Omega)$ we have

$$\int_{\Omega} \nabla v \cdot \nabla \varphi \, dx \le 0$$

and $v \leq 0$ q.e. on Γ_D . Then, $v \leq 0$ a.e. in Ω .

Proof. Let us write $v = v_+ - v_-$, where $v_+ = \max\{v, 0\}$ and $v_- = -\min\{v, 0\}$. Since $v_+ \in H^1(\Omega)$, $v_+ = 0$ q.e. on Γ_D , and $v_+ \ge 0$, we can take v_+ as a test function to obtain

$$\int_{\Omega} \nabla v_{+} \cdot \nabla v_{+} \, dx - \int_{\Omega} \nabla v_{-} \cdot \nabla v_{+} \, dx = \int_{\Omega} \nabla v \cdot \nabla v_{+} \, dx \le 0$$

As $v_{-} = 0$ on the support of u_{+} , the strong locality of the Dirichlet energy implies that $(\nabla v_{-}, \nabla v_{+})_{L^{2}(\Omega)} = 0$. Therefore, $\|\nabla v_{+}\|_{L^{2}(\Omega)} \leq 0$, which implies that $v_{+} = \text{const} = 0$.

Corollary 4.4 (Comparison principle). Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain with $\partial \Omega = \Gamma_D \cup \Gamma_N$. Moreover, suppose that Γ_D is closed and of positive capacity. Let $v_1, v_2 \in H^1(\Omega)$ satisfy

$$\begin{cases} -\Delta v_1 \leq -\Delta v_2 & \text{in } \Omega \\ v_1 \leq v_2 & \text{in } \Gamma_D \\ \partial_{\nu} v_1 \leq \partial_{\nu} v_2 & \text{in } \Gamma_N \end{cases}$$

which means, for every nonnegative $\varphi \in H^1_{\Gamma_D}(\Omega)$ we have

$$\int_{\Omega} \nabla v_1 \cdot \nabla \varphi \, dx \le \int_{\Omega} \nabla v_2 \cdot \nabla \varphi \, dx$$

and $v_1 \leq v_2$ q.e. on Γ_D . Then, $v_1 \leq v_2$ a.e. in Ω .

4.2. Almost optimal Hölder regularity for mixed BVPs. We now restrict our attention to the special case that is related to our situation of Signorini-type problems. That means, we fix $\Omega = B_1^+ \subset \mathbb{R}^n$ with $n \geq 2$, $\Gamma_N = \{(x', x_{n-1}, 0) : x_{n-1} > 0\} \cap B_1$, and $\Gamma_D = \partial B_1^+ \setminus \Gamma_N = F \cup (\partial B_1)^+$ for $F \coloneqq \{(x', x_{n-1}, 0) : x_{n-1} \leq 0\} \cap B_1$. Here we denote $x = (x', x_{n-1}, x_n) \in \mathbb{R}^n$ and use the cylindrical coordinates $x' \in \mathbb{R}^{n-2}$, $r = \sqrt{x_{n-1}^2 + x_n^2}$, and

$$\theta = \theta(x_{n-1}, x_n) = \begin{cases} \arccos(x_{n-1}/r) & \text{if } x_n \ge 0\\ 2\pi - \arccos(x_{n-1}/r) & \text{if } x_n < 0. \end{cases}$$

We also assume that the Dirichlet data ψ is zero on the horizontal hyperplane $\{x_n = 0\}$. Then, we are concerned with a weak solution $v \in H^1(B_1^+) \cap C(\overline{B_1^+})$ to the mixed BVP

(4.2)
$$\begin{cases} -\Delta v = f \quad \text{in } B_1^+ \\ v = 0 \quad \text{on } F \\ \partial_{\nu} v = 0 \quad \text{on } \Gamma_N. \end{cases}$$

Here $\vec{\nu} = (0, \dots, 0, -1)$ on Γ_N and $f \in H^1(B_1^+) \cap L^{\infty}(B_1^+)$.

Let us make a couple of comments on solutions to (4.2). First of all, since we are interested in the regularity of v in $\overline{B_{1/2}^+}$, we do not specify a boundary behavior of v on $(\partial B_1)^+$. Next, we can show that v is continuous in $\overline{B_{1/2}^+}$ as follows: By standard regularity theory v is continuous in the non-intersection region $\overline{B_{1/2}^+} \setminus (\Gamma_D \cap \overline{\Gamma_N})$, see [GT01, Chapter 6] for instance. Moreover, v is continuous at $x_0 \in I_{1/2} := \Gamma_D \cap \overline{\Gamma_N} \cap \overline{B_{1/2}} = \{(z', 0, 0) : |z'| \le 1/2\}$ by using the barrier constructed in [Lie86,Lie89], since the straight Γ_N -wedge condition (defined in [Lie86]) is satisfied at $x_0 \in I_{1/2}$. We note that this condition does not hold when we reach the boundary ∂B_1 , for example at $x_0 = (\pm 1, \cdots, 0, 0) \in \partial B_1 \cap \Gamma_D \cap \overline{\Gamma_N}$ when $n \ge 3$. Finally, we would like to point out that the assumption $v \in C(\overline{B_1^+})$ is not restrictive. In fact, a solution v to (4.2) is continuous and so bounded in $\overline{B_r^+}$ for any $r \in (0, 1)$, by repeating the previous argument. Therefore, we may consider a slightly smaller ball than B_1 , if necessary.

Our goal in this subsection is to derive the almost optimal Hölder regularity of solutions in this special case, that is, we will show that $v \in C^{1/2-\varepsilon}(\overline{B^+_{1/4}})$ for any $\varepsilon \in (0, 1/2)$. Let us first prove that the pointwise regularity of v at the intersection region $I_{1/2}$ is $C^{1/2-\varepsilon}$ for any $\varepsilon \in (0, 1/2)$.

Lemma 4.5 (Intersection points). Let $\varepsilon \in (0, 1/2)$. Let $v \in H^1(B_1^+) \cap C(\overline{B_1^+})$ be a weak solution to (4.2), where $f \in H^1(B_1^+) \cap L^{\infty}(B_1^+)$. Then, v is $C^{1/2-\varepsilon}$ at $z \in I_{1/2}$; that is, there exists a constant C > 0 depending only on n and ε such that

$$|v(x)| \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x - z|^{1/2 - \varepsilon}$$

for any $x = (x', x_{n-1}, x_n) \in B^+_{1/2}(z)$.

Proof. We may assume that z = 0 by applying the standard translation and scaling argument, if necessary. We define an auxiliary function h given by

$$h(x', x_{n-1}, x_n) \coloneqq r^{1/2-\varepsilon} \cos\left[\left(\frac{1-\varepsilon}{2}\right)\theta\right] \quad \text{for } x_n \ge 0.$$

Then, a direct calculation shows that

$$\begin{pmatrix} -\Delta h = \left(\frac{\varepsilon}{2} - \frac{3\varepsilon^2}{4}\right) r^{-3/2-\varepsilon} \cos\left[\left(\frac{1-\varepsilon}{2}\right)\theta\right] & \text{in } B_1^+ \\ h = r^{1/2-\varepsilon} \sin(\varepsilon \pi/2) \ge 0 & \text{on } F \\ \partial_{\nu} h = \partial_{\theta} h = 0 & \text{on } \Gamma_N, \end{cases}$$

and $r^{1/2-\varepsilon}\sin(\varepsilon\pi/2) \le h(x) \le r^{1/2-\varepsilon}$ for $\theta \in [0,\pi]$. We note that $\varepsilon/2 - 3\varepsilon^2/4 > 0$ for any $\varepsilon \in (0,1/2)$. We now construct a barrier function (supersolution)

$$w(x) \coloneqq \left(\|v\|_{L^{\infty}(B_{1}^{+})} + \|f\|_{L^{\infty}(B_{1}^{+})} \right) \left(C_{1}h(x) + C_{2}|x|^{2} \right)$$

for some $C_1, C_2 > 0$. Then, there exist constants $C_1, C_2 > 0$ depending only on n and ε such that

$$\begin{cases} -\Delta w \ge \|f\|_{L^{\infty}(B_{1}^{+})} & \text{in } B_{1}^{+} \\ w \ge 0 & \text{on } F \\ w \ge \|v\|_{L^{\infty}(B_{1}^{+})} & \text{on } (\partial B_{1})^{+} \\ \partial_{\nu}w = 0 & \text{on } \Gamma_{N}. \end{cases}$$

Therefore, an application of the comparison principle (Corollary 4.4) between v and w yields that $v \leq w$ in B_1^+ . In particular, we obtain

$$v(x) \le w(x) \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x|^{1/2-\varepsilon},$$

where C > 0 depends only on n and ε . By considering -v, -f instead of v, f, we arrive at the desired estimate.

We next extend this pointwise regularity to any point in $B'_{1/4} = \{(x', x_{n-1}, 0) \in B_{1/4}\}.$

Lemma 4.6 (Pointwise boundary regularity). Let $\varepsilon \in (0, 1/2)$. Let $v \in H^1(B_1^+) \cap C(\overline{B_1^+})$ be a weak solution to (4.2), where $f \in H^1(B_1^+) \cap L^{\infty}(B_1^+)$. Then, v is $C^{1/2-\varepsilon}$ at $z \in B'_{1/4}$; that is, there exists a constant C > 0 depending only on n and ε such that

$$|v(x) - v(z)| \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x - z|^{1/2 - \varepsilon}$$

for any $x = (x', x_{n-1}, x_n) \in B^+_{1/4}(z)$.

Proof. If $z_{n-1} = 0$, then the desired estimate follows from Lemma 4.5. We assume that $z_{n-1} > 0$, that is, $z \in \Gamma_N$ with $r := z_{n-1} \in (0, 1/4)$. By applying Lemma 4.5 for (z', 0, 0), we observe that

(4.3)
$$|v(x)| \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x - (z', 0, 0)|^{1/2 - \varepsilon} \quad \text{for } x \in B_{1/2}^+(z', 0, 0).$$

We now choose $x \in B_{1/4}^+(z)$ and split two cases as follows.

(i) $(|x-z| \le r/2)$ In $B^+_{3r/4}(z)$, v solves the Poisson equation with zero Neumann boundary condition. We thus apply the boundary estimate developed in [GT01, Section 6.7] to obtain

$$|v(x) - v(z)| \le C\left(\frac{\|v\|_{L^{\infty}(B_{3r/4}^{+}(z))}}{r^{1/2-\varepsilon}} + r^{2-(1/2-\varepsilon)}\|f\|_{L^{\infty}(B_{1}^{+})}\right)|x - z|^{1/2-\varepsilon}.$$

On the other hand, (4.3) together with the triangle inequality gives that

$$\|v\|_{L^{\infty}(B^{+}_{3r/4}(z))} \leq C\left(\|v\|_{L^{\infty}(B^{+}_{1})} + \|f\|_{L^{\infty}(B^{+}_{1})}\right) r^{1/2-\varepsilon},$$

which implies that

$$|v(x) - v(z)| \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x - z|^{1/2 - \varepsilon}.$$

(*ii*) (|x - z| > r/2) In this case, (4.3) again shows that

$$\begin{aligned} |v(x) - v(z)| &\leq |v(x)| + |v(z)| \leq C \left(\|v\|_{L^{\infty}(B_{1}^{+})} + \|f\|_{L^{\infty}(B_{1}^{+})} \right) \left(|x - z|^{1/2 - \varepsilon} + r^{1/2 - \varepsilon} \right) \\ &\leq C \left(\|v\|_{L^{\infty}(B_{1}^{+})} + \|f\|_{L^{\infty}(B_{1}^{+})} \right) |x - z|^{1/2 - \varepsilon}. \end{aligned}$$

The other case $z_{n-1} < 0$ follows from the same argument; just replace the Neumann problem by the Dirichlet problem.

We finally combine the interior regularity and the pointwise boundary regularity by following the argument in [Wan92, Theorem 3.1].

Theorem 4.7 (Global regularity). Let $\varepsilon \in (0, 1/2)$. Let $v \in H^1(B_1^+) \cap C(\overline{B_1^+})$ be a weak solution to (4.2), where $f \in H^1(B_1^+) \cap L^{\infty}(B_1^+)$. Then, $v \in C^{1/2-\varepsilon}(\overline{B_{1/4}^+})$; that is, there exists a constant C > 0 depending only on n and ε such that

$$|v(x) - v(y)| \le C \left(\|v\|_{L^{\infty}(B_1^+)} + \|f\|_{L^{\infty}(B_1^+)} \right) |x - y|^{1/2 - \varepsilon}$$

for any $x, y \in B_{1/4}^+$.

Proof. Since the proof is essentially the same as in Lemma 4.6, we only overview the proof. If $x_n = 0$ or $y_n = 0$, then it is a consequence of Lemma 4.6. Otherwise, we let $\hat{x} := (x', x_{n-1}, 0) \in B'_{1/4}$ with $r = x_n \in (0, 1/4)$. We utilize the estimate obtained in Lemma 4.6 for \hat{x} and split two cases depending on the distance between x and y. In particular, if |x - z| < r/2, we recall the interior regularity of solutions to the Poisson equation; see [CC95, Proposition 4.10] for instance.

4.3. Almost optimal regularity for Signorini-type problems. We now consider the special situation provided in Section 4.2 (with $f \equiv 0$) and claim that a weak solution u of the associated Signorini-type problem belongs to $C^{1/2-\varepsilon}(\overline{B_{1/4}^+})$ for any $\varepsilon > 0$. In the following theorem, we may assume $u \in L^{\infty}(B_1)$ by repeating a similar argument illustrated at the beginning of Section 4.2.

Theorem 4.8 (Almost optimal regularity). Let $\Omega = B_1 \subset \mathbb{R}^n$ with $n \geq 2$ and $F = \{(x', x_{n-1}, 0) : x_{n-1} \leq 0\} \cap B_1$. If u is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem, then $u \in C^{1/2-\varepsilon}(\overline{B_{1/4}})$ for any $\varepsilon > 0$. Moreover, there exists a constant C > 0 depending only on n and ε such that

$$\|u\|_{C^{1/2-\varepsilon}(\overline{B_{1/4}})} \le C \|u\|_{L^{\infty}(B_1)}.$$

Proof. We first assume that u is symmetric with respect to the hyperplane $\{x_n = 0\}$. We will construct an extended obstacle $\overline{\Psi}$ defined in $B_{3/4}$ and understand u as a solution to the classical obstacle problem with the obstacle $\overline{\Psi}$. For this purpose, we let $\eta \in C_c^{\infty}(B_1)$ be a cutoff function such that $\eta = 1$ in $B_{3/4}$ and $0 \le \eta \le 1$. Then, Theorem 4.2 guarantees the existence of a unique weak solution $\Psi \in H^1(B_1^+)$ of the following mixed BVP:

$$\begin{cases} -\Delta \Psi = 0 & \text{in } B_1^+ \\ \Psi = -\|u\|_{L^{\infty}(B_1)} \cdot (1-\eta) & \text{on } \Gamma_D = F \cup (\partial B_1)^+ \\ \partial_{\nu} \Psi = 0 & \text{on } \Gamma_N. \end{cases}$$

We observe that $\Psi = 0$ on $F \cap B_{3/4}$ and by the comparison principle Corollary 4.4,

$$-\|u\|_{L^{\infty}(B_1)} \le \Psi \le u \text{ in } B_1^+$$

In particular, u can be understood as a solution to the classical $\mathcal{K}_{u,\overline{\Psi},B_{3/4}}(B_{3/4})$ -obstacle problem, where

$$\overline{\Psi}(x', x_n) \coloneqq \begin{cases} \Psi(x', x_n) & \text{if } x_n \ge 0\\ \Psi(x', -x_n) & \text{if } x_n < 0. \end{cases}$$

Since the almost optimal estimate Theorem 4.7 for Ψ in $B^+_{3/4}$ yields that

$$\|\Psi\|_{C^{1/2-\varepsilon}(\overline{B_{1/4}^+})} \le C \|u\|_{L^{\infty}(B_1)}$$

the desired estimate follows from [Caf98, Theorem 2].

In the case that u is not symmetric with respect to $\{x_n = 0\}$, we let $w \in H^1(B_1)$ be the weak solution to the Dirichlet boundary value problem

$$\begin{cases} -\Delta w = 0 & \text{in } B_1 \\ w = g & \text{on } \partial B_1 \end{cases}$$

where $g(x, y) = (u(x', x_n) - u(x', -x_n))/2$. Then, it is easy to check that $\bar{u} \coloneqq u - w$ becomes a symmetric solution to the $\mathcal{K}_{\bar{u},0,F}(B_1)$ -obstacle problem, which finishes the proof.

Remark 4.9. In the articles [ACM22, AC22, AC23] the authors proved a Boyarski–Meyer estimate for solutions to mixed boundary value problems of the form

(4.4)
$$\begin{cases} -\mathcal{L}u = \operatorname{div} H & \operatorname{in} \Omega \\ u = 0 & \operatorname{on} \Gamma_D \\ \partial_{\nu} u = 0 & \operatorname{on} \Gamma_N, \end{cases}$$

where \mathcal{L} is an elliptic operator in divergence form, the *p*-Laplacian or $p(\cdot)$ -Laplacian, $\partial\Omega = \Gamma_D \cup \Gamma_N$, and the Dirichlet part of the boundary Γ_D satisfies a variant of the capacity density condition (3.1). The authors proved increased integrability of the gradient of the solution if H has higher integrability, more precisely, if $H \in (L^{p'(\cdot)(1+\delta_0)}(\Omega))^n$, $\delta_0 > 0$, then $\nabla u \in (L^{p(\cdot)(1+\delta)}(\Omega))^n$ for some $\delta_0 > \delta > 0$. In particular, from standard embedding results, the solution to (4.4) in the *p*-Laplacian case is Hölder continuous if p = n. Thus, this result goes in a similar direction as our results due to the connection of Signorini-type problems and mixed BVPs. Although such result does not give the (almost) optimal regularity, it allows for a more general treatment of operators and obstacle/Dirichlet sets F (including fractals) and may be compared to Theorem 2.12.

4.4. Generalization to non-zero obstacles. We also extend the results of Section 4 to include non-zero obstacles ψ as follows:

Theorem 4.10. Let $\Omega = B_1 \subset \mathbb{R}^n$, $n \geq 2$, $F = \{(x', x_{n-1}, 0) \mid x_{n-1} \leq 0\} \cap B_1$, and $\psi \in C^{\beta_0}(F)$ for some $\beta_0 \in (0, 1)$. Moreover, assume there exists an extension $\Psi_0 \in H^1(B_1)$ of ψ . If $u \in H^1(\Omega)$ solves the $\mathcal{K}_{u,\psi,F}(\Omega)$ -obstacle problem, then u is C^{β} -regular at $x_0 = 0$ for some $\beta \in (0, 1/2)$ such that $\beta \leq \beta_0$.

Sketch of the proof. This lemma follows from repeating the arguments in Section 4 with several modifications. We summarize the main strategy as follows. Here we may assume that u is bounded in $L^{\infty}(B_1)$ and symmetric with respect to the hyperplane $\{x_n = 0\}$ as usual.

- (i) We extend the obstacle ψ to Ψ on the domain $B_{3/4}$ by solving the associated mixed BVP. Then, u can be understood as the solution to the $\mathcal{K}_{u,\Psi,B_{3/4}}(B_{3/4})$ -obstacle problem, that is, a classical obstacle problem in $B_{3/4}$.
- (ii) We show the almost optimal Hölder regularity of Ψ by constructing appropriate barrier functions.
- (*iii*) We transport this regularity to u by means of [Caf98, Theorem 2].

Since the case of $n \ge 3$ can be proved in a similar way to the case of n = 2 and the steps (i), (iii) do not require essential changes, we focus on the step (ii) when n = 2. In particular, we present the main difference arising from the proof of Lemma 4.5, where the choice of Hölder exponent $\beta \in (0, 1/2) \cap (0, \beta_0]$ can be explained.

To be precise, we let $v \in H^1(B_1^+) \cap C(\overline{B_1^+})$ be a weak solution to the mixed BVP

$$\begin{cases} -\Delta v = f & \text{in } B_1^+ \\ v = \psi & \text{on } F \\ \partial_\nu v = 0 & \text{on } \Gamma_N. \end{cases}$$

Without loss of generality, we may assume that $\psi(0) = 0$.

We choose $\varepsilon \coloneqq 1/2 - \beta > 0$ in the proof of Lemma 4.5 and recall an auxiliary function h given by

$$h(x_1, x_2) \coloneqq r^{\beta} \cos\left[\left(\frac{1/2+\beta}{2}\right)\theta\right] \quad \text{for } x_2 \ge 0 \text{ and } r = \sqrt{x_1^2 + x_2^2}$$

so that $h(x) = C_{\beta} r^{\beta}$ on F for some constant $C_{\beta} > 0$. Moreover, we construct a modified barrier function

$$w(x) \coloneqq \left(\|v\|_{L^{\infty}(B_{1}^{+})} + \|f\|_{L^{\infty}(B_{1}^{+})} + \|\psi\|_{C^{\beta_{0}}(F)} \right) \left(C_{1}h(x) + C_{2}r^{2} \right)$$

for some $C_1, C_2 > 0$. Then, there exist constants $C_1, C_2 > 0$ depending only on β such that

$$\begin{cases} -\Delta w \ge -\|f\|_{L^{\infty}(B_{1}^{+})} & \text{in } B_{1}^{+} \\ w \ge v & \text{on } F \\ w \ge \|v\|_{L^{\infty}(B_{1}^{+})} & \text{on } (\partial B_{1})^{+} \\ \partial_{\nu}w = 0 & \text{on } \Gamma_{N}. \end{cases}$$

In fact, it follows from $\beta \leq \beta_0$ that

$$v(x) = \psi(x) \le \|\psi\|_{C^{\beta}(F)} r^{\beta_{0}} \le \|\psi\|_{C^{\beta}(F)} r^{\beta} \le w(x) \text{ on } F,$$

when we choose $C_1 > 0$ sufficiently large. Therefore, an application of the comparison principle between v and w yields that $v \leq w$ in B_1^+ . In other words, we obtain

$$v(x) \le w(x) \le C \left(\|v\|_{L^{\infty}(B_{1}^{+})} + \|f\|_{L^{\infty}(B_{1}^{+})} + \|\psi\|_{C^{\beta_{0}}(F)} \right) r^{\beta},$$

where C > 0 depends only on β . By considering -v, -f instead of v, f, we arrive at the desired estimate.

We deduce the desired conclusion after repeating the remaining part of Section 4.

5. Optimal Hölder regularity in the half-line obstacle case in two dimensions

In this section we focus on the two dimensional Signorini-type problem in the unit ball $B_1 \subset \mathbb{R}^2$ and obstacle region $F = \{(x_1, 0) \mid x_1 \leq 0\} \cap B_1$ being the left half of the horizontal line in B_1 . Compared to Section 4, we restrict our attention to n = 2 but develop the optimal $C^{1/2}$ -regularity in Theorem 5.11 and provide blowup profiles in Theorem 5.10.

Let $u \in H^1(B_1)$ be a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. From Theorem 2.7 we know that u is harmonic, therefore smooth, away from the contact set $\Lambda(u)$. In particular, this includes the right half $B'_1 \setminus F$ of the horizontal line in B_1 . For any $x_0 \in F$ that is neither (0,0) nor (-1,0), u is a local solution to the zero thin obstacle problem. More precisely, there exists $\delta > 0$ such that u solves the $\mathcal{K}_{u,0,B'_{\delta}(x_0)}(B_{\delta}(x_0))$ -obstacle problem, where $B'_{\delta}(x_0) = \{x \in B_{\delta}(x_0) \mid x_2 = 0\}$ as always. Then, by known results for the thin obstacle problem or Signorini problem (see for example [AC04, Sal12, PSU12]), u is Lipschitz in an open neighbourhood along x_0 and $C^{1,1/2}$ -regular from both sides up to x_0 .

Hence, it suffices to study the regularity at $x_0 = 0$. In Theorem 5.11 we prove the $C^{1/2}$ -regularity of u in $x_0 = 0$. The proof of Theorem 5.11 is based on the approach developed in [CSS08, ACS08], which is also available for obstacle problems with respect to the fractional Laplacian. The main strategy is to use Almgren's frequency function, [Alm00],

$$N(r,u) \coloneqq \frac{r \int_{B_r} |\nabla u|^2 \, dx}{\int_{\partial B_r} u^2 \, d\sigma}.$$

In order to show the monotonicity of $r \mapsto N(r, u)$ in Theorem 5.5, we first show a variant of Rellich's formula, [Rel40], in Proposition 5.3, which in the case of a smooth function φ reads

(5.1)
$$\int_{\partial B_r} |\nabla \varphi|^2 \, d\sigma = \frac{n-2}{r} \int_{B_r} |\nabla \varphi|^2 \, dx + 2 \int_{\partial B_r} (\partial_\nu \varphi)^2 \, d\sigma - \frac{2}{r} \int_{B_r} (x \cdot \nabla \varphi) \Delta \varphi \, dx.$$

This formula turned out to be the main difficulty in the proof of the optimal regularity. In fact, it is the main reason that we are restricted to the two dimensional setting here, see Remark 5.4. In the zero thin obstacle problem case, (5.1) can be proved as follows: First, one shows the $C^{1,\alpha}$ -regularity of u for some $\alpha > 0$. Then, one uses that Δu can be seen as a measure supported on the horizontal line in B_1 . The $C^{1,\alpha}$ -regularity of u can then be used to show that (5.1) holds for $\varphi = u$ as the derivative of u is Hölder continuous. In particular, the last term on the right-hand side of (5.1) vanishes as $\partial_{\nu} u \Delta u = 0$ in the sense of measures. See the proof of [ACS08, Lemma 1] or [Sal12, Theorem 2.5.1].

Nevertheless, in the setting of Theorem 5.11 we can only use the $C^{1,\alpha}$ -regularity away from 0. Thus, the idea is to apply Rellich's formula away from 0 in $B_r \setminus B_{\varepsilon}$ and then take the limit $\varepsilon \to 0$. The problematic term that arises is then of the form

(5.2)
$$\varepsilon \int_{\partial B_{\varepsilon}} |\nabla u|^2 \, dx.$$

To show that (5.2) converges to 0 for $\varepsilon \to 0$ we use the harmonicity away from B'_1 and the C^{α} -regularity from Theorem 2.12 of u to get the estimate $|\nabla u| \leq C\varepsilon^{\alpha-1}$ on ∂B_{ε} .

Then, with the help of Almgren's frequency function we can do a blowup analysis in Theorem 5.10 which proves the desired $C^{1/2}$ -regularity in Theorem 5.11. In particular, in Theorem 5.10 we show that any blowup u_0 of u at $x_0 = 0 \in \Lambda(u)$ can be written in polar coordinates as

$$u_0(r,\theta) = r^{\kappa} \cos(\kappa\theta)$$
 for some $\kappa \in 2\mathbb{N}$ or $\kappa \in \mathbb{N}_0 + \frac{1}{2}$

up to a multiplicative constant.

5.1. **Preliminary gradient estimates.** We start with two simple estimates to control the gradient of harmonic functions and solutions to thin obstacle problems. Both of them hold for any dimension.

Lemma 5.1. Let $n \ge 2$, $x_0 \in \mathbb{R}^n$, and r > 0. If u is harmonic in $B_r(x_0)$, then

$$|\nabla u(x_0)| \le \frac{2\sqrt{n}}{r} \sup_{\partial B_{r/2}(x_0)} |u|$$

Proof. Since u is harmonic in $B_r(x_0)$, $\partial_i u$ is also harmonic. Thus, by the mean value property and integration by parts we have

$$\partial_i u(x_0) = \oint_{B_{r/2}(x_0)} \partial_i u \, dx = \frac{2^n}{|B_1| r^n} \int_{\partial B_{r/2}(x_0)} u \frac{x_i}{|x|} \, d\sigma,$$

it. \Box

which implies the statement.

Lemma 5.2. Let $B_1 \subset \mathbb{R}^n$ with $n \geq 2$, $u \in H^1(B_1)$, and $x \in B'_1$. If there exists R > 0 such that u is a solution to the zero thin obstacle problem in $B_R(x)$, then

$$\sup_{B_{r/2}^{\pm}(x)} |\nabla u| \le \frac{C}{r} \sup_{B_{3r/4}(x)} |u| \quad \text{for any } r \in (0, R],$$

for some constant C > 0 depending only on n and R.

Proof. If $v \in H^1(B_1)$ is a solution to the zero thin obstacle problem, then $v \in C^{1,\alpha}(B_{3/4}^+)$, $\alpha > 0$, with bound

$$\|v\|_{C^{1,\alpha}(B^+_{3/4})} \le C \, \|v\|_{L^2(B_{3/4})} \le C \sup_{B_{3/4}} |v|.$$

See for example [Caf79, Theorem 1]. Since u is a local solution, by translation and scaling

$$v(y) \coloneqq u(ry + x) \in H^1(B_1)$$

is a solution to the zero thin obstacle problem. Thus, the statement follows from the relation

$$\nabla v(y)| = r|\nabla u(ry + x)|.$$

5.2. Rellich-type formula.

Proposition 5.3 (Rellich-type formula). Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$ and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. Then,

(5.3)
$$\int_{\partial B_r} |\nabla u|^2 \, d\sigma = 2 \int_{\partial B_r} (\partial_\nu u)^2 \, d\sigma \quad \text{for every } 0 < r < 1.$$

Proof. We start as in the proof of [ACS08, (6)] or [Sal12, (13)]. Let

$$V(x) \coloneqq x |\nabla u|^2 - 2(x \cdot \nabla u) \nabla u \quad \text{for } x \in B_1 \setminus \Lambda(u).$$

As long as we are away from $\Lambda(u)$, V is C^1 as u is harmonic and therefore smooth. In particular,

div
$$V = (n-2)|\nabla u|^2 = 0$$
 in $B_1 \setminus \Lambda(u)$.

As discussed before, for any $z \in F \setminus \{0\}$, u is in $C^{1,1/2}(B_{\delta}(z)^+ \cup B'_{\delta}(z))$ for some $\delta > 0$. In particular, V belongs to $C(B_{\delta}(z)^+ \cup B'_{\delta}(z))$. For $0 < \delta < \varepsilon$ let

$$F_{\delta} \coloneqq \{ x \in B_1 \mid \operatorname{dist}(x, F) < \delta \}.$$

Then, the divergence theorem implies

(5.4)
$$0 = \int_{B_r \setminus (B_\varepsilon \cup F_\delta)} \operatorname{div} V \, dx = \int_{\partial (B_r \setminus (B_\varepsilon \cup F_\delta))} V \cdot \vec{\nu} \, d\sigma$$

We separate the surface integral into the following disjoint sets:

$$\partial(B_r \setminus (B_{\varepsilon} \cup F_{\delta})) = \partial B_r \setminus F_{\delta} \cup (\partial F_{\delta} \cap B_r) \setminus B_{\varepsilon} \cup \partial B_{\varepsilon} \setminus F_{\delta} \eqqcolon A_1 \cup A_2 \cup A_3$$

(i) On A_1 we have

$$V \cdot \vec{\nu} = r |\nabla u|^2 - 2r (\partial_{\nu} u)^2$$

which implies

(5.5)
$$\lim_{\varepsilon \to 0} \lim_{\delta \to 0} \int_{A_1} V \cdot \vec{\nu} \, d\sigma = r \int_{\partial B_r} |\nabla u|^2 - 2(\partial_{\nu} u)^2 \, d\sigma.$$

By comparing (5.4) and (5.5) with (5.3), it remains to show that remaining integrals for A_2 and A_3 converge to 0.

(ii) On A_2 let us consider $A_2^+ = A_2 \cap B_1^+$ and $A_2^- = A_2 \cap B_1^-$. Since every $x \in A_2^+$ is of the form $x = (x_1, \delta)$ and $\vec{\nu}(x) = (0, -1)$, we have $V \cdot \vec{\nu} = -\delta |\nabla u|^2 + 2(x \cdot \nabla u)\partial_{x_2} u$. Since V is uniformly continuous up to $B'_r \setminus B_{\varepsilon}$, we have

$$\lim_{\delta \to 0} V(x_1, \delta) \cdot \vec{\nu} = 2\left(\begin{pmatrix} x_1 \\ 0 \end{pmatrix} \cdot \nabla u(x_1, 0) \right) \partial_{x_2} u(x_1, 0).$$

Similarly, for $x = (x_1, -\delta) \in A_2^-$ we have

$$\lim_{\delta \to 0} V(x_1, -\delta) \cdot \vec{\nu} = -2\left(\binom{x_1}{0} \cdot \nabla u(x_1, 0)\right) \partial_{x_2} u(x_1, 0).$$

Thus,

$$\lim_{\delta \to 0} \int_{A_2} V \cdot \vec{\nu} \, d\sigma = \lim_{\delta \to 0} \int_{A_2^+} V \cdot \vec{\nu} \, d\sigma + \int_{A_2^-} V \cdot \vec{\nu} \, d\sigma = 0.$$

(iii) On $A_3 = \partial B_{\varepsilon} \setminus F_{\delta}$ we again use the regularity of V on $\partial B_{\varepsilon} \cap \overline{F_{\delta}}$ to deduce

$$\lim_{\delta \to 0} \int_{A_3} V \cdot \vec{\nu} \, d\sigma = \int_{\partial B_{\varepsilon}} V \cdot \vec{\nu} \, d\sigma.$$

On ∂B_{ε} we have the bound

$$|V \cdot \vec{\nu}| \le 3\varepsilon |\nabla u|^2.$$

Thus, it is enough to show that

(5.6)
$$\lim_{\varepsilon \to 0} \varepsilon \int_{\partial B_{\varepsilon}} |\nabla u|^2 \, d\sigma = 0.$$

Note, if there is $\alpha \in (0,1)$ and a subset $D \subset \partial B_{\varepsilon}$ such that

(5.7) $|\nabla u| \le C\varepsilon^{\alpha-1}$ on *D* for some constant *C* independent of ε ,

then we conclude

$$\varepsilon \int_D |\nabla u|^2 \, d\sigma \le \varepsilon^{1+2(\alpha-1)} \sigma(\partial B_\varepsilon) = \varepsilon^{2\alpha},$$

which converges to 0. Thus, if (5.7) holds for a finite partition of ∂B_{ε} , then (5.6) follows.

We show (5.7) in the two cases:

$$D_1 \coloneqq \{x \in B_1 : x_1 > 0 \text{ or } 2|x_2| \ge |x_1|\} \text{ and } D_2 \coloneqq \{x \in B_1 : x_1 < 0 \text{ and } 2|x_2| < |x_1|\}.$$

Assume $x \in D_1 \cap \partial B_{\varepsilon}$. In the case $x_1 \ge 0$, u is harmonic in $B_{\varepsilon}(x)$ and thus Lemma 5.1 applied to u - u(x) and additionally the Hölder estimate in Theorem 2.12 for some $\alpha \in (0, 1)$ imply

$$|\nabla u(x)| \le \frac{4}{\varepsilon} \sup_{\partial B_{\varepsilon/2}(x)} |u| \le C\varepsilon^{\alpha-1}$$

for some constant C > 0 depending on n, α and $||u||_{L^{\infty}(B_{3/4})}$. In the case $x_1 < 0$, u is harmonic in $B_{|x_2|}(x)$ as dist $(x, F) = |x_2|$. As above, we have

$$|\nabla u(x)| \le C |x_2|^{\alpha - 1} \le 5C\varepsilon^{\alpha - 1}$$

since $\varepsilon^2 = |x_1|^2 + |x_2|^2 \le 5|x_2|^2$.

Next, assume $x \in D_2 \cap \partial B_{\varepsilon}$. Set $x' := (x_1, 0)$. Note that

$$|x - x'|^2 = |x_2|^2 < \frac{1}{4}|x_1|^2.$$

If $B_{|x_1|}(x') \cap \Lambda(u) = \emptyset$, then u is harmonic in $B_{|x_1|/2}(x) \subset B_{|x_1|}(x')$. Thus, Lemma 5.1, Theorem 2.12, and the estimate $\varepsilon^2 = |x_1|^2 + |x_2|^2 < \frac{5}{4}|x_1|^2$ imply

$$|\nabla u(x)| \le \frac{C}{|x_1|} \sup_{\partial B_{|x_1|/4}(x)} |u - u(x)| \le C\varepsilon^{\alpha - 1}$$

for some constant C > 0. In the case $B_{|x_1|}(x') \cap \Lambda(u) \neq \emptyset$, fix some $z \in B_{|x_1|}(x') \cap \Lambda(u)$. Then, we use that u is a solution to the zero thin obstacle problem in $B_{|x_1|}(x')$ and $x \in B_{|x_1|/2}(x')$. Thus, Lemma 5.2 and Theorem 2.12 imply

$$|\nabla u(x)| \le \frac{C}{|x_1|} \sup_{B_{\frac{3}{4}|x_1|}(x')} |u| = \frac{C}{|x_1|} \sup_{B_{\frac{3}{4}|x_1|}(x')} |u - u(z)| \le C|x_1|^{\alpha - 1} \le C\varepsilon^{\alpha - 1}$$

for some constant C > 0. This finishes the proof.

Remark 5.4. Let us comment on the dimensional constraint in Proposition 5.3.

(i) Assume that $B_1 \subset \mathbb{R}^n$ for $n \geq 3$. If F is the left half of the hyperplane, that is, $F = \{x \in B_1 \mid x_n = 0\}$ $0, x_{n-1} \leq 0$ as in Section 4, then the critical set for the regularity is $\mathcal{C} := \{x \in B_1 \mid x_n = x_{n-1} = 0\}$. We can repeat the proof with $C_{\varepsilon} = \{x \mid \operatorname{dist}(x, \mathcal{C}) < \varepsilon\}$ instead of B_{ε} up to the estimates on A_2 . On $A_3 = \partial \mathcal{C}_{\varepsilon}$ however, we only have the estimate

$$|V \cdot \vec{\nu}| \le 4(\varepsilon + |x'|) |\nabla u|^2.$$

Thus, we lose the linear factor of ε in (5.6). Hence, it is not enough to show $|\nabla u| \leq C\varepsilon^{\alpha-1}$ for some $\alpha > 0$.

(ii) If one replaces $B_1 \subset \mathbb{R}^n$ by the cylinder $Z_1 \coloneqq (-1,1)^{n-2} \times B_1^{\mathbb{R}^2}(0) \subset \mathbb{R}^n, n \geq 3$, then one may choose

$$V(x) \coloneqq (x - x') |\nabla u|^2 - 2((x - x') \cdot \nabla u) \nabla u,$$

where $x' = (x_1, ..., x_{n-2}, 0, 0) \in Z_1$. Then, for $F = \{x \in Z_1 \mid x_n = 0, x_{n-1} \leq 0\}$ the proof in this case is analogous to Proposition 5.3.

5.3. Almgren's frequency function. We are going to utilize Almgren's frequency function as in [ACS08], see also [PSU12, Sal12].

Theorem 5.5. Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, and assume that $u \in H^1(B_R)$, R > 0, is a solution to the $\mathcal{K}_{u,0,F}(B_R)$ -obstacle problem. Then,

$$N(r) = N(r, u) \coloneqq \frac{r \int_{B_r} |\nabla u|^2 \, dx}{\int_{\partial B_r} u^2 \, d\sigma}$$

is nondecreasing for 0 < r < R. Moreover, $N(r, u) \equiv \kappa \in \mathbb{R}_+$ for 0 < r < R if and only if u is homogenous of degree κ in B_R , that is,

$$x \cdot \nabla u - \kappa u = 0$$
 in B_R .

Proof. We follow the proof of [ACS08, Lemma 1]. Let

$$D(r)\coloneqq \int_{B_r} |\nabla u|^2\,dx \quad \text{and} \quad H(r)\coloneqq \int_{\partial B_r} u^2\,d\sigma.$$

Since u is a local solution to the zero thin obstacle problem at every $x_0 \in F \setminus \{0\}$, u is $C^{1,1/2}$ at every such point seen from B_R^+ and B_R^- . Thus, we have

$$D'(r) = \int_{\partial B_r} |\nabla u|^2 \, d\sigma \quad \text{and} \quad H'(r) = \frac{n-1}{r} H(r) + 2 \int_{\partial B_r} u \partial_\nu u \, d\sigma,$$

which immediately imply that

(5.8)
$$\frac{d}{dr}\log N(r) = \frac{2-n}{r} + \frac{\int_{\partial B_r} |\nabla u|^2 \, d\sigma}{\int_{B_r} |\nabla u|^2} - 2\frac{\int_{\partial B_r} u \partial_\nu u \, d\sigma}{\int_{\partial B_r} u^2 \, d\sigma}.$$

The monotonicity of N follows from $\frac{d}{dr} \log N \ge 0$. An application of Rellich's formula (5.3) to rewrite D'(r) and (3.2) to rewrite D(r) in (5.8) implies

$$\frac{d}{dr}\log N(r) = 2\left(\frac{\int_{\partial B_r} \partial_\nu u^2 \, d\sigma}{\int_{\partial B_r} u \partial_\nu u \, d\sigma} - \frac{\int_{\partial B_r} u \partial_\nu u \, d\sigma}{\int_{\partial B_r} u^2 \, d\sigma}\right).$$

г		
L		
L		

We see that $\frac{d}{dr} \log N(r) \ge 0$ is equivalent to the Cauchy–Schwarz inequality

(5.9)
$$\int_{\partial B_r} u \partial_{\nu} u \, d\sigma \le \left(\int_{\partial B_r} u^2 \, d\sigma \right)^{\frac{1}{2}} \left(\int_{\partial B_r} \partial_{\nu} u^2 \, d\sigma \right)^{\frac{1}{2}}.$$

Thus, N is nondecreasing. Moreover, N'(r) = 0 if and only if equality holds in (5.9). Hence, Cauchy-Schwarz implies that N is constant if and only if u and $\partial_{\nu} u$ are linearly dependent for every r.

Let us now assume that $N(r) = \kappa \in \mathbb{R}_+$ for every r. Thus, there exists $f : (0, R) \to \mathbb{R}$ such that $\partial_{\nu} u(x) = f(|x|)u(x)$. A simple calculation using (3.2) shows that $\kappa = rf(r)$, which implies

$$0 = r\partial_{\nu}u(x) - rf(r)u(x) = \nabla u \cdot x - \kappa u(x).$$

In the contrary case, we get that $\partial_{\nu} u = \frac{\kappa}{r} u$ which implies that u solves the ordinary differential equation

$$\frac{d}{dr}u(x) = \nabla u(x) \cdot \frac{x}{r} = \frac{\kappa}{r}u(x).$$

This further implies that u is of the form

 $u(x) = r^{\kappa}g(\theta)$ for any x with polar representation (r, θ) ,

for a function $g: \partial B_1 \to \mathbb{R}$. This implies that $N \equiv \kappa$.

Remark 5.6. The dimensional constraint in Theorem 5.5 is only due to Rellich's formula in Proposition 5.3.

The main application of Theorem 5.5 is the following growth estimate for solutions at the origin, which is the same as [PSU12, Lemma 9.14]:

Lemma 5.7 (Growth estimate). Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem with $0 \in \Lambda(u)$. Let $\kappa := N(0+, u)$. Then, there exists a constant $C = C(n, \kappa, \|u\|_{L^{\infty}(B_{3/4})})$ such that

$$\sup_{B} |u| \le Cr^{\kappa} \quad for \ every \ 0 < r < 1/2.$$

Hence, to prove the optimal regularity it remains to show a lower bound on $\kappa = N(0+, u)$.

5.4. Blowup analysis and optimal regularity. We now consider rescalings of u_r , r > 0, of u given by

(5.10)
$$u_r(x) := \frac{u(rx)}{\left(r^{1-n} \int_{\partial B_r} u^2\right)^{1/2}}.$$

Note that u_r is normalized in the sense of

(5.11)
$$\|u_r\|_{L^2(\partial B_1)} = 1.$$

Let us now assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem for $F = \{(x_1, 0) \mid x_1 \leq 0\}$ and additionally assume that $0 \in \Lambda(u)$. Then, we investigate the blowups of u at the origin, that is, the limits of u_r for a subsequence of $r \searrow 0$. The existence of a blowup is guaranteed by the following result.

Proposition 5.8. Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. Then, the family $\{u_r\}_{r>0}$ of rescalings (5.10) is uniformly bounded in $H^1 \cap C^{\alpha}_{\text{loc}}$, for some $\alpha > 0$, in the following sense: For any $r_0 > 0$ and $0 < Rr_0 < 1$ there exists a constant $C(r_0) > 0$ depending only on r_0 such that

$$||u_r||_{H^1(B_R)} + ||u_r||_{C^{\alpha}(B_{R/2})} \le C(r_0) \text{ for every } r \ge r_0.$$

The proof of Proposition 5.8 is standard and follows mainly from Theorem 5.5 and Theorem 2.12. For a proof, see for example [ACS08] or [PSU12, page 177]. For any R > 0 and some (small) $\alpha > 0$ we can use Proposition 5.8 (with double the radius and a slightly bigger Hölder exponent) to get a subsequence $u_j \coloneqq u_{r_j}$ and a limit $u_0 \in H^1(B_R) \cap C^{\alpha}(B_R)$ such that

(5.12) $\begin{aligned} u_j \to u_0 & \text{weakly in } H^1(B_R), \\ u_j \to u_0 & \text{in } L^2(\partial B_1), \\ u_j \to u_0 & \text{in } C^{\alpha}(B_R). \end{aligned}$

Note, by extracting further subsequences of $\{r_j\}_j$ for an increasing sequence of R, we actually have $u_0 \in H^1_{loc}(\mathbb{R}^n) \cap C^{\alpha}_{loc}(\mathbb{R}^n)$. Moreover, there exists a subsequence of $\{r_j\}_j$, which we keep denoting by $\{r_j\}_j$, such that the convergence in (5.12) holds for any R > 0. We call u_0 the blowup (at $x_0 = 0$) of u.

Lemma 5.9 (Homogeneity of blowups). Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem and let u_0 be a blowup of u. Then, u_0 is a non-zero global solution to the F-Signorini problem, that is, u_0 solves the $\mathcal{K}_{u_0,0,F}(B_R)$ -obstacle problem for every R > 0. Moreover, u_0 is homogenous of degree $\kappa = N(0+, u)$ in B_R for every R > 0.

Proof. From the weak H^1 -convergence of the u_j , it is clear that u_0 is harmonic in $\mathbb{R}^n \setminus F$ and superharmonic in \mathbb{R}^n . Moreover, by the C^{α} -convergence of u_j , we have $u_0 \geq 0$ on F as clearly $u_r \geq 0$ on $F \cap B_R$. On the other hand, for any $x_0 \in F \setminus \Lambda(u_0)$, the uniform convergence implies that $x_0 \in F \setminus \Lambda(u_j)$ for all sufficiently large j. In particular, such u_j are harmonic in x_0 and therefore u_0 is harmonic in x_0 . Thus, u_0 satisfies the Euler-Lagrange equations (2.4) and hence Lemma 2.10 implies that u_0 is a global solution. In particular, u_0 is non-zero due to (5.11).

As in the proof of [PSU12, Proposition 9.5] we conclude that $H(r, u_0) = \int_{\partial B_r} u_0^2 d\sigma > 0$ for any r > 0. From the Gauss-Green formula in Lemma 3.5, for every r > 0 we have

$$D(r, u_0) = \int_{B_r} |\nabla u_0|^2 = \int_{\partial B_r} u_0 \partial_\nu u_0 = \lim_{j \to \infty} \int_{\partial B_r} u_j \partial_\nu u_j = \lim_{j \to \infty} \int_{B_r} |\nabla u_j|^2.$$

Note, the limit holds due to the $C^{1,\alpha}$ -regularity of u_j away from 0, for some $\alpha > 0$, by being a (local) solution to a thin obstacle problem. Thus, together with the uniform convergence $u_j \to u_0$ and the fact that $H(r, u_0) > 0$, we conclude for every r > 0

$$N(r, u_0) = \lim_{j \to \infty} N(r, u_j) = \lim_{j \to \infty} N(rr_j, u) = \kappa.$$

Finally, Theorem 5.5 implies the homogeneity of u_0 .

Theorem 5.10 (Classification of blowups). Let n = 2, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, assume that $u \in H^1(B_1)$ is a symmetric solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem and let u_0 be a blowup of u with homogeneity $\kappa := N(0+, u)$. In particular, the following values of κ are possible:

- (i) In general, $\kappa \in \frac{1}{2}\mathbb{N}$.
- (ii) If $0 \in \Lambda(u)$ is an isolated contact point, that is, there exists $\delta > 0$ such that $B_{\delta}(0) \cap \Lambda(u) = \{0\}$, then $\kappa \in 2\mathbb{N}$.
- (iii) If $0 \in \Lambda(u)$ is an accumulation point of $\Lambda(u)$, then $\kappa \in \mathbb{N}_0 + \frac{1}{2}$.

In every of the previous cases, u_0 is of the form

$$u_0(r,\theta) = \begin{cases} r^{\kappa} \cos(\kappa\theta) & \text{if } \kappa \in 2\mathbb{N} \text{ or } \kappa \in 2\mathbb{N}_0 + \frac{3}{2} \\ -r^{\kappa} \cos(\kappa\theta) & \text{if } \kappa \in 2\mathbb{N}_0 + \frac{1}{2}. \end{cases}$$

up to a positive multiplicative constant.

Proof. First, we determine the possible values of $\kappa = N(0+, u)$. Since n = 2, we can compute several candidates for global solutions u_0 as in [PSU12, Page 178]. More precisely, by using polar coordinates and the homogeneity of u_0 we have

(5.13)
$$u_0(r,\theta) = r^{\kappa} \Phi(\theta) \quad \text{for } (r,\theta) \in [0,+\infty) \times [-\pi,\pi),$$

for some $\Phi : [-\pi, \pi) \to \mathbb{R}$. Note, to derive (5.13) we use the homogeneity of u_0 from Lemma 5.9 in the upper and lower half-space to deduce (5.13) there and conclude by continuity. In particular, since $u_0 \in C^{\alpha}_{\text{loc}}(\mathbb{R}^2)$ for some $\alpha > 0$, we immediately see that $\kappa \ge \alpha > 0$. Using the harmonicity of u_0 in the upper and lower half-space we see that Φ is of the form

$$\Phi(\theta) = a\sin(\kappa|\theta|) + b\cos(\kappa\theta)$$

for some $a, b \in \mathbb{R}$.

Next, using the symmetry one checks that u_0 satisfies the boundary conditions

- (a) $u_0 \partial_{\nu} u_0 = 0$ on $\{(x_1, 0) \mid x_1 < 0\};$
- (b) $\partial_{\nu} u_0 = 0$ on the $\{(x_1, 0) \mid x_1 > 0\}$.

We conclude that Φ is of the form

$$\Phi(\theta) = b\cos(\kappa\theta).$$

In particular, the condition (a) implies that $\kappa \in \frac{1}{2}\mathbb{N}$. Using the additional restrictions $u_0 \ge 0$ and $\partial_{\nu} u_0 \ge 0$ on F, there exists $m \in \mathbb{N}_0$ such that κ can only be of the following form:

$$\kappa = \begin{cases} 2m, 2m + 3/2 & \text{if } b > 0\\ 2m + 1, 2m + 1/2 & \text{if } b < 0. \end{cases}$$

At last, we want to rule out the values $\kappa = 2m$ or $\kappa = 2m + 1$. Let us assume that $0 \in \Lambda(u)$ is an isolated contact point, that is, there exists $\delta > 0$ such that $B_{\delta}(0) \cap \Lambda(u) = \{0\}$. In this case, u is harmonic in the punctured disk $B_{\delta}(0) \setminus \{0\}$. By the removable singularity theorem for harmonic functions, we conclude that u is also harmonic in x = 0. However, then u_0 is harmonic in x = 0 and in particular smooth. This is only possible if $\kappa \in 2\mathbb{N}$.

Next, assume that $0 \in \Lambda(u)$ is an accumulation point of $\Lambda(u)$. Then, there exists a sequence of decreasing radii $r_k > 0$ converging to 0 such that $u(-r_k/2, 0) = 0$ for every k. Note, by repeating the construction of a blowup based on the family $\{u_{r_k}\}_k$, there exists a subsequence, which will also be denoted by $\{r_k\}_k$, such that u_{r_k} converges to a blowup \tilde{u}_0 in the sense of (5.12). However, from the previous calculation, we can see that \tilde{u}_0 has to be of the same form as u_0 , hence $\tilde{u}_0 = \tilde{b}u_0$, for some $\tilde{b} \in \mathbb{R}$. Moreover, from the definition of the rescalings (5.10) we see that $u(-r_k/2, 0) = 0$ implies that $u_{r_k}(-1/2, 0) = 0$ and therefore $u_0(-1/2, 0) = 0$. Thus, $\cos(\kappa\pi) = 0$, which only allows for $\kappa \in \mathbb{N}_0 + \frac{1}{2}$.

Theorem 5.11 (Optimal regularity). Let n = 2, $F = \{(x_1, 0) | x_1 \leq 0\}$, and assume that $u \in H^1(B_1)$ is a solution to the $\mathcal{K}_{u,0,F}(B_1)$ -obstacle problem. Then, $u \in C^{1/2}_{\text{loc}}(B_1)$.

Proof. Clearly, if $0 \notin \Lambda(u)$, then u is smooth in a small neighbourhood of $x_0 = 0$ and therefore there is nothing to do. Assume $0 \in \Lambda(u)$. In the case when u is symmetric, the result follows from the growth estimate Lemma 5.7 and the classification of blowups Theorem 5.10, which implies $\kappa = N(0+, u) \ge 1/2$. If u is not symmetric, then consider $w \in H^1(B_1)$ as the weak solution to the boundary value problem

$$\begin{cases} -\Delta w = 0 & \text{in } B_1 \\ w = g & \text{on } \partial B_1, \end{cases}$$

where g(x,y) = (u(x,y) - u(x,-y))/2. It is easy to check that $v \coloneqq u - w$ is a symmetric solution to the $\mathcal{K}_{v,0,F}(B_1)$ -obstacle problem. Thus, $v \in C^{1/2}_{\text{loc}}(B_1)$ and since w is smooth, we have $u \in C^{1/2}_{\text{loc}}(B_1)$.

Remark 5.12. We expect that one can extend the approach in Section 5 to include sufficiently regular obstacle data ψ following the ideas in [CSS08]. Let us give some instructions. For this, fix an obstacle ψ with extension $\Psi \in H^1(B_1)$, let $u \in H^1(B_1)$ solve the $\mathcal{K}_{u,\psi,F}(B_1)$ -obstacle problem, and set $v := u - \Psi$. Then, v satisfies the inhomogeneous zero-obstacle problem (3.15) for $f := \Delta \Psi$. First, one shows Rellich's formula for v:

$$\int_{\partial B_r} |\nabla v|^2 \, d\sigma = 2 \int_{\partial B_r} (\partial_\nu v)^2 \, d\sigma + \frac{2}{r} \int_{B_r} (x \cdot \nabla v) f \, dx.$$

This can then be used to prove the monotonicity of the modified Almgren's frequency

$$N^{f}(r) \coloneqq (1 + C_0 r) \frac{r \int_{B_r} |\nabla v|^2 - f v \, dx}{\int_{\partial B_r} u^2 \, d\sigma}$$

for a sufficiently large constant $C_0 > 0$, similar to [CSS08, Theorem 3.1 and Appendix]. Then, as in [CSS08, Section 6], one constructs a blowup v_0 from the sequence v_r , defined as in (5.10), using N^f . Note that

$$-\Delta v_r(x) = \frac{r^2 f(rx)}{\int_{\partial B_r} v^2 \, d\sigma}$$

which converges to 0 for $r \to 0$, at least for Hölder regular Ψ and bounded $f = \Delta \Psi$. Note, Hölder regularity of Ψ implies Hölder regularity of v due to Theorem 2.12. Thus, the blowup v_0 is a global solution to the homogeneous zero-obstacle Signorini problem as in Lemma 5.9. Hence, the homogeneity of the blowup $\kappa := N^f(0+)$ is bounded from below by $N(r, v_0) \geq 1/2$. One concludes the optimal regularity similar to [CSS08, Theorem 6.7].

6. Application: Obstacles with jump-type discontinuities

Let $\Omega \subset \mathbb{R}^n$ be open and $F \subset \Omega$ be relatively closed. Moreover, assume that there is a finite partition of F, that is, there is a positive integer N, relatively closed subsets $F_i \subset \Omega$, $i = 1, \ldots, N$, such that $F = \bigcup_{i=1}^N F_i$, and $F_i \cap F_j$ is at most a set of codimension 2 for each $i \neq j$. For the obstacle, let $\psi_i \in C(F_i)$ and consider $\psi : F \to \mathbb{R}$ as the glued obstacle:

(6.1)
$$\psi(x) = \sup_{j:x \in F_i \cap F_j} \psi_j(x) \quad \text{if } x \in F_i.$$

For any $x \in F$ let us denote $J_x := \{j : x \in F_j\}.$

Theorem 6.1. Let Ω , F and ψ be as described above. Assume that $u \in H^1(\Omega)$ is a solution to the $\mathcal{K}_{u,\psi,F}(\Omega)$ obstacle problem. Fix some $x_0 \in F$. If there exists i and $\delta > 0$ such that $x_0 \in F_i$ and

(6.2)
$$\psi_i(x_0) > \sup_{j \in J_{x_0} \setminus \{i\}} \sup_{F_j \cap B_{\delta}(x_0)} \psi_j,$$

then there exists $0 < \delta' \leq \delta$ such that u is a solution to the $\mathcal{K}_{u,\psi_i,F_i}(B_{\delta'}(x_0))$ -obstacle problem.

Proof. By definition of ψ in (6.1) it is clear that ψ is upper semi-continuous on F. Thus, Theorem 2.7 implies that u is continuous in Ω . Then, the obstacle condition $u \ge \psi$, (6.2), and the continuity of u and the ψ_j 's imply that there exists $0 < \delta' \le \delta$ such that

(6.3)
$$u > \psi_j$$
 in $F_j \cap B_{\delta'}$ for every $j \neq i$.

To see this, assume the contrary. Then, there exists a sequence δ_n converging to 0, $j_n \in J_{x_0} \setminus \{i\}$, and $x_n \in F_{j_n} \cap B_{\delta_n}(x_0)$ such that $u(x_n) = \psi_j(x_n)$. However, as $x_n \to x_0$ and u is continuous, we then have

$$\lim_{n \to \infty} \psi_{j_n}(x_n) = \lim_{n \to \infty} u(x_n) = u(x_0) \ge \psi_i(x_0),$$

which contradicts (6.2).

Now, let us check the Euler-Lagrange equations (2.4) with respect to $B_{\delta'}(x_0)$, F_i , and ψ_i . Clearly, u is superharmonic in $B_{\delta'}(x_0)$ by Proposition 2.8. Due to (6.3), we have $\Lambda(u) \cap B_{\delta'} = \{x \in F_i \mid u(x) = \psi_i(x)\} \cap B_{\delta'}(x_0)$. Thus, u is harmonic in $B_{\delta'}(x_0) \cap \{x \in F_i \mid u(x) = \psi_i(x)\}$ by Theorem 2.7. At last, $u \ge \psi_i$ on F_i clearly holds. Hence, Lemma 2.10 implies that u solves the $\mathcal{K}_{u,\psi_i,F_i}(B_{\delta'}(x_0))$ -obstacle problem. \Box

Next, we present implications of Theorem 6.1 for thin obstacle problems within the settings of our regularity results for Signorini-type problems. The optimal regularity result in Theorem 5.11 implies:

Corollary 6.2. Let $\Omega = B_1 \subset \mathbb{R}^2$, $F = B'_1$, and $\psi : B'_1 \to \mathbb{R}$ be given by

$$\psi(x_1,0) \coloneqq \mathbb{1}_{(-1,0]}(x_1)$$

Then, any solution $u \in H^1(B_1)$ to the $\mathcal{K}_{u,\psi,B'}(B_1)$ -obstacle problem is $C^{1/2}$ -regular at $x_0 = 0$.

Proof. Clearly, every assumption in Theorem 6.1 is satisfied for obvious choices of F_i and ψ_i . Thus, there is $\delta' > 0$ such that u solves the $\mathcal{K}_{u,1,(-\delta',0]}(B_{\delta'}(0))$ -obstacle problem. It is easy to see that $v \coloneqq u-1$ solves the $\mathcal{K}_{v,0,(-\delta',0]}(B_{\delta'}(0))$ -obstacle problem. Theorem 5.11 implies that v is $C_{\text{loc}}^{1/2}(B_{\delta'}(0))$. Hence, u is $C^{1/2}$ -regular at $x_0 = 0$.

Remark 6.3. As discussed in Remark 5.12, we expect that a variant of the regularity result Theorem 5.11 for non-zero obstacles hold. Hence, a variant of Corollary 6.2 for obstacles of the form

$$\psi(x_1,0) = \psi_1(x_1) \mathbb{1}_{(-1,0]}(x_1) + \psi_2(x_1) \mathbb{1}_{(0,1)}(x_1),$$

where ψ_1, ψ_2 are sufficiently regular and satisfy $\psi_1(0) > \psi_2(0)$, holds.

The almost optimal regularity result in Theorem 4.10 implies:

Corollary 6.4. Let $\Omega = B_1 \subset \mathbb{R}^n$, $n \ge 2$, and $F = \{(x', x_{n-1}, 0) \mid 0 \le x_{n-1} < 1\}$. Assume that $\psi : B'_1 \to \mathbb{R}$ is of the form

$$\psi(x',0) = \psi_1(x') \mathbb{1}_F(x') + \psi_2(x') \mathbb{1}_{B_1' \setminus F}(x'),$$

for ψ_1 , ψ_2 upper semi-continuous. If ψ_1 is in $C^{\beta_1}(F)$, for some $\beta_1 \in (0,1)$, has an extension as in Theorem 4.10, and satisfies

$$\psi_1(0) > \sup_{B_\delta \cap (B_1' \setminus F)} \psi_2$$

for some $\delta > 0$, then any solution $u \in H^1(B_1)$ to the $\mathcal{K}_{u,\psi,B'_1}(B_1)$ -obstacle problem is C^{β} regular at $x_0 = 0$, for any $\beta \in (0, 1/2)$ such that $\beta \leq \beta_1$.

The optimal regularity result in Theorem 3.9 implies:

Corollary 6.5. Let $\Omega = B_1 \subset \mathbb{R}^n$, $n \geq 2$. Assume that ψ is given as in (6.1) for a finite partition F_j of B'_1 and $\psi_j : F_j \to \mathbb{R}$ satisfy the assumptions in Theorem 3.9. If there exists i and $x_0 \in F_i$ such that (6.2) holds, then any solution $u \in H^1(B_1)$ to the $\mathcal{K}_{u,\psi,B'_1}(B_1)$ -obstacle problem, whose contact set satisfies the capacity density condition (3.1) at $x_0 \in \Lambda(u)$, is $C^{1/2}$ -regular at x_0 .

At last, let us comment further on the Lipschitz regularity result obtained in [Kin71]. For this, let us show the following:

Lemma 6.6. Let $\Omega = B_1 \subset \mathbb{R}^2$, $F = \{(x_1, 0) \mid x_1 \leq 0\}$, and $\psi \in C^{1,\alpha}(F)$, $\alpha \in (0, 1)$, that is nonnegative and satisfies $\psi(0) = 0$. Then, there exists a $C^{1,\alpha}$ -extension $\Psi : B'_1 \to \mathbb{R}$ of ψ such that any solution $u \in H^1_0(B_1)$ to the $\mathcal{K}_{0,\psi,F}(B_1)$ -obstacle problem is a solution to the $\mathcal{K}_{0,\Psi,B'_1}(B_1)$ -obstacle problem.

Proof. Let us choose Ψ by

$$\Psi(x_1) \coloneqq \begin{cases} \psi(x_1) & \text{if } x_1 \le 0\\ \psi'(0)x_1 & \text{if } x_1 > 0. \end{cases}$$

Clearly, $\Psi \in C^{1,\alpha}(B'_1)$. Since ψ is nonnegative and u = 0 on ∂B_1 , the maximum principle implies that u is nonnegative. Thus, $u \ge \Psi$ on B'_1 . In particular

$$\{x \in F \mid u(x) = \psi(x)\} = \{x \in B'_1 \mid u(x) = \Psi(x)\}.$$

Hence, u satisfies the Euler-Lagrange equations (2.4) with respect to Ψ and therefore u solves the $\mathcal{K}_{0,\Psi,B'_1}(B_1)$ obstacle problem.

The setting considered in Lemma 6.6 coincides with the one in [Kin71, Theorem 2], where the author proved Lipschitz regularity of the solution u. However, Lemma 6.6 shows that this solution is actually a solution to a thin obstacle problem. Thus, the combination of Lemma 6.6 and the regularity result in [Ric78] imply that the solution u is Lipschitz in B_1 and is $C^{1,\beta}$, for $\beta = \min\{1/2, \alpha\}$, in the upper and lower half ball up to the horizontal line.

Note, Lemma 6.6 is based on the same observation as Theorem 6.1, that is: Any change of the obstacle that happens below a certain threshold, which depends on the boundary data, does not change the solution of the associated obstacle problem.

References

- [AC04] I. Athanasopoulos and L. A. Caffarelli. Optimal regularity of lower dimensional obstacle problems. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 310(Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 35 [34]):49–66, 226, 2004. reprinted in Journal of Mathematical Sciences, 132:274–284, 2006.
- [AC22] Yu. A. Alkhutov and A. G. Chechkina. Many-dimensional Zaremba problem for an inhomogeneous p-Laplace equation. Dokl. Math., 106(1):243–246, 2022. Translation of Dokl. Akad. Nauk 505 (2022), 37–41.
- [ACF84] Hans Wilhelm Alt, Luis A. Caffarelli, and Avner Friedman. Variational problems with two phases and their free boundaries. Trans. Amer. Math. Soc., 282(2):431–461, 1984.
- [ACM22] Yurij A. Alkhutov, Gregory A. Chechkin, and Vladimir G. Maz'ya. Boyarsky-Meyers estimate for solutions to Zaremba problem. Arch. Ration. Mech. Anal., 245(2):1197–1211, 2022.
- [ACS08] I. Athanasopoulos, L. A. Caffarelli, and S. Salsa. The structure of the free boundary for lower dimensional obstacle problems. Amer. J. Math., 130(2):485–498, 2008.
- [Ada82] David R. Adams. Capacity and the obstacle problem. Appl. Math. Optim., 8(1):39–57, 1982.
- [AH96] David R. Adams and Lars Inge Hedberg. Function spaces and potential theory, volume 314 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1996.
- [AKSZ07] Hiroaki Aikawa, Tero Kilpeläinen, Nageswari Shanmugalingam, and Xiao Zhong. Boundary Harnack principle for *p*-harmonic functions in smooth Euclidean domains. *Potential Anal.*, 26(3):281–301, 2007.
- [Alm00] Frederick J. Almgren, Jr. Almgren's big regularity paper, volume 1 of World Scientific Monograph Series in Mathematics. World Scientific Publishing Co., Inc., River Edge, NJ, 2000. Q-valued functions minimizing Dirichlet's integral and the regularity of area-minimizing rectifiable currents up to codimension 2, With a preface by Jean E. Taylor and Vladimir Scheffer.

- [BB11] Anders Björn and Jana Björn. Nonlinear potential theory on metric spaces, volume 17 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2011.
- [BC24] Sun-Sig Byun and Yumi Cho. Calderón-Zygmund estimates for nonlinear elliptic obstacle problems with log-BMO matrix weights. Discrete Contin. Dyn. Syst. Ser. B, 29(12):4772–4792, 2024.
- [BCO16] Sun-Sig Byun, Yumi Cho, and Jihoon Ok. Global gradient estimates for nonlinear obstacle problems with nonstandard growth. Forum Math., 28(4):729–747, 2016.
- [BDM11] Verena Bögelein, Frank Duzaar, and Giuseppe Mingione. Degenerate problems with irregular obstacles. J. Reine Angew. Math., 650:107–160, 2011.
- [BL74] Alain Bensoussan and Jacques-Louis Lions. Contrôle impulsionnel et contrôle continu. Méthode des inéquations quasi variationnelles non linéaires. C. R. Acad. Sci. Paris Sér. A, 278:675–679, 1974.
- [BL75] A. Bensoussan and J.-L. Lions. Optimal impulse and continuous control: method of nonlinear quasi-variational inequalities. Trudy Mat. Inst. Steklov., 134:5–22, 407, 1975.
- [BL24] Anna Kh Balci and Ho-Sik Lee. Zaremba problem with degenerate weights. arXiv preprint arXiv:2403.19813, 2024.
- [BS93] J. Banasiak and J. Szczepaniak. On regularity of solutions to inner obstacle problems. Z. Anal. Anwendungen, 12(3):401–404, 1993.
- [Caf77] Luis A. Caffarelli. The regularity of free boundaries in higher dimensions. Acta Math., 139(3-4):155–184, 1977.
- [Caf79] L. A. Caffarelli. Further regularity for the Signorini problem. Comm. Partial Differential Equations, 4(9):1067–1075, 1979.
- [Caf98] L. A. Caffarelli. The obstacle problem revisited. J. Fourier Anal. Appl., 4(4-5):383–402, 1998.
- [CC95] L. A. Caffarelli and X. Cabré. Fully nonlinear elliptic equations, volume 43 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1995.
- [Cho91] Hi Jun Choe. A regularity theory for a general class of quasilinear elliptic partial differential equations and obstacle problems. Arch. Rational Mech. Anal., 114(4):383–394, 1991.
- [CJK02] Luis A. Caffarelli, David Jerison, and Carlos E. Kenig. Some new monotonicity theorems with applications to free boundary problems. Ann. of Math. (2), 155(2):369–404, 2002.
- [CK80] L. A. Caffarelli and D. Kinderlehrer. Potential methods in variational inequalities. J. Analyse Math., 37:285–295, 1980.
- [CR77] L. A. Caffarelli and N. M. Rivière. Asymptotic behaviour of free boundaries at their singular points. Ann. of Math. (2), 106(2):309–317, 1977.
- [CSS08] Luis A. Caffarelli, Sandro Salsa, and Luis Silvestre. Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian. *Invent. Math.*, 171(2):425–461, 2008.
- [DL76] G. Duvaut and J.-L. Lions. Inequalities in mechanics and physics, volume 219 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin-New York, 1976. Translated from the French by C. W. John.
- [EH10] Michela Eleuteri and Jens Habermann. Calderón-Zygmund type estimates for a class of obstacle problems with p(x) growth. J. Math. Anal. Appl., 372(1):140–161, 2010.
- [Ele07] Michela Eleuteri. Regularity results for a class of obstacle problems. Appl. Math., 52(2):137–170, 2007.
- [Fal14] Kenneth Falconer. Fractal geometry. John Wiley & Sons, Ltd., Chichester, third edition, 2014. Mathematical foundations and applications.
- [Fic64] Gaetano Fichera. Problemi elastostatici con vincoli unilaterali: Il problema di Signorini con ambigue condizioni al contorno. Atti Accad. Naz. Lincei Mem. Cl. Sci. Fis. Mat. Natur. Sez. Ia (8), 7:91–140, 1963/64.
- [FM79] Jens Frehse and Umberto Mosco. Variational inequalities with one-sided irregular obstacles. Manuscripta Math., 28(1-3):219–233, 1979.
- [FM82] Jens Frehse and Umberto Mosco. Irregular obstacles and quasivariational inequalities of stochastic impulse control. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 9(1):105–157, 1982.
- [FM84] J. Frehse and U. Mosco. Wiener obstacles. In Nonlinear partial differential equations and their applications. Collège de France seminar, Vol. VI (Paris, 1982/1983), volume 109 of Res. Notes in Math., pages 225–257. Pitman, Boston, MA, 1984.
- [FOT11] Masatoshi Fukushima, Yoichi Oshima, and Masayoshi Takeda. Dirichlet forms and symmetric Markov processes, volume 19 of De Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, extended edition, 2011.
- $[FR16] Xavier Fernández-Real. C^{1, \alpha} estimates for the fully nonlinear Signorini problem. Calc. Var. Partial Differential Equations, 55(4): Art. 94, 20, 2016.$
- [FR22] Xavier Fernández-Real. The thin obstacle problem: a survey. Publ. Mat., 66(1):3–55, 2022.
- [Fre72] Jens Frehse. On the regularity of the solution of a second order variational inequality. Boll. Un. Mat. Ital. (4), 6:312–315, 1972.
- [Fre75] Jens Frehse. Two dimensional variational problems with thin obstacles. Math. Z., 143(3):279–288, 1975.
- [Fre77] Jens Frehse. On Signorini's problem and variational problems with thin obstacles. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 4(2):343–362, 1977.
- [Fri82] Avner Friedman. Variational principles and free-boundary problems. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1982. Pure and Applied Mathematics.
- [FRRO22] Xavier Fernández-Real and Xavier Ros-Oton. Regularity theory for elliptic PDE, volume 28 of Zurich Lectures in Advanced Mathematics. EMS Press, Berlin, 2022.
- [Giu71] Enrico Giusti. Superfici minime cartesiane con ostacoli discontinui. Arch. Rational Mech. Anal., 40:251–267, 1971.
- [Gri21] Antonio Giuseppe Grimaldi. Regularity results for solutions to a class of obstacle problems. Nonlinear Anal. Real World Appl., 62:Paper No. 103377, 26, 2021.

- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [HKM06] Juha Heinonen, Tero Kilpeläinen, and Olli Martio. Nonlinear potential theory of degenerate elliptic equations. Dover Publications, Inc., Mineola, NY, 2006. Unabridged republication of the 1993 original.
- [HLW14] Yongpan Huang, Dongsheng Li, and Lihe Wang. Boundary behavior of solutions of elliptic equations in nondivergence form. Manuscripta Math., 143(3-4):525–541, 2014.
- [Ior82] Iordan V. Iordanov. Solutions höldériennes d'inéquations variationnelles à contraintes discontinues. I. Serdica, 8(3):296–306, 1982.
- [Ior83] Iordan V. Iordanov. Solutions Höldériennes d'inéquations variationnelles à contraintes discontinues. II. Serdica, 9(3):289–300, 1983.
- [Ior73] Iordan Iordanov. Regularity of the solution of a certain elliptic variational inequality with two constraints. Annuaire Univ. Sofia Fac. Math. Méc., 67:169–190, 1972/73.
- [JK95] David Jerison and Carlos E. Kenig. The inhomogeneous Dirichlet problem in Lipschitz domains. J. Funct. Anal., 130(1):161–219, 1995.
- [JOS03] Sławomir Jagodziński, Anna Olek, and Kuba Szczepaniak. Lipschitz character of solutions to the double inner obstacle problems. Irish Math. Soc. Bull., (52):11–17, 2003.
- [JOS05] Sławomir Jagodziński, Anna Olek, and Kuba Szczepaniak. Inner obstacle problem: convergence of the solutions for impediments with varying domains. Int. J. Pure Appl. Math., 24(2):265–270, 2005.
- [JOS08] Sławomir Jagodziński, Anna Olek, and Kuba Szczepaniak. Lipschitz character of solutions to the inner obstacle problems. Irish Math. Soc. Bull., (61):15–27, 2008.
- [JOS09] Sławomir Jagodziński, Anna Olek, and Kuba Szczepaniak. Regularity of the solutions to the inner obstacle problems, identification of the inner obstacles with the Dirichlet boundary problems. Int. J. Math. Anal. (Ruse), 3(21-24):1003– 1010, 2009.
- [Kin71] David Kinderlehrer. Variational inequalities with lower dimensional obstacles. Israel J. Math., 10:339–348, 1971.
- [KK73] L. I. Kamynin and B. N. Khimchenko. Lipschitz boundary estimates for a solution of a second order elliptic-parabolic equation. Soviet Math. Dokl., 14:1407–1412, 1973.
- [KK74] L. I. Kamynin and B. N. Khimchenko. A maximum principle and Lipschitz boundary estimates for the solution of a second order elliptic-parabolic equation. *Sibirsk. Mat. Ž.*, 15:343–367, 461, 1974.
- [KN77] D. Kinderlehrer and L. Nirenberg. Regularity in free boundary problems. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 4(2):373–391, 1977.
- [KPS15] Herbert Koch, Arshak Petrosyan, and Wenhui Shi. Higher regularity of the free boundary in the elliptic Signorini problem. Nonlinear Anal., 126:3–44, 2015.
- [KS80] David Kinderlehrer and Guido Stampacchia. An introduction to variational inequalities and their applications, volume 88 of Pure and Applied Mathematics. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [Lan72] N. S. Landkof. Foundations of modern potential theory, volume Band 180 of Die Grundlehren der mathematischen Wissenschaften. Springer-Verlag, New York-Heidelberg, 1972. Translated from the Russian by A. P. Doohovskoy.
- [Lew72] Hans Lewy. On the coincidence set in variational inequalities. J. Differential Geometry, 6:497–501, 1972.
- [Lew68] Hans Lewy. On a variational problem with inequalities on the boundary. J. Math. Mech., 17:861–884, 1967/68.
 [Lie86] Gary M. Lieberman. Mixed boundary value problems for elliptic and parabolic differential equations of second order.
- J. Math. Anal. Appl., 113(2):422–440, 1986. [Lie89] Gary M. Lieberman. Optimal Hölder regularity for mixed boundary value problems. J. Math. Anal. Appl., 143(2):572–586, 1989.
- [LS67] J.-L. Lions and G. Stampacchia. Variational inequalities. Comm. Pure Appl. Math., 20:493–519, 1967.
- [LS69] Hans Lewy and Guido Stampacchia. On the regularity of the solution of a variational inequality. Comm. Pure Appl. Math., 22:153–188, 1969.
- [LS71] Hans Lewy and Guido Stampacchia. On existence and smoothness of solutions of some non-coercive variational inequalities. Arch. Rational Mech. Anal., 41:241–253, 1971.
- [Mon03] R. Monneau. On the number of singularities for the obstacle problem in two dimensions. J. Geom. Anal., 13(2):359–389, 2003.
- [Mos86] Umberto Mosco. Pointwise potential estimates for elliptic obstacle problems. In Nonlinear functional analysis and its applications, Part 2 (Berkeley, Calif., 1983), volume 45, Part 2 of Proc. Sympos. Pure Math., pages 207–217. Amer. Math. Soc., Providence, RI, 1986.
- [Mos87] Umberto Mosco. Wiener criterion and potential estimates for the obstacle problem. Indiana Univ. Math. J., 36(3):455–494, 1987.
- [MR10] Vladimir Maz'ya and Jürgen Rossmann. *Elliptic equations in polyhedral domains*, volume 162 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [MZ91] J. H. Michael and William P. Ziemer. Existence of solutions to obstacle problems. *Nonlinear Anal.*, 17(1):45–71, 1991.
- [Ok17] Jihoon Ok. Gradient continuity for nonlinear obstacle problems. *Mediterr. J. Math.*, 14(1):Paper No. 16, 24, 2017.
- [PSU12] Arshak Petrosyan, Henrik Shahgholian, and Nina Uraltseva. Regularity of free boundaries in obstacle-type problems, volume 136 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012.
- [Rel40] Franz Rellich. Darstellung der Eigenwerte von $\Delta u + \lambda u = 0$ durch ein Randintegral. Math. Z., 46:635–636, 1940.
- [Ric78] David Joseph Allyn Richardson. VARIATIONAL PROBLEMS WITH THIN OBSTACLES. ProQuest LLC, Ann Arbor, MI, 1978. Thesis (Ph.D.)–The University of British Columbia (Canada).

- [Saf08] Mikhail V Safonov. Boundary estimates for positive solutions to second order elliptic equations. arXiv preprint arXiv:0810.0522, 2008.
- [Sal12] Sandro Salsa. The problems of the obstacle in lower dimension and for the fractional Laplacian. In Regularity estimates for nonlinear elliptic and parabolic problems, volume 2045 of Lecture Notes in Math., pages 153–244. Springer, Heidelberg, 2012.
- [Ser64] James Serrin. Local behavior of solutions of quasi-linear equations. Acta Math., 111:247–302, 1964.
- [Sha68] Eliahu Shamir. Regularization of mixed second-order elliptic problems. Israel J. Math., 6:150–168, 1968.
- [Shi92] Norio Shimakura. Partial differential operators of elliptic type, volume 99 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1992. Translated and revised from the 1978 Japanese original by the author.
- [Sig59] A. Signorini. Questioni di elasticità non linearizzata e semilinearizzata. Rend. Mat. e Appl. (5), 18:95–139, 1959.
- [Sta64] Guido Stampacchia. Formes bilinéaires coercitives sur les ensembles convexes. C. R. Acad. Sci. Paris, 258:4413–4416, 1964.
- [SV72] Guido Stampacchia and Alfonso Vignoli. A remark on variational inequalities for a second order nonlinear differential operator with non Lipschitz obstacles. *Boll. Un. Mat. Ital.* (4), 5:123–131, 1972.
- [TNH23] Minh-Phuong Tran, Thanh-Nhan Nguyen, and Phuoc-Nguyen Huynh. Calderón-Zygmund type estimates for singular quasilinear elliptic obstacle problems with measure data. *Studia Math.*, 271(3):287–319, 2023.
- [Wan92] Lihe Wang. On the regularity theory of fully nonlinear parabolic equations. II. Comm. Pure Appl. Math., 45(2):141– 178, 1992.

¹Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Seoul 08826, Republic of Korea

Email address: kiahm@snu.ac.kr

²School of Mathematics, Korea Institute for Advanced Study, Seoul 02455, Republic of Korea Email address: sechan@kias.re.kr

³FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 100131, 33501 BIELEFELD, GERMANY *Email address:* wschefer@math.uni-bielefeld.de