THE SEA ALGORITHM FOR ENDOMORPHISMS OF SUPERSINGULAR ELLIPTIC CURVES

TRAVIS MORRISON, LORENZ PANNY, JANA SOTÁKOVÁ, AND MICHAEL WILLS

ABSTRACT. For a prime p > 3 and a supersingular elliptic curve E defined over \mathbb{F}_{p^2} with $j(E) \notin \{0, 1728\}$, consider an endomorphism α of E represented as a composition of L isogenies of degree at most d. We prove that the trace of α may be computed in $O(n^4(\log n)^2 + dLn^3)$ bit operations, where $n = \log(p)$, using a generalization of the SEA algorithm for computing the trace of the Frobenius endomorphism of an ordinary elliptic curve. When $L \in O(\log p)$ and $d \in O(1)$, this complexity matches the heuristic complexity of the SEA algorithm. Our theorem is unconditional, unlike the complexity analysis of the SEA algorithm, since the kernel of an arbitrary isogeny of a supersingular elliptic curve is defined over an extension of constant degree, independent of p. We also provide practical speedups, including a fast algorithm to compute the trace of α modulo p.

1. INTRODUCTION

Let p be a prime and let E be a supersingular elliptic curve defined over \mathbb{F}_{p^2} . An important subroutine in some algorithms for computing the endomorphism ring of E involves computing the trace of an endomorphism of E. Indeed, the trace pairing on $\operatorname{End}(E)$ determines the structure of $\operatorname{End}(E)$ equipped with the quadratic form deg as a quadratic lattice up to isometry and hence its isomorphism class as an order in a quaternion algebra. More concretely, an efficient algorithm for computing the trace of an endomorphism of E yields an efficient algorithm for deciding whether 4 endomorphisms are a \mathbb{Z} -basis for $\operatorname{End}(E)$: The endomorphisms $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ span $\operatorname{End}(E)$ if and only if $\det(\operatorname{tr}(\alpha_i \alpha_i)) = p^2$.

We can compute the trace of $\alpha \in \operatorname{End}(E)$ using a generalization of Schoof's algorithm [20], as observed by Kohel in his thesis [12, Theorem 81]. We will assume the degree of α is smooth and thus α may be represented as a sequence of small degree isogenies. To compute tr α , we compute tr $\alpha \mod \ell$ for enough small primes $\ell \leq B$ such that $\prod_{\ell < B} \ell > 4\sqrt{\deg \alpha}$. We can then recover tr α using the Chinese remainder theorem. For a prime ℓ , if $E[\ell] \subseteq \ker \alpha$, then tr $\alpha \equiv 0 \pmod{\ell}$. Otherwise, there exists a $P \in E[\ell]$ such that $\alpha(P)$ is nonzero, and to compute $t_{\ell} = \operatorname{tr} \alpha \mod \ell$ it suffices to find a $0 \leq c < \ell$ such that and $\alpha^2(P) + [\deg \alpha]P = c\alpha(P)$. If such a P exists, then $t_{\ell} = c$. And to check whether such a P exists, we can compute the restrictions of $\alpha^2 + [\deg \alpha]$ and $[c] \circ \alpha$ to $E[\ell]$. This involves computing the coordinate functions of $\alpha|_{E[\ell]}$ and $[1]|_{E[\ell]}$ and then performing arithmetic in $\operatorname{End}(E[\ell])$ using the group law on E to compute

$$\begin{aligned} \alpha^{2} \big|_{E[\ell]} &= \left(\alpha \big|_{E[\ell]} \right)^{2}, \quad [d] \big|_{E[\ell]} = d[1] \big|_{E[\ell]}, \quad (c\alpha) \big|_{E[\ell]} = c \left(\alpha \big|_{E[\ell]} \right), \\ \text{and} \left(\alpha^{2} + [d] \right) \big|_{E[\ell]} &= \alpha^{2} \big|_{E[\ell]} + [d] \big|_{E[\ell]}. \end{aligned}$$

These restrictions can be represented as (a(x), b(x)y) where $a, b \in \mathbb{F}_q[x]/(\psi_\ell)$ where ψ_ℓ is the ℓ th division polynomial of E.

To speed up this computation, which involves polynomial arithmetic modulo the degree $(\ell^2 - 1)/2$ polynomial ψ_{ℓ} , we might look for inspiration in the SEA algorithm, and specifically to Elkies' improvement to Schoof's algorithm, for computing the trace of the Frobenius endomorphism π_E of an ordinary curve E. Elkies' idea is to note that, when an odd prime ℓ splits in $\mathbb{Z}[\pi_E]$, E admits a rational ℓ -isogeny $\phi: E \to E'$ and thus the division polynomial ψ_{ℓ} has a nontrivial factor $h \in \mathbb{F}_q[x]$. Indeed, the polynomial h is the kernel polynomial of ϕ and has degree $(\ell - 1)/2$. Such a prime is called an Elkies prime for E. Asymptotically, 50% of the primes are Elkies primes for E. Since ker ϕ is defined over \mathbb{F}_q , it is invariant under π_E so $\pi_E|_{E[\ell]}$ has a nontrivial eigenvalue $\lambda \in \mathbb{Z}/\ell\mathbb{Z}$. Then $t_{\ell} \equiv \lambda + q/\lambda \pmod{\ell}$. Since deg h is smaller than deg ψ_{ℓ} by a factor on the order of ℓ , computing $\pi_E|_{\ker \phi}$ is about ℓ times faster than computing $\pi_E|_{E[\ell]}$. This gives a linear speedup in computing t_{ℓ} whenever ℓ is an Elkies prime. Primes ℓ that are not Elkies primes are called Atkin primes due to an improvement of Atkin for speeding up the computation of t_{ℓ} . Atkin's method does not give an asymptotic speedup, and not even the Generalized Riemann

Date: January 28, 2025.

²⁰²⁰ Mathematics Subject Classification. 11Y16, 68W30, 14K02.

Key words and phrases. Supersingular elliptic curve, SEA algorithm, isogeny, endomorphism, trace.

Hypothesis is strong enough to prove that there are enough small Elkies primes for E so that the SEA algorithm is asymptotically faster than Schoof's algorithm, as shown by Shparlinski and Sutherland [23, Corollary 15]. In practice, there are enough Elkies primes and the SEA algorithm performs much better than one is able to prove. Atkin primes are skipped entirely in Sutherland's point-counting records using the SEA algorithm [26] together with his algorithm for computing *instantiated modular polynomials* [27]. Finally, Shparlinski and Sutherland [22] show that when one averages over elliptic curves over \mathbb{F}_q , there are as many Elkies primes as Atkin primes $\ell < L$ so long as $L \geq (\log p)^{\epsilon}$ for fixed $\epsilon > 0$ and sufficiently large q, and in another direction, the same authors show in [23] that for a fixed elliptic curve E over \mathbb{Q} and for almost all primes p of good reduction for E, the SEA algorithm is faster than Schoof's algorithm for computing the number of points of the reduction of E modulo p.

Returning to our setup, and with Elkies' improvement to Schoof's algorithm and the above discussion in mind, we now leverage the assumption that E is supersingular and defined over \mathbb{F}_{p^2} and that $j(E) \notin \{0, 1728\}$. Then every isogeny of E defined over $\overline{\mathbb{F}_{p^2}}$ is actually defined over \mathbb{F}_{p^2} ; we prove a more general version of this in Proposition 2.2. Therefore every prime ℓ is an Elkies prime for E! Let α be an arbitrary endomorphism of E and let $h \in \mathbb{F}_{p^2}[x]$ be the kernel polynomial of an ℓ -isogeny ψ of E. There is no reason for ker ψ to be stable under α , i.e., α is not necessarily an endomorphism of ker ψ , unlike the Frobenius endomorphism. However, we can still compute restrictions of $c\alpha, \alpha^2$, $[\deg \alpha]$, add them together, and test for equality using arithmetic in the ring $\mathbb{F}_{p^2}[x]/(h(x))$. One can interpret this as performing arithmetic in Hom(ker $\psi, E[\ell]$), see Section 3.1. This allows us to compute $t_{\ell} := \operatorname{tr} \alpha \mod \ell$ as in the SEA algorithm. We show in Theorem 3.10 that when α is the composition of $L \in O(\log p)$ isogenies of degree $d \in O(1)$ and $\ell \in O(\log p)$ that t_{ℓ} can be computed with an expected $O(n^3(\log n)^3)$ bit operations, where $n = \lceil \log p \rceil$. In Theorem 4.1, we show that tr α can be computed with an expected $O(n^4(\log n)^2)$ bit operations.

First, we stress that this result is unconditional and matches the heuristic complexity analysis of the SEA algorithm: as discussed above, heuristics beyond the Generalized Riemann Hypothesis are required to prove that the SEA algorithm is asymptotically faster than Schoof's algorithm [23, pp. 2–3]. Next, we observe that under our assumption on α , the input has size $O((\log p)^2)$, so this is actually a quasiquadratic algorithm. Finally, our assumption that the separable degree of α is smooth of size $p^{O(1)}$ is reasonable since the best algorithms for generating endomorphisms of a supersingular elliptic curve will satisfy this assumption [4, 6, 8].

For another work involving computing the trace of an endomorphism of E other than the Frobenius endomorphism, see [16], in which Morain–Scribot–Smith compute point counts on curves E/\mathbb{F}_{p^2} with a d-isogeny $\phi: E \to E^{(p)}$ by computing the trace of $\psi = \pi \circ \phi$ modulo Elkies primes. They observe that for an Elkies prime ℓ such that E has precisely two rational ℓ -isogenies with kernels K_1 and K_2 , each kernel K_i is invariant under ψ . For another related work, see [2], where Couveignes–Morain use the existence of an ℓ^n -isogeny cycle in a modified SEA algorithm to compute the trace of Frobenius modulo ℓ^n (while we are modifying the SEA algorithm to compute the trace of a cycle!).

Outline. In Section 2, we prove that the kernel of every isogeny of a supersingular elliptic curve E over \mathbb{F}_q is defined over an extension of degree at most 3. This result implies we may use Elkies' method to compute the trace of an endomorphism of a supersingular curve modulo ℓ for every odd prime ℓ . In Section 3, we give algorithms for computing the trace of an endomorphism α modulo an odd prime $\ell \neq p$ by restricting to the kernel K of a rational ℓ -isogeny, even when α does not fix K, and analyze its complexity. We discuss how arithmetic with restrictions of endomorphisms to kernels is performed, i.e., how to represent, add together, and test equality of elements in Hom(ker $\phi, E[\ell]$). The results and analysis in this section apply to any elliptic curve over a finite field of characteristic p > 3 with a rational isogeny of degree ℓ . In 3.2 we show how to compute the trace of a separable endomorphism modulo p, the characteristic, using the action of endomorphisms on invariant differentials. In 4 we prove our main Theorem, Theorem 4.1, on the complexity of the SEA algorithm for supersingular endomorphisms. We discuss some timings of our implementation¹ in Section 5 and conclude in Section 6 with a discussion of how the ideas in Section 3.1 might apply to the SEA algorithm for computing the trace of Frobenius modulo Atkin primes, an idea already explored by Dewaghe [5].

Acknowledgements. We thank François Morain who informed us that the idea of applying Elkies' algorithm for Atkin primes ℓ for an ordinary curve E that a rational ℓ -isogeny over a small but nontrivial extension of \mathbb{F}_q is due to Dewaghe [5]. We also thank Andrew Sutherland for helpful discussions, and the

¹https://github.com/travismo/beyond-the-SEA/

Park City Mathematics Institute, where this work began. This material is based upon work supported by the National Science Foundation under Grant No. 2340564. This research was funded in part by the Commonwealth of Virginia's Commonwealth Cyber Initiative (CCI). The first author acknowledges support of the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université), and LabEx CARMIN (ANR-10-LABX-59-01).

2. Background

In this section, we fix notation and cover some of the relevant background on computational aspects of elliptic curves over finite fields.

2.1. Isogenies of elliptic curves. Let p > 3 be a prime, let $q = p^n$, and let E be an elliptic curve defined over \mathbb{F}_q . Throughout, we will assume p > 3, so E is given by a short Weierstrass equation $E: y^2 = x^3 + Ax + B$ where $A, B \in \mathbb{F}_q$ and $4A^3 + 27B^2 \neq 0$. If E' is another elliptic curve, an *isogeny* $E \to E'$ is a non-constant rational map that maps the identity on E to the identity on E'. If ϕ satisfies $\pi^n \circ \phi = \phi \circ \pi^n$, where π is the q-power Frobenius endomorphism, we will say ϕ is an isogeny of E defined over \mathbb{F}_{q^n} . The *degree* of an isogeny ϕ is its degree as a rational map, and ϕ is separable if it is separable as a rational map. If ϕ is separable, then $\deg \phi = \# \ker \phi$. For every isogeny $\phi: E \to E'$, there is a unique *dual* isogeny $\hat{\phi}: E' \to E$ such that $\hat{\phi} \circ \phi = [\deg \phi]$ and $\phi \circ \hat{\phi} = [\deg \phi]$. An *endomorphism* of E is either an isogeny or the constant 0-map [0] from E to E. The degree map is a positive-definite quadratic form on $\operatorname{End}(E)$. The *trace* of an endomorphism $\alpha \in \operatorname{End}(E)$ is the integer tr α such that $\alpha + \hat{\alpha} = [\operatorname{tr} \alpha]$, and the bilinear pairing on $\operatorname{End}(E)$ corresponding to the degree map is $(\alpha, \beta) \mapsto \operatorname{tr} \alpha \hat{\beta}$.

For purposes of calculation and algorithms, we will assume our isogenies are in standard form:

Definition 2.1. Let E and E' be elliptic curves in short Weierstrass form and let E be given by $y^2 = x^3 + Ax + B$. Let $\phi: E \to E'$ be a separable ℓ -isogeny. We say that ϕ is in **standard form** if

$$\phi(x,y) = \left(\frac{u(x)}{v(x)}, cy\frac{s(x)}{t(x)}\right)$$

for polynomials $u, v, s, t \in k[x]$ where

(1) v(x) is the polynomial

$$v(x) = \prod_{\substack{Q \in \ker \phi \\ Q \neq 0}} (x - x_Q),$$

(2) s(x)/t(x) = (u(x)/v(x))'.

By Proposition 4.1 of [1], every separable isogeny may be written in standard form. Suppose E is the curve $y^2 = x^3 + Ax + B$, that ϕ is separable and has degree ℓ , and let $-\sigma$ be the coefficient of $x^{\ell-1}$ in v(x). By Proposition 4.1 of [1], if c = 1 the polynomial u(x) satisfies the relation

$$\frac{u(x)}{v(x)} = \ell x - \sigma - (3x^2 + A)\frac{v'(x)}{v(x)} - 2(x^3 + Ax + B)\left(\frac{v'(x)}{v(x)}\right)'.$$

The kernel polynomial of an isogeny ϕ is the monic polynomial that vanishes precisely once at the collection of x-coordinates of the nonzero points in ker ϕ . If h(x) is the kernel polynomial of $\phi(x)$ and E is given by $y^2 = f(x)$, then v(x) above is

$$v(x) = \frac{h(x)^2}{\gcd(f(x), h(x))}.$$

In particular, when ϕ has odd degree, we have $h(x)^2 = v(x)$. The isogeny ϕ is normalized if $\phi^*\omega' = \omega$ where ω, ω' are the invariant differentials dx/2y on E and E' respectively. Equivalently, ϕ is normalized if the constant c in the standard form of ϕ is 1. A normalized isogeny ϕ is defined over \mathbb{F}_q if and only if ker ϕ is $\operatorname{Gal}(\overline{\mathbb{F}_q})$ -stable, and this holds if and only if $h(x) \in \mathbb{F}_q[x]$. Since v(x) is determined by the kernel of ϕ , if ϕ is normalized the polynomial u(x) (and hence the polynomials s and t) are completely determined by ker ϕ and the equation $y^2 = x^3 + Ax + B$ defining E. Every finite subgroup K of $E(\overline{\mathbb{F}_q})$ determines an isogeny $E \to E/K$; here E/K is the unique elliptic curve E' such that the separable isogeny $E \to E'$ with kernel K is normalized. While we do not need it, there are formulas for the coefficients giving a Weierstrass equation defining E/K from the data E and K; see the discussion following [1, Proposition 4.1] in Section 4.1. The constant c in the standard form results from post-composition of the normalized isogeny $E \to E/K$ with the isomorphism $(x, y) \mapsto (c^{-2}x, c^{-3}y)$. 2.2. Supersingular elliptic curves. An elliptic curve E defined over \mathbb{F}_q is supersingular if its geometric endomorphism algebra is a quaternion algebra. Equivalently, the trace of Frobenius $t_E \coloneqq q + 1 - \# E(\mathbb{F}_q)$ is congruent to 0 modulo p. This restricts the possibilities for how π_E can act on the ℓ -torsion of E. This, in turn, restricts the minimal field of definition of an isogeny between two supersingular elliptic curves. Every supersingular elliptic curve E over \mathbb{F}_{p^2} with $j(E) \notin \{0, 1728\}$ has all its isogenies defined over \mathbb{F}_{p^2} . We'll now show more generally that a supersingular curve over \mathbb{F}_q has all its isogenies defined over an extension of degree at most 3.

Lemma 2.2. Let $q = p^e$ be a power of a prime p > 3 and let E be a supersingular elliptic curve defined over \mathbb{F}_q . Then if $\psi: E \to E'$ is any isogeny of E, its kernel is defined over the extension \mathbb{F}_{q^m} with m = 1, 2, or 3. If $j(E) \notin \{0, 1728\}$, then we can take m = 1 if e is even and m = 2 if e is odd.

Proof. According to [19, Theorem 4.2(ii) and (iii)], the q-power Frobenius π of E satisfies a polynomial equation $\pi^2 - k\sqrt{q}\pi + q = 0$, where k = 0 for odd e and $k \in \{0, \pm 1, \pm 2\}$ for even e. A calculation shows that $\pi^4 - (k^2 - 2)q\pi^2 + q^2 = 0$ and $\pi^6 - (k^2 - 3)kq^{3/2}\pi^3 + q^3 = 0$.

- If k = 0, then $0 = \pi^2 + q$, hence $\pi^2 = -q \in \mathbb{Z}$. Use m = 2. If $k \in \{\pm 1\}$, then $0 = \pi^6 \pm 2q\sqrt{q}\pi^3 + q^3 = (\pi^3 \pm q\sqrt{q})^2$, hence $\pi^3 = \mp q\sqrt{q} \in \mathbb{Z}$. Use m = 3. If $k \in \{\pm 2\}$, then $0 = \pi^4 2q\pi^2 + q^2 = (\pi^2 q)^2$, hence $\pi^2 = q \in \mathbb{Z}$. Use m = 2.

Therefore, for all respective exponents m, we have $\pi^m \in \mathbb{Z}$ and thus $\pi^m(\ker \psi) = \ker \psi$.

Finally, for $j(E) \notin \{0, 1728\}$, set m = 1 if e is even and m = 2 if e is odd. Then $\sqrt{q}^m \in \mathbb{Z}$. Since π^m and $[\sqrt{q}^m]$ are both purely inseparable endomorphisms of degree q^m , there exists an automorphism $\zeta \in \operatorname{Aut}(E)$ such that $\pi^m = \zeta[\sqrt{q}^m]$. But the assumption $j(E) \notin \{0, 1728\}$ implies $\operatorname{Aut}(E) = \{[\pm 1]\}$, so $\pi^m \in \mathbb{Z}$ and thus $\pi^m(\ker \psi) = \ker \psi$. \square

In particular, Lemma 2.2 reveals that every isogeny between supersingular elliptic curves over \mathbb{F}_q is defined over \mathbb{F}_{q^6} : This is the smallest extension of \mathbb{F}_q containing both \mathbb{F}_{q^2} and \mathbb{F}_{q^3} .

2.3. Arithmetic in $\mathbb{F}_q[x]/(h(x))$. Just as in Schoof's algorithm, we will reduce computing the trace of an endomorphism of an elliptic curve E to various algebraic operations over finite fields. For details on the complexity of arithmetic over finite fields, see $[30, \S8, 11]$ and for an introductory overview, see [25,Lecture 3]. Let M(n) denote the number of bit operations required to multiply two *n*-bit integers. Let $q = p^e$ be a power of an odd prime and let $n = \lfloor \log q \rfloor$. Elements of \mathbb{F}_q can be multiplied by lifting to polynomials of $\mathbb{F}_p[x]$ of degree e-1 and then multiplied with Kronecker substitution [30, § 8.4] requiring $O(\mathsf{M}(d(\log p + \log e)))$ bit operations. This cost simplifies to $O(\mathsf{M}(de)) \subseteq O(\mathsf{M}(n))$ under the assumption that $\log e \in O(\log p)$. We can compute an inverse of an element in \mathbb{F}_q^{\times} with $O(\mathsf{M}(n) \log n)$ bit operations using fast Euclidean division [30, §11.1] and Kronecker substitution. Let f, g, h be polynomials in $\mathbb{F}_q[x]$ of degree at most $d \in O(\log n)$ and let $R = \mathbb{F}_q[x]/(h(x))$. The product $fg \mod h$ can be computed with $O(\mathsf{M}(dn))$ bit operations Kronecker substitution [30, Corollary 8.28]. We can test if $f \in \mathbb{R}^{\times}$, i.e., if gcd(f,h) = 1, and if so compute $f^{-1} \mod h$ with $O(\mathsf{M}(dn)\log(dn))$ bit operations using fast Euclidean division and Kronecker substitution. A root of f in \mathbb{F}_q can be found with Rabin's probabilistic algorithm [17] using $O(n\mathsf{M}(dn))$ bit operations, again using Kronecker substitution and our assumption that $\log d \in O(n)$. The bound $\mathsf{M}(n) \in O(n \log n)$ holds by [10]. We summarize this in the following table which will be used in the complexity analysis in Section 3.1.

Proposition 2.3. Let p be an odd prime, let $e \in O(\log p)$ be an integer, let $q = p^e$, and set $n \coloneqq \lceil \log q \rceil$. Let d be a positive integer and assume $d \in O(\log n)$. Let $a, b \in \mathbb{F}_q^{\times}$ and let $f, g, h \in \mathbb{F}_q[x]$ be polynomials of degree at most d. The following table gives the bit complexity of computing products and inverses of elements of $\mathbb{F}_q[x]$ and $\mathbb{F}_q[x]/(h(x))$ as well as root-finding in \mathbb{F}_q .

Operation	bit complexity
ab	$O(n \log n)$
a^{-1}	$O(n(\log n)^2)$
$fg \mod h$	$O(dn\log(dn))$
$f^{-1} \bmod h$	$O(dn(\log(dn))^2)$
f(r) = 0	$O(n^2 \log n)$

3. Computing the trace modulo a prime

Suppose α is an endomorphism of an elliptic curve E defined over \mathbb{F}_q . Schoof's algorithm, originally designed to compute the trace of the Frobenius endomorphism π_E , computes tr α by computing $t_{\ell} :=$ tr $\alpha \mod \ell$ for enough primes $\ell \in O(\log p)$ in order to recover tr α with the Chinese Remainder Theorem. This is possible since $\mathbb{Z}[\alpha]$ is an imaginary quadratic order, so we have the bound $|\operatorname{tr} \alpha| < 2(\deg \alpha)^{1/2}$. Suppose now *E* is supersingular, defined over \mathbb{F}_{p^2} , and that $j(E) \notin \{0, 1728\}$. Then

$$t_E \coloneqq p^2 + 1 - \#E(\mathbb{F}_q) = \pm p$$

is the trace of π_E , and π_E is [p] or [-p]. This implies every isogeny of E is defined over \mathbb{F}_{p^2} (see 2.2); every odd prime is an Elkies prime for E! We prove in Theorem 3.10 that t_{ℓ} may be computed in $O(n^3(\log n)^3)$ time, assuming $\log \deg \alpha \in O(n)$ and α is represented by a sequence of isogenies of degree bounded by a constant. Note that this result is *unconditional*—we do not require GRH since all primes are Elkies primes. The method is discussed in Section 3.1.

Another advantage in our case is that we can actually compute the trace of α modulo p. This does not impact the asymptotic complexity but gives a dramatic speedup in practice. We discuss this in Section 3.2. Another practical speedup is that $\#E(\mathbb{F}_q)$ is easily computable since we assume E is supersingular. Thus whenever ℓ divides $\#E(\mathbb{F}_q)$, we can compute t_{ℓ} working directly with a rational point of order ℓ . Again, this does not provide an asymptotic speedup, but the idea is helpful in practice.

3.1. Computing the trace modulo ℓ . In this section, we consider the computation of t_{ℓ} for an odd prime $\ell \neq p$. We suppose E (not necessarily supersingular) has a rational ℓ -isogeny ψ and compute t_{ℓ} for an endomorphism α . We discuss how to perform arithmetic with restrictions of isogenies to ker ψ , even isogenies which do not fix ker ψ .

3.1.1. Arithmetic in Hom $(E, E)_h$. Suppose E is an elliptic curve and $\psi: E \to E/\ker \psi$ is a separable isogeny. Consider the exact sequence

$$0 \longrightarrow \ker \psi \longrightarrow E \longrightarrow E/\ker \psi \longrightarrow 0.$$

Taking Hom (\cdot, E') (whereby we mean homomorphisms of elliptic curves, i.e., isogenies and the zero map), the isogeny ψ induces a pullback map

$$\psi^* \colon \operatorname{Hom}(E/\ker\psi, E') \to \operatorname{Hom}(E, E')$$

 $\rho \mapsto \rho \circ \psi.$

We also have a map $\operatorname{Hom}(E, E') \to \operatorname{Hom}(\ker \psi, E')$ given by restriction to $\ker \psi$. Because an isogeny $\phi: E \to E'$ is zero on $\ker \psi$ if and only if it factors through ψ , we have an exact sequence

$$0 \longrightarrow \operatorname{Hom}(E/\operatorname{ker} \psi, E') \xrightarrow{\psi^*} \operatorname{Hom}(E, E') \xrightarrow{\cdot \mid_{\operatorname{ker} \psi}} \operatorname{Hom}(\operatorname{ker} \psi, E').$$

Note that $I := \psi^* \operatorname{Hom}(E/\ker \psi, E) = \operatorname{Hom}(E/\ker \psi, E)\psi$ is a left ideal of $\operatorname{End}(E)$, so its image under the restriction to ker ψ is isomorphic the quotient $\operatorname{End}(E)/I$. Suppose for the moment E is ordinary. In the context of Schoof's algorithm, the isogeny ψ is the multiplication-by- ℓ map $[\ell]$ on E, the ideal I is $\ell \operatorname{End}(E)$, and $\operatorname{End}(E[\ell]) = \operatorname{End}(E)/\ell \operatorname{End}(E)$ is a two-dimensional algebra over $\mathbb{Z}/\ell\mathbb{Z}$. The trace of an endomorphism is computed by computing the characteristic equation of its image in $\operatorname{End}(E[\ell])$. Algorithmically, arithmetic in $\operatorname{End}(E[\ell])$ requires polynomial arithmetic in $\mathbb{F}_p[x]$ modulo the ℓ th division *polynomial.* For a prime ℓ such that E admits a rational ℓ -isogeny, a so-called Elkies prime, the isogeny ψ is a rational ℓ -isogeny and $\operatorname{End}(\ker\psi) = \operatorname{End}(E)/I \cong \mathbb{Z}/\ell\mathbb{Z}$. Computing the trace modulo ℓ requires arithmetic in $\mathbb{F}_p[x]/(h(x))$ where h is the kernel polynomial of ψ . One can solve for the eigenvalue λ of π_E restricted to ker ψ and recover the trace as $t_{\ell} = \lambda + q/\lambda \mod \ell$. In both cases, the endomorphism π_E fixes ker ψ and one can interpret the algorithm as performing arithmetic in End(ker ψ). But as far as asymptotics are concerned, the fact that π_E fixes a one dimensional subspace of $E[\ell]$ is irrelevant to the speedup. To be more precise, one could compute t_{ℓ} by computing the characteristic equation of π_E as in Schoof's algorithm with the same time complexity. The crucial fact is that ker ψ is cut out by a degree $O(\ell)$ polynomial, a factor of ℓ smaller than the degree of the division polynomial. This is where the asymptotic speedup comes from in Elkies' improvement to Schoof's algorithm for computing t_{ℓ} .

We now explain how to explicitly calculate with elements of $\operatorname{Hom}(\ker \psi, E')$. Let q be a power of a prime p > 3 and let E and E' be elliptic curves defined over \mathbb{F}_q by $y^2 = f_E(x)$ and $y^2 = f_{E'}(x)$ respectively on affine points. Suppose $\psi \colon E \to E/\ker \psi$ is an \mathbb{F}_q -isogeny of prime degree $\ell \neq p$, so $\ker \psi$ is a subvariety of E cut out by its kernel polynomial h(x); more precisely, $\ker \psi$ is given by

$$\ker \psi = \operatorname{Proj} \frac{\mathbb{F}_q[X, Y, Z]}{(Y^2 Z - F_E(X, Z), H(X, Z))}_{5}$$

where F_E and H are the homogenizations of f_E and h respectively. Note that the affine points of ker ψ are given by Spec R, where

$$R = \frac{\mathbb{F}_q[x, y]}{(y^2 - f_E(x), h(x))} \cong \frac{\left(\mathbb{F}_q[x]/(h(x))\right)[y]}{(y^2 - f_E(x))}$$

Given an \mathbb{F}_q -isogeny $\phi : E \to E'$, the closed embedding $\iota : \ker \psi \hookrightarrow E$ allows us to form the restriction $\phi |_{\ker \psi}$ by the composition $\phi \circ \iota$. Since we assume that $\deg \psi$ is prime, the intersection $\ker \psi \cap \ker \phi$ is either trivial or equal to $\ker \psi$. If $\ker \psi \cap \ker \phi = \{0_E\}$, then $\phi |_{\ker \psi}$ restricts to a morphism of affine varieties from Spec R to the affine points of E; concretely, if ϕ is given in standard form as $\phi(x, y) = \left(\frac{u(x)}{v(x)}, \frac{s(x)}{t(x)}y\right)$, then v and t are invertible in $\mathbb{F}_q[x]/(h(x))$, and $\phi |_{\ker \psi}$ is given on affine points by

$$\phi\big|_{\ker\psi}(x,y) = \left(u(x)v(x)^{-1}, s(x)t(x)^{-1}y\right) \in \mathbb{A}^2(R).$$

In this way, such a ϕ induces an affine *R*-point of E', which we denote by ϕ_h . If ker $\psi \subseteq \ker \phi$, we set $\phi_h = 0_{E'} \in E'(R)$. We define

$$\operatorname{Hom}(E, E')_h \coloneqq \{\phi_h \mid \phi : E \to E' \text{ an isogeny}\} \subseteq E'(R).$$

Note that $\operatorname{Hom}(E, E')_h$ and $\iota^*(\operatorname{Hom}(E, E'))$ are in bijection via the map $\iota^* \phi \mapsto \phi_h$.

The following lemma will be used to get a distinguished representative for non-zero ϕ_h .

Lemma 3.1. Let q be a power of a prime p > 3, let E be an elliptic curve over \mathbb{F}_q , let $\psi: E \to E/\ker \psi$ be a separable \mathbb{F}_q -isogeny of odd prime degree with kernel polynomial $h(x) \in \mathbb{F}_q[x]$. Let $\phi: E \to E'$ be an isogeny defined over \mathbb{F}_q with ker $\phi \cap \ker \psi = \{0\}$.

(a) For any $a, b \in \mathbb{F}_q[x]$, we have that $\phi_h = (a(x), b(x)y) \in \mathbb{A}(R)$ if and only if for all $P = (x_P, y_P) \in \ker \psi \setminus \{0\}$,

$$\phi(P) = (a(x_P), b(x_P)y_P).$$

(b) There exist unique a(x), $b(x) \in \mathbb{F}_q[x]$ with deg $a < \deg h$ and deg $b < \deg h$ satisfying the above.

Proof. Suppose that $a(x), b(x) \in \mathbb{F}_q[x]$ are such that $\phi_h = (a(x), b(x)y)$ (under the map $\mathbb{F}_q[x, y] \to R$). We first show that $\phi(P) = (a(x_P), b(x_P)y_P)$ for any nonzero $P \in \ker \psi$. Write

$$\phi(x,y) = \left(\frac{u(x)}{v(x)}, \frac{s(x)}{t(x)}y\right)$$

Then $a(x)v(x) \equiv u(x) \pmod{h(x)}$ so there exists $g(x) \in \mathbb{F}_q[x]$ such that $a(x)v(x) = u(x) + g_1(x)h(x)$. Since h(x) cuts out ker ψ , for any affine $P \in \ker \psi$ we have $h(x_P) = 0$. Therefore

$$a(x_P)v(x_P) = u(x_P) + h(x_P)g_2(x_P) = u(x_P)$$

Because ker $\phi \cap \ker \psi = \{0\}$, we have $v(x_P) \neq 0$, and we get $a(x_P) = u(x_P)/v(x_P) = x_{\phi(P)}$. Similarly there exists a polynomial $g_2(x)$ such that $b(x)t(x) = s(x) + g_2(x)h(x)$ so for each affine point P of ker ψ we have

$$b(x_P)t(x_P)y_P = (s(x_P) + h(x_P)g_2(x_P))y_P = s(x_P)y_P$$

Using again that $t(x_P) \neq 0$ and $h(x_P) = 0$, we see $b(x_P) = y_{\phi(P)}$. We conclude $(a(x_P), b(x_P)y_P) = \phi(P)$ for each affine $P \in \ker \psi$.

Suppose $a(x), b(x) \in \mathbb{F}_q[x]$ satisfy $(a(x_P), b(x_P)y_P) = \phi(P)$ for all affine $P \in \ker \psi$. Let $\phi_1(x), \phi_2(x) \in \mathbb{F}_q[x]$ satisfy $\phi_h = (\phi_1(x), \phi_2(x)y)$. Then $\phi_1(x) - a(x)$ is zero at x_P for all affine $P \in \ker \psi$ and therefore vanishes at all the roots of h(x) in $\overline{\mathbb{F}_q}$. Since h(x) is separable, we conclude h(x) divides $a(x) - \phi_1(x)$. We also have $(\phi_2(x_P) - b(x_P))y_P = 0$ for all affine $P \in \ker \psi$. Since $\deg \psi$ is an odd prime, we have $\ker \psi \cap E[2] = 0$, so y_P is nonzero for each P. Therefore $\phi_2(x_P) - b(x_P) = 0$ and again we conclude h(x) divides $\phi_2(x) - b(x)$. Thus as elements of R we have $a(x) = \phi_1(x)$ and $b(x) = \phi_2(x)y$.

There is a unique polynomial $a(x) \in \overline{\mathbb{F}_q}[x]$ of degree less than deg h interpolating the deg h many pairs $(x_P, x_{\phi(P)})$. Since ker $\psi \cap E[2] = 0$, for any affine point $P \in \ker \psi$ we have $y_P \neq 0$. Thus there is a unique polynomial $b(x) \in \overline{\mathbb{F}_q}[x]$ of degree less than deg h interpolating the deg h many pairs $(x_P, y_{\phi(P)}y_P^{-1})$. By the first part of the proof, the x and y coordinates of ϕ_h also respectively interpolate the sets of points $\{(x_P, x_{\phi(P)} : P \neq 0 \in \ker \psi\}$ and $\{(x_P, y_{\phi(P)}y_P^{-1})\}$, so the polynomials a(x) and b(x)y give the x and y coordinates of ϕ_h .

The previous lemma allows us make the following definition:

Definition 3.2. Let q be a power of a prime p, let E and E' be elliptic curves over \mathbb{F}_q , let ker $\psi \subseteq E$ be the kernel of a separable isogeny of odd prime degree, and let h be the kernel polynomial of ψ . For $\phi_h \in \text{Hom}(E, E')_h$ non-zero, the *standard form* of ϕ_h is the unique representative of the form (a(x), b(x)y) where $a(x), b(x)y \in \mathbb{F}_q[x, y]$ with deg a and deg b both less than deg h.

Since ker ψ is cyclic of prime order, any point in ker ψ generates the whole group. This allows for interpolation of polynomials in some cases to be done with a single point, and this strategy allows us to show the following:

Lemma 3.3. Let q be a power of a prime p > 3, let E be an elliptic curve over \mathbb{F}_q , let $\psi: E \to E/\ker \psi$ be a separable \mathbb{F}_q -isogeny of odd prime degree with kernel polynomial $h(x) \in \mathbb{F}_q[x]$. Let $\phi, \rho: E \to E'$ be isogenies defined over \mathbb{F}_q with $\ker \phi \cap \ker \psi = \ker \rho \cap \ker \psi = \{0\}$. Let $\phi_h = (a(x), b(x)y)$ be the standard form of ϕ_h and let $\rho_h = (c(x), d(x)y)$ be the standard form of ρ_h .

(a) Either a(x) = c(x) or a(x) - c(x) is invertible in R.

(b) If a(x) = c(x), then either b(x) = d(x) or b(x) = -d(x).

(c) Finally, b(x)y is invertible in R.

Proof.

- (a) We will show that a(x) = c(x) if $a(x_P) = c(x_P)$ for a single nonzero $P \in \ker \psi$. Suppose there exists a nonzero $P \in \ker \psi$ such that $a(x_P) = c(x_P)$. By Lemma 3.1, we have $x(\phi(P)) = x(\rho(P))$. This implies $\phi(P) = \pm \rho(P)$. Since P generates ker ψ , it follows that $x(\phi(Q)) = x(\rho(Q))$ for all nonzero $Q \in \ker \psi$. By the uniqueness part of Lemma 3.1, we have a(x) = c(x). Thus if $a(x) \neq c(x)$, the polynomial a(x) - c(x) does not vanish at any root of h(x). This implies gcd(a(x) - c(x), h(x)) = 1so a(x) - c(x) is invertible in R.
- (b) Assume a(x) = c(x). As above, this implies $\phi(Q) = \pm \rho(Q)$ for all nonzero $Q \in \ker \psi$. Assume $b(x) \neq d(x)$. Again by Lemma 3.1, this implies $b(x_P) \neq d(x_P)$ for some nonzero $P \in \ker \psi$. Then $\phi(P) = -\rho(P)$, so $\phi|_{\ker \psi} = (-\rho)|_{\ker \psi}$. Again using Lemma 3.1, we get b(x) = -d(x).
- (c) Since $\phi_h = (a(x), b(x)y)$ is nonzero, $\phi(P) \neq 0$ for all nonzero $P \in \ker \psi$. Since ψ has odd degree, the nonzero points in $\ker \psi$ have odd order, so $\phi(P)$ also has odd order. In particular, $y(\phi(P)) = b(x_P)y_P \neq 0$, so $b(x_P) \neq 0$. Thus $b(x_P) \neq 0$ for all nonzero $P \in \ker \psi$ so gcd(b(x), h(x)) = 1. Thus b(x) has an inverse $b(x)^{-1}$ in R. Again, the assumption that ψ has odd degree implies $f_{E_1}(x)$ is invertible modulo h(x) as well (since f_{E_1} vanishes precisely at the coordinates of the nonzero 2-torsion of E and hence is nonzero at any root of h). Then the inverse of b(x)y in R is $b(x)^{-1}f_{E_1}(x)^{-1}y$. \Box

Despite the fact that elements of $\operatorname{Hom}(E_1, E_2)_h$ are defined over a ring with zero divisors, our assumption that ker ψ is cyclic of prime order allows for several of the usual operations on points of elliptic curves to be translated to $\operatorname{Hom}(E_1, E_2)_h \subseteq E(R)$. The following proposition checks that the usual algorithms (found in [24, Group Law Algorithm 2.3], for example) are well-defined and satisfy the expected relations. It also shows that we can compute the restriction $(\phi_2 \circ \phi_1)_h$ of a composition of isogenies $\phi_2 \circ \phi_1$ by evaluating ϕ_2 at $(\phi_1)_h$.

Proposition 3.4. Let q be a power of a prime p > 3, let E_1 and E_2 be elliptic curves over \mathbb{F}_q , and let $\psi: E_1 \to E_1 / \ker \psi$ be a separable \mathbb{F}_q -isogeny of odd prime degree with kernel polynomial $h(x) \in \mathbb{F}_q[x]$.

- (a) The usual formulas for the geometric group law on E_2 make the set $\operatorname{Hom}(E_1, E_2)_h$ into a group. The map $\operatorname{Hom}(E_1, E_2) \to \operatorname{Hom}(E_1, E_2)_h$ is a group homomorphism: for $\phi_h, \rho_h \in \operatorname{Hom}(E_1, E_2)_h$, we have that $\phi_h + \rho_h = (\phi + \rho)_h$.
- (b) Let E_3 be another elliptic curve over \mathbb{F}_q , and let $\phi_2 : E_2 \to E_3$ an isogeny with standard form $\phi_2 = (\frac{u}{n}, \frac{s}{2}y)$. The usual formulas for evaluation

$$\phi_2(x,y) = \begin{cases} 0 & \text{if } v(x) = 0; \\ (u(x)v(x)^{-1}, s(x)t(x)^{-1}y) & \text{otherwise.} \end{cases}$$

give a well-defined map from $\text{Hom}(E_1, E_2)_h$ to $\text{Hom}(E_1, E_3)_h$ satisfying $\phi_2((\phi_1)_h) = (\phi_2 \circ \phi_1)_h$ for $(\phi_1)_h \in \text{Hom}(E_1, E_2)_h$. More precisely, to compute $\phi_2((\phi_1)_h)$, we evaluate ϕ_2 at (a(x), b(x)y) and the arithmetic is performed in R.

Proof.

(a) If $\phi_h = (a(x), b(x)y)$ is nonzero, define $-\phi_h$ to be (a(x), -b(x)y). Then Lemma 3.1 implies $-\phi_h = (-\phi)_h$. If either ϕ_h or ρ_h is equal to zero, or if $\phi_h = -\rho_h$, then the sum $\phi_h + \rho_h$ is set accordingly as either ρ_h or ϕ_h or 0 directly, and the sum is well-defined.

Suppose ϕ_h and ρ_h are nonzero with $\phi_h \neq -\rho_h$, and write $\phi_h = (a(x), b(x)y)$ and $\rho_h = (c(x), d(x)y)$ in standard form. Suppose E is given by the equation $y^2 = f_{E_1}(x)$. If a(x) = c(x), then since $\phi_h \neq -\rho_h$ we have $b(x) \neq -d(x)$. In this case, by Lemma 3.3 we get that b(x) = d(x), and that b(x)yis invertible in R. Define $m(x) \coloneqq \frac{1}{2}f'(a(x))f(x)^{-1} \in R$. Otherwise, $a(x) \neq c(x)$ so a(x) - c(x) is invertible in R. In this case, define $m(x) \coloneqq (b(x) - d(x))(a(x) - c(x))^{-1} \in R$. We claim that in either case $m(x_P)y_P$ is the slope of the line through $\phi(P)$ and $\rho(P)$ for all nonzero $P \in \ker \psi$. Indeed, any rational function on E which is regular at the nonzero points of ker ψ determines an element of R. We have evaluation maps at $P \in \ker \psi$ for such functions as well as for elements in R. The claim follows because evaluation of a function on E regular at $P \in \ker \psi$ agrees with the evaluation of its image in R, since $h(x_P) = 0$. The idea is that R is the coordinate ring of ker $\psi - 0$; it is $\mathbb{F}_q[E]$ modulo the ideal of functions vanishing on ker $\psi - \{0\}$ and this ideal is principle, generated by h(x). The quotient map $\mathbb{F}_q[E] \to R$ corresponds to the inclusion ker $\psi - \{0\} \hookrightarrow E - \{0\}$, so the two evaluation maps agree.

The claim implies that if we define $r(x) \coloneqq m(x)f(x) - a(x) - c(x)$ and $s(x) \coloneqq m(x)(r(x) - a(x)) + b(x)$, then

$$(r(x_P), s(x_P)y_P) = (\phi + \rho)(P)$$

holds for all nonzero $P \in \ker \psi$. Then by Lemma 3.1, we get that $(\phi + \rho)_h = \phi_h + \rho_h$.

(b) Write the isogeny $\phi_2(x) = (\frac{u}{v}, \frac{s}{t}y)$ in standard form, and let $\phi_1 \in \text{Hom}(E_1, E_2)$ with $(\phi_1)_h = (a(x), b(x)y)$ in standard form. Since ker ψ is of prime order, we have that either $\phi_1(\ker\psi) \subseteq \ker\phi_2$ or that $\phi_1(\ker\psi) \cap \ker\phi_2 = \{0\}$. In the former case we have that $v(x_Q) = 0$ for all $Q \in \phi_1(\ker(\psi))$; equivalently, $v(x(\phi_1(P))) = 0$ for all $P \in \ker\psi$, so by Lemma 3.1 we have $v(a(x_P)) = 0$ for all nonzero P in ker ψ so v(a(x)) = 0 in R. Therefore the evaluation formula is well-defined as $\phi_2((\phi_1)_h) = 0$. In the latter case, v(x) is nonzero at $a(x_P)$ for all $P \in \ker\psi$, so v(a(x)) is nonzero at x_P for all $P \in \ker\psi$. Again, this implies v(a(x)) is invertible in R, and again the evaluation formula is well-defined. The same argument above together with part (a) of Lemma 3.1 imply the equality $\phi_2((\phi_1)_h) = (\phi_2 \circ \phi_1)_h$ holds, since the coordinates of $\phi_2((\phi_1)_h)$ interpolate the values of $\phi_2 \circ \phi_1$ at the nonzero points in ker ψ .

The following proposition shows that the remainder modulo ℓ of the trace of an endomorphism of E can be determined using arithmetic in $\text{Hom}(E, E)_h$.

Proposition 3.5. Let p > 3 be prime, let E be an elliptic curve over \mathbb{F}_q , let ker $\psi \subseteq E[\ell]$ be the kernel of an ℓ -isogeny with kernel polynomial h(x), and let $\alpha \in End(E)$. If α_h is nonzero and the integer $0 \leq t \leq \ell - 1$ satisfies

$$(\alpha^2)_h + [\deg \alpha]_h = t\alpha_h$$

then tr $\alpha \equiv t \pmod{\ell}$. If ψ_1 and ψ_2 are two ℓ -isogenies of E with distinct kernel polynomials h_1 and h_2 such that α_{h_1} and α_{h_2} are zero, then tr $\alpha \equiv 0 \pmod{\ell}$.

Proof. First suppose α_h is nonzero. Then for any nonzero point $P \in \ker \psi$ we have $\alpha(P) \neq 0$. Assume that $(\alpha^2)_h + [\deg \alpha]_h = t\alpha_h$. Proposition 3.1 implies $(\alpha^2 + [\deg \alpha])(P) = t\alpha(P)$ for any nonzero $P \in \ker \psi$. This equation also holds when replacing t with tr α , so subtracting the two equations yields $0 = (\operatorname{tr} \alpha - t)\alpha(P)$. Since $\alpha(P) \neq 0$ and P has order ℓ , the point $\alpha(P)$ has order ℓ too. Therefore tr $\alpha \equiv t \pmod{\ell}$.

If α_{h_i} is zero for two distinct kernel polynomials h_1, h_2 then $\alpha(E[\ell]) = 0$ because the ℓ -torsion is generated by the corresponding kernels and α is zero on the generators. This implies there exists $\beta \in \text{End}(E)$ such that $\alpha = \ell\beta$, so

$$\operatorname{tr} \alpha = \operatorname{tr}(\ell\beta) = \ell \operatorname{tr} \beta \equiv 0 \pmod{\ell}.$$

Proposition 3.4 also gives a concrete method for computing with restrictions, and yields the following complexities:

Corollary 3.6. Let q be a power of a prime p > 3, let E_1 and E_2 be elliptic curves over \mathbb{F}_q , and let $\psi: E_1 \to E_1 / \ker \psi$ be a separable \mathbb{F}_q -isogeny of odd prime degree ℓ with kernel polynomial $h(x) \in \mathbb{F}_q[x]$. Let $n = \lceil \log q \rceil$ and assume $\log \ell \in O(n)$.

- (a) Given $\phi_h, \rho_h \in \text{Hom}(E_1, E_2)_h$ in standard form, the standard form of $(\phi + \rho)_h$ can be computed with $O(\ell n(\log \ell)^2)$ bit operations. When $\ell \in O(n)$, this simplifies to $O(n^2(\log n)^2)$.
- (b) Given $\phi_h \in \text{Hom}(E_1, E_2)_h$ in standard form and an integer $1 \le c < \ell 1$, the standard form of $c\phi_h$ can be computed with $O(\ell n (\log \ell)^2)$ bit operations. When $\ell \in O(n)$, this simplifies to $O(n^2 (\log n)^2)$.

(c) Let E_3 be another elliptic curve over \mathbb{F}_q . Given an isogeny $\phi_2 \colon E_2 \to E_3$ in standard form of degree d, and $(\phi_1)_h \in \operatorname{Hom}(E_1, E_2)_h$ in standard form, one may compute the standard form of $(\phi_2 \circ \phi_1)_h$ in $O(d\ell n \log(\ell) + \ell n (\log \ell)^2)$ bit operations. When $\ell \in O(n)$, this simplifies to $O(dn^2 \log n + n^2 (\log n)^2)$.

Proof. By Proposition 3.4 part (a), we can compute $\alpha_h + \rho_h$ and $c\alpha_h$ using the group law applied to R-points on E. Two points in E(R) can be added in $O(\mathsf{M}(\ell n) \log \ell)$ bit operations, the cost of an inversion in R. We can compute $c\alpha_h$ in the same time with the usual double-and-add approach, using projective coordinates so that at most one inversion is required. We can compute $(\alpha_2 \circ \alpha_1)_h$ by evaluating ϕ_2 at $(\phi_1)_h$ by Proposition 3.4 part (b); this can be done using naive modular composition with O(d) multiplications and O(1) inversions in R for a total cost of $O(d\ell n \log(\ell) + \ell n (\log \ell)^2)$ bit operations. \Box

We will also need to compute compositions in longer chains:

Corollary 3.7. Let q be a power of a prime p, and let E_1, \ldots, E_{L+1} elliptic curves over \mathbb{F}_q with $E = E_1 = E_{L+1}$ connected by isogenies $\phi_i : E_i \to E_{i+1}$ for $i = 1, \ldots, L$. Let $\psi : E_1 \to E_1 / \ker \psi$ be a separable \mathbb{F}_q -isogeny of odd prime degree ℓ with kernel polynomial $h(x) \in \mathbb{F}_q[x]$.

Suppose that the ϕ_i are all given in standard form, let d be an upper bound on the degree of all ϕ_i , and assume $\log \ell \in O(\log p)$. Let $n = \lceil \log q \rceil$. The standard form of the restriction $(\phi_L \circ \cdots \circ \phi_1)_h \in$ $\operatorname{Hom}(E, E)_h$ can be computed in

$$O(dL\ell n \log \ell + \ell n (\log \ell)^2)$$

bit operations. In the case that $d \in O(1)$ and $L, \ell \in O(n)$, the total complexity is $O(n^3 \log n)$.

Proof. Write each ϕ_i as $\phi_i(x, y) = \left(\frac{u_i(x)}{v_i(x)}, \frac{s_i(x)}{t_i(x)}y\right)$ in standard form, and suppose we have already computed $(\phi_{n-1} \circ \cdots \circ \phi_1)_h$ for some $n = 1, \ldots, e$ in the form

$$\left(\frac{a(x)}{b(x)}, \frac{c(x)}{d(x)}y\right)$$

for some $u, v, s, t \in \mathbb{F}_q[x]$ reduced modulo h(x), where we take the n = 1 case to be $(\mathrm{id}_{E_1})_h = (x, y)$. Note that this is not the standard form of this restriction, but would become so after computing the divisions modulo h(x). Let $d_i = \deg \phi_i = \deg u_i = \deg v_i + 1$. Since $s_i/t_i = (u_i/v_i)'$, we have $\deg s_i = \deg t_i$. We then have by Proposition 3.4(b) that

$$\begin{aligned} (\phi_n \circ \dots \circ \phi_1)_h &= \left(\frac{u_n(\frac{a(x)}{b(x)})}{v_n(\frac{a(x)}{b(x)})}, \ \frac{s_n(\frac{a(x)}{b(x)})}{t_n(\frac{a(x)}{b(x)})} \frac{c(x)}{d(x)}y \right) \\ &= \left(\frac{U_n(a(x), b(x))}{b(x)V_n(a(x), b(x))}, \ \frac{S_n(a(x), b(x))}{T_n(a(x), b(x))} \frac{c(x)}{d(x)}y \right), \end{aligned}$$

where the polynomials $U_n, V_n, S_n, T_n \in \mathbb{F}_q[X, Z]$ denote the homogenizations of u_n, v_n, s_n and t_n respectively Note that we have degree bounds

$$\deg U_n, \deg V_n, \deg S_n, \deg T_n \leq d$$

and

$$\deg a, \deg b, \deg c, \deg d \le \deg h \in O(\ell),$$

and so $-d \leq r_x, r_y \leq d$. The polynomial compositions and multiplications shown above may thus be computed modulo modulo h(x) in $O(d\mathsf{M}(\ell \log p))$ bit operations. Applying this iteratively, we find that computing $(\phi_L \circ \cdots \circ \phi_1)_h$ as $(\frac{a(x)}{b(x)}, \frac{c(x)}{d(x)}y)$ takes $O(dL\mathsf{M}(\ell \log p))$ bit operations, and performing the two inversions modulo h(x) to arrive at the standard form takes an additional $O(\mathsf{M}(\ell \log p) \log \ell)$ bit operations, yielding the stated complexity.

The method outlined in Corollary 3.7 gives an asymptotic time savings over computing the standard forms of the restrictions of the intermediate compositions when $d \in O(1)$ and $L \in O(n)$. Essentially, we are working with projective coordinates, allowing us to compute the restriction of the chain of isogenies with just 2 inversions in R instead of 2L inversions in R.

Our algorithm for computing the trace modulo ℓ requires the kernel polynomial of an ℓ -isogeny of E, just as in the SEA algorithm. The idea is to first compute the instantiated modular polynomial $\Phi_{\ell}(j(E), Y)$ with Sutherland's algorithm [27], find a root j' of $\Phi_{\ell}(j(E), Y)$ to get the j-invariant of a curve E' which is ℓ -isogenous to E, and then computing the corresponding ℓ -isogeny using Elkies' algorithm [7]. **Theorem 3.8** (Sutherland [27]). Assume the generalized Riemann Hypothesis. Let E be an elliptic curve over \mathbb{F}_q and let $n = \lceil \log q \rceil$. The instantiated modular polynomial $\Phi_{\ell}(j(E), Y)$ can be computed in expected $O(\ell^2 B(\log B)^2)$ time and $O(\ell n + \ell^2 \log B)$ space, where $B \in O(\ell \log \ell + n)$. When $\ell \in O(n)$, the expected runtime is $O(n^3(\log n)^3)$ and the space complexity is $O(n^2 \log n)$.

Proof. This result is due to Sutherland [27, Theorem 4]. The algorithm for computing $\Phi_{\ell}(j(E), Y)$ with the stated complexity is [27, Algorithm 1]. An inspection of the proof of [27, Theorem 4] shows that the stated runtime of $O(\ell^2 B(\log B)^2 \log B)$ can be replaced with $O(\ell^2 B(\log B)^2)$ after the replacement of Schönhage and Strassen's bound $\mathsf{M}(n) \in O(n \log n \log n)$ [18] with Harvey and van der Hoeven's result $\mathsf{M}(n) \in O(n \log n)$ [10].

We also will make use of Elkies' algorithm for computing the kernel polynomial of an ℓ -isogeny. This algorithm uses the modular polynomial $\Phi_{\ell}(X, Y)$; given a nonsingular zero (j, j'), the algorithm first computes models E and E' with j-invariants j and j' and then computes the kernel polynomial. The algorithm fails when (j, j') is a singular point (but see [15] for an extension in this case). For our purposes, we will just prove that if p is sufficiently large relative to ℓ and if j = j(E) is not 0 or 1728, then there is at most one singular point of the form (j(E), j') of multiplicity at most 2.

Lemma 3.9. Let E be a supersingular elliptic curve defined over \mathbb{F}_{p^2} and assume $j(E) \notin \{0, 1728\}$. Let $\ell < (p/4)^{1/4}$ be a prime. Then there is at most one root \tilde{j} of $\Phi_{\ell}(j, Y)$ such that the multiplicity of \tilde{j} is larger than 1, and its multiplicity is at most 2.

Proof. Assume $\phi_{\ell}(j(E), Y)$ has a multiple root j of multiplicity m. Then there are m cyclic ℓ -isogenies $\phi_i \colon E \to E'$ with $j(E') = j', 1 \leq i \leq m$, and with distinct kernels. Any pair of distinct ℓ -isogenies $\phi_i \neq \phi_j$ gives rise to a ℓ^2 -endomorphism $\alpha_{ij} \coloneqq \widehat{\phi_j}\phi_i$ of E. Since we assume ϕ_i and ϕ_j have distinct kernels and that $j(E) \notin \{0, 1728\}$, we have that $\phi_j \neq u\phi_i v$ for any automorphisms $u \in \operatorname{Aut}(E')$ and $v \in \operatorname{Aut}(E) = \{[\pm 1]\}$. Thus α_{ij} is a cyclic ℓ^2 -endomorphism of E. The discriminant D_{ij} of α_{ij} satisfies $-4\ell^2 \leq D_{ij} < 0$, so by [11, Theorem 2'] and our hypothesis that $\ell < (p/4)^{1/4}$, any two α_{ij} must commute and therefore be integral elements of norm ℓ^2 all contained in the same imaginary quadratic field. We then are left with counting the maximal number of elements of norm ℓ^2 in a single quadratic imaginary order contained in $\operatorname{End}(E)$. Let $\alpha = \alpha_{12}$ and let $K = \mathbb{Q}(\alpha)$ and let \mathcal{O}_K be the ring of integers of K.

The order $\mathbb{Z}[\alpha]$ is maximal at ℓ , i.e., the conductor of $\mathbb{Z}[\alpha]$ is coprime to ℓ , because disc α is coprime to ℓ . Indeed, if $\ell \mid \text{disc } \alpha$ then $\ell \mid \text{tr } \alpha$ since deg $\alpha = \ell^2$. This implies $\alpha^2 = \ell(\alpha + \ell)$, so $\alpha^2(E[\ell]) = 0$. This would imply $\phi_2 \phi_1 = u\ell$ for some $u \in \text{Aut}(E')$, which would imply

$$\phi_1\widehat{\phi_2} = \ell u = \phi_1 \circ \widehat{\phi_1} \circ u.$$

This would imply $\phi_2 = u^{-1} \circ \phi_1$, contradicting the assumption that ϕ_1 and ϕ_2 have distinct kernels. Let $\mathcal{O} = \mathbb{Z}[\alpha]$.

We claim that the prime ℓ splits in \mathcal{O} . Since the conductor of \mathcal{O} is coprime to ℓ , the ideal $\ell \mathcal{O}$ factors uniquely into a product of primes of \mathcal{O} coprime to the conductor of \mathcal{O} , and the primes \mathfrak{p} appearing in this factorization are in bijection with the primes $\mathfrak{p}\mathcal{O}_K$ appearing in the factorization of $\ell \mathcal{O}_K$. Let \mathfrak{l} be a prime above ℓ in \mathcal{O} . If ℓ ramifies in \mathcal{O}_K then since $\operatorname{nrd} \alpha = \ell^2$ we have

$$\alpha \mathcal{O}_K = \mathfrak{l}^2 \mathcal{O}_K = \ell \mathcal{O}_K$$

implying

$$\alpha \mathcal{O} = \mathfrak{l}^2 = \ell \mathcal{O}$$

which shows $\alpha = u\ell$ for some $u \in \mathcal{O}^{\times} \subseteq \operatorname{Aut}(E) = \{ [\pm 1] \}$, contradicting the fact that ker α is cyclic. Similarly ℓ is not inert, i.e., $\mathfrak{l} \neq \ell \mathcal{O}$, because again this would imply $\alpha \mathcal{O} = \ell^2 \mathcal{O}$ which would imply α does not have cyclic kernel. So ℓ must split in \mathcal{O} as $\ell \mathcal{O} = \mathfrak{l}$, and $\alpha \mathcal{O} = \mathfrak{l}^2$.

The only ideals of norm ℓ^2 are $\ell \mathcal{O}, \mathfrak{l}^2, \overline{\mathfrak{l}}^2$, hence the only elements of norm ℓ^2 are $\pm \ell, \pm \alpha, \pm \widehat{\alpha}$. Thus there exist at most two cyclic ℓ -isogenies of E which extend to cyclic endomorphisms of E of degree ℓ^2 .

We are now ready to prove the main theorem of this section on the complexity of computing the trace reduced modulo ℓ of a supersingular endomorphism.

Theorem 3.10. Let $\alpha = \phi_L \circ \cdots \circ \phi_1$ be an endomorphism of a supersingular elliptic curve E defined over \mathbb{F}_q , let $n = \lceil \log p \rceil$, and let $\ell \in O(n)$ be an odd prime Let $d = \max\{\deg \phi_i\}$. Then $t_\ell := \operatorname{tr} \alpha \mod \ell$ can be computed in an expected $O(n^3(\log n)^3 + dLn^2 \log n)$ bit operations. Assuming GRH, the space complexity is $O(\ell^3 \log \ell + dLn + \ell n)$. The time and (conditional) space complexities simplify to $O(n^3(\log n)^3)$ and $O(n^3 \log n)$, respectively, when $d \in O(1)$ and $L, \ell \in O(n)$. *Proof.* All isogenies of E will be defined over an extension of degree either 1, 2, or 3 of \mathbb{F}_q , and the cost of arithmetic operations in this extension increase only by a constant factor. We will proceed with assuming E already has all its isogenies defined over \mathbb{F}_q .

To compute a kernel polynomial for an ℓ -isogeny ψ of E, we first compute $\Phi_{\ell}(X,Y)$ a cost of $O(\ell^3(\log \ell)^3)$ bit operations; this can be done using the algorithm of Kunzweiler and Robert [13, Theorem 5.2, Remark 5.3]. With Φ_{ℓ} we can compute $\varphi_{\ell}(Y) \coloneqq \Phi_{\ell}(j(E), Y)$ by evaluating at X = j(E)in time $O(\ell^2 M(\ell \log \ell + n)) \subseteq O(n^3 \log n)$. Alternatively, one could first compute the instantiated modular polynomial $\Phi(Y) := \Phi_{\ell}(j(E), Y)$, along with the partial derivatives $\Phi_X(Y)$ and $\Phi_{XX}(Y)$, in $O((\log q)^3(\log q)^3)$ expected time and $O(\ell^3 \log \ell) \subseteq O(n^3 \log n)$ space under GRH [27, Algorithm 1, Theorem 4]. Next, compute a kernel polynomial h_1 . To do this, compute a random root $j_1 \in \mathbb{F}_{p^2}$ of $\Phi_{\ell}(j(E), Y)$ in expected $O(\ell n^2 \log(\ell n)) \subseteq O(n^3 \log n)$ time. The root j_1 that is necessarily the *j*-invariant of a supersingular elliptic curve E_1 with the property that there exists an ℓ -isogeny $\psi: E \to E_1$, and this isogeny is defined over \mathbb{F}_{p^2} since E is supersingular. Assuming for the moment $j(E_1)$ is a simple (i.e., multiplicity 1) root of $\Phi(Y)$, we proceed to compute h_1 by first computing a model for the curve E_1 such that the isogeny $\psi_1: E \to E_1$ is normalized, meaning $\phi^* \omega_1 = \omega$ where ω, ω_1 are the invariant differentials dx/2y of E and E_1 respectively. This is done via Elkies' algorithm [7]; see also [20, Section 7] and [9, Chapter 25, Algorithm 28] for details. The model E_1 can be computed with $O(\ell)$ operations in \mathbb{F}_{p^2} . The isogeny $\psi: E \to E'$ can then be computed in $O(\mathsf{M}(\ell) \log \ell)$ operations in \mathbb{F}_{p^2} with the algorithm of Bostan, Morain, Salvy and Schost [1, Theorem 1]. If j_1 is not a simple root of $\Phi(Y)$, we compute random roots of $\Phi(Y)/(Y-j_1)^e$, where e is the multiplicity of j_1 as a root of $\Phi(Y)$, until finding a simple root. Assuming $\ell \leq p/4$, there are at most 2 roots of $\Phi(Y)$ which are not simple roots, counted with multiplicity, by Proposition 3.9.

We then compute α_{h_1} , requiring $O(dL\ell n \log \ell + \ell n (\log \ell)^2)$ bit operations. If $\alpha_{h_1} = 0$, we repeat the above, computing a second kernel polynomial h_2 and the restriction α_{h_2} . If $\alpha_{h_2} = 0$ a second time, then tr $\alpha \equiv 0 \pmod{\ell}$ by Proposition 3.5. So suppose $\alpha_h \neq 0$ for $h = h_1$ or h_2 . We can then compute $(\alpha^2)_h$, $[\deg \alpha]_h$, and $(\alpha^2)_h + [\deg \alpha]_h$. The cost is dominated by the $O(de\ell n \log \ell + \ell n (\log \ell)^2)$ bit operations required to compute α_h . If $(\alpha^2)_h + [\deg \alpha]_h = 0$ we return 0; otherwise we compute $c\alpha_h$ for increasing $c = 1, 2, \ldots, \ell - 1$ until $c\alpha_h = (\alpha^2)_h + [\deg \alpha]_h$, requiring $O(\ell^2 n (\log \ell)^2)$ operations by Corollary 3.6(a). The algorithm is correct by Proposition 3.5.

3.2. Computing the trace modulo p. In this section, we give a fast algorithm for computing the trace modulo p of a separable endomorphism of an elliptic curve over a field of characteristic p. If $E \cong \mathbb{C}/\Lambda$ is an elliptic curve over \mathbb{C} , then $\alpha \in \text{End}(E)$ lifts to a map $\mathbb{C} \to \mathbb{C}$ given by multiplication by a complex number a. The complex number $a \in \mathbb{C}$ satisfies deg $\alpha = a\overline{a}$ and tr $\alpha = a + \overline{a}$, where $\overline{}$ is complex conjugation. The element a is the scaling factor in \mathbb{C} such that $\alpha^*\omega = a\omega$, where ω is any invariant differential on E. Thus tr α may be computed by computing the trace of the scaling factor of α^2 . We'll adapt this idea to the case that α is a separable endomorphism of a supersingular elliptic curve.

Let E be a supersingular elliptic curve defined over \mathbb{F}_{p^2} and let ω be an invariant differential on E. We have a ring homomorphism

$$\operatorname{End}(E) \to \mathbb{F}_{p^2}$$

 $\alpha \mapsto a_{\alpha},$

where $a_{\alpha} \in \mathbb{F}_{p^2}$ satisfies

$$\alpha^*\omega = a_\alpha\omega.$$

Proposition 3.11. Let q be a power of a prime p and let E be an elliptic curve over \mathbb{F}_q . Suppose $\alpha \in \text{End}(E)$. Then modulo p we have

$$\operatorname{tr} \alpha \equiv \begin{cases} a_{\alpha} + a_{\alpha}^{-1} \operatorname{deg} \alpha & \text{if } \alpha \text{ is separable;} \\ a_{\widehat{\alpha}} & \text{if } \alpha \text{ is inseparable.} \end{cases}$$

Suppose now that E is supersingular. Then the above simplifies to

$$\operatorname{r} \alpha \equiv \operatorname{Tr}_{\mathbb{F}_{p^2}/\mathbb{F}_p}(a_\alpha) \pmod{p},$$

where $\operatorname{Tr}_{\mathbb{F}_{n^2}/\mathbb{F}_p} a = a + a^p$ is the field trace of the extension \mathbb{F}_{p^2} of \mathbb{F}_p .

Proof. As stated above, the map $\alpha \mapsto a_{\alpha}$ is a ring homomorphism. The endomorphism α satisfies its characteristic polynomial $\chi_{\alpha}(x) := x^2 - (\operatorname{tr} \alpha)x + \operatorname{deg} \alpha$, so

$$a_{\alpha}^2 - (\operatorname{tr} \alpha)a_{\alpha} + \deg \alpha = 0$$

²The idea to compute tr α from the action of α on an invariant differential appears to be known to experts; see [3].

holds as an identity in \mathbb{F}_q . Similarly, $a_{\widehat{\alpha}}$ is also a root of χ_{α} in \mathbb{F}_q . The kernel of the map $\alpha \mapsto a_{\alpha}$ consists precisely of the inseparable endomorphisms of E [24, Theorem III.5.5(b)], so $a_{\alpha} = 0$ for all inseparable endomorphisms. If α is inseparable, the polynomial $\chi_{\alpha} \mod p$ reduces to $x(x - \operatorname{tr} \alpha) \mod p$. Since $a_{\alpha} = 0$ is one root of χ_{α} in \mathbb{F}_q , the other root is $a_{\widehat{\alpha}} = \operatorname{tr} \alpha$. If α is separable, then $a_{\alpha} \neq 0 \in \mathbb{F}_q$. Since the map $\alpha \mapsto a_{\alpha}$ is a ring homomorphism, we have

$$a_{\alpha}a_{\widehat{\alpha}} = a_{\deg\alpha} = \deg\alpha$$

as an identity in \mathbb{F}_q , from which the congruence in the proposition follows.

Now assume that E is supersingular. We will show

$$\operatorname{tr} \alpha \equiv \operatorname{Tr}_{\mathbb{F}_{p^2}/\mathbb{F}_p}(a_\alpha) \pmod{p}.$$

That $a_{\alpha} \in \mathbb{F}_{p^2}$ is clear since a_{α} satisfies a quadratic polynomial with coefficients in \mathbb{F}_p . If $\alpha = m \in \mathbb{Z}$, then $\widehat{m} = m$ and $[m]^* \omega = m \omega$, as ω is an invariant differential. Thus

$$a_{\widehat{m}} = a_m = m \equiv m^p \pmod{p}$$
.

Now assume that $\alpha \notin \mathbb{Z}$. Then the prime p cannot split in $\mathbb{Z}[\alpha]$, so the polynomial χ_{α} has either zero or one roots in \mathbb{F}_p . Suppose first that χ_{α} has no roots in \mathbb{F}_p , i.e., p is inert in $\mathbb{Z}[\alpha]$. Then $\chi_{\alpha} \mod p$ is the minimal polynomial of a_{α} , so

$$\operatorname{tr} \alpha \equiv \operatorname{Tr}_{\mathbb{F}_2/\mathbb{F}_p} a_\alpha \pmod{p}$$

as desired. If instead χ_{α} has exactly one root $a_{\alpha} = a_{\widehat{\alpha}} = m \in \mathbb{F}_p$, we conclude

$$\operatorname{tr} \alpha \equiv a_{\operatorname{tr} \alpha} \equiv a_{\alpha} + a_{\widehat{\alpha}} \equiv 2m \equiv m + m^p \equiv \operatorname{Tr}_{\mathbb{F}_p^2/\mathbb{F}_p} a_{\alpha} \pmod{p}.$$

Remark 3.12. It is not true that the reduced trace of α is congruent modulo p to the field trace of a_{α} when α is an endomorphism of an ordinary elliptic curve E. For example, let $\alpha = \pi_E$ be the Frobenius endomorphism of E. Then $a_{\pi_E} = 0$ and thus has trace zero but tr $\alpha \not\equiv 0 \pmod{p}$ since E is ordinary.

Remark 3.13. When E is supersingular, Proposition 3.11 shows the map $\alpha \mapsto a_{\alpha}$ is a homomorphism of algebras with involution, where the involution on $\operatorname{End}(E)$ is the dual map and the involution on \mathbb{F}_{p^2} is the *p*-power Frobenius automorphism, i.e., the generator of $\operatorname{Gal}(\mathbb{F}_{p^2}/\mathbb{F}_p)$. More precisely, we claim that $a_{\widehat{\alpha}} = a_{\alpha}^p$. This can be seen directly via the Deuring Lifting Theorem [14, Theorem 14]: given the endomorphism $\alpha \in \operatorname{End}(E)$, there is a number field L, an elliptic curve A over L with endomorphism ϕ and a prime \mathfrak{P} of L above p such that A and ϕ reduce modulo \mathfrak{P} to E and α , respectively. There is an isomorphism $\operatorname{End}(A) \to \mathcal{O}$ where \mathcal{O} is an imaginary quadratic order given by $\theta \mapsto a_{\theta}$. This isomorphism respects the involutions on $\operatorname{End}(A)$ and \mathcal{O} : we have $a_{\widehat{\theta}} = \overline{a_{\theta}}$. Let $\mathfrak{p} = \mathfrak{P} \cap \mathcal{O}$. Suppose the prime p is inert in \mathcal{O} , so the Frobenius element at p corresponds to complex conjugation. This implies

$$a_{\widehat{\theta}} = \overline{a_{\theta}} \equiv a_{\theta}^p \pmod{\mathfrak{p}}$$

We also have $a_{\phi} \equiv a_{\alpha} \pmod{\mathfrak{p}}$ since ϕ reduces to α modulo \mathfrak{P} . Together, the congruences imply $a_{\widehat{\alpha}} \equiv a_{\alpha}^p \pmod{\mathfrak{p}}$ as claimed. If p is ramified in \mathcal{O} , the same conclusion holds, since

$$a_{\widehat{\theta}} = \overline{a_{\theta}} \equiv a_{\theta} \equiv a_{\theta}^p \pmod{\mathfrak{p}}.$$

When α is inseparable, so too is $\hat{\alpha}$ since E is supersingular. Then $a_{\alpha} = a_{\hat{\alpha}} = 0$, so the claim holds in this case as well.

Proposition 3.11 suggests a simple approach to computing the trace of a separable endomorphism α modulo p: compute the action of α on the space of invariant differentials. In particular, compute $a_{\alpha} \in \mathbb{F}_{p^2}$ and then compute $\operatorname{Tr}_{\mathbb{F}_{p^2}/\mathbb{F}_p} a_{\alpha} = a_{\alpha} + a_{\alpha}^p$. Recall that we assume α is represented by a sequence of isogenies, which in turn are represented by rational maps. If $\alpha = \phi_L \circ \cdots \circ \phi_1$ where $\phi_i \colon E_i \to E_{i+1}$ are isogenies, and if we choose the standard invariant differential $\omega_i = dx/2y$ on each E_i , then $\phi_i^* \omega_{i+1} = c_i \omega_i$ for some $c_i \in \mathbb{F}_{p^2}$ for $i = 1, \ldots, L - 1$. Then

$$a_{\alpha} = \prod_{i=1}^{L} c_i.$$

Thus to compute a_{α} we must compute c_i .

Lemma 3.14. Let E and E' be elliptic curves in short Weierstrass form and let E be given by $y^2 = x^3 + Ax + B$. Let $\phi: E \to E'$ be an ℓ -isogeny in standard form $\phi(x, y) = (u/v, cys/t)$. Then the leading coefficients of u(x) and s(x) are both equal to c^2 . In particular, if c_x and c_y are the leading coefficients of u(x) and cs(x), respectively, then $c = c_x^{-1}c_y$.

Proof. The proof involves simply inspecting the arguments in [29] and [1, Proposition 4.1]. First assume that ϕ is normalized, i.e., c = 1. We claim that the leading coefficient of u(x) is 1. Let $F = \ker \phi$ and let $F^+ \subseteq F$ contain precisely one of Q or -Q for all nonzero $Q \in F$. Vèlu's formulas [29] express u(x)/v(x) as

$$I_x(x) \coloneqq x + \sum_{Q \in F^+} \left(\frac{t_Q}{x - x_Q} + 4 \frac{x_Q^3 + Ax_Q + B}{(x - x_Q)^2} \right),$$

where $t_Q = 3x_Q^2 + A$ if Q has order 2 and otherwise $t_Q = 2(3x_Q^2 + A)$. When written as a rational function, the denominator of I_x is v(x). Note when written with common denominator v(x), the terms $\frac{t_Q}{x-x_Q}$ and $4\frac{x_Q^3 + Ax_Q + B}{(x-x_Q)^2}$ will have numerators of degree $\ell - 2$ and $\ell - 1$, respectively, since the degree of v(x) is $\ell - 1$. Thus the top degree term in the numerator of I_x will be $x \cdot x^{\ell-1} = x^{\ell}$, so we conclude u(x) is monic.

Since $I_y(x) = (u(x)/v(x))'$, and since u and v are monic of respective degrees ℓ and $\ell-1$, the numerator of $I_y(x) = (u(x)/v(x))'$ has leading term equal to

$$x^{\ell-1} \cdot \ell x^{\ell-1} - x^{\ell} \cdot (\ell-1) x^{\ell-2} = (\ell - (\ell-1)) x^{2\ell-2} = x^{2\ell-2}.$$

This settles the case when ϕ is normalized.

Suppose now $\phi: E \to E'$ defined by $(x, y) \mapsto (u(x)/v(x), c(u(x)/v(x))'y)$, where c is arbitrary. Consider the isomorphism $\iota: E' \to E''$ given by $(x, y) \mapsto (c^{-2}x, c^{-3}y)$. Then $\iota^* \omega_{E''} = c \omega_{E'}$. Thus $\phi' = \iota \circ \phi$ is normalized:

$$(\iota\phi)^*\omega_{E''} = \phi^*\iota^*\omega_{E''} = \phi^*c\omega_{E'} = c \cdot c^{-1}\omega_E = \omega_E$$

The coordinate functions of ϕ' are

$$\bigg(\frac{c^{-2}u(x)}{v(x)}, c^{-3}y\bigg(\frac{u(x)}{v(x)}\bigg)'\bigg).$$

The leading coefficient of $c^{-2}N(x)$ must be 1, so the leading coefficient of u(x) must be c^2 . Similarly the leading coefficient of the numerator of $c^{-3}(u(x)/v(x))'$ must be 1 (and the denominator is clearly monic), so the leading coefficient of the numerator of (u(x)/v(x))' must be c^3 .

Lemma 3.14 implies that, given an isogeny $\phi(x, y) = (I_x, cyI_y)$ in standard form, we may calculate c by finding the leading coefficients c_x and c_y of the numerators of I_x and cI_y and then calculating $c = c_x^{-1}c_y$.

Proposition 3.15. Let p > 3 be a prime and let E be a supersingular elliptic curve defined over \mathbb{F}_q . Let $\alpha \in \operatorname{End}(E)$ be an endomorphism. Let $\alpha = \phi_L \circ \cdots \circ \phi_1 \circ \pi_{p^r}$ where ϕ_i are isogenies of degree at most d defined over \mathbb{F}_q and π_{p^r} is the p^r -power Frobenius. Let $(\phi_x(x), c_i y \phi_y(x))$ be a representation of ϕ in standard form. Let $n = \lceil \log q \rceil$. Then tr $\alpha \mod p$ can be computed with $O(Ldn + Ln \log n + n(\log n)^2)$ bit operations.

Proof. If α is inseparable, i.e., $r \ge 1$, then tr $\alpha \equiv 0 \pmod{p}$ by Proposition 3.11. Suppose r = 0 so α is separable. For $i = 1, 2, \ldots, e$, let $c_{i,x}$ and $c_{i,y}$ be the leading coefficients of the numerators of $\phi_{i,x}$ and $c\phi_{i,y}$ respectively. By Lemma 3.14, we have $c_i = c_{i,x}^{-1}c_{i,y}$. We then have $a_\alpha = \prod_i c_i = (\prod_i c_{i,x})^{-1}(\prod_i c_{i,y})$. Extracting the coefficients $c_{i,x}$ and $c_{i,y}$ from $\phi_{i,x}$ and $\phi_{i,y}$ can be done in time O(Ldn), i.e., the time required to read in the input. Finally a_α may be computed with O(L) multiplications and one inversion in \mathbb{F}_{p^2} at a cost of $O(Ldn + LM(n) + M(n) \log n) \subseteq O(Ln \log n + n(\log n)^2)$ bit operations. This correctly computes tr α mod p by Proposition 3.11.

Remark 3.16. In practice, an endomorphism α will be represented by a sequence of L isogenies of small (likely prime) degree; presumably these isogenies are computed by Vélu's or Kohel's formulas and therefore are normalized. Thus one can expect the first L - 1 will be normalized and ϕ_L is not. Thus we can find a_{α} by calculating the normalizing coefficient of the final isogeny ϕ_L .

3.3. Computing with rational points. We can exploit the fact that E is supersingular in another. For simplicity, assume E is defined over \mathbb{F}_{p^2} and that $j(E) \notin \{0, 1728\}$. Then $\pi_E = [p]$ or [-p] so $\#E(\mathbb{F}_{p^2})) = (p-1)^2$ or $(p+1)^2$ and as abelian groups we have $E(\mathbb{F}_{p^2}) \cong \mathbb{Z}/(p-1) \times \mathbb{Z}/\mathbb{Z}(p-1)$ or $\mathbb{Z}/(p+1)\mathbb{Z} \times \mathbb{Z}/(p+1)\mathbb{Z}$ accordingly [19, Lemma 4.8]. We can easily decide which is the case given E, for instance by testing whether a random point has order dividing p+1. Assume $\#E(\mathbb{F}_{p^2}) = (p+1)^2$.

- Suppose $\ell^e \mid (p+1)$. Then we can compute the trace t_ℓ as follows:
- (1) Find a point P of order ℓ^e ;
- (2) compute $Q = \alpha(P)$ and $R = (\alpha^2 + \deg(\alpha))(P)$;

- (3) Find $d = \text{dlog}_Q(R)$;
- (4) Then $d \equiv \operatorname{tr} \alpha \pmod{\ell^{e'}}$, where $\ell^{e'}$ is the order of $\alpha(P)$. (Note $e' \leq e$.)

Even better, one can compute $t \mod \ell^e$ using the discrete logarithm between Q = P and $R = \alpha(P) + \hat{\alpha}(P)$. In a group of order ℓ^k , discrete logarithms can be computed with $O(k \log \ell \log \log(\ell^k))$ group operations using the Pohlig–Hellman algorithm as described in [21, § 11.2.3]. Since $\ell^k \in O(p)$ this requires $O(n \log n)$ group operations totaling $O(n^2(\log n)^2)$ bit operations, where $n = \lceil \log p \rceil$. Therefore, this procedure is faster than the algorithm in 3.1 for primes ℓ dividing p-1. Moreover, we recover more information than before: we can recover tr α modulo the order of P.

When $\ell \mid (p-1)$, the quadratic twist of E has a point of order ℓ^k for some $k \ge 1$. Conjugating α by a twisting isomorphism preserves its trace, so we can again compute the trace of α modulo a power of ℓ efficiently by solving a discrete logarithm between points of order ℓ^k on the twist. In isogeny-based cryptography, one often uses primes such that $p^2 - 1$ has a large smooth factor. Therefore, this trick will be particularly useful for cases of practical interest.

4. Computing the trace

We now describe our algorithm for computing the trace of an endomorphism $\alpha \in \text{End}(E)$ of a supersingular elliptic curve E defined over \mathbb{F}_q where q is a power of a prime p > 3. As in the SEA algorithm, we compute tr α by computing $t_{\ell} \coloneqq \text{tr } \alpha \mod \ell$ for enough primes ℓ in order to recover tr α with the Chinese Remainder Theorem.

Theorem 4.1. Let $\alpha = \phi_L \circ \cdots \circ \phi_1$ be a separable endomorphism of a supersingular elliptic curve E defined over \mathbb{F}_q with $j(E) \notin \{0, 1728\}$. Let $n = \lceil \log q \rceil$. Assume that $L \log d \in O(n)$. Then tr α can be computed with $O(n^4(\log n)^2 + dLn^3)$ bit operations. When $d \in O(1)$ and $L \in O(n)$, the complexity is $O(n^4(\log n)^2)$.

Proof. Let $t_{\ell} \coloneqq \operatorname{tr} \alpha \mod \ell$. We need t_{ℓ} for $\ell < B$ such that $\prod_{\ell \leq B} \ell > 4d^{L/2} > 4\sqrt{\deg \alpha}$. By the Prime Number Theorem, we may take $B \in O(L \log d) \subseteq O(n)$, so the largest prime used is O(n). We can compute $t_{\ell} = \operatorname{tr} \alpha \mod \ell$ in time $O(n^3(\log n)^3 + den^2 \log n)$ by Theorem 3.10. The number of primes used is $\pi(B) \in O(n/\log n)$. The total cost is $O(n^3(\log n)^2 + dLn^3)$.

Remark 4.2. If $\alpha \in \operatorname{End}(E)$ is not separable, then α factors as $\alpha = \alpha_s \circ \pi_{p^r} = \phi_L \circ \cdots \circ \phi_1 \circ \pi_{p^r}$. We can compute $t_{\ell} = \operatorname{tr} \alpha \mod \ell$ by computing the restriction of π_{p^r} to ker ψ where ψ is an ℓ -isogeny as in the SEA algorithm, i.e., by computing x^{p^r} modulo the kernel polynomial h of ψ with the square-and-multiply algorithm and then computing $\alpha_h = \alpha_s(\pi_{p^r})$ as in Section 3.1. This has the same complexity (see e.g. [23, Theorem 13]) under the assumption that $r \in O(L)$. Thus we can compute the trace of any endomorphism of a supersingular elliptic curve in time $O(n^3(\log n)^2 + dLn^3)$ assuming $L \log d \in O(n)$ and $r \in O(L)$.

5. Timings

Our implementation of our algorithm in Sage [28] is available at https://github.com/travismo/ beyond-the-SEA/. To demonstrate the asymptotic speedup from Elkies primes and the practical speedup from computing the trace modulo p, we ran the following experiment. For each bit length $b \in \{16, \ldots, 32\}$, we computed a random b-bit prime p, 5 supersingular elliptic curves E/\mathbb{F}_{p^2} , and for each curve, we computed an endomorphism of degree 2^L for $L = 4\lceil \log_2 p \rceil$ using the cycle-finding algorithm in [6]. We then computed the trace of α with Schoof's algorithm (i.e., using division polynomials), with the SEA-algorithm (i.e., using kernel polynomials of ℓ -isogenies), and then the SEA algorithm with the trace-modulo-p algorithm, and finally the SEA algorithm, trace-modulo-p, and using points whenever $\ell \mid \#E(\mathbb{F}_{p^2})$. We observe a substantial speedup from using kernel polynomials rather than division polynomials. We also see the speedup offered from computing the trace modulo p: we very efficiently get about $\log p$ bits of the trace. Since the input endomorphism always has degree $2^{4\lceil \log p \rceil}$, and we need to compute the trace modulo $N = p \prod_{\ell < B} \ell$ so that $N > 4 \cdot 2^{2\lceil \log p \rceil}$ when using the trace-modulo-p information makes the cost of a 32-bit instance roughly as expensive as a 16-bit instance that does not use the modulo-p information and computes the trace modulo $N' = \prod_{\ell < B} \ell$ so that $N' > 4 \cdot 2^{2\lceil \log p \rceil}$. The data in the second figure supports comparison.




FIGURE 1. All four methods

FIGURE 2. The same timings for methods 2 through 4

6. Dewaghe's algorithm for Elkies' method applied to Atkin primes

We conclude with a discussion connecting the material in Section 3.1 with the method of Dewaghe for extending Elkies' method for computing $t_{\ell} = \operatorname{tr} \pi_E \mod \ell$ for an ordinary curve E and Atkin prime ℓ for E. For simplicity, suppose E is defined over \mathbb{F}_p . Additionally assume $j(E) \notin \{0, 1728\}$. Define $\varphi_{\ell}(x) \coloneqq \Phi_{\ell}(j(E), x)$ and suppose φ_{ℓ} has no roots in \mathbb{F}_p so E has no \mathbb{F}_p -rational ℓ -isogenies. Then ℓ is not an Elkies prime for E, but it could be that it is an Elkies prime for E after a small extension of the base field. School proves in [20, Proposition 6.2(iii)] (under our assumptions that E is ordinary and $j(E) \notin \{0, 1728\}$) that if φ_{ℓ} has no roots in \mathbb{F}_p , it factors into $(\ell + 1)/r$ factors of degree r, where r is the order of the image of π_E in PGL($E[\ell]$). The integer r can be computed by computing $\operatorname{gcd}(x^{p^i} - x, \varphi_{\ell}(x))$ for $i = 1, 2, \ldots, \ell + 1$ as is done in Atkin's modification of School's algorithm [20, §6].

Let f be an irreducible factor of degree r of $\varphi(x)$ and let $\mathbb{F}_{p^r} = \mathbb{F}_p[t]/(f(t))$. The division polynomial ψ_{ℓ} has a factor h of degree $(\ell - 1)/2$ over \mathbb{F}_{p^r} corresponding to the kernel polynomial of a \mathbb{F}_{p^r} -rational ℓ -isogeny with kernel K. It is not the case that π_E induces an endomorphism of K, but we can still use the algorithm in 3.1 to compute $t_{\ell} \coloneqq \operatorname{tr} \pi_E \mod \ell$. We may compute $t_{\ell} = \operatorname{tr} \pi_E \mod \ell$ by, as before, computing c such that

$$\pi_E^2\big|_{\ker\phi} + p\big|_{\ker\phi} = c\pi_E\big|_{\ker\phi}.$$

In the worst case, this is no faster than Schoof's algorithm for computing t_{ℓ} : the degree of f can be $\ell + 1$, and the degree of h will be $(\ell - 1)/2$, so the polynomial arithmetic in $\mathbb{F}_{p^r}[x]/(h)$ will be as expensive as the arithmetic in $\mathbb{F}_p[x]/(\psi_{\ell})$. But it is possible that $\varphi_{\ell}(x)$ can have irreducible factors of degree strictly less than $\ell + 1$.

Because E has a \mathbb{F}_{p^r} -rational ℓ -isogeny with kernel polynomial h, there is a factor of ψ_{ℓ} of degree $r(\ell-1)/2$ in $\mathbb{F}_p[x]$ given by $g = \prod_{i=1}^r h^{(p^i)}$. Then the roots of g are the x-coordinates of affine points in the orbit of K under π_E . Let V be the subscheme of E cut out by g. Then one may compute t_{ℓ} by computing $c = t_{\ell}$ such that $\pi_E^2|_V + q|_V = c\pi_E|_V$. At a high level, this is the method of Dewaghe [5, Section 4] for applying Elkies' method to Atkin primes.

References

- A. Bostan, F. Morain, B. Salvy, and É. Schost. Fast algorithms for computing isogenies between elliptic curves. Math. Comp., 77(263):1755–1778, 2008.
- [2] Jean-Marc Couveignes and Fran, cois Morain. Schoof's algorithm and isogeny cycles. In Algorithmic number theory (Ithaca, NY, 1994), volume 877 of Lecture Notes in Comput. Sci., pages 43–58. Springer, Berlin, 1994.
- [3] John Cremona. Comment to sage math ticket $\#35546,\,2023.$
- [4] Christina Delfs and Steven D. Galbraith. Computing isogenies between supersingular elliptic curves over \mathbb{F}_p . Des. Codes Cryptography, 78(2):425–440, February 2016.
- [5] L. Dewaghe. Remarks on the Schoof-Elkies-Atkin algorithm. Math. Comp., 67(223):1247–1252, 1998.
- [6] Kirsten Eisenträger, Sean Hallgren, Chris Leonardi, Travis Morrison, and Jennifer Park. Computing endomorphism rings of supersingular elliptic curves and connections to path-finding in isogeny graphs. In ANTS XIV-Proceedings of the Fourteenth Algorithmic Number Theory Symposium, pages 215–232, 2020.

- [7] Noam D. Elkies. Elliptic and modular curves over finite fields and related computational issues. In Computational perspectives on number theory (Chicago, IL, 1995), volume 7 of AMS/IP Stud. Adv. Math., pages 21–76. Amer. Math. Soc., Providence, RI, 1998.
- [8] Jenny Fuselier, Annamaria Iezzi, Mark Kozek, Travis Morrison, and Changningphaabi Namoijam. Computing supersingular endomorphism rings using inseparable endomorphisms, 2023.
- [9] Steven D. Galbraith. Mathematics of public key cryptography. Cambridge University Press, Cambridge, 2012.
- [10] David Harvey and Joris van der Hoeven. Integer multiplication in time O(n log n). Ann. of Math. (2), 193(2):563-617, 2021.
- [11] Masanobu Kaneko. Supersingular j-invariants as singular moduli mod p. Osaka J. Math., 26(4):849–855, 1989.
- [12] David Kohel. Endomorphism rings of elliptic curves over finite fields. PhD thesis, University of California, Berkeley, 1996.
- [13] Sabrina Kunzweiler and Damien Robert. Computing modular polynomials by deformation. To appear in the Proceedings of the Sixteenth Algorithmic Number Theory Symposium, 2024.
- [14] Serge Lang. Elliptic functions, volume 112 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2 edition, 1987. With an appendix by J. Tate.
- [15] William E. Mahaney and Travis Morrison. Computing isogenies at singular points of the modular polynomial, 2024.
- [16] Fran, cois Morain, Charlotte Scribot, and Benjamin Smith. Computing cardinalities of Q-curve reductions over finite fields. LMS J. Comput. Math., 19:115–129, 2016.
- [17] Michael O. Rabin. Probabilistic algorithms in finite fields. SIAM J. Comput., 9(2):273–280, 1980.
- [18] A. Schönhage and V. Strassen. Schnelle Multiplikation großer Zahlen. Computing, 7:281–292, 1971.
- [19] René Schoof. Nonsingular plane cubic curves over finite fields. J. Combin. Theory Ser. A, 46(2):183–211, 1987.
- [20] René Schoof. Counting points on elliptic curves over finite fields. volume 7, pages 219–254. 1995. Les Dix-huitièmes Journées Arithmétiques (Bordeaux, 1993).
- [21] Victor Shoup. A computational introduction to number theory and algebra. Cambridge University Press, Cambridge, 2 edition, 2009.
- [22] Igor E. Shparlinski and Andrew V. Sutherland. On the distribution of Atkin and Elkies primes. Found. Comput. Math., 14(2):285–297, 2014.
- [23] Igor E. Shparlinski and Andrew V. Sutherland. On the distribution of Atkin and Elkies primes for reductions of elliptic curves on average. LMS J. Comput. Math., 18(1):308–322, 2015.
- [24] Joseph H. Silverman. The arithmetic of elliptic curves. Springer, New York, 2009.
- [25] Andrew Sutherland. 18.783 Elliptic Curves. Massachusetts Institute of Technology: MIT OpenCourseWare. License: Creative Commons BY-NC-SA.
- [26] Andrew Sutherland. Genus 1 point counting records over prime fields, 2010.
- [27] Andrew V. Sutherland. On the evaluation of modular polynomials. In ANTS X Proceedings of the Tenth Algorithmic Number Theory Symposium, volume 1 of Open Book Ser., pages 531–555. Math. Sci. Publ., Berkeley, CA, 2013.
- [28] The Sage Developers. SageMath, the Sage Mathematics Software System (Version 10.5), 2024.
- [29] Jacques Vélu. Isogénies entre courbes elliptiques. C. R. Acad. Sci. Paris Sér. A-B, 273:A238–A241, 1971.
- [30] Joachim von zur Gathen and Jürgen Gerhard. Modern computer algebra. Cambridge University Press, Cambridge, 3 edition, 2013.

TRAVIS MORRISON, VIRGINIA TECH, USA Email address: tmo@vt.edu

LORENZ PANNY, TECHNISCHE UNIVERSITÄT MÜNCHEN, GERMANY *Email address:* lorenz@yx7.cc

JANA SOTÁKOVÁ, UNIVERSITY OF AMSTERDAM, NETHERLANDS *Email address:* ja.sotakova@gmail.com

MICHAEL WILLS, UNIVERSITY OF VIRGINIA, USA *Email address*: muf5cz@virginia.edu