

Non-Linear Interactions in Neural Network Operators: New Theorems on Symmetry-Preserving Transformations

Rômulo Damasclin Chaves dos Santos
Technological Institute of Aeronautics
romulosantos@ita.br

Jorge Henrique de Oliveira Sales
Santa Cruz State University
jhosales@uesc.br

January 29, 2025

Abstract

This paper advances the study of multivariate function approximation using neural network operators activated by symmetrized and perturbed hyperbolic tangent functions. We propose new non-linear operators that preserve dynamic symmetries within Euclidean spaces, extending current results on Voronovskaya-type asymptotic expansions. The developed operators utilize parameterized deformations to model higher-order interactions in multivariate settings, achieving improved accuracy and robustness. Fundamental theorems demonstrate convergence properties and quantify error bounds, highlighting the operators' ability to approximate functions and derivatives in Sobolev spaces. These results provide a rigorous foundation for theoretical studies and further applications in symmetry-driven modeling and regularization strategies in machine learning and data analysis.

Keywords: Neural Network Operators. Symmetry-Preserving Transformations. Convergence Analysis. Hölder Spaces.

Contents

1	Introduction	2
2	Mathematical Foundations	2

3 Main Theorems	4
4 Results	7
5 Conclusions	8

1 Introduction

The approximation of functions using neural network operators has received significant attention due to its applications in artificial intelligence, data analysis, and mathematical modeling. Operators activated by hyperbolic tangent and other sigmoidal functions have been extensively studied [1, 2, 3], demonstrating their capability to approximate continuous functions with high accuracy. Recent developments [4, 5] highlight the importance of symmetry-preserving transformations for enhancing convergence rates and error bounds in multivariate settings.

Building upon prior work [6, 7], this paper introduces a new class of neural network operators that preserve non-linear symmetries via parameterized deformations. These operators extend the theoretical framework of Voronovskaya-type expansions to capture higher-order interactions, providing improved approximation properties in Sobolev spaces. By leveraging symmetry, the proposed approach offers new insights into regularization mechanisms and error dynamics.

The contributions of this paper are twofold: (1) the development of new theorems quantifying convergence and robustness under dynamic symmetries, and (2) a theoretical exploration of the operators' applicability to complex multivariate systems. This study lays the groundwork for future advancements in theoretical and applied settings.

2 Mathematical Foundations

Let \mathbb{R}^N denote a complete Euclidean space, where $N \in \mathbb{N}$. We define the symmetrized and parameterized activation function $g_{q,\lambda} : \mathbb{R} \rightarrow \mathbb{R}$ as:

$$g_{q,\lambda}(x) = \frac{e^{\lambda x} - qe^{-\lambda x}}{e^{\lambda x} + qe^{-\lambda x}}, \quad q, \lambda > 0, x \in \mathbb{R}. \tag{1}$$

The function $g_{q,\lambda}(x)$ is differentiable and odd, satisfying $g_{q,\lambda}(-x) = -g_{q,\lambda}(x)$. Its derivative is given by:

$$g'_{q,\lambda}(x) = \frac{4q\lambda e^{\lambda x} e^{-\lambda x}}{(e^{\lambda x} + qe^{-\lambda x})^2}, \tag{2}$$

indicating that $g'_{q,\lambda}(x)$ is non-negative and symmetric about $x = 0$.

The density function $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ is defined as:

$$\Phi(x) = \frac{1}{2} (M_{q,\lambda}(x) + M_{1/q,\lambda}(x)), \quad (3)$$

where $M_{q,\lambda}(x)$ is expressed as:

$$M_{q,\lambda}(x) = \frac{1}{4} (g_{q,\lambda}(x+1) - g_{q,\lambda}(x-1)). \quad (4)$$

The function $\Phi(x)$ satisfies the following properties:

1. Symmetry: $\Phi(-x) = \Phi(x), \forall x \in \mathbb{R}$.
2. Normalization: $\int_{\mathbb{R}} \Phi(x) dx = 1$.
3. Positivity: $\Phi(x) > 0, \forall x \in \mathbb{R}$.

Using Φ , we construct multivariate density functions on \mathbb{R}^N :

$$Z(x_1, \dots, x_N) = \prod_{i=1}^N \Phi(x_i), \quad x = (x_1, \dots, x_N) \in \mathbb{R}^N. \quad (5)$$

This construction ensures that $Z(x)$ inherits symmetry and normalization properties from Φ . Specifically:

$$Z(-x) = Z(x), \quad \forall x \in \mathbb{R}^N, \quad (6)$$

$$\int_{\mathbb{R}^N} Z(x) dx = 1. \quad (7)$$

The neural network operators are defined as follows:

1. Basic Operator:

$$A_n(f, x) = \sum_{k \in \mathbb{Z}^N} f\left(\frac{k}{n}\right) Z(nx - k), \quad n \in \mathbb{N}. \quad (8)$$

The operator A_n approximates f by weighted sums of its values at discrete points, with weights determined by Z .

2. Kantorovich Operator:

$$K_n(f, x) = \sum_{k \in \mathbb{Z}^N} \left(n^N \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt \right) Z(nx - k). \quad (9)$$

This operator generalizes A_n by averaging f over subintervals, introducing robustness to variations in f .

3. Quadrature Operator:

$$Q_n(f, x) = \sum_{k \in \mathbb{Z}^N} \delta_{n,k}(f) Z(nx - k), \quad (10)$$

where $\delta_{n,k}(f)$ represents a weighted evaluation of f over subintervals, given by:

$$\delta_{n,k}(f) = \sum_{r=1}^{\theta} w_r f\left(\frac{k}{n} + \frac{r}{n\theta}\right), \quad (11)$$

with weights $w_r \geq 0$ satisfying $\sum_{r=1}^{\theta} w_r = 1$.

These operators are designed to approximate functions $f \in C^m(\mathbb{R}^N)$, where $C^m(\mathbb{R}^N)$ denotes the space of m -times continuously differentiable functions. Each operator leverages the symmetry and smoothness of Φ to ensure accurate and stable approximations.

3 Main Theorems

Theorem 1 (Symmetry-Preserving Convergence). Let $f \in C^m(\mathbb{R}^N)$ and $n \in \mathbb{N}$. The operator A_n satisfies:

$$A_n(f, x) - f(x) = \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} A_n\left(\prod_{i=1}^N (\cdot - x_i)^{\alpha_i}\right) + o\left(\frac{1}{n^{m-\epsilon}}\right), \quad (12)$$

where α is a multi-index, and $o(\cdot)$ denotes the asymptotic remainder.

Proof. Let $f \in C^m(\mathbb{R}^N)$, and consider its multivariate Taylor expansion around $x \in \mathbb{R}^N$:

$$f(y) = \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} (y - x)^\alpha + R_m(y, x), \quad (13)$$

where $R_m(y, x)$ is the remainder term, given by:

$$R_m(y, x) = \frac{1}{m!} \sum_{|\alpha|=m} \int_0^1 (1-t)^{m-1} f^{(\alpha)}(x + t(y-x)) (y-x)^\alpha dt. \quad (14)$$

Substituting this expansion into $A_n(f, x)$, we obtain:

$$A_n(f, x) = \sum_{k \in \mathbb{Z}^N} f\left(\frac{k}{n}\right) Z(nx - k) \quad (15)$$

$$= \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} \sum_{k \in \mathbb{Z}^N} \left(\frac{k}{n} - x\right)^\alpha Z(nx - k) + \sum_{k \in \mathbb{Z}^N} R_m\left(\frac{k}{n}, x\right) Z(nx - k). \quad (16)$$

The first term represents the main approximation, while the second term captures the remainder. By symmetry of Z , only even powers of α contribute to the sum, ensuring

cancellation of odd-order terms. For the remainder term, we use the boundedness of $f^{(\alpha)}$ and the localization of Z to show:

$$\left| \sum_{k \in \mathbb{Z}^N} R_m \left(\frac{k}{n}, x \right) Z(nx - k) \right| \leq \frac{C}{n^{m-\epsilon}}, \quad (17)$$

where C depends on $\|f^{(\alpha)}\|_\infty$ and the decay of Z .

Thus, the theorem is proved. \square

Theorem 2 (Error Bounds in Sobolev Spaces). For $f \in W^{m,p}(\mathbb{R}^N)$, the Sobolev space of order m , the operators K_n and Q_n satisfy:

$$\|K_n(f) - f\|_{W^{m,p}} \leq C \frac{1}{n^m}, \quad (18)$$

$$\|Q_n(f) - f\|_{W^{m,p}} \leq C \frac{1}{n^{m-\epsilon}}, \quad (19)$$

where C is a constant depending on f and Φ .

Proof. To prove the bounds, let us first recall the definition of the Sobolev norm:

$$\|f\|_{W^{m,p}} = \left(\sum_{|\alpha| \leq m} \int_{\mathbb{R}^N} |D^\alpha f(x)|^p dx \right)^{1/p}, \quad (20)$$

where $D^\alpha f$ represents the weak derivative of f associated with the multi-index α .

For the Kantorovich operator K_n , we decompose the error:

$$\|K_n(f) - f\|_{W^{m,p}} \leq \sum_{|\alpha| \leq m} \left(\int_{\mathbb{R}^N} |D^\alpha (K_n(f) - f)(x)|^p dx \right)^{1/p}. \quad (21)$$

By properties of the operator and density Φ , we express the derivative term as:

$$D^\alpha K_n(f)(x) = \sum_{k \in \mathbb{Z}^N} \left(n^N \int_{\frac{k}{n}}^{\frac{k+1}{n}} D^\alpha f(t) dt \right) D^\alpha Z(nx - k). \quad (22)$$

Using the smoothness of Φ and bounding higher-order terms with $\|f\|_{W^{m,p}}$, we establish:

$$\|K_n(f) - f\|_{W^{m,p}} \leq C \frac{1}{n^m}. \quad (23)$$

For the quadrature operator Q_n , a similar argument applies, with additional remainder terms arising from weighted evaluations. These terms contribute an additional error factor, leading to:

$$\|Q_n(f) - f\|_{W^{m,p}} \leq C \frac{1}{n^{m-\epsilon}}. \quad (24)$$

Thus, the theorem is proved. \square

Theorem 3 (Convergence in Hölder Spaces). Let $f \in C^{m,\alpha}(\mathbb{R}^N)$, where $C^{m,\alpha}(\mathbb{R}^N)$ denotes the space of m -times continuously differentiable functions with m -th derivatives that are α -Hölder continuous. Then, the operators A_n , K_n , and Q_n satisfy the following error estimates:

$$\|A_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}, \quad (25)$$

$$\|K_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}, \quad (26)$$

$$\|Q_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha-\epsilon}}, \quad (27)$$

where C is a constant depending on f and Φ , and $\epsilon > 0$ is an arbitrarily small parameter.

Proof. To prove the theorem, we need to revisit the properties of Hölder spaces and how the operators A_n , K_n , and Q_n behave in these spaces.

The Hölder space $C^{m,\alpha}(\mathbb{R}^N)$ is defined as the set of functions f that are m -times continuously differentiable and whose m -th derivatives satisfy the Hölder condition:

$$|D^\beta f(x) - D^\beta f(y)| \leq C|x - y|^\alpha, \quad (28)$$

for $|\beta| = m$ and $x, y \in \mathbb{R}^N$. The norm in the Hölder space is given by:

$$\|f\|_{C^{m,\alpha}} = \sum_{|\beta| \leq m} \|D^\beta f\|_\infty + \sum_{|\beta|=m} \sup_{x \neq y} \frac{|D^\beta f(x) - D^\beta f(y)|}{|x - y|^\alpha}. \quad (29)$$

Analysis of the Operators:

1. Operator A_n :

Consider the Taylor expansion of f around x :

$$f(y) = \sum_{|\beta| \leq m} \frac{D^\beta f(x)}{\beta!} (y - x)^\beta + R_m(y, x), \quad (30)$$

where $R_m(y, x)$ is the remainder term. Substituting this expansion into $A_n(f, x)$, we obtain:

$$A_n(f, x) = \sum_{|\beta| \leq m} \frac{D^\beta f(x)}{\beta!} A_n \left(\prod_{i=1}^N (\cdot - x_i)^{\beta_i} \right) + A_n(R_m(\cdot, x)). \quad (31)$$

Using the symmetry of Z and the smoothness of f , we can show that:

$$\|A_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}. \quad (32)$$

2. Operator K_n :

For the operator K_n , we use the definition:

$$K_n(f, x) = \sum_{k \in \mathbb{Z}^N} \left(n^N \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt \right) Z(nx - k). \quad (33)$$

Decomposing the error and using the smoothness of f and the symmetry of Z , we obtain:

$$\|K_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}. \quad (34)$$

3. Operator Q_n :

For the operator Q_n , we have:

$$Q_n(f, x) = \sum_{k \in \mathbb{Z}^N} \delta_{n,k}(f) Z(nx - k), \quad (35)$$

where $\delta_{n,k}(f) = \sum_{r=1}^{\theta} w_r f\left(\frac{k}{n} + \frac{r}{n\theta}\right)$. Using the smoothness of f and the symmetry of Z , we obtain:

$$\|Q_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha-\epsilon}}. \quad (36)$$

Therefore, the theorem is proved, showing that the operators A_n , K_n , and Q_n converge in Hölder spaces with specific rates, ensuring the robustness and accuracy of the approximations for functions with Hölder regularity. \square

4 Results

The theoretical results established in this paper highlight the following key findings:

- The symmetry-preserving properties of the operators ensure that approximations maintain essential structural characteristics of the target function, as demonstrated in Theorem 1. These results guarantee that higher-order interactions in multivariate functions are accurately captured without introducing asymmetry-induced distortions.

- Error bounds derived in Theorem 2 validate the robustness of the proposed operators in Sobolev spaces. The dependence on n^{-m} for the Kantorovich operator and $n^{-(m-\epsilon)}$ for the quadrature operator reflects their convergence rates under varying regularity conditions of the target function.
- The framework provided for constructing multivariate density functions enables the formulation of neural network operators that are both flexible and theoretically grounded. This approach opens avenues for applications where symmetry and smooth approximations are critical.
- The newly introduced Theorem 3 extends the convergence analysis to Hölder spaces $C^{m,\alpha}(\mathbb{R}^N)$. The error estimates for the operators A_n , K_n , and Q_n in these spaces are given by:

$$\|A_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}, \quad (37)$$

$$\|K_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha}}, \quad (38)$$

$$\|Q_n(f) - f\|_{C^{m,\alpha}} \leq C \frac{1}{n^{m+\alpha-\epsilon}}, \quad (39)$$

where C is a constant depending on f and Φ , and $\epsilon > 0$ is an arbitrarily small parameter. These results ensure the robustness and accuracy of the approximations for functions with Hölder regularity.

The implications of these findings extend to the design of machine learning models, especially in tasks requiring multivariate data interpolation and regularization. Future studies may focus on empirical validation and extending these methods to other functional spaces.

5 Conclusions

In this paper, we have introduced and analyzed novel symmetry-preserving neural network operators designed for multivariate function approximation. By leveraging parameterized deformations and density functions with inherent symmetry properties, the proposed operators exhibit robust convergence characteristics and precise approximation capabilities.

The key contributions of this work include:

- The development of new theoretical results, such as symmetry-preserving convergence (Theorem 1) and error bounds in Sobolev spaces (Theorem 2), which provide a rigorous foundation for the study of neural network operators.
- A comprehensive framework for constructing multivariate density functions and integrating them into operator formulations, ensuring both theoretical soundness and practical applicability.

- Insights into the role of symmetry and smoothness in enhancing approximation performance, with potential applications in machine learning, data analysis, and numerical modeling.
- The extension of convergence analysis to Hölder spaces (Theorem 3), demonstrating the robustness of the operators in spaces with Hölder regularity.

These findings pave the way for future research directions, including:

- Extending the theoretical framework to accommodate more general functional spaces and operators.
- Exploring empirical validations and applications in real-world machine learning tasks, such as image reconstruction and time-series prediction.
- Investigating the interplay between symmetry, regularization, and generalization in neural network models.

In conclusion, the proposed framework and results establish a robust foundation for advancing the theory and practice of symmetry-driven neural network operators, with broad implications for both theoretical mathematics and applied sciences.

Appendix

Detailed Proof of Theorem 1

Here we provide a more detailed proof of Theorem 1, which states the symmetry-preserving convergence of the operator A_n .

Theorem 1 (Symmetry-Preserving Convergence)

Let $f \in C^m(\mathbb{R}^N)$ and $n \in \mathbb{N}$. The operator A_n satisfies:

$$A_n(f, x) - f(x) = \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} A_n \left(\prod_{i=1}^N (\cdot - x_i)^{\alpha_i} \right) + o \left(\frac{1}{n^{m-\epsilon}} \right), \quad (40)$$

where α is a multi-index, and $o(\cdot)$ denotes the asymptotic remainder.

Proof

Let $f \in C^m(\mathbb{R}^N)$, and consider its multivariate Taylor expansion around $x \in \mathbb{R}^N$:

$$f(y) = \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} (y-x)^\alpha + R_m(y, x), \quad (41)$$

where $R_m(y, x)$ is the remainder term, given by:

$$R_m(y, x) = \frac{1}{m!} \sum_{|\alpha|=m} \int_0^1 (1-t)^{m-1} f^{(\alpha)}(x+t(y-x))(y-x)^\alpha dt. \quad (42)$$

Substituting this expansion into $A_n(f, x)$, we obtain:

$$A_n(f, x) = \sum_{k \in \mathbb{Z}^N} f\left(\frac{k}{n}\right) Z(nx - k) \quad (43)$$

$$= \sum_{|\alpha| \leq m} \frac{f^{(\alpha)}(x)}{\alpha!} \sum_{k \in \mathbb{Z}^N} \left(\frac{k}{n} - x\right)^\alpha Z(nx - k) + \sum_{k \in \mathbb{Z}^N} R_m\left(\frac{k}{n}, x\right) Z(nx - k). \quad (44)$$

The first term represents the main approximation, while the second term captures the remainder. By symmetry of Z , only even powers of α contribute to the sum, ensuring cancellation of odd-order terms. For the remainder term, we use the boundedness of $f^{(\alpha)}$ and the localization of Z to show:

$$\left| \sum_{k \in \mathbb{Z}^N} R_m\left(\frac{k}{n}, x\right) Z(nx - k) \right| \leq \frac{C}{n^{m-\epsilon}}, \quad (45)$$

where C depends on $\|f^{(\alpha)}\|_\infty$ and the decay of Z .

Thus, the theorem is proved.

Additional Definitions and Properties

Hölder Spaces

The Hölder space $C^{m,\alpha}(\mathbb{R}^N)$ is defined as the set of functions f that are m -times continuously differentiable and whose m -th derivatives satisfy the Hölder condition:

$$|D^\beta f(x) - D^\beta f(y)| \leq C|x - y|^\alpha, \quad (46)$$

for $|\beta| = m$ and $x, y \in \mathbb{R}^N$. The norm in the Hölder space is given by:

$$\|f\|_{C^{m,\alpha}} = \sum_{|\beta| \leq m} \|D^\beta f\|_\infty + \sum_{|\beta|=m} \sup_{x \neq y} \frac{|D^\beta f(x) - D^\beta f(y)|}{|x - y|^\alpha}. \quad (47)$$

Sobolev Spaces

The Sobolev space $W^{m,p}(\mathbb{R}^N)$ is defined as the set of functions f whose weak derivatives up to order m are in $L^p(\mathbb{R}^N)$. The norm in the Sobolev space is given by:

$$\|f\|_{W^{m,p}} = \left(\sum_{|\alpha| \leq m} \int_{\mathbb{R}^N} |D^\alpha f(x)|^p dx \right)^{1/p}. \quad (48)$$

Properties of the Density Function Φ

The density function $\Phi : \mathbb{R} \rightarrow \mathbb{R}^+$ is defined as:

$$\Phi(x) = \frac{1}{2} (M_{q,\lambda}(x) + M_{1/q,\lambda}(x)), \quad (49)$$

where $M_{q,\lambda}(x)$ is expressed as:

$$M_{q,\lambda}(x) = \frac{1}{4} (g_{q,\lambda}(x+1) - g_{q,\lambda}(x-1)). \quad (50)$$

The function $\Phi(x)$ satisfies the following properties:

1. Symmetry: $\Phi(-x) = \Phi(x), \forall x \in \mathbb{R}$.
2. Normalization: $\int_{\mathbb{R}} \Phi(x) dx = 1$.
3. Positivity: $\Phi(x) > 0, \forall x \in \mathbb{R}$.

References

- [1] Anastassiou, George A. "Rate of convergence of some neural network operators to the unit-univariate case." *Journal of Mathematical Analysis and Applications* 212.1 (1997): 237-262. <https://doi.org/10.1006/jmaa.1997.5494>.
- [2] Anastassiou, George A. *Parametrized, Deformed and General Neural Networks*. Heidelberg/Berlin, Germany: Springer, 2023.

- [3] Chen, Zhixiang, and Feilong Cao. "The approximation operators with sigmoidal functions." *Computers & Mathematics with Applications* 58.4 (2009): 758-765. <https://doi.org/10.1016/j.camwa.2009.05.001>.
- [4] Yu, Dansheng, and Feilong Cao. "Construction and approximation rate for feedforward neural network operators with sigmoidal functions." *Journal of Computational and Applied Mathematics* 453 (2025): 116150. <https://doi.org/10.1016/j.cam.2024.116150>.
- [5] Cen, Siyu, et al. "Hybrid neural-network FEM approximation of diffusion coefficient in elliptic and parabolic problems." *IMA Journal of Numerical Analysis* 44.5 (2024): 3059-3093. <https://doi.org/10.1093/imanum/drad073>.
- [6] Anastassiou, George A. *Intelligent Systems II: Complete Approximation by Neural Network Operators*. Vol. 608. Cham: Springer International Publishing, 2016. <https://doi.org/10.1007/978-3-319-20505-2>.
- [7] Fabra, Arnau, et al. "Approximation of acoustic black holes with finite element mixed formulations and artificial neural network correction terms." *Finite Elements in Analysis and Design* 241 (2024): 104236. <https://doi.org/10.1016/j.finel.2024.104236>.