Symmetry-induced fragmentation and dissipative time crystal

Haowei $Li¹$ and Wei $Yi^{1, 2, 3, 4, 5, *}$

¹*CAS Key Laboratory of Quantum Information, University of Science and Technology of China, Hefei 230026, China*

²*Anhui Province Key Laboratory of Quantum Network,*

University of Science and Technology of China, Hefei 230026, China

³*CAS Center For Excellence in Quantum Information and Quantum Physics, Hefei 230026, China*

⁴*Hefei National Laboratory, University of Science and Technology of China, Hefei 230088, China*

⁵*5Anhui Center for Fundamental Sciences in Theoretical Physics,*

University of Science and Technology of China, Hefei 230026, China

We propose a general protocol for engineering dissipative time crystals in quantum open systems through symmetry-induced fragmentation and ergodicity breaking. Building upon a $U(1)$ -symmetry-induced Liouvillespace fragmentation, we devise a generic Liouvillian possessing a series of non-dissipative eigenmodes with purely imaginary eigenvalues, which give rise to long-time oscillations breaking both ergodicity and timetranslation symmetry. Illustrating our proposal using a dissipative lattice model, we show that, even when the *U*(1) symmetry is broken, a prethermal time-crystal behavior survives, with distinct oscillation frequencies at different times of the steady-state approaching dynamics. The richly structured prethermal time-crystal dynamics derive from Fermi statistics and the Liouvillian skin effect of our model, under which excitations above the boundary-localized dark states can be mapped to the irreducible representations of the permutation group. The branching rules of the permutation group then dictate that the Liouvillian gaps acquired by the non-dissipative eigenmodes should be higher-order powers of the symmetry-breaking perturbation, leading to the prethermal time-crystal dynamics.

Introduction.— Time crystals possess peculiar spatiotemporal structures that simultaneously break ergodicity and time-translation symmetry, exhibiting self-organized timeperiodic dynamics [\[1](#page-4-1)[–4\]](#page-4-2). Over the past decade, various types of time crystals have been proposed and experimentally observed, and a common theme is the breaking of ergodicity. For instance, in discrete time crystals [\[3](#page-4-3)[–26\]](#page-5-0), where the discrete time-translation symmetry of a periodically driven system (or Floquet system) is spontaneously broken, the system breaks ergodicity either by resorting to the many-body localization [\[5–](#page-4-4)[12\]](#page-4-5), or, in an approximate fashion, through the Floquet prethermalization [\[13](#page-4-6)[–18\]](#page-5-1). In the latter case, subharmonic oscillations occur within an intermediate prethermal time window, before the system finally gives in to the Floquet heating. In continuous time crystals that arise in the non-equilibrium dynamics of dissipative systems [\[27–](#page-5-2) [38\]](#page-5-3), the ergodicity is broken either through stable limit cycles [\[27](#page-5-2)[–30\]](#page-5-4) or emergent symmetries in the thermodynamic limit [\[31](#page-5-5)[–35\]](#page-5-6). But ergodicity of a quantum many-body system can also be broken through other exotic mechanisms, including the recently discovered quantum many-body scars [\[39–](#page-5-7) [45\]](#page-5-8) and Hilbert-space fragmentation [\[46–](#page-5-9)[53\]](#page-5-10). Whether timecrystalline order can be stabilized under these novel circumstances is an important but open question.

In this work, we show that dissipative time crystals can emerge in quantum open systems through a symmetryinduced Liouville-space fragmentation. We propose a general framework where the *U*(1) symmetry of a designed Liouvillian gives rise to fragmentation, thus breaking ergodicity. As a result, the Liouvillian develops a series of non-dissipative eigenmodes with purely imaginary eigenvalues that are responsible for persistent oscillations at long times, a key signature of dissipative time crystals. We then focus on a dissipative lattice model of fermions as an illustrating exam-

ple, and further demonstrate that the system is survived by prethermal time-crystal dynamics under symmetry-breaking perturbations. In particular, with the onset of perturbations, the eigenvalue of each non-dissipative eigenmode acquires a small real part (the Liouvillian gap of the eigenmode), whose magnitude depends on the imaginary component of the eigenvalue. This gives rise to a rich structure in the prethermal timecrystal dynamics—the system oscillates with distinct frequencies at different time scales of the steady-state approaching dynamics, as different eigenmodes take turns to be dominant. Detailed analysis reveals that both the robustness and structure of the prethermal time crystal are predicated by the Liouvillian skin effect and Fermi statistics. Under the Liouvillian skin effect [\[54–](#page-5-11)[64\]](#page-6-0), the dark states of the quantum jump operators become boundary localized, forming a real-space Fermi sea. As a consequence, fermionic excitations above the dark states can be mapped to the irreducible representations of the permutation group. Using the branching rules of the permutation group, we show that the non-dissipative eigenmodes become weakly dissipative, as they acquire eigenvalues that exhibit higher-order power-law scalings with the symmetry-breaking perturbation. Hence, it is this algebraic structure that protects the prethermal time-crystal behavior. Our work illustrates, for the first time, that symmetry-induced fragmentation can stabilize time-crystal dynamics in quantum open systems. It further highlights the pivotal role of non-Hermitian phenomena, such as the Liouvillian skin effect, in the dynamics of manybody quantum open systems.

General framework.— We start from a general Lindblad master equation (setting $\hbar = 1$)

$$
\frac{d\rho}{dt} = \mathcal{L}[\rho] = -i[\hat{H}, \rho] + \sum_{\mu} \gamma_{\mu} \left(2\hat{K}_{\mu}\rho \hat{K}_{\mu}^{\dagger} - \left\{ \hat{K}_{\mu}^{\dagger} \hat{K}_{\mu}, \rho \right\} \right), \quad (1)
$$

where ρ is the density matrix of the open system, $\mathcal L$ is the

FIG. 1. Symmetry-induced fragmentation and time crystal. (a) Block-diagonaized Liouvillian matrix. The red boxes indicate the diagonal symmetry sectors with $f_i = f_j$, the inset shows a zoomedin view of the matrix elements (tinged according to the color bar) within a symmetry sector. (b) The Liouvillian spectrum on the complex plane. The inset shows the non-dissipative eigenmodes on the imaginary axis. (c) The long-time dynamics of $\langle \hat{c}_3^{\dagger} \hat{c}_2 + \hat{c}_2^{\dagger} \hat{c}_3 \rangle$. (d) Two periods of the oscillatory dynamics starting at $t_0\gamma = 10^5$ (up-
per panel) and $t_0\gamma = 10^7$ (lower panel), respectively marked by the per panel) and $t_0\gamma = 10^7$ (lower panel), respectively marked by the blue and green vertical lines in (c). For all figures, we take $s = 0$. blue and green vertical lines in (c). For all figures, we take $s = 0$, $\Delta/\gamma = 0.2, L = 8$, and $N = 4$.

Liouvillian superoperator, \hat{H} is the coherent Hamiltonian, and \hat{K}_{μ} are a set of quantum jump operators. We focus on the age where \hat{K}_{μ} are a set of quantum jump operators. We focus on the case where $\mathcal L$ possesses a $U(1)$ symmetry, defined as $[L, \mathcal{U}^{\hat{A}}] = 0$ for the symmetry superoperator $\mathcal{U}^{\hat{A}}$, where $\mathcal{U}^{\hat{A}}$ (o) $-\frac{e^{i\theta \hat{A}}Qe^{-i\theta \hat{A}}}$ for $\theta \in [0, 2\pi)$ and a Hermitian operator $\mathcal{U}^{\hat{A}}[\rho] = e^{i\theta \hat{A}} \rho e^{-i\theta \hat{A}}$ for $\theta \in [0, 2\pi)$ and a Hermitian operator \hat{A} . Note that this is a natural extension of the Z₂ symmetry \hat{A} . Note that this is a natural extension of the Z_2 symmetry discussed in Ref. [\[65\]](#page-6-1), where θ takes only discrete values of $\{0, \pi\}$. It follows that \mathcal{L} and $\mathcal{U}^{\hat{A}}$ can be simultaneously diagonalized which is more transparent by vectorizing diagonalized, which is more transparent by vectorizing the density matrix ρ and rewriting the Liouvillian as: \mathcal{L} = $-i\left[\hat{H}\otimes\mathbb{I}-\mathbb{I}\otimes\hat{H}^T+\sum_{\mu}\gamma_{\mu}(2\hat{K}_{\mu}\otimes\hat{K}_{\mu}^*-\hat{K}_{\mu}^{\dagger}\hat{K}_{\mu}\otimes\mathbb{I}-\mathbb{I}\otimes\hat{K}_{\mu}^T\hat{K}_{\mu}^*\right].$
In this enlarged Hilbert space, the Liouvillian *f*, can be cast In this enlarged Hilbert space, the Liouvillian $\mathcal L$ can be cast into a block-diagonal matrix, where the diagonal blocks correspond to symmetry sectors with distinct eigenvalues of $U^{\hat{A}}$. The Liouville space is thus fragmented under the $U(1)$ symmetry, and dynamics within a given symmetry sector is decoupled from others, breaking the ergodicity.

We now modify the coherent Hamiltonian $\hat{H}^{\prime} = \hat{H} + \Delta \hat{A}$, where Δ is a real parameter, leading to a new Liouvillian \mathcal{L}' , Apparently, \mathcal{L}' has the same $U(1)$ symmetry, satifying $[\mathcal{L}', \hat{\mathcal{U}}^{\hat{A}}] = 0$. Nevertheless, the Liouvillian spectrum and steady-state degeneracy are significantly modified under \hat{I}' . steady-state degeneracy are significantly modified under L' . Consider a steady state ρ^a of L that is also an eigenstate of $qI^{\hat{A}}$ We then have $qI^{\hat{A}}I^{\alpha q} = e^{i\theta a} \rho^a$ where a denotes the $U^{\hat{A}}$. We then have $U^{\hat{A}}[\rho^a] = e^{i\theta a} \rho^a$, where *a* denotes the difference between two eigenvalues of \hat{A} . Since $U^{\hat{A}}[\rho^a] =$ difference between two eigenvalues of \hat{A} . Since $\mathcal{U}^{\hat{A}}[\rho^a] =$ $[\hat{A}, \rho^a] = a\rho^a$, which leads to $\mathcal{L}'[\rho^a] = -i\Delta a$. Hence, any \hat{A} + *i* $\theta[A, \rho^a]$ + $\frac{(i\theta)^2}{2}[A, [A, \rho^a]] + \cdots$ by definition, we have

ρ L′ , acquiring a purely imaginary eigenvalue −*i*∆*a*. Typically, ρ^a with $a \neq 0$ is driven out of the steady-state subspace of the steady states of $\mathcal L$ can have support on multiple symmetry sectors (or consist of eigenmodes with different *a*). Thus, a series of eigenmodes would emerge on the imaginary axis of the Liouvillian spectrum of \mathcal{L}' , which reduce the steady-state degeneracy, suggesting a steady-state phase transition [\[65\]](#page-6-1). More importantly, these eigenmodes are non-dissipative and give rise to persistent oscillations at long times, breaking the time-translation symmetry.

Here some remarks are in order. If the eigenvalues of \hat{A} are commensurate, one can always rescale \hat{A} , leading to integer eigenvalues. In this case, the long-time oscillations can be characterized by a single frequency $\Delta/2\pi$, corresponding to a time-crystalline order. Otherwise, when the eigenvalues of \hat{A} are non-commensurate, the long-time oscillations consist of multiple non-commensurate frequencies, corresponding to a quasi time crystals. Thus, dissipative (quasi) time crystals can be engineered based on the symmetry-induced fragmentation of the Liouville space.

Symmetry-induced dissipative time crystal.— To illustrate the general framework above, we consider, as an example, a one-dimensional dissipative lattice model of fermions, with the Liouvillian

$$
\mathcal{L}[\rho] = -i[\hat{H}, \rho] + \gamma \left(2\hat{K}\rho \hat{K}^{\dagger} - \left\{\hat{K}^{\dagger}\hat{K}, \rho\right\}\right),\tag{2}
$$

$$
\hat{H} = \gamma \hat{K}^{\dagger} \hat{K} + \Delta \hat{B}.
$$
 (3)

Here \hat{c}^{\dagger}_{i} $\int_{i}^{T} (\hat{c}_j)$ is the fermion creation (annihilation) operator on site *j*, $\hat{B} = \sum_{j=1}^{L} j \hat{c}_j^{\dagger}$ $\int_{i}^{\tau} \hat{c}_j$, *L* is the total number of lattice sites, and the jump operator $\hat{K} = \sum_{j=1}^{L-1} \hat{c}_j^{\dagger}$ $\int_{i}^{T} \hat{c}_{j+1}$ collectively shifts the atoms leftward (toward larger site index). The model can be implemented in an atom-cavity hybrid configuration, and ex-hibits the Liouvillian skin effect [\[59–](#page-6-2)[64\]](#page-6-0)—the geometry of the Liouvillian spectrum on the complex plane depends on the boundary condition, and the steady states are localized toward the left boundary as opposed to evenly distributed under the periodic boundary condition [\[64,](#page-6-0) [66\]](#page-6-3).

For the remainder of this work, we only consider the open boundary condition, under which $[\hat{K}, \hat{K}^{\dagger}] \neq 0$, and both the dark states of \hat{K} and the steady states of \hat{L} accumulate to the dark states of \hat{K} and the steady states of \hat{L} accumulate to the left boundary. In the light of the Liouvillian skin effect, we introduce an anti-symmetrized excitation basis in reference to the maximally localized state where all *N* fermions occupy the left-most *N* sites, forming a Fermi sea in the real space. Specifically, starting from the maximally localized state (denoted as $| \rangle$), the basis $|k_{N_e}, \ldots, k_2, k_1 \rangle$ denotes a sequential
excitation of k_1, k_2, \ldots, k_k (N is the total excitation numexcitation of k_1 , k_2 , up to k_{N_e} (N_e is the total excitation number), where the operation k_i involves moving the $(N + 1 - i)$ th fermion (from the left) k_i sites toward the right. We then have $k_1 \geq k_2 \geq k_3 \cdots \geq k_{N_e}$ due to the Pauli exclusion, and, as a convention, we omit any vanishing k_i in the notation.

When $\Delta = 0$, the Liouvillian possesses the $U(1)$ symmetry with $\mathcal{U}^{\hat{B}}[\rho] = e^{i\theta \hat{B}} \rho e^{-i\theta \hat{B}}$, and $\theta \in [0, 2\pi)$. It is
straightforward to show that the excitation basis states destraightforward to show that the excitation basis states defined above are eigenstates of \hat{B} , with $\hat{B}|k_{N_e}, \dots, k_2, k_1\rangle =$

 $B|k_{N_e}, \ldots, k_2, k_1\rangle$, and the integer eigenvalue $B = \sum_{i=1}^{N_e} k_i + N(N+1)/2$. Denoting the class of basis states with eigenvalue $N(N+1)/2$. Denoting the class of basis states with eigenvalue *B* as $|\varphi^{B}$, we have $\mathcal{U}^{\hat{B}}[\rho^{B_1,B_2}] = e^{i\theta f} \rho^{B_1,B_2}$, where $\rho^{B_1,B_2} =$
 $|\varphi^{B_1}\rangle/\langle \varphi^{B_2}|$ and $f = B_1 - B_2$. Importantly the Liouvillian f . $|\varphi^{B_1}\rangle\langle\varphi^{B_2}|$ and $f = B_1 - B_2$. Importantly, the Liouvillian \mathcal{L}
is block-diagonal in the basis of vectorized $I_0^{B_1, B_2}$, with each is block-diagonal in the basis of vectorized $\{\rho^{B_1,B_2}\}\$, with each diagonal block labeled by f [see Fig. 1(a)], and characterized diagonal block labeled by *f* [see Fig. [1\(](#page-1-0)a)], and characterized by the corresponding eigenvalue $e^{i\theta f}$ of $\mathcal{U}^{\hat{B}}$. Since f takes integer values in the range of $[-N(L - N), N(L - N)]$, there are altogether $2N(L - N) + 1$ diagonal blocks. Such a symmetryinduced fragmentation also leaves its mark in the steady-state subspace. Specifically, a general steady state of $\mathcal L$ can be expressed in the basis of $\rho_s^{B_1,B_2} = |D^{B_1}\rangle\langle D^{B_2}|$, where the dark
states $|D^{B_1}\rangle$ satisfy $\hat{\mathcal{K}}|D^{B_1}\rangle - 0$ and $\hat{\mathcal{R}}|D^{B_1}\rangle - \mathcal{R}|D^{B_1}\rangle$. Just as α^a states $|D^{B_i}\rangle$ satisfy $\hat{K}|D^{B_i}\rangle = 0$ and $\hat{B}|D^{B_i}\rangle = B_i|D^{B_i}\rangle$. Just as ρ^a *^Bi*⟩. Just as ρ in the general framework, $\rho_s^{B_1,B_2}$ is a simultaneous eigenvector
of Γ and $\mathcal{H}^{\hat{B}}$ with $\Gamma[\rho_s^{B_1,B_2}] = 0$ and $\mathcal{H}^{\hat{B}}[\rho_s^{B_1,B_2}] = a^{i\theta f} \rho_s^{B_1,B_2}$ of L and $\mathcal{U}^{\hat{B}}$, with $\mathcal{L}[\rho_{s}^{B_1,B_2}] = 0$ and $\mathcal{U}^{\hat{B}}[\rho_{s}^{B_1,B_2}] = e^{i\theta f} \rho_{s}^{B_1,B_2}$.
In other words, a steady state is generally a superposition of In other words, a steady state is generally a superposition of eigenmodes $\rho_s^{B_1,B_2}$ from different symmetry sectors (or different diagonal blocks of Γ) ent diagonal blocks of \mathcal{L}).

Now with $\Delta > 0$, following the results in the general framework, we have $\mathcal{L}[\rho_s^{B_1,B_2}] = -i\Delta f \rho_s^{B_1,B_2}$. As illustrated in Fig. 1(b) since $f \in \mathbb{Z}$ a series of equidistant eigenmodes Fig. [1\(](#page-1-0)b), since $f \in \mathbb{Z}$, a series of equidistant eigenmodes emerge along the imaginary axis, their eigenvalues being multiples of *i* Δ . For *L*, $N \gg 1$, the number of these non-dissipative modes is exponentially large in *N* [\[64\]](#page-6-0). These eigenmodes are responsible for the time-crystalline order, manifesting themselves as persistent oscillations with frequency $\Delta/2\pi$ in the long-time dynamics. This is apparent in Fig. $1(c)(d)$ $1(c)(d)$, where we plot the evolution of the correlation function $\langle \hat{c}_3^\dagger \rangle$ $\frac{1}{3}\hat{c}_2 + \hat{c}_2^{\dagger}$ $\begin{matrix} 7 \\ 2 \end{matrix}$ ĉ₃ \rangle , for an initial state of an equal-weight superposition of all excitation basis states. Note that, following the general framework, one can also devise dissipative time crystals where the symmetry generator \hat{A} involves many-body interactions [\[66\]](#page-6-3).

Breaking the symmetry: prethermal time crystal.— An outstanding feature of time crystals is their robustness against perturbations. While symmetry is crucial for the time crystal discussed above, it is natural to question its fate in the presence of symmetry-breaking perturbations.

For this purpose, we consider adding a symmetry-breaking term $\hat{H}_1 = s(\hat{K} + \hat{K}^{\dagger})$ to the coherent Hamiltonian, where $s > 0$. As a result, the Liouvillian becomes tri-block diagonal, as illustrated in Fig. [2\(](#page-2-0)a). While the previously nondissipative eigenmodes no longer stay on the imaginary axis, they do not deviate much on the complex plane even for a considerably large symmetry-breaking term [see Fig. [2\(](#page-2-0)b)]. More quantitatively, we plot the real and imaginary components of the eigenvalues of the perturbed eigenmodes as functions of *s* in Fig. $2(c)$ $2(c)$ and (d), respectively. From the numerical fit, we find that the Liouvillian eigenvalue λ_f nearby $-i\Delta f$ satisfies a higher-order power-law scaling

$$
\lambda_f + i\Delta f \sim s^p,\tag{4}
$$

with the exponent p dependent on f , or the location of the eigenmode along the imaginary axis. While the small real components suggest the domination of the long-time dynamics by these eigenmodes, the small deviation from −*i*∆*f* implies negligible frequency shift of the oscillations.

FIG. 2. Prethermal time crystal. (a) Tri-diagonal Liouvillian matrix under symmetry-breaking perturbations. The red and green boxes respectively indicate the diagonal $(f_i = f_j)$ and sub-diagonal $(|f_i - f_j| = 1)$ blocks. (b) The Liouvillian spectrum and its zoomed-in view (inset) near the imaginary axis. (c)(d) The real and imaginary eigenvalue components of the weakly-dissipative eigenmodes [in the inset of (b)], with respect to their corresponding unperturbed values (−*i*∆*f*). The dashed lines are numerical fits with the power-law scaling $\lambda_f + i\Delta f \sim s^p$, where $p = 2, 4, 6, 8$ from top to bottom. For (a)(b), we take $I = 8$, $N = 4$, and $s/\gamma = 0$?; while for (c)(d), $I = 10$ and we take $L = 8$, $N = 4$, and $s/\gamma = 0.2$; while for (c)(d), $L = 10$ and *N* = 5. For all calculations, we set Δ/γ = 0.2.

To understand the higher-order scaling behavior, we adopt the Liouvillian perturbation formalism developed in Ref. [\[64\]](#page-6-0), and rewrite the Liovillian as $\mathcal{L} = \mathcal{L}_0 + s\mathcal{L}_1$, where $\mathcal{L}_1[\rho] =$ $-i[\hat{K} + \hat{K}^{\dagger}, \rho]$. Denoting $\rho_0 = \rho_s^{B_1, B_2}$ as the steady state of \mathcal{L}_0
in the symmetry sector labeled by $f = B_1 - B_2$ we further in the symmetry sector labeled by $f = B_1 - B_2$, we further construct the density matrix in a perturbative fashion

$$
\rho = \rho_0 + s^n \sum_n \rho_n,\tag{5}
$$

where ρ_n is to be determined. In the following, we first show that an exact non-dissipative eigenmode of $\mathcal L$ can be constructed following Eq. [\(5\)](#page-2-1) in the thermodynamic limit $L, N \rightarrow \infty$. This would help us to understand the more relevant case of finite *L* and *N*.

A key observation here is that each excitation basis state $|k_{N_e}, \ldots, k_1\rangle$ can be mapped to an irreducible representation
of the permutation group. This is because the operations of the permutation group. This is because the operations $\hat{K}|k_{N_e}, \dots, k_1\rangle$ and $\hat{K}^{\dagger}|k_{N_e}, \dots, k_1\rangle$ respectively correspond to the restriction and induction of the irreducible representation with restriction and induction of the irreducible representation with the partition $[k_{N_e}, \ldots, k_1]$ [\[66\]](#page-6-3). Such a structure is illustrated in Fir 3(a) where the expansion coefficients (marked in red) of Fig. [3\(](#page-3-0)a), where the expansion coefficients (marked in red) of $(\hat{K}^{\dagger})^n$ \rangle over the excitation basis correspond to the dimensions of the corresponding irreducible representation [\[66\]](#page-6-3). The correspondence enables the representation of the excitation basis using the Young diagrams, and, crucially, the branching rules

FIG. 3. Mapping to the permutation group. (a) Excited states, in the excitation basis, generated from the dark state $|\rangle$ by \hat{K}^{\dagger} . The expansion coefficients (red) coincide with the dimensions of the corresponding irreducible representation (of the permutation group). An example of the Young-diagram correspondence of the excitation basis is given in the upper right corner, with which the structure in (a) corresponds to the Young's lattice [\[66\]](#page-6-3). (b) Relation of the exponent $p = 2l^{D}$ ^{*D_B* $)$ +2 with *B* for *L* = 10 and *N* = 5, obtained analytically from} the group analysis. (c) Maximum exponent p_m for eigenmodes characterized by f , numerically calculated for different *N* and $L = 10$. For (b)(c), we take $s/\gamma = 0.2$, and $\Delta/\gamma = 0.2$.

of the permutation group guarantees the relation [\[66\]](#page-6-3)

$$
\hat{K}(\hat{K}^{\dagger})^{n} \mid \ \rangle = n(\hat{K}^{\dagger})^{n-1} \mid \ \rangle, \tag{6}
$$

which, as illustrated in Fig. $3(a)$ $3(a)$, is mapped to the restriction of the irreducible representations. Furthermore, we observe from simple examples and numerical calculations that, if we replace $|\rangle$ with other dark states $|D\rangle$ of \hat{K} , the mapping above still holds, and we still have $\hat{K}(\hat{K}^{\dagger})^n|D\rangle = n(\hat{K}^{\dagger})^{n-1}|D\rangle$ [\[66\]](#page-6-3).

With these understandings, we have $\hat{K}(\hat{K}^{\dagger})^n|D^{B_{1,2}}\rangle$ = $n(\hat{K}^{\dagger})^{n-1} |D^{B_{1,2}}\rangle$ for any $n \in \mathbb{N}^+$. Then, by setting $\rho_n = \sum_{j=0}^n \alpha_n^{(j)} \hat{K}^{\dagger j} |D^{B_1} \rangle \langle D^{B_2} | \hat{K}^{n-j}$, where $\alpha_n^{(j)} =$
 $\frac{a_n^{(j)} - a_{n-1}^{(j)}}{a_n^{(j)}} = \frac{a_n^{(j)} - a_{n-1}^{(j)}}{a_n^{(j)}}$ $i^{n-2j} s^n/((1-i)\gamma - i\Delta)^{n-j}((1+i)\gamma + i\Delta)^j(n-j)!j!$, we have the recurrence relation [\[66\]](#page-6-3)

$$
\mathcal{L}_0[\rho_n] + i\Delta f \rho_n = -s \mathcal{L}_1[\rho_{n-1}],\tag{7}
$$

which holds for all $n \in \mathbb{N}^+$ given $L, N \to \infty$. It is then straight-
forward to show that $0 = \alpha_0 + s^n \sum_{i=1}^{\infty} a_i$ is an exact nonforward to show that $\rho = \rho_0 + s^n \sum_{n=1}^{\infty} \rho_n$ is an exact non-
dissinative eigenmode of f with the eigenvalue $-i\Delta f$ since dissipative eigenmode of $\mathcal L$ with the eigenvalue $-i\Delta f$, since the perturbative expansion cancels at every order of *s*, thanks to Eq. [\(7\)](#page-3-1).

$ D_1\rangle$, $ D_2\rangle$ of the eigenmode $ D_1\rangle\langle D_2 $	f	Ν	$(l_N^{ D_1\rangle}, l_L^{ D_1\rangle}$	p_m
$ 111\rangle - 22\rangle + 3\rangle$ $ 11\rangle - 2\rangle$	1	3	(0, 4)	2
		4	(1, 3)	$\overline{4}$
		5	(2, 2)	6
$ 11\rangle - 2\rangle$ \vert)	2	$\overline{2}$	(0, 6)	$\overline{2}$
		3	(1, 5)	$\overline{4}$
		$\overline{4}$	(2, 4)	6
		5	(3, 3)	8
$ 111\rangle - 22\rangle + 3\rangle$ \vert)	3	3	(0, 4)	\overline{c}
		4	(1, 3)	$\overline{4}$
		5	(2, 2)	6
$ 22\rangle - 13\rangle + 4\rangle$ \vert)	4	$\overline{2}$	(0, 4)	$\overline{2}$
		3	(1, 3)	$\overline{4}$
		$\overline{4}$	(2, 2)	6
		5	(3, 1)	$\overline{4}$
$ 122\rangle - 113\rangle + 23\rangle + 2 14\rangle - 2 5\rangle$	5	3	(0, 2)	\overline{c}
\mapsto		4	(1, 1)	4
$ 44\rangle - 134\rangle + 224\rangle + 1133\rangle - 1223\rangle$	5	5	(1, 1)	4
$ 111\rangle - 22\rangle + 3\rangle$				
$ 33\rangle - 24\rangle + 15\rangle - 6\rangle$	6	$\overline{2}$	(0, 2)	$\overline{2}$
		3	(1, 1)	$\overline{\mathcal{L}}$
$ 123\rangle - 114\rangle + 24\rangle - 222\rangle - 2 33\rangle$	6	4	(1, 2)	$\overline{4}$
\mapsto		5	(2,1)	4

TABLE I. Analytic results from the permutation-group analysis, for *L* = 10 and $|f| \le 6$.

For finite *L* or *N*, however, the mapping ceases to apply at a sufficiently large *n*, because of the limited lattice size or atom number. Hence a cutoff in Eq. [\(7\)](#page-3-1) and the perturbative series [\(5\)](#page-2-1) becomes necessary. Generally, for any dark state $|D\rangle$ of \hat{K} , we define $l_N^{|D\rangle} = \max\{j|\langle D|\hat{c}_j^{\dagger}$ $\int_{j}^{T} \hat{c}_j |D\rangle = 1$, labeling the site index to the left of which the Fermi sea is intact (without hole excitations). We also define $l_L^{|D\rangle}$ = $\min\{j|\langle D|\hat{c}_{L-j}^{\dagger}c_{L-j}|D\rangle > 0\}$, labeling the site index to the right of which there are no fermion excitations. We then find of which there are no fermion excitations. We then find that the relation $\hat{K}(\hat{K}^{\dagger})^n|D\rangle = n(\hat{K}^{\dagger})^{n-1}|D\rangle$ holds only up to $n \leq l^{(D)} = \min(l^{(D)}_N, l^{(D)}_L)$. It follows that, for the steady state ρ_0 of \mathcal{L}_0 , by defining $l = \min(l^{|D^{B_1}\rangle}, l^{|D^{B_2}\rangle})$, the truncated pertur-
bative series $Q = Q_0 + s^n \nabla^l$, Q_0 is an approximate eigenbative series $\rho = \rho_0 + s^n \sum_{n=1}^l \rho_n$ is an approximate eigen-
mode of f, with the approximate eigenvalue $\lambda_{\alpha} + i\Delta f$. mode of \mathcal{L} , with the approximate eigenvalue $\lambda_f + i\Delta f =$ $Tr(\mathcal{L}[\rho]\rho) + i\Delta f \sim s^p$, where $p = 2l + 2$ [\[64,](#page-6-0) [66\]](#page-6-3).
The nower-law scaling of the eigenvalues suggest

The power-law scaling of the eigenvalues suggest that these eigenmodes are weakly dissipative, which gives rise to rich prethermal dynamics. For instance, when *L* = 2*N*, the steady state $\rho_s = |11\rangle\langle |-|2\rangle\langle |$ of \mathcal{L}_0 acquires an eigen-
value $\lambda + i2\Delta \approx s^{2N-2}$ under the symmetry-breaking perturvalue $\lambda + i2\Delta \sim s^{2N-2}$ under the symmetry-breaking pertur-
bation (see Table I). The thermalization time of the mode is bation (see Table. [I\)](#page-3-2). The thermalization time of the mode is therefore $\sim s^{2-2N}$, which increases exponentially with increasing *N*, and manifests itself in the dynamics as oscillations at the frequency Δ/π . Thus, depending on the initial state, a prethermal time-crystalline behavior emerges, with distinct oscillatory frequencies at different times of the steady-state approaching dynamics.

In Fig. [3\(](#page-3-0)b), we show the analytically calculated exponent *p* for all dark states in a chain of $L = 10$ at half filling $N = 5$. Generally, *p* increases with decreasing *B*, meaning localized states (with smaller *B*) tend to be more robust dy-

FIG. 4. Stage-wise dynamics of the prethermal time crystal. (a) Time evolution of $P_{|f|}$ (see main text for definition). (b) Numerically simulated evolution of $\langle \hat{K}_2 + \hat{K}_2^{\dagger} \rangle$, where $\hat{K}_2 = \sum_{j=1}^{L-2} \hat{c}_j^{\dagger} \hat{c}_{j+2}$. In (a)(b), stages with different dominant eigenmodes are separated by gray vertical lines. (c) Oscillations zoomed in, starting at different times $t_0\gamma = 10^3$ (upper), $t_0\gamma = 10^5$ (middle) and $t_0\gamma = 10^7$ (lower), which are labeled
with vertical lines in (a)(b) with the corresponding colors. The oscilwith vertical lines in (a)(b) with the corresponding colors. The oscillation period is given by $T_f = 2\pi/f\Delta$, with $f = 1, 3, 2$, respectively, for upper, middle and lower panels, consistent with (a). For all figures, we take $L = 10$, $N = 4$, $s/\gamma = 0.2$, and $\Delta/\gamma = 0.2$.

namically, as they feature a smaller Liouvillian gap $\text{Re}\lambda_f$. On
the other hand, for a given f, there can be many weakly disthe other hand, for a given f , there can be many weakly dissipative eigenmodes with different exponents *p*. We thus define p_m as the maximum among them. In Fig. [3\(](#page-3-0)c), we plot p_m as functions of *f* and *N* for a lattice with $L = 10$, which are obtained by numerically fitting the Liouvillian spectrum. The numerically calculated p_m (and the corresponding eigenmodes) are consistent with the analytical results, tabulated in Table. [I.](#page-3-2) Generally, for a fixed *N* (fixed *f*), *p^m* becomes larger for smaller *f* (larger *N*). This means that: i) in a system with fixed atom number, oscillations with smaller frequencies dominate at later times; ii) with increasing atom number, oscillations with the same frequency emerge later.

Prethermal dynamics.— We numerically confirm the conclusions above by simulating the open-system dynamics for $L = 10$ and $N = 4$ in Fig. [4.](#page-4-7) We initialize the system such that the initial density matrix has support on eigenmodes with $f = \pm 1$ ($p_m = 2$), $f = \pm 2$ ($p_m = 6$), and $f = \pm 3$ $(p_m = 4)$. The overlap of the time-dependent density matrix with eigenmodes nearby the eigenvalue −*i*∆*f* is defined as $P_{|f|} = \sum_{|B_1 - B_2| = |f|} \text{Tr}(\rho \rho_s^{B_1, B_2})$. With different p_m , the eigen-
modes dominate at different times, leading to the stage-wise modes dominate at different times, leading to the stage-wise oscillatory dynamics in Fig. $4(a)(b)$ $4(a)(b)$. The oscillations within different time scales feature distinct frequencies dictated by *f* [see Fig. [4\(](#page-4-7)c)], consistent with the theoretical predictions in Table. [I.](#page-3-2)

Discussion.— We show that time crystals can be engineered

based on a *U*(1)-symmetry-induced fragmentation of the Liouville space. Combined with the Liouvillian skin effect, such a scheme further opens up the possibility of generating prethermal time crystals in the symmetry-broken regime, where an elegant permutation-group representation of the fermionic excitations plays a key role. Such a connection essentially relies on the limited Hilbert space available to the excitations of identical fermions above the real-space Fermi sea. For future studies, it would be interesting to explore what new perspectives can derive from other forms of symmetries.

We thank Heran Wang and Zhong Wang for helpful discussions. This work is supported by the National Natural Science Foundation of China (Grant No. 12374479), and by the Innovation Program for Quantum Science and Technology (Grant No. 2021ZD0301205).

[∗] wyiz@ustc.edu.cn

- [1] F. Wilczek, Quantum Time Crystals, Phys. Rev. Lett. 109, 160401 (2012).
- [2] K. Sacha and J. Zakrzewski, Time crystals: a review, Rep. Prog. Phys. 81, 016401 (2018).
- [3] M. P. Zaletel, M. Lukin, C. Monroe, C. Nayak, F. Wilczek, and N. Y. Yao, Colloquium: Quantum and classical discrete time crystals, Rev. Mod. Phys. 95, 031001 (2023).
- [4] D. V. Else, C. Monroe, C. Nayak, and N. Y. Yao, Discrete time crystals, Annu. Rev. Condens. Matter Phys. 11, 467 (2020).
- [5] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, Phase Structure of Driven Quantum Systems, Phys. Rev. Lett. 116, 250401 (2016).
- [6] D. V. Else, B. Bauer, and C. Nayak, Floquet Time Crystals, Phys. Rev. Lett. 117, 090402 (2016).
- [7] S. Choi, J. Choi, R. Landig, G. Kucsko, H. Zhou, J. Isoya, F. Jelezko, S. Onoda, H. Sumiya, V. Khemani, C. von Keyserlingk, N. Y. Yao, E. Demler, and M. D. Lukin, Observation of discrete time-crystalline order in a disordered dipolar many-body system, Nature 543, 221 (2017).
- [8] J. Randall, C. E. Bradley, F. V. van der Gronden, A. Galicia, M. H. Abobeih, M. Markham, D. J. Twitchen, F. Machado, N. Y. Yao, and T. H. Taminiau, Many-body localized discrete time crystal with a programmable spin-based quantum simulator, Science 374, 1474 (2021).
- [9] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I.-D. Potirniche, A. C. Potter, A. Vishwanath, et al., Observation of a discrete time crystal, Nature 543, 217 (2017).
- [10] P. Frey and S. Rachel, Realization of a discrete time crystal on 57 qubits of a quantum computer, Sci. Adv. 8, eabm7652 (2022).
- [11] X. Mi *et al.*, Observation of Time-Crystalline Eigenstate Order on a Quantum Processor, Nature 601, 531-536 (2022).
- [12] S. Liu, S.-X. Zhang, C.-Y. Hsieh, S. Zhang, and H. Yao, Discrete Time Crystal Enabled by Stark Many-Body Localization, Phys. Rev. Lett. 130, 120403 (2023).
- [13] D. V. Else, B. Bauer, and C. Nayak, Prethermal Phases of Matter Protected by Time-Translation Symmetry, Phys. Rev. X 7, 011026 (2017).
- [14] T.-S. Zeng and D. N. Sheng, Prethermal time crystals in a onedimensional periodically driven Floquet system, Phys. Rev. B 96, 094202 (2017).
- [15] A. Kyprianidis, F. Machado, W. Morong, P. Becker, K. S.

Collins, D. V. Else, L. Feng, P. W. Hess, C. Nayak, G. Pagano *et al.*, Observation of a prethermal discrete time crystal, Science 372, 1192 (2021).

- [16] F. Machado, Q. Zhuang, N. Y. Yao, and M. P. Zaletel, Absolutely Stable Time Crystals at Finite Temperature, Phys. Rev. Lett. 131, 180402 (2023).
- [17] M. Yue and Z. Cai, Prethermal Time-Crystalline Spin Ice and Monopole Confinement in a Driven Magnet, Phys. Rev. Lett. 131, 056502 (2023).
- [18] D. Vu and S. Das Sarma, Dissipative Prethermal Discrete Time Crystal, Phys. Rev. Lett. 130, 130401 (2023).
- [19] A. Pizzi, D. Malz, G. De Tomasi, J. Knolle, and A. Nunnenkamp, Time crystallinity and finite-size effects in clean Floquet systems, Phys. Rev. B 102, 214207 (2020).
- [20] W. Deng and Z.-C. Yang, Using models with static quantum many-body scars to generate time-crystalline behavior under periodic driving, Phys. Rev. B 108, 205129 (2023).
- [21] Z. Gong, R. Hamazaki, and M. Ueda, Discrete Time-Crystalline Order in Cavity and Circuit QED Systems, Phys. Rev. Lett. 120, 040404 (2018).
- [22] K. Chinzei and T. N. Ikeda, Time Crystals Protected by Floquet Dynamical Symmetry in Hubbard Models, Phys. Rev. Lett. 125, 060601 (2020).
- [23] X. Yang and Z. Cai, Dynamical transitions and critical behavior between discrete time crystal phases, Phys. Rev. Lett. 126, 020602 (2021).
- [24] Z.-A. Hu, B. Fu, X. Li, and S.-Q. Shen, Solvable model for discrete time crystal enforced by nonsymmorphic dynamical symmetry, Phys. Rev. Res. 5, L032024 (2023).
- [25] P. Xu and T.-S. Deng, Boundary discrete time crystals induced by topological superconductors in solvable spin chains, Phys. Rev. B 107, 104301 (2023).
- [26] F. M. Gambetta, F. Carollo, M. Marcuzzi, J. P. Garrahan, and I. Lesanovsky, Discrete Time Crystals in the Absence of Manifest Symmetries or Disorder in Open Quantum Systems, Phys. Rev. Lett. 122, 015701 (2019).
- [27] H. Kessler, J. G. Cosme, M. Hemmerling, L. Mathey, and A. Hemmerich, Emergent limit cycles and time crystal dynamics in an atom-cavity system, Phys. Rev. A 99, 053605 (2019).
- [28] P. Kongkhambut *et al.*, Observation of a Continuous Time Crystal, Science 377, 670-673 (2022).
- [29] R. Mattes, I. Lesanovsky, and F. Carollo, Entangled time-crystal phase in an open quantum light-matter system, Phys. Rev. A 108, 062216 (2023).
- [30] X. Wu, Z. Wang, F. Yang, R. Gao, C. Liang, M. K. Tey, X. Li, T. Pohl, and L. You, Dissipative time crystal in a strongly interacting Rydberg gas, Nature Physics 20, 1389-1394 (2024).
- [31] F. Iemini, A. Russomanno, J. Keeling, M. Schirò, M. Dalmonte, and R. Fazio, Boundary Time Crystals, Phys. Rev. Lett. 121, 035301 (2018).
- [32] G. Piccitto, M. Wauters, F. Nori, and N. Shammah, Symmetries and conserved quantities of boundary time crystals in generalized spin models, Phys. Rev. B 104, 014307 (2021).
- [33] L. F. dos Prazeres, L. da S. Souza, and F. Iemini, Boundary time crystals in collective *d*-level systems, Phys. Rev. B 103, 184308 (2021).
- [34] G. Passarelli, P. Lucignano, R. Fazio, and A. Russomanno, Dissipative time crystals with long-range Lindbladians, Phys. Rev. B 106, 224308 (2022).
- [35] A. Cabot, L. S. Muhle, F. Carollo, and I. Lesanovsky, Quantum trajectories of dissipative time crystals, Phys. Rev. A 108, L041303 (2023).
- [36] C. Booker, B. Buča, and D. Jaksch, Non-stationarity and dissipative time crystals: spectral properties and finite-size effects,

New J. Phys. 22, 085007 (2020).

- [37] K. Bull, A. Hallam, Z. Papić, and I. Martin, Tuning between Continuous Time Crystals and Many-Body Scars in Long-Range *XYZ* Spin Chains, Phys. Rev. Lett. 129, 140602 (2022).
- [38] M. Krishna, P. Solanki, M. Hajdušek, and S. Vinjanampathy, Measurement-Induced Continuous Time Crystals, Phys. Rev. Lett. 130, 150401 (2023).
- [39] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papic, Weak ergodicity breaking from quantum many-body ´ scars, Nature Physics 14, 745-749 (2018).
- [40] H. Keßler, J. G. Cosme, C. Georges, L. Mathey, and A. Hemmerich, From a continuous to a discrete time crystal in a dissipative atom-cavity system, New J. Phys. 22, 085002 (2020).
- [41] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, Nat. Phys. 17, 675 (2021).
- [42] Z. Papić, Weak ergodicity breaking through the lens of quantum entanglement, in *Entanglement in Spin Chains: From Theory to Quantum Technology Applications*, edited by A. Bayat, S. Bose, and H. Johannesson (Springer International Publishing, Cham, 2022), pp. 341-395.
- [43] S. Moudgalya, B. A. Bernevig, and N. Regnault, Quantum many-body scars and Hilbert space fragmentation: A review of exact results, Rep. Prog. Phys. 85, 086501 (2022).
- [44] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, Quantum many-body scars: A quasiparticle perspective, Annu. Rev. Condens. Matter Phys. 14, 443 (2023).
- [45] Q. Hummel, K. Richter, and P. Schlagheck, Genuine Many-Body Quantum Scars along Unstable Modes in Bose-Hubbard Systems, Phys. Rev. Lett. 130, 250402 (2023).
- [46] V. Khemani, M. Hermele, and R. Nandkishore, Localization from Hilbert space shattering: From theory to physical realizations, Phys. Rev. B 101, 174204 (2020).
- [47] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, Ergodicity breaking arising from Hilbert space fragmentation in dipole-conserving Hamiltonians, Phys. Rev. X 10, 011047 (2020).
- [48] Z.-C. Yang, F. Liu, A. V. Gorshkov, and T. Iadecola, Hilbertspace fragmentation from strict confinement, Phys. Rev. Lett. 124, 207602 (2020).
- [49] C. M. Langlett and S. Xu, Hilbert space fragmentation and exact scars of generalized Fredkin spin chains, Phys. Rev. B 103, L220304 (2021).
- [50] F. M. Surace, P. P. Mazza, G. Giudici, A. Lerose, A. Gambassi, and M. Dalmonte, Hilbert space fragmentation in a 2D quantum spin system with subsystem symmetries, SciPost Phys. 13, 098 (2022).
- [51] A. Yoshinaga, H. Hakoshima, and T. Imoto, Hilbert space fragmentation in open quantum systems, Phys. Rev. Res. 5, 043239 (2023).
- [52] G. Francica and L. Dell'Anna, Hilbert space fragmentation in a long-range system, Phys. Rev. B 108, 045127 (2023).
- [53] A. Chattopadhyay, B. Mukherjee, K. Sengupta, and A. Sen, Strong Hilbert space fragmentation via emergent quantum drums in two dimensions, SciPost Phys. 14, 146 (2023).
- [54] S. Yao and Z. Wang, Edge States and Topological Invariants of Non-Hermitian Systems, Phys. Rev. Lett. 121, 086803 (2018).
- [55] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal bulk-boundary correspondence in non-Hermitian systems, Phys. Rev. Lett. 121, 026808 (2018).
- [56] V. M. Martinez Alvarez, J. E. Barrios Vargas, and L. E. F. Foa Torres, Non-Hermitian robust edge states in one dimension: Anomalous localization and eigenspace condensation at exceptional points, Phys. Rev. B 97, 121401(R) (2018).
- [57] Z. Yang, K. Zhang, C. Fang, and J. Hu, Non-Hermitian bulk-

boundary correspondence and auxiliary generalized Brillouin zone theory, Phys. Rev. Lett. 125, 226402 (2020).

- [58] X. Zhang, T. Zhang, M. H. Lu, and Y. F. Chen, A review on non-Hermitian skin e ffect, Adv. Phys. X 7, 1 (2022).
- [59] F. Song, S. Yao, and Z. Wang, Non-Hermitian Skin E ffect and Chiral Damping in Open Quantum Systems, Phys. Rev. Lett. 123, 170401 (2019).
- [60] T. Haga, M. Nakagawa, R. Hamazaki, and M. Ueda, Liouvillian Skin E ffect: Slowing Down of Relaxation Processes without Gap Closing, Phys. Rev. Lett. 127, 070402 (2021).
- [61] F. Yang, Q.-D. Jiang, and E. J. Bergholtz, Liouvillian skin e ffect in an exactly solvable model, Phys. Rev. Res. 4, 023160 (2022).
- [62] S. Hamanaka, K. Yamamoto, and T. Yoshida, Interactioninduced Liouvillian skin effect in a fermionic chain with a twobody loss, Phys. Rev. B 108, 155114 (2023).
- [63] Z. Wang, Y. Lu, Y. Peng, R. Qi, Y. Wang, and J. Jie, Accelerating relaxation dynamics in open quantum systems with Liouvillian skin effect, Phys. Rev. B 108, 054313 (2023).
- [64] H. Li, H. Wu, W. Zheng, and W. Yi, Many-body non-Hermitian skin e ffect under dynamic gauge coupling, Phys. Rev. Res. 5 , 033173 (2023).
- [65] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, Spectral theory of Liouvillians for dissipative phase transitions, Phys. Rev. A 98, 042118 (2018).
- [66] See Supplemental Material for details, which includes Refs. [\[67](#page-6-4), [68\]](#page-6-5).
- [67] B. E. Sagan, The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions (2nd Ed.), Graduate Texts in Math. 203, Springer, New York, 2001.
- [68] A. M. Vershik, A new approach to the representation theory of the symmetric groups. III. Induced representations and the Frobenius-Young correspondence, Mosc. Math. J. 6, 567 (2006).

Supplemental Material for "Symmetry-induced fragmentation and dissipative time crystal"

In this Supplemental Material, we provide details on an alternative example of symmetry-induced fragmentation and time crystal, the implementation scheme of the model in the main text, the Liouvillian skin effect of the model, the mapping between excitations and irreducible representations of the permutation group, and details on the Liouvillian perturbation approach.

An alternative example of symmetry-induced fragmentation and time crystal

In the main text, we devise a dissipative time crystal based on the $U(1)$ symmetry with a single-particle symmetry generator. Here we illustrate a more involved case where the symmetry generator features many-body interactions. We consider the following Liouvillian in a one-dimensional spin-1/2 chain under the periodic boundary condition (setting $\hbar = 1$)

$$
\frac{d\rho}{dt} = \mathcal{L}[\rho] = -i[\hat{H}, \rho] + \sum_{j=1}^{N} \gamma \left(2\hat{L}_{j}\rho \hat{L}_{j}^{\dagger} - \left\{ \hat{L}_{j}^{\dagger} \hat{L}_{j}, \rho \right\} \right),\tag{S1}
$$

where the coherent Hamiltonian and jump operators read

$$
\hat{H} = \frac{U}{4} \sum_{j} (\sigma_{j}^{z} + 1)(\sigma_{j+1}^{z} + 1), \quad \hat{L}_{j} = \frac{\gamma}{4} (1 - \sigma_{j-1}^{z}) \sigma_{j}^{+} (1 - \sigma_{j+1}^{z}).
$$
\n(S2)

Here *j* indicates the site index of the spin, *N* is the lattice size, and *U* and γ are the interaction and dissipation rates, respectively. The Liouvillian [\(S1\)](#page-7-0) has the following *U*(1) symmetry

$$
[\mathcal{L}, \mathcal{U}^{\hat{H}}] = 0, \quad \text{with} \quad \mathcal{U}^{\hat{H}}[\rho] = e^{i\theta \hat{H}} \rho e^{-i\theta \hat{H}}, \tag{S3}
$$

where $\theta \in [0, 2\pi)$, and \hat{H} also plays the role of the symmetry generator.

The eigenstates of \hat{H} are product states of spin-up $|\uparrow\rangle$ and spin-down $|\downarrow\rangle$ states along the chain. For a given product state, its eigenvalue is determined by counting the number of adjacent spin-up pairs |↑↑⟩. The eigenvalue thus takes integer values in the range of [0, *N*]. Following the derivations in the main text, a series of equidistant Liouvillian eigenmodes should emerge on the imaginary axis with $\lambda_n = inU$ ($n \in \mathbb{Z}$). This is illustrated in Fig. [S1\(](#page-1-0)a) for a chain with $N = 6$. These non-dissipative eigenmodes give rise to persistent oscillations in the long-time dynamics. This is shown in Fig. $S1(b)(c)$ $S1(b)(c)$. For our numerical simulations, we initialize the system in a pure state, with the density matrix $\rho(t = 0) = (\phi_1) \langle \phi_1 | + |\phi_1 \rangle \langle \phi_2 | + |\phi_2 \rangle \langle \phi_1 | + |\phi_2 \rangle \langle \phi_2 |)/2$, where $|\phi_1\rangle = | \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \rangle$, and $|\phi_2\rangle = | \downarrow \uparrow \uparrow \uparrow \uparrow \uparrow \rangle$. Since $|\phi_1\rangle$ and $|\phi_2\rangle$ are eigenstates of \mathcal{U}^H with eigenvalues $e^{2i\theta}$ and $e^{-2i\theta}$, respectively, the oscillation frequency is U/π cons oscillation frequency is U/π , consistent with the numerical results in Fig. $S1(b)(c)$ $S1(b)(c)$.

FIG. S1. (a) Liouvillian spectrum for Eq. [\(S1\)](#page-7-0) with $N = 6$. (b) The long-time dynamics of $\langle \sigma_1^x \rangle$. (c) A zoomed-in view of (b) over four four spectrols with different starting times of $t_0 \times 10^5$ (blue) and $t_0 \times$ periods, with different starting times of $t_0\gamma = 10^5$ (blue) and $t_0\gamma 10^7$ (green), respectively, as marked by the vertical dashed lines in (b). For all simulations we take $U = \gamma$ simulations, we take $U = \gamma$.

Implementing the tilted lattice model

The tilted lattice model in the main text can be implemented using an atom-cavity hybrid setup, similar to the one in Ref. [\[1\]](#page-12-0). As illustrated in Fig. [S2,](#page-2-0) spinless fermions in a tilted one-dimensional lattice are coupled to a cavity. While the detuning δ between adjacent sites suppresses direct inter-site hopping, two Raman-processes are introduced to induce laser-assisted hopping. One of the Raman processes is assisted by the cavity field (red dashed arrow). Given a lossy cavity, dynamics of the fermions adiabatically follows that of the cavity field, and the fermions are effectively subject to a cavity-dependent dynamic gauge potential. An additional two-photon detuning ∆ is introduced for both Raman processes, which is absent in Ref. [\[1\]](#page-12-0).

Adiabatically eliminating the cavity field and taking the tight-binding approximation, the atomic density matrix ρ is governed by the Lindblad master equation (with $\hbar = 1$)

$$
\frac{d\rho}{dt} = \mathcal{L}[\rho] = -i[\hat{H}, \rho] + \gamma \left(2\hat{K}\rho \hat{K}^{\dagger} - \left\{ \hat{K}^{\dagger} \hat{K}, \rho \right\} \right),\tag{S4}
$$

with the coherent Hamiltonian

$$
\hat{H} = -\frac{\Delta_c}{\kappa} \gamma \hat{K}^\dagger \hat{K} + s(\hat{K} + \hat{K}^\dagger) + \Delta \hat{B}.
$$
 (S5)

Here $\gamma = \kappa \lambda^2/(\Delta_c^2 + \kappa^2)$, the quantum jump operator $\hat{K} = \sum_{j=1}^{L-1} \hat{c}_j^{\dagger}$
A $/\kappa = -1$ we get the Liouvillian Eqs. (3)(4) of the main text \int_{i}^{\dagger} , Δ_c is the cavity detuning, and $\hat{B} = \sum_{j=1}^{L} j \hat{c}_j^{\dagger}$ ^{*f*}_{*c*}^{*j*}*c*_{*j*}**.** Setting $\Delta_c/\kappa = -1$, we get the Liouvillian Eqs. (3)(4) of the main text.

Liouvillian skin effect of the tilted lattice model

The titled lattice model Eqs. (3)(4) in the main text exhibits the Liouvillian skin effect. This is manifest in the difference of Liouvillian spectra under different boundary conditions, the boundary localization of the steady states under the open boundary condition, and atomic density dynamics.

First, in Fig. [S3\(](#page-3-0)a), we show the Liouvillian spectra under both the periodic boundary condition (blue) and the open boundary condition (red). The eigenspectra are drastically different under different boundary conditions, which is a key signature of the Liouvillian skin effect. In particular, under the periodic boundary condition, there are no non-dissipative eigenmodes on the imaginary axis, meaning the time crystal behavior only exists under the open boundary condition.

Second, under the open boundary condition, all dark states of \hat{K} become boundary localized, as are the steady states of the Liouvillian. In Fig. [S3\(](#page-3-0)b), we plot the spatial distribution of the dark states (solid lines), which are all localized toward the left boundary. Whereas under the open boundary, the distribution is uniform (black dashed line).

The Liouvillian skin effect also manifests as the boundary-approaching dynamics. In Fig. [S3\(](#page-3-0)c), we show the time evolution of $\langle B \rangle$ – $(N + 1)N/2$, with the system initialized in an equal-weight superposition of all excitation basis [which is the same as Fig. 1(c)(d) in the main text]. Here $\langle \hat{B} \rangle$ is an indication of the extent of localization of the time-evolved state: the state is more localized with smaller $\langle \hat{B} \rangle$. Evidently, $\langle \hat{B} \rangle$ rapidly decreases toward the steady-state value in the dynamics, which also indicates the localized nature of the steady states.

FIG. S2. Schematics for implementing the tilted lattice model in a atom-cavity hybrid setup. Here $\Omega_{1,2,3}$ are the single-photon Rabi frequencies of the laser and cavity couplings, ω_p is the frequency of the pumping laser (marked in green), ω_c is the cavity-field frequency (marked in red
dashed) $\Delta = \omega - \omega$ is the cavity detuning and *k* is the cavity decay ra dashed), $\Delta_c = \omega_c - \omega_p$ is the cavity detuning, and κ is the cavity decay rate.

FIG. S3. (a) Liouvillian spectra under the open boundary condition (red) and the periodic boundary condition (blue), respectively. (b) Spatial distribution of all the dark states under the open boundary condition (solid lines) and the periodic boundary condition (black dashed line). (c) Evolution of $\langle \hat{B} \rangle$. For all figures, we take *L* = 8, *N* = 4, and Δ/γ = 0.2.

Mapping between excitations with the permutation-group representation

We start from the following relations

$$
\hat{K}|k_{N_e}, k_{N_e-1}, \dots k_1\rangle = \sum_{i=1}^{N_e} (1 - \delta_{k_i, k_{i+1}})|p_{N_e}^{(i)}, p_{N_e-1}^{(i)}, \dots, p_1^{(i)}\rangle, \tag{S6}
$$

$$
\hat{K}^{\dagger}|k_{N_e},k_{N_e-1},\ldots,k_1\rangle = \sum_{i=1}^{N_e+1} (1-\delta_{k_{i-1},k_i})|q_{N_e+1}^{(i)},q_{N_e}^{(i)},\ldots,q_1^{(i)}\rangle.
$$
 (S7)

Here we set $k_{N_e+1} = 0$, $p_j^{(i)} = k_j - \delta_{i,j}$, $q_j^{(i)} = k_j + \delta_{i,j}$, and formally require $\delta_{k_0,k_1} = 0$. We also omit vanishing values of $p_j^{(i)}$ and $q_j^{(i)}$ after the calculation, following the convention of the excitation basis defined in the main text. Note that the Kronecker-delta functions $\delta_{i,j}$ are introduced to enforce the Pauli exclusion.

Expanding $(\hat{K}^{\dagger})^n$ onto the excitation state basis, we have $(\hat{K}^{\dagger})^n$ $\rangle = \sum_k b_{n,k} |\phi_n^k\rangle$, where $b_{n,k}$ are the expansion coefficients,
d $|d^k\rangle = |k_1|$, $|k_2|$, \ldots $|k_3|$ with the additional condition and $|\phi_n^k\rangle = |k_{N_e}, k_{N_e-1} \dots k_1\rangle$ with the additional condition $\sum_{i=1}^{N_e} k_i = n$. We further use the notation $r \nearrow k$ to denote the condition $\langle A_r^r | \hat{\mathcal{L}} \rangle$ = 1. From Eq. (57), we find $\langle \phi_{n-1}^r | \hat{K} | \phi_n^k \rangle = 1$. From Eq. [\(S7\)](#page-9-0), we find

$$
b_{n,k} = \sum_{r \nearrow k} b_{n-1,r}.\tag{S8}
$$

In the following, we show that the state $|\phi_n^k\rangle$ can be mapped to an irreducible representation of the permutation group. And, \hat{k} and \hat{k}^{\dagger} in Eqs. (S6) and (S7) respectively correspond to the restriction and crucially, the action of \hat{K} and \hat{K}^{\dagger} in Eqs. [\(S6\)](#page-9-1) and [\(S7\)](#page-9-0) respectively correspond to the restriction and induction of the irreducible representation. With this mapping, we shall prove the relation $nb_{n-1,k} = \sum_{k \nearrow r} b_{n,r}$.

First, let us briefly review the permutation group. The permutation group considers the permutation of *n* individual numbers in an array. The so-called cycle notation (a_1, a_2, \ldots, a_n) describes the permutation of the elements from the positions $a_1, a_2, a_3, \ldots, a_n$ to new positions $a_n, a_1, a_2, \ldots, a_{n-1}$. Any permutation can be denoted using multiples of the cycle notation. For example, a permutation from "123456" to "653124" can be denoted as $(3)(25)(146)$.

Consider an irreducible representation of the permutation group for a set of *n* numbers. The representation is indexed by a partition of *n*. Here different partitions indicate different ways to perform the permutation. For example, with $n = 6$, the partition [1, 2, 3] indicates the permutation with the cycle notation $(a_1)(a_2, a_3)(a_4, a_5, a_6)$. In the literature, the Young diagram is often used to denote the irreducible representation. For an irreducible representation with the partition $[\phi_n^k] = [k_{N_e}, k_{N_e-1} \dots k_1]$
satisfying $k_1 \ge k_2 \ge \dots \ge k_M$ and $\sum k_i = n$, the Young diagram consists of a total of satisfying $k_1 \geq k_2 \geq \cdots \geq k_{N_e}$ and $\sum_j k_j = n$, the Young diagram consists of a total of *n* layered boxes, with k_j boxes in the *j*th layer (counting from the bottom layer). See Fig. [S4](#page-4-7) for an illustration.

FIG. S4. Young's lattice, showing the irreducible representations of the permutation group for different *n*. The numbers in red indicate the dimensions of each irreducible representation of the corresponding Young diagram. The arrows indicate the induction of the corresponding representation.

FIG. S5. Excitations generated by *K*ˆ † from different dark states. (a) The initial dark state is |2⟩−|11⟩. (b) The initial dark state is |3⟩−|12⟩+|111⟩.

$$
\text{Res}[k_{N_e}, k_{N_e-1}, \dots k_1] = \bigoplus_{i=1}^{N_e} (1 - \delta_{k_i, k_{i+1}}) [p_{N_e}^{(i)}, p_{N_e-1}^{(i)}, \dots, p_1^{(i)}],
$$
(S9)

$$
\text{Ind}[k_{N_e}, k_{N_e-1}, \dots k_1] = \bigoplus_{i=1}^{N_e+1} [1 - \delta_{k_{i-1}, k_i}] q_{N_e+1}^{(i)}, q_{N_e}^{(i)}, \dots q_1^{(i)}].
$$
\n(S10)

These are the so-called branching rules of the permutation group. Here $p_j^{(i)}$, $q_j^{(i)}$ have the same definitions as those in Eqs. [\(S6\)](#page-9-1) and [\(S7\)](#page-9-0). Importantly, the forms of the restricted and induced representations in Eqs. [\(S9\)](#page-11-0) and [\(S10\)](#page-11-1) are very similar to those of the excitation basis states in Eqs. [\(S6\)](#page-9-1) and [\(S7\)](#page-9-0). It is in this sense that we map the states $|\phi_n^k\rangle$ to the irreducible representations of the permutation room. Under such a manning, the potation r. λ_k indicates tha of the permutation group. Under such a mapping, the notation $r \nearrow k$ indicates that the representation $[\phi_n^k]$ is in the induced
representations of $[\phi_n^k]$. I (or equivalently $[\phi_n^k]$ is in the restricted representatio representations of $[\phi_{n-1}^k]$ (or, equivalently, $[\phi_{n-1}^k]$ is in the restricted representations of $[\phi_n^k]$).
We then use c_{n+1} odenote the dimension of the irreducible representation $[\phi_1^k]$. From the

We then use $c_{n,k}$ to denote the dimension of the irreducible representation $[\phi_n^k]$. From the branching rules of the permutation ν group, we have [\[2,](#page-12-1) [3\]](#page-12-2)

$$
c_{n,k} = \sum_{r \nearrow k} c_{n-1,r}, \quad n c_{n-1,r} = \sum_{r \nearrow k} c_{n,k}.
$$
 (S11)

From the first relation in Eq. [\(S11\)](#page-11-2), we have $b_h^k = c_h^k$, which can be proved by induction from $b_{0,0} = c_{0,0} = 1$. And with the second relation in Eq. [\(S11\)](#page-11-2), we immediately have $nb_{n-1,k} = \sum_{k \nearrow r} b_{n,r}$.
Finally as a lost suppose in the main text (Fig. (S) therein) we h

Finally, as a key expression in the main text ([Eq. (6)] therein), we have

$$
\hat{K}(\hat{K}^{\dagger})^{n} \mid \ \rangle = \sum_{k} b_{n,k} \hat{K} |\phi_{n}^{k}\rangle = n \sum_{r} b_{n-1,r} |\phi_{n-1}^{r}\rangle = n(\hat{K}^{\dagger})^{(n-1)} \mid \ \rangle. \tag{S12}
$$

We plot the permutation-group representations for different *n* in the form of Young's lattice in Fig. [S4,](#page-4-7) labeled by the irreducible-representation dimensions for each Young diagram (which can be calculated through the hook length formula). Compared to Fig. 3(a) of the main text, it is clear that an irreducible representation of the parition $[k_{N_e}, k_{N_e-1}, \ldots, k_1]$ (in Fig. [S4\)](#page-4-7) can
be manned to the excitation basis state $[k_{N_e}, k_{N_e-1}]$ (in Fig. 3(a) of the main be mapped to the excitation basis state $|k_{N_e}, k_{N_e-1}, \ldots k_1\rangle$ [in Fig. 3(a) of the main text]. The expansion coefficients in Fig. 3(a), defined as h_e , above are identical with the corresponding irreducible representat defined as $b_{n,k}$ above, are identical with the corresponding irreducible representation dimensions $c_{n,k}$.

In Fig. $S5(a)(b)$ $S5(a)(b)$, we show the excitations, generated by \hat{K}^{\dagger} , from the dark states $|D\rangle = |2\rangle - |11\rangle$ and $|D\rangle = |3\rangle - |12\rangle +$ $|111\rangle$, respectively. In either case, $(\hat{K}^{\dagger})^n|D\rangle$ can be expanded onto the excitation basis that can be mapped to the irreducible representations of the permutation group. And the expansion coefficients are also identical to the corresponding dimensions of the irreducible representation. Hence, we surmise that, for any dark state $|D\rangle$ of \hat{K} , the relation $\hat{K}(\hat{K}^{\dagger})^n|D\rangle = n(\hat{K}^{\dagger})^{(n-1)}|D\rangle$ still holds. This conjecture is consistent with our numerical results throughout the work [and particularly in Fig. 3(c) of the main text].

From the analysis above, we observe that the mapping between the excitation basis and the permutation-group representations essentially relies on the fermion statistics and the limited Hilbert space for the excited states (above dark states). The latter derives from the Liouvillian skin effect in our model, which leads to boundary localized dark states that can be visualized as the real-space Fermi sea. We expect similar structures can be implemented in the momentum space (or a more general parameter space) of identical fermions, so that prethermal time crystals can be devised even in the absence of the skin effect.

Liouvillian perturbation approach

In this section, we outline the Liouvillain perturbation approach that leads to Eq. (7) of the main text. Following Ref. [\[1\]](#page-12-0), we set $\mathcal{L} = \mathcal{L}_0 + s\mathcal{L}_1$, with

$$
\mathcal{L}_0[\rho] = -i[\Delta\hat{B} + \gamma \hat{K}^\dagger \hat{K}, \rho] + \gamma (2\hat{K}\rho \hat{K}^\dagger - \hat{K}^\dagger \hat{K}\rho - \rho \hat{K}^\dagger \hat{K}),
$$
\n(S13)

$$
\mathcal{L}_1[\rho] = -i[\hat{K} + \hat{K}^\dagger, \rho]. \tag{S14}
$$

For a non-dissipative eigenmode $\rho_0 = |D^{B_1}\rangle\langle D^{B_2}|$ with $\mathcal{L}_0[\rho_0] = -i\Delta f \rho_0$, we set

$$
\rho_n = \sum_{j=0}^n \alpha_n^{(j)} \rho_n^j, \quad \text{with} \quad \rho_n^j = \hat{K}^{\dagger j} \rho_0 \hat{K}^{n-j}, \tag{S15}
$$

with the coefficients

$$
\alpha_n^{(j)} = \frac{i^{n-2j} s^n}{((1-i)\gamma - i\Delta)^{n-j}((1+i)\gamma + i\Delta)^j (n-j)! j!}.
$$
\n(S16)

We therefore have

$$
\mathcal{L}_0[\rho_n] + i\Delta f \rho_n = -\sum_{j=0}^n ((i\gamma + i\Delta)(2j - n) + \gamma n) \alpha_n^{(j)} \rho_n^j + 2\gamma \sum_{j=1}^{n-1} j(n - j) \alpha_n^{(j)} \rho_{n-2}^{j-1},
$$
\n(S17)

$$
-s\mathcal{L}_1[\rho_{n-1}] = is \sum_{j=1}^n \alpha_{n-1}^{(j-1)} \rho_n^j - is \sum_{j=0}^{n-1} \alpha_{n-1}^{(j)} \rho_n^j + is \sum_{j=1}^{n-1} \left(j\alpha_{n-1}^{(j)} - (n-j)\alpha_{n-1}^{(j-1)}\right) \rho_{n-2}^{j-1}.
$$
 (S18)

Note that Eqs. [\(S17\)](#page-12-3) and [\(S18\)](#page-12-4) hold for $n \ge 2$. For $n = 1$, the two equations still hold if we set the second term on the right of each to be zero. Plugging Eq. $(S16)$ into Eqs. $(S17)$ and $(S18)$, we have

$$
\mathcal{L}_0[\rho_n] + i\Delta f \rho_n = -s\mathcal{L}_1[\rho_{n-1}],\tag{S19}
$$

holding for $n \in \mathbb{N}^+$. Therefore, $\rho = \rho_0 + s^n \sum_{n=1}^{\infty} \rho_n$ is a non-dissipative eigenmode of $\mathcal L$ with $\mathcal L[\rho] = -i\Delta f$.
However, when the lattice length *L* or the atom number *N* is finite, the relation $\hat K(\hat K^$

However, when the lattice length *L* or the atom number *N* is finite, the relation $\hat{K}(\hat{K}^{\dagger})^n|D^{B_{1,2}}\rangle = n(\hat{K}^{\dagger})^{(n-1)}|D^{B_{1,2}}\rangle$ only holds up to a certain order $n \leq l$. This is because the excitation basis $|k_{N_e}, k_{N_e-1}, \dots, k_1\rangle$ must satisfy $N_e \leq N$ and $k_1 \leq L - N$ for the L and $\epsilon^{(i)} = 1$ and $\epsilon^{(i)} = k + 1$ annear in the expression of \hat{K}^{\dagger} finite *L* or *N*. However, terms with $q_N^{(i)}$ $\binom{i}{N_e+1} = 1$ and $q_1^{(i)}$ $\hat{K}^{\dagger} | k_{N_e}, k_{N_e-1}, \ldots, k_1 \rangle$. To make the $\hat{K}^{\dagger} | k_{N_e}, k_{N_e-1}, \ldots, k_1 \rangle$. To make the expression of $\hat{K}^{\dagger}|k_{N_e}, k_{N_e-1}, \dots, k_1\rangle$ consistent with Eq. [\(S7\)](#page-9-0), we must have $N_e \le N-1$ and $k_1 \le L-N-1$. Similarly, to make the expression of $(\hat{K}^{\dagger})^n | k_{N_e}, k_{N_e-1}, \dots k_1$ consistent with Eq. [\(S7\)](#page-9-0), we must have $n \le N - N_e$ and $n \le L - N - k_1$. Here $N - N_e$
indicates the number of consecutively occupied sites from the left (the real-space Fermi se indicates the number of consecutively occupied sites from the left (the real-space Fermi sea), and *L*−*N* −*k*¹ indicates the number of consecutively unoccupied sites from the right. In this context, since any dark state $|D\rangle$ can be expanded onto the excitation basis, we define $l_N^{\mid D \rangle} = \max\{j \mid \langle D | \hat{c}_j^{\dagger} \rangle\}$ $\int_a^{\infty} c_j |D\rangle = 1$, labeling the site index to the left of which the Fermi sea is intact (without hole excitations). We also define $l_L^{(D)} = \min\{j|\langle D|\hat{c}_{L-j}^{\dagger}c_{L-j}|D\rangle > 0\}$, labeling the site index to the right of which there are no fermion excitations. We then find that the relation $\hat{K}(\hat{K}^{\dagger})^n|D\rangle = n(\hat{K}^{\dagger})^{n-1}|D\rangle$ holds only up to $n \le l^{|D\rangle} = \min(l^{|D\rangle}, l^{|D\rangle}$. It follows that the expression $\mathcal{L}_0[\rho_n] + i\Delta f \rho_n = -s\mathcal{L}_1[\rho_{n-1}]$ also only holds for $n \leq l$, where $l = \min(l^{|D^B|})$, $l^{|D^B|}$) (note that $\rho_0 = |D^B| \setminus \langle D^B|$).
Then the constructed density matrix $\rho = \rho_0 + \nabla^l$ as can be regard Then, the constructed density matrix $\rho = \rho_0 + \sum_{n=1}^{l} \rho_n$ can be regarded as an approximate eigenmode of \mathcal{L} , with the approximate eigenvalue calculated using $Tr(\rho f[\alpha])$ eigenvalue calculated using $Tr(\rho \mathcal{L}[\rho]).$

Starting from the expression $\mathcal{L}[\rho] = -i\Delta f \rho + s^{l+1} \mathcal{L}_1[\rho_l]$, we have

$$
\operatorname{Tr}(\rho \mathcal{L}[\rho]) + i\Delta f = \sum_{n=0}^{l} s^{l+n+1} \operatorname{Tr}(\rho_n \mathcal{L}_1[\rho_l]).
$$
\n
$$
(S20)
$$

Now for each $\rho_0 = |D^{B_1}\rangle\langle D^{B_2}|$ with $B_1 \neq B_2$, we have $\text{Tr}(\rho_n^j \rho_n^k) = 0$. This is because $\text{Tr}(\rho_n^j \rho_n^k) = \text{Tr}(\hat{\mathcal{K}}^{j} \cup D^{B_1}\rangle\langle D^{B_2}|\hat{\mathcal{K}}^{(n-j)}\hat{\mathcal{K}}^{(n-j)}\hat{\mathcal{K}}^{(n-j)}\hat{\mathcal{K}}^{(n-j)}\hat{\mathcal{K}}^{(n-j)}\hat{\$ Now for each $\rho_0 = |D^{B_1}\rangle\langle D^{B_2}|$ with $B_1 \neq B_2$, we have $Tr(\rho_n^j \rho_m^k) = 0$. This is because $Tr(\rho_n^j \rho_m^k) = Tr(\hat{K}^{\dagger j}|D^{B_1}\rangle\langle D^{B_2}|\hat{K}^{(n-j)}\hat{K}^{\dagger k}|D^{B_1}\rangle\langle D^{B_2}|\hat{K}^{(n-j)}\hat{K}^{\dagger k}|D^{B_1}\rangle$. And when $n-j \leq$ whereas for $n - j > k$, we have $\langle D^{B_2} | \hat{K}^{(n-j)} \hat{K}^{(k)} | D^{B_1} \rangle \propto \langle D^{B_2} | \hat{K}^{(n-k-j)} | D^{B_1} \rangle = 0$. From $\text{Tr}(\rho_n^j \rho_m^k) = 0$, we then have $Tr(\rho_n L_1[\rho_l]) = 0$ for $n \le l$. Therefore, $Tr(\rho L[\rho]) + i\Delta f$ is at least of the order $p = 2l + 2$ in *s*. The resulting expression $\lambda \epsilon + i\Delta f \propto s^{2l+2}$ is consistent with the numerical results in Fig. 3(c) of the main text $\lambda_f + i\Delta f \sim s^{2l+2}$ is consistent with the numerical results in Fig. 3(c) of the main text.

[∗] wyiz@ustc.edu.cn

^[1] H. Li, H. Wu, W. Zheng, and W. Yi, Many-body non-Hermitian skin effect under dynamic gauge coupling, Phys. Rev. Res. 5, 033173 (2023)

^[2] B. E. Sagan, The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions (2nd Ed.), Graduate Texts in Math. 203, Springer, New York, 2001.

^[3] A. M. Vershik, A new approach to the representation theory of the symmetric groups. III. Induced representations and the Frobenius-Young correspondence, Mosc. Math. J. 6, 567 (2006).