

# ON THE KUMMER PRO-ÉTALE COHOMOLOGY OF $\mathbb{B}_{\mathrm{dR}}$

XINYU SHAO

ABSTRACT. We investigate  $p$ -adic cohomologies of log rigid analytic varieties over a  $p$ -adic field. For a log rigid analytic variety  $X$  defined over a discretely valued field, we compute the Kummer pro-étale cohomology of  $\mathbb{B}_{\mathrm{dR}}^+$  and  $\mathbb{B}_{\mathrm{dR}}$ . When  $X$  is defined over  $\mathbb{C}_p$ , we introduce a logarithmic  $B_{\mathrm{dR}}^+$ -cohomology theory, serving as a deformation of log de Rham cohomology. Additionally, we establish the log de Rham-étale comparison in this setting and prove the degeneration of both the Hodge-Tate and Hodge-log de Rham spectral sequences for  $X$  defined over a discretely valued field or admitting a proper semistable formal model over  $\mathrm{Spf}(\mathcal{O}_C)$ .

## CONTENTS

1. Introduction	1
1.1. Background	2
1.2. Kummer pro-étale cohomology of $\mathbb{B}_{\mathrm{dR}}$	3
1.3. Geometric $B_{\mathrm{dR}}^+$ -cohomology and comparison theorem	4
1.4. Acknowledgments	5
1.5. Notation and conventions	5
2. Logarithmic de Rham cohomology	6
3. Period sheaves	9
3.1. Definitions	9
3.2. Local study of $\mathbb{B}_{\mathrm{dR}}$ and $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}},\log}$	12
4. Kummer pro-étale cohomology of $\mathbb{B}_{\mathrm{dR}}^+$ and $\mathbb{B}_{\mathrm{dR}}$	14
5. Logarithmic $B_{\mathrm{dR}}^+$ -cohomology for rigid analytic varieties	20
5.1. Preliminary: The $L\eta_f$ functor	20
5.2. Logarithmic $B_{\mathrm{dR}}^+$ -cohomology	22
5.3. Comparison with Kummer pro-étale cohomology	25
5.4. Degeneration of (log) Hodge-Tate spectral sequence	26
References	31

## 1. INTRODUCTION

Let  $\mathcal{O}_K$  be a complete discrete valuation ring with fraction field  $K$  of characteristic 0 and with perfect residue field  $k$  of characteristic  $p$ . Let  $W(k)$  be the ring of Witt vectors of  $k$  with fraction field  $F$  (therefore  $W(k) = \mathcal{O}_F$ ). Let  $\bar{K}$  be an algebraic closure of  $K$  and  $C$  be its  $p$ -adic completion, and let  $\mathcal{O}_{\bar{K}}$  be the integer closure of  $\mathcal{O}_K$  in  $\bar{K}$ . Let  $W(\bar{k})$  be the ring of Witt vectors of  $k$  with

fraction field  $\check{F}$  (therefore  $\check{F}$  is the  $p$ -adic completion of  $F^{\text{nr}}$ , the maximal unramified extension of  $K$  in  $\bar{K}$ ) and  $W(\bar{k}) = \mathcal{O}_{\check{F}}$ . Let  $\phi$  be the absolute Frobenius on  $W(\bar{k})$ . Set  $\mathcal{G} = \text{Gal}(\bar{K}/K)$ .

We will denote by  $\mathcal{O}_K$  and  $\mathcal{O}_K^\times$ , depending on the context, the scheme  $\text{Spec}(\mathcal{O}_K)$  or the formal scheme  $\text{Spf}(\mathcal{O}_K)$  with the trivial log structure and the canonical (i.e., associated to the closed point) log structure respectively.

**1.1. Background.** Before presenting the main theorem, we review some recent advancements in  $p$ -adic Hodge theory.

The aim of  $p$ -adic Hodge theory is to establish a Hodge-type decomposition over  $p$ -adic fields, analogous to the classical Hodge theory developed by Deligne and Hodge. For smooth algebraic varieties, such a decomposition was first achieved in [Fal88]. In the case of proper smooth rigid analytic varieties, Scholze extended this framework using perfectoid geometry, leading to the following results in [Sch13a].

**Theorem 1.1.** [Sch13a, Corollary 1.8] *For any proper smooth rigid-analytic variety  $X$  defined over  $K$ , the Hodge-de Rham spectral sequence*

$$E_1^{i,j} := H^j(X, \Omega_X^i) \Rightarrow H_{\text{dR}}^{i+j}(X)$$

*degenerates at  $E_1$ , and there is a Hodge-Tate decomposition*

$$H_{\text{ét}}^i(X_C, \mathbb{Q}_p) \otimes_K C \simeq \bigoplus_{j=0}^i H^{i-j}(X, \Omega_X^j) \otimes_K C(-j).$$

*The  $p$ -adic étale cohomology  $H_{\text{ét}}^i(X_C, \mathbb{Q}_p)$  is de Rham, with associated filtered  $K$ -vector space  $H_{\text{dR}}^i(X)$ .*

Following Scholze's celebrated work on proper smooth rigid analytic varieties, numerous generalizations have emerged. The work by Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu in [DLLZ23a] (see also the work of Shizhang Li and Xuanyu Pan in [LP19]) extended the above comparison theorem to almost proper rigid analytic varieties  $X$ , which means  $X$  can be written as  $\bar{X} - Z$ , with  $\bar{X}$  being a proper smooth variety and  $Z$  a Zariski closed subset in  $\bar{X}$ . In [DLLZ23a], they further constructed the Simpson and Riemann-Hilbert correspondences for such varieties. Their approach involves developing the theory of log adic spaces, enabling a natural log structure on  $\bar{X}$  induced by the open immersion  $\bar{X} - Z \hookrightarrow \bar{X}$ . By employing resolution of singularities,  $Z$  can be assumed to be a strictly normal crossing divisor on  $\bar{X}$ , allowing the reduction of the problem on  $X$  to one on  $\bar{X}$ , where methods akin to those in [Sch13a] apply.

Another direction of generalization involves exploring general rigid analytic varieties without requiring  $X$  to be proper or smooth. This was studied in [Bos23a] and [Bos23b]. In [Bos23a, Theorem 1.1.8], Guido Bosco computed the pro-étale cohomology of the period sheaf  $\mathbb{B}_{\text{dR}}$  for smooth rigid analytic varieties over a discretely valued field  $K$ . Within the framework of condensed mathematics, he computed the pro-étale cohomology of  $\mathbb{B}_{\text{dR}}$  via the Poincaré lemma, which, for proper smooth rigid analytic varieties over  $K$ , reproduces results from [Sch13a, Theorem 7.11]. In [Bos23b], by using éh-topology, Guido Bosco showed it is also possible to work with non-smooth rigid analytic varieties over an algebraically closed complete field  $C$ . It is worth noting that most of

the results in [Bos23a] and [Bos23b] are valid only within the framework of condensed mathematics, recently developed by Dustin Clausen and Peter Scholze in [CS19].

**1.2. Kummer pro-étale cohomology of  $\mathbb{B}_{\mathrm{dR}}$ .** This article computes the Kummer pro-étale cohomology of  $\mathbb{B}_{\mathrm{dR}}$  for log-smooth rigid analytic varieties over a discrete field  $K$ , extending the results of [Bos23a, Theorem 1.1.8].

In algebraic geometry, for a given smooth algebraic variety  $X$ , we typically consider a compactification  $\overline{X}$  such that  $\overline{X} - X$  is a strict normal crossing divisor. This setup allows  $\overline{X}$  to carry a natural log structure that is log-smooth, thereby reducing many problems to the proper case where cohomology theory behaves more predictably. However, in rigid analytic geometry, compactifying a rigid analytic variety is generally not feasible (while it can be done in the category of adic spaces, the resulting space no longer remains a rigid analytic variety, introducing new complications). This is why, in [DLLZ23a], the authors restrict themselves considering almost proper rigid analytic varieties, which can be associated with a smooth proper rigid analytic variety endowed with a well-behaved log structure after resolution of singularities.

To extend the known comparison theorems for rigid analytic varieties, it is natural to introduce log structures on general (not necessarily proper) rigid analytic varieties. For pro-étale cohomology, recent works by Pierre Colmez, Gabriel Dospinescu, and Wiesława Nizioł [CN17], [CDN20], [CN20], [CN25] provide a detailed description of the pro-étale cohomology for smooth rigid analytic varieties, thanks to the comparison with syntomic cohomology. However, for non-quasi-compact rigid analytic varieties, computing étale cohomology remains challenging. By equipping a quasi-compact rigid analytic variety with a log structure derived from a strict normal crossing divisor, we can apply results on pro-étale cohomology to compute the étale cohomology of the trivial locus, which is itself not quasi-compact. Further exploration in log geometry, therefore, holds potential to clarify significant aspects of the étale cohomology of rigid analytic varieties.

The main theorem of this article is as follows (see Theorem 4.6 for a more general statement).

**Theorem 1.2.** *Let  $X$  be a log smooth rigid analytic variety defined over  $K$ , with fine saturated log-structure. Define*

$$\mathrm{R}\Gamma_{\mathrm{logdR}}(X_{\mathbb{B}_{\mathrm{dR}}}) := \mathrm{R}\Gamma(X, \Omega_X^{\bullet, \mathrm{log}} \otimes_K^{\blacksquare} B_{\mathrm{dR}}).$$

(i) *We have a  $\mathcal{G}_K$ -equivariant, compatible with filtrations, natural isomorphisms in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ :*

$$\mathrm{R}\Gamma_{\mathrm{prokét}}(X_C, \mathbb{B}_{\mathrm{dR}}) \simeq \mathrm{R}\Gamma_{\mathrm{logdR}}(X_{\mathbb{B}_{\mathrm{dR}}}).$$

(ii) *Assume  $X$  is connected and paracompact. Then, for each  $r \in \mathbb{Z}$ , we have a  $\mathcal{G}_K$ -equivariant natural quasi-isomorphism in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ :*

$$\mathrm{R}\Gamma_{\mathrm{prokét}}(X_C, \mathrm{Fil}^r \mathbb{B}_{\mathrm{dR}}) \simeq \mathrm{Fil}^r(\mathrm{R}\Gamma_{\mathrm{logdR}}(X) \otimes_K^{L, \blacksquare} B_{\mathrm{dR}}),$$

where  $\mathrm{R}\Gamma_{\mathrm{logdR}}(X)$  is the condensed log de Rham cohomology complex in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$  (see Definition 2.1).

This result generalizes [Bos23a, Theorem 1.1.8]. By comparing these two results, we can relate the pro-Kummer étale cohomology and the pro-étale cohomology of  $\mathbb{B}_{\mathrm{dR}}$  when the log-structure of  $X$  is coming from a strictly normal crossing divisor.

**Corollary 1.3.** *Let  $X$  be a smooth rigid analytic variety over  $K$ ,  $D \subset X$  is a strictly normal crossing divisor, and  $U = X - D$ . Endow  $X$  with the log-structure coming from  $D$ . Then we have a (non filtered!) natural quasi-isomorphisms in  $D(\text{Mod}_K^{\text{cond}})$ :*

$$\text{R}\Gamma_{\underline{\text{prokét}}}(X_C, \mathbb{B}_{\text{dR}}) \simeq \text{R}\Gamma_{\underline{\text{proét}}}(U_C, \mathbb{B}_{\text{dR}}).$$

It is unclear whether this corollary can be proved directly, as computations for Kummer (pro)étale cohomology typically require case-by-case treatment. For instance, if  $X$  is quasi-compact and  $D \subset X$  is a strictly normal crossing divisor,  $U = X - D$ , we then have

$$\text{R}\Gamma_{\underline{\text{prokét}}}(X_C, \mathbb{Q}_p) \simeq \text{R}\Gamma_{\underline{\text{két}}}(X_C, \mathbb{Q}_p) \simeq \text{R}\Gamma_{\underline{\text{ét}}}(U_C, \mathbb{Q}_p),$$

which follows from [DLLZ23b, Corollary 6.3.4]. Note that  $\text{R}\Gamma_{\underline{\text{ét}}}(U_C, \mathbb{Q}_p) \neq \text{R}\Gamma_{\underline{\text{proét}}}(U_C, \mathbb{Q}_p)$  in general. For example, when  $U = \mathbb{A}^1$ ,  $H_{\underline{\text{ét}}}^1(\mathbb{A}_C^1, \mathbb{Q}_p) = 0$  but  $H_{\underline{\text{proét}}}^1(\mathbb{A}_C^1, \mathbb{Q}_p)$  is very large, as indicated by the main theorem of [CDN20].

**1.3. Geometric  $B_{\text{dR}}^+$ -cohomology and comparison theorem.** Now let  $X$  be a log smooth rigid analytic variety over  $C$ . Since the algebra  $B_{\text{dR}}^+$  lacks a natural  $C$ -structure, establishing a meaningful comparison theorem between the log de Rham cohomology and the Kummer étale cohomology necessitates the introduction of a new cohomology theory,  $\text{R}\Gamma_{B_{\text{dR}}^+}(X)$ . This theory serves as a deformation of log de Rham cohomology via the map  $B_{\text{dR}}^+ \rightarrow C$ .

For rigid analytic varieties without log structures, various approaches have been proposed to construct geometric  $B_{\text{dR}}^+$ -cohomology theories. For example:

- In [BMS18], Bhargav Bhatt, Matthew Morrow, and Peter Scholze introduce  $\text{R}\Gamma_{\text{cris}}(X/B_{\text{dR}}^+)$  using the concept of “very small objects.”
- In [Guo21], Haoyang Guo constructs  $\text{R}\Gamma(X/\Sigma_{\text{inf}}, \mathcal{O}_{X/\Sigma_{\text{inf}}})$  by studying the infinitesimal site over  $B_{\text{dR}}^+$ .
- In [CN25], Pierre Colmez and Wiesława Nizioł define  $\text{R}\Gamma_{\text{dR}}(X/B_{\text{dR}}^+)$  via local semistable formal models of  $X$ .

Notably, all these definitions are shown to be equivalent by [CN25, Proposition 3.27]. In [Bos23b], by using the  $L\eta_f$ -functor, Guido Bosco introduces  $\text{R}\Gamma_{B_{\text{dR}}^+}(X)$  via period sheaves, and proves that it agrees with the geometric  $B_{\text{dR}}^+$ -cohomology.

In this article, we adopt Bosco’s approach to define  $\text{R}\Gamma_{B_{\text{dR}}^+}(X)$ , as we believe it provides the most direct framework for defining logarithmic  $B_{\text{dR}}^+$ -cohomology with minimal prerequisites. After establishing such a theory, we will prove the following comparison theorem.

**Theorem 1.4.** *Let  $X$  be a proper log smooth rigid analytic varieties over  $C$ . Then there are cohomology groups  $H_{B_{\text{dR}}^+}^i(X)$  which come with a canonical filtered isomorphism*

$$H_{\text{két}}^i(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \simeq H_{B_{\text{dR}}^+}^i(X) \otimes_{B_{\text{dR}}^+} B_{\text{dR}}.$$

When  $X$  comes from  $X_0$  over a discrete valued field  $K$ , this isomorphism agrees with the comparison isomorphism (see [DLLZ23a, Theorem 1.1])

$$H_{\text{két}}^i(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \simeq H_{\text{logdR}}^i(X_0) \otimes_K B_{\text{dR}},$$

under a canonical identification

$$H_{B_{\text{dR}}^+}^i(X) = H_{\log\text{dR}}^i(X_0) \otimes_K B_{\text{dR}}^+.$$

Moreover, if  $X$  comes from  $X_0$  over a discrete valued field  $K$  or admits a proper semistable formal model over  $\text{Spf}(\mathcal{O}_C)$ ,  $H_{\log\text{dR}}^i(X/B_{\text{dR}}^+)$  is a finite free  $B_{\text{dR}}^+$ -module, and we have

(i) The Hodge–de Rham spectral sequence

$$E_1^{ij} := H^j(X, \Omega_X^{\log, i}) \Rightarrow H_{\log\text{dR}}^{i+j}(X)$$

degenerates at  $E_1$ .

(ii) The Hodge–Tate spectral sequence

$$E_2^{ij} := H^j(X, \Omega_X^{\log, i})(-j) \Rightarrow H_{\text{két}}^{i+j}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C$$

degenerates at  $E_2$ .

We are currently unable to prove the degeneration of the Hodge–Tate and Hodge–log de Rham spectral sequences in full generality. This limitation arises from our approach, which is similar to those in [BMS18] and [Guo19], as it relies on the existence of a  $B_{\text{dR}}^+/t^2$ -lifting of  $X$ . Addressing the following question would be good enough to (partially) resolve this issue:

**Question 1.5.** *Let  $X$  be a proper smooth rigid analytic varieties over  $C$ , and let  $D \hookrightarrow X$  be a strictly normal crossing divisor. Does the pair  $(X, D)$  admits a log smooth lifting  $(\mathbb{X}, \mathbb{D})$  along  $B_{\text{dR}}^+/t^2 \rightarrow C$ ?*

We note that by using a spreading-out technique of Conrad–Gabber [CG], such a lifting always exists for  $X$  and  $D$ , as proved in [Guo19, Proposition 7.4.4].

These results serve as an initial step toward the study of rational  $p$ -adic Hodge theory for logarithmic rigid analytic varieties. In fact, we can define the syntomic cohomology for logarithmic rigid analytic varieties by adopting a similar framework. We will explore these ideas in detail in a forthcoming paper.

**1.4. Acknowledgments.** The author would like to sincerely thank Wiesława Nizioł for her insightful discussions, and for posing key questions that shaped this article. The article is part of the author’s Ph.D. thesis.

**1.5. Notation and conventions.** In this article, We freely use the language of condensed mathematics as developed in [CS19]. To avoid set-theoretic issues, we fix an uncountable strong limit cardinal  $\kappa$ , which means  $\kappa$  is uncountable and for all  $\lambda < \kappa$ , also  $2^\lambda < \kappa$ .

We adopt the language of adic spaces as developed in [Hub96]. A rigid analytic variety is defined as a quasi-separated adic space locally of finite type over  $\text{Spa}(L, \mathcal{O}_L)$  for a  $p$ -adic field  $L$ . All rigid analytic spaces considered will be over  $K$  or  $C$ . We assume all rigid analytic varieties are separated, taut, and countable at infinity. We denote by  $\text{Sm}_L$  the category of smooth rigid analytic varieties (or smooth dagger varieties) over  $L$ . Moreover, we always assume analytic adic spaces are  $\kappa$ -small, meaning the cardinality of their underlying topological spaces is less than  $\kappa$ .

If  $\mathcal{A}$  is an abelian category, unless stated otherwise, we always work with derived stable  $\infty$ -category  $D(\mathcal{A})$ .

We also use the theory of log adic space. For definitions and properties of log adic space, we refer the reader to [DLLZ23b]. A (pre)log-structure on a condensed ring is simply a (pre)log-structure on the underlying ring.

## 2. LOGARITHMIC DE RHAM COHOMOLOGY

In this section, we study the de Rham cohomology for logarithmic rigid analytic varieties. We will show most of the known results continue to hold in the condensed mathematics setting.

Suppose  $X$  is a log smooth rigid analytic varieties over  $L = K$  or  $C$ , recall that in [DLLZ23b, Construction 3.3.2], one has logarithmic differential étale sheaf  $\Omega_X^{1,\log}$  (also denote by  $\Omega_X^{\log}$ ), locally defined by the universal objects of derivations. When  $X$  is log smooth, by [DLLZ23b, Lemma 3.3.15]  $\Omega_X^{1,\log}$  is locally free of finite rank. For any  $i \geq 1$ , let  $\Omega_X^{i,\log} := \bigwedge^i \Omega_X^{1,\log}$ .

For any affinoid étale morphism  $U \rightarrow X$ ,  $\Omega_X^{i,\log}(U)$  is a finite  $\mathcal{O}_U(U)$ -module, therefore it has a natural structure of  $K$ -Banach space. This leads to the following definition.

**Definition 2.1.** Suppose  $X$  is a log smooth rigid analytic varieties over  $L$ , the logarithmic de Rham cohomology of  $X$  is defined to be

$$\mathrm{R}\Gamma_{\underline{\log\mathrm{dR}}}(X) := \mathrm{R}\Gamma\left(X, \underline{\Omega_X^{\bullet,\log}}\right)$$

in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ .

Recall that the usual de Rham cohomology for log rigid analytic varieties is defined by

$$\mathrm{R}\Gamma_{\log\mathrm{dR}}(X) := \mathrm{R}\Gamma\left(X, \Omega_X^{\bullet,\log}\right).$$

When  $H_{\log\mathrm{dR}}^i(X)$  is finite dimensional (for example, when  $X$  is log smooth and proper), it carries a natural topology from the topology of  $L$ . In this case, the structure of  $\mathrm{R}\Gamma_{\underline{\log\mathrm{dR}}}(X)$  is simple.

**Lemma 2.2.** [Bos23a, Lemma 1.5.11] *Suppose  $X$  is log smooth and proper, then we have*

$$\underline{H_{\log\mathrm{dR}}^i(X)} = \underline{H_{\log\mathrm{dR}}^i(X)}$$

for all  $i \geq 0$ .

We are mainly interested in the case where the log-structure of  $X$  comes from a strictly normal crossing divisor  $D \subset X$ . Following [Gro66b], [Kie67] and [Del70], we give the following definition.

**Definition 2.3.** Suppose  $X$  is a smooth rigid analytic variety over  $L$ ,  $D \subset X$  is a strictly normal crossing divisor,  $U = X - D$ . Let  $i \geq 1$ .

(i) We define  $\Omega_X^i(*D)$  to be the differential  $i$ -forms on  $X$  with meromorphic poles along  $D$ . In other words, let  $\mathcal{I}$  be the ideal sheaf associated to the closed immersion  $D \hookrightarrow X$ , then

$$\Omega_X^i(*D) := \varinjlim_n \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}^n, \Omega_X^i).$$

(ii) We define  $\Omega_X^i(\log D)$  to be the differential  $i$ -forms on  $X$  with logarithmic poles along  $D$ . In other words,  $\Omega_X^i(\log D)$  contains the elements  $\omega$  of  $\Omega_X^i(*D)$  such that  $\omega$  and  $d\omega$  have a pole of order at most 1.

We can also give a local description of the structure of  $\Omega_X^i(\log D)$ . By the existence of tubular neighborhood, [Kie67, Theorem 1.18], locally  $X$  can be written as  $V := \text{Sp}(A \langle x_1, \dots, x_d \rangle)$ , and  $D$  is defined by the equation  $x_1 \cdots x_d = 0$ , where  $A$  is a smooth affinoid  $L$ -algebra. Therefore,  $\Omega_X^i(\log D)(V)$  are generated by forms

$$\omega \cdot (x_{i_1} \cdots x_{i_q})^{-1} dx_{i_1} \cdots dx_{i_q}$$

for  $\omega \in \Omega_A^{i-q}$  and  $1 \leq i_1 < \cdots < i_q \leq d$ .

**Lemma 2.4.** *We have natural isomorphisms*

$$\Omega_X^{i,\log} \xrightarrow{\simeq} \Omega_X^i(\log D),$$

for each  $i \geq 1$ , where the log-structure of  $X$  is given by the open immersion  $U \hookrightarrow X$ .

*Proof.* We can check the lemma locally. By base change, we may assume  $V = \text{Sp}(L \langle x_1, \dots, x_d \rangle)$  as above, and  $D$  is defined by the equation  $x_1 \cdots x_d = 0$ , and the log-structure of  $V$  is defined by the open immersion  $V - D \hookrightarrow V$ . We need to construct a natural isomorphism

$$\Omega_X^{1,\log}(V) \xrightarrow{\simeq} \Omega_X^1(\log D)(V).$$

According to [DLLZ23b, Proposition 3.2.25] (taking  $P = \emptyset$  and  $Q = \mathbb{Z}_{\geq 0}^d = \bigoplus_{i=1}^d \mathbb{Z}_{\geq 0} e_i$ ), we have

$$\Omega_X^{1,\log}(V) \simeq \bigoplus_{i=1}^d L \langle x_1, \dots, x_d \rangle e_i.$$

We can then construct the natural isomorphism  $\Omega_X^{1,\log}(V) \xrightarrow{\simeq} \Omega_X^1(\log D)(V)$  by sending  $e_i$  to  $x_i^{-1} dx_i$ .  $\square$

Now we endow  $X$  with the log-structure defined by the open immersion  $U \hookrightarrow X$ . Since  $X$  is log smooth,  $\Omega_X^i(\log D)$  is a coherent sheaf, therefore have a natural structure of  $K$ -Banach space, and we have a natural condensed sheaf of  $\mathcal{O}_X$ -module  $\underline{\Omega_X^i(\log D)}$ .

Choose a basis  $\mathcal{B}$  of  $X$  of the form  $V = \text{Sp}(A \langle x_1, \dots, x_d \rangle)$  as above, and we endow

$$\Omega_X^i(*D)(V) \simeq \varinjlim_n \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}^n, \Omega_X^i)(V)$$

with the direct limit topology. Then  $\Omega_X^i(*D)$  is a sheaf of topological abelian groups on  $\mathcal{B}$ . We denote by  $\underline{\Omega_X^i(*D)}$  the condensed sheaf associated to  $\Omega_X^i(*D)$ .

**Lemma 2.5.** *We have*

$$\underline{\Omega_X^i(*D)} \simeq \varinjlim_n \underline{\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}^n, \Omega_X^i)}.$$

**Remark 2.6.** This Lemma does not hold automatically, as the condensification functor does not commute with colimits in general.

*Proof.* Denote by

$$\mathcal{F}_n := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}^n, \Omega_X^i).$$



By definition, we need to show the natural morphism

$$\varinjlim_n \mathcal{F}_n \rightarrow \varinjlim_n \mathcal{F}_n = \underline{\Omega}_X^i(*D)$$

is an isomorphism. To see this, we use [BS15, Lemma 4.3.7]. It suffices to show for a basis  $\mathcal{B}$  of  $X$  of the form  $V = \mathrm{Sp}(A\langle x_1, \dots, x_d \rangle)$  as above,  $\mathcal{F}_n(V) \rightarrow \mathcal{F}_{n+1}(V)$  is a closed immersion. Since the sheaf of ideal  $\mathcal{I}$  is generated by  $x_1 \cdots x_d = 0$ , and  $\Omega_X^i$  is locally free of finite rank, by shrinking  $V$  (refining  $\mathcal{B}$ ) and base change we are reduced to showing the map of topological abelian groups

$$L\langle x_1, \dots, x_d \rangle \xrightarrow{x_1 \cdots x_d} L\langle x_1, \dots, x_d \rangle$$

is a closed immersion, which is easy to check: in fact, if  $f \in L\langle x_1, \dots, x_d \rangle$  is not in the ideal  $(x_1 \cdots x_d)$ , then for  $N$  suffices large,  $f + p^N \mathcal{O}_L\langle x_1, \dots, x_d \rangle \cap (x_1 \cdots x_d) = \emptyset$ .  $\square$

**Remark 2.7.** In fact, denote by  $j : U \hookrightarrow X$ , it is easy to see that we have

$$\Omega_X^i(\log D)(V) \hookrightarrow \Omega_X^i(*D)(V) \hookrightarrow j_* \Omega_U^i(V)$$

as topological abelian groups.

We want to compare the hypercohomology of  $\Omega_X^\bullet(\log D)$  and the de Rham cohomology of  $U$ . This leads to the following theorem.

**Theorem 2.8.** [Kie67, Theorem 2.3] *Suppose  $X$  is a smooth rigid analytic varieties over  $L$ ,  $D \subset X$  is a strictly normal crossing divisor,  $U = X - D$ . Then we have (non-filtered) quasi-isomorphisms in  $D(\mathrm{Mod}_L^{\mathrm{cond}})$ :*

$$\mathrm{R}\Gamma(X, \underline{\Omega}^\bullet(\log D)) \xrightarrow{\simeq} \mathrm{R}\Gamma(X, \underline{\Omega}^\bullet(*D)) \xrightarrow{\simeq} \mathrm{R}\Gamma_{\mathrm{dR}}(U).$$

*Proof.* Take  $V = \mathrm{Sp}(A\langle x_1, \dots, x_d \rangle)$  as above, define

$$V^\circ := \{v \in V \text{ such that } |x_1(v)| < 1, \dots, |x_d(v)| < 1\}.$$

Consider the natural maps

$$\Omega^\bullet(\log D)(V^\circ) \xrightarrow{s} \Omega^\bullet(*D)(V^\circ) \xrightarrow{r} j_* \Omega_U^\bullet(V^\circ),$$

where  $j$  is the open immersion  $j : U \hookrightarrow X$ . In [Kie67, Theorem 2.3], Kiehl constructed a map  $\mathrm{Res} : j_* \Omega_U^\bullet(V^\circ) \rightarrow \Omega^\bullet(\log D)(V^\circ)$  and showed that  $r \circ s$  and  $\mathrm{Res} \circ r$  are both null-homotopic. Therefore, to show the same is true for condensed cohomology groups, it enough for us to check that  $s, r$  and  $\mathrm{Res}$  are continuous, which is clear from the construction. This concludes the proof.  $\square$

This theorem also allows us to describe the logarithmic de Rham cohomology for Stein varieties with log-structures. We begin with the following lemma.

**Lemma 2.9.** [GeK04, Corollary 3.2] *Let  $X$  be a smooth Stein variety over  $L$ ,  $D \subset X$  is a strictly normal crossing divisor, let  $U = X - D$ . Endow  $X$  with the log-structure given by the open immersion  $U \hookrightarrow X$ . Then the differentials  $d : \Omega^{i-1, \log}(X) \rightarrow \Omega^{i, \log}(X)$  have closed images.*

*Proof.* The proof of [GeK04, Corollary 3.2] goes through by using Theorem 2.8, [GeK04, Theorem 3.1], and the fact that  $\Omega^{i, \log}(X)$  are coherent sheaves over  $X$  for all  $i \geq 0$ .  $\square$



With the notations in the above lemma, for  $X$  a smooth Stein variety, endowing  $\Omega^{i,\log}(X)^{d=0} \subset \Omega^{i,\log}(X)$  with the subspace topology, and  $H_{\log\text{dR}}^i(X) = \Omega^{i,\log}(X)^{d=0}/d\Omega^{i-1,\log}(X)$  with the induced quotient topology, we have

**Proposition 2.10.** [Bos23a, Lemma 1.5.13] *With the above notations, for all  $i \geq 0$ , we have*

$$H_{\log\text{dR}}^i(X) = \underline{H_{\log\text{dR}}^i(X)} = \underline{\Omega^{i,\log}(X)^{d=0}/d\Omega^{i-1,\log}(X)}.$$

*Proof.* Since  $\Omega^{i,\log}(X)^{d=0}$ ,  $d\Omega^{i-1,\log}(X)$  and  $H_{\log\text{dR}}^i(X)$  are all  $L$ -Fréchet spaces, this proposition follows from [Bos23a, Lemma 1.5.9] and [Bos23a, Lemma 1.A.33]  $\square$

### 3. PERIOD SHEAVES

We review the definition of period sheaves for log adic space and their local properties, which follow from [DLLZ23a].

Suppose  $X$  is an analytic fs log adic space over  $\text{Spa}(\mathbb{Q}_p, \mathbb{Z}_p)$ .

**3.1. Definitions.** We begin with the basic definition.

**Definition 3.1.** The following are defined to be sheaves on  $X_{\text{prokét}}$ :

(i) We define  $\mathbb{A}_{\text{inf}} := W(\widehat{\mathcal{O}}_{X_{\text{prokét}}}^+)$  and  $\mathbb{B}_{\text{inf}} := \mathbb{A}_{\text{inf}}[1/p]$ , where the latter is equipped with a natural map  $\theta : \mathbb{B}_{\text{inf}} \rightarrow \widehat{\mathcal{O}}_{X_{\text{prokét}}}^+$ .

(ii) We define the positive de Rham sheaf  $\mathbb{B}_{\text{dR}}^+ := \lim_{n \in \mathbb{N}} B_{\text{inf},X}/(\ker \theta)^n$ , with filtration given by  $\text{Fil}^r \mathbb{B}_{\text{dR}}^+ := (\ker \theta)^r \mathbb{B}_{\text{dR}}^+$ .

(iii) We define the de Rham sheaf  $\mathbb{B}_{\text{dR}} := \mathbb{B}_{\text{dR}}^+[1/t]$ , where  $t$  is a generator of  $\text{Fil}^1 \mathbb{B}_{\text{dR}}^+$ . The filtration of  $\mathbb{B}_{\text{dR}}$  is given by  $\text{Fil}^r \mathbb{B}_{\text{dR}} := \sum_{j \in \mathbb{Z}, r+j \geq 0} t^{-j} (\ker \theta)^{r+j} \mathbb{B}_{\text{dR}}^+$ .

The following proposition shows that Kummer pro-étale locally these period sheaves are given by the relative period rings.

**Proposition 3.2.** [DLLZ23a, Proposition 2.2.4] *Suppose  $U \in X_{\text{prokét}}$  is a log affinoid perfectoid object, with associated perfectoid space  $\widehat{U} = \text{Spa}(R, R^+)$ , then*

(i) *For  $\mathcal{F} \in \{\widehat{\mathcal{O}}^+, \widehat{\mathcal{O}}^{b,+}, \mathbb{A}_{\text{inf}}, \mathbb{B}_{\text{inf}}, \mathbb{B}_{\text{dR}}^+, \mathbb{B}_{\text{dR}}\}$ , we have  $\mathcal{F}(U) = \mathcal{F}(R, R^+)$ .*

(ii)  *$H^j(U, \mathbb{B}_{\text{dR}}^+) = 0$  and  $H^j(U, \mathbb{B}_{\text{dR}}) = 0$  for  $j > 0$ .*

All these period rings carry a natural condensed sheaf structure defined as follows: given  $X$  an analytic log adic space over  $\text{Spa}(C, \mathcal{O}_C)$  and  $\mathcal{F} \in \{\widehat{\mathcal{O}}^+, \widehat{\mathcal{O}}^{b,+}, \mathbb{A}_{\text{inf}}, \mathbb{B}_{\text{inf}}, \mathbb{B}_{\text{dR}}^+, \mathbb{B}_{\text{dR}}\}$ , denoting  $\mathcal{B}$  the basis of  $X_{\text{prokét}}$  consisting of log affinoid perfectoid objects  $U \in X_{\text{prokét}}$ , then  $\mathcal{F}(U)$  for  $U \in \mathcal{B}$  carry a natural structure of topological abelian groups, by giving  $\widehat{\mathcal{O}}^+(U)$  the  $p$ -adic topology, and endowing all other period sheaves with induced topology, as explained in [Sch13a, Corollary 6.6]. We can then associate to  $\mathcal{F}$  a condensed abelian sheaf  $\underline{\mathcal{F}}$  on  $X_{\text{prokét}}$ , such that  $\underline{\mathcal{F}}(U) = \underline{\mathcal{F}}(U)$  for all log affinoid perfectoid objects  $U \in \mathcal{B}$ .

We pass now to condensed Kummer pro-étale cohomology. We begin with the definition of condensed Kummer pro-étale cohomology.

**Definition 3.3.** Let  $X$  be a analytic log adic space over  $\mathrm{Spa}(\mathbb{Z}_p, \mathbb{Z}_p)$ . Denote by  $*_{\kappa\text{-proét}}$  the site of  $\kappa$ -small profinite sets, with coverings given by finite families of jointly surjective maps. There is a natural morphism of sites

$$f_{\mathrm{cond}} : X_{\mathrm{prokét}} \rightarrow *_{\kappa\text{-proét}}$$

defined by sending a profinite set  $S$  to  $X \times S \in X_{\mathrm{prokét}}$ . Let  $\mathcal{F}$  be a sheaf of abelian groups on  $X_{\mathrm{prokét}}$ , the condensed Kummer pro-étale cohomology of  $\mathcal{F}$  is then defined by

$$\mathrm{R}\Gamma_{\underline{\mathrm{prokét}}}(X, \mathcal{F}) := \mathrm{R}f_{\mathrm{cond},*}\mathcal{F}.$$

Denote by  $H_{\underline{\mathrm{prokét}}}^i(X, \mathcal{F}) := \mathrm{R}^i f_{\mathrm{cond},*}\mathcal{F}$  its  $i$ -th cohomology group valued in  $\mathrm{CondAb}$ .

**Remark 3.4.** When  $X$  is an analytic log adic space over  $\mathrm{Spa}(C, \mathcal{O}_C)$ , denote by  $f : X \rightarrow \mathrm{Spa}(C, \mathcal{O}_C)$  the structure map, then the above definition coincidence with the derived pushforward of  $f$ , i.e. we have  $\mathrm{R}\Gamma_{\underline{\mathrm{prokét}}}(X, \mathcal{F}) = \mathrm{R}f_*\mathcal{F}$ .

**Proposition 3.5.** [Sch13a, Corollary 6.6] *Let  $X$  be an analytic log adic space over  $\mathrm{Spa}(C, \mathcal{O}_C)$  and  $\mathcal{F} \in \{\widehat{\mathcal{O}}^+, \widehat{\mathcal{O}}^{b,+}, \mathbb{A}_{\mathrm{inf}}, \mathbb{B}_{\mathrm{inf}}, \mathbb{B}_{\mathrm{dR}}^+, \mathbb{B}_{\mathrm{dR}}\}$ . For any log affinoid perfectoid object  $U \in \mathcal{B}$  and profinite set  $S \in *_{\kappa\text{-proét}}$ , we have*

$$\mathcal{F}(U \times S) = \mathcal{C}(S, \mathcal{F}(U)).$$

In particular, we have

$$H_{\underline{\mathrm{prokét}}}^i(U, \mathcal{F}) = H_{\mathrm{prokét}}^i(U, \underline{\mathcal{F}})$$

for all  $i \geq 0$ .

Now let  $X$  be a locally noetherian fs log adic space over  $\mathrm{Spa}(K, \mathcal{O}_K)$ . We recall the definition of  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}, \log}$ , a log version of the geometric de Rham period sheaves  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}}$ .

Let  $U = \lim_{i \in I} U_i \in X_{\mathrm{prokét}}$  be a log affinoid perfectoid object, with  $U_i = (\mathrm{Spa}(R_i, R_i^+), \mathcal{M}_i, \alpha_i)$ . For each  $i \in I$ , and with associated perfectoid space  $\widehat{U} = \mathrm{Spa}(R_\infty, R_\infty^+)$ , where  $(R_\infty, R_\infty^+)$  is the  $p$ -adic completion of  $\lim_{i \in I} (R_i, R_i^+)$ , which is perfectoid. For each  $i \in I$ , write  $M_i := \mathcal{M}_i(U_i)$ . By [DLLZ23b, Theorem 5.4.3],  $(\widehat{\mathcal{O}}(U), \widehat{\mathcal{O}}^+(U)) = (R_\infty, R_\infty^+)$ , and the tilt of  $(\widehat{\mathcal{O}}(U), \widehat{\mathcal{O}}^+(U))$  is  $(R_\infty^{b,+}, R_\infty^b)$ . Define  $M := \mathcal{M}_{X_{\mathrm{prokét}}}(U) = \lim_{i \in I} M_i$ , and  $M^b := \mathcal{M}_{X_{\mathrm{prokét}}^b}(U) = \lim_{a \rightarrow a^p} M$ .

**Definition 3.6.** (i) The period sheaf  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}, \log}^+$  is defined to be the sheaf associated to the presheaf

$$U \mapsto \varinjlim_{i \in I} (R_i^+ \widehat{\otimes}_{W(k)} \mathbb{A}_{\mathrm{inf}}(U)) [1/p] \left[ \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]}, (m_i, m_i^b) \in M_i \times_M M^b \right]^{\wedge_{\ker \theta_{\log}}}.$$

Here  $\widehat{\otimes}$  is the  $p$ -adic completion of the tensor product,  $[\alpha^b(m^b)]$  is obtained by the multiplicative map

$$R^{+b} \rightarrow W(R^{+b})[1/p]/\xi^r : f \mapsto [f]$$

induced by  $R^{+b} \rightarrow W(R^{+b})$ , for each  $r \geq 1$ , and

$$\theta_{\log} : (R_i^+ \widehat{\otimes}_{W(k)} \mathbb{A}_{\mathrm{inf}}(U)) [1/p] \left[ \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]}, (m_i, m_i^b) \in M_i \times_M M^b \right] \rightarrow R_\infty$$

is induced by the map  $R_i^+ \rightarrow R_\infty^+$ ,  $\theta : A_{\text{inf}}(U) \rightarrow R_\infty^+$  and  $\theta_{\log} \left( \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]} \right) = 1$  for any  $(m_i, m_i^b) \in M_i \times_M M^b$ . “ $\wedge_{\ker \theta_{\log}}$ ” means the completion with respect to the  $\ker \theta_{\log}$ -adic topology. We equip  $\mathcal{O}_{\text{dR}, \log}^+$  with the filtration  $\text{Fil}^r \mathcal{O}_{\text{dR}, \log}^+ := (\ker \theta_{\log})^r \mathcal{O}_{\text{dR}, \log}^+$ .

(ii)  $\mathcal{O}_{\text{dR}, \log}$  is the completion of the sheaf  $\mathcal{O}_{\text{dR}, \log}^+[1/t]$  with respect to the filtration defined by  $\text{Fil}^r \mathcal{O}_{\text{dR}, \log}^+[1/t] := \sum_{j \in \mathbb{Z}, r+j \geq 0} t^{-j} (\ker \theta_{\log})^{r+j} \mathcal{O}_{\text{dR}, \log}^+$ .

Similar to  $\mathcal{O}_{\text{dR}}^+$ , one can equip  $\mathcal{O}_{\text{dR}, \log}^+$  with a natural log connection

$$\nabla : \mathcal{O}_{\text{dR}, \log}^+ \rightarrow \mathcal{O}_{\text{dR}, \log}^+ \otimes_{\mathcal{O}_{X_{\text{prokét}}}} \Omega_X^{\log}$$

as follows (By abuse of notation, we still denote by  $\Omega_X^{\log}$  the pullback of log differential sheaf  $\Omega_X^{\log}$  to  $X_{\text{prokét}}$ ). We may write  $\mathcal{O}_{\text{dR}, \log}^+(U)$  as

$$\mathcal{O}_{\text{dR}, \log}^+(U) = \varinjlim_{i \in I} \widehat{S}_i = \varinjlim_{i \in I} \varprojlim_{r, s} (S_{i, r} / (\ker \theta_{\log})^s).$$

Here

$$S_{i, r} = (R_i^+ \widehat{\otimes}_{W(k)} (\mathbb{A}_{\text{inf}}(U)) / \xi^r) [1/p] \left[ \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]}, (m_i, m_i^b) \in M_i \times_M M^b \right],$$

and  $\widehat{S}_i := \varprojlim_{r, s} (S_{i, r} / (\ker \theta_{\log})^s)$ . Then there is a unique  $\mathbb{B}_{\text{dR}}^+(U) / \xi^r$ -linear log connection

$$\nabla_{i, r} : S_{i, r} \rightarrow S_{i, r} \otimes_{R_i} \Omega_X^{\log}(U_i)$$

extending  $d : R_i \rightarrow \Omega_X^{\log}(U_i)$  and  $\delta : M_i \rightarrow \Omega_X^{\log}(U_i)$  such that

$$\nabla_{i, r} \left( \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]} \right) = \frac{\alpha_i(m_i)}{[\alpha^b(m^b)]} \delta(a)$$

for any  $(m_i, m_i^b) \in M_i \times_M M^b$ . Then the definition of  $\nabla_{i, r}$  gives

$$\nabla_{i, r} ((\ker \theta_{\log})^s) \subset (\ker \theta_{\log})^{s-1} \otimes_{R_i} \Omega_X^{\log}(U_i)$$

for all  $s \geq 1$ . Then the  $\mathbb{B}_{\text{dR}}^+$ -linear log connection

$$\nabla : \mathcal{O}_{\text{dR}, \log}^+ \rightarrow \mathcal{O}_{\text{dR}, \log}^+ \otimes_{\mathcal{O}_{X_{\text{prokét}}}} \Omega_X^{\log}$$

is obtained by taking

$$\nabla := \varinjlim_{i \in I} \varprojlim_r \left( \nabla_{i, r}^{\wedge_{\ker \theta_{\log}}} \right).$$

By inverting  $t$ ,  $\nabla$  further extends to a  $\mathbb{B}_{\text{dR}}$ -linear log connection

$$\nabla : \mathcal{O}_{\text{dR}, \log} \rightarrow \mathcal{O}_{\text{dR}, \log} \otimes_{\mathcal{O}_{X_{\text{prokét}}}} \Omega_X^{\log},$$

satisfying

$$\nabla (\text{Fil}^r \mathcal{O}_{\text{dR}, \log}) \subset (\text{Fil}^{r-1} \mathcal{O}_{\text{dR}, \log}) \otimes_{\mathcal{O}_{X_{\text{prokét}}}} \Omega_X^{\log}$$

for all  $r \in \mathbb{Z}$ .

**3.2. Local study of  $\mathbb{B}_{\text{dR}}$  and  $\mathcal{O}_{\mathbb{B}_{\text{dR},\text{log}}}$ .** We briefly discuss the local properties of  $\mathbb{B}_{\text{dR}}$  and  $\mathcal{O}_{\mathbb{B}_{\text{dR},\text{log}}}$ , as studied in [DLLZ23a, 2.3]. Let  $P$  be a toric monoid (i.e. a fs sharp monoid). Denote by

$$\mathbb{E} := \text{Spa}(K\langle P \rangle, \mathcal{O}_K\langle P \rangle)$$

with the log-structure induced by the natural map  $P \rightarrow K\langle P \rangle$ .  $\mathbb{E}$  admits a pro-Kummer étale cover as follows. For each  $m \in \mathbb{Z}_{>0}$ , let  $\frac{1}{m}P$  be the toric monoid such that  $P \hookrightarrow \frac{1}{m}P$  can be identified with the multiple map  $[m] : P \rightarrow P$ . Let  $P_{\mathbb{Q}_{\geq 0}} := \lim_m \left( \frac{1}{m}P \right)$  and  $P_{\mathbb{Q}}^{\text{gp}} := (P_{\mathbb{Q}_{\geq 0}})^{\text{gp}} \simeq P^{\text{gp}} \otimes \mathbb{Q}$ . Denote by

$$\mathbb{E}_{m,C} := \text{Spa} \left( K \left\langle \frac{1}{m}P \right\rangle, \mathcal{O}_K \left\langle \frac{1}{m}P \right\rangle \right)$$

equipped with the log-structure induced by  $\frac{1}{m}P \rightarrow K\langle \frac{1}{m}P \rangle$ . The morphism  $\mathbb{E}_m \rightarrow \mathbb{E}$  is a finite Kummer étale cover with Galois group

$$\Gamma_m = \text{Hom}(P^{\text{gp}}, \boldsymbol{\mu}_m),$$

where  $\boldsymbol{\mu}_m$  is the  $m$ -th roots of unity in  $C$ . Denote the log affinoid perfectoid object by

$$\widetilde{\mathbb{E}}_C := \lim_m \mathbb{E}_{m,C} \in \mathbb{E}_{C,\text{prokét}},$$

where for  $m|m'$ , the transition maps are induced by  $\frac{1}{m}P \hookrightarrow \frac{1}{m'}P$ . Then we have the associated perfectoid space

$$\widehat{\mathbb{E}}_C := \text{Spa}(C\langle P_{\mathbb{Q}_{\geq 0}} \rangle, \mathcal{O}_C\langle P_{\mathbb{Q}_{\geq 0}} \rangle).$$

The morphism  $\widehat{\mathbb{E}}_C \rightarrow \mathbb{E}_C$  is a Galois pro-finite Kummer étale cover with Galois group

$$\Gamma = \text{Hom}(P_{\mathbb{Q}_{\geq 0}}^{\text{gp}}/P^{\text{gp}}, \boldsymbol{\mu}_{\infty}),$$

where  $\boldsymbol{\mu}_{\infty} := \cup_{m \in \mathbb{N}} \boldsymbol{\mu}_m$ . The natural action of the Galois group  $\Gamma$  on  $\mathcal{O}_C\langle P_{\mathbb{Q}_{\geq 0}} \rangle$  is given by

$$\gamma(T^a) = \gamma(a)T^a$$

for all  $\gamma \in \Gamma$  and  $a \in P_{\mathbb{Q}_{\geq 0}}$ , where  $T^a$  is the corresponding element of  $a$  in  $\mathcal{O}_C\langle P_{\mathbb{Q}_{\geq 0}} \rangle$ .

Suppose  $X = \text{Spa}(R, R^+)$  is an affinoid fs log adic space of finite type over  $\text{Spa}(K, \mathcal{O}_K)$ , with a strictly étale morphism  $X \rightarrow \mathbb{E}$ , which can be written as a composite of rational embedding and finite étale maps. By pulling back to  $X$  we get

$$\widetilde{X}_C := X_C \times_{\mathbb{E}_C} \widehat{\mathbb{E}}_C = \text{Spa}(R_{\infty}, R_{\infty}^+),$$

which is a Galois pro-finite Kummer étale cover with Galois group  $\Gamma$ .

Let  $\mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P]]$  be the sheaf of monoid algebras. For  $a \in P$ , denote by  $e^a$  the image of  $a$  in  $\mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P]]$  via the natural morphism  $P \rightarrow \mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P]]$ . Let  $\mathfrak{m} \subset \mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P]]$  be the sheaf of ideals generated by  $\{e^a - 1\}_{a \in P}$ , and denote by

$$\mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P - 1]] := \varprojlim_r (\mathbb{B}_{\text{dR}}|_{\widetilde{X}}[[P]]/\mathfrak{m}^r).$$

Consider the monoid homomorphism (with respect to the additive structure on  $\mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]]$ )

$$P \rightarrow \mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]] : a \mapsto \log(e^a) := \sum_{l=1}^{\infty} (-1)^{l-1} \frac{1}{l} (e^a - 1)^l,$$

which uniquely extends to a group homomorphism

$$P^{\text{gp}} \rightarrow \mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]] : a \mapsto y_a := \log(e^{a^+}) - \log(e^{a^-}),$$

where  $a = a^+ - a^-$  for  $a^+, a^- \in P$ . Choose a  $\mathbb{Z}$ -basis  $\{a_1, \dots, a_n\}$  of  $P^{\text{gp}}$ , and for each  $j = 1, 2, \dots, n$ , denote by  $y_j := y_{a_j}$ , then we have a canonical isomorphism of  $\mathbb{B}_{\text{dR}}|_{\tilde{X}}$ -algebras:

$$\mathbb{B}_{\text{dR}}|_{\tilde{X}}[[y_1, \dots, y_n]] \xrightarrow{\cong} \mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]] : y_j \mapsto y_j := y_{a_j},$$

matching the ideals  $(y_1, \dots, y_n)^r$  and  $(\xi, y_1, \dots, y_n)^r$  of the source with the ideals  $\mathfrak{m}^r$  and  $(\xi, \mathfrak{m})^r$  of the target, respectively, for all  $r \in \mathbb{Z}$ .

Now, similar to [Sch13a, Proposition 6.10], we can give a local description of  $\mathcal{O}\mathbb{B}_{\text{dR}, \log}^+$ . Recall that we denote by  $U = \varprojlim_{i \in I} U_i \in X_{\text{prokét}}/\tilde{X}$  a log affinoid perfectoid object, with  $U_i = (\text{Spa}(R_i, R_i^+), \mathcal{M}_i, \alpha_i)$ . Consider the map (note that the sheaf  $\mathcal{M}^b|_{\tilde{X}}$  is generated by  $P_{\mathbb{Q}_{\geq 0}}$  and  $\varprojlim_{f_i \rightarrow f^p} \mathcal{O}_{X_{\text{prokét}}}^{\times}|_{\tilde{X}}$ , therefore  $\alpha^b(a)$  makes sense)

$$(\mathbb{B}_{\text{dR}}^+(U)/\xi^r)[P] \rightarrow S_{i,r} : e^a \mapsto \frac{\alpha(a)}{[\alpha^b(a)]}, \text{ for all } a \in P,$$

which sends  $(\xi, \mathfrak{m})$  to  $\ker(\theta_{\log})$ , so this map induces a map  $\mathbb{B}_{\text{dR}}\tilde{X}[[P-1]] \rightarrow \hat{S}_i$ . After taking completion and sheafification, we obtain a map

$$\mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]] \rightarrow \mathcal{O}\mathbb{B}_{\text{dR}, \log}^+|_{\tilde{X}} \quad (3.1)$$

on  $X_{\text{prokét}}/\tilde{X}$ . The map is compatible with filtrations on both sides, where the filtration on  $\mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]]$  is given by  $\text{Fil}^r \mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]] := (\xi, \mathfrak{m})^r \mathbb{B}_{\text{dR}}|_{\tilde{X}}[[P-1]]$  for all  $r \in \mathbb{Z}$ . We have the following proposition.

**Proposition 3.7.** [DLLZ23b, Proposition 2.3.15] *The map (3.1) is an isomorphism of filtered sheaves.*

**Remark 3.8.** In fact, for  $U = \varprojlim_{i \in I} U_i$  as above, the natural map  $\hat{S}_i \rightarrow \mathcal{O}\mathbb{B}_{\text{dR}, \log}^+(U)$  is already an isomorphism.

**Remark 3.9.** We can also describe the log connection of  $\mathcal{O}\mathbb{B}_{\text{dR}, \log}^+$  using this isomorphism. In fact, choose  $\mathbb{Z}$ -basis  $\{a_1, \dots, a_n\}$  of  $P^{\text{gp}}$ , write  $a_j = a_j^+ - a_j^-$  for  $j = 1, 2, \dots, n$ , then  $y_j = b_j^+ - b_j^- := \log(e^{a_j^+}) - \log(e^{a_j^-})$ , and the isomorphism sends  $y_j$  to  $\frac{\alpha(b_j^+)}{[\alpha^b(b_j^+)]} - \frac{\alpha(b_j^-)}{[\alpha^b(b_j^-)]}$ . Therefore,

$$\nabla(y_j) = \nabla\left(\frac{\alpha(b_j^+)}{[\alpha^b(b_j^+)]}\right) - \nabla\left(\frac{\alpha(b_j^-)}{[\alpha^b(b_j^-)]}\right) = \delta(a_j^+) - \delta(a_j^-) = \delta(a_j).$$

We note that, over  $\tilde{X}$ , the log connection of  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$  is compatible with the canonical one of the polynomial algebra  $\mathbb{B}_{\mathrm{dR}}|_{\tilde{X}}[[y_1, \dots, y_n]]$ , since  $\Omega_X^{\mathrm{log}} \simeq \bigoplus_{j=1}^n \mathcal{O}_X \delta(a_j)$  by [DLLZ23b, Theorem 3.3.17, Corollary 3.3.18, Proposition 3.2.25, and Corollary 3.2.29].

**Corollary 3.10.** [DLLZ23b, Corollary 2.3.16] *The isomorphism 3.1 induces isomorphisms*

$$\mathrm{Fil}^r \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+ \simeq t^r \mathbb{B}_{\mathrm{dR}}^+ \{W_1, \dots, W_n\}$$

over  $X_{\mathrm{prokét}}/\hat{X}$ , for all  $r \in \mathbb{Z}$ . Here  $\mathbb{B}_{\mathrm{dR}}^+ \{W_1, \dots, W_n\}$  is the ring of power series that are  $t$ -adically convergent, and

$$W_j = t^{-1}y_j$$

for each  $1 \leq j \leq n$ . In particular,

$$\mathrm{gr}^r \mathcal{O}\mathbb{B}_{\mathrm{dR},\log} \simeq t^r \hat{\mathcal{O}}_{X_{\mathrm{prokét}}} [W_1, \dots, W_n],$$

for all  $r \in \mathbb{Z}$ .

The above discussion allows us to deduce the Poincaré lemma for  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$  and  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}$  with log connections.

**Proposition 3.11.** [DLLZ23a, Corollary 2.4.2] *Let  $X$  be a log smooth rigid analytic variety defined over  $K$ , with fine log-structure, and is of dimension  $n$ . Then, we have an exact sequence of sheaves on  $X_{\mathrm{prokét}}$ :*

$$0 \rightarrow \mathbb{B}_{\mathrm{dR}}^+ \rightarrow \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+ \xrightarrow{\nabla} \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+ \otimes_{\mathcal{O}_X} \Omega_X^{\mathrm{log},1} \xrightarrow{\nabla} \dots \xrightarrow{\nabla} \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+ \otimes_{\mathcal{O}_X} \Omega_X^{\mathrm{log},n} \rightarrow 0.$$

The exact sequence of sheaves also holds when we replace  $\mathbb{B}_{\mathrm{dR}}^+$  and  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$  with  $\mathbb{B}_{\mathrm{dR}}$  and  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}$  respectively. For  $r \in \mathbb{Z}$  we also have compatible exact sequences of sheaves on  $X_{\mathrm{prokét}}$ :

$$0 \rightarrow \mathrm{Fil}^r \mathbb{B}_{\mathrm{dR}} \rightarrow \mathrm{Fil}^r \mathcal{O}\mathbb{B}_{\mathrm{dR},\log} \xrightarrow{\nabla} (\mathrm{Fil}^{r-1} \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}) \otimes_{\mathcal{O}_X} \Omega_X^{\mathrm{log},1} \xrightarrow{\nabla} \dots \xrightarrow{\nabla} (\mathrm{Fil}^{r-n} \mathcal{O}\mathbb{B}_{\mathrm{dR},\log}) \otimes_{\mathcal{O}_X} \Omega_X^{\mathrm{log},n} \rightarrow 0.$$

#### 4. KUMMER PRO-ÉTALE COHOMOLOGY OF $\mathbb{B}_{\mathrm{dR}}^+$ AND $\mathbb{B}_{\mathrm{dR}}$

We will establish an analog version of [Bos23a, Theorem 1.8] for log adic spaces. Moreover, we will also construct a fully faithful functor from the category of filtered  $\mathcal{O}_X$ -modules with integrable connections to the category of  $\mathbb{B}_{\mathrm{dR}}$ -local systems, thereby generalizing [Sch13a, Theorem 7.6].

We start with the definition of local systems for log adic spaces. Let  $X$  be a log smooth rigid analytic variety over  $K$ , equipped with a fine log structure.

**Definition 4.1.** (i) A  $\mathbb{B}_{\mathrm{dR}}^+$ -local system is a sheaf of  $\mathbb{B}_{\mathrm{dR}}^+$ -module  $\mathbb{M}^+$  that is locally on  $X_{\mathrm{prokét}}$  free of finite rank.

(ii) An  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$ -module with integrable log connection is a sheaf of  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$ -module  $\mathcal{M}$  that is locally on  $X_{\mathrm{prokét}}$  free of finite rank, together with an integrable log connection

$$\nabla_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M} \otimes_{\mathcal{O}_X} \Omega_X^{\mathrm{log}},$$

satisfying the Leibniz rule with respect to the derivation  $\nabla$  of  $\mathcal{O}\mathbb{B}_{\mathrm{dR},\log}^+$ .

**Theorem 4.2.** *The functor*

$$\mathbb{M}^+ \mapsto (\mathcal{M}, \nabla_{\mathcal{M}}) := \left( \mathbb{M}^+ \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log}^+, \text{id} \otimes \nabla \right)$$

*gives an equivalence between the category of  $\mathbb{B}_{\text{dR}}^+$ -local systems and the category of  $\mathcal{O}\mathbb{B}_{\text{dR},\log}^+$ -modules with integrable log connection. The inverse is given by  $(\mathcal{M}, \nabla_{\mathcal{M}}) \mapsto \mathbb{M}^+ := (\mathcal{M})^{\nabla_{\mathcal{M}}=0}$ .*

*Proof.* It is clear that for a  $\mathbb{B}_{\text{dR}}^+$ -local system  $\mathbb{M}^+$ ,

$$\left( \mathbb{M}^+ \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log}^+ \right)^{\nabla=0} = \mathbb{M}^+.$$

We need to show that for  $(\mathcal{M}, \nabla)$  a  $\mathcal{O}\mathbb{B}_{\text{dR},\log}^+$ -module with integrable log connection, the natural morphism

$$(\mathcal{M})^{\nabla_{\mathcal{M}}=0} \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log}^+ \xrightarrow{\simeq} \mathcal{M},$$

is an isomorphism. The problem is local, so we can use Proposition 3.7 and reduce to the case that  $X = \text{Spa}(R, R^+) \rightarrow \mathbb{E}$  is a strictly étale morphism, which can be written as a composite of rational embedding and finite étale maps. Now suppose  $M$  is a locally free  $\mathbb{B}_{\text{dR}}^+(R^\infty)[[y_1, \dots, y_n]]$ -module, with an integrable log connection  $\nabla_M$ , using Remark 3.9 and [Kat70, Proposition 8.9], we have  $M \simeq M^{\nabla=0} \otimes_{\mathbb{B}_{\text{dR}}^+(R^\infty)} \mathbb{B}_{\text{dR}}^+(R^\infty)[[y_1, \dots, y_n]]$ , which concludes the proof.  $\square$

**Definition 4.3.** [Sch13a, Definition 7.4 and Definition 7.5] (i) A filtered  $\mathcal{O}_X$ -module with integrable log connection is a locally free  $\mathcal{O}_X$ -module  $\mathcal{E}$  on  $X$  (with étale, pro-étale or analytic topology, since they are all equivalent), together with a separated and exhaustive decreasing filtration  $\text{Fil}^r \mathcal{E}, i \in \mathbb{Z}$ , by locally direct summands, and an integrable log connection  $\nabla$  satisfying the Griffiths transversality.

(ii) We say  $\mathcal{E}$  and an  $\mathcal{O}\mathbb{B}_{\text{dR},\log}^+$ -module with integrable log connection  $\mathcal{M}$  are associated if there is an isomorphism of sheaves on  $X_{\text{prokét}}$

$$\mathcal{M} \otimes_{\mathcal{O}\mathbb{B}_{\text{dR},\log}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log} \simeq \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\text{dR},\log},$$

compatible with filtrations and connections, where the filtration on the left side is the one on  $\mathcal{O}\mathbb{B}_{\text{dR},\log}$ , i.e.,

$$\text{Fil}^r (\mathcal{M} \otimes_{\mathcal{O}\mathbb{B}_{\text{dR},\log}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log}) := \sum_{j \in \mathbb{Z}} t^{-j} (\ker \theta_{\log})^{r+j} \mathcal{M}.$$

**Theorem 4.4.** (i) *If  $\mathcal{M}$  is an  $\mathcal{O}\mathbb{B}_{\text{dR},\log}^+$ -module with integrable log connection and horizontal section  $\mathbb{M}^+$ , which is associated to a filtered  $\mathcal{O}_X$ -module with integrable connection  $\mathcal{E}$ , then*

$$\mathbb{M}^+ = \text{Fil}^0 (\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\text{dR},\log})^{\nabla=0}.$$

*Similarly, one can reconstruct  $\mathcal{E}$  with filtration and log connection by*

$$\mathcal{E}_{\text{ét}} \simeq \nu_* \left( \mathbb{M}^+ \otimes_{\mathbb{B}_{\text{dR}}^+} \mathcal{O}\mathbb{B}_{\text{dR},\log} \right)^{\nabla=0},$$

*where  $\nu$  is the morphism of site  $X_{\text{prokét}} \rightarrow X_{\text{ét}}$ .*

(ii) *If  $\mathcal{E}$  is an  $\mathcal{O}_X$ -module with integrable log connection, the sheaf*

$$\mathbb{M}^+ = \text{Fil}^0 (\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{B}_{\text{dR},\log})^{\nabla=0}$$



is a  $\mathbb{B}_{\mathrm{dR}}^+$ -local system such that  $\mathcal{E}$  is associated to  $\mathcal{M} = \mathbb{M}^+ \otimes_{\mathbb{B}_{\mathrm{dR}}^+} \mathcal{O}_{\mathbb{B}_{\mathrm{dR}}, \log}$ .

In particular, the functor sending  $\mathcal{E}$  to  $\mathbb{M}^+$  is a fully faithful functor from the category of filtered  $\mathcal{O}_X$ -module with integrable log connection to the category of  $\mathbb{B}_{\mathrm{dR}}^+$ -local systems.

*Proof.* The proof goes as the same way as the proof of [Sch13a, Theorem 7.6], by replacing  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}}$  with  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}, \log}$  and  $\Omega_X$  with  $\Omega_X^{\log}$ .  $\square$

**Definition 4.5.** Let  $X$  be a log smooth rigid-analytic variety over  $K$  with fine log-structure. Let  $(\mathcal{E}, \nabla, \mathrm{Fil}^\bullet)$  be a filtered  $\mathcal{O}_X$ -module with integrable log connection. We define the log de Rham complex of  $X$  by

$$\mathrm{logdR}_X^\mathcal{E} := \left[ \mathcal{E} \xrightarrow{\nabla} \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_X^{\log, 1} \xrightarrow{\nabla} \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_X^{\log, 2} \xrightarrow{\nabla} \dots \right].$$

We equip  $\mathrm{logdR}_X^\mathcal{E}$  with the filtration given by

$$\mathrm{Fil}^r \mathrm{logdR}_X^\mathcal{E} := \left[ \mathrm{Fil}^r \mathcal{E} \xrightarrow{\nabla} \mathrm{Fil}^{r-1} \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_X^{\log, 1} \xrightarrow{\nabla} \mathrm{Fil}^{r-2} \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_X^{\log, 2} \xrightarrow{\nabla} \dots \right]$$

for  $r \in \mathbb{Z}$ .

(i) For  $i \geq 0$ , the condensed de Rham cohomology group  $H_{\mathrm{logdR}}^i(X, \mathcal{E})$  with coefficient is defined to be the  $i$ -th cohomology group of the complex

$$\mathrm{R}\Gamma_{\mathrm{logdR}}(X, \mathcal{E}) := \mathrm{R}\Gamma(X, \mathrm{logdR}_X^\mathcal{E})$$

of  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ .

(ii) We define the complex of  $D(\mathrm{Mod}_K^{\mathrm{cond}})$

$$\mathrm{R}\Gamma_{\mathrm{logdR}}(X_{B_{\mathrm{dR}}}, \mathcal{E}) := \mathrm{R}\Gamma(X, \mathrm{logdR}_X^\mathcal{E} \otimes_K^\blacksquare B_{\mathrm{dR}}),$$

and we endow it with the filtration induced from the tensor product filtration.

We will prove the following theorem.

**Theorem 4.6.** *Let  $X$  be a log smooth rigid analytic variety defined over  $K$ , with fine saturated log-structure. Let  $(\mathcal{E}, \nabla, \mathrm{Fil}^\bullet)$  be a filtered  $\mathcal{O}_X$ -module with integrable log connection, with associated  $\mathbb{B}_{\mathrm{dR}}^+$ -local system  $\mathbb{M}^+$ . Denote by  $\mathbb{M} := \mathbb{M}^+[1/t]$ .*

(i) *We have a  $\mathcal{G}_K$ -equivariant, compatible with filtrations, natural quasi-isomorphisms in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ :*

$$\mathrm{R}\Gamma_{\mathrm{prokét}}(X_C, \mathbb{M}) \simeq \mathrm{R}\Gamma_{\mathrm{logdR}}(X_{B_{\mathrm{dR}}}, \mathcal{E}).$$

(ii) *Assume  $X$  is connected and paracompact. Then, for each  $r \in \mathbb{Z}$ , we have a natural  $\mathcal{G}_K$ -equivariant quasi-isomorphisms in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ :*

$$\mathrm{R}\Gamma_{\mathrm{prokét}}(X_C, \mathrm{Fil}^r \mathbb{M}) \simeq \mathrm{Fil}^r(\mathrm{R}\Gamma_{\mathrm{logdR}}(X, \mathcal{E}) \otimes_K^{L, \blacksquare} B_{\mathrm{dR}}),$$

where  $\mathrm{R}\Gamma_{\mathrm{logdR}}(X)$  is the log de Rham cohomology complex in  $D(\mathrm{Mod}_K^{\mathrm{cond}})$ .

To prove the theorem, we need the following proposition, which comes from the local study of  $\mathbb{B}_{\mathrm{dR}}$  and  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}, \log}$ , as Section 3. We follow the notations in 3. Suppose that  $X = \mathrm{Spa}(R, R^+)$  is an affinoid fs log adic space of finite type over  $\mathrm{Spa}(K, \mathcal{O}_K)$ , with a strictly étale morphism  $X \rightarrow \mathbb{E}$ ,

which can be written as a composite of rational embeddings and finite étale maps. By pulling back  $\widehat{\mathbb{E}}_C \rightarrow \mathbb{E}_C$  to  $X$  we get

$$\widetilde{X}_C := X_C \times_{\mathbb{E}_C} \widehat{\mathbb{E}}_C = \text{Spa}(R_\infty, R_\infty^+).$$

**Proposition 4.7.** *Suppose  $S \in *_{\kappa, \text{proét}}$ , then we have*

$$H_{\text{prokét}}^i(X_C \times S, \text{gr}^j \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}) = \begin{cases} \mathcal{C}^0(S, R \widehat{\otimes}_K C(j)) & \text{if } i = 0 \\ 0 & \text{if } i > 0 \end{cases},$$

where  $j \in \mathbb{Z}$  is the Tate twist.

*Proof.* By twisting we can reduce to the case  $j = 0$ . When  $S = *$ , this is [DLLZ23a, Lemma 3.3.15]. For the general case, since  $\widehat{\mathbb{E}}_C \rightarrow \mathbb{E}_C$  is a Galois pro-finite Kummer étale cover with Galois group  $\Gamma$ , the Cartan–Leray spectral sequence associated to the affinoid perfectoid  $\Gamma$ -cover  $\widetilde{X}_C \times S \rightarrow X_C \times S$ , combined with the vanishing theorem for  $\text{gr}^0 \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}$  gives

$$H_{\text{prokét}}^i(X_C \times S, \text{gr}^0 \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}) = H_{\text{cont}}^i(\Gamma, \text{gr}^0 \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}(X_C \times S))$$

for  $i \geq 0$ . Note that  $X_C \times S = \text{Spa}(\mathcal{C}^0(S, R_\infty), \mathcal{C}^0(S, R_\infty^+))$ . The same proof as in [DLLZ23a, Lemma 3.3.15] gives  $H_{\text{prokét}}^i(X_C \times S, \text{gr}^0 \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}) = 0$  for  $i > 0$  and

$$H_{\text{prokét}}^0(X_C \times S, \text{gr}^0 \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}) = \mathcal{C}^0(S, R) \widehat{\otimes}_K C.$$

The proposition then follows from the fact that  $\mathcal{C}^0(S, R) \widehat{\otimes}_K C \simeq \mathcal{C}^0(S, R \widehat{\otimes}_K C)$ , which follows from [PGS10, Corollary 10.5.4].  $\square$

**Proposition 4.8.** *Let  $X$  be a log smooth rigid analytic variety defined over  $K$ , with fine log-structure. Let  $(\mathcal{E}, \nabla, \text{Fil}^\bullet)$  be a filtered  $\mathcal{O}_X$ -module with integrable log connection, with associated  $\mathbb{B}_{\text{dR}}^+$ -local system  $\mathbb{M}^+ := \text{Fil}^0(\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log})^{\nabla=0}$ , and  $\mathbb{M} = \mathbb{M}^+[1/t]$ . Denote by  $\lambda$  the morphism of sites*

$$X_{\text{prokét}}/X_C \simeq X_{C, \text{prokét}} \rightarrow X_{C, \text{ét}, \text{cond}}.$$

Then, we have a natural quasi-isomorphism of complexes of sheaves valued in  $\text{Mod}_K^{\text{cond}}$  on  $X_{C, \text{ét}}$ :

$$(R\lambda_* \mathbb{M})^\nabla \simeq \underline{\text{logdR}}_X^\mathcal{E} \otimes_K^\blacksquare B_{\text{dR}}.$$

Moreover, the quasi-isomorphism is compatible with filtrations, where the filtration on the right side is given by the tensor product filtration.

*Proof.* Suppose  $X$  is of dimension  $n$  and connected. Denote by  $\mathcal{M}$  the  $\mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}^+$ -module with integrable log connection associated to  $(\mathcal{E}, \nabla, \text{Fil}^\bullet)$ , and

$$\mathcal{M}' := \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}^+} \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log} \simeq \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}_{\mathbb{B}_{\text{dR}}, \log}.$$

The logarithmic Poincaré lemma gives an exact sequence of sheaves on  $X_{C, \text{prokét}}$ :

$$0 \rightarrow \mathbb{M} \rightarrow \mathcal{M}' \xrightarrow{\nabla} \mathcal{M}' \otimes_{\mathcal{O}_X} \Omega_X^{\text{log}, 1} \xrightarrow{\nabla} \dots \xrightarrow{\nabla} \mathcal{M}' \otimes_{\mathcal{O}_X} \Omega_X^{\text{log}, n} \rightarrow 0,$$

which is also exact after taking  $\text{Fil}^r$ :

$$0 \rightarrow \text{Fil}^r \mathbb{M} \rightarrow \text{Fil}^r \mathcal{M}' \xrightarrow{\nabla} (\text{Fil}^{r-1} \mathcal{M}') \otimes_{\mathcal{O}_X} \Omega_X^{\text{log}, 1} \xrightarrow{\nabla} \dots \xrightarrow{\nabla} (\text{Fil}^{r-n} \mathcal{M}') \otimes_{\mathcal{O}_X} \Omega_X^{\text{log}, n} \rightarrow 0.$$

Therefore, we have a natural quasi-isomorphism between  $(R\lambda_*\mathbb{M})^\nabla$  and

$$\mathcal{C}^\bullet := R\lambda_* \left[ \mathcal{M}' \xrightarrow{\nabla} \mathcal{M}' \otimes_{\mathcal{O}_X} \Omega_X^{\log,1} \xrightarrow{\nabla} \dots \xrightarrow{\nabla} \mathcal{M}' \otimes_{\mathcal{O}_X} \Omega_X^{\log,n} \right]^\nabla,$$

which is compatible with filtrations.

We claim we have a natural filtered quasi-isomorphism

$$\underline{\log\mathrm{dR}}_X^\mathcal{E} \otimes_K^\blacksquare \underline{B}_{\mathrm{dR}} \xrightarrow{\simeq} \mathcal{C}^\bullet \quad (4.1)$$

of complexes of sheaves on  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$  (as explained in the Lemma 2.5 of [LZ17], we can regard these sheaves as sheaves over  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$ ). To construct this morphism, it suffices to define a natural morphism  $\mathcal{O}_X \otimes_K^\blacksquare \underline{B}_{\mathrm{dR}} \rightarrow (\lambda_*\mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}})^\nabla$  of sheaves on  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$ , which is compatible with filtration. This is already constructed in the proof of [Bos23a, Corollary 6.12] (also in [LZ17, Lemma 3.7] and [DLLZ23a, Lemma 3.3.2]), by noting that the pushforward of  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}}$  along the natural morphism of sites  $X_{C,\mathrm{prok}\acute{\mathrm{e}}\mathrm{t}} \rightarrow X_{C,\mathrm{pro}\acute{\mathrm{e}}\mathrm{t}}$  is  $\mathcal{O}_{\mathbb{B}_{\mathrm{dR}}}$ .

As explained in [Bos23a, Corollary 6.12], since the filtration on  $\underline{\log\mathrm{dR}}_X^\mathcal{E} \otimes_K^\blacksquare \underline{B}_{\mathrm{dR}}$  and  $\mathcal{C}^\bullet$  is exhaustive and complete, by [BMS19, Lemma 5.2], it suffices to show the natural morphism 4.1 is a quasi-isomorphism on graded pieces, which means that for any  $r \in \mathbb{Z}$ , we have a natural isomorphism

$$\mathrm{gr}^j(\underline{\log\mathrm{dR}}_X^\mathcal{E} \otimes_K^\blacksquare \mathrm{gr}^j \underline{B}_{\mathrm{dR}}) \rightarrow \mathrm{gr}^j \mathcal{C}^\bullet$$

on  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$ . Since (4.1) is a filtered morphism, it suffices to show that for any locally free  $\mathcal{O}_X$ -module  $\mathcal{F}$  on  $X_{\acute{\mathrm{e}}\mathrm{t}}$  of finite rank, we have a natural quasi-isomorphism

$$\underline{\mathcal{F}} \otimes_K^\blacksquare \underline{B}_{\mathrm{dR}} \xrightarrow{\simeq} R\lambda_* (\nu^* \mathcal{F} \otimes_{\mathcal{O}_X} \mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}})^\nabla$$

on  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$ , for any  $r \in \mathbb{Z}$ . Write  $\lambda : X_{C,\mathrm{prok}\acute{\mathrm{e}}\mathrm{t}} \rightarrow X_{C,\acute{\mathrm{e}}\mathrm{t},\mathrm{cond}}$  as

$$X_{C,\mathrm{prok}\acute{\mathrm{e}}\mathrm{t}} \xrightarrow{\alpha} X_{C,\mathrm{pro}\acute{\mathrm{e}}\mathrm{t}} \xrightarrow{\lambda'} X_{C,\acute{\mathrm{e}}\mathrm{t},\mathrm{cond}},$$

and write  $\nu : X_{\mathrm{prok}\acute{\mathrm{e}}\mathrm{t}} \rightarrow X_{\acute{\mathrm{e}}\mathrm{t}}$  as

$$X_{\mathrm{prok}\acute{\mathrm{e}}\mathrm{t}} \rightarrow X_{\mathrm{pro}\acute{\mathrm{e}}\mathrm{t}} \xrightarrow{\nu'} X_{\acute{\mathrm{e}}\mathrm{t}}.$$

According to the proof of [Bos23a, Corollary 6.12], we have a natural quasi-isomorphism

$$\underline{\mathcal{F}} \otimes_K^\blacksquare \mathrm{gr}^j \underline{B}_{\mathrm{dR}} \xrightarrow{\simeq} R\lambda'_* (\nu'^* \mathcal{F} \otimes_{\mathcal{O}_X} \mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR}}})^\nabla$$

on  $X_{C,\acute{\mathrm{e}}\mathrm{t}}$ , for any  $r \in \mathbb{Z}$ . Therefore, it suffices to show that we have a natural quasi-isomorphism

$$\nu'^* \mathcal{F} \otimes_{\mathcal{O}_X} \mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR}}} \xrightarrow{\simeq} R\alpha_* (\nu^* \mathcal{F} \otimes_{\mathcal{O}_X} \mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}})$$

on  $X_{C,\mathrm{pro}\acute{\mathrm{e}}\mathrm{t}}$ , for any  $r \in \mathbb{Z}$ . Since  $\mathcal{F}$  is locally free, it suffices to show that

$$\mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR}}} \rightarrow R\alpha_* \mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}}$$

is an isomorphism for any  $r \in \mathbb{Z}$ , which follows from the local description of  $\mathrm{gr}^j \mathcal{O}_{\mathbb{B}_{\mathrm{dR},\log}}$ , i.e., Corollary 3.10 (Note that  $R\alpha_* \mathbb{B}_{\mathrm{dR},\log} = \mathbb{B}_{\mathrm{dR}}$ ).  $\square$

Now we can prove Theorem 4.6.

*Proof of theorem 4.6.* The morphism of sites  $\lambda : X_{C, \text{prokét}} \rightarrow X_{C, \text{ét, cond}}$  factors as

$$\lambda : X_{C, \text{prokét}} \xrightarrow{\mu} X_{C, \text{prokét, cond}} \rightarrow X_{C, \text{ét, cond}}.$$

We have

$$\text{R}\Gamma_{\text{prokét}}(X_C, \mathbb{M}) \simeq \text{R}\Gamma_{\text{prokét, cond}}(X_C, R\mu_* \mathbb{M}) \simeq \text{R}\Gamma_{\text{ét, cond}}(X_C, R\lambda_* \mathbb{M}) \simeq \text{R}\Gamma_{\text{ét}}(X_C, (R\lambda_* \mathbb{M})^\nabla).$$

Let  $\epsilon : X_{C, \text{ét}} \rightarrow X_{\text{ét}}$  be the base change morphism, then the above property, combined with [Bos23a, Lemma 5.6] and [Bos23a, Lemma 6.13], we have

$$\text{R}\Gamma_{\text{prokét}}(X_C, \mathbb{M}) \simeq \text{R}\Gamma_{\text{ét}}(X_C, (R\lambda_* \mathbb{M})^\nabla) \simeq \text{R}\Gamma_{\text{ét}}(X_C, \underline{\text{logdR}}_X^\mathcal{E} \otimes_K^\blacksquare B_{\text{dR}}),$$

which conclude the proof of (i).

For the second part, it suffices to show that the natural morphism

$$\text{Fil}^r(\text{R}\Gamma_{\text{logdR}}(X, \mathcal{E}) \otimes_K^{L, \blacksquare} B_{\text{dR}}) \rightarrow \text{R}\Gamma(X, \text{Fil}^r(X_C, \underline{\text{logdR}}_X^\mathcal{E} \otimes_K^\blacksquare B_{\text{dR}})) \simeq \text{R}\Gamma_{\text{prokét}}(X_C, \text{Fil}^r \mathbb{M})$$

is a quasi-isomorphism. We may assume  $X$  is connected and paracompact, then the claim follows from [Bos23a, Theorem 5.20].  $\square$

**Remark 4.9.** If  $X$  is a quasi-compact smooth rigid analytic varieties over  $K$ , we have a  $\mathcal{G}_K$ -equivariant, compatible with filtrations, natural isomorphisms in  $D(\text{Mod}_K^{\text{cond}})$ :

$$\begin{aligned} \text{R}\Gamma_{\text{prokét}}(X_C, \mathbb{M}) &\simeq \text{colim}_{j \in \mathbb{N}} \text{R}\Gamma_{\text{prokét}}(X_C, \text{Fil}^{-j} \mathbb{M}) \\ &\simeq \text{colim}_{j \in \mathbb{N}} \text{Fil}^{-j}(\text{R}\Gamma_{\text{logdR}}(X, \mathcal{E}) \otimes_K^{L, \blacksquare} B_{\text{dR}}) \\ &\simeq \text{R}\Gamma_{\text{logdR}}(X, \mathcal{E}) \otimes_K^{L, \blacksquare} B_{\text{dR}}. \end{aligned}$$

The first isomorphism follows from the fact that the site  $X_{\text{prokét}}$  is coherent, see [DLLZ23b, Proposition 5.1.5]. The second one is the above theorem. The last one follows from the fact that  $\otimes_K^{L, \blacksquare}$  commutes with filtered colimits, and filtered colimits are exact in  $\text{Mod}_K^{\text{cond}}$ .

**Remark 4.10.** This theorem allows us to compare the pro étale cohomology and pro Kummer étale cohomology of period sheaves. Suppose that  $X$  is a smooth rigid analytic varieties over  $K$ ,  $D \subset X$  is a strictly normal crossing divisor, and  $U = X - D$ . Endow  $X$  with the log-structure coming from  $D$ . Then the canonical morphism

$$\text{R}\Gamma_{\text{prokét}}(X, \mathbb{M}) \xrightarrow{\simeq} \text{R}\Gamma_{\text{proét}}(U, \mathbb{M})$$

is a (non filtered) quasi-isomorphisms in  $D(\text{Mod}_K^{\text{cond}})$ . To see this, we can assume that  $X$  is quasi-compact, then the claim follows from the above remark, Theorem 2.8 and [Bos23a, Remark 6.14].

**Remark 4.11.** One can define filtered  $\mathcal{O}_{X_{\text{két}}}$ -modules with integrable connections over  $X_{\text{két}}$  in the same way, leading to analogous results. In fact, by [DLLZ23b, Definition 4.3.6], a coherent  $\mathcal{O}_{X_{\text{két}}}$ -module is locally the inverse image of a coherent sheaf on the analytic site of  $X$ .

**Remark 4.12.** If  $X$  is a log smooth, proper rigid analytic varieties over  $K$ ,  $(\mathcal{E}, \nabla, \text{Fil}^\bullet)$  be a filtered  $\mathcal{O}_X$ -module with integrable log connection, then by Lemma 2.2 and the finiteness of coherent sheaf

cohomology on proper rigid analytic varieties, we have  $H_{\log dR}^i(X, \mathcal{E}) = \underline{H}_{\log dR}^i(X, \mathcal{E})$  for all  $i \geq 0$ . Therefore

$$H_{\log dR}^i(X, \mathcal{E}) \otimes_K^{L\blacksquare} B_{dR} = \underline{H}_{\log dR}^i(X, \mathcal{E}) \otimes_K B_{dR}.$$

In particular, this recovers the comparison theorem in [DLLZ23a, Theorem 3.2.7], where they generalize [Sch13a, Corollary 1.8].

## 5. LOGARITHMIC $B_{dR}^+$ -COHOMOLOGY FOR RIGID ANALYTIC VARIETIES

In this chapter, we focus on logarithmic rigid analytic varieties over  $C$ . As outlined in the introduction, establishing a meaningful log de Rham-étale comparison theorem requires developing a  $B_{dR}^+$ -cohomology theory for such varieties. To achieve this, we adopt the approaches presented in [BMS18] and [Bos23b].

This chapter is dedicated to proving Theorem 1.4, which is essential in formulating the semistable conjecture for the geometric case as stated in [Sha25]. To maintain focus, we limit our exploration of logarithmic  $B_{dR}^+$ -cohomology to the essentials. A more comprehensive study of logarithmic  $B_{dR}^+$ -cohomology, along with logarithmic  $B$ -cohomology, will be presented in a subsequent article.

**5.1. Preliminary: The  $L\eta_f$  functor.** We review a construction introduced in [BO78] and generalized in [BMS18], which is essential for the construction of logarithmic  $B_{dR}^+$ -cohomology.

Let  $(T, \mathcal{O}_T)$  be a ringed topos. Denote by  $D(\mathcal{O}_T)$  (resp.  $K(\mathcal{O}_T)$ ) the derived category (resp. homotopy category) of  $\mathcal{O}_T$ -modules. Let  $(f) \subset \mathcal{O}_T$  be an invertible ideal sheaf.

**Definition 5.1.** [BO78, Definition 8.6] Let  $M^\bullet \in K(\mathcal{O}_T)$  be a  $f$ -torsion-free complex of  $\mathcal{O}_T$ -modules. Define a new  $f$ -torsion-free complex  $\eta_f M^\bullet \in K(\mathcal{O}_T)$  as

$$\eta_f(M^\bullet)^i := \{x \in f^i M^i \mid dx \in f^{i+1} M^{i+1}\}$$

for all  $i \in \mathbb{Z}$ .

The definition of  $\eta_f$  depends only on the ideal sheaf  $(f)$ , and it is independent of the generator of  $(f)$ . According to [BO78, Proposition 8.19],  $\eta_f$  factors canonically over a (not triangulated!) functor

$$L\eta_f : D(\mathcal{O}_T) \rightarrow D(\mathcal{O}_T)$$

on derived categories.

We now define a filtration on  $L\eta_f M$ . We refer the reader to [BMS19, Chapter 5] for notations and basic proportions on filtered derived  $\infty$ -category of  $R$ -modules. Recall that the filtered derived  $\infty$ -category of  $\mathcal{O}_T$ -modules is defined to be

$$DF(\mathcal{O}_T) := \text{Fun}(\mathbb{Z}^{\text{op}}, D(\mathcal{O}_T)).$$

Let  $F \in DF(\mathcal{O}_T)$ , we denote by

$$\text{gr}^i(F) := F(i)/F(i+1)$$

the  $i$ -th graded piece of  $F$ .

**Definition 5.2.** Let  $M \in D(\mathcal{O}_T)$ , we define  $\text{Fil}^* L\eta_f M$  a filtration on  $L\eta_f M$  as follows: choose a representation  $M^\bullet \in K(\mathcal{O}_T)$  of  $M$  such that each  $M^i$  is  $f$ -torsion-free. Then the  $i$ -th level of  $\text{Fil}^* L\eta_f M$  is given by

$$\text{Fil}^i L\eta_f M := f^i M^\bullet \cap \eta_f M^\bullet.$$

To understand the filtration on  $L\eta_f M$ , we recall the following definition introduced in [BMS19], an elaboration of [Bei87, Appendix A].

**Definition 5.3.** [BMS19, Definition 5.3] Let  $DF^{\leq 0}(\mathcal{O}_T) \subset DF(\mathcal{O}_T)$  denote the full subcategory consisting of objects  $F$  such that  $\text{gr}^i(F) \in D^{\leq i}(\mathcal{O}_T)$  for all  $i$ . Dually,  $DF^{\geq 0}(\mathcal{O}_T) \subset DF(\mathcal{O}_T)$  is the full subcategory consisting of objects  $F$  such that  $F(i) \in D^{\geq i}(\mathcal{O}_T)$  for all  $i$ . We call the pair  $(DF^{\leq 0}(\mathcal{O}_T), DF^{\geq 0}(\mathcal{O}_T))$  the Beilinson  $t$ -structure on  $DF(\mathcal{O}_T)$ .

This definition is justified by Theorem 5.4 below.

**Theorem 5.4.** [BMS19, Theorem 5.4] (1) The Beilinson  $t$ -structure  $(DF^{\leq 0}(\mathcal{O}_T), DF^{\geq 0}(\mathcal{O}_T))$  is a  $t$ -structure on  $DF(\mathcal{O}_T)$ .

(2) Let  $\tau_B^{\leq 0}$  denote the connective cover functor associated with the  $t$ -structure from (1). Then, there exists a natural isomorphism

$$\text{gr}^i \circ \tau_B^{\leq 0}(-) \simeq \tau^{\leq i} \circ \text{gr}^i(-).$$

(3) The heart of the Beilinson  $t$ -structure, defined as

$$DF(\mathcal{O}_T)^\heartsuit := DF^{\leq 0}(\mathcal{O}_T) \cap DF^{\geq 0}(\mathcal{O}_T),$$

which is equivalent to the abelian category  $\text{Ch}(\mathcal{O}_T)$  of chain complexes of  $\mathcal{O}_T$ -modules. This equivalence can be described as follows: for  $F \in DF(\mathcal{O}_T)$ , its 0-th cohomology  $H_B^0(F)$  in the Beilinson  $t$ -structure corresponds to the chain complex  $(H^\bullet(\text{gr}^\bullet(F)), d)$ , where  $d$  is the boundary map induced by the exact triangle

$$\text{gr}^{i+1}(F) = F(i+1)/F(i+2) \rightarrow F(i)/F(i+2) \rightarrow F(i)/F(i+1) = \text{gr}^i(F).$$

We cite the following proposition of [BMS19], which describe the structure of the filtration of the décalage functor.

**Proposition 5.5.** [BMS19, Proposition 5.8] Fix  $M \in D(\mathcal{O}_T)$ . Let  $f^* \otimes M$  be the  $f$ -adic filtration on  $M$ , i.e., the  $i$ -th level of the filtration of  $K$  is  $f^i \otimes M$  with obvious maps. Then  $L\eta_f M$  is filtered isomorphic with  $\tau_B^{\leq 0}(f^* \otimes M)$  in  $DF(\mathcal{O}_T)$ .

We list some results of  $L\eta_f$ , which will be used in the rest of this chapter.

**Proposition 5.6.** (1)[BMS18, Lemma 6.10] If  $M \in D^{\geq 0}(\mathcal{O}_T)$  such that  $H^0(M)[f] = 0$ . Then there is a canonical map  $\text{Fil}^* L\eta_f M \rightarrow f^* \otimes M$  in  $DF(\mathcal{O}_T)$ .

(2)[BMS18, Proposition 6.12] Let  $M \in D(\mathcal{O}_T)$ , construct a complex  $H^\bullet(M/(t))$  with terms

$$H^i(M/(t)) := H^i(M/L(t)) \otimes_{\mathcal{O}_T} (t^i),$$

and with differential induced by the boundary map corresponding to the short exact sequence

$$0 \rightarrow (t)/(t^2) \rightarrow \mathcal{O}_T/(t^2) \rightarrow \mathcal{O}_T/(t) \rightarrow 0.$$

Then there is a natural quasi-isomorphism

$$L\eta_f M \otimes_{\mathcal{O}_T}^L \mathcal{O}_T/(t) \xrightarrow{\simeq} H^\bullet(M/(t)).$$

(3)[BMS18, Lemma 6.14] Let  $u : (T, \mathcal{O}_T) \rightarrow (T', \mathcal{O}_{T'})$  be a flat map of ringed topoi. Let  $(f) \subset \mathcal{O}_T$  be an invertible ideal sheaf with  $(f') := u^*(f) \subset \mathcal{O}_{T'}$ , which is still invertible. Then the diagram

$$\begin{array}{ccc} D(\mathcal{O}_T) & \xrightarrow{u^*} & D(\mathcal{O}_{T'}) \\ \downarrow L\eta_f & & \downarrow L\eta_{f'} \\ D(\mathcal{O}_T) & \xrightarrow{u^*} & D(\mathcal{O}_{T'}) \end{array}$$

commutes, i.e., there is a natural quasi-isomorphism  $L\eta_{f'} u^* M \rightarrow u^* L\eta_f M$  in  $D(\mathcal{O}_{T'})$  for all  $M \in D(\mathcal{O}_T)$ .

*Proof.* The compatibility for filtration of (1) follows from the corresponding proof in [BMS18].  $\square$

**5.2. Logarithmic  $B_{\mathrm{dR}}^+$ -cohomology.** Similar to [Bos23b], we can define the logarithmic  $B_{\mathrm{dR}}^+$ -cohomology by using the  $L\eta_f$  functor.

**Definition 5.7.** Let  $X$  be a log smooth rigid analytic varieties over  $C$ . We denote by  $\lambda : X_{\mathrm{prok\acute{e}t}} \rightarrow X_{\acute{e}t, \mathrm{cond}}$  the natural morphism of sites. We define the logarithmic  $B_{\mathrm{dR}}^+$ -cohomology of  $X$  as the complex in  $D(\mathrm{Mod}_{B_{\mathrm{dR}}^+}^{\mathrm{solid}})$ :

$$\mathrm{R}\Gamma_{B_{\mathrm{dR}}^+}(X) := \mathrm{R}\Gamma_{\acute{e}t, \mathrm{cond}}(X, L\eta_t R\lambda_* \mathbb{B}_{\mathrm{dR}}^+).$$

We endow  $\mathrm{R}\Gamma_{B_{\mathrm{dR}}^+}(X)$  with the filtration given by Definition 5.2.

**Remark 5.8.** By introducing the logarithmic infinitesimal sites, one is able to define  $\mathrm{R}\Gamma_{\mathrm{logdR}}(X/B_{\mathrm{dR}}^+)$ , and there should be a canonical filtered quasi-isomorphism:

$$\mathrm{R}\Gamma_{B_{\mathrm{dR}}^+}(X) \simeq \mathrm{R}\Gamma_{\mathrm{logdR}}(X/B_{\mathrm{dR}}^+).$$

**Proposition 5.9.** Let  $X$  be a log smooth rigid analytic varieties of dimension  $d$  over  $C$ , then  $R^i \lambda_* \mathbb{B}_{\mathrm{dR}}^+ = 0$  for all  $i > d$ .

*Proof.* The problem is local, following the notations in Section 3, we may assume  $X = \mathrm{Spa}(R, R^+)$  is an affinoid log rigid space over  $C$ , with a strictly étale morphism  $X \rightarrow \mathbb{E}$ , which can be written as a composite of rational embeddings and finite étale maps. By pulling back  $\widehat{\mathbb{E}} \rightarrow \mathbb{E}$  to  $X$  we get

$$\widetilde{X} := X \times_{\mathbb{E}} \widehat{\mathbb{E}} = \mathrm{Spa}(R_\infty, R_\infty^+),$$

a Galois pro-finite Kummer étale cover with Galois group  $\Gamma$ . The Cartan–Leray spectral sequence gives an quasi-isomorphism

$$\mathrm{R}\Gamma_{\mathrm{cond}}(\Gamma, H^0(\widetilde{X}, \mathbb{B}_{\mathrm{dR}})) \xrightarrow{\simeq} \mathrm{R}\Gamma_{\mathrm{prok\acute{e}t}}(X, \mathbb{B}_{\mathrm{dR}}),$$

which concludes the proof since  $\Gamma \simeq \widehat{\mathbb{Z}}(1)^d$  has cohomological dimension  $d$  by [Bos23a, B.3].  $\square$

This proposition allows us to deduce the following two propositions.



**Proposition 5.10.** *The natural map*

$$L\eta_t R\lambda_* \mathbb{B}_{\text{dR}} \rightarrow R\varprojlim_m L\eta_t R\lambda_*(\mathbb{B}_{\text{dR}}/\text{Fil}^m)$$

*is a quasi-isomorphism compatible with filtration.*

*Proof.* The proof is similar to [Bra18, Lemma 4.3] or [Bos23b, Lemma 2.41].  $\square$

**Proposition 5.11.** *If  $X$  is an affinoid log smooth rigid analytic variety over  $C$ , the natural morphism*

$$L\eta_t \text{R}\Gamma_{\text{prokét}}(X, \mathbb{B}_{\text{dR}}^+) \rightarrow \text{R}\Gamma_{\text{ét, cond}}(X, L\eta_t R\lambda_* \mathbb{B}_{\text{dR}}^+) = \text{R}\Gamma_{B_{\text{dR}}^+}(X)$$

*in  $D(\text{Mod}_{B_{\text{dR}}^+}^{\text{solid}})$  is a filtered quasi-isomorphism, where the filtration on both sides are given by the filtration of  $L\eta_t$  defined above.*

*Proof.* The proof is similar to [Bra18, Proposition 3.11] or [Bos23b, Proposition 2.42].  $\square$

**Theorem 5.12.** *Let  $X$  be a log smooth rigid analytic varieties over  $K$  then we have a natural isomorphism in  $D(\text{Mod}_K^{\text{solid}})$ :*

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X_C) \xrightarrow{\simeq} \text{R}\Gamma_{\text{logdR}}(X) \otimes_K^{L\blacksquare} B_{\text{dR}}^+,$$

*compatible with filtration and  $G_K$ -action.*

*Proof.* We may assume that  $X$  is affinoid. In this case, we have

$$\begin{aligned} \text{R}\Gamma_{B_{\text{dR}}^+}(X_C) &\simeq L\eta_t \text{R}\Gamma_{\text{prokét}}(X_C, \mathbb{B}_{\text{dR}}^+) \\ &\simeq L\eta_t \text{R}\Gamma_{\text{ét}}(X_C, (R\lambda_* B_{\text{dR}}^+)^{\vee}) \\ &\simeq L\eta_t \text{R}\Gamma_{\text{ét}}(X, \text{Fil}^0(\text{logdR}_X^{\mathcal{E}} \otimes_K^{\blacksquare} B_{\text{dR}})) \\ &\simeq \left( \underline{\mathcal{O}}_X(X) \rightarrow \underline{\Omega}_X^{1, \text{log}}(X) \rightarrow \underline{\Omega}_X^{2, \text{log}}(X) \rightarrow \dots \right) \otimes_K^{L\blacksquare} B_{\text{dR}}^+ \\ &\simeq \text{R}\Gamma_{\text{logdR}}(X) \otimes_K^{L\blacksquare} B_{\text{dR}}^+, \end{aligned}$$

where the first quasi-isomorphism follows from Proposition 5.11, the second quasi-isomorphism follows from the same argument as in the proof of Theorem 4.6, the third quasi-isomorphism follows from Proposition 4.8, the fourth quasi-isomorphism follows from [Bos23a, Lemma 6.13] and [Bos23a, Theorem 5.20], and the last quasi-isomorphism follows from [Bos23a, Lemma 5.6]. This concludes the proof.  $\square$

When  $X$  is defined over  $\mathbb{C}$ , our construction of the  $B_{\text{dR}}^+$ -cohomology theory indeed provides a deformation of the logarithmic de Rham cohomology. Moreover, we can describe its filtration in a manner analogous to [CN25, Proposition 3.13] in the rigid analytic setting, as follows.

**Theorem 5.13.** (1) *Let  $X$  be a log smooth rigid analytic varieties over  $C$ , then we have a natural isomorphism in  $D(\text{Mod}_C^{\text{solid}})$ :*

$$\theta : \text{R}\Gamma_{B_{\text{dR}}^+}(X) \otimes_{B_{\text{dR}}^+}^{L\blacksquare} C \xrightarrow{\simeq} \text{R}\Gamma_{\text{logdR}}(X).$$

(2) More generally, for  $r \geq 0$ , we have a natural distinguished triangle in  $D(\text{Mod}_C^{\text{solid}})$ :

$$\text{Fil}^{r-1} \text{R}\Gamma_{B_{\text{dR}}^+}(X) \xrightarrow{t} \text{Fil}^r \text{R}\Gamma_{B_{\text{dR}}^+}(X) \xrightarrow{\theta} \text{Fil}^r \text{R}\Gamma_{\log\text{dR}}(X).$$

(3) For  $r \geq 0$ , we have a natural distinguished triangle in  $D(\text{Mod}_C^{\text{solid}})$ :

$$\text{Fil}^{r+1} \text{R}\Gamma_{B_{\text{dR}}^+}(X) \rightarrow \text{Fil}^r \text{R}\Gamma_{B_{\text{dR}}^+}(X) \rightarrow \text{R}\Gamma(X, \tau^{\leq r} R\lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}(r)).$$

*Proof.* We have

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X) \otimes_{B_{\text{dR}}^+}^{L\blacksquare} C \simeq \text{R}\Gamma_{\text{ét,cond}}(X, L\eta_t R\lambda_* \mathbb{B}_{\text{dR}}^+ \otimes_{B_{\text{dR}}^+}^{L\blacksquare} (B_{\text{dR}}^+/t)).$$

We are reduced to showing

$$\left( L\eta_t R\lambda_* \mathbb{B}_{\text{dR}}^+ \otimes_{B_{\text{dR}}^+}^{L\blacksquare} (B_{\text{dR}}^+/t) \right)^\blacktriangledown \simeq \underline{\Omega_X^{\bullet,\log}}.$$

By Proposition 5.6 (2), we have

$$\begin{aligned} \left( L\eta_t R\lambda_* \mathbb{B}_{\text{dR}}^+ \otimes_{B_{\text{dR}}^+}^{L\blacksquare} (B_{\text{dR}}^+/t) \right)^\blacktriangledown &\simeq H^\bullet((R\lambda_* \mathbb{B}_{\text{dR}}^+)/t)^\blacktriangledown \simeq H^\bullet(R\lambda_*(\mathbb{B}_{\text{dR}}^+/t))^\blacktriangledown \simeq H^\bullet(R\lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}})^\blacktriangledown \\ &\simeq \left( R^\bullet \lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}} \right)^\blacktriangledown \otimes_{B_{\text{dR}}^+} \text{Fil}^\bullet B_{\text{dR}}^+ / \text{Fil}^{\bullet+1} B_{\text{dR}}^+ \\ &\simeq \underline{\Omega_X^{\bullet,\log}(-\bullet)} \otimes_C C(\bullet) \\ &\simeq \underline{\Omega_X^{\bullet,\log}(-\bullet)}, \end{aligned}$$

where the second-to-last isomorphism follows from the proposition below. This proves (1). Then (2) follows from [Wu24, Proposition 3.4], and (3) follows from [Wu24, Lemma 3.3].  $\square$

**Remark 5.14.** We will see that when  $X$  descends to  $K$ , we have a natural quasi-isomorphism

$$\tau^{\leq r} R\lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}(r) \simeq \bigoplus_{i \leq r} \Omega_X^{i,\log}(r-i)[-i],$$

and in general, a choice of a log smooth  $B_{\text{dR}}^+/t^2$ -lift  $\mathbb{X}$  of  $X$  via the map  $B_{\text{dR}}^+/t^2 \rightarrow C$  gives such a quasi-isomorphism.

**Proposition 5.15.** *Let  $X$  be a log smooth rigid analytic varieties over  $C$ , then we have a natural isomorphism*

$$\left( R^i \lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}} \right)^\blacktriangledown \simeq \underline{\Omega_X^{i,\log}(-i)}$$

for all  $i \geq 0$ .

*Proof.* The same argument as in the proof of [Sch13b, Proposition 3.23] applies, with a log-version of [Sch13b, Lemma 3.24] stated below.  $\square$

**Lemma 5.16.** *Consider the exact sequence (for example, see [KN99, Proposition 2.3])*

$$0 \rightarrow \mathbb{Z}_p(1) \rightarrow \varprojlim_{\times p} \nu^{-1} \mathcal{M}_X^{gp} \rightarrow \nu^{-1} \mathcal{M}_X^{gp} \rightarrow 0$$

on  $X_{\text{prokét}}$ . Here  $\nu$  is the natural morphism of sites  $\nu : X_{\text{prokét}} \rightarrow X_{\text{ét}}$ . It induces a boundary map  $\mathcal{M}_X^{gp} = \nu_* \nu^{-1} \mathcal{M}_X^{gp} \rightarrow R^1 \lambda_* \mathbb{Z}_p(1)$ . Then there is a unique  $\underline{\mathcal{O}}_X$ -linear map  $\underline{\Omega}_X^{\text{log}} \rightarrow R^1 \lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}(1)^\nabla$  such that the diagram

$$\begin{array}{ccc} \mathcal{M}_X^{gp} & \longrightarrow & R^1 \nu_* \mathbb{Z}_p(1) \\ \downarrow \delta & & \downarrow \\ \underline{\Omega}_X^{\text{log}}(*) & \longrightarrow & R^1 \lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}(1)^\nabla(*) \end{array}$$

commutes. This map is an isomorphism.

*Proof.* Similar to [Sch13b, Lemma 3.24].  $\square$

Similar to the log de Rham cohomology, we can compare the logarithmic  $B_{\text{dR}}^+$ -cohomology with the  $B_{\text{dR}}^+$ -cohomology of the trivial locus when the log structure comes from a strictly normal crossing divisor.

**Proposition 5.17.** *Let  $X$  be a log smooth rigid analytic varieties over  $C$ , with log-structure giving by a strictly normal crossing divisor  $D \subset X$ . Denote by  $U := X - D$ . Then, we have a natural (non filtered!) isomorphism in  $D(\text{Mod}_{B_{\text{dR}}^+}^{\text{solid}})$ :*

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X) \simeq \text{R}\Gamma_{\text{dR}}(U/B_{\text{dR}}^+).$$

*Proof.* The problem is local, so we may assume  $X$  can be descent to  $X_0$  over a discrete valued field  $K$  by the lemma below. Then the claim follows from Theorem 5.12, [Bos23b, Theorem 5.1] and Theorem 2.8.  $\square$

**Lemma 5.18.** *If  $X$  is a log smooth rigid analytic varieties over  $C$  of dimension  $d$ , with log-structure giving by a strictly normal crossing divisor  $D \subset X$ . Then étale locally  $X$  has a basis of rigid analytic varieties of the form  $\text{Sp}(A_C \langle x_1, \dots, x_d \rangle)$  with log structure given by the normal crossing divisor  $x_1 \cdots x_d = 0$ , where  $\text{Sp}(A)$  is a smooth rigid analytic variety over  $K$  for some discrete valued field  $K$  in  $C$ .*

*Proof.* By the existence of tubular neighborhood, [Kie67, Theorem 1.18], locally  $X$  can be written as  $V := \text{Sp}(R \langle x_1, \dots, x_d \rangle)$ , and  $D$  is defined by the equation  $x_1 \cdots x_d = 0$ . Then the claim follows from Temkin's alteration theorem [Tem17, Theorem 3.3.1].  $\square$

**5.3. Comparison with Kummer pro-étale cohomology.** The following comparison theorem can be easily deduced from our construction for logarithmic  $B_{\text{dR}}^+$ -cohomology.

**Theorem 5.19.** *Let  $X$  be a log smooth proper rigid analytic varieties over  $C$ . Then we have a canonical filtered isomorphism*

$$H_{\text{két}}^i(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \simeq H_{B_{\text{dR}}^+}^i(X) \otimes_{B_{\text{dR}}^+} B_{\text{dR}}.$$

Here, the filtration on  $H_{B_{\text{dR}}^+}^i(X)$  is defined by

$$\text{Fil}^* H_{B_{\text{dR}}^+}^i(X) := \text{Im}(H^i(\text{Fil}^* \text{R}\Gamma_{B_{\text{dR}}^+}(X)) \rightarrow H_{B_{\text{dR}}^+}^i(X)).$$

Moreover, when  $X$  descends to  $X_0$  over a discrete valued field  $K$ , this isomorphism agrees with the comparison isomorphism

$$H_{\text{két}}^i(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \simeq H_{\text{logdR}}^i(X_0) \otimes_K B_{\text{dR}},$$

under a canonical identification in Theorem 5.12.

*Proof.* We will construct a morphism:

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X) \rightarrow \text{R}\Gamma_{\text{prokét}}(X, \mathbb{B}_{\text{dR}}^+) \simeq \text{R}\Gamma_{\text{két}}(X, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} B_{\text{dR}}^+,$$

where the last isomorphism follows from [DLLZ23a, Lemma 3.6.1] (which also works over  $C$ , by the same proof as presented in [Sch13a, Theorem 8.4]). Then we will prove that such a morphism is a quasi-isomorphism after inverting  $t$ .

By Proposition 5.6 (1) and Proposition 5.11, we have a natural morphism

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X) = \text{R}\Gamma_{\text{ét,cond}}(X, L\eta_t R\lambda_* \mathbb{B}_{\text{dR}}^+) \rightarrow \text{R}\Gamma_{\text{ét,cond}}(X, R\lambda_* \mathbb{B}_{\text{dR}}^+) = \text{R}\Gamma_{\text{prokét}}(X, \mathbb{B}_{\text{dR}}^+),$$

which gives the desired morphism.

To show that such a morphism is a quasi-isomorphism after inverting  $t$ , it suffices to prove it locally so we can assume that  $X$  is affinoid. By Proposition 5.6 (3), considering the base change  $B_{\text{dR}}^+ \rightarrow B_{\text{dR}}$ , we have a quasi-isomorphism

$$\text{R}\Gamma_{B_{\text{dR}}^+}(X) \left[ \frac{1}{t} \right] \simeq L\eta_t \left( \text{R}\Gamma_{\text{prokét}}(X, \mathbb{B}_{\text{dR}}^+) \left[ \frac{1}{t} \right] \right) \simeq \text{R}\Gamma_{\text{prokét}}(X, \mathbb{B}_{\text{dR}}),$$

which is clearly compatible with filtration. This concludes the proof.

When  $X$  descends to  $X_0$  over a discrete valued field  $K$ , the compatibility stated in the theorem is clear due to Theorem 5.12.  $\square$

**5.4. Degeneration of (log) Hodge–Tate spectral sequence.** This section is devoted to proving the following theorem, as promised in the introduction. For convenience, we will only work with usual cohomology groups in this section. However, clearly all results can be extended to condensed cohomology group without any difficulty.

Recall that we denote by  $\nu$  the natural morphism of sites  $\nu : X_{\text{prokét}} \rightarrow X_{\text{ét}}$ .

**Theorem 5.20.** *Let  $X$  be a proper log smooth rigid analytic variety over  $C$  of dimension  $d$ . If  $X$  admits a log smooth  $B_{\text{dR}}^+/t^2$ -lift  $\mathbb{X}$  via the map  $B_{\text{dR}}^+/t^2 \rightarrow C$ , then*

(i) *The Hodge–log de Rham spectral sequence*

$$E_1^{ij} := H^j(X, \Omega_X^{\text{log}, i}) \Rightarrow H_{\text{logdR}}^{i+j}(X)$$

*degenerates at  $E_1$ .*

(ii) *The Hodge–Tate spectral sequence*

$$E_2^{ij} := H^j(X, \Omega_X^{\text{log}, i})(-j) \Rightarrow H_{\text{két}}^{i+j}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C$$

*degenerates at  $E_2$ .*

*Moreover,  $H_{\text{logdR}}^i(X/B_{\text{dR}}^+)$  is a finite free  $B_{\text{dR}}^+$ -module.*

The key step in the proof is to demonstrate the existence of a splitting

$$\bigoplus_{i \geq 0} \Omega_X^{i, \log}(-i)[-i] \xrightarrow{\sim} R\nu_* \hat{\mathcal{O}}_{X_{\mathrm{prokét}}}.$$

For rigid analytic varieties, this is shown in [Guo19, Proposition 7.2.5] by choosing a smooth  $B_{\mathrm{dR}}^+/t^2$ -lifting  $\mathbb{X}$  of  $X$ . Such a lifting always exists when  $X$  is proper. Additionally, the splitting is functorial depending on the choice of lifting.

We now turn our attention to the logarithmic case, discussing conditions under which such a splitting exists.

5.4.1.  $\mathbb{B}_{\mathrm{dR}}^+/t^2$ -liftings of  $X$ . As demonstrated in [BMS18] or [Guo19], a key step in the proof of the theorem is to establish the existence of a  $\mathbb{B}_{\mathrm{dR}}^+/t^2$ -lifting for proper  $X$ . For log adic space, in general such a lifting is more difficult to construct. We will show the following proposition.

**Proposition 5.21.** *Let  $X$  be a quasi-compact log smooth rigid analytic varieties over  $C$ . Then  $X$  admits a log smooth  $B_{\mathrm{dR}}^+/t^2$ -lift  $\mathbb{X}$  via the map  $B_{\mathrm{dR}}^+/t^2 \rightarrow C$  in the following cases:*

- (1)  $X$  descends to  $X_0$  over a discrete valued field  $K$ ;
- (2)  $X$  is the analytification of a finite type log algebraic variety  $X^{\mathrm{alg}}$  log smooth over  $C$ ;
- (3)  $X$  admits a proper log smooth model over  $\mathrm{Spf}(\mathcal{O}_C)$  or  $\mathrm{Spf}(\mathcal{O}_C^\omega)$ . Here  $\mathrm{Spf}(\mathcal{O}_C^\omega)$  is the formal log scheme  $\mathrm{Spf}(\mathcal{O}_C)$  with log-structure induced by  $\mathbb{N} \rightarrow \mathcal{O}_C, 1 \mapsto \omega$  for some pseudo-uniformizer  $\omega \in \mathcal{O}_C$ . In particular, when  $X$  admits a proper semistable formal model over  $\mathrm{Spf}(\mathcal{O}_C)$ .

*Proof.* (1) is clear.

For (2), denote by  $\underline{U}^{\mathrm{alg}} \subset \underline{X}^{\mathrm{alg}}$  the trivial locus of the underlying space  $\underline{X}^{\mathrm{alg}}$ . By [Gro66a, Proposition 8.9.1] there exists a finitely generated field  $A$  over  $\mathbb{Q}$  such that  $\underline{U}^{\mathrm{alg}} \subset \underline{X}^{\mathrm{alg}}$  descends to  $\underline{U}_0^{\mathrm{alg}} \rightarrow \underline{X}_0^{\mathrm{alg}}$  over  $A$  (which is still an open immersion by faithful flat descent), where  $\underline{X}_0^{\mathrm{alg}}$  is finite type over  $A$ . This induces the compactifying log-structure on  $X_0^{\mathrm{alg}}$ , which is still log smooth. By embedding a transcendental basis of  $A$  over  $\mathbb{Q}$  into  $\mathbb{Q}_p$ , we can embed  $A$  into a finite  $\mathbb{Q}_p$  extension  $K$ . This reduces to (1).

For (3), choosing an inclusion of field  $\check{F} \rightarrow C$ , we will show that there exists a smooth rigid space  $\mathcal{S}$  over  $\check{F}$ , such that  $X$  can be lifted to a proper log smooth  $\check{F}$ -rigid space  $X_A$  over  $\mathcal{S}$  with a map  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow \mathcal{S}$ . Then since  $B_{\mathrm{dR}}^+/t^2 \rightarrow C$  has a natural  $\check{F}$ -structure, by formal smoothness the map  $\mathrm{Spa}(C, \mathcal{O}_C) \rightarrow \mathcal{S}$  can be lifted to a map  $\mathrm{Spa}(B_{\mathrm{dR}}/t^2, B_{\mathrm{dR}}^+/t^2) \rightarrow \mathcal{S}$ . This will finish the proof by taking the base change of  $X_A$ .

We will prove the case when  $X$  admits a proper log smooth model over  $\mathrm{Spf}(\mathcal{O}_C^\omega)$ ; the proof for  $X$  over  $\mathrm{Spf}(\mathcal{O}_C)$  is similar and easier. Let  $\mathfrak{X}$  be a proper log smooth model over  $\mathrm{Spf}(\mathcal{O}_C^\omega)$  with generic fiber  $X$ . Denote by  $\mathfrak{X}_s$  its special fiber. Note that  $\mathfrak{X}_s$  is log smooth over  $\mathrm{Spec}(\bar{k})^0$ . Here  $\mathrm{Spec}(\bar{k})^0$  is the log scheme  $\mathrm{Spec}(\bar{k})$  with log-structure induced by  $\mathbb{N} \rightarrow \bar{k}, 1 \mapsto 0$ .

Denote by  $\mathcal{C}_{\mathcal{O}_{\check{F}}}[[\mathbb{N}]]$  the category of complete local artin  $\mathcal{O}_{\check{F}}[[\mathbb{N}]$ -algebras with residue field  $\bar{k}$ . Consider the log smooth deformation functor

$$\mathbf{LD} : \mathcal{C}_{\mathcal{O}_{\check{F}}}[[\mathbb{N}]] \rightarrow \mathbf{Set}$$

defined by sending  $A \in \mathcal{C}_{\mathcal{O}_{\check{F}}}[[\mathbb{N}]]$  to the isomorphism class of log smooth lifting of  $\mathfrak{X}_s \rightarrow \mathrm{Spec}(\bar{k})^0$  on  $A$ . According to [Kat96, Theorem 8.7], the log smooth deformation functor  $\mathbf{LD}$  has a versal

deformation, i.e., a complete artinian local  $\mathcal{O}_{\check{F}}[[\mathbb{N}]]$ -algebra  $A$  with the residue field  $\bar{k}$ , with the natural log-structure, a proper log smooth adic formal scheme  $\mathfrak{X}_A$  over  $A$  where  $A$  is topologized by powers of its maximal ideal, deforming  $\mathfrak{X}_s \rightarrow \mathrm{Spec}(\bar{k})^0$  such that the induced classifying map

$$h_A : \mathrm{Hom}_{\mathcal{O}_{\check{F}}[[\mathbb{N}]]}(A, -) \rightarrow \mathbf{LD}$$

is formally smooth. According to the proof of [BMS18, Proposition 13.15], we can extend  $h_A, \mathbf{LD}$  to the ind-completion of  $\mathcal{C}_{\mathcal{O}_{\check{F}}[[\mathbb{N}]}}$ . This category includes  $\mathcal{O}_C/\omega^k$  for all  $k \geq 1$ , where  $\mathcal{O}_C/\omega^k$  is regarded as a  $\mathcal{O}_{\check{F}}[[\mathbb{N}]]$ -algebra via  $\mathbb{N} \rightarrow \mathcal{O}_C/\omega^k, 1 \mapsto \omega$ . Denote by  $\mathfrak{X}_k$  the reduction of  $\mathfrak{X}$  module  $\omega^k$ . Since we have a canonical map  $\eta_0 : A \rightarrow \bar{k}$  keeping log-structures, and an isomorphism  $\psi_0 : \eta_0^* \mathfrak{X}_A \simeq \mathfrak{X}_s$ , applying the formal smoothness of  $h_A$  (on the ind-completion of  $\mathcal{C}_{\mathcal{O}_{\check{F}}[[\mathbb{N}]}}$ ) to the  $\mathcal{O}_C/\omega \rightarrow \bar{k}$ , we can choose a map  $\eta_1 : A \rightarrow \mathcal{O}_C/\omega$  lifting  $\eta_0$  and an isomorphism  $\psi_1 : \eta_1^* \mathfrak{X}_A \simeq \mathfrak{X}_1$  of log  $\mathcal{O}_C/\omega$ -schemes lifting  $\psi_0$ . We can then construct  $\psi_n, \eta_n$  inductively, taking the inverse limit, and we can show that there exist a  $\mathcal{O}_{\check{F}}[[\mathbb{N}]]$ -algebra map  $\eta : A \rightarrow \mathcal{O}_C$  and an isomorphism  $\eta^* \mathfrak{X}_A \simeq \mathfrak{X}$  as formal log  $\mathcal{O}_C$ -schemes. Here we view  $\mathcal{O}_C$  as  $\mathcal{O}_{\check{F}}[[\mathbb{N}]]$ -algebra via  $\mathbb{N} \rightarrow \mathcal{O}_C, 1 \mapsto \omega$ .

Finally, after inverting  $p$  we have a log  $\check{F}$ -rigid space  $X_A$  log smooth over  $\mathcal{S} := \mathrm{Spa}(A, A[1/p])$ . We can shrink  $\mathcal{S}$  to a suitable small closed subset, such that  $\mathcal{S}$  is smooth over  $\check{F}$ . This concludes the proof.  $\square$

5.4.2. *Splitting of the (log) Hodge–Tate map.* We will prove the following proposition.

**Proposition 5.22.** *Let  $X$  be a log smooth rigid analytic varieties over  $C$ . Then a choice of log smooth  $B_{\mathrm{dR}}^+/t^2$ -lift  $\mathbb{X}$  of  $X$  gives a natural splitting of the (log) Hodge-Tate map*

$$\mathrm{HT} : H_{\mathrm{prok\acute{e}t}}^1(X, \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}}) \rightarrow H^0(X, \Omega_X^{\mathrm{log}}(-1)).$$

Here the map HT is induced by the spectral sequence

$$E_2^{ij} := H_{\acute{e}t}^j(X, R^i \nu_* \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}}) \Rightarrow H_{\mathrm{prok\acute{e}t}}^{i+j}(X, \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}}).$$

We prove this proposition in the fashion of [Heu25, Proposition 2.15]. Via the homeomorphism  $|\mathbb{X}| = |X|$ , we may regard  $\mathcal{O}_{\mathbb{X}}$  as the sheaf on  $X_{\acute{e}t}$ . Define

$$L_{\mathbb{X}} := \left\{ \begin{array}{l} \text{homomorphisms } \tilde{\varphi} \text{ of} \\ \text{sheaves of } B_{\mathrm{dR}}^+/t^2\text{-algebras} \\ \text{on } X_{\mathrm{prok\acute{e}t}} \text{ such that the} \\ \text{right square is commuta-} \\ \text{tive:} \end{array} \left. \begin{array}{ccc} \nu^{-1} \mathcal{O}_{\mathbb{X}} & \xrightarrow{\tilde{\varphi}} & \mathbb{B}_{\mathrm{dR}}^+/t^2 \\ \downarrow & & \downarrow \\ \nu^{-1} \mathcal{O}_X & \longrightarrow & \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}} \end{array} \right\}.$$

Note that  $L_{\mathbb{X}}$  is a sheaf on  $X_{\mathrm{prok\acute{e}t}}$  as a subsheaf of  $\mathcal{H}om(\lambda^{-1} \mathcal{O}_{\mathbb{X}}, \mathbb{B}_{\mathrm{dR}}^+/t^2)$ . We have

**Lemma 5.23.**  *$L_{\mathbb{X}}$  is a pro-Kummer étale torsor under  $\nu^* \Omega_X^{\mathrm{log}}(-1)^\vee$ .*

*Proof.* Similar to the proof of [Heu25, Lemma 2.13] by using the theory of infinitesimal liftings for log smooth morphism, see [Kat89, Proposition 3.14].  $\square$

*Proof of Propostion 5.22.* For  $\theta \in H^0(X, \Omega_X^{\mathrm{log}}(-1))$ , define a map

$$s_{\mathbb{X}} : H^0(X, \Omega_X^{\mathrm{log}}(-1)) \rightarrow H_{\mathrm{prok\acute{e}t}}^1(X, \widehat{\mathcal{O}}_{X_{\mathrm{prok\acute{e}t}}})$$

by

$$s_{\mathbb{X}} : \theta \mapsto L_{\mathbb{X}, \theta} := L_{\mathbb{X}} \times \nu^* \Omega_X^{\log}(-1)^\vee \nu^* \mathcal{O}_X$$

via identifying  $\Omega_X^{\log}(-1)^\vee \rightarrow \mathcal{O}_X$ . We need to check  $\text{HT}(s_{\mathbb{X}}(\theta)) = \theta$ .

The problem is local, so we may assume  $X = \text{Spa}(R, R^+)$  is an affinoid log rigid space over  $C$ , with a strictly étale morphism  $X \rightarrow \mathbb{E} = \text{Spa}(C\langle P \rangle, \mathcal{O}_C\langle P \rangle)$ , which can be written as a composite of rational embeddings and finite étale maps. Let  $T^a$  be the induced coordinate on  $X$ , for any  $a \in P$ . Let  $\tilde{X} = \text{Spa}(R_\infty, R_\infty^+) \rightarrow X$  be the Galois pro-finite Kummer étale cover with Galois group  $\Gamma$  as before. Choose a  $\mathbb{Z}$ -basis  $\{a_1, \dots, a_n\}$  of  $P^{\text{gp}}$ , where  $d = \dim X$ . Then  $\delta(a_1), \dots, \delta(a_d)$  form a basis of  $\Omega_R^{\log}$ . Denote by  $\partial_1, \dots, \partial_d$  its dual basis in  $\Omega_R^{\log, \vee}$ .

For any lift  $\mathbb{X} = \text{Spa}(R', R'^+)$ , by formal log smoothness there exists a lift of the map  $C\langle P \rangle \rightarrow R$  induced by  $X \rightarrow \mathbb{E}$  to an étale morphism

$$B_{\text{dR}}^+/t^2\langle P \rangle \rightarrow R'$$

of  $B_{\text{dR}}^+/t^2$ -algebras. Any choice of such a lift induces a section of  $L_{\mathbb{X}}(\tilde{X})$  as follows: the morphism

$$B_{\text{dR}}^+/t^2\langle P \rangle \rightarrow B_{\text{dR}}^+/t^2\langle P_{\mathbb{Q}>0} \rangle : T^a \rightarrow [T^{a/p^\infty}] \text{ for any } a \in P$$

extends by formal log étaleness to a unique map  $R' \rightarrow B_{\text{dR}}^+/t^2\langle P_{\mathbb{Q}>0} \rangle$  lifting the map  $R \rightarrow R_\infty$ . The following lemma describe the  $\Gamma$ -action on  $L_{\mathbb{X}}(\tilde{X})$ .

**Lemma 5.24.** *Write  $a_i = a_i^+ - a_i^-$  with  $a_i^+, a_i^- \in P$  for  $i = 1, 2, \dots, d$ . Let  $c_i^+, c_i^- : \Gamma \rightarrow \mathbb{Z}_p(1) = \varprojlim_n \mu_{p^n}$  be the maps such that for any  $\gamma \in \Gamma$ , we have*

$$\gamma[T^{a_i^+}/p^\infty] = [c_i^+(\gamma)][T^{a_i^+}/p^\infty], \gamma[T^{a_i^-}/p^\infty] = [c_i^-(\gamma)][T^{a_i^-}/p^\infty].$$

*Then under the identification  $L_{\mathbb{X}}(\tilde{X}) = \Omega_R^{\log}(-1)^\vee \otimes_R R_\infty$  induced by  $R' \rightarrow B_{\text{dR}}^+/t^2\langle P_{\mathbb{Q}>0} \rangle$ , the  $\Gamma$ -action on the right is given by the continuous 1-cocycle*

$$\Gamma \rightarrow \Omega_R^{\log, \vee}(1), \gamma \mapsto \sum_{i=1}^d \frac{c_i^+(\gamma)}{c_i^-(\gamma)} \partial_i.$$

*Proof.* The proof is similar to [Heu25, Lemma 2.16], let  $R_\infty\{1\} := \ker(B_{\text{dR}}/t^2(R_\infty) \rightarrow R_\infty)$ . For any  $\sum x_i \partial_i \in \text{Hom}(\Omega_R^{\log}, R_\infty\{1\})$ , denote by  $x_i^+ := x_i \partial_i(\delta(a_i^+))$ ,  $x_i^- := x_i \partial_i(\delta(a_i^-))$  for all  $i$ , then the corresponding element of be the lifting  $L_{\mathbb{X}}(R_\infty)$  is given by sending  $T^{a_i^+} \mapsto [T^{a_i^+}/p^\infty] + x_i^+$  and  $T^{a_i^-} \mapsto [T^{a_i^-}/p^\infty] + x_i^-$ . Then the cocycle induced by  $\gamma$  can be described as:

$$\begin{aligned} \gamma([T^{a_i^+}/p^\infty] + x_i^+) - [T^{a_i^+}/p^\infty] &= ([c_i^+(\gamma)] - 1)[T^{a_i^+}/p^\infty] + \gamma x_i^+, \\ \gamma([T^{a_i^-}/p^\infty] + x_i^-) - [T^{a_i^-}/p^\infty] &= ([c_i^-(\gamma)] - 1)[T^{a_i^-}/p^\infty] + \gamma x_i^-. \end{aligned}$$

This concludes the proof, since we can identify  $[\epsilon] - 1$  by  $t$  in  $B_{\text{dR}}/t^2$ , for  $\epsilon = (1, \dots) \in \mathbb{Z}_p(1)$ .  $\square$

For any  $i = 1, \dots, d$ , consider the  $\mathcal{O}(1)$  torsor  $L_{\mathbb{X}} \times \nu^* \Omega_X^{\log, \vee}(1) \nu^* \mathcal{O}_X(1)$  induced by  $\delta(a_i) : \Omega_X^{\log, \vee} \rightarrow \mathcal{O}_X$ . We need to check its image under  $\text{HT}(1)$  is  $\delta(a_i)$ . This follows from the commutative diagram in Lemma 5.16:  $\delta(a_i)$  maps to  $\text{HT}(1)^{-1} \delta(a_i)$  through the map over the lower left corner, and it maps to the class defined by the 1-cocycle  $\Gamma \rightarrow R(1) : \gamma \rightarrow c_i^+(\gamma)/c_i^-(\gamma)$  through the map over the



upper right corner, which is exactly the cocycle associated to  $L_{\mathbb{X}} \times \nu^* \Omega_X^{\log, \vee}(1) \nu^* \mathcal{O}_X(1)$  by the lemma above. This concludes the proof.  $\square$

5.4.3. *Splitting of  $R\nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$ .* The splitting of the Hodge–Tate map induces a natural decomposition of  $R\lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$  as follows.

**Proposition 5.25.** *Let  $X$  be a log smooth rigid analytic varieties over  $C$ . Then a choice of log smooth  $B_{\text{dR}}^+/t^2$ -lift  $\mathbb{X}$  of  $X$  gives a natural splitting of  $R\nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}$  as  $\bigoplus_{i \geq 0} \Omega_X^{i, \log}(-i)[-i]$  in the derived category.*

*Proof.* The splitting of the (log) Hodge–Tate map gives a natural isomorphism

$$\mathcal{O}_X \oplus \Omega_X^{1, \log}(-1)[-1] \xrightarrow{\sim} \tau^{\leq 1} R\nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}.$$

The quotient map

$$\left(\Omega_X^{1, \log}\right)^{\otimes i} \rightarrow \bigwedge^i \Omega_X^{1, \log} = \Omega_X^{i, \log}$$

has a canonical  $\mathcal{O}_X$ -linear section  $s_i$  given by

$$\omega_1 \wedge \dots \wedge \omega_i \mapsto \frac{1}{i!} \sum_{\sigma \in S_i} \text{sgn}(\sigma) \omega_{\sigma(1)} \otimes \dots \otimes \omega_{\sigma(i)}.$$

Therefore, we can construct a map

$$\Omega_X^{i, \log}(-i)[-i] \rightarrow (\Omega_X^{1, \log}(-1)[-1])^{\otimes \circ_X i} \rightarrow (R\lambda_* \widehat{\mathcal{O}}_{X_{\text{prokét}}})^{\otimes \circ_X i} \rightarrow R\nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}},$$

which induces a natural map

$$\bigoplus_{i \geq 0} \Omega_X^{i, \log}(-i)[-i] \rightarrow R\nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}.$$

The map constructed above is a quasi-isomorphism by (the construction of) Proposition 5.15.  $\square$

5.4.4. *Proof of Theorem 5.20.* With all these preparations, the proof for Theorem 5.20 is then followed by counting dimensions.

We start with (2). The spectral sequence

$$E_2^{ij} := H_{\text{ét}}^j(X, R^i \nu_* \widehat{\mathcal{O}}_{X_{\text{prokét}}}) \Rightarrow H^{i+j}(X_{\text{prokét}}, \widehat{\mathcal{O}}_{X_{\text{prokét}}}) \simeq H_{\text{két}}^{i+j}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C$$

converges, where the last isomorphism follows from the primitive comparison theorem in [DLLZ23b, Theorem 6.2.1]. The claim then follows from Proposition 5.25 by counting dimensions.

To show (1), we need to show

$$\sum_{i=1}^d \dim_C H^j(X, \Omega_X^{i, \log}) = \dim_C H_{\text{logdR}}^{i+j}(X).$$

By (2), we have

$$\sum_{i=1}^d \dim_C H^j(X, \Omega_X^{i, \log}) = \dim_{\mathbb{Q}_p} H_{\text{két}}^{i+j}(X, \mathbb{Q}_p).$$

On the other hands, Theorem 5.19 shows

$$\begin{aligned} \dim_{\mathbb{Q}_p} H_{\text{két}}^{i+j}(X, \mathbb{Q}_p) &= \dim_{B_{\text{dR}}} H_{B_{\text{dR}}}^{i+j}(X)(*) \begin{bmatrix} 1 \\ t \end{bmatrix} \\ &= \dim_{B_{\text{dR}}} H_{B_{\text{dR}}}^{i+j}(X)(*)/torsion \\ &\leq \dim_{B_{\text{dR}}} H_{B_{\text{dR}}}^{i+j}(X)(*)/t \\ &\leq \dim_C H_{\text{logdR}}^{i+j}(X), \end{aligned}$$

where the second equality holds since  $H_{B_{\text{dR}}}^i(X)(*)/torsion$  is a finite rank torsion-free module over  $B_{\text{dR}}^+$  without nontrivial divisible submodule, so it must be free by [Rot60, Theorem 3]. The last inequality follows from Theorem 5.13, which induces a short exact sequence

$$0 \rightarrow H_{B_{\text{dR}}}^{i+j}(X)(*)/t \rightarrow H_{\text{logdR}}^{i+j}(X) \rightarrow H_{B_{\text{dR}}}^{i+j+1}(X)(*)[t] \rightarrow 0.$$

This proves (1) as all inequalities must be equality.

Finally, the above computation also shows  $H_{\text{logdR}}^i(X/B_{\text{dR}}^+)$  is a finite free  $B_{\text{dR}}^+$ -module as it is torsion-free. This finish the proof.

## REFERENCES

- [Bei87] A. A. Beilinson. On the derived category of perverse sheaves. In *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 27–41. Springer, Berlin, 1987.
- [BMS18] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Integral  $p$ -adic Hodge theory. *Publ. Math. Inst. Hautes Études Sci.*, 128:219–397, 2018.
- [BMS19] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Topological Hochschild homology and integral  $p$ -adic Hodge theory. *Publ. Math. Inst. Hautes Études Sci.*, 129:199–310, 2019.
- [BO78] Pierre Berthelot and Arthur Ogus. *Notes on crystalline cohomology*. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1978.
- [Bos23a] Guido Bosco. On the  $p$ -adic pro-étale cohomology of drinfeld symmetric spaces, 2023, 2110.10683.
- [Bos23b] Guido Bosco. Rational  $p$ -adic hodge theory for rigid-analytic varieties, 2023, 2306.06100.
- [Bra18] Arthur-César Le Bras. Overconvergent relative de rham cohomology over the fargues-fontaine curve, 2018, 1801.00429.
- [BS15] Bhargav Bhatt and Peter Scholze. The pro-étale topology for schemes. *Astérisque*, (369):99–201, 2015.
- [CDN20] Pierre Colmez, Gabriel Dospinescu, and Wiesława Nizioł. Cohomology of  $p$ -adic Stein spaces. *Invent. Math.*, 219(3):873–985, 2020.
- [CG] Brian Conrad and Ofer Gabber. Spreading out of rigid-analytic varieties.
- [CN17] Pierre Colmez and Wiesława Nizioł. Syntomic complexes and  $p$ -adic nearby cycles. *Invent. Math.*, 208(1):1–108, 2017.
- [CN20] Pierre Colmez and Wiesława Nizioł. On  $p$ -adic comparison theorems for rigid analytic varieties, I. *Münster J. Math.*, 13(2):445–507, 2020.
- [CN25] Pierre Colmez and Wiesława Nizioł. On the cohomology of  $p$ -adic analytic spaces, I: The basic comparison theorem. *J. Algebraic Geom.*, 34(1):1–108, 2025.
- [CS19] Dustin Clausen and Peter Scholze. Lectures on condensed mathematics. 2019.
- [Del70] Pierre Deligne. *Équations différentielles à points singuliers réguliers*, volume Vol. 163 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1970.
- [DLLZ23a] Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu. Logarithmic Riemann-Hilbert correspondences for rigid varieties. *J. Amer. Math. Soc.*, 36(2):483–562, 2023.

- [DLLZ23b] Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu. Logarithmic adic spaces: some foundational results. In *p-adic Hodge theory, singular varieties, and non-abelian aspects*, Simons Symp., pages 65–182. Springer, Cham, [2023] ©2023.
- [Fal88] Gerd Faltings. *p*-adic Hodge theory. *J. Amer. Math. Soc.*, 1(1):255–299, 1988.
- [GeK04] Elmar Große-Klönne. De Rham cohomology of rigid spaces. *Math. Z.*, 247(2):223–240, 2004.
- [Gro66a] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III. *Inst. Hautes Études Sci. Publ. Math.*, (28):255, 1966.
- [Gro66b] A. Grothendieck. On the de Rham cohomology of algebraic varieties. *Inst. Hautes Études Sci. Publ. Math.*, (29):95–103, 1966.
- [Guo19] Haoyang Guo. Hodge-tate decomposition for non-smooth spaces, 2019.
- [Guo21] Haoyang Guo. Crystalline cohomology of rigid analytic spaces, 2021, 2112.14304.
- [Heu25] Ben Heuer. A *p*-adic simpson correspondence for smooth proper rigid varieties, 2025, 2307.01303.
- [Hub96] Roland Huber. *Étale cohomology of rigid analytic varieties and adic spaces*. Aspects of Mathematics, E30. Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [Kat70] Nicholas M. Katz. Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin. *Inst. Hautes Études Sci. Publ. Math.*, (39):175–232, 1970.
- [Kat89] Kazuya Kato. Logarithmic structures of Fontaine-Illusie. In *Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988)*, pages 191–224. Johns Hopkins Univ. Press, Baltimore, MD, 1989.
- [Kat96] Fumiharu Kato. Log smooth deformation theory. *Tohoku Math. J. (2)*, 48(3):317–354, 1996.
- [Kie67] Reinhardt Kiehl. Die de Rham Kohomologie algebraischer Mannigfaltigkeiten über einem bewerteten Körper. *Inst. Hautes Études Sci. Publ. Math.*, (33):5–20, 1967.
- [KN99] Kazuya Kato and Chikara Nakayama. Log Betti cohomology, log étale cohomology, and log de Rham cohomology of log schemes over  $\mathbf{C}$ . *Kodai Math. J.*, 22(2):161–186, 1999.
- [LP19] Shizhang Li and Xuanyu Pan. Logarithmic de Rham comparison for open rigid spaces. *Forum Math. Sigma*, 7:Paper No. e32, 53, 2019.
- [LZ17] Ruochuan Liu and Xinwen Zhu. Rigidity and a Riemann-Hilbert correspondence for *p*-adic local systems. *Invent. Math.*, 207(1):291–343, 2017.
- [PGS10] C. Perez-Garcia and W. H. Schikhof. *Locally convex spaces over non-Archimedean valued fields*, volume 119 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2010.
- [Rot60] Joseph J. Rotman. A note on completions of modules. *Proc. Amer. Math. Soc.*, 11:356–360, 1960.
- [Sch13a] Peter Scholze. *p*-adic Hodge theory for rigid-analytic varieties. *Forum Math. Pi*, 1:e1, 77, 2013.
- [Sch13b] Peter Scholze. Perfectoid spaces: A survey, 2013, 1303.5948.
- [Sha25] Xinyu Shao. Hyodo-kato cohomology in rigid geometry: some foundational results. 2025.
- [Tem17] Michael Temkin. Altered local uniformization of Berkovich spaces. *Israel J. Math.*, 221(2):585–603, 2017.
- [Wu24] Zhiyou Wu. A note on Hodge-Tate spectral sequences. *Math. Proc. Cambridge Philos. Soc.*, 176(3):625–642, 2024.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

*Email address:* xshao@imj-prg.fr