

$\otimes$ -FROBENIUS FUNCTORS AND EXACT MODULE CATEGORIES

DAVID JAKLITSCH AND HARSHIT YADAV

ABSTRACT. We call a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between finite tensor categories  $\otimes$ -Frobenius if the left and right adjoints of  $F$  are isomorphic as  $\mathcal{C}$ -bimodule functors. We provide various characterizations of  $\otimes$ -Frobenius functors such as the unimodularity of the centralizer  $\mathcal{Z}({}_F\mathcal{D}_F)$ . We analyze the procedure of pulling back actions of tensor categories on module categories along tensor functors, and show that properties like pivotality and unimodularity are preserved if and only if  $F$  is  $\otimes$ -Frobenius. These results allow the transport of Frobenius algebras under certain conditions. We apply the results to internal natural transformations and examples coming from Hopf algebras.

## CONTENTS

1. Introduction	2
2. Background	5
2.1. Monoidal categories and functors	5
2.2. Finite multitensor categories and module categories	6
2.3. Nakayama functors and relative Serre functors	8
3. $\otimes$ -Frobenius functors	10
3.1. Perfect tensor functors and relative modular objects	10
3.2. Center of a perfect tensor functor	12
3.3. The properties of $\otimes$ -Frobenius functors	15
4. Twisted module categories	18
4.1. Exactness of module categories	19
4.2. Action of central objects and relative Serre functors	21
4.3. Unimodular structures	23
4.4. Pivotal structures and sphericity	24
5. Applications to Hopf algebras	28
5.1. $\otimes$ -Frobenius functors from maps of Hopf algebras	28
5.2. Unimodular and pivotal comodule algebras	31
6. Applications to internal natural transformations	35
6.1. Exactness and unimodularity of the $\mathcal{Z}(\mathcal{C})$ -module category $\text{Func}(\mathcal{M}, \mathcal{N})$	35
6.2. Pivotal setting	38
6.3. Examples from Hopf algebras	39
References	44

## 1. INTRODUCTION

Tensor categories have emerged as fundamental structures in the mathematical study of quantum field theory. They also offer a unified framework for a wide array of mathematical domains, including subfactor theory of von Neumann algebras, vertex operator algebras and Hopf algebras. In this article, we concentrate on the class of finite tensor categories and our results are particularly relevant beyond the semisimple setting.

A typical approach to understanding an algebraic structure is via its representations. Module categories play this central role in the theory of tensor categories. About two decades ago, the work of Etingof and Ostrik [EO04] showed that studying all module categories over a finite tensor category is generally intractable, suggesting a focus on the specific class of *exact module categories*.

Physically, if  $\mathcal{C}$  is a semisimple tensor category associated with a 2-dimensional rational conformal field theory (CFT), semisimple (exact)  $\mathcal{C}$ -module categories supply the data necessary for obtaining the full local conformal field theory [FRS02]. Extending beyond rational CFTs to the finite non-semisimple setting, Fuchs and Schweigert [FS21a] proposed the class of *pivotal exact  $\mathcal{C}$ -module categories* (explored in [Sch13, Shi23a]) as the required data for constructing the full CFT. Since then, additional classes of module categories, such as *unimodular* [Yad23] and *spherical* [FGJS22], have been introduced, which hold significance for the construction of topological quantum field theories (TQFTs) [FGJS24].

The primary objective of this article is to construct new examples of unimodular, pivotal, and spherical  $\mathcal{C}$ -module categories. To achieve this, we investigate the process of twisting module actions by tensor functors. Specifically, given a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , any  $\mathcal{D}$ -module category  $\mathcal{M}$  can be regarded as a  $\mathcal{C}$ -module category by pulling back the  $\mathcal{D}$ -action along  $F$ . We establish necessary and sufficient conditions to ensure that  $F$  preserves the exactness, unimodularity, pivotality and sphericity of module categories under twisting. This is formalized through the introduction of a new concept,  $\otimes$ -Frobenius functors, presented in this paper.

$\otimes$ -Frobenius functors are tensor functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  that admit isomorphic adjoints internal to the 2-category  $\mathbf{Bimod}(\mathcal{C})$  of  $\mathcal{C}$ -bimodules. This is an instance of ambidextrous adjunction which has been studied in prior works such as [Lau06, Str04] and more recently in [FLP24a]. We study  $\otimes$ -Frobenius functors and provide various characterizations along with examples coming from Hopf algebras. We also apply our results to show that, under certain conditions, the objects called *internal natural transformations* (introduced in [FS21b]) have the structure of a (symmetric) Frobenius algebra inside the Drinfeld center of a tensor category. These are relevant for the construction of modular tensor categories and for the description of boundary conditions in CFT [FRS02].

### Main Results

We refer the reader to §2 for the background definitions. A functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between tensor categories is called *perfect* iff it preserves projective objects and *Frobenius* in case its left and right adjoints are isomorphic. These are properties independent of a monoidal structure on  $F$ . A functor is called *Frobenius monoidal* if it admits a lax monoidal and an oplax monoidal structure compatible in a way akin to the product and coproduct of a Frobenius algebra. In the presence of a tensor (that is, a  $\mathbb{k}$ -linear, faithful and strong monoidal) structure on  $F$ , the Frobenius property on  $F$  is related to  $F^{ra}$  admitting a Frobenius monoidal structure. In Proposition 3.7 we show that if  $F^{ra}$  is

Frobenius monoidal, then  $F$  is Frobenius. Conversely if  $F$  is Frobenius and  $F^{ra}$  is exact and faithful, then  $F^{ra}$  is Frobenius monoidal.

A tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  induces a  $\mathcal{C}$ -bimodule category structure on  $\mathcal{D}$  (which we denote as  ${}_F\mathcal{D}_F$ ). Then  $F$  can be seen as a  $\mathcal{C}$ -bimodule functor. Applying the center construction (see §3.2) to this bimodule functor we obtain a tensor functor  $\mathcal{Z}(F): \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{Z}({}_F\mathcal{D}_F)$ . Moreover, the right adjoint  $F^{ra}$  of  $F$  is also a  $\mathcal{C}$ -bimodule functor and its center  $\mathcal{Z}(F^{ra})$  is a lax monoidal functor that is the right adjoint of  $\mathcal{Z}(F)$ .

We call  $F$   $\otimes$ -Frobenius if its adjoints are isomorphic as  $\mathcal{C}$ -bimodule functors. As the main result of §3, we obtain various characterizations of  $\otimes$ -Frobenius functors, relating them to Frobenius functors, Frobenius monoidal functors and unimodular tensor categories:

**Theorem A.** (3.18) *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. Then the following are equivalent:*

- (i)  $F$  is  $\otimes$ -Frobenius.
- (ii) The tensor functor  $\mathcal{Z}(F): \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{Z}({}_F\mathcal{D}_F)$  is Frobenius.
- (iii) The tensor category  $\mathcal{Z}({}_F\mathcal{D}_F)$  is unimodular.
- (iv) The lax monoidal functor  $\mathcal{Z}(F^{ra}): \mathcal{Z}({}_F\mathcal{D}_F) \rightarrow \mathcal{Z}(\mathcal{C})$  is (braided) Frobenius monoidal.

Here the equivalence of parts (i), (ii) and (iii) is proved by utilizing results from [Shi17], and the equivalence with (iv) follows from [FLP24a].

While tensor functors that are Frobenius are not in general  $\otimes$ -Frobenius, using this result, we show that central tensor functors are Frobenius if and only if they are  $\otimes$ -Frobenius (Proposition 3.20). Furthermore, we prove that any tensor functor between separable fusion categories is  $\otimes$ -Frobenius (Theorem 3.23), thereby answering [FLP24b, Questions 5.22 & 5.23].

In §4, we apply the results of §3 to twisted actions on module categories. A tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  induces on a left  $\mathcal{D}$ -module category  $\mathcal{M}$  a left  $\mathcal{C}$ -module action via:

$$c \triangleright^F m := F(c) \triangleright m$$

for every  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$ . We denote the resulting  $\mathcal{C}$ -module category as  ${}_F\mathcal{M}$ .

When  $\mathcal{M}$  is exact as a  $\mathcal{D}$ -module category, we show that  ${}_F\mathcal{M}$  is exact as a  $\mathcal{C}$ -module category if and only if  $F$  is perfect. For perfect functors, the *relative modular object*  $\chi_F \in \mathcal{D}$ , introduced in [Shi17], ‘measures’ how much  $F^{la}$  and  $F^{ra}$  differ from one another. Moreover,  $\chi_F$  is equipped with a  $F$ -half-braiding  $\sigma: \chi_F \otimes F(c) \rightarrow F(c) \otimes \chi_F$  such that  $(\chi_F, \sigma) \in \mathcal{Z}({}_F\mathcal{D}_F)$ . In particular, using  $\sigma$ ,  $\chi_F$  induces a  $\mathcal{C}$ -module functor

$$\chi_F \triangleright -: {}_F\mathcal{M} \rightarrow {}_F\mathcal{M}.$$

We show that the pair  $(\chi_F, \sigma)$  is the relative modular object of  $\mathcal{Z}(F)$  (Proposition 3.13). From this, it follows that  $\chi_F \triangleright -$  is trivial as a  $\mathcal{C}$ -module functor if and only if  $F$  is  $\otimes$ -Frobenius.

To understand properties like pivotality, unimodularity, etc. of a  $\mathcal{C}$ -module category the main hurdle is to determine its so called relative Serre functor. In Proposition 4.14, we provide a formula for the relative Serre functor of the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  and show that it is described using the functor  $\chi_F \triangleright -$  and the relative Serre functor  $\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}$  of the  $\mathcal{D}$ -module category  $\mathcal{M}$  as follows:

$$\mathbb{S}_{{}_F\mathcal{M}}^{\mathcal{C}} \cong \chi_F \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}.$$

See §4.2 for details. This allows us to obtain the following result:

**Theorem B.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a tensor functor and  $\mathcal{M}$  a  $\mathcal{D}$ -module category. The table below summarizes the assumptions needed to transfer a property on  $\mathcal{M}$  to the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$ . The ‘result’ column shows that said property is transferred iff  $F$  obeys the respective assumptions.*

Reference	Assumptions			Result	
	$\mathcal{C}, \mathcal{D}$	$\mathcal{M}$	$F$	$F$	${}_F\mathcal{M}$
Prop. 4.5	FTC	exact	tensor	perfect $\iff$	exact
Prop. 4.20	FTC	unimodular	perfect	$\otimes$ -Frobenius $\iff$	unimodular
Prop. 4.24	pivotal	pivotal	perfect, pivotal	$\otimes$ -Frobenius $\iff$	pivotal
Prop 4.33	spherical	spherical	perfect, pivotal	$\otimes$ -Frobenius $\iff$	spherical

A consequence of these results is that: if  $F$  is a perfect tensor functor, then its right adjoint  $F^{\text{ra}}$  preserves exact algebras. Furthermore, if  $\mathcal{C}$  and  $\mathcal{D}$  are pivotal and  $F$  is a pivotal  $\otimes$ -Frobenius functor, then  $F^{\text{ra}}$  preserves exact symmetric Frobenius algebras.

In §5, we illustrate our results for the categories  $\text{Rep}(H)$  of finite-dimensional representations of a finite-dimensional Hopf algebra  $H$ . We show that the tensor functor  $F_{\mathbf{f}}: \text{Rep}(H) \rightarrow \text{Rep}(H')$ , induced by a bialgebra map  $\mathbf{f}: H' \rightarrow H$ , is  $\otimes$ -Frobenius if and only if  $\mathbf{f}(g_{H'}) = g_H$  and  $\alpha_H \circ \mathbf{f} = \alpha_{H'}$ , where  $g$  and  $\alpha$  stand for the distinguished grouplike elements of a Hopf algebra and its dual (Proposition 5.3). We provide examples of such bialgebra maps and explain how our results generalize some recent findings in [FLP24b] obtained using different techniques.

Pivotal (and unimodular)  $\text{Rep}(H)$ -module categories are understood in terms of  $H$ -comodule algebras equipped with certain elements called pivotal (and unimodular) elements. We also provide consequences of our results in §4 and interpret them in terms of these elements.

In §6, we provide another application of Theorem B to generalize some algebraic results obtained in [FS21b] inspired by logarithmic CFTs. For a  $\mathcal{C}$ -module category  $\mathcal{M}$ , let  $\mathcal{C}_{\mathcal{M}}^*$  denote the category of  $\mathcal{C}$ -module endofunctors of  $\mathcal{M}$ . There is a perfect tensor functor  $\Psi: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}_{\mathcal{M}}^*$  which allows us to consider  $\mathcal{C}_{\mathcal{M}}^*$ -module categories as  $\mathcal{Z}(\mathcal{C})$ -module categories. We prove that  $\Psi$  is  $\otimes$ -Frobenius if and only if  $\mathcal{M}$  is unimodular (Proposition 6.9). Given a finite tensor category  $\mathcal{C}$  and  $\mathcal{M}$  and  $\mathcal{N}$  exact left  $\mathcal{C}$ -module categories, we show that the  $\mathcal{C}_{\mathcal{M}}^*$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is exact, thereby generalizing [FS21b, Thm. 18]. A similar statement holds for unimodular module categories. If, in addition,  $\mathcal{C}$  is pivotal,  $\mathcal{M}$  and  $\mathcal{N}$  are pivotal  $\mathcal{C}$ -modules and  $\mathcal{M}$  is unimodular, then  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is a pivotal  $\mathcal{Z}(\mathcal{C})$ -module category. Furthermore,  $\Psi^{\text{ra}}$  is a pivotal Frobenius monoidal functor.

For  $G_1, G_2 \in \text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ , we show that the object  $\underline{\text{Nat}}(G_1, G_2) \in \mathcal{Z}(\mathcal{C})$  of internal natural transformations [FS21b] is given by  $\Psi^{\text{ra}}(G_1^{\text{ra}} \circ G_2)$ . Thus, Theorem B above gives criteria for when  $\underline{\text{Nat}}(G, G)$  is a symmetric Frobenius algebra in  $\mathcal{Z}(\mathcal{C})$ , generalizing the results of [FS21b]. When  $\mathcal{C} = \text{Rep}(H)$  for a finite dimensional Hopf algebra  $H$  and  $\mathcal{M} = \text{Rep}(L)$  for an exact left  $H$ -comodule algebra, we explicitly describe the algebra structure of  $\underline{\text{Nat}}(P, P)$  and give conditions on  $H$  and  $L$  that ensure  $\underline{\text{Nat}}(P, P)$  is a (symmetric) Frobenius algebra, for  $P \in \text{Func}_{\text{Rep}(H)}(\text{Rep}(L), \text{Rep}(L))$ .

## Acknowledgments

D.J. is supported by The Research Council of Norway - project 324944. H.Y. is supported by a start-up grant from the University of Alberta and an NSERC Discovery Grant. We would like to thank J. Fuchs and C. Schweigert for discussions that encouraged us to start this project. We also thank K. Shimizu for a helpful email correspondence and pointing out Example 5.13.

## 2. BACKGROUND

In this article, we will work over an algebraically closed field  $\mathbb{k}$ .  $\text{Vec}$  will denote the category of finite-dimensional  $\mathbb{k}$ -vector spaces. Given a category  $\mathcal{X}$ , we will denote its opposite category as  $\mathcal{X}^{\text{op}}$ . For a morphism  $f: x \rightarrow x'$  in  $\mathcal{X}$ , the corresponding morphism in  $\mathcal{C}^{\text{op}}$  is denoted as  $f^{\text{op}}: x' \rightarrow x$ . Unless stated otherwise, we will follow the same conventions as in [EGNO16].

### 2.1. Monoidal categories and functors

Owing to MacLane's coherence theorem, we assume that all monoidal categories are strict. A generic monoidal category is denoted as a triple  $(\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}})$  where  $\otimes_{\mathcal{C}}$  stands for the tensor product and  $\mathbb{1}_{\mathcal{C}}$  for the unit object. The *monoidal opposite*  $\overline{\mathcal{C}}$  of  $\mathcal{C}$  is the monoidal category having the same underlying category as  $\mathcal{C}$ , but with reversed tensor product, i.e.  $c \otimes_{\overline{\mathcal{C}}} d := d \otimes_{\mathcal{C}} c$ . Moreover, the opposite category  $\mathcal{C}^{\text{op}}$  is also endowed with a monoidal structure given by the same tensor product and unit as  $\mathcal{C}$ .

A monoidal category  $\mathcal{C}$  is called *rigid* if every object  $c \in \mathcal{C}$  admits left and right duals. We use the following conventions for dualities: the right dual  $c^{\vee}$  of an object  $c \in \mathcal{C}$  comes equipped with evaluation and coevaluation morphisms

$$\text{ev}_c: c^{\vee} \otimes c \longrightarrow \mathbb{1} \quad \text{and} \quad \text{coev}_c: \mathbb{1} \longrightarrow c \otimes c^{\vee},$$

and the left dual  ${}^{\vee}c$  of  $c \in \mathcal{C}$  comes with evaluation and coevaluation morphisms

$$\widetilde{\text{ev}}_c: c \otimes {}^{\vee}c \longrightarrow \mathbb{1} \quad \text{and} \quad \widetilde{\text{coev}}_c: \mathbb{1} \longrightarrow {}^{\vee}c \otimes c.$$

This matches the conventions in [FSS20]. A rigid monoidal category is called *pivotal* if it is equipped with a *pivotal structure*, that is, a monoidal natural isomorphism  $\mathfrak{p}: \text{id}_{\mathcal{C}} \xrightarrow{\sim} (-)^{\vee\vee}$ .

An *algebra* in  $\mathcal{C}$  is a triple  $(A, m: A \otimes A \rightarrow A, u: \mathbb{1} \rightarrow A)$  where  $m$  satisfies  $m \circ (\text{id}_A \otimes m) = m \circ (m \otimes \text{id}_A)$  and  $u$  satisfies  $m \circ (\text{id}_A \otimes u) = \text{id}_A = m \circ (u \otimes \text{id}_A)$ . A *coalgebra* in  $\mathcal{C}$  is a triple  $(A, \Delta: A \rightarrow A \otimes A, \varepsilon: A \rightarrow \mathbb{1})$  such that  $(A, \Delta^{\text{op}}, \varepsilon^{\text{op}})$  is an algebra in  $\mathcal{C}^{\text{op}}$ . A *Frobenius algebra* is a tuple  $(A, m, u, \Delta, \varepsilon)$  where  $(A, m, u)$  forms an algebra,  $(A, \Delta, \varepsilon)$  forms a coalgebra and  $m, \Delta$  satisfy:

$$(m \otimes \text{id}_A) \circ (\text{id}_A \otimes \Delta) = \Delta \circ m = (\text{id}_A \otimes m) \circ (\Delta \otimes \text{id}_A).$$

Let  $\mathcal{C}$  be a pivotal category with pivotal structure  $\mathfrak{p}$ . A *symmetric Frobenius algebra* is a Frobenius algebra  $(A, m, u, \Delta, \varepsilon)$  satisfying:

$$(\varepsilon m \otimes \text{id}_{A^{\vee}})(\text{id}_A \otimes \text{coev}_A) = (\mathfrak{p}_{\vee A} \otimes \varepsilon m)(\widetilde{\text{coev}}_A \otimes \text{id}_A).$$

Let  $\mathcal{C}$  and  $\mathcal{D}$  be monoidal categories. A (*lax*) *monoidal functor* is a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  along with a morphism  $F_0: \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$  and a natural transformation  $F_2(c, c'): F(c) \otimes_{\mathcal{D}} F(c') \rightarrow F(c \otimes_{\mathcal{C}} c')$  for  $c, c' \in \mathcal{C}$  obeying certain associativity and unitality conditions. In the case that  $F_0$  and  $F_2(c, c')$  are isomorphisms we call  $F$  *strong monoidal*.

An *oplax monoidal functor* is a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  equipped with a morphism  $F^0: F(\mathbb{1}_{\mathcal{C}}) \rightarrow \mathbb{1}_{\mathcal{D}}$  and a natural transformation  $F^2(c, c'): F(c \otimes_{\mathcal{C}} c') \rightarrow F(c) \otimes_{\mathcal{D}} F(c')$  such that  $(F^{\text{op}}, (F^0)^{\text{op}}, (F^2)^{\text{op}})$  is lax monoidal.

A tuple  $(F, F_0, F_2, F^0, F^2)$  is called *Frobenius monoidal* if  $(F, F_0, F_2)$  is lax monoidal,  $(F, F^0, F^2)$  is oplax monoidal and for all  $c, c', c'' \in \mathcal{C}$ , the following conditions are fulfilled [DP08]:

$$(\text{id}_{F(c)} \otimes_{\mathcal{D}} F_2(c', c'')) (F^2(c, c') \otimes_{\mathcal{D}} \text{id}_{F(c'')}) = F^2(c, c' \otimes_{\mathcal{C}} c'') F_2(c \otimes_{\mathcal{C}} c', c''),$$

$$(F_2(c, c') \otimes_{\mathcal{D}} \text{id}_{F(c'')}) (\text{id}_{F(c)} \otimes_{\mathcal{D}} F^2(c', c'')) = F^2(c \otimes_{\mathcal{C}} c', c'') F_2(c, c' \otimes_{\mathcal{C}} c'').$$

Any strong monoidal functor  $(F, F_0, F_2)$  is Frobenius monoidal with  $F^0 = F_0^{-1}$  and  $F^2 = F_2^{-1}$ .

A Frobenius monoidal functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between rigid monoidal categories preserves dualities, that is, we have natural isomorphisms  $\zeta_c : F(c^\vee) \xrightarrow{\sim} F(c)^\vee$ . Let  $(\mathcal{C}, \mathfrak{p}^c)$  and  $(\mathcal{D}, \mathfrak{p}^d)$  be two pivotal monoidal categories. We call it *pivotal* if it satisfies:

$$\zeta_c^\vee \circ \mathfrak{p}_{F(c)}^{\mathcal{D}} = \zeta_{c^\vee} \circ F(\mathfrak{p}^c) : F(c) \rightarrow F(c^\vee)^\vee \quad (2.1)$$

where  $\zeta_c : F(c^\vee) \xrightarrow{\sim} F(c)^\vee$  is the canonical isomorphism associated to  $F$ . By applying  $\zeta$  twice, we obtain a natural isomorphism  $\xi_c : F(c^{\vee\vee}) \xrightarrow{\sim} F(c)^{\vee\vee}$ .

Let  $(F, F_0, F_2)$  be a strong monoidal functor from  $\mathcal{C}$  to  $\mathcal{D}$  with a right adjoint  $F^{\text{ra}}$  with unit  $\eta$  and counit  $\varepsilon$ . The monoidal structure of  $F$  induces a lax monoidal structure on  $F^{\text{ra}}$  via

$$F_2^{\text{ra}}(d, d') := F^{\text{ra}}(\varepsilon_d \otimes \varepsilon_{d'}) \circ F^{\text{ra}}(F_2(F^{\text{ra}}(d), F^{\text{ra}}(d'))^{-1}) \circ \eta_{F^{\text{ra}}(d) \otimes F^{\text{ra}}(d')}, \quad F_0^{\text{ra}} = F^{\text{ra}}(F^0) \circ \eta_{\mathbb{1}_{\mathcal{C}}}.$$

By means of  $\eta$ ,  $\varepsilon$  and  $F_2$ , we can form two natural transformations called *left and right coHopf operators*, respectively [BLV11]:

$$h_{d,c}^l : F^{\text{ra}}(d) \otimes_{\mathcal{C}} c \rightarrow F^{\text{ra}}(d \otimes_{\mathcal{D}} F(c)) \quad \text{and} \quad h_{d,c}^r : c \otimes_{\mathcal{C}} F^{\text{ra}}(d) \rightarrow F^{\text{ra}}(F(c) \otimes_{\mathcal{D}} d). \quad (2.2)$$

In a similar manner, when  $F$  admits a left adjoint  $F^{\text{la}}$ , we can equip  $F^{\text{la}}$  with an oplax monoidal structure  $((F^{\text{la}})^2, (F^{\text{la}})^0)$ . We use the unit, counit of the adjunction  $F^{\text{la}} \dashv F$  and  $F_2$  to define natural transformations called *left and right Hopf operators*, respectively [BLV11]:

$$H_{d,c}^l : F^{\text{la}}(d \otimes_{\mathcal{D}} F(c)) \rightarrow F^{\text{la}}(d) \otimes_{\mathcal{C}} c \quad \text{and} \quad H_{d,c}^r : F^{\text{la}}(F(c) \otimes_{\mathcal{D}} d) \rightarrow c \otimes_{\mathcal{C}} F^{\text{la}}(d). \quad (2.3)$$

If  $\mathcal{C}$  and  $\mathcal{D}$  are rigid, the maps  $h^l, h^r$  and  $H^l, H^r$  are all isomorphisms. We note that in references such as [FLP24a, FLP24b, FLP24c], the co/Hopf operators are also called projection formula morphisms.

## 2.2. Finite multitensor categories and module categories

A *finite  $\mathbb{k}$ -linear category* is an abelian category that is equivalent to the category of finite-dimensional representations of a finite dimensional  $\mathbb{k}$ -algebra. All categories considered in this article will be finite  $\mathbb{k}$ -linear and all functors are assumed to be additive and  $\mathbb{k}$ -linear, unless otherwise stated. Every  $\mathbb{k}$ -linear functor between finite  $\mathbb{k}$ -linear categories that is left (right) exact admits a left (right) adjoint [DSPS19, Cor. 1.9]. The *image* of a  $\mathbb{k}$ -linear functor  $F : \mathcal{X} \rightarrow \mathcal{Y}$ , denoted by  $\text{im}(F)$ , is the full subcategory of  $\mathcal{Y}$  consisting of subquotients of objects in the essential image of  $F$ . A  $\mathbb{k}$ -linear functor  $F : \mathcal{X} \rightarrow \mathcal{Y}$  is called *surjective* iff its image is equivalent to  $\mathcal{Y}$ .

A *finite multitensor category* is a finite abelian  $\mathbb{k}$ -linear, rigid monoidal category with  $\mathbb{k}$ -bilinear tensor product. If, in addition,  $\text{End}_{\mathcal{C}}(\mathbb{1}) \cong \mathbb{k}$  holds, we call  $\mathcal{C}$  a *finite tensor category*.

A *tensor functor* is a  $\mathbb{k}$ -linear, exact, and strong monoidal functor. It is established that tensor functors between finite tensor categories are faithful [Del07, Corollaire 2.10(ii)]. The image  $\text{im}(F)$  of a tensor functor  $F$  is a tensor subcategory of  $\mathcal{D}$ . A fully faithful tensor functor is called *injective*.

A *left module category* over a finite tensor category  $\mathcal{C}$  is a finite  $\mathbb{k}$ -linear category  $\mathcal{M}$  equipped with a module action functor  $\triangleright : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$  that is  $\mathbb{k}$ -bilinear and exact in each variable, and natural isomorphisms  $x \triangleright (y \triangleright m) \cong (x \otimes y) \triangleright m$  (for  $x, y \in \mathcal{C}$ ,  $m \in \mathcal{M}$ ) that satisfy certain coherence conditions. Right module categories and bimodule categories are similarly defined [EGNO16, Def.

7.1.7]. A module category  ${}_c\mathcal{M}$  is called *exact* iff for any  $m \in \mathcal{M}$  and any projective object  $p \in \mathcal{C}$ ,  $p \triangleright m$  is projective in  $\mathcal{M}$ . A module category is called *indecomposable* iff it is not equivalent to a direct sum of two non-trivial module categories. A  $(\mathcal{C}, \mathcal{D})$ -bimodule category is *exact* iff it is exact as left  $\mathcal{C}$ -module category and also exact as a right  $\mathcal{D}$ -module category. We call an algebra  $A \in \mathcal{C}$  *exact* (resp. *indecomposable*) if the left  $\mathcal{C}$ -module category  $\text{mod}_A(\mathcal{C})$  of right  $A$ -modules in  $\mathcal{C}$  is exact (resp. indecomposable).

A  $\mathcal{C}$ -module functor between two  $\mathcal{C}$ -module categories  $\mathcal{M}$  and  $\mathcal{N}$  consists of a functor  $H: \mathcal{M} \rightarrow \mathcal{N}$  endowed with a *module structure*, i.e. a natural isomorphism

$$\varphi_{c,m}: H(c \triangleright m) \xrightarrow{\sim} c \triangleright H(m)$$

for  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$  satisfying the appropriate pentagon axioms. Module functors can be composed: given  $\mathcal{C}$ -module categories  $\mathcal{M}$ ,  $\mathcal{N}$  and  $\mathcal{L}$  and  $\mathcal{C}$ -module functors  $H_1: \mathcal{M} \rightarrow \mathcal{N}$  and  $H_2: \mathcal{N} \rightarrow \mathcal{L}$  the composition functor  $H_2 \circ H_1: \mathcal{M} \rightarrow \mathcal{L}$  inherits a  $\mathcal{C}$ -module functor structure via

$$H_2 \circ H_1(c \triangleright m) \xrightarrow{H_2(\varphi_{c,m}^1)} H_2(c \triangleright H_1(m)) \xrightarrow{\varphi_{c,H_1(m)}^2} c \triangleright H_2(H_1(m)) \quad (2.4)$$

where the isomorphism  $\varphi^i$  is the module structure of  $H_i$  for  $i = 1, 2$ .

A *module natural transformation* between module functors is a natural transformation between the underlying functors commuting with the respective module structures. Given  $\mathcal{C}$ -module categories  $\mathcal{M}$  and  $\mathcal{N}$ ,  $\text{Func}(\mathcal{M}, \mathcal{N})$  denotes the category with  $\mathcal{C}$ -module functors between  $\mathcal{M}$  and  $\mathcal{N}$  as objects and module natural transformations as morphisms.

Let  $\mathcal{C}$  be a finite tensor category and  $\mathcal{M}$  an indecomposable exact left  $\mathcal{C}$ -module category. The *dual tensor category* of  $\mathcal{C}$  with respect to  $\mathcal{M}$  is the finite tensor category  $\mathcal{C}_{\mathcal{M}}^* := \text{Func}(\mathcal{C}, \mathcal{M})$  of  $\mathcal{C}$ -module endofunctors with tensor product given by composition of module functors  $H \otimes H' = H \circ H'$  and monoidal unit  $\text{id}_{\mathcal{M}}$ . Furthermore, given exact  $\mathcal{C}$ -module categories  $\mathcal{M}$  and  $\mathcal{N}$ , the category of module functors  $\text{Func}(\mathcal{M}, \mathcal{N})$  is endowed with the structure of a  $(\mathcal{C}_{\mathcal{N}}^*, \mathcal{C}_{\mathcal{M}}^*)$ -bimodule category, via composition of module functors.

Let  $\mathcal{M}_{\mathcal{D}}$  be a right module category and  ${}_{\mathcal{D}}\mathcal{N}$  a left module category. A *balanced functor* is a bilinear functor  $F: \mathcal{M} \times \mathcal{N} \rightarrow \mathcal{Y}$  into a  $\mathbb{k}$ -linear category  $\mathcal{Y}$  equipped with a natural isomorphism

$$F(m \triangleleft d, n) \xrightarrow{\sim} F(m, d \triangleright n)$$

for  $d \in \mathcal{D}$ ,  $m \in \mathcal{M}$  and  $n \in \mathcal{N}$ , obeying a pentagon coherence condition.

The *relative Deligne product* of  $\mathcal{M}_{\mathcal{D}}$  and  ${}_{\mathcal{D}}\mathcal{N}$  is a  $\mathbb{k}$ -linear category  $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}$  equipped with a right exact  $\mathcal{D}$ -balanced functor  $\boxtimes_{\mathcal{D}}: \mathcal{M} \times \mathcal{N} \rightarrow \mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}$  such that for every  $\mathbb{k}$ -linear category  $\mathcal{Y}$  the functor

$$\text{Rex}(\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}, \mathcal{Y}) \xrightarrow{\sim} \text{Bal}^{\text{re}}(\mathcal{M} \times \mathcal{N}, \mathcal{Y}), \quad F \mapsto F \circ \boxtimes_{\mathcal{D}}$$

is an equivalence of categories, where  $\text{Bal}^{\text{re}}$  denotes the category of right exact balanced functors.

An exact module category  ${}_{\mathcal{D}}\mathcal{M}$  becomes an exact  $(\mathcal{D}, \overline{\mathcal{D}_{\mathcal{M}}^*})$ -bimodule category with action of the dual tensor category  $\mathcal{D}_{\mathcal{M}}^*$  given by module functor evaluation [EGNO16, Lemma 7.12.7]. The category of  $\mathcal{D}$ -module functors  $\overline{\mathcal{M}} := \text{Func}_{\mathcal{D}}(\mathcal{M}, \mathcal{D})$  is naturally endowed with the structure of an exact  $(\overline{\mathcal{D}_{\mathcal{M}}^*}, \mathcal{D})$ -bimodule category via

$$F \triangleright H \triangleleft d := H \circ F(-) \otimes d$$

for  $F \in \mathcal{D}_{\mathcal{M}}^*$ ,  $H \in \overline{\mathcal{M}}$  and  $d \in \mathcal{D}$ . There are bimodule equivalences

$$\mathcal{M} \boxtimes_{\mathcal{D}_{\mathcal{M}}^*} \overline{\mathcal{M}} \xrightarrow{\sim} \mathcal{D} \quad \text{and} \quad \overline{\mathcal{M}} \boxtimes_{\mathcal{D}} \mathcal{M} \xrightarrow{\sim} \mathcal{D}_{\mathcal{M}}^* \quad (2.5)$$

of  $\mathcal{D}$ -bimodule categories and of  $\mathcal{D}_{\mathcal{M}}^*$ -bimodule categories, respectively [DSPS18, Prop. 2.4.10].

Let  $\mathcal{M}$  be a  $\mathcal{C}$ -module category. For an object  $m \in \mathcal{M}$  the action functor  $- \triangleright m : \mathcal{C} \rightarrow \mathcal{M}$  is an exact functor and therefore it admits a right adjoint  $\underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{C}}(m, -)$ , i.e. there are natural isomorphisms

$$\text{Hom}_{\mathcal{M}}(c \triangleright m, n) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(c, \underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{C}}(m, n))$$

for  $c \in \mathcal{C}$  and  $m, n \in \mathcal{M}$ . This extends to a left exact functor

$$\underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{C}}(-, -) : \mathcal{M}^{\text{opp}} \times \mathcal{M} \rightarrow \mathcal{C}$$

called the *internal Hom functor* of  ${}_{\mathcal{C}}\mathcal{M}$ . A module category is exact iff its internal Hom is an exact functor [EGNO16, Cor. 7.9.6 & Prop. 7.9.7].

**Lemma 2.1.** *Let  $\mathcal{M}$  be a  $(\mathcal{C}, \mathcal{D})$ -bimodule and  $\mathcal{N}$  a left  $\mathcal{D}$ -module.*

(i) [Sch15, Prop. 4.15 (3)] *The internal Hom for the left  $\mathcal{C}$ -module  $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}$  is given by:*

$$\underline{\text{Hom}}_{\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}}^{\mathcal{C}}(m \boxtimes_{\mathcal{D}} n, m' \boxtimes_{\mathcal{D}} n') \cong \underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{C}}(m \triangleleft {}^{\vee}\underline{\text{Hom}}_{\mathcal{N}}^{\mathcal{D}}(n, n'), m')$$

(ii) *If  $\mathcal{M}$  and  $\mathcal{N}$  are exact, then  $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}$  is an exact left  $\mathcal{C}$ -module category.*

*Proof.* Part (i) follows from the following sequence of isomorphisms

$$\begin{aligned} \text{Hom}_{\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}}(c \triangleright (m \boxtimes_{\mathcal{D}} n), m' \boxtimes_{\mathcal{D}} n') &\cong \text{Hom}_{\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}}((c \triangleright m) \boxtimes_{\mathcal{D}} n, m' \boxtimes_{\mathcal{D}} n') \\ &\cong \text{Hom}_{\mathcal{D}}(\mathbf{1}_{\mathcal{D}}, \underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{D}}(c \triangleright m, m') \otimes \underline{\text{Hom}}_{\mathcal{N}}^{\mathcal{D}}(n, n')) \\ &\cong \text{Hom}_{\mathcal{D}}({}^{\vee}\underline{\text{Hom}}_{\mathcal{N}}^{\mathcal{D}}(n, n'), \underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{D}}(c \triangleright m, m')) \\ &\cong \text{Hom}_{\mathcal{M}}((c \triangleright m) \triangleleft {}^{\vee}\underline{\text{Hom}}_{\mathcal{N}}^{\mathcal{D}}(n, n'), m'). \\ &\cong \text{Hom}_{\mathcal{C}}(c, \underline{\text{Hom}}_{\mathcal{M}}^{\mathcal{C}}(m \triangleleft {}^{\vee}\underline{\text{Hom}}_{\mathcal{N}}^{\mathcal{D}}(n, n'), m')). \end{aligned}$$

The fact that the actions are exact functors and assuming the internal homs of  $\mathcal{M}$  and  $\mathcal{N}$  to be exact implies that  $\underline{\text{Hom}}_{\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N}}^{\mathcal{C}}(-, -)$  is exact, which leads to the second statement.  $\square$

### 2.3. Nakayama functors and relative Serre functors

Any finite  $\mathbb{k}$ -linear category  $\mathcal{X}$  becomes a left  $\text{Vec}$ -module category via the action  $\blacktriangleright$  defined by:

$$\text{Hom}_{\mathbb{k}}(V, \text{Hom}_{\mathcal{X}}(x, y)) \cong \text{Hom}_{\mathcal{X}}(V \blacktriangleright x, y), \quad (V \in \text{Vec}, x, y \in \mathcal{X}).$$

The left and right exact Nakayama functors of  $\mathcal{X}$  are defined as the image of the identity functor under the Eilenberg-Watts correspondence [FSS20, Def. 3.14], explicitly given by the (co)ends

$$\mathbb{N}_{\mathcal{X}}^l(x) := \int_{y \in \mathcal{X}} \text{Hom}_{\mathcal{X}}(y, x) \blacktriangleright y, \quad \mathbb{N}_{\mathcal{X}}^r(x) := \int^{y \in \mathcal{X}} \text{Hom}_{\mathcal{X}}(x, y)^* \blacktriangleright y.$$

The right exact Nakayama functor is equipped with a family of natural isomorphisms

$$\mathbf{n}_F : \mathbb{N}_{\mathcal{Y}}^r \circ F \xrightarrow{\sim} F^{\text{rra}} \circ \mathbb{N}_{\mathcal{X}}^r \quad (2.6)$$

for every right exact functor  $F : \mathcal{X} \rightarrow \mathcal{Y}$  admitting a double-right adjoint. There are similar natural isomorphisms involving the left exact Nakayama functor.



Given a finite tensor category  $\mathcal{C}$ , the Nakayama functors of the  $\mathbb{k}$ -linear category underlying  $\mathcal{C}$  determine *distinguished invertible objects*

$$D_c := \mathbb{N}_c^l(\mathbf{1}) \quad \text{and} \quad D_c^{-1} := \mathbb{N}_c^r(\mathbf{1}). \quad (2.7)$$

These come with a monoidal natural isomorphism

$$\mathcal{R}_c : D_c \otimes - \xrightarrow{\sim} (-)^{v^v v^v} \otimes D_c \quad (2.8)$$

called the *Radford isomorphism*. A finite tensor category  $\mathcal{C}$  is *unimodular* iff  $D_c$  is isomorphic to the monoidal unit  $\mathbf{1}$ .

Given a  $\mathcal{C}$ -module category  $\mathcal{M}$ , the natural isomorphisms (2.6) endow the Nakayama functor of  $\mathcal{M}$  with the structure of a twisted module functor of the form

$$\mathfrak{n}_{c,m} : \mathbb{N}_{\mathcal{M}}^r(c \triangleright m) \xrightarrow{\sim} {}^{v^v}c \triangleright \mathbb{N}_{\mathcal{M}}^r(m). \quad (2.9)$$

Similarly, the left exact Nakayama functor becomes a twisted module functor with the appropriate appearance of double-duals.

Let  $\mathcal{C}$  be a finite tensor category and  $\mathcal{M}$  a left  $\mathcal{C}$ -module category. A (right) *relative Serre functor* of  $\mathcal{M}$  [FSS20, Def. 4.22] is an endofunctor  $\mathbb{S}_{\mathcal{M}}^c : \mathcal{M} \rightarrow \mathcal{M}$  equipped with a natural isomorphism

$$\phi_{m,n} : \underline{\mathrm{Hom}}_{\mathcal{M}}^c(m, n)^v \xrightarrow{\sim} \underline{\mathrm{Hom}}_{\mathcal{M}}^c(n, \mathbb{S}_{\mathcal{M}}^c(m)), \quad (m, n \in \mathcal{M}).$$

A relative Serre functor  $(\mathbb{S}_{\mathcal{M}}^c, \phi)$  exists iff  $\mathcal{M}$  is exact [FSS20]. In this case, it is unique up to a unique isomorphism [Shi23a, Lem. 3.5] and it is an equivalence of categories. The (left) relative Serre functor [FSS20, Def. 4.22], denoted by  $\overline{\mathbb{S}}_{\mathcal{M}}^c : \mathcal{M} \rightarrow \mathcal{M}$ , serves as quasi-inverse of  $\mathbb{S}_{\mathcal{M}}^c$ . Relative Serre functors come with a twisted module functor structure, i.e. there are natural isomorphisms

$$\mathfrak{s}_{c,m} : \mathbb{S}_{\mathcal{M}}^c(c \triangleright m) \xrightarrow{\sim} c^{v^v} \triangleright \mathbb{S}_{\mathcal{M}}^c(m) \quad (2.10)$$

for  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$ , obeying an appropriate pentagon axiom. Furthermore, given a  $\mathcal{C}$ -module functor  $H : \mathcal{M} \rightarrow \mathcal{N}$  there is a natural isomorphism

$$\mathfrak{s}_H : \mathbb{S}_{\mathcal{N}}^c \circ H \xrightarrow{\sim} H^{r^r a} \circ \mathbb{S}_{\mathcal{M}}^c \quad (2.11)$$

of twisted module functors.

The relative Serre functors of an exact  $(\mathcal{C}, \mathcal{D})$ -bimodule category  $\mathcal{M}$  are related to its Nakayama functors by the isomorphisms

$$D_c \triangleright \mathbb{N}_{\mathcal{M}}^r \cong \mathbb{S}_{\mathcal{M}}^c \quad \text{and} \quad D_c^{-1} \triangleright \mathbb{N}_{\mathcal{M}}^l \cong \overline{\mathbb{S}}_{\mathcal{M}}^c \quad (2.12)$$

of twisted bimodule functors [FSS20, Thm. 4.26]. Similarly, there exist isomorphisms of twisted bimodule functors for the relative Serre functors of  $\mathcal{M}_{\mathcal{D}}$ . These isomorphisms lead to an extension of Radford's theorem to module categories:

**Theorem 2.2.** [FGJS22, Cor. 4.16] *Let  ${}_c\mathcal{M}$  be an exact module category. There exists a natural isomorphism*

$$\mathcal{R}_{\mathcal{M}} : D_c \triangleright - \xrightarrow{\cong} \mathbb{S}_{\mathcal{M}}^c \circ \mathbb{S}_{\mathcal{M}}^c \triangleleft D_{c^*} = D_{c^*} \circ \mathbb{S}_{\mathcal{M}}^c \circ \mathbb{S}_{\mathcal{M}}^c \quad (2.13)$$

*of twisted bimodule functors.*

### 3. $\otimes$ -FROBENIUS FUNCTORS

A  $\mathbb{k}$ -linear functor with isomorphic adjoints is called a Frobenius functor. In the case of a tensor functor, its adjoint functors are endowed with additional structure, namely the structure of bimodule functors. In this section, we introduce the notion of  $\otimes$ -Frobenius tensor functors, which account for the additional data. We provide a list of characterizations and show some relations to Frobenius monoidal functors.

#### 3.1. Perfect tensor functors and relative modular objects

The relative modular function of a Hopf algebra extension studied in [FMS97] finds a categorical analogue in the notion of the relative modular object of a tensor functor defined in [Shi17]. Under certain conditions, this object measures the failure to which a tensor functor preserves the distinguished invertible object (2.7) and the Radford isomorphism of a finite tensor category. We first need the notion of a perfect tensor functor:

**Definition 3.1.** [BN14, §2.1] A tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between multitensor categories is called *perfect* iff it admits a right adjoint that is exact.

Being perfect is a property of a tensor functor. In the finite case, this notion has several characterizations listed in the lemma that follows.

**Lemma 3.2.** [Shi17, Lemma 4.3] *Let  $\mathcal{C}$  and  $\mathcal{D}$  be finite multitensor categories and  $F: \mathcal{C} \rightarrow \mathcal{D}$  a tensor functor. Then the following are equivalent:*

- (i)  $F$  admits a double-right adjoint.
- (ii)  $F$  admits a double-left adjoint.
- (iii)  $F$  preserves projective objects, i.e.  $F(p) \in \mathcal{D}$  is projective for every projective  $p \in \mathcal{C}$ .
- (iv) There exists an object  $\chi_F \in \mathcal{D}$  such that  $F^{\text{la}} \cong F^{\text{ra}}(- \otimes \chi_F) \cong F^{\text{ra}}(\chi_F \otimes -)$  as  $\mathbb{k}$ -linear functors.
- (v) There exists  $c \in \mathcal{C}$  such that  $F(c) \in \mathcal{D}$  is projective.
- (vi)  $F$  is perfect.

*Proof.* The equivalence (i)  $\iff$  (ii)  $\iff$  (iii)  $\iff$  (iv) is proven in [Shi17, Lemma 4.3] where it is also shown that the object  $\chi_F$  can be realized as  $F^{\text{rra}}(\mathbf{1})$  (note that  $\chi_F$  of this paper is  $\chi_F^*$  of [Shi17]). As  $\mathcal{C}$  is finite and therefore admits projective objects, (iii)  $\implies$  (v) is immediate. Using the same argument as in [EG17, Lemma 2.4], (v)  $\implies$  (iii) follows. Finally, since  $\mathcal{C}$  and  $\mathcal{D}$  are finite, (vi) is equivalent to (i).  $\square$

A first consequence of Lemma 3.2 is that surjective tensor functors are perfect, since they preserve projective objects [EO04, Thm. 2.5]. The converse is not true: for instance, consider any tensor functor  $F$  between fusion categories that is not surjective (like  $\text{Vec}_H \subset \text{Vec}_G$  for a proper subgroup  $H \subset G$ ), then  $F$  preserves projectives, but is not surjective. In fact, any tensor functor from a non-semisimple tensor category to a semisimple tensor category is perfect. Also notice that not every tensor functor is perfect: for a finite tensor category  $\mathcal{C}$  that is not semisimple, the tensor functor  $\text{Vec} \rightarrow \mathcal{C}$  given by  $\mathbb{k} \mapsto \mathbf{1}$  does not preserve projective objects.

**Definition 3.3.** [Shi17, Def. 4.4] The object  $\chi_F \in \mathcal{D}$  from Lemma 3.2 (iv) is called *the relative modular object of  $F$* .

**Proposition 3.4.** [Shi17] *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. The relative modular object of  $F$  obeys the following properties:*

- (i)  $\chi_F$  is an invertible object in  $\mathcal{D}$ .
- (ii) There is a natural isomorphism  $F^{\text{rra}} \cong F(-) \otimes \chi_F$  of  $\mathbb{k}$ -linear functors.
- (iii) There is a natural isomorphism  $F^{\text{lla}} \cong F(-) \otimes \chi_F^{-1}$  of  $\mathbb{k}$ -linear functors.
- (iv) There is an isomorphism  $\chi_F \cong F^{\text{rra}}(\mathbf{1}_{\mathcal{C}})$ .
- (v) There is an isomorphism  $\chi_F \cong F(D_{\mathcal{C}}) \otimes D_{\mathcal{D}}^{-1}$ .

*Proof.* Part (i) is proven in [Shi17, Prop. 4.7(2)]. By taking right adjoints to the isomorphism  $F^{\text{lla}} \cong F^{\text{ra}}(- \otimes \chi_F)$  from Lemma 3.2(iv), we obtain that

$$F \cong (F^{\text{lla}})^{\text{ra}} \cong (F^{\text{ra}} \circ (- \otimes \chi_F))^{\text{ra}} \cong (- \otimes \chi_F)^{\text{ra}} \circ F^{\text{rra}} \cong F^{\text{rra}}(-) \otimes \chi_F^{-1}.$$

Tensoring by  $\chi_F$  on both sides yields (ii). Part (iii) is proved in a similar manner. Lastly, since  $F(\mathbf{1}_{\mathcal{C}}) \cong \mathbf{1}_{\mathcal{D}}$ , part (iv) follows by (ii). According to [FSS20, Rem. 4.17]  $F^{\text{lla}}(D_{\mathcal{C}}) \cong D_{\mathcal{D}}$ . The isomorphism in (v) is then obtained by using (iii).  $\square$

**Definition 3.5.** [CMZ97] A ( $\mathbb{k}$ -linear) functor  $F: \mathcal{X} \rightarrow \mathcal{Y}$  is called *Frobenius* if it admits a two-sided adjoint, i.e. it admits both a left adjoint and a right adjoint and  $F^{\text{la}} \cong F^{\text{ra}}$ .

Given an adjoint pair  $F \dashv G$ , note that  $F$  is Frobenius if and only if  $G$  is Frobenius. Another immediate consequence of Definition 3.5 is that a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  that is Frobenius must be perfect as well:  $F$  admits a double-right adjoint, namely  $F^{\text{rra}} \cong F$ . However, the converse is not true as shown by the following example.

*Example 3.6.* For a finite tensor category  $\mathcal{C}$ , the forgetful functor  $U: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$  is surjective and therefore perfect. However,  $U$  is Frobenius if and only if  $\mathcal{C}$  is unimodular [Shi16, Thm. 4.10].

The following result relates the property of  $F$  being Frobenius and  $F^{\text{ra}}$  being Frobenius monoidal.

**Proposition 3.7.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a tensor functor between tensor categories admitting a right adjoint  $F^{\text{ra}}: \mathcal{D} \rightarrow \mathcal{C}$ .*

- (i) *If the lax functor  $F^{\text{ra}}$  admits a Frobenius monoidal structure, then  $F$  must be Frobenius.*
- (ii) *If  $F^{\text{ra}}$  is exact and faithful, then  $F$  being Frobenius implies that the induced lax and oplax structures turn  $F^{\text{ra}}$  into a Frobenius monoidal functor.*

*Proof.* The left and right adjoints of  $F$  are related by  $F^{\text{la}}(d) \cong {}^{\vee}(F^{\text{ra}}(d^{\vee}))$  [Shi17, Lem. 4.1]. As  $F^{\text{ra}}$  is Frobenius monoidal it preserves duals. Thus, there is a natural isomorphism

$$F^{\text{la}}(d) \cong {}^{\vee}(F^{\text{ra}}(d^{\vee})) \cong {}^{\vee}(F^{\text{ra}}(d)^{\vee}) \cong F^{\text{ra}}(d),$$

thereby proving that  $F$  is Frobenius. To prove (ii), notice that  $F$  being Frobenius implies that  $F^{\text{ra}}(\mathbf{1}_{\mathcal{D}})$  is a Frobenius algebra in  $\mathcal{C}$ : according to [BN11, Prop. 6.1],  $\mathcal{D}$  can be identified with the category  $\text{mod}_A(\mathcal{C})$  with  $A = F^{\text{ra}}(\mathbf{1}_{\mathcal{D}})$ . Under this identification,  $F^{\text{ra}}: \text{mod}_A(\mathcal{C}) \rightarrow \mathcal{C}$  is the forgetful functor. By [Shi16, Lemma 2.3], we get that  $A$  is a Frobenius algebra iff  $F^{\text{ra}}$  is a Frobenius functor iff  $F$  is a Frobenius functor. Then, as explained in [Yad24, §3.1], we can apply [Bal17, Prop. 4.5] to obtain that  $F^{\text{ra}}$  is endowed with a Frobenius monoidal structure.  $\square$

**Question 3.8.** Note that by [BN11, Lem. 3.1] the assumption that  $F^{\text{ra}}$  is faithful is equivalent to asking that  $F$  is dominant, that is, every object in  $\mathcal{D}$  is a quotient of  $F(c)$  for some  $c \in \mathcal{C}$ . Does the statement in Proposition 3.7 (ii) hold without assuming that  $F$  is dominant?

**Proposition 3.9.** Given a perfect tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , the following hold:

- (i)  $F$  is Frobenius if and only if  $\chi_F \cong \mathbb{1}_{\mathcal{D}}$ .
- (ii) If  $\mathcal{C}$  is unimodular, then  $F$  is Frobenius if and only if  $\mathcal{D}$  is unimodular.
- (iii) If  $\mathcal{C}$  is semisimple, then  $F$  is Frobenius.

*Proof.* The first statement is proven in [Shi17, Prop. 4.7(i)]. If  $\mathcal{C}$  is unimodular, by Proposition 3.4 (v),  $D_{\mathcal{D}}^{-1} \cong \chi_F$ . Then, (ii) follows from (i). For (iii), note that when  $\mathcal{C}$  is semisimple,  $\mathbb{1}_{\mathcal{C}}$  is projective. As  $F$  preserves projectives,  $\mathbb{1}_{\mathcal{D}} \cong F(\mathbb{1}_{\mathcal{C}})$  is projective. Consequently,  $\mathcal{D}$  is semisimple. It follows that both  $\mathcal{C}$  and  $\mathcal{D}$  are unimodular. Therefore,  $\chi_F \cong \mathbb{1}_{\mathcal{D}}$  and by part (i),  $F$  is Frobenius.  $\square$

*Remark 3.10.* The unimodularity of  $\mathcal{C}$  in Proposition 3.9 (ii) is crucial. As Example 5.5 shows, it might be the case that  $F$  is Frobenius and  $\mathcal{D}$  is unimodular, but  $\mathcal{C}$  is not unimodular.

Perfect, Frobenius and surjective are properties of a tensor functor that behave well under composition of tensor functors.

**Proposition 3.11.** Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{B} \rightarrow \mathcal{C}$  be tensor functors, we have that:

- (i) If  $F$  and  $G$  are perfect, their composition  $F \circ G: \mathcal{B} \rightarrow \mathcal{D}$  is perfect and its relative modular object is given by

$$\chi_{F \circ G} \cong F(\chi_G) \otimes \chi_F.$$

- (ii) If  $F$  and  $G$  are Frobenius, the composition  $F \circ G$  is Frobenius.
- (iii) If  $F$  and  $G$  are surjective, the composition  $F \circ G$  is surjective.

*Proof.* If each  $F$  and  $G$  admit double-adjoints, then their composition also admits double-adjoints and from Lemma 3.2 it follows that  $F \circ G$  is perfect showing (i). We can compute relative modular objects in terms of double-adjoints according to Proposition 3.4(ii):

$$\chi_{F \circ G} \cong (F \circ G)^{\text{rra}}(\mathbb{1}_{\mathcal{B}}) \cong (F^{\text{rra}} \circ G^{\text{rra}})(\mathbb{1}_{\mathcal{B}}) \cong F^{\text{rra}}(\chi_G) \cong F(\chi_G) \otimes \chi_F.$$

Assertion (ii) follows from (i) and Proposition 3.9(i). Now, in order to prove (iii), notice that since  $F$  is surjective, any  $z \in \mathcal{D}$  is a subquotient of  $F(y)$  for some  $y \in \mathcal{C}$ . Also, since  $G$  is surjective,  $y$  is a subquotient of  $G(x)$  for some  $x \in \mathcal{B}$ . Now, left exactness of  $F$  implies that it preserves kernels (hence subobjects). Also, right exactness of  $F$  implies that it preserves cokernels (hence quotient objects). Hence,  $F$  preserves subquotients. Thus,  $F(y)$  is a subquotient of  $FG(x)$ . Since the subquotient of a subquotient of some object  $p$  is also a subquotient of  $p$  [EGNO16, Ex. 1.3.6], we conclude that  $z$  is a subquotient of  $GF(x)$ .  $\square$

### 3.2. Center of a perfect tensor functor

In this section, we recall the notion of the center  $\mathcal{Z}(F)$  of a tensor functor  $F$ , and describe its relative modular object in case it is perfect.

Let  $\mathcal{C}$  be a finite tensor category and  $\mathcal{M}$  a  $\mathcal{C}$ -bimodule category. The *center of  $\mathcal{M}$* , denoted by  $\mathcal{Z}(\mathcal{M})$ , is the category whose objects are pairs  $(m, \gamma)$ , where  $m \in \mathcal{M}$  and  $\gamma_c: m \triangleleft c \rightarrow c \triangleright m$  is

natural isomorphism obeying a hexagon axiom. The hom spaces of  $\mathcal{Z}(\mathcal{M})$  are given by

$$\mathrm{Hom}_{\mathcal{Z}(\mathcal{M})}((m, \gamma), (m', \gamma')) := \{f \in \mathrm{Hom}_{\mathcal{M}}(m, m') \mid (f \otimes \mathrm{id}_c) \circ \gamma_c = \gamma'_c \circ (\mathrm{id}_c \otimes f) \ \forall c \in \mathcal{C}\}.$$

The center construction extends to a strict pseudo-functor [Shi20, §3.6], [FLP24c, Lem. 4.3]

$$\mathcal{Z} : \mathbf{Bimod}(\mathcal{C}) \longrightarrow \mathrm{FinCat}_{\mathbb{k}}, \quad \mathcal{M} \longmapsto \mathcal{Z}(\mathcal{M}) \quad (3.1)$$

from the 2-category of  $\mathcal{C}$ -bimodule categories, bimodule functors and bimodule natural transformations to the 2-category of finite  $\mathbb{k}$ -linear categories,  $\mathbb{k}$ -linear functors and natural transformations.

Let  $F, G : \mathcal{C} \longrightarrow \mathcal{D}$  be tensor functors between finite tensor categories. We use the notation  ${}_F\mathcal{D}_G$  to denote the  $\mathcal{C}$ -bimodule category  $\mathcal{D}$  with left and right actions given by

$$c \triangleright d \triangleleft c' := F(c) \otimes d \otimes G(c').$$

The  $F$ -centralizer of  $\mathcal{D}$  is defined as the center  $\mathcal{Z}({}_F\mathcal{D}_F)$  of the  $\mathcal{C}$ -bimodule category  ${}_F\mathcal{D}_F$ . Explicitly, its objects are pairs  $(d, \gamma)$  consisting of an object  $d \in \mathcal{D}$  and a  $F$ -half-braiding  $\gamma$ , i.e. a natural isomorphism  $\gamma_c : d \otimes F(c) \rightarrow F(c) \otimes d$  for  $c \in \mathcal{C}$ , obeying the respective hexagon axiom. The  $F$ -centralizer of  $\mathcal{D}$  is endowed with a monoidal structure defined via

$$(d, \gamma) \otimes (d', \gamma') = (d \otimes d', \rho)$$

where the  $F$ -half-braiding  $\rho_c$  is the composition

$$d \otimes d' \otimes F(c) \xrightarrow{\mathrm{id} \otimes \gamma'_c} d \otimes F(c) \otimes d' \xrightarrow{\gamma_c \otimes \mathrm{id}} F(c) \otimes d \otimes d'$$

and the monoidal unit  $\mathbb{1}_{\mathcal{Z}({}_F\mathcal{D}_F)}$  is given by  $\mathbb{1}_{\mathcal{D}}$  together with the trivial half-braiding. Moreover,  $\mathcal{Z}({}_F\mathcal{D}_F)$  is rigid [Maj91, Thm. 3.3] and finite [Shi23b, Thm. 3.8], therefore  $\mathcal{Z}({}_F\mathcal{D}_F)$  is a finite tensor category. By considering the identity tensor functor  $F = \mathrm{id}_{\mathcal{C}}$ , we recover the *Drinfeld center*  $\mathcal{Z}(\mathcal{C})$  of  $\mathcal{C}$ , which is further endowed with the structure of a braiding.

Now, the functor  $F$  can be seen as a  $\mathcal{C}$ -bimodule functor  $F : \mathcal{C} \longrightarrow {}_F\mathcal{D}_F$ . The center construction (3.1) provides a  $\mathbb{k}$ -linear functor

$$\mathcal{Z}(F) : \mathcal{Z}(\mathcal{C}) \longrightarrow \mathcal{Z}({}_F\mathcal{D}_F), \quad (c, \nu) \longmapsto (F(c), \gamma)$$

where  $\gamma_{c'}$  is given by the composition

$$F(c) \otimes F(c') \xrightarrow{F_2} F(c \otimes c') \xrightarrow{F(\nu_{c'})} F(c' \otimes c) \xrightarrow{F_2^{-1}} F(c') \otimes F(c).$$

In addition, the strong monoidal structure of  $F$  induces a strong monoidal structure on  $\mathcal{Z}(F)$ .

**Lemma 3.12.** *Let  $F : \mathcal{C} \longrightarrow \mathcal{D}$  be a tensor functor between finite tensor categories.*

- (i)  $\mathcal{Z}({}_F\mathcal{D}_F)$  is a finite tensor category.
- (ii)  $\mathcal{Z}(F) : \mathcal{Z}(\mathcal{C}) \longrightarrow \mathcal{Z}({}_F\mathcal{D}_F)$  is a tensor functor.
- (iii)  $\mathcal{Z}(F^{\mathrm{ra}})$  is right adjoint to  $\mathcal{Z}(F)$ .

*Proof.* As  $\mathcal{C}$  and  $\mathcal{D}$  are finite and  $F$  is exact, it admits a right adjoint  $F^{\mathrm{ra}}$  which is endowed with a lax monoidal structure such that the adjunction  $F \dashv F^{\mathrm{ra}}$  is monoidal. Then, by [FLP24c, Prop. 4.8],  $\mathcal{Z}(F^{\mathrm{ra}})$  is right adjoint to  $\mathcal{Z}(F)$  and the adjunction  $\mathcal{Z}(F) \dashv \mathcal{Z}(F^{\mathrm{ra}})$  is monoidal. Since  $\mathcal{Z}(F)$  is a monoidal functor between rigid categories that admits a right adjoint, then it also admits a left adjoint. As  $\mathcal{Z}(\mathcal{C})$  and  $\mathcal{Z}({}_F\mathcal{D}_F)$  are finite, it follows that  $\mathcal{Z}(F)$  is exact and thus a tensor functor.  $\square$

**Proposition 3.13.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. The tensor functor  $\mathcal{Z}(F)$  is perfect. Moreover,  $\chi_F$  is endowed with a  $F$ -half-braiding*

$$\sigma_c: \chi_F \otimes F(c) \xrightarrow{\sim} F(c) \otimes \chi_F, \quad \text{for } c \in \mathcal{C}, \quad (3.2)$$

such that the object  $\chi_{\mathcal{Z}(F)} := (\chi_F, \sigma)$  in  $\mathcal{Z}({}_F\mathcal{D}_F)$  is the relative modular object of  $\mathcal{Z}(F)$ .

*Proof.* Since  $F$  is perfect, it admits a double-right adjoint  $F^{rra}$ . This functor can be seen as a  $\mathcal{C}$ -bimodule functor  $F^{rra}: \mathcal{C} \rightarrow {}_F\mathcal{D}_F$ . As pseudo-functors preserve internal adjunctions, it follows that, under the center construction (3.1), the functor  $\mathcal{Z}(F^{rra})$  serves as double-right adjoint to  $\mathcal{Z}(F)$ , and thus  $\mathcal{Z}(F)$  is perfect. By the definition of the center construction (3.1), we have that the functor  $\mathcal{Z}(F^{rra})$  is explicitly given by

$$(c', \nu) \in \mathcal{Z}(\mathcal{C}) \mapsto (F^{rra}(c'), \gamma^{c'}) \in \mathcal{Z}({}_F\mathcal{D}_F)$$

where the  $F$ -half-braiding  $\gamma^{c'}$  is defined using the  $\mathcal{C}$ -bimodule structure of  $F^{rra}$  as the composition

$$\gamma^{c'}: F^{rra}(c') \otimes F(c) \xrightarrow{\sim} F^{rra}(c' \otimes c) \xrightarrow{F^{rra}(\nu_c)} F^{rra}(c \otimes c') \xrightarrow{\sim} F(c) \otimes F^{rra}(c').$$

According to Proposition 3.4(iv), we have that  $\chi_F \cong F^{rra}(\mathbb{1}_{\mathcal{C}})$ . Then the  $F$ -half-braiding associated to  $c' = \mathbb{1}_{\mathcal{C}}$  provides the desired  $\sigma$ . Again by Proposition 3.4(iv), the relative modular object of  $\mathcal{Z}(F)$  can be obtained by evaluation at the monoidal unit, thus

$$\chi_{\mathcal{Z}(F)} \cong \mathcal{Z}(F^{rra})(\mathbb{1}_{\mathcal{Z}(\mathcal{C})}) = (F^{rra}(\mathbb{1}_{\mathcal{C}}), \sigma) \cong (\chi_F, \sigma)$$

which completes the proof.  $\square$

The pair  $\chi_{\mathcal{Z}(F)} = (\chi_F, \sigma)$  measures the extend to which the Radford isomorphism is preserved under  $F$ , in the following sense:

**Proposition 3.14.** [Shi23b, Thm. 4.4] *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. There exists an isomorphism*

$$F(D_{\mathcal{C}}) \xrightarrow{\sim} \chi_F \otimes D_{\mathcal{D}} \quad (3.3)$$

fulfilling the commutativity of the following diagram

$$\begin{array}{ccccc} F(D_{\mathcal{C}} \otimes c) & \xrightarrow{\cong} & F(D_{\mathcal{C}}) \otimes F(c) & \xrightarrow{(3.3)} & \chi_F \otimes D_{\mathcal{D}} \otimes F(c) \\ F(\mathcal{R}_{\mathcal{C}}) \downarrow & & & & \downarrow \text{id} \otimes \mathcal{R}_{\mathcal{D}} \\ F(c^{\vee\vee\vee\vee} \otimes D_{\mathcal{C}}) & & & & \chi_F \otimes F(c)^{\vee\vee\vee\vee} \otimes D_{\mathcal{D}} \\ \cong \downarrow & & & & \downarrow \cong \\ F(c^{\vee\vee\vee\vee}) \otimes F(D_{\mathcal{C}}) & \xrightarrow{(3.3)} & F(c^{\vee\vee\vee\vee}) \otimes \chi_F \otimes D_{\mathcal{D}} & \xleftarrow{\sigma_c^{\vee 4} \otimes \text{id}} & \chi_F \otimes F(c^{\vee\vee\vee\vee}) \otimes D_{\mathcal{D}} \end{array} \quad (3.4)$$

where  $\mathcal{R}_{\mathcal{C}}$  (resp.  $\mathcal{R}_{\mathcal{D}}$ ) is the Radford isomorphism, and  $\sigma$  is the isomorphism (3.2).

*Proof.* A similar statement is proven in [Shi23b, Thm. 4.4], where an isomorphism of the form  $F(D_{\mathcal{C}}) \xrightarrow{\sim} D_{\mathcal{D}} \otimes \chi_F$  is considered instead and the diagram (3.4) is appropriately adapted. In fact, we can obtain diagram (3.4) from the diagram in [Shi23b, Thm. 4.4] by means of the Radford

isomorphism  $(\mathcal{R}_{\mathcal{D}})_{\chi}: D_{\mathcal{D}} \otimes_{\chi_F} \cong \chi_F \otimes D_{\mathcal{D}}$ . It is enough to check that the following diagram commutes

$$\begin{array}{ccccc}
 \chi_F \otimes D_{\mathcal{D}} \otimes F(c) & \xrightarrow{\text{id} \otimes \mathcal{R}_{\mathcal{D}}} & \chi_F \otimes F(c^{\vee\vee\vee\vee}) \otimes D_{\mathcal{D}} & \xrightarrow{\sigma \otimes \text{id}} & F(c^{\vee\vee\vee\vee}) \otimes \chi_F \otimes D_{\mathcal{D}} \\
 \uparrow \mathcal{R}_{\mathcal{D}} \otimes \text{id} & & \swarrow \mathcal{R}_{\mathcal{D}} & & \swarrow \mathcal{R}_{\mathcal{D}} \\
 D_{\mathcal{D}} \otimes \chi_F \otimes F(c) & \xrightarrow{\text{id} \otimes \sigma} & D_{\mathcal{D}} \otimes F(c) \otimes \chi_F & \xrightarrow{\mathcal{R}_{\mathcal{D}} \otimes \text{id}} & F(c^{\vee\vee\vee\vee}) \otimes D_{\mathcal{D}} \otimes \chi_F \\
 & & & & \uparrow \text{id} \otimes \mathcal{R}_{\mathcal{D}}
 \end{array}$$

where the identification  $F(c^{\vee\vee\vee\vee}) \cong F(c)^{\vee\vee\vee\vee}$  is implicit. The commutativity of the triangles follow from the monoidality of the Radford isomorphism and the remaining middle square commutes due to naturality of the  $F$ -half-braiding  $\sigma$ .  $\square$

### 3.3. The properties of $\otimes$ -Frobenius functors

A tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  can be viewed a  $\mathcal{C}$ -bimodule functor  $F: \mathcal{C} \rightarrow {}_F\mathcal{D}_F$ . Moreover, it admits left and right adjoints that are also  $\mathcal{C}$ -bimodule functors [DSPS19, Cor. 2.13]. This motivates the following stronger version of Definition 3.5 that accounts for the additional structure.

**Definition 3.15.** A tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between finite multitensor categories is called  $\otimes$ -Frobenius iff there exists a  $\mathcal{C}$ -bimodule natural isomorphism  $F^{\text{la}} \cong F^{\text{ra}}: {}_F\mathcal{D}_F \rightarrow \mathcal{C}$ .

The notion of a  $\otimes$ -Frobenius tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  is an instance of an ambi-adjunction (see [FLP24a, Def. 2.7] for a definition) internal to the 2-category  $\mathbf{Bimod}(\mathcal{C})$ . Clearly, a bimodule natural isomorphism  $F^{\text{la}} \cong F^{\text{ra}}$  is in particular a natural isomorphism. Therefore, any  $\otimes$ -Frobenius functor is Frobenius and thus perfect. The converse does not hold, an example of a (surjective) Frobenius functor that is not  $\otimes$ -Frobenius can be found in Example 5.5. It is worth pointing out that  $\otimes$ -Frobenius is a property of a tensor functor  $F$ , in the sense that it asks for the existence and not for a specific choice of bimodule isomorphism between the left and right adjoint functors of  $F$ .

*Remark 3.16.* Given  $\otimes$ -Frobenius functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{B} \rightarrow \mathcal{C}$ , the composition  $F \circ G$  is also  $\otimes$ -Frobenius. Indeed, if  $F \cong F^{\text{ra}}$  and  $G \cong G^{\text{ra}}$  as bimodule functors, then  $F \circ G \cong F^{\text{ra}} \circ G^{\text{ra}} \cong (F \circ G)^{\text{ra}}$  as bimodule functors. The second isomorphism is obtained by the uniqueness of adjoints in the 2-category  $\mathbf{Bimod}(\mathcal{C})$ .

Another consequence of being  $\otimes$ -Frobenius (and thus Frobenius) is the following: as  $F$  is strong monoidal,  $F^{\text{ra}}$  is endowed with a lax monoidal structure and  $F^{\text{la}}$  with an oplax monoidal structure. Using the isomorphism  $F^{\text{la}} \cong F^{\text{ra}}$ ,  $F^{\text{ra}}$  also inherits an oplax structure. Next, directly following the results in [FLP24a], we obtain that these two structures form a Frobenius monoidal functor.

**Proposition 3.17.** [FLP24a, Thm. 3.18] *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a  $\otimes$ -Frobenius functor. Then  $F^{\text{ra}}$  is endowed with a Frobenius monoidal structure.*

*Proof.* Since  $F$  is  $\otimes$ -Frobenius,  $F^{\text{la}} \cong F^{\text{ra}}$  in the 2-category  $\mathbf{Bimod}(\mathcal{C})$ . Thus, by the implication (2)  $\implies$  (1) in [FLP24a, Thm. 3.17], the Hopf operators (2.3) of  $F^{\text{ra}} \dashv F$  and the coHopf operators (2.2) of  $F \dashv F^{\text{ra}}$  satisfy  $H_{d,c}^l = (h_{d,c}^l)^{-1}$  and  $H_{d,c}^r = (h_{d,c}^r)^{-1}$ . By [FLP24a, Thm. 3.18], the second relation implies the claim.  $\square$

We show next that the notion of a  $\otimes$ -Frobenius tensor functor is closely related to relative modular objects. To make this relation precise, we need the following additional fact. The  $F$ -centralizer of  $\mathcal{D}$  can be realized as the category of  $\mathcal{C}$ -bimodule functors from  $\mathcal{C}$  to  ${}_F\mathcal{D}_F$ , denoted by  $\text{Func}_{|\mathcal{C}}(\mathcal{C}, {}_F\mathcal{D}_F)$ . Explicitly, there is an equivalence of categories [Shi23b, Lemma 3.1]

$$\text{Func}_{|\mathcal{C}}(\mathcal{C}, {}_F\mathcal{D}_F) \xrightarrow{\sim} \mathcal{Z}({}_F\mathcal{D}_F), \quad H \mapsto (H(\mathbb{1}), \gamma) \quad (3.5)$$

where

$$\gamma_c: H(\mathbb{1}) \otimes F(c) = H(\mathbb{1}) \triangleleft^F c \xrightarrow{\sim} H(c) \xrightarrow{\sim} c \triangleright^F H(\mathbb{1}) = F(c) \otimes H(\mathbb{1})$$

is given by the bimodule functor structure of  $H$ .

The following assignment serves as a quasi-inverse of (3.5). For  $(d, \gamma) \in \mathcal{Z}({}_F\mathcal{D}_F)$  consider the functor  $F(-) \otimes d$  together with  $\mathcal{C}$ -bimodule functor structure defined via

$$F(c_1 \otimes c' \otimes c_2) \otimes d \xrightarrow{F_2} F(c_1) \otimes F(c') \otimes F(c_2) \otimes d \xrightarrow{\gamma^{-1}} F(c_1) \otimes F(c') \otimes d \otimes F(c_2).$$

Now, using the equivalence (3.5) and our previous results, we obtain the following a list of characterizations of the notion of a  $\otimes$ -Frobenius functors.

**Theorem 3.18.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. The following are equivalent:*

- (i)  $F$  is  $\otimes$ -Frobenius.
- (ii) There is a natural isomorphism  $F \cong F^{\text{rra}}$  of  $\mathcal{C}$ -bimodule functors.
- (iii) The relative modular object  $\chi_{\mathcal{Z}(F)} = (\chi_F, \sigma)$  is trivial in  $\mathcal{Z}({}_F\mathcal{D}_F)$ , i.e.  $\chi_F \cong \mathbb{1}_{\mathcal{D}}$  and  $\sigma_c = \text{id}_{F(c)}$  for all  $c \in \mathcal{C}$ .
- (iv) The tensor functor  $\mathcal{Z}(F)$  is Frobenius.
- (v) The finite tensor category  $\mathcal{Z}({}_F\mathcal{D}_F)$  is unimodular.
- (vi) The lax monoidal functor  $\mathcal{Z}(F^{\text{ra}})$  admits a Frobenius monoidal structure.

*Proof.* Since  $F$  is perfect, it admits double-adjoints. Then, by taking adjoints in  $\mathbf{Bimod}(\mathcal{C})$ , there exists a  $\mathcal{C}$ -bimodule natural isomorphism  $F^{\text{la}} \cong F^{\text{ra}}$  if and only if there exists a bimodule natural isomorphism  $F \cong F^{\text{rra}}$ , which proves (i)  $\iff$  (ii).

To prove (ii)  $\iff$  (iii), notice that the category equivalence (3.5) assigns  $(\chi_F, \sigma)$  to  $F^{\text{rra}}$  as defined in Proposition 3.13. Moreover, (3.5) sends  $F$  to the unit of  $\mathcal{Z}({}_F\mathcal{D}_F)$ . Then,  $F \cong F^{\text{rra}}$  in  $\text{Func}_{|\mathcal{C}}(\mathcal{C}, {}_F\mathcal{D}_F)$  if and only if  $\mathbb{1} \cong (\chi_F, \sigma)$  in  $\mathcal{Z}({}_F\mathcal{D}_F)$ .

Since, by Proposition 3.13,  $(\chi_F, \sigma)$  is the relative modular object of  $\mathcal{Z}(F)$ , the equivalence (iii)  $\iff$  (iv) follows by Proposition 3.9(i). Next, note that

$$(\chi_F, \sigma) = \chi_{\mathcal{Z}(F)} \cong F(D_{\mathcal{Z}(\mathcal{C})}) \otimes D_{\mathcal{Z}({}_F\mathcal{D}_F)}^{-1} \cong D_{\mathcal{Z}({}_F\mathcal{D}_F)}^{-1},$$

where the last isomorphism holds because  $\mathcal{Z}(\mathcal{C})$  is unimodular. This implies that (iv)  $\iff$  (v).

(i)  $\implies$  (vi) follows by a direct application of [FLP24a, Thm. 3.17] and [FLP24a, Prop. 4.15]. The implication (vi)  $\implies$  (iv) follows by Proposition 3.7.  $\square$

Note that there is a strong monoidal functor  $\text{res}: \mathcal{Z}(\mathcal{D}) \rightarrow \mathcal{Z}({}_F\mathcal{D}_F)$  given by restricting the half-braiding to objects of the form  $F(c)$ . Using this, we obtain the lax monoidal functor:

$$\mathcal{Z}(F^{\text{ra}})': \quad \mathcal{Z}(\mathcal{D}) \xrightarrow{\text{res}} \mathcal{Z}({}_F\mathcal{D}_F) \xrightarrow{\mathcal{Z}(F^{\text{ra}})} \mathcal{Z}(\mathcal{C}). \quad (3.6)$$



Since  $F \dashv F^{ra}$  is a monoidal adjunction between rigid monoidal categories, by [FLP24c, Cor. 4.11], the functor  $\mathcal{Z}(F^{ra})'$  becomes a braided lax monoidal functor. Then we obtain the following result which may be considered as a generalization of Proposition 3.17.

**Corollary 3.19.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a  $\otimes$ -Frobenius functor. Then, the induced functor  $\mathcal{Z}(F^{ra})'$  from (3.6) is endowed with a (braided) Frobenius monoidal structure.*

*Proof.* The functor  $\text{res}$  is strong monoidal and therefore Frobenius monoidal. Moreover, by Theorem 3.18,  $\mathcal{Z}(F^{ra})$  is Frobenius monoidal. As Frobenius monoidal functors compose, the claim follows.  $\square$

Let  $\mathcal{B}$  be a braided finite tensor category and  $\mathcal{D}$  a finite tensor category. A *central tensor functor* consists of a tensor functor  $F: \mathcal{B} \rightarrow \mathcal{D}$  together with a braided functor  $\tilde{F}: \mathcal{B} \rightarrow \mathcal{Z}(\mathcal{D})$  and a monoidal natural isomorphism  $F \cong U \circ \tilde{F}$ , where  $U: \mathcal{Z}(\mathcal{D}) \rightarrow \mathcal{D}$  is the forgetful functor.

**Proposition 3.20.** *Let  $\mathcal{B}$  be a braided finite tensor category,  $\mathcal{D}$  a finite tensor category and  $F: \mathcal{B} \rightarrow \mathcal{D}$  a central tensor functor. Then, the following are equivalent:*

- (i)  $F$  is  $\otimes$ -Frobenius.
- (ii)  $F$  is Frobenius.
- (iii) The lax monoidal functor  $F^{ra}$  admits a Frobenius monoidal structure.

*Proof.* That  $F$  is  $\otimes$ -Frobenius automatically implies that  $F$  is Frobenius. To prove the converse, we make use of [Shi23b, Thm. 5.6] that states that for  $b \in \mathcal{B}$ , the associated component of the  $F$ -half-braiding  $\sigma_b: \chi_F \otimes F(b) \xrightarrow{\sim} F(b) \otimes \chi_F$  agrees with the half-braiding of  $\tilde{F}(b)$  coming from the central structure of  $F$ . According to Proposition 3.9, we have that  $\chi_F \cong \mathbb{1}$ . Since the component of the half-braiding of  $\tilde{F}(b)$  associated to  $\mathbb{1}$  is the identity, it follows that  $\sigma_b = \text{id}_{F(b)}$ . We conclude, by means of Theorem 3.18, that  $F$  is  $\otimes$ -Frobenius. This establishes (i)  $\iff$  (ii). Proposition 3.17 states that (i)  $\implies$  (iii) and Proposition 3.7 (i) shows (iii)  $\implies$  (ii).  $\square$

*Example 3.21.* For a finite tensor category  $\mathcal{C}$ , consider the central functor  $U: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$ . Then:

$$\mathcal{C} \text{ is unimodular} \stackrel{[\text{Shi16}]}{\iff} U \text{ is Frobenius} \stackrel{\text{Prop. 3.20}}{\iff} U \text{ is } \otimes\text{-Frobenius.}$$

This generalizes an observation from the Hopf algebra case in [FLP24b, Corollary 5.1].

*Remark 3.22.* Proposition 3.20 implies that a braided tensor functor that is Frobenius is  $\otimes$ -Frobenius, as well. This has also been observed in [FLP24a]. Later in Corollary 4.19 and Propositions 4.20 and 4.24 we provide additional criteria in terms of module categories that allow to show that a functor is  $\otimes$ -Frobenius.

A semisimple finite tensor category is called a *fusion category*. Moreover, a fusion category is called *separable* if its global dimension is nonzero. If  $\text{char}(\mathbb{k}) = 0$ , then all fusion categories are separable according to [ENO05, Thm. 2.3]. This definition of separability is equivalent [DSPS18, Thm. 2.6.7] to the alternate notion given in [DSPS18] in terms of algebras.

**Theorem 3.23.** *If  $\mathcal{C}$  and  $\mathcal{D}$  are separable fusion categories and  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a tensor functor, then  $F$  is  $\otimes$ -Frobenius.*

*Proof.* We have the following equivalence of categories:

$$\mathcal{Z}({}_F\mathcal{D}_F) \simeq \text{Fun}_{\mathcal{C}|\mathcal{C}}(\mathcal{C}, {}_F\mathcal{D}_F) \simeq \text{Fun}_{\mathcal{C}\boxtimes\overline{\mathcal{C}}}(\mathcal{C}, {}_F\mathcal{D}_F) \simeq \overline{\mathcal{C}} \boxtimes_{\mathcal{C}\boxtimes\overline{\mathcal{C}}} {}_F\mathcal{D}_F.$$

As the categories  $\mathcal{C}$  and  ${}_F\mathcal{D}_F$  are semisimple, they are separable  $\mathcal{C}$ -bimodule categories because of [DSPS18, Prop. 2.5.10]. Moreover, the relative Deligne product of separable bimodules is separable [DSPS18, Thm. 2.5.5]. Since, by [DSPS18, Prop. 2.5.3], separable module categories are semisimple, we obtain that  $\mathcal{Z}({}_F\mathcal{D}_F)$  is semisimple and, in particular, unimodular. The claim follows from Theorem 3.18.  $\square$

*Remark 3.24.* Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a strong monoidal  $\mathbb{k}$ -linear functor between separable fusion categories. Assuming that  $F$  admits adjoints, then it is exact and thus a tensor functor. By Theorem 3.23, we have that  $F$  is  $\otimes$ -Frobenius. We can conclude from Corollary 3.19 that the functor  $\mathcal{Z}(F^{\text{ra}})'$  from (3.6) is Frobenius monoidal, thereby positively answering [FLP24b, Questions 5.22 & 5.23] with  $\mathbb{k}$  having arbitrary characteristic. It is worth pointing out that our current argument requires  $\mathbb{k}$  to be algebraically closed.

## 4. TWISTED MODULE CATEGORIES

Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a tensor functor between finite tensor categories. Given a left  $\mathcal{D}$ -module category  $\mathcal{M}$ , we obtain a left  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  by pulling the  $\mathcal{D}$ -action back along  $F$ , i.e.  ${}_F\mathcal{M}$  consists of the same underlying category as  $\mathcal{M}$ , but with the action defined by

$$c \triangleright^F m := F(c) \triangleright m$$

for every  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$ . We refer to the so obtained  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  as a *F-twisted module category*. Given a  $\mathcal{D}$ -module functor  $H: \mathcal{M} \rightarrow \mathcal{N}$ , denote by

$$F^*(H): {}_F\mathcal{M} \rightarrow {}_F\mathcal{N} \tag{4.1}$$

the  $\mathcal{C}$ -module functor given by  $H$  as a functor and with module structure

$$\varphi_{c,m}^* := \varphi_{F(c),m}: H(F(c) \triangleright m) \xrightarrow{\sim} F(c) \triangleright H(m).$$

A module functor involving twisted module categories is also called a *twisted module functor*.

*Example 4.1.* Consider the double-dual tensor functor  $\mathbb{D} := (-)^{\vee\vee}: \mathcal{D} \rightarrow \mathcal{D}$ . The relative Serre functor of a  $\mathcal{D}$ -module category  $\mathcal{M}$

$$\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}: \mathcal{M} \rightarrow {}_{\mathbb{D}}\mathcal{M} \tag{4.2}$$

becomes a twisted  $\mathcal{D}$ -module functor with constraints given by (2.10).

A monoidal natural transformation  $\alpha: F \Rightarrow G$  between tensor functors from  $\mathcal{C}$  to  $\mathcal{D}$  determines a  $\mathcal{C}$ -module equivalence  $\text{id}^\alpha: {}_F\mathcal{M} \xrightarrow{\sim} {}_G\mathcal{M}$ . This is defined by the assignment  $\text{id}^\alpha(m) = m$  together with the  $\mathcal{C}$ -module structure

$$\alpha_c \triangleright \text{id}_m = (\text{id}^\alpha)_{c,m}: \text{id}^\alpha(c \triangleright^F m) = F(c) \triangleright m \rightarrow G(c) \triangleright m = c \triangleright^G \text{id}^\alpha(m),$$

for  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$ . Notice that since  $\mathcal{C}$  is rigid, every such  $\alpha$  is necessarily invertible.

*Example 4.2.* Given a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  and a  $\mathcal{D}$ -module category  $\mathcal{M}$ , we can use the monoidal natural isomorphism  $\xi_{\mathcal{C}}: F(c^{\vee\vee}) \rightarrow F(c)^{\vee\vee}$  [NS07, Lem. 1.1] to get an equivalence between the  $\mathcal{C}$ -module categories  ${}_F(\mathbb{D}\mathcal{M})$  and  ${}_{\mathbb{D}}({}_F\mathcal{M})$ . In the following, we will often use this identification.

The  $\mathcal{C}$ -module category  ${}_F\mathcal{D}$  captures the procedure of twisting actions along  $F$ . For any  $\mathcal{D}$ -module category  $\mathcal{M}$  the assignment

$${}_F\mathcal{D} \boxtimes_{\mathcal{D}} \mathcal{M} \xrightarrow{\sim} {}_F\mathcal{M}, \quad d \boxtimes m \mapsto d \triangleright m \tag{4.3}$$

is an equivalence of  $\mathcal{C}$ -module categories.

The internal Hom of a  $F$ -twisted module category  ${}_F\mathcal{M}$  can be described in terms of the internal Hom of the original module category  $\mathcal{M}$  by the formula

$$\underline{\mathrm{Hom}}_{{}_F\mathcal{M}}^{\mathcal{C}}(m, n) \cong F^{\mathrm{ra}}(\underline{\mathrm{Hom}}_{\mathcal{M}}^{\mathcal{D}}(m, n)). \tag{4.4}$$

This description will be useful to investigate the exactness of twisted module categories.

*Remark 4.3.* If  ${}_F\mathcal{M}$  is an indecomposable  $\mathcal{C}$ -module category, then  $\mathcal{M}$  is indecomposable as a  $\mathcal{D}$ -module category. Indeed, assume that we have a non-trivial  $\mathcal{D}$ -module category decomposition,

$$\mathcal{M} \simeq \bigoplus_i \mathcal{M}_i$$

then by twisting the components of  $\mathcal{M}$ , we obtain a non-trivial decomposition

$${}_F\mathcal{M} \simeq \bigoplus_i {}_F\mathcal{M}_i$$

as  $\mathcal{C}$ -module categories. The converse does not hold. Consider a non-trivial finite group  $G$  and the unique tensor functor  $\iota: \mathbf{Vec} \rightarrow \mathbf{Vec}_G$ . The regular module category  $\mathbf{Vec}_G$  is an indecomposable  $\mathbf{Vec}_G$ -module category, but  ${}_{\iota}\mathbf{Vec}_G = \mathbf{Vec}_G$  is decomposable as a  $\mathbb{k}$ -linear category.

### 4.1. Exactness of module categories

Tensor categories are regarded as the categorification of rings, with exact module categories playing the role of projective modules. A classical result concerning the preservation of projectivity under the pullback of ring actions via homomorphisms states: let  $f: A \rightarrow B$  be a ring homomorphism. If  $B$  is projective as an  $A$ -module then, a  $B$ -module  $M$  is projective if and only if  $M$  is projective as an  $A$ -module. This result finds an analog in the categorical setting of twisting actions along tensor functors  $F: \mathcal{C} \rightarrow \mathcal{D}$ .

**Proposition 4.4.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. A  $\mathcal{D}$ -module category  $\mathcal{M}$  is exact if and only if the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is exact.*

*Proof.* Since,  $F$  is perfect,  $F(p) \in \mathcal{D}$  is projective for any projective  $p \in \mathcal{C}$ . If  $\mathcal{M}$  is exact as  $\mathcal{D}$ -module, then for every  $m \in \mathcal{M}$ , we have that  $p \triangleright^F m = F(p) \triangleright m$  is projective in  $\mathcal{M}$ , and thus  ${}_F\mathcal{M}$  is an exact  $\mathcal{C}$ -module. For the converse, pick a non-zero projective object  $p \in \mathcal{C}$ . Since  $F$  is faithful, it reflects zero objects and thus  $F(p)$  is non-zero [Shi16, Lemma 5.1]. Assuming that  ${}_F\mathcal{M}$  is an exact  $\mathcal{C}$ -module,  $F(p) \triangleright m$  is projective for all  $m \in \mathcal{M}$ . The claim follows by [EG17, Lemma 2.4].  $\square$

This means that twisting by perfect functors preserves exactness of module categories. As we see next, the converse also holds, i.e. if a tensor functor preserves exactness, it must be perfect.

**Proposition 4.5.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a tensor functor and  $\mathcal{M}$  an exact  $\mathcal{D}$ -module category. The  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is exact if and only if the tensor functor  $F$  is perfect.*

*Proof.* First, notice that  ${}_F\mathcal{D}$  is an exact  $\mathcal{C}$ -module category if and only if  $F$  is perfect. Indeed, if  $F$  is perfect, it preserves projective objects according to Lemma 3.2, and then  ${}_F\mathcal{D}$  is exact. On the other hand, exactness of  ${}_F\mathcal{D}$  implies that  $F^{\text{ra}} \cong \underline{\text{Hom}}_{{}_F\mathcal{D}}(\mathbb{1}, -)$  is exact, and therefore  $F$  admits a double right adjoint, which means that  $F$  is perfect.

Now, for the case of an arbitrary exact  $\mathcal{D}$ -module category  $\mathcal{M}$ , again preservation of projectives by  $F$  implies that  ${}_F\mathcal{M}$  is exact. If  ${}_F\mathcal{M}$  is exact, then it becomes an exact  $(\mathcal{C}, \overline{\mathcal{D}}_{\mathcal{M}}^*)$ -bimodule. Since  $\overline{\mathcal{M}}$  is an exact left  $\overline{\mathcal{D}}_{\mathcal{M}}^*$ -module, Lemma 2.1(ii) implies that

$${}_F\mathcal{M} \boxtimes_{\overline{\mathcal{D}}_{\mathcal{M}}^*} \overline{\mathcal{M}} \stackrel{(4.3)}{\simeq} {}_F\mathcal{D} \boxtimes_{\mathcal{D}} \mathcal{M} \boxtimes_{\overline{\mathcal{D}}_{\mathcal{M}}^*} \overline{\mathcal{M}} \stackrel{(2.5)}{\simeq} {}_F\mathcal{D} \boxtimes_{\mathcal{D}} \mathcal{D} \simeq {}_F\mathcal{D}$$

is an exact  $\mathcal{C}$ -module category, and thus  $F$  is perfect.  $\square$

**Corollary 4.6.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a tensor functor between finite tensor categories. The lax monoidal functor  $F^{\text{ra}}: \mathcal{D} \rightarrow \mathcal{C}$  preserves exactness of algebras if and only if  $F$  is perfect.*

*Proof.* Given an exact algebra  $A \in \mathcal{D}$ , consider the module category  $\mathcal{M} = \text{mod}_A(\mathcal{D})$ . Since,  ${}_F\mathcal{M} \simeq \text{mod}_{F^{\text{ra}}(A)}(\mathcal{C})$ , Proposition 4.5 implies that  $F^{\text{ra}}(A)$  is exact if and only if  $F$  is perfect.  $\square$

*Remark 4.7.* The corestriction of a tensor functor  $F$  to its image is surjective and therefore perfect. Thus, by Corollary 4.6,  $F^{\text{ra}}(A)$  is exact in  $\mathcal{C}$  for any exact algebra  $A \in \text{im}(F)$ . For instance, since  $\mathbb{1}_{\mathcal{D}} \in \text{im}(F)$  is an exact algebra, we have that  $F^{\text{ra}}(\mathbb{1}_{\mathcal{D}})$  is an exact algebra in  $\mathcal{C}$ .

For a finite tensor category  $\mathcal{C}$ , let  $\mathbf{Mod}^{\text{ex}}(\mathcal{C})$  denote the 2-category with exact left  $\mathcal{C}$ -module categories as objects,  $\mathcal{C}$ -module functors as 1-morphisms and module natural transformations as 2-morphisms.

**Corollary 4.8.** *A perfect tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  between finite tensor categories, induces a pseudo-functor*

$$\mathbf{F}^* : \mathbf{Mod}^{\text{ex}}(\mathcal{D}) \rightarrow \mathbf{Mod}^{\text{ex}}(\mathcal{C}), \quad \mathcal{M} \mapsto {}_F\mathcal{M}, \quad H \mapsto F^*(H) \quad (4.5)$$

*between their 2-categories of exact module categories. The pseudo-functor (4.5) is pseudo-naturally equivalent to the pseudo-functor  ${}_F\mathcal{D} \boxtimes_{\mathcal{D}} -: \mathbf{Mod}^{\text{ex}}(\mathcal{D}) \rightarrow \mathbf{Mod}^{\text{ex}}(\mathcal{C})$ .*

*Proof.* By Proposition 4.5,  $F$ -twisted module categories are exact. A routine check shows that  $F^*(H_2 \circ H_1) = F^*(H_2) \circ F^*(H_1)$ . The equivalences (4.3) assemble into the desired pseudo-natural equivalence in the second part of the statement.  $\square$

*Remark 4.9.* Let  $\mathcal{C}$  and  $\mathcal{D}$  be finite tensor categories.

- (a) Pairs  $(\mathcal{M}, F)$  consisting of an exact  $\mathcal{D}$ -module category  $\mathcal{M}$  and a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  such that  ${}_F\mathcal{M}$  is exact were first studied in [EO04, §3.5] under the name of *exact pair*. Proposition 4.5 states that  $(F, \mathcal{M})$  is an exact pair if and only if  $F$  is perfect.
- (b) For any exact  $\mathcal{D}$ -module category  $\mathcal{M}$ , the assignment (4.1) extends to a tensor functor

$$F^* : \mathcal{D}_{\mathcal{M}}^* \rightarrow \mathcal{C}_{F\mathcal{M}}^*, \quad (4.6)$$

called the *dual* of  $F$  with respect to  $\mathcal{M}$ . If  $F$  is perfect, then the dual  $F^*$  is perfect: indeed,  ${}_F\mathcal{M}$  is exact as a  $\mathcal{C}$ -module category by Proposition 4.5. Moreover,  ${}_F\mathcal{M}$  is exact as a  $\mathcal{C}_{F^*\mathcal{M}}^*$ -module category according to [EGNO16, Lemma 7.12.7]. Now, we have that  ${}_{F^*}({}_F\mathcal{M}) = \mathcal{M}$  as a left  $\mathcal{D}_{\mathcal{M}}^*$ -module category, which is exact again by [EGNO16, Lemma 7.12.7]. Thus, Proposition 4.5 implies that  $F^*$  is perfect.

## 4.2. Action of central objects and relative Serre functors

Given tensor functors  $F, G: \mathcal{C} \rightarrow \mathcal{D}$  and  $\mathcal{M}$  a  $\mathcal{D}$ -module category, there is a  $\mathbb{k}$ -linear functor

$$\mathcal{Z}({}_F\mathcal{D}_G) \rightarrow \text{Func}({}_G\mathcal{M}, {}_F\mathcal{M}), \quad (d, \gamma) \mapsto (d \triangleright -, \varphi) \quad (4.7)$$

where the module functor structure  $\varphi$  on  $d \triangleright -$  is defined by the composition

$$d \triangleright (c \triangleright^G m) = d \otimes G(c) \triangleright m \xrightarrow{\gamma} F(c) \otimes d \triangleright m = c \triangleright^F (d \triangleright m).$$

For an object  $(d, \gamma) \in \mathcal{Z}({}_F\mathcal{D}_G)$  and a  $\mathcal{C}$ -module functor  $H: \mathcal{N} \rightarrow {}_G\mathcal{M}$ , we use the notation  $d \triangleright H$  to denote the composite  $\mathcal{C}$ -module functor from  $\mathcal{N} \rightarrow {}_G\mathcal{M} \rightarrow {}_F\mathcal{M}$ .

*Example 4.10.* Let  $\mathcal{C}$  be a finite tensor category and let  $\mathbb{D} = (-)^{\vee\vee}$  and  $\overline{\mathbb{D}} = {}^{\vee\vee}(-)$  denote the double dual functors. The Radford isomorphism (2.8) together with the distinguished invertible object  $D_{\mathcal{C}}$  forms an object in  $\mathcal{Z}({}_{\mathbb{D}}\mathcal{C}_{\overline{\mathbb{D}}})$ . Therefore, for any  $\mathcal{C}$ -module category  $\mathcal{M}$ , there is a  $\mathcal{C}$ -module functor  $D_{\mathcal{C}} \triangleright -: {}_{\mathbb{D}}\mathcal{M} \rightarrow {}_{\mathbb{D}}\mathcal{M}$ .

Consider a perfect tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  and a  $\mathcal{D}$ -module category  $\mathcal{M}$ . By Proposition 3.13, we have that the relative modular object  $\chi_{\mathcal{Z}(F)} = (\chi_F, \sigma)$  is an object in  $\mathcal{Z}({}_F\mathcal{D}_F)$  and thus its action becomes a  $\mathcal{C}$ -module endofunctor on  ${}_F\mathcal{M}$ :

$$\chi_F \triangleright -: {}_F\mathcal{M} \rightarrow {}_F\mathcal{M} \quad (4.8)$$

with  $\mathcal{C}$ -module functor structure given by

$$\chi_F \triangleright (c \triangleright^F m) = \chi_F \otimes F(c) \triangleright m \xrightarrow{\sigma_c \triangleright \text{id}} F(c) \otimes \chi_F \triangleright m = c \triangleright^F (\chi_F \triangleright m)$$

where  $\sigma_c: \chi_F \otimes F(c) \xrightarrow{\sim} F(c) \otimes \chi_F$  is the  $F$ -half-braiding of  $\chi_F$ . Trivializing such module functors provides a characterization of  $\otimes$ -Frobenius.

**Lemma 4.11.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor. The following are equivalent:*

- (i)  $F$  is  $\otimes$ -Frobenius.
- (ii) The  $\mathcal{C}$ -module functor  $\chi_F \otimes -: {}_F\mathcal{D} \rightarrow {}_F\mathcal{D}$  is isomorphic to  $\text{id}_{{}_F\mathcal{D}}$ .
- (iii) For every  $\mathcal{D}$ -module  $\mathcal{M}$ , the  $\mathcal{C}$ -module functor  $\chi_F \triangleright -: {}_F\mathcal{M} \rightarrow {}_F\mathcal{M}$  is isomorphic to  $\text{id}_{{}_F\mathcal{M}}$ .
- (iv) There exists a  $\mathcal{D}$ -module category  $\mathcal{M}$ , such that the  $\mathcal{C}$ -module functor  $\chi_F \triangleright -: {}_F\mathcal{M} \rightarrow {}_F\mathcal{M}$  is isomorphic to  $\text{id}_{{}_F\mathcal{M}}$ .

*Proof.* (i)  $\implies$  (ii) follows from the definition of  $\chi_F \otimes -$ . To prove the converse, assume  $\chi_F \otimes - \cong \text{id}_{{}_F\mathcal{D}}$ . It follows by applying at the unit object that  $\chi_F \cong \mathbb{1}_{\mathcal{D}}$ . Moreover,  $\sigma$  is trivial since it coincides with the  $\mathcal{C}$ -module structure of the identity functor  $\text{id}_{{}_F\mathcal{D}}$ , and thus, by Theorem 3.18,  $F$  is  $\otimes$ -Frobenius.

Now, given a  $\mathcal{D}$ -module category  $\mathcal{M}$ , the diagram of  $\mathcal{C}$ -module equivalences

$$\begin{array}{ccc}
 {}_F\mathcal{D} \boxtimes_{\mathcal{D}} \mathcal{M} & \xrightarrow{\chi_F \otimes - \boxtimes_{\mathcal{D}} \text{id}_{\mathcal{M}}} & {}_F\mathcal{D} \boxtimes_{\mathcal{D}} \mathcal{M} \\
 \downarrow (4.3) & & \downarrow (4.3) \\
 {}_F\mathcal{M} & \xrightarrow{\chi_F \triangleright -} & {}_F\mathcal{M}
 \end{array}$$

commutes up to natural isomorphism (given by the module associativity constraints  $(\chi_F \otimes d) \triangleright m \cong \chi_F \triangleright (d \triangleright m)$  of  $\mathcal{M}$ ). Thus, a trivialization of  $\chi_F \otimes -$  provides a trivialization of  $\chi_F \triangleright -$  and vice versa, thereby proving (ii)  $\iff$  (iii)  $\iff$  (iv).  $\square$

*Remark 4.12.* For any tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , Proposition 3.14 provides an isomorphism

$$(F(D_{\mathcal{C}}), F(\mathcal{R}_{\mathcal{C}})) \xrightarrow{\sim} (\chi_F \otimes D_{\mathcal{D}}, (\sigma \otimes \text{id}) \circ (\text{id} \otimes \mathcal{R}_{\mathcal{D}}))$$

of objects in  $\mathcal{Z}(\mathbb{D} \circ \mathbb{D} \mathcal{D}_F)$ . This data can be re-written as an isomorphism

$$(F(D_{\mathcal{C}}) \otimes D_{\mathcal{D}}^{-1}, (F(\mathcal{R}_{\mathcal{C}}) \otimes \text{id}) \circ (\text{id} \otimes \mathcal{R}_{\mathcal{D}}^{-1})) \xrightarrow{\sim} (\chi_F, \sigma) = \chi_{\mathcal{Z}(F)}$$

of objects in  $\mathcal{Z}({}_F\mathcal{D}_F)$ . Under the equivalence (4.7), we obtain an isomorphism

$$(F(D_{\mathcal{C}}) \triangleright -) \circ F^*(D_{\mathcal{D}}^{-1} \triangleright -) \cong \chi_F \triangleright - : {}_F\mathcal{M} \rightarrow {}_F\mathcal{M}. \quad (4.9)$$

of  $\mathcal{C}$ -module functors, for any  $\mathcal{D}$ -module category  $\mathcal{M}$ .

A consequence of Proposition 4.5 is that the  $F$ -twisted module category arising from an exact module category is exact and thus admits a relative Serre functor. In order to describe the relative Serre functor of  ${}_F\mathcal{M}$ , we first describe its Nakayama functor as a  $\mathcal{C}$ -module functor.

**Lemma 4.13.** *Given a tensor functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  and a  $\mathcal{D}$ -module category  $\mathcal{M}$ , we have that  $\mathbb{N}_{F\mathcal{M}} = F^*(\mathbb{N}_{\mathcal{M}})$  as twisted  $\mathcal{C}$ -module functors.*

*Proof.* By the definition of  $F^*$  we have equality as functors and on both sides the twisted module functor structure is given by the isomorphism  $\mathbf{n}_{F(c),m}$  from (2.9).  $\square$

The following proposition relates the relative Serre functor and the relative modular object of  $F$ .

**Proposition 4.14.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor and  $\mathcal{M}$  an exact  $\mathcal{D}$ -module category. There is a natural isomorphism*

$$\chi_F \triangleright F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \cong \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}} \quad (4.10)$$

of twisted  $\mathcal{C}$ -module functors, where  $\chi_F \triangleright -$  is the  $\mathcal{C}$ -module functor from (4.8).

*Proof.* On one hand we have an isomorphism of twisted  $\mathcal{C}$ -module functors from (2.12)

$$\mathbb{N}_{F\mathcal{M}} \cong D_{\mathcal{C}}^{-1} \triangleright^F \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}} \quad (4.11)$$

and on the other hand an isomorphism of twisted  $\mathcal{D}$ -module functors from (2.12)

$$D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}} \cong \mathbb{N}_{\mathcal{M}}. \quad (4.12)$$

Moreover,  $\mathbb{N}_{F\mathcal{M}} = F^*(\mathbb{N}_{\mathcal{M}})$ , by Lemma 4.13. Thus, by composing (4.11) with the image of (4.12) under  $F^*$  we obtain an isomorphism of twisted  $\mathcal{C}$ -module functors

$$F^*(D_{\mathcal{D}}^{-1} \triangleright -) \circ F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) = F^*(D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \cong D_{\mathcal{C}}^{-1} \triangleright^F \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}} = F(D_{\mathcal{C}}^{-1}) \triangleright \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}}$$

where the first equality is the functoriality of  $F^*$  and the last one is the definition of the  $F$ -twisted action. Post-composing both sides with the module functor  $F(D_C) \triangleright -$  and using the isomorphism (4.9) we obtain the desired result.  $\square$

We spell out some details that are implicit in the above result. Since the relative Serre functor is unique up to unique isomorphism, from here on, we can fix  $\mathbb{S}_{F\mathcal{M}}^c = \chi_F \triangleright F^*(\mathbb{S}_{\mathcal{M}}^D)$ . The twisted left  $\mathcal{C}$ -module structure of  $\mathbb{S}_{F\mathcal{M}}^c$  is given by the composition

$$\mathfrak{s}_{c,m}^F : \chi_F \triangleright \mathbb{S}_{\mathcal{M}}^D(F(c) \triangleright m) \xrightarrow{\text{id} \triangleright \mathfrak{s}_{F(c),m}} \chi_F \triangleright (F(c)^{\vee\vee} \triangleright \mathbb{S}_{\mathcal{M}}^D(m)) \xrightarrow{\sigma_{c^{\vee\vee}} \triangleright \text{id}} (F(c^{\vee\vee}) \otimes \chi_F) \triangleright \mathbb{S}_{\mathcal{M}}^D(m)$$

### 4.3. Unimodular structures

**Definition 4.15.** [Yad23, Def. 3.2] Let  $\mathcal{C}$  be a finite tensor category, and  $\mathcal{M}$  an exact  $\mathcal{C}$ -module category. A *unimodular structure* on  $\mathcal{M}$  consists of a module natural isomorphism

$$\tilde{\mathfrak{u}} : \text{id}_{\mathcal{M}} \xrightarrow{\cong} D_{\mathcal{C}\mathcal{M}}^*$$

A module category endowed with a unimodular structure is called a *unimodular module category*. We also say that  $\mathcal{M}$  is *unimodular* iff  $\mathcal{M}$  admits a unimodular structure.

*Remark 4.16.* Notice that for a module category  ${}_c\mathcal{M}$  to be unimodular  $\mathcal{C}$  is not required to be unimodular. The dual tensor category  $\mathcal{C}_{\mathcal{M}}^*$  must be unimodular instead [Yad23, Rem. 3.7].

As we have seen, the relative modular object of a perfect tensor functor measures its failure of preserving distinguished invertible objects and, in particular, its failure of preserving unimodularity. It turns out, that the relative modular object also relates the distinguished invertible objects of the dual tensor categories with respect to an exact module category.

**Proposition 4.17.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor and  $\mathcal{M}$  an exact  $\mathcal{D}$ -module category. There is a natural isomorphism of  $\mathcal{C}$ -module functors*

$$\chi_F \triangleright F^*(D_{\mathcal{D}\mathcal{M}}^{-1}) \cong D_{\mathcal{C}\mathcal{M}}^{-1} \tag{4.13}$$

relating the distinguished invertible objects of the dual categories of  $\mathcal{M}$  and  ${}_F\mathcal{M}$  and the  $\mathcal{C}$ -module functor  $\chi_F \triangleright -$  defined in (4.8).

*Proof.* The desired natural isomorphism arises as the composition

$$\begin{aligned} D_{\mathcal{C}\mathcal{M}}^{-1} &\cong D_{\mathcal{C}}^{-1} \triangleright^F \mathbb{S}_{F\mathcal{M}}^c \circ \mathbb{S}_{F\mathcal{M}}^c = F(D_{\mathcal{C}}^{-1}) \triangleright \mathbb{S}_{F\mathcal{M}}^c \circ \mathbb{S}_{F\mathcal{M}}^c \\ &\cong \chi_F \triangleright F^*(D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^D \circ \mathbb{S}_{\mathcal{M}}^D) \cong \chi_F \triangleright F^*(D_{\mathcal{D}\mathcal{M}}^{-1}) \end{aligned}$$

where the first isomorphism comes from (2.13), next is the definition of the  $F$ -twisted action, the second to last isomorphism is (4.10) and lastly we use isomorphism (2.13) once more.  $\square$

*Remark 4.18.* Because of Proposition 3.4(v), the isomorphism (4.13) gives a description of the relative modular object of the tensor functor (4.6). Explicitly, we have an isomorphism

$$\chi_{F^*} \cong \chi_F \triangleright - : {}_F\mathcal{M} \rightarrow {}_F\mathcal{M}. \tag{4.14}$$

of  $\mathcal{C}$ -module functors.

**Corollary 4.19.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor and  $\mathcal{M}$  an exact  $\mathcal{D}$ -module category. The tensor functor*

$$F^* : \mathcal{D}_{\mathcal{M}}^* \longrightarrow \mathcal{C}_{F\mathcal{M}}^*$$

*is Frobenius if and only if  $F$  is  $\otimes$ -Frobenius.*

*Proof.* Lemma 4.11 and Proposition 3.9 provide characterizations of the properties  $\otimes$ -Frobenius and Frobenius in terms of relative modular objects. The result follows from isomorphism (4.14).  $\square$

Twisting module actions by perfect tensor functors that are  $\otimes$ -Frobenius allows for the transfer of unimodular structures on module categories, as we see next.

**Proposition 4.20.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor and  $\mathcal{M}$  a unimodular  $\mathcal{D}$ -module category. Assume that  $F$  is  $\otimes$ -Frobenius, then the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is endowed with a unimodular structure via*

$${}_F\tilde{\mathbf{u}} : \text{id}_{\mathcal{M}} = F^*(\text{id}_{\mathcal{M}}) \xrightarrow{F^*(\tilde{\mathbf{u}})} F^*(D_{\mathcal{D}_{\mathcal{M}}^*}) \cong D_{\mathcal{C}_{F\mathcal{M}}^*}.$$

*Moreover,  $F$  is  $\otimes$ -Frobenius if and only if the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is unimodular.*

*Proof.* By Lemma 4.11,  $F$  is  $\otimes$ -Frobenius if and only if  $\chi_F \triangleright -$  is trivial. The result follows from Proposition 4.17.  $\square$

*Remark 4.21.* Proposition 4.20 recovers the result in [FLP24b, Thm. 4.21] as a special case, where an inclusion of Hopf algebras with unimodular duals is considered. See Remark 5.7 for more details.

*Remark 4.22.* We call a tensor category  $\mathcal{C}$  *weakly unimodular* iff there exists a unimodular tensor category in the Morita class of  $\mathcal{C}$ , or equivalently if  $\mathcal{C}$  admits a unimodular module category. Proposition 4.20 shows that there are no  $\otimes$ -Frobenius functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  from a tensor category  $\mathcal{C}$  that is not weakly unimodular into a tensor category  $\mathcal{D}$  that is weakly unimodular. For instance, take  $\mathcal{C}$  to be the tensor category of representations of the Taft Hopf algebra. In [Yad23], it is proven that  $\mathcal{C}$  does not admit unimodular module categories. Thus, we can conclude that  $\mathcal{C}$  does not admit a  $\otimes$ -Frobenius functor into any weakly unimodular category.

#### 4.4. Pivotal structures and sphericity

Let  $\mathcal{C}$  be a pivotal finite tensor category and  $\mathcal{M}$  an exact left  $\mathcal{C}$ -module category. The pivotal structure  $\mathfrak{p}: \text{id}_{\mathcal{C}} \xrightarrow{\sim} (-)^{\vee\vee}$  turns the relative Serre functor of  $\mathcal{M}$  into a  $\mathcal{C}$ -module functor via

$$\mathbb{S}_{\mathcal{M}}^{\mathcal{C}}(c \triangleright m) \xrightarrow{(2.10)} c^{\vee\vee} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{C}}(m) \xrightarrow{\mathfrak{p}_c \triangleright \text{id}} c \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{C}}(m)$$

for  $c \in \mathcal{C}$  and  $m \in \mathcal{M}$ .

**Definition 4.23.** [Shi17, Def. 3.11] A *pivotal structure* on an exact left  $\mathcal{C}$ -module category  $\mathcal{M}$  is a module natural isomorphism

$$\tilde{\mathfrak{p}} : \text{id}_{\mathcal{M}} \xrightarrow{\sim} \mathbb{S}_{\mathcal{M}}^{\mathcal{C}}.$$

A module category together with a pivotal structure is called a *pivotal module category*. A module category  $\mathcal{M}$  is *pivotal* in case there exists a pivotal structure for  $\mathcal{M}$ .

Just as a perfect functor  $F$  preserves exactness of module categories, requiring  $F$  to be  $\otimes$ -Frobenius ensures the transfer of pivotal structures on module categories.



**Proposition 4.24.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a pivotal tensor functor between pivotal tensor categories and  $\mathcal{M}$  a pivotal  $\mathcal{D}$ -module category. In case  $F$  is  $\otimes$ -Frobenius, the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is endowed with a pivotal structure via*

$${}_F\tilde{\mathfrak{p}}: \text{id}_{\mathcal{M}} = F^*(\text{id}_{\mathcal{M}}) \xrightarrow{F^*(\tilde{\mathfrak{p}})} F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \cong \mathbb{S}_{{}_F\mathcal{M}}^{\mathcal{C}}. \quad (4.15)$$

Furthermore,  $F$  is  $\otimes$ -Frobenius if and only if the  $\mathcal{C}$ -module category  ${}_F\mathcal{M}$  is pivotal.

*Proof.* By Proposition 4.14, the pivotal structures of  $\mathcal{M}$  and  ${}_F\mathcal{M}$  furnish a trivialization of  $\chi_F \triangleright -$ . Lemma 4.11 ensures that  $F$  is  $\otimes$ -Frobenius. The converse follows similarly.  $\square$

*Remark 4.25.* In the setting of Proposition 4.24, the tensor functor  $F^*: \mathcal{D}_{\mathcal{M}}^* \rightarrow \mathcal{C}_{{}_F\mathcal{M}}^*$  is pivotal. The 2-categorical extension of this fact is proven in Proposition 4.30.

**Theorem 4.26.** *Let  $\mathcal{C}$  and  $\mathcal{D}$  be pivotal tensor categories and  $F: \mathcal{C} \rightarrow \mathcal{D}$  a pivotal tensor functor.*

- (i) *Assume that  $F$  is  $\otimes$ -Frobenius, then the lax monoidal functor  $F^{\text{ra}}: \mathcal{D} \rightarrow \mathcal{C}$  preserves exact symmetric Frobenius algebras.*
- (ii) *If  $\mathcal{C}$  is endowed with a braiding,  $F$  has the structure of a central functor and  $F$  is Frobenius, then  $F^{\text{ra}}(\mathbb{1})$  is a haploid commutative exact symmetric Frobenius algebra.*

*Proof.* Given an exact algebra  $A \in \mathcal{D}$ , pivotal structures on the module category  $\mathcal{M} = \text{mod}_A(\mathcal{D})$  correspond to symmetric Frobenius structures on  $A$  [Shi24, Thm. 6.4]. From Corollary 4.6 we know that  $F^{\text{ra}}$  preserves exact algebras, thus  $F^{\text{ra}}(A)$  is exact. Moreover, since  $F$  is  $\otimes$ -Frobenius, Proposition 4.24 implies that  ${}_F\mathcal{M} = \text{mod}_{F^{\text{ra}}(A)}(\mathcal{C})$  inherits a pivotal structure, which in turn induces a symmetric Frobenius structure on  $F^{\text{ra}}(A)$ .

For assertion (ii), observe first that by Proposition 3.20, the fact that  $F$  is Frobenius implies it is indeed  $\otimes$ -Frobenius. Consequently, the claim follows from part (i) and [EGNO16, Prop. 8.8.8].  $\square$

*Remark 4.27.* A non-degenerate ribbon finite tensor category is called a modular tensor category (MTC). By [SY24, Thm. 5.21(d)], given a MTC  $\mathcal{C}$  and a haploid commutative exact symmetric Frobenius algebra  $A$  in  $\mathcal{C}$ , then the category of local modules  $\mathcal{C}_A^{\text{loc}}$  is also a MTC. Thus, the above result shows that central tensor functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  that are Frobenius and pivotal can be used to obtain new MTCs. Let  $A = F^{\text{ra}}(\mathbb{1}_{\mathcal{D}})$  and denote by  $G: \mathcal{C} \rightarrow \text{im}(F) \cong \text{mod}_A(\mathcal{C})$  the restriction of  $F$  to its image. Then,  $G$  is also central and we can describe the category  $\mathcal{C}_A^{\text{loc}}$  as the Müger centralizer  $\mathcal{Z}_{(2)}(\mathcal{C} \subset \mathcal{Z}(\text{im}(F)))$ .

*Remark 4.28.* In [Shi24], the notions of relative Serre functors and pivotal structures are extended to module categories that are not necessarily exact. We expect Propositions 4.17 and 4.24 to generalize to this setting. Moreover, by [Shi24, Thm. 6.4], symmetric Frobenius algebra structures on a (not necessarily exact) algebra  $A$  in  $\mathcal{C}$  are in correspondence with pivotal structures on the  $\mathcal{C}$ -module category  $\text{mod}_A(\mathcal{C})$ . Using this, Theorem 4.26 can be generalized to non-exact algebras.

The previous results have a 2-categorical description. In order to fully reformulate them, we need some additional structure: given a 2-category  $\mathcal{B}$ , the existence of dualities for 1-morphisms extends to a pseudo-functor

$$\begin{aligned} (-)^{\vee}: \mathcal{B} &\longrightarrow \mathcal{B}^{\text{op,op}}, \\ x &\longmapsto x, \\ (a: x \rightarrow y) &\longmapsto (a^{\vee}: y \rightarrow x). \end{aligned}$$

A *pivotal structure* on a 2-category  $\mathcal{B}$  with dualities is a pseudo-natural equivalence

$$\mathbf{P} : \text{id}_{\mathcal{B}} \xrightarrow{\sim} (-)^{\vee\vee}. \quad (4.16)$$

A 2-category together with a pivotal structure is called a *pivotal 2-category*.

**Definition 4.29.** Let  $\mathcal{B}_1$  and  $\mathcal{B}_2$  be pivotal 2-categories. A *pivotal pseudo-functor* consists of a pseudo-functor  $\mathcal{F} : \mathcal{B}_1 \rightarrow \mathcal{B}_2$  admitting an invertible modification filling the diagram:

$$\begin{array}{ccc} & & \mathcal{F} \circ (-)^{\vee\vee} \\ & \nearrow \text{id}_{\mathcal{F}} \circ \mathbf{P}^{\mathcal{B}_1} & \parallel \sim \\ \mathcal{F} & & \\ & \searrow \mathbf{P}^{\mathcal{B}_2} \circ \text{id}_{\mathcal{F}} & \downarrow \sim \\ & & (-)^{\vee\vee} \circ \mathcal{F} \end{array} \quad (4.17)$$

Denote by  $\mathbf{Mod}^{\text{piv}}(\mathcal{C})$  the 2-category that has pivotal  $\mathcal{C}$ -module categories as objects,  $\mathcal{C}$ -module functors as 1-morphisms and module natural transformations as 2-morphisms. Pivotal modules are exact and therefore every module functor between pivotal modules have adjoints. These turn  $\mathbf{Mod}^{\text{piv}}(\mathcal{C})$  into a 2-category with dualities for 1-morphisms. Moreover,  $\mathbf{Mod}^{\text{piv}}(\mathcal{C})$  is endowed with a pivotal structure (4.16) with components given for any object  $\mathcal{M}$  by  $\mathbf{P}_{\mathcal{M}} := \text{id}_{\mathcal{M}}$  and for a 1-morphism  $H : {}_c\mathcal{N}_1 \rightarrow {}_c\mathcal{N}_2$  by

$$\mathbf{P}_H : H \xrightarrow{\text{id} \circ \tilde{\mathbf{p}}_1} H \circ \mathbb{S}_{\mathcal{N}_1}^c \xrightarrow{(2.11)} \mathbb{S}_{\mathcal{N}_2}^c \circ H^{\text{lla}} \xrightarrow{(\tilde{\mathbf{p}}_2)^{-1} \text{oid}} H^{\text{lla}},$$

where  $\tilde{\mathbf{p}}_i$  are the pivotal structures of the module categories  $\mathcal{N}_i$ .

**Proposition 4.30.** A pivotal tensor functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between pivotal finite tensor categories, that is  $\otimes$ -Frobenius, induces a pivotal pseudo-functor

$$F^* : \mathbf{Mod}^{\text{piv}}(\mathcal{D}) \rightarrow \mathbf{Mod}^{\text{piv}}(\mathcal{C}), \quad (\mathcal{M}, \tilde{\mathbf{p}}) \mapsto ({}_F\mathcal{M}, {}_F\tilde{\mathbf{p}}), \quad H \mapsto F^*(H) \quad (4.18)$$

between their pivotal 2-categories of pivotal module categories, where  ${}_F\tilde{\mathbf{p}}$  is given by (4.15).

*Proof.* From (4.5) we have that  $F$ -twisting of module categories extends to a pseudo-functor. By Proposition 4.24 pivotal structures on  $\mathcal{D}$ -module categories induce pivotal structures on the associated  $F$ -twisted  $\mathcal{C}$ -module categories, and thus the assignment in (4.18) is well-defined. It remains to be checked that the pseudo-functor  $F^*$  is pivotal. First, on objects we have that  $\mathbf{P}_{\mathcal{M}}^{\mathcal{B}_1} = \text{id}_{\mathcal{M}} = \mathbf{P}_{\mathcal{M}}^{\mathcal{B}_2}$ . Now, given a module functor  $H : {}_d\mathcal{N}_1 \rightarrow {}_d\mathcal{N}_2$  the diagram

$$\begin{array}{ccccc} & & F^*(H) \circ F^*(\mathbb{S}_{\mathcal{N}_1}^d) & \xrightarrow{(2.11)} & F^*(\mathbb{S}_{\mathcal{N}_2}^d) \circ F^*(H^{\text{lla}}) \\ & \nearrow F^*(\text{id} \circ \tilde{\mathbf{p}}_1) & \parallel (4.10) & & \searrow F^*(\tilde{\mathbf{p}}_2^{-1} \text{oid}) \\ F^*(H) & & & & F^*(H^{\text{lla}}) \\ & \searrow \text{id} \circ {}_F\tilde{\mathbf{p}}_1 & \parallel (4.10) & & \nearrow {}_F\tilde{\mathbf{p}}_2^{-1} \text{oid} \\ & & F^*(H) \circ \mathbb{S}_{{}_F\mathcal{N}_1}^c & \xrightarrow{(2.11)} & \mathbb{S}_{{}_F\mathcal{N}_2}^c \circ F^*(H^{\text{lla}}) \end{array}$$

commutes by the definition of the transported pivotal structures on  ${}_F\mathcal{N}_1$  and  ${}_F\mathcal{N}_2$  and the fact that (4.10) is an isomorphism of twisted module functors. It follows that the diagram (4.17) commutes on the nose (i.e. the identity modification is a filler), and thus (4.18) is pivotal.  $\square$

We now turn to investigate the preservation of sphericity. Given a unimodular finite tensor category  $\mathcal{C}$ , a pivotal structure  $\mathbf{p}: \text{id}_{\mathcal{C}} \xrightarrow{\sim} (-)^{\vee\vee}$  on  $\mathcal{C}$  is called *spherical* iff  $\mathbf{p}^{\vee\vee} \circ \mathbf{p} = \mathcal{R}_{\mathcal{C}}$ . In analogy, the sphericity property for a pivotal module category can be defined in terms of the module Radford isomorphism (2.13).

**Definition 4.31.** [FGJS22, Def. 5.22]

Let  $\mathcal{M}$  be a pivotal module category over a (unimodular) spherical tensor category  $\mathcal{D}$ . Consider further an isomorphism  $\tilde{\mathbf{u}}_{\mathcal{D}}: \mathbb{1} \xrightarrow{\sim} D_{\mathcal{D}}^{-1}$  and a unimodular structure  $\tilde{\mathbf{u}}_{\mathcal{M}}: \text{id}_{\mathcal{M}} \xrightarrow{\sim} D_{\mathcal{D}\mathcal{M}}^{-1}$  on  $\mathcal{M}$ .

The pivotal module  ${}_{\mathcal{D}}\mathcal{M}$  is called  $(\tilde{\mathbf{u}}_{\mathcal{D}}, \tilde{\mathbf{u}}_{\mathcal{M}})$ -spherical iff the diagram

$$\begin{array}{ccc} \mathbb{S}_{\mathcal{M}}^{\mathcal{D}} & \xrightarrow{\overline{\mathcal{R}}_{\mathcal{M}}} & \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}} \\ & \swarrow \tilde{\mathbf{p}} & \searrow \tilde{\mathbf{p}} \\ & \text{id}_{\mathcal{M}} & \end{array}$$

commutes, where  $\tilde{\mathbf{p}}$  is the pivotal structure of  $\mathcal{M}$ , and  $\overline{\mathcal{R}}_{\mathcal{M}}$  is the composition

$$\mathbb{S}_{\mathcal{M}}^{\mathcal{D}} \xrightarrow{\tilde{\mathbf{u}}_{\mathcal{D}} \triangleright \text{id}} D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}} \xrightarrow{\mathcal{R}_{\mathcal{M}}} D_{\mathcal{D}\mathcal{M}}^{-1} \circ \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}} \xleftarrow{\tilde{\mathbf{u}}_{\mathcal{M}} \text{oid}} \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}}, \quad (4.19)$$

and  $\mathcal{R}_{\mathcal{M}}$  is the module Radford isomorphism (2.13).

The following technical lemma shows the extension to which the procedure of  $F$ -twisting module categories preserves the module Radford isomorphism.

**Lemma 4.32.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a perfect tensor functor and  $\mathcal{M}$  an exact  $\mathcal{D}$ -module category. Then the following diagram commutes.*

$$\begin{array}{ccc} D_{\mathcal{C}}^{-1} \triangleright^F \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}} & \xrightarrow{(3.3) \triangleright (4.10)} & F^*(D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \\ \mathcal{R}_{F\mathcal{M}} \Downarrow & & \Downarrow F^*(\mathcal{R}_{\mathcal{M}}) \\ D_{\mathcal{C}^*F\mathcal{M}}^{-1} \circ \overline{\mathbb{S}}_{F\mathcal{M}}^{\mathcal{C}} & \xrightarrow{(4.13) \circ (4.10)} & F^*(D_{\mathcal{D}^*F\mathcal{M}}^{-1} \circ \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \end{array}$$

*Proof.* The commutativity follows after noticing that we can rewrite the diagram as

$$\begin{array}{ccccc} F(D_{\mathcal{C}}^{-1}) \triangleright \mathbb{S}_{F\mathcal{M}}^{\mathcal{C}} & \xrightarrow{(4.10)} & F(D_{\mathcal{C}}^{-1}) \otimes \chi_F^{-1} \triangleright F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) & \xrightarrow{(3.3)} & F^*(D_{\mathcal{D}}^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \\ \mathcal{R}_{F\mathcal{M}} \Downarrow & \cong \searrow & \downarrow \cong & \cong \swarrow & \Downarrow F^*(\mathcal{R}_{\mathcal{M}}) \\ & & \mathbb{N}_{F\mathcal{M}} = F^*(\mathbb{N}_{\mathcal{M}}) & & \\ & \cong \swarrow & \downarrow \cong & \cong \searrow & \\ D_{\mathcal{C}^*F\mathcal{M}}^{-1} \circ \overline{\mathbb{S}}_{F\mathcal{M}}^{\mathcal{C}} & \xrightarrow{(4.10)} & D_{\mathcal{C}^*F\mathcal{M}}^{-1} \circ \chi_F^{-1} \triangleright F^*(\overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}}) & \xrightarrow{(4.13)} & F^*(D_{\mathcal{D}^*F\mathcal{M}}^{-1} \circ \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) \end{array} \quad (4.20)$$

where each of the triangles commutes due to the definition of the isomorphisms involved.  $\square$

**Proposition 4.33.** *Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a pivotal tensor functor between (unimodular) spherical tensor categories such that  $F$  is  $\otimes$ -Frobenius, and let  $\mathcal{M}$  be a pivotal  $\mathcal{D}$ -module. Given a trivialization  $\tilde{\mathbf{u}}_{\mathcal{C}}: \mathbb{1} \xrightarrow{\sim} D_{\mathcal{C}}$  and a unimodular structure  $\tilde{\mathbf{u}}_{\mathcal{M}}$  on  $\mathcal{M}$ , if  $\mathcal{M}$  is  $(F(\tilde{\mathbf{u}}_{\mathcal{C}}), \tilde{\mathbf{u}}_{\mathcal{M}})$ -spherical, then  ${}_{F}\mathcal{M}$  is  $(\tilde{\mathbf{u}}_{\mathcal{C}}, {}_{F}\tilde{\mathbf{u}}_{\mathcal{M}})$ -spherical.*

*Proof.* From Proposition 4.20 it follows that  $F^*(\tilde{\mathbf{u}}_{\mathcal{M}})$  is a unimodular structure on  ${}_F\mathcal{M}$ . The assertion holds due to the commutativity of the following diagram.

$$\begin{array}{ccccc}
\mathbb{S}_{F\mathcal{M}}^c & \xrightarrow{\tilde{\mathbf{u}}_c \triangleright \text{id}} & D_c^{-1} \triangleright^F \mathbb{S}_{F\mathcal{M}}^c & \xrightarrow{\mathcal{R}_{F\mathcal{M}}} & D_{C^*}^{-1} \circ \overline{\mathbb{S}}_{F\mathcal{M}}^c & \xleftarrow{F^*(\tilde{\mathbf{u}}_{\mathcal{M}})\text{oid}} & \overline{\mathbb{S}}_{F\mathcal{M}}^c \\
(4.10) \downarrow & & \downarrow (4.10) & & \downarrow (4.10) & & \downarrow (4.10) \\
F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) & \xrightarrow{\tilde{\mathbf{u}}_c \triangleright \text{id}} & F(D_c^{-1}) \triangleright F^*(\mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) & & D_{C^*}^{-1} \circ F^*(\overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}}) & \xleftarrow{F^*(\tilde{\mathbf{u}}_{\mathcal{M}})\text{oid}} & F^*(\overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}}) \\
\uparrow F^*(\tilde{\mathbf{p}}) & \searrow F^*(F(\tilde{\mathbf{u}}_c) \triangleright \text{id}) & \downarrow (3.3) & & \downarrow (4.13) & \swarrow F^*(\tilde{\mathbf{u}}_{\mathcal{M}})\text{oid} & \downarrow F^*(\tilde{\mathbf{p}}) \\
F^*(\tilde{\mathbf{p}}) & & F^*(D_D^{-1} \triangleright \mathbb{S}_{\mathcal{M}}^{\mathcal{D}}) & \xrightarrow{F^*(\mathcal{R}_{\mathcal{M}})} & F^*(D_{\mathcal{D}^*}^{-1} \circ \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{D}}) & & \\
\parallel & & & & & & \parallel \\
\text{id}_{\mathcal{M}} & \xlongequal{\hspace{10em}} & & & & & \text{id}_{\mathcal{M}}
\end{array}$$

where the upper left and right squares commute due to naturality of  $\triangleright$  and  $\circ$ , respectively. The middle diagram commutes according to Lemma 4.32. The bottom diagram is the condition that  $\mathcal{M}$  is  $(F(\tilde{\mathbf{u}}_c), \tilde{\mathbf{u}}_{\mathcal{M}})$ -spherical. Finally, the remaining triangles commute from the definitions of  $F(\tilde{\mathbf{u}}_c) : \mathbb{1} \xrightarrow{\sim} F(D_c) \cong D_D$  and  ${}_F\tilde{\mathbf{u}}_{\mathcal{M}}$ , respectively.  $\square$

## 5. APPLICATIONS TO HOPF ALGEBRAS

In this section, we illustrate the results from the previous section for tensor categories coming from Hopf algebras. Let  $(H, m, u, \Delta, \varepsilon, S)$  be a finite-dimensional Hopf algebra over  $\mathbb{k}$ . We will use the Sweedler notation  $\Delta(h) = h_1 \otimes h_2$  for the coproduct. Since  $H$  is finite-dimensional, the antipode is invertible and we denote its inverse as  $\overline{S}$ . We refer the reader to [Rad11] for basics of Hopf algebras.

Given a map  $\mathbf{f} : A \rightarrow B$  of  $\mathbb{k}$ -algebras and  $M$  a left  $B$ -module, we use the notation  $M_{\mathbf{f}}$  to denote the left  $A$ -module  $M$  with action defined by  $a \cdot_{\mathbf{f}} m := \mathbf{f}(a) \cdot m$  for  $m \in M$  and  $a \in A$ . Note that for algebra maps  $\mathbf{f} : A \rightarrow B$  and  $\mathbf{g} : B \rightarrow C$  and a  $C$ -module  $M$ , we have that  $(M_{\mathbf{g}})_{\mathbf{f}} = M_{\mathbf{g}\mathbf{f}}$  as left  $A$ -modules.

### 5.1. $\otimes$ -Frobenius functors from maps of Hopf algebras

In this section, we discuss when tensor functors arising from bialgebra maps between two Hopf algebras are  $\otimes$ -Frobenius and illustrate results obtained in §3 with examples.

#### 5.1.1. Preliminaries about the category $\text{Rep}(H)$

The category  $\text{Rep}(H)$  of finite dimensional representations of  $H$  is a finite tensor category. For  $X, Y \in \text{Rep}(H)$ , recall that  $X \otimes Y = X \otimes_{\mathbb{k}} Y$  and  $X^{\vee} = X^* := \text{Hom}(X, \mathbb{k})$  as  $\mathbb{k}$ -vector spaces. The  $H$ -action is given by

$$h \cdot (x \otimes_{\mathbb{k}} y) = h_1 \cdot x \otimes_{\mathbb{k}} h_2 \cdot y, \quad h \cdot \langle f, - \rangle = \langle f, S(h) - \rangle$$

for  $x \in X, y \in Y, f \in X^*$  and  $h \in H$ . The unit object is the one-dimensional  $H$ -representation  $\mathbb{k}_{\varepsilon}$ .

For  $X \in \text{Rep}(H)$ , using the canonical isomorphism  $\phi_X : X \rightarrow X^{\vee\vee}$  and thinking of elements in  $X^{**} = X^{\vee\vee}$  as  $\phi_X(x)$  for some  $x \in X$  we get that the  $H$ -action is given by  $h \cdot \phi_X(x) = \phi_X(S^2(h) \cdot x)$ . Thus,  $X^{\vee\vee} \cong X_{S^2}$ . In a similar manner,  $X^{\vee\vee\vee\vee} \cong X_{S^4}$ .

Let  $\Lambda \in H$  be a non-zero left integral of  $H$ , that is an element satisfying  $h\Lambda = \langle \varepsilon, h \rangle \Lambda$ . Then, the modular function  $\alpha : H \rightarrow \mathbb{k}$  of  $H$  is defined to be the unique algebra map satisfying  $\Lambda h = \langle \alpha, h \rangle \Lambda$  for all  $h \in H$ . Set  $\bar{\alpha} := \alpha \circ S$ , this is the inverse of  $\alpha$  in the algebra  $H^*$ . This matches the conventions used in prior work [Yad23] whose results we will use next. One can calculate that  $D_{\text{Rep}(H)} = \mathbb{N}_{\text{Rep}(H)}^l(\mathbb{k}_\varepsilon) = \mathbb{k}_{\bar{\alpha}}$  (see [Yad23, §4.2] for details). We call  $H$  *unimodular* if  $\alpha = \varepsilon$ . The modular object of  $\text{Rep}(H)$  is  $\alpha_{\text{Rep}(H)} := \mathbb{k}_\alpha$ . Clearly,  $D_{\text{Rep}(H)} = \alpha_{\text{Rep}(H)}^\vee$ .

A map  $\lambda : H \rightarrow \mathbb{k}$  is called a *right cointegral* of  $H$  if it satisfies  $\langle \lambda, h_1 \rangle h_2 = \langle \lambda, h \rangle 1_H$  for all  $h \in H$ . Such a cointegral always exists and from here on we fix a left integral  $\Lambda$  and a right cointegral  $\lambda$  satisfying  $\langle \lambda, \Lambda \rangle = 1$ . There exists a unique element  $g_H \in H$  satisfying

$$h_1 \langle \lambda, h_2 \rangle = \langle \lambda, h \rangle g_H. \quad (5.1)$$

We will use the notation  $\overline{g_H} = g_H^{-1}$ . Using this data, we can describe the Radford isomorphism (2.8)  $D_{\text{Rep}(H)} \otimes X \rightarrow X^{\vee\vee\vee\vee} \otimes D_{\text{Rep}(H)}$  of  $\text{Rep}(H)$  explicitly as follows

$$\mathbb{k}_{\bar{\alpha}} \otimes X \xrightarrow{\mathcal{R}_X} X_{S^4} \otimes \mathbb{k}_{\bar{\alpha}}, \quad 1 \otimes_{\mathbb{k}} x \mapsto g_H \cdot x \otimes_{\mathbb{k}} 1 \quad (5.2)$$

and can be derived using [Yad23, Prop. 4.23]. We will also need its variant corresponding to  $D^* \otimes X^{\vee\vee\vee\vee} \rightarrow X \otimes D^*$  given by:

$$\mathbb{k}_\alpha \otimes X_{S^4} \xrightarrow{\mathcal{R}'_X} X \otimes \mathbb{k}_\alpha, \quad 1 \otimes_{\mathbb{k}} x \mapsto \overline{g_H} \cdot x \otimes_{\mathbb{k}} 1. \quad (5.3)$$

A *pivotal element* of a Hopf algebra is a grouplike element  $g_{\text{piv}} \in H$  satisfying  $g_{\text{piv}} h g_{\text{piv}}^{-1} = S^2(h)$  for all  $h \in H$ . Pivotal structures on  $\text{Rep}(H)$  are in bijection with pivotal elements on  $H$ . Given a pivotal element  $g_{\text{piv}}$ , the pivotal structure  $\mathfrak{p} : X \rightarrow X^{\vee\vee} = X_{S^2}$  is given by  $x \mapsto g_{\text{piv}} \cdot x$ .

### 5.1.2. ⊗-Frobenius functors induced by bialgebra maps

Let  $\mathfrak{f} : H' \rightarrow H$  be a bialgebra map between two finite-dimensional Hopf algebras over  $\mathbb{k}$ . This means in particular that  $\mathfrak{f}$  commutes with the antipodes, that is,

$$\mathfrak{f} \circ S_{H'} = S_H \circ \mathfrak{f}. \quad (5.4)$$

Moreover,  $\mathfrak{f}$  induces a tensor functor  $F_{\mathfrak{f}} : \text{Rep}(H) \rightarrow \text{Rep}(H')$  defined via the assignment

$$X \mapsto X_{\mathfrak{f}}, \quad (f : X \rightarrow Y) \mapsto (f : X_{\mathfrak{f}} \rightarrow Y_{\mathfrak{f}})$$

called *the restriction of scalars* functor. We call  $\mathfrak{f}$  *perfect* iff the tensor functor  $F_{\mathfrak{f}}$  is perfect. Using Lemma 3.2, one can obtain alternate characterizations of perfect bialgebra maps of which we mention one:

**Lemma 5.1.** *A bialgebra map  $\mathfrak{f} : H' \rightarrow H$  is perfect if and only if  $H_{\mathfrak{f}}$  is a projective  $H'$ -module.*

*Remark 5.2.* As Hopf algebras are free (and therefore projective) over their Hopf subalgebras [NZ89],  $\mathfrak{f}$  is perfect if  $H'$  is a Hopf subalgebra of  $H$ , that is,  $\mathfrak{f}$  is injective.

Let  $\mathfrak{f} : H' \rightarrow H$  be a bialgebra map. Define the *relative modular function*  $\chi_{\mathfrak{f}} : H' \rightarrow \mathbb{k}$  by  $\chi_{\mathfrak{f}}(h') = \langle \alpha_H, \mathfrak{f}(h'_1) \rangle \langle \alpha_{H'}, S(h'_2) \rangle$ . The relative modular object of the perfect tensor functor  $F_{\mathfrak{f}}$  is determined by the regular modular function of  $\mathfrak{f}$  as we show next.

**Proposition 5.3.** *Let  $\mathbf{f} : H' \rightarrow H$  be a perfect bialgebra map.*

- (i) *The relative modular object of  $F_{\mathbf{f}}$  is the  $H'$ -representation  $\chi_{F_{\mathbf{f}}} \cong \mathbb{k}_{\chi_{\mathbf{f}}}$ . The  $F_{\mathbf{f}}$ -half-braiding of  $\chi_{F_{\mathbf{f}}}$  is given for  $X \in \text{Rep}(H)$  by the isomorphism*

$$\begin{aligned} \sigma_X : \quad \chi_{F_{\mathbf{f}}} \otimes F_{\mathbf{f}}(X) &\longrightarrow F_{\mathbf{f}}(X) \otimes \chi_{F_{\mathbf{f}}}, \\ 1 \otimes_{\mathbb{k}} x &\longmapsto \overline{g_H} \mathbf{f}(g_{H'}) \cdot x \otimes_{\mathbb{k}} 1 \end{aligned} \quad (5.5)$$

- (ii)  *$F_{\mathbf{f}}$  is Frobenius iff  $\chi_{\mathbf{f}} = \varepsilon_{H'}$  iff  $\alpha_H \circ \mathbf{f} = \alpha_{H'}$ .*

- (iii) *The restriction of scalars functor  $F_{\mathbf{f}}$  satisfies*

$$F_{\mathbf{f}} \text{ is } \otimes\text{-Frobenius} \iff \chi_{\mathbf{f}} = \varepsilon_{H'} \text{ and } \mathbf{f}(g_{H'}) = g_H \iff \alpha_H \circ \mathbf{f} = \alpha_{H'} \text{ and } \mathbf{f}(g_{H'}) = g_H.$$

- (iv) *In case that  $H' \subseteq H$  is a Hopf subalgebra, then  $F_{\mathbf{f}}$  is Frobenius if and only if  $\alpha_H(h') = \alpha_{H'}(h')$  for all  $h' \in H'$ . If, in addition,  $g_H = g_{H'}$ , then  $F_{\mathbf{f}}$  is  $\otimes$ -Frobenius.*

*Proof.* By [Shi17], we know that  $\chi_{F_{\mathbf{f}}} = F_{\mathbf{f}}(\mathbb{k}_{\alpha_H}) \otimes (\mathbb{k}_{\alpha_{H'}})^* = \mathbb{k}_{\alpha_H \mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}}$ . Now, consider  $X \in \text{Rep}(H)$ . Then, the commutative diagram (3.4) in Proposition 3.14 gives a description of  $\sigma_X$ . Namely, it is given by the following compositions of Radford isomorphisms

$$\begin{aligned} \mathbb{k}_{\alpha_H \mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} \otimes X_{\mathbf{f}} &\xrightarrow{\text{id} \otimes \mathcal{R}_{X_{\mathbf{f}}}} \mathbb{k}_{\alpha_H \mathbf{f}} \otimes X_{\mathbf{f} S_{H'}^4} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} \\ &= \mathbb{k}_{\alpha_H \mathbf{f}} \otimes X_{S_H^4 \mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} = (\mathbb{k}_{\alpha_H} \otimes X_{S_H^4})_{\mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} \\ &\xrightarrow{(\mathcal{R}'_X)_{\mathbf{f}} \otimes \text{id}} (X \otimes \mathbb{k}_{\alpha_H})_{\mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} = X_{\mathbf{f}} \otimes \mathbb{k}_{\alpha_H \mathbf{f}} \otimes \mathbb{k}_{\overline{\alpha}_{H'}} \end{aligned}$$

Using the description of  $\mathcal{R}$  (5.2) and  $\mathcal{R}'$  (5.3) provided earlier, the above map is given by

$$1 \otimes_{\mathbb{k}} 1 \otimes_{\mathbb{k}} x \mapsto 1 \otimes_{\mathbb{k}} (g_{H'} \cdot_{\mathbf{f}} x) \otimes_{\mathbb{k}} 1 = 1 \otimes_{\mathbb{k}} \mathbf{f}(g_{H'}) \cdot x \otimes_{\mathbb{k}} 1 \mapsto \overline{g_H} \mathbf{f}(g_{H'}) \cdot x \otimes_{\mathbb{k}} 1 \otimes_{\mathbb{k}} 1.$$

Using part (i) and Proposition 3.9, the first equivalence is clear. For the second equivalence, note that we can rewrite the relative modular function as

$$\chi_{\mathbf{f}} = (\alpha_H \circ \mathbf{f}) \times \overline{\alpha}_{H'} \in H^*.$$

As  $\varepsilon \in H^*$  is its unit, we obtain that  $\chi_{\mathbf{f}} = \varepsilon$  if and only if  $(\alpha_H \circ \mathbf{f}) \times \overline{\alpha}_{H'} = \varepsilon$  if and only if  $\alpha_H \circ \mathbf{f} = (\overline{\alpha}_{H'})^{-1} = \alpha_{H'}$ .

By the equivalence (i)  $\iff$  (iii) of Theorem 3.18,  $F_{\mathbf{f}}$  is  $\otimes$ -Frobenius if and only if  $\chi_{\mathbf{f}} \cong \mathbb{1}$  as an object of  $\text{Rep}(H)$  and the half-braiding is the identity map. By part (i), the former is equivalent to  $\chi_{\mathbf{f}} = \varepsilon_{H'}$  and the latter to  $\mathbf{f}(g_{H'}) = g_H$ . Thus, by part (ii), we are done. Part (iii) is a straightforward consequence of (ii).  $\square$

Lastly, we discuss the pivotality of the functor  $F_{\mathbf{f}}$ . Let  $(H', g'_{\text{piv}})$  and  $(H, g_{\text{piv}})$  be pivotal Hopf algebras. We call a bialgebra map  $\mathbf{f} : H' \rightarrow H$  *pivotal* if it satisfies  $\mathbf{f}(g'_{\text{piv}}) = g_{\text{piv}}$ .

**Lemma 5.4.** *The tensor functor  $F_{\mathbf{f}}$  between the pivotal categories  $\text{Rep}(H')$  and  $\text{Rep}(H)$  is pivotal if and only if  $\mathbf{f}$  is pivotal.*

*Proof.* Being a tensor functor,  $F_{\mathbf{f}}$  yields a canonical isomorphism

$$\zeta_X : (X^{\vee})_{\mathbf{f}} = F_{\mathbf{f}}(X^{\vee}) \longrightarrow F_{\mathbf{f}}(X)^{\vee} = (X_{\mathbf{f}})^{\vee}.$$

From (5.4), one can check that  $\zeta_X(f) = f$  for all  $f \in X^{\vee}$ . Since the isomorphisms  $\zeta_X$  are just identity maps, this follows from the definition (2.1) of functor pivotality.  $\square$

### 5.1.3. Examples and remarks

*Example 5.5.* For a Hopf algebra  $H$ , take  $\mathbf{f}$  to be the unit  $u : \mathbb{k} \rightarrow H$  of  $H$ . Then,  $\mathbf{f}$  is a bialgebra map whose induced functor is just the fiber functor  $F_{\mathbf{f}} : \text{Rep}(H) \rightarrow \text{Vec}$ .

- (i) As  $\alpha_H$  is an algebra map, it satisfies  $\alpha_H(1_H) = 1$ . Therefore,  $\alpha_H|_{\mathbb{k}} = \alpha_{\mathbb{k}}$ . This shows that the fiber functor  $F_{\mathbf{f}}$  is always a Frobenius functor.
- (ii) Now, note that  $g_{\mathbb{k}} = 1$ . Thus,  $F_{\mathbf{f}}$  is  $\otimes$ -Frobenius if and only if  $g_H = \mathbf{f}(g_{\mathbb{k}}) = 1_H$ . This is the case if and only if  $H^*$  is unimodular.

For instance,  $H = u_q(\mathfrak{sl}_2)^*$  satisfies this because  $H^* \cong u_q(\mathfrak{sl}_2)$  is unimodular. Thus, the fiber functor  $F : \text{Rep}(u_q(\mathfrak{sl}_2)^*) \rightarrow \text{Vec}$  is  $\otimes$ -Frobenius. On the other hand, the fiber functor  $F : \text{Rep}(u_q(\mathfrak{sl}_2)) \rightarrow \text{Vec}$  is Frobenius, but not  $\otimes$ -Frobenius because  $u_q(\mathfrak{sl}_2)^*$  is not unimodular.

*Remark 5.6.* Let  $\mathbf{f} : H' \rightarrow H$  be a bialgebra map such that  $\chi_{\mathbf{f}} = \varepsilon$ . If  $H$  is unimodular, then so is  $H'$ . This follows from Proposition 3.9(ii) or can be checked directly. However, the converse is not true as illustrated by the following example. Take  $\mathbf{f} = u : \mathbb{k} \rightarrow H$  for a non-unimodular Hopf algebra  $H$  (like the Taft algebra or  $u_q(\mathfrak{sl}_2)^*$ ). By Example 5.5,  $F_{\mathbf{f}}$  is Frobenius (and thus,  $\chi_{\mathbf{f}} = \varepsilon$ ) even though  $H$  is not unimodular.

*Remark 5.7.* A bialgebra map  $\mathbf{f} : H' \rightarrow H$  is called a *Frobenius extension* if  $\mathbf{f}$  is injective and  $F_{\mathbf{f}}$  is Frobenius. In [FLP24b], the concept of a *central Frobenius extension* for Hopf algebras  $H' \subset H$  is introduced. In the finite-dimensional setting, a Frobenius extension  $H' \subset H$  is central Frobenius if and only if  $F_{\mathbf{f}}$  is  $\otimes$ -Frobenius. Indeed, combining [FLP24b, Thm. 4.9] and [FLP24a, Thm. 3.17], the extension  $H \subset H'$  is central Frobenius if and only if the left and right adjoints of  $F_{\mathbf{f}} : \text{Rep}(H) \rightarrow \text{Rep}(H')$  are isomorphic as  $\text{Rep}(H)$ -bimodule functors. This is precisely the definition of being  $\otimes$ -Frobenius. Using this observation, various results in [FLP24b] can be obtained as corollaries of our results in §3 and §4. For instance, [FLP24b, Thm. 4.21], is a special case of Proposition 4.20.

## 5.2. Unimodular and pivotal comodule algebras

In this section, we apply the results from §4 to module categories over Hopf algebras. We start recalling the notions of exact, unimodular and pivotal module categories in this setting.

### 5.2.1. Preliminaries

Left module categories over  $\mathcal{C} = \text{Rep}(H)$  are understood in terms of left  $H$ -comodule algebras  $(L, \delta : L \rightarrow H \otimes_{\mathbb{k}} L)$ . Namely, the category  $\text{Rep}(L)$  is endowed with a left  $\text{Rep}(H)$ -module category structure, defined as follows: for  $X \in \text{Rep}(H)$  and  $M \in \text{Rep}(L)$ ,  $X \triangleright M \in \text{Rep}(L)$  is just  $X \otimes_{\mathbb{k}} M$  as a vector space with  $L$ -action given by

$$l \cdot (x \otimes_{\mathbb{k}} m) = l_{-1} \cdot x \otimes_{\mathbb{k}} l_0 \cdot m.$$

where we have used the Sweedler notation  $\delta(l) = l_{-1} \otimes_{\mathbb{k}} l_0$ . We call  $L$  *exact* (resp. *indecomposable*) if the  $\text{Rep}(H)$ -module category  $\text{Rep}(L)$  is exact (resp. indecomposable). According to [AM07],  $\mathcal{M} = \text{Rep}(L)$  is an exact left  $\mathcal{C}$ -module category if and only if  $L$  is  $H$ -simple from the right and the space of  $H$ -coinvariants of  $L$  is  $\mathbb{k}$ . Moreover, by [Skr07], exact  $H$ -comodule algebras are known to be Frobenius algebras. Thus, we have the following additional information:

- $\lambda_L : L \rightarrow \mathbb{k}$  a linear map called *Frobenius form* and a pair of bases  $\{a^i\}$  and  $\{b_j\}$  of  $L$  that satisfy  $\langle \lambda_L, a^i b_j \rangle = \delta_{i,j}$ .

- An algebra automorphism  $\nu \equiv \nu_L : L \rightarrow L$  called *Nakayama automorphism* satisfying  $\langle \lambda_L, ab \rangle = \langle \lambda_L, \nu(b)a \rangle$  for all  $a, b \in L$ .

With these data at hand, we can give an explicit description of the relative Serre and Nakayama functor of the exact  $\text{Rep}(H)$ -module category  $\text{Rep}(L)$  (see [Yad23, §4] for proofs). Let  $M \in \text{Rep}(L)$ ,  $X \in \text{Rep}(H)$ ,  $x \in X$ ,  $m \in M$  and  $a \in L$ .

- The Nakayama functor is given by  $\mathbb{N}(M) = M_\nu$  with twisted module functor structure
 
$$\mathfrak{n}_{X,M} : (X \triangleright M)_\nu \rightarrow {}^{\vee\vee}X \triangleright M_\nu, \quad x \otimes_{\mathbb{k}} m \mapsto \langle \lambda_L, a_0^i \rangle \phi_X(S^{-1}(a_{-1}^i) \cdot x) \otimes_{\mathbb{k}} \nu(b_i) \cdot m.$$
- The relative Serre functor is given by  $\mathbb{S}(M) = M_{\nu'_H}$  where  $\nu'_H(a) = \langle \alpha, S(a_{-1}) \rangle \nu(a_0)$  with
 
$$\mathfrak{s}_{X,M} : (X \triangleright M)_{\nu'_H} \rightarrow X^{\vee\vee} \triangleright M_{\nu'_H}, \quad x \otimes_{\mathbb{k}} m \mapsto \langle \lambda_L, a_0^i \rangle \phi_X(g_H S^{-1}(a_{-1}^i) \cdot x) \otimes_{\mathbb{k}} \nu(b_i) \cdot m.$$

Here we have used the Einstein convention to suppress summation symbols over  $i$ .

The distinguished invertible object of the dual tensor category  $\text{Rep}(H)_{\text{Rep}(L)}^*$  is given by the composition  $\mathbb{S} \circ \mathbb{N}$ . When this is trivial,  $\text{Rep}(H)_{\text{Rep}(L)}^*$  is a unimodular tensor category and we call  $L$  unimodular. From [Yad23], we get that  $L$  is unimodular if and only if it admits an invertible element  $\tilde{g}_{\text{uni}} \in L$  (called *unimodular element*) that satisfies

$$\tilde{g}_{\text{uni}} a \tilde{g}_{\text{uni}}^{-1} = \nu'_H(\nu(a)), \quad (5.6)$$

$$1_H \otimes \tilde{g}_{\text{uni}} = \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle g_H S^{-3}(a_{-1}^j) S^{-1}(a_{-1}^i) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \delta(\tilde{g}_{\text{uni}}). \quad (5.7)$$

Given a Hopf algebra  $H$  with a pivotal element  $g_{\text{piv}}$ , we call a left  $H$ -comodule algebra pivotal if it admits an invertible element  $\tilde{g}_{\text{piv}} \in L$  (called *pivotal element*) that satisfies

$$\tilde{g}_{\text{piv}} a \tilde{g}_{\text{piv}}^{-1} = \nu'_H(a), \quad g_{\text{piv}} \otimes \tilde{g}_{\text{piv}} = \left( \langle \lambda_L, a_0^i \rangle g_H S^{-1}(a_{-1}^i) \otimes_{\mathbb{k}} \nu(b_i) \right) \delta(\tilde{g}_{\text{piv}}). \quad (5.8)$$

Then, pivotal structures on  $\text{Rep}(L)$  are in bijection with pivotal elements of  $L$  [Shi23a].

*Remark 5.8.* For an indecomposable exact left  $H$ -comodule algebra  $L$ , the space of pivotal or unimodular elements is at most one-dimensional. Thus, one is usually only interested in whether such an element exists or not. Our results in the next section provide a way to answer this.

### 5.2.2. Twisting module categories obtained using comodule algebras

In this section, we record some consequences of the results obtained in §4. As before,  $\mathfrak{f} : H' \rightarrow H$  is a bialgebra map. Let  $L$  be a left  $H'$ -comodule algebra with coaction  $\delta : L \rightarrow H' \otimes L$ . Define the map  $\delta_{\mathfrak{f}}$  by the composition:

$$L \xrightarrow{\delta} H' \otimes L \xrightarrow{\mathfrak{f} \otimes \text{id}_L} H \otimes L.$$

Explicitly, it is given for  $a \in L$  by the formula  $\delta_{\mathfrak{f}}(a) = \mathfrak{f}(a_{-1}) \otimes_{\mathbb{k}} a_0$ . As a consequence of Proposition 4.4, we obtain:

**Lemma 5.9.** *Let  $\mathfrak{f} : H' \rightarrow H$  be a perfect bialgebra map. Then  $(L, \delta)$  is an exact  $H'$ -comodule algebra if and only if  $(L, \delta_{\mathfrak{f}})$  is an exact  $H$ -comodule algebra.*

Given a left  $H$ -comodule algebra  $L$ , it is in general difficult to tell whether  $L$  is pivotal or unimodular. Next, we provide some criteria using results from §4.

**Proposition 5.10.** *Let  $\mathfrak{f} : H' \rightarrow H$  be a bialgebra map such that  $\chi_{\mathfrak{f}} = \varepsilon_{H'}$  and  $\mathfrak{f}(g_{H'}) = g_H$ .*



- (i) If  $(L, \delta)$  is a unimodular  $H'$ -comodule algebra with unimodular element  $\tilde{g}_{\text{uni}}$ , then  $(L, \delta_{\mathbf{f}})$  is a unimodular  $H$ -comodule algebra with the same unimodular element.
- (ii) Suppose that  $H$  and  $H'$  admit pivotal elements  $g_{\text{piv}}$  and  $g'_{\text{piv}}$  respectively, and  $\mathbf{f}(g'_{\text{piv}}) = g_{\text{piv}}$ . If  $(L, \delta)$  is a pivotal  $H'$ -comodule algebra with pivotal element  $\tilde{g}_{\text{piv}}$ , then  $(L, \delta_{\mathbf{f}})$  is a pivotal  $H$ -comodule algebra with the same pivotal element.

*Proof.* Because of the assumption that  $\chi_{\mathbf{f}} = \varepsilon$  and  $\mathbf{f}(g_{H'}) = g_H$ , by Proposition 5.3,  $F_{\mathbf{f}}$  is  $\otimes$ -Frobenius.

(i) As  $\tilde{g}_{\text{uni}}$  is a unimodular element of the  $H'$ -comodule algebra  $L$ , it satisfies (5.6) and (5.7). By Proposition 4.20, it follows that  $\text{Rep}(L)$  is also a unimodular  $\text{Rep}(H)$ -module category. In other words,  $(L, \delta_{\mathbf{f}})$  is a unimodular  $H$ -comodule algebra. In fact, we will show that  $\tilde{g}_{\text{uni}}$  is a unimodular element for it. First note that

$$\begin{aligned} \nu'_H(\nu(a)) &= \langle \alpha_H, S_H(\mathbf{f}(a_{-1})) \rangle \nu^2(a_0) \stackrel{(5.4)}{=} \langle \alpha_H, \mathbf{f}(S_{H'}(a_{-1})) \rangle \nu^2(a_0) \\ &\stackrel{\text{Prop. 5.3(iii)}}{=} \langle \alpha_{H'}, S_{H'}(a_{-1}) \rangle \nu^2(a_0) = \nu'_{H'}(\nu(a)) = \tilde{g}_{\text{uni}} a \tilde{g}_{\text{uni}}^{-1}. \end{aligned}$$

Thus,  $\tilde{g}_{\text{uni}}$  satisfies the first condition (5.6) of being a unimodular element for  $H$ . Also,

$$\begin{aligned} 1_H \otimes_{\mathbb{k}} \tilde{g}_{\text{uni}} &= (\mathbf{f} \otimes_{\mathbb{k}} \text{id}) [1_{H'} \otimes_{\mathbb{k}} \tilde{g}_{\text{uni}}] \\ \stackrel{(5.7) \text{ for } L, H'}{=} &= (\mathbf{f} \otimes_{\mathbb{k}} \text{id}) \left[ \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle g_{H'} S_{H'}^{-3}(a_{-1}^j) S_{H'}^{-1}(a_{-1}^i) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \delta(\tilde{g}_{\text{uni}}) \right] \\ &= \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle \mathbf{f}(g_{H'} S_{H'}^{-3}(a_{-1}^j) S_{H'}^{-1}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \left( (\mathbf{f} \otimes_{\mathbb{k}} \text{id}) \circ \delta \right) (\tilde{g}_{\text{uni}}) \\ \stackrel{(\mathbf{f} \text{ alg. map})}{=} &= \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle \mathbf{f}(g_{H'}) \mathbf{f}(S_{H'}^{-3}(a_{-1}^j)) \mathbf{f}(S_{H'}^{-1}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \delta_{\mathbf{f}}(\tilde{g}_{\text{uni}}) \\ \stackrel{(5.4)}{=} &= \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle \mathbf{f}(g_{H'}) S_H^{-3}(\mathbf{f}(a_{-1}^j)) S_H^{-1}(\mathbf{f}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \delta_{\mathbf{f}}(\tilde{g}_{\text{uni}}) \\ &= \left( \langle \lambda_L, a_0^i \rangle \langle \lambda_L, a_0^j \rangle g_H S_H^{-3}(\mathbf{f}(a_{-1}^j)) S_H^{-1}(\mathbf{f}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_j b_i) \right) \delta_{\mathbf{f}}(\tilde{g}_{\text{uni}}) \end{aligned}$$

Comparing with (5.7), this shows that  $\tilde{g}_{\text{uni}}$  also satisfies the second condition. Thus, it is a unimodular element of  $L$  considered as an  $H$ -comodule algebra.

(ii) The assumptions of ensure that  $F$  is  $\otimes$ -Frobenius and pivotal. Thus, the claim that  $(L, \delta_{\mathbf{f}})$  is a pivotal  $H$ -comodule algebra is a direct consequence of Proposition 4.24. We will show that  $\tilde{g}_{\text{piv}}$  serves as pivotal element. By a similar calculation as in part (i), we obtain that  $\nu'_H(a) = \tilde{g}_{\text{piv}} a \tilde{g}_{\text{piv}}^{-1}$  for all  $a \in L$ . Furthermore,

$$\begin{aligned} g_{\text{piv}} \otimes_{\mathbb{k}} \tilde{g}_{\text{piv}} &= \mathbf{f}(g'_{\text{piv}}) \otimes_{\mathbb{k}} \tilde{g}_{\text{piv}} = (\mathbf{f} \otimes_{\mathbb{k}} \text{id})(g'_{\text{piv}} \otimes_{\mathbb{k}} \tilde{g}_{\text{piv}}) \\ \stackrel{(5.8) \text{ for } L, H'}{=} &= (\mathbf{f} \otimes_{\mathbb{k}} \text{id}) \left[ \left( \langle \lambda_L, a_0^i \rangle g'_H S_H^{-1}(a_{-1}^i) \otimes_{\mathbb{k}} \nu(b_i) \right) \delta(\tilde{g}_{\text{piv}}) \right] \\ &= \langle \lambda_L, a_0^i \rangle \left( \mathbf{f}(g'_H S_H^{-1}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_i) \right) (\mathbf{f} \otimes_{\mathbb{k}} \text{id}) \delta(\tilde{g}_{\text{piv}}) \\ \stackrel{(\mathbf{f} \text{ alg. map})}{=} &= \langle \lambda_L, a_0^i \rangle \left( \mathbf{f}(g'_H) \mathbf{f}(S_H^{-1}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_i) \right) \delta_{\mathbf{f}}(\tilde{g}_{\text{piv}}) \\ \stackrel{(\mathbf{f}(g_{H'})=g_H, (5.4))}{=} &= \left( \langle \lambda_L, a_0^i \rangle g_H S_H^{-1}(\mathbf{f}(a_{-1}^i)) \otimes_{\mathbb{k}} \nu(b_i) \right) \delta_{\mathbf{f}}(\tilde{g}_{\text{piv}}) \end{aligned}$$

Comparing with (5.8), this shows that  $\tilde{g}_{\text{piv}}$  satisfies the two conditions needed for it to be a pivotal element of the  $H$ -comodule algebra  $(L, \delta_{\mathbf{f}})$ .  $\square$

**Corollary 5.11.** *Let  $H' \subset H$  be a Hopf subalgebra so that  $\alpha_H|_{H'} = \alpha_{H'}$  and  $g_H = g_{H'}$ .*

- (i) *Any unimodular  $H'$ -comodule algebra is also a unimodular  $H$ -comodule algebra with the same unimodular element.*
- (ii) *If  $g_{\text{piv}} \in H'$  is a pivotal element for both  $H$  and  $H'$ , then a pivotal  $H'$ -comodule algebra is also a pivotal  $H$ -comodule algebra with the same pivotal element.*

*Proof.* The statement is an instance of Proposition 5.10 for an inclusion of Hopf algebras.  $\square$

### 5.2.3. Examples

We close the section with an example illustrating multiple results in the case of  $\mathcal{C} = \text{Rep}(u_q(\mathfrak{sl}_2))$ . Fix an odd integer  $n = 2h + 1$  for  $h > 0$  and let  $q$  be a primitive  $n$ -th root of unity. The Hopf algebra  $u_q(\mathfrak{sl}_2)$  is generated, as an algebra, by  $E$ ,  $F$  and  $K$  subject to the relations

$$K^n = 1, \quad E^n = F^n = 0, \quad KE = q^2EK, \quad KF = q^{-2}FK, \quad EF - FE = \frac{K - K^{-1}}{q - q^{-1}}.$$

The coalgebra structure is given by  $\varepsilon(K) = 1$ ,  $\varepsilon(E) = \varepsilon(F) = 0$  and

$$\Delta(K) = K \otimes K, \quad \Delta(E) = E \otimes K + 1 \otimes E, \quad \Delta(F) = F \otimes 1 + K^{-1} \otimes F.$$

The antipode, on the generators, assigns the following values:  $S(K) = K^{-1}$ ,  $S(E) = -EK^{-1}$  and  $S(F) = -KF$ . Also, the element  $\Lambda = (\sum_{i=0}^{n-1} K^i)E^{n-1}F^{n-1}$  is a left integral. From this description, one obtains that the modular function  $\alpha : u_q(\mathfrak{sl}_2) \rightarrow \mathbb{k}$  is given by  $\alpha(E) = \alpha(F) = 0$  and  $\alpha(K) = 1$ . Moreover, the distinguished grouplike element is given by  $g_{u_q(\mathfrak{sl}_2)} = K^2$ . Lastly,  $S^2(h) = KhK^{-1}$  for all  $h \in u_q(\mathfrak{sl}_2)$ , so  $g_{\text{piv}} = K$  is a pivotal element of  $u_q(\mathfrak{sl}_2)$ .

*Example 5.12.* If one considers the subalgebra of  $u_q(\mathfrak{sl}_2)$  generated by  $E$  and  $K$ , one obtains the Taft (Hopf) algebra  $T := T_n(q^2)$ . The right modular function  $\alpha_T$  is given by  $\alpha(E) = 0$  and  $\alpha(K) = q^{-2}$ . As  $\alpha|_T \neq \alpha_T$ , the functor  $F : \text{Rep}(u_q(\mathfrak{sl}_2)) \rightarrow \text{Rep}(T)$  is not Frobenius.

*Example 5.13.* If one considers the subalgebra of  $u_q(\mathfrak{sl}_2)$  generated by  $K$ , one obtains a Hopf subalgebra isomorphic to the group algebra,  $T' := \mathbb{k}\mathbb{Z}_n$ , of  $\mathbb{Z}_n$ . The right modular function  $\alpha_{T'}$  is given by  $\alpha_{T'}(K) = 1$ . As  $\alpha|_{T'} = \alpha_{T'}$ , the restriction of scalars functor  $F : \text{Rep}(u_q(\mathfrak{sl}_2)) \rightarrow \text{Rep}(\mathbb{k}\mathbb{Z}_n)$  is a Frobenius functor. However,  $g_{u_q(\mathfrak{sl}_2)} = K^2$  and  $g_{\mathbb{k}\mathbb{Z}_n} = 1$ . Thus,  $F$  is not  $\otimes$ -Frobenius.

Recall that indecomposable exact  $\mathbb{k}\mathbb{Z}_n$ -comodule algebras are of the form  $\mathbb{k}L_\psi$  with coaction  $z \mapsto K^{n/d} \otimes z$  where  $L = \langle z \rangle \cong \mathbb{Z}_d$  (for  $d$  a divisor of  $n$ ) is a subgroup of  $\mathbb{Z}_n$  and  $\psi$  is a 2-cocycle of  $L$  satisfying  $d(\psi) = 0$  [Ost03]. But the only possible 2-cocycle is the trivial one. Thus, the comodule algebras  $L$  can be identified with Hopf subalgebras of  $\mathbb{k}\mathbb{Z}$  generated by  $K^{n/d}$ . Moreover, as  $\mathbb{k}\mathbb{Z}_n$  is semisimple, every exact comodule algebra is unimodular. Since, the restriction of scalars functor  $F : \text{Rep}(u_q(\mathfrak{sl}_2)) \rightarrow \text{Rep}(\mathbb{k}\mathbb{Z}_n)$  is not  $\otimes$ -Frobenius, using Proposition 4.20, we can conclude that the aforementioned  $\mathbb{k}\mathbb{Z}_n$ -comodule algebras are not unimodular as  $u_q(\mathfrak{sl}_2)$ -comodule algebras.

Furthermore,  $K$  is a pivotal element for  $\mathbb{k}\mathbb{Z}_n$ . By [Shi23a, Remark 4.28], the comodule algebras  $L = \langle K^{n/d} \rangle$  are pivotal if and only if the pivotal element  $K \in L$ . However, this is not possible unless  $L = \mathbb{k}\mathbb{Z}_n$ . In this case, by Proposition 4.24, we get that  $\mathbb{k}\mathbb{Z}_n$  is not a pivotal  $u_q(\mathfrak{sl}_2)$ -comodule algebra.

## 6. APPLICATIONS TO INTERNAL NATURAL TRANSFORMATIONS

In [FS21b] the authors studied an internalized version of the vector space of natural transformations between two functors. Given a finite tensor category  $\mathcal{C}$ , an appropriate internal Hom involving two module functors  $H, H' : {}_c\mathcal{M} \rightarrow {}_c\mathcal{N}$  gives an object inside the Drinfeld center of  $\mathcal{C}$  called the *object of internal natural transformations* from  $H$  to  $H'$ . Under certain assumptions, this object is endowed with the structure of a symmetric Frobenius algebra internal to  $\mathcal{Z}(\mathcal{C})$ . In this section, the results from §4 allow us to reprove and strengthen many statements about internal natural transformations.

### 6.1. Exactness and unimodularity of the $\mathcal{Z}(\mathcal{C})$ -module category $\text{Func}(\mathcal{M}, \mathcal{N})$

Let  $\mathcal{C}$  be a finite tensor category and  $\mathcal{M}$  and  $\mathcal{N}$  exact  $\mathcal{C}$ -module categories. Recall, that the category of module functors  $\text{Func}(\mathcal{M}, \mathcal{N})$  becomes an exact  $(\mathcal{C}_{\mathcal{N}}^*, \mathcal{C}_{\mathcal{M}}^*)$ -bimodule category, with action given by composition of module functors (2.4). The internal Hom's can be described using adjoint functors:

**Lemma 6.1.** *For  $H, H' \in \text{Func}(\mathcal{M}, \mathcal{N})$ , we have that the internal Hom's are given by*

$$\underline{\text{Hom}}_{\text{Func}(\mathcal{M}, \mathcal{N})}^{\mathcal{C}_{\mathcal{N}}^*}(H, H') \cong H' \circ H^{\text{la}} \quad \text{and} \quad \underline{\text{Hom}}_{\text{Func}(\mathcal{M}, \mathcal{N})}^{\overline{\mathcal{C}_{\mathcal{M}}^*}}(H, H') \cong H^{\text{ra}} \circ H'.$$

*Proof.* For  $F \in \mathcal{C}_{\mathcal{M}}^*$ , observe that there is a natural isomorphism

$$\varphi : \text{Hom}_{\text{Func}(\mathcal{M}, \mathcal{N})}(H \circ F, H') \cong \text{Hom}_{\text{Func}(\mathcal{M}, \mathcal{M})}(F, H^{\text{ra}} \circ H').$$

The assignment  $\varphi$  and its inverse are defined via the formulas

$$\varphi(\alpha) = (\text{id}_{H^{\text{ra}}} \circ \alpha)(\eta \circ \text{id}_F) : F \Rightarrow H^{\text{ra}} \circ H', \quad \varphi^{-1}(\beta) = (\varepsilon \circ \text{id}_F)(\text{id}_H \circ \beta) : H \circ F \Rightarrow H'.$$

where  $\eta : \text{id}_{\mathcal{M}} \Rightarrow H^{\text{ra}} \circ H$  and  $\varepsilon : H \circ H^{\text{ra}} \Rightarrow \text{id}_{\mathcal{N}}$  denote the unit and counit of the adjunction  $H \dashv H^{\text{ra}}$ . Since  $\alpha, \eta$  are  $\mathcal{C}$ -module natural transformations, we have that  $\varphi(\alpha)$  is a  $\mathcal{C}$ -module functor. The statement for the  $\mathcal{C}_{\mathcal{N}}^*$ -internal Hom follows analogously.  $\square$

Given an exact  $\mathcal{C}$ -module category  $\mathcal{N}$  we have a tensor functor

$$\begin{aligned} \Psi_{\mathcal{N}} : \quad \mathcal{Z}(\mathcal{C}) &\longrightarrow \mathcal{C}_{\mathcal{N}}^* \\ (c, \nu) &\longmapsto c \triangleright - \end{aligned} \tag{6.1}$$

where the half-braiding  $\nu$  endows the functor  $c \triangleright - : \mathcal{N} \rightarrow \mathcal{N}$  with a  $\mathcal{C}$ -module structure. The tensor structure on the functor  $\Psi_{\mathcal{N}}$  is given by the module associativity constraints of  $\mathcal{N}$

$$c \triangleright (d \triangleright n) \cong (c \otimes d) \triangleright n.$$

*Remark 6.2.* In [DKR15, §3.1], the tensor functor (6.1) is called  $\alpha$ -induction. The  $\mathcal{Z}(\mathcal{C})$ -action on  $\text{Func}(\mathcal{M}, \mathcal{N})$  is also studied there. However, in the literature,  $\alpha$ -induction typically assumes that  $\mathcal{C}$  is braided, a condition we do not impose. Consequently, we avoid this terminology in this article.

For an exact  $\mathcal{C}$ -module category  $\mathcal{M}$ , the same assignment given by (6.1) can be made into a tensor functor whose target is the monoidal opposite of the dual tensor category with respect to  $\mathcal{M}$

$$\Phi_{\mathcal{M}} : \quad \mathcal{Z}(\mathcal{C}) \longrightarrow \overline{\mathcal{C}_{\mathcal{M}}^*},$$

but with tensor structure defined by half-braidings

$$c \triangleright (d \triangleright n) \cong (c \otimes d) \triangleright n \xrightarrow{\nu_d^c} (d \otimes c) \triangleright n.$$

Now, we can consider the twist of the  $\mathcal{C}_\mathcal{N}^*$ -module category  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  by  $\Psi_\mathcal{N}$ . Explicitly, the  $\mathcal{Z}(\mathcal{C})$ -action is given by

$$\begin{aligned} \triangleright^{\Psi_\mathcal{N}} : \mathcal{Z}(\mathcal{C}) \times \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N}) &\rightarrow \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N}) \\ ((c, \nu), H) &\mapsto c \triangleright H \end{aligned}$$

the composition of the  $\mathcal{C}$ -module functors  $c \triangleright - : \mathcal{N} \rightarrow \mathcal{N}$  and  $H$ , and module structure (2.4). Similarly, if we consider  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  as a (left)  $\overline{\mathcal{C}_\mathcal{M}^*}$ -module category, we obtain a  $\mathcal{Z}(\mathcal{C})$ -module structure by pulling back the action along  $\Phi_\mathcal{M}$ :

$$\begin{aligned} \triangleright^{\Phi_\mathcal{M}} : \mathcal{Z}(\mathcal{C}) \times \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N}) &\rightarrow \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N}) \\ ((c, \nu), H) &\mapsto H(c \triangleright -) \end{aligned} .$$

These two actions of the Drinfeld center  $\mathcal{Z}(\mathcal{C})$  on  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  are equivalent.

**Lemma 6.3.**  $\Psi_\mathcal{N}\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  and  $\Phi_\mathcal{M}\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  are equivalent as  $\mathcal{Z}(\mathcal{C})$ -module categories.

*Proof.* We have for every  $(c, \nu) \in \mathcal{Z}(\mathcal{C})$  and  $H \in \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  a module natural isomorphism

$$(c, \nu) \triangleright^{\Psi_\mathcal{N}} H = c \triangleright H \xrightarrow{\varphi_{c, -}} H(c \triangleright -) = (c, \nu) \triangleright^{\Phi_\mathcal{M}} H$$

given by the  $\mathcal{C}$ -module functor structure of  $H$ . A routine check shows that these isomorphisms furnish a  $\mathcal{Z}(\mathcal{C})$ -module functor structure on the identity functor of the category  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$ .  $\square$

*Remark 6.4.* Notice that the functor  $\Psi_\mathcal{N}$  (6.1) can be factorized as  $U \circ S$ , where  $U : \mathcal{Z}(\mathcal{C}_\mathcal{N}^*) \rightarrow \mathcal{C}_\mathcal{N}^*$  is the forgetful functor, and

$$\begin{aligned} S : \mathcal{Z}(\mathcal{C}) &\longrightarrow \mathcal{Z}(\mathcal{C}_\mathcal{N}^*), \\ (c, \nu) &\longmapsto c \triangleright - \end{aligned} \tag{6.2}$$

where  $c \triangleright - : \mathcal{N} \rightarrow \mathcal{N}$  is the  $\mathcal{C}$ -module functor assigned by (6.1), and the  $\mathcal{C}_\mathcal{N}^*$ -half-braiding on  $c \triangleright -$  is defined for  $H \in \mathcal{C}_\mathcal{N}^*$  by the  $\mathcal{C}$ -module structure  $\varphi_{c, -} : c \triangleright - \circ H \xrightarrow{\sim} H \circ c \triangleright -$  of  $H$ . Moreover,  $S$  is braided and thus endows  $\Psi_\mathcal{N}$  with the structure of a central tensor functor.

The internal (co)Hom's of the  $\mathcal{Z}(\mathcal{C})$ -module category  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  can be described in terms of a (co)end as proved in [FS21b]. In the case that  $\mathcal{C} = \text{Vec}$ , this construction recovers, by the Yoneda lemma, the vector space of natural transformations between the functors involved.

**Definition 6.5.** [FS21b, Def. 4] Let  $\mathcal{M}$  and  $\mathcal{N}$  be exact  $\mathcal{C}$ -module categories. Given  $H, H' \in \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$ , their object of *internal natural transformations* is defined as the internal hom

$$\underline{\text{Nat}}_{(\mathcal{M}, \mathcal{N})}(H, H') := \underline{\text{Hom}}_{\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})}^{\mathcal{Z}(\mathcal{C})}(H, H').$$

When clear from the context, we will drop the indices  $\mathcal{M}, \mathcal{N}$ .

In view of Lemma 6.3, the  $\mathcal{Z}(\mathcal{C})$ -module structure on  $\text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$  can be obtained by twisting either via the perfect tensor functor  $\Psi_\mathcal{N}$  or via  $\Phi_\mathcal{M}$ . As a consequence, the internal natural transformations object can also be described in terms of either of these perfect tensor functors.

**Lemma 6.6.** *Let  $\mathcal{M}$  and  $\mathcal{N}$  be exact  $\mathcal{C}$ -module categories. There is a natural isomorphism*

$$\underline{\text{Nat}}_{(\mathcal{M}, \mathcal{N})}(H, H') \cong \Phi_\mathcal{M}^{\text{ra}}(H^{\text{ra}} \circ H') \cong \Psi_\mathcal{N}^{\text{ra}}(H' \circ H^{\text{la}})$$

for  $H, H' \in \text{Func}_\mathcal{C}(\mathcal{M}, \mathcal{N})$ .

*Proof.* The result follows from Lemma 6.1 and isomorphism (4.4).  $\square$

**Lemma 6.7.** *The functor (6.1) is surjective. Thus, the tensor functors  $\Psi_{\mathcal{N}}$  and  $\Phi_{\mathcal{M}}$  are perfect.*

*Proof.* In view of Remark 6.4, we have a factorization  $U \circ S$ . According to [Shi20, Thm. 3.13]  $S$  is in particular an equivalence of  $\mathbb{k}$ -linear categories. Furthermore, by [EGNO16, Cor. 7.13.11], the functor  $U$  is surjective. Thus, it follows by Proposition 3.11 (iii), that (6.1) is surjective. Since, composition of tensor functors is a tensor functor, considering appropriately adapted tensor structures on (6.2) implies that  $\Psi_{\mathcal{N}}$  and  $\Phi_{\mathcal{M}}$  are perfect.  $\square$

We obtain the following result generalizing [FS21b, Thm. 18], where we do not require the additional assumption that  $\mathcal{C}$  is pivotal.

**Theorem 6.8.** *Let  $\mathcal{C}$  be a finite tensor category, and  $\mathcal{M}$  and  $\mathcal{N}$  be exact  $\mathcal{C}$ -module categories. The  $\mathcal{Z}(\mathcal{C})$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is exact.*

*Proof.* By [EGNO16, Prop. 7.12.14],  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is an exact left  $\mathcal{C}_{\mathcal{N}}^*$ -module category. Also, by Lemma 6.7,  $\Psi_{\mathcal{N}}$  is a perfect tensor functor. It follows from Proposition 4.5 that  $\Psi_{\mathcal{N}} \text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is an exact left  $\mathcal{Z}(\mathcal{C})$ -module category.  $\square$

**Proposition 6.9.** *The tensor functor  $\Psi_{\mathcal{N}}$  is  $\otimes$ -Frobenius iff  $\mathcal{N}$  is unimodular iff  $\Psi_{\mathcal{N}}^{\text{ra}}$  is Frobenius monoidal. Similarly, the tensor functor  $\Phi_{\mathcal{M}}$  is  $\otimes$ -Frobenius iff  $\mathcal{M}$  is unimodular iff  $\Phi_{\mathcal{M}}^{\text{ra}}$  is Frobenius monoidal.*

*Proof.* From Remark 6.4 we have that  $\Psi_{\mathcal{N}} = U \circ S$ . Thus, according to Proposition 3.11, the relative modular object of  $\Psi_{\mathcal{N}}$  can be computed as  $U(\chi_S) \circ \chi_U$ . But, since  $S$  is an equivalence  $\chi_S \cong \text{id}_{\mathcal{N}}$ , and since  $\mathcal{Z}(\mathcal{C}_{\mathcal{N}}^*)$  is unimodular,  $\chi_U \cong D_{\mathcal{C}_{\mathcal{N}}^*}^{-1}$  by Proposition 3.14. It follows that the relative modular object of  $\Psi_{\mathcal{N}}$  is  $D_{\mathcal{C}_{\mathcal{N}}^*}^{-1}$ , which together with Proposition 3.9 proves that  $\Psi_{\mathcal{N}}$  is Frobenius iff  $\mathcal{N}$  is unimodular. Since  $\Psi_{\mathcal{N}}$  is a central tensor functor, Proposition 3.20 ensures that  $\Psi_{\mathcal{N}}$  is  $\otimes$ -Frobenius iff  $\Psi_{\mathcal{N}}$  is Frobenius iff  $\Psi_{\mathcal{N}}^{\text{ra}}$  is Frobenius monoidal.  $\square$

**Proposition 6.10.** *Let  $\mathcal{C}$  be a finite tensor category, and  $\mathcal{M}$  and  $\mathcal{N}$  be unimodular  $\mathcal{C}$ -module categories. The  $\mathcal{Z}(\mathcal{C})$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is unimodular.*

*Proof.*  $\Psi_{\mathcal{N}}$  is perfect by Lemma 6.7. Now, assuming that  $\mathcal{N}$  is unimodular, Proposition 6.9 implies that  $\Psi_{\mathcal{N}}$  is  $\otimes$ -Frobenius. From [FGJS22, Thm. 3.11] the dual-tensor category of  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  with respect to  $\mathcal{C}_{\mathcal{N}}^*$  is equivalent to  $\mathcal{C}_{\mathcal{M}}^*$ . This means that a unimodular structure on  ${}_{\mathcal{C}}\mathcal{M}$  induces a unimodular structure on the  $\mathcal{C}_{\mathcal{N}}^*$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ . Thus, by Proposition 4.20 we can pull back the unimodular structure along  $\Psi_{\mathcal{N}}$  to obtain a unimodular structure on the  $\mathcal{Z}(\mathcal{C})$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ .  $\square$

**Proposition 6.11.** *Let  $\mathcal{C}$  be a finite tensor category and  $\mathcal{M}$  and  $\mathcal{N}$  be exact  $\mathcal{C}$ -module categories. Additionally, assume that either  $\mathcal{M}$  or  $\mathcal{N}$  is unimodular.*

- (i) *Let  $H \in \text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  and  $H^{\text{ra}} \cong H^{\text{la}}$  a  $\mathcal{C}$ -module natural isomorphism. Then the algebra  $\underline{\text{Nat}}_{(\mathcal{M}, \mathcal{N})}(H, H)$  is endowed with the structure of a Frobenius algebra in  $\mathcal{Z}(\mathcal{C})$ .*
- (ii) *The algebra  $\underline{\text{Nat}}(\text{id}_{\mathcal{M}}, \text{id}_{\mathcal{M}})$  has the structure of a commutative Frobenius algebra in  $\mathcal{Z}(\mathcal{C})$ .*

*Proof.* The assumption that  $H^{\text{la}} \cong H^{\text{ra}}$  as  $\mathcal{C}$ -module functors implies that the composition  $H^{\text{ra}} \circ H$  has a canonical structure of a Frobenius algebra in  $\mathcal{C}_{\mathcal{N}}^*$ . Moreover,  $\mathcal{M}$  being unimodular implies that  $\Phi_{\mathcal{M}}^{\text{ra}}$  is Frobenius monoidal according to Proposition 6.9. Thus it preserves Frobenius algebras.

Consequently,  $\Phi_{\mathcal{M}}^{\text{ra}}(H^{\text{ra}} \circ H)$  inherits a Frobenius algebra structure. The statement follows from Lemma 6.6. The commutativity in assertion (ii) follows from [EGNO16, Prop. 8.8.8]. A similar argument shows the claim if we assume  $\mathcal{N}$  to be unimodular instead.  $\square$

## 6.2. Pivotal setting

Next we explore the implications on the pivotal setting. Let  $\mathcal{C}$  be a pivotal finite tensor category. The pivotal structure  $\mathfrak{p}_{\mathcal{C}}: c \xrightarrow{\sim} c^{\vee\vee}$  on  $\mathcal{C}$  induces a pivotal structure on  $\mathcal{Z}(\mathcal{C})$  via  $\mathfrak{p}_{(\mathcal{C}, \sigma)} := \mathfrak{p}_{\mathcal{C}}$  [EGNO16, Ex. 7.13.6]. This choice on the Drinfeld center renders the forgetful functor  $U: \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$  a pivotal tensor functor.

**Lemma 6.12.** *Given a pivotal  $\mathcal{C}$ -module category  $\mathcal{N}$ , the tensor functor  $\Psi_{\mathcal{N}}$  is pivotal.*

*Proof.* As described in Remark 6.4 we have a factorization  $\Psi_{\mathcal{N}} = U \circ S$ . By [FGJS22, Prop. 5.15],  $S$  is pivotal, which leads to the result.  $\square$

Now, consider pivotal  $\mathcal{C}$ -module categories  $\mathcal{M}$  and  $\mathcal{N}$ . The relative Serre functors of the  $(\mathcal{C}_{\mathcal{N}}^*, \mathcal{C}_{\mathcal{M}}^*)$ -bimodule category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  can be expressed in terms of those of  $\mathcal{M}$  and  $\mathcal{N}$  according to [FGJS22, Lemma 5.3] and [FGJS22, Cor. 4.12] by

$$\mathbb{S}_{\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})}^{\mathcal{C}_{\mathcal{N}}^*}(H) \cong \overline{\mathbb{S}}_{\mathcal{N}}^{\mathcal{C}} \circ H \circ \mathbb{S}_{\mathcal{M}}^{\mathcal{C}}, \quad \overline{\mathbb{S}}_{\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})}^{\mathcal{C}_{\mathcal{M}}^*}(H) \cong \mathbb{S}_{\mathcal{N}}^{\mathcal{C}} \circ H \circ \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{C}}$$

for  $H \in \text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ .

**Proposition 6.13.** *Let  $\mathcal{M}$  and  $\mathcal{N}$  pivotal  $\mathcal{C}$ -module categories.*

(i)  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  inherits the structure of a pivotal  $(\mathcal{C}_{\mathcal{N}}^*, \mathcal{C}_{\mathcal{M}}^*)$ -bimodule category via the isomorphisms

$$\begin{aligned} P_H : H &\xrightarrow{\tilde{\mathfrak{q}}^{-1} \circ \text{id} \circ \tilde{\mathfrak{p}}} \overline{\mathbb{S}}_{\mathcal{N}}^{\mathcal{C}} \circ H \circ \mathbb{S}_{\mathcal{M}}^{\mathcal{C}} \cong \mathbb{S}_{\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})}^{\mathcal{C}_{\mathcal{N}}^*}(H) \\ Q_H : H &\xrightarrow{\tilde{\mathfrak{q}} \circ \text{id} \circ \tilde{\mathfrak{p}}^{-1}} \mathbb{S}_{\mathcal{N}}^{\mathcal{C}} \circ H \circ \overline{\mathbb{S}}_{\mathcal{M}}^{\mathcal{C}} \cong \overline{\mathbb{S}}_{\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})}^{\mathcal{C}_{\mathcal{M}}^*}(H) \end{aligned}$$

where  $\tilde{\mathfrak{p}}$  and  $\tilde{\mathfrak{q}}$  are the pivotal structures of  ${}_{\mathcal{C}}\mathcal{M}$  and  ${}_{\mathcal{C}}\mathcal{N}$  respectively.

(ii) For  $H \in \text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$ , the compositions  $H \circ H^{\text{la}}$  and  $H^{\text{ra}} \circ H$  are symmetric Frobenius algebras in  $\mathcal{C}_{\mathcal{N}}^*$  and  $\mathcal{C}_{\mathcal{M}}^*$ , respectively.

*Proof.* Part (i) follows by applying [FGJS22, Prop. 5.4] to the bimodule categories  ${}_{\mathcal{C}}\mathcal{M}_{\overline{\mathcal{C}_{\mathcal{M}}^*}}$  and  ${}_{\mathcal{C}}\mathcal{N}_{\overline{\mathcal{C}_{\mathcal{N}}^*}}$ , that are pivotal bimodules in view of [FGJS22, Lemma 5.8 (i)]. By [Shi23a, Thm. 3.1], the internal ends, which are  $H \circ H^{\text{la}}$  and  $H^{\text{ra}} \circ H$  due to Lemma 6.1, are symmetric Frobenius algebras.  $\square$

**Proposition 6.14.** *Let  $\mathcal{C}$  be a pivotal finite tensor category, and  $\mathcal{M}$  and  $\mathcal{N}$  pivotal  $\mathcal{C}$ -module categories. Additionally, assume that either  $\mathcal{M}$  or  $\mathcal{N}$  or both are unimodular. The  $\mathcal{Z}(\mathcal{C})$ -module category  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is endowed with a pivotal structure.*

*Proof.* Since  $\mathcal{N}$  is unimodular, by Proposition 6.9,  $\Psi_{\mathcal{N}}$  is  $\otimes$ -Frobenius. On the other hand,  $\Psi_{\mathcal{N}}$  is pivotal. As the  $\mathcal{C}_{\mathcal{N}}^*$ -module  $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{N})$  is pivotal, the claim follows from Proposition 4.24.  $\square$

In the pivotal setting, the objects of internal natural transformations are not only algebras, but symmetric Frobenius algebras.

**Corollary 6.15.** *Let  $\mathcal{C}$  be a pivotal finite tensor category and  $\mathcal{M}, \mathcal{N}$  be pivotal  $\mathcal{C}$ -module categories. Assume that  $\mathcal{N}$  (or  $\mathcal{M}$ ) is unimodular.*

- (i) *The algebra  $\underline{\text{Nat}}(H, H)$  has the structure of a symmetric Frobenius algebra in  $\mathcal{Z}(\mathcal{C})$  for any  $H \in \text{Func}(\mathcal{M}, \mathcal{N})$ .*
- (ii)  *$\underline{\text{Nat}}(\text{id}_{\mathcal{M}}, \text{id}_{\mathcal{M}})$  has the structure of a commutative symmetric Frobenius algebra in  $\mathcal{Z}(\mathcal{C})$ .*
- (iii) *The functors  $\Psi_{\mathcal{N}}^{ra}$  and  $\Phi_{\mathcal{M}}^{ra}$  are pivotal Frobenius monoidal.*

*Proof.* By Lemma 6.6,  $\underline{\text{Nat}}(H, H) = \Psi_{\mathcal{N}}^{ra}(H \circ H^{\text{la}})$ . So part (i) follows by combining Proposition 6.13(ii) which tells that  $H \circ H^{\text{la}}$  is symmetric Frobenius and Theorem 4.26(i) which tells that  $\Psi_{\mathcal{N}}^{ra}$  preserves symmetric Frobenius algebras. Part (ii) follows in a similar manner using Theorem 4.26(ii). Lastly, by [Yad24, Thm. 1.2(ii)], part (ii) implies part (iii).  $\square$

Under the additional requirement that  $\mathcal{C}$  is unimodular and spherical, any pivotal  $\mathcal{C}$ -module category is unimodular [FGJS22, Thm. 5.18].

**Corollary 6.16.** *Let  $\mathcal{C}$  be a spherical (unimodular) tensor category and  $\mathcal{M}, \mathcal{N}$  be pivotal  $\mathcal{C}$ -module categories. Then  $\text{Func}(\mathcal{M}, \mathcal{N})$  is a pivotal  $\mathcal{Z}(\mathcal{C})$ -module category.*

*Proof.* According to [FGJS22, Thm. 5.18],  $\mathcal{C}_{\mathcal{N}}^*$  and  $\mathcal{C}_{\mathcal{M}}^*$  are automatically unimodular under unimodularity and sphericity of  $\mathcal{C}$ . Thus,  $\mathcal{N}$  is unimodular as a  $\mathcal{C}$ -module category and the claim follows by Proposition 6.14.  $\square$

*Remark 6.17.* In the situation of Corollary 6.16, a trivialization  $\tilde{u}_{\mathcal{C}} : \mathbb{1} \xrightarrow{\sim} D_{\mathcal{C}}$  together with the pivotal structure of  $\mathcal{M}$  form a unimodular structure on  $\mathcal{M}$  via the composition

$$\tilde{u}_{\mathcal{M}}^{\tilde{p}} : \text{id}_{\mathcal{M}} \xrightarrow{\tilde{p}^2} (\mathbb{S}_{\mathcal{M}}^{\mathcal{C}})^2 \xrightarrow{\tilde{u}_{\mathcal{C}} \triangleright \text{id}} D_{\mathcal{C}}^{-1} \triangleright (\mathbb{S}_{\mathcal{M}}^{\mathcal{C}})^2 \xrightarrow{(2.13)} D_{\mathcal{C}_{\mathcal{M}}^*}^{-1}.$$

With this choice of unimodular structure,  $\mathcal{M}$  is  $(\tilde{u}_{\mathcal{C}}, \tilde{u}_{\mathcal{M}}^{\tilde{p}})$ -spherical. Moreover, the pivotal  $(\mathcal{C}_{\mathcal{N}}^*, \mathcal{C}_{\mathcal{M}}^*)$ -bimodule category  $\text{Func}(\mathcal{M}, \mathcal{N})$  becomes  $(\tilde{u}_{\mathcal{N}}^{\tilde{q}}, \tilde{u}_{\mathcal{M}}^{\tilde{p}})$ -spherical. It follows from Proposition 4.33 that  $\text{Func}(\mathcal{M}, \mathcal{N})$  is  $(\tilde{u}_{\mathcal{N}}^{\tilde{q}}, \tilde{u}_{\mathcal{M}}^{\tilde{p}})$ -spherical as a  $\mathcal{Z}(\mathcal{C})$ -bimodule category.

### 6.3. Examples from Hopf algebras

Let  $(H, m, u, \Delta, \varepsilon, S)$  be a finite-dimensional Hopf algebra over  $\mathbb{k}$ , and  $L$  a left  $H$ -comodule algebra. Setting  $\mathcal{C} = \text{Rep}(H)$  and  $\mathcal{N} = \text{Rep}(L)$ , we get that  $\mathcal{N}$  is a left  $\mathcal{C}$ -module category. From here on, we assume that  $L$  is *exact* (that is, the  $\mathcal{C}$ -module category  $\mathcal{N}$  is exact) and indecomposable.

We first recall some definitions. A (left-left-) *Yetter-Drinfeld  $H$ -module* is a left  $H$ -module  $X$  endowed with a left  $H$ -comodule structure  $\delta$  denoted  $x \mapsto x_{-1} \otimes x_0$ , such that

$$(h \cdot x)_{-1} \otimes (h \cdot x)_0 = h_1 x_{-1} S(h_3) \otimes h_2 x_0$$

holds for all  $h \in H, x \in X$ . We denote by  ${}^H_H \mathcal{YD}$  the category of finite-dimensional Yetter-Drinfeld modules.  ${}^H_H \mathcal{YD}$  inherits a tensor structure defined via

$$(X, \cdot_X, \delta_X) \otimes (Y, \cdot_Y, \delta_Y) := (X \otimes Y, \cdot, \delta)$$

where  $h \cdot (x \otimes y) := h_1 \cdot_X x \otimes h_2 \cdot_Y y$  and  $\delta(x \otimes y) := x_{-1} y_{-1} \otimes x_0 \otimes y_0$ . The unit object is  $(\mathbb{k}, \varepsilon, u)$ . Moreover, the tensor category of Yetter-Drinfeld modules is naturally endowed with a braiding

defined by  $c_{X,Y}(x \otimes y) = x_{-1} \cdot y \otimes x_0$ . Furthermore, there is an equivalence

$$\mathcal{Z}(\mathcal{C}) \xrightarrow{\sim} {}^H_H\mathcal{YD} \quad (6.3)$$

of braided tensor categories [EGNO16, Ex. 8.5.6].

Let  ${}^H\mathfrak{M}$  denote the tensor category of finite-dimensional  $H$ -comodules and  ${}^H_L\mathfrak{M}_L$  the category of  $L$ -bimodules in  ${}^H\mathfrak{M}$ . Explicitly, an object of  ${}^H_L\mathfrak{M}_L$  is a finite-dimensional  $L$ -bimodule  $(P, \prec, \succ)$  equipped with a left  $H$ -coaction  $\delta : P \rightarrow H \otimes P$  obeying that

$$\delta(apb) = a_{-1}p_{-1}b_{-1} \otimes a_0p_0b_0$$

for all  $a, b \in L$  and  $p \in P$ . The category  ${}^H_L\mathfrak{M}_L$  is endowed with a tensor structure given by

$$(P, \prec_P, \succ_P, \rho_P) \otimes (Q, \prec_Q, \succ_Q, \rho_Q) := (P \otimes_L Q, \prec, \succ, \rho)$$

where  $l \succ (p \otimes_L q) \prec l' := (l \succ_P p) \otimes_L (q \prec_Q l')$  and  $\rho(p \otimes_L q) := p_{-1}q_{-1} \otimes (p_0 \otimes_L q_0)$ . This tensor category realizes the dual tensor category of  $L$ . More precisely, there is an equivalence of tensor categories [AM07, Prop. 1.23]

$$\mathcal{C}_{\mathcal{N}}^* = \text{Func}_{\mathcal{C}}(\mathcal{N}, \mathcal{N}) \xrightarrow{\sim} {}^H_L\mathfrak{M}_L \quad (6.4)$$

where  $\mathcal{C} = \text{Rep}(H)$  and  $\mathcal{N} = \text{Rep}(L)$ .

Any  $P \in {}^H_L\mathfrak{M}_L$  is in particular a finitely generated projective left  $L$ -module. Thus, there exist elements  $b_1, \dots, b_k \in P$  and  $b^1, \dots, b^k \in \text{Hom}_L(P, L)$  such that for any  $p \in P$ , we have that  $b^i(p) \cdot b_i = p$ , where we have used Einstein's summation convention.

**Proposition 6.18.** *Let  $H$  be a finite-dimensional Hopf algebra, and  $L$  an exact  $H$ -comodule algebra. For every  $P \in {}^H_L\mathfrak{M}_L$ , the object  ${}^\dagger P$  given by*

- The vector space  $\text{Hom}_L(P, L)$ .
- The  $L$ -bimodule structure defined by the formula

$$(l \succ f \prec l')(p) := f(p \prec_P l) \cdot l'.$$

- The  $H$ -coaction  $\delta : {}^\dagger P \rightarrow H \otimes {}^\dagger P$  denoted  $\delta(f) = f_{-1} \otimes f_0$  and defined via

$$f_{-1} \otimes f_0(p) := S(p_{-1})f(p_0)_{-1} \otimes f(p_0)_0,$$

is left dual to  $P$  in the tensor category  ${}^H_L\mathfrak{M}_L$ , with

$$\begin{aligned} \widetilde{\text{ev}}_P : P \otimes_L {}^\dagger P &\longrightarrow L & \text{and} & & \widetilde{\text{coev}}_P : L &\longrightarrow {}^\dagger P \otimes_L P \\ p \otimes f &\longmapsto f(p) & & & l &\longmapsto b^i \otimes b_i \prec_P l \end{aligned} \quad (6.5)$$

as evaluation and coevaluation morphisms, where  $b^i$  and  $b_i$  is a pair of dual bases over  $L$ .

*Proof.* By [Shi23a, Lemma 4.8], the above data yields an object of  ${}^H_L\mathfrak{M}_L$ . Note that in [Shi23a] this object is denoted  $P^\dagger$ . Moreover, in [Shi23a, Rem. 4.9] it is pointed out that the functor  ${}^\dagger(-)$  is the counterpart of taking right-adjoints, up to the equivalence (6.4). Since, in our setting, (6.4) is a tensor equivalence, duals are preserved leading to the result.  $\square$

The next key ingredient in the study of internal natural transformations is the tensor functor  $\Psi_{\mathcal{N}}$  from (6.1). Under the equivalences (6.3) and (6.4) this tensor functor descends to

$$\Psi : {}^H_H\mathcal{YD} \longrightarrow {}^H_L\mathfrak{M}_L, \quad (X, \cdot, \delta) \longmapsto (X \otimes L, \prec, \succ, \rho) \quad (6.6)$$



where  $L$  acts on the left and right of  $X \otimes L$  via  $s \succ (x \otimes l) \prec t := s_{-1} \cdot x \otimes s_0 l t$ . The left  $H$ -comodule structure on  $X \otimes L$  is given by  $\rho(m \otimes l) := m_{-1} l_{-1} \otimes m_0 \otimes l_0$ . The tensor structure on the functor (6.6) is explicitly given by the natural transformation

$$\Psi_2(X, Y) : \begin{array}{ccc} \Psi(X) \otimes_L \Psi(Y) = (X \otimes L) \otimes_L (Y \otimes L) & \rightarrow & X \otimes Y \otimes L = \Psi(X \otimes Y) \\ x \otimes l \otimes_L y \otimes l' & \mapsto & x \otimes l_{-1} \cdot y \otimes l_0 l' \end{array}, \quad (6.7)$$

and the isomorphism  $\Psi_0$  defined as

$$\Psi_0 : \begin{array}{ccc} \mathbb{1}_{L^H \mathfrak{M}_L} = L & \longrightarrow & \mathbb{k} \otimes L = \Psi(\mathbb{1}_{H^H \mathcal{YD}}) \\ l & \longmapsto & 1 \otimes l \end{array}.$$

One can verify that  $\Psi_2$  is invertible with inverse  $\Psi_2(X, Y)^{-1}(x \otimes y \otimes l) := x \otimes 1_L \otimes_L y \otimes l$ . A routine check shows that  $(\Psi, \Psi_2, \Psi_0)$  is indeed a strong monoidal functor.

The right adjoint functor to (6.6)

$$\Psi^{\text{ra}} : \begin{array}{ccc} H^H \mathfrak{M}_L & \longrightarrow & H^H \mathcal{YD} \end{array} \quad (6.8)$$

finds an explicit description in the following theorem from [BM21].

**Theorem 6.19.** [BM21, Thm. 4.11] *For every  $(P, \prec, \succ, \rho) \in H^H \mathfrak{M}_L$ , there is an isomorphism*

$$\Psi^{\text{ra}}(P) \cong (S^L(H, P), *, \rho)$$

of Yetter-Drinfeld modules, where

- The vector space  $S^L(H, P) := \{f \in \text{Hom}_L(H \otimes L, P) \mid f(h \otimes l) = f(h \otimes 1) \prec l, \forall h \in H, l \in L\}$ .
- The  $H$ -action  $*$  :  $H \otimes S^L(H, P) \rightarrow S^L(H, P)$  is defined by the formula

$$h * f(h' \otimes l) := f(h' h \otimes l).$$

- The  $H$ -coaction  $\delta : S^L(H, P) \rightarrow H \otimes S^L(H, P)$ , denoted by  $\delta(f) = f_{-1} \otimes f_0$ , is defined by

$$f_{-1} \otimes f_0(h \otimes l) := S(h_1) f(h_2 \otimes 1)_{-1} h_3 \otimes f(h_2 \otimes 1)_0 \cdot l.$$

For our purposes, we will use a more convenient description of the functor  $\Psi^{\text{ra}}$ .

**Definition 6.20.** Let  $H$  be a finite-dimensional Hopf algebra, and  $L$  an exact  $H$ -comodule algebra. For every  $P \in H^H \mathfrak{M}_L$ , we define the triple  $(T^L(H, P), \star, \rho)$  consisting of:

- The underlying vector space  $\{g \in \text{Hom}(H, P) \mid l \succ g(h) = g(l_{-1} h) \prec l_0, \forall l \in L\}$ .
- The  $H$ -module structure  $\star : H \otimes T^L(H, P) \rightarrow T^L(H, P)$  defined via

$$(h \star f)(h') := f(h' h).$$

- The  $H$ -comodule structure  $\rho : T^L(H, P) \rightarrow H \otimes T^L(H, P)$  given by

$$\rho(f)(h) \equiv f_{-1} \otimes f_0(h) := S(h_1) f(h_2)_{-1} h_3 \otimes f(h_2)_0.$$

The construction from Definition 6.20 extends to a functor

$$T^L(H, -) : \begin{array}{ccc} H^H \mathfrak{M}_L & \longrightarrow & H^H \mathcal{YD} \end{array} \quad (6.9)$$

which turns out to be right-adjoint to (6.8), as well.

**Proposition 6.21.** *For  $P \in {}^H_L\mathfrak{M}_L$ , there is an isomorphism*

$$\Psi^{\text{ra}}(P) \cong T^L(H, P)$$

*of Yetter-Drinfeld modules.*

*Proof.* The vector spaces  $T^L(H, P)$  and  $S^L(H, P)$  are in bijection via the maps

$$\begin{aligned} \alpha : S^L(H, P) &\rightarrow T^L(H, P) & \beta : T^L(H, P) &\rightarrow S^L(H, P) \\ \alpha(f)(h) &:= f(h \otimes 1_L) & \beta(g)(h \otimes l) &:= g(h) \prec l. \end{aligned}$$

It is straightforward to see that  $\alpha, \beta$  are well-defined. Further, for  $f \in S^L(H, P)$ ,

$$\beta(\alpha(f))(h \otimes l) = \alpha(f)(h) \prec l = f(h \otimes 1_L) \prec l = f(h \otimes l).$$

Similarly, for  $g \in T^L(H, P)$ , we have that

$$\alpha(\beta(g))(h) = \beta(g)(h \otimes 1_L) = g(h) \prec 1_L = g(h).$$

Therefore,  $\alpha$  and  $\beta$  are mutual inverses. Next, we show that  $\star$  is obtained by transporting  $*$  along the isomorphism  $\alpha$ .

$$\alpha(h * f)(h') = (h * f)(h' \otimes 1_L) = f(h'h \otimes 1_L) = \alpha(f)(h'h) = [h * \alpha(f)](h').$$

Similarly, the following calculation shows that  $\rho$  is obtained by transporting  $\delta$  along  $\alpha$ :

$$\begin{aligned} \alpha(f)_{-1} \otimes \alpha(f)_0(h) &= S(h_1)\alpha(f)(h_2)_{-1}h_3 \otimes \alpha(f)(h_2)_0 \\ &= S(h_1)f(h_2 \otimes 1_L)_{-1}h_3 \otimes f(h_2 \otimes 1_L)_0 \\ &= f_{-1} \otimes f_0(h \otimes 1_L) = f_{-1} \otimes \alpha(f)(h). \end{aligned}$$

Thus,  $T^L(H, P)$  is a Yetter-Drinfeld module and it is isomorphic to  $S^L(H, P)$ . By Theorem 6.19, the assertion follows.  $\square$

**Corollary 6.22.** *The object of internal natural transformations from  $P \in {}^H_L\mathfrak{M}_L$  to  $Q \in {}^H_L\mathfrak{M}_L$  is given by*

$$\underline{\text{Nat}}(P, Q) \cong T^L(H, {}^\dagger P \otimes_L Q)$$

*as a Yetter-Drinfeld module.*

*Proof.* It follows from Lemma 6.6 and Proposition 6.21.  $\square$

The tensor structure (6.7) of the functor  $\Psi$  endows its right adjoint functor (6.9) with a lax monoidal structure. The following proposition provides an explicit description.

**Proposition 6.23.** *The functor  $T^L(H, -)$  inherits a lax monoidal structure with constraints*

$$\begin{aligned} \mu_2(P, Q) : T^L(H, P) \otimes T^L(H, Q) &\longrightarrow T^L(H, P \otimes_L Q), & \mu_0 : \mathbb{k} &\rightarrow T^L(H, L) \\ f \otimes g &\longmapsto f \diamond g & 1 &\mapsto \varepsilon_H 1_L \end{aligned}$$

*where  $(f \diamond g)(h) = f(h_1) \otimes_L g(h_2)$  for all  $P, Q \in {}^H_L\mathfrak{M}_L$ .*

*Proof.* Using [BM21, Prop. 4.9], we get that the component maps of the unit of the adjunction  $\Psi \dashv \Psi^{\text{ra}}$  are given by

$$\eta_X^r : (X, \cdot, \delta) \rightarrow \Psi^{\text{ra}}\Psi(X) = (S^L(H, X \otimes L), *, \rho) \quad \text{where,} \quad \eta_X^r(x)(h \otimes l) := h \cdot x \otimes l.$$

Similarly, the components of the counit are given by

$$\varepsilon_P^r: \Psi \Psi^{\text{ra}}(P) = S^L(H, P) \otimes L \rightarrow P \quad \text{where,} \quad \varepsilon_P^r(\alpha \otimes l) = \alpha(1_H \otimes l).$$

The lax monoidal constraints  $\mu_2$  and  $\mu_0$  of  $\Psi^{\text{ra}}$  are defined as follows: for  $P, Q \in {}^H_L \mathfrak{M}_L$ , the component maps  $\mu_2(P, Q)$  are the compositions

$$\mu_2(P, Q) = \Psi^{\text{ra}}(\varepsilon_P^r \otimes_L \varepsilon_Q^r) \circ \Psi^{\text{ra}} \Psi_2^{-1}(\Psi^{\text{ra}}(P), \Psi^{\text{ra}}(Q)) \circ \eta_{\Psi^{\text{ra}}(P) \otimes \Psi^{\text{ra}}(Q)}^r$$

Explicitly, for  $f \in T^L(H, P)$  and  $g \in T^L(H, Q)$ , we get that

$$\begin{aligned} f \otimes g & \xrightarrow{\eta_{\Psi^{\text{ra}}(P) \otimes \Psi^{\text{ra}}(Q)}^r} [h \mapsto h \cdot (f \otimes g) \otimes 1_L = h_1 \cdot f \otimes h_2 \cdot g \otimes 1_L] \\ & \xrightarrow{\Psi^{\text{ra}} \Psi_2^{-1}(\Psi^{\text{ra}}(P), \Psi^{\text{ra}}(Q))} [h \mapsto (h_1 \cdot f \otimes 1_L) \otimes_L (h_2 \cdot g \otimes 1_L)] \\ & \xrightarrow{\Psi^{\text{ra}}(\varepsilon_P^r \otimes_L \varepsilon_Q^r)} [h \mapsto (h_1 \cdot f)(1_L) \prec 1_L \otimes_L (h_2 \cdot g)(1_L) \prec 1_L] \\ & = [h \mapsto f(h_1) \otimes_L g(h_2) =: f \diamond g] \end{aligned}$$

We also obtain that

$$\mu_0 = \Psi^{\text{ra}}(\Psi_0^{-1}) \circ \eta_{\mathbb{1}}^r: \mathbb{1}_{{}^H_H \mathcal{YD}} = \mathbb{k} \longrightarrow T^L(H, L) = \Psi^{\text{ra}}(L).$$

which, by the definitions of  $\Psi_0$  and  $\eta^r$ , is explicitly given by  $\mu_0 1(h) = \varepsilon(h) 1_L$ .  $\square$

The objects of internal endonatural transformations inherit the structure of an algebra internal to  ${}^H_H \mathcal{YD}$  in the following way:

**Proposition 6.24.** *For  $P \in {}^H_L \mathfrak{M}_L$ , the object  $\text{Nat}(P, P) \cong T^L(H, \dagger P \otimes_L P)$  is endowed with the structure of an algebra with multiplication map*

$$\begin{aligned} m_P: T^L(H, \dagger P \otimes_L P) \otimes T^L(H, \dagger P \otimes_L P) & \longrightarrow T^L(H, \dagger P \otimes_L P) \\ f \otimes g & \longmapsto f \cdot g \end{aligned} \tag{6.10}$$

where

$$f \cdot g(h) := (\text{id}_{\dagger P} \otimes_L \widetilde{\text{ev}}_P \otimes_L \text{id}_P) \circ (f \diamond g)(h) = f(h_1)^i \otimes_L \widetilde{\text{ev}}_P(f(h_1)_i \otimes_L g(h_2)^j) \otimes_L g(h_2)_j.$$

Here  $p^i \otimes_L p_j$  denotes a decomposition in pure tensors for  $p \in \dagger P \otimes_L P$ .

*Proof.* The object  $\dagger P \otimes_L P$  is an algebra in  ${}^H_L \mathfrak{M}_L$  by means of the evaluation and coevaluation morphisms (6.5). By transporting this algebra structure along the lax monoidal functor  $T^L(H, -)$  from Proposition 6.23 we obtain the multiplication on internal natural transformations.  $\square$

In the particular case of  $P = L$  the monoidal unit in  ${}^H_L \mathfrak{M}_L$ . We obtain on the Yetter-Drinfeld module

$$T^L(H, L) = \{f \in \text{Hom}(H, L) \mid lf(h) = f(l_{-1}h)l_0, \forall h \in H, l \in L\}.$$

It is an algebra with the multiplication map given by

$$\begin{aligned} m_L: T^L(H, L) \otimes T^L(H, L) & \longrightarrow T^L(H, L) \\ f \otimes g & \longmapsto (h \mapsto f(h_1)g(h_2)) \end{aligned}$$

and the unit map  $u: \mathbb{k} \rightarrow T^L(H, L)$ ,  $1 \mapsto \varepsilon_H(-) 1_L$ . In fact,  $(T^L(H, L), m, u)$  is a commutative algebra in  ${}^H_H \mathcal{YD}$ .

**Proposition 6.25.** *Let  $L$  be a unimodular exact left  $H$ -comodule algebra,  $T^L(H, L)$  is a commutative Frobenius algebra in  ${}^H_H\mathcal{YD}$ . If, in addition,  $H$  is pivotal and  $L$  is pivotal,  $T^L(H, L)$  is a commutative symmetric Frobenius algebra in  ${}^H_H\mathcal{YD}$ .*

*Proof.* The statement is an immediate consequence of Proposition 6.11(ii) and Corollary 6.15(ii).  $\square$

*Remark 6.26.* Suppose that  $L$  is an indecomposable exact left  $H$ -comodule algebra. Then

$$\mathrm{Hom}_{\mathcal{Z}(\mathcal{C})}(\Psi^{\mathrm{ra}}(L), \mathbf{1}_{\mathcal{Z}(\mathcal{C})}) \cong \mathrm{Hom}_{\mathcal{E}\mathfrak{M}_L}(L, \Psi(\mathbf{1}_{\mathcal{Z}(\mathcal{C})})) \cong \mathrm{Hom}_{\mathcal{E}\mathfrak{M}_L}(L, L) \cong \mathbb{k}$$

where the last isomorphism holds because  $L$  is indecomposable. When  $L$  is unimodular,  $T^L(H, L)$  is a Frobenius algebra and its Frobenius form belongs to  $\mathrm{Hom}_{\mathcal{Z}(\mathcal{C})}(\Psi^{\mathrm{ra}}(L), \mathbf{1}_{\mathcal{Z}(\mathcal{C})})$ . Also note that any scalar multiple of a Frobenius form is also a Frobenius form. Thus, any nonzero map  $\lambda': \Psi^{\mathrm{ra}}(L) = T^L(H, L) \rightarrow \mathbb{k} \in {}^H_H\mathcal{YD}$  will be a Frobenius form.

**Question 6.27.** *How is the Frobenius form of the algebra  $T^L(H, L)$  explicitly described in terms of the unimodular element  $g_{\mathrm{uni}}$  of a unimodular left  $H$ -comodule algebra  $L$ ?*

*Example 6.28.* The algebra  $L = \mathbb{k}$  is a left  $H$ -comodule algebra with the trivial coaction  $1 \mapsto 1_H \otimes 1$ . For simplicity, denote  $\hat{H} \equiv T^{\mathbb{k}}(H, \mathbb{k})$ . As an object of  ${}^H_H\mathcal{YD}$ , we have that  $\hat{H} \cong H^*$ , with left  $H$ -action  $\star$  given by  $(h \star f)(h') = f(h'h)$ . The left  $H$ -coaction  $\rho$  of  $\hat{H}$  is given by  $f \mapsto f_{-1} \otimes f_0$  where,

$$f_{-1}f_0(h) = S(h_1)f(h_2)_{-1}h_3f(h_2)_0 = S(h_1)f(h_2)h_3.$$

Consider the functor  $\Psi : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}_{\mathcal{N}}^*$ . For  $\mathcal{N} = \mathrm{Vec}_{\mathbb{k}}$ ,  $\mathcal{C}_{\mathcal{N}}^* \cong \mathrm{Rep}(H^*)$ . Thus, by [Yad23, Theorem 3.15], we have that  $\Psi^{\mathrm{ra}}(\mathrm{id}_{\mathcal{N}})$  is a Frobenius algebra in  $\mathcal{Z}(\mathcal{C}) = {}^H_H\mathcal{YD}$  if and only if  $\mathrm{Rep}(H^*)$  is unimodular if and only if  $H^*$  is a unimodular Hopf algebra. Any nonzero  $\lambda_{\hat{H}} : \hat{H} \rightarrow \mathbb{k}$  is given by  $\lambda_{\hat{H}}(f) = f(h)$  for some  $h' \in H$ . Additionally,  $\lambda_{\hat{H}}$  has to be a right  $H$ -module map. Namely,

$$\langle f, h'h \rangle = \langle h \star f, h' \rangle = \langle \lambda_{\hat{H}}, h \star f \rangle = \varepsilon(h) \langle \lambda_{\hat{H}}, f \rangle = \varepsilon(h) \langle f, \Lambda \rangle \quad \forall h \in H.$$

This forces  $h'$  to be a right integral of  $H$ . As the space of right integrals is 1-dimensional, using Remark 6.26, the Frobenius form is given by  $\lambda_{\hat{H}}(f) = f(\Lambda)$  where  $\Lambda$  is a right integral of  $H$ .

*Example 6.29.*  $L = H$  is also a left  $H$ -comodule algebra with coaction  $h \mapsto h_1 \otimes h_2$ . Then

$$T^H(H, H) = \{f : H \rightarrow H \mid hf(h') = f(h_1h')h_2\}$$

Note that  $T^H(H, H)$  is Frobenius if and only if  $\mathcal{C}_{\mathcal{C}}^* \cong \mathcal{C}$  is unimodular if and only if  $H$  is unimodular. In this case, a Frobenius form is given by  $\langle \lambda', f \rangle = \langle \lambda, f(1_H) \rangle$  where  $\lambda$  is a right cointegral of  $H$ . This is because  $\lambda'$  satisfies  $\langle \lambda', h \star f \rangle = \varepsilon(h) \langle \lambda', h \rangle$  and  $f_{-1} \langle \lambda', f_0 \rangle = 1_H \langle \lambda', f \rangle$  making it a morphism in  ${}^H_H\mathcal{YD}$ .

## REFERENCES

- [AM07] Nicolás Andruskiewitsch and Juan Martín Mombelli. On module categories over finite-dimensional Hopf algebras. *Journal of Algebra*, 314(1):383–418, 2007.
- [Bal17] Adriana Balan. On Hopf adjunctions, Hopf monads and Frobenius-type properties. *Applied Categorical Structures*, 25(5):747–774, 2017.
- [BLV11] Alain Bruguières, Steve Lack, and Alexis Virelizier. Hopf monads on monoidal categories. *Advances in Mathematics*, 227(2):745–800, 2011.

- [BM21] Noelia Bortolussi and Martín Mombelli. The character algebra for module categories over Hopf algebras. In *Colloquium Mathematicum*, volume 165, pages 171–197. Instytut Matematyczny Polskiej Akademii Nauk, 2021.
- [BN11] Alain Bruguières and Sonia Natale. Exact sequences of tensor categories. *International Mathematics Research Notices*, 2011(24):5644–5705, 2011.
- [BN14] Alain Bruguières and Sonia Natale. Central exact sequences of tensor categories, equivariantization and applications. *Journal of the Mathematical Society of Japan*, 66(1):257–287, 2014.
- [CMZ97] S Caenepeel, G Militaru, and Shenglin Zhu. Doi-Hopf modules, Yetter-Drinfel’d modules and Frobenius type properties. *Transactions of the American Mathematical Society*, 349(11):4311–4342, 1997.
- [Del07] Pierre Deligne. Catégories tannakiennes. *The Grothendieck Festschrift: A Collection of Articles Written in Honor of the 60th Birthday of Alexander Grothendieck*, pages 111–195, 2007.
- [DKR15] Alexei Davydov, Liang Kong, and Ingo Runkel. Functoriality of the center of an algebra. *Advances in Mathematics*, 285:811–876, 2015.
- [DP08] Brian Day and Craig Pastro. Note on Frobenius monoidal functors. *New York J. Math*, 14:733–742, 2008.
- [DSPS18] Christopher Douglas, Christopher Schommer-Pries, and Noah Snyder. Dualizable tensor categories. *Memiors of the American Mathematical Society*, 2018.
- [DSPS19] Christopher Douglas, Christopher Schommer-Pries, and Noah Snyder. The balanced tensor product of module categories. *Kyoto Journal of Mathematics*, 59(1):167–179, 2019.
- [EG17] Pavel Etingof and Shlomo Gelaki. Exact sequences of tensor categories with respect to a module category. *Advances in Mathematics*, 308:1187–1208, 2017.
- [EGNO16] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*, volume 205. American Mathematical Soc., 2016.
- [ENO05] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik. On fusion categories. *Annals of Mathematics*, pages 581–642, 2005.
- [EO04] Pavel Etingof and Viktor Ostrik. Finite tensor categories. *Moscow Mathematical Journal*, 4(3):627–654, 2004.
- [FGJS22] Jürgen Fuchs, Cesar Galindo, David Jaklitsch, and Christoph Schweigert. Spherical Morita contexts and relative Serre functors. *Kyoto J. Math.*, 2022. [arXiv:2207.07031](https://arxiv.org/abs/2207.07031).
- [FGJS24] Jürgen Fuchs, Cesar Galindo, David Jaklitsch, and Christoph Schweigert. A manifestly Morita-invariant construction of Turaev-Viro invariants. *preprint*, 2024. [arXiv:2407.10018](https://arxiv.org/abs/2407.10018).
- [FLP24a] Johannes Flake, Robert Laugwitz, and Sebastian Posur. Frobenius monoidal functors from ambidjunctions and their lifts to Drinfeld centers. *preprint*, 2024. [arXiv:2410.08702](https://arxiv.org/abs/2410.08702).
- [FLP24b] Johannes Flake, Robert Laugwitz, and Sebastian Posur. Frobenius monoidal functors induced by Frobenius extensions of Hopf algebras. *preprint*, 2024. [arXiv:2412.15056](https://arxiv.org/abs/2412.15056).
- [FLP24c] Johannes Flake, Robert Laugwitz, and Sebastian Posur. Projection formulas and induced functors on centers of monoidal categories. *preprint*, 2024. [arXiv:2402.10094](https://arxiv.org/abs/2402.10094).
- [FMS97] Davida Fischman, Susan Montgomery, and Hans-Jürgen Schneider. Frobenius extensions of subalgebras of Hopf algebras. *Transactions of the American Mathematical Society*, 349(12):4857–4895, 1997.
- [FRS02] Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. TFT construction of RCFT correlators I: Partition functions. *Nuclear Physics B*, 646(3):353–497, 2002.
- [FS21a] Jürgen Fuchs and Christoph Schweigert. Bulk from boundary in finite CFT by means of pivotal module categories. *Nuclear Physics B*, 967:115392, 2021.
- [FS21b] Jürgen Fuchs and Christoph Schweigert. Internal natural transformations and Frobenius algebras in the Drinfeld center. *Transformation Groups*, pages 1–36, 2021.
- [FSS20] Jürgen Fuchs, Gregor Schaumann, and Christoph Schweigert. Eilenberg-Watts calculus for finite categories and a bimodule Radford  $S^4$  theorem. *Transactions of the American Mathematical Society*, 373(1):1–40, 2020.
- [Lau06] Aaron Lauda. Frobenius algebras and ambidextrous adjunctions. *Theory and Applications of Categories*, 16(4):84–122, 2006.
- [Maj91] Shahn Majid. Representations, duals and quantum doubles of monoidal categories. In *Proceedings of the Winter School "Geometry and Physics"*, pages 197–206. Circolo Matematico di Palermo, 1991.

- [NS07] Siu-Hung Ng and Peter Schauenburg. Higher Frobenius-Schur indicators for pivotal categories. *Hopf algebras and generalizations*, 441:63–90, 2007.
- [NZ89] Warren D Nichols and M Bettina Zoeller. A Hopf algebra freeness theorem. *American Journal of Mathematics*, 111(2):381–385, 1989.
- [Ost03] Victor Ostrik. Module categories, weak hopf algebras and modular invariants. *Transformation groups*, 8:177–206, 2003.
- [Rad11] David E Radford. *Hopf algebras*, volume 49. World Scientific, 2011.
- [Sch13] Gregor Schaumann. Traces on module categories over fusion categories. *Journal of Algebra*, 379:382–425, 2013.
- [Sch15] Gregor Schaumann. Pivotal tricategories and a categorification of inner-product modules. *Algebras and Representation Theory*, 18(6):1407–1479, 2015.
- [Shi16] Kenichi Shimizu. On unimodular finite tensor categories. *International Mathematics Research Notices*, 2017(1):277–322, 2016.
- [Shi17] Kenichi Shimizu. The relative modular object and Frobenius extensions of finite Hopf algebras. *Journal of Algebra*, 471:75–112, 2017.
- [Shi20] Kenichi Shimizu. Further results on the structure of (co) ends in finite tensor categories. *Applied Categorical Structures*, 28(2):237–286, 2020.
- [Shi23a] Kenichi Shimizu. Relative Serre functor for comodule algebras. *Journal of Algebra*, 634:237–305, 2023.
- [Shi23b] Kenichi Shimizu. Ribbon structures of the Drinfeld center of a finite tensor category. *Kodai Mathematical Journal*, 46(1):75–114, 2023.
- [Shi24] Kenichi Shimizu. Quasi-frobenius algebras in finite tensor categories. *preprint*, 2024. [arXiv:2402.02929](https://arxiv.org/abs/2402.02929).
- [Skr07] Serge Skryabin. Projectivity and freeness over comodule algebras. *Transactions of the American Mathematical Society*, 359(6):2597–2623, 2007.
- [Str04] Ross Street. Frobenius monads and pseudomonoids. *Journal of mathematical physics*, 45(10):3930–3948, 2004.
- [SY24] Kenichi Shimizu and Harshit Yadav. Commutative exact algebras and modular tensor categories. *preprint*, 2024. [arXiv:2408.06314](https://arxiv.org/abs/2408.06314).
- [Yad23] Harshit Yadav. On unimodular module categories. *Advances in Mathematics*, 432:109264, 2023.
- [Yad24] Harshit Yadav. Frobenius monoidal functors from (co)Hopf adjunctions. *Proceedings of the American Mathematical Society*, 152(02):471–487, 2024.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, 0316 OSLO, NORWAY  
*Email address:* [dajak@uio.no](mailto:dajak@uio.no)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, T6G 2G1, CANADA  
*Email address:* [hyadav3@ualberta.ca](mailto:hyadav3@ualberta.ca)