

ON THE REGULARITY OF NONDEGENERATE HYPO-ANALYTIC STRUCTURES OF HYPERSURFACE TYPE

ILYA KOSOVSKIY AND VINÍCIUS NOVELLI

ABSTRACT. For a smooth, non-degenerate locally integrable structure of hypersurface type on a manifold M , we provide necessary and sufficient conditions for it to be equivalent, near a point, to a real-analytic locally integrable structure (*the analytic regularizability*), generalizing a recent result of Zaitsev and the first author [14].

First, we discover, in our setting, a (previously unknown) invariant CR submanifold Σ in M of hypersurface type, which we call *the central submanifold*. We prove that the analytic regularizability of M is equivalent to that of the associated CR manifold Σ . Furthermore, as a byproduct of our construction, we show that the central manifold construction reduces the whole (smooth or analytic) equivalence problem for nondegenerate structures with the Levi positivity condition to that of the associated central manifolds, i.e. to CR geometry.

Second, we make use of a classical construction due to Marson [15] and show that sufficient for the analytic regularizability of M is the analytic regularizability of the CR manifold \tilde{M} associated with M in the sense of Marson.

We show applications of both regularizability conditions to classes of locally integrable structures.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. Nondegenerate locally integrable structures.	3
2.2. The method of associated differential equations	5
2.3. Condition (E)	6
3. Special coordinates for non-degenerate structures	7
4. The central manifold and the regularizability	10
5. Additional sufficient conditions: The external CR structure	13
6. An application to rigid structures	15
References	16

1. INTRODUCTION

This work is concerned with *locally integrable structures* of hypersurface type. These are locally integrable systems of complex vector fields on a smooth manifold for which the rank of the characteristic set does not exceed one. Within this class are, in particular, CR structures on real hypersurfaces in a complex manifold and (locally integrable) Mizohata structures. The theory of locally integrable structures has a long and rich history (see for instance the classical sources Berhanu-Cordaro-Hounie [4], Treves [18], Cordaro-Treves [8], Baouendi-Chang-Treves [1]). Such structures can be of several regularity levels (C^k , smooth, Gevrey, real-analytic), and as a rule, the higher the regularity is, the more techniques are available to understand them. For example, in the real-analytic case we have such powerful techniques as the *external complexification*, *Segre varieties* (see, for example, Baouendi-Ebenfelt-Rothschild [2]), and *normal forms* (e.g. Chern-Moser [7], Kolar-Kossovskiy-Zaitsev [11]). The

¹MSC 35F05, 32V05, 32V25

Date: January 28, 2025. *Corresponding author: viniciusnovelli@usp.br.

Key words and phrases. Locally Integrable Structures, CR Geometry, Mizohata Structures.

latter techniques are available solely in the analytic case, which makes the existence of coordinates where a structure becomes real-analytic a crucial property.

This inspires the following widely open question, which is our main concern in this paper:

Question 1.1. *Let (M, \mathcal{V}, p) be a germ of a smooth locally integrable structure (of hypersurface type) near a point $p \in M$. Find necessary and sufficient conditions for \mathcal{V} to be equivalent to a germ of a real-analytic locally integrable structure.*

In the situation when the regularity property from 1.1 holds, we say that \mathcal{V} is *analytically regularizable* near $p \in M$.

We shall point out some subtlety with the notion of equivalence between integrable structures. In this paper, we consider it in the weak sense: an *equivalence between structures* is a germ of a diffeomorphism (of certain regularity) for which the derivative preserves the bundle. There is also the notion of *hypo-analytic equivalence* between these structures (once a hypo-analytic structure is fixed, defined by a system of first integrals). We do not pursue this problem in this paper. These notions will be recalled in Section 2.

For some classes of structures, Question 1.1 is well understood. For instance, if the structure \mathcal{V} is *elliptic* at p (meaning that the characteristic set at this point is trivial), then, by a theorem of Nirenberg [16], one can find coordinates $(x_1, \dots, x_m, y_1, \dots, y_m, t_1, \dots, t_n)$ near p such that a complete system of first integrals for \mathcal{V} is given by $z_j = x_j + iy_j$, $j = 1, \dots, m$. In particular, all such structures are analytically regularizable. Another case where the result is known is that of Mizohata structures (when the locally integrable structure has corank 1), when an argument using Morse's lemma yields coordinates (x, t_1, \dots, t_n) such that the first integral is of the form $x + i\mathcal{L}(t)$, where \mathcal{L} is a non-degenerate quadratic form in t (see Theorem VII.1.1 in [18]). In particular, they are also analytically regularizable.

To the best of our knowledge, for other classes of structures, the regularizability problem as stated in 1.1 was widely open until the recent work [14] by Zaitsev and the first author. They considered the case of smooth strictly pseudoconvex hypersurfaces in \mathbb{C}^n . They introduce a property (called *condition (E)*), which we review in detail in Section 2 and mention now that this property is a holomorphic extension property for certain smooth function associated with a real hypersurface M (*the Φ -function*). The main result of [14] is at a glance the following.

Theorem 1.1 (Kossovskiy, Zaitsev [14]). *Let $M \subset \mathbb{C}^n$ be a smooth, strictly pseudoconvex hypersurface, and let $p \in M$. Then M is CR equivalent near p to a real-analytic hypersurface if and only if M satisfies condition (E).*

The goal of this paper is to extend this result to a larger class of locally integrable structures (of hypersurface type). One can view, through the *local fine embedding* (see Section 2), such a structure as a smooth family of Levi non-degenerate hypersurfaces, and the issue is to find a smooth family of normalizing transformations satisfying the additional equations coming from the extra parameters.

Our technique yields two Levi non-degenerate hypersurfaces associated to \mathcal{V} . One (which is intrinsic) consists of the set of points for which the structure is not elliptic. This submanifold, $\Sigma \subset M$, which we refer to as the *central submanifold of \mathcal{V}* , inherits a structure of the ambient space that of a Levi non-degenerate hypersurface, and yields a necessary condition for regularizability of \mathcal{V} . We state the first main result, and the details can be found in Section 4.

Theorem 1.2. *Let (M, \mathcal{V}, p) be a germ of a nondegenerate locally integrable structure of hypersurface type. Assume the Levi form at p is definite. Then, \mathcal{V} is analytically regularizable if and only if the central CR hypersurface Σ satisfies condition (E) at p .*

On the other hand, one can fix a set of first integrals (induced by a particular system of coordinates) and construct, following the idea of Marson in [15], an associated (external) hypersurface, which in the positive-definite Levi form case is a strictly pseudoconvex CR hypersurface. Our second main result is that regularizability of this external hypersurface entails regularizability of \mathcal{V} . The details can be found in Section 5.

Theorem 1.3. *Let (M, \mathcal{V}, p) be a germ of a nondegenerate locally integrable structure of hypersurface type. Assume that the Levi form at p is definite. Fix a system of coordinates given by the fine local embedding near p and let M^\bullet be the external hypersurface associated to M at p in these coordinates. Then, if M^\bullet satisfies condition (E), then \mathcal{V} is analytically regularizable at p .*

Finally, we outline that, as a bi-product of our considerations, we show the following remarkable property: *if a given nondegenerate structure with a definite Levi form is rigid, then it is analytically regularizable if and only if its central manifold is actually analytic.* This result is new even in the case of CR structures. Details can be found in Section 6.

Remark 1.1. In an upcoming paper of Zaitsev and the first author [10], Condition (E) above is replaced by a series of bounds for certain derivatives of the defining function of the hypersurface. For each given hypersurface, these bounds can be directly verified, which further emphasizes the effectiveness of the regularizability conditions for locally integrable structures introduced in the paper.

Remark 1.2. In fact, the construction of the central manifold associated with a locally integrable structure in our paper gives consequence going much beyond the regularizability problem. Namely (see Theorem 4.3 below), it reduces the *entire* (smooth or analytic) equivalence problem for nondegenerate structures with the Levi positivity to that for the associated central manifolds. The latter makes then possible the application of the whole machinery of CR geometry (e.g. normal forms [7, 11] and methods of *Parabolic Geometries* [5]) for the classification of the involutive structures under discussion.

Acknowledgements

The first author was supported by the Sustech University internal Grant Y01286147, NSFC grant K24281006, Austrian Science Fund Grant P34369, and GACR Grant GC22-15012. The second author was supported by FAPESP, grant 2023/17607-7.

The authors would like to thank Paulo Cordaro for numerous fruitful discussions during the preparation of the paper.

2. PRELIMINARIES

2.1. Nondegenerate locally integrable structures. A *smooth locally integrable structure* is the datum of a smooth N -dimensional manifold M and a smooth complex vector subbundle $\mathcal{V} \subset \mathbb{C}TM$ (of complex rank n) such that the Lie bracket of two local sections is still a section, and such that the orthogonal $T' := \mathcal{V}^\perp \subset \mathbb{C}T^*M$ (for the duality between vector fields and one-forms) is locally generated by exact one-forms. A distribution u defined on an open set of M is a *solution* of \mathcal{V} if du is a local section of T' . A family of complex-valued functions Z_1, \dots, Z_m , where $m = N - n$, is a *complete set of first-integrals* for \mathcal{V} over an open set $U \subset M$ if dZ_1, \dots, dZ_m forms a basis of T' over U .

Given a point $p \in M$, the *characteristic set* of \mathcal{V} at p is the space $T' \cap T_p^*M =: T_p^\circ$ of the real covectors of T' . We say the structure is *of hypersurface type* if

$$\dim_{\mathbb{R}} T_p^\circ \leq 1, \quad p \in M.$$

It is important to note that, in general, the association

$$M \ni p \mapsto T_p^\circ$$

does *not* define a vector bundle over M . This is true in the case of CR structures, but we shall encounter structures (of Mizohata type) for which the dimension of T_p° is not locally constant (it is, however, an upper semicontinuous function of p).

The results of this paper are local, so we will always reason in a neighborhood of a distinguished point, which we henceforth refer to as the origin and denote by 0 . When $\dim T_0^\circ = 0$, the structure is *elliptic* in a neighborhood of 0 , which is not an interesting case for the problems discussed here. We will therefore work under the hypothesis that

$$\dim_{\mathbb{R}} T_0^\circ = 1.$$

We can select a solution w in a neighborhood of 0 such that dw spans T_0° at the origin. Select $m-1 =: \nu$ solutions z_1, \dots, z_ν in some open neighborhood of 0 such that $dz_1 \wedge \dots \wedge dz_\nu \wedge dw \neq 0$ at 0.

It is a matter of linear algebra (see [18] or [4]) to see that this is equivalent to

$$dz_1 \wedge \dots \wedge dz_\nu \wedge d\bar{z}_1 \wedge \dots \wedge d\bar{z}_\nu \wedge d(\operatorname{Re} w) \neq 0, \quad d(\operatorname{Im} w) = 0$$

at the origin.

We can conveniently take $x_i = \operatorname{Re} z_i$, $y_j = \operatorname{Im} z_j$ and $s = \operatorname{Re} w$ as coordinates (for $i, j \in \{1, \dots, \nu\}$) and adjoin to them coordinates $t_1, \dots, t_{n'}$ ($n' := n - \nu$), obtaining a complete system of coordinates for M near 0. Adding appropriate constants allows us to assume that this system is centered at 0, and $\operatorname{Im} w(0) = 0$. We conclude that the differentials of the functions

$$z_j = x_j + iy_j, \quad j = 1, \dots, \nu, \quad w = s + i\phi(z, \bar{z}, s, t) \quad (2.1)$$

span the bundle T' in a neighborhood of 0, and moreover we have $\phi(0) = 0$ and $d\phi(0) = 0$. The characteristic set is spanned at the origin by $ds|_0$. We can then construct a frame for \mathcal{V} near 0 in the following way:

$$L_j = \frac{\partial}{\partial \bar{z}_j} - i \frac{\phi_{\bar{z}_j}}{1 + i\phi_s} \frac{\partial}{\partial s}, \quad j = 1, \dots, \nu, \quad L_{\nu+l} = \frac{\partial}{\partial t_l} - i \frac{\phi_{t_l}}{1 + i\phi_s} \frac{\partial}{\partial s}, \quad l = 1, \dots, n'. \quad (2.2)$$

One very important invariant of \mathcal{V} is the *Levi form*, which we discuss next.

Definition 2.1. Given a point $p \in M$ and a characteristic covector $\omega \in T_p^\circ \setminus \{0\}$, consider

$$\begin{aligned} \mathcal{B}_{\omega,p} : \mathcal{V}_p \times \mathcal{V}_p &\rightarrow \mathbb{C} \\ (v, w) &\mapsto \frac{1}{2i} \omega \left([L_1|_p, \overline{L_2|_p}] \right), \end{aligned}$$

where L_1, L_2 are local sections of \mathcal{V} such that $L_1|_p = v$, $L_2|_p = w$. The value $\mathcal{B}_{\omega,p}(v, w)$ does not depend on the sections L_1, L_2 chosen, and defines a hermitian form on \mathcal{V}_p . The associated quadratic form

$$\mathcal{L}_{\omega,p}(v) := \mathcal{B}_{\omega,p}(v, v), \quad v \in \mathcal{V}_p,$$

is called the *Levi form* of \mathcal{V} at (ω, p) .

In the case of structures of hypersurface type, since $\dim_{\mathbb{R}} T_0^\circ = 1$, it is customary to suppress the dependence on the characteristic covector, and speak only of the Levi form of the structure at 0. In the coordinates described in 2.1, the Levi form (at the characteristic vector $ds|_0$) can be represented by the following $n \times n$ matrix:

$$\mathcal{L}_p = \begin{pmatrix} \left(\frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} (0) \right)_{\nu \times \nu} & \left(\frac{\partial^2 \phi}{\partial z_j \partial t_k} (0) \right)_{\nu \times n'} \\ \left(\frac{\partial^2 \phi}{\partial t_j \partial \bar{z}_k} (0) \right)_{n' \times \nu} & \left(\frac{\partial^2 \phi}{\partial t_j \partial t_k} (0) \right)_{n' \times n'} \end{pmatrix}. \quad (2.3)$$

The structures that we shall work with in this paper are defined, then, as follows.

Definition 2.2. A (*Levi*-)nondegenerate structure of hypersurface type \mathcal{V} at $0 \in M$ is a locally integrable structure of hypersurface type such that the Levi form is nondegenerate at 0. A nondegenerate structure of hypersurface type is one which is of this form for all $p \in M$.

The two main examples of structures of this kind are the following:

Example 2.1 (Levi nondegenerate hypersurfaces). If $n' = 0$, i.e., there are no variables t_j , then the structure is a Levi nondegenerate real hypersurface $M \subset \mathbb{C}^{n+1}$, given in a neighborhood of zero by $\operatorname{Im} w = \phi(z, \bar{z}, \operatorname{Re} w)$, where coordinates in \mathbb{C}^{n+1} are $(z, w) \in \mathbb{C}^n \times \mathbb{C}$.

Example 2.2 (Mizohata structures). If $\nu = 0$, i.e., there are no variables z_j , then the structure is a locally integrable Mizohata structure (see [18]), with coordinates s, t_1, \dots, t_n and first integral $Z(s, t) = s + i\phi(s, t)$.

A general nondegenerate structure of hypersurface type can be seen as a superposition of Mizohata structures and CR hypersurfaces of complex space.

2.2. The method of associated differential equations. The study of the relationship between the geometry of real hypersurfaces in \mathbb{C}^2 and 2nd order ODEs

$$w'' = \Phi(z, w, w'). \quad (2.1)$$

goes back to Segre [17] and Cartan [6], see also Webster [19] More generally, the geometry of a real hypersurface in \mathbb{C}^{n+1} , $n \geq 1$, is related to that of a complete second order system of PDEs

$$w_{z_k z_l} = \Phi_{kl}(z_1, \dots, z_n, w, w_{z_1}, \dots, w_{z_n}), \quad \Phi_{kl} = \Phi_{lk}, \quad k, l = 1, \dots, n, \quad (2.2)$$

Moreover, *in the real-analytic case*, this relation becomes more explicit by means of the Segre family. Namely, to any real-analytic Levi-nondegenerate hypersurface $M \subset \mathbb{C}^{n+1}$, $n \geq 1$, one can uniquely associate a holomorphic ODE (2.1) ($n = 1$) or a holomorphic PDE system (2.2) ($n \geq 2$), whose solutions are precisely the Segre varieties. The Segre family of M plays a role of a “mediator” between the hypersurface and the associated differential equations. For recent work on associated differential equations in the degenerate setting, see e.g. the papers [13, 12] of the first author with Lamel and Shafikov.

The associated differential equation procedure is particularly simple in the case of a Levi-nondegenerate hypersurface in \mathbb{C}^2 . In this case the Segre family is an anti-holomorphic 2-parameter family of complex holomorphic curves. It then follows from the standard ODE theory that there exists a unique ODE (2.1), for which the Segre varieties are precisely the graphs of solutions. This ODE is called *the associated ODE*.

In general, both right hand sides in (2.1),(2.2) appear as functions determining the 2-jet of a Segre variety at a given point as a function of the 1-jet at the same point. More explicitly, we use coordinates

$$(z, w) = (z_1, \dots, z_n, w) \in \mathbb{C}^n \times \mathbb{C} = \mathbb{C}^{n+1}.$$

Fix a real-analytic hypersurface $M \subset \mathbb{C}^{n+1}$ passing through the origin, and choose a sufficiently small neighborhood U of the origin. When M is Levi-nondegenerate, we can associate a *complete second order system of holomorphic PDEs* to M , which is uniquely determined by the condition that the differential equations are satisfied by all the graphing functions $h(z, \zeta) = w(z)$ of the Segre varieties $\{Q_\zeta\}_{\zeta \in U}$ of M in a neighbourhood of the origin.

To be more precise, we consider a so-called *complex defining equation* (see, e.g., [3]),

$$w = \rho(z, \bar{z}, \bar{w}),$$

of M near the origin, which one obtains by substituting $u = \frac{1}{2}(w + \bar{w})$, $v = \frac{1}{2i}(w - \bar{w})$ into a real-analytic defining equation and solving for w as function of (z, \bar{z}, \bar{w}) by the implicit function theorem. The Segre variety Q_x of a point

$$x = (a, b) \in U, \quad a \in \mathbb{C}^n, b \in \mathbb{C},$$

is now given as the graph of the function

$$w(z) = \rho(z, \bar{a}, \bar{b}), \quad (2.3)$$

where we slightly abuse the notation using the same letter w for both the last coordinate and the function $w(z)$ defining a Segre variety. Differentiating (2.3) we obtain

$$w_{z_j} = \rho_{z_j}(z, \bar{a}, \bar{b}), \quad j = 1, \dots, n. \quad (2.4)$$

Considering (2.3) and (2.4) as a holomorphic system of equations with the unknowns \bar{a}, \bar{b} , in view of the Levi-nondegeneracy of M , an application of the implicit function theorem yields holomorphic functions A_1, \dots, A_n, B such that (2.3) and (2.4) are solved by

$$\bar{a}_j = A_j(z, w, w'), \quad \bar{b} = B(z, w, w'),$$

where we write

$$w' = (w_{z_1}, \dots, w_{z_n}).$$

The implicit function theorem applies here because the Jacobian of the system coincides with the Levi determinant of M for $(z, w) \in M$ ([3]). Differentiating (2.3) twice and substituting the above solution for \bar{a}, \bar{b} finally yields

$$w_{z_k z_l} = \rho_{z_k z_l}(z, A(z, w, w'), B(z, w, w')) =: \Phi_{kl}(z, w, w'), \quad k, l = 1, \dots, n, \quad (2.5)$$

or, more invariantly, i.e. independent of the coordinate choice,

$$j_{(z,w)}^2 Q_x = \Phi(x, j_{(z,w)}^1 Q_x). \quad (2.6)$$

Now (2.5) (or (2.6)) is the desired complete system of holomorphic second order PDEs denoted by $\mathcal{E} = \mathcal{E}(M)$.

Definition 2.3. We call $\mathcal{E} = \mathcal{E}(M)$ the system of PDEs associated with M .

2.3. Condition (E). Our main results rely on a new *analytic regularizability* condition for hypersurfaces of low regularity. In the C^∞ case, such condition was introduced in [14] and called *Condition (E)* (“E” for *extension*). We shall give below both an invariant and a coordinate-based formulations of it. For the basic concepts in CR Geometry (such as Segre varieties and formal submanifolds) we refer to [3], and for *jet bundles* and related concepts to [5].

Let

$$\pi: J^{1,n} \rightarrow \mathbb{C}^{n+1},$$

be the bundle of 1-jets of complex hypersurfaces of \mathbb{C}^{n+1} , which is a projective holomorphic fiber bundle over \mathbb{C}^{n+1} with the fiber isomorphic to \mathbb{P}^n , and let $M \subset \mathbb{C}^{n+1}$, $n \geq 1$, be (for the moment) a (C^∞) smooth *strictly pseudoconvex* real hypersurface. Then the complex tangent bundle $T^{\mathbb{C}}M$ induces the natural smooth (global) embedding

$$\varphi: M \rightarrow J^{1,n}, \quad x \mapsto (x, T_x^{\mathbb{C}}M).$$

The image

$$\varphi(M) =: M_J \subset J^{1,n}$$

is a smooth totally real $(2n+1)$ -dimensional real submanifold in the $(2n+1)$ -dimensional complex manifold $J^{1,n}$ by an observation of Webster [19]. Next, associated with M is the smooth (weakly) pseudoconvex real hypersurface

$$\widehat{M} := \pi^{-1}(M) \subset J^{1,n}.$$

The manifold M_J is a smooth real submanifold in \widehat{M} . Note that \widehat{M} itself is locally CR-equivalent to $M \times \mathbb{C}^n$ (and thus is *holomorphically degenerate*, see [3]). In what follows we denote by U^+ the *pseudoconvex side of M* , by which we mean the subset of an open neighborhood U of a point of M defined by $\rho < 0$, i.e.

$$U^+ = U \cap \{\rho < 0\},$$

where ρ is a local defining function of M in U with $d\rho \neq 0$ and the complex hessian satisfying $\partial\bar{\partial}\rho(X, \bar{X}) \geq 0$ for $X \in T^{1,0}M$. Given such U^+ , we write

$$\widehat{U}^+ := \pi^{-1}(U^+).$$

Then \widehat{M} is obviously (weakly) pseudoconvex and \widehat{U}^+ is the pseudoconvex side of \widehat{M} . Since all our considerations are local, the exact choice of neighborhoods of the reference point won't play any role.

We next fix a point $p \in M$ along with the corresponding point

$$\widehat{p} := (\pi|_{M_J})^{-1}(p) \in M_J \subset \widehat{M} \subset J^{1,n}.$$

Since M is smooth, we may consider at each point $q \in M$ near p , its formal complexification at q as a formal complex hypersurface in $\mathbb{C}^{n+1} \times \overline{\mathbb{C}^{n+1}}$ obtained by complexifying the formal Taylor series at q of the defining function ρ . In this way, the formal Segre variety Q_q of M is defined as a formal complex hypersurface defined by power series in $\mathbb{C}[[Z - q]]$, where $Z = (z, w) \in \mathbb{C}^n \times \mathbb{C}$ as before. Then the map 2-jets

$$q \in M \mapsto j_q^2 Q_q \in J^{2,n} \quad (2.1)$$

induces a *smooth embedding* of M into the bundle

$$J^{2,n} = J^{2,n}(\mathbb{C}^{n+1})$$

of 2-jets of complex hypersurfaces in \mathbb{C}^{n+1} . The space $J^{2,n}$ is canonically a fiber bundle

$$\pi_1^2: J^{2,n} \rightarrow J^{1,n}$$

over the 1-jet bundle $J^{1,n}$. The 2-jet embedding (2.1) defines a canonical section of π_1^2 ,

$$s_1^2: M_J \rightarrow J^{2,n}. \quad (2.2)$$

Now we recall our analytic regularizability condition for a smooth strictly pseudoconvex hypersurface [14]:

Definition 2.4. We say that M satisfies Condition (E) at $p \in M$ if the canonical section s_1^2 given by (2.2) extends holomorphically and smoothly up to the boundary to a neighborhood of $\widehat{U}^+ \cup \widehat{M}$.

We next give an (equivalent to the above) coordinate-based formulation of Condition (E). Let $M \subset \mathbb{C}^{n+1}$ be a smooth hypersurface with the defining equation

$$\rho(Z, \bar{Z}) = 0, \quad Z = (z, w) = (z_1, \dots, z_n, w) \in \mathbb{C}^{n+1}, \quad (2.3)$$

and $p \in M$ a fixed point where $\rho_w(p, \bar{p}) \neq 0$. Then the formal Segre variety at a point $q = (\tilde{q}, q_{n+1}) \in M$ near p is a graph of a function $w(z)$ (considered as a formal power series in $(z - \tilde{q})$). Then the 2-jets (2.1) are represented by either the scalar function Φ defined pointwise as $w''(z)$ for $z = \tilde{q}$ (case $n = 1$), or the symmetric matrix function $\Phi = (\Phi_{ij})$, $i, j = 1, \dots, n$, defined pointwise as the collection of the partial derivatives $w_{z_i z_j}$ for $z = \tilde{q}$ (the general case $n \geq 1$). It is possible to verify that, in turn, for $n = 1$ we have

$$\Phi = \frac{1}{(\rho_w)^3} \begin{vmatrix} \rho & \rho_z & \rho_w \\ \rho_z & \rho_{zz} & \rho_{zw} \\ \rho_w & \rho_{zw} & \rho_{ww} \end{vmatrix}, \quad (2.4)$$

and for $n > 1$ we have

$$\Phi_{ij} = \frac{1}{(\rho_w)^3} \begin{vmatrix} \rho & \rho_{z_j} & \rho_w \\ \rho_{z_i} & \rho_{z_i z_j} & \rho_{z_i w} \\ \rho_w & \rho_{z_j w} & \rho_{ww} \end{vmatrix}, \quad i, j = 1, \dots, n. \quad (2.5)$$

(To obtain (2.4),(2.5), one has to differentiate the identity (2.3) once, assuming w to be a function of z , express all the w_{z_j} in terms of the 1-jet of ρ , and then differentiate (2.3) once more to obtain $w_{z_i z_j} = \Phi_{ij}$ in terms of the 2-jet of ρ .) Both the scalar function (2.4) and the matrix valued function (2.5) can be considered as either smooth functions on the strictly pseudoconvex hypersurface M or as that on the totally real submanifold $M_J \subset J^{1,n}$ introduced above.

In terms of the Φ -function (2.4)-(2.5), Condition (E) reads in an equivalent form as follows:

Definition 2.5. We say that M satisfies Condition (E) at p , if the function Φ defined on M_J by either (2.4) or (2.5) extends holomorphically and smoothly up to the boundary to a neighborhood of \widehat{p} in the pseudoconvex side $\widehat{U}^+ \cup \widehat{M}$.

It is obvious that 2.5 is equivalent to 2.4. Now the analytic regularizability of a hypersurface can be characterized in terms of Condition (E), as stated in Theorem 1.1.

3. SPECIAL COORDINATES FOR NON-DEGENERATE STRUCTURES

Let (M, \mathcal{V}) be a non-degenerate locally integrable structure of hypersurface type. We denote by Σ° the closure of $T^\circ \setminus \{0\}$ in T^*M . This is a conic subset, and we denote by Σ the image of Σ° by the standard projection $\pi: T^*M \rightarrow M$. This is precisely the set of non-elliptic points of \mathcal{V} . We shall determine equations for these points in terms of the local coordinates of the previous section.

Theorem 3.1. *Let (M, \mathcal{V}) be a smooth, non-degenerate structure of hypersurface type. Then, $\Sigma^\circ \subset T^*M$ is a conic symplectic submanifold of dimension $2\nu + 2$, whose base projection Σ is a closed $(2\nu + 1)$ -dimensional submanifold of M . In the local coordinates 2.1, the set $U \cap \Sigma$ is described by the equations*

$$\frac{\partial \phi}{\partial t_j}(z, \bar{z}, s, t) = 0, \quad j = 1, \dots, n'.$$

Proof. We shall first prove that Σ is a smooth $(2\nu + 1)$ -dimensional manifold. We work in the coordinates 2.1. Assume that $(z, \bar{z}, s, t) \in \Sigma \cap U$. Then, there is a non-trivial linear combination

$$0 \neq v = \sum_{j=1}^{\nu} (a_j + ib_j) dz_j + (\mu + i\nu) (ds + id\phi) \in T'_{(z, \bar{z}, s, t)}, \quad a_j, b_j, \mu, \nu \in \mathbb{R},$$

such that $\text{Im } v = 0$. Computing this imaginary part, we obtain

$$\text{Im } v = \sum_{j=1}^{\nu} (b_j + \mu \phi_{x_j}) dx_j + \sum_{j=1}^{\nu} (a_j + \mu \phi_{y_j}) dy_j + \sum_{l=1}^{n'} \mu \phi_{t_l} dt_l + (\nu + \mu \phi_s) ds.$$

Since $v \neq 0$, we must have $\mu \neq 0$, which entails $\phi_{t_l} = 0$ for all $l = 1, \dots, n'$. We conclude that $d_t \phi(z, \bar{z}, s, t) = 0$. Conversely, suppose that $d_t \phi(z, \bar{z}, s, t) = 0$. Then, one can take $\mu = 1$, $\nu = -\phi_s(z, \bar{z}, s, t)$, $a_j = -\phi_{y_j}(z, \bar{z}, s, t)$ and $b_j = -\phi_{x_j}(z, \bar{z}, s, t)$. Therefore, the covector

$$v = -2i \sum_{j=1}^{\nu} \phi_{z_j}(z, \bar{z}, s, t) dz_j + (1 - i\phi_s(z, \bar{z}, s, t)) dw$$

is non-trivial and real, so $(z, \bar{z}, s, t) \in \Sigma \cap U$.

Since \mathcal{V} is non-degenerate, the matrix $\left(\frac{\partial^2 \phi}{\partial t_l \partial t_{l'}} \right)$, $1 \leq l, l' \leq n'$, is non-singular, which implies the map

$$(z, \bar{z}, s, t) \mapsto (z, \bar{z}, s, \partial_t \phi)$$

is a diffeomorphism of a neighborhood of 0 in U onto another such neighborhood. We conclude that Σ is a smooth submanifold of dimension $N - n' = 2\nu + n' + 1 - n' = 2\nu + 1$. It is also closed, since the property of being elliptic is an open condition.

From non-degeneracy, the forms

$$dx_j, dy_j, ds, d(\partial_{t_l} \phi), \quad j = 1, \dots, \nu, \quad l = 1, \dots, n'$$

make up a basis of T^*M near an open neighborhood U of $p \in M$. If σ denotes the tautological one-form of T^*M , we can write

$$\sigma = \sum_{j=1}^{\nu} a_j dx_j + b_j dy_j + \xi ds + d(\partial_t \phi \cdot \eta).$$

The equations for Σ^0 in π^*U are given by $\partial_t \phi = \eta = 0$, so the pullback of σ to Σ^0 is given by $\sum_{j=1}^{\nu} a_j dx_j + b_j dy_j + \xi ds$. Since the 2-form

$$\sum_{j=1}^{\nu} da_j \wedge dx_j + db_j \wedge dy_j + d\xi \wedge ds$$

is non-degenerate, Σ^0 is a symplectic submanifold of T^*M (of dimension $2\nu + 2$). \square

The submanifold Σ inherits an involutive structure from \mathcal{V} . We briefly recall the main notions (see Sections I.3 and I.4 in [18]):

Definition 3.1. Let (M, \mathcal{V}) be a smooth locally integrable structure and $\Sigma \subset M$ an embedded smooth submanifold. Assume that the dimension of

$$\mathbb{C}T_p\Sigma \cap \mathcal{V}_p$$

is constant for $p \in \Sigma$. Then, we say that Σ is *compatible* with the structure \mathcal{V} , and

$$\mathcal{V}_\Sigma = \bigcup_{p \in M} (\mathbb{C}T_p\Sigma \cap \mathcal{V}_p)$$

defines a smooth locally integrable structure Σ , which we call the *induced locally integrable structure*.

Remark 3.1. If Σ is a compatible submanifold, then the orthogonal

$$T'_\Sigma = \bigcup_{p \in M} T'_{\Sigma, p} \subset \mathbb{C}T^*\Sigma$$

is given by

$$T'_{\Sigma, p} = \left\{ \omega|_{\mathbb{C}T_p\Sigma}; \omega \in T'_p \right\}.$$

A related notion is that of a *non-characteristic submanifold*:

Definition 3.2. Let (M, \mathcal{V}) be a smooth involutive structure and $\Sigma \subset M$ an embedded smooth submanifold. We say Σ is *non-characteristic* if, for every $p \in \Sigma$, the map

$$\begin{aligned} T_p^\circ &\rightarrow T_p^*\Sigma \\ v &\mapsto v|_{T_p\Sigma} \end{aligned}$$

is injective.

We now apply these ideas to the structure \mathcal{V} at hand. Since the vector fields $\partial/\partial t_j$ are transversal to the submanifold Σ , we can write it locally as graph $t = F(z, \bar{z}, s)$, for some smooth function F fixing the origin. Introducing the change of variables

$$t \mapsto t - F(z, \bar{z}, s),$$

the submanifold Σ is defined, near 0, by the equations

$$t_j = 0, \quad j = 1, \dots, n'.$$

In these new coordinates, a system of first integrals for \mathcal{V} is given by

$$z_j = x_j + iy_j, \quad j = 1, \dots, \nu, \quad w = s + i\phi(z, \bar{z}, s, t), \quad (3.1)$$

where $\phi(0) = 0$, $d\phi(0) = 0$ and $\phi_{t_j}(z, \bar{z}, s, 0) \equiv 0$ for all $j = 1, \dots, n'$ and sufficiently small (z, \bar{z}, s) .

Now we obtain the following:

Proposition 3.1. *Let (M, \mathcal{V}) be a smooth non-degenerate structure of hypersurface type. Let Σ be the submanifold constructed in 3.1. Then, Σ is a non-characteristic compatible submanifold of M . Moreover, the induced involutive structure on Σ is that of a locally integrable Levi-nondegenerate CR structure of hypersurface type.*

Proof. Let $p \in \Sigma$. Working in the coordinates 3.1 centered at p , we have ds as a basis for T_p° , and clearly the curve $s \mapsto (0, 0, s, 0)$ is contained in Σ . In particular, the restriction $ds|_{T_p\Sigma}$ does not vanish identically, which implies Σ is non-characteristic. Compatibility follows from the expression of the vector fields 2.2 (which remain of the same form after the coordinate change 3.1), from which we see that L_j , $j = 1, \dots, \nu$, are precisely (complex) tangent to Σ . In particular, the structure is CR Levi-nondegenerate of hypersurface type. \square

Applying Morse's lemma (with parameters), one can simplify the coordinates 3.1 even further.

Theorem 3.2. *Let (M, \mathcal{V}) be a non-degenerate smooth structure of hypersurface type. Then, given a point $p \in M$, there are coordinates (z, \bar{z}, s, t) centered at p , a non-degenerate quadratic form \mathcal{L} in \mathbb{R}^n and smooth functions*

$$z_j = x_j + iy_j, \quad j = 1, \dots, \nu, \quad w = s + i(\phi(z, \bar{z}, s) + \mathcal{L}(t)) \quad (3.2)$$

with $\phi(0) = 0$, $d\phi(0)$ such that $\{dz_1, \dots, dz_\nu, dw\}$ span T' in a neighborhood of 0. Moreover, the structure \mathcal{V} induces on the submanifold Σ given by $\{t = 0\}$ an invariantly defined locally integrable Levi-nondegenerate CR structure of hypersurface type, equivalent to the hypersurface given by the defining function

$$\rho(z, \bar{z}, w, \bar{w}) = \operatorname{Im} w - \phi(z, \bar{z}, \operatorname{Re} w)$$

in $\mathbb{C}^{\nu+1}$.

Proof. Again, we work locally as in 3.1. We can apply Morse's lemma with parameters (see Lemma C.6.1 in [9]). If the neighborhood of 0 is sufficiently small, the only solution of $d_t\phi(z, \bar{z}, s, t) = 0$ is $t = 0$. Therefore, there is a smooth family of diffeomorphisms $t \mapsto G(z, \bar{z}, s, t)$ (with (z, \bar{z}, s) as parameters) and a non-degenerate quadratic form \mathcal{L} in \mathbb{R}^n such that

$$\phi(z, \bar{z}, s, t) = \phi(z, \bar{z}, s, 0) + \mathcal{L}(G(z, \bar{z}, s, t)).$$

Considering the change of variables

$$(z, \bar{z}, s, t) \mapsto (z, \bar{z}, s, G(z, \bar{z}, s, t)),$$

we obtain the result. \square

The CR submanifold Σ will play an important role in what follows, so we give it a special name.

Definition 3.3. Let (M, \mathcal{V}) be a nondegenerate locally integrable structure of hypersurface type. We call Σ the *central CR manifold* of \mathcal{V} . Given $p \in \Sigma$, we define $m(t)$ to be the signature of the quadratic form $\mathcal{L}(t)$.

4. THE CENTRAL MANIFOLD AND THE REGULARIZABILITY

The notion of mappings between integrable structures we consider in this work is the following:

Definition 4.1. Let (M, \mathcal{V}) and (N, \mathcal{W}) be two formally integrable structures over smooth manifolds M, N . A *mapping* (or morphism) between them is a smooth map $f : M \rightarrow N$ such that $(df)_p \mathcal{V}_p \subset \mathcal{W}_{f(p)}$ for all $p \in M$.

Remark 4.1. We use the same notation df for the complexification $1 \otimes df$ of df acting on the complexified tangent spaces.

In the locally integrable case, we can restate this notion in terms of the first integrals.

Proposition 4.1. *Let (M, \mathcal{V}) and (N, \mathcal{W}) be locally integrable structures over smooth manifolds M, N . Let $f : M \rightarrow N$ be a smooth map. Then, f is a mapping between \mathcal{V} and \mathcal{W} if and only if for every solution $u \in C^\infty(U)$ of \mathcal{W} (defined on an open subset $U \subset N$), the pullback $u \circ f$ is a solution of \mathcal{V} over $f^{-1}(U)$.*

We make the simple remark that a diffeomorphism $f : M \rightarrow N$ which is a mapping from \mathcal{V} to \mathcal{W} is not, in general, an equivalence of integrable structures. It is the case, however, if both of them have the same rank.

Now we specialize to the case of nondegenerate structures of hypersurface type. The first observation is that a map between structures induces a CR map between the central CR hypersurfaces.

Proposition 4.2. *Let (M_1, \mathcal{V}_1) and (M_2, \mathcal{V}_2) be two non-degenerate structures of hypersurface type, and let $\Sigma_i \subset M_i$ denote the central CR structure associated to \mathcal{V}_i , $i = 1, 2$. If $f : M_1 \rightarrow M_2$ is a smooth mapping between \mathcal{V}_1 and \mathcal{V}_2 , then f restricts to a CR map $\Sigma_1 \rightarrow \Sigma_2$.*

Proof. It is clear that $f(\Sigma_1) \subset \Sigma_2$, since the derivative of f is a real mapping. Moreover, since $\mathcal{V}_{\Sigma_i, p} = \mathbb{C}T_p\Sigma_i \cap (\mathcal{V}_i)_p$, the derivative of f maps $\mathcal{V}_{\Sigma_1, p}$ into $\mathcal{V}_{\Sigma_2, f(p)}$, and is therefore a morphism from \mathcal{V}_{Σ_1} into \mathcal{V}_{Σ_2} . Since they are CR structures, this is precisely the definition of a CR map. \square

We can use the central CR hypersurface and the signature $m(t)$ to give a necessary condition for equivalence of non-degenerate structures of hypersurface type.

Theorem 4.1. *Let (M_i, \mathcal{V}_i) be a (resp. smooth or analytic) nondegenerate locally integrable structure of hypersurface type, with central CR hypersurfaces Σ_i , $i = 1, 2$. Let $p \in \Sigma_1$, $q \in \Sigma_2$. Then, if there is a (resp. smooth or analytic) local equivalence between \mathcal{V}_1 and \mathcal{V}_2 (from a neighborhood of p to a neighborhood of q), then*

- (1) Σ_1 is locally CR equivalent to Σ_2 (from a neighborhood of p in Σ_1 to one of p_2 in Σ_2).
- (2) $m(p) = m(q)$.

Proof. This follows immediately from Proposition 4.2. \square

It turns out that the converse is also true for definite structures.

Theorem 4.2. *Let (M_i, \mathcal{V}_i) be a (resp. smooth or analytic) nondegenerate locally integrable structure of hypersurface type, with central CR hypersurface Σ_i , $i = 1, 2$. Let $p \in \Sigma_1$, $q \in \Sigma_2$. Assume furthermore that both (M_i, \mathcal{V}_i) are positive. Then, if there is a (resp. smooth or analytic) local equivalence F between (Σ_1, p) and (Σ_2, q) . then there it is the restriction onto Σ_1 of a (resp. smooth or analytic) local equivalence \tilde{F} between germs of \mathcal{V}_1 and \mathcal{V}_2 at p, q respectively.*

Proof. We deal with the case of smooth F , and discuss the analytic one in the end of the proof.

In view of the previous sections, we can obtain coordinates of the following kind for \mathcal{V}_1 : there is an open set

$$U = W \times I \subset \mathbb{R}^{2\nu+1+n'},$$

where $W \subset \mathbb{R}^{2\nu+1}$ is an open neighborhood of 0 and $I = (-\delta, \delta)^{n'} \subset \mathbb{R}^{n'}$. According to (3.2), we have coordinates (x, y, s, t) in $W \times I$, $z = x + iy \in \mathbb{C}^\nu$, $s \in \mathbb{R}$, $t \in I$, such that the first integrals of \mathcal{V}_1 are given by

$$z = z(x, y, s, t) = x + iy, \quad w = s + i \left(\phi(z, \bar{z}, s) + \frac{|t|^2}{2} \right), \quad |t|^2 = t_1^2 + \dots + t_{n'}^2,$$

where $\phi \in C^\infty(W)$ is a smooth, real-valued function such that $\phi(0) = 0$ and $d\phi(0) = 0$. The central manifold Σ_1 is characterized by setting $t = 0$ in our coordinates. In this way,

$$\Sigma_1 = \{(z, w) \in \Omega; \operatorname{Im} w = \phi(z, \bar{z}, \operatorname{Re} w)\}$$

is a strictly pseudoconvex hypersurface in $\mathbb{C}^{\nu+1}$, where $\Omega \subset \mathbb{C}^{\nu+1}$ is an appropriate neighborhood of the origin (as follows for the positivity of definitivity of the Levi form). We can also think of it as the locally integrable structure on W defined by the first integrals z and $w = s + i\phi(z, \bar{z}, s)$.

We perform the same procedure for \mathcal{V}_2 , which we express with a prime ' on top of all the objects, and assume we are given a CR equivalence $F : W \rightarrow W'$ between Σ_1 and Σ_2 . Our goal is to obtain an equivalence $F^\bullet : U \rightarrow U'$ between \mathcal{V}_1 and \mathcal{V}_2 (maybe after contraction of the neighborhoods about 0).

We can assume the following: let

$$\Omega_+ = \{(z_1, z_2) \in \Omega; \operatorname{Im} w > \phi(z, \bar{z}, \operatorname{Re} w)\}$$

be the pseudoconvex side of Σ_1 (analogously for the second structure, with a prime '). Then, there is a map

$$\tilde{F} : \Omega_+ \cup \Sigma_1 \rightarrow \Omega'_+ \cup \Sigma_2$$

which is biholomorphic as a map $\Omega_+ \rightarrow \Omega'_+$, extends smoothly to $\Omega_+ \cup \Sigma_1$, and its restriction to Σ_1 is the CR equivalence $\Sigma_1 \rightarrow \Sigma_2$. We can smoothly extend \tilde{F} to be a smooth diffeomorphism between full neighborhoods of Σ_1 and Σ_2 (biholomorphic only on the side Ω_+).

Let $\tilde{F}(z, w) = (f(z, w), g(z, w)) \in \mathbb{C}^\nu \times \mathbb{C}$. In order to simplify notation, we write

$$z := x + iy, \quad w := s + i \left(\phi(z, \bar{z}, s) + \frac{|t|^2}{2} \right), \quad \zeta := s + i\phi(z, \bar{z}, s), \quad (x, y, s, t) \in U.$$

In view of the discussion above, we have the basic identity for \tilde{F} :

$$\operatorname{Im} g(z, \zeta) = \phi'(f(z, \zeta), \overline{f(z, \zeta)}, \operatorname{Re} g(z, \zeta)).$$

Note that the function $\rho(z, \bar{z}, \eta, \bar{\eta}) := \operatorname{Im} g(z, \eta) - \phi'(f(z, \eta), \overline{f(z, \eta)}, \operatorname{Re} g(z, \eta))$ is a *smooth defining function* for the hypersurface Σ_1 in a neighborhood of the origin. In particular, it develops into the following identity:

$$\operatorname{Im} g(z, \eta) - \phi'(f(z, \eta), \overline{f(z, \eta)}, \operatorname{Re} g(z, \eta)) = \lambda(z, \eta)(\operatorname{Im} \eta - \phi(z, \bar{z}, \operatorname{Re} \eta)), \quad (4.1)$$

for (z, η) in a full neighborhood of the origin, where λ is a smooth non-vanishing function. Note that $\lambda(0) = 1$, so contracting the neighborhoods about the origin allows us to assume $\lambda > 0$. Then, we set

$$F^\bullet(x, y, s, t) = (X(x, y, s, t), Y(x, y, s, t), S(x, y, s, t), T(x, y, s, t)),$$

where

$$\begin{aligned} X(x, y, s, t) &= \operatorname{Re} f(z, w), & Y(x, y, s, t) &= \operatorname{Im} f(z, w), \\ S(x, y, s, t) &= \operatorname{Re} g(z, w), & T(x, y, s, t) &= t\sqrt{\lambda(z, w)} \end{aligned}$$

for $(x, y, s, t) \in U$ (everywhere here $w = s + i(\phi(z, \bar{z}, s) + |t|^2/2)$ and $z = x + iy$). Observe that

$$(z, w) = (z, \zeta + i|t|^2/2) \in \Omega_+,$$

therefore the expression for F^\bullet is well-defined. In view of the vanishing of $\lambda - 1$ on Σ , F^\bullet is smooth diffeomorphism in a neighborhood of the origin. Now, to show it defines a map of $\mathcal{V}_1 \rightarrow \mathcal{V}_2$, we have to prove that the pullback of the first integrals of \mathcal{V}_2 are solutions of \mathcal{V}_1 . Indeed,

$$((F^\bullet)^* z')(x, y, s, t) = X + iY = f(z, w),$$

which is a holomorphic function of the first integrals (z, w) , so it is a solution. For w' , we have

$$((F^\bullet)^* w')(x, y, s, t) = S + i \left(\phi'(X, Y, S) + \frac{|T|^2}{2} \right) = g(z, w)$$

(as follows from (4.1)), which is again a holomorphic function of the first integrals (z, w) . We conclude that F^\bullet is an equivalence between \mathcal{V}_1 and \mathcal{V}_2 .

In the analytic case, following the proof, we easily see that λ above can be chosen analytic (since the extension \tilde{F} becomes actually holomorphic by the reflection principle), thus the equivalence map F^\bullet is analytic too. \square

Theorems 4.1 and 4.2 now imply the following result, which is interesting even independently of the regularizability problem.

Theorem 4.3. *Let (M_i, \mathcal{V}_i) be a (resp. smooth or analytic) nondegenerate locally integrable structure of hypersurface type, with central CR hypersurfaces Σ_i , $i = 1, 2$. Let $p \in \Sigma_1$, $q \in \Sigma_2$. Assume furthermore that both (M_i, \mathcal{V}_i) are positive. Then, the structures \mathcal{V}_1 and \mathcal{V}_2 at p, q respectively are (resp. smoothly or analytically) equivalent if and only if the germs (Σ_1, p) and (Σ_2, q) of their central manifolds are (resp. smoothly or analytically) equivalent*

Theorem 4.3 immediately implies Theorem 1.2 from the Introduction, characterizing the analytic regularizability of structures in terms of their central manifolds.

We give below an example of using the central manifold for investigating the regularizability of nondegenerate structures.

Example 4.1. Consider the structure \mathcal{V} defined in a neighborhood of \mathbb{R}^4 (with variables (x, y, s, t)), with the first integrals

$$Z = z, \quad W = s + i(h(|z|^2) + t^2), \quad h \in C^\omega(U \setminus \{0\}) \cap C^\infty(U), \quad h \notin C^\omega(U), \quad U = (-a, a) \subset \mathbb{R}$$

where the function h satisfies

$$h(0) = h'(0) = 0, \quad h''(0) \neq 0.$$

In this case, $\nu = n' = 1$ and the vector fields that span the structure are

$$L_1 = \frac{\partial}{\partial \bar{z}} - izh'(|z|^2) \frac{\partial}{\partial s}, \quad L_2 = \frac{\partial}{\partial t} - 2it \frac{\partial}{\partial s}.$$

The central manifold Σ is given by

$$\{\operatorname{Im} w = h(|z|^2)\} \subset \mathbb{C}^2.$$

Following the elimination procedure in Section 2, we conclude that the Segre varieties of Σ near points with $z \neq 0$ satisfy the ODE

$$z^2 w'' = \psi(zw'),$$

where the function

$$\psi \in C^\omega(U \setminus \{0\}) \cap C^\infty(U), \quad \psi(0) = \psi'(0) = 0,$$

is related with h via

$$\xi^2 h''(\xi) = \psi(\xi h'(\xi)), \quad \xi \in U.$$

Considering the latter identity as an ODE for h with ψ fixed, we conclude that $\psi \notin C^\omega(U)$ (otherwise h would be analytic at 0 too as a solution of an analytic ODE). This means that Σ does not satisfy Condition (E), and so V is not analytically regularizable by Theorem 1.2.

5. ADDITIONAL SUFFICIENT CONDITIONS: THE EXTERNAL CR STRUCTURE

In this section, we work on sufficient conditions to ensure analytic regularizability of (M, \mathcal{V}) . We first describe a procedure introduced by Marson [15] that yields, in the coordinate system just introduced, an associated (generic) CR manifold to (M, \mathcal{V}) . The main idea is to formally add new variables x^\bullet that will turn the non-CR variables t_j into imaginary parts of new complex variables.

We maintain the notation of the previous section. Let $X \subset \mathbb{R}^{n'}$ be a neighborhood of 0 and let $U \subset \mathbb{R}^N$ be an open neighborhood of the origin for which we have the coordinates described in Sections 2 and 3. Set $U^\bullet = X \times U \subset \mathbb{R}^{n'+N}$. Let $M^\bullet = Z^\bullet(U^\bullet)$ be the submanifold parametrized by the mapping

$$Z^\bullet : U^\bullet \rightarrow \mathbb{C}^{m+n'}$$

given by

$$\begin{cases} Z_j^\bullet(x, y, x^\bullet, t, s) = x_j^\bullet + it_j, & j = 1, \dots, n', \\ Z_{n'+j}^\bullet(x, y, x^\bullet, t, s) = Z_j(x, y) = x_j + iy_j, & j = 1, \dots, \nu, \\ W^\bullet(x, y, s, t) = W(x, y, s, t) = s + i\phi(x, y, s, t). \end{cases}$$

This map is easily seen to have full rank, and induces a CR structure \mathcal{V}^\bullet on U^\bullet , with dual structure bundle T'^\bullet being generated by

$$\{dZ_1^\bullet, \dots, dZ_{m+n'}^\bullet\}.$$

We write $z_l^\bullet = x_l^\bullet + it_l$ and $z_j = x_j + iy_j$ for all $1 \leq l \leq n'$ and $1 \leq j \leq \nu$. Then, we can consider the vector fields

$$\begin{cases} L_l^\bullet = \frac{\partial}{\partial z_l^\bullet} - i \frac{\phi_{\bar{z}_l^\bullet}}{1 + \phi_s} \frac{\partial}{\partial s}, & l = 1, \dots, n', \\ L_j = \frac{\partial}{\partial z_j} - i \frac{\phi_{\bar{z}_j}}{1 + \phi_s} \frac{\partial}{\partial s}, & j = 1, \dots, \nu. \end{cases} \quad (5.1)$$

It's clear that $L_k^\bullet Z_l^\bullet = L_j Z_l^\bullet = 0$ for all $1 \leq k \leq n'$ and $1 \leq l \leq \nu$ (we maintain, for simplicity, the name L_j for the second set of vector fields, but formally they are the pullback of the vector fields L_j in U by the projection $X \times U \rightarrow U$). Therefore,

$$L_1^\bullet, \dots, L_{n'}^\bullet, L_1, \dots, L_\nu \text{ generate } \mathcal{V}^\bullet \text{ over } U^\bullet.$$

The new structure obtained has the same rank as the original structure, but we remark that it is *not invariant*, meaning that it depends on the particular coordinates (x, y, s, t) chosen.

We can describe the image of Z^\bullet explicitly as

$$M^\bullet = \left\{ (z^\bullet, z, w) \in \mathbb{C}^{n'} \times \mathbb{C}^\nu \times \mathbb{C}; \operatorname{Im} w = \phi(z, \bar{z}, \operatorname{Im} z^\bullet, \operatorname{Re} w) \right\}$$

This hypersurface is called an *external CR structure* associated to (\mathcal{V}, p) . We shall denote the Levi form of \mathcal{V}^\bullet (at the characteristic covector $ds|_0$) by $\mathcal{L}^\bullet(\zeta, \zeta^\bullet)$, where $\zeta \in \mathbb{C}^\nu$ and $\zeta^\bullet \in \mathbb{C}^{n'}$. From Section 2, the Levi form is represented by the matrix

$$\begin{pmatrix} \left(\frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} \right)_{1 \leq j, k \leq \nu} & \left(\frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_r^\bullet} \right)_{\substack{1 \leq j \leq \nu \\ 1 \leq r \leq n'}} \\ \left(\frac{\partial^2 \phi}{\partial z_l^\bullet \partial \bar{z}_k} \right)_{\substack{1 \leq l \leq n' \\ 1 \leq k \leq \nu}} & \left(\frac{\partial^2 \phi}{\partial z_l^\bullet \partial \bar{z}_r^\bullet} \right)_{1 \leq l, r \leq n'} \end{pmatrix},$$

which is given by

$$\begin{pmatrix} \left(\frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} \right)_{1 \leq j, k \leq \nu} & \frac{i}{2} \left(\frac{\partial^2 \phi}{\partial z_j \partial \bar{t}_r} \right)_{\substack{1 \leq j \leq \nu \\ 1 \leq r \leq n'}} \\ -\frac{i}{2} \left(\frac{\partial^2 \phi}{\partial t_l \partial \bar{z}_k} \right)_{\substack{1 \leq l \leq n' \\ 1 \leq k \leq \nu}} & \frac{1}{4} \left(\frac{\partial^2 \phi}{\partial t_l \partial \bar{t}_r} \right)_{1 \leq l, r \leq n'}. \end{pmatrix}$$

Therefore, if $\mathcal{L}(\zeta, \varpi)$ is the Levi form of \mathcal{V} at the covector $ds|_0$, we conclude that

$$\mathcal{L}^\bullet(\zeta, \zeta^\bullet) = \mathcal{L}(\zeta, (2i)^{-1} \zeta^\bullet).$$

In particular, we obtain the following

Proposition 5.1. *If \mathcal{V} is a nondegenerate structure of hypersurface type such that the Levi form is positive at $ds|_0$, then the external CR hypersurface is strictly pseudoconvex at the origin.*

We can now proceed with the proof of Theorem 1.3. We work in the coordinates described in Sections 2 and 3. Recall that we have, in a neighborhood of the origin, a smooth real-valued function ϕ vanishing at 0, with derivatives vanishing at 0 such that

$$\begin{cases} Z_j(x, y) = x_j + iy_j = z_j, & j = 1, \dots, \nu, \\ W(x, y, s, t) = s + i\phi(x, y, s, t) \end{cases}$$

and such that dZ_j and dW span, in a neighborhood of 0, the bundle T' . We have a smooth, strictly pseudoconvex hypersurface

$$M^\bullet = \left\{ (z^\bullet, z, w) \in \mathbb{C}^{n'} \times \mathbb{C}^\nu \times \mathbb{C}; \operatorname{Im} w = \phi(z, \bar{z}, \operatorname{Im} z^\bullet, \operatorname{Re} w) \right\} \quad (5.2)$$

which we assume satisfies condition (E). From Theorem 1.1, there is a CR diffeomorphism $f : M^\bullet \rightarrow N$ (near the origin), where (N^\bullet, q) is a real-analytic hypersurface in $\mathbb{C}^{n'+\nu+1}$.

We make use of the Abelian subalgebra $\mathfrak{a} \subset \text{aut}(M^\bullet, p)$ generated by the real vector fields

$$\left\{ \frac{\partial}{\partial z_l^\bullet} + \overline{\frac{\partial}{\partial z_l^\bullet}} \right\}, \quad l = 1, \dots, n'.$$

Its presence implies that the real-analytic hypersurface N^\bullet has an abelian subalgebra $\tilde{\mathfrak{a}} \subset \text{hol}(N^\bullet, q)$ of real dimension n' . Since N^\bullet is real-analytic, there are germs $V_1, \dots, V_{n'}$ of holomorphic vector fields in $\mathbb{C}^{n'+\nu+1}$ such that

$$\text{span}\{\text{Re } V_1, \dots, \text{Re } V_{n'}\} = \tilde{\mathfrak{a}}$$

and $\text{Re } V_1, \dots, \text{Re } V_{n'}$ are tangent to N^\bullet . Applying the holomorphic version of Frobenius' theorem, we can find holomorphic coordinates $(Z^\bullet, Z, W) \in \mathbb{C}^{n'+\nu+1}$ near q and a real-analytic function ψ (vanishing, along with its derivatives at the origin) such that N^\bullet is given by

$$\left\{ (Z^\bullet, Z, W) \in \mathbb{C}^{n'+\nu+1} : \text{Im } W = \psi(Z, \bar{Z}, \text{Im } Z^\bullet, \text{Re } W) \right\}$$

near q . Moreover, since f maps \mathfrak{a} to $\tilde{\mathfrak{a}}$, we conclude that f is of the form

$$f(z^\bullet, z, w) = (z^\bullet + f_1(z, w), f_2(z, w)).$$

We now consider the real-analytic locally integrable structure (N, \mathscr{W}) given by the first integrals

$$\begin{cases} \tilde{Z}_j(x, y, s, t) = x_j + iy_j, & j = 1, \dots, \nu, \\ \tilde{W}(x, y, s, t) = s + i\psi(x, y, s, t). \end{cases}$$

Then, the induced map $F : M \rightarrow N$, given by

$$F(x, y, s, t) = (x, y, \text{Re } f_2(Z(x, y), W(x, y, s, t)), t + \text{Im } f_1(Z(x, y), W(x, y, s, t))),$$

defines an isomorphism between the structures \mathscr{V} and \mathscr{W} , for the pullback of the first integrals \tilde{Z}_j and \tilde{W}_j are solutions of \mathscr{V} . We conclude that M is equivalent (near p) to a real-analytic structure. \square

6. AN APPLICATION TO RIGID STRUCTURES

We start with a

Definition 6.1. A locally integrable structure (2.2) is called *rigid*, if the respective function ϕ satisfies $\phi_s = 0$.

This definition generalizes the important class of rigid CR structures in CR geometry (see, e.g., [3]).

We first establish the following, somewhat surprising, result on the analytic regularizability of rigid CR structures.

Theorem 6.1. *Let $M \subset \mathbb{C}^{n+1}$ be a rigid strictly pseudoconvex CR-hypersurface*

$$M = \{\text{Im } w = H(z, \bar{z})\}, \quad (z, w) \in U \times \mathbb{C},$$

where $U \subset \mathbb{C}^n$ is a domain and $H \in C^\infty(U)$. Then M is analytically regularizable at a point $p \in M$ if and only if M is actually analytic near p .

Proof. We have to prove the necessity, since the sufficiency is trivial. We assume, without loss of generality, $p = 0$, $H(0) = 0$, $dH(0) = 0$. According to Section 2, M satisfies Condition (E), which means that the associated Φ -function (2.4) or (2.4) extends holomorphically to a domain $\hat{\Omega}^+ \subset J^1(\mathbb{C}^n, \mathbb{C})$ of the kind $\Omega^+ \times G$, where G is a neighborhood of the origin in \mathbb{C}^n and Ω^+ is a one-sided pseudoconvex neighborhood of M near 0. The fact that M is rigid means that it is invariant under real shifts $w \mapsto w + a$, $a \in \mathbb{R}$, hence the associated system (2.5) has its defining function Φ independent of w . Hence, being constant on the intersection of each line $z = \text{const}, \zeta = \text{const}$ with $\hat{\Omega}^+$ (for constant chosen close to the origin), Φ obviously extends holomorphically to a (full) neighborhood of the origin. This means that the Segre family of M is analytic near 0, and so is M . \square

Combining now Theorem 6.1 with Theorem 1.2, we arrive to the following regularizability theorem.

Theorem 6.2. *Let \mathcal{V} be a smooth rigid nondegenerate structure with a definite Levi form. Then \mathcal{V} is analytically regularizable at a point if and only if its central manifold is actually analytic near p .*

REFERENCES

- [1] M. S. Baouendi, C. H. Chang, and F. Treves. “Microlocal hypo-analyticity and extension of CR functions”. In: *Journal of Differential Geometry* 18.3 (1983), pp. 331–391.
- [2] M. S. Baouendi, P. Ebenfelt, and L. P. Rothschild. *Real Submanifolds in Complex Space and Their Mappings (PMS-47)*. Princeton University Press, 1999.
- [3] M. Salah Baouendi, Peter Ebenfelt, and Linda Preiss Rothschild. *Real submanifolds in complex space and their mappings*. Vol. 47. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1999, pp. xii+404. ISBN: 0-691-00498-6. DOI: 10.1515/9781400883967. URL: <https://doi.org/10.1515/9781400883967>.
- [4] S. Berhanu, P. D. Cordaro, and J. Hounie. *An Introduction to Involutive Structures*. New Mathematical Monographs. Cambridge University Press, 2014.
- [5] Andreas Čap and Jan Slovák. *Parabolic geometries. I*. Vol. 154. Mathematical Surveys and Monographs. Background and general theory. American Mathematical Society, Providence, RI, 2009, pp. x+628. ISBN: 978-0-8218-2681-2. DOI: 10.1090/surv/154. URL: <https://doi.org/10.1090/surv/154>.
- [6] Élie Cartan. “Sur la géométrie pseudo-conforme des hypersurfaces de l’espace de deux variables complexes II”. In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2)* 1.4 (1932), pp. 333–354. ISSN: 0391-173X. URL: http://www.numdam.org/item?id=ASNSP_1932_2_1_4_333_0.
- [7] S.S. Chern and J. K. Moser. “Real hypersurfaces in complex manifolds”. In: *Acta Math.* 133 (1974), pp. 219–271.
- [8] P. Cordaro and F. Treves. “Homology and cohomology in hypo-analytic structures of the hypersurface type”. In: *Journal of Geometric Analysis* 1.1 (1991), pp. 39–70.
- [9] Lars Hörmander. *The Analysis of Linear Partial Differential Operators III: Pseudo-Differential Operators*. Springer Berlin Heidelberg, 2007.
- [10] D.Zaitsev I. Kossovskiy. “Sphericity and Analyticity of a Real Hypersurface in Low Regularity.” In: *To appear* (2025).
- [11] Martin Kolar, Ilya Kossovskiy, and Dmitri Zaitsev. “Normal forms in Cauchy-Riemann geometry”. In: *Contemp. Math.* 681 (2017), pp. 153–177.
- [12] I. Kossovskiy and R. Shafikov. “Analytic differential equations and spherical real hypersurfaces”. In: *J. Differential Geom.* 102.1 (2016), pp. 67–126. ISSN: 0022-040X. URL: <http://projecteuclid.org/euclid.jdg/1452002878>.
- [13] I. Kossovskiy and R. Shafikov. “Divergent CR-equivalences and meromorphic differential equations”. In: *J. Eur. Math. Soc. (JEMS)* 18.12 (2016), pp. 2785–2819. ISSN: 1435-9855. DOI: 10.4171/JEMS/653. URL: <https://doi.org/10.4171/JEMS/653>.
- [14] I. Kossovskiy and D. Zaitsev. “Real-analytic coordinates for smooth strictly pseudoconvex CR-structures”. In: *Math. Res. Lett.* 29.5 (2022), pp. 1461–1484.
- [15] M. E. Marson. “Wedge extendability for hypo-analytic structures”. In: *Comm. Partial Differential Equations* 17.3-4 (1992), pp. 579–592.
- [16] L. Nirenberg. *A complex Frobenius theorem*. Sem. analytic functions 1, 172-189 (1958). 1958.
- [17] Beniamino Segre. “Questioni geometriche legate colla teoria delle funzioni di due variabili complesse.” Italian. In: *Rend. Sem. Mat. Roma, II. Ser.* 7.2 (1932), pp. 59–107.
- [18] F. Treves. *Hypo-Analytic Structures: Local Theory*. Princeton University Press, 1992.
- [19] S. M. Webster. “On the mapping problem for algebraic real hypersurfaces”. In: *Invent. Math.* 43.1 (1977), pp. 53–68. ISSN: 0020-9910. DOI: 10.1007/BF01390203. URL: <https://doi.org/10.1007/BF01390203>.

DEPARTMENT OF MATHEMATICS & INTERNATIONAL CENTER OF MATHEMATICS, SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY, SHENZHEN, CHINA

DEPARTMENT OF MATHEMATICS, MASARYK UNIVERSITY, BRNO, CZECHIA

INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, VIENNA UNIVERSITY OF TECHNOLOGY, VIENNA, AUSTRIA
Email address: ilyakos@sustech.edu.cn

DEPARTMENT OF MATHEMATICS, INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO, SÃO CARLOS, BRAZIL
Email address: viniciusnovelli@usp.br