# $L^2$ decay estimates of weak solutions to 3D fractional MHD equations in exterior domains

Zhi-Min Chen, Bo-Qing Dong, Qiuyue Zhang

School of Mathematical Sciences, Shenzhen University, Shenzhen 518060, China

### Abstract

Consider three-dimensional fractional MHD equations in an exterior domain with the Dirichlet boundary condition assumed. Asymptotic behaviours of weak solutions to the three-dimensional exterior fractional MHD equations are studied.  $L^2$  decay estimates of the weak solutions are obtained.

*Keywords:* Fractional MHD equations, asymptotic behaviours, weak solutions, exterior domains.

## 1. Introduction

The traditional form of incompressible magnetohydrodynamic (MHD) equations is expressed as (cf. [7, 11])

$$\partial_t u - \Delta u + \nabla (p + \frac{1}{2}|B|^2) = -u \cdot \nabla u + B \cdot \nabla B, \qquad (1)$$

$$\partial_t B - \Delta B = B \cdot \nabla u - u \cdot \nabla B, \qquad (2)$$

$$\nabla \cdot u = 0, \ \nabla \cdot B = 0, \tag{3}$$

for unknown velocity field  $u = (u_1, u_2, u_3)$ , magnetic field  $B = (B_1, B_2, B_3)$ and pressure p.

Mathematical investigations of the MHD equations are mainly assuming the fluid motion in whole spaces so that Fourier transform can be adopted. If the fluid domain  $\Omega$  involves a boundary  $\partial \Omega \neq \emptyset$ , mathematical difficulties

*Email addresses:* zmchen@szu.edu.cn (Zhi-Min Chen), bqdong@szu.edu.cn (Bo-Qing Dong), qyzhang0722@163.com (Qiuyue Zhang)

<sup>2020</sup> Mathematics subject classification. 35B40, 35D30, 35Q35, 76D99.

arise in the understanding of the magnetic equation (2) (cf. [11, 6]), since a suitable boundary condition with respect to the divergence-free condition  $\nabla \cdot B = 0$  and the integral involving the Laplacian  $-\Delta B = \nabla \times \nabla \times B$  is

$$(\nabla \times B) \times n = 0 \text{ and } B \cdot n = 0 \text{ on } \partial\Omega,$$
 (4)

where n the normal vector field of  $\partial \Omega$ .

In the present study, the smooth fluid domain  $\Omega$  is exterior to the bounded domain  $\mathbb{R}^3 \setminus \Omega$ . For simplicity of analysis, the Dirichlet boundary condition

$$u = 0, B = 0 \text{ on } \partial\Omega,$$
 (5)

is adopted and thus the Stokes operator

$$\mathcal{A} = -P\Delta$$

with the domain

$$D(\mathcal{A}) = \left\{ w \in W^{2,2}(\Omega)^3; \ \nabla \cdot w = 0, \ w|_{\partial\Omega} = 0 \right\},$$

is assumed. Here P is the Helmholtz projection operator mapping  $L^2(\Omega)^3$  onto the space

$$L^2_{\sigma}(\Omega)^3 =$$
 the completion of  $C^{\infty}_{0,\sigma}(\Omega)^3$  in  $L^2(\Omega)^3$ ,

where  $C_{0,\sigma}^{\infty}(\Omega)^3$  denotes the set of all divergence-free and compactly supported smooth vector fields on  $\Omega$ .

The corresponding exterior MHD problem with respect to the Dirichlet boundary condition was studied by Liu and Han [8]. However there is little literature on the exterior MHD problem involving fractional Stokes operators in exterior domains. Therefore, it is the purpose of the present study to provide a primary understanding of the fractional MHD equations

$$\partial_t u + \mathcal{A}^{\alpha} u = -P(u \cdot \nabla u) + P(B \cdot \nabla B), \tag{6}$$

$$\partial_t B + \mathcal{A}^\beta B = -P(u \cdot \nabla B) + P(B \cdot \nabla u) \tag{7}$$

in the exterior domain  $\Omega$ , for  $0 < \alpha, \beta \leq 1$ . The fractional operators are denoted by the spectral presentation

$$\mathcal{A}^{\kappa} = \int_0^\infty \lambda^{\kappa} dE_{\lambda},\tag{8}$$

with the domain

 $D(\mathcal{A}^{\kappa}) =$  the completion of  $D(\mathcal{A})$  under the norm  $\|\mathcal{A}^{\kappa}u\|_{L^{2}} + \|u\|_{L^{2}}$ . (9)

Here  $E_{\lambda}$  represents the spectral resolution of the unit determined by the operator  $\mathcal{A}$  (see Yosida [16, page 313, Theorem 1]).

**Definition 1.1.** Let  $(u_0, B_0) \in L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3$  and  $0 < \alpha, \beta \leq 1$ . A vector field (u, B) is called a weak solution of (6)-(7) associated with the initial condition

$$u|_{t=0} = u_0, \ B|_{t=0} = B_0, \tag{10}$$

if, for given T > 0, (u, B) satisfies the following the conditions

$$(u,B) \in L^{\infty}\left(0,T; L^{2}_{\sigma}(\Omega)^{3} \times L^{2}_{\sigma}(\Omega)^{3}\right) \cap L^{2}(0,T; D(\mathcal{A}^{\frac{\alpha}{2}}) \times D(\mathcal{A}^{\frac{\beta}{2}})),$$

$$\int_{0}^{T} (-\langle u(t), \partial_{t}\phi(t)\rangle + \langle \mathcal{A}^{\frac{\alpha}{2}}u(t), \mathcal{A}^{\frac{\alpha}{2}}\phi(t)\rangle)dt - \langle u(0), \phi(0)\rangle$$
$$= \int_{0}^{T} \langle u(t) \otimes u(t), \nabla\phi(t)\rangle dt - \int_{0}^{T} \langle B(t) \otimes B(t), \nabla\phi(t)\rangle dt, \quad (11)$$

and

$$\int_{0}^{T} (-\langle B(t), \partial_{t}\psi(t)\rangle + \langle \mathcal{A}^{\frac{\beta}{2}}B(t), \mathcal{A}^{\frac{\beta}{2}}\psi(t)\rangle)dt - \langle B(0), \psi(0)\rangle$$
$$= \int_{0}^{T} \langle u(t) \otimes B(t), \nabla\phi(t)\rangle dt - \int_{0}^{T} \langle B(t) \otimes u(t), \nabla\phi(t)\rangle dt, \quad (12)$$

where  $\langle \cdot, \cdot \rangle$  represents the inner product of  $L^2_{\sigma}(\Omega)^3$  and  $(\phi, \psi)$  denotes a test function in

$$C^{1}([0,T]; C^{\infty}_{0,\sigma}(\Omega)^{3} \times C^{\infty}_{0,\sigma}(\Omega)^{3})$$
 with  $\phi(T) = 0$  and  $\psi(T) = 0$ .

We are now in the position to state the main result.

**Theorem 1.1.** Let  $\Omega$  be a smooth exterior domain of  $\mathbb{R}^3$ ,  $\frac{3}{4} < \alpha$ ,  $\beta \leq 1$  and  $(u_0, B_0) \in L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3$ . Then (6)-(7) admit a global weak solution so that

$$\|(u(t), B(t))\|_{L^2} \to 0 \quad as \quad t \to \infty.$$

$$\tag{13}$$

Moreover, if the linear analytic semigroup has the algebraic decay property

$$\|(e^{-t\mathcal{A}^{\alpha}}u_{0}, e^{-t\mathcal{A}^{\beta}}B_{0})\|_{L^{2}} \le Ct^{-\gamma}, \quad for \ t > 0,$$
(14)

for  $0 < \gamma < \frac{1}{2}$  when  $\max\{\alpha, \beta\} = 1$  and  $0 < \gamma \leq \frac{1}{2}$  when  $\max\{\alpha, \beta\} < 1$ , then the nonlinear solution (u, B) has the same algebraic decay property

$$\|(u(t), B(t))\|_{L^2} \le Ct^{-\gamma}, \quad for \ t > 0.$$
 (15)

Theorem 1.1 will be proved by employing the spectral representation of the exterior Stokes operator in the  $L^2$  space. The exterior Navier-Stokes flows and related problems were studied by Masuda [9], Miyakawa [10], Borchers and Miyakawa [1, 2, 3], Chen [5] and Chen, Kagei and Miyakawa [4]. Especially, when B = 0 and  $\alpha = 1$ , Theorem 1.1 reduces to the  $L^2$  decay result of the exterior Navier-Stokes flows given by Miyakawa [2].  $L^2$  decay estimates of weak solutions to the MHD equations ( $\alpha = \beta = 1$ ) in whole spaces were studied by Schonbek *et al* [13] by applying the classic Fourier splitting method from Schonbek [12].

The proof of Theorem 1.1 is shown in the following two sections. Section 2 is a collection of preliminary lemmas for the proof. Section 3 consists of three subsections: existence of weak solutions, proof of assertion (13) and proof of assertion (15).

## 2. Preliminaries

Let C and  $C_n$  be generic constants, which are independent of the quantities  $u, B, \phi, \psi, t > 0$  and  $x \in \Omega$ , but they may depend on initial vector field  $(u_0, B_0)$ . However,  $C_n$  depends on the integer  $n \ge 1$ , while C is independent of  $n \ge 1$ .

Following Sobolev imbedding is fundamental important in our analysis.

**Lemma 2.1.** (Miyakawa [10, Theorem 2.4]) Let  $0 \leq \kappa \leq 1$  and  $H^{\frac{\kappa}{2}}$  denote the completion of  $D(\mathcal{A})$  with respect to the norm  $\|\mathcal{A}^{\frac{\kappa}{2}}w\|_{L^2}$ . Then the Sobolev imbedding

$$\|w\|_{L^r} \le 2\|\mathcal{A}^{\frac{\kappa}{2}}w\|_{L^2}, \text{ for } w \in H^{\frac{\kappa}{2}},$$
 (16)

holds true with respect to

$$\frac{1}{r} = \frac{1-\kappa}{2} + \frac{\kappa}{6}$$

**Lemma 2.2.** If  $0 < \kappa \leq 1$  and  $w \in L^2_{\sigma}(\Omega)^3$ , then we have

$$\|\mathcal{A}^{\kappa}e^{-t\mathcal{A}^{\kappa}}w\|_{L^{2}} \le t^{-1}\|w\|_{L^{2}}; \quad \|e^{-t\mathcal{A}^{\kappa}}w\|_{L^{2}} \le \|w\|_{L^{2}}, \tag{17}$$

$$\|\nabla e^{-t\mathcal{A}^{\kappa}}w\|_{L^{2}} \le Ct^{-\frac{1}{2\kappa}}\|w\|_{L^{2}}.$$
(18)

$$\lim_{t \to \infty} \|e^{-t\mathcal{A}^{\kappa}}w\|_{L^2} = 0,$$
(19)

This lemma is a simple consequence of the spectral representation ( see, for example, Yosida [16, Page 313, Theorem 1]) and the property of  $\mathcal{A}^{\kappa}$  being positive and a self-adjoint operator.

Equation (17) is obtained as

$$\|\mathcal{A}^{\kappa n} e^{-t\mathcal{A}^{\kappa}} w\|_{L^{2}}^{2} = \int_{0}^{\infty} \lambda^{2n\kappa} e^{-2t\lambda^{\kappa}} d\|E_{\lambda} w\|_{L^{2}}^{2} \le t^{-2n} \|w\|_{L^{2}}^{2}, \ n = 0, 1, \quad (20)$$

which also ensures the analytic property of the semigroup  $e^{-t\mathcal{A}^{\kappa}}$ .

Similar to (20), we have

$$\|\nabla e^{-t\mathcal{A}^{\kappa}}w\|_{L^{2}} = \|\mathcal{A}^{\frac{1}{2}}e^{-t\mathcal{A}^{\kappa}}w\|_{L^{2}} \leq Ct^{-\frac{1}{2\kappa}}\|w\|_{L^{2}}.$$
 (21)

and thus obtain (18).

Decay property (19) readily follows from (17), (18) and the density of  $C^{\infty}_{\sigma}(\Omega)^3$  in  $L^2_{\sigma}(\Omega)^3$ .

## 3. Proof of Theorem 1.1

## 3.1. Existence of weak solutions

With the use of the mollification operator

$$J_n = n(n+\mathcal{A})^{-1},\tag{22}$$

for positive integers n, we construct approximation solutions to (6)-(7) by solving the equations

$$\partial_t u_n + \mathcal{A}^{\alpha} u_n = P(J_n B_n \cdot \nabla B_n - J_n u_n \cdot \nabla u_n), \quad u_n(0) = J_n u_0, \tag{23}$$

$$\partial_t B_n + \mathcal{A}^\beta B_n = P(J_n B_n \cdot \nabla u_n - J_n u_n \cdot \nabla B_n), \quad B_n(0) = J_n B_0.$$
(24)

By the spectral representation and for  $w \in L^2_{\sigma}(\Omega)^3$ , we have

$$\|J_n w\|_{L^2} \le \|w\|_{L^2},\tag{25}$$

$$\lim_{n \to \infty} \|J_n w - w\|_{L^2} = 0,$$
(26)

and, for a constant  $C_n$  dependent of n and  $\frac{3}{4} < \kappa \leq 1$ ,

$$\|J_n w\|_{L^{\infty}} \le C_n \|w\|_{L^2}, \ \|\mathcal{A}^{\kappa} J_n w\|_{L^2} \le C_n \|w\|_{L^2},$$
(27)

after the use of Sobolev imbedding.

Now we show the existence of weak solutions approached by approximate solutions in the space

$$L^{\infty}(0,T; W^{1,2}_{0,\sigma}(\Omega)^3 \times W^{1,2}_{0,\sigma}(\Omega)^3),$$

where  $W_{0,\sigma}^{1,2}(\Omega)^3$  is the closure of  $C_{0,\sigma}^{\infty}(\Omega)^3$  in the Sobolev space  $W^{1,2}(\Omega)^3$ .

**Theorem 3.1.** For  $\frac{3}{4} < \alpha$ ,  $\beta \leq 1$  and  $(u_0, B_0) \in L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3$ , then (6)-(7) admit a weak solution.

Let us begin with the unique existence of the approximate solutions.

**Proposition 3.1.** For  $\frac{3}{4} < \alpha$ ,  $\beta \leq 1$ ,  $(u_0, B_0) \in L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3$ , T > 0and an integer  $n \geq 1$ , then there exits a unique vector field  $(u_n(t), B_n(t))$  on [0, T] so that

$$(u_n, B_n) \in L^{\infty}(0, T; W^{1,2}_{0,\sigma}(\Omega)^3 \times W^{1,2}_{0,\sigma}(\Omega)^3)$$

and solves the integral equations

$$u_{n}(t) = e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0} + \int_{0}^{t} e^{-(t-s)\mathcal{A}^{\alpha}}P(J_{n}B_{n}(s)\cdot\nabla B_{n}(s) - J_{n}u_{n}(s)\cdot\nabla u_{n}(s))ds, (28)$$
  
$$B_{n}(t) = e^{-t\mathcal{A}^{\beta}}J_{n}B_{0} + \int_{0}^{t} e^{-(t-s)\mathcal{A}^{\beta}}P(J_{n}B_{n}(s)\cdot\nabla u_{n}(s) - J_{n}u_{n}(s)\cdot\nabla B_{n}(s))ds. (29)$$

*Proof.* To apply the contraction mapping principle with respect to a fixed integer  $n \ge 1$  and a bound M > 0, we define the complete metric space

$$X_T = \{ (u, B) \in L^{\infty}(0, T; W^{1,2}_{0,\sigma}(\Omega)^3 \times W^{1,2}_{0,\sigma}(\Omega)^3); \\ (u(0), B(0)) = (J_n u_0, J_n B_0), \ \| (u, B) \|_{X_T} \le M \}$$

with the norm

$$\|(u,B)\|_{X_T} = \underset{0 \le t \le T}{\operatorname{ess\,sup}}(\|(u(t),B(t))\|_{L^2} + \|\nabla(u(t),B(t))\|_{L^2}),$$

and the operators

$$\Phi_n(u,B)(t) = e^{-t\mathcal{A}^{\alpha}}J_nu_0 + \int_0^t e^{-(t-s)\mathcal{A}^{\alpha}}P(J_nB\cdot\nabla B - J_nu\cdot\nabla u)ds,$$
  
$$\Psi_n(u,B)(t) = e^{-t\mathcal{A}^{\beta}}J_nB_0 + \int_0^t e^{-(t-s)\mathcal{A}^{\beta}}P(J_nB\cdot\nabla u - J_nu\cdot\nabla B)ds.$$

Employing Lemma 2.2 and (25)-(27), we obtain that

$$\begin{aligned} \|e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0}\|_{L^{2}} + \|\nabla e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0}\|_{L^{2}} + \|e^{-t\mathcal{A}^{\beta}}J_{n}B_{0}\|_{L^{2}} + \|\nabla e^{-t\mathcal{A}^{\beta}}J_{n}B_{0}\|_{L^{2}} \\ &\leq \|u_{0}\|_{L^{2}} + \|B_{0}\|_{L^{2}} + \|e^{-t\mathcal{A}^{\alpha}}\mathcal{A}^{\frac{1}{2}}J_{n}u_{0}\|_{L^{2}} + \|e^{-t\mathcal{A}^{\beta}}\mathcal{A}^{\frac{1}{2}}J_{n}B_{0}\|_{L^{2}} \\ &\leq C_{n}(\|u_{0}\|_{L^{2}} + \|B_{0}\|_{L^{2}}) \leq \frac{M}{4}, \end{aligned}$$
(30)

provided that M is sufficiently large.

Therefore, for  $(u, B) \in X_T$ , we use (25), (27) and Lemma 2.2 to produce

$$\begin{split} \|\Phi_{n}(u,B)(t)\|_{L^{2}} + \|\nabla\Phi_{n}(u,B)(t)\|_{L^{2}} \\ &\leq \|e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0}\|_{L^{2}} + \int_{0}^{t}\|e^{-(t-s)\mathcal{A}^{\alpha}}P(J_{n}B\cdot\nabla B - J_{n}u\cdot\nabla u)\|_{L^{2}}ds \\ &+ \|\nabla e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0}\|_{L^{2}} + \int_{0}^{t}\|\nabla e^{-(t-s)\mathcal{A}^{\alpha}}P(J_{n}B\cdot\nabla B - J_{n}u\cdot\nabla u)\|_{L^{2}}ds \\ &\leq \frac{M}{4} + C\int_{0}^{t}(1 + (t-s)^{-\frac{1}{2\alpha}})(\|J_{n}u\cdot\nabla u\|_{L^{2}} + \|J_{n}B\cdot\nabla B\|_{L^{2}})ds \\ &\leq \frac{M}{4} + C\int_{0}^{t}(1 + (t-s)^{-\frac{1}{2\alpha}})(\|J_{n}u\|_{L^{\infty}}\|\nabla u\|_{L^{2}} + \|J_{n}B\|_{L^{\infty}}\|\nabla B\|_{L^{2}})ds \\ &\leq \frac{M}{4} + C_{n}(T + T^{1-\frac{1}{2\alpha}})M^{2}, \end{split}$$

or

$$\|\Phi_n(u,B)(t)\|_{L^2} + \|\nabla\Phi_n(u,B)(t)\|_{L^2} \le \frac{M}{2},\tag{31}$$

for  $0 < t \leq T$  and T sufficiently small.

Similarly, for (u, B),  $(u', B') \in X_T$ , we have the contraction property

$$\begin{split} \|\Phi_{n}(u,B)(t) - \Phi_{n}(u',B')(t)\|_{L^{2}} + \|\nabla\Phi_{n}(u,B)(t) - \nabla\Phi_{n}(u',B')(t)\|_{L^{2}} \\ &\leq C_{n}(T+T^{1-\frac{1}{2\alpha}})M\|(u,B) - (u',B')\|_{X_{T}} \\ &\leq \frac{1}{4}\|(u,B) - (u',B')\|_{X_{T}}, \end{split}$$
(32)

when T is sufficiently small.

Arguing in the same manner, we have (31)-(32) with  $\Phi_n$  replaced by  $\Psi_n$ . Thus  $(\Phi_n, \Psi_n)$  is a contraction operator mapping  $X_T$  into itself, and so  $(\Phi_n, \Psi_n)$  admits a unique fixed point  $(u_n, B_n) \in X_T$  satisfying integral equations (28)-(29). We thus obtain local existence result. This integral equation solution also solves the differential equations (23) and (24). This is confirmed by the  $L^2$  theory of linear parabolic equation given by Simon [14], which also ensures that

$$(u_n, B_n) \in L^2\left(0, T; D(\mathcal{A}^{\alpha}) \times D(\mathcal{A}^{\beta})\right), (\partial_t u_n, \partial_t B_n) \in L^2\left(0, T; L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3\right).$$

It is readily seem that the local solution can be extended globally, if

$$\|(u_n, B_n)\|_{X_T} < \infty, \tag{33}$$

whenever  $(u_n, B_n)$  solves (23) and (24) on the open time interval [0, T) for T > 0. Indeed, taking the inner product of (23)-(24) with  $(u_n, B_n)$  and integrating parts by the divergence free condition of  $J_n u$  and  $J_n B$ , we have

$$\langle \partial_t u_n, u_n \rangle + \langle \partial_t B_n, B_n \rangle + \langle \mathcal{A}^{\alpha} u_n, u_n \rangle + \langle \mathcal{A}^{\beta} B_n, B_n \rangle = 0.$$
(34)

Hence, we have the energy inequality, after the integration with respect to t and the use of (25),

$$\|u_n(t)\|_{L^2}^2 + \|B_n(t)\|_{L^2}^2 + 2\int_0^t (\|\mathcal{A}^{\frac{\alpha}{2}}u_n\|_{L^2}^2 + \|\mathcal{A}^{\frac{\beta}{2}}B_n\|_{L^2}^2)ds \leq \|J_nu_0\|_{L^2}^2 + \|J_nB_0\|_{L^2}^2 \leq \|u_0\|_{L^2}^2 + \|B_0\|_{L^2}^2.$$
(35)

This shows the uniform boundedness of  $||(u_n, B_n)(t)||_{L^2}$ .

To obtain the boundedness of  $\|\nabla(u_n, B_n)(t)\|_{L^2}$ , we apply (18), (27) and (35) into (28)-(29) to produce

$$\begin{aligned} \|\nabla(u_n, B_n)(t)\|_{L^2} &\leq C_n \|(u_0, B_0)\|_{L^2} \\ &+ C_n \int_0^t ((t-s)^{-\frac{1}{2\alpha}} + (t-s)^{-\frac{1}{2\beta}}) \|(u_n, B_n)\|_{L^2} \|\nabla(u_n, B_n)\|_{L^2} ds \\ &\leq C_n + C_n \int_0^t ((t-s)^{-\frac{1}{2\alpha}} + (t-s)^{-\frac{1}{2\beta}}) \|\nabla(u_n, B_n)\|_{L^2} ds. \end{aligned}$$

Hence, using the Gronwall inequality, we have the desired bound

$$\|\nabla(u_n, B_n)(t)\|_{L^2} \leq C_n \exp(C_n T^{1-\frac{1}{2\alpha}} + C_n T^{1-\frac{1}{2\beta}}), \ 0 < t < T.$$
(36)

Therefore, the combination of (35) and (36) gives (33). Thus the local solution  $(u_n, B_n)$  can be extended to the time interval [0, T] for any T > 0. The proof of Proposition 3.1 is complete.

**Proof of Theorem 3.1.** To obtain the weak solution existence, we need to study the compactness of the sequence  $(u_n, B_n)$ . For  $\phi \in W^{1,2}_{0,\sigma}(\Omega)^3$ , taking the inner product of (28) with  $\phi$  and integrating by parts, we have

$$\begin{aligned} \langle \partial_t u_n, \phi \rangle &= -\langle \mathcal{A}^{\alpha} u_n, \phi \rangle + \langle J_n B_n \cdot \nabla B_n - J_n u_n \cdot \nabla u_n, \phi \rangle \\ &= -\langle \mathcal{A}^{\alpha - \frac{1}{2}} u_n, \mathcal{A}^{\frac{1}{2}} \phi \rangle - \langle J_n u_n \otimes u_n, \nabla \phi \rangle + \langle J_n B_n \otimes B_n, \nabla \phi \rangle \\ &\leq (\|\mathcal{A}^{\alpha - \frac{1}{2}} u_n\|_{L^2} + \|J_n u_n\|_4 \|u_n\|_4 + \|J_n B_n\|_4 \|B_n\|_4) \|\nabla \phi\|_{L^2}. \end{aligned}$$
(37)

It follows from Hölder inequality and Lemma 2.1 that

$$\|w\|_{L^4} \le \|w\|_{L^2}^{1-\frac{3}{4\kappa}} \|w\|_{L^{1/(\frac{1}{2}-\frac{\kappa}{3})}}^{\frac{3}{4\kappa}} \le C \|w\|_{L^2}^{1-\frac{3}{4\kappa}} \|\mathcal{A}^{\frac{\kappa}{2}}w\|_{L^2}^{\frac{3}{4\kappa}}, \text{ for } \kappa > \frac{3}{4}.$$
(38)

Therefore, by (25), (35) and an interpolation inequality, we see that (37) becomes

$$\begin{aligned} \|\partial_{t}u_{n}\|_{W^{-1,2}} &\leq C \|u_{n}\|_{L^{2}}^{\frac{1}{\alpha}-1} \|\mathcal{A}^{\frac{\alpha}{2}}u_{n}\|_{L^{2}}^{2-\frac{1}{\alpha}} \\ &+ C \|u_{n}\|_{L^{2}}^{2-\frac{3}{2\alpha}} \|\mathcal{A}^{\frac{\alpha}{2}}u_{n}\|_{L^{2}}^{\frac{3}{2\alpha}} + C \|B_{n}\|_{L^{2}}^{2-\frac{3}{2\beta}} \|\mathcal{A}^{\frac{\beta}{2}}B_{n}\|_{L^{2}}^{\frac{3}{2\beta}} \\ &\leq C (\|\mathcal{A}^{\frac{\alpha}{2}}u_{n}\|_{L^{2}}^{2-\frac{1}{\alpha}} + \|\mathcal{A}^{\frac{\alpha}{2}}u_{n}\|_{L^{2}}^{\frac{3}{2\alpha}} + \|\mathcal{A}^{\frac{\beta}{2}}B_{n}\|_{L^{2}}^{\frac{3}{2\beta}}). \end{aligned} (39)$$

Arguing in the same manner as in the derivation of (39), we have

$$\|\partial_t B_n\|_{W^{-1,2}} \leq C(\|\mathcal{A}^{\frac{\beta}{2}}B_n\|_{L^2}^{2-\frac{1}{\beta}} + \|\mathcal{A}^{\frac{\alpha}{2}}u_n\|_{L^2}^{\frac{3}{2\alpha}} + \|\mathcal{A}^{\frac{\beta}{2}}B_n\|_{L^2}^{\frac{3}{2\beta}}).$$
(40)

Thus, the combination of (35), (39) and (40) yields the boundedness of

$$\partial_t u_n, \partial_t B_n \in L^{\kappa}(0, T; W^{-1,2}(\Omega)^3) \text{ for } \kappa = \min\{\frac{2}{2 - \frac{1}{\alpha}}, \frac{2}{2 - \frac{1}{\beta}}, \frac{4\alpha}{3}, \frac{4\beta}{3}\}$$
 (41)

uniformly with respect to all  $n \ge 1$  for any fixed T > 0. It follows from (35) that the sequence  $(u_n, B_n)$  is uniformly bounded with respect to integers  $n \ge 1$  in the space

$$L^{\infty}(0,T; L^{2}_{\sigma}(\Omega)^{3} \times L^{2}_{\sigma}(\Omega)^{3}) \cap L^{2}(0,T; D(\mathcal{A}^{\frac{\alpha}{2}}) \times D(\mathcal{A}^{\frac{\beta}{2}})).$$
(42)

Therefore,  $(u_n, B_n)$  admits a subsequence, denoted again by  $(u_n, B_n)$ , converging to an element (u, B) in the space given by (42) in the following sense

$$(u_n, B_n) \to (u, B)$$
 weak-star in  $L^{\infty}(0, T; L^2_{\sigma}(\Omega)^3 \times L^2_{\sigma}(\Omega)^3)$   
 $(u_n, B_n) \to (u, B)$  weakly in  $L^2(0, T; D(\mathcal{A}^{\frac{\alpha}{2}}) \times D(\mathcal{A}^{\frac{\beta}{2}})).$ 

Furthermore, due to the compact result from Temam [15, Theorem 2.1 in page 271], we see that

$$(u_n, B_n) \to (u, B)$$
 strongly in  $L^2(0, T; L^2(K)^3 \times L^2(K)^3)$ ,

for any compact set  $K \subset \Omega$ . This convergence implies that (u, B) is a desired weak solution solving (11) and (12).

The proof of Theorem 3.1 is complete.

#### 3.2. Proof of assertion (13)

From the convergence of the sequence  $(u_n, B_n)$  in the previous subsection, we see that

$$\|(u,B)(t)\|_{L^2} \le \liminf_{n \to \infty} \|(u_n,B_n)(t)\|_{L^2},\tag{43}$$

for t > 0. Thus it suffices to show corresponding  $L^2$  decay estimates of the approximate solutions  $(u_n, B_n)$  uniformly with respect to  $n \ge 1$ . Therefore, for simplicity of notation, we omit the subscript n for the approximate solutions. Thus approximate equations (28) and (29) are rewritten, respectively, as

$$u(t) = e^{-t\mathcal{A}^{\alpha}}J_{n}u_{0} + \int_{0}^{t} e^{-(t-s)\mathcal{A}^{\alpha}}P(J_{n}B\cdot\nabla B - J_{n}u\cdot\nabla u)ds, \qquad (44)$$

$$B(t) = e^{-t\mathcal{A}^{\beta}}J_nB_0 + \int_0^t e^{-(t-s)\mathcal{A}^{\beta}}P(J_nB\cdot\nabla u - J_nu\cdot\nabla B)ds.$$
(45)

According to the proof of (35), energy inequality (35) can be rephrased as

$$\|u(t)\|_{L^{2}}^{2} + \|B(t)\|_{L^{2}}^{2} + \int_{s}^{t} (\|\mathcal{A}^{\frac{\alpha}{2}}u(r)\|_{L^{2}}^{2} + \|\mathcal{A}^{\frac{\beta}{2}}B(r)\|_{L^{2}}^{2})dr$$
  
$$\leq \|u(s)\|_{L^{2}}^{2} + \|B(s)\|_{L^{2}}^{2}, \qquad (46)$$

for  $0 \leq s < t$ .

Let us begin with the estimate of the nonlinear integrand of (44) in the  $L^2$  norm. Applying the divergence-free property and integrating by parts, we take a test function  $\phi \in L^2_{\sigma}(\Omega)^3$  to obtain

$$\langle e^{-t\mathcal{A}^{\alpha}} P(J_n B \cdot \nabla B - J_n u \cdot \nabla u), \phi \rangle$$
  
=  $\langle J_n u \otimes u, \nabla e^{-t\mathcal{A}^{\alpha}} \phi \rangle - \langle J_n B \otimes B, \nabla e^{-t\mathcal{A}^{\alpha}} \phi \rangle,$  (47)

Applying Hölder inequality to the previous identity and employing the estimates (18), (25) and (38), we have

$$\begin{aligned} \|e^{-t\mathcal{A}^{\alpha}}P(J_{n}B\cdot\nabla B-J_{n}u\cdot\nabla u)\|_{L^{2}} \\ &\leq Ct^{-\frac{1}{2\alpha}}(\|J_{n}u\|_{L^{4}}\|u\|_{L^{4}}+\|J_{n}B\|_{L^{4}}\|B\|_{L^{4}}) \\ &\leq Ct^{-\frac{1}{2\alpha}}(\|u\|_{L^{2}}^{2-\frac{3}{2\alpha}}\|\mathcal{A}^{\frac{\alpha}{2}}u\|_{L^{2}}^{\frac{3}{2\alpha}}+\|B\|_{L^{2}}^{2-\frac{3}{2\beta}}\|\mathcal{A}^{\frac{\beta}{2}}B\|_{L^{2}}^{\frac{3}{2\beta}}). \end{aligned}$$
(48)

With the application of (48), the  $L^2$  estimate of (44) becomes

$$\|u(t)\|_{L^{2}} - \|e^{-t\mathcal{A}^{\alpha}}u_{0}\|_{L^{2}}$$

$$\leq C \int_{0}^{t} (t-s)^{-\frac{1}{2\alpha}} (\|u(s)\|_{L^{2}}^{2-\frac{3}{2\alpha}} \|\mathcal{A}^{\frac{\alpha}{2}}u(s)\|_{L^{2}}^{\frac{3}{2\alpha}} + \|B(s)\|_{L^{2}}^{2-\frac{3}{2\beta}} \|\mathcal{A}^{\frac{\beta}{2}}B(s)\|_{L^{2}}^{\frac{3}{2\beta}}) ds.$$

$$(49)$$

Arguing in the same manner with the derivation of (49), we have the  $L^2$  estimate of (45) derived as

$$\|B(t)\|_{L^{2}} - \|e^{-t\mathcal{A}^{\beta}}B_{0}\|_{L^{2}}$$

$$\leq C \int_{0}^{t} (t-s)^{-\frac{1}{2\beta}} \|u(s)\|_{L^{2}}^{1-\frac{3}{4\alpha}} \|\mathcal{A}^{\frac{\alpha}{2}}u(s)\|_{L^{2}}^{\frac{3}{4\alpha}} \|B(s)\|_{L^{2}}^{1-\frac{3}{4\beta}} \|\mathcal{A}^{\frac{\beta}{2}}B(s)\|_{L^{2}}^{\frac{3}{4\beta}} ds.$$

$$(50)$$

Integrating (49)-(50) respectively, we have

After changing the integration order and integrating with respect to r, we have

$$\int_{0}^{t} \|u(s)\|_{L^{2}} ds + \int_{0}^{t} \|B(s)\|_{L^{2}} ds - \int_{0}^{t} \|e^{-s\mathcal{A}^{\alpha}} u_{0}\|_{L^{2}} ds - \int_{0}^{t} \|e^{-s\mathcal{A}^{\beta}} B_{0}\|_{L^{2}} ds 
\leq Ct^{1-\frac{1}{2\alpha}} \int_{0}^{t} (\|u(s)\|_{L^{2}}^{2-\frac{3}{2\alpha}} \|\mathcal{A}^{\frac{\alpha}{2}} u(s)\|_{L^{2}}^{\frac{3}{2\alpha}} + \|B(s)\|_{L^{2}}^{2-\frac{3}{2\beta}} \|\mathcal{A}^{\frac{\beta}{2}} B(s)\|_{L^{2}}^{\frac{3}{2\beta}} ds 
+ Ct^{1-\frac{1}{2\beta}} \int_{0}^{t} \|u(s)\|_{L^{2}}^{1-\frac{3}{4\alpha}} \|\mathcal{A}^{\frac{\alpha}{2}} u(s)\|_{L^{2}}^{\frac{3}{4\alpha}} \|B(s)\|_{L^{2}}^{1-\frac{3}{4\beta}} \|\mathcal{A}^{\frac{\beta}{2}} B(s)\|_{L^{2}}^{\frac{3}{4\beta}} ds.$$
(51)

By using Hölder inequality in (51) and then taking energy inequality (46) into account, the  $L^2$  estimate becomes

$$\int_{0}^{t} \|u(s)\|_{L^{2}} ds + \int_{0}^{t} \|B(s)\|_{L^{2}} ds - \int_{0}^{t} \|e^{-s\mathcal{A}^{\alpha}}u_{0}\|_{L^{2}} ds - \int_{0}^{t} \|e^{-s\mathcal{A}^{\beta}}B_{0}\|_{L^{2}} ds \\
\leq Ct^{1-\frac{1}{2\alpha}} t^{1-\frac{3}{4\alpha}} \left(\int_{0}^{t} \|u(s)\|_{L^{2}}^{\frac{8\alpha}{3}-2} \|\mathcal{A}^{\frac{\alpha}{2}}u(s)\|_{L^{2}}^{2} ds\right)^{\frac{3}{4\alpha}} \\
+ Ct^{1-\frac{1}{2\alpha}} t^{1-\frac{3}{4\beta}} \left(\int_{0}^{t} \|B(s)\|_{L^{2}}^{\frac{8\beta}{3}-2} \|\mathcal{A}^{\frac{\beta}{2}}B(s)\|_{L^{2}}^{2} ds\right)^{\frac{3}{4\beta}} \\
+ Ct^{1-\frac{1}{2\beta}} t^{1-\frac{3}{8\alpha}-\frac{3}{8\beta}} \left(\int_{0}^{t} \|u\|_{L^{2}}^{\frac{8\alpha}{3}-2} \|\mathcal{A}^{\frac{\alpha}{2}}u\|_{L^{2}}^{2} ds\right)^{\frac{3}{8\alpha}} \left(\int_{0}^{t} \|B\|_{L^{2}}^{\frac{8\beta}{3}-2} \|\mathcal{A}^{\frac{\beta}{2}}B\|_{L^{2}}^{2} ds\right)^{\frac{3}{8\beta}} \\
\leq C[t^{2-\frac{5}{4\alpha}} + t^{2-\frac{1}{2\alpha}-\frac{3}{4\beta}} + t^{2-\frac{3}{8\alpha}-\frac{7}{8\beta}}]\|(u_{0}, B_{0})\|_{L^{2}}^{2}.$$
(52)

On the other hand, it follows from (46) that

$$\|(u,B)(t)\|_{L^2} \le \frac{1}{t} \int_0^t \|(u,B)(s)\|_{L^2} ds.$$
(53)

Therefore, equation (52) becomes

$$\begin{aligned} \|(u,B)(t)\|_{L^{2}} &\leq \frac{1}{t} \int_{0}^{t} \|e^{-s\mathcal{A}^{\alpha}}u_{0}\|_{L^{2}}ds + \frac{1}{t} \int_{0}^{t} \|e^{-s\mathcal{A}^{\beta}}B_{0}\|_{L^{2}}ds \\ &+ C(t^{1-\frac{5}{4\alpha}} + t^{1-\frac{1}{2\alpha}-\frac{3}{4\beta}} + t^{1-\frac{3}{8\alpha}-\frac{7}{8\beta}}) \\ &\leq \frac{1}{t} \int_{0}^{t} \|e^{-s\mathcal{A}^{\alpha}}u_{0}\|_{L^{2}}ds + \frac{1}{t} \int_{0}^{t} \|e^{-s\mathcal{A}^{\beta}}B_{0}\|_{L^{2}}ds + Ct^{-\frac{1}{4\alpha}}, \end{aligned}$$
(54)

since t > 1 is assumed due to the energy inequality (46) and

$$\frac{5}{4\alpha} - 1 \ge \frac{1}{4\alpha}, \quad \frac{1}{2\alpha} + \frac{3}{4\beta} - 1 \ge \frac{1}{4\alpha}, \quad \frac{3}{8\alpha} + \frac{7}{4\beta} - 1 \ge \frac{1}{4\alpha}, \tag{55}$$

due to  $\alpha, \beta \leq 1$ . This shows the desired  $L^2$  decay (13) due to Lemma 2.2.

## 3.3. Proof of assertion (15)

By assumption (14), equation (54) implies

$$\|(u,B)(t)\|_{L^2} \le Ct^{-\gamma} + Ct^{-\frac{1}{4\alpha}}.$$
(56)

Here, we only consider the case  $t \ge 1$ , due to the uniform bound

$$||(u, B)(t)||_{L^2} \le ||(u_0, B_0)||_{L^2}$$

given by (46). Hence, we obtain (15) when  $\gamma \leq \frac{1}{4\alpha}$ . For  $\gamma > \frac{1}{4\alpha}$ , since  $1 \geq \alpha > \frac{3}{4}$ , we set p, q, r > 1 so that

$$\frac{1}{p} + \frac{3}{4\alpha} = 1, \ \frac{1}{q} + \frac{3}{4\beta} = 1, \ \frac{1}{r} + \frac{3}{8\alpha} + \frac{3}{8\beta} = 1.$$
(57)

By (14), (53), (46) and Hölder inequality, estimate (51) becomes

$$\begin{split} \|(u,B)(t)\|_{L^{2}} &\leq Ct^{-\gamma} + Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} \|u\|_{L^{2}}^{(2-\frac{3}{2\alpha})p} ds \right)^{\frac{1}{p}} \left( \int_{0}^{t} \|\mathcal{A}^{\frac{\alpha}{2}}u\|_{L^{2}}^{2} ds \right)^{\frac{3}{4\alpha}} \\ &+ Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} \|B\|_{L^{2}}^{(2-\frac{3}{2\beta})q} ds \right)^{\frac{1}{q}} \left( \int_{0}^{t} \|\mathcal{A}^{\frac{\beta}{2}}B\|_{L^{2}}^{2} ds \right)^{\frac{3}{4\beta}} \\ &+ Ct^{-\frac{1}{2\beta}} \left( \int_{0}^{t} \|u\|_{L^{2}}^{(1-\frac{3}{4\alpha})r} \|B\|_{L^{2}}^{(1-\frac{3}{4\beta})r} ds \right)^{\frac{1}{r}} \left( \int_{0}^{t} \|\mathcal{A}^{\frac{\alpha}{2}}u\|_{L^{2}}^{2} ds \right)^{\frac{3}{8\alpha}} \left( \int_{0}^{t} \|\mathcal{A}^{\frac{\beta}{2}}B\|_{L^{2}}^{2} ds \right)^{\frac{3}{8\beta}} \\ &\leq Ct^{-\gamma} + Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} \|u\|_{L^{2}}^{(2-\frac{3}{2\alpha})p} ds \right)^{\frac{1}{p}} + Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} \|B\|_{L^{2}}^{(2-\frac{3}{2\beta})q} ds \right)^{\frac{1}{q}} \\ &+ Ct^{-\frac{1}{2\beta}} \left( \int_{0}^{t} \|u\|_{L^{2}}^{(1-\frac{3}{4\alpha})r} \|B\|_{L^{2}}^{(1-\frac{3}{4\beta})r} ds \right)^{\frac{1}{r}}. \end{split}$$

The desired decay estimate will be derived by a boot strap iteration scheme. To do so, we begin with the step  $\gamma_1 = \frac{1}{4\alpha}$  so that

$$||(u,B)(t)||_{L^2} \le Ct^{-\gamma_1}.$$
(59)

Assuming the existence of  $\gamma_n$  so that  $0 < \gamma_n < \gamma \leq \frac{1}{2}$  for a positive integer n and there holds the decay estimate

$$||(u,B)(t)||_{L^2} \le Ct^{-\gamma_n},$$
(60)

we will show that the previous estimate can be upgrade to the one involving a power to be defined as  $-\gamma_{n+1}$ . Indeed, the application of (57) and (60) into (58), we have

$$\begin{aligned} \|(u,B)(t)\|_{L^{2}} &\leq Ct^{-\gamma} + Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} s^{-\gamma_{n}(2-\frac{3}{2\alpha})p} ds \right)^{\frac{1}{p}} + Ct^{-\frac{1}{2\alpha}} \left( \int_{0}^{t} s^{-\gamma_{n}(2-\frac{3}{2\beta})q} ds \right)^{\frac{1}{q}} \\ &+ Ct^{-\frac{1}{2\beta}} \left( \int_{0}^{t} s^{-\gamma_{n}(1-\frac{3}{4\alpha})r} s^{-\gamma_{n}(1-\frac{3}{4\beta})r} ds \right)^{\frac{1}{r}} \\ &\leq C[t^{-\gamma} + t^{-\frac{1}{2\alpha}} (t^{\frac{1}{p}} - \gamma_{n}(2-\frac{3}{2\alpha}) + t^{\frac{1}{q}} - \gamma_{n}(2-\frac{3}{2\beta})) + t^{-\frac{1}{2\beta}} t^{\frac{1}{r}} - \gamma_{n}(2-\frac{3}{4\alpha}-\frac{3}{4\beta})] \\ &= C[t^{-\gamma} + t^{1-\frac{5}{4\alpha}} - \gamma_{n}(2-\frac{3}{2\alpha}) + t^{1-\frac{2}{4\alpha}} - \frac{3}{4\beta} - \gamma_{n}(2-\frac{3}{2\beta}) + t^{1-\frac{3}{8\alpha}} - \frac{7}{8\beta} - \gamma_{n}(2-\frac{3}{4\alpha}-\frac{3}{4\beta})], \quad (61) \end{aligned}$$

since  $0 < 2\gamma_n < 1$ ,  $\alpha, \beta > \frac{3}{4}$  and

$$\gamma_n(2-\frac{3}{2\alpha})p = \gamma_n(2-\frac{3}{2\beta})q = \gamma_n(2-\frac{3}{4\alpha}-\frac{3}{4\beta})r = 2\gamma_n < 1.$$

due to (57). Rewrite (61) as

$$\|(u,B)(t)\|_{L^2} \leq C(t^{-\gamma} + t^{-a_n} + t^{-b_n} + t^{-c_n}),$$
(62)

with

$$a_n = (\frac{5}{4\alpha} - 1) + \gamma_n (2 - \frac{3}{2\alpha}),$$
  

$$b_n = (\frac{2}{4\alpha} + \frac{3}{4\beta} - 1) + \gamma_n (2 - \frac{3}{2\beta}),$$
  

$$c_n = \frac{3}{8\alpha} + \frac{7}{8\beta} - 1 + \gamma_n (2 - \frac{3}{4\alpha} - \frac{3}{4\beta}).$$

We see that

$$a_n - b_n = \frac{3}{4\alpha} - \frac{3}{4\beta} - 2\gamma_n(\frac{3}{4\alpha} - \frac{3}{4\beta}) = \frac{3}{4}(\frac{1}{\alpha} - \frac{1}{\beta})(1 - 2\gamma_n),$$
(63)

$$a_n - c_n = \frac{7}{8\alpha} - \frac{7}{8\beta} - \gamma_n (\frac{3}{4\alpha} - \frac{3}{4\beta}) = \frac{3}{4} (\frac{1}{\alpha} - \frac{1}{\beta}) (\frac{7}{6} - \gamma_n),$$
(64)

$$b_n - c_n = \frac{1}{8\alpha} - \frac{1}{8\beta} + \gamma_n (\frac{3}{4\alpha} - \frac{3}{4\beta}) = \frac{3}{4} (\frac{1}{\alpha} - \frac{1}{\beta}) (\frac{1}{6} + \gamma_n).$$
(65)

This implies

$$a_n \ge b_n \ge c_n$$
, when  $\frac{1}{\alpha} - \frac{1}{\beta} > 0$ .

Hence we set

$$\gamma_{n+1} = c_n, \text{ for } \frac{1}{\alpha} - \frac{1}{\beta} > 0.$$
(66)

This together with (62) gives

$$||(u,B)(t)||_{L^2} \le Ct^{-\gamma} + Ct^{-\gamma_{n+1}}.$$
(67)

The definition of  $\gamma_{n+1}$  shows

$$\gamma_{n+1} = \left(\frac{3}{8\alpha} + \frac{7}{8\beta} - 1\right) \sum_{m=0}^{n-1} \left(2 - \frac{3}{4\alpha} - \frac{3}{4\beta}\right)^m + \gamma_1 \left(2 - \frac{3}{4\alpha} - \frac{3}{4\beta}\right)^n.$$

The limit of this summation is, as  $n \to \infty$ ,

$$\frac{\frac{3}{8\alpha} + \frac{7}{8\beta} - 1}{1 - (2 - \frac{3}{4\alpha} - \frac{3}{4\beta})} = \frac{\frac{3}{8\alpha} + \frac{7}{8\beta} - 1}{\frac{3}{4\alpha} + \frac{3}{4\beta} - 1} \begin{cases} = \frac{1}{2}, \text{ when } \beta = 1, \\ > \frac{1}{2}, \text{ when } \beta < 1. \end{cases}$$

Hence, there is a positive integer  $n_0$  so that  $\gamma_{n_0+1} \geq \gamma$ . Thus (67) becomes

$$\|(u,B)(t)\|_{L^2} \le Ct^{-\gamma} + Ct^{-\gamma_{n_0+1}} \le Ct^{-\gamma}.$$
(68)

On the other hand, when  $\frac{1}{\alpha} - \frac{1}{\beta} \leq 0$ , we see from (63)-(65) that

$$a_n \le b_n \le c_n. \tag{69}$$

We thus define  $\gamma_{n+1} = a_n$  and so

$$\gamma_{n+1} = \left(\frac{5}{4\alpha} - 1\right) + \gamma_n \left(2 - \frac{3}{2\alpha}\right)$$
$$= \left(\frac{5}{4\alpha} - 1\right) \sum_{m=0}^{n-1} \left(2 - \frac{3}{2\alpha}\right)^m + \gamma_1 \left(2 - \frac{3}{2\alpha}\right)^n$$

The limit of this summation is, as  $n \to \infty$ ,

$$\left(\frac{5}{4\alpha} - 1\right) \sum_{m=0}^{\infty} (2 - \frac{3}{2\alpha})^m = \frac{5 - 4\alpha}{6 - 4\alpha} \begin{cases} = \frac{1}{2}, \text{ when } \alpha = 1, \\ > \frac{1}{2}, \text{ when } \alpha < 1. \end{cases}$$
(70)

This together with (62) and (69) implies the existence of an integer  $n_0 \ge 1$ so that  $\gamma_{n_0+1} \ge \gamma$  and hence

$$\|(u,B)(t)\|_{L^2} \le C(t^{-\gamma} + t^{-\gamma_{n_0+1}}) \le Ct^{-\gamma}.$$
(71)

This gives (15) and completes the proof of Theorem 1.1.

Acknowledgment. This research is supported by The Shenzhen Natural Science Fund of China (Stable Support Plan Program No. 20220805175116001).

## References

- [1] W. Borchers, T. Miyakawa, Algebraic  $L^2$  decay for Navier-Stokes flows in exterior domains. Acta Math. 165 (1990), 189-227.
- [2] W. Borchers, T. Miyakawa, L<sup>2</sup>-Decay for Navier-Stokes Flows in Unbounded Domains, with Application to Exterior Stationary Flows. Arch. Rational Mech. Anal. 118 (1992), 273-295.
- [3] W. Borchers, T. Miyakawa, On stability of exterior stationary Navier-Stokes flows. Acta Math. 174 (1995), 311-382.
- [4] Z. M. Chen, Y. Kagei, T. Miyakawa, Remarks on stability of purely conductive steady states to the exterior Boussinesq problem, Adv. Math. Sci. Appl. 1 (1992), 411-430.
- [5] Z. M. Chen, Solutions of the stationary and nonstationary navier-stokes equations in exterior domains, Pacific J. Math. 159 (1993), 227-240.
- [6] Q. Duan, Y. Xiao, Z. Xin, On the vanishing dissipation limit for the incompressible MHD equations on bounded domains, Sci. China Math. 65 (2022), 31-50.
- [7] L. Landau, E. Lifchitz, Electrodynamique des milieux continus, Physique thkorique, tome VIII, MIR, Moscow, 1969.
- [8] Z. Liu, P. Han, Decay for turbulent solutions of the magnetohydrodynamic equations in an exterior domain. J. Math. Phys. 61,091506 (2020).

- [9] K. Masuda, On the stability of incompressible viscous fluid motions past objects. J. Math. Soc. Japan, Vol. 27, No. 2, 1975.
- [10] T. Miyakawa, On nonstationary solutions of the Navier-Stokes equations in an exterior domain. Hiroshima Math. J. 12 (1982), 115-140.
- [11] M. Sermange, R. Temam, Some Mathematical questions related to the MHD equations, Comm. Pure Appl. Math., 36 (1983), 636-664
- [12] M. E. Schonbek, L<sup>2</sup> decay for weak solutions of the Navier-Stokes equations, Arch. Rational Mech. Anal. 88 (1985), 209-222.
- [13] M. E. Schonbek, T. P. Schonbek, S. Endre, Large-time behaviour of solutions to the magnetohydrodynamics equations, Math. Ann., 304 (1996), 717-756.
- [14] L. de Simon, Un'applicazione della teoria degli integrali singolari allo studio delle equazioni differenziali lineari astratte del primo ordine. Rend. Sem. Mat. Univ. Padova 34 (1964), 205-223.
- [15] R. Teman, Navier-Stokes Equations, North-Holland, Amsterdam, 1977.
- [16] K. Yosida, Functional Analysis, Sixth Edition, Springer, New York, 1980.