

Categorification of Biquandle Arrow Weight Invariants via Quivers

Sam Nelson*

Migiwa Sakurai†

Abstract

Introduced in [8], biquandle arrow weights invariants are enhancements of the biquandle counting invariant for oriented virtual and classical knots defined from biquandle-colored Gauss diagrams using a tensor over an abelian group satisfying certain properties. In this paper we categorify the biquandle arrow weight polynomial invariant using biquandle coloring quivers, obtaining new infinite families of polynomial invariants of oriented virtual and classical knots.

KEYWORDS: Biquandles, homsets, enhancements, categorification, virtual knots, Gauss diagrams, biquandle arrow weights, quivers

2020 MSC: 57K12

1 Introduction

Biquandles are algebraic structures encoding the Reidemeister moves in knot theory. Associated to every oriented virtual knot K is a *fundamental biquandle* $\mathcal{B}(K)$ which is conjectured to be a complete invariant up to a type of reversed mirror image [5]. Given a finite biquandle X , the set of biquandle homomorphisms from $\mathcal{B}(K)$ to X , $\text{Hom}(\mathcal{B}(K), X)$, is an invariant of classical and virtual knots known as the *biquandle homset invariant*. Elements of the homset can be represented as virtual knot diagrams with elements of X assigned to the semiarcs or as Gauss diagrams with elements of X assigned to the segments of the circle between arrows satisfying the biquandle coloring condition; these X -colored Gauss diagrams are analogous to matrices representing linear transformations with respect to a choice of basis.

In [8] the authors introduced an infinite family of enhancements of the biquandle homset invariant via structures called *biquandle arrow weights* which are analogous to biquandle cocycles arising not from crossings in a virtual knot diagram but from arrow crossings in X -colored Gauss diagrams.

In [4] the first author introduced a categorification of the quandle homset invariant known as the *quandle colorings quiver* associated to each subset of the set of endomorphisms of the coloring quandle X . In [2, 3, 7] this idea was applied to categorify other homset enhancements including the quandle cocycle enhancement, the quandle module enhancement, the psyquandle homset invariant and more. From these categorifications, new polynomial invariants were defined via decategorification including two-variable polynomial invariants and enhancements of the in-degree polynomial decategorification.

In this paper we apply the same idea to categorify the biquandle arrow weight invariant, in the process obtaining new polynomial invariants of virtual knots via decategorification. The paper is organized as follows. In Section 2 we briefly review the basics of biquandles, Gauss diagrams and arrow weights. In Section 3 we introduce biquandle arrow weight quivers and obtain some new polynomial invariants of classical and virtual knots via decategorification. We observe that this is an infinite family of invariants with several choices of parameters available including target biquandle X , abelian group A , biquandle arrow weight W and subset S of the set of endomorphisms of X . In Section 4 we collect some examples to illustrate the computation of the new invariants and make tables of the invariant values for a sampling of virtual knots for some choices of parameters (X, A, W, S) . We conclude in Section 5 with some questions for future research.

*Email: Sam.Nelson@cmc.edu. Partially supported by Simons Foundation collaboration grant 702597.


†Email: migiwa@shibaura-it.ac.jp

This paper, including all text, illustrations, and python code for computations, was written strictly by the authors without the use of generative AI in any form.

2 Review of Biquandles, Gauss Diagrams and Arrow Weights

In order to introduce our new categorification of the Biquandle Arrow Weight invariant, it will be useful to briefly recall the original invariant. More detail can be found in our earlier paper [8].

A *biquandle* is a set X with two binary operations $\triangleright, \bar{\triangleright}$ satisfying certain axioms (see [5] or [8]) derived from the Reidemeister moves. More precisely, a *biquandle coloring* or an X -*coloring* of an oriented knot or link diagram D by a biquandle X assigns an element of an element of X to each semiarc in D so that at every crossing we have the condition




called the *biquandle coloring condition*.

The biquandle axioms are then the conditions required such that given any biquandle coloring of a diagram before a Reidemeister move, there exists a unique coloring of the diagram after the move which agrees with the original coloring outside the neighborhood of the move. Then, it follows that the number of colorings of an oriented knot or link diagram by a finite biquandle X is a (possibly zero) integer-valued invariant, denoted $\Phi_X^{\mathbb{Z}}$, called the *biquandle counting invariant*.


In particular, associated to every oriented classical or virtual knot K there is a *fundamental biquandle* $\mathcal{B}(K)$ defined from any diagram of L with a presentation given by generators associated to semiarcs and relations at the crossings. An X -coloring then determines and is determined by a unique *biquandle homomorphism* $f : \mathcal{B}(K) \rightarrow X$. Then the sets $\text{Hom}(\mathcal{B}(K), X)$ of biquandle homomorphisms (known as the *biquandle homset*) is an invariant of K . More precisely, choosing a different diagram for K changes the way each homset element is represented, with X -colored diagrams representing the same homset element iff they are related by X -colored Reidemeister moves. Choosing a diagram for K is analogous to choosing input and output bases for vector spaces in linear algebra, with X -colored diagrams representing homset elements analogously to matrices representing linear transformations with Reidemeister moves playing the role of change of basis matrices.

Let K be a virtual knot and D a virtual knot diagram of an oriented virtual knot K . Then, D may be regarded as the image $f(\mathbb{S}^1)$ of a generic immersion $f : \mathbb{S}^1 \rightarrow \mathbb{R}^2$. Recall that a *Gauss diagram* for D is the preimage of D with chords, each of which connects the preimages of each real crossing. We can specify crossing information of each real crossing on the corresponding chord by directing the chord toward the undercrossing point and assigning each chord with the sign (i.e., the local writhe) of the crossing as shown:




A *biquandle coloring* of a Gauss diagram by a biquandle X is then an assignment of an element of X to

each segment of the circle between the arrowheads and tails such that at every arrow we have as shown:



In particular, to each arrow in a biquandle-colored Gauss diagram we associate an ordered pair (x, y) of biquandle elements together with a sign. The set of biquandle colorings of a Gauss diagram D representing a virtual knot K by a biquandle X can be identified with the homset $\text{Hom}(\mathcal{B}(K), X)$ with each X -coloring representing a homset element and with a choice of diagram D being analogous to a choice of basis for a vector space.

Finally, recall from [8] that a *biquandle arrow weight* is a function ϕ sending pairs of pairs of biquandle elements to elements in an abelian group A satisfying axioms which can be found in [8]. The biquandle arrow weight axioms are chosen so that the sum Σ_D of $\phi((a, b), (c, d))$ values over the set of arrow crossings with arrows labeled (a, b) and (c, d) and sign given by the product of arrow signs over the set of arrows in an X -colored Gauss diagram



is unchanged by Gauss diagram Reidemeister moves.

Given a finite biquandle X and biquandle arrow weight W with values in an abelian group A , for any oriented classical or virtual knot K the multiset of values over the set of X -colorings of any Gauss diagram D representing K is an invariant of knots, known as the *biquandle arrow weight multiset*, denoted $\Phi_X^{W, M}(K)$. We can convert this multiset into a “polynomial” form by summing over the multiset a formal variable raised to the “power” of each multiset element, obtaining the *biquandle arrow weight polynomial* $\Phi_X^W(K)$. We note that depending on A , the “polynomial” may be regarded as a bookkeeping device rather than a true polynomial.

We then have:

Theorem 1. ([8]) *The biquandle arrow weight multiset and polynomial are invariants of classical and virtual knots.*


Example 1. Let X be the biquandle specified by the operation tables

\triangleright	1	2	$\bar{\triangleright}$	1	2
1	2	2	1	2	2
2	1	1	2	1	1

and let $A = \mathbb{Z}_{16}$. Then we compute via `python` that we have biquandle arrow weight given by

$$\begin{bmatrix} \begin{bmatrix} 0 & 4 \\ 4 & 0 \end{bmatrix} & \begin{bmatrix} 4 & 0 \\ 8 & 12 \end{bmatrix} \\ \begin{bmatrix} 4 & 8 \\ 0 & 12 \end{bmatrix} & \begin{bmatrix} 0 & 12 \\ 12 & 0 \end{bmatrix} \end{bmatrix}$$

among others. The virtual trefoil knot 2.1 then has two X -colorings



yielding biquandle arrow weight multiset $\Phi_X^{W,M}(2.1) = \{8, 8\}$ and polynomial $2u^8$.

3 Biquandle Arrow Weight Quivers


We can now introduce our infinite family of categorifications of the biquandle arrow weight invariants.

Let K be an oriented virtual knot represented by a Gauss diagram D (see [5, 6]). Let X be a finite biquandle, A an abelian group and β a biquandle arrow weight for X with coefficients in A .

Recall that a map $\phi : X \rightarrow X$ is a *biquandle endomorphism* if for all $x, x' \in X$ we have

$$\phi(x \triangleright x') = \phi(x) \triangleright \phi(x') \text{ and } \phi(x \bar{\triangleright} x') = \phi(x) \bar{\triangleright} \phi(x').$$

Given an X -coloring of D , if we apply ϕ to all of the semiarc colors the result is also an X -coloring of D .



It then follows, as in [4], that the directed graph or *quiver* with a vertex for each X -coloring of D and directed edge connecting each coloring to its image under each $\phi \in S$ is unchanged by Reidemeister moves

and hence defines an invariant of virtual knots, which we call the *biquandle coloring quiver* $BCB_{X,S}(K)$. When $S = \text{Hom}(X, X)$ is the complete set of endomorphisms, we say that $BCB_{X,S}(K)$ is the *full quiver*.

Since the vertices in the quiver represent the elements of the biquandle homset, given a biquandle arrow weight W we can associate to each vertex in the quiver the corresponding Σ_D value to obtain a *weighted quiver* $\mathcal{Q}_{X,W,S}(K)$, i.e. a quiver weighted with elements of A at the vertices. We then have our main result:

Proposition 2. *The weighted quiver $\mathcal{Q}_{X,W,S}(K)$ is an invariant of oriented classical and virtual knots.*

Proof. All of the ingredients are already invariant under classical and virtual Reidemeister moves. \square

Quivers are categories with vertices as objects and directed paths as morphisms. Thus, this weighted quiver categorifies the biquandle arrow weight polynomial, which we can now regard as a decategorification by summing over the vertices terms of the form $u^{w(v)}$ where $w(v)$ is the weight at each vertex.

We additionally define two standard polynomial decategorifications, yielding new polynomial invariants of classical and virtual knots.

Definition 1. Let X be a finite biquandle, $S \subset \text{Hom}(X, X)$ a set of biquandle endomorphisms, A an abelian group and W a biquandle arrow weight with A coefficients. Then for any oriented classical or virtual knot or link K represented by a Gauss diagram D , we define

- *The arrow weight in-degree polynomial* to be the sum, over all vertices v in the vertex set V of the quiver, of products of a formal variable u to the power of the vertex weight $w(v) = \Sigma_D$ at the vertex v times a formal variable w to the power of the in-degree of the vertex v (the number of arrows directed in to v),

$$\Phi_{X,W,S}^{\text{deg}_+}(K) = \sum_{v \in V} u^{w(v)} w^{\text{deg}_+(v)}$$

and we further define

- *The arrow weight two-variable polynomial* to be the sum, over all edges e in the edge set E of the quiver, of products of a formal variable s to the power of the vertex weight at the source vertex $S(e)$ times a formal variable t to the power of the vertex weight at the target vertex $T(e)$,

$$\Phi_{X,W,S}^2(K) = \sum_{e \in E} s^{S(e)} t^{T(e)}.$$

Proposition 3. *The polynomials $\Phi_{X,W,S}^{\text{deg}_+}(K)$ and $\Phi_{X,W,S}^2(K)$ are invariants of oriented classical and virtual knots.*

Proof. These are determined by the biquandle arrow weighted quiver, which is invariant. \square

We can also make use of the weighting of the vertices to define a new invariant quiver-valued invariant from which we can extract new polynomial invariants as decategorifications.

Definition 2. Let X be a biquandle, W a biquandle arrow weight with coefficients in an abelian group A , and $S \subset \text{Hom}(X, X)$ a set of biquandle endomorphisms. Then let us form the quotient $\mathcal{Q}_{X,W,S}^Q(K)$ of the weighted quiver $\mathcal{Q}_{X,W,S}(K)$ by identifying the sets of vertices which share a common weight. We will call this the *arrow weighted quotient quiver*.

Proposition 4. *The arrow weighted quotient quiver $\mathcal{Q}_{X,W,S}^Q(K)$ is an invariant of classical and virtual knots.*

Proof. The arrow weighted quiver $\mathcal{Q}_{X,W,S}(K)$ is already unchanged by Reidemeister moves before taking the quotient; it follows that the quotient is also unchanged by Reidemeister moves. \square

One easy polynomial invariant we can obtain from $\mathcal{Q}_{X,W,S}^Q(K)$ is defined by letting each vertex contribute its weight and number of loops:

Definition 3. Let X be a biquandle, W a biquandle arrow weight with coefficients in an abelian group A , and $S \subset \text{Hom}(X, X)$ a set of biquandle endomorphisms. Let $\mathcal{Q}_{X,W,S}^Q(K)$ be the resulting arrow weighted quotient quiver, and let its vertex set be V . Then the *arrow weighted quotient quiver loop polynomial* is

$$\Phi_{X,W,S}^{Q,L}(K) = \sum_{v \in V} L(v)x^{W(v)}$$


where $W(v)$ is the biquandle arrow weight at the vertex v and $L(v)$ is the number of loops in the quiver based at v .

Corollary 5. *The arrow weighted quotient quiver loop polynomial $\Phi_{X,W,S}^{Q,L}(K)$ is an invariant of classical and virtual knots.*

4 Examples

In this section we compute some examples of biquandle arrow weight quivers and their decategorification polynomial invariants. We remark that these are toy examples in the sense that the true power of this infinite family of invariants lies in using larger biquandles and coefficient groups than we are currently able to compute.

Example 2. Let X , A and W be the biquandle and biquandle arrow weight from Example 1. Then the virtual knot 2.1 has biquandle arrow weight quiver



with respect to the full endomorphism set $S = \{[1, 2], [2, 1]\}$ where we specify an endomorphism $\sigma : X \rightarrow X$ where $X = \{x_1, \dots, x_n\}$ with a vector $\sigma = [\sigma(x_1), \dots, \sigma(x_n)]$ of the images of elements of X .

Example 3. Let X be the biquandle given by the operation tables

\triangleright	1	2	3	\triangleright	1	2	3
1	3	3	3	1	3	3	3
2	1	1	1	2	1	1	1
3	2	2	2	3	2	2	2

with endomorphism set $S = \{[3 \ 1 \ 2], [1 \ 2 \ 3], [2 \ 3 \ 1]\}$. We then compute that the 4-tensor

$$\left[\begin{array}{c} \left[\begin{array}{ccc} 0 & 1 & 3 \\ 3 & 0 & 1 \\ 1 & 3 & 0 \end{array} \right] \\ \left[\begin{array}{ccc} 3 & 4 & 0 \\ 0 & 3 & 4 \\ 4 & 0 & 3 \end{array} \right] \\ \left[\begin{array}{ccc} 1 & 0 & 4 \\ 4 & 1 & 0 \\ 0 & 4 & 1 \end{array} \right] \end{array} \right] \left[\begin{array}{c} \left[\begin{array}{ccc} 1 & 0 & 4 \\ 4 & 1 & 0 \\ 0 & 4 & 1 \end{array} \right] \\ \left[\begin{array}{ccc} 0 & 1 & 3 \\ 3 & 0 & 1 \\ 1 & 3 & 0 \end{array} \right] \\ \left[\begin{array}{ccc} 3 & 4 & 0 \\ 0 & 3 & 4 \\ 4 & 0 & 3 \end{array} \right] \end{array} \right] \left[\begin{array}{c} \left[\begin{array}{ccc} 3 & 4 & 0 \\ 0 & 3 & 4 \\ 4 & 0 & 3 \end{array} \right] \\ \left[\begin{array}{ccc} 1 & 0 & 4 \\ 4 & 1 & 0 \\ 0 & 4 & 1 \end{array} \right] \\ \left[\begin{array}{ccc} 0 & 1 & 3 \\ 3 & 0 & 1 \\ 1 & 3 & 0 \end{array} \right] \end{array} \right]$$

defines a biquandle arrow weight with coefficients in $A = \mathbb{Z}_8$.

Computing the in-degree polynomials, we obtain the table

$\Phi_{X,W,S}^{\text{deg}_+}(K)$	K
$3w^3$	3.3, 3.4, 3.5, 3.6, 3.7, 4.1, 4.2, 4.3, 4.6, 4.7, 4.8, 4.12, 4.16, 4.21, 4.25, 4.31, 4.36, 4.43, 4.46, 4.53, 4.55, 4.56, 4.59, 4.65, 4.71, 4.73, 4.75, 4.76, 4.77, 4.85, 4.86, 4.89, 4.90, 4.92, 4.95, 4.96, 4.98, 4.99, 4.100, 4.101, 4.104, 4.105, 4.106, 4.107, 4.108
$3uw^3$	4.22, 4.62, 4.83, 4.87, 4.88
$3u^2w^3$	4.13, 4.15, 4.17, 4.20, 4.23, 4.29, 4.32, 4.34, 4.38, 4.41, 4.49, 4.51, 4.58, 4.61, 4.67, 4.72
$3u^3w^3$	4.24, 4.26, 4.42, 4.47, 4.80
$3u^4w^3$	2.1, 3.1, 3.2, 4.4, 4.5, 4.9, 4.14, 4.27, 4.30, 4.37, 4.44, 4.48, 4.52, 4.54, 4.60, 4.64, 4.69, 4.74, 4.82, 4.84, 4.91, 4.94, 4.102
$3u^5w^3$	4.66, 4.79, 4.93
$3u^6w^3$	4.10, 4.11, 4.18, 4.19, 4.33, 4.35, 4.40, 4.50, 4.57, 4.63, 4.68, 4.70
$3u^7w^3$	4.28, 4.39, 4.45, 4.78, 4.81, 4.97, 4.103

Since all of these virtual knots have the same unweighted in-degree polynomial, $3w^3$, this example shows that our new construction is a proper enhancement of the in-degree polynomial.


Example 4. Let X be the biquandle specified by the operation tables

\triangleright	1	2	3	4	\triangleright	1	2	3	4
1	1	3	4	2	1	1	1	1	1
2	2	4	3	1	2	4	4	4	4
3	3	1	2	4	3	2	2	2	2
4	4	2	1	3	4	3	3	3	3

Let $S = \{[1, 4, 2, 3], [1, 1, 1, 1], [1, 3, 4, 2], [1, 2, 3, 4]\} \subset \text{Hom}(X, X)$ and let W be the biquandle arrow weight with coefficient group $A = \mathbb{Z}_6$ specified by the 4-tensor


$$\left[\begin{array}{c} \left[\begin{array}{cccc} 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \\ 0 & 3 & 0 & 3 \end{array} \right] \left[\begin{array}{cccc} 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \\ 3 & 0 & 0 & 3 \end{array} \right] \left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \left[\begin{array}{cccc} 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \end{array} \right] \end{array} \right]$$

Then the virtual knots 3.1 and 3.3 have biquandle arrow weight quivers



with respect to X, S and W . In particular we note that the coloring quivers without the arrow weights are isomorphic, showing that the biquandle arrow weight quiver is a proper enhancement of the biquandle coloring quiver.

Example 5. Continuing with the virtual knots, biquandle, arrow weight and endomorphism set from Example 4, we obtain the quotient quivers



$$\Phi_{X,W,S}^{Q,L}(3.1)$$

$$\Phi_{X,W,S}^{Q,L}(3.3)$$

and quotient quiver loop polynomials $\Phi_{X,W,S}^{Q,L}(3.1) = 10x^0 + 1x^3$ and $\Phi_{X,W,S}^{Q,L}(3.3) = 6x^0 + 4x^3$.

Example 6. We computed the biquandle arrow weight 2-variable polynomials for a choice of orientation for each of the virtual knots with up to 4 classical crossings at [1] for the biquandle with operation tables

$$\begin{array}{c|ccc} \succeq & 1 & 2 & 3 \\ \hline 1 & 3 & 3 & 3 \\ 2 & 1 & 1 & 1 \\ 3 & 2 & 2 & 3 \end{array}, \begin{array}{c|ccc} \bar{\succeq} & 1 & 2 & 3 \\ \hline 1 & 3 & 3 & 3 \\ 2 & 1 & 1 & 1 \\ 3 & 2 & 2 & 3 \end{array},$$

arrow weight over $A = \mathbb{Z}_3$ given by the operation tensor

$$\left[\begin{array}{c} \left[\begin{array}{ccc} 0 & 1 & 2 \\ 2 & 0 & 1 \\ 1 & 2 & 0 \end{array} \right] \\ \left[\begin{array}{ccc} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{array} \right] \\ \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \end{array} \right] \left[\begin{array}{c} \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \\ \left[\begin{array}{ccc} 0 & 1 & 2 \\ 2 & 0 & 1 \\ 1 & 2 & 0 \end{array} \right] \\ \left[\begin{array}{ccc} 2 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 2 & 0 \end{array} \right] \\ \left[\begin{array}{ccc} 2 & 0 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 2 \end{array} \right] \end{array} \right] \left[\begin{array}{c} \left[\begin{array}{ccc} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{array} \right] \\ \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \\ \left[\begin{array}{ccc} 0 & 1 & 2 \\ 2 & 0 & 1 \\ 1 & 2 & 0 \end{array} \right] \end{array} \right]$$

and endomorphism set

$$S = \{[3, 2, 1], [1, 2, 3], [2, 3, 1]\}.$$

The results are in the table.

$\Phi_{X,W,S}^2(K)$	K
9	2.1, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 4.12, 4.14, 4.16, 4.21, 4.25, 4.26, 4.27, 4.28, 4.30, 4.31, 4.36, 4.37, 4.43, 4.44, 4.45, 4.46, 4.47, 4.48, 4.52, 4.53, 4.54, 4.55, 4.56, 4.59, 4.60, 4.64, 4.65, 4.69, 4.71, 4.73, 4.74, 4.75, 4.76, 4.77, 4.80, 4.81, 4.82, 4.83, 4.84, 4.85, 4.86, 4.87, 4.88, 4.89, 4.90, 4.91, 4.92, 4.93, 4.94, 4.95, 4.96, 4.97, 4.98, 4.99, 4.100, 4.101, 4.102, 4.103, 4.104, 4.105, 4.106, 4.107, 4.108
$9st$	4.11, 4.17, 4.22, 4.23, 4.32, 4.61, 4.62, 4.63, 4.66, 4.67, 4.68, 4.79
$9s^2t^2$	4.10, 4.13, 4.15, 4.18, 4.19, 4.20, 4.24, 4.29, 4.33, 4.34, 4.35, 4.38, 4.39, 4.40, 4.41, 4.42, 4.49, 4.50, 4.51, 4.57, 4.58, 4.70, 4.72, 4.78

Example 7. We computed the biquandle arrow weight quotient quiver loop polynomials for a choice of orientation for each of the virtual knots with up to 4 classical crossings at [1] for the biquandle with operation table

\triangleright	1	2	3	4	\triangleright	1	2	3	4
1	2	2	2	2	1	2	4	2	4
2	3	3	3	3	2	1	3	1	3
3	4	4	4	4	3	4	2	4	2
4	1	1	1	1	4	3	1	3	1

arrow weight over $A = \mathbb{Z}_4$ given by the tensor

$$\left[\begin{array}{c} \left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \\ \left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \\ \left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \\ \left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{array} \right]$$

and endomorphism set

$$S = \{[4, 1, 2, 3], [1, 2, 3, 4], [2, 3, 4, 1], [3, 4, 1, 2]\}.$$

The results are in the table.

$\Phi_{X,W,S}^{Q,L}(K)$	K
$4x^2 + 4$	2.1, 3.2, 3.3, 3.4, 4.3, 4.4, 4.5, 4.6, 4.9, 4.11, 4.13, 4.14, 4.15, 4.17, 4.18, 4.19, 4.20, 4.22, 4.23 4.24, 4.25, 4.26, 4.27, 4.28, 4.29, 4.30, 4.31, 4.33, 4.34, 4.35, 4.37, 4.38, 4.39, 4.40, 4.42, 4.44, 4.45, 4.46, 4.48, 4.49, 4.51, 4.52, 4.54, 4.57, 4.60, 4.61, 4.62, 4.63, 4.64, 4.66, 4.67, 4.69, 4.74, 4.78, 4.79, 4.81, 4.82, 4.83, 4.84, 4.87, 4.88, 4.92, 4.93, 4.94, 4.95, 4.97, 4.101, 4.103, 4.104
16	3.1, 3.6, 4.1, 4.8, 4.10, 4.16, 4.32, 4.41, 4.43, 4.47, 4.50, 4.55, 4.56, 4.58, 4.59, 4.68, 4.71, 4.72, 4.73, 4.75, 4.76, 4.77, 4.80, 4.89, 4.90, 4.96, 4.98, 4.99, 4.102, 4.105, 4.107, 4.108
$16x^2$	3.5, 3.7, 4.2, 4.7, 4.12, 4.21, 4.36, 4.53, 4.65, 4.70, 4.85, 4.86, 4.91, 4.100, 4.106

5 Questions

As mentioned in Section 4, faster search and computation algorithms for finding biquandle arrow weight quivers with respect to biquandles with larger cardinality and larger finite or infinite coefficient groups are needed. Our current custom `python` code can handle finding “toy” examples with biquandles of small cardinality and with small coefficient groups such as \mathbb{Z}_n for small n , but the true power of this infinite family of invariants awaits faster and more efficient algorithms for finding biquandle arrow weights and quivers.

What is the relationship between biquandle arrow weight quivers and (bi)quandle cocycle quivers as defined in [3]? Indeed, what is the algebraic structure in general of biquandle arrow weights, and how are they related to (bi)quandle cohomology?

What is the geometric meaning of a biquandle arrow weight and its associated quivers?

What other decategorifications are possible? What about further enhancements of the weighted quiver structure? How about functors to other categories?

References

- [1] D. Bar-Natan. The knot atlas http://katlas.org/wiki/Main_Page.
- [2] J. Cenicerros, A. Christiana, and S. Nelson. Psyquandle coloring quivers. *arXiv:2107.05668*, 2021.
- [3] K. Cho and S. Nelson. Quandle cocycle quivers. *Topology Appl.*, 268:106908, 10, 2019.
- [4] K. Cho and S. Nelson. Quandle coloring quivers. *J. Knot Theory Ramifications*, 28(1):1950001, 12, 2019.
- [5] M. Elhamdadi and S. Nelson. *Quandles—an introduction to the algebra of knots*. Student Mathematical Library. American Mathematical Society, 2015.
- [6] M. Goussarov, M. Polyak, and O. Viro. Finite-type invariants of classical and virtual knots. *Topology*, 39:1045–1068, 2000.
- [7] K. Istambouli and S. Nelson. Quandle module quivers. *J. Knot Theory Ramifications*, 29(12):2050084, 14, 2020.
- [8] S. Nelson and M. Sakurai. Biquandle arrow weight enhancements. *Internat. J. Math.*, 34(8):Paper No. 2350046, 13, 2023.

DEPARTMENT OF MATHEMATICAL SCIENCES
CLAREMONT MCKENNA COLLEGE
850 COLUMBIA AVE.
CLAREMONT, CA 91711

GRADUATE SCHOOL OF ENGINEERING AND SCIENCE, COLLEGE OF ENGINEERING,
SHIBAURA INSTITUTE OF TECHNOLOGY
307 FUKASAKU, MINUMA-KU, SAITAMA-SHI, SAITAMA, 337-8570, JAPAN