MULTIPLICATIVE DEPENDENCE IN LINEAR RECURRENCE SEQUENCES

ATTILA BÉRCZES, LAJOS HAJDU, ALINA OSTAFE, AND IGOR E. SHPARLINSKI

ABSTRACT. For a wide class of integer linear recurrence sequences $(u(n))_{n=1}^{\infty}$, we give an upper bound on the number of *s*-tuples $(n_1, \ldots, n_s) \in (\mathbb{Z} \cap [M+1, M+N])^s$ such that the corresponding elements $u(n_1), \ldots, u(n_s)$ in the sequence are multiplicatively dependent.

CONTENTS

1.1. Motivation and set-up11.2. Notation31.3. Main results32. Preliminaries42.1. Arithmetic properties of linear recurrence sequences42.2. Vertex covers63. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11References11	1. Introduction	1
1.3. Main results32. Preliminaries42.1. Arithmetic properties of linear recurrence sequences42.2. Vertex covers63. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	1.1. Motivation and set-up	1
2. Preliminaries42.1. Arithmetic properties of linear recurrence sequences42.2. Vertex covers63. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	1.2. Notation	3
2.1. Arithmetic properties of linear recurrence sequences42.2. Vertex covers63. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	1.3. Main results	3
2.2. Vertex covers63. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	2. Preliminaries	4
3. Proofs73.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	2.1. Arithmetic properties of linear recurrence sequences	4
3.1. Proof of Theorem 1.173.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	2.2. Vertex covers	6
3.2. Proof of Theorem 1.393.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	3. Proofs	7
3.3. Proof of Theorem 1.5104. Possible applications of our approach11Acknowledgement11	3.1. Proof of Theorem 1.1	7
4. Possible applications of our approach11Acknowledgement11	3.2. Proof of Theorem 1.3	9
Acknowledgement 11	3.3. Proof of Theorem 1.5	10
	4. Possible applications of our approach	11
References 11	Acknowledgement	11
	References	11

1. INTRODUCTION

1.1. Motivation and set-up. Let $\mathbf{u} = (u(n))_{n=1}^{\infty}$ be an integer linear recurrence sequence of order $d \ge 1$, that is, a sequence of integers satisfying a relation of the form

$$u(n+d) = c_{d-1}u(n+d-1) + \ldots + c_0u(n), \qquad n = 1, 2, \ldots,$$

²⁰²⁰ Mathematics Subject Classification. 11B37, 11D61.

Key words and phrases. Multiplicative dependence, linear recurrence sequences.

and not satisfying any shorter relation. In this case

$$f(X) = X^{d} - c_{d-1}X^{n+d-1} - \dots - c_{0} \in \mathbb{Z}[X]$$

is called the characteristic polynomial of **u**.

Recently there have been several works [3-6, 9-11, 13] investigating multiplicative relations of the form

(1.1)
$$u(n_1)^{k_1} \dots u(n_s)^{k_s} = 1.$$

However, these papers consider certain special cases. The works [6, 11, 13] are limited to the case of binary (that is, of order d = 2) linear recurrence sequences and also assume that the exponents k_1, \ldots, k_s are *fixed* non-zero integers, while the papers [3, 4, 9, 10] concern specific sequences. Under these restrictions, the mentioned papers contain several finiteness results. Finally, the recent work [5] concerns linear recurrence sequences of arbitrary order – however, under a rather restrictive condition on the coefficients c_i defining the generating relation.

Here we are interested in the case of general sequences of arbitrary order $d \ge 2$ and also we do not fix the exponents k_1, \ldots, k_s . Thus, we study *s*-tuples $(u(n_1), \ldots, u(n_s))$, which are *multiplicatively dependent* (m.d.), where, as usual, we say that the nonzero complex numbers $\gamma_1, \ldots, \gamma_s$ are m.d. if there exist integers k_1, \ldots, k_s , not all zero, such that

$$\gamma_1^{k_1} \dots \gamma_s^{k_s} = 1.$$

However, instead of finiteness results, we give an upper bound on the density of such s-tuples.

More precisely, for $M \ge 0$ and $N \ge 1$, we are interested in the following quantity

$$\mathsf{M}_s(M,N) = \sharp\{(n_1,\ldots,n_s) \in (\mathbb{Z} \cap [M+1,M+N])^s : u(n_1),\ldots,u(n_s) \text{ are m.d.}\}.$$

To estimate $M_s(M, N)$ we also study

$$\mathsf{M}_{s}^{*}(M,N) = \sharp\{(n_{1},\ldots,n_{s}) \in (\mathbb{Z} \cap [M+1,M+N])^{s}: u(n_{1}),\ldots,u(n_{s}) \text{ are } \mathsf{m.d. of maximal rank}\},\$$

where the maximality of the rank for m.d. of $(u(n_1), \ldots, u(n_s))$ means that no sub-tuple is m.d. In particular, this implies that if one has a m.d. (1.1) of maximal rank, then $k_1 \cdots k_s \neq 0$.

We can then estimate $M_s(M, N)$ via the inequality

(1.2)
$$\mathsf{M}_{s}(M,N) \leqslant \sum_{t=1}^{s} \binom{s}{t} \mathsf{M}_{t}^{*}(M,N) N^{s-t}.$$

1.2. Notation. We recall that the notations U = O(V), $U \ll V$ and $V \gg U$ are equivalent to $|U| \leq cV$ for some positive constant c, which throughout this work, may depend only on the integer parameter s and the sequence **u**.

It is convenient to denote by $\log_k x$ the k-fold iterated logarithm, that is, for $x \ge 1$ we set

$$\log_1 x = \log x$$
 and $\log_k = \log_{k-1} \max\{\log x, 2\}, k = 2, 3, \dots$

1.3. Main results. We say that the sequence **u** is *non-degenerate* if there are no roots of unity among the ratios of distinct roots of f. We say that the sequence **u** has a dominant root, if its characteristic polynomial f has a root λ with

$$|\lambda| > \max\{|\mu|: f(\mu) = 0, \ \mu \neq \lambda\}.$$

Furthermore, we say that \mathbf{u} is *simple* if f has no multiple roots.

Theorem 1.1. Let **u** be a simple non-degenerate sequence of order $d \ge 2$. For any fixed $s \ge 1$, uniformly over $M \ge 0$, we have

$$\mathsf{M}^*(M, N) \leq N^{s(1-1/(4d-3))+o(1)}$$
.

Analysing the proof of Theorem 1.1, one can see that for M = 0 we can drop o(1) in the bound.

Remark 1.2. Considering s-tuples with $n_1 = n_2$ we see that

(1.3)
$$\mathsf{M}_{s}(M,N) \ge N^{s-1}$$

Therefore, it is impossible to derive a bound of the same type as in Theorem 1.1 for $M_s(M, N)$.

When M is (exponentially) large compared to N, we get the following bound, which improves Theorem 1.1 for s < 4d - 3.

Theorem 1.3. Let **u** be a simple non-degenerate sequence of order $d \ge 2$ with a dominant root and let

$$M \ge \exp(N \log_3 N / \log_2 N).$$

Then, for any fixed $s \ge 1$, uniformly over M, we have

$$\mathsf{M}^*_s(M,N) \leqslant N^{s-1+o(1)}.$$

Remark 1.4. The condition on M in Theorem 1.3 is chosen to achieve the strongest possible bound. Examining its proof one can see that for s < 4d - 3 one can also improve Theorem 1.1 for $M \ge \exp(N^{\eta})$ with any $\eta > s/(4d - 3)$ (but only for sequences with a dominant root). From the definition of m.d. of maximal rank, we have $\mathsf{M}_1^*(M, N) = O(1)$, see [1, Lemma 2.1]. Hence, we see from (1.2) that in applying Theorem 1.1 to bounding $\mathsf{M}_s(M, N)$ the case of s = 2 becomes the bottleneck. Thus, we now investigate this case separately.

Theorem 1.5. Let **u** be a simple non-degenerate sequence of order $d \ge 2$ with a dominant root. Uniformly over $M \ge 0$, we have

$$\mathsf{M}_{2}^{*}(M, N) = N + O(1).$$

Since, as we have mentioned, $M_1^*(M, N) = O(1)$, the bounds of Theorems 1.1 and 1.5 inserted in (1.2) imply that if **u** is a simple nondegenerate sequence of order $d \ge 2$ with a dominant root then

$$\mathsf{M}_{s}(M, N) \ll N^{s-3/(4d-3)+o(1)},$$

where the bottleneck comes from the bound on $M_3^*(M, N)$.

If $M \ge \exp(N \log_3 N / \log_2 N)$, then using instead Theorem 1.3, one obtains the upper bound

$$\mathsf{M}_s(M,N) \ll N^{s-1+o(1)},$$

which matches the trivial lower bound (1.3).

2. Preliminaries

2.1. Arithmetic properties of linear recurrence sequences. In this section we collect various results about the arithmetic properties of a linear recurrence sequence that we need for our main results. These include:

- a lower bound of square-free parts of elements in **u**,
- a bound for the number of elements in **u** that are S-units,
- various results on congruences with elements in **u**,
- a result on the finiteness of perfect powers in **u**.

Some of these are obtained under the condition that \mathbf{u} has a dominant root.

We start with a lower bound of Stewart [17, Theorem 1] on the square-free part of elements in a linear recurrence.

For any integer m, we define rad(m) to be the largest square-free factor of m.

Lemma 2.1. Let **u** be a simple non-degenerate sequence of order $d \ge 2$ with a dominant root. Then there exist constants C_1 and C_2 , which are effectively computable only in terms of **u**, such that if $n \ge C_2$, then

$$\operatorname{rad}(u(n)) > n^{C_1(\log_2 n)/\log_3 n}$$

We also need the following upper bound from [15, Theorem 1 and Corollary] on the number of terms of \mathbf{u} composed out of primes from a given set. We note that the condition of the exponential growth of the terms of \mathbf{u} , assumed in [15], is now known to hold for non-degenerate recurrence sequences, see [8, 14]. Hence we have the following result.

Lemma 2.2. Let **u** be a non-degenerate sequence of order $d \ge 2$ and let S be an arbitrary set of r primes. Then, for $M \ge 0$, the number A(S; M, N) of terms $u(M + 1), \ldots, u(M + N)$, composed exclusively of primes from S, satisfies

$$A(S; M, N) \ll \begin{cases} rNM^{-1}\log(N+M) & \text{for } M \ge 1, \\ r(\log N)^2 & \text{for } M = 0. \end{cases}$$

We now present two results regarding solutions to certain congruences with elements in a linear recurrence sequence. We start with a result, which follows from [15, Lemma 2 and Lemma 3].

Lemma 2.3. Let **u** be a non-degenerate sequence of order $d \ge 2$ and let $m \ge 1$ be an integer. Then we have

$$\sharp\{n \in \mathbb{Z} \cap [M+1, M+N] : u(n) \equiv 0 \pmod{m}\} \ll N/\log m + 1.$$

The second bound that we need holds modulo primes and follows from [2, Lemma 6]. In [2] it is formulated only for the interval [1, N], however the result is uniform with respect to the sequence **u** and hence it holds uniformly with respect to M, too.

Let $\overline{\mathbb{F}}_p$ be the algebraic closure of the finite field \mathbb{F}_p of p elements.

Lemma 2.4. Let **u** be a simple sequence of order $d \ge 2$. Let $M \ge 0$ and $N \ge 1$. Let p be a prime, and set

$$\varrho_p = \min_{1 \le i < j \le d} r_{ij},$$

where r_{ij} is the multiplicative order of λ_i/λ_j in $\overline{\mathbb{F}}_p$, with the λ_i being the roots of the characteristic polynomial of **u**. Then we have

$$\sharp\{n \in \mathbb{Z} \cap [M+1, M+N] : u(n) \equiv 0 \pmod{p}\} \ll N(N^{-1} + \varrho_p^{-1})^{1/(d-1)}$$

The following result is certainly well-known and is based on classical ideas of Hooley [12], however for completeness we present a short proof.

Lemma 2.5. For $R \ge 2$ we consider the set

$$\mathcal{W}(R) = \{ p \ prime : \ \varrho_p \leq R \}.$$

Then $\sharp \mathcal{W}(R) \ll R^2 / \log R$.

Proof. Write $\lambda_1, \ldots, \lambda_q$ for the distinct roots of the characteristic polynomial of **u**.

For $R \ge 2$, let

$$Q(R) = \prod_{\rho \leqslant R} \prod_{1 \leqslant i < j \leqslant q} \operatorname{Nm}_{K/\mathbb{Q}}(\lambda_i^{\rho} - \lambda_j^{\rho}),$$

where $\operatorname{Nm}_{K/\mathbb{Q}}$ is the norm from the splitting field K of f to \mathbb{Q} . Note that $Q(R) \neq 0$ because λ_i/λ_j is not a root of unity and since λ_i and λ_j are algebraic integers we also have $Q(R) \in \mathbb{Z}$.

Clearly, for any prime p with $\rho_p \leq R$ we have $p \mid Q(R)$, hence $\sharp \mathcal{W}(R) \leq \omega(Q(R))$, where $\omega(k)$ is the number of prime divisors of an integer $k \geq 1$. As clearly $\omega(k)! \leq k$, by the Stirling formula we get

$$\sharp \mathcal{W}(R) \ll \frac{\log Q(R)}{\log \log Q(R)}.$$

Since obviously $\log Q(R) \ll R^2$, the result follows.

Finally, we need a result on the finiteness of perfect powers in linear recurrence sequences with a dominant root. The most general and convenient form for us, which is built on several previous results in this direction, is given by of Bugeaud and Kaneko [7, Theorem 1.1].

Lemma 2.6. Let **u** be a simple non-degenerate sequence of order $d \ge 2$ with a dominant root. Then the equation $u(n) = m^k$ has only finitely many solutions in integer $k \ge 2$, $m \ne 0$, $n \ge 1$.

2.2. Vertex covers. We need the following graph-theoretic result.

Lemma 2.7. Let G be a graph with vertex set \mathcal{V} , having no isolated vertex. Put $\ell = \sharp \mathcal{V}$. Then there exists $\mathcal{V}_1 \subseteq \mathcal{V}$ with $\sharp \mathcal{V}_1 \leq \ell/2$ such that for any $v_2 \in \mathcal{V}_2 = \mathcal{V} \setminus \mathcal{V}_1$ there exists a vertex $v_1 \in \mathcal{V}_1$ which is a neighbour of v_2 .

Proof. The statement must be well-known, but we give a simple proof. If \tilde{G} is a graph (without isolated vertices) obtained from G by omitting some edges, and the statement is valid for \tilde{G} , then the statement is obviously valid for G. Let \tilde{G} be a forest graph (that is, a graph without cycles) obtained from G by omitting some edges, such that the number of connected components of G and \tilde{G} are the same. Then \tilde{G} is a bipartite graph, so the statement is clearly valid for it. Hence the result follows.

3. Proofs

3.1. **Proof of Theorem 1.1.** Suppose that for some $n_1, \ldots, n_s \in [M + 1, M + N]$ the terms $u(n_1), \ldots, u(n_s)$ are m.d. of maximal rank, that is, we have (1.1) with some nonzero integers k_1, \ldots, k_s .

Choose a positive real number $R \ge 2$ to be specified later, and let $\mathcal{W}(R)$ be as in Lemma 2.5.

Write t for the number of indices $i = 1, \ldots, s$ for which $u(n_i)$ has a prime divisor $p_i \notin \mathcal{W}(R)$, and let r = s - t for the number of indices i with $u(n_i)$ having all prime divisors in $\mathcal{W}(R)$. Without loss of generality, we may assume that the corresponding integers are n_1, \ldots, n_t , and n_{t+1}, \ldots, n_s , respectively.

By Lemmas 2.2 and 2.5, for $M \ge 1$, the number K_1 of such *r*-tuples $(n_{t+1}, \ldots, n_s) \in [M+1, M+N]^r$ satisfies

(3.1)
$$K_1 \ll \left(\frac{R^2 N \log(N+M)}{M \log R}\right)^r.$$

If M = 0, then we have the bound

(3.2)
$$K_1 \ll \left(\frac{R^2(\log N)^2}{\log R}\right)^r.$$

We assume that such an r-tuple (n_{t+1}, \ldots, n_s) is fixed.

Consider the *t*-tuples $(n_1, \ldots, n_t) \in [M+1, M+N]^t$. Recall that for any $1 \leq i \leq t$, there is a prime $p_i \notin \mathcal{W}(R)$ such that $p_i \mid u(n_i)$.

Define the graph \mathcal{G} whose vertices are $u(n_1), \ldots, u(n_t)$, and connect the vertices $u(n_i)$ and $u(n_j)$ precisely when $gcd(u(n_i), u(n_j))$ has a prime divisor outside $\mathcal{W}(R)$. Observe that as $u(n_1), \ldots, u(n_s)$ are **m.d.** of maximal rank, \mathcal{G} has no isolated vertex. Thus, by Lemma 2.7, there exists a subset \mathcal{I} of $\{1, \ldots, t\}$ with

$$(3.3) m = \sharp \mathcal{I} \leqslant |t/2|$$

such that for any j with

$$j \in \{n_1, \ldots, n_t\} \setminus \mathcal{I}$$

the vertex $u(n_i)$ is connected with some $u(n_i)$ in \mathcal{G} , for some $i \in \mathcal{I}$.

Without loss of generality we may assume that $\mathcal{I} = \{1, \ldots, m\}$. Trivially, the number K_2 of such *m*-tuples $(n_1, \ldots, n_m) \in [M+1, M+N]^m$ satisfies

$$(3.4) K_2 \ll N^m$$

We now fix such an *m*-tuple. For $\ell = t - m$, we now count the number K_3 of the remaining ℓ -tuples $(n_{m+1}, \ldots, n_t) \in [M+1, M+N]^{\ell}$. Since each $u(n_j)$ with $m+1 \leq j \leq t$ has a common prime factor $p \notin \mathcal{W}(R)$

with $u(n_i)$ for some $1 \leq i \leq m$, by Lemma 2.4 we obtain that n_j comes from a set \mathcal{N} of cardinality

$$\# \mathcal{N} \ll N \left(N^{-1} + \varrho_p^{-1} \right)^{1/(d-1)} \leqslant N \left(N^{-1} + R^{-1} \right)^{1/(d-1)}.$$

Thus we obtain

(3.5)
$$K_3 \leq (\sharp \mathcal{N})^{\ell} \ll \left(N \left(N^{-1} + R^{-1} \right)^{1/(d-1)} \right)^{t-m}.$$

We consider now two cases based on $M \leq N \log N$ or $M > N \log N$. If $M \leq N \log N$, then

$$\mathsf{M}_{s}^{*}(M, N) \leqslant \mathsf{M}_{s}^{*}(0, 2N \log N),$$

therefore we reduce to counting s-tuples in the interval $[0, 2N \log N]^s$.

Putting together the bounds (3.2), (3.4) and (3.5) (with N replaced by $2N \log N$), for some non-negative integer $t \leq s$ and r = s - t, we obtain

(3.6)

$$M_{s}^{*}(M, N) \leq K_{1}K_{2}K_{3} \\ \ll \left(R^{2}(\log N + \log \log N)^{2}/\log R\right)^{r} (N \log N)^{m} \\ \left(N \log N \left(N^{-1} + R^{-1}\right)^{1/(d-1)}\right)^{t-m} \\ \leq N^{t+o(1)}R^{2r} \left(\left(N^{-1} + R^{-1}\right)^{1/(d-1)}\right)^{t/2},$$

where the last inequality comes from (3.3).

Letting $R = N^{\eta}$ with some $0 < \eta < 1/2$, we obtain

$$\mathsf{M}^*_s(M,N) \ll N^{t+2\eta r - \eta t/(2(d-1)) + o(1)} = N^{2\eta s + (1-2\eta)t - \eta t/(2(d-1)) + o(1)} + O(1) + O(1)$$

Writing t = zs (and noting that $0 \le z \le 1$), the exponent of the last term above (omitting the expression o(1)) is given by

$$f_{\eta}(z) = \frac{s}{2(d-1)}((2d - 4d\eta + 3\eta - 2)z + 4\eta(d-1)).$$

So taking

$$\eta = \frac{2(d-1)}{4d-3},$$

(to make $f_{\eta}(z)$ a constant), we obtain

$$\mathsf{M}^*_s(M,N) \ll N^{2\eta s + o(1)} = N^{s - s/(4d - 3) + o(1)},$$

which concludes this case.

If $M > N \log N$, then the bound (3.1) becomes

$$K_1 \ll (R^2/\log R)^r.$$

Putting this together with (3.4) and (3.5), we obtain (3.6) without the $(\log N)^2$ factor, that is,

$$\begin{aligned} \mathsf{M}_{s}^{*}(M,N) &\leq K_{1}K_{2}K_{3} \\ &\ll (R^{2}/\log R)^{r}N^{m} \left(N\left(N^{-1}+R^{-1}\right)^{1/(d-1)}\right)^{t-m} \\ &\ll N^{t}R^{2r} \left(\left(N^{-1}+R^{-1}\right)^{1/(d-1)}\right)^{t/2}. \end{aligned}$$

Using the same discussion and choice of η as above, we conclude the proof.

Remark 3.1. Clearly in (3.6) we can replace t/2 with $\lfloor t/2 \rfloor$ but this does not change the optimal choice of η and thus the final bound.

3.2. **Proof of Theorem 1.3.** Let $(n_1, \ldots, n_s) \in [M+1, M+N]^s$ such that $u(n_1), \ldots, u(n_s)$ is m.d. of maximal rank, which implies that there exist integers $k_i \neq 0, i = 1, \ldots, s$, such that (1.1) holds. We can rewrite this relation as

(3.7)
$$\prod_{i\in\mathcal{I}}u(n_i)^{k_i}=\prod_{j\in\mathcal{J}}u(n_j)^{k_j},\quad k_i,k_j>0,$$

where $\mathcal{I} \cup \mathcal{J} = \{1, \ldots, s\}, \mathcal{I} \neq \emptyset, \mathcal{J} \neq \emptyset, \mathcal{I} \cap \mathcal{J} = \emptyset$. Let $I = \sharp \mathcal{I}$ and $J = \sharp \mathcal{J}$, and thus, I + J = s.

Fix one of $2^s - 2$ possible choices of the sets \mathcal{I} and \mathcal{J} as above. Fix n_i , $i \in \mathcal{I}$, trivially in $O(N^I)$ ways. Then, the square-free part $rad(u(n_i))$ of $u(n_i)$ is fixed for each $i \in \mathcal{I}$.

We may also assume that $n_i \ge C_2$, $i \in \mathcal{I}$, with C_2 as in Lemma 2.1, since this condition is violated only for $O(N^{s-1})$ choices of (n_1, \ldots, n_s) , which is admissible. By Lemma 2.1, for $n_i \in [M + 1, M + N]$, one has

(3.8)
$$\operatorname{rad}(u(n_i)) > n_i^{c(\log_2 n_i)/\log_3 n_i} \gg M^{c(\log_2 M)/\log_3(M+N)}.$$

For $i \in \mathcal{I}$, from (3.7) we see

$$\operatorname{rad}(u(n_i)) \mid \prod_{j \in \mathcal{J}} u(n_j).$$

This implies that there is a factorisation $\operatorname{rad}(u(n_i)) = d_1 \cdots d_J$ such that for each positive integer d_ℓ there exists $j \in \mathcal{J}$ such that $d_\ell \mid u(n_j)$. Let $\ell, 1 \leq \ell \leq J$, be such that $d_\ell \geq \operatorname{rad}(u(n_i))^{1/J}$, and

(3.9)
$$u(n_i) \equiv 0 \pmod{d_\ell}.$$

From (3.8) we have

(3.10)
$$d_{\ell} > M^{c_0(\log_2 M)/\log_3(M+N)}$$

with $c_0 = c/J \ge c/s$.

Using now Lemma 2.3, the inequality (3.10) and the fact that $J \leq s$, the number of $n_j \in [M + 1, M + N]$ satisfying the congruence (3.9) is

$$O\left(N/\log d_{\ell}+1\right) = O\left(N\frac{\log_3(M+N)}{\log M \log_2 M}+1\right)$$

Therefore, using the trivial bound N^{J-1} for the number of the remaining choices of n_j with $j \in \mathcal{J}$, we obtain that the total number of $n_j \in [M+1, M+N], j \in \mathcal{J}$, is

$$O\left(N^{J}\frac{\log_{3}(M+N)}{\log M \log_{2} M} + N^{J-1}\right).$$

Thus we obtain that

$$\mathsf{M}_{s}^{*}(M, N) \ll N^{s} \frac{\log_{3}(M+N)}{\log M \log_{2} M} + N^{s-1}.$$

Choosing $M \ge \exp(N \log_3 N / \log_2 N)$, we conclude the proof.

3.3. **Proof of Theorem 1.5.** Clearly for s = 2 we have to count integers $M + 1 \le m, n \le M + N$, with

(3.11)
$$u(m)^a = u(n)^b$$

for some positive integers a and b, where without loss of generality we can assume that gcd(a, b) = 1. We also notice that since the relation (3.11) is of maximal rank, neither $u(m) = \pm 1$ nor $u(n) = \pm 1$ holds.

Since **u** has a dominant root, |u(n)| grows monotonically with n, provided that n is large enough. Hence there are N + O(1) solutions $(m, n) \in [M + 1, M + N]^2$ with a = b = 1.

Now we count pairs (m, n) for which (3.11) holds with some $(a, b) \neq (1, 1)$.

We observe that if a > 1 then u(n) is the *a*-th power and by Lemma 2.6 there are O(1) such values of *n*. For b > 1 the argument also applies to *m*. Hence the total contribution from such solutions, over all a, b > 1, is O(1).

If a > 1 and b = 1 then again we see that there are O(1) such values of n. From this we easily derive that a = O(1), and hence we obtain O(1) possible values for m. So the contribution of such solutions to (3.11) is also O(1) only.

The case of a = 1 and b > 1 is completely analogous, which concludes the proof.

4. Possible applications of our approach

Our approach works for many other integer sequences $(a(n))_{n=1}^{\infty}$, provided the following information is available:

- (i) there are good bounds on the number of solutions to congruences $a(n) \equiv 0 \pmod{q}$, $1 \leq n \leq N$, in a broad range of positive integers q (or even just prime q = p) and N;
- (ii) there are good bounds (or known finiteness) on the number of perfect powers among $a(n), 1 \leq n \leq N$.

For example, using results of [16], coupled with the finiteness result of Lemma 2.6, one can estimate the number of multiplicatively dependent *s*-tuples from values of linear recurrence sequences at polynomial values of the argument $(u(F(n)))_{n=1}^{\infty}$, where $F \in \mathbb{Z}[X]$.

Acknowledgement

During this work, A.B. and L.H. were supported, in part, by the NKFIH grants 130909 and 150284 and A.O. and I.S. by the Australian Research Council Grant DP230100530. A.O. gratefully acknowledges the hospitality and support of the University of Debrecen, where this work was initiated, and the Max Planck Institute for Mathematics and Institut des Hautes Études Scientifiques, where this work has been carried out.

References

- F. Amoroso and E. Viada, 'On the zeros of linear recurrence sequences', Acta Arith., 147 (2011), 387–396. 4
- [2] W. D. Banks, J. B. Friedlander, S. V. Konyagin and I. E. Shparlinski, 'Incomplete exponential sums and Diffie-Hellman triples', *Math. Proc. Camb. Phil.* Soc., 140 (2006), 193–206. 5
- [3] H. Batte, M. Ddamulira, J. Kasozi and F. Luca, 'Multiplicative independence in the sequence of k-generalized Lucas numbers', *Indag. Math.*, in press, DOI: 10.1016/j.indag.2024.09.002. 2
- [4] M. Behera and P. K. Ray, 'Multiplicative dependence between Padovan and Perrin numbers', Bol. Soc. Mat. Mexicana, 29 (2023), 52. 2
- [5] M. Behera and P. K. Ray, 'On multiplicative dependent vectors with coordinates from certain higher order recurrences', Int. J. Number Theory, <u>20</u> (2024), 2491–2507. 2
- [6] A. Bérczes and V. Ziegler, 'On geometric progressions on Pell equations and Lucas sequences', Glas. Mat., 48 (2013), 1–22. 2
- [7] Y. Bugeaud and H. Kaneko, 'On perfect powers in linear recurrence sequences of integers', Kyushu J. Math., 73 (2019), 221–227. 6
- [8] J.-H. Evertse, 'On sums of S-units and linear recurrences', Compositio Math., 53 (1984), 225–244. 5

- C. A. Gómez, J. C. Gómez and F. Luca, 'Multiplicative dependence between k-Fibonacci and k-Lucas numbers', Period. Math. Hungar., 81 (2020), 217–233.
- [10] C. A. Gómez Ruiz and F. Luca, 'Multiplicative independence in k-generalized Fibonacci sequences', Lith. Math. J., 56 (2016), 503–517. 2
- [11] C. A. Gómez Ruiz and F. Luca, 'Multiplicative Diophantine equations with factors from different Lucas sequences', J. Number Theory, 170 (2017), 282– 301. 2
- [12] C. Hooley, 'Artin's conjecture for primitive roots', J. Reine Angew. Math., 225 (1967), 209–220. 5
- [13] F. Luca and V. Ziegler, 'Multiplicative relations on binary recurrences', Acta Arith., 161 (2013), 183–199. 2
- [14] A. J. van der Poorten and H. P. Schlickewei, 'Additive relations in fields', J. Austral Math. Soc., 51 (1991), 154–170. 5
- [15] I. E. Shparlinski, 'Some arithmetic properties of recurrence sequences', Math. Notes, 47 (1990), 612–617 (translated from Matem. Zametki). 5
- [16] I. E. Shparlinski and U. Zannier, 'Arithmetic properties of quadratic exponential polynomials', New York J. Math., 25 (2019) 207–218. 11
- [17] C. L. Stewart, 'On the greatest square-free factor of terms of a linear recurrence sequence', *Diophantine Equations*, Tata Inst. Fund. Res. Stud. Math. 20, Tata Inst. Fund. Res., Mumbai, 2008, 257–264. 4

INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN, P. O. BOX 400, H-4002 DEBRECEN, HUNGARY

Email address: berczesa@science.unideb.hu

Count István Tisza Foundation, Institute of Mathematics, University of Debrecen, P. O. Box 400, H-4002 Debrecen, Hungary, and HUN-REN-DE Equations, Functions, Curves and their Applications Research Group

Email address: hajdul@science.unideb.hu

School of Mathematics and Statistics, University of New South Wales, Sydney NSW 2052, Australia

Email address: alina.ostafe@unsw.edu.au

School of Mathematics and Statistics, University of New South Wales, Sydney NSW 2052, Australia

Email address: igor.shparlinski@unsw.edu.au