

The nonmodal kinetic theory of the macroscale convective flows of magnetized plasma, generated by the inhomogeneous microturbulence

V. V. Mikhailenko,^{1, a)} V. S. Mikhailenko,^{1, b)} and Hae June Lee^{2, c)}

¹⁾*Plasma Research Center, Pusan National University, Busan 46241, South Korea*

²⁾*Department of Electrical Engineering, Pusan National University, Busan 46241, South Korea*

(Dated: 30 January 2025)

In this paper, we develop the nonmodal kinetic theory of the macroscale convective flows of magnetized plasma, which stem from the average motion of ions and electrons in the electric field of the spatially inhomogeneous microturbulence. This theory bases on the two-scales approach to the solution of the Vlasov-Poisson system of equations for magnetized plasma, in which the solutions depend simultaneously on micro and macro scales. The developed theory predicts the generation of the sheared poloidal convective flow and of the radial compressed flow with radial flow velocity gradient. It was found that the macroscale (radial) inhomogeneity of the spectral intensity of the microturbulence is the condition necessary for the development of the two-dimensional non-diffusive convective plasma flows. The developed theory includes the theory of the evolution of the microscale turbulence in the sheared-compressed convective flows, formed by the microturbulence, and the theory of the slow macroscale evolution of a bulk of plasma by the compressed-sheared convective flows.

PACS numbers: 52.35.Ra, 52.35.Kt

^{a)}E-mail: vladimir@pusan.ac.kr

^{b)}E-mail: vsmikhailenko@pusan.ac.kr

^{c)}E-mail: haejune@pusan.ac.kr

I. INTRODUCTION

A common feature of the contemporary tokamaks is their operation in the regimes of the enhanced confinement, in which the suppression of the drift instabilities and turbulence by the poloidal sheared flow and the formation of the transport barriers with steep density profile (commonly referred to as the pedestal) in the edge region inside the last closed flux surface (LCFS) occur. The formation of the pedestal introduces in the edge region the radial spatial scale intermediate between macroscale commensurable with the plasma core density and temperature inhomogeneity lengths, and the microscale commensurable with wavelength of the microscale turbulence in the pedestal and the ion Larmor radius. This intermediate spatial scale is referred to as the mesoscale. It was observed essentially in all ion cyclotron (IC) and fast wave (FW) plasma heating experiments in tokamaks with applied MWs of IC or FW power¹⁻⁴ the anomalously large lost of the RF power, applied for the electron heating and current drive in the plasma core. The experiments reported lost of 50% or more of the launched RF power in the edge region and presumably in the cool, low density scrape-off layer (SOL) plasma adjacent to the FW antenna. This general experimental result displays a strong obstacle in sustaining high performance discharges in a tokamak configuration not predicted by the linear theory of the propagation and absorption of the IC and FW wave in the edge of tokamak plasma, which claimed that the injection of FW may be efficient for RF heating and current drive in the steady-state noninductive tokamak operation. It was assumed⁵⁻⁸ that the nonlinear excitation of the parametric microscale IC quasimode decay instability⁵ following IC or FW injection, which was observed in SOL^{9,10} and predicted theoretically^{5,6} and the turbulent heating of ions by the parametric turbulence^{10,11}, powered by the parametric instabilities, are the basic nonlinear processes responsible for the anomalous absorption of the applied RF power in SOL plasma and of the anomalous heating of ions in SOL. The excitation of the parametric microturbulence and the formation of the stationary microscale spectrum, resulted from the nonlinear saturation of these instabilities, and the processes of the turbulent heating of the plasma, resulted from the interactions of ions and electrons with microscale turbulence, are the local processes, which occur on the spatial scales across the magnetic field much less than the spatial inhomogeneity of a tokamak edge plasma and of the applied FW field. The analysis of the turbulent heating of ions by the parametric turbulence, performed in Refs.^{12,13} on the base

of the quasilinear theory for the ion and electron distribution functions, revealed that the absorption of RF energy by ions and electrons by the developed microscale IC turbulence in the cold low density SOL plasma is a weak effect, which cannot be responsible for the observed large loss of the RF power in SOL.

It was found in the FW heating experiments^{3,4} on the National Spherical Torus eXperiment (NSTX) that a large part of the FW power loss flows as the RF-produced heat flux in the SOL mostly along open field lines and deposited to the divertor. The formation and propagation of such a flow is a nonlocal process which is missed completely in the quasilinear theory of the anomalous diffusive transport. The kinetic theory of the non-diffusive convective flows, driven by the electrostatic IC parametric turbulence, was developed in Refs.¹⁴⁻¹⁶. This theory revealed that the large scale spatial (radial) inhomogeneity of the spectral intensity of the microturbulence is the condition necessary for the development of the two-dimensional non-diffusive convective flows. It was derived in Ref.¹⁴, that the velocities of the radial and poloidal convective flows are proportional to the gradient of the spectral intensity of the microscale turbulence and the radial velocity of the convective flow is in the direction opposite to this gradient. In the tokamak plasmas, this convective flow moves outward of pedestal to the SOL region. It was concluded^{14,15} that the dynamical processes in the pedestal and near SOL are heavily determined by the convective flows in this region by the inhomogeneous microturbulence and by interplay of the convective flows with the sheared poloidal edge flow of a tokamak plasma, which is the inherent component of the pedestal development and sustainment.

The nonlocal approach to the kinetic theory of the convective flows, generated by the spatially inhomogeneous microturbulence is presented in Sec. II. This theory involves two disparate spatial scales: microscale, commensurable with the wavelength of the microscale instabilities and of the ion gyroradius, and a much larger scale commensurable with the spatial inhomogeneity of the plasma density and temperature and of the inhomogeneity of FW wave. For FW plasma heating of the edge tokamak plasma this large scale is considered as the mesoscale. A brief outline of the kinetic theory of the mesoscale non-diffusive convective flows, which stem from the averaged motion of ions and electrons in the electric field of the inhomogeneous edge microturbulence, is presented in Sec. III. It was found¹⁴⁻¹⁶ that the mesoscale convective flow may be responsible for the observed transport^{3,4} of the plasma thermal energy from the strongly inhomogeneous high-density hot pedestal plasma

to the lower density cold near-SOL plasma.

The nonlocal two-scales approach to the theory of the macroscale convective flows, which develop in the bulk of plasma by the radially inhomogeneous microturbulence, is presented in Sec. IV. This two-scales theory is the extension of the theory of the mesoscale convection flows, developed in Refs.^{14–16}, on the plasma macroscales comparable with scales of the bulk of plasma. In that Section, we derive the basic equations, which determine the formation of the macroscale spatially inhomogeneous compressed-sheared convective flows by the inhomogeneous microturbulence, the back-action of the inhomogeneous convected flows on the microturbulence, and the slow macroscale evolution of a bulk of plasma in the compressed-sheared convective flows. The detailed analysis of the temporal evolution of the microscale turbulence in the inhomogeneous convected flows is presented in Sec.V. In that Section, the integral equation, which governs the nonmodal temporal evolution of the microscale potential in the macroscale convective flows is derived. The theory of the macroscale evolution of the ion component of a plasma in the convected flows is presented in Sec. VI. This section contains 1) the nonmodal quasilinear theory, which governs the temporal evolution of the ion distribution function resulted from the interactions of ions with ensemble of sheared-compressed microscale waves with random phases, 2) the self-consistent theory of the temporal evolution of the macroscale electrostatic potential of the plasma respond on the macroscale convective flow. This equation, is the basic equation of the stability theory of the convective flows against the development of the secondary macroscale instabilities of a plasma with inhomogeneous convective flows. Conclusions are presented in Sec. VII.

II. THE NONLOCAL APPROACH TO THE THEORY OF THE CONVECTIVE FLOWS, GENERATED BY THE SPATIALLY INHOMOGENEOUS MICROTURBULENCE

The theory of the convective flows in a weakly inhomogeneous plasma, driven by the spatially inhomogeneous electrostatic microturbulence, developed in Refs.^{14–16}, bases on the two-scales approach to the solution of the Vlasov-Poisson system of equations, in which the solutions depend simultaneously on widely different scales. It accounts for the evolution on a slow and on a fast time-scales, and on a small and on a large spatial scales.

This theory was developed in Refs.^{14–16} in a slab geometry approximation, in which the

coordinates x, y, z for the microscale fast variations are viewed as corresponding to the radial, poloidal and toroidal directions, respectively, of the toroidal coordinate system. The large scale coordinates X, Y and the long time T of the slow variations of the plasma parameters across the magnetic field are used here to distinguish them from the short scale (microscale) variables x, y and fast time t by the relations $X = \varepsilon x$, $Y = \varepsilon y$, $T = \varepsilon t$, where $\varepsilon \ll 1$ is a small parameter. Within a slab geometry approximation, the Vlasov equation for the velocity distribution function F_α of α plasma species ($\alpha = i$ for ions and $\alpha = e$ for electrons), which accounts for the evolution of the plasma turbulence on the small and large scales, in FW electric field $\mathbf{E}_0(X, t)$ has a form

$$\begin{aligned} & \frac{\partial F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)}{\partial t} + \varepsilon \frac{\partial F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)}{\partial T} \\ & + \mathbf{v} \cdot \left(\frac{\partial F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)}{\partial \mathbf{r}} + \varepsilon \frac{\partial F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)}{\partial X} \right) \\ & + \frac{e_\alpha}{m_\alpha} \left(\mathbf{E}_0(t, X) + \mathbf{E}(\mathbf{r}, t, X) + \frac{1}{c} [\mathbf{v} \times \mathbf{B}_0] \right) \frac{\partial F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)}{\partial \mathbf{v}} = 0, \end{aligned} \quad (1)$$

where $\mathbf{E}(\mathbf{r}, t, X)$ is the field of the electrostatic response of a plasma on the propagating FW, both inhomogeneous on the large radial spatial scale X ; \mathbf{B}_0 is the confined magnetic field, directed along z axes. $\mathbf{E}(\mathbf{r}, t, X)$ field is determined by the Poisson equation

$$\nabla_{\mathbf{r}} \cdot \mathbf{E}(\mathbf{r}, t, X) = 4\pi \sum_{\alpha=i,e} e_\alpha \int f_\alpha(\mathbf{v}, \mathbf{r}, t, X) d\mathbf{v}, \quad (2)$$

in which f_α is the fluctuating part of the distribution function F_α , $f_\alpha = F_\alpha - F_{0\alpha}$, where $F_{0\alpha}$ is the equilibrium distribution function. It was found in Ref.¹⁴ that the spatially inhomogeneous FW field $\mathbf{E}_0(t, X)$ may be excluded from the Vlasov equation (1) for $F_\alpha(\mathbf{v}, \mathbf{r}, t, X, T)$ by the transformation of the velocity \mathbf{v} and position \mathbf{r} and X variables to new velocity \mathbf{v}_α and position \mathbf{r}_α , X_α , determined in the reference flow, which moves relative to the laboratory frame with velocity $\mathbf{V}_\alpha(t, X)$ of α species particle in $\mathbf{E}_0(t, X)$ and confined \mathbf{B}_0 fields. In such a reference flow for ions, FW electric field is presented in the Vlasov equation (1) for $F_i(\mathbf{v}_i, \mathbf{r}_i, t, X_i)$ for the microscale perturbations only in terms on the order of $|R_{ix}/L_E| \ll 1$ (where R_{ix} is the ion displacement in the FW field and L_E is a spatial scale of the FW inhomogeneity), which may be neglected. Without these terms, the Vlasov equation for $F_i(\mathbf{v}_i, \mathbf{r}_i, t, X_i)$ with great accuracy has a form as for a steady plasma in the uniform magnetic field \mathbf{B}_0 without FW field, and contains the variable X_i as a parameter, i. e.

$$\frac{\partial F_i(\mathbf{v}_i, \mathbf{r}_i, t, X_i)}{\partial t} + \mathbf{v}_i \cdot \frac{\partial F_i}{\partial \mathbf{r}_i} + \frac{e_i}{m_i c} [\mathbf{v}_i \times \mathbf{B}_0] \frac{\partial F_i}{\partial \mathbf{v}_i}$$

$$+\frac{e_i}{m_i}\mathbf{E}_i(\mathbf{r}_i, t, X_i)\frac{\partial F_i(\mathbf{v}_i, \mathbf{r}_i, t, X_i)}{\partial \mathbf{v}_i}=0. \quad (3)$$

In this equation, $\mathbf{E}_i(\mathbf{r}_i, t, X_i)$ is the electric field of the electrostatic two dimensional microturbulence, which is the microscale response of a plasma on a large scale plasma ion-homogeneities. The saturation of the microscale instability with the frequency $\omega(\mathbf{k})$ and the growth rate $\gamma(\mathbf{k})$, which occurs at time $t \gtrsim \gamma^{-1}(\mathbf{k}) \gtrsim 2\pi|\omega^{-1}(\mathbf{k})|$, is followed by the formation at the fast time t the steady level of the spatially inhomogeneous along X_i microturbulence. At that stage, the electric field \mathbf{E}_i of the electrostatic two dimensional microturbulence, directed almost across the magnetic field \mathbf{B}_0 , may be presented in the ion reference flow in the form, that includes the microscale \mathbf{r}_i and the large scale X_i variables, i. e.

$$\mathbf{E}_i(\mathbf{r}_i, X_i, t) = \frac{1}{2(2\pi)^2} \int d\mathbf{k} [\mathbf{E}_i(\mathbf{k}, X_i) e^{i\psi(\mathbf{r}_i, t)} + \mathbf{E}_i^*(\mathbf{k}, X_i) e^{-i\psi(\mathbf{r}_i, t)}], \quad (4)$$

with phase

$$\psi(\mathbf{r}_i, t) = -\omega(\mathbf{k})t + \mathbf{k}\mathbf{r}_i + \theta(\mathbf{k}), \quad (5)$$

changed on the microscale \mathbf{r}_i and fast time t , i. e. as a linear superposition of the electric fields of perturbations which has a form of the plane waves with frequencies $\omega(\mathbf{k})$, the wave vectors \mathbf{k} directed almost across the magnetic field, and with position dependent on X_i amplitudes $\mathbf{E}_i(\mathbf{k}, X_i)$. For the treatment of the slow evolution of a plasma on the large scales, the Vlasov equation (3) should be considered in variables X_i, Y_i, T ,

$$\begin{aligned} \frac{\partial F_i(\mathbf{v}_i, X_i, Y_i, T)}{\partial T} + v_{ix} \frac{\partial F_i}{\partial X_i} + v_{iy} \frac{\partial F_i}{\partial Y_i} + \frac{\omega_{ci}}{\varepsilon} v_{iy} \frac{\partial F_i}{\partial v_{ix}} - \frac{\omega_{ci}}{\varepsilon} v_{ix} \frac{\partial F_i}{\partial v_{iy}} \\ + \frac{e_i}{\varepsilon m_i} \mathbf{E}_i(X_i, \varepsilon^{-1}X_i, \varepsilon^{-1}Y_i, \varepsilon^{-1}T) \frac{\partial F_i}{\partial \mathbf{v}_i} = 0. \end{aligned} \quad (6)$$

In a plasma with turbulent electric field $\mathbf{E}_i(\mathbf{r}_i, X_i, t)$, the velocity \mathbf{v}_i of ions is the total velocity of the ion thermal motion and of the ion motion in the turbulent electric field (4) and in the magnetic field \mathbf{B}_0 . It was found in Refs.^{14,15}, that some two-dimensional spatially inhomogeneous microturbulence-associated reference flow, which moves with velocity $\tilde{\mathbf{U}}_i(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ relative to the laboratory frame, may be determined, in which the ion velocity \mathbf{v}_i at time T is equal to

$$\mathbf{v}_i = \tilde{\mathbf{v}}_i + \tilde{\mathbf{U}}_i(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon), \quad (7)$$

where $\tilde{\mathbf{v}}_i$ is the thermal velocity of an ion in the reference flow, and \tilde{X}_i and \tilde{Y}_i are the coordinates of the ion position, determined in the reference flow. These co-ordinates are coupled with X_i, Y_i co-ordinates by the relations

$$X_i = \tilde{X}_i + \int_{T_0}^T \tilde{U}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T_1 \right) dT_1 = \tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \quad (8)$$

$$Y_i = \tilde{Y}_i + \int_{T_0}^T \tilde{U}_{iy} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T_1 \right) dT_1 = \tilde{Y}_i + \tilde{R}_{iy} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \quad (9)$$

The inverse transform $(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T) \rightarrow (\mathbf{v}_i, X_i, Y_i, T)$ is determined by relations

$$\tilde{\mathbf{v}}_i = \mathbf{v}_i - \tilde{\mathbf{V}}_i(X_i, Y_i, T, \varepsilon), \quad (10)$$

$$\tilde{X}_i = X_i - \int_{T_0}^T \tilde{V}_{ix} \left(X_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T_1 \right) dT_1 = X_i - R_{ix} \left(X_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right), \quad (11)$$

$$\tilde{Y}_i = Y_i - \int_{T_0}^T \tilde{V}_{iy} \left(X_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T_1 \right) dT_1 = Y_i - R_{iy} \left(X_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right). \quad (12)$$

In this reference flow, the electrostatic potential $\varphi(\mathbf{r}_i, t, X_i)$, which determines the electrostatic field of the microscale turbulence in the laboratory frame, $\mathbf{E}_i(\mathbf{r}_i, t, X_i) = -\nabla_{\mathbf{r}_i} \varphi(\mathbf{r}_i, t, X_i)$, is presented in the form

$$\varphi(\mathbf{r}_i, X_i, Y_i, t) = \tilde{\varphi}_i \left(\tilde{\mathbf{r}}_i, \tilde{X}_i + \tilde{R}_{ix}, t \right) + \Phi_i \left(\tilde{X}_i, \tilde{Y}_i, T \right), \quad (13)$$

where $\tilde{\varphi}_i$ is the electrostatic potential of the microscale turbulence,

$$\tilde{\mathbf{E}}_i \left(\tilde{\mathbf{r}}_i, \tilde{X}_i + \tilde{R}_{ix}, t \right) = -\nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_i \left(\tilde{\mathbf{r}}_i, \tilde{X}_i + \tilde{R}_{ix}, t \right); \quad (14)$$

$\Phi_i \left(\tilde{X}_i, \tilde{Y}_i, T \right)$ is the potential of the large scale plasma response on the formation and slow evolution of the large scale plasma inhomogeneities, observed in the reference flow.

$$\bar{\mathbf{E}}_i \left(\tilde{X}_i, \tilde{Y}_i, T \right) = -\nabla \Phi_i \left(\tilde{X}_i, \tilde{Y}_i, T \right). \quad (15)$$

The Vlasov equation (6) for the ion distribution function $F_i \left(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T \right)$ has a form

$$\frac{\partial F_i \left(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T \right)}{\partial T} + \tilde{v}_{ix} \frac{\partial F_i}{\partial \tilde{X}_i} + \tilde{v}_{iy} \frac{\partial F_i}{\partial \tilde{Y}_i}$$

$$\begin{aligned}
& - \left(\tilde{v}_{ix} + \tilde{U}_{ix} \left(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon \right) \right) \int_{T_0}^T \frac{\partial \tilde{V}_{ix}}{\partial X_i} dT_1 \frac{\partial F_i}{\partial \tilde{X}_i} - \left(\tilde{v}_{iy} + \tilde{U}_{iy} \left(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon \right) \right) \int_{T_0}^T \frac{\partial \tilde{V}_{iy}}{\partial X_i} dT_1 \frac{\partial F_i}{\partial \tilde{Y}_i} \\
& + \left(\frac{\omega_{ci}}{\varepsilon} \tilde{v}_{iy} - \tilde{v}_{ix} \frac{\partial \tilde{V}_{ix}}{\partial X_i} \right) \frac{\partial F_i}{\partial \tilde{v}_{ix}} - \left(\frac{\omega_{ci}}{\varepsilon} + \frac{\partial \tilde{V}_{iy}}{\partial X_i} \right) \tilde{v}_{ix} \frac{\partial F_i}{\partial \tilde{v}_{iy}} - \frac{e_i}{m_i} \tilde{\nabla} \Phi \left(\tilde{X}_i, \tilde{Y}_i, T \right) \\
& \quad - \left[\frac{\partial \tilde{U}_{ix} \left(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon \right)}{\partial T} - \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{iy} \right. \\
& \quad \left. - \frac{e_i}{\varepsilon m_i} \tilde{E}_{ix} \left(\tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right) \right] \frac{\partial F_i}{\partial \tilde{v}_{ix}} \\
& \quad - \left[\frac{\partial \tilde{U}_{iy} \left(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon \right)}{\partial T} + \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{ix} \right. \\
& \quad \left. - \frac{e_i}{\varepsilon m_i} \tilde{E}_{iy} \left(\tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right) \right] \frac{\partial F_i}{\partial \tilde{v}_{iy}} = 0. \quad (16)
\end{aligned}$$

In Eq. (16), the identities $\tilde{\mathbf{V}}_i(\mathbf{r}_i, X_i, t) = \tilde{\mathbf{U}}_i(\tilde{\mathbf{r}}_i, \tilde{X}_i, t)$ and

$$\frac{\partial \tilde{\mathbf{V}}_i(X_i, Y_i, T)}{\partial T} + \tilde{V}_{ix}(X_i, Y_i, T) \frac{\partial \tilde{\mathbf{V}}_i(X_i, Y_i, T)}{\partial X_i} = \frac{\partial \tilde{\mathbf{U}}_i(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)}{\partial T}, \quad (17)$$

which follow from Eqs. (7)-(12), were used.

The Vlasov equation (16) for $\bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, t)$, and the similar equation for $\bar{F}_e(\tilde{\mathbf{v}}_e, \tilde{X}_e, \tilde{Y}_e, t)$, and the Poisson equation for the potential $\Phi_i(\tilde{X}_i, \tilde{Y}_i, t)$,

$$\begin{aligned}
\frac{\partial^2 \Phi_i(\tilde{X}_i, \tilde{Y}_i, t)}{\partial^2 \tilde{X}_i} + \frac{\partial^2 \Phi_i(\tilde{X}_i, \tilde{Y}_i, t)}{\partial^2 \tilde{Y}_i} &= -4\pi \left(e_i \int d\mathbf{v}_i \bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, t) \right. \\
&\quad \left. - |e| \int d\mathbf{v}_e \bar{F}_e(\tilde{\mathbf{v}}_e, \tilde{X}_e, \tilde{Y}_e, t) \right), \quad (18)
\end{aligned}$$

compose the Vlasov-Poisson system, which governs the kinetic large scale evolution of a plasma under the action of the spatially inhomogeneous microturbulence.

III. THE MESOSCALE NON-DIFFUSIVE CONVECTIVE FLOWS

The first task in the solution of Eq. (16) is determining the velocity $\tilde{\mathbf{U}}_i(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ of the reference flow. The proper selection of $\tilde{\mathbf{U}}_i(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ provides the opportunity for the development of the kinetic theories of the mesoscale as well as the macroscale non-diffusive

convective flows. Equation (16) displays that selection of the velocities $\tilde{U}_{ix}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ and $\tilde{U}_{iy}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ as a solution of the equations

$$\begin{aligned} & \frac{\partial \tilde{U}_{ix}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)}{\partial T} - \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{iy} \\ &= \frac{e_i}{\varepsilon m_i} \tilde{E}_{ix} \left(\tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right), \end{aligned} \quad (19)$$

$$\begin{aligned} & \frac{\partial \tilde{U}_{iy}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)}{\partial T} + \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{ix} \\ &= \frac{e_i}{\varepsilon m_i} \tilde{E}_{iy} \left(\tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right), \end{aligned} \quad (20)$$

gives the ion Vlasov equation which contains the microscale electric field $\tilde{\mathbf{E}}$ only in terms on the order of the ratio of the ion displacement R_{ix} in that electric field to the mesoscale inhomogeneity length L_E of the $\tilde{\mathbf{E}}_i$ field. The solution to Eqs. (19), (20) for velocities $\tilde{U}_{ix}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ and $\tilde{U}_{iy}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon)$ were derived in Ref.¹⁵ using the approximation

$$\begin{aligned} & \tilde{\mathbf{E}}_i \left(\tilde{X}_i + \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right), \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right) \\ &= \tilde{\mathbf{E}}_{i0} \left(\tilde{X}_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right) + \tilde{\mathbf{E}}_{i1} \left(\tilde{X}_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right), \end{aligned} \quad (21)$$

where

$$\tilde{\mathbf{E}}_{i1} \left(\tilde{X}_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} T \right) = \frac{\partial \tilde{\mathbf{E}}_{i0}}{\partial \tilde{X}_i} \tilde{R}_{ix} \left(\tilde{X}_i, \varepsilon^{-1} \tilde{X}_i, \varepsilon^{-1} \tilde{Y}_i, \varepsilon^{-1} T \right). \quad (22)$$

which is valid for the small displacement, $|\tilde{\mathbf{R}}_i| \ll L_E$, of an ion in the inhomogeneous electric field $\tilde{\mathbf{E}}_i$. At time $T \gtrsim \gamma^{-1}$, electric field $\tilde{\mathbf{E}}_{i0}$ becomes the random function of the initial phase $\theta(\mathbf{k})$ with zero mean value, i. e. $\langle \tilde{\mathbf{E}}_{i0} \rangle = 0$. By averaging of the Vlasov equation (18) over the ensemble of the initial phases of the microscale perturbations, the equation, which determines the long time evolution on the mesoscales of the ensemble averaged ion distribution function $\bar{F}_i = \bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T)$,

$$\begin{aligned} & \frac{\partial \bar{F}_i}{\partial T} + \tilde{v}_{ix} \frac{\partial \bar{F}_i}{\partial \tilde{X}_i} + \tilde{v}_{iy} \frac{\partial \bar{F}_i}{\partial \tilde{Y}_i} - \bar{U}_{ix}(\tilde{X}_i) \frac{\partial \bar{F}_i}{\partial \tilde{X}_i} - \bar{U}_{iy}(\tilde{X}_i) \frac{\partial \bar{F}_i}{\partial \tilde{Y}_i} \\ &+ v_{iz} \frac{\partial \bar{F}_i}{\partial Z_i} + \frac{1}{\varepsilon} \omega_{ci} \tilde{v}_{iy} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{ix}} - \frac{1}{\varepsilon} \omega_{ci} \tilde{v}_{ix} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{iy}} - \frac{e_i}{m_i} \frac{\partial \Phi_i(\tilde{X}_i, \tilde{Y}_i, Z_i, T)}{\partial \tilde{X}_i} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{ix}} \\ &- \frac{e_i}{m_i} \frac{\partial \Phi_i(\tilde{X}_i, \tilde{Y}_i, Z_i, T)}{\partial \tilde{Y}_i} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{iy}} - \frac{e_i}{m_i} \frac{\partial \Phi_i(\tilde{X}_i, \tilde{Y}_i, Z_i, T)}{\partial Z_i} \frac{\partial \bar{F}_i}{\partial v_{iz}} = 0, \end{aligned} \quad (23)$$

was derived in Ref.¹⁵. In this equation, the velocities $\bar{U}_{ix}(\tilde{X}_i)$ and $\bar{U}_{iy}(\tilde{X}_i)$,

$$\bar{U}_{ix}(\tilde{X}_i) = \left\langle \tilde{U}_{ix}^{(0)}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon) \int_0^T \frac{\partial}{\partial X_i} \tilde{U}_{ix}^{(0)}(X_i, Y_i, T_1, \varepsilon) dT_1 \right\rangle, \quad (24)$$

$$\bar{U}_{iy}(\tilde{X}_i) = \left\langle \tilde{U}_{iy}^{(0)}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon) \int_0^T \frac{\partial}{\partial X_i} \tilde{U}_{iy}^{(0)}(X_i, Y_i, T_1, \varepsilon) dT_1 \right\rangle, \quad (25)$$

are the velocities of the ion convective flows relative to the laboratory frame, as the average velocities of the ion motion in the electric field of the microturbulence inhomogeneous on the mesoscale.

The system of equations which contains Eq. (23) for $\bar{F}_i = \bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T)$, the similar equation for the electron distribution function $\bar{F}_e = \bar{F}_e(\tilde{\mathbf{v}}_e, \tilde{X}_e, \tilde{Y}_e, T)$, and the Poisson equation for the potential $\Phi_i(\tilde{X}_i, \tilde{Y}_i, Z_i, T)$

$$\begin{aligned} & \frac{\partial^2 \Phi_i}{\partial^2 \tilde{X}_i} + \frac{\partial^2 \Phi_i}{\partial^2 \tilde{Y}_i} \\ & = -4\pi \left(e_i \int d\tilde{\mathbf{v}}_i \bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T) - |e| \int d\tilde{\mathbf{v}}_e \bar{F}_e(\tilde{\mathbf{v}}_e, \tilde{X}_e, \tilde{Y}_e, T) \right), \end{aligned} \quad (26)$$

was investigated in details Refs.^{15,16}, where the simplest expansion for the flow velocities

$$\bar{U}_{ix}(\tilde{X}_i) = \bar{U}_{ix}^{(0)}(\tilde{X}_i^{(0)}) + \bar{U}'_{ix}(\tilde{X}_i^{(0)}) (\tilde{X}_i - \tilde{X}_i^{(0)}), \quad (27)$$

$$\bar{U}_{iy}(\tilde{X}_i) = \bar{U}_{iy}^{(0)}(\tilde{X}_i^{(0)}) + \bar{U}'_{iy}(\tilde{X}_i^{(0)}) (\tilde{X}_i - \tilde{X}_i^{(0)}). \quad (28)$$

was used with the uniform velocity compressing rate, $\bar{U}'_{ix}(\tilde{X}_i^{(0)}) = \text{const}$, and the uniform velocity shearing rate, $\bar{U}'_{iy}(\tilde{X}_i^{(0)}) = \text{const}$.

IV. THE THEORY OF THE MACROSCALE CONVECTIVE FLOWS

In Eqs. (24), (25), the electric field $\tilde{\mathbf{E}}_i$ of the microscale turbulence is considered as not changed with time by the convective flow, formed by the inhomogeneous microscale turbulence itself. However, in the flow with spatially inhomogeneous flow velocity, any perturbation, which before the development of the flow has a plane wave structure, experiences the continuous distortion in the flow and becomes the sheared-compressed mode with time dependent structure^{15,16}. This effect may be negligible small for the mesoscale convective

flow, but it is important for the macroscale convective flows, in which the evolution of the microturbulence in the convective flows occurs on the long macroscales during a long time. In this section, we develop the theory of the generation of the macroscale convective flows by the microturbulence, which accounts also the macroscale nonlinear back-reaction effects of the convection flows on the temporal evolution of the microturbulence in the spatially inhomogeneous convective flow.

We consider Eq. (16) for the ion distribution function $F_i = F_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T, \varepsilon^{-1}X_i, \varepsilon^{-1}Y_i, \varepsilon^{-1}T)$ with accounting for the expansion (21),

$$\begin{aligned}
& \frac{\partial F_i}{\partial T} + \tilde{v}_{ix} \frac{\partial F_i}{\partial \tilde{X}_i} + \tilde{v}_{iy} \frac{\partial F_i}{\partial \tilde{Y}_i} \\
& - \left(\tilde{v}_{ix} + \tilde{U}_{ix}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon) \right) \int_{T_0}^T \frac{\partial \tilde{V}_{ix}}{\partial \tilde{X}_i} dT_1 \frac{\partial F_i}{\partial \tilde{X}_i} - \left(\tilde{v}_{iy} + \tilde{U}_{iy}(\tilde{X}_i, \tilde{Y}_i, T, \varepsilon) \right) \int_{T_0}^T \frac{\partial \tilde{V}_{iy}}{\partial \tilde{X}_i} dT_1 \frac{\partial F_i}{\partial \tilde{Y}_i} \\
& + \left(\frac{\omega_{ci}}{\varepsilon} \tilde{v}_{iy} - \tilde{v}_{ix} \frac{\partial \tilde{V}_{ix}}{\partial \tilde{X}_i} \right) \frac{\partial F_i}{\partial \tilde{v}_{ix}} - \left(\frac{\omega_{ci}}{\varepsilon} + \frac{\partial \tilde{V}_{iy}}{\partial \tilde{X}_i} \right) \tilde{v}_{ix} \frac{\partial F_i}{\partial \tilde{v}_{iy}} \\
& + \frac{e_i}{m_i} \left(\frac{1}{\varepsilon} \tilde{\mathbf{E}}_{i0}(\tilde{X}_i, \varepsilon^{-1}\tilde{X}_i, \varepsilon^{-1}\tilde{Y}_i, \varepsilon^{-1}T) - \tilde{\nabla}\Phi(\tilde{X}_i, \tilde{Y}_i, T) \right) \frac{\partial F_i}{\partial \tilde{\mathbf{v}}_i} \\
& - \left[\frac{\partial \tilde{U}_{ix}(\tilde{X}_i, T, \varepsilon)}{\partial T} - \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{iy} - \frac{e_i}{\varepsilon m_i} \left(\tilde{E}_{i1x}(\tilde{X}_i, \varepsilon^{-1}X_i, \varepsilon^{-1}Y_i, \varepsilon^{-1}T) \right) \right] \frac{\partial F_i}{\partial \tilde{v}_{ix}} \\
& - \left[\frac{\partial \tilde{U}_{iy}(\tilde{X}_i, T, \varepsilon)}{\partial T} + \frac{\omega_{ci}}{\varepsilon} \tilde{U}_{ix} - \frac{e_i}{\varepsilon m_i} \left(\tilde{E}_{i1y}(\tilde{X}_i, \varepsilon^{-1}X_i, \varepsilon^{-1}Y_i, \varepsilon^{-1}T) \right) \right] \frac{\partial F_i}{\partial \tilde{v}_{iy}} = 0. \quad (29)
\end{aligned}$$

In Eq. (29), we will present the ion distribution function F_i in the form

$$F_i = \bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T) + f_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \varepsilon^{-1}X_i, \varepsilon^{-1}Y_i, \varepsilon^{-1}Z_i, \varepsilon^{-1}T), \quad (30)$$

where $\bar{F}_i = \langle F_i \rangle$ is the averaged F_i over the ensemble of the initial phases, and f_i is the microscale perturbation of F_i with $\langle f_i \rangle = 0$. In the averaged Eq. (29), $\langle \tilde{E}_{i0x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle = \langle \tilde{E}_{i0y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle = 0$, and the velocities $\bar{U}_{ix}(\tilde{X}_i, t) = \langle \tilde{U}_{ix}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle$ and $\bar{U}_{iy}(\tilde{X}_i, t) = \langle \tilde{U}_{iy}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle$ are determined by the equations

$$\frac{\partial \bar{U}_{ix}}{\partial t} - \omega_{ci} \bar{U}_{iy} = \frac{e_i}{m_i} \langle \tilde{E}_{i1x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle, \quad (31)$$

$$\frac{\partial \bar{U}_{iy}}{\partial t} + \omega_{ci} \bar{U}_{ix} = \frac{e_i}{m_i} \langle \tilde{E}_{i1y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \rangle. \quad (32)$$

With averaged over the fast time $t \gg \omega_{ci}^{-1}$ solutions to Eqs. (31) and (32) for the velocities of the reference flow, $\langle \bar{U}_{ix}(\tilde{X}_i, t) \rangle_t = \bar{U}_{ix}(\tilde{X}_i)$ and $\langle \bar{U}_{iy}(\tilde{X}_i, t) \rangle_t = \bar{U}_{iy}(\tilde{X}_i)$, derived in Appendix A, and with accounting for that for a tokamak plasma $\omega_{ci} \gg \epsilon |\frac{\partial \bar{U}_{iy}}{\partial \tilde{X}_i}|, \epsilon |\frac{\partial \bar{U}_{ix}}{\partial \tilde{X}_i}|$, Eq. (29) obtains a simple form:

$$\begin{aligned} \frac{\partial F_i}{\partial T} + \tilde{v}_{ix} \left(1 - \int_{T_0}^T \frac{\partial \bar{U}_{ix}(\tilde{X}_i)}{\partial \tilde{X}_i} dT_1 \right) \frac{\partial F_i}{\partial \tilde{X}_i} + \left(\tilde{v}_{iy} - \tilde{v}_{ix} \int_{T_0}^T \frac{\partial \bar{U}_{iy}(\tilde{X}_i)}{\partial \tilde{X}_i} dT_1 \right) \frac{\partial F_i}{\partial \tilde{Y}_i} \\ + \tilde{v}_{iy} \frac{\omega_{ci}}{\epsilon} \frac{\partial F_i}{\partial \tilde{v}_{ix}} - \tilde{v}_{ix} \frac{\omega_{ci}}{\epsilon} \frac{\partial F_i}{\partial \tilde{v}_{iy}} \\ + \left(\frac{e_i}{\epsilon m_i} \tilde{\mathbf{E}}_{i0}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) - \frac{e_i}{m_i} \nabla \Phi(\tilde{X}_i, \tilde{Y}_i, Z, T) \right) \frac{\partial F_i}{\partial \tilde{\mathbf{v}}_i} = 0, \end{aligned} \quad (33)$$

in which the terms on the order of $O\left(|\tilde{E}_{i0}|^4\right)$ are neglected.

For deriving the simplest solution to Eq. (33) for $\bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T)$ and for $f_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \epsilon^{-1}X_i, \epsilon^{-1}Y_i, \epsilon^{-1}Z_i, \epsilon^{-1}T)$ we use the expansions for the velocities

$$\bar{U}_{ix}(\tilde{X}_i) = \bar{U}_{ix}^{(0)}(\tilde{X}_i^{(0)}) + \bar{U}'_{ix}(\tilde{X}_i^{(0)}) (\tilde{X}_i - \tilde{X}_i^{(0)}), \quad (34)$$

$$\bar{U}_{iy}(\tilde{X}_i) = \bar{U}_{iy}^{(0)}(\tilde{X}_i^{(0)}) + \bar{U}'_{iy}(\tilde{X}_i^{(0)}) (\tilde{X}_i - \tilde{X}_i^{(0)}). \quad (35)$$

In what follows, we consider the case of the uniform velocity compressing rate, $\bar{U}'_{ix}(\tilde{X}_i^{(0)}) = \text{const}$, and of the uniform velocity shearing rate, $\bar{U}'_{iy}(\tilde{X}_i^{(0)}) = \text{const}$, and put $\tilde{X}_i^{(0)} = 0$. With expansions (34) and (35), the equation for $\bar{F}_i(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T)$, which determines the slow macroscale evolution of F_i , is derived by averaging of Eq. (33) over the ensemble of the initial phases,

$$\begin{aligned} \frac{\partial \bar{F}_i}{\partial T} + \tilde{v}_{ix} (1 - \bar{U}'_{ix} T) \frac{\partial \bar{F}_i}{\partial \tilde{X}_i} + (\tilde{v}_{iy} - \tilde{v}_{ix} \bar{U}'_{iy} T) \frac{\partial \bar{F}_i}{\partial \tilde{Y}_i} \\ + \tilde{v}_{iy} \frac{\omega_{ci}}{\epsilon} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{ix}} - \tilde{v}_{ix} \frac{\omega_{ci}}{\epsilon} \frac{\partial \bar{F}_i}{\partial \tilde{v}_{iy}} \\ - \frac{e_i}{m_i} \nabla \Phi(\tilde{X}_i, \tilde{Y}_i, Z, T) \frac{\partial \bar{F}_i}{\partial \tilde{\mathbf{v}}_i} = - \frac{e_i}{\epsilon m_i} \left\langle \tilde{\mathbf{E}}_{i0}(\tilde{\mathbf{r}}_i, t, \tilde{X}_i, T) \frac{\partial f_i}{\partial \tilde{\mathbf{v}}_i} \right\rangle. \end{aligned} \quad (36)$$

This equation accounts for the well known quasilinear effect of the microscale turbulence on the resonant ions, which is responsible for the local processes of the anomalous diffusion and anomalous heating of the resonant ions. Also, it accounts for the nonlinear back-reaction effect on the bulk of ions the spatially inhomogeneous sheared-compressed flows, which are the average effect of the motion of the nonresonant ions in the electric field of

the microturbulence inhomogeneous on the macroscale. This effect consists in the formation of the macroscale response of the plasma on the spatially inhomogeneous convective flows. Solution of this equation is presented in Sec. VI.

The fast microscale evolution of F_i is determined by equation for

$$f_i \left(\tilde{\mathbf{v}}_i, \tilde{X}_i, \varepsilon^{-1} X_i, \varepsilon^{-1} Y_i, \varepsilon^{-1} Z_i, \varepsilon^{-1} T \right) = f_i \left(\tilde{\mathbf{v}}_i, \tilde{\mathbf{r}}_i, t, \tilde{X}_i \right),$$

$$\frac{\partial f_i}{\partial t} + \tilde{v}_{ix} (1 - \bar{U}'_{ix} t) \frac{\partial f_i}{\partial \tilde{x}_i} + (\tilde{v}_{iy} - \tilde{v}_{ix} \bar{U}'_{iy} t) \frac{\partial f_i}{\partial \tilde{y}_i} + \omega_{ci} \tilde{v}_{iy} \frac{\partial f_i}{\partial \tilde{v}_{ix}} - \omega_{ci} \tilde{v}_{ix} \frac{\partial f_i}{\partial \tilde{v}_{iy}}$$

$$- \frac{e_i}{m_i} \nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_{i0} \left(\tilde{\mathbf{r}}_i, \tilde{X}_i, t \right) \frac{\partial \bar{F}_i \left(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, T \right)}{\partial \tilde{\mathbf{v}}_i} = 0, \quad (37)$$

where $\tilde{\varphi}_{i0}$ is the electrostatic potential of the microscale turbulence,

$$\tilde{\mathbf{E}}_{i0} \left(\tilde{\mathbf{r}}_i, \tilde{X}_i, t \right) = -\nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_{i0} \left(\tilde{\mathbf{r}}_i, \tilde{X}_i, t \right). \quad (38)$$

and the prime in \bar{U}'_{ix} , \bar{U}'_{iy} in Eq. (37) denotes the derivatives of \bar{U}_{ix} , \bar{U}_{iy} over the microscale co-ordinate $\tilde{x}_i = \frac{\tilde{X}_i}{\varepsilon}$. In Eq. (37), the variables $\tilde{X}_i, \tilde{Y}_i, Z, T$ enter as the parameters. Equations (36) and (37) presents the two-scale expansion of the Vlasov equation in the frame of references co-moving with the ion convective flow, formed by the average motion of ions in the microscale turbulence inhomogeneous along the coordinate \tilde{X}_i . The system of Eqs. (36), (37) for \bar{F}_i and for f_i , and the similar decomposition on \bar{F}_e and for f_e for the electron Vlasov equation, and the Poisson equations for the macroscale potential Φ and for the potential of the microscale turbulence φ compose the two-scale Vlasov-Poisson system, which describe the back-reaction effects of the convection flows on the microturbulence (Eq. (37)) and the macroscale plasma respond on the development convective flows in plasma (Eq. (36)).

V. THE EVOLUTION OF THE MICROSCALE TURBULENCE IN THE MACROSCALE CONVECTIVE FLOWS

In the guiding center coordinates \hat{x}_i, \hat{y}_i , determined by the relations

$$\tilde{x}_i = \hat{x}_i - \frac{\hat{v}_{i\perp}}{\omega_{ci}} (1 - \bar{U}'_{ix} t) \sin(\phi_1 - \omega_{ci} t) + O\left(\frac{\bar{U}_{ix}}{\omega_{ci}}\right), \quad (39)$$

$$\tilde{y}_i = \hat{y}_i + \frac{\hat{v}_{i\perp}}{\omega_{ci}} \cos(\phi_1 - \omega_{ci} t) + \frac{\hat{v}_{i\perp}}{\omega_{ci}} \sin(\phi_1 - \omega_{ci} t) \bar{U}'_{iy} t + O\left(\frac{\bar{U}_{iy}}{\omega_{ci}}\right), \quad (40)$$

the Vlasov equation (37) for $f_i \left(\hat{v}_{i\perp}, \phi_1, v_z, \hat{x}_i, \hat{y}_i, z, \hat{X}_i, t \right)$ has a form

$$\frac{\partial f_i}{\partial t} = \frac{e_i}{m_i} \left[-\frac{\omega_{ci}}{\hat{v}_{i\perp}} \frac{\partial \tilde{\varphi}_{i0}}{\partial \phi_1} \frac{\partial \bar{F}_{i0}}{\partial \hat{v}_{i\perp}} + \frac{1}{\omega_{ci}} (1 - \bar{U}'_{ix} t) \frac{\partial \tilde{\varphi}_{i0}}{\partial \hat{y}_i} \frac{\partial \bar{F}_{i0}}{\partial \hat{X}_i} + \frac{\partial \tilde{\varphi}_{i0}}{\partial z_i} \frac{\partial \bar{F}_{i0}}{\partial v_{iz}} \right], \quad (41)$$

where the potential $\tilde{\varphi}_{i0}$ is equal to

$$\begin{aligned}\tilde{\varphi}_{i0}(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t) &= \frac{1}{(2\pi)^3} \int dk_{\tilde{x}_i} dk_{\tilde{y}_i} dk_z \tilde{\varphi}_{i0}(\tilde{\mathbf{k}}_i, \tilde{X}_i, t) e^{ik_{\tilde{x}_i}\tilde{x}_i + ik_{\tilde{y}_i}\tilde{y}_i + ik_z z} \\ &= \frac{1}{(2\pi)^3} \int dk_{\tilde{x}_i} dk_{\tilde{y}_i} dk_z \tilde{\varphi}_{i0}(\tilde{\mathbf{k}}_i, \hat{X}_i, t) \\ &\times \sum_{n=-\infty}^{\infty} J_n \left(\frac{\hat{k}_{i\perp}(t) \hat{v}_{i\perp}}{\omega_{ci}} \right) e^{ik_{\tilde{x}_i}\tilde{x}_i + ik_{\tilde{y}_i}\tilde{y}_i + ik_z z - in(\phi_1 - \omega_{ci}t - \chi_i(t))},\end{aligned}\quad (42)$$

where $\tilde{\mathbf{k}}_i = (k_{\tilde{x}_i}, k_{\tilde{y}_i}, k_z)$ and $\hat{k}_{i\perp}(t)$ and $\chi_i(t)$ are determined by the relations

$$\hat{k}_{i\perp}^2(t) = (k_{\tilde{x}_i} - (k_{\tilde{x}_i} \bar{U}'_{ix} + k_{\tilde{y}_i} \bar{U}'_{iy})t)^2 + k_{\tilde{y}_i}^2, \quad \sin \chi_i(t) = \frac{k_{\tilde{y}_i}}{k_{i\perp}(t)}.\quad (43)$$

The solution to Eq. (41) with nonmodal microscale potential (42),

$$\begin{aligned}f_i(\hat{v}_{i\perp}, \phi_1, v_z, \hat{x}_i, \hat{y}_i, z, \hat{X}_i, t) &= i \frac{e_i}{m_i} \frac{1}{(2\pi)^3} \int dk_{\tilde{x}_i} dk_{\tilde{y}_i} dk_z \\ &\sum_{n_1=-\infty}^{\infty} \int_{t_0}^t dt_1 \tilde{\varphi}_{i0}(\tilde{\mathbf{k}}_i, \hat{X}_i, t_1) J_{n_1} \left(\frac{\hat{k}_{i\perp}(t_1) \hat{v}_{i\perp}}{\omega_{ci}} \right) \\ &\times \left[\frac{n_1 \omega_{ci}}{\hat{v}_{i\perp}} \frac{\partial \bar{F}_{i0}}{\partial \hat{v}_{i\perp}} + \frac{k_{\tilde{y}_i}}{\omega_{ci}} (1 - \bar{U}'_{ix} t_1) \frac{\partial \bar{F}_{i0}}{\partial \hat{X}_i} + k_z \frac{\partial \bar{F}_{i0}}{\partial v_{iz}} \right] \\ &\times e^{ik_{\tilde{x}_i}\hat{x}_i + ik_{\tilde{y}_i}\hat{y}_i + ik_z z + ik_z v_{iz} t_1 - in_1(\phi_1 - \omega_{ci}t_1 - \chi_i(t_1))}.\end{aligned}\quad (44)$$

displays the effects of the sheared-compressed flow on the temporal evolution of the perturbation f_i of the ion distribution function. Equation (44) predicts that the nonmodal effects reveal in two effects. The first one is the time dependence of the argument $\hat{k}_{i\perp}(t_1) \hat{v}_{i\perp} / \omega_{ci}$ of the Bessel function J_n . The time dependence of the finite Larmor radius effect was found¹⁷ to be the basic linear mechanism of the interactions of ions rotating on the Larmor orbit with wave in the sheared flow across the magnetic field, where the wave experiences permanent distortion by the sheared flow. Equation (44) extends this nonmodal effect on the interaction of ions with wave in the sheared-compressed convective flow. The second is a new nonmodal time dependent effect of the accelerated compressed flow along X_i on the ion drift along coordinate Y_i .

For the low frequency electrostatic perturbations, for which $\frac{\partial \tilde{\varphi}_{i0}}{\partial t} \ll \omega_{ci} \tilde{\varphi}_{i0}$, only the terms with $n = n_1 = 0$ should be retained in summations over n and n_1 in Eqs. (42) and (44). The Fourier transformed over coordinates \hat{x}_i, \hat{y}_i low frequency density perturbation $n_i(\tilde{\mathbf{k}}_i, \tilde{X}_i, t) = \int d\hat{\mathbf{v}}_i f_i(\hat{v}_{i\perp}, v_z, \tilde{\mathbf{k}}_i, \tilde{X}_i, t)$, of ions with the Maxwellian distribution

$\bar{F}_{i0}(\mathbf{v}_i, \hat{X}_i, T)$ with inhomogeneous ions density and ion temperature,

$$\bar{F}_{i0}(\mathbf{v}_e, \hat{X}_i, T) = \frac{n_{i0}(\hat{X}_i)}{(2\pi v_{Ti}^2(\hat{X}_i))^{3/2}} \exp\left(-\frac{v_{i\perp}^2 + v_z^2}{2v_{Ti}^2(\hat{X}_i)}\right), \quad (45)$$

was found in the form

$$\begin{aligned} n_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t) &= -\frac{e_i}{T_i} n_{0i}(\hat{X}_i) \int_{t_0}^t dt_1 \frac{d}{dt_1} \psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1) \\ &+ \frac{e_i}{T_i} n_{0i}(\hat{X}_i) \int_{t_0}^t dt_1 \frac{d}{dt_1} \left[\psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1) I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \right. \\ &\quad \left. \times e^{-\frac{1}{2}\rho_i^2(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) - \frac{1}{2}k_z^2 v_{Ti}^2(t-t_1)^2} \right] \\ &+ \frac{e_i}{T_i} n_{0i}(\hat{X}_i) \int_{t_0}^t dt_1 \psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1) I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) e^{-\frac{1}{2}\rho_i^2(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) - \frac{1}{2}k_z^2 v_{Ti}^2(t-t_1)^2} \\ &\quad \times \left(ik_{\tilde{y}_i} v_{di} (1 - \eta_i) (1 - \bar{U}'_{ix} t) - k_z^2 v_{Ti}^2 (t - t_1) - \frac{i}{2} k_{\tilde{y}_i} v_{di} \eta_i k_z^2 v_{Ti}^2 (t - t_1)^2 \right) \\ &+ i \frac{e_i}{T_i} n_{0i}(\hat{X}_i) k_{\tilde{y}_i} v_{di} \eta_i \int_{t_0}^t dt_1 \psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1) e^{-\frac{1}{2}\rho_i^2(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) - \frac{1}{2}k_z^2 v_{Ti}^2(t-t_1)^2} \\ &\quad \times \left[\left(1 - \frac{1}{2}\rho_i^2(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \right) I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \right. \\ &\quad \left. + \rho_i^2 k_{i\perp}(t) k_{i\perp}(t_1) I_1(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \right]. \end{aligned} \quad (46)$$

In Equation (46), $\psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t) = \varphi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t) \Theta(t - t_0)$, where $\Theta(t - t_0)$ is the unit-step Heaviside function (it is equal to zero for $t < t_0$ and is equal to unity for $t \geq t_0$).

The perturbation $f_e(\tilde{\mathbf{v}}_e, \mathbf{r}_e, \tilde{X}_e, t)$ of the electron distribution function $\bar{F}_{e0}(\mathbf{v}_e, \hat{X}_e, T)$, is determined by the equation

$$\frac{\partial f_e}{\partial t} = \frac{e}{m_e} \left[\frac{1}{\omega_{ce}} \frac{\partial \varphi_{e0}}{\partial \tilde{y}_e} \frac{\partial \bar{F}_{e0}}{\partial \tilde{X}_e} + \frac{\partial \varphi_{e0}}{\partial z_e} \frac{\partial \bar{F}_{e0}}{\partial v_{ez}} \right], \quad (47)$$

where $\varphi_{e0} = \varphi_{e0}(\mathbf{r}_e, \tilde{X}_e, t)$. The solution $f_e(\hat{v}_{e\perp}, v_z, k_{\tilde{x}_e}, k_{\tilde{y}_e}, k_z, \tilde{X}_e, t)$ of Eq. (43), Fourier transformed over \tilde{x}_e, \tilde{y}_e ,

$$\begin{aligned} f_e(\hat{v}_{e\perp}, v_z, k_{\tilde{x}_e}, k_{\tilde{y}_e}, k_z, \tilde{X}_e, t) &= i \frac{e}{m_e} \int dt_1 \varphi_{e0}(\mathbf{k}_e, \tilde{X}_e, t_1) \\ &\quad \times \left[\frac{k_{\tilde{y}_e}}{\omega_{ce}} \frac{\partial \bar{F}_{e0}}{\partial \tilde{X}_e} + k_z \frac{\partial \bar{F}_{e0}}{\partial v_{ez}} \right] e^{-ik_z v_{ez}(t-t_1)}, \end{aligned} \quad (48)$$

determines the temporal evolution of the separate spatial Fourier harmonic of the perturbation $f_e(\hat{v}_{e\perp}, v_z, k_{\tilde{x}_e}, k_{\tilde{y}_e}, k_z, \tilde{X}_e, t)$ in the laboratory frame.

The separate harmonic of the long wavelength, $k_{e\perp}\rho_e \ll 1$, electron density perturbation $n_e(\hat{\mathbf{k}}_e, \tilde{X}_e, t_1) = \int d\hat{\mathbf{v}}_e f_e(\hat{\mathbf{v}}_e, \hat{\mathbf{k}}_e, \tilde{X}_e, t)$ for the Maxwellian distribution of electrons, with inhomogeneous density and with uniform temperature,

$$\bar{F}_{e0}(\mathbf{v}_e, \hat{X}_e, T) = \frac{n_{e0}(\hat{X}_e)}{(2\pi v_{Te}^2)^{3/2}} \exp\left(-\frac{v_{i\perp}^2 + v_z^2}{2v_{Te}^2}\right), \quad (49)$$

is given approximately by the relation

$$\begin{aligned} n_e(\mathbf{k}_e, \hat{X}_e, t) &= i\frac{e}{T_e} n_{0e}(\hat{X}_e) \int_{t_0}^t dt_1 \left[-\frac{d\psi_e(\mathbf{k}_e, \hat{X}_e, t_1)}{dt_1} \right. \\ &\left. + \left(\frac{d\psi_e(\mathbf{k}_e, \hat{X}_e, t_1)}{dt_1} + ik_{\tilde{y}_e} v_{de} \psi_e(\mathbf{k}_e, \hat{X}_e, t_1) \right) e^{-\frac{1}{2}k_z^2 v_{Te}^2 (t-t_1)^2} \right]. \end{aligned} \quad (50)$$

In this equation, $\psi_e(\hat{\mathbf{k}}_e, \tilde{X}_e, t) = \varphi_e(\hat{\mathbf{k}}_e, \tilde{X}_e, t) \Theta(t - t_0)$.

The Poisson equation for the potential of the microscale plasma turbulence we derive here for the potential $\varphi_{e0}(\tilde{x}_e, \tilde{y}_e, z, t)$, determined in variables \tilde{x}_e, \tilde{y}_e of the laboratory frame,

$$\begin{aligned} \frac{\partial^2 \psi_e(\tilde{x}_e, \tilde{y}_e, z, \tilde{X}_e, t)}{\partial^2 \tilde{x}_e} + \frac{\partial^2 \psi_e(\tilde{x}_e, \tilde{y}_e, z, \tilde{X}_e, t)}{\partial^2 \tilde{y}_e} + \frac{\partial^2 \psi_e(\tilde{x}_e, \tilde{y}_e, \tilde{X}_e, z, t)}{\partial^2 \tilde{z}_e} \\ = -4\pi \left[e_i n_i(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t) - |e| n_e(\tilde{x}_e, \tilde{y}_e, z, \tilde{X}_e, t) \right]. \end{aligned} \quad (51)$$

The Fourier transform of Eq. (51) over \tilde{x}_e and \tilde{y}_e ,

$$(k_{\tilde{x}_e}^2 + k_{\tilde{y}_e}^2) \psi_e(k_{\tilde{x}_e}, k_{\tilde{y}_e}, \tilde{X}_e, t) = 4\pi e_i n_i^{(e)}(\mathbf{k}_e, \tilde{X}_e, t) + 4\pi e n_e(\mathbf{k}_e, \tilde{X}_e, t). \quad (52)$$

contains the Fourier transform $n_i^{(e)}(\mathbf{k}_e, \tilde{X}_e, t)$, of the perturbation of the ion density $n_i(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t)$ performed over \tilde{x}_e and \tilde{y}_e , i. e

$$\begin{aligned} n_i^{(e)}(\mathbf{k}_e, \tilde{X}_e, t) &= \int d\tilde{x}_e \int d\tilde{y}_e n_i(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t) e^{-ik_{\tilde{x}_e} \tilde{x}_e - ik_{\tilde{y}_e} \tilde{y}_e} \\ &= \int d\tilde{x}_e \int d\tilde{y}_e n_i(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t) e^{-ik_{\tilde{x}_e} \tilde{x}_i - ik_{\tilde{y}_e} \tilde{y}_i - ik_{\tilde{x}_e} (\tilde{x}_e - \tilde{x}_i) - ik_{\tilde{y}_e} (\tilde{y}_e - \tilde{y}_i)} \\ &= \int d\tilde{x}_i \int d\tilde{y}_i n_i(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t) \left| \frac{\partial(\tilde{x}_e, \tilde{y}_e)}{\partial(\tilde{x}_i, \tilde{y}_i)} \right| e^{-ik_{\tilde{x}_e} \tilde{x}_i - ik_{\tilde{y}_e} \tilde{y}_i - ik_{\tilde{x}_e} (\tilde{x}_e - \tilde{x}_i) - ik_{\tilde{y}_e} (\tilde{y}_e - \tilde{y}_i)}. \end{aligned} \quad (53)$$

It follows from Eqs. (8) and (12) that

$$\tilde{x}_e = \tilde{x}_i (1 + \bar{U}'_{ix} t) + \bar{U}_{ix}^{(0)}(0) t, \quad (54)$$

$$\tilde{x}_i = \frac{\tilde{x}_e - \bar{U}_{ix}^{(0)} t}{1 + \bar{U}'_{ix} t}, \quad (55)$$

and

$$\tilde{y}_e = \tilde{y}_i + \left(\bar{U}_{iy}^{(0)} + \bar{U}'_{iy} \tilde{x}_i \right) t, \quad (56)$$

$$\tilde{y}_i = \tilde{y}_e - \frac{\bar{U}'_{iy} t}{1 + \bar{U}'_{ix} t} \tilde{x}_e - \bar{U}_{iy}^{(0)} t + \frac{\bar{U}'_{iy} \bar{U}_{iy}^{(0)} t^2}{1 + \bar{U}'_{ix} t}. \quad (57)$$

With Eqs. (54) and (56), Eq. (53) becomes

$$\begin{aligned} n_i^{(e)} \left(\mathbf{k}_e, \tilde{X}_e, t \right) &= \int d\tilde{x}_i \int d\tilde{y}_i n_i \left(\tilde{x}_i, \tilde{y}_i, z, \tilde{X}_i, t \right) |1 + \bar{U}'_{ix} t| \\ &\times \exp \left[-ik_{\tilde{x}_e} \left(\tilde{x}_i + \left(\bar{U}_{ix}^{(0)} + \bar{U}'_{ix} \tilde{x}_i \right) t \right) - ik_{\tilde{y}_e} \left(\tilde{y}_i + \left(\bar{U}_{iy}^{(0)} + \bar{U}'_{iy} \tilde{x}_i \right) t \right) \right] \\ &= e^{-ik_{\tilde{x}_e} \bar{U}_{ix}^{(0)} t - ik_{\tilde{y}_e} \bar{U}_{iy}^{(0)} t} |1 + \bar{U}'_{ix} t| n_i \left(k_{\tilde{x}_e} (1 + \bar{U}'_{ix} t) + k_{\tilde{y}_e} \bar{U}'_{iy} t, k_{\tilde{y}_e}, t \right). \end{aligned} \quad (58)$$

Equation (46) for $n_i \left(\hat{\mathbf{k}}_i, \tilde{X}_i, t \right)$ contains the Fourier transform $\psi_i \left(\hat{\mathbf{k}}_i, \tilde{X}_i, t \right)$ of the potential $\psi_i \left(\tilde{x}_i, \tilde{y}_i, \tilde{X}_i, t_1 \right)$. The connection relation of $\psi_i \left(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1 \right)$ with $\psi_e \left(\hat{\mathbf{k}}_e, \tilde{X}_e, t_1 \right)$ follows from the relation

$$\begin{aligned} \psi_i \left(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1 \right) &= \int d\tilde{x}_i \int d\tilde{y}_i \psi_i \left(\tilde{x}_i, \tilde{y}_i, \tilde{X}_i, t_1 \right) e^{-ik_{\tilde{x}_i} \tilde{x}_i - ik_{\tilde{y}_i} \tilde{y}_i} \\ &= \int d\tilde{x}_e \int d\tilde{y}_e \psi_e \left(\tilde{x}_e, \tilde{y}_e, \tilde{X}_e, t_1 \right) \left| \frac{\partial \left(\tilde{x}_i(t_1), \tilde{y}_i(t_1) \right)}{\partial \left(\tilde{x}_e, \tilde{y}_e \right)} \right| \\ &\quad \times e^{-ik_{\tilde{x}_i} \tilde{x}_e - ik_{\tilde{y}_i} \tilde{y}_e - ik_{\tilde{x}_i} (\tilde{x}_i - \tilde{x}_e) - ik_{\tilde{y}_i} (\tilde{y}_i - \tilde{y}_e)}. \\ &= \frac{1}{4\pi^2} \frac{1}{|1 + \bar{U}'_{ix} t_1|} \int dk_{\tilde{x}_e} \int dk_{\tilde{y}_e} \psi_e \left(k_{\tilde{x}_e}, k_{\tilde{y}_e}, \tilde{X}_i, t_1 \right) \\ &\times \int d\tilde{x}_e \int d\tilde{y}_e \exp \left[i \left(k_{\tilde{x}_e} - k_{\tilde{x}_i} \right) \tilde{x}_e + i \left(k_{\tilde{y}_e} - k_{\tilde{y}_i} \right) \tilde{y}_e - ik_{\tilde{x}_i} \left(\tilde{x}_i - \tilde{x}_e \right) - ik_{\tilde{y}_i} \left(\tilde{y}_i - \tilde{y}_e \right) \right]. \end{aligned} \quad (59)$$

The integrating of Eq. (59) over \tilde{x}_e, \tilde{y}_e , in which the relations

$$\tilde{x}_i(t_1) - \tilde{x}_e = -\frac{\bar{U}_{ix}^{(0)} t_1}{1 + \bar{U}'_{ix} t_1} - \frac{\bar{U}'_{ix} t_1}{1 + \bar{U}'_{ix} t_1} \tilde{x}_e = b_{0x}(t_1) + b_{1x}(t_1) \tilde{x}_e, \quad (60)$$

$$\tilde{y}_i(t_1) - \tilde{y}_e = -\bar{U}_{iy}^{(0)} t_1 + \frac{\bar{U}'_{iy} \bar{U}_{ix}^{(0)} t_1^2}{1 + \bar{U}'_{ix} t_1} - \frac{\bar{U}'_{iy} t_1}{1 + \bar{U}'_{ix} t_1} \tilde{x}_e = b_{0y}(t_1) + b_{1y}(t_1) \tilde{x}_e, \quad (61)$$

are employed, gives the relation

$$\begin{aligned} \psi_i \left(k_{\tilde{x}_i}, k_{\tilde{y}_i}, \tilde{X}_i, t_1 \right) &= \frac{1}{|1 + \bar{U}'_{ix} t_1|} e^{-ik_{\tilde{x}_i} b_{0x}(t_1) - ik_{\tilde{y}_i} b_{0y}(t_1)} \\ &\times \psi_e \left(k_{\tilde{x}_i} (1 + b_{1x}(t_1)) + k_{\tilde{y}_i} b_{1x}(t_1), k_{\tilde{y}_i}, \tilde{X}_i, t_1 \right). \end{aligned} \quad (62)$$

It follows from Eq. (58) that the wave numbers of $n_i^{(e)}$, which are conjugate with co-ordinates \tilde{x}_i, \tilde{y}_i , are $k_{\tilde{x}_e} + (k_{\tilde{x}_e} \bar{U}'_{ix} + k_{\tilde{y}_e} \bar{U}'_{iy}) t$ and $k_{\tilde{y}_e}$. The applying this result to Eq. (59) gives the presentation of the Fourier transform $\psi_i(k_{\tilde{x}_i}, k_{\tilde{y}_i}, \tilde{X}_i, t_1)$ in variables $k_{\tilde{x}_e}, k_{\tilde{y}_e}$,

$$\begin{aligned} \psi_i(k_{\tilde{x}_i}, k_{\tilde{y}_i}, \tilde{X}_i, t_1) &= \frac{1}{|1 + \bar{U}'_{ix} t_1|} \exp \left[ik_{\tilde{x}_e} \bar{U}_{ix}^{(0)} t_1 \left(1 + \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) \right. \\ &\quad \left. + ik_{\tilde{y}_e} \bar{U}_{iy}^{(0)} t_1 \left(1 + \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) \right] \\ &\times \psi_e \left(k_{\tilde{x}_e} \left(1 + \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) + k_{\tilde{y}_e} \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1}, k_{\tilde{y}_e}, \tilde{X}_i, t_1 \right) \\ &= \psi_i^{(e)}(k_{\tilde{x}_e}, k_{\tilde{y}_e}, \tilde{X}_i, t, t_1). \end{aligned} \quad (63)$$

Equation (63) displays, that the separate spatial Fourier harmonic $\psi_i(k_{\tilde{x}_i}, k_{\tilde{y}_i}, \tilde{X}_i, t_1)$ of the electrostatic potential, determined in the frame of references, which moves with velocities (31), (32) is perceived in the electron (laboratory) frame as the Doppler-shifted continuously sheared and compressed mode with time-dependent wave vectors.

For the deriving the ion density perturbation $n_i^{(e)}$, determined by Eq. (55), the potential $\psi_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t_1)$ in Eq. (46), which determines $n_i(\hat{\mathbf{k}}_i, \tilde{X}_i, t)$, should be changed on $\psi_i^{(e)}$ given by Eq. (63).

$$\begin{aligned} n_i^{(e)}(\mathbf{k}_e, \tilde{X}_e, t) &= -\frac{e_i}{T_i} n_{0i}(\hat{X}_i) \int_{t_0}^t dt_1 \frac{d}{dt_1} \left\{ \left| \frac{1 + \bar{U}'_{ix} t}{1 + \bar{U}'_{ix} t_1} \right| \right. \\ &\times \exp \left[-ik_{\tilde{x}_e} \bar{U}_{ix}^{(0)} t_1 \left(1 - \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) - ik_{\tilde{y}_e} \bar{U}_{iy}^{(0)} t_1 \left(1 - \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) \right] \\ &\times \psi_e \left(k_{\tilde{x}_e} \left(1 + \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) + k_{\tilde{y}_e} \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1}, k_{\tilde{y}_e}, \tilde{X}_i, t_1 \right) \\ &\times \left[1 - I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) e^{-\frac{1}{2}(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \rho_i^2 - \frac{1}{2} k_z^2 v_{Ti}^2 (t-t_1)^2} \right] \left. \right\} \\ &\quad + \frac{e_i}{T_i} n_{0i}(\hat{X}_i) \int_{t_0}^t dt_1 \frac{d}{dt_1} \left| \frac{1 + \bar{U}'_{ix} t}{1 + \bar{U}'_{ix} t_1} \right| \\ &\times \exp \left[-ik_{\tilde{x}_e} \bar{U}_{ix}^{(0)} t_1 \left(1 - \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) - ik_{\tilde{y}_e} \bar{U}_{iy}^{(0)} t_1 \left(1 - \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) \right] \\ &\quad \times e^{-\frac{1}{2}(k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \rho_i^2 - \frac{1}{2} k_z^2 v_{Ti}^2 (t-t_1)^2} \\ &\times \psi_e \left(k_{\tilde{x}_e} \left(1 + \frac{\bar{U}'_{ix}(t-t_1)}{1 + \bar{U}'_{ix} t_1} \right) + k_{\tilde{y}_e} \frac{\bar{U}'_{iy}(t-t_1)}{1 + \bar{U}'_{ix} t_1}, k_{\tilde{y}_e}, \tilde{X}_i, t_1 \right) \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \left(ik_{\tilde{y}_e} v_{di} (1 - \eta_i) (1 - \bar{U}'_{ix} t_1) - k_z^2 v_{Te}^2 (t - t_1) - \frac{ik_{\tilde{y}_e} v_{di} \eta_i}{2} k_z^2 v_{Ti}^2 (t - t_1)^2 \right) \right. \\
& \times I_0 (k_{i\perp} (t) k_{i\perp} (t_1) \rho_i^2) + ik_{\tilde{y}_e} v_{di} \eta_i \left[\left(1 - (k_{i\perp}^2 (t) + k_{i\perp}^2 (t_1)) \frac{\rho_i^2}{2} \right) I_0 (k_{i\perp} (t) k_{i\perp} (t_1) \rho_i^2) \right. \\
& \left. \left. + k_{i\perp} (t) k_{i\perp} (t_1) \rho_i^2 I_1 (k_{i\perp} (t) k_{i\perp} (t_1) \rho_i^2) \right] \right\} \quad (64)
\end{aligned}$$

where

$$\begin{aligned}
k_{i\perp}^2 (t) &= (k_{\tilde{x}_e} - (k_{\tilde{x}_e} \bar{U}'_{ix} + k_{\tilde{y}_e} \bar{U}'_{iy}) \bar{U}'_{iy} t^2)^2 + k_{\tilde{y}_e}^2, \\
k_{i\perp}^2 (t_1) &= (k_{\tilde{x}_e} - (k_{\tilde{x}_e} \bar{U}'_{ix} + k_{\tilde{y}_e} \bar{U}'_{iy}) \bar{U}'_{iy} t_1^2)^2 + k_{\tilde{y}_e}^2. \quad (65)
\end{aligned}$$

It follows from Eqs. (60) and (61), that for the spatially uniform flow, for which $\bar{U}'_{ix} = \bar{U}'_{iy} = 0$, the spatial Fourier harmonics of the electrostatic potential $\psi_i (\mathbf{k}_i, \tilde{X}_i, t)$ and of the perturbation of the ion density $n_i (\mathbf{k}_i, \tilde{X}_i, t)$ are perceived in the electron frame as the Doppler-shifted modes

$$\psi_i (\mathbf{k}_i, t_1) = \exp \left(ik_x \bar{U}_{ix}^{(0)} t_1 + ik_y \bar{U}_{iy}^{(0)} t_1 \right) \psi_i^{(e)} (\mathbf{k}_e, t_1), \quad (66)$$

$$n_i (\mathbf{k}_e, t) = \exp \left(-ik_x \bar{U}_{ix}^{(0)} t - ik_y \bar{U}_{iy}^{(0)} t \right) n_i^{(e)} (\mathbf{k}_i, t), \quad (67)$$

and the Poisson equation for the Fourier transform $\psi_e (\mathbf{k}_e, \omega)$ gives the well known dispersion equation

$$1 + \varepsilon_e (\mathbf{k}_e, \omega) + \varepsilon_i \left(k_e, \omega - k_{ex} \bar{U}_{ix}^{(0)} - k_{ey} \bar{U}_{iy}^{(0)} \right) = 0, \quad (68)$$

where ε_e and ε_i are the linear electron and ion dielectric permittivities, as for the instabilities driven by the spatially uniform current with current velocities $\bar{\mathbf{U}}_{ix}^{(0)}$ and $\bar{\mathbf{U}}_{iy}^{(0)}$. It follows from Eq. (64), that the modal time dependence $\sim e^{-i\omega(\mathbf{k}_i)t}$ of the potential ψ_e exists only at the initial stage of the potential evolution when the sheared-compressed effects of the convected flow are negligible small, i. e. when $\bar{U}'_{ix} (t - t_0) \ll 1$, $\bar{U}'_{iy} (t - t_0) \ll 1$. At a longer time, the time dependence of the potential ψ_e becomes very different from a canonical modal form. The nonmodal evolution of the potential ψ_e can be investigated employing the Poisson equation Eq. (49) for the potential ψ_e with perturbation of the electron density determined by (47) and with perturbation of the ion density $n_i^{(e)} (\mathbf{k}_e, \tilde{X}_e, t)$ determined by (61). This equation for a short time, for which $\bar{U}'_{ix} (t - t_0) < 1$ and $\bar{U}'_{iy} (t - t_0) < 1$, corresponding the initial stage of the nonmodal evolution of the perturbation of ions $n_i^{(e)} (\mathbf{k}_e, X_i, t)$ and of the

potential ψ_e , has a form

$$\begin{aligned}
k_e^2 \lambda_{De}^2 \psi_e (\mathbf{k}_e, \tilde{X}_e, t) &= \int_{t_0}^t dt_1 \left\{ -\tau \frac{d\psi_e}{dt_1} + \tau e^{-\frac{1}{2} k_z^2 v_{Te}^2 (t-t_1)^2} \right. \\
&\times \left[\frac{d\psi_e}{dt_1} + \tau \left(ik_{ye} v_{de} - k_z^2 v_{Te}^2 (t-t_1) - \frac{ik_{ye} v_{de} \eta_e}{2} k_z^2 v_{Te}^2 (t-t_1)^2 \right) \psi_e (\mathbf{k}_e, \tilde{X}_e, t_1) \right] \Big\} \\
&+ \int_{t_0}^t dt_1 \left\{ -|1 + \bar{U}'_{ix}(t-t_1)| \exp \left[-i \left(k_{\tilde{X}_e} \bar{U}_{ix}^{(0)} + k_{\tilde{y}_e} \bar{U}_{iy}^{(0)} \right) (t-t_1) \right] \right. \\
&\times \left[1 - I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \exp \left(-\frac{1}{2} (k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \rho_i^2 - \frac{1}{2} k_z^2 v_{Ti}^2 (t-t_1)^2 \right) \right] \\
&\times \psi_e \left(k_{\tilde{X}_e} (1 + \bar{U}'_{ix}(t-t_1)) + k_{\tilde{y}_e} \bar{U}'_{iy}(t-t_1), k_{\tilde{y}_e}, \tilde{X}_e, t \right) \Big\} \\
&+ \int_{t_0}^t dt_1 |1 + \bar{U}'_{ix}(t-t_1)| e^{-ik_{\tilde{X}_e} \bar{U}_{ix}^{(0)}(t-t_1) - ik_{\tilde{y}_e} \bar{U}_{iy}^{(0)}(t-t_1)} \\
&\times \exp \left(-\frac{1}{2} (k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \rho_i^2 - \frac{1}{2} k_z^2 v_{Ti}^2 (t-t_1)^2 \right) \\
&\times \psi_e \left(k_{\tilde{X}_e} (1 + \bar{U}'_{ix}(t-t_1)) + k_{\tilde{y}_e} \bar{U}'_{iy}(t-t_1), k_{\tilde{y}_e}, \tilde{X}_e, t \right) \\
&\times \left\{ \left[ik_{\tilde{y}_e} v_{di} (1 - \eta_i) (1 - \bar{U}'_{ix} t_1) - k_z^2 v_{Ti}^2 (t-t_1) - \frac{i}{2} k_{\tilde{y}_e} v_{di} \eta_i k_z^2 v_{Ti}^2 (t-t_1)^2 \right. \right. \\
&\quad \left. \left. + ik_{\tilde{y}_e} v_{di} \eta_i \left(1 - \frac{1}{2} (k_{i\perp}^2(t) + k_{i\perp}^2(t_1)) \rho_i^2 \right) \right] I_0(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \right. \\
&\quad \left. + ik_{\tilde{y}_e} v_{di} \eta_i k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2 I_1(k_{i\perp}(t) k_{i\perp}(t_1) \rho_i^2) \right\}. \tag{69}
\end{aligned}$$

In Eq.(69), $T = T_i/T_e$, $\eta_i = d \ln T_i / d \ln n_i$, $\eta_e = d \ln T_e / d \ln n_e$, $v_{d\alpha} = (cT_\alpha / eB) d \ln n_{0\alpha} / dx$ is the ion(electron) ($\alpha = i(e)$) diamagnetic velocity, and I_0 and I_1 are the modified Bessel functions of the first kind and orders 0 and 1, respectively. For the steady plasmas, Eq. (69) becomes the integral equations of the convolution type, which can be solved by using various kinds of integral transform. In the $t_0 \rightarrow -\infty$ limit explored by the eigenmode analysis, Eq.(69) has the solution of the form $\varphi(\mathbf{k}, \omega) \varepsilon(\mathbf{k}, \omega) = 0$ for the Fourier transformed over time variable potential $\varphi(\mathbf{k}, \omega)$, where¹⁸

$$\begin{aligned}
\varepsilon(\mathbf{k}, \omega) &= 1 + T + i \sqrt{\frac{\pi}{2}} \frac{(\omega - k_y v_{di} (1 - \frac{\eta_i}{2}))}{k_z v_{Ti}} W(z_i) I_0(k_\perp^2 \rho_i^2) e^{-\rho_i^2 k_\perp^2} \\
&\quad - z_i \frac{k_y v_{di} \eta_i}{\sqrt{2} k_z v_{Ti}} \left(1 + i \sqrt{\frac{\pi}{2}} z_i W(z_i) \right) I_0(k_\perp^2 \rho_i^2) e^{-\rho_i^2 k_\perp^2} \\
&\quad + i \sqrt{\frac{\pi}{2}} \frac{k_y v_{di} \eta_i}{k_z v_{Ti}} W(z_i) k_\perp^2 \rho_i^2 e^{-k_\perp^2 \rho_i^2} (I_0(k_\perp^2 \rho_i^2) - I_1(k_\perp^2 \rho_i^2))
\end{aligned}$$

$$+iT\sqrt{\frac{\pi}{2}}\frac{(\omega - k_y v_{de})}{k_z v_{Te}}W(z_e), \quad (70)$$

and $W(z_{i(e)}) = e^{-z_{i(e)}^2} \left(1 + (2i/\sqrt{\pi}) \int_0^{z_{i(e)}} e^{t^2} dt \right)$ is the complex error function with argument $z_{i(e)} = \omega/\sqrt{2}k_z v_{Ti(e)}$. The solution $\omega(\mathbf{k})$ of the dispersion equation $\varepsilon(\mathbf{k}, \omega) = 0$ reveals the ion temperature gradient instability¹⁸, which develops due to inverse ion Landau damping. The maximum growth rate $\gamma(\mathbf{k})$ of that instability is of the order of the frequency. It attains for the perturbations, which have the phase velocity along the magnetic field of the order of the ion thermal velocity, i.e.

$$\omega(\mathbf{k}) \sim \gamma(\mathbf{k}) \sim k_z v_{Ti}, \quad (71)$$

and the normalized wave number component across the magnetic field, $k_{\perp}\rho_i$, comparable with the unity. It follows from Ref.¹⁹, that the kinetic ion temperature gradient instability of the steady plasmas may be saturated due to the scattering of ions by the ensemble of the unstable waves on a very low level for the root-mean-square (rms) magnitude $\tilde{\varphi} = (\int |\varphi(\mathbf{k}, \omega)|^2 d\mathbf{k}d\omega)^{1/2}$ of the perturbed electrostatic potential

$$\frac{e|\tilde{\varphi}|}{T_i} \sim \frac{k_{0z}}{k_{0\perp}} \sim \frac{\omega(\mathbf{k}_0)}{\omega_{ci}}, \quad (72)$$

where the estimates $k_{0z} \sim \omega(\mathbf{k}_0)/v_{Ti}$ and $k_{0\perp} \sim \rho_i^{-1}$ were used. It is assumed in Eq. (72) that the saturated spectrum of fluctuations of $\varphi(\mathbf{k})$ is peaked near the linearly most unstable wave vector, \mathbf{k}_0 .

VI. THE MACROSCALE EVOLUTION OF THE COMPRESSED-SHEARED CONVECTIVE FLOWS

The macroscale evolution of bulk of ions in the compressed-sheared plasma convective flow is determined by Eq. (33), which with velocity variables $\tilde{v}_{i\perp}$ and ϕ , for which $\tilde{v}_{ix} = \tilde{v}_{i\perp} \cos \phi$ and $\tilde{v}_{iy} = \tilde{v}_{i\perp} \sin \phi$, has a form

$$\begin{aligned} \frac{\partial \bar{F}_i}{\partial T} + \tilde{v}_{i\perp} \cos \phi (1 - \bar{U}'_{ix} T) \frac{\partial \bar{F}_i}{\partial \tilde{X}_i} + (\tilde{v}_{i\perp} \sin \phi - \tilde{v}_{i\perp} \cos \phi \bar{U}'_{iy} T) \frac{\partial \bar{F}_i}{\partial \tilde{Y}_i} - \frac{1}{\varepsilon} \omega_{ci} \frac{\partial \bar{F}_i}{\partial \phi} \\ + \tilde{v}_{iz} \frac{\partial \bar{F}_i}{\partial Z} - \frac{e_i}{m_i} \nabla \Phi(\tilde{X}_i, \tilde{Y}_i, T) \frac{\partial \bar{F}_i}{\partial \tilde{\mathbf{v}}_i} - \frac{e_i}{\varepsilon m_i} \left\langle \nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_{i0}(\tilde{\mathbf{r}}_i, \tilde{X}_i, T, \varepsilon) \frac{\partial f_i}{\partial \tilde{\mathbf{v}}_i} \right\rangle = 0. \end{aligned} \quad (73)$$

In the guiding center co-ordinates \hat{X}_i and \hat{Y}_i , determined by the relations,

$$\tilde{X}_i = \hat{X}_i - \varepsilon \frac{v_{i\perp}}{\omega_{ci}} \sin \left(\phi_1 - \frac{1}{\varepsilon} \omega_{ci} T \right) (1 - \bar{U}'_{ix} T) + O(\varepsilon^2), \quad (74)$$

$$\tilde{Y}_i = \hat{Y}_i + \varepsilon \frac{v_{i\perp}}{\omega_{ci}} \cos \left(\phi_1 - \frac{1}{\varepsilon} \omega_{ci} T \right) + \varepsilon \frac{v_{i\perp}}{\omega_{ci}} \sin \left(\phi_1 - \frac{1}{\varepsilon} \omega_{ci} T \right) \bar{U}'_{iy} T + O(\varepsilon^2), \quad (75)$$

with $v_{i\perp} = \hat{v}_{i\perp}$ and $\phi = \phi_1 - \frac{1}{\varepsilon} \omega_{ci} T$, Eq. (73) for $\bar{F}_i(\hat{v}_{i\perp}, \phi, \hat{X}_i, \hat{Y}_i, T, \varepsilon)$, averaged over the fast time $t = \frac{T}{\varepsilon}$, becomes

$$\begin{aligned} & \frac{\partial \bar{F}_i}{\partial T} - \frac{e_i}{m_i} \left\{ \frac{1}{\varepsilon} \frac{\omega_{ci}}{\hat{v}_{i\perp}} \left(\frac{\partial \Phi_i}{\partial \hat{v}_{i\perp}} \frac{\partial \bar{F}_i}{\partial \phi} - \frac{\partial \Phi_i}{\partial \phi} \frac{\partial \bar{F}_i}{\partial \hat{v}_{i\perp}} \right) + \frac{\partial \Phi}{\partial Z} \frac{\partial \bar{F}_i}{\partial v_z} \right. \\ & \left. + \frac{\varepsilon}{\omega_{ci}} \left((1 - \bar{U}'_{ix} T) \frac{\partial \Phi_i}{\partial \hat{Y}_i} \frac{\partial \bar{F}_i}{\partial \hat{X}_i} - \frac{\partial \Phi_i}{\partial \hat{X}_i} \frac{\partial \bar{F}_i}{\partial \hat{Y}_i} - \frac{1}{2} \bar{U}'_{ix} T \frac{\partial \Phi_i}{\partial \hat{Y}_i} \frac{\partial \bar{F}_i}{\partial \hat{Y}_i} \right) \right\} \\ & - \frac{1}{\varepsilon} \frac{e_i}{m_i} \left\langle \nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_{i0}(\tilde{\mathbf{r}}_i, \tilde{X}_i, T, \varepsilon) \frac{\partial f_i}{\partial \tilde{\mathbf{v}}_i} \right\rangle = 0. \end{aligned} \quad (76)$$

The solution to Eq. (76) we find in the form

$$\bar{F}_i(\hat{v}_{i\perp}, \phi, v_z, \hat{X}_i, \hat{Y}_i, Z_i, T) = \bar{F}_{i0}(\hat{v}_{i\perp}, v_z, \hat{X}_i, Z_i, T) + \bar{F}_{i1}(\hat{v}_{i\perp}, \phi, v_z, \hat{X}_i, \hat{Y}_i, z_i, T, \varepsilon), \quad (77)$$

where \bar{F}_{i0} is the equilibrium ion distribution function inhomogeneous along co-ordinate \hat{X}_i .

It is determined by the quasilinear equation

$$\frac{\partial \bar{F}_{i0}}{\partial T} = \frac{1}{\varepsilon} \frac{e_i}{m_i} \left\langle \nabla_{\tilde{\mathbf{r}}_i} \tilde{\varphi}_{i0}(\tilde{\mathbf{r}}_i, \tilde{X}_i, T, \varepsilon) \frac{\partial f_i}{\partial \tilde{\mathbf{v}}_i} \right\rangle. \quad (78)$$

Employing Eq. (38) for $f_i(\hat{v}_{i\perp}, \phi_1, v_z, \hat{x}_i, \hat{y}_i, z, \hat{X}_i, t)$ and Eq. (39) for $\tilde{\varphi}_{i0}(\tilde{\mathbf{r}}_i, \tilde{X}_i, T, \varepsilon)$ in Eq. (78), we derived the quasilinear equation for \bar{F}_{i0} ,

$$\begin{aligned} \frac{\partial \bar{F}_{i0}}{\partial T} &= \frac{e_i^2}{m_i^2} \sum_{n=-\infty}^{\infty} \int_{T_0}^T dT_1 \int d\mathbf{k} \left(\frac{k_y}{\omega_{ci}} (1 - \bar{U}'_{ix} t) \frac{\partial}{\partial X} + \frac{n\omega_{ci}}{\hat{v}_{i\perp}} \frac{\partial}{\partial \hat{v}_{i\perp}} + k_z \frac{\partial}{\partial v_z} \right) \\ & \times J_n \left(\frac{k_{i\perp}(T) \hat{v}_{i\perp}}{\omega_{ci}} \right) J_n \left(\frac{k_{i\perp}(T_1) \hat{v}_{i\perp}}{\omega_{ci}} \right) \left\langle \tilde{\varphi}_{i0}(\tilde{\mathbf{k}}_i, \tilde{X}_i, t) \tilde{\varphi}_{i0}(\tilde{\mathbf{k}}_i, \tilde{X}_i, t_1) \right\rangle \\ & \times \left(\frac{k_y}{\omega_{ci}} (1 - \bar{U}'_{ix} t_1) \frac{\partial \bar{F}_{i0}}{\partial X} + \frac{n\omega_{ci}}{\hat{v}_{i\perp}} \frac{\partial \bar{F}_{i0}}{\partial \hat{v}_{i\perp}} + k_z \frac{\partial \bar{F}_{i0}}{\partial v_z} \right) \end{aligned} \quad (79)$$

The function $\bar{F}_{i1}(\hat{v}_{i\perp}, \phi, v_z, \hat{X}_i, \hat{Y}_i, z_i, T, \varepsilon)$ is the perturbation of \bar{F}_{i0} , caused by the electrostatic potential $\Phi_i(\tilde{X}_i, \tilde{Y}_i, T)$ of the plasma respond on the sheared-compressed convective flows. The equation for \bar{F}_{i1} ,

$$\begin{aligned} & \frac{\partial}{\partial T} \bar{F}_{i1}(\hat{v}_{i\perp}, \phi, v_z, \hat{X}_i, \hat{Y}_i, z_i, T, \varepsilon) \\ &= \frac{e_i}{m_i} \left\{ \frac{\varepsilon}{\omega_{ci}} (1 - \bar{U}'_{ix} T) \frac{\partial \Phi_i}{\partial \hat{Y}_i} \frac{\partial \bar{F}_{i0}}{\partial \hat{X}_i} - \frac{\omega_{ci}}{\varepsilon} \frac{1}{\hat{v}_{i\perp}} \frac{\partial \Phi_i}{\partial \phi} \frac{\partial \bar{F}_{i0}}{\partial \hat{v}_{i\perp}} + \frac{\partial \Phi_i}{\partial Z} \frac{\partial \bar{F}_{i0}}{\partial v_z} \right\}, \end{aligned} \quad (80)$$

which follows from Eqs. (70) and (72). The solution to Eq. (80) may be derived by the Fourier transforming of Eq. (80) over \hat{X}_i, \hat{Y}_i and employing the relation

$$\begin{aligned}\Phi_i(\tilde{X}_i, \tilde{Y}_i, Z, T) &= \int \Phi_i(\mathbf{K}, T) \exp\left(iK_{\tilde{X}_i}\tilde{X}_i + iK_{\tilde{Y}_i}\tilde{Y}_i + iK_z z\right) dK_{\tilde{X}_i} dK_{\tilde{Y}_i} dK_z \\ &= \int \Phi_i(\mathbf{K}, T) \exp\left(iK_{\tilde{X}_i}\hat{X}_i + iK_{\tilde{Y}_i}\hat{Y}_i + iK_z z_1\right) \\ &\quad \times \sum_{n=-\infty}^{\infty} J_n\left(\frac{K_{i\perp}(T)\hat{v}_{i\perp}}{\omega_{ci}}\right) e^{-in(\phi - \frac{1}{\varepsilon}\omega_{ci}T - \chi_i(T))} dK_{\tilde{X}_i} dK_{\tilde{Y}_i} dK_z\end{aligned}\quad (81)$$

where $K_{i\perp}(T)$ and $\chi_i(T)$ are determined by the relations

$$K_{i\perp}^2(T) = (K_{\tilde{X}_i} - (K_{\tilde{X}_i}\bar{U}'_{ix} + K_{\tilde{Y}_i}\bar{U}'_{iy})T)^2 + K_{\tilde{Y}_i}^2, \quad \sin\chi_i(T) = \frac{K_{\tilde{Y}_i}}{K_{i\perp}(T)}. \quad (82)$$

This solution for $\bar{F}_{i1}(\hat{v}_{i\perp}, v_z, K_{\tilde{X}_i}, K_{\tilde{Y}_i}, K_z, T)$ for slow, $\frac{\partial\Phi_i}{\partial T} \ll \omega_{ci}\Phi_i$, macroscale perturbations has a form

$$\begin{aligned}\bar{F}_{i1}(\hat{v}_{i\perp}, v_z, K_{\tilde{X}_i}, K_{\tilde{Y}_i}, K_z, T) &= i\frac{e_i}{m_i} \int dT_1 \Phi_i(\mathbf{K}_i, T_1) J_0\left(\frac{K_{i\perp}(T)\hat{v}_{i\perp}}{\omega_{ci}}\right) J_0\left(\frac{K_{i\perp}(T_1)\hat{v}_{i\perp}}{\omega_{ci}}\right) \\ &\quad \times \left[\frac{K_{\tilde{Y}_i}}{\omega_{ci}}(1 - \bar{U}'_{ix}T_1)\frac{\partial\bar{F}_{i0}}{\partial\tilde{X}_i} + K_z\frac{\partial\bar{F}_{i0}}{\partial v_{iz}}\right] e^{-iK_z v_{iz}(t-t_1)},\end{aligned}\quad (83)$$

As it follows from Eqs. (A6) and (A7), the electron convective velocities are negligible small and are assumed to be equal to zero. The electron Vlasov equation for the average electron distribution function $\bar{F}_e(\hat{v}_{e\perp}, \phi, v_z, \hat{X}_e, \hat{Y}_e, z_e, T, \varepsilon)$ in the electron guiding center coordinates $\hat{X}_e \approx \tilde{X}_e, \hat{Y}_e \approx \tilde{Y}_e$ for $\tilde{X}_e \gg \rho_e$ and $\tilde{Y}_e \gg \rho_e$ becomes

$$\begin{aligned}\frac{\partial\bar{F}_e}{\partial T} - \frac{e}{m_e} \left\{ \frac{1}{\varepsilon} \frac{\omega_{ce}}{\hat{v}_{e\perp}} \left(\frac{\partial\Phi_e}{\partial\hat{v}_{e\perp}} \frac{\partial\bar{F}_e}{\partial\phi} - \frac{\partial\Phi_e}{\partial\phi} \frac{\partial\bar{F}_e}{\partial\hat{v}_{e\perp}} \right) + \frac{\partial\Phi}{\partial Z} \frac{\partial\bar{F}_e}{\partial v_z} \right. \\ \left. + \frac{\varepsilon}{\omega_{ci}} \left(\frac{\partial\Phi_e}{\partial\hat{Y}_e} \frac{\partial\bar{F}_e}{\partial\hat{X}_e} - \frac{\partial\Phi_e}{\partial\hat{X}_e} \frac{\partial\bar{F}_e}{\partial\hat{Y}_e} \right) \right\} \\ + \frac{1}{\varepsilon} \frac{e}{m_e} \left\langle \tilde{\mathbf{E}}_{e0}(X_e, \varepsilon^{-1}X_e, \varepsilon^{-1}Y_e, Z_e, \varepsilon^{-1}T) \frac{\partial f_e}{\partial\tilde{\mathbf{v}}_e} \right\rangle = 0.\end{aligned}\quad (84)$$

The solution to Eq. (84) for the electron distribution function \bar{F}_e we derive in the form (77) applied for \bar{F}_i ,

$$\bar{F}_e(\hat{v}_{e\perp}, \phi, \hat{X}_e, \hat{Y}_e, T, \varepsilon) = \bar{F}_{e0}(\hat{v}_{e\perp}, v_{ez}, \hat{X}_e, T) + \bar{F}_{e1}(\hat{v}_{e\perp}, \phi, v_{ez}, \hat{X}_e, \hat{Y}_e, Z_e, T, \varepsilon), \quad (85)$$

in which we distinguish the equilibrium electron distribution function $\bar{F}_{e0}(\hat{v}_{e\perp}, v_{ez}, \hat{X}_e, T)$, determined by the quasilinear equation

$$\frac{\partial\bar{F}_{e0}}{\partial T} = -\frac{1}{\varepsilon} \frac{e}{m_e} \left\langle \tilde{\mathbf{E}}_{e0}(\hat{X}_e, \hat{x}_e, \hat{y}_e, z_e, \varepsilon^{-1}T) \frac{\partial f_e}{\partial\tilde{\mathbf{v}}_e} \right\rangle, \quad (86)$$

and the perturbation $\bar{F}_{e1}(\hat{v}_{e\perp}, \phi, v_z, \hat{X}_e, \hat{Y}_e, z_e, T, \varepsilon)$ of \bar{F}_{e0} , which stems from the formation of the self-consistent potential Φ_e of the plasma response on the formation of the convective flows. The equation for \bar{F}_{e1} follows from Eqs. (84) - (86) and for the perturbations, for which $\frac{\partial \Phi_e}{\partial T} \ll \omega_{ce} \Phi_e$, has a form

$$\frac{\partial}{\partial T} \bar{F}_{e1}(\hat{v}_{e\perp}, \phi, v_z, \hat{X}_e, \hat{Y}_e, Z, T, \varepsilon) = \frac{e}{m_e} \left\{ \frac{\varepsilon}{\omega_{ce}} \frac{\partial \Phi_e}{\partial \hat{Y}_e} \frac{\partial \bar{F}_{e0}}{\partial \hat{X}_e} + \frac{\partial \Phi_e}{\partial Z} \frac{\partial \bar{F}_{e0}}{\partial v_z} \right\}. \quad (87)$$

The solution to Eq. (87), which determines the evolution of the separate spatial long wavelength, $K_{e\perp} \rho_e \ll 1$, macroscale Fourier harmonic $\bar{F}_{e1}(\hat{\mathbf{v}}_e, K_{\hat{X}_e}, K_{\hat{Y}_e}, Z, T)$ of the perturbation $\bar{F}_{e1}(\hat{\mathbf{v}}_e, \hat{X}_e, \hat{Y}_e, Z, T)$,

$$\begin{aligned} \bar{F}_{e1}(\hat{v}_{e\perp}, v_z, K_{\hat{X}_e}, K_{\hat{Y}_e}, K_z, T) &= i \frac{e}{\varepsilon m_e} \int dT_1 \Phi(\mathbf{K}_e, T_1) \\ &\times \left[\frac{K_{\hat{Y}_e}}{\omega_{ce}} \frac{\partial \bar{F}_{i0}}{\partial \hat{X}_e} + K_z \frac{\partial \bar{F}_{e0}}{\partial v_{ez}} \right] e^{-iK_z v_{iz}(t-t_1)}, \end{aligned} \quad (88)$$

was derived by the Fourier transforming of Eq. (87) over \hat{X}_e, \hat{Y}_{ei} . In Eq. (82), the Fourier transformation of the potential $\Phi(\tilde{X}_e, \tilde{Y}_e, Z_e, T)$ over coordinates \tilde{X}_e, \tilde{Y}_e ,

$$\Phi_e(\tilde{X}_e, \tilde{Y}_e, Z, T) = \int \Phi_e(\mathbf{K}_e, T) \exp\left(iK_{\tilde{X}_e} \hat{X}_e + iK_{\tilde{Y}_e} \hat{Y}_e + iK_z z\right) dK_{\tilde{X}_e} dK_{\tilde{Y}_e} dK_z, \quad (89)$$

was used. In Eq. (81), the potential Φ_i of the macroscale plasma response on the formation of the compressed-sheared convective flow is determined by the Poisson equation,

$$\begin{aligned} &\frac{\partial^2 \Phi_i(\tilde{X}_i, \tilde{Y}_i, \tilde{Z}_i, t)}{\partial^2 \tilde{X}_i} + \frac{\partial^2 \Phi_i(\tilde{X}_i, \tilde{Y}_i, \tilde{Z}_i, t)}{\partial^2 \tilde{Y}_i} + \frac{\partial^2 \Phi_i(\tilde{X}_i, \tilde{Y}_i, \tilde{Z}_i, t)}{\partial^2 \tilde{Z}_i} \\ &= -4\pi \left(e_i \int d\mathbf{v}_i \bar{F}_{i1}(\tilde{\mathbf{v}}_i, \tilde{X}_i, \tilde{Y}_i, \tilde{Z}_i, t) - |e| \int d\mathbf{v}_e \bar{F}_{e1}(\tilde{\mathbf{v}}_e, \tilde{X}_e, \tilde{Y}_e, \tilde{Z}_i, t) \right). \end{aligned} \quad (90)$$

Eq. (83) for \bar{F}_{i1} and Eq. (88) for \bar{F}_{e1} , and the Poisson equation (90) compose the system, which governs the kinetic macroscale evolution of a plasma under the action of the spatially inhomogeneous microturbulence.

VII. CONCLUSIONS

In this paper, we present the two-scale non-modal approach to the kinetic theory of the microscale turbulence of a plasma, inhomogeneous on the macroscales across the confined magnetic field. This approach reveals the effect of the formation of the macroscale convective

flows of such a plasma caused by the interaction of ions with microscale turbulence. The flow velocities are found as the average velocities of the motion of ions and electrons in the spatially inhomogeneous microturbulence and are proportional to the gradient of the spectral intensity of the electric field of the microturbulence. It was found (Eqs. (A6) and (A7)) that the velocities of the electron convective flows are negligible small relative to the ion flow velocities and, therefore, the convective motion of electron can be neglected. This result predicts that the macroscale convective flow transports mostly the ions. For the ion flow, generated by the low frequency microturbulence with radially decreasing spectral intensity, Eq. (A6) predicts that the radial velocity of the ion flow is directed outward of the plasma core to the edge of the tokamak plasma. This result reveals the necessity in the investigations of the temporal evolution of the macroscale convective flow of ions in the edge region of the tokamak plasma, investigation of the radial accumulation of ions and formation of the radially inhomogeneous mesoscale ion density growth, and formation of the localized radial electric field -the precursor of the formation of the mesoscale poloidal sheared flow.

Any microscale perturbation in the radially inhomogeneous flows, which before the development of the sheared-compressed flow had a plane wave structure, experiences the continuous distortion in the flow and become the sheared-compressed mode with time dependent structure. This distortion grows with time and forms a time-dependent nonmodal process, which affects the microturbulence and the average ion distribution function. The derived quasilinear equation (79), which determines the nonmodal evolution of the average ion distribution function, resulted from the interactions of ions with ensemble of the microscale sheared-compressed waves, and the integral equation (69), for the electrostatic potential for the microscale perturbations, which determines the macroscale nonlinear back-reaction of the sheared-compressed convective flows on the microscale perturbations, are the basic equations which determined the long time macroscale evolution of the plasma with inhomogeneous microturbulence. This work was supported by National R& D Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (Grant No. NRF-2022R1A2C1012808126)

ACKNOWLEDGMENTS

This work was supported by National R&D Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (Grant No. NRF-2022R1A2C1012808126)

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Appendix A: Solutions to Eqs. (31) and (32) for $\bar{U}_{ix}(X_i, t)$ and $\bar{U}_{iy}(X_i, t)$

Direct integration of Eqs. (31), (32) gives

$$\begin{aligned} \bar{U}_{ix}(\tilde{X}_i, t) &= \frac{e_i}{m_i} \int_0^t dt_1 \left[\langle \tilde{E}_{i1x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_1) \rangle \cos \omega_{ci}(t - t_1) \right. \\ &\quad \left. + \langle \tilde{E}_{i1y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_1) \rangle \sin \omega_{ci}(t - t_1) \right], \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} \bar{U}_{iy}(\tilde{X}_i, t) &= \frac{e_i}{m_i} \int_0^t dt_1 \left[- \langle \tilde{E}_{i1x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_1) \rangle \sin \omega_{ci}(t - t_1) \right. \\ &\quad \left. + \langle \tilde{E}_{i1y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_1) \rangle \cos \omega_{ci}(t - t_1) \right], \end{aligned} \quad (\text{A2})$$

where

$$\tilde{E}_{i1x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) = \frac{\partial}{\partial \tilde{X}_i} \left(\tilde{E}_{i0x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \right) \cdot \tilde{R}_{ix}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t), \quad (\text{A3})$$

$$\tilde{E}_{i1y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) = \frac{\partial}{\partial \tilde{X}_i} \left(\tilde{E}_{i0y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) \right) \cdot \tilde{R}_{ix}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t), \quad (\text{A4})$$

$$\begin{aligned} \tilde{R}_{ix}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t) &= \int_0^t dt_1 \tilde{U}_{ix}^{(0)}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_1) \\ &= \frac{e_i}{m_i} \int_0^t dt_1 \int_0^{t_1} dt_2 \left[\langle \tilde{E}_{i0x}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_2) \rangle \cos \omega_{ci}(t_1 - t_2) \right. \\ &\quad \left. + \langle \tilde{E}_{i0y}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t_2) \rangle \sin \omega_{ci}(t_1 - t_2) \right], \end{aligned} \quad (\text{A5})$$

where $\tilde{\mathbf{E}}_{i0}(\tilde{X}_i, \tilde{x}_i, \tilde{y}_i, t)$ is the electric field $\mathbf{E}_i(\mathbf{r}_i, X_i, t)$, determined in $\tilde{\mathbf{r}}_i, \tilde{X}_i$ coordinates. The integration of Eqs. (A1), (A2) with accounting for Eqs. (A3) - (A5) and averaging over

the fast time $t \gg \omega_{ci}^{-1}$ velocities $\bar{U}_{ix}(\tilde{X}_i, t)$ and $\bar{U}_{iy}(\tilde{X}_i, t)$ gives for $\bar{U}_{ix}(\tilde{X}_i)$ and $\bar{U}_{iy}(\tilde{X}_i)$ solutions

$$\begin{aligned} \bar{U}_{ix}(\tilde{X}_i) &= \langle \bar{U}_{ix}(\tilde{X}_i, t) \rangle_t \\ &= \frac{e_i^2}{m_i^2} \frac{1}{\omega_{ci}} \frac{1}{(2\pi)^3} \int d\mathbf{k}_i \frac{\partial}{\partial \tilde{X}_i} \left(\tilde{E}_{i0y}(\tilde{X}_i, \mathbf{k}_i) \right) \tilde{E}_{i0x}(\tilde{X}_i, \mathbf{k}_i) \frac{1}{(\omega_{ci}^2 - \omega^2(\mathbf{k}_i))}, \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} \bar{U}_{iy}(\tilde{X}_i) &= \langle \bar{U}_{iy}(\tilde{X}_i, t) \rangle_t \\ &= \frac{e_i^2}{m_i^2} \frac{1}{\omega_{ci}} \frac{1}{(2\pi)^3} \int d\mathbf{k}_i \frac{\partial}{\partial \tilde{X}_i} \left(\tilde{E}_{i0x}(\tilde{X}_i, \mathbf{k}_i) \right)^2 \frac{1}{(\omega_{ci}^2 - \omega^2(\mathbf{k}_i))}. \end{aligned} \quad (\text{A7})$$

The velocities $\bar{U}_{ex}(\tilde{X}_e)$ and $\bar{U}_{ey}(\tilde{X}_e)$ of the electron convective flows are determined by Eqs. (A6), (A7) with ion species index changed on the electron species index. It follows from Eqs. (A6), (A7) that the electron convective velocities are in ω_{ce}/ω_{ci} times less than the ion convective velocities $\bar{U}_{ix}(\tilde{X}_i)$ and $\bar{U}_{iy}(\tilde{X}_i)$ and, therefore, the convective motion of electrons may be neglected. This result predicts that the macroscale convective flow, generated by the low frequency microscale turbulence with radially decreasing spectral density, transports mostly the ions with velocity directed outward of the plasma core to the edge of the tokamak plasma. This result reveals the necessity in the investigation of the temporal evolution of the macroscale convective flow of ions in the edge region of the tokamak plasma, investigation of the radial accumulation of ions and formation of the radially inhomogeneous mesoscale ion density growth, and formation of the localized radial electric field - the precursors of the formation of the mesoscale poloidal sheared flow.

REFERENCES

- ¹C. K. Phillips, R. E. Bell, L. A. Berry, P. T. Bonoli, R. W. Harvey, J. C. Hosea, E. F. Jaeger, B. P. LeBlanc, P. M. Ryan, G. Taylor, E. J. Valeo, J. B. Wilgen, J. R. Wilson, J. C. Wright, H. Yuh and the NSTX Team, "Spectral effects on fast wave core heating and current drive", Nuclear Fusion **49**, 075015 (2009).
- ²G. Taylor, R. E. Bell, J. C. Hosea, B. P. LeBlanc, C. K. Phillips, M. R. Wilson, J-W. Ahn, G. Chen, D. L. Green, E. F. Jaeger, R. Maingi, P. M. Ryan, J. B. Wilgen, W. W. Heidbrink, D. Liu, P. T. Bonoli, T. Brecht, M. Choi, and R. W. Harvey, "Advances in high-harmonic fast wave physics in the National Spherical Torus Experiment", Phys. Plasmas **17**, 056114 (2010).

- ³R. J. Perkins, J. C. Hosea, G. J. Kramer, J.-W. Ahn, R. E. Bell, A. Diallo, S. Gerhardt, T. K. Grey, D. L. Green, E. F. Jaeger, M. A. Jaworski, B. P. LeBlanc, A. McLean, R. Maingi, C. K. Phillips, L. Roquemore, P. M. Ryan, S. Sabbagh, G. Taylor, J. R. Wilson, "High-harmonic fast-wave power flow along magnetic field lines in the scrape-off Layer of NSTX", *Phys. Rev. Lett.* **109**, 045001 (2012).
- ⁴R. J. Perkins, J.-W. Ahn, R. E. Bell, A. Diallo, S. Gerhardt, T. K. Gray, D. L. Green, E. F. Jaeger, J. C. Hosea, M. A. Jaworski, B. P. LeBlanc, G. J. Kramer, A. McLean, R. Maingi, C. K. Phillips, M. Podest'a, L. Roquemore, P. M. Ryan, S. Sabbagh, F. Scotti, G. Taylor, J. R. Wilson, "Fast-wave power flow along SOL field lines in NSTX and the associated power deposition profile across the SOL in front of the antenna. *Nucl. Fusion* **53**, 083025 (2013).
- ⁵M. Porkolab, "Parametric processes in magnetically confined CTR plasmas", *Nucl. Fusion* **18**, 367 (1978).
- ⁶M. Porkolab, "Parametric instabilities in the tokamak edge plasma in the ion cyclotron heating regimes", *Fusion Engineering and Design*, **12**, 93 (1990).
- ⁷M. Porkolab, R. I. Pinsky, "Parametric instabilities during high power helicon wave injection on DIII-D," *EPJ Web Conf.* **157**, 03042 (2017).
- ⁸M. Porkolab, "Parametric instabilities in the tokamak edge plasma in the ion cyclotron heating regime", *Fusion Eng. Des.* **12**, 93 (1990).
- ⁹J. R. Wilson, S. Bernabei, T. Biewer, S. Diem, J. Hosea, B. LeBlanc, C. K. Phillips, P. Ryan, and D. W. Swain. Parametric Decay During HHFW on NSTX. *AIP Conference Proceedings* **787**, 66 (2005).
- ¹⁰T. M. Biewer, R. E. Bell, S. J. Diem, C. K. Phillips, and R. J. Wilson, "Edge ion heating by launched high harmonic fast waves in the National Spherical Torus Experiment", *Phys. Plasmas* **12**, 056108 (2005).
- ¹¹J. C. Rost, M. Porkolab, R. L. Boivin, "Edge ion heating and parametric decay during injection of ion cyclotron resonance frequency power on the Alcator C-Mod tokamak" *Phys. Plasmas* **9**, 1262 (2002).
- ¹²V. S. Mikhailenko, V. V. Mikhailenko, Hae June Lee. "Ion cyclotron parametric turbulence and the anomalous heating of ions in the tokamak edge plasma in the fast wave heating regime", *Phys. Plasmas* **27**, 052508 (2020).

- ¹³V. V. Mikhailenko, Hae June Lee, V. S. Mikhailenko. "The ion cyclotron parametric instabilities and the anomalous heating of ions in the scrape-off layer of tokamak plasma in the high harmonic fast wave heating regime", *Phys. Plasmas* **31**, 032307 (2024).
- ¹⁴V. S. Mikhailenko, V. V. Mikhailenko, Hae June Lee. "The ion cyclotron parametric instabilities and anomalous convective transport of the inhomogeneous plasma in front of the fast wave antenna". *Phys. Plasmas* **28**, 042304 (2021).
- ¹⁵V. S. Mikhailenko, V. V. Mikhailenko, Hae June Lee, "Anomalous convective transport of the tokamak edge plasma, caused by the inhomogeneous ion cyclotron parametric turbulence", *Phys. Plasmas* **29**, 072301 (2022).
- ¹⁶V. S. Mikhailenko, V. V. Mikhailenko, Hae June Lee, "Nonmodal kinetic theory of the stability of the compressed – sheared plasma flows generated by the inhomogeneous microscale turbulence in the tokamak edge plasma", *Phys. Plasmas* **30**, 052302 (2023).
- ¹⁷V. S. Mikhailenko, V. V. Mikhailenko, K. N. Stepanov. "Renormalized non-modal theory of the kinetic drift instability of plasma shear flow", *Phys. Plasmas* **18**, 062103 (2011).
- ¹⁸A. B. Mikhailovski, *Theory of Plasma Instabilities*, Volume 2: Instabilities of an inhomogeneous plasma (Springer-Verlag, 2013).
- ¹⁹V. V. Mikhailenko, V. S. Mikhailenko, Hae June Lee, "Renormalized theory of ion temperature gradient instability of the magnetic-field-aligned plasma shear flow with hot ions", *Phys. Plasmas* **22**, 102308 (2015).