

# SELF-IMPROVING PROPERTIES OF WEIGHTED NORM INEQUALITIES ON METRIC MEASURE SPACES

JUHA KINNUNEN, JUHA LEHRBÄCK, ANTTI V. VÄHÄKANGAS, AND DACHUN YANG

ABSTRACT. This work discusses self-improving properties of the Muckenhoupt condition and weighted norm inequalities for the Hardy–Littlewood maximal function on metric measure spaces with a doubling measure. Our main result provides direct proofs of these properties by applying a Whitney covering argument and a technique inspired by the Calderón–Zygmund decomposition. In particular, this approach does not rely on reverse Hölder inequalities.

## 1. INTRODUCTION

Harmonic analysis on metric measure spaces, and more broadly on spaces of homogeneous type, continues to be an active area of research. Coifman and Weiss provided a systematic account of many aspects of harmonic analysis on spaces of homogeneous type in [2]. The theory of weighted norm inequalities and Muckenhoupt weights has been explored in greater detail in works such as those by Strömberg and Torchinsky in [12] and by Genebashvili, Gogatishvili, Kokilashvili and Krbec in [5]. Muckenhoupt weights exhibit a remarkable self-improving property: if a weight belongs to a certain Muckenhoupt class  $A_p$ ,  $1 < p < \infty$ , then it also belongs to another Muckenhoupt class  $A_{p-\varepsilon}$  with a slightly smaller index, see [12, Lemma 8, p. 5]. This self-improving nature of Muckenhoupt weights arises from the fact that these weights satisfy a reverse Hölder inequality, see [12, Theorem 15, p. 9] for details.

Reverse Hölder inequalities are powerful tools in the study of weighted norm inequalities, but the purpose of this work is to discuss an alternative approach to the self-improving property of the Muckenhoupt weights that entirely avoids relying on reverse Hölder inequalities. Thus the novelty in this work lies in the methodology rather than in the actual results.

Instead of reverse Hölder inequalities, we utilize the close connection between Muckenhoupt weights and weighted norm inequalities for the Hardy–Littlewood maximal function, see [3, 4]. More specifically, let  $(X, d, \mu)$  be a metric measure space equipped with a doubling measure  $\mu$  and let  $w$  be a weight in  $X$ ; see Sections 2 and 3 for the definitions. Then the following assertions are equivalent, for  $1 < p < \infty$ .

(A) There exists a constant  $C > 0$  such that

$$\int_X (M^* f(x))^p w(x) d\mu(x) \leq C \int_X |f(x)|^p w(x) d\mu(x) \quad (1.1)$$

for every  $f \in L^\infty(X)$  with a bounded support.

(B) The weight  $w$  belongs to the Muckenhoupt class  $A_p$ .

(C) There exists a constant  $C > 0$  such that inequality (1.1) holds for every  $f \in L^p(w d\mu)$ .

In (1.1), the (noncentered) Hardy–Littlewood maximal function of  $f$  is defined as

$$M^* f(x) = \sup_{B \ni x} \frac{1}{\mu(B)} \int_B |f(y)| d\mu(y),$$

*Date:* January 30, 2025.

*2020 Mathematics Subject Classification.* 42B25, 42B35, 46E30, 30L99.

*Key words and phrases.* Maximal function, Muckenhoupt weight, analysis on metric spaces.

where the supremum is taken over all balls  $B$  in  $X$  containing  $x$ . It was shown in [2] that  $M^*$  is a bounded operator on  $L^p(X)$  for every  $1 < p \leq \infty$  and of weak type  $(1, 1)$  under the assumption that  $\mu$  is a doubling measure; see also [12, Theorem 3, p. 3]. Observe that assertion (C) above states that the operator  $M^*$  is bounded on  $L^p(w d\mu)$ , for  $1 < p < \infty$ .

The equivalence of the assertions (B) and (C) is well known, see [12, Theorem 9, p. 5]. For our purposes, it is convenient to consider also assertion (A), which is a priori slightly weaker than (C). That (A) implies (B) is shown in Lemma 4.2, and the more involved implication from (B) to (C) is proved in Theorem 4.5 by adapting an argument of Lerner [10]. This approach is direct and, in particular, does not refer to reverse Hölder inequalities. Finally, the implication from (C) to (A) is a triviality.

Now, assume that assertion (C) holds, i.e., that  $M^*$  is bounded on some  $L^p(w d\mu)$ ,  $1 < p < \infty$ , where  $w$  is a weight and  $\mu$  a doubling measure in  $X$ . By the Marcinkiewicz interpolation theorem and the boundedness of the maximal operator  $M^*$  on  $L^\infty(w d\mu)$ , it follows that then  $M^*$  is also bounded on  $L^{p+\varepsilon}(w d\mu)$  for every  $\varepsilon > 0$ . On the other hand, by using the equivalence of the assertions (B) and (C) and the known self-improving property of the Muckenhoupt  $A_p$  condition, there exists some  $\varepsilon > 0$  such that  $M^*$  is bounded on  $L^{p-\varepsilon}(w d\mu)$  as well.

The main goal in this work is to give a direct and essentially self-contained proof for this self-improving property of the weighted norm inequalities. In particular, we show in Lemma 5.6 that if assertion (C) holds, then there exist  $\varepsilon > 0$  and  $C > 0$  such that

$$\int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \leq C \int_X |f(x)|^{p-\varepsilon} w(x) d\mu(x) \quad (1.2)$$

for every  $f \in L^\infty(X)$  with a bounded support. The proof of Lemma 5.6 uses a Whitney covering argument and a technique inspired by the Calderón–Zygmund decomposition, based on an unpublished note on Muckenhoupt weights by Keith and Zhong. This approach extends the corresponding Euclidean result with the Lebesgue measure [8, Theorem 8.11] to metric spaces and it does not rely on reverse Hölder inequalities. Similar ideas have been applied in various contexts, for example, in Keith and Zhong [7], Kinnunen, Lehrbäck, Vähäkangas and Zhong [9] and Lewis [11].

By the equivalence of the assertions (A), (B) and (C), we obtain from Lemma 5.6 an independent proof for the self-improving property of the Muckenhoupt classes, see Theorem 5.11, and we also conclude that the improved weighted norm inequality (1.2) holds for all functions  $f \in L^{p-\varepsilon}(w d\mu)$ , see Corollary 5.13.

Before the statements and proofs of our main results and tools in Sections 4 and 5, we recall preliminaries on metric spaces and on Muckenhoupt weights in Sections 2 and 3, respectively. In Section 3 we also prove some basic properties of the Muckenhoupt classes for the sake of completeness.

## Acknowledgements

The authors would like to thank Xiao Zhong for interesting discussions on the topics of the paper. Dachun Yang is supported by the National Natural Science Foundation of China (Grant numbers 12431006 and 12371093).

## 2. PRELIMINARIES

Throughout the paper, we assume that  $X = (X, d, \mu)$  is a metric measure space equipped with a metric  $d$  and a positive complete Borel measure  $\mu$  such that  $0 < \mu(B) < \infty$  for all open balls  $B \subset X$ , where

$$B = B(x, r) = \{y \in X : d(y, x) < r\}$$

with  $x \in X$  and  $r > 0$ . The space  $X$  is separable under the above assumptions, we refer to [1, Proposition 1.6]. We also assume throughout that  $\#X \geq 2$  and that the measure  $\mu$  is

doubling, that is, there exists a constant  $c_\mu > 1$ , called the doubling constant of  $\mu$ , such that

$$\mu(2B) \leq c_\mu \mu(B) \quad (2.1)$$

for all balls  $B = B(x, r)$  in  $X$ . Here we use for  $0 < t < \infty$  the notation  $tB = B(x, tr)$ . We let

$$\int_A f(y) d\mu(y) = \frac{1}{\mu(A)} \int_A f(y) d\mu(y)$$

be the integral average of  $f \in L^1(A)$  over a measurable set  $A \subset X$  with  $0 < \mu(A) < \infty$ . We say that a function  $f: X \rightarrow \mathbb{R}$  has a bounded support, if  $\{x \in X : f(x) \neq 0\}$  is contained in some ball  $B$  of  $X$ . The characteristic function of a set  $E \subset X$  is denoted by  $\mathbf{1}_E$ ; that is,  $\mathbf{1}_E(x) = 1$  if  $x \in E$  and  $\mathbf{1}_E(x) = 0$  if  $x \in X \setminus E$ .

### 3. MUCKENHOUPT WEIGHTS

For convenience, we recall the standard definition and elementary properties of Muckenhoupt weights in the setting of metric measure spaces. We follow [4] and also refer to [8].

**Definition 3.1.** A measurable function  $w: X \rightarrow \mathbb{R}$  is a weight in  $X$ , if  $w$  is integrable on all balls of  $X$  and  $w(x) > 0$  for almost every  $x \in X$ . If  $w$  is a weight in  $X$  and  $E \subset X$  is a measurable set, we write

$$w(E) = \int_E w(x) d\mu(x). \quad (3.2)$$

We say that a weight  $w$  in  $X$  is doubling, if there exists a constant  $c_w \geq 1$  such that

$$0 < w(B(x, 2r)) \leq c_w w(B(x, r)) < \infty$$

for all  $x \in X$  and  $r > 0$ .

**Definition 3.3.** A weight  $w$  in  $X$  belongs to the Muckenhoupt class  $A_p$ , for  $1 < p < \infty$ , if there exists a constant  $C$  such that

$$\int_B w(x) d\mu(x) \left( \int_B w(x)^{\frac{1}{1-p}} d\mu(x) \right)^{p-1} \leq C \quad (3.4)$$

for every ball  $B \subset X$ . The smallest possible constant  $C$  in (3.4) is called the  $A_p$  constant of  $w$ , and it is denoted by  $[w]_{A_p}$ .

If  $1 < p < q < \infty$ , then Hölder's inequality shows that  $A_p \subset A_q$ . Next we recall some other basic properties of Muckenhoupt weights that will be applied later. We begin with a duality statement.

**Lemma 3.5.** *Let  $1 < p < \infty$  and write  $p' = \frac{p}{p-1}$ . Then  $w \in A_p$  if and only if  $w^{1-p'} \in A_{p'}$ .*

*Proof.* Assume that  $w \in A_p$ . Then

$$\begin{aligned} & \int_B w(x)^{1-p'} d\mu(x) \left( \int_B (w(x)^{1-p'})^{\frac{1}{1-p'}} d\mu(x) \right)^{p'-1} \\ &= \int_B w(x)^{1-p'} d\mu(x) \left( \int_B w(x) d\mu(x) \right)^{\frac{1}{p-1}} \leq [w]_{A_p}^{\frac{1}{p-1}} \end{aligned}$$

for every ball  $B \subset X$ . This shows that  $w^{1-p'} \in A_{p'}$ . Conversely, if  $w^{1-p'} \in A_{p'}$  then  $w = (w^{1-p'})^{1-p} \in A_p$  by the same argument.  $\square$

Next we consider a weighted Hölder inequality for integral averages.

**Lemma 3.6.** *Let  $1 < p < \infty$  and assume that  $w \in A_p$ . Let  $B \subset X$  be a ball and  $f \in L^1(B)$  be a nonnegative function. Then*

$$\int_B f(x) d\mu(x) \leq [w]_{A_p}^{\frac{1}{p}} \left( \frac{1}{w(B)} \int_B f(x)^p w(x) d\mu(x) \right)^{\frac{1}{p}}.$$

*Proof.* Hölder's inequality and (3.4) imply

$$\begin{aligned} \int_B f(x) d\mu(x) &\leq \frac{1}{\mu(B)} \left( \int_B w(x)^{\frac{1}{1-p}} d\mu(x) \right)^{\frac{p-1}{p}} \left( \int_B f(x)^p w(x) d\mu(x) \right)^{\frac{1}{p}} \\ &= \left( \frac{w(B)}{\mu(B)} \right)^{\frac{1}{p}} \left( \int_B w(x)^{\frac{1}{1-p}} d\mu(x) \right)^{\frac{p-1}{p}} \left( \frac{1}{w(B)} \int_B f(x)^p w(x) d\mu(x) \right)^{\frac{1}{p}} \\ &\leq [w]_{A_p}^{\frac{1}{p}} \left( \frac{1}{w(B)} \int_B f(x)^p w(x) d\mu(x) \right)^{\frac{1}{p}}. \quad \square \end{aligned}$$

Since the underlying measure  $\mu$  is doubling, an application of Lemma 3.6 shows that a Muckenhoupt weight in  $X$  is doubling.

**Lemma 3.7.** *Let  $1 < p < \infty$  and assume that  $w \in A_p$ . Then  $w$  is doubling with a constant  $c_w = C(c_\mu, p, [w]_{A_p})$ .*

*Proof.* Let  $x \in X$  and  $r > 0$ . The doubling condition in (2.1) and Lemma 3.6 with  $f = \mathbf{1}_{B(x,r)}$  give

$$\begin{aligned} \frac{1}{c_\mu} &\leq \frac{\mu(B(x,r))}{\mu(B(x,2r))} = \int_{B(x,2r)} f(y) d\mu(y) \\ &\leq [w]_{A_p}^{\frac{1}{p}} \left( \frac{1}{w(B(x,2r))} \int_{B(x,2r)} f(y)^p w(y) d\mu(y) \right)^{\frac{1}{p}} \\ &\leq [w]_{A_p}^{\frac{1}{p}} \left( \frac{w(B(x,r))}{w(B(x,2r))} \right)^{\frac{1}{p}}. \end{aligned}$$

The claim follows by raising both sides to power  $p$  and reorganizing the terms.  $\square$

#### 4. MAXIMAL FUNCTION ESTIMATES

Let  $w$  be a weight in  $X$ . For  $1 \leq p < \infty$ , the weighted  $L^p$ -norm of a measurable function  $f$  on  $X$  is defined as

$$\|f\|_{L^p(w d\mu)} = \left( \int_X |f(x)|^p w(x) d\mu(x) \right)^{\frac{1}{p}},$$

and for  $p = \infty$  we define

$$\|f\|_{L^\infty(w d\mu)} = \inf \{ c \geq 0 : |f(x)| \leq c \text{ } w\text{-a.e. } x \in X \}.$$

If  $\|f\|_{L^p(w d\mu)} < \infty$ , we write  $f \in L^p(w d\mu)$ , and in the case  $w = 1$  we use the notation  $L^p(d\mu) = L^p(X)$ .

**Definition 4.1.** Let  $w$  be a doubling weight in  $X$ . For a measurable function  $f$  on  $X$ , the weighted maximal function  $M^{*,w}f$  is defined by

$$M^{*,w}f(x) = \sup_{B \ni x} \frac{1}{w(B)} \int_B |f(y)| w(y) d\mu(y),$$

where the supremum is taken over all balls  $B$  in  $X$  with  $x \in B$ . When  $w = 1$ , we obtain the noncentered maximal function  $M^*f = M^{*,1}f$ .

The following simple lemma gives a sufficient condition for the Muckenhoupt property. Observe that the class of test functions is restricted to essentially bounded functions with a bounded support. This restriction is convenient for later purposes.

**Lemma 4.2.** *Let  $1 < p < \infty$  and assume that  $w$  is a weight in  $X$ . Assume that there exists a constant  $C_1$  such that*

$$\int_X (M^* f(x))^p w(x) d\mu(x) \leq C_1 \int_X f(x)^p w(x) d\mu(x) \quad (4.3)$$

for every nonnegative  $f \in L^\infty(X)$  with a bounded support. Then  $w \in A_p$  and  $[w]_{A_p} \leq C_1$ .

*Proof.* Fix a ball  $B \subset X$ . Let  $\varepsilon > 0$  and  $f = (\varepsilon + w)^{\frac{1}{1-p}} \mathbf{1}_B$ . Observe that  $f \in L^\infty(X)$  and  $f$  has a bounded support. Clearly

$$\int_B (\varepsilon + w(y))^{\frac{1}{1-p}} d\mu(y) = \int_B |f(y)| d\mu(y) \leq M^* f(x)$$

for every  $x \in B$ , and thus

$$\int_B w(x) d\mu(x) \left( \int_B (\varepsilon + w(y))^{\frac{1}{1-p}} d\mu(y) \right)^p \leq \int_B (M^* f(x))^p w(x) d\mu(x).$$

By the assumption, we have

$$\begin{aligned} \int_B (M^* f(x))^p w(x) d\mu(x) &\leq \frac{1}{\mu(B)} \int_X (M^* f(x))^p w(x) d\mu(x) \\ &\leq C_1 \frac{1}{\mu(B)} \int_X f(x)^p w(x) d\mu(x) \\ &\leq C_1 \int_B (\varepsilon + w(x))^{\frac{p}{1-p}} (\varepsilon + w(x)) d\mu(x) \\ &= C_1 \int_B (\varepsilon + w(x))^{\frac{1}{1-p}} d\mu(x). \end{aligned}$$

This gives

$$\int_B w(x) d\mu(x) \left( \int_B (\varepsilon + w(y))^{\frac{1}{1-p}} d\mu(y) \right)^{p-1} \leq C_1$$

for every  $\varepsilon > 0$ . Fatou's lemma implies

$$\begin{aligned} \int_B w(x) d\mu(x) \left( \int_B w(x)^{\frac{1}{1-p}} d\mu(x) \right)^{p-1} \\ \leq \liminf_{\varepsilon \rightarrow 0^+} \int_B w(x) d\mu(x) \left( \int_B (\varepsilon + w(x))^{\frac{1}{1-p}} d\mu(x) \right)^{p-1} \leq C_1, \end{aligned}$$

and the claim follows.  $\square$

Next we aim to show a strengthened converse of Lemma 4.2, see Theorem 4.5 below. Since a doubling weight  $w$  in  $X$  can be interpreted as a doubling measure as in Definition 3.1, we have the following weighted version of the Hardy–Littlewood maximal function theorem, see for instance [1, Theorem 3.13].

**Lemma 4.4.** *Let  $1 < p < \infty$  and let  $w$  be a doubling weight in  $X$ , with a constant  $c_w$ . Then there exists a constant  $C = C(p, c_w)$  such that*

$$\|M^{*,w} f\|_{L^p(w d\mu)} \leq C \|f\|_{L^p(w d\mu)}$$

for every  $f \in L^p(w d\mu)$ .

The proof of the weighted norm inequality in Theorem 4.5 is adapted from [10]. We also refer to [4, Theorem 8.19].

**Theorem 4.5.** *Let  $1 < p < \infty$  and  $w \in A_p$ . There exists a constant  $C = C(c_\mu, p, [w]_{A_p})$  such that*

$$\int_X (M^* f(x))^p w(x) d\mu(x) \leq C \int_X |f(x)|^p w(x) d\mu(x)$$

for every  $f \in L^p(w d\mu)$ .

*Proof.* Define  $\sigma = w^{1-p'} = w^{\frac{1}{1-p}}$ . Lemma 3.5 implies that  $\sigma \in A_{p'}$ , where  $p' = \frac{p}{p-1}$ . Let  $f \in L^p(w d\mu)$ . Let  $x \in X$  and consider a ball  $B = B(x_0, r)$  such that  $x \in B$ . Then

$$\int_B |f(y)| d\mu(y) = A \left( \frac{\mu(B)}{w(B)} \left( \frac{1}{\sigma(B)} \int_B |f(y)| d\mu(y) \right)^{p-1} \right)^{\frac{1}{p-1}},$$

where

$$A = \left( \frac{w(B)\sigma(B)^{p-1}}{\mu(B)^p} \right)^{\frac{1}{p-1}} \leq [w]_{A_p}^{\frac{1}{p-1}}.$$

Let  $z \in B$ . Then

$$\frac{1}{\sigma(B)} \int_B |f(y)| d\mu(y) = \frac{1}{\sigma(B)} \int_B |f(y)\sigma(y)^{-1}| \sigma(y) d\mu(y) \leq M^{*,\sigma}(f\sigma^{-1})(z).$$

Hence we have

$$\begin{aligned} \int_B |f(y)| d\mu(y) &\leq [w]_{A_p}^{\frac{1}{p-1}} \left( \frac{1}{w(B)} \int_B (M^{*,\sigma}(f\sigma^{-1})(z))^{p-1} w(z)^{-1} w(z) d\mu(z) \right)^{\frac{1}{p-1}} \\ &\leq [w]_{A_p}^{\frac{1}{p-1}} (M^{*,w}((M^{*,\sigma}(f\sigma^{-1}))^{p-1} w^{-1})(x))^{\frac{1}{p-1}}. \end{aligned}$$

By taking supremum over all balls  $B$  as above, it follows that

$$M^* f(x) \leq [w]_{A_p}^{\frac{1}{p-1}} (M^{*,w}((M^{*,\sigma}(f\sigma^{-1}))^{p-1} w^{-1})(x))^{\frac{1}{p-1}}$$

for all  $x \in X$ . Observe that  $w \in A_p$  and  $\sigma \in A_{p'}$  are doubling weights by Lemma 3.7, with constants  $c_w$  and  $c_\sigma$  depending only on  $c_\mu$ ,  $p$  and  $[w]_{A_p}$ . Lemma 4.4 implies

$$\begin{aligned} \int_X (M^* f(x))^p w(x) d\mu(x) &\leq [w]_{A_p}^{\frac{p}{p-1}} \int_X (M^{*,w}((M^{*,\sigma}(f\sigma^{-1}))^{p-1} w^{-1})(x))^{\frac{p}{p-1}} w(x) d\mu(x) \\ &\leq C(c_\mu, p, [w]_{A_p}) \int_X (M^{*,\sigma}(f\sigma^{-1})(x))^p \sigma(x) d\mu(x) \\ &\leq C(c_\mu, p, [w]_{A_p}) \int_X |f(x)|^p w(x) d\mu(x), \end{aligned}$$

where the facts  $\sigma = w^{\frac{1}{1-p}}$  and  $w = \sigma^{1-p}$  are used.  $\square$

## 5. SELF-IMPROVEMENT RESULTS

We need the following Whitney type covering theorem, see [6, Proposition 4.1.15] and [2].

**Lemma 5.1.** *Assume that  $\Omega \neq \emptyset$  is an open subset of  $X$  such that  $X \setminus \Omega \neq \emptyset$ . There exist balls  $B_i = B(x_i, r_i) \subset \Omega$ ,  $i \in \mathbb{N}$ , with*

$$r_i = \frac{1}{8} \text{dist}(x_i, X \setminus \Omega), \quad (5.2)$$

such that  $\Omega = \bigcup_{i=1}^{\infty} B_i$  and

$$\sum_{i=1}^{\infty} \mathbf{1}_{B_i}(x) \leq C(c_\mu) \quad (5.3)$$

for every  $x \in \Omega$ .

Let  $f \in L^\infty(X)$  be a nonnegative function with a bounded support. Let  $t > 0$  be such that

$$E_t = \{x \in X : M^*f(x) > t\} \neq X,$$

where  $M^*f$  is the noncentered maximal function of  $f$  as in Definition 4.1. Then  $E_t \subsetneq X$  is an open set. We define a function  $f_t$  as below.

- If  $E_t \neq \emptyset$ , consider a Whitney type cover  $\{B_i : i \in \mathbb{N}\}$  of  $E_t$  as in Lemma 5.1, and let

$$f_t(x) = \begin{cases} f(x), & x \in X \setminus E_t, \\ \sup_{B_i \ni x} \int_{B_i} f(y) d\mu(y), & x \in E_t, \end{cases} \quad (5.4)$$

where the supremum is taken over all balls  $B_i$ ,  $i \in \mathbb{N}$ , such that  $x \in B_i$ .

- If  $E_t = \emptyset$ , let  $f_t(x) = f(x)$  for every  $x \in X$ .

**Lemma 5.5.** *Assume that  $f \in L^\infty(X)$  is a nonnegative function with a bounded support. Let  $t > 0$  and let  $E_t \subsetneq X$  and  $f_t$  be as defined above. Then the following assertions hold:*

- $f_t(x) \leq C(c_\mu)t$  for almost every  $x \in X$ .
- $M^*f(x) \leq C(c_\mu)M^*f_t(x)$  for every  $x \in X \setminus E_t$ .

*Proof.* By the Lebesgue differentiation theorem [6, Section 3.4], we have

$$f_t(x) = f(x) \leq M^*f(x) \leq t$$

for almost every  $x \in X \setminus E_t$ . If  $E_t = \emptyset$ , then (a) holds. If  $E_t \neq \emptyset$  and  $x \in E_t$ , then consider any ball  $B_i$ ,  $i \in \mathbb{N}$ , such that  $x \in B_i$ . Recall that  $E_t$  is the union of such balls. By (5.2) and (2.1), there exists a ball  $B \subset X$  such that  $B \setminus E_t \neq \emptyset$ ,  $B_i \subset B$  and  $\mu(B) \leq c_\mu^4 \mu(B_i)$ . For any  $z \in B \setminus E_t$  we have

$$\int_{B_i} f(y) d\mu(y) \leq c_\mu^4 \int_B f(y) d\mu(y) \leq c_\mu^4 M^*f(z) \leq c_\mu^4 t.$$

By taking supremum over all balls  $B_i$  as above, we see that  $f_t(x) \leq c_\mu^4 t$  for every  $x \in E_t$ , and thus (a) holds also in the case  $E_t \neq \emptyset$ .

To prove part (b), we may clearly assume that  $E_t \neq \emptyset$ . Let  $x \in X \setminus E_t$ . Let  $z \in X$  and  $r > 0$  be such that  $x \in B(z, r)$ . Then

$$\begin{aligned} \int_{B(z,r)} f(y) d\mu(y) &= \int_{B(z,r) \setminus E_t} f(y) d\mu(y) + \int_{B(z,r) \cap E_t} f(y) d\mu(y) \\ &\leq \int_{B(z,r) \setminus E_t} f(y) d\mu(y) + \sum_{i=1}^{\infty} \int_{B(z,r) \cap B_i} f(y) d\mu(y). \end{aligned}$$

Using the fact that  $0 \leq f = f_t$  in  $X \setminus E_t$ , we get

$$\int_{B(z,r) \setminus E_t} f(y) d\mu(y) \leq \int_{B(z,2r) \setminus E_t} f(y) d\mu(y) = \int_{B(z,2r) \setminus E_t} f_t(y) d\mu(y).$$

Next we consider any ball  $B_i$  satisfying  $B(z,r) \cap B_i \neq \emptyset$ , if such  $B_i$  exists. Since  $x \in B(z,r) \setminus E_t$ , by definition (5.2) of the radius  $r_i$ , we have  $B_i \subset B(z,2r)$ . Thus (5.4) implies

$$\begin{aligned} \int_{B(z,r) \cap B_i} f(y) d\mu(y) &\leq \int_{B_i} f(y) d\mu(y) = \int_{B_i} \int_{B_i} f(y) d\mu(y) d\mu(z) \\ &\leq \int_{B_i} f_t(z) d\mu(z) = \int_{B(z,2r) \cap B_i} f_t(y) d\mu(y). \end{aligned}$$

By collecting the estimates above and using (5.3), we obtain

$$\begin{aligned} \int_{B(z,r)} f(y) d\mu(y) &\leq \int_{B(z,2r) \setminus E_t} f_t(y) d\mu(y) + \sum_{i=1}^{\infty} \int_{B(z,2r) \cap B_i} f_t(y) d\mu(y) \\ &\leq C(c_\mu) \int_{B(z,2r)} f_t(y) d\mu(y), \end{aligned}$$

and consequently

$$\int_{B(z,r)} f(y) d\mu(y) \leq C(c_\mu) \int_{B(z,2r)} f_t(y) d\mu(y) \leq C(c_\mu) M^* f_t(x).$$

By taking supremum over all balls  $B(z, r)$  as above, we conclude that (b) holds.  $\square$

Next we state and prove our main results concerning the self-improving properties of weighted norm inequalities for the non-centered maximal function. To emphasize the key aspects of the arguments, we first formulate this property only in terms of essentially bounded functions with a bounded support. We will later strengthen this result and recover it for all  $f \in L^{p-\varepsilon}(w d\mu)$ , see Corollary 5.13.

**Lemma 5.6.** *Let  $1 < p < \infty$  and let  $w$  be a weight in  $X$ . Assume that there exists a constant  $C_1$  such that*

$$\int_X (M^* f(x))^p w(x) d\mu(x) \leq C_1 \int_X |f(x)|^p w(x) d\mu(x) \quad (5.7)$$

for all  $f \in L^p(w d\mu)$ . Then there exist constants  $\varepsilon = \varepsilon(c_\mu, p, C_1)$  and  $C = C(c_\mu, p, C_1)$  such that  $0 < \varepsilon < p - 1$  and

$$\int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \leq C \int_X f(x)^{p-\varepsilon} w(x) d\mu(x)$$

for all nonnegative  $f \in L^\infty(X)$  with a bounded support.

*Proof.* Fix a nonnegative function  $f \in L^\infty(X)$  with a bounded support. Let  $t > 0$  be such that  $E_t = \{x \in X : M^* f(x) > t\} \subsetneq X$  and let  $f_t$  be as in (5.4). By using assumption (5.7), we get

$$\int_X (M^* f_t(x))^p w(x) d\mu(x) \leq C_1 \int_X f_t(x)^p w(x) d\mu(x).$$

Employing also Lemma 5.5, we obtain

$$\begin{aligned} \int_{X \setminus E_t} (M^* f(x))^p w(x) d\mu(x) &\leq C(c_\mu, p) \int_{X \setminus E_t} (M^* f_t(x))^p w(x) d\mu(x) \\ &\leq C(c_\mu, p) \int_X (M^* f_t(x))^p w(x) d\mu(x) \\ &\leq C(c_\mu, p, C_1) \left( \int_{X \setminus E_t} f(x)^p w(x) d\mu(x) + t^p w(E_t) \right). \end{aligned} \quad (5.8)$$

We consider parameters  $t_0 > 0$  and  $0 < \varepsilon < p - 1$ . We will choose the latter parameter more specifically at the end of the proof. We have

$$\begin{aligned} &\int_{t_0}^{\infty} t^{-1-\varepsilon} \int_{X \setminus E_t} (M^* f(x))^p w(x) d\mu(x) dt \\ &\leq C(c_\mu, p, C_1) \left( \int_{t_0}^{\infty} t^{-1-\varepsilon} \int_{X \setminus E_t} f(x)^p w(x) d\mu(x) dt + \int_{t_0}^{\infty} t^{p-1-\varepsilon} w(E_t) dt \right). \end{aligned} \quad (5.9)$$



By Fubini's theorem,

$$\begin{aligned}
 & \int_{t_0}^{\infty} t^{-1-\varepsilon} \int_{X \setminus E_t} (M^* f(x))^p w(x) d\mu(x) dt \\
 &= \int_X (M^* f(x))^p \left( \int_{t_0}^{\infty} \mathbf{1}_{X \setminus E_t}(x) t^{-1-\varepsilon} dt \right) w(x) d\mu(x) \\
 &= \int_X (M^* f(x))^p \left( \int_{\max\{t_0, M^* f(x)\}}^{\infty} t^{-1-\varepsilon} dt \right) w(x) d\mu(x) \\
 &= \frac{1}{\varepsilon} \int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x).
 \end{aligned}$$

In a similar way, we obtain

$$\begin{aligned}
 \int_{t_0}^{\infty} t^{-1-\varepsilon} \int_{X \setminus E_t} f(x)^p w(x) d\mu(x) dt &= \frac{1}{\varepsilon} \int_X f(x)^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x) \\
 &\leq \frac{1}{\varepsilon} \int_X f(x)^{p-\varepsilon} w(x) d\mu(x),
 \end{aligned}$$

where the inequality holds since  $0 \leq f(x) \leq \max\{t_0, M^* f(x)\}$  for almost every  $x \in X$ .

The last term in (5.9) can be estimated as

$$\begin{aligned}
 \int_{t_0}^{\infty} t^{p-1-\varepsilon} w(E_t) dt &= \int_X \left( \int_{t_0}^{\infty} \mathbf{1}_{E_t}(x) t^{p-1-\varepsilon} dt \right) w(x) d\mu(x) \\
 &= \int_{E_{t_0}} \left( \int_{t_0}^{M^* f(x)} t^{p-1-\varepsilon} dt \right) w(x) d\mu(x) \\
 &\leq \frac{1}{p-\varepsilon} \int_{E_{t_0}} (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \\
 &\leq \frac{1}{p-\varepsilon} \int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x).
 \end{aligned}$$

Multiplying of the obtained inequalities by  $\varepsilon > 0$  and using (5.9) gives

$$\begin{aligned}
 \int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x) &\leq C(c_\mu, p, C_1) \int_X f(x)^{p-\varepsilon} w(x) d\mu(x) \\
 + C(c_\mu, p, C_1) \frac{\varepsilon}{p-\varepsilon} \int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x).
 \end{aligned} \tag{5.10}$$

We fix  $0 < \varepsilon < p - 1$ , only depending on  $c_\mu$ ,  $p$  and  $C_1$ , such that

$$C(c_\mu, p, C_1) \frac{\varepsilon}{p-\varepsilon} \leq \frac{1}{2}.$$

Recall that  $f \in L^\infty(X)$  has a bounded support and  $w$  is integrable on balls. Thus the last term on the right-hand side of (5.10) is finite by the assumption in (5.7) and the fact that  $t_0 > 0$ . Hence, we may absorb this term to the left-hand side of (5.10), and we obtain the estimate

$$\int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x) \leq C(c_\mu, p, C_1) \int_X f(x)^{p-\varepsilon} w(x) d\mu(x)$$

that holds uniformly for all  $t_0 > 0$ . Finally, by Fatou's lemma,

$$\begin{aligned}
 \int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) &\leq \liminf_{t_0 \rightarrow 0^+} \int_X (M^* f(x))^p (\max\{t_0, M^* f(x)\})^{-\varepsilon} w(x) d\mu(x) \\
 &\leq C(c_\mu, p, C_1) \int_X f(x)^{p-\varepsilon} w(x) d\mu(x),
 \end{aligned}$$

and this concludes the proof.  $\square$

We can now prove a self-improving property of Muckenhoupt weights.

**Theorem 5.11.** *Let  $1 < p < \infty$  and assume that  $w \in A_p$ . Then there exist  $\varepsilon = \varepsilon(c_\mu, p, [w]_{A_p})$  such that  $0 < \varepsilon < p - 1$  and  $w \in A_{p-\varepsilon}$  and  $[w]_{A_{p-\varepsilon}} \leq C(c_\mu, p, [w]_{A_p})$ . Moreover,*

$$\int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \leq C(c_\mu, p, [w]_{A_p}) \int_X |f(x)|^{p-\varepsilon} w(x) d\mu(x) \quad (5.12)$$

for all  $f \in L^{p-\varepsilon}(w d\mu)$ .

*Proof.* Theorem 4.5 and Lemma 5.6 combined show that there are constants  $\varepsilon = \varepsilon(c_\mu, p, [w]_{A_p})$  and  $C_1 = C(c_\mu, p, [w]_{A_p})$  such that  $0 < \varepsilon < p - 1$  and

$$\int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \leq C_1 \int_X f(x)^{p-\varepsilon} w(x) d\mu(x)$$

for all nonnegative  $f \in L^\infty(X)$  with a bounded support. Lemma 4.2 implies that  $w \in A_{p-\varepsilon}$  and  $[w]_{A_{p-\varepsilon}} \leq C_1 = C(c_\mu, p, [w]_{A_p})$ . Finally, the inequality in (5.12) holds for all  $f \in L^{p-\varepsilon}(w d\mu)$  by Theorem 4.5.  $\square$

By combining the results in Lemma 4.2 and Theorem 5.11, we obtain the following stronger variant of Lemma 5.6, in which the improved version of the weighted norm inequality holds for all functions  $f \in L^{p-\varepsilon}(w d\mu)$ .

**Corollary 5.13.** *Let  $1 < p < \infty$  and let  $w$  be a weight in  $X$ . Assume that there exists a constant  $C_1$  such that*

$$\int_X (M^* f(x))^p w(x) d\mu(x) \leq C_1 \int_X f(x)^p w(x) d\mu(x)$$

for all nonnegative functions  $f \in L^\infty(X)$  with a bounded support. Then there exist constants  $\varepsilon = \varepsilon(c_\mu, p, C_1)$  and  $C = C(c_\mu, p, C_1)$  such that  $0 < \varepsilon < p - 1$  and

$$\int_X (M^* f(x))^{p-\varepsilon} w(x) d\mu(x) \leq C \int_X |f(x)|^{p-\varepsilon} w(x) d\mu(x)$$

for all  $f \in L^{p-\varepsilon}(w d\mu)$ .

## REFERENCES

- [1] A. Björn and J. Björn. *Nonlinear potential theory on metric spaces*, volume 17 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2011.
- [2] R. R. Coifman and G. Weiss. *Analyse harmonique non-commutative sur certains espaces homogènes*, volume Vol. 242 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1971. Étude de certaines intégrales singulières.
- [3] D. V. Cruz-Uribe, J. M. Martell, and C. Pérez. *Weights, extrapolation and the theory of Rubio de Francia*, volume 215 of *Operator Theory: Advances and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [4] J. García-Cuerva and J. L. Rubio de Francia. *Weighted norm inequalities and related topics*, volume 116 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1985. Notas de Matemática [Mathematical Notes], 104.
- [5] I. Genebashvili, A. Gogatishvili, V. Kokilashvili, and M. Krbec. *Weight theory for integral transforms on spaces of homogeneous type*, volume 92 of *Pitman Monographs and Surveys in Pure and Applied Mathematics*. Longman, Harlow, 1998.
- [6] J. Heinonen, P. Koskela, N. Shanmugalingam, and J. T. Tyson. *Sobolev spaces on metric measure spaces*, volume 27 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2015. An approach based on upper gradients.
- [7] S. Keith and X. Zhong. The Poincaré inequality is an open ended condition. *Ann. of Math. (2)*, 167(2):575–599, 2008.
- [8] J. Kinnunen, J. Lehrbäck, and A. Vähäkangas. *Maximal function methods for Sobolev spaces*, volume 257 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2021.

- [9] J. Kinnunen, J. Lehrbäck, A. V. Vähäkangas, and X. Zhong. Maximal function estimates and self-improvement results for Poincaré inequalities. *Manuscripta Math.*, 158(1-2):119–147, 2019.
- [10] A. K. Lerner. An elementary approach to several results on the Hardy-Littlewood maximal operator. *Proc. Amer. Math. Soc.*, 136(8):2829–2833, 2008.
- [11] J. L. Lewis. On very weak solutions of certain elliptic systems. *Comm. Partial Differential Equations*, 18(9-10):1515–1537, 1993.
- [12] J.-O. Strömberg and A. Torchinsky. *Weighted Hardy spaces*, volume 1381 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989.

(J.K.) DEPARTMENT OF MATHEMATICS, AALTO UNIVERSITY, P.O. BOX 11100, FI-00076 AALTO UNIVERSITY, FINLAND

*Email address:* juha.k.kinnunen@aalto.fi

(J.L.) UNIVERSITY OF JYVASKYLA, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35, FI-40014 UNIVERSITY OF JYVASKYLA, FINLAND

*Email address:* juha.lehrback@jyu.fi

(A.V.V.) UNIVERSITY OF JYVASKYLA, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35, FI-40014 UNIVERSITY OF JYVASKYLA, FINLAND

*Email address:* antti.vahakangas@iki.fi

(D.Y.) LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS (MINISTRY OF EDUCATION OF CHINA), SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, BEIJING 100875, THE PEOPLE'S REPUBLIC OF CHINA

*Email address:* dcyang@bnu.edu.cn