Unifying Inflation, dark energy and dark matter with a scalar field and exotic fermions

Eduardo Guendelman^{1, 2, 3, *} and Ramón Herrera^{4, †}

¹Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva, Israel. ²Frankfurt Institute for Advanced Studies (FIAS),

Ruth-Moufang-Strasse 1, 60438 Frankfurt am Main, Germany.

³Bahamas Advanced Study Institute and Conferences, 4A Ocean Heights,

Hill View Circle, Stella Maris, Long Island, The Bahamas.

⁴Instituto de Física, Pontificia Universidad Católica de Valparaíso,

Avenida Brasil 2950, Casilla 4059, Valparaíso, Chile.

In this paper we consider a new approach to unify inflation and the late universe with dark energy and dark matter formulated in a model that includes a non-Riemannian metric independent measure and a scalar field with spontaneously broken scale symmetry. Here first of all inflation is possible, which is then followed by a reheating oscillating period and this leads to the formation of all kind of particles, including fermions, which as the universe expands can contribute to the dark energy and the to the dark matter of the universe. During the inflationary epoch, we find different constraints on the parameter space associated to the effective potential of the scalar field from the observational data. After reheating the scalar field retraces its trajectory in field space but now the scalar field potential can be drastically modified by the effect of the fermions. In this sense, the present dark energy with its very small value in comparison to the inflationary phase which can be adjusted by choosing appropriately the parameter space of couplings of the Riemannian and non Riemannian measures to the fermions.

I. INTRODUCTION

The accelerated expansion of the present universe [1] revived the idea of Einstein of introducing a cosmological constant term, or of Dark Energy (DE) of the late universe, for reviews of DE, from the theory point of view see [2]. One of the interesting aspects that have been pointed out is the coincidence, at least in orders of magnitude between Dark Energy and Dark Matter (DM) [3], in fact in this paper we will find that DE and DM may have a common origin. One approach to DE is to assume that the accelerated expansion is driven by a scalar field with a potential rather than just a cosmological constant [4], in our case, a scalar field will play a role, but fermionic matter will also be important.

Of course a much larger cosmological constant is thought to be present in the early universe, driving an even much faster accelerated expansion of the universe, the inflationary phase [5], which solves many puzzles and is though to be also the source of the primordial perturbations. The inflation then ends at some point and this is then followed by particle creation which produces all the matter in the universe, after 13 billion years we have then again a universe dominated by something the cosmological constant mentioned at the beginning which is much smaller than the inflationary cosmological constant.

On the other hand, sterile neutrinos can be DM and these particles are the basis for some models as also for the generation of DE[6]. An idea has been suggested in Refs.[7, 8] in which a coupling of the neutrinos with a scalar field produces a dark energy density that depends on the neutrino mass. In this approximation the neutrino mass is related to the density of the background nonrelativistic neutrinos in a power-law relation. We mention that in the model described in Ref. [7], the energy density associated to the neutrinos is small in relation to the energy density in the total dark energy sector.

The generation of dark energy by fermions is studied in the context of the Two Measures Theory, where the traditional measure of integration $\sqrt{-g}$ is used in some parts of the Lagrangian of the theory, but is replaced by a metric independent measure Φ in other parts of the Lagrangian [9], [10], [11]. then to this idea, the requirement of scale invariance was added im [12], where the spontaneous scale symmetry was was generated by the integration of the degrees of freedom defining the modified measure. Scale invariance modified measures models for Braneworlds were studied in Ref.[13].

The subject of unifying the early inflation with the late DE was first studied in the Two Measure Theory in [14] by introducing curvature square terms, then this approach was used to obtain emergent universe solutions that evolve

^{*} guendel@bgu.ac.il

[†] ramon.herrera@pucv.cl

from an emergent non singular phase to inflation and then to the present DE phase, [15], and then have shown that similar effect can take place even without curvature square terms [16] by showing that the two measures theory can induce K essence terms when considering the Einstein frame, that can also support the emergent solutions and then combined effects of the induced K essence terms and curvature square terms to obtain an Emergent Cosmology, Inflation and Dark Energy were considered in [17]. The models [14], [15], [16] and [17], are non oscillating ones, so to account for reheating after inflation, a curvaton field must be added, this has been done in [18], [19] These kind of models provide also the possibility to incorporate inflation, early dark energy and late dark energy, which can be of use in the resolution of the H_0 tension [20], which was then generalized [21] to account for the non singular origin of the universe, the H_0 problem and the DM as originating from the scalar field due to its K essence behavior.

In a previous publication, dark energy generation from fermions was considered in the Two Measure Theory[22], but inflation was not studied, so it was not a full description of the full history of the universe. Furthermore, the model studied here not only accounts for both inflation and dark energy, but also the effective potential has an oscillating region that allows particle creation and reheating, which allows for the creation of the fermions themselves that will ultimately lead to a modification of the effective potential, which in the same region in scalar field space where inflation took place, now provides an energy density of the order of $(10^{-3}ev)^4$ corresponding to the DE for the late universe, while the fermions provide the DM as well.

In this article, we want to describe the unification of the early and present universe. To describe the early universe we assume an inflationary stage dominates by a scalar field from an effective potential without the present of fermions. Besides, we want to explore the possibility that the matter produced in the reheating after inflation can have a role in the modification of the cosmological constant of the early universe. We explore in particular the possibility that fermions, like neutrinos, or sterile neutrinos could modify the original inflaton field potential generating a new effective potential, and this same inflaton field could then drive the late universe expansion. In addition, we can account also for the dark matter of the present universe from these fermions assuming that this matter is also produced during the reheating mechanism.

We organize our paper as follows: In Section II, we provide a brief analysis of our model. In this section, we also present the basic equations in the Einstein frame associated with the scalar field and primordial fermions. In Section III, we study the inflationary stage of our model. Under the slow-roll approximation, we derive the observational parameters, such as the scalar spectral index, the power spectrum, and the tensor-to-scalar ratio. From these observables, we obtain different constraints on the parameter space of our model. Additionally, in Sections IV and V, we analyze dark energy and dark matter from fermions. Here, we show that it is possible to obtain fermions associated with DE and DM from different energy-momentum tensors. In Section VI, we present our conclusions. Finally, we include Appendix A for a discussion of the spin connection and the Einstein frame. We chose units in which $c = \hbar = 1$.

II. GENERAL DESCRIPTION

In this section, we will present a brief analysis of the implications of considering both the scalar field and fermion fields. Thus, in order to describe our model, we consider that the general action can be written as [22]

$$S = \int d^4x e^{\alpha\phi/M_p} (\Phi + b\sqrt{-g}) \left[-\frac{1}{\kappa} R(\omega, e) + \frac{1}{2} g^{\mu\nu} \phi_{,\mu} \phi_{,\nu} \right] - \int d^4x e^{2\alpha\phi/M_p} [\Phi V_1 + \sqrt{-g} V_2] - \int d^4x \sqrt{-g} \frac{1}{4} g^{\alpha\beta} g^{\mu\nu} F_{\alpha\mu} F_{\beta\nu} + \int d^4x e^{\alpha\phi/M_p} (\Phi + k\sqrt{-g}) \frac{i}{2} \sum_i \overline{\Psi}_i \left(\gamma^a e^{\mu}_a \overrightarrow{\nabla}^{(i)}_{\mu} - \overleftarrow{\nabla}^{(i)}_{\mu} \gamma^a e^{\mu}_a \right) \Psi_i - \int d^4x e^{\frac{3}{2}\alpha\phi/M_p} \left[(\Phi + h_E \sqrt{-g}) \mu_E \overline{E} E + (\Phi + h_n \sqrt{-g}) \mu_n \overline{n} n \right],$$
(1)

in which the measure fields φ_a are utilized for definition of the measure $\Phi = \varepsilon^{\mu\nu\alpha\beta}\varepsilon_{abcd}\partial_{\mu}\varphi_{a}\partial_{\nu}\varphi_{b}\partial_{\alpha}\varphi_{c}\partial_{\beta}\varphi_{d}$. Therefore, the quantity Ψ_i (i = n, E neutrinos, electrons) is the general notation for the primordial fermion fields n and E; the variable $F_{\alpha\beta}$ is defined as $F_{\alpha\beta} = \partial_{\alpha}A_{\beta} - \partial_{\beta}A_{\alpha}$ and the parameters μ_n and μ_E are the mass parameters. Besides, we have $\overrightarrow{\nabla}_{\mu}^{(n)} = \overrightarrow{\partial} + \frac{1}{2}\omega_{\mu}^{cd}\sigma_{cd}, \ \overrightarrow{\nabla}_{\mu}^{(E)} = \overrightarrow{\partial} + \frac{1}{2}\omega_{\mu}^{cd}\sigma_{cd} + ieA_{\mu}; R(\omega, V) = e^{a\mu}e^{b\nu}R_{\mu\nu ab}(\omega)$ is the scalar curvature; e^{μ}_{a} and ω_{μ}^{ab} are the vierbein and spin-connection; $g^{\mu\nu} = e^{\mu}_{a}e^{\nu}_{b}\eta^{ab}$ and $R_{\mu\nu ab}(\omega) = \partial_{\mu}\omega_{\nu ab} + \omega^{c}_{\mu a}\omega_{\nu cb} - (\mu \leftrightarrow \nu)$. The quantities V_1 and V_2 are constants with the dimensionality $(mass)^4$. In addition, the constants b, k, h_n, h_E are non specified dimensionless real parameters. In relation to the parameter α , we have that it is a real parameter which is positive and dimensionless. We will determine the relative signs of the kinetic terms and mass terms in the above lagrangian

3

comparing only the terms that are multiplied by the conventional measure, and since k will be determined to negative, we will choose the corresponding coefficients of the mass terms, like μ_E and μ_n to be negative.

We note that the general action given by Eq.(1) is invariant under the global scale transformations

$$e^{a}_{\mu} \to e^{\theta/2} e^{a}_{\mu}, \quad \omega^{\mu}_{ab} \to \omega^{\mu}_{ab}, \quad \varphi_{a} \to \lambda_{ab} \varphi_{b} \quad A_{\alpha} \to A_{\alpha},$$

$$\phi \to \phi - \frac{M_{p}}{\alpha} \theta, \quad \Psi_{i} \to e^{-\theta/4} \Psi_{i}, \quad \overline{\Psi}_{i} \to e^{-\theta/4} \overline{\Psi}_{i}, \quad \det(\lambda_{ab}) = e^{2\theta}.$$
 (2)

In relation to the two measure theory we have that it is a generally coordinate invariant theory, where the action becomes [9]-[11]

$$S = \int L_1 \Phi d^4 x + \int L_2 \sqrt{-g} d^4 x, \qquad (3)$$

where there are two Lagrangians L_1 and L_2 and two measures of integration: the standard $\sqrt{-g}$ and the new one Φ . Here, we have that the Lagrangian L_1 satisfies the relation [9]

$$B_a^{\mu}\partial_{\mu}L_1 = 0$$
, and then $L_1 = M^4 = \text{const.}$ (4)

where $B_a^{\mu} = \varepsilon^{\mu\nu\alpha\beta}\varepsilon_{abcd}\partial_{\nu}\varphi_b\partial_{\alpha}\varphi_c\partial_{\beta}\varphi_d$ and since the determinant of the matrix B_a^{μ} is proportional to the measure, it follows that for a non singular measure we have that $\partial_{\mu}L_1 = 0$.

From the action given by Eq.(1), we have a first order formalism that contain terms proportional to $\partial_{\mu}\chi$, where $\chi = \Phi/\sqrt{-g}$. Thus, we obtain that the space-time non-Riemannian and equations of motion - non canonical.

In order to obtain the Einstein frame, we can define a new set of variables (ϕ and A_{μ} remain unchanged) such that

$$\tilde{e}_{a\mu} = e^{\frac{i}{2}\alpha\phi/M_p} (\chi + b)^{1/2} e_{a\mu}, \quad \tilde{g}_{\mu\nu} = e^{\alpha\phi/M_p} (\chi + b) g_{\mu\nu},$$

$$\Psi'_i = e^{-\frac{1}{4}\alpha\phi/M_p} \frac{(\chi + k)^{1/2}}{(\chi + b)^{3/4}} \Psi_i, \quad \text{with} \quad i = n, E.$$
(5)

Note that $\tilde{e}_{a\mu}$, $\tilde{g}_{\mu\nu}$, n' and E' are invariant under the scale transformations (2).

In the Einstein frame the equations take the standard General Relativity form given by

$$G_{\mu\nu}(\tilde{g}_{\alpha\beta}) = \frac{\kappa}{2} T^{eff}_{\mu\nu},\tag{6}$$

in which the tensor $G_{\mu\nu}(\tilde{g}_{\alpha\beta})$ corresponds to the Einstein tensor in the Riemannian space-time with the metric $\tilde{g}_{\mu\nu}$ and the quantity $\kappa = 8\pi/(M_p^2)$ with M_p the Planck mass. Also, the energy-momentum tensor $T_{\mu\nu}^{eff}$ is defined as

$$T_{\mu\nu}^{eff} = \phi_{,\mu}\phi_{,\nu} - \frac{1}{2}\tilde{g}_{\mu\nu}\tilde{g}^{\alpha\beta}\phi_{,\alpha}\phi_{,\beta} + \tilde{g}_{\mu\nu}V_{eff}(\phi;\chi) + T_{\mu\nu}^{(em)} + T_{\mu\nu}^{(ferm,can)} + T_{\mu\nu}^{(ferm,noncan)},$$
(7)

where the effective potential associated to scalar field and χ is given by

$$V_{eff}(\phi,\chi) = \frac{b\left(M^4 e^{-2\alpha\phi/M_p} + V_1\right) - V_2}{(\chi+b)^2}.$$
(8)

Besides, the quantity $T_{\mu\nu}^{(em)}$ corresponds to the canonical energy momentum tensor of the electromagnetic field and the tensor $T_{\mu\nu}^{(ferm,can)}$ represents the canonical energy momentum tensor for primordial fermions n' and E' in curved space-time. The quantity $T_{\mu\nu}^{(ferm,noncan)}$ denotes the noncanonical contribution of the fermions into the energy momentum tensor and it is defined as[22]

$$T_{\mu\nu}^{(ferm,noncan)} = -\tilde{g}_{\mu\nu}\Lambda_{dyn}^{(ferm)},\tag{9}$$

where $\Lambda_{dyn}^{(ferm)}$ is given by

$$\Lambda_{dyn}^{(ferm)} \equiv Z_n(\chi)m_n(\chi)\overline{n'}n' + Z_E(\chi)m_E(\chi)\overline{E'}E', \qquad (10)$$

and $Z_i(\chi)$ and $m_i(\chi)$ (i = n', E') are defined as

$$Z_i(\chi) \equiv \frac{(\chi - \chi_1^{(i)})(\chi - \chi_2^{(i)})}{2(\chi + k)(\chi + h_i)}, \qquad m_i(\chi) = \frac{\mu_i(\chi + h_i)}{(\chi + k)(\chi + b)^{1/2}},$$
(11)

where

$$\chi_{1,2}^{(i)} = \frac{1}{2} \left[k - 3h_i \pm \sqrt{(k - 3h_i)^2 + 8b(k - h_i) - 4kh_i} \right].$$
(12)

It is important to mention that the noncanonical contribution given by $T^{(ferm,noncan)}_{\mu\nu}$ of the fermions into the energy momentum tensor produces a cosmological constant like term but it is proportional to fermion densities $\overline{\Psi}'_i \Psi'_i$ (i = n', E'), and this is why we will refer to it as "dynamical fermionic Λ term".

The scalar field χ given by $\chi = \Phi/\sqrt{-g}$, is determined in terms of the scalar field ϕ and $\overline{\Psi}'_i \Psi'_i$ (i = n', E') by the following equation[22]

$$\frac{1}{(\chi+b)^2} \left\{ (b-\chi) \left[M^4 e^{-2\alpha\phi/M_p} + V_1 \right] - 2V_2 \right\} = \Lambda_{dyn}^{(ferm)}.$$
(13)

In the following, we will study the inflationary stage in which we will consider the absence of fermions during this epoch.

III. INFLATION

In this section we will analyze the inflationary era where we do not consider fermions so that $\Lambda_{dyn}^{(ferm)} = 0$. Thus, from Eq.(13) we find that the quantity χ in terms of the scalar field becomes

$$\chi = \chi_0(\phi) \equiv b - \frac{2V_2}{V_1 + M^4 e^{-2\alpha\phi/M_p}}.$$
(14)

Now replacing this equation into Eq.(8) we obtain that the effective potential of the scalar field ϕ in absence of fermions results

$$V_{eff}(\phi) \equiv V_{eff}(\phi;\chi_0)|_{\overline{\psi'}\psi'=0} = \frac{[V_1 + M^4 e^{-2\alpha\phi/M_p}]^2}{4[b\left(V_1 + M^4 e^{-2\alpha\phi/M_p}\right) - V_2]}.$$
(15)




FIG. 1: A schematic representation of the effective potential $V(\phi)$ as a function of the scalar field ϕ , given by Eq. (15). The flat region corresponds to the inflationary phase described by Eq. (16). At the minimum, the mechanism of reheating, and also the production of fermions occurs.

In order to analyze the inflationary period, we can consider that the scalar field start from very high positive values $(\phi \to +\infty)$, with a flat region given by

$$V_{eff}(\phi) \simeq \frac{V_1^2}{4(bV_1 - V_2)}, \text{ with } bV_1 > V_2.$$
 (16)

Fig.1 shows the effective potential as a function of the scalar field defined by Eq.(15). For this potential, we note that the flat region refers to the inflationary phase defined by Eq.(16). Furthermore, at the minimum of this effective potential, the mechanism of reheating occurs. This process involves the perturbative decay of the oscillating field ϕ at the end of the inflationary scenario, which generates the reheating of the universe[23]. In this context, various

particles, including fermions, are produced at the minimum of the effective potential. Here we have used the values; $\alpha = 1, V_1 = (1.1/3)M_p^4, V_2 = (1/3)M_p^4, M = 0.01M_p$ and b = 1, respectively.

Now, expanding the effective potential (15), we obtain that the next term is given by

$$V_{eff}(\phi) \simeq \frac{V_1^2}{4(bV_1 - V_2)} + \frac{V_1 M^4}{2(bV_1 - V_2)} \left[1 - \frac{bV_1}{2(bV_1 - V_2)} \right] e^{-2\alpha\phi/M_p} + \dots,$$
(17)

with $bV_1/2 < V_2$, in order to obtain a decrease of the effective potential as we move to smaller values of the scalar field during the inflationary epoch. However, we also have that $bV_1 > V_2$, then the condition on the parameter V_2 becomes $bV_1/2 < V_2 < bV_1$.

This choice $bV_1/2 < V_2$ differs from that of Ref.[22], where inflation was not considered and the evolution was in the other direction. In contrast, here, the high values of the scalar field will be used twice, once for inflation and then, after reheating, the same scalar field goes back in the other direction, but now feeling a modified potential modified by the particles created during the reheating.

In the context of the slow roll approximation and considering a flat Friedmann-Roberson-Walker metric, we have that the equations of motion are reduced to

$$3H^2 = \kappa \rho \simeq \kappa V_{eff}, \text{ and } 3H\dot{\phi} \simeq -\frac{\partial V_{eff}}{\partial \phi},$$
(18)

where $H = \dot{a}/a$ denotes the Hubble parameter and the quantity *a* the scale factor. Here, in the absence of fermions, we have considered Eq.(6) together with the conservation of the energy-momentum tensor, defined by Eq. (7).

Thus, under the slow roll approximation from Eq.(17), we find that the scalar field in terms of the cosmological time can be written as

$$e^{\alpha\phi/M_p}\sqrt{(V_{01}+V_{02}e^{-2\alpha\phi/M_p})}\left(e^{\alpha\phi/M_p}+\frac{\sqrt{V_{02}}\operatorname{ArcSinh}[e^{\alpha\phi/M_p}\sqrt{V_{01}/V_{02}}]}{\sqrt{V_{01}}\sqrt{1+e^{2\alpha\phi/M_p}V_{01}/V_{02}}}\right)=\frac{4V_{02}(\alpha/M_p)^2}{\sqrt{3\kappa}}t+C_0,\qquad(19)$$

where C_0 corresponds to an integration constant. Here the constants V_{01} and V_{02} are defined as

$$V_{01} = \frac{V_1^2}{4(bV_1 - V_2)}, \quad \text{and} \quad V_{02} = \frac{V_1 M^4}{2(bV_1 - V_2)} \left[1 - \frac{bV_1}{2(bV_1 - V_2)} \right] < 0, \tag{20}$$

respectively. From Eq.(19), we note that for $\phi \to \infty$, the solution for the scalar field in terms of time is approximated to

$$e^{2\alpha\phi/M_p} \sim \frac{4V_{02}(\alpha/M_p)^2}{\sqrt{3\kappa V_{01}}}t + C'_0,$$

where $C'_0 = C_0 / \sqrt{V_{01}}$.

By introducing the slow roll parameters ϵ and η , we have

$$\epsilon = \frac{1}{2\kappa} \left[\frac{\partial V_{eff} / \partial \phi}{V_{eff}} \right]^2, \quad \text{and} \quad \eta = \frac{1}{\kappa} \left[\frac{\partial^2 V_{eff} / \partial \phi^2}{V_{eff}} \right]. \tag{21}$$

Thus, considering that the end of inflation taken place when $\epsilon(\phi = \phi_e) = 1$, we find the value at the end of inflation becomes

$$e^{-2\alpha\phi_e/M_p} = -\left[\frac{V_{01}}{V_{02}}\right] \frac{1}{\left[\frac{\sqrt{2}\alpha}{\sqrt{\kappa}M_p} + 1\right]}.$$
(22)

Here ϕ_e denotes the value of the scalar field at the end of inflationary epoch and recalled that $V_{02} < 0$.

Also, the number of e-folds N between two different values of the times t and t_e results

$$N = \int_{t}^{t_{e}} H \, dt \simeq \kappa \int_{\phi_{e}}^{\phi} \frac{V_{eff}}{(\partial V_{eff}/\partial \phi)} \, d\phi, \tag{23}$$

where t_e denotes the end of the inflationary epoch. Thus, we find that the number of e-folds as a function of the scalar field can be written as

$$N = \left(\frac{\kappa M_p}{2\alpha |V_{02}|}\right) \left(\left[\frac{M_p V_{01} e^{2\alpha \phi/M_p}}{2\alpha} + V_{02} \phi\right] - \left[\frac{M_p V_{01} e^{2\alpha \phi_e/M_p}}{2\alpha} + V_{02} \phi_e\right] \right).$$
(24)

In this context, we can determine the scalar field as a function of the number of e- folds as

$$\phi(N) = \frac{\tilde{N}}{V_{02}} - \frac{M_p \text{ProducLog}[(V_{01}/V_{02})e^{2\alpha N/(M_p V_{02})}]}{2\alpha},$$
(25)

in which the quantity \tilde{N} is defined as

$$\tilde{N} = \left(\frac{2\alpha|V_{02}|}{\kappa M_p}\right) N + \left[\frac{M_p V_{01} e^{2\alpha\phi_e/M_p}}{2\alpha} + V_{02}\phi_e\right],$$

where the scalar field at the end of inflation ϕ_e is given by Eq.(22). Besides, the ProductLog function corresponds to the product logarithm, also called the Omega function or Lambert W function. This function is a multivalued function, and it satisfy the converse relation of the function $f(w) = we^w$, see Ref.[24]

On the other hand, the power spectrum \mathcal{P}_R of the curvature perturbations, in the context of the slow roll approximation becomes[25]

$$\mathcal{P}_R = \left(\frac{H^2}{\dot{\phi}^2}\right) \left(\frac{H}{2\pi}\right)^2 \simeq \frac{\kappa^3}{12\pi^2} \left[\frac{V_{eff}^3}{(\partial V_{eff}/\partial \phi)^2}\right],\tag{26}$$

and the scalar spectral index n_s results [25]

$$n_s - 1 = -6\epsilon + 2\eta,\tag{27}$$

and the tensor to scalar ratio r is defined as [25]

$$r = \frac{8}{\kappa} \left[\frac{\partial V_{eff} / \partial \phi}{V_{eff}} \right]^2.$$
⁽²⁸⁾

From Eq.(26) we find that the scalar power spectrum results

$$\mathcal{P}_R \simeq k_1 e^{4\alpha\phi/M_p} \left[V_{01} + V_{02} e^{-2\alpha\phi/M_p} \right]^3,$$
(29)

where the constant k_1 is defined as

$$k_1 = \frac{\kappa^3 M_p^2}{48\pi^2 \alpha^2 V_{02}^2}$$

Also, we determine that the scalar spectral index in terms of the scalar field from Eq.(27) results

$$n_s - 1 = -\left[\frac{4\alpha^2 |V_{02}|}{\kappa M_p^2}\right] (|V_{02}| + 2V_{01}e^{2\alpha\phi/M_p}) (V_{01} + V_{02}e^{2\alpha\phi/M_p})^{-2}.$$
(30)

From Eq. (28) we find that the tensor to scalar ratio r as a function of the scalar field becomes

$$r = r_0 e^{-4\alpha\phi/M_p} \left(V_{01} + V_{02} e^{2\alpha\phi/M_p} \right)^{-2}, \tag{31}$$

where the constant r_0 is defined as

$$r_0 = \frac{32\alpha^2 V_{02}^2}{\kappa M_p^2}$$

In particular assuming the value of the parameter $\alpha = 1$ together with the cosmological parameters $\mathcal{P}_R = 2.2 \times 10^{-9}$ and $n_s = 0.968$ at N = 60 (see [26]), then numerically we obtain that the parameters V_{01} and V_{02} result

$$V_{01} \simeq 4,034 \times 10^{-4} M_p^4$$
, and $|V_{02}| = 5,286 \times 10^{-5} M_p^4$, (32)

respectively. In addition, using these values, we find that from these values the tensor to scalar ratio r at N = 60, becomes $r \simeq 0.01299$.

7

In relation to the original parameters, we note that by substituting the constraints given by Eq.(32) into Eq. (20), we find that the parameter M, obtained from the condition (4), can be expressed in terms of the parameters V_1 and b as follows:

$$M \simeq \left(\frac{6.3 \times 10^{-3} V_1}{\frac{b}{125 V_1} - 1}\right)^{1/4},\tag{33}$$

with the condition $b > 125V_1$ to ensure a real solution for the parameter M. Besides, we note that for the special situation in which the ratio $b/125V_1 \gg 1$, the parameter M becomes $M \simeq V_1^{1/2} b^{-1/4}$. Additionally, we find that the equation for V_2 in terms of V_1 and b is given by

$$V_2 \simeq bV_1 - 625V_1^2. \tag{34}$$

Here we need to consider that $b > 625V_1$ to obtain a positive value of the parameter V_2 . Thus, this condition on the parameter b is more strong that the condition $b > 125V_1$ obtained from Eq.(33) to find a real solution of the parameter M. As before, here we have considered Eqs.(20) and (32), respectively. In addition, we note that for the case in which the parameter $V_1 \ll 1$, we have that $V_2 \sim b V_1$.

After of the inflationary stage, the scalar field oscillates above the minimum of the effective potential and during this process occurs the particles production and in particular the fermions. In particular, we will look at the effect of the fermions as they get diluted and how they affect the dark energy in the late universe.

In the following, we will analyze the effect of the fermions as dark energy and dark matter.

IV. DARK ENERGY FROM FERMIONS

In this section we will study the accelerated expansion of the present universe produced from the effective potential associated to the fermions generated during the inflationary stage.

For dust, the neutrino has zero momenta, and then we have $\overline{n}'n' = u^{\dagger}u$ where u corresponds to the large component of the Dirac spinor n'. From the space components of the 4-current $\tilde{e}^{\mu}_{a}\overline{n}'\gamma^{a}n'$ equal zero and assuming the 4-current conservation we have $\overline{n}'n' = u^{\dagger}u = \frac{const}{a^{3}}$ where a = a(t) as before is the scale factor. Thus, from Eq. (13) we have

$$\frac{(b-\chi)\left[M^4 e^{-2\alpha\phi/M_p} + V_1\right] - 2V_2}{(\chi+b)^{3/2}} = \frac{(\chi-\chi_1^{(N)})(\chi-\chi_2^{(N)})}{(\chi+k)^2}\mu_N\frac{const}{a^3}.$$
(35)

We note that there is a solution where the decaying fermion contribution $u^{\dagger}u \sim \frac{const}{a^3}$ is not trivial to the right hand side of Eq.(35). In this sense, to generate the dark energy from the fermions, we can consider the asymptotic behavior $\chi \to -k$ in such a way that $(\chi + k)^{-2} \propto a^3$, then the r.h.s. of (35) became a constant. We note that the described regime denotes a very unexpected state of the primordial fermions.

To obtain the total effective potential $U_{eff}(\phi)$ in terms of the scalar field associated with fermions during the current universe, we have the following

$$U_{eff} = V_{eff} - \Lambda_{dyn}^{(ferm)} \simeq \frac{V_2 - kV_1}{(b-k)^2} - \frac{k}{(b-k)^2} M^4 e^{-2\alpha\phi/M_p} + \mathcal{O}\left(\chi + k\right).$$
(36)

Here we have utilized Eqs.(8) and (13) together with $\chi \to -k$. Notice that this vacuum energy can be very small if -k is very big.

We note from this total effective potential U_{eff} we have a modified flat region for positive large ϕ associated to dark energy at the present given by

$$U_{eff}(\phi \to \infty) = U_{DE} \simeq \frac{V_2 - kV_1}{(b-k)^2}.$$
 (37)

In addition, we can define the density parameter associated to the dark energy $\Omega_{DE} = \kappa \rho_{DE}/3H^2 \simeq \kappa U_{eff}/3H^2$. Thus, evaluating the parameter Ω_{DE} at the present time i.e., Ω_{DE_0} , we find

$$\Omega_{DE_0} \simeq \frac{\kappa}{3H_0^2} \left(\frac{V_2 - kV_1}{(b-k)^2} \right).$$
(38)

By considering that the Hubble parameter at the present time, $H_0 \approx 10^{-61} M_p$, we obtain the constraint

$$\frac{V_2 - kV_1}{(b-k)^2} \approx 2 \times 10^{-122} \frac{M_p^2}{\kappa},\tag{39}$$

where we have used that at the present the parameter $\Omega_{DE_0} \simeq 0.7$.

Besides, for very large values of k and considering Eq.(37), we can obtain an approximation for the parameter k and it becomes of order

$$|k| \sim \frac{V_1}{U_{DE}} \sim 10^{119}.$$
(40)

Here, we have assumed that $V_1 \sim V_2$, $b \sim \mathcal{O}(1)$, and, from Eq.(32) $V_{01} \sim V_1 \sim 10^{-4} M_p^4$, obtained during the inflationary epoch, together with the value of the dark energy at the present epoch, $U_{DE} \sim 10^{-123} M_p^4$.

V. DARK MATTER FROM FERMIONS

In this section, we will analyze the dark matter associated with the high-density fermions (localized) in our present universe generated during the inflationary epoch.

From the constraint equation given by Eq.(13), the right hand size is proportional to the fermions densities. In this context, $\Lambda_{dyn}^{(ferm)}$ given by Eq.(10) becomes very large unless the coefficient Z_i defined by Eq.(11) goes to zero. Thus, to obtain the coefficient $Z_i \to 0$, we need to consider that $\chi \to \chi_{1,2}^{(i)}$ where $\chi_{1,2}^{(i)}$ are defined by Eq.(12). By considering that in the studied regime masses (11) of the non-relativistic primordial fermions are very large,

By considering that in the studied regime masses (11) of the non-relativistic primordial fermions are very large, we can neglect their kinetic energy. Thus, after averaging on the microscopic contribution to the energy-momentum tensor $T^{(n,can)}_{\mu\nu}$ we can written [22]

$$< T^{(n',can)}_{\mu\nu} >_{cosm.av.} \simeq \delta^0_\mu \delta^0_\nu \frac{h_n - k}{(b-k)^{1/2}} \mu_n \frac{c}{(\chi+k)a^3},$$
(41)

where c is a constant. In order to obtain that the coefficient $Z_i \to 0$ and considering only the neutrinos (i = n'), in which $\chi \to \chi_{1,2}^{n'}$ and then we have that the canonical energy momentum tensor for fermions n' results

$$< T_{\mu\nu}^{(n',can)} >_{cosm.av.} \simeq \delta_{\mu}^{0} \delta_{\nu}^{0} \frac{h_{n} - k}{(b-k)^{1/2}} \mu_{n} \frac{c_{1,2}}{(\chi_{1,2}^{n'} + k)a^{3}} \propto \frac{1}{a^{3}}.$$
(42)

Here we note that there are two cases for the canonical energy momentum associated to the coefficients $\chi_{1,2}^{n'}$ (solutions) defined by Eq.(12).

Thus, for the first solution $\chi_1^{n'}$ from Eq.(42) we find that $\langle T_{\mu\nu}^{(n',can)} \rangle_{cosm.av.}$ becomes

$$< T^{(n',can)}_{\mu\nu} >_{cosm.av.} \simeq \delta^0_\mu \delta^0_\nu \frac{h_n - k}{(b-k)^{1/2}} \mu_n \frac{c_1}{(\chi^{n'}_1 + k)a^3} \propto \frac{1}{a^3}.$$
 (43)

In the case when consider the second solution for the coefficient $\chi_2^{n'}$ we have that the canonical energy momentum tensor for fermions is given by

$$< T^{(n',can)}_{\mu\nu} >_{cosm.av.} \simeq \delta^0_\mu \delta^0_\nu \frac{h_n - k}{(b-k)^{1/2}} \mu_n \frac{c_2}{(\chi^{n'}_2 + k)a^3} \propto \frac{1}{a^3}.$$
 (44)

In this context, we consider the energy density related to dark matter to be defined as $\rho_{DM} = \langle T_{00}^{(n',can)(1)} \rangle_{cosm.av.} + \langle T_{00}^{(n',can)(2)} \rangle_{cosm.av.}$. In this sense, we find that the energy density associated to dark matter for large-k can be written as

$$\rho_{DM} \simeq \frac{h_n - k}{(b-k)^{1/2}} \mu_n \left[\frac{c_1}{(\chi_1^{n'} + k)} + \frac{c_2}{(\chi_2^{n'} + k)} \right] \frac{1}{a^3} \propto \frac{1}{a^3}.$$
(45)

Defining the density parameter related to the dark matter $\Omega_{DM} = \kappa \rho_{DM}/3H^2$, then we find that at the present time the density parameter becomes

$$\Omega_{DM_0} = \kappa \frac{h_n - k}{3H_0^2 (b - k)^{1/2}} \mu_n \left[\frac{c_1}{(\chi_1^{n'} + k)} + \frac{c_2}{(\chi_2^{n'} + k)} \right],\tag{46}$$

where we have used that scale factor at present $a_0 = 1$. Now considering the density parameter $\Omega_{DM_0} \simeq 0.3$ and the Hubble parameter in the present $H_0 \approx 10^{-61} M_p$, we have

$$\frac{h_n - k}{(b-k)^{1/2}} \mu_n \left[\frac{c_1}{(\chi_1^{n'} + k)} + \frac{c_2}{(\chi_2^{n'} + k)} \right] = \frac{3H_0^2 \Omega_{DM_0}}{\kappa} \approx \left(\frac{10^{-122} M_p^2}{\kappa} \right),\tag{47}$$

here we have utilized the value of k given by Eq.(40).

By assuming that the parameter k is very large and the mass parameter μ_n negative, we find that the constraint given by Eq.(47) is reduced to

$$|\mu_n| \left[\frac{c_1}{2} + c_2\right] \approx \left(\frac{10^{-62} M_p^2}{\kappa}\right). \tag{48}$$

Here we have used that for large-k, the first solution of Eq.(12) corresponds to $\chi_1^{n'} \to k$ and the second solution $\chi_2^{n'} \to 0$. In addition, we have considered the valor of k found in Eq.(40).

VI. DISCUSSION ON POSSIBLE FURTHER DEVELOPMENTS

In this paper we have discussed a unified picture of inflation and the late DE and DM period of the universe based on a scalar field together with the creation of fermions that in a model with a dilaton field and spontaneously broken scale invariance in the context of a Two Measures Theory.

During the inflationary stage the scalar field produces the accelerated expansion of the universe before the production of exotic fermions. In inflation we have found different constraints related to the effective potential of the scalar field in absence of fermions from the observational parameters.

In the case of the late universe, the fermions, when diluted can have exotic behavior and can substantially modified the scalar field potential, reducing the very large inflation to the very low DE in the late universe. In order to describe the late universe, we have considered that after reheating the scalar field retraces its trajectory in field space but now the scalar field potential can be drastically modified by the effect of the fermions. From the noncanonical contribution of the fermions associated to the energy momentum tensor we have obtained a part of the dark energy responsible of the present accelerated universe. The other contribution to the dark energy is related to the scalar field contribution which is also modified by the fermions because of the solution of the constraint equation given by Eq.(35). We have considered that the present dark energy with its very small value in comparison to the inflationary phase which can be adjusted by choosing appropriately the parameter space of couplings of the Riemannian and non Riemannian measures to the fermions.

To generate the dark matter we have considered the canonical contribution of fermions related to energy momentum tensor. In these clumped regions, we have that the matter is more dense. For the dark matter we have assumed two solutions associated to the constraint $\chi \to \chi_{1,2}^{n'}$ that contributes to the energy density of the dark matter ρ_{DM} . In the limit in which the parameter k is very large, we have obtained a constraint on the mass parameter and the coefficients c_1 and c_2 from the present density parameter Ω_{DM_0} , see Eq.(48).

Of course in the future, there are many issues to be studied, for example, what are these fermions?, may be sterile neutrinos?. How this theory works for structure formation?. Since the final value of the DE takes some time as the universe needs to expand so that the asymptotic value of the DE sets in, does this evolution of the DE helps to explain the H_0 tension through some kind of Early DE scenario?.

ACKNOWLEDGEMENTS

EG acknowledges the Pontificia Universidad Católica de Valparaíso for hospitality. He also acknowledges COST action CA23130 - Bridging high and low energies in search of quantum gravity (BridgeQG) and CosmoVerse • COST Action CA21136 Addressing observational tensions in cosmology with systematics and fundamental physics for finantial support.

Appendix A: Connection in the original and Einstein frames

In this appendix, we show the dependence of the spin connection ω_{μ}^{ab} on e_{μ}^{a} , χ , Ψ and $\overline{\Psi}$. Thus, varying the Eq.(1) with respect to ω_{μ}^{ab} and considering

$$R(V,\omega) \equiv -\frac{1}{4\sqrt{-g}} \varepsilon^{\mu\nu\alpha\beta} \varepsilon_{abcd} e^c_{\alpha} e^d_{\beta} R^{ab}_{\mu\nu}(\omega) \tag{A1}$$

we find

$$\varepsilon^{\mu\nu\alpha\beta}\varepsilon_{abcd}\left[(\chi+b)e^{c}_{\alpha}D_{\nu}e^{d}_{\beta} + \frac{1}{2}e^{c}_{\alpha}e^{d}_{\beta}\left(\chi_{,\nu} + \frac{\alpha}{M_{p}}\phi_{,\nu}\right)\right] + \frac{\kappa}{4}\sqrt{-g}(\chi+k)e^{c\mu}\varepsilon_{abcd}\overline{\Psi}\gamma^{5}\gamma^{d}\Psi = 0, \tag{A2}$$

in which

$$D_{\nu}e_{a\beta} \equiv \partial_{\nu}e_{a\beta} + \omega^{d}_{\nu a}e_{d\beta}.$$
 (A3)

The solution of Eq. (A2) can be written as

$$\omega_{\mu}^{ab} = \omega_{\mu}^{ab}(e) + K_{\mu}^{ab}(e, \overline{\Psi}, \Psi) + K_{\mu}^{ab}(\chi, \phi), \tag{A4}$$

where

$$\omega_{\mu}^{ab}(e) = e_{\alpha}^{a} e^{b\nu} \{^{\alpha}_{\mu\nu}\} - e^{b\nu} \partial_{\mu} e_{\nu}^{a}, \tag{A5}$$

and it corresponds to the Riemannian part of the spin-connection. Besides, the quantity

$$K^{ab}_{\mu}(e,\overline{\Psi},\Psi) = \frac{\kappa}{8} \frac{\chi+k}{\chi+b} \eta_{cn} e_{d\mu} \varepsilon^{abcd} \overline{\Psi} \gamma^5 \gamma^n \Psi, \tag{A6}$$

denoted the fermionic contribution that differs from the standard expression [27] by the factor $\frac{\chi+k}{\chi+b_g}$. Also, the parameter $K^{ab}_{\mu}(\chi,\phi)$ is defined as

$$K^{ab}_{\mu}(\chi,\phi) = \frac{1}{2(\chi+b)} \left(\chi_{,\alpha} + \frac{\alpha}{M_p}\phi_{,\alpha}\right) \left(e^a_{\mu}e^{b\alpha} - e^b_{\mu}e^{a\alpha}\right),\tag{A7}$$

and it corresponds to the non-Riemannian part of the spin-connection obtained by specific features of TMT.

In the Einstein frame, we determine that the variables defined by Eq.(5), the spin-connection read

$$\omega_{\mu}^{\prime ab} = \omega_{\mu}^{ab}(\tilde{e}) + \frac{\kappa}{8} \eta_{cn} \tilde{e}_{d\mu} \varepsilon^{abcd} \overline{\Psi}^{\prime} \gamma^5 \gamma^n \Psi^{\prime}, \tag{A8}$$

which is equivalent to the spin-connection of the Einstein-Cartan space-time[27] with the vierbein \tilde{e}^a_{μ} . In the solution of this paper, the last term given by Eq.(A8) is zero.

A. G. Riess et al., Astron. J. 116, 1009 (1998); S. Perlmutter et al., Astrophys. J. 517, 565 (1999); N. Bahcall, J.P. Ostriker, S.J. Perlmutter and P.J. Steinhardt, Science 284, 1481 (1999); D.N. Spergel, et al. [WMAP collaboration], Astrophys. J. Suppl. 148, 175 (2003); A. C. S. Readhead, et al. Astrophys. J. 609, 498 (2004); J. H. Goldstein et al., Astrophys.J. 599, 773-785 (2003); R. Rebolo, et al., astro-ph/0402466; M. Tegmark, et al. Phys.Rev. D69, 103501 (2004); E. Hawkins et al., Mon.Not.Roy.Astron.Soc. 346, 78 (2003); W. L. Freedman, et al., Astrophys.J. 553, 47 (2001); R. Daly, S.G. Djorgovski, Astrophys.J. 597, 9, (2003).

 ^[2] S.M.Carroll, Living Rev. Rel. 4, 1 (2001); A. Vilenkin, hep-th/0106083; P.J.E. Peebles and B. Ratra, Rev. Mod. Phys. 75, 559 (2003); T. Padmanabhan, Phys. Rep. 380, 235 (2003); V. Sahni, AIP Conf. Proc. 782, 166 (2005).

^[3] I. Zlatev, L Wang and P. Steinhardt, Phys. Rev. Lett. 82, 896 (1999).

^[4] C. Wetterich, Nucl. Phys. B302, 668 (1988); B. Ratra and P.J.E. Peebles, Phys. Rev. D37, 3406 (1988); P.J.E. Peebles and B. Ratra, Astrophys. J. 325, L17 (1988); R. Caldwell, R. Dave and P. Steinhardt, Phys. Rev. Lett. 80, 1582 (1998); N. Weiss, Phys. Lett. B197, 42 (1987); Y. Fujii and T. Nishioka, Phys. Rev. D42, 361 (1990); M.S. Turner and M. White, Phys. Rev. D56, R4439 (1997); P. Ferreira and M.Joyce, Phys. Rev. Lett. 79, 4740 (1997); Phys. Rev. D58, 023503 (1998).
E. Copeland, A. Liddle and D. Wands, Phys. Rev. D57, 4686 (1998); P. Steinhardt, L Wang and I. Zlatev, Phys. Rev. D59, 123504 (1999).

- [5] Alan H. Guth, Phys.Rev.D 23 (1981) 347-356; Adv.Ser.Astrophys.Cosmol. 3 (1987) 139-148, A. Starobinsky, Phys.Lett. B 91, 99-102 (1980), E.W. Kolb and M.S. Turner, "The Early Universe", Addison Wesley (1990);
 A. Linde, "Particle Physics and Inflationary Cosmology", Harwood, Chur, Switzerland (1990);
 A. Guth, "The Inflationary Universe", Vintage, Random House (1998);
 S. Dodelson, "Modern Cosmology", Acad. Press (2003);
 S. Weinberg, "Cosmology", Oxford Univ. Press (2008), V. Mukhanov, "Physical Foundations of Cosmology", Cambridge
- S. Weinberg, "Cosmology", Oxford Univ. Press (2008), V. Mukhanov, "Physical Foundations of Cosmology", Cambridge Univ. Press (2005).
- [6] P.Q. Hung, hep-ph/0010126; P.Q. Hung and H. Pas, astro-ph/0311131; A. Singh, *Phys. Rev.* D52, 6700 (1995); M. Blasone, A. Capolupo, S. Capozzielo, S. Carloni, G. Vitiello, *Phys. Lett.* A323, 182 (2004); X. Zhang, hep-ph/0410292; X. Bi, B. Feng, H. Li and X. Zhang, hep-ph/0412002; A.W. Brookfield, C.van de Bruck, D.F. Mota, D. Tocchini-Valentini, astro-ph/050349; V. Barger, P. Huber and D. Marfatia, hep-ph/0502196; M. Cirelli, M.C. Gonzalez-Garcia and C. Pena-Garay, hep-ph/0503028; X.-f. Zhang, H. Li, Y.-S. Piao and X. Zhang, astro-ph/0501652.
- [7] R. Fardon, A. E. Nelson, N. Weiner *JCAP* **0410**, 005 (2004).
- [8] R.D. Peccei, *Phys. Rev.* D71, 023527 (2005).
- [9] E.I. Guendelman and A.B. Kaganovich, Phys. Rev. D53, 7020 (1996); Mod. Phys. Lett. A12, 2421 (1997); Phys. Rev. D55, 5970 (1997); Mod. Phys. Lett. A12, 2421 (1997); Phys. Rev. D55, 5970 (1997); ibid. D57, 7200 (1998); Mod. Phys. Lett. A13, 1583 (1998).
- [10] E.I. Guendelman and A.B. Kaganovich, Phys. Rev. D56, 3548 (1997).
- [11] E.I. Guendelman and A.B. Kaganovich, Phys. Rev. D60, 065004 (1999).
- [12] E.I. Guendelman, Mod. Phys. Lett. A14, 1043 (1999); Class. Quant. Grav. 17, 361 (2000); gr-qc/9906025; Mod. Phys. Lett. A14, 1397 (1999); gr-qc/9901067; hep-th/0106085; Found. Phys. 31, 1019 (2001);
- [13] E.I. Guendelman, Phys. Lett. B412, 42 (1997); E.I. Guendelman, gr-qc/0303048; E.I. Guendelman and E. Spallucci, hep-th/0311102.
- [14] E.I. Guendelman and O. Katz, Class. Quant. Grav., 20, 1715 (2003).
- [15] Sergio del Campo, Eduardo I. Guendelman, Ramon Herrera, Pedro Labraña. Emerging Universe from Scale Invariance, JCAP 06 (2010) 026 • e-Print: 1006.5734 [astro-ph.CO]
- [16] Sergio del Campo, Eduardo I. Guendelman, Alexander B. Kaganovich, Ramon Herrera, Pedro Labrana, Emergent Universe from Scale Invariant Two Measures Theory, Phys.Lett.B 699 (2011) 211-216 • e-Print: 1105.0651 [astro-ph.CO]
- [17] Eduardo Guendelman, Ramón Herrera, Pedro Labrana, Emil Nissimov, Svetlana Pacheva. Emergent Cosmology, Inflation and Dark Energy Gen.Rel.Grav. 47 (2015) 2, 10 • e-Print: 1408.5344 [gr-qc]
- [18] Eduardo I. Guendelman, Ramón Herrera, Curvaton reheating mechanism in a scale invariant two measures theory Gen.Rel.Grav. 48 (2016) 1, 3 • e-Print: 1511.08645 [gr-qc]; S. del Campo and R. Herrera, Phys. Rev. D 76, 103503 (2007) doi:10.1103/PhysRevD.76.103503 [arXiv:0710.5524 [astro-ph]]; S. del Campo, R. Herrera, J. Saavedra, C. Campuzano and E. Rojas, Phys. Rev. D 80, 123531 (2009) doi:10.1103/PhysRevD.80.123531 [arXiv:0912.4721 [astro-ph.CO]].
- [19] Eduardo I. Guendelman, Ramon Herrera, Pedro Labrana, Instant preheating in a scale invariant two measures theory Phys.Rev.D 103 (2021) 123515 • e-Print: 2005.14151 [gr-qc]
- [20] Eduardo Guendelman, Ramón Herrera, David Benisty, Unifying inflation with early and late dark energy with multiple fields: Spontaneously broken scale invariant two measures theory, Phys.Rev.D 105 (2022) 12, 124035 • e-Print: 2201.06470 [gr-qc]
- [21] Eduardo Guendelman, Ramon Herrera, Unification: Emergent universe followed by inflation and dark epochs from multifield theory, Annals Phys. 460 (2024) 169566 • e-Print: 2301.10274 [gr-qc]
- [22] E. I. Guendelman and A. B. Kaganovich, Int. J. Mod. Phys. A 21, 4373-4406 (2006) doi:10.1142/S0217751X06032538 [arXiv:gr-qc/0603070 [gr-qc]].
- [23] L. Abbott, E. Farhi, and M. B. Wise, Particle Production in the New Inflationary Cosmology, Phys.Lett. B 117, 29 (1982);
 A. Dolgov and A. D. Linde, Baryon Asymmetry in Inflationary Universe, Phys.Lett. B 116,329 (1982).
- [24] L. Surhone, M. Timplendon and S. Marseken, "Wright Omega Function: Mathematics, Lambert W Function, Continuous Function, Analytic Function, Differential Equation, Separation or Variables", Betascript Publishing (2010).
- [25] X. Chen, M. X. Huang, S. Kachru and G. Shiu, JCAP 0701 (2007) 002; B. A. Bassett, S. Tsujikawa and D. Wands, Rev. Mod. Phys. 78, (2006) 537-589.
- [26] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [27] F.W. Hehl, P. v. d. Heyde and G. Kerlick, *Reviews of Modern Physics* 48, 393 (1976); V. de Sabbata and M. Gasperini, *Introduction to Gravitation*, (World Scientific Publishing, Singapure, 1985).