

Joint Pricing and Resource Allocation: An Optimal Online-Learning Approach

Jianyu Xu^{1,2}, Xuan Wang², Yu-Xiang Wang³, and Jiashuo Jiang²

¹Carnegie Mellon University, PA

²Hong Kong University of Science and Technology, Hong Kong SAR

³University of California San Diego, CA

January 31, 2025

Abstract

We study an online learning problem on dynamic pricing and resource allocation, where we make joint pricing and inventory decisions to maximize the overall net profit. We consider the stochastic dependence of demands on the price, which complicates the resource allocation process and introduces significant non-convexity and non-smoothness to the problem. To solve this problem, we develop an efficient algorithm that utilizes a “Lower-Confidence Bound (LCB)” meta-strategy over multiple OCO agents. Our algorithm achieves $\tilde{O}(\sqrt{Tmn})$ regret (for m suppliers and n consumers), which is *optimal* with respect to the time horizon T . Our results illustrate an effective integration of statistical learning methodologies with complex operations research problems.

1 Introduction

The problem of *dynamic pricing* examines strategies of setting and adjusting prices in response to varying customer behaviors and market conditions. The mainstream of existing works on dynamic pricing, including Kleinberg and Leighton (2003); Broder and Rusmevichientong (2012); Cohen et al. (2020); Wang et al. (2021b), focuses on the estimation of *demand* curves while putting aside the decisions on the *supply* side. Another series of literature, including Besbes and Zeevi (2009); Chen et al. (2019, 2021a); Keskin et al. (2022), takes supply and inventories into account. However, these works simplify the supply cost as uniform and static, underestimating the difficulty of allocating products through sophisticated supply chains among multiple parties such as factories, warehouses, and retailers.

On the other hand, the problem of *resource allocation* – to serve different demand classes with various types of resources – presents a complex challenge within the field of operations research. Analogous to online dynamic pricing, the recent proliferation of e-platforms has magnified the importance of developing online allocation algorithms that efficiently manage supply and demand on the fly while maximizing cumulative utilities. However, traditional approaches in resource allocation are insufficient in depicting scenarios where the demand is stochastic and dependent on the price. Therefore, it is critical to develop price-dependent online allocation models and methodologies that can simultaneously learn the demand curve and optimize the joint decisions on price, inventory, and allocation.

This work introduces a novel framework for tackling the online pricing and allocation problem with an emphasis on learning under uncertainty. More specifically, we consider a problem setting where both the price and inventory decisions are made at the beginning of each time period, followed by inventory allocation based on the realized price-dependent stochastic demands during that period. We summarize the proposed framework as follows:

Pricing and Allocation. For $t = 1, 2, \dots, T$:

1. Determine the inventories of m suppliers as $\vec{I} := [I_1, I_2, \dots, I_m]^\top$ and incur an immediate inventory cost $\sum_{i=1}^m \gamma_i \cdot I_i$.
2. Propose a price p_t for all n consumers.
3. Based on the price p_t , consumers generate their demands as $\vec{D} := [D_1, D_2, \dots, D_n]^\top$.
4. We allocate inventories \vec{I} to satisfy demands \vec{D} . The allocation from Supplier i to Consumer j is denoted as $X_{i,j}$. The total supplying cost is $\sum_{i=1}^m \sum_{j=1}^n C_{i,j} \cdot X_{i,j}$.
5. We receive a payment from the consumers as $\sum_{i=1}^m \sum_{j=1}^n p_t \cdot X_{i,j}$ in total.

As described above, we assume that the inventories are *perishable* and the leftover inventory cannot be carried over to the following period. Furthermore, we assume that the price is *identical* for all consumers. We formalize the above process as solving the following two-stage stochastic programming problem:

$$\min_{p, \vec{I}} \langle \vec{\gamma}, \vec{I} \rangle + \mathbb{E}_{\vec{D}} [g(\vec{I}, p, \vec{D}) | p].$$

Here $g(\vec{I}, p, \vec{D})$ is the minimum *negative net profit* under the best allocation scheme given inventories \vec{I} , price p , and realized demand \vec{D} (see Eq. (2) for its definition).

1.1 Summary of Results.

In this work, we establish a novel framework of solving online pricing and allocation problem under demand uncertainties. Our main contributions are threefold:

1. **Algorithmic Design against Non-Convexity.** We propose an efficient online-learning algorithm for the (price,inventory) joint decision problem, which is highly *non-convex*. To navigate to the global optimal decisions among many sub-optimal, our algorithm incorporates an optimistic meta-algorithm with multiple online convex optimization (OCO) agents working locally.
2. **Closed-Form Allocation Scheme.** We present an algorithm that outputs a price-dependent *closed-form* solution to the optimal allocation problem.
3. **Regret Analysis.** Our algorithm achieves $\tilde{O}(\sqrt{Tmn})$ regret, which is *optimal* with respect to T as it matches the information-theoretic lower bound in (Broder and Rusmevichientong, 2012).

1.2 Technical Novelty.

To the best of our knowledge, we are the first to study dynamic pricing and inventory control under the framework of online resource allocation with uncertainty. Some existing works, such as Wang et al. (2021b); Chen and Gallego (2021), focus on related topics. However, their approaches are not capable of overcoming the significant challenges in our problem setting, including the local convexity, multiple suboptimals, and non-smoothness. As a consequence, we develop new techniques to solve the problem.

Confidence bound in vertical and horizontal directions. According to our problem setting, we suffer from *bandit feedback* and *non-convexity* with respect to price p . Existing methodologies in dynamic pricing mostly adopt continuum-armed bandits to resolve the challenges, nonetheless leading to $O(T^{2/3})$ regret that is sub-optimal in our setting. In this work, we design an $O(\sqrt{T})$ -regret algorithm, which makes a careful and novel use of the piecewise-convex property of the objective function. Specifically, we assign an *agent* to work on each convex interval, conducting *online convex optimization* (OCO) in local domains. Each agent maintains two high-probability confidence bounds: (1) A *vertical* bound on the *value* of local optimal, updated at the end of each *sub-epoch*. (2) A *horizontal* bound on the *location* of local optimal, updated at the end of each *epoch*.

Lower-confidence-bound (LCB) strategy over agents. In order to distinguish among every local-optimality iterated by a local OCO agent, we develop an LCB meta-algorithm over those agents. With the help of the vertical bound of each agent, we estimate the least possible value in each convex interval within reasonable error, and select the most optimistic agent (i.e., the one with the least lower bound) to make decisions in the following sub-epoch.

Closed-form solution of the second-stage LP. In the second stage when inventories are allocated to fulfill the realized demands, we solve a linear program (LP) that maximizes the net profit under the corresponding demand and supply constraints. In this work, we develop an efficient polynomial-time algorithm that achieves a closed-form solution to this LP. (On the contrary, an LP is generally not guaranteed to have closed-form solutions.) We also show the dependence of this solution on the price p , from which we derive the piecewise-convex property of the objective function.

The remainder of this paper is organized as follows. In Section 2, we review the relevant literature, contrasting existing approaches with ours. Section 3 lays out the preliminaries, including notation and problem setup. In Section 4, we introduce our learning-based algorithm and highlight its novel structures. Section 5 provides the regret analysis and theoretical guarantees. We finally discuss potential extensions in Section 6 and conclude this paper in Section 7.

2 Related Works

In this section, we present a review of the pertinent literature on pricing, allocation, and online convex optimization (OCO) problems, aiming to position our work within the context of related studies.

Dynamic Pricing. Quantitative research on dynamic pricing dates back to Cournot (1897) and has attracted significant attention in the field of machine learning (Leme et al., 2021; Xu and Wang, 2021; Jia et al., 2022; Choi et al., 2023; Simchi-Levi and Wang, 2023). For single-product pricing problems, the crux is to learn the demand curve and approach the optimal price. Under the assumptions of bandit feedback and k^{th} -smooth demand curves, Wang et al. (2021a) achieves an $O(T^{\frac{k+1}{2k+1}})$ regret. However, their methodologies are not applicable to our setting: Our objective function is only Lipschitz continuous, leading to $k = 1$ and an $O(T^{2/3})$ sub-optimal regret. In contrast, the piecewise convex property in our problem enables advanced methods to achieve a better regret. Another stream of works considers the heterogeneity of pricing processes, which includes item-wise features (Javanmard and Nazerzadeh, 2019; Cohen et al., 2020; Luo et al., 2021; Xu and Wang, 2022; Fan et al., 2021; Luo et al., 2022; Tullii et al., 2024), heteroscedasticity of valuations (Wang et al., 2021a; Ban and Keskin, 2021; Xu and Wang, 2024), time-

instationarity (Leme et al., 2021; Baby et al., 2023) and price discrimination (Chen et al., 2021c; Cohen et al., 2021; Eyster et al., 2021; Cohen et al., 2022; Xu et al., 2023; Karan et al., 2024). However, most of them are analyzing the differences (and their down-streaming effects) either in price or in demands, instead of the allocation process as we are concerned in this paper.

Pricing and Inventory Co-Decisions. The incorporation of inventory constraints into dynamic pricing problems began with the work of Besbes and Zeevi (2009) which assumed a fixed initial stock, and decisions of replenishment were later allowed in Chen et al. (2019). More recent studies, including Chen et al. (2020); Keskin et al. (2022), assumed perishable goods and took inventory costs into account. The stream of work by Chen et al. (2021a, 2023) further assumed the inventory-censoring effect on demands. However, none of them consider the heterogeneity of supply, nor the impact of prices on the allocation process. In our work, we not only model the inventory cost of each warehouse individually, but also depict the unit supplying cost from Warehouse i to Consumer j as a unique coefficient $C_{i,j}$.

Resource Allocation. There is a broad literature on the study of resource allocation and various policies have been derived under various settings (e.g. Reiman and Wang 2008; Jasin and Kumar 2012; Ferreira et al. 2018; Asadpour et al. 2020; Bumpensanti and Wang 2020; Vera and Banerjee 2019; Jiang et al. 2022). Notably, the intersection of pricing and resource allocation has also been studied, for example, in Chen et al. (2021b) and Vera et al. (2021). However, previous works have primarily focused on either the allocation decision or the pricing decision separately. In contrast, in our paper, we consider a two-stage process where we first make the pricing decision which affects the demand, and then make the allocation decision. This feature distinguishes our paper from previous works on (price-based) resource allocation.

Online Convex Optimization (OCO). OCO models a scenario where decisions are made iteratively, facing a series of convex loss functions, with the objective of minimizing cumulative regret over time (Shalev-Shwartz et al., 2012) or within certain budgets (Jenatton et al., 2016). In our work, we adopt zeroth-order methods in Agarwal et al. (2011) when we iterate within each local convexity interval. For a detailed review of classic and contemporary results on OCO, we kindly refer readers to Hazan (2016).

3 Preliminaries

In this section, we rigorously define the problem we are studying. We firstly formulate the offline version of the problem as a two-stage stochastic program in Section 3.1. The properties of its solution are presented in Section 3.2. We then develop the formulation

of the online version in Section 3.3, where the demand parameters are unknown. Finally, we present assumptions that are crucial to our online algorithm design by the end of this section.

3.1 Offline Problem Setting

We consider the following scenario where a retail company makes their decisions with the goal of maximizing its net profit. Suppose the company produces and sells identical products, and it has m warehouses and n retailers. In general, this company faces the following three problems on inventory, pricing and allocation:

1. What are the appropriate quantities each warehouse should load?
2. What is the optimal price that retailers should set?
3. How to allocate inventories from warehouses to stores in the face of heterogeneous supply costs?

To address these questions, we model the problem as a two-stage stochastic program.

(i) In Stage 1, the company makes *inventory* decisions $\vec{I} = [I_1, I_2, \dots, I_m]^\top$, where I_i represents the inventory level of Warehouse i . Each unit of inventory at Warehouse i incurs an inventory cost γ_i .

In addition, the company decides a uniform *price* p for the products.

(ii) In Stage 2, a stochastic demand $\vec{D} = [D_1, D_2, \dots, D_n]^\top$ is generated based on the price p , where D_j represents the demand at Retailer j . Then the products are allocated from warehouses to stores in order to fulfill the realized demand. Each unit of supply from Warehouse i to Retailer j incurs an allocation cost $C_{i,j}$, and each unit of fulfilled demand increases the total revenue by p .

The company aims to make the best (inventory, price) decisions that maximize their net profit, which can be formulated as the following optimization problem (where we equivalently minimize the negative net profit):

$$\min_{p, \vec{I}} \langle \vec{\gamma}, \vec{I} \rangle + \mathbb{E}_{\vec{D}} [g(\vec{I}, p, \vec{D}) | p] \quad (1)$$

where

$$\begin{aligned}
g(\vec{I}, p, \vec{D}) &= \min_{\mathbf{X} \in \mathbb{R}_+^{m \times n}} \sum_{i=1}^m \sum_{j=1}^n (-p + C_{i,j}) X_{i,j} \\
\text{s.t. } & X_{i,j} \geq 0, \forall i \in [m], j \in [n] \\
& \sum_{i=1}^m X_{i,j} \leq D_j, \forall j \in [n] \\
& \sum_{j=1}^n X_{i,j} \leq I_i, \forall i \in [m].
\end{aligned} \tag{2}$$

Here $X_{i,j}$ represents the quantity of inventories allocated from Warehouse i to Retailer j , and $g(\vec{I}, p, \vec{D})$ is the optimal objective value of the second-stage problem, which optimally allocates inventory \vec{I} to demand \vec{D} based on price p and cost parameters $\{C_{i,j}\}$.

It is worth noting that the distribution of demand \vec{D} is dependent on the price p , and the minimization over p and \vec{I} takes this into account. However, the solution to $g(\vec{I}, p, \vec{D})$ is not relevant to this dependence, as it is solved after the realization of \vec{D} .

We denote an optimal solution to Eq. (1) as (p^*, \vec{I}^*) , and denote an optimal solution to Eq. (2) as \mathbf{X}^* . Since Eq. (2) is a linear program, we may directly solve it with any off-the-shelf optimization tool. However, in order to solve Eq. (1), we have to know the distribution of \vec{D} and how it is dependent on the price p , both of which are not directly accessible from the seller's side as they do not have the full knowledge of the entire market. In the next subsection, we will discuss how we can “learn” the demand distributional function under mild assumptions.

3.2 Solution to $g(\vec{I}, p, \vec{D})$

Before we study the online problem, we first provide a solution to the optimization problem in Eq. (2) and investigate the properties of $g(\vec{I}, p, \vec{D})$. We firstly propose a greedy algorithm to solve Eq. (2).

We show the correctness of Algorithm 1 as follows.

Theorem 3.1. *The matrix \mathbf{X} returned by Algorithm 1 is an optimal solution to the optimization problem in Eq. (2).*

While the complete proof of Theorem 3.1 is relegated to Appendix A.1, we provide a high-level idea of the proof below. Notice that Eq. (2) presents a linear program (LP), and therefore strong duality holds. The proof is conducted in the following steps:

1. Construct the dual problem of Eq. (2).

Algorithm 1 Greedy $g(\vec{I}, p, \vec{D})$ solver

- 1: **Input:** Parameters $C_{i,j}, i \in [m], j \in [n]$; Variable $\vec{I} \in \mathbb{R}^m, p \in \mathbb{R}^+, \vec{D} \in \mathbb{R}^n$.
- 2: Sort $\{(i, j)\}_{i=1, j=1}^{i=m, j=n}$ tuples into $\{(i_k, j_k)\}_{k=1}^{mn}$ according to Algorithm 2.
- 3: Denote $C_{i_0, j_0} := 0$ and $C_{i_{mn+1}, j_{mn+1}} := p_{\max}$ as a complementary definition of $\{C_{i_k, j_k}\}$.
- 4: Find $K \in \{0\} \cup [mn]$ such that

$$C_{i_K, j_K} < p \leq C_{i_{K+1}, j_{K+1}} \quad (3)$$

- 5: Let $\tilde{I} \leftarrow \vec{I}, \tilde{D} \leftarrow \vec{D}, \mathbf{X} \leftarrow 0^{m \times n}$.
 - 6: **for** Epoch $k = 1, 2, 3, \dots, K$ **do**
 - 7: Set $X_{i_k, j_k} \leftarrow \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\}$.
 - 8: Update $\tilde{I}_{i_k} \leftarrow \tilde{I}_{i_k} - X_{i_k, j_k}$.
 - 9: Update $\tilde{D}_{j_k} \leftarrow \tilde{D}_{j_k} - X_{i_k, j_k}$.
 - 10: **end for**
 - 11: **RETURN** \mathbf{X} .
-

Algorithm 2 Sort tuples $\{(i, j)\}_{i=1, j=1}^{i=m, j=n}$ as follows:

- 1: **Input:** $\{(i, j)\}$ tuples.
 - 2: **for** each different pairs of tuples (i, j) and (i', j') **do**
 - 3: If $C_{i,j} < C_{i',j'}$, then $(i, j) \prec (i', j')$.
 - 4: If $C_{i,j} = C_{i',j'}$ and $i > i'$, then $(i, j) \prec (i', j')$.
 - 5: If $C_{i,j} = C_{i',j'}$, $i = i'$ and $j > j'$, then $(i, j) \prec (i', j')$.
 - 6: **end for**
 - 7: **Output:** $\{(i_k, j_k)\}_{k=1}^{mn}$.
-

2. Propose an algorithm that solves this dual problem and propose a feasible dual solution.
3. Show that the value of this dual solution equals that of the primal solution provided by Algorithm 1.

Next, we show the marginal convexity of $g(\vec{I}, p, \vec{D})$.

Lemma 3.2. *The function $g(\vec{I}, p, \vec{D})$ defined in Eq. (2) is marginally convex on \vec{I} and on \vec{D} .*

The key to proving Lemma 3.2 is to show that $g(\vec{I}, p, \vec{D})$ is the piecewise maximum over a group of linear functions. Please refer to Appendix A.2 for details.

3.3 Online Problem Setting

Due to insufficient knowledge of the actual demand distribution, the company might propose (p, \vec{I}) pairs that are suboptimal, leading to lower net profits compared to the optimal solution. However, the company has observations on the realized demand at each store, which enables them to estimate demand and subsequently enhance their decision-making. In what follows, we study the online decision-making problem of setting prices and managing inventory.

Denote p_t , \vec{I}_t and \vec{D}_t as the price, inventory and realized demand in each time period $t = 1, 2, \dots, T$, respectively. We make the following semi-parametric assumption on the demand model.

Assumption 3.3. Assume the realized demand is *linear* and *noisy*. Specifically, assume

$$\vec{D}_t = \vec{a} - \vec{b} \cdot p + \vec{N}_t. \quad (4)$$

Here $\vec{a}, \vec{b}, \vec{N}_t \in \mathbb{R}^n$ are the base demand, the price sensitivity parameter, and the market noise of the retailers' demand, respectively. Assume \vec{a}, \vec{b} are fixed, and \vec{N}_t are samples drawn from *identical and independent distributions* (i.i.d.) over time t , such that $\mathbb{E}[\vec{N}_t] = \vec{0}$.

Given the linear-and-noisy demand model in (4), we define the cost function that we aim to minimize. Denote

$$\begin{aligned} Q(\vec{I}, p) &:= \langle \vec{\gamma}, \vec{I} \rangle + \mathbb{E}_{\vec{N}_t}[g(\vec{I}, p, \vec{a} - \vec{b} \cdot p + \vec{N}_t)] \\ Q_t(\vec{I}, p) &:= \langle \vec{\gamma}, \vec{I} \rangle + g(\vec{I}, p, \vec{a} - \vec{b} \cdot p + \vec{N}_t) \end{aligned} \quad (5)$$

Based on the definition of functions Q and Q_t above, we know that $Q_t(\vec{I}, p)$ (and therefore $Q(\vec{I}, p)$) is marginally convex on \vec{I} , since $g(\vec{I}, p, \vec{D})$ is marginally convex on \vec{I} (according to Lemma 3.2) and $\langle \vec{\gamma}, \vec{I} \rangle$ is linear. Also, we have the following properties.

Lemma 3.4. Denote $C_{i_0, j_0} = 0$ and $C_{i_{mn+1}, j_{mn+1}} = p_{\max}$. For any $K \in \{0\} \cup [mn]$, function $Q_t(\vec{I}, p)$ is Lipschitz and marginally convex on p in range $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$.

We defer the proof of Lemma 3.4 to Appendix A.3. Furthermore, we get following results (in a non-trivial way).

Lemma 3.5. Define an optimistic cost function $W(p)$:

$$W(p) := \min_{\vec{I} \in \mathbb{R}_+^m} Q(\vec{I}, p). \quad (6)$$

$W(p)$ is Lipschitz. We denote its Lipschitz coefficient as L_W for future use. Also, for any $K \in \{0\} \cup [mn]$, function $W(p)$ is convex in range $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$.

The proof of Lemma 3.5 is relegated to Appendix A.4. Finally, we define the *regret* as the relative loss on net profit compared to that achieved by optimal decisions.

Algorithm 3 LCB Meta Algorithm

- 1: **Input:** m, n, T , sorted supply costs $\{C_{i_k, j_k}\}_{k=0}^{mn+1}$.
 - 2: **Initialization:** Agents \mathcal{A}_K (defined in Algorithm 4) and parameters $\hat{W}_K := 0, \Delta_K := +\infty, LCB_K := -\infty, K = 0, 1, 2, \dots, mn$, global parameters δ_K, n_0 .
 - 3: **while** $t \leq T$ **do**
 - 4: Let $\hat{K} := \operatorname{argmin}_{K \in [mn] \cup 0} LCB_K$.
 - 5: Run agent $\mathcal{A}_{\hat{K}}$ for one sub-epoch, and get updated \hat{W}_K and Δ_K
 - 6: Update $LCB_{K'} \leftarrow \hat{W}_{K'} - 34 \cdot \Delta_{K'}$ for all $K' = 0, 1, 2, \dots, mn$.
 - 7: **end while**
-

Definition 3.6 (Regret). At each time $t = 1, 2, \dots, T$, denote \vec{I}_t and p_t as the inventory and price decisions, respectively. Define

$$\text{Reg} := \sum_{t=1}^T Q(\vec{I}_t, p_t) - \min_{\vec{I}, p} Q(\vec{I}, p) \quad (7)$$

as the regret of decision sequence $\{(\vec{I}_t, p_t)\}_{t=1}^T$.

Before we conclude this section, we present some assumptions that are crucial to our algorithm design.

Assumption 3.7 (Boundedness). We assume boundedness on the norms of $\vec{\gamma}, \vec{a}, \vec{b}, \vec{I}$ and on price p . Specifically, there exist constants $\gamma_{\max}, a_{\max}, b_{\max}, I_{\max}, p_{\max}$ such that $\|\vec{\gamma}\|_{\infty} \leq \gamma_{\max}, \|\vec{a}\|_{\infty} \leq a_{\max}, \|\vec{b}\|_{\infty} \leq b_{\max}, \|\vec{I}\|_1 \leq I_{\max}$ and $p \in [0, p_{\max}]$. Without loss of generality, we assume $\gamma_{\max}, a_{\max}, b_{\max}, I_{\max}, p_{\max} \geq 1$.

Assumption 3.8 (Knowledge over parameters). We have *full* knowledge on the problem parameters $\vec{\gamma}, \{C_{i,j}\}_{i=1, j=1}^{m,n}$ and the boundedness parameters $\gamma_{\max}, a_{\max}, b_{\max}, I_{\max}, p_{\max}$ before $t = 0$. We have *no* prior knowledge on the model parameters \vec{a}, \vec{b} nor any information about the distribution of \vec{N}_t .

4 Algorithm

In this section, we present an online learning algorithm that proposes asymptotically optimal (inventory, price) decisions.

4.1 Algorithm Design Overview

We design a hierarchical algorithm to solve this problem. Algorithm 3 serves as the main algorithm, where we initialize $(mn + 1)$ agents \mathcal{A}_K for $K = 0, 1, \dots, mn$ and run each of

Algorithm 4 Optimization Agent K

- 1: **Input:** Price interval $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$, parameter δ_K, n_0, L_W .
 - 2: **Output:** \hat{W}_K, Δ_K (at the end of each sub-epoch)
 - 3: **Initialization:** $L_{K,1} = C_{i_K, j_K}, U_{K,1} = C_{i_{K+1}, j_{K+1}}, \hat{W}_K = 0, \Delta_K = +\infty$, and $a_{K,1} = \frac{3L_{K,1} + U_{K,1}}{4}, c_{K,1} = \frac{L_{K,1} + U_{K,1}}{2}, b_{K,1} = \frac{L_{K,1} + 3U_{K,1}}{4}$.
 - 4: **for** $p = a_{K,1}, b_{k,1}, c_{k,1}$, **do**
 - 5: Propose inventory $\vec{I}_0 := [1, 1, \dots, 1]^\top$ and price p .
 - 6: Record the marginal function of \vec{I} as $Q_0(\vec{I}, p)$.
 - 7: Find $I_{K,1,0}(p) := \operatorname{argmin}_{\vec{I}} Q_0(\vec{I}, p)$.
 - 8: **end for**
 - 9: **STAGE 1: Search for optimality asymptotically**
 - 10: (See Algorithm 5 for details)
 - 11: **STAGE 2: Complementary sampling to enhance Δ_K**
 - 12: (See Algorithm 6 for details)
 - 13: **STAGE 3: Pure local-exploitation**
 - 14: (See Algorithm 7 for details)
-

them for a period of time. The design of each agent \mathcal{A}_K is defined as Algorithm 4, which contains the following three stages:

- (i) In Stage 1 (presented as Algorithm 5), we search for the local optimal decision within each \mathcal{A}_K 's domain.
- (ii) In Stage 2 (presented as Algorithm 6), we gather sufficient number of samples to ensure the confidence bound of \mathcal{A}_K is converging at a proper rate.
- (iii) In Stage 3 (presented as Algorithm 7), we do pure exploitation on the local optimal of \mathcal{A}_K , while also updating the confidence bound.

In the following subsections, we will introduce each component of our algorithm design (including the meta-algorithm and each OCO agent) in details.

4.2 Meta-Algorithm: a Lower-Confidence-Bound (LCB) Strategy

Since we have full-information feedback over any decision w.r.t. \vec{I} , we may always propose greedy inventories without causing bias. However, we only have bandit feedback w.r.t. the price p , as we have no direct feedback on the prices we are not proposing. Therefore, we conduct online learning on the optimistic cost function $W(p) = \min_{\vec{I}} Q(\vec{I}, p)$.

Due to the piecewise convexity of $W(p)$, we divide the price range $[0, p_{\max}]$ into $(mn + 1)$ intervals $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}], K = 0, 1, \dots, mn$. Within each interval, we initialize an OCO agent \mathcal{A}_K that is responsible for converging to the local optimal. However, we cannot run

Algorithm 5 Agent \mathcal{A}_K Stage 1

```
1: for Epoch  $\tau = 1, 2, \dots, O(\log T)$ , do
2:   Let  $a_{K,\tau} = \frac{3L_{K,1} + U_{K,1}}{4}$ ,  $c_{K,\tau} = \frac{L_{K,1} + U_{K,1}}{2}$ ,  $b_{K,\tau} = \frac{L_{K,1} + 3U_{K,1}}{4}$ .
3:   for Sub-epoch  $s = 1, 2, \dots$  do
4:     Let sub-epoch length  $n_s := 2^s$ 
5:     Define a flag  $:= 0$  for error-bar update.
6:     for  $\hat{p}_\tau = a_{K,\tau}, b_{K,\tau}, c_{K,\tau}$  do
7:       for  $t = 1, 2, \dots, n_s$  do
8:         Propose decisions  $(\vec{I}_t, p_t) = (\vec{I}_{K,\tau,s-1}(\hat{p}_\tau), \hat{p}_\tau)$ .
9:         Observe and record the marginal function  $Q_t(\vec{I}, \hat{p}_\tau)$  with respect to  $\vec{I}$ .
10:      end for
11:      Define an aggregated function  $Q_{K,\tau,s}(\vec{I}, \hat{p}_\tau) := \frac{1}{n_s} \cdot \sum_{t=1}^{n_s} Q_t(\vec{I}, \hat{p}_\tau)$ .
12:      Define the empirical optimal inventory
13:         $I_{K,\tau,s}(\hat{p}_\tau) := \operatorname{argmin}_{\vec{I}} Q_{K,\tau,s}(\vec{I}, \hat{p}_\tau)$ .
14:      Denote  $\hat{Q}_{K,\tau,s,\hat{p}_\tau} := Q_{k,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau)$ , and  $\Delta_{K,\tau,s} := \frac{\delta_K}{2\sqrt{n_s}}$ .
15:    end for
16:    if  $\hat{Q}_{K,\tau,s,a_{K,\tau}} > \hat{Q}_{K,\tau,s,b_{K,\tau}} + 4\Delta_{K,\tau,s}$  then
17:      Update  $L_{K,\tau+1} \leftarrow a_{K,\tau}$ ,  $U_{K,\tau+1} \leftarrow U_{K,\tau}$ , flag  $\leftarrow 1$ .
18:    else if  $\hat{Q}_{K,\tau,s,a_{K,\tau}} < \hat{Q}_{K,\tau,s,b_{K,\tau}} - 4\Delta_{K,\tau,s}$  then
19:      Update  $L_{K,\tau+1} \leftarrow L_{K,\tau}$ ,  $U_{K,\tau+1} \leftarrow b_{K,\tau}$ , flag  $\leftarrow 1$ .
20:    else if  $\hat{Q}_{K,\tau,s,c_{K,\tau}} < \hat{Q}_{K,\tau,s,a_{K,\tau}} - 4\Delta_{K,\tau,s}$  then
21:      Update  $L_{K,\tau+1} \leftarrow a_{K,\tau}$ ,  $U_{K,\tau+1} \leftarrow U_{K,\tau}$ , flag  $\leftarrow 1$ .
22:    else if  $\hat{Q}_{K,\tau,s,c_{K,\tau}} < \hat{Q}_{K,\tau,s,b_{K,\tau}} - 4\Delta_{K,\tau,s}$  then
23:      Update  $L_{K,\tau+1} \leftarrow L_{K,\tau}$ ,  $b_{K,\tau+1} \leftarrow U_{K,\tau}$ , flag  $\leftarrow 1$ .
24:    end if
25:    if flag  $== 1$  then
26:      if  $U_{K,\tau+1} - L_{K,\tau+1} > \frac{1}{T}$  then
27:        Continue to Epoch  $\tau + 1$  (without updating  $\hat{W}_K$  or  $\Delta_K$ ).
28:      else
29:        Set  $\hat{p}_K^* \leftarrow C_{K,\tau}$ ,  $\hat{I}_{K,0}^* \leftarrow \vec{I}_{K,\tau,s}(C_{K,\tau})$ .
30:        Break to STAGE 2 (without updating  $\hat{W}_K$  or  $\Delta_K$ ).
31:      end if
32:    else if  $\Delta_{K,\tau,s-1} < \Delta_K$  then
33:      Update  $\Delta_K \leftarrow \Delta_{K,\tau,s-1}$ ,  $\hat{W}_K \leftarrow \min_{\hat{p}_\tau \in \{a_{K,\tau}, c_{K,\tau}, b_{K,\tau}\}} \hat{Q}_{K,\tau,s-1,\hat{p}_\tau}$ .
34:    end if
35:  end for
```

Algorithm 6 Agent \mathcal{A}_K Stage 2 (a sub-epoch totally)

- 1: **Initialization:** Set $\hat{p}_K^* := C_{K,\tau}, \hat{I}_{K,0}^* = \vec{I}_{K,\tau,s}(C_{K,\tau})$ from Stage 1, and $\check{C}_K := \log_{4/3} p_{\max} + 1$.
 - 2: **if** $\Delta_K < +\infty$ **then**
 - 3: Define $N_{K,2} := \frac{4\delta_K^2}{\Delta_K}$
 - 4: **else**
 - 5: Define $N_{K,2} := n_0 = 6(\log_{4/3} T + \check{C}_K)$
 - 6: **end if**
 - 7: Let $r_K := \log_2(N_{K,2})$
 - 8: **for** $r = 1, 2, \dots, r_K$ **do**
 - 9: **for** $t = 1, 2, \dots, m_r := 2^r$ **do**
 - 10: Propose decisions $(\vec{I}_t, p_t) = (\hat{I}_{K,r-1}^*, \hat{p}_K^*)$.
 - 11: Observe and record the marginal function $Q_t(\vec{I}, \hat{p}_K^*)$ with respect to \vec{I} .
 - 12: **end for**
 - 13: Set $\hat{I}_{K,r}^* := \operatorname{argmin}_{\vec{I}} \bar{Q}_{K,r}(\vec{I}, \hat{p}_K^*)$, where $\bar{Q}_{K,r}(\vec{I}, \hat{p}_K^*) := \frac{1}{m_r} \cdot \sum_{t=1}^T Q_t(\vec{I}, \hat{p}_K^*)$.
 - 14: **end for**
 - 15: Denote $\vec{I}_K^* := \hat{I}_{K,r_K}^*, \hat{Q}_K^* := \bar{Q}_{K,r_K}(\vec{I}_K^*, \hat{p}_K^*)$.
 - 16: **Update** $\hat{W}_K \leftarrow \hat{Q}_K^* - L_W \cdot \frac{1}{T}$ **and** $\Delta_K \leftarrow \frac{\delta_K}{\sqrt{N_{K,2}}}$.
-

multiple OCO agents simultaneously. Therefore, we require a meta-algorithm that serves as a manager over these agents and determine which \mathcal{A}_K to run at each time, so as to locate the optimal price with the least cumulative regret.

To achieve this, we develop a lower-confidence-bound (LCB) meta-algorithm as shown in Algorithm 3. We firstly ask each \mathcal{A}_K agent to maintain a confidence bound $[\hat{W}_K - 34\Delta_K, \hat{W}_K + 34\Delta_K]$ of its local optimal. Given this, the meta-algorithm then selects the agent K that *minimizes* the lower confidence bound. As we further show that $\Delta_K \approx O(\sqrt{1/T_K})$ where T_K is the total time periods that \mathcal{A}_K has been running so far, we may upper bound the cumulative regret as $O(\sqrt{Tmn})$.

4.3 Agent \mathcal{A}_K : a Zeroth-Order Optimizer

As described in Section 4.2, we divide the price range $[0, p_{\max}]$ into $(mn + 1)$ intervals $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}], K = 0, 1, 2, \dots, mn$, within each of which the objective function $W(p)$ is convex. We then assign an agent \mathcal{A}_K to each interval, conducting online convex optimization (OCO) locally. We require the agent \mathcal{A}_K to learn and converge to the local optimal $p_K^* := \operatorname{argmin}_{p \in [C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]} W(p)$ over time, while also maintaining a valid error bar $[\hat{W}_K - 34\Delta_K, \hat{W}_K + 34\Delta_K]$ that contains $W(p_K^*)$ with high probability. To achieve

Algorithm 7 Agent \mathcal{A}_K Stage 3 (each t as a sub-epoch)

- 1: Inherit $\hat{p}_K^*, \hat{I}_{K,r_K}^*, \bar{Q}_{K,r_K}(\vec{I}, \hat{p}_K^*)$ from Algorithm 6.
 - 2: Denote $\vec{I}_K^* := \hat{I}_{K,r_K}^*, \hat{Q}_K^* := \bar{Q}_{K,r_K}(\vec{I}_K^*, p_K^*)$.
 - 3: Let $N_{K,3} \leftarrow N_{K,2}$ as its initialization.
 - 4: **while** $t \leq T$ **do**
 - 5: Propose decisions $(\vec{I}_t, p'_t) = (\vec{I}_K^*, \hat{p}_K^*)$.
 - 6: Update $\hat{Q}_K^* \leftarrow \frac{N_{K,3} \hat{Q}_K^* + Q_t(\vec{I}_t, p'_t)}{N_{K,3} + 1}$, and $N_{K,3} \leftarrow N_{K,3} + 1$
 - 7: **Update** $\hat{W}_K \leftarrow \hat{Q}_K^* - L_W \cdot \frac{1}{T}$ **and** $\Delta_K \leftarrow \frac{\delta_K}{\sqrt{N_{K,3}}}$.
 - 8: **end while**
-

the optimal regret, we rely on the following properties of \mathcal{A}_K :

- (a) The cumulative sub-regret of \mathcal{A}_K , i.e. performance suboptimality compared with $W(p_K^*)$, is bounded by $\tilde{O}(\sqrt{T_K})$ as an optimal rate of OCO (if we have run \mathcal{A}_K for T_K times so far).
- (b) The error bar Δ_K is bounded by $\tilde{O}(\sqrt{\frac{1}{T_K}})$ as a requirement of the meta-algorithm.

In this work, we propose Algorithm 4 as the framework of \mathcal{A}_K . In what follows, we elaborate each component of \mathcal{A}_K 's algorithmic design in detail.

Horizontal search space for p_K^* . In the design of Stage 1 algorithm as presented in Algorithm 5, we adopt the framework of zeroth-order online convex optimization. Specifically, we establish an epoch-based update rule of the search space of local optimal p_K^* . The search space (interval) for Epoch $\tau = 1, 2, \dots$ is denoted as $[L_{K,\tau}, U_{K,\tau}]$. Within each epoch, we divide the time horizon into a series of *doubling sub-epochs* to gather samples for $W(a), W(b), W(c)$ where a, b, c are the three quarter points. By the end of each sub-epoch, we update the estimates and examine whether their estimation error bar is *separable* according to certain rules. As we keep doubling the size of sub-epochs, the estimation error bars are shrinking exponentially until they are separated. Then we reduce the search space by one quarter and proceed to Epoch $\tau + 1$ Sub-Epoch 1. When the search space is as sufficiently small as $O(1/T)$, we stop searching and proceed to Stage 2.

Vertical uncertainty bound for $W(p_K^*)$. In Stage 1, we maintain an error bar Δ_K as the confidence bound of estimating each local optimal $W(p_K^*)$. We show that the error bar has a size of $\tilde{O}(\frac{1}{\sqrt{T_K}})$ if we have run \mathcal{A}_K for T_K times so far. In addition to the statistical concentrations, another intuition of this fact comes from Lemma A.9: If we cannot distinguish among the value of each quaternary of a convex function due to their uncertainty, then we may construct a comparable uncertainty bound for the optimal.

Complementary sampling to enhance Δ_K . It is worth noting that $\Delta_K < +\infty$ does not exist for granted even after Stage 1. This is because we cannot update Δ_K when the search space $[L_K, U_K]$ is updated, i.e., no simultaneous “horizontal converging” and “vertical converging”. As a consequence, if we are very “lucky” that we can always reduce the search space in the *first* sub-epoch of every epoch until $U_{K,\tau} - L_{K,\tau} \leq 1/T$, then we will have $\Delta_K = +\infty$ until Stage 2. We settle this issue in two approaches: (1) We upper bound the time periods before any $\Delta_K < +\infty$ by $O(\log T)$ based on the Pigeon-Hole Theorem. (2) We have Stage 2 (Algorithm 6) as a complementary sampling stage without causing excessive regret. By the end of Stage 2, we will have an ideal error bar for each agent \mathcal{A}_K .

Pure local exploitation contributing to global LCB. From Agent \mathcal{A}_K ’s perspective, it runs pure exploitation in Stage 3 (Algorithm 7) without causing extra sub-regret. However, it still keeps updating the estimates of \hat{W}_K and Δ_K to facilitate the LCB meta-algorithm.

4.4 Technical Novelty

To the best of our knowledge, we are the first to propose a methodology undergoing “horizontal-and-vertical” convergence simultaneously. In contrast, existing works adopt either “vertical convergence” such as bandits algorithms (which allow non-convexity of objective functions but cannot achieve $O(\sqrt{T})$ regret even with smoothness assumptions), or “horizontal convergence” which is applicable to many online planning and optimization scenarios but requires global convexity assumptions.

5 Analysis

In this section, we provide the theoretical analysis on the performance of Algorithm 3. We firstly propose our main theorem that upper bounds the cumulative regret.

Theorem 5.1 (Regret). *Let $n_0 = 6(\log_{4/3} T + \check{C}_K)$ where $\check{C}_K := \log_{4/3} p_{\max} + 1$ and $\delta_K = \sqrt{2 \log \frac{48(2mn+1)T}{\epsilon}} \cdot \max\{p_{\max}, \gamma_{\max}\} I_{\max}$. Algorithm 3 guarantees an $\tilde{O}(\sqrt{Tmn} + mn)$ regret with probability at least $1 - \epsilon$. Here $\tilde{O}(\cdot)$ omits the dependence on $\log \frac{1}{\epsilon}$ and $\log T$.*

This regret rate is near-optimal with respect to T , as it matches the information-theoretic lower bound of $\Omega(\sqrt{T})$ (see Broder and Rusmevichientong, 2012, Theorem 3.1), which describes a special case as $m = n = 1$ and $\gamma_1 = C_{1,1} = 0$ in our setting.

5.1 Sub-Regret and Confidence Bound of \mathcal{A}_K

In this subsection, we present two lemmas that show the convergence of each agent \mathcal{A}_K . Specifically, Lemma 5.2 shows the “horizontal convergence” of (inventory, price) decisions

towards the local optimal. Lemma 5.3 shows the “vertical convergence” of estimation error Δ_K such that we are maintaining and updating a valid lower-confidence bound for the majority of time.

Lemma 5.2 (Sub-regret of every \mathcal{A}_K). *For agent \mathcal{A}_K defined as Algorithm 4 that has been running for T_K time periods so far, the cumulative sub-regret is bounded by:*

$$SReg_K = \sum_{t_K=1}^{T_K} Q(\vec{I}_{t_K}, P_{t_K}) - W(p_K^*) = \tilde{O}(\sqrt{T_K}). \quad (8)$$

The proof of Lemma 5.2 is relegated to Appendix A.5.

Lemma 5.3 (Validity of Δ_K). *For any agent \mathcal{A}_K has been running for T_K time periods with $T_K \geq 6(\log_{4/3} T + \check{C}_K)$, we have $\Delta_K = \tilde{O}(\frac{1}{\sqrt{T_K}})$.*

The proof of Lemma 5.3 is presented in Appendix A.6. From Lemma 5.3, we directly get the following Corollary 5.4.

Corollary 5.4. *After at most $N_0 := 6(mn + 1)(\log_{4/3} T + \check{C}_K)$ time periods, there does not exist any $K \in [mn] \cup \{0\}$ such that $\Delta_K = +\infty$.*

Furthermore, due to the piecewise convexity of $W(p)$ and the convergence rate of Δ_K , combining with Corollary 5.4, we have the following lemma.

Lemma 5.5. *At any time $t > N_0 := 6(mn + 1)(\log_{4/3} T + \check{C}_K)$, we have*

$$LCB_K \leq W(p_K^*) \text{ and } LCB_K \geq W(p_K^*) - 35\Delta_K - \frac{2L_W}{T}. \quad (9)$$

Here L_W is the Lipschitz coefficient of $W(p)$.

The proof details of Lemma 5.5 is displayed in Appendix A.7.

5.2 Overall Regret

With the help of all the lemmas presented above, we are now ready to provide an upper bound on the total regret.

$$\begin{aligned}
\text{Reg}_T &:= \sum_{K=0}^{mn} \text{SReg}_K + \sum_{t=1}^T W(P_{K_t}^*) - W(P^*) \\
&\leq \sum_{K=0}^{mn} \tilde{O}(\sqrt{T_K}) + \sum_{t=N_0}^T (35\Delta_K + \frac{2L_W}{T}) + 6(mn+1)(\log_{4/3} T + \check{C}_K) \cdot a_{\max} \cdot p_{\max} \\
&\leq \tilde{O}\left(\sqrt{\left(\sum_{K=0}^{mn} 1\right)\left(\sum_{K=0}^{mn} T_K\right)} + \sum_{t=N_0}^T \Delta_{K_t} + T \cdot \frac{L_W}{T} + mn\right) \\
&= \tilde{O}\left(\sqrt{mn \cdot T} + \sum_{K=0}^{mn} \sum_{t_K=1}^{T_K} \frac{1}{\sqrt{t_K}} + mn\right) \\
&= \tilde{O}\left(\sqrt{Tmn} + \sum_{K=0}^{mn} 2\sqrt{T_K} + mn\right) \\
&= \tilde{O}\left(\sqrt{Tmn} + mn\right).
\end{aligned} \tag{10}$$

6 Discussion

Here we discuss some potential extensions of our work, serving as heuristics towards a broader field of research.

Generalization to non-linear demands. We assume the demand \vec{D} is a linear function of price p , which is a widely-used assumption (see [LaFrance, 1985](#)). Meanwhile, we still want to generalize our methodologies to a broader family of non-linear demands. Notice that the second-stage allocation problem defined by Eq. (2) does not involve the formulation of demand w.r.t. p . Therefore, we may still divide the price space into $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$ intervals, and run an individual online optimization agent within each interval. With a similar analysis, we can achieve an $\tilde{O}(T^\alpha(mn)^{1-\alpha})$ regret, where $\alpha \geq 1/2$ is dependent on the demand family we assume. On the other hand, by selecting $m = n = 1$ and $C_{i,j} = 0$, we may have a lower bound at $\Omega(T^\alpha)$.

Generalization to censored demands. In this work, we consider a warehouse-retailer setting where the demand orders are realized and informed to the suppliers *before* they are served. However, there exists another supply-demand relationship, such as groceries and wholesales, where the realized demands are revealed only *after* the resources are delivered from the supply side to the demand side as a preparation. In that case, we should estimate the prospective demand and carefully balance the allocation among individuals in each

side respectively, which goes much beyond a greedy allocation scheme as we solve Eq. (2). Besides, the realized demand might be *censored* when supply shortage occurs, making the problem more challenging. Therefore, we expect future investigations toward that new problem.

Pricing and service fairness. Our model maintains fairness in the pricing process by offering the same price to all consumers. However, while the greedy policy for resource allocation is reasonable, widely adopted, and analytically optimal, it leads to differentiated service levels among consumers. We anticipate future research focused on ensuring fairness in service levels during resource allocation.

7 Conclusion

In this paper, we study an online learning problem under the framework of pricing and allocation, where we make joint pricing and inventory decisions and allocate supplies to fulfill demands over time. To solve this non-convex problem, we propose a hierarchical algorithm which incorporates an LCB meta-algorithm over multiple local OCO agents. Our analysis shows that it guarantees an $\tilde{O}(\sqrt{Tmn} + mn)$ regret, which is optimal with respect to T as it matches the existing lower bound. Our work sheds light on the cross-disciplinary research of machine learning and operations research, especially from an online perspective.

References

- Agarwal, A., Foster, D. P., Hsu, D. J., Kakade, S. M., and Rakhlin, A. (2011). Stochastic convex optimization with bandit feedback. *Advances in Neural Information Processing Systems*, 24.
- Asadpour, A., Wang, X., and Zhang, J. (2020). Online resource allocation with limited flexibility. *Management Science*, 66(2):642–666.
- Baby, D., Xu, J., and Wang, Y.-X. (2023). Non-stationary contextual pricing with safety constraints. *Transactions on Machine Learning Research*.
- Ban, G.-Y. and Keskin, N. B. (2021). Personalized dynamic pricing with machine learning: High-dimensional features and heterogeneous elasticity. *Management Science*, 67(9):5549–5568.
- Besbes, O. and Zeevi, A. (2009). Dynamic pricing without knowing the demand function: Risk bounds and near-optimal algorithms. *Operations Research*, 57(6):1407–1420.
- Broder, J. and Rusmevichientong, P. (2012). Dynamic pricing under a general parametric choice model. *Operations Research*, 60(4):965–980.

- Bumpensanti, P. and Wang, H. (2020). A re-solving heuristic with uniformly bounded loss for network revenue management. *Management Science*, 66(7):2993–3009.
- Chen, B., Chao, X., and Ahn, H.-S. (2019). Coordinating pricing and inventory replenishment with nonparametric demand learning. *Operations Research*, 67(4):1035–1052.
- Chen, B., Chao, X., and Shi, C. (2021a). Nonparametric learning algorithms for joint pricing and inventory control with lost sales and censored demand. *Mathematics of Operations Research*, 46(2):726–756.
- Chen, B., Chao, X., and Wang, Y. (2020). Data-based dynamic pricing and inventory control with censored demand and limited price changes. *Operations Research*, 68(5):1445–1456.
- Chen, B., Wang, Y., and Zhou, Y. (2023). Optimal policies for dynamic pricing and inventory control with nonparametric censored demands. *Management Science*.
- Chen, N. and Gallego, G. (2021). Nonparametric pricing analytics with customer covariates. *Operations Research*.
- Chen, Q., Jasin, S., and Duenyas, I. (2021b). Joint learning and optimization of multi-product pricing with finite resource capacity and unknown demand parameters. *Operations Research*, 69(2):560–573.
- Chen, X., Zhang, X., and Zhou, Y. (2021c). Fairness-aware online price discrimination with nonparametric demand models. *arXiv preprint arXiv:2111.08221*.
- Choi, Y.-G., Kim, G.-S., Choi, Y., Cho, W., Paik, M. C., and Oh, M.-h. (2023). Semi-parametric contextual pricing algorithm using cox proportional hazards model. In *International Conference on Machine Learning*, pages 5771–5786. PMLR.
- Cohen, M. C., Elmachtoub, A. N., and Lei, X. (2022). Price discrimination with fairness constraints. *Management Science*.
- Cohen, M. C., Lobel, I., and Paes Leme, R. (2020). Feature-based dynamic pricing. *Management Science*, 66(11):4921–4943.
- Cohen, M. C., Miao, S., and Wang, Y. (2021). Dynamic pricing with fairness constraints. *Available at SSRN 3930622*.
- Cournot, A. A. (1897). *Researches into the Mathematical Principles of the Theory of Wealth*. Macmillan.
- Eyster, E., Madarász, K., and Michailat, P. (2021). Pricing under fairness concerns. *Journal of the European Economic Association*, 19(3):1853–1898.

- Fan, J., Guo, Y., and Yu, M. (2021). Policy optimization using semiparametric models for dynamic pricing. *arXiv preprint arXiv:2109.06368*.
- Ferreira, K. J., Simchi-Levi, D., and Wang, H. (2018). Online network revenue management using thompson sampling. *Operations research*, 66(6):1586–1602.
- Hazan, E. (2016). Introduction to online convex optimization. *Foundations and Trends in Optimization*, 2(3-4):157–325.
- Jasin, S. and Kumar, S. (2012). A re-solving heuristic with bounded revenue loss for network revenue management with customer choice. *Mathematics of Operations Research*, 37(2):313–345.
- Javanmard, A. and Nazerzadeh, H. (2019). Dynamic pricing in high-dimensions. *The Journal of Machine Learning Research*, 20(1):315–363.
- Jenatton, R., Huang, J., and Archambeau, C. (2016). Adaptive algorithms for online convex optimization with long-term constraints. In *International Conference on Machine Learning*, pages 402–411. PMLR.
- Jia, H., Shi, C., and Shen, S. (2022). Online learning and pricing with reusable resources: Linear bandits with sub-exponential rewards. In *International Conference on Machine Learning*, pages 10135–10160. PMLR.
- Jiang, J., Ma, W., and Zhang, J. (2022). Degeneracy is ok: Logarithmic regret for network revenue management with indiscrete distributions. *arXiv preprint arXiv:2210.07996*.
- Karan, A., Balepur, N., and Sundaram, H. (2024). Designing fair systems for consumers to exploit personalized pricing. *arXiv preprint arXiv:2409.02777*.
- Keskin, N. B., Li, Y., and Song, J.-S. (2022). Data-driven dynamic pricing and ordering with perishable inventory in a changing environment. *Management Science*, 68(3):1938–1958.
- Kleinberg, R. and Leighton, T. (2003). The value of knowing a demand curve: Bounds on regret for online posted-price auctions. In *IEEE Symposium on Foundations of Computer Science (FOCS-03)*, pages 594–605. IEEE.
- LaFrance, J. T. (1985). Linear demand functions in theory and practice. *Journal of Economic theory*, 37(1):147–166.
- Leme, R. P., Sivan, B., Teng, Y., and Worah, P. (2021). Learning to price against a moving target. In *International Conference on Machine Learning*, pages 6223–6232. PMLR.
- Luo, Y., Sun, W. W., et al. (2021). Distribution-free contextual dynamic pricing. *arXiv preprint arXiv:2109.07340*.

- Luo, Y., Sun, W. W., and Liu, Y. (2022). Contextual dynamic pricing with unknown noise: Explore-then-ucb strategy and improved regrets. In *Advances in Neural Information Processing Systems*.
- Reiman, M. I. and Wang, Q. (2008). An asymptotically optimal policy for a quantity-based network revenue management problem. *Mathematics of Operations Research*, 33(2):257–282.
- Shalev-Shwartz, S. et al. (2012). Online learning and online convex optimization. *Foundations and Trends® in Machine Learning*, 4(2):107–194.
- Simchi-Levi, D. and Wang, C. (2023). Pricing experimental design: causal effect, expected revenue and tail risk. In *International Conference on Machine Learning*, pages 31788–31799. PMLR.
- Tullii, M., Gaucher, S., Merlis, N., and Perchet, V. (2024). Improved algorithms for contextual dynamic pricing. *arXiv preprint arXiv:2406.11316*.
- Vera, A. and Banerjee, S. (2019). The bayesian prophet: A low-regret framework for online decision making. *ACM SIGMETRICS Performance Evaluation Review*, 47(1):81–82.
- Vera, A., Banerjee, S., and Gurvich, I. (2021). Online allocation and pricing: Constant regret via bellman inequalities. *Operations Research*, 69(3):821–840.
- Wang, H., Talluri, K., and Li, X. (2021a). On dynamic pricing with covariates. *arXiv preprint arXiv:2112.13254*.
- Wang, Y., Chen, B., and Simchi-Levi, D. (2021b). Multimodal dynamic pricing. *Management Science*.
- Xu, J., Qiao, D., and Wang, Y.-X. (2023). Doubly fair dynamic pricing. In *International Conference on Artificial Intelligence and Statistics*, pages 9941–9975. PMLR.
- Xu, J. and Wang, Y.-X. (2021). Logarithmic regret in feature-based dynamic pricing. *Advances in Neural Information Processing Systems*, 34.
- Xu, J. and Wang, Y.-X. (2022). Towards agnostic feature-based dynamic pricing: Linear policies vs linear valuation with unknown noise. *International Conference on Artificial Intelligence and Statistics (AISTATS)*.
- Xu, J. and Wang, Y.-X. (2024). Pricing with contextual elasticity and heteroscedastic valuation. In *Forty-first International Conference on Machine Learning*.

A Proof Details

A.1 Proof of Theorem 3.1

Proof. We conduct the proof according to the proof sketch displayed after Theorem 3.1 in the main paper.

Firstly, we construct the dual problem of Eq. (2). Here we denote $\tilde{C}_{i,j}(p) := -p + C_{i,j}$, and we know that the sorting of $\tilde{C}_{i,j}$ is identical to that of $C_{i,j}$ for any $p \in \mathbb{R}$. The Lagrangian can be written as

$$\begin{aligned} L(X, A, \mu, \lambda) &:= \sum_{i=1}^m \sum_{j=1}^n \tilde{C}_{i,j} X_{i,j} - \sum_{i=1}^m \sum_{j=1}^n A_{i,j} X_{i,j} + \sum_{j=1}^n \mu_j (\sum_{i=1}^m X_{i,j} - D_j) + \sum_{i=1}^m \lambda_i (\sum_{j=1}^n X_{i,j} - I_i) \\ &= \sum_i \sum_j (\tilde{C}_{i,j} - A_{i,j} + \mu_j + \lambda_i) X_{i,j} - \sum_j \mu_j D_j - \sum_i \lambda_i I_i. \end{aligned} \quad (11)$$

Here $A \in \mathbb{R}_+^{m \times n}$, $\mu \in \mathbb{R}_+^n$, $\lambda \in \mathbb{R}_+^m$. Therefore, the primal problem can be written as $\min_X \max_{A, \mu, \lambda} L(X, A, \mu, \lambda)$, and the dual problem is:

$$\begin{aligned} &\max_{A, \mu, \lambda} \min_X L(X, A, \mu, \lambda) \\ &= \begin{cases} -\sum_j \mu_j D_j - \sum_i \lambda_i I_i & \text{if } \tilde{C}_{i,j} - A_{i,j} + \mu_j + \lambda_i = 0 \text{ or } X_{i,j} = 0, \forall i \in [m], j \in [n] \\ -\infty & \text{otherwise.} \end{cases} \end{aligned} \quad (12)$$

This comes from the complementary slackness. Since $\tilde{C}_{i_k, j_k} < 0$ for $k \leq K$, the condition of $\tilde{C}_{i_k, j_k} - A_{i_k, j_k} + \mu_{j_k} + \lambda_{i_k} = 0$ is not necessarily satisfiable for those $k \in [K]$. Given this, the dual problem can be equivalently written as

$$\begin{aligned} &-\min_{\mu, \lambda} \sum_{j=1}^n \mu_j D_j + \sum_{i=1}^m \lambda_i I_i \\ &\text{s.t. } \tilde{C}_{i_k, j_k} + \mu_{j_k} + \lambda_{i_k} \geq 0, \forall k \leq K. \end{aligned} \quad (13)$$

Denote $h(\mu, \lambda; \vec{I}, \vec{D}) := \sum_{j=1}^n \mu_j D_j + \sum_{i=1}^m \lambda_i I_i$. Also, define a new equation

$$\tilde{h}(\mu, \lambda; \vec{I}, \vec{D}, \vec{Q}) := \sum_{j=1}^n \mu_j \tilde{D}_j + \sum_{i=1}^m \lambda_i \tilde{I}_i + \sum_{k=1}^{mn} Q_k (\lambda_{i_k} + \mu_{j_k}). \quad (14)$$

By initializing $\vec{Q} = 0$ and $\vec{I} = \vec{I}, \vec{D} = \vec{D}$, we know that $h(\mu, \lambda; \vec{I}, \vec{D}) = \tilde{h}(\mu, \lambda; \vec{I}, \vec{D}, \vec{Q})$. In the following, we conduct a series of equivalent transformations on \tilde{h} , which will further induce an algorithm to solve the dual problem.

Notice that

Algorithm 8 Equivalent transformation on \tilde{h}

Initialization: $\vec{Q} = 0, \vec{I} = \vec{I}, \vec{D} = \vec{D}$.
for $k = 1, 2, \dots, K$, **do**
 Set $Q_k \leftarrow \min\{\tilde{D}_{j_k}, \tilde{I}_{i_k}\}$.
 Update $\tilde{I}_{i_k} \leftarrow \tilde{I}_{i_k} - Q_k$.
 Update $\tilde{D}_{j_k} \leftarrow \tilde{D}_{j_k} - Q_k$.
end for

- i The value of $\tilde{h}(\mu, \lambda; \vec{I}, \vec{D}, \vec{Q})$ is unchanged over the processing of Algorithm 8.
- ii The $\{Q_k\}_{k=1}^{mn}$ series we get from Algorithm 8 is identical to the $\{X_{i_k, j_k}\}_{k=1}^{mn}$ series from Algorithm 1.

In the following, we focus on proving this lemma

Lemma A.1 (Feasible dual solutions). *The equation set of $\lambda \in \mathbb{R}_+^m, \mu \in \mathbb{R}_+^n$*

$$\begin{cases} \mu_{j_k} + \lambda_{i_k} = -\tilde{C}_{i_k, j_k}, & \text{for } k \in \{k \leq K, Q_k \neq 0\} \\ \mu_{j_k} = \lambda_{i_k} = 0, & \text{otherwise.} \end{cases} \quad (15)$$

has a feasible solution.

Before proving Lemma A.1, we firstly propose and prove the following series of lemma.

Lemma A.2. *After the operations of Algorithm 8, the total number of k such that $Q_k \neq 0$ does not exceed $\min\{K, m + n\}$.*

Proof of Lemma A.2. According to Algorithm 8, the operation at time k leads to either $\tilde{I}_{i_k} = 0$ or $\tilde{D}_{j_k} = 0$. On the one hand, there are at most k times of operations. On the other hand, there are at most $(m + n)$ number of \tilde{I}_j and \tilde{D}_j that can be reduced to 0. ■

Lemma A.3. *For $k_1, k_2 \in [K], k_1 < k_2$, s.t. $Q_{k_1} \neq 0, Q_{k_2} \neq 0, i_{k_1} = i_{k_2}$, then for any $k > k_1$ we have either $Q_k = 0$ or $j_k \neq j_{k_1}$.*

Similarly, for $k_3, k_4 \in [K], k_3 < k_4$, s.t. $Q_{k_3} \neq 0, Q_{k_4} \neq 0, j_{k_3} = j_{k_4}$, then for any $k > k_3$ we have either $Q_{k_3} = 0$ or $i_k \neq i_{k_3}$.

Proof of Lemma A.3. We only show the first case here, and the second case is by symmetry. Consider $\tilde{I}_{i_{k_2}}$ by the start of period $k = k_2$: Since $Q_{k_2} \neq 0$, we have $\tilde{I}_{i_{k_2}} > 0$ at that time. Also, since $k_1 < k_2$, we have $\tilde{I}_{i_{k_1}} > 0$ by the end of period $k = k_1$. As a result, we have

$\tilde{D}_{j_{k_1}} = 0$ by the end of period $k = k_1$. Therefore, for any $k > k_1$ such that $j_k = j_{k_1}$, we have $\tilde{D}_{j_k} = 0$ and therefore $Q_k = 0$. \blacksquare

Lemma A.4. *Define a bipartite graph $G(V, E)$, where $V = V_1 \cup V_2, V_1 = [m], V_2 = [n]$ and $E := \{(i_k, j_k) \text{ for all } k \text{ s.t. } Q_k \neq 0\}$. There exists a series of independent trees $\{T_l\}_{l=1}^t$ such that $G = T_1 \cup T_2 \cup \dots \cup T_t$. Here two independent trees $T_i, T_j \subset G$ satisfies $T_i \cap T_j = \emptyset$.*

Proof of Lemma A.4. We sufficiently show that there exists no loop in this bipartite graph $G(V, E)$. In fact, according to Lemma A.3, for any edge $(i_{k_1}, j_{k_1}) \in E$, either i_{k_1} or j_{k_1} is a leaf node. On the other hand, there exists no leaf node in any loop. Therefore, no loop exists in this graph. \blacksquare

With the help of Lemma A.2, Lemma A.3, Lemma A.4, we may process to prove Lemma A.1.

Proof of Lemma A.1. Based on Lemma A.4, the set of tuples $\{(i_k, j_k), k \in [K] \text{ s.t. } Q_k \neq 0\}$ can be separated as a series of disjoint "tree" sets. Therefore, we only need to prove Lemma A.1 holds on a connected "tree" set. Denote this tree as $T := G_T(V_T, E_T)$, where $t = |T|$ is the size (i.e. number of nodes) of tree T , $E_T := \{(i_{k_\tau}, j_{k_\tau}), \tau = 1, 2, \dots, t, 1 \leq k_1 \leq k_2 \leq \dots \leq k_t \leq K\}$, and $V_T := \{i_{k_\tau}\}_{\tau=1}^t \cup \{j_{k_\tau}\}_{\tau=1}^t$. In the following, we show that there exists $\{\mu_{j_{k_\tau}}\}_{\tau=1}^t \geq 0, \{\lambda_{i_{k_\tau}}\}_{\tau=1}^t \geq 0$ such that $\mu_{j_{k_\tau}} + \lambda_{i_{k_\tau}} = -\tilde{C}_{i_{k_\tau}, j_{k_\tau}}$ for $\tau = 1, 2, \dots, t$.

We prove this proposition by induction on t (denoted as the size of tree T).

- (i) When $t = 1$, the equation set reduces to $\mu_{j_{k_1}} + \lambda_{i_{k_1}} = -\tilde{C}_{i_{k_1}, j_{k_1}}$. A feasible solution is $\mu_{j_{k_1}} = -\tilde{C}_{i_{k_1}, j_{k_1}}$ and $\lambda_{i_{k_1}} = 0$.
- (ii) Assume that the proposition holds for $t \leq N$ for some positive integer N . In the following, we show that it also holds for $t = N + 1$.

Consider (i_{k_1}, j_{k_1}) which is the edge in the first place. According to Lemma A.3, we show that either i_{k_1} or j_{k_1} is a leaf of tree T : In fact, if i_{k_1} is not a leaf, then $\exists \tau_1 \geq 2$ s.t. $i_{k_{\tau_1}} = i_{k_1}$ and $j_{k_{\tau_1}} \neq j_{k_1}$, and therefore $k_{\tau_1} \geq k_1$. According to Lemma A.3, for any $k > k_1, Q_k \neq 0$, we have $j_k \neq j_{k_1}$. As a result, we have $j_{k_\tau} \neq j_{k_1}$ for any $\tau \in [t], \tau \neq 1$. Similarly, if j_{k_1} is not a leaf, then i_{k_1} must be a leaf. Without loss of generality, assume j_{k_1} is a leaf, and therefore i_{k_1} is not a leaf (otherwise T is not connected).

Denote $T' := G(V_T \setminus \{j_{k_1}\}, E_T \setminus \{(i_{k_1}, j_{k_1})\})$ as the rest of the tree T excluding j_{k_1} , and we have $|T'| = N$ and $i_{k_1} \in T'$. According to the induction hypothesis that holds for $t = N$. the equation set defined on tree T' has a feasible solution. As a result, there exists $\tau^* \geq 2, \tau^* \in [t]$ such that $i_{k_1} = i_{k_{\tau^*}}$. Notice that $\lambda_{i_{k_1}} = \lambda_{i_{k_{\tau^*}}} \leq -\tilde{C}_{i_{k_{\tau^*}}, j_{k_{\tau^*}}} \leq -\tilde{C}_{i_{k_1}, j_{k_1}}$. Therefore, we know that $-\tilde{C}_{i_{k_1}, j_{k_1}} - \lambda_{i_{k_{\tau^*}}} \geq 0$. By letting

$\mu^{j_{k_1}} = -\tilde{C}_{i_{k_1}, j_{k_1}} - \lambda_{i_{k_\tau^*}}$ and combine it with the solutions derived by induction hypothesis, we have construct a feasible solution to the equation set as $t = N + 1$.

By the principle of induction, we have proven Lemma A.1. \blacksquare

Now we are able to get back to the main stream of the proof. Denote the solutions to Eq. (15) as (μ^*, λ^*) , and we have

$$\begin{aligned} h(\mu^*, \lambda^*; \vec{I}, \vec{D}) &= \tilde{h}(\mu^*, \lambda^*; \vec{I}, \vec{D}, \vec{Q}) \\ &= 0 + 0 + \sum_{k=1}^K Q_k \cdot (-\tilde{C}_{i_k, j_k}) \\ &= - \sum_{k=1}^K X_{i_k, j_k} \tilde{C}_{i_k, j_k}. \end{aligned} \tag{16}$$

Therefore, we get a feasible solution to the dual problem shown in Eq. (13) whose value equals $-h(\mu^*, \lambda^*; \vec{I}, \vec{D}) = \sum_{k=1}^K X_{i_k, j_k} \tilde{C}_{i_k, j_k}$. This further equals the value of the primal solution to Eq. (2) proposed by Algorithm 1. Due to the duality, we know that the primal solution is optimal. \blacksquare

A.2 Proof of Lemma 3.2

Notice that Eq. (2) defines a linear programming which contains a matrix variable with constraints on the (weighted) sum of each row and each column. Therefore, we may prove a generalized version of Lemma 3.2, which is defined as follows.

Lemma A.5. *Given parameters $c \in \mathbb{R}^s$, $A \in \mathbb{R}_+^{m \times s}$, $B \in \mathbb{R}_+^{n \times s}$, define the following optimization problem*

$$\begin{aligned} g(\vec{I}, \vec{D}) &:= \min_{x \in \mathbb{R}} c^\top x \\ \text{s.t. } &x \succeq 0 \\ &Ax \preceq \vec{I} \\ &Bx \preceq \vec{D}. \end{aligned} \tag{17}$$

It holds that $g(\vec{I}, \vec{D})$ is convex w.r.t. $[\vec{I}; \vec{D}]$.

Proof of Lemma A.5. Consider the Lagrangian of Eq. (17):

$$L(x, \mu, \lambda, \eta; \vec{I}, \vec{D}) := c^\top x - \mu^\top x + \lambda^\top (Ax - \vec{I}) + \eta^\top (Bx - \vec{D}) = (c - \mu + A^\top \lambda + B^\top \eta)^\top x - \lambda^\top \vec{I} - \eta^\top \vec{D}. \tag{18}$$

Therefore, we have

$$\begin{aligned}
g(\vec{I}, \vec{D}) &:= \min_x \max_{\mu, \lambda, \eta \geq 0} L(x, \mu, \lambda, \eta; \vec{I}, \vec{D}) = \max_{\mu, \lambda, \eta \geq 0} \min_x L(x, \mu, \lambda, \eta; \vec{I}, \vec{D}) \\
&= \max_{\mu, \lambda, \eta \geq 0} \min_x (c - \mu + A^\top \lambda + B^\top \eta)^\top x - \lambda^\top \vec{I} - \eta^\top \vec{D} \\
&= \max_{\mu, \lambda, \eta \geq 0} -\lambda^\top \vec{I} - \eta^\top \vec{D} \\
&\quad \text{s.t. } c - \mu + A^\top \lambda + B^\top \eta = 0.
\end{aligned} \tag{19}$$

Here the second line is due to the strong duality of linear programming. Since the last line indicates that $g(\vec{I}, \vec{D})$ can be represented as the piecewise max of linear functions (which is convex), we know that $g(\vec{I}, \vec{D})$ is also convex w.r.t. $[\vec{I}; \vec{D}]$ jointly. \blacksquare

A.3 Proof of Lemma 3.4

At the beginning, we define a few key quantities and notations. We denote $\tilde{C}_{i,j}(p) := -p + C_{i,j}$ as we did in Appendix A.1. Also, we adopt the notations \tilde{I}_{i_k} and \tilde{D}_{j_k} in Algorithm 1 as the remaining inventories and demands at each step k of allocating X_{i_k, j_k} . Moreover, we denote $\{X_{i_k, j_k}^*\}_{k=1}^K$ as the optimal solution to Eq. (2) generated by Algorithm 1. Given Algorithm 1 and Theorem 3.1, we know that $g(\vec{I}, p, \vec{D})$ can be written as the following form

$$g(\vec{I}, p, \vec{D}) = \sum_{k=1}^K \tilde{C}_{i_k, j_k}(p) \cdot \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\} = \sum_{k=1}^K \tilde{C}_{i_k, j_k}(p) \cdot X_{i_k, j_k}^*. \tag{20}$$

We firstly show that $Q_t(\vec{I}, p)$ is Lipschitz, which is sufficient to show that $g(\vec{I}, p, \vec{D})$ is Lipschitz. In fact, for any $\Delta \vec{I} \geq 0$, we have $g(\vec{I} + \Delta \vec{I}, p, \vec{D}) \geq g(\vec{I}, p, \vec{D}) - \Delta \vec{I} \cdot C_{i_1, j_1}$ and $g(\vec{I} + \Delta \vec{I}, p, \vec{D}) \leq g(\vec{I}, p, \vec{D})$. Also, for any $\Delta \vec{D} \geq 0$, we have $g(\vec{I}, p, \vec{D} + \Delta \vec{D}) \geq g(\vec{I}, p, \vec{D}) - \Delta \vec{I} \cdot C_{i_1, j_1}$ and $g(\vec{I}, p, \vec{D} + \Delta \vec{D}) \leq g(\vec{I}, p, \vec{D})$. Moreover, for any $\Delta p \geq 0$, we have $g(\vec{I}, p + \Delta p, \vec{D}) \geq g(\vec{I}, p, \vec{D}) - \Delta p \cdot \|\vec{I}\|_1$ and $g(\vec{I}, p + \Delta p, \vec{D}) \leq g(\vec{I}, p, \vec{D}) + \Delta p \cdot \|\vec{I}\|_1$.

Since $Q_t(\vec{I}, p) = \langle \vec{\gamma}, \vec{I} \rangle + g(\vec{I}, p, \vec{D}_t(p))$ where $\vec{D}_t(p) = \vec{a} - \vec{b} \cdot p + \vec{N}_t$, we have

$$\begin{aligned}
\frac{\partial Q_t(\vec{I}, p)}{\partial p} &= \sum_{k=1}^K \frac{\partial \tilde{C}_{i_k, j_k}(p)}{\partial p} \cdot \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\} + \sum_{k=1}^K \tilde{C}_{i_k, j_k} \cdot \frac{\partial \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\}}{\partial p} \\
&= \sum_{k=1}^K (-1) \cdot \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\} + \left\langle \frac{\partial g(\vec{I}, p, \vec{D})}{\partial \vec{D}} \Big|_{\vec{D}=\vec{D}_t(p)}, \frac{\partial \vec{D}_t(p)}{\partial p} \right\rangle \\
&= - \sum_{k=1}^K \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\} + \left\langle \frac{\partial g(\vec{I}, p, \vec{D})}{\partial \vec{D}} \Big|_{\vec{D}=\vec{D}_t(p)}, -\vec{b} \right\rangle \\
\frac{\partial^2 Q_t(\vec{I}, p)}{\partial p^2} &= -2 \sum_{k=1}^K \frac{\partial \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\}}{\partial p} + \langle \mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D})^\top (-\vec{b}), -\vec{b} \rangle \\
&= -2 \sum_{k=1}^K \frac{\partial \min\{\tilde{I}_{i_k}, \tilde{D}_{j_k}\}}{\partial p} + \vec{b}^\top \mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D}) \vec{b} \\
&= 2 \left\langle \frac{\partial \sum_{k=1}^K -X_{i_k, j_k}^*}{\partial \vec{D}}, -\vec{b} \right\rangle + \vec{b}^\top \mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D}) \vec{b}, p \in [C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}].
\end{aligned} \tag{21}$$

Here $\mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D})$ represents the partial Hessian matrix of $g(\vec{I}, p, \vec{D})$ with respect to \vec{D} . As we have shown in Lemma 3.2 that g is marginally convex on \vec{D} , we know that $\mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D}) \succeq 0$ and therefore $\vec{b}^\top \mathcal{H}_{\vec{D}} g(\vec{I}, p, \vec{D}) \vec{b} \geq 0$. Since $\vec{b} \geq 0$ as we have assumed, it is sufficient to show $\sum_{k=1}^K \sum_{k=1}^K -X_{i_k, j_k}^*$ is monotonically non-increasing with respect to \vec{D} .

Now we Consider the following optimization problem

$$\begin{aligned}
R(\vec{I}, \vec{D}) &:= \min_{X \in \mathbb{R}^{m \times n}} \sum_{k=1}^K -\mathbb{1}[\tilde{C}_{i_k, j_k} < p] \cdot X_{i_k, j_k} \\
\text{s.t.} \quad &X_{i, j} \geq 0 \\
&\sum_{i=1}^m X_{i, j} \leq D_j, \forall j \in [n] \\
&\sum_{j=1}^n X_{i, j} \leq I_i, \forall i \in [m].
\end{aligned} \tag{22}$$

According to Lemma A.5, we know that $R(\vec{I}, \vec{D})$ is a convex function w.r.t. $[\vec{I}; \vec{D}]$. Also, if we denote $\mathbb{1}_{i_k, j_k} := \mathbb{1}[\tilde{C}_{i_k, j_k} < p]$, we will see $-\mathbb{1}_{i_k, j_k} \leq -\mathbb{1}_{i_{k+1}, j_{k+1}}, \forall k \in [mn - 1]$. This indicates the solution to Eq. (2) generated by Algorithm 1 is also an optimal solution to Eq. (22).

Consider an arbitrary $\Delta \vec{D} \geq \vec{0}$. Since the optimal solution X^* to $R(\vec{I}, \vec{D})$ is a feasible solution to $R(\vec{I}, \vec{D} + \Delta \vec{D})$ (as the constraints get loose), we know that $R(\vec{I}, \vec{D}) \geq R(\vec{I}, \vec{D} +$

$\Delta\vec{D}$). Therefore, we have $\sum_{k=1}^K -X_{i_k, j_k}^*$ is monotonically non-increasing with respect to \vec{D} . This holds the proof.

A.4 Proof of Lemma 3.5

We firstly propose and prove a lemma that shows the number of singularities of function g . Denote $\vec{I}^*(p) := \operatorname{argmin}_{\vec{I}} Q(\vec{I}, p)$, and we have

Lemma A.6. *For any noise vector $\vec{N}_t \in \mathbb{R}^n$, there exists at most 3^{m+n} number of $p \in [C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$ such that $g(\vec{I}^*(p), p, \vec{a} - \vec{b} \cdot p + \vec{N}_t)$ is non-smooth with respect to p .*

Proof of Lemma A.6. Recall that $g(\vec{I}, p, \vec{D}_t(p)) = \sum_{k=1}^K (-p + C_{i_k, j_k}) \cdot X_{i_k, j_k}^*$ where $X_{i_k, j_k} = \min_{\vec{I}_{i_k}, \vec{D}_{j_k}(p)}$. Therefore, the non-smoothness comes from the assignment of X_{i_k, j_k} , which further comes from $\vec{D}_{j_k}(p)$.

To facilitate the proof, we slightly enhance the lemma by letting $\vec{I} = \vec{u} + \vec{v} \cdot p$, and $\vec{D} = \vec{a} - \vec{b} \cdot p$ and prove the lemma for any $\vec{u}, \vec{v}, \vec{a}, \vec{b} \geq 0$. (Here we merge \vec{N}_t into \vec{a} as it is fixed.)

We prove this lemma by induction on $m + n$.

- (i) When $m + n = 2$ i.e. $m = n = 1$, we have $g(\vec{I}, p, \vec{D}) = (-p + C_{1,1}) \min\{u_1 + v_1 p, a_1 - b_1 p\}$, which has $1 \leq 3^{m+n}$ singularities at most.
- (ii) Assume that the lemma holds for $\forall m, n, m + n \leq Z$ for some integer $Z \geq 2$. Now, consider the case when $m + n = Z + 1$, and we have

$$\begin{aligned}
& g(\vec{I}, p, \vec{D}) \\
&= (-p + C_{i_1, j_1}) \cdot \min\{u_{i_1} + v_{i_1} p, a_{j_1} - b_{j_1} \cdot p\} + \sum_{k=2}^K (-p + C_{i_k, j_k}) \cdot \min\{\vec{I}_{i_k}(p), \vec{D}_{j_k}(p)\} \\
&= \mathbf{1}[u_{i_1} + v_{i_1} p \leq a_{j_1} - b_{j_1} \cdot p] \left((-p + C_{i_1, j_1})(u_{i_1} + v_{i_1} p) + g(\vec{I}(p) - (u_{i_1} + v_{i_1} p)\vec{e}_{i_1}, p, \vec{D}(p) - (u_{i_1} + v_{i_1} p)\vec{e}_{j_1}) \right) \\
&\quad + \mathbf{1}[u_{i_1} + v_{i_1} p > a_{j_1} - b_{j_1} \cdot p] \left((-p + C_{i_1, j_1})(a_{j_1} - b_{j_1} p) + g(\vec{I}(p) - (a_{j_1} - b_{j_1} p)\vec{e}_{i_1}, p, \vec{D}(p) - (a_{j_1} + b_{j_1} p)\vec{e}_{j_1}) \right). \tag{23}
\end{aligned}$$

Here $\vec{e}_i := [0, 0, \dots, 0, 1, 0, \dots, 0]^\top$ is an indicator vector which has only one "1". Notice that $g(\vec{I}(p) - (u_{i_1} + v_{i_1} p)\vec{e}_{i_1}, p, \vec{D}(p) - (u_{i_1} + v_{i_1} p)\vec{e}_{j_1})$ and $g(\vec{I}(p) - (a_{j_1} - b_{j_1} p)\vec{e}_{i_1}, p, \vec{D}(p) - (a_{j_1} + b_{j_1} p)\vec{e}_{j_1})$ is discussed in $m + n = Z$. Therefore, from the inductive hypothesis, we know that the total number of singularities is no more than $1 + 3^Z + 3^Z \leq 3^{Z+1}$.

According to the principle of induction, we have proved this lemma valid. ■

According to the definition of $W(p)$ given by Eq. (6), we have

$$\begin{aligned} W'(p) &= \frac{d Q(\vec{I}^*(p), p)}{d p} \\ &= \frac{\partial Q(\vec{I}^*, p)}{\partial p} + \frac{\partial Q(\vec{I}, p)}{\partial I} \Big|_{\vec{I}=\vec{I}^*(p)} \cdot \frac{d \vec{I}^*(p)}{d p}. \end{aligned} \quad (24)$$

According to the definition of $\vec{I}^*(p)$, we have $\frac{\partial Q(\vec{I}, p)}{\partial \vec{I}} = 0$ at $\vec{I} = \vec{I}^*(p)$. Now we show that $\frac{\partial Q(\vec{I}^*(p), p)}{\partial p}$ is a monotonically increasing function of p . According to the definition of Q , we have

$$\begin{aligned} &Q(\vec{I}^*(p), p) \\ &= \mathbb{E}[Q_t(\vec{I}^*(p), p)] \\ &= \mathbb{E}[g(\vec{I}^*, p, \vec{a} - \vec{b} \cdot p + \vec{N}_t) + \langle \vec{\gamma}, \vec{I}^* \rangle] \\ &= \int_{N_1} \int_{N_2} \dots \int_{N_n} g(\vec{I}^*, p, \vec{a} - \vec{b} \cdot p + [N_1, N_2, \dots, N_n]^\top) \cdot \rho_{\vec{N}_t}(N_1, N_2, \dots, N_n) dN_1 dN_2 \dots dN_n + \langle \vec{\gamma}, \vec{I}^* \rangle \\ &= \lim_{\Delta \rightarrow 0^+} \lim_{C_j = C \rightarrow +\infty} \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \left(g(\vec{I}^*, p, \vec{a} - \vec{b} \cdot p + [-C_1 + l_1\Delta, -C_2 + l_2\Delta, \dots, -C_n + l_n\Delta]^\top) + \langle \vec{\gamma}, \vec{I}^* \rangle \right) \\ &\quad \cdot \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right]. \end{aligned} \quad (25)$$

Denote

$$\begin{aligned} \check{Q}_\Delta(\vec{I}^*, p) &:= \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \left(g(\vec{I}^*, p, \vec{a} - \vec{b} \cdot p + [-C_1 + l_1\Delta, -C_2 + l_2\Delta, \dots, -C_n + l_n\Delta]^\top) + \langle \vec{\gamma}, \vec{I}^* \rangle \right) \\ &\quad \cdot \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right]. \end{aligned} \quad (26)$$

Due to the property that the limit operation preserves convexity, it is sufficient to show that $\check{Q}_\Delta(\vec{I}^*, p)$ is convex with respect to p . According to Lemma A.6, we know that $\check{Q}_\Delta(\vec{I}^*, p)$ has at most $E := (\frac{2C}{\Delta})^n \cdot 3^{m+n}$ non-smooth singularities. Without loss of generality, denote them as

$$C_{i_K, j_K} \leq P_1 < P_2 < \dots < P_E \leq C_{i_{K+1}, j_{K+1}}.$$

Also denote $P_0 := C_{i_K, j_K}$ and $P_{E+1} := C_{i_{K+1}, j_{K+1}}$. Now we propose another two lemmas.

Lemma A.7. For $p \in (P_e, P_{e+1})$, $e = 0, 1, 2, \dots, E$, we show that $\frac{\partial \check{Q}(\vec{I}^*, p)}{\partial p}$ is monotonically increasing on p .

Proof of Lemma A.7. Notice that

$$\begin{aligned} \frac{\partial}{\partial p} \check{Q}_\Delta(\vec{I}^*, p) &= \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \frac{\partial g(\vec{I}^*, p, \vec{a} - \vec{b}p + \vec{N}_t)}{\partial p} \Big|_{\vec{N}_t} \cdot \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right] \\ &= \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \frac{\partial Q_t(\vec{I}^*, p)}{\partial p} \Big|_{\vec{N}_t} \cdot \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right]. \end{aligned} \quad (27)$$

Here $\vec{N}_t := [-C_1 + l_1\Delta, -C_2 + l_2\Delta, \dots, -C_n + l_n\Delta]^\top$. Now we consider the monotonicity of $\frac{\partial Q_t(\vec{I}^*, p)}{\partial p}$ on each (P_e, P_{e+1}) interval. Since there exist no singularities in this interval, we know that $Q_t(\vec{I}^*, p) \in \mathbb{C}^2$ in this range, and therefore we have

$$\begin{aligned} \frac{d}{d p} \frac{\partial Q_t(\vec{I}^*, p)}{\partial p} &:= \left\langle \frac{\partial Q_t(\vec{I}^*, p)}{\partial I \partial p}, \frac{d \vec{I}^*(p)}{d p} \right\rangle + \frac{\partial^2 Q_t(\vec{I}^*, p)}{\partial p^2} \\ &= \left\langle \frac{\partial}{\partial p} \frac{\partial Q_t(\vec{I}^*, p)}{\partial \vec{I}} \Big|_{\vec{I}=\vec{I}^*(p)}, \frac{d \vec{I}^*(p)}{d p} \right\rangle + \frac{\partial^2 Q_t(\vec{I}^*, p)}{\partial p^2} \\ &= \left\langle \vec{0}, \frac{d \vec{I}^*(p)}{d p} \right\rangle + \frac{\partial^2 Q_t(\vec{I}^*, p)}{\partial p^2} \\ &= \frac{\partial^2 Q_t(\vec{I}^*, p)}{\partial p^2} \geq 0. \end{aligned} \quad (28)$$

Here the second line that we swap the sequence of derivatives is due to the smoothness within the (P_e, P_{e+1}) smooth interval, and the last line is from Lemma 3.4 which shows the marginal convexity of $Q_t(\vec{I}, p)$ w.r.t. p . Therefore, we have proved the lemma. \blacksquare

Lemma A.8. At each P_e for $e = 0, 1, 2, \dots, E$, we have $W'(P_e^-) \leq W'(P_e^+)$.

Proof of Lemma A.8. We firstly consider $W'(P_e^-)$. According to the proof of Lemma A.7,

we have

$$\begin{aligned}
W'(P_e^-) &= \lim_{p \rightarrow P_e^-} W'(p) \\
&= \lim_{p \rightarrow P_e^-} \frac{\partial}{\partial p} \lim_{\Delta \rightarrow 0^+} \lim_{C \rightarrow +\infty} \check{Q}_\Delta(\vec{I}^*, p) \\
&= \lim_{p \rightarrow P_e^-} \lim_{\Delta \rightarrow 0^+} \lim_{C \rightarrow +\infty} \frac{\partial}{\partial p} \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} Q_t(\vec{I}^*, p) |_{\vec{N}_t} \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right] \\
&= \lim_{\Delta \rightarrow 0^+} \lim_{C \rightarrow +\infty} \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right] \cdot \lim_{p \rightarrow P_e^-} \frac{\partial Q_t(\vec{I}^*, p)}{\partial p} \\
&\leq \lim_{\Delta \rightarrow 0^+} \lim_{C \rightarrow +\infty} \sum_{l_1=1}^{\frac{2C_1}{\Delta}} \sum_{l_2=1}^{\frac{2C_2}{\Delta}} \dots \sum_{l_n=1}^{\frac{2C_n}{\Delta}} \Pr\left[\bigcap_{j=1}^n (\vec{N}_t(j) \in [-C_j + (l_j - 1)\Delta, -C_j + l_j\Delta])\right] \cdot \lim_{p \rightarrow P_e^+} \frac{\partial Q_t(\vec{I}^*, p)}{\partial p} \\
&= \lim_{p \rightarrow P_e^+} \frac{\partial}{\partial p} \lim_{\Delta \rightarrow 0^+} \lim_{C \rightarrow +\infty} \check{Q}_\Delta(\vec{I}^*, p) \\
&= \lim_{p \rightarrow P_e^+} W'(p) \\
&= W'(p_e^+).
\end{aligned} \tag{29}$$

Here the fourth and the sixth lines are due to the Moore-Osgood theorem of exchanging limits, as $Q_t(\vec{I}^*, p) \in \mathbb{C}^2$ when $p \in (P_{e-1}, P_e)$ and $p \in (P_e, P_{e+1})$ respectively. The fifth line comes from Lemma 3.4: as $Q_t(\vec{I}^*, p)$ is convex w.r.t. p , the left derivatives of $\frac{\partial Q_t(\vec{I}^*, p)}{\partial p}$ should not exceed its right derivatives at any point p . \blacksquare

Applying Lemma A.7 on Eq. (24), we know that $W(p)$ is convex within each smooth interval (P_e, P_{e+1}) . Also, from Lemma A.8, we know that $W(p)$ is convex at any singularity P_e as its left derivatives does not exceed its right derivatives. Combining those two properties, we know that $Q(\vec{I}^*, p)$ is convex w.r.t. p . This ends the proof.

A.5 Proof of Lemma 5.2

The main idea of this proof originates from OCO with zeroth-order (bandit) feedback, as is displayed in Agarwal et al. (2011). Specifically, we conduct the proof in the following steps:

- (a) When an agent \mathcal{A}_K is in Stage 1, Epoch τ and Sub-Epoch s , then we sequentially show that

- (i) The aggregated function $Q_{K,\tau,s}(\vec{I}, p)$ is concentrated to $Q(\vec{I}, p)$ for the three proposed $p = \hat{p}_\tau \in \{a_{K,\tau}, b_{K,\tau}, c_{K,\tau}\}$ and for any \vec{I} , with $O(1/\sqrt{n_s})$ error.
 - (ii) The $\hat{Q}_{K,\tau,s,p}$, which takes the empirical optimal inventory decision, is concentrated to $Q(\vec{I}^*(p), p) = W(p)$ at those proposed prices $p = \hat{p}_\tau \in \{a_{K,\tau}, b_{K,\tau}, c_{K,\tau}\}$, with $O(1/\sqrt{n_s})$ error.
 - (iii) According to the convexity of $W(p)$, we upper bound the sub-regret per round $W(p) - W(p_K^*)$ by $O(1/\sqrt{n_s})$, where p_K^* is the local optimal price.
 - (iv) We show that the total number of epochs in Stage 1 is $O(\log T)$. According to the doubling lengths of n_s , the total sub-regret of \mathcal{A}_K is $\tilde{O}(\sqrt{T_K})$ by the time when \mathcal{A} has proposed T_K pairs of decisions (\vec{I}_t, p_t) .
- (b) When an agent \mathcal{A}_K reaches Stage 2 or 3, we know that the search space $[L_K, U_K]$ is smaller than $1/T$. Given that the Q_t functions (and therefore $W(p)$) are Lipschitz, we may upper bound the sub-regret per step as $O(1/T)$ and the total sub-regret as $O(1)$.

Before we get to proof details, we propose a lemma that generally holds for convex functions.

Lemma A.9. *Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a L -Lipschitz convex function. Denote $f(x^*) := \min_{x \in [a, b]} f(x)$, $x_1 = \frac{3a+b}{4}$, $x_2 = \frac{a+b}{2}$, $x_3 = \frac{a+3b}{4}$. Assume there exists some fixed constants A and $\Delta > 0$ such that $f(x_i) \in [A - \Delta, A + \Delta]$, $i = 1, 2, 3$, then we have*

$$\max\{f(x_1), f(x_2), f(x_3)\} - f(x^*) \leq 4\Delta. \quad (30)$$

Please kindly find the proof of Lemma A.9 in Appendix A.8.

Now we return to the main proof. We firstly propose the following concentration lemma

Lemma A.10. *For Agent \mathcal{A}_K running in Epoch τ Sub-Epoch s , and $\forall \vec{I} \in \mathbb{R}_+$, symbolic variable $\hat{p}_\tau \in \{a_{K,\tau}, b_{K,\tau}, c_{K,\tau}\}$, we have*

$$|Q_{K,\tau,s}(\vec{I}, \hat{p}_\tau) - Q(\vec{I}, \hat{p}_\tau)| \leq \frac{C_c}{2\sqrt{n_s}} \quad (31)$$

with probability $\Pr \geq 1 - \hat{\epsilon}$. Here $C_c := \sqrt{2 \log \frac{2}{\hat{\epsilon}}} \cdot Q_{\max}$ and $Q_{\max} := \max\{p_{\max}, \gamma_{\max}\} I_{\max}$.

We will specify the value of $\hat{\epsilon}$ as a function of ϵ by the end of this proof. We defer the proof of Lemma A.10 to Appendix A.9. From Lemma A.10, we may get the following corollary.

Corollary A.11. For Agent \mathcal{A}_K running in Epoch τ Sub-Epoch s , and symbolic variable $\hat{p}_\tau \in \{a_{K,\tau}, b_{K,\tau}, c_{K,\tau}\}$ we have

$$\begin{aligned} |\hat{Q}_{K,\tau,s,\hat{p}_\tau} - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau)| &\leq \frac{C_c}{2\sqrt{n_s}}, \text{ and} \\ 0 \leq Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) &\leq \frac{C_c}{\sqrt{n_s}} \end{aligned} \quad (32)$$

with probability $\Pr \geq 1 - 4T\hat{\epsilon}$.

Proof of Corollary A.11. For each fixed tuple $(K, \tau, s, \hat{p}_\tau)$, according to Lemma A.10, we have

$$\begin{aligned} \hat{Q}_{K,\tau,s,\hat{p}_\tau} - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) &= Q_{K,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q_{K,\tau,s}(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) + Q_{K,\tau,s}(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) \\ &\leq 0 + \frac{C_c}{2\sqrt{n_s}} = \frac{C_c}{2\sqrt{n_s}}. \\ \hat{Q}_{K,\tau,s,\hat{p}_\tau} - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) &= Q_{K,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) + Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) \\ &\geq -\frac{C_c}{2\sqrt{n_s}} + 0 = -\frac{C_c}{2\sqrt{n_s}} \end{aligned} \quad (33)$$

with probability $\Pr \geq 1 - 2\hat{\epsilon}$. Here the first inequality comes from the optimality of $I_{K,\tau,s}(\hat{p}_\tau)$ over $Q_{K,\tau,s}(\cdot, \hat{p}_\tau)$ as well as the concentration of $Q_{K,\tau,s}(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau)$ towards $Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau)$. The second inequality comes from the concentration of $Q_{K,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau)$ towards $Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau)$ as well as the optimality of $\vec{I}^*(\hat{p}_\tau)$ over $Q(\cdot, \hat{p}_\tau)$.

Also, we have

$$\begin{aligned} &Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q(I^*(\hat{p}_\tau), \hat{p}_\tau) \\ &= Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q_{K,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) \\ &\quad + Q_{K,\tau,s}(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q_{K,\tau,s}(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) \\ &\quad + Q_{K,\tau,s}(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) \\ &\leq \frac{C_c}{2\sqrt{n_s}} + 0 + \frac{C_c}{2\sqrt{n_s}} \\ &= \frac{C_c}{\sqrt{n_s}} \end{aligned} \quad (34)$$

with probability $\Pr \geq 1 - 2\hat{\epsilon}$. Here the third line comes from the concentrations (the first and the third term) as well as the optimality of $I_{K,\tau,s}(\hat{p}_\tau)$ over $Q_{K,\tau,s}(\cdot, \hat{p}_\tau)$. Besides, the other side that $Q(I_{K,\tau,s}(\hat{p}_\tau), \hat{p}_\tau) - Q(I^*(\hat{p}_\tau), \hat{p}_\tau)$ is due to the optimality of $\vec{I}^*(\hat{p}_\tau)$ over $Q(\cdot, \hat{p}_\tau)$.

Since the combination of $(K, \tau, s, \hat{p}_\tau)$ is unique, and the total number of combinations is exactly T , we apply the union bound of probability and get that Eq. (32) holds for all $(K, \tau, s, \hat{p}_\tau)$ tuples with probability $\Pr \geq 1 - 4T\hat{\epsilon}$. \blacksquare

Combining Corollary A.11 with Lemma A.9, we have the following corollary

Corollary A.12. *Define a flag as shown in Algorithm 5, and define $p_K^* := \operatorname{argmin}_{p \in [C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]} W(p)$. When $\text{flag} == 0$ by the end of Sub-Epoch s of Epoch τ , we have*

$$|W(\hat{p}_\tau) - W(p_K^*)| \leq 16 \cdot \frac{C_c}{\sqrt{n_s}} \quad (35)$$

holds for $\hat{p}_\tau = a_{K,\tau}, b_{K,\tau}, c_{K,\tau}$ with probability $\Pr \geq 1 - 24T\hat{\epsilon}$.

Proof of Corollary A.12. When $\text{flag} == 0$, according to Algorithm 5, we know that

$$\begin{aligned} |\hat{Q}_{K,\tau,s,a_{K,\tau}} - \hat{Q}_{K,\tau,s,b_{K,\tau}}| &\leq 4\Delta_{K,\tau,s} \\ \hat{Q}_{K,\tau,s,a_{K,\tau}} - \hat{Q}_{K,\tau,s,c_{K,\tau}} &\leq 4\Delta_{K,\tau,s} \\ \hat{Q}_{K,\tau,s,b_{K,\tau}} - \hat{Q}_{K,\tau,s,c_{K,\tau}} &\leq 4\Delta_{K,\tau,s}. \end{aligned} \quad (36)$$

Also, according to the convexity of $W(P)$ in $[C_{i_K, j_K}, C_{i_{K+1}, j_{K+1}}]$, we know that $W(c_{K,\tau}) \leq \frac{W(a_{K,\tau}) + W(b_{K,\tau})}{2} \leq \max\{W(a_{K,\tau}), W(b_{K,\tau})\}$. Without loss of generality, assume $W(a_{K,\tau}) \geq W(b_{K,\tau})$, and then we have

$$\begin{aligned} \hat{Q}_{K,\tau,s,c_{K,\tau}} - \hat{Q}_{K,\tau,s,a_{K,\tau}} &= Q(I_{K,\tau,s}(c_{K,\tau}), c_{K,\tau}) - Q(\vec{I}^*(c_{K,\tau}), c_{K,\tau}) \\ &\quad + W(c_{K,\tau}) - W(a_{K,\tau}) \\ &\quad + Q(\vec{I}^*(a_{K,\tau}), a_{K,\tau}) - Q(I_{K,\tau,s}(a_{K,\tau}), a_{K,\tau}) \\ &\leq \frac{C_c}{\sqrt{n_s}} + 0 + \frac{C_c}{\sqrt{n_s}} \\ &= \frac{2C_c}{\sqrt{n_s}} \leq 4\Delta_{K,\tau,s}. \end{aligned} \quad (37)$$

Therefore we know that $|\hat{Q}_{K,\tau,s,b_{K,\tau}} - \hat{Q}_{K,\tau,s,c_{K,\tau}}| \leq 4\Delta_{K,\tau,s}$. Combining Corollary A.11, we have

$$\begin{aligned} |W(a_{K,\tau}) - W(b_{K,\tau})| &\leq |Q(\vec{I}^*(a_{K,\tau}), a_{K,\tau}) - Q(I_{K,\tau,s}(a_{K,\tau}), a_{K,\tau})| \\ &\quad + |Q(I_{K,\tau,s}(a_{K,\tau}), a_{K,\tau}) - Q(I_{K,\tau,s}(b_{K,\tau}), b_{K,\tau})| \\ &\quad + |Q(I_{K,\tau,s}(b_{K,\tau}), b_{K,\tau}) - Q(\vec{I}^*(b_{K,\tau}), b_{K,\tau})| \\ &\leq \frac{C_c}{\sqrt{n_s}} + 4\Delta_{K,\tau,s} + \frac{C_c}{\sqrt{n_s}} \\ &\leq \frac{4C_c}{\sqrt{n_s}}. \end{aligned} \quad (38)$$

And also

$$\begin{aligned}
|W(a_{K,\tau}) - W(c_{K,\tau})| &\leq |Q(\vec{I}^*(a_{K,\tau}), a_{K,\tau}) - Q(I_{K,\tau,s}(a_{K,\tau}), a_{K,\tau})| \\
&\quad + |Q(I_{K,\tau,s}(a_{K,\tau}), a_{K,\tau}) - Q(I_{K,\tau,s}(b_{K,\tau}), c_{K,\tau})| \\
&\quad + |Q(I_{K,\tau,s}(c_{K,\tau}), b_{K,\tau}) - Q(\vec{I}^*(b_{K,\tau}), c_{K,\tau})| \\
&\leq \frac{C_c}{\sqrt{n_s}} + 4\Delta_{K,\tau,s} + \frac{C_c}{\sqrt{n_s}} \\
&\leq \frac{4C_c}{\sqrt{n_s}}.
\end{aligned} \tag{39}$$

By applying Lemma A.9 with $\Delta = \frac{4C_c}{\sqrt{n_s}}$, we show that the lemma holds with $\Pr \geq 1 - 24T\hat{\epsilon}$ (since we have used Corollary A.11 for 6 times). \blacksquare

Finally, we show that upper bounds the total number of epochs in which \mathcal{A}_K is running, and we first denote this number as M_K . In fact, from the design of *Algorithm 5*, we know that by the end of each epoch, we have $U_{K,\tau+1} - L_{K,\tau+1} = \frac{3}{4}(U_{K,\tau} - L_{K,\tau})$, i.e. the length of search space $[L_{K,\tau}, U_{K,\tau}]$ reduces by 1/4. Since $L_{K,1} = C_{i_K, j_K}, U_{K,1} = C_{i_{K+1}, j_{K+1}}$, we have

$$\begin{aligned}
M_K &\leq \log_{3/4} \frac{U_{K,M_K} - L_{K,M_K}}{U_{K,1} - L_{K,1}} \\
&= \log_{4/3} \frac{U_{K,1} - L_{K,1}}{U_{K,M_K} - L_{K,M_K}} \\
&\leq \log_{4/3} \frac{C_{i_{K+1}, j_{K+1}} - C_{i_K, j_K}}{\frac{1}{T} \cdot \frac{3}{4}} \\
&= \log_{4/3}(C_{i_{K+1}, j_{K+1}}) + \log_{4/3} T + 1 \\
&\leq \log_{4/3} p_{\max} + \log_{4/3} T + 1.
\end{aligned} \tag{40}$$

Denote $\check{C}_K := \log_{4/3} p_{\max} + 1$, and we have $M_K \leq \log_{4/3} T + \check{C}_K$

With all properties above, we may derive the total sub-regret for \mathcal{A}_K . Firstly, the cumulative sub-regret in Epoch τ Sub-Epoch s is

$$\begin{aligned}
& \text{SubReg}(\mathcal{A}_K, \tau, s) \\
& := n_s \cdot \sum_{\hat{p}_\tau = a_{K,\tau}, b_{K,\tau}, c_{K,\tau}} Q(\vec{I}_{K,\tau,s-1}(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(p_K^*), p_K^*) \\
& \leq n_s \cdot \sum_{\hat{p}_\tau = a_{K,\tau}, b_{K,\tau}, c_{K,\tau}} |Q(\vec{I}_{K,\tau,s-1}(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau)| + |Q(\vec{I}^*(\hat{p}_\tau), \hat{p}_\tau) - Q(\vec{I}^*(p_K^*), p_K^*)| \\
& \leq n_s \cdot \sum_{\hat{p}_\tau = a_{K,\tau}, b_{K,\tau}, c_{K,\tau}} \frac{C_c}{\sqrt{n_s}} + 20 \cdot \frac{C_c}{\sqrt{n_s}} \\
& = 63C_c\sqrt{n_s} \\
& = 63C_c \cdot 2^{s/2}.
\end{aligned} \tag{41}$$

Secondly, denote the number of sub-epochs in Epoch τ as S_τ and the length of Epoch τ as T_τ (therefore we know that $T_\tau = 3 \cdot 2^{S_\tau+1} - 1$), and the cumulative sub-regret in Epoch τ is bounded by

$$\begin{aligned}
\text{SubReg}(\mathcal{A}_K, \tau) & := \sum_{s=1}^{S_\tau} \text{SubReg}(\mathcal{A}_K, \tau, s) \\
& \leq \sum_{s=1}^{S_\tau} 63C_c \cdot 2^{s/2} \\
& \leq 63C_c \cdot \frac{1}{\sqrt{2}-1} 2^{\frac{S_\tau+1}{2}} \\
& \leq 200C_c \cdot \sqrt{\frac{T_\tau+1}{3}} \\
& \leq 200C_c \sqrt{T_\tau}.
\end{aligned} \tag{42}$$

Thirdly, we may calculate the total sub-regret of \mathcal{A}_K as

$$\begin{aligned}
SReg_K &:= \sum_{\tau=1}^{M_K} SubReg(\mathcal{A}_K, \tau) \\
&\leq \sum_{\tau=1}^{M_K} 200C_c \sqrt{T_\tau} \\
&\leq 200C_c \cdot \sqrt{\left(\sum_{\tau=1}^{M_K} (\sqrt{T_\tau})^2 \right) \left(\sum_{\tau=1}^{M_K} 1 \right)} \\
&= 200C_c \left(\sum_{\tau=1}^{M_K} T_\tau \right) (M_K) \\
&\leq 200C_c \cdot \sqrt{T_K \cdot M_K} \\
&\leq 200C_c \sqrt{T_K (\log_{4/3} T + \check{C}_K)} \\
&= \tilde{O}(\sqrt{T_K}).
\end{aligned} \tag{43}$$

This rate holds with $\Pr \geq 1 - 24T\hat{\epsilon}$ for each $K \in [2mn + 1]$. Let $\hat{\epsilon} := \frac{\epsilon}{24 \cdot (2mn+1)T}$ so that $C_c = \delta_K = \sqrt{2 \log \frac{48(2mn+1)T}{\epsilon}} \cdot \max\{p_{\max}, \gamma_{\max}\} \cdot I_{\max}$, and we complete the proof of Lemma 5.2.

A.6 Proof of Lemma 5.3

We analyze the behavior of Δ_K by considering the current stage of \mathcal{A}_K .

1. If \mathcal{A}_K is currently in Stage 1. Suppose \mathcal{A}_K has played for τ_K epochs. Since each epoch reduces the price interval $[L_{K,\tau}, U_{K,\tau}]$ to its 3/4, we know that $\tau_K \leq M_K \leq \log_{4/3} T + \check{C}_K$ where $C_K := \log_{4/3} p_{\max} + 1$ (see Eq. (40)).

According to Pigeon-Hole Theorem, at least the longest epoch $\hat{\tau}$ has been played for $\frac{T_K}{\tau_K} \geq \frac{6(\log_{4/3} T + C_K)}{\log_{4/3} T + C_K} = 6$ times. As a result, at least **two** sub-epochs have been reached in this epoch. Since we update Δ_K by the end of each sub-epoch (except for the last sub-epoch of each epoch), at least one of these two sub-epochs leads to an update on Δ_K . As a result, $\Delta_K < +\infty$ after this update, and after $T_K \geq 6(\log_{4/3} T + C_K)$ time periods.

Denote the length of this epoch $\hat{\tau}$ as $H_{K,\hat{\tau}}$, and we know that $H_{K,\hat{\tau}} \geq \frac{T_K}{\log_{4/3} T + C_K}$. Also, we denote the length of each sub-epoch of Epoch $\hat{\tau}$ as $H_{K,\hat{\tau},s}$, $s = 1, 2, \dots, S_{\hat{\tau}}$, where $S_{\hat{\tau}}$ is denoted as the number of sub-epochs of Epoch $\hat{\tau}$. Given those definitions,

we have

$$\begin{aligned}
H_{K,\hat{\tau},S_{\hat{\tau}}-1} &\geq \sum_{s=1}^{S_{\hat{\tau}}-2} H_{K,\hat{\tau},s} \\
H_{K,\hat{\tau},S_{\hat{\tau}}-1} &\geq \frac{H_{K,\hat{\tau},S_{\hat{\tau}}}}{2}.
\end{aligned} \tag{44}$$

As a consequence, we have

$$H_{K,\hat{\tau},S_{\hat{\tau}}-1} \geq \frac{H_{K,\hat{\tau}}}{4} \geq \frac{T_K}{4(\log_{4/3} T + C_K)}. \tag{45}$$

Since we still can update Δ_K by the end of Epoch $\hat{\tau}$ Sub-Epoch $S_{\hat{\tau}} - 1$, we may upper-bound Δ_K in the following approach

$$\begin{aligned}
\Delta_K &= \frac{\delta_K}{2\sqrt{n_{S_{\hat{\tau}}-1}}} \\
&= \frac{\delta_K}{2 \cdot \sqrt{H_{K,\hat{\tau},S_{\hat{\tau}}-1}/3}} \\
&\leq \frac{\delta_K}{\frac{2}{\sqrt{3}} \cdot \sqrt{\frac{T_K}{4(\log_{4/3} T + C_K)}}} \\
&= \sqrt{\frac{1}{T_K}} \cdot \delta_K \cdot \sqrt{3(\log_{4/3} T + C_K)} \\
&= \tilde{O}\left(\sqrt{\frac{1}{T_K}}\right).
\end{aligned} \tag{46}$$

2. If \mathcal{A}_K reaches Stage 2. Since we only run Stage 2 for once without stopping, updating \hat{W}_K, Δ_K or switching agents, we assume that T_K reaches the end of Stage 2 without loss of generality. We firstly upper and lower bound the length of Stage 2. Denote $T_{K,1}$ as the time periods that \mathcal{A}_K spent on Stage 1, and $T_{K,2} := T_K - T_{K,1}$ as the time periods that \mathcal{A}_K has spent on Stage 2 so far. Remember that the purpose of conducting Stage 2 is to guarantee a Δ_K that is comparable to $\sqrt{\frac{1}{T_K}}$, and at the end of Stage 2 we reduce Δ_K to its half comparing to the one we have by the end of Stage 1 (if not $+\infty$). Therefore, we have

$$2T_{K,1} \geq 2H_{K,\hat{\tau}} \geq N_{K,2} \geq H_{K,\hat{\tau}} \geq \frac{T_{K,1}}{\log_{4/3} T + C_K}. \tag{47}$$

Here the first inequality represents that \mathcal{A}_K runs $T_{K,1}$ time periods in Stage 1, including $H_{K,\hat{\tau}}$ time periods in Stage 1 Epoch $\hat{\tau}$. The second and third inequalities hold because we get comparable Δ_K in Stage 1 and in Stage 2, and the best Δ_K we got in

Stage 1 is on the longest sub-epoch, which is $\hat{\tau}$. The last inequality is from the proof shown in Case 1 (when \mathcal{A}_K reaches Stage 1).

Also, since Stage 2 applies a "Doubling Trick", we have

$$\begin{aligned} T_{K,2} &\leq 2^1 + 2^2 + \dots + N_{K,2} \\ &\leq 2N_{k,2}, \\ \Rightarrow T_{K,2} &\leq 4T_{K,1} \\ \Rightarrow T_{K,1} &\leq T_K \leq 5T_{K,1}. \end{aligned} \tag{48}$$

As a result, we have

$$\Delta_K = \frac{\delta_K}{\sqrt{N_{K,2}}} \leq \frac{\delta_K}{\sqrt{\frac{T_{K,1}}{\log_{4/3} T + C_K}}} \leq \frac{\delta_K (\log_{4/3} T + C_K)}{\sqrt{\frac{T_K}{5}}} = \tilde{O}\left(\frac{1}{\sqrt{T_K}}\right). \tag{49}$$

3. If \mathcal{A}_K reaches Stage 3. Denote $T_{K,3} := T_K - T_{K,1} - T_{K,2}$ as the time periods that \mathcal{A}_K has spent on Stage 3 so far. According to Algorithm 7, we know that $\Delta_K = \frac{\delta_K}{\sqrt{N_{K,3}}} = \frac{\delta_K}{\sqrt{N_{K,2} + T_{K,3}}} = \tilde{O}\left(\frac{1}{\sqrt{N_{K,2} + T_{K,3}}}\right)$. Also, since

$$\begin{aligned} N_{K,2} + T_{K,3} &\geq \frac{1}{2} \cdot N_{K,2} + \frac{1}{2} \cdot N_{K,2} + T_{K,3} \\ &\geq \frac{1}{2} \cdot \frac{T_{K,1}}{\log_{4/3} T + C_K} + \frac{1}{2} \cdot \frac{T_{K,2}}{2} + T_{K,3} \\ &\geq \frac{T_{K,1} + T_{K,2} + T_{K,3}}{2(\log_{4/3} T + C_K)} \\ &= \frac{T_K}{2(\log_{4/3} T + C_K)}. \end{aligned} \tag{50}$$

Therefore, we have $\Delta_K \leq \tilde{O}\left(\frac{1}{\sqrt{\frac{T_K}{2(\log_{4/3} T + C_K)}}}\right) = \tilde{O}\left(\frac{1}{\sqrt{T_K}}\right)$. This ends the proof of Lemma 5.3.

A.7 Proof of Lemma 5.5

We consider each case where \mathcal{A}_K is in Stage 1,2,3, respectively.

1. If \mathcal{A}_K is in Stage 1. When updating Δ_K , we know that $flag == 0$ at Stage 1 Epoch τ Sub-Epoch $s-1$ according to Algorithm 5. Denote $\hat{p} := \operatorname{argmin}_{\hat{p}_\tau \in \{a_{K,\tau}, c_{K,\tau}, b_{K,\tau}\}} \hat{Q}_{K,\tau,s-1,\hat{p}_\tau}$,

and we have that

$$\begin{aligned}
LCB_K &= \hat{W}_K - 34\Delta_K \\
&= \hat{Q}_{K,\tau,s-1}(I_{K,\tau,s-1}(\hat{p}), \hat{p}) - 32 \cdot \frac{\delta_K}{2\sqrt{n_{s-1}}} \\
&= \hat{Q}_{K,\tau,s-1}(I_{K,\tau,s-1}(\hat{p}), \hat{p}) - W(\hat{p}) + W(\hat{p}) - 17 \cdot \frac{\delta_K}{\sqrt{n_{s-1}}} \quad (51) \\
&\leq \frac{C_c}{\sqrt{n_{s-1}}} - \frac{\delta_K}{\sqrt{n_{s-1}}} + W(\hat{p}) - 16 \cdot \frac{\delta_K}{\sqrt{n_{s-1}}} \\
&\leq 0 + W(p_K^*).
\end{aligned}$$

Here the fourth line comes from Corollary A.11 and the last line is an application of $C_c = \delta_K$ and Corollary A.12.

On the other hand, the lower bound LCB_K is not too faraway from $W(p_K^*)$ since we have:

$$\begin{aligned}
&LCB_K \\
&= \min\{Q_{K,\tau,s-1,a_{K,\tau}}, Q_{K,\tau,s-1,c_{K,\tau}}, Q_{K,\tau,s-1,b_{K,\tau}}\} - 34\Delta_K \\
&= \min\{Q_{K,\tau,s-1,a_{K,\tau}} - \frac{C_c}{2\sqrt{n_{s-1}}}, Q_{K,\tau,s-1,c_{K,\tau}} - \frac{C_c}{2\sqrt{n_{s-1}}}, Q_{K,\tau,s-1,b_{K,\tau}} - \frac{C_c}{2\sqrt{n_{s-1}}}\} - 33 \cdot \frac{C_c}{2\sqrt{n_{s-1}}} \\
&\geq \min\{W(a_{K,\tau}), W(b_{K,\tau}), W(c_{K,\tau})\} - 33 \cdot \frac{C_c}{2\sqrt{n_{s-1}}} \\
&\geq W(p_K^*) - 33\Delta_K. \quad (52)
\end{aligned}$$

2. If \mathcal{A}_K is in Stage 2 and Stage 3, we may consider them altogether as the only update of LCB as well as Δ_K occurs by the end of Stage 2, which is also the 0-th time period

of Stage 3. In this case, we have

$$\begin{aligned}
& \hat{W}_K - W(p_K^*) \\
&= (\hat{W}_K - \hat{Q}_K^*) + (Q_{K,r_K}(\vec{I}_K^*, \hat{p}_K^*) - Q_{K,r_K}(\vec{I}^*(\hat{p}_K^*), \hat{p}_K^*)) \\
&\quad + (Q_{K,r_K}(\vec{I}^*(\hat{p}_K^*), \hat{p}_K^*) - Q(\vec{I}^*(\hat{p}_K^*), \hat{p}_K^*)) + (W(\hat{p}_K^*) - W(p_K^*)) \\
&\leq 0 + 0 + \frac{\delta_K}{2\sqrt{N_{K,2} + T_{K,3}}} + \frac{L_W}{T} \\
&\leq \Delta_K + \frac{L_W}{T}. \\
& W(p_K^*) - \hat{W}_K \\
&= (W(p_K^*) - W(\hat{p}_K^*)) + (W(\hat{p}_K^*) - Q(\vec{I}_K^*, \hat{p}_K^*)) + (Q(\vec{I}_K^*, \hat{p}_K^*) - Q_{K,r_K}(\vec{I}_K^*, \hat{p}_K^*)) + (\hat{Q}_K^* - \hat{W}_K) \\
&\leq 0 + 0 + \frac{\delta_K}{2\sqrt{N_{K,2} + T_{K,3}}} + \frac{L_W}{T} \\
&\leq \Delta_K + \frac{L_W}{T}.
\end{aligned} \tag{53}$$

Since $LCB_K = \hat{W}_K - 34\Delta_K - \frac{L_W}{T}$, we have

$$\begin{aligned}
LCB_K &\leq W(p_K^*) + \Delta_K + \frac{L_W}{T} - 34\Delta_K - \frac{L_W}{T} \\
&\leq W(p_K^*). \\
LCB_K &\geq W(p_K^*) - \Delta_K - \frac{L_W}{T} - 34\Delta_K - \frac{L_W}{T} \\
&\geq W(p_K^*) - 35\Delta_K - \frac{2L_W}{T}.
\end{aligned} \tag{54}$$

Combining the two cases listed above, we have proved this lemma.

A.8 Proof of Lemma A.9

Proof. Denote $f_1 := f(x_1)$, $f_2 := f(x_2)$, $f_3 := f(x_3)$. Then we prove this lemma by cases where x^* locates.

1. When $x^* \in [a, x_1]$, we denote $\epsilon := \frac{x_2 - x_1}{x_2 - x^*}$. We know that $\epsilon \in [\frac{1}{2}, 1]$, and then we have $\epsilon f(x^*) + (1 - \epsilon)f(x_2) \geq f(\epsilon x^* + (1 - \epsilon)x_2) = f(x_1)$ due to the convexity of $f(x)$. As a result, we have:

$$\begin{aligned}
f(x^*) &\geq \frac{f_1 - (1 - \epsilon)f_2}{\epsilon} \\
&= f_2 + \frac{f_1 - f_2}{\epsilon} \\
&= f_2 - \frac{f_2 - f_1}{\epsilon}.
\end{aligned} \tag{55}$$

If $f_1 \geq f_2$, then we have $f(x^*) \geq f_2 \geq A - \Delta = A + \Delta - 2\Delta \geq \max\{f_1, f_2, f_3\} - 2\Delta$.
 Otherwise $f_1 < f_2$, then we have

$$\begin{aligned}
 f(x^*) &\geq f_2 - \frac{f_2 - f_1}{\epsilon} \\
 &\geq f_2 - \frac{f_2 - f_1}{\frac{1}{2}} \\
 &= 2f_1 - f_2 \\
 &\geq 2(A - \Delta) - (A + \Delta) \\
 &= A - 3\Delta \\
 &= A + \Delta - 4\Delta \\
 &\geq \max\{f_1, f_2, f_3\} - 4\Delta.
 \end{aligned}$$

2. When $x^* \in (x_1, x_2]$, let $\epsilon = \frac{x_3 - x_2}{x_3 - x^*}$, and the proof goes the same way as in (1).
3. When $x^* \in (x_2, x_3]$, we let $\epsilon = \frac{x_2 - x_1}{x^* - x_1}$ and we know that $\epsilon \in [\frac{1}{2}, 1]$. Since $x_2 = \epsilon \cdot x^* + (1 - \epsilon)x_1$, we have $\epsilon f(x^*) + (1 - \epsilon)f(x_1) \geq f(x_2)$ according to the convexity of $f(x)$. Therefore, we have:

$$\begin{aligned}
 f(x^*) &\geq \frac{f_2 - (1 - \epsilon)f_1}{\epsilon} \\
 &= f_1 + \frac{f_2 - f_1}{\epsilon} \\
 &= f_1 - \frac{f_1 - f_2}{\epsilon}.
 \end{aligned} \tag{56}$$

If $f_1 \leq f_2$, then we have $f(x^*) \geq f_1 \geq A - \Delta = A + \Delta - 2\Delta \geq \max\{f_1, f_2, f_3\} - 2\Delta$.
 Otherwise $f_1 > f_2$, then we have

$$\begin{aligned}
 f(x^*) &\geq f_1 - \frac{f_1 - f_2}{\epsilon} \\
 &\geq f_1 - \frac{f_1 - f_2}{\frac{1}{2}} \\
 &= 2f_2 - f_1 \\
 &\geq 2(A - \Delta) - (A + \Delta) \\
 &= A - 3\Delta \\
 &= A + \Delta - 4\Delta \\
 &\geq \max\{f_1, f_2, f_3\} - 4\Delta.
 \end{aligned}$$

4. When $x^* \in (x_3, b]$, let $\epsilon = \frac{x_3 - x_2}{x^* - x_2}$, and the proof goes the same way as (3).

■

A.9 Proof of Lemma A.10

Proof. Notice that

$$-p_{\max} I_{\max} \leq -p \cdot \|\vec{I}\|_1 \leq Q_t(\vec{I}, p) \leq \langle \vec{\gamma}, \vec{I} \rangle \leq \gamma_{\max} I_{\max} \quad (57)$$

Denote $Q_{\max} := \max\{p_{\max}, \gamma_{\max}\} I_{\max}$. By applying Hoeffding's Inequality to $\forall \vec{I}, \hat{p}_\tau \in \{a_{K,\tau}, b_{K,\tau}, c_{K,\tau}\}$, we have

$$\begin{aligned} \Pr[|Q_{K,\tau,s}(\vec{I}, \hat{p}_\tau) - Q(\vec{I}, \hat{p}_\tau)| \geq x] &< 2 \exp\left\{-\frac{2x^2 n_s}{Q_{\max}^2}\right\} \\ \Rightarrow |Q_{K,\tau,s}(\vec{I}, \hat{p}_\tau) - Q(\vec{I}, \hat{p}_\tau)| &\leq \frac{Q_{\max} \sqrt{2 \log \frac{2}{\hat{\epsilon}}}}{2\sqrt{n_s}} \text{ with } \Pr \geq 1 - \hat{\epsilon}. \end{aligned} \quad (58)$$

Let $C_c = Q_{\max} \sqrt{2 \log \frac{2}{\hat{\epsilon}}}$ and this completes the proof. ■