# Nonlocal prior mixture-based Bayesian wavelet regression

Nilotpal Sanyal

Department of Mathematical Sciences, University of Texas at El Paso

#### Abstract

We propose a novel Bayesian wavelet regression approach using a three-component spike-and-slab prior for wavelet coefficients, combining a point mass at zero, a moment (MOM) prior, and an inverse moment (IMOM) prior. This flexible prior supports small and large coefficients differently, offering advantages for highly dispersed data where wavelet coefficients span multiple scales. The IMOM prior's heavy tails capture large coefficients, while the MOM prior is better suited for smaller non-zero coefficients. Further, our method introduces innovative hyperparameter specifications for mixture probabilities and scaling parameters, including generalized logit, hyperbolic secant, and generalized normal decay for probabilities, and double exponential decay for scaling. Hyperparameters are estimated via an empirical Bayes approach, enabling posterior inference tailored to the data. Extensive simulations demonstrate significant performance gains over two-component wavelet methods. Applications to electroencephalography and noisy audio data illustrate the method's utility in capturing complex signal characteristics. We implement our method in an R package NLPwavelet.

*Keywords:* Three component spike-and-slab prior; wavelet analysis; nonlocal prior; generalized logit decay; hyperbolic secant decay; generalized normal decay.

### 1 Introduction

We focus on spike-and-slab mixture models for wavelet-based Bayesian nonparametric regression. There are several existing approaches that consider for the wavelet coefficients various spike-and-slab mixture models such as mixtures of two Gaussian distributions with different standard deviations (Chipman et al., 1997), mixtures of a Gaussian and a point mass at zero (Clyde et al., 1998; Abramovich et al., 1998), mixtures of a heavy-tailed distribution and a point mass at zero (Johnstone and Silverman, 2005), and mixtures of a nonlocal prior and a point mass at zero (Sanyal and Ferreira, 2017). In contrast to the other methods that used traditional local priors for the wavelet coefficients, Sanyal and Ferreira (2017) pioneered the use of nonlocal priors (Johnson and Rossell, 2010) for wavelet regression. Specifically, they used two different nonlocal priors, namely the moment prior (MOM) and the inverse moment (IMOM) prior in their mixture model. In this work, we flexibly extend all the previous approaches by proposing a three-component spike-andslab mixture model for the wavelet coefficients where along with a point mass for the spike part, a mixture of nonlocal priors is used to model the slab component. We introduce novel hyperparameter specifications that are shown to provide improved estimates in extensive simulation experiments with highly dispersed data.

In the Bayesian paradigm, nonlocal priors have been shown to encourage model selection parsimony and selective shrinkage for spurious coefficients (Johnson and Rossell, 2012; Rossell and Telesca, 2017). Further, previous work (Sanyal et al., 2019) has observed, in the context of high-dimensional genomics data, that different nonlocal priors provide better support for large and small regression coefficients. In wavelet regression, the wavelet coefficients at multiple resolution levels capture both location and scale characteristics of the underlying function (Mallat, 2008). If the underlying function is highly dispersed, its energy is spread across a wide range of location or scale components. For such a function, while most of the wavelet coefficients will likely be small or near-zero, non-zero wavelet coefficients will span across multiple scales. In other words, no single scale will dominate the wavelet coefficients, as different scales will capture different portions of the signal's energy. This will lead to significant coefficients at both coarse and fine scales. We conjecture that, if the underlying function is highly dispersed, a mixture of non-local priors will provide better support to the distribution of the wavelet coefficients compared to individual non-local priors. Specifically, in this work, we consider for the wavelet coefficients a prior that is a mixture of a point mass at zero, a MOM prior, and an IMOM prior. The IMOM prior has heavier tails that is expected to provide better support for larger coefficients whereas the MOM prior is expected to provide better support for smaller non-zero coefficients.

In Bayesian wavelet regression, the probability weights associated with different component distributions of a spike-and-slab mixture prior, henceforth called *mixture probabilities*, and the scaling parameters of the component distributions are often governed by multiple hyperparameters (Abramovich et al., 1998; Clyde et al., 1998; Sanyal and Ferreira, 2017). For the mixture probabilities at different resolution levels, previous work considered exponential decay specification (Abramovich et al., 1998), Bernoulli distribution (Clyde et al., 1998), and logit specification (Sanyal and Ferreira, 2017). Further, for the scaling parameters, exponential decay (Abramovich et al., 1998) and polynomial decay specifications (Sanyal and Ferreira, 2017) have been considered. In this work, we propose several novel specifications that flexibly model the variations in the mixture probabilities and scaling components. For the mixture probabilities, we consider a generalized logit decay, a hyperbolic secant decay, and a generalized normal decay, whereas for the scaling parameter, we consider a double exponential decay. Each of these specifications are controlled by a few hyperparameters. Following an empirical-Bayes approach, we estimate the hyperparameters from the data and develop posterior inference conditional on the hyperparameter estimates.

Through extensive simulation studies, we assess the performance gains of our approach

under the different hyperparameter specifications, comparing it to two-component spikeand-slab prior-based wavelet methods. Finally, applications to real-world data, including electroencephalography data from a meditation study and audio data from a noisy musical recording, illustrate the practical utility of the proposed method.

In what follows, Section 2 describes the proposed Bayesian hierarchical model along with the hyperparameter specifications and Section 3 describes the empirical Bayes inference procedure. Subsequently, Section 4 describes the simulation experiment and results. The real data applications appear in Section 5 and Section 6. Finally, Section 7 concludes with an overall discussion of our work and relevant remarks. Theoretical proofs are included in the Appendix.

### 2 Bayesian hierarchical model

#### **Observation model**

Suppose  $y_1, \ldots, y_n$  represent *n* noisy observations from an unknown function f(t), which we aim to estimate. We consider the observation model  $y_i = f(t_i) + \epsilon_i$ ,  $i = 1, \ldots, n$ , where  $t_i = i/n$  represents the equispaced sampling points, and the errors  $\epsilon_i, \ldots, \epsilon_n$  are assumed to be independent and identically distributed (i.i.d.) normal random variables,  $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$ , with unknown variance  $\sigma^2$ . Let  $\mathbf{y} = (y_1, \ldots, y_n)$  denote the vector of observations,  $\mathbf{f} = (f(t_1), \ldots, f(t_n))$  the vector of true functional values, and  $\boldsymbol{\epsilon} = (\epsilon_1, \ldots, \epsilon_n)$ the vector of errors. The observation model can be expressed in matrix form as:

$$\mathbf{y} = \mathbf{f} + \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_n), \tag{1}$$

where  $\mathbf{I}_n$  is the *n*-dimensional identity matrix. In wavelet regression, the goal is to estimate the function f(t) from the observations  $\mathbf{y}$  by decomposing f(t) into wavelet basis functions. This decomposition allows us to exploit the multiscale nature of wavelets to capture both local and global features of the function. A chief feature of wavelet regression is its ability to adapt to different levels of smoothness and to handle noisy data efficiently, making it especially useful in non-parametric function estimation problems.

#### Wavelet coefficient model

We represent  $\mathbf{f}$  using an orthogonal wavelet basis matrix as  $\mathbf{f} = \mathbf{W}\mathbf{d}$  (Donoho and Johnstone, 1995), where  $\mathbf{W}$  is the orthogonal basis matrix and  $\mathbf{d}$  is a vector whose elements include the scaling coefficient at the coarsest resolution level along with the wavelet coefficients at all resolution levels. Let  $\hat{\mathbf{d}} = \mathbf{W}^T \mathbf{y}$  denote the vector of empirical wavelet coefficients. Since  $\mathbf{W}$  is orthogonal, we can express  $\hat{\mathbf{d}}$  as  $\hat{\mathbf{d}} = \mathbf{d} + \boldsymbol{\epsilon}^*$ , where  $\boldsymbol{\epsilon}^* = \mathbf{W}^T \boldsymbol{\epsilon}$ represents the transformed error vector with  $\boldsymbol{\epsilon}^* \sim \mathcal{N}(0, \sigma^2 \mathbf{I}_n)$ .

Suppose  $d_{lj}$  denotes the wavelet coefficient at position j and resolution level l, with  $\hat{d}_{lj}$  defined similarly for the empirical wavelet coefficients. Then, the model in terms of individual coefficients is given by

$$\widehat{d}_{lj} = d_{lj} + \epsilon_{lj}^*, \quad \epsilon_{lj}^* \sim N(0, \sigma^2)$$
(2)

#### Mixture prior for the wavelet coefficients

For the wavelet coefficient  $d_{lj}$ , we consider a spike-and-slab mixture prior that is a mixture of three components—a MOM prior, an IMOM prior, and a point mass at zero—given by

$$d_{lj}|\gamma_l^{(1)},\gamma_l^{(2)},\tau_l^{(1)},\tau_l^{(2)},\sigma^2,r,\nu \sim \gamma_l^{(1)}MOM\left(\tau_l^{(1)},r,\sigma^2\right) + \left(1-\gamma_l^{(1)}\right)\gamma_l^{(2)}IMOM\left(\tau_l^{(2)},\nu,\sigma^2\right) \\ + \left(1-\gamma_l^{(1)}\right)\left(1-\gamma_l^{(2)}\right)\delta_0(\cdot), \quad 0 < \gamma_l^{(1)},\gamma_l^{(2)} < 1, \quad (3)$$

where  $\gamma_1^{(l)}$  and  $\gamma_l^{(2)}$  are mixture probabilities, the *MOM* prior with order *r* and variance component  $\tau_l^{(1)}\sigma^2$ , involving the scale parameter  $\tau_l^{(1)}$ , has the density function

$$mom(d_{lj}|\tau_l^{(1)}, r, \sigma^2) = \widetilde{M}_r \left(\tau_l^{(1)} \sigma^2\right)^{-r-1/2} d_{lj}^{2r} \exp\left(-\frac{d_{lj}^2}{2\tau_l^{(1)} \sigma^2}\right), \quad r > 1, \tau_l^{(1)} > 0, \\ \widetilde{M}_r = \frac{(2\pi)^{-1/2}}{(2r-1)!!},$$

with  $(2r-1)!! = 1 \times 3 \times \ldots \times (2r-1)$ , and the *IMOM* prior with shape parameter  $\nu$  and

variance component  $\tau_l^{(2)}\sigma^2$ , involving the scale parameter  $\tau_l^{(2)}$ , has the density function

$$imom(d_{lj}|\tau_l^{(2)},\nu,\sigma^2) = \frac{\left(\tau_l^{(2)}\sigma^2\right)^{\nu/2}}{\Gamma(\nu/2)} |d_{lj}|^{-\nu-1} \exp\left(-\frac{\tau_l^{(2)}\sigma^2}{d_{lj}^2}\right), \quad \nu > 1, \tau_l^{(2)} > 0.$$

The conditional density of the wavelet coefficient  $d_{lj}$  given that  $d_{lj} \neq 0$  can be expressed as

$$\pi(d_{lj}|d_{lj} \neq 0) = \frac{\gamma_l^{(1)}}{\gamma_l^{(1)} + \left(1 - \gamma_l^{(1)}\right)\gamma_l^{(2)}} \widetilde{M}_r \left(\tau_l^{(1)}\sigma^2\right)^{-r-1/2} d_{lj}^{2r} \exp\left(-\frac{d_{lj}^2}{2\tau_l^{(1)}\sigma^2}\right)$$
$$\frac{\left(1 - \gamma_l^{(1)}\right)\gamma_l^{(2)}}{\gamma_l^{(1)} + \left(1 - \gamma_l^{(1)}\right)\gamma_l^{(2)}} \frac{\left(\tau_l^{(2)}\sigma^2\right)^{\nu/2}}{\Gamma(\nu/2)} |d_{lj}|^{-\nu-1} \exp\left(-\frac{\tau_l^{(2)}\sigma^2}{d_{lj}^2}\right).$$

Figure 1 shows plots of our proposed three-component spike-and-slab mixture model (solid line) for  $d_{lj}$  with  $\gamma_{1l} = \gamma_{2l} = 0.25$ ,  $\tau_{1l} = \tau_{2l} = 0.2$ ,  $r = \nu = 1$ , and  $\sigma = 1$  along with two-component spike-and-slab mixture models based on MOM (dashed line) and IMOM (dotted line) priors with  $\gamma_l = 0.5$ ,  $\tau_l = 1$ , and  $\sigma = 1$ .

#### Hyperparameter specifications

The mixture prior given in (3) depends on the mixture probabilities  $\gamma_1^{(l)}$  and  $\gamma_l^{(2)}$ , scale parameters  $\tau_l^{(1)}$  and  $\tau_l^{(1)}$ , order r, and shape parameter  $\nu$ . Previous work in wavelet regression considered two-component spike-and-slab mixture priors, where the mixture probabilities were specified using exponential decay specification (Abramovich et al., 1998), Bernoulli distribution (Clyde et al., 1998), and logit specification (Sanyal and Ferreira, 2017). In this work, we examine three specifications for the mixture probabilities that flexibly model the variations in the mixture probabilities and to the best of our knowledge are hitherto ununsed in wavelet regression. We compare these novel specifications with the previously proposed logit specification for  $\gamma_l^{(1)}$  and  $\gamma_l^{(2)}$ , given by  $\gamma_l^{(1)} = \exp(\theta_1^{\gamma} - \theta_2^{\gamma} l) / \{1 + \exp(\theta_1^{\gamma} - \theta_2^{\gamma} l)\}, \gamma_l^{(2)} = \exp(\theta_3^{\gamma} - \theta_4 l) \{1 + \exp(\theta_3^{\gamma} - \theta_4^{\gamma} l)\}$ , where  $\theta_1^{\gamma}, \theta_3^{\gamma} \in \mathbb{R}, \ \theta_2^{\gamma}, \theta_4^{\gamma} > 0$  (Sanyal and Ferreira, 2017). Our novel specifications are described as follows:



Figure 1: Plots of our proposed three-component spike-and-slab mixture model (solid line) with  $\gamma_{1l} = \gamma_{2l} = 0.25$ ,  $\tau_{1l} = \tau_{2l} = 0.2$ ,  $r = \nu = 1$ , and  $\sigma = 1$  along with two-component spike-and-slab mixture models based on MOM (dashed line) and IMOM (dotted line) priors with  $\gamma_l = 0.5$ ,  $\tau_l = 1$ , and  $\sigma = 1$ .

(a) Generalized logit or Richards decay specifications, given by

$$\begin{split} \gamma_l^{(1)} &= \frac{1}{[1 + \exp\{-(\theta_1^{\gamma} - \theta_2^{\gamma}l)\}]^{\theta_3^{\gamma}}}, \quad \theta_1^{\gamma} \in \mathbb{R}, \ \theta_2^{\gamma}, \theta_3^{\gamma} > 0\\ \gamma_l^{(2)} &= \frac{1}{[1 + \exp\{-(\theta_4^{\gamma} - \theta_5^{\gamma}l)\}]^{\theta_6^{\gamma}}}, \quad \theta_4^{\gamma} \in \mathbb{R}, \ \theta_5^{\gamma}, \theta_6^{\gamma} > 0; \end{split}$$

(b) Hyperbolic secant decay specifications, given by

$$\gamma_l^{(1)} = \frac{2}{\pi} \arctan\left[\exp\left(\frac{\pi}{2} \left(\theta_1^{\gamma} - \theta_2^{\gamma} l\right)\right)\right], \quad \theta_1^{\gamma} \in \mathbb{R}, \ \theta_2^{\gamma} > 0$$
  
$$\gamma_l^{(2)} = \frac{2}{\pi} \arctan\left[\exp\left(\frac{\pi}{2} \left(\theta_3^{\gamma} - \theta_4^{\gamma} l\right)\right)\right], \quad \theta_3^{\gamma} \in \mathbb{R}, \ \theta_4^{\gamma} > 0;$$

(c) Generalized normal decay specifications, given by

$$\gamma_l^{(1)} = \frac{1}{2} + \operatorname{sign}(\theta_1^{\gamma} - l) \frac{1}{2\Gamma(1/\theta_2^{\gamma})} \gamma \left( 1/\theta_2^{\gamma}, \left| \frac{\theta_1^{\gamma} - l}{\theta_3^{\gamma}} \right|^{\theta_2^{\gamma}} \right), \quad \theta_1^{\gamma} \in \mathbb{R}, \ \theta_2^{\gamma}, \theta_3^{\gamma} > 0$$
$$\gamma_l^{(2)} = \frac{1}{2} + \operatorname{sign}(\theta_4^{\gamma} - l) \frac{1}{2\Gamma(1/\theta_5^{\gamma})} \gamma \left( 1/\theta_5^{\gamma}, \left| \frac{\theta_4^{\gamma} - l}{\theta_6^{\gamma}} \right|^{\theta_5^{\gamma}} \right), \quad \theta_4^{\gamma} \in \mathbb{R}, \ \theta_5^{\gamma}, \theta_6^{\gamma} > 0$$

For scale parameters of spike-and-slab mixture priors, previously considered specifications include exponential decay (Abramovich et al., 1998) and polynomial decay (Sanyal and Ferreira, 2017). Here, for  $\tau_l^{(1)}$  and  $\tau_l^{(2)}$ , we consider the polynomial decay specification given by  $\tau_l^{(1)} = \theta_1^{\tau} l^{-\theta_2^{\tau}}, \tau_l^{(2)} = \theta_3^{\tau} l^{-\theta_4^{\tau}}$  with  $\theta_1^{\tau}, \theta_2^{\tau}, \theta_3^{\tau}, \theta_4^{\tau} > 0$ . In addition, for modeling the variations in the scale parameters more flexibly, we novelly propose

(d) Double exponential decay specifications, given by

$$\begin{aligned} \tau_l^{(1)} &= \theta_1^\tau \exp(-\theta_2^\tau l) + \theta_3^\tau \exp(-\theta_4^\tau l), \quad \theta_1^\tau, \theta_2^\tau, \theta_3^\tau, \theta_4^\tau > 0 \\ \tau_l^{(2)} &= \theta_5^\tau \exp(-\theta_6^\tau l) + \theta_7^\tau \exp(-\theta_8^\tau l), \quad \theta_5^\tau, \theta_6^\tau, \theta_7^\tau, \theta_8^\tau > 0 \end{aligned}$$

Each of the above existing and novel specifications are controlled by a few hyperparameters. Note that, the hyperparameters for the mixture probabilities are superscripted with  $\gamma$  and those for the scale parameters are superscripted with  $\tau$ . Figure 2 shows the plots of the different specifications for the mixture probabilities (left) and scale parameters (right)



Figure 2: Plots of the different specifications considered for the mixture probabilities  $\gamma_l^{(1)}$ and  $\gamma_l^{(2)}$  (logit, generalized logit, hyperbolic secant, and generalized normal) and scale parameters  $\tau_l^{(1)}$  and  $\tau_l^{(2)}$  (polynomial decay and double exponential decay) against resolution level, with specified values of the hyperparameters.

against resolution level, with specific values of the hyperparameters. In our simulation studies described in Section 4, we analyze each simulated dataset using  $4 \times 2 = 8$  configurations arising out of the combinations of above specifications for the mixture probabilities and the scale parameters. Let  $\boldsymbol{\theta}$  generically denote the set of all hyperparameters for any given configuration.

## 3 Inference

For inference in our proposed Bayesian hierarchical wavelet regression model based on three-component spike-and-slab mixture priors, we adopt the empirical Bayes approach (Clyde and George, 2000; Sanyal and Ferreira, 2017). This methodology estimates the hyperparameters from the data and performs posterior inference conditioned on these estimated hyperparameters. For simplicity, in our simulation analysis, we set r = 1 and  $\nu = 1$ . Further, we estimate the error variance  $\sigma^2$  using the median absolute deviation estimator (Donoho et al., 1995)  $\hat{\sigma} = 0.6745^{-1} \operatorname{median}_j(|\hat{d}_{Lj} - \operatorname{median}_j(\hat{d}_{Lj})|)$ , which is an well-established practice in wavelet regression (Abramovich et al., 1998; Clyde et al., 1998; Clyde and George, 2000; Johnstone and Silverman, 2005).

#### Hyperparameter estimation

Let  $\delta(x)$  denote the value of the point mass function  $\delta(0)$  at x. The following result is used to obtain the hyperparameter estimates.

**Result 1** Integrating out  $d_{lj}$  from the wavelet coefficient model (2) using the mixture prior of the wavelet coefficients in (3) and using the Laplace approximation for the IMOM prior component, we get the marginal distribution of the empirical wavelet coefficients,  $\hat{d}_{lj}$ , as

$$\pi(\widehat{d}_{lj}|\sigma^{2},\boldsymbol{\theta},r,\nu) \approx \gamma_{l}^{(1)} \left(1+\tau_{l}^{(1)}\right)^{-r} M_{r}^{*}(\widehat{d}_{lj},\tau_{l}^{(1)},\sigma^{2}) \phi\left(\widehat{d}_{lj};0,\sigma^{2}\left(1+\tau_{l}^{(1)}\right)\right) + \left(1-\gamma_{l}^{(1)}\right) \gamma_{l}^{(2)} \frac{\left(\tau_{l}^{(2)}\sigma^{2}\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)} \phi\left(\widehat{d}_{lj};0,\sigma^{2}\right) \sqrt{2\pi}\sigma_{*}h(d_{lj}^{*}(\widehat{d}_{lj})) + \left(1-\gamma_{l}^{(1)}\right) \left(1-\gamma_{l}^{(2)}\right) \phi\left(\widehat{d}_{lj};0,\sigma^{2}\right),$$
(4)

where, in the offshoot of the MOM component,

$$M_r^{\star}\left(\widehat{d}_{lj},\tau_l,\sigma^2\right) = \frac{1}{(2r-1)!!} \sum_{i=0}^r \frac{(2r)!}{(2i)!(r-i)!2^{r-i}} \left(\sqrt{\frac{\tau_l^{(1)}}{1+\tau_l^{(1)}}} \frac{\widehat{d}_{lj}}{\sigma}\right)^{2i},$$

and in the offshoot of the IMOM component,

$$h(d_{lj}) = |d_{lj}|^{-(\nu+1)} \exp\left\{-\frac{1}{2\sigma^2} \left(d_{lj}^2 - 2d_{lj}\widehat{d}_{lj}\right) - \frac{\tau_l \sigma^2}{d_{lj}^2}\right\},\,$$

 $d_{lj}^*(\widehat{d}_{lj})$  is the global maxima of  $h(d_{lj})$ , and  $\sigma_*^2 = -1/L_h''(d_{lj}^*(\widehat{d}_{lj}))$ , with  $L_h(d_{lj}) = \log(h(d_{lj}))$ .

The proof of Result 1 is given in the Appendix. Using (4), the marginal likelihood function

is approximately given by

$$\begin{split} \prod_{l} \prod_{j} \left[ \gamma_{l}^{(1)} \left( 1 + \tau_{l}^{(1)} \right)^{-r} M_{r}^{*}(\widehat{d}_{lj}, \tau_{l}^{(1)}, \sigma^{2}) \phi \left( \widehat{d}_{lj}; 0, \sigma^{2} \left( 1 + \tau_{l}^{(1)} \right) \right) \right. \\ \left. + \left( 1 - \gamma_{l}^{(1)} \right) \gamma_{l}^{(2)} \frac{\left( \tau_{l}^{(2)} \sigma^{2} \right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)} \phi \left( \widehat{d}_{lj}; 0, \sigma^{2} \right) \sqrt{2\pi} \sigma_{*} h(d_{lj}^{*}(\widehat{d}_{lj})) \right. \\ \left. + \left( 1 - \gamma_{l}^{(1)} \right) \left( 1 - \gamma_{l}^{(2)} \right) \phi \left( \widehat{d}_{lj}; 0, \sigma^{2} \right) \right]. \end{split}$$

This is a function only of the data and the hyperparameters. Hence, we maximize this function with respect to the hyperparameters to obtain their estimates,  $\hat{\theta}$ . With  $\theta = \hat{\theta}$ , the prior distribution in (3) is fully known.

#### Posterior distribution

The following two results provide the posterior estimates of the wavelet coefficients.

**Result 2** The conditional posterior density of the wavelet coefficient  $d_{lj}$ , given that  $d_{lj} \neq 0$ and the hyperparameter estimates  $\hat{\theta}$ , by using the Laplace approximation for the IMOM prior component, can be expressed as

$$\begin{aligned} \pi(d_{lj}|d_{lj} \neq 0, \sigma^2, \boldsymbol{\theta}, r, \nu, \mathbf{y}) = & \frac{p_{lj}^{(1)}}{p_{lj}^{(1)} + p_{lj}^{(2)}} \frac{\widetilde{M}_r}{M_r^* \left(\widehat{d}_{lj}, \tau_l, \sigma^2\right)} d_{lj}^{2r} \exp\left\{-\frac{1}{2\sigma^2 \frac{\tau_l^{(1)}}{\left(1 + \tau_l^{(1)}\right)}} \left(d_{lj} - \frac{\tau_l^{(1)}}{1 + \tau_l^{(1)}}\widehat{d}_{lj}\right)^2\right\} \\ &+ \frac{p_{lj}^{(2)}}{p_{lj}^{(1)} + p_{lj}^{(2)}} \phi\left(d_{lj}|d_{lj}^*(\widehat{d}_{lj}), \sigma_*^2\right),\end{aligned}$$

where

$$p_{lj}^{(1)} = \frac{O_{lj}^{(1)}}{1 + O_{lj}^{(1)} + O_{lj}^{(2)}}, \qquad p_{lj}^{(2)} = \frac{O_{lj}^{(2)}}{1 + O_{lj}^{(1)} + O_{lj}^{(2)}},$$
$$O_{lj}^{(1)} = \frac{\gamma_l^{(1)}}{\left(1 - \gamma_l^{(1)}\right) \left(1 - \gamma_l^{(2)}\right)} \left(1 + \tau_l^{(1)}\right)^{-r-1/2} M_r^* \exp\left\{\frac{1}{2\sigma^2} \frac{\tau_l^{(1)}}{1 + \tau_l^{(1)}} \hat{d}_{lj}^2\right\},$$

and

$$O_{lj}^{(2)} = \frac{\gamma_l^{(2)}}{1 - \gamma_l^{(2)}} \frac{\left(\tau_l^{(2)} \sigma^2\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)} \sqrt{2\pi} \sigma_* h(d_{lj}^*(\widehat{d}_{lj})).$$

The proof is given in the Appendix.

**Result 3** The posterior expectation of the wavelet coefficients  $d_{lj}$  is

$$\bar{d}_{lj} = E(d_{lj}|y) = p_{lj}^{(1)} \frac{M_r^{**}}{M_r^*} \sqrt{\frac{\tau_l^{(1)}}{1 + \tau_l^{(1)}}\sigma + p_{lj}^{(2)}d_{lj}^*(\hat{d}_{lj})},$$

where

$$M_r^{\star\star}\left(\widehat{d}_{lj},\tau_l,\sigma^2\right) = \frac{1}{(2r-1)!!} \sum_{i=1}^{r+1} \frac{(2r+1)!}{(2i-1)!(r+1-i)!2^{r+1-i}} \left(\sqrt{\frac{\tau_l^{(1)}}{1+\tau_l^{(1)}}} \frac{\widehat{d}_{lj}}{\sigma}\right)^{2i-1}.$$

The proof of Result 3 is immediate from the proof of Result 2 and hence is ommitted. Let  $\mathbf{\bar{d}} = d_{lj}$ : l = 1, ..., L, j = 1..., J denote the vector of posterior means of the wavelet coefficients. We use  $\mathbf{\bar{d}}$  in the inverse discrete wavelet transform to obtain the posterior mean of the unknown function f, given by

$$\bar{f} = E(\mathbf{f}|\mathbf{y}) = WE(\mathbf{d}|\mathbf{y}) = W\bar{\mathbf{d}}.$$

### 4 Simulation study

In this section, we conduct an extensive simulation analysis to comparatively evaluate the performance of the proposed method with different hyperparameter configurations. We consider three well-known test functions proposed by Donoho and Johnstone (1994)—*blocks*, *bumps*, and *doppler*—that are used as standard test functions in the wavelet literature. However, we modify the coefficients used by Donoho and Johnstone (1994) so as to obtain more highly dispersed signals. We define our test functions as:

$$f_{blocks}(t) = \sum h_j K(t - t_j)$$
, where  $K(t) = \{1 + \text{sign}(t)\}/2$ ,

(tj) = (0.1, 0.13, 0.15, 0.23, 0.25, 0.40, 0.44, 0.65, 0.76, 0.78, 0.81),(hj) = (4, -8, 3, -4, 8, -4.2, 2.1, 4.3, -6.1, 2.1, -4.7).

$$f_{bumps}(t) = \sum h_j K((t - t_j)/w_j)$$
, where  $K(t) = (1 + |t|)^{-4}$ ,

$$(t_j) = (0.1, 0.13, 0.15, 0.23, 0.25, 0.40, 0.44, 0.65, 0.76, 0.78, 0.81),$$
  

$$(h_j) = (2, 10, 1, 4, 8, 4.2, 2.1, 4.3, 1.1, 3.1, 8.2),$$
  

$$(w_j) = (0.005, 0.005, 0.006, 0.01, 0.01, 0.03, 0.01, 0.01, 0.005, 0.008, 0.005).$$

$$f_{doppler}(t) = \{t(1-t)\}^{1/2} \sin\{2\pi(1+\epsilon)/(t+\epsilon)\}, \epsilon = 0.01.$$

In addition, we consider three different linear combinations of these functions—*lcomb1*, *lcomb2*, and *lcomb3*—that represent curves that combine various features—such as, blockiness, bumpiness, and changing frequency—in different proportions, given by

$$f_{lcomb1}(t) = 0.4 f_{blocks}(t) + 0.4 f_{bumps}(t) + 0.2 f_{doppler}(t)$$
  
$$f_{lcomb2}(t) = 0.4 f_{blocks}(t) + 0.2 f_{bumps}(t) + 0.4 f_{doppler}(t)$$
  
$$f_{lcomb3}(t) = 0.2 f_{blocks}(t) + 0.4 f_{bumps}(t) + 0.4 f_{doppler}(t).$$

Figure 3 shows the plots of the test functions blocks, bumps, and doppler using Donoho-Johnstone specifications (dashed line) and our specifications (solid line), and *lcomb1*, *lcomb2*, and *lcomb3* based on our specifications, all evaluated at 1024 equally spaced in (0,1). We evaluate each considered test function at n = 512, 1024, 2048, and 4096 equidistant points in the interval (0,1) and add random Gaussian noise with mean 0 to generate data with signal-to-noise ratio, SNR = 3, 5, and 7. For each combination of n and SNR, we consider 10 replications. The simulated datasets are analyzed using the proposed methodology with 8 different hyperparameter configurations described in Section 2. For comparison, we also analyzed the datasets using individual MOM and IMOM prior-based two-component mixture models (Sanyal and Ferreira, 2017). Thus, a total of 24 analysis methods were applied to each dataset. For wavelet transformation, we consider the Daubechies least asymmetric wavelet with six vanishing moments and periodic boundary conditions. Wavelet computations are implemented using the R package *wavethresh* (Nason, 2024).

Table 1 presents, for each test function and analysis method, the number of (n, SNR) combinations where the method achieved the lowest MSE. For each test function, the



Figure 3: Plots of the test functions *blocks*, *bumps*, and *doppler* using the Donoho-Johnstone (DJ) specifications (dashed line) and our specifications (solid line), and three linear combinations of them—*lcomb1*, *lcomb2*, and *lcomb3*—based on our specifications, all evaluated at 1024 equally spaced in (0,1).

Table 1: Method comparison: Data were simulated for each test function across 12 combinations of sample size (n) and signal-to-noise ratio (SNR), with 10 replications per combination. A total of 24 analysis methods were applied to each dataset—8 methods with MOM-based two-component mixture prior, 8 methods with IMOM-based two-component mixture prior, 8 methods with IMOM-based two-component mixture prior. The average mean squared error (MSE) was computed for each method across the replications. The table presents, for each test function and analysis method, the number of (n, SNR) combinations where the method achieved the lowest MSE. For each test function, the method with the highest frequency of best performance is highlighted in bold.

Method	blocks	bumps	doppler	lcomb1	lcomb2	lcomb3	Total
mom-logit-polynom	0	0	0	0	0	0	0
$\operatorname{mom-logit-doubleexp}$	1	0	0	0	0	0	1
mom-genlogit-polynom	0	0	0	0	0	0	0
$\operatorname{mom-genlogit-doubleexp}$	1	0	0	0	0	0	1
mom-hypsec-polynom	1	0	0	0	1	0	2
mom-hypsec-doubleexp	2	0	0	0	0	0	2
mom-gennormal-polynom	0	0	0	0	0	0	0
$\operatorname{mom-gennormal-doubleexp}$	0	0	0	0	0	0	0
imom-logit-polynom	0	0	0	0	2	1	3
$\operatorname{imom-logit-doubleexp}$	0	1	0	0	0	0	1
imom-genlogit-polynom	0	0	0	1	0	0	1
$\operatorname{imom-genlogit-doubleexp}$	0	0	0	0	0	0	0
imom-hypsec-polynom	0	0	1	0	1	0	2
imom-hypsec-doubleexp	0	0	0	0	0	0	0
imom-gennormal-polynom	0	0	2	1	0	2	5
$imom\-gennormal\-doubleexp$	0	0	0	0	0	0	0
mixture-logit-polynom	0	5	0	1	3	0	9
mixture-logit-doubleexp	0	0	0	2	1	0	3
mixture-genlogit-polynom	3	1	0	0	1	1	6
mixture-genlogit-doubleexp	2	0	0	0	0	0	2
mixture-hypsec-polynom	1	0	4	2	1	0	8
mixture-hypsec-doubleexp	1	1	0	1	2	2	7
mixture-gennormal-polynom	0	4	3	1	0	2	10
$mixture\mbox{-gennormal-doubleexp}$	0	0	2	3	0	4	9
Total	12	12	12	12	12	12	72

method with the highest frequency of best performance (i.e., lowest MSE) is highlighted in bold. We observe the following:

- (a) For every function, the highest frequency of best performance was shown by a threecomponent mixture method, which is proposed in the current work.
- (b) Out of all 8 three-component mixture methods, the one with generalized normal specification for the mixture probabilities and polynomial decay specification for the scale parameters (*mixture-gennormal-polynom*) showed the maximum number of best performances in total (10 times) for all the test functions. This is followed by the method using generalized normal specification for the mixture probabilities and double exponential decay specification for the scale parameters (*mixture-gennormal-doubleexp*), as well as the method using logit specification for the mixture probabilities and polynomial decay specification for the scale parameters (*mixture-logit-polynom*), each showing best performance 9 times in total for all the test functions.
- (c) Considering only the test functions *lcomb1*, *lcomb2*, and *lcomb3*, that represent signals with mixed characteristics in various proportions, the *mixture-gennormal-doubleexp* method showed maximum number of best performances (7 times).

For a real data, SNR and the original function will not be known. So, next, in Figure 4, we show, for each of the 8 three-component mixture methods, MSE for different sample sizes (n), averaged over the 3 SNRs and 6 test functions considered in our study. Overall, the method with generalized logit and double exponential decay specifications (*mixture-genlogit-doubleexp*) showed best performance, followed by the method with hyperbolic secant and double exponential decay specifications (*mixture-hypsec-doubleexp*). Notably, the methods with generalized normal specifications, which showed the best performance in Table 1, showed worse performance in Figure 4 that used averaged MSEs. This implies the methods with generalized normal specifications, although most often provide



Figure 4: For the proposed three-component mixture-based methods, MSE for different sample sizes (n), averaged over 3 SNRs and 6 test functions.

best denoising, may sometimes produce large bias that negatively affects their overall average performance. So, the results obtained by their usage should be validated by field knowledge or external means (such as, auditory assessment for sound signals).

### 5 EEG meditation study

We demonstrate the utility and flexibility of the proposed methodolology through the analysis of electroencephalogram (EEG) data from a meditation study, investigating the relation between mind wandering and meditation practice. Detailed information about the study is provided in Brandmeyer and Delorme (2018), and the dataset is available on the *Open-Neuro* platform (see the Supplementary Appendix A for link). The meditation experiment involved 24 subjects—12 experienced meditators (10 males, 2 females) and 12 novices (2 males, 10 females). Participants meditated while being interrupted approximately every two minutes to report their level of concentration and mind wandering via three probing questions. Each participant completed 2 to 3 sessions lasting 45 to 90 minutes, with a minimum of 30 probes per participant. EEG data were collected using a 64-channel Biosemi system (channels A1–A32 and B1–B32) with a Biosemi 10-20 head cap montage at a sampling rate of 2048 Hz, providing spatial information about brain activity across different scalp regions. The dataset available on *OpenNeuro* has already been downsampled to 256 Hz.

For our analysis, we focused on data from session 1 of four participants: two expert meditators (one male and one female, identified as subjects 10 and 5 in the original dataset, respectively) and two novices (one male and one female, identified as subjects 23 and 14, respectively). The data were average-referenced to mitigate common noise and artifacts by calculating the mean signal across all electrodes at each time point and subtracting it from the signal at each electrode. Following this, a 2 Hz high-pass filter was applied using an infinite impulse response (IIR) filter with a transition bandwidth of 0.7 Hz and an order of 6. Denoting the time of probe 1 by t, we analyzed the EEG signal within the 16-second interval (t - 8, t + 8). With a sampling rate of 256 Hz, this interval comprised  $256 \times 16=4096$  data points, representing noisy observations of the underlying signal. We analyzed the EEG data using the *mixture-genlogit-doubleexp*, *mixture-hypsec-doubleexp*, and *mixture-gennormal-polynom* methods (see Section 4), employing Daubechies least asymmetric wavelet transform with 6 vanishing moments and periodic boundary conditions.

Figure 5 presents the plots of the posterior means of the EEG signal (in slategray) based on the mixture-genlogit-doubleexp (left column), mixture-hypsec-doubleexp (middle column), and mixture-gennormal-polynom (right column) methods, suporimposed on the observed data (in black), obtained during the 16-second interval from the A10 channel of the 4 considered participants. Similar plots for some other channels are presented in the Supplementary Appendix B. The plots clearly indicate that our method with all the considered hyperparameter configurations has yielded significantly denoised estimates, mixture-gennormal-polynom providing the strongest shrinkage.

### 6 Musical sound study

To further demonstrate the utility of the proposed method, we consider analyzing musical sounds that often exhibit sudden frequency changes over short periods, resulting in highly dispersed signals. For this study, we focus on a vocal music recording from 1934, originally published on a 78 RPM record of Hindustani classical music (one of India's two classical music traditions) and performed by eminent Ustad Amir Khan. A noisy copy of this recording was sourced from YouTube (link provided in the Supplementary Appendix A). From the recording, we extracted a 15-second segment representing a wide frequency range and saved it as a 16-bit WAV audio file, which represents noisy data.

For our analysis, we divided each of the two audio file channels (left and right) into sections of 4096 data points to enhance computational efficiency. Each section was independently analyzed using the *mixture-genlogit-doubleexp* and *mixture-gennormal-polynom* methods (see Section 4) employing the Daubechies coiflets wavelet transform with 5 vanishing moments and periodic boundary conditions. The posterior estimates of the individual sections were then combined to reconstruct the posterior estimate of the entire audio segment.

In Figure 6, we present the posterior mean of the right-channel signal of the selected audio segment (in slate gray), suporimposed on the noisy data (in black). The audio files corresponding to these posterior estimates along with the original data and posterior estimates using the hard thresholding rule have been provided in the Supplementary Material for auditory comparison and assessment. The posterior estimates show signifcant denoising and precise recovery of the signal. There is a slight systematic noise present in the posterior audio files which is due to analyzing the whole segment in disjoint sections for computational convenience. That can be mitigated by analyzing larger sections, or ana-



Figure 5: Plots of the posterior means of the EEG signal (in slate gray) based on the imomlogit-polynom (left column), mixture-genlogit-doubleexp (middle column), and mixturegennorm-polynom (right column) methods, suporimposed on the observed data (in black), obtained during the 16-second interval (t - 8, t + 8), t being probe 1 onset time, from the A10 channel of the 4 considered participants.



Figure 6: Plots of the posterior means (in slate gray) of the right-channel signal of the chosen audio segment of the vocal music recording, superimposed on the noisy data (in red).

lyzing partially overlapping sections with weighted averaging (e.g., cross-fading or Hann windowing) to smoothly combine overlapping regions and reduce boundary artifacts.

### 7 Discussion

In this work, we proposed the use of nonlocal prior mixtures for wavelet-based nonparametric function estimation. The main innovations of our methodology are as follows:

- (a) We introduce a three-component spike-and-slab prior for the wavelet coefficients. This structure is particularly suited for modeling highly dispersed signals. The slab component is a mixture of two nonlocal priors—the MOM (moment) and IMOM (inverse moment) priors, which offers enhanced adaptability to signal characteristics.
- (b) We propose flexible and previously unexplored hyperparameter specifications. These include generalized logit (or Richards), hyperbolic secant, and generalized normal de-

cay specifications for the mixture probabilities, as well as a double exponential decay structure for the scale parameter. These enhancements provide improved flexibility and accuracy in modeling complex signal patterns as demonstrated in our simulation study.

We implement our methodology within the R programming language (R Core Team, 2024) as a package named NLPWavelet. The R functions of the package along with the data analysis code for the simulation study and real data analyses are provided among the Supplementary Material.

In the simulation study, using more dispersed versions of the Donoho-Johnstone test functions and various linear combinations of them, we have compared the performance of the proposed approach with existing two-component spike-and-slab mixture prior, and demonstrated the superior flexibility of the proposed approach. Further, analysis using several novel hyperparameter configurations provided valuable insights into the relative advantages. Analysis of the real datasets illustrate the practical utility of the proposed methodology.

A practical issue with the proposed approach is increased computational cost. However, with high-performance computing resources, that can be mitigated. An obvious extension of the proposed approach is to adapt it to multidimensional wavelet regression, such 2D oe 3D image processing tasks. Another possible avenue is to develop fully Bayesian hierarchical models by specifying prior distributions for the hyperparameters and employing Markov chain Monte Carlo tools or variational methods for posterior inference. In addition, one can explore combining wavelet decompositions with local polynomial regression to mitigate the boundary bias issues. Further, adapting our methodology to non-Gaussian or skewed data can be an interesting enterpise. We leave all these to future research.

# Appendix

### Proof of Result 1

From the wavelet coefficient model (2) and the mixture prior of the wavelet coefficients in (3), using the Laplace approximation for the IMOM prior component, we get the marginal distribution of the empirical wavelet coefficients,  $\hat{d}_{lj}$ , as

$$\begin{split} &\pi(\widehat{d}_{lj}|\sigma^{2},\boldsymbol{\theta},r,\nu) \\ &= \int \pi(\widehat{d}_{lj}|d_{lj},\sigma^{2},\boldsymbol{\theta},r,\nu)\pi(d_{lj}|\sigma^{2},\boldsymbol{\theta},r,\nu)dd_{lj} \\ &= \int (2\pi\sigma^{2})^{-\frac{1}{2}} \exp\left\{-\frac{(\widehat{d}_{lj}-d_{lj})^{2}}{2\sigma^{2}}\right\} \left[\gamma_{l}^{(1)}\widetilde{M_{r}}\left(\tau_{l}^{(1)}\sigma^{2}\right)^{-r-\frac{1}{2}}d_{lj}^{2r}\exp\left(-\frac{d_{lj}^{2}}{2\tau_{l}^{(1)}\sigma^{2}}\right) \\ &\quad + \left(1-\gamma_{l}^{(1)}\right)\gamma_{l}^{(2)}\frac{\left(\tau_{l}^{(2)}\sigma^{2}\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)}|d_{lj}|^{-\nu-1}\exp\left(-\frac{\tau_{l}^{(2)}\sigma^{2}}{d_{lj}^{2}}\right) + \left(1-\gamma_{l}^{(1)}\right)\left(1-\gamma_{l}^{(2)}\right)\delta(d_{lj})\right] dd_{lj} \\ &= (2\pi\sigma^{2})^{-\frac{1}{2}}\left[\gamma_{l}^{(1)}\widetilde{M_{r}}\left(\tau_{l}^{(1)}\sigma^{2}\right)^{-r-\frac{1}{2}}\int d_{lj}^{2r}\exp\left\{-\frac{1}{2\sigma^{2}}\left((\widehat{d}_{lj}-d_{lj})^{2}+\frac{d_{lj}^{2}}{\tau_{l}^{(1)}}\right)\right\} dd_{lj} \\ &\quad + \left(1-\gamma_{l}^{(1)}\right)\gamma_{l}^{(2)}\frac{\left(\tau_{l}^{(2)}\sigma^{2}\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)}\int |d_{lj}|^{-\nu-1}\exp\left\{-\frac{(\widehat{d}_{lj}-d_{lj})^{2}}{2\sigma^{2}}-\frac{\tau_{l}^{(2)}\sigma^{2}}{d_{lj}^{2}}\right\} dd_{lj} \\ &\quad + \left(1-\gamma_{l}^{(1)}\right)\left(1-\gamma_{l}^{(2)}\right)\exp\left(-\frac{\widehat{d}_{lj}}{2\sigma^{2}}\right)\right] \\ &= (2\pi\sigma^{2})^{-\frac{1}{2}}\left[\gamma_{l}^{(1)}\widetilde{M_{r}}\left(\tau_{l}^{(1)}\sigma^{2}\right)^{-r-\frac{1}{2}}\exp\left\{-\frac{\widehat{d}_{lj}^{2}}{2\sigma^{2}\left(1+\tau_{l}^{(1)}\right)}\right\}\left(2\pi\sigma^{2}\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\right)^{1/2}\left(\sigma^{2}\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\right)^{r} \\ &\quad \sum_{i=0}^{r}\frac{(2r)!}{(2i)!(r-i)!2^{r-1}}\left(\frac{\frac{\tau_{l}^{(1)}}{\sqrt{\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}}}\right)^{2i}+\left(1-\gamma_{l}^{(1)}\right)\gamma_{l}^{(2)}\frac{\left(\tau_{l}^{(2)}\sigma^{2}\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)}\exp\left(-\frac{\widehat{d}_{lj}^{2}}{2\sigma^{2}}\right)\right] \\ &\quad (\text{using Laplace approximation) \\ &= \gamma_{l}^{(1)}\left(1+\tau_{l}^{(1)}\right)^{-r}M_{r}^{*}\left(\widehat{d}_{ij},\tau_{l}^{(1)},\sigma^{2}\right)\phi\left(\widehat{d}_{ij};0,\sigma^{2}\right). \end{split}$$

where

$$M_{r}^{\star}\left(\widehat{d}_{lj},\tau_{l},\sigma^{2}\right) = \frac{1}{(2r-1)!!} \sum_{i=0}^{r} \frac{(2r)!}{(2i)!(r-i)!2^{r-i}} \left(\sqrt{\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}} \frac{\widehat{d}_{lj}}{\sigma}\right)^{2i},$$
$$h(d_{lj}) = |d_{lj}|^{-(\nu+1)} \exp\left\{-\frac{1}{2\sigma^{2}} \left(d_{lj}^{2} - 2d_{lj}\widehat{d}_{lj}\right) - \frac{\tau_{l}\sigma^{2}}{d_{lj}^{2}}\right\},$$

 $d_{lj}^*(\widehat{d}_{lj})$  is the global maxima of  $h(d_{lj})$ , and  $\sigma_*^2 = -1/L_h''(d_{lj}^*(\widehat{d}_{lj}))$ , with  $L_h(d_{lj}) = \log(h(d_{lj}))$ .

# 7.1 Proof of Result 2

$$\pi\left(d_{lj}|\sigma^2,\boldsymbol{\theta},r,\nu,\mathbf{y}\right) \propto \pi\left(\mathbf{y}|d_{lj},\sigma^2,\boldsymbol{\theta},r,\nu\right) \ \pi\left(d_{lj}|\sigma^2,\boldsymbol{\theta},r,\nu\right).$$

The proportionality constant is

$$\begin{split} C &= \int \pi \left( \mathbf{y} | d_{lj}, \sigma^2, \boldsymbol{\theta}, r, \nu \right) \pi \left( d_{lj} | \sigma^2, \boldsymbol{\theta}, r, \nu \right) dd_{lj} \\ &= (2\pi\sigma^2)^{-\frac{1}{2}} \left[ \gamma_l^{(1)} \left( 1 + \tau_l^{(1)} \right)^{-r - \frac{1}{2}} M_r^* \left( \widehat{d}_{lj}, \tau_l^{(1)}, \sigma^2 \right) \exp \left\{ -\frac{\widehat{d}_{lj}^2}{2\sigma^2 \left( 1 + \tau_l^{(1)} \right)} \right\} \\ &+ \left( 1 - \gamma_l^{(1)} \right) \gamma_l^{(2)} \frac{\left( \tau_l^{(2)} \sigma^2 \right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)} \exp \left( -\frac{\widehat{d}_{lj}^2}{2\sigma^2} \right) \sqrt{2\pi} \sigma_* h(d_{lj}^*(\widehat{d}_{lj})) \\ &+ \left( 1 - \gamma_l^{(1)} \right) \left( 1 - \gamma_l^{(2)} \right) \exp \left( -\frac{\widehat{d}_{lj}^2}{2\sigma^2} \right) \right] \\ &= (2\pi\sigma^2)^{-\frac{1}{2}} \left( 1 - \gamma_l^{(1)} \right) \left( 1 - \gamma_l^{(2)} \right) \exp \left( -\frac{\widehat{d}_{lj}^2}{2\sigma^2} \right) \left[ O_{lj}^{(1)} + O_{lj}^{(2)} + 1 \right], \end{split}$$

Thus,

$$\begin{split} &\pi\left(d_{ij}|\sigma^{2},\boldsymbol{\theta},r,\nu,\mathbf{y}\right) \\ &= \frac{1}{C}\pi\left(\mathbf{y}|d_{ij},\sigma^{2},\boldsymbol{\theta},r,\nu\right)\pi\left(d_{ij}|\sigma^{2},\boldsymbol{\theta},r,\nu\right) \\ &= \frac{1}{(2\pi\sigma^{2})^{-\frac{1}{2}}\left(1-\gamma_{l}^{(1)}\right)\left(1-\gamma_{l}^{(2)}\right)\exp\left(-\frac{\hat{d}_{ij}^{2}}{2\sigma^{2}}\right)\left[O_{ij}^{(1)}+O_{ij}^{(2)}+1\right]} \\ &\left(2\pi\sigma^{2})^{-\frac{1}{2}}\left[\gamma_{l}^{(1)}\widetilde{M_{r}}\left(\tau_{l}^{(1)}\sigma^{2}\right)^{-r-\frac{1}{2}}\exp\left\{-\frac{\hat{d}_{ij}^{2}}{2\sigma^{2}\left(1+\tau_{l}^{(1)}\right)}\right\}d_{lj}^{2r}\exp\left\{-\frac{1}{2\sigma^{2}\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\left(d_{ij}-\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\hat{d}_{lj}\right)^{2}\right\} \\ &+\left(1-\gamma_{l}^{(1)}\right)\gamma_{l}^{(2)}\frac{\left(\tau_{l}^{(2)}\sigma^{2}\right)^{\frac{\nu}{2}}}{\Gamma(\nu/2)}\exp\left(-\frac{\hat{d}_{ij}^{2}}{2\sigma^{2}}\right)\sqrt{2\pi}\sigma_{*}h(d_{ij}^{*}(\hat{d}_{ij}))\phi\left(d_{ij}|d_{ij}^{*}(\hat{d}_{ij}),\sigma_{*}^{2}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij})\right. \\ &=\frac{O_{lj}^{(1)}}{O_{lj}^{(1)}+O_{lj}^{(2)}+1}\frac{\widetilde{M_{r}}}{M_{r}^{*}\left(\hat{d}_{ij},\tau,\sigma^{2}\right)}d_{ij}^{2r}\exp\left\{-\frac{1}{2\sigma^{2}\frac{\tau_{l}^{(1)}}{(1+\tau_{l}^{(1)})}}\left(d_{ij}-\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\hat{d}_{ij}\right)^{2}\right\} \\ &+\frac{O_{lj}^{(2)}}{O_{lj}^{(1)}+O_{lj}^{(2)}+1}\phi\left(d_{ij}|d_{ij}^{*}(\hat{d}_{ij}),\sigma_{*}^{2}\right)+\frac{1}{O_{lj}^{(1)}+O_{lj}^{(2)}+1}\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}) \\ &=p_{lj}^{(1)}\frac{\widetilde{M_{r}}}{M_{r}^{*}\left(\hat{d}_{ij},\tau_{l},\sigma^{2}\right)}d_{lj}^{2r}\exp\left\{-\frac{1}{2\sigma^{2}\frac{\tau_{l}^{(1)}}{(1+\tau_{l}^{(1)})}}\left(d_{ij}-\frac{\tau_{l}^{(1)}}{1+\tau_{l}^{(1)}}\hat{d}_{ij}\right)^{2}\right\}+p_{lj}^{(2)}\phi\left(d_{ij}|d_{ij}^{*}(\hat{d}_{ij}),\sigma_{*}^{2}\right) \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}-p_{lj}^{(2)}\right)\exp\left\{\frac{\hat{d}_{ij}d_{ij}}{\sigma^{2}}-\frac{d_{ij}^{2}}{2\sigma^{2}}\right\}\delta(d_{ij}), \\ &+\left(1-p_{lj}^{(1)}$$

from which Result 2 follows.

## References

- Abramovich, F., Sapatinas, T., and Silverman, B. (1998). "Wavelet thresholding via a Bayesian approach." Journal of the Royal Statistical Society – Series B, 60, 4, 725–749.
- Brandmeyer, T. and Delorme, A. (2018). "Reduced mind wandering in experienced meditators and associated EEG correlates." *Experimental brain research*, 236, 2519–2528.

Chipman, H. A., Kolaczyk, E. D., and McCulloch, R. E. (1997). "Adaptive Bayesian

Wavelet Shrinkage." Journal of the American Statistical Association, 92, 440, 1413–1421.

- Clyde, M. and George, E. I. (2000). "Flexible empirical Bayes estimation for wavelets." Journal of the Royal Statistical Society – Series B, 62, 4, 681–698.
- Clyde, M., Parmigiani, G., and Vidakovic, B. (1998). "Multiple shrinkage and subset selection in wavelets." *Biometrika*, 85, 2, 391–401.
- Donoho, D. and Johnstone, J. (1994). "Ideal spatial adaptation by wavelet shrinkage." Biometrika, 81, 3, 425–455.
- Donoho, D. L. and Johnstone, I. M. (1995). "Adapting to Unknown Smoothness via Wavelet Shrinkage." Journal of the American Statistical Association, 90, 432, 1200–1224.
- Donoho, D. L., Johnstone, I. M., Kerkyacharian, G., and Picard, D. (1995). "Wavelet Shrinkage: Asymptopia?" Journal of the Royal Statistical Society – Series B, 57, 2, pp. 301–369.
- Johnson, V. and Rossell, D. (2010). "On the use of non-local prior densities in Bayesian hypothesis tests." Journal of the Royal Statistical Society Series B, 72, 2, 143–170.
- (2012). "Bayesian model selection in high-dimensional settings." Journal of the American Statistical Association, 107, 498, 649–660.
- Johnstone, I. M. and Silverman, B. W. (2005). "Empirical Bayes Selection of Wavelet Thresholds." *The Annals of Statistics*, 33, 4, pp. 1700–1752.
- Mallat, S. (2008). A Wavelet Tour of Signal Processing, Third Edition: The Sparse Way.3rd ed. USA: Academic Press, Inc.
- Nason, G. (2024). *wavethresh: Wavelets Statistics and Transforms*. R package version 4.7.3.

- R Core Team (2024). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rossell, D. and Telesca, D. (2017). "Nonlocal priors for high-dimensional estimation." Journal of the American Statistical Association, 112, 517, 254–265.
- Sanyal, N. and Ferreira, M. A. (2017). "Bayesian wavelet analysis using nonlocal priors with an application to FMRI analysis." Sankhya B, 79, 2, 361–388.
- Sanyal, N., Lo, M.-T., Kauppi, K., Djurovic, S., Andreassen, O. A., Johnson, V. E., and Chen, C.-H. (2019). "GWASinlps: non-local prior based iterative SNP selection tool for genome-wide association studies." *Bioinformatics*, 35, 1, 1–11.