## A UNIFIED PERSPECTIVE ON THE DYNAMICS OF DEEP TRANSFORMERS

VALÉRIE CASTIN\*, PIERRE ABLIN<sup>†</sup>, JOSÉ A. CARRILLO<sup>‡</sup>, AND GABRIEL PEYRÉ\*

Abstract. Transformers, which are state-of-the-art in most machine learning tasks, represent the data as sequences of vectors called tokens. This representation is then exploited by the attention function, which learns dependencies between tokens and is key to the success of Transformers. However, the iterative application of attention across layers induces complex dynamics that remain to be fully understood. To analyze these dynamics, we identify each input sequence with a probability measure and model its evolution as a Vlasov equation called Transformer PDE, whose velocity field is non-linear in the probability measure. Our first set of contributions focuses on compactly supported initial data. We show the Transformer PDE is well-posed and is the mean-field limit of an interacting particle system, thus generalizing and extending previous analysis to several variants of self-attention: multi-head attention, L2 attention, Sinkhorn attention, Sigmoid attention, and masked attention—leveraging a conditional Wasserstein framework. In a second set of contributions, we are the first to study non-compactly supported initial conditions, by focusing on Gaussian initial data. Again for different types of attention, we show that the Transformer PDE preserves the space of Gaussian measures, which allows us to analyze the Gaussian case theoretically and numerically to identify typical behaviors. This Gaussian analysis captures the evolution of data anisotropy through a deep Transformer. In particular, we highlight a clustering phenomenon that parallels previous results in the non-normalized discrete case.

Key words. Transformers, self-attention, interacting particle systems, gradient flows, Vlasov equations

MSC codes. 35Q68, 68T07, 35B40

1. Introduction. Transformers, introduced in [52], are extremely successful deep learning models, which have reached the state of the art in a wide variety of tasks, from natural language processing to computer vision. A key feature of Transformers is that each data point (image, sentence) is mapped to a sequence of vectors  $(x_1,\ldots,x_n) \in (\mathbb{R}^d)^n$ , called *tokens*, before being processed by the model. Each sequence of tokens is then processed by a succession of layers containing a self-attention block and a multi-layer perceptron (MLP), interleaved with a layer normalization (LayerNorm) operation. The core component of the Transformer architecture is really self-attention: contrary to the MLP and LayerNorm, which are applied token-wise, self-attention makes all tokens interact, which allows the model to learn complex dependencies between them. In this work, we aim to model the evolution of tokens as they go through a deep Transformer, and to identify typical behaviors of the dynamics, to improve our understanding of how Transformers process data. We consider a simplified model with only attention blocks, i.e., without MLPs and layer normalizations; note that adding LayerNorm in the analysis has been studied in [27, 8] and leads to very different dynamics, as tokens are constrained to evolve on a sphere.

Variants of self-attention. The most wide-spread version of self-attention is the one introduced in [52], which we call Softmax self-attention, but several variants have been proposed over the years to improve specific behaviors of self-attention, such as its robustness [35, 47] or its computational complexity [34, 59, 46]. We adapt our

<sup>\*</sup>CNRS and Ecole Normale Supérieure PSL, Paris, France. Emails: valerie.castin@ens.psl.eu (corresponding author), gabriel.peyre@ens.psl.eu.

<sup>&</sup>lt;sup>†</sup>Apple, Paris, France. Email: p\_ablin@apple.com.

 $<sup>^{\</sup>ddagger} \rm Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK. Email: carrillo@maths.ox.ac.uk.$ 

analysis to each of the following variants, that we present in more detail in Section 2: L2 self-attention [35], Sinkhorn self-attention [47], self-attention without Softmax [50], ReLU [59] and Sigmoid [46] self-attention. Finally, several self-attention blocks are typically linearly combined to obtain multi-head attention, and possibly masked in Transformer decoders [52]—we include these two cases in our analysis, building on the framework introduced in [19] for masked self-attention.

Transformers as interacting particle systems. Transformers are deep models: several residual self-attention layers are stacked (typically 10 to 100 [39]) and interleaved with non-linear operations, which results in an involved architecture whose properties are only partially understood. To uncover the mechanisms by which Transformers process data, three recent papers [47, 28, 27] propose to model the behavior of tokens going through the Transformer architecture as an interacting particle system, ruled by a system of ordinary differential equations (ODEs)—thus adapting the theory of neural ODEs [20, 58, 30] to Transformers. This allows for a mathematical study of the dynamics of tokens  $(x_1(t), \ldots, x_n(t))$  as they are processed by a Transformer, where the time variable t corresponds to layer position in the model. Let us focus on the simplest of these models, introduced in [47], where only residual single-head self-attention layers are stacked, and assume that the parameters of the attention layers are continuous functions of time. Denoting  $(x_1(0), \ldots, x_n(0)) \in (\mathbb{R}^d)^n$  the input tokens, the dynamics of tokens as they go through the model reads

(1.1) 
$$\dot{x}_i(t) = \Gamma_{t,X(t)}(x_i(t)) \qquad 1 \le i \le n$$

where  $X(t) := (x_1(t), \ldots, x_n(t))$  and  $\Gamma_{t,X}$  is the attention vector field, defined for  $X \in (\mathbb{R}^d)^n$  and three time-depending matrices  $Q(t), K(t) \in \mathbb{R}^{k \times d}$  and  $V(t) \in \mathbb{R}^{d \times d}$  with  $k \leq d$  as

$$\Gamma_{t,X} \colon x \in \mathbb{R}^d \mapsto \sum_{j=1}^n p_j(t,x)V(t)x_j \in \mathbb{R}^d$$

where  $p_j(t,x) \coloneqq \exp(Q(t)x \cdot K(t)x_j) / \sum_{\ell=1}^n \exp(Q(t)x \cdot K(t)x_\ell)$  for Softmax (i.e., vanilla) self-attention (see Section 2 for the other types of attention),  $\cdot$  denoting the Euclidean scalar product in  $\mathbb{R}^d$ . The authors in [28] provide a thorough study of the dynamics (1.1) in the case of time-independent matrices Q, K, V, showing the emergence of clusters after properly rescaling tokens.

The Transformer PDE. The dynamics (1.1) can be generalized as the following PDE on probability measures [47, 60, 28, 8]:

(1.2) 
$$\partial_t \mu + \operatorname{div}(\mu \Gamma_\mu) = 0$$

where tokens are now represented by a probability measure  $\mu \in \mathcal{P}(\mathbb{R}^d)$  and

$$\Gamma_{\mu} \colon x \in \mathbb{R}^d \mapsto \int Vy \,\kappa_{\mu}(x, y) \,\mathrm{d}\mu(y)$$

with  $Q(t), K(t) \in \mathbb{R}^{k \times d}, V(t) \in \mathbb{R}^{d \times d}$  and  $\kappa_{\mu}(x, y) \coloneqq \exp(Qx \cdot Ky) / \int \exp(Qx \cdot Ky) d\mu(y)$  for Softmax self-attention (see Section 2 for the other types of attention). We do not write explicitly the dependency of  $\kappa_{\mu}$  in t to lighten notations. The case (1.1) of discrete initial data  $(x_1(0), \ldots, x_n(0))$  corresponds to plugging the empirical measure  $\frac{1}{n} \sum_{i=1}^{n} \delta_{x_i(0)}$  in Equation (1.2). This viewpoint allows us to take possibly

continuous probability measures as initial data, thus generalizing the Transformer model to inputs with infinitely many tokens. The well-posedness of the evolution (1.2), which is a Vlasov-type equation [55, 24] or an aggregation-type equation [29, 3], is a non-trivial problem, as the velocity field  $\Gamma_{\mu}$  is typically non-linear in  $\mu$  and associated with a kernel  $\kappa_{\mu}(x, y)$  that is unbounded, since we do not normalize dynamics as opposed to [8], and cannot be rewritten as a function of |x - y|. This rules out more traditional approaches, presented for instance in [24, 9, 11, 42]. In the case of Softmax self-attention, when the initial data is constrained in a compact set and Q, K, V are constant over time (i.e., across layers), Equation (1.2) is the mean-field limit of the dynamics (1.1) [28], in the sense that Equation (1.2) is well-posed for compactly supported initial data, and satisfies a stability estimate of the form

## $W_2(\mu(t),\nu(t)) \le C(t,R_0)W_2(\mu_0,\nu_0)$

where  $R_0$  is the radius of any compact set containing the support of  $\mu_0$  and  $\nu_0$ , and  $W_2$  is the Wasserstein distance between measures [54, 49]. In the first part of this work (Section 3), we extend this result to time-depending parameters and to the variants of self-attention mentioned above. We are also the first to study Equation (1.2) for non-compactly supported initial data, by focusing on the case of Gaussian probability measures (Section 4). The Gaussian case is special, as we show that Equation (1.2) preserves Gaussians for several self-attention variants mentioned above. This property is observed in several algorithms in sampling and optimization [26, 18, 15]. This allows us to summarize the evolution on Gaussians as two matrix ordinary differential equations (ODEs) connecting the expectation and the covariance matrix of the Gaussian data, and to study its well-posedness and behavior theoretically and numerically.

Gradient flow structure of the PDE. To equip a partial differential equation with a gradient flow structure can be a useful tool to prove convergence and properties of the limit. Several works have tackled this question for the Transformer PDE. [47] show that Equation (1.2) is not a Wasserstein gradient flow, whereas the Sinkformer PDE is, allowing them to highlight a diffusive behavior when the parameter  $\varepsilon$  associated to Sinkhorn tends to zero. [27] study an alternative to the Transformer PDE where tokens are of unit norm, thus modelling the effect of LayerNorm, and leverage the gradient flow structure of this projected dynamics to prove the emergence of one single cluster when  $t \to \infty$ . They also introduce a modified metric on  $(\mathbb{S}^{d-1})^n$ , which equips the original dynamics (1.1) with a non-Euclidean gradient flow structure, and is a particular case of the more general framework introduced in [38]. In Section 5. we propose a generalization of this metric from tokens to probability measures, formally showing that it equips the Transformer PDE (1.2) associated with Softmax self-attention with a gradient flow structure. Adapting metrics to convert similar nonlocal PDEs to gradient flows has been a very fruitful approach in different settings [16, 13, 7]. In fact, this has very recently been developed for the normalized dynamics similar to (1.2) associated to (1.1) on the sphere in [8]. In the case of Gaussian initial data (Section 4), the gradient flow structure of the Sinkformer PDE [47] leads to a Bures-Wasserstein gradient flow. The Bures-Wasserstein distance, first introduced in the field of quantum information geometry [6], appears in several recent works in machine learning [36, 22, 37].

Our main contributions can be summarized as:

- i) If  $\Gamma_{\mu}$  is the velocity field of one of the following variants of self-attention, in its single-head *or* multi-head form, and in its masked *or* unmasked form: Softmax self-attention, L2 self-attention, Sinkhorn self-attention, Sigmoid self-attention, then we show that for any compactly supported initial data  $\mu \in \mathcal{P}_c(\mathbb{R}^d)$ , the evolution (1.2) is well-posed. The masked self-attention case leverages the conditional optimal transport framework introduced in [19]. We also derive a stability estimate with respect to the initial data, which proves that Equation (1.2) is the mean-field limit of the dynamical system (1.1). (Section 3)
- ii) For Softmax, L2, Sinkhorn and linear self-attention, we show that initial data that are Gaussians stay Gaussian along the dynamics (1.2). This allows us to derive explicit ODEs on the mean and the covariance matrix of these Gaussian solutions. For Softmax self-attention, we demonstrate under certain assumptions on the parameters that the limiting covariance of the solution is rank-deficient, which mimics the clustering phenomenon observed in [28]. (Sections 4.1, 4.3 and 4.4)
- iii) With a numerical study of the covariance ODEs in the Gaussian case, we point out that rank-deficiency of the limiting covariance generically holds even without the assumptions of our theoretical analysis, for Softmax self-attention as well as L2 self-attention. Moreover, we identify a range of typical behaviors in dimension 2, divided in three categories: (i) convergence, (ii) divergence at  $t \to +\infty$  and (iii) divergence in finite time. We prove that L2 self-attention cannot lead to finite-time divergence, contrary to Softmax self-attention. (Section 4.5)
- iv) Finally, we generalize results in [27] by introducing a twisted metric on the space of probability measures, which equips the Softmax Transformer PDE (1.2) with a gradient flow structure. We prove the non geodesic convexity of the associated functional for the non-normalized dynamics. This complements very recent results in [8] for the normalized dynamics. We also reformulate the covariance ODE for Sinkhorn attention in the Gaussian case as a Bures-Wasserstein gradient flow. (Section 5)

The code for our figures is at github.com/vcastin/2025-transformers-PDEs.

**1.1. Notations.**  $\mathbb{N}^*$  is the set of positive natural numbers. We denote  $\cdot$  the Euclidean scalar product and  $|\cdot|$  the Euclidean norm.  $\|\cdot\|_2$  is the associated operator norm.  $\mathcal{P}(\mathbb{R}^d)$  (resp.  $\mathcal{P}_c(\mathbb{R}^d)$ ) is the set of probability measures (resp. compactly supported probability measures) on  $\mathbb{R}^d$ . For all  $p \geq 1$ , the *p*-Wasserstein space (resp. distance) is denoted  $\mathcal{P}_p(\mathbb{R}^d)$  (resp.  $W_p$ ). The rank of a matrix M is denoted rk M. We denote  $B_R$  or  $\overline{B}(0, R)$  the closed ball of center 0 and of radius R. The support of a probability distribution  $\mu$  is denoted  $\sup \mu$ . The Gaussian distribution of mean  $\alpha \in \mathbb{R}^d$  and of variance  $\Sigma \in \mathbb{R}^{d \times d}$  is denoted  $\mathcal{N}(\alpha, \Sigma)$ , and  $\mathcal{N}(0, \Sigma)$  is denoted  $G_{\Sigma}$ . The set of  $d \times d$  real symmetric matrices is denoted  $\mathcal{S}_d$  and the subset of  $\mathcal{S}_d$  of positive semi-definite (resp. positive definite) matrices is denoted  $\mathcal{S}_d^+$  (resp.  $\mathcal{S}_d^{++}$ ). For any invertible matrix  $A \in \mathbb{R}^{d \times d}$ , we denote  $A^{-\top} := (A^{-1})^{\top}$ . We denote  $\mathcal{C}^{\infty}(\mathbb{R}^d, \mathbb{R})$  the space of infinitely differentiable functions, and  $\mathcal{C}_0^{\infty}(\mathbb{R}^d, \mathbb{R})$  the space of  $\mathcal{C}^{\infty}$  functions that tend to 0 at infinity.

2. Some Variants of Self-Attention. Several variants of self-attention and masked self-attention [52] have been introduced over the years to improve some properties of the attention map, such as its robustness [35, 47] or its computational complexity [34, 59, 46]. Our approach provides a unified framework to compare the

behavior of a Transformer model for each variant: we model any Transformer model (Encoder-only or Decoder-only), which is the composition of several residual attention layers interleaved with projections and multi-layer perceptrons—that we do not include here, by a PDE of the form

(2.1) 
$$\partial_t \mu + \operatorname{div}(\mu \Gamma_\mu) = 0$$

where the velocity field  $\Gamma_{\mu} \colon \mathbb{R}^{\delta} \to \mathbb{R}^{\delta}$  depends on the data  $\mu \in \mathcal{P}(\mathbb{R}^{\delta})$  and on the chosen variant of self-attention. For unmasked self-attention,  $\delta$  is equal to the dimension d of each token. In the case of masked self-attention, we have  $\delta = d + 1$  following [19], as explained in Subsection 2.6. This approach generalizes the case a finite number of tokens, which corresponds to  $\mu$  being an empirical measure  $\sum_{i=1}^{n} \delta_{x_i}$ . Indeed, casting this formula for  $\mu$  in Equation (2.1) leads back to the discrete ODE system (1.1).

Here, we list all the attention variants covered by our framework. Let  $k, d \in \mathbb{N}^*$  with  $k \leq d$  and matrices  $Q, K \in \mathbb{R}^{k \times d}$  and  $V \in \mathbb{R}^{d \times d}$ . We denote  $A \coloneqq K^{\top}Q$ .

**2.1. Softmax Single-Headed Self-Attention.** We call Softmax self-attention the original definition of self-attention in [52]. For any integer  $n \in \mathbb{N}^*$  and any vectors  $x_1, \ldots, x_n \in \mathbb{R}^d$ , Softmax self-attention with parameters (Q, K, V) maps the sequence  $X = (x_1, \ldots, x_n) \in (\mathbb{R}^d)^n$  to

(2.2) 
$$f^{(\mathrm{SM})}(x_1,\ldots,x_n) \coloneqq \left(V \sum_{j=1}^n P_{ij} x_j\right)_{1 \le i \le n} \in (\mathbb{R}^d)^n,$$

with 
$$P_i \coloneqq \operatorname{softmax} \left( (Qx_i \cdot Kx_j)_{1 \le j \le n} \right)$$

where  $\operatorname{softmax}(w) \coloneqq (\exp(w_i) / \sum_{j=1}^d \exp(w_j))_{1 \le i \le n}$ . Note that, as we do not consider training, we absorbed the usual normalizing factor  $1 / \sqrt{k}$  in Q and K. Now, Equation (2.2) can be rewritten as

(2.3) 
$$f(X) = (\Gamma_X^{(SM)}(x_1), \dots, \Gamma_X^{(SM)}(x_n)),$$

with

$$\Gamma_X^{(\mathrm{SM})} \colon x \in \mathbb{R}^d \mapsto \frac{\sum_{i=1}^n e^{Qx \cdot Kx_i} Vx_i}{\sum_{i=1}^n e^{Qx \cdot Kx_i}}$$

Equation (2.3) naturally leads to the PDE viewpoint (2.1) by defining, for any compactly supported probability measure  $\mu \in \mathcal{P}_c(\mathbb{R}^d)$ , the velocity field

$$\Gamma^{(\mathrm{SM})}_{\mu} \colon x \in \mathbb{R}^d \mapsto \frac{\int V y \, e^{Qx \cdot Ky} \, \mathrm{d}\mu(y)}{\int e^{Qx \cdot Ky} \, \mathrm{d}\mu(y)}$$

This generalization of Softmax self-attention to probability measures appears first in [57], and later, in this specific form, in [47]. Note that Softmax self-attention does not depend separately on Q and K, but only on the product  $A := K^{\top}Q \in \mathbb{R}^{d \times d}$ . Remarkably,  $\Gamma_{\mu}^{(\text{SM})}$  is still well-defined when  $\mu$  is Gaussian, and even has a closed form (see Section 4.1).

**2.2. Linear Self-Attention.** Several variants of self-attention, called *linear* variants, reduce the computational complexity of attention from quadratic to linear in the number of tokens n [34, 21, 45, 51, 50]. The simplest version of linear attention replaces the Softmax by the identity in Equation (2.2), and is widely used in the field of in-context learning [25, 56, 40, 48]. It is defined as

$$f^{(\mathrm{lin})}(x_1,\ldots,x_n) \coloneqq (\Gamma_X^{(\mathrm{lin})}(x_1),\ldots,\Gamma_X^{(\mathrm{lin})}(x_n)),$$

with

$$\Gamma_X^{(\mathrm{lin})}(x) \colon x \in \mathbb{R}^d \mapsto V \frac{1}{n} \sum_{j=1}^n x_j x_j^\top A x,$$

which corresponds to the velocity field

$$\Gamma^{(\mathrm{lin})}_{\mu} \colon x \in \mathbb{R}^d \mapsto \int V y(Qx \cdot Ky) \,\mathrm{d}\mu(y).$$

Let us point out that a slight modification of this attention without Softmax [50] can be obtained as a second-order expansion of Softmax self-attention in the regime where  $A := K^{\top}Q$  is very small. Denoting  $f_{\varepsilon}^{(SM)}$  the Softmax attention map with parameters  $(\varepsilon Q, K, V/\varepsilon)$ , we have for any  $X = (x_1, \ldots, x_n) \in (\mathbb{R}^d)^n$  that

$$f_{\varepsilon}^{(\mathrm{SM})}(X)_{i} = \frac{1}{\varepsilon n} \sum_{j=1}^{n} V x_{j} + V \left( \frac{1}{n} \sum_{j=1}^{n} x_{j} x_{j}^{\top} - \frac{1}{n^{2}} \sum_{\ell=1}^{n} x_{\ell} \sum_{\ell=1}^{n} x_{\ell}^{\top} \right) A x_{i} + o_{\varepsilon \to 0}(\varepsilon),$$

with  $A := K^{\top}Q$ . We denote

$$f^{(\mathrm{lin},\varepsilon)}(x_1,\ldots,x_n) \coloneqq (\Gamma_X^{(\mathrm{lin},\varepsilon)}(x_1),\ldots,\Gamma_X^{(\mathrm{lin},\varepsilon)}(x_n))$$

with

$$\Gamma_X^{(\mathrm{lin},\varepsilon)} \colon x \in \mathbb{R}^d \mapsto \frac{1}{n\varepsilon} \sum_{j=1}^n V x_j + V \left( \frac{1}{n} \sum_{j=1}^n x_j x_j^\top - \frac{1}{n^2} \sum_{\ell=1}^n x_\ell \sum_{\ell=1}^n x_\ell^\top \right) A x_\ell$$

the variant of linear attention obtained by expanding the Softmax attention map. The second sum corresponds exactly to attention without Softmax. Note that this attention variant has the following simple expression for any compactly supported or Gaussian probability measure  $\mu$  of expectation  $\alpha$  and covariance  $\Sigma$ :

$$\Gamma^{(\mathrm{lin},\varepsilon)}_{\mu} \colon x \in \mathbb{R}^d \mapsto \frac{1}{\varepsilon} V \alpha + V \Sigma A x.$$

This expression is the same as for Softmax self-attention up to a rescaling of A (see Subsection 4.1). Therefore, all the results we state for Sotmax self-attention in the Gaussian case also hold for this particular choice of linear self-attention.

**2.3.** L2 Self-Attention. For any integer  $n \in \mathbb{N}^*$  and vectors  $x_1, \ldots, x_n \in \mathbb{R}^d$ , L2 self-attention with parameters (Q, K, V) maps the sequence  $X = (x_1, \ldots, x_n) \in (\mathbb{R}^d)^n$  to

$$f^{(L2)}(x_1,\ldots,x_n) \coloneqq \left(V \sum_{j=1}^n P_{ij} x_j\right)_{1 \le i \le n} \in (\mathbb{R}^d)^n,$$

with  $P_{ij} := e^{-|Qx_i - Kx_j|^2} / \sum_{\ell=1}^n e^{-|Qx_i - Kx_\ell|^2}$ . L2 self-attention was introduced, in its discrete form, in [35], for its Lipschitz continuity. Following the same approach as for traditional self-attention, we define the velocity field associated with L2 selfattention, for any compactly supported probability measure  $\mu \in \mathcal{P}_c(\mathbb{R}^d)$ , as

$$\Gamma^{(L2)}_{\mu} \colon x \in \mathbb{R}^d \mapsto \frac{\int Vy \, e^{-|Qx - Ky|^2} \, \mathrm{d}\mu(y)}{\int e^{-|Qx - Ky|^2} \, \mathrm{d}\mu(y)}.$$

Note that  $\Gamma_{\mu}^{(L2)}$  depends on Q, K and not only on their product  $A = K^{\top}Q$ . Like Softmax self-attention, it can be computed in closed form when  $\mu$  is Gaussian (Section 4.3).

**2.4. Sinkhorn Self-Attention.** Sinkhorn self-attention has been introduced in [47]. We only give its definition on probability measures for conciseness. For any compactly supported or Gaussian probability measure  $\mu \in \mathcal{P}_c(\mathbb{R}^d)$ , the velocity field associated to Sinkhorn self-attention with the parameter  $\varepsilon > 0$  and the cost  $c_{\varepsilon}(x, y) \coloneqq \frac{1}{2\varepsilon} |Qx - Ky|^2$  is defined as

$$\Gamma_{\mu,\varepsilon}^{(\mathrm{sink})} \colon x \in \mathbb{R}^d \mapsto \frac{1}{\varepsilon} \int Vy \, \kappa_{\mu,\varepsilon}^{\infty}(x,y) \, \mathrm{d}\mu(y)$$

where  $\kappa_{\mu,\varepsilon}^{\infty}$  is obtained by performing the Sinkhorn-Knopp algorithm on  $\kappa_{\mu,\varepsilon}^{0} \coloneqq e^{-c_{\varepsilon}}$ , i.e.  $\kappa_{\mu,\varepsilon}^{\infty}(x,y)$  is the limit of the following sequence:

(2.4) 
$$\kappa_{\mu,\varepsilon}^{\ell+1}(x,y) = \begin{cases} \frac{\kappa_{\mu,\varepsilon}^{\ell}(x,y)}{\int \kappa_{\mu,\varepsilon}^{\ell}(x,y') \, \mathrm{d}\mu(y')} & \text{if } \ell \text{ is even,} \\ \frac{\kappa_{\mu,\varepsilon}^{\ell}(x,y)}{\int \kappa_{\mu,\varepsilon}^{\ell}(x',y) \, \mathrm{d}\mu(x')} & \text{if } \ell \text{ is odd.} \end{cases}$$

This algorithm—which, in its discrete version, normalizes the kernel matrix  $\kappa$  iteratively row-wise and column-wise—outputs a bi-stochastic kernel:  $\int \kappa_{\mu,\varepsilon}^{\infty}(z,y) d\mu(z) = \int \kappa_{\mu,\varepsilon}^{\infty}(x,z) d\mu(z) = 1$  for all  $x, y \in \mathbb{R}^d$ .

The Sinkhorn algorithm has deep connections with entropic optimal transport (EOT). Indeed, we have the following reinterpretation of Sinkhorn self-attention. Let  $\mu$  and  $\nu$  be two probability measures on  $\mathbb{R}^d$ , either supported in a compact set or Gaussian. Keeping the notation  $c_{\varepsilon}(x, y) := \frac{1}{2\varepsilon} |Qx - Ky|^2$ , consider the entropic optimal transport problem

(2.5) 
$$OT_{\varepsilon}(\mu,\nu) \coloneqq \min_{\pi \in \Pi(\mu,\nu)} \int c_{\varepsilon}(x,y) \, \mathrm{d}\pi(x,y) + \mathrm{KL}(\pi \| \mu \otimes \nu),$$

where  $\Pi(\mu, \nu)$  is the set of couplings between  $\mu$  and  $\nu$  and  $\operatorname{KL}(\alpha \| \beta) \coloneqq \int_{\mathcal{X}} \log\left(\frac{\mathrm{d}\alpha}{\mathrm{d}\beta}\right) \mathrm{d}\alpha$ is the Kullback-Leibler divergence, which is infinite if  $\alpha$  is not absolutely continuous with respect to  $\beta$ . Assuming that  $\mu = \nu$  and denoting  $\mathrm{d}\pi^*(x, y) \coloneqq \kappa_{\mu,\varepsilon}^{\infty}(x, y) \mathrm{d}(\mu \otimes \mu)$ where  $\kappa_{\mu,\varepsilon}^{\infty}$  is obtained as above with Sinkhorn, it is well-known [43, 32] that the coupling  $\pi^*$  is the unique solution of the entropic optimal transport problem (2.5). Therefore,  $\kappa_{\mu,\varepsilon}^{\infty}$  is the density of the optimal coupling  $\pi^*$  with respect to the measure  $\mu \otimes \mu$ . Note that, like for Softmax self-attention,  $\kappa_{\mu,\varepsilon}^{\infty}$  only depends on the product  $A = K^{\top}Q$  (see [47]). We also study the Gaussian case for Sinkhorn self-attention in Section 4.4. **2.5.** Unnormalized Self-Attention. A natural family of self-attention variants consists of suppressing the normalization factor in the definition of  $\Gamma$ , and possibly changing the exponential that appears in the Softmax for a smoother function such as identity [50] (which falls back on linear attention, Section 2.2), ReLU [59] or Sigmoid [46]. In Section 3, we will consider the following variants:

$$\begin{split} &\Gamma^{(\text{exp})}_{\mu} \colon x \in \mathbb{R}^{d} \mapsto \int Vy \ e^{Qx \cdot Ky} \, \mathrm{d}\mu(y), \\ &\Gamma^{(\text{ReLU})}_{\mu} \colon x \in \mathbb{R}^{d} \mapsto \int Vy \ \text{ReLU}(Qx \cdot Ky) \, \mathrm{d}\mu(y), \\ &\Gamma^{(\sigma)}_{\mu} \colon x \in \mathbb{R}^{d} \mapsto \int Vy \ \sigma(Qx \cdot Ky) \, \mathrm{d}\mu(y), \end{split}$$

where ReLU:  $x \in \mathbb{R} \mapsto \max(x, 0)$  and  $\sigma: z \in \mathbb{R} \mapsto (1+e^{-z})^{-1}$  is the Sigmoid function. It is easy to see that none of these attention variants preserve the Gaussianity of the data, as  $\Gamma_{\mu}$  is not an affine function when  $\mu$  is Gaussian.

**2.6.** Masked Self-Attention. Softmax self-attention, introduced at the beginning of Section 2.1, is used in Encoders, which appear in Encoder-only [23] and Encoder-Decoder Transformers [52]. In Decoders, however, what is used is masked self-attention, which takes into account the sequential nature of the inputs. In its discrete version, masked self-attention maps the sequence  $X = (x_1, \ldots, x_n) \in (\mathbb{R}^d)^n$ to  $(f^{(m)}(X)_1, \ldots, f^{(m)}(X)_n) \in (\mathbb{R}^d)^n$  such that

(2.6) 
$$f^{(\mathrm{m})}(X)_i \coloneqq f^{(\mathrm{SM})}(x_1, \dots, x_i),$$

where  $f^{(SM)}$  is Softmax self-attention, defined in Equation (2.2). Similarly, one can define a masked attention map for any discrete self-attention variant. Masked selfattention is crucial for next-token prediction tasks in NLP and time series: at test time, it is used to generate data in an auto-regressive fashion, i.e., each newly generated token contributes to the computation of the next one; and during training, it prevents the model to cheat in its next-token prediction task by making use of tokens it has to predict.

Equation (2.6) cannot be directly generalized to probability measures, as masked attention makes use of the order of tokens, which is lost when representing a sequence  $(x_1, \ldots, x_n)$  by the associated empirical measure  $\frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ . To include masked selfattention in our unified framework, we leverage a contribution in [19], which extends masked self-attention to probability measures by adding a *position coordinate* in [0, 1] to the input space: this coordinate specifies the order of tokens. The ambient space of the Transformer PDE (2.1) then becomes the space of compactly supported probability measures  $\bar{\mu}$  on the product space  $[0, 1] \times \mathbb{R}^d$ . Let  $d \in \mathbb{N}^*$  and  $Q, K \in \mathbb{R}^{k \times d}$ and  $V \in \mathbb{R}^{d \times d}$ . The velocity field associated with masked Softmax self-attention is defined, for any compactly supported probability measure  $\bar{\mu} \in \mathcal{P}_c([0, 1] \times \mathbb{R}^d)$ , as

(2.7) 
$$\Gamma_{\bar{\mu}}^{(\mathrm{m})}(\sigma, x) \coloneqq \left(0, \frac{\int_{[0,\sigma] \times \mathbb{R}^d} V y \, e^{Ax \cdot y} \, \mathrm{d}\bar{\mu}(\tau, y)}{\int_{[0,\sigma] \times \mathbb{R}^d} e^{Ax \cdot y} \, \mathrm{d}\bar{\mu}(\tau, y)}\right)$$

with  $A \coloneqq K^{\top}Q$ . In what follows, we call *time marginal* the first marginal of  $\bar{\mu}$ , defined as

(2.8) 
$$d\theta(s) \coloneqq \int_{\mathbb{R}^d} d\bar{\mu}(s, x).$$

 $\theta$  is therefore a probability measure on [0, 1]. We call space marginal and denote  $\mu$  the marginal of  $\bar{\mu}$  in  $\mathbb{R}^d$ , defined as

$$\mathrm{d}\mu(x) \coloneqq \int_{[0,1]} \,\mathrm{d}\bar{\mu}(s,x)$$

 $\mu$  is therefore a probability measure on  $\mathbb{R}^d$ . For a finite number of tokens, Equation (2.6) can be cast in the framework (2.7) by choosing  $\bar{\mu} = \frac{1}{n} \sum_{i=1}^{n} \delta_{(i/n,x_i)}$ , for instance. Equation (2.7) then provides a natural generalization of masked Softmax self-attention to continuous distributions of tokens. Moreover, this idea, originally applied to Softmax self-attention, can be used for any of the attention variants mentioned above.

Note that with this choice of velocity field, the time marginal of  $\bar{\mu}(t)$  stays constant along the dynamics (2.1). However, Equation (2.7) does not allow for a study of the Gaussian case—e.g., when the space marginal is Gaussian—as the masked Transformer PDE does not preserve this structure.

2.7. Multi-Head Self-Attention. The attention functions presented above provide different ways of learning dependencies—for example, semantic dependencies—between tokens. In practice, to increase the expressive power of attention and learn dependencies at different scales in the text or image, several *attention heads* with different parameters are linearly combined to obtain what is called multi-head attention. If  $\Gamma$  is the velocity field of a (possibly masked) attention function—Softmax, linear, L2, Sigmoid..., the associated multi-head velocity field takes the form

$$\Gamma^{(\mathrm{MH})}_{\mu} \coloneqq \sum_{h=1}^{H} \Gamma^{(h)}_{\mu}$$

for H the number of heads, which has to be a divisor of d, and where the parameters of  $\Gamma_{\mu}^{(h)}$  are  $Q^{(h)}, K^{(h)} \in \mathbb{R}^{d/H,d}$  and  $V^{(h)} \in \mathbb{R}^{d \times d}$ . Note that multi-head attention usually involves matrices  $W^{(h)}$  that multiply each term  $\Gamma_{\mu}^{(h)}$  for  $1 \leq h \leq H$ . As we do not consider training, we absorb  $W^{(h)}$  in  $V^{(h)}$ , so that our matrices  $V^{(h)}$  are  $d \times d$ . Moreover, it is practical to represent the list of matrices  $(Q^{(h)}, K^{(h)})_{1 \leq h \leq H} \in (\mathbb{R}^{d/H \times d})^H$  by two square matrices  $Q, K \in \mathbb{R}^{d \times d}$ , where Q (resp. K) is obtained by stacking the  $Q^{(h)}$  (resp.  $K^{(h)}$ ) row-wise:

(2.9) 
$$Q \coloneqq \begin{pmatrix} Q^{(1)} \\ \vdots \\ Q^{(H)} \end{pmatrix} \quad \text{and} \quad K \coloneqq \begin{pmatrix} K^{(1)} \\ \vdots \\ K^{(H)} \end{pmatrix}.$$

In the rest of the paper, we will mostly state results for Softmax multi-head attention, simply denoted  $\Gamma_{\mu}^{(MH)}$ , but they immediately extend to the other types of attention we consider. Moreover, we have a closed form for  $\Gamma_{\mu}^{(MH)}$  when  $\mu$  is Gaussian, which directly comes from the single-head formula (see Section 4).

**3. Well-Posedness for a Compactly Supported Initial Condition.** We first investigate the behavior of the Transformer PDE

(3.1) 
$$\partial_t \mu + \operatorname{div}(\mu \Gamma_\mu) = 0$$

when the initial condition is compactly supported, for all variants of self-attention presented in Section 2. For any  $p \ge 1$ , we will be looking for solutions of Equation

(3.1), as curves  $\mu \in \mathcal{C}([0,T], \mathcal{P}_p(\mathbb{R}^d))$ , continuous with the topology induced by the Wasserstein distance  $W_p$  on  $\mathcal{P}_p(\mathbb{R}^d)$ —which makes it a complete space. We say that  $\mu \in \mathcal{C}([0,T], \mathcal{P}_p(\mathbb{R}^d))$  is a (weak) solution of Equation (3.1) with initial data  $\mu_0$  if for any  $\mathcal{C}^{\infty}$  function  $\psi \in \mathcal{C}_0^{\infty}([0,\infty) \times \mathbb{R}^d)$  that tends to zero at infinity we have

$$\int_0^T \int_{\mathbb{R}^d} \left( \frac{\partial \psi}{\partial t} + \Gamma_\mu \cdot \nabla_x \psi \right) \, \mathrm{d}\mu(t) = \int_{\mathbb{R}^d} \psi(T, \cdot) \, \mathrm{d}\mu(T) - \int_{\mathbb{R}^d} \psi(0, \cdot) \, \mathrm{d}\mu_0$$

We focus on proving the well-posedness of the PDE (3.1), i.e., existence and uniqueness of a global solution, and deriving a stability estimate. The case of Softmax selfattention has already been studied in the particular case of constant parameters, while the other cases are completely new. Our proof, which relies on a fixed-point argument, is different and technically simpler than the one in [28] since it relies on classical Dobrushin type estimates [24, 9].

**3.1. Unmasked Self-Attention.** Let us first study Equation (3.1) for our attention variants in their unmasked form. We have the following result.

THEOREM 3.1. Let  $d, k \in \mathbb{N}^*$  with  $k \leq d$  and  $p \geq 1$ . Let  $Q, K: [0, +\infty) \to \mathbb{R}^{k \times d}$ and  $V: [0, +\infty) \to \mathbb{R}^{d \times d}$  be three continuous maps, modeling the evolution of parameters Q, K, V across layers of the Transformer. We set  $\varepsilon = 1$  for Sinkformer attention, by absorbing the  $\varepsilon$  in Q, K, V. Let  $\mu_0 \in \mathcal{P}_c(\mathbb{R}^d)$  be a compactly supported initial condition. Then, for any choice of velocity field in  $\{\Gamma^{(SM)}, \Gamma^{(L2)}, \Gamma^{(sink)}, \Gamma^{(\sigma)}, \Gamma^{(MH)}\}$ , and denoting  $\Gamma_{\mu}(t, \cdot) \coloneqq \Gamma_{\mu(t)}(\cdot)$  the attention map associated to Q(t), K(t), V(t) (with the convention (2.9) and k = d for multi-head attention), the Transformer PDE

(3.2) 
$$\partial_t \mu + \operatorname{div}(\mu \Gamma_\mu) = 0$$

with initial condition  $\mu_0$  has a unique global weak solution  $\mu \in \mathcal{C}([0, +\infty), \mathcal{P}_c(\mathbb{R}^d))$ , with  $\mathcal{P}_c(\mathbb{R}^d)$  equipped with the p-Wasserstein distance  $W_p$ . Moreover, let  $R_0$  be the smallest radius such that  $\operatorname{Supp} \mu_0 \subset B_{R_0}$ , and define

$$R(t) \coloneqq \exp\left(\int_0^t \|V(s)\|_2 \, \mathrm{d}s\right) R_0$$

for  $t \geq 0$ . Then, the solution  $\mu$  satisfies

$$\operatorname{Supp} \mu(t) \subset B_{R(t)}$$

for all  $t \ge 0$ . Finally, we have the following stability estimate. For all  $R_0 > 0$  and t > 0, there exists a constant  $C(t, R_0)$  depending only on t and  $R_0$  (and on the choice of  $\Gamma_{\mu}$ ), such that for any initial conditions  $\mu_0$  and  $\nu_0$  supported in  $B_{R_0}$ , and denoting  $\mu$  and  $\nu$  the associated global solutions of Equation (3.2), we have

$$W_p(\mu(t), \nu(t)) \le C(t, R_0) W_p(\mu_0, \nu_0).$$

Therefore, for compactly supported initial data, the Transformer PDE is wellposed: it has a unique global solution on [0,T] with initial data  $\mu_0$ , whose support typically grows exponentially with time. Moreover, it generalizes the discrete system (1.1), which corresponds to the initial data  $\mu_0 := \frac{1}{n} \sum_{i=1}^n \delta_{x_i(0)}$ , and is the meanfield limit of this interacting particle system on any compact time interval [0,T]: if  $(\mu^{\eta}(0))_{\eta}$  is a sequence of empirical measures such that  $W_p(\mu^{\eta}(0), \mu(0)) \to_{\eta \to \infty} 0$ , then the distance  $W_p(\mu^{\eta}(t), \mu(t))$  tends to 0 as well when  $\eta \to \infty$  for any  $0 \le t \le T$ ; and this holds for Softmax, L2, Sinkhorn and Sigmoid self-attention and their multihead version. However, the constant  $C(t, R_0)$  is exponential in R(t), which is itself exponential in t. When t becomes large, the stability estimate is therefore very loose.

*Proof.* Let us detail the proof of Theorem 3.1 in the simpler case where the parameters Q, K, V are constant over time. The general case is deferred to Appendix B.4.

Step 1: defining the flow  $\phi_t(\mu)(x)$ . For all T > 0, denote  $\mathcal{C}_{co}([0,T], \mathcal{P}_c(\mathbb{R}^d))$  the set of equi-compactly supported curves, i.e., of continuous curves  $\mu \in \mathcal{C}([0,T], \mathcal{P}_c(\mathbb{R}^d))$ such that for any compact time interval  $[s,t] \subset [0,T]$ , there exists a compact set  $K \subset \mathbb{R}^d$  verifying:  $\forall \tau \in [s,t]$ ,  $\operatorname{Supp}(\mu(\tau)) \subset K$ . Let  $\mu \in \mathcal{C}_{co}([0,T], \mathcal{P}_c(\mathbb{R}^d))$  and  $x \in \mathbb{R}^d$ , and consider the Cauchy problem

(3.3) 
$$\begin{cases} \dot{r}(t) = \Gamma_{\mu}(t, r(t)) \text{ in } 0 \le t \le T \\ r(0) = x. \end{cases}$$

We prove the following estimates for each self-attention type (see Appendix B.1). If  $\operatorname{Supp} \mu \subset B_R$ , then

- (i)  $\sup_{x \in \mathbb{R}^d} |\Gamma_\mu(x)| \le ||V||_2 R$ ,
- (ii)  $\sup_{x \in \mathbb{R}^d} \|D_x \Gamma_\mu\|_2 \le \|V\|_2 \|A\|_2 R^2$ ,
- (iii)  $|\Gamma_{\mu}(x) \Gamma_{\nu}(x)| \le c(x, R)W_p(\mu, \nu),$

where c(x, R) is a continuous function that depends on x, R and Q, K, V. Then, traditional Cauchy theory tells us that problem (3.3) has a unique global solution. Indeed, denote R > 0 a radius such that  $\operatorname{Supp} \mu(t) \subset B_R$  for all  $0 \leq t \leq T$ . Such a radius exists because  $\mu$  is equi-compactly supported. Equations (ii) and (iii) ensure that the map  $(t, x) \in [0, T] \times \mathbb{R}^d \mapsto \Gamma_{\mu(t)}(x)$  is continuous. Thanks to Equation (ii), we also have, for all  $t \in [0, T]$ :

$$\left|\Gamma_{\mu(t)}(x) - \Gamma_{\mu(t)}(y)\right| \le \|V\|_2 \|A\|_2 R^2 |x - y|$$

Finally, Equation (i) ensures that  $\Gamma_{\mu(t)}$  grows less than linearly with |x|—in fact, it is bounded with respect to |x|—so that there is a unique global solution. We can therefore define the associated flow  $\phi_t(\mu)(x)$ , such that the solution r of Problem (3.3) satisfies  $r(t) = \phi_t(\mu)(x)$ .

Step 2: fixed point argument on [0,T]. Let us set any initial condition  $\bar{\mu}_0 \in \mathcal{P}_c(\mathbb{R}^d)$ , and a time T > 0 to be chosen later. Define the map

$$\mathcal{F} \colon \mathcal{C}_{\rm co}([0,T],\mathcal{P}_c(\mathbb{R}^d)) \to \mathcal{C}_{\rm co}([0,T],\mathcal{P}_c(\mathbb{R}^d))$$
$$\mu \mapsto \phi_t(\mu)_{\sharp} \bar{\mu}_0$$

and equip the space  $\mathcal{C}_{co}([0,T],\mathcal{P}_c(\mathbb{R}^d))$  with the distance

$$\mathcal{D}_{p,T}(\mu,\nu) \coloneqq \max_{0 \le t \le T} W_p(\mu(t),\nu(t))$$

We look for a complete space X preserved by  $\mathcal{F}$  and such that  $\mathcal{F}$  is a contraction on X, to apply a fixed point argument and show the existence and uniqueness of a solution to Equation (3.2) in [0,T]. Let X be the set of curves  $\mu \in \mathcal{C}_{co}([0,T],\mathcal{P}_c(\mathbb{R}^d))$ satisfying

$$\operatorname{Supp} \mu(t) \subset \overline{B}(0, e^{\|V\|_2 t} R_0)$$

for all  $0 \leq t \leq T$ , where  $R_0$  is the smallest radius such that  $\operatorname{Supp} \overline{\mu}_0 \subset B_{R_0}$ . We show (see Appendix B.2) that  $\mathcal{F}(X) \subset X$ , that  $(X, \mathcal{D}_{p,T})$  is complete and that

$$\mathcal{D}_{p,T}(\mathcal{F}(\mu),\mathcal{F}(\nu)) \le f(T,R_0)\mathcal{D}_{p,T}(\mu,\nu)$$

for any  $\mu, \nu \in X$ , where  $f(T, R_0)$  is a continuous function such that  $f(T, R_0) \to_{T \to 0} 0$ . Let us then choose T > 0 small enough to have  $f(T, R_0) < 1$ , so that  $\mathcal{F}$  becomes a strict contraction. According to Banach fixed-point theorem,  $\mathcal{F}$  has a unique fixed point  $\mu$ . Equivalently, Equation (3.2) with initial data  $\bar{\mu}_0$  has a unique solution in [0, T].

Step 3: constructing a global solution. By repeating Step 2 with the updated initial condition  $\phi_T(\mu)_{\sharp}\overline{\mu}_0$ , we can extend  $\mu$  to further times, and so on. However, the time interval that we add at each step depends on  $R_0$ , which grows as we iterate the argument—and the length of the time interval shrinks accordingly. Assume by contradiction that this method does not allow us to extend  $\mu$  beyond some finite limiting time  $T_{\text{lim}}$ . It is easy to check that  $\mu$  satisfies, for all  $t \in [0, T_{\text{lim}})$ :

$$\operatorname{Supp} \mu(t) \subset \overline{B}(0, e^{\|V\|_2 t} R_0).$$

where  $R_0$  is the smallest radius such that  $\operatorname{Supp} \overline{\mu}_0 \subset B_{R_0}$ . Denote  $R(t) := e^{\|V\|_2 t} R_0$ for all  $0 \le t \le T_{\lim}$ , and choose T' > 0 small enough so that

$$f(T', R(T_{\lim})) < 1.$$

By using the same arguments as above, we can extend the restriction  $\mu_{|[0,T_{\text{lim}}-T'/2]}$  to the time interval  $[0, T_{\text{lim}} + T'/2]$ , which is a contradiction. Therefore,  $\mu$  can be extended to arbitrarily large times by iterating the fixed point argument, and so there exists a unique global solution to Problem (3.2) with initial solution  $\bar{\mu}_0$ .

Step 4: stability estimates. We derive the stability estimates in Appendix B.3 adapting the traditional Dobrushin's method [24, 29, 11], building on Equations (i), (ii), (iii).

Theorem 3.1 does not cover the unnormalized attentions  $\Gamma_{\mu}^{(\text{exp})}$ ,  $\Gamma_{\mu}^{(\text{lin})}$  and  $\Gamma_{\mu}^{(\text{ReLU})}$ . Indeed, for these three types of attention and assuming that  $\mu$  is supported in  $B_R$ , the velocity field  $\Gamma$  cannot be bounded by R up to a constant factor. We rather have that

$$\sup_{x \in B_R} \left| \Gamma_{\mu}^{(\exp)}(x) \right| \le \|V\|_2 R e^{\|A\|_2 R^2}$$
$$\sup_{x \in B_R} \left| \Gamma_{\mu}^{(\ln)}(x) \right| \le \|V\|_2 \|A\|_2 R^3$$
$$\sup_{x \in B_R} \left| \Gamma_{\mu}^{(\text{ReLU})}(x) \right| \le \|V\|_2 \|A\|_2 R^3,$$

and when  $V = I_d$ , each bound is reached for a well-chosen Dirac measure. Therefore, Lemma B.8 does not hold, and we have no guarantee that  $\mu$  stays compactly supported across the Transformer PDE dynamics, which makes the analysis more difficult—this is left for future work.

**3.2. Masked Self-Attention.** The case of masked self-attention is special: indeed, finding an estimate of the form

$$\left|\Gamma_{\bar{\mu}}^{(\mathrm{m})}(s,x) - \Gamma_{\bar{\nu}}^{(\mathrm{m})}(s,x)\right| \le C(x)W_2(\bar{\mu},\bar{\nu})$$

with the framework introduced in Section 2.6, appears to be a hard problem even when the supports of  $\mu$  and  $\nu$  are constrained in a compact set and have the same time marginal (defined in Equation (2.8)). Note that when the supports of  $\mu$  and  $\nu$  are not restricted to a compact set, we prove that no such estimate can hold (see Appendix B.5). A key remark to circumvent this issue is to notice that if  $\bar{\mu}(t)$  is the solution of Equation (2.1) initialized at  $\bar{\mu}_0 \in \mathcal{P}_c([0,1] \times \mathbb{R}^d)$ , then the time marginal of  $\bar{\mu}(t)$  is constant over time. For any probability distribution  $\theta \in \mathcal{P}([0,1])$ , let us denote  $\mathcal{P}_c^{\theta}([0,1] \times \mathbb{R}^d)$  the set of compactly supported probability measures whose time marginal is equal to  $\theta$ . The idea is to equip this space with an alternative to the Wasserstein distance, the conditional Wasserstein distance.

DEFINITION 3.2 (Conditional Wasserstein distance [31]). Let  $d \in \mathbb{N}^*$  and  $\bar{\mu}, \bar{\nu} \in \mathcal{P}^{\theta}_{c}([0,1] \times \mathbb{R}^d)$ . The conditional Wasserstein distance between  $\bar{\mu}$  and  $\bar{\nu}$  is defined as

$$d(\bar{\mu},\bar{\nu}) \coloneqq \left(\int_0^1 W_2(\mu^\tau,\nu^\tau)^2 \,\mathrm{d}\theta(\tau)\right)^{1/2}$$

where we have written

$$\mathrm{d}\bar{\mu}(\tau, x) =: \mathrm{d}\theta(\tau) \,\mathrm{d}\mu^{\tau}(x)$$

with the disintegration theorem.

Remark 3.3. One can also define the conditional p-Wasserstein distance for  $p \ge 1$  as

$$d(\bar{\mu}, \bar{\nu}) \coloneqq \left( \int_0^1 W_p(\mu^{\tau}, \nu^{\tau})^p \,\mathrm{d}\theta(\tau) \right)^{1/p}$$

with the same notations as in Definition 3.2. All our proofs adapt directly to this case without changing the constant factors, as the estimates rely on a  $W_1$  estimate.

The idea of Definition 3.2 is to constrain the transport plans so that they preserve the time marginal. This allows us to prove the following well-posedness result.

THEOREM 3.4. Let  $k, d \in \mathbb{N}^*$ . Let  $Q, K: [0, +\infty) \to \mathbb{R}^{k \times d}$  and  $V: [0, +\infty) \to \mathbb{R}^{d \times d}$  be three continuous maps, modeling the evolution of parameters Q, K, V across layers of the Transformer. Let  $\overline{\mu}_0 \in \mathcal{P}_c([0,1] \times \mathbb{R}^d)$  be a compactly supported initial condition. Let  $\Gamma \in \{\Gamma^{(SM)}, \Gamma^{(L2)}, \Gamma^{(sink)}, \Gamma^{(\sigma)}, \Gamma^{(MH)}\}$  be an unmasked velocity field associated with the parameters Q(t), K(t), V(t) (with the convention (2.9) and k = d for multi-head attention), and denote  $\Gamma_{\overline{\mu}}(t) \coloneqq \Gamma^{(m)}_{\overline{\mu}(t)}$  the corresponding masked attention map. Then, the masked Transformer PDE

(3.4) 
$$\partial_t \bar{\mu} + \operatorname{div}(\bar{\mu}\Gamma_{\bar{\mu}}) = 0$$

with initial condition  $\bar{\mu}_0$  has a unique global weak solution  $\bar{\mu} \in \mathcal{C}([0, +\infty), \mathcal{P}_c([0, 1] \times \mathbb{R}^d))$ , with  $\mathcal{P}_c([0, 1] \times \mathbb{R}^d)$  equipped with the conditional Wasserstein distance d. Moreover, let  $R_0$  be the smallest radius such that  $\operatorname{Supp} \mu_0 \subset B_{R_0}$ , where  $\mu_0$  is the space marginal of  $\bar{\mu}_0$ , and define

$$R(t) \coloneqq e^{\int_0^t \|V(s)\|_2 \,\mathrm{d}s} R_0$$

for  $t \geq 0$ . Then, the space marginal  $\mu$  of the solution  $\bar{\mu}$  satisfies

$$\operatorname{Supp} \mu(t) \subset B_{R(t)}$$

for all  $t \ge 0$ . Finally, we have the following stability estimate. For all  $R_0 > 0$  and t > 0, there exists a constant  $C(t, R_0)$  depending only on t and  $R_0$  such that for any initial conditions  $\bar{\mu}_0$  and  $\bar{\nu}_0$  supported in  $[0, 1] \times B_{R_0}$ , and denoting  $\bar{\mu}$  and  $\bar{\nu}$  the associated global solutions of Equation (3.4), we have

$$d(\bar{\mu}(t), \bar{\nu}(t)) \leq C(t, R_0) d(\bar{\mu}_0, \bar{\nu}_0).$$

*Proof.* It suffices to do the proof for single-head masked attention, as we have the same type of estimates on  $\Gamma$  for single-head and multi-head masked attention. The proof follows the same steps as for Theorem 3.1, using that  $\mathcal{P}_p^{\theta}([0,1] \times \mathbb{R}^d)$  equipped with the distance d is a complete space [2] and leveraging the following estimates, that derive from the same computations as for unmasked self-attention:

(i) 
$$\sup_{x \in \mathbb{R}^d} \left| \Gamma_{\bar{\mu}}^{(\text{mask})}(x) \right| \le \|V\|_2 R,$$

(ii) 
$$\sup_{x \in \mathbb{R}^d} \left\| D_x \Gamma_{\bar{\mu}}^{(\text{mask})} \right\|_2 \le \|V\|_2 \|A\|_2 R^2$$
,

(iii)  $\left| \Gamma_{\bar{\mu}}^{(\text{mask})}(x) - \Gamma_{\bar{\nu}}^{(\text{mask})}(x) \right| \le c(x, R) d(\bar{\mu}, \bar{\nu}),$ 

where c(x, R) is a continuous function that depends on x, R and Q, K, V.

4. Clustering For Gaussian Initial Data. Let us now tackle the particular case where the initial data is a Gaussian measure. This is an over-simplistic modelization of real data; still, it provides a simple example of anisotropic data, which allows us to study the evolution of the anisotropy across the dynamics. Indeed, the Gaussian case has the nice property that several types of unmasked self-attention introduced in Section 2 have a closed form when  $\mu$  is Gaussian—more precisely,  $\Gamma_{\mu}$  becomes an affine function. This implies that Gaussian input measures stay Gaussian during the dynamics  $\partial_t \mu + \operatorname{div}(\mu \Gamma_{\mu}) = 0$ , so that we can summarize the evolution as a system of ordinary differential equations linking the expectation and the covariance matrix of  $\mu$ , which can then be studied more easily than the initial PDE.

**4.1. Softmax Self-Attention.** Let us start with Softmax self-attention. We have the following closed form for the velocity field  $\Gamma_{\mu}^{(\text{SM})}$  when  $\mu$  is a Gaussian probability measure. Recall that  $A := K^{\top}Q$ .

LEMMA 4.1. Let  $\mu = \mathcal{N}(\alpha, \Sigma)$  be a Gaussian measure on  $\mathbb{R}^d$ . Then for all  $x \in \mathbb{R}^d$  it holds

$$\Gamma^{(\mathrm{SM})}_{\mu}(x) = V(\alpha + \Sigma A x)$$

Note that this covers the case of linear attention as well by absorbing the  $\varepsilon$  in the matrix A (see Subsection 2.2). When the initial data is Gaussian, we can then rewrite the dynamics as follows.

PROPOSITION 4.2. Let  $k, d \in \mathbb{N}^*$ . Let  $Q, K : [0, +\infty) \to \mathbb{R}^{k \times d}$  and  $V : [0, +\infty) \to \mathbb{R}^{d \times d}$  be three continuous maps, modeling the evolution of parameters Q, K, V across layers of the Transformer. Consider a Gaussian initial condition  $\mu_0 = \mathcal{N}(\alpha_0, \Sigma_0)$  with  $\alpha_0 \in \mathbb{R}^d$  and  $\Sigma_0 \in \mathbb{R}^{d \times d}$  positive definite. Then, the Transformer PDE

$$\partial_t \mu + \operatorname{div}(\mu \Gamma^{(\mathrm{SM})}_{\mu}) = 0$$

associated to Softmax self-attention with parameters Q(t), K(t), V(t) has a unique maximal solution  $\mu$  on  $[0, t_{\max})$ , such that  $\mu(t)$  is Gaussian for all  $t \in [0, t_{\max})$ . Moreover, denoting  $\mu(t) =: \mathcal{N}(\alpha(t), \Sigma(t))$  and  $A := K^{\top}Q$ , we have

(4.1) 
$$\dot{\Sigma} = V \Sigma A \Sigma + \Sigma A^{\top} \Sigma V^{\top}.$$

and

$$\dot{\alpha} = V(I_d + \Sigma A)\alpha.$$

The proof is in Appendix C.1. In the rest of the section, let us assume to simplify the problem that Q, K, V are constant over time. The following paragraphs analyze Equation (4.1), i.e., the evolution of the covariance matrix of  $\mu$ , highlighting a clustering phenomenon and characterizing the stationary points of the dynamics under two different assumptions—providing a general characterization remains an open problem.

Limiting points have low rank when the parameters commute. Let us first look at Equation (4.1) in dimension 1, writing (s, a, v) for  $(\Sigma, A, V)$  in this case.

PROPOSITION 4.3. Let  $a, v \in \mathbb{R}$  and  $s_0 \in \mathbb{R}^*_+$ . Consider the Equation (4.1) in dimension 1, which reads

$$\dot{s} = 2avs^2$$

with initial condition  $s(0) = s_0$ . There is a unique maximal solution to this equation, defined as

$$s(t) = (s_0^{-1} - 2vat)^{-1}$$

for  $t \in [0, t_{\max})$ , where  $t_{\max} := (2vas_0)^{-1}$  if av > 0, and  $t_{\max} := +\infty$  if  $av \le 0$ .

In dimension 1, the dynamics on Gaussians are therefore simple: the solution either i) goes to zero or ii) blows up, i.e., becomes infinite, in finite time. Translating these behaviors in terms of the mass repartition induced by the Gaussian measure gives either i) clustering of the mass towards a single Dirac when  $t \to +\infty$  or ii) escaping of the mass to infinity in finite time. We can extend this analysis to higher dimensions under a commutation assumption.

**PROPOSITION 4.4.** Consider the matrix-valued differential equation

(4.2) 
$$\dot{\Sigma} = V \Sigma A \Sigma + \Sigma A^{\top} \Sigma V^{\top}.$$

Assume that the matrix V commutes with  $\Sigma_0$  and  $VA + A^{\top}V^{\top}$ . Then, Equation (4.2) with initial condition  $\Sigma_0 \succ 0$  has a unique maximal solution, defined as

$$\Sigma(t) = \left(\Sigma_0^{-1} - t(VA + A^{\top}V^{\top})\right)^{-1}$$

for all  $t \ge 0$  such that the matrix  $\Sigma_0^{-1} - t(VA + A^{\top}V^{\top})$  is invertible. Moreover, the behavior of the solution therefore depends on the sign of the eigenvalues of  $VA + A^{\top}V^{\top}$  as follows.

- 1. If  $VA + A^{\top}V^{\top} \leq 0$ , then Equation (4.2) has a global solution. Moreover, the matrix  $\Sigma(t)$  converges to a limit  $\Sigma^* \succeq 0$  satisfying  $\lambda_i(\Sigma^*) = 0$  for all i such that  $\lambda_i(VA + A^{\top}V^{\top}) < 0$ . Therefore, the mass concentrates on an affine subspace of  $\mathbb{R}^d$ , of dimension equal to the multiplicity of 0 as an eigenvalue of  $VA + A^{\top}V^{\top}$ .
- 2. If  $VA + A^{\top}V^{\top}$  has a positive eigenvalue, then the largest eigenvalue of  $\Sigma(t)$  goes to  $+\infty$  in finite time, so the mass escapes to infinity in finite time.

Remark 4.5. Note that in the even simpler case where  $\Sigma_0$  and  $VA + A^{\top}V^{\top}$  commute, the limit of  $\Sigma(t)$  becomes explicit. Indeed, as these matrices are codiagonalizable we get

$$\lambda_i(\Sigma(t)) = \left(\lambda_i(\Sigma_0)^{-1} - t\lambda_i(VA + A^\top V^\top)\right)^{-1}$$

In this case, if  $VA + A^{\top}V^{\top} \leq 0$  then  $\Sigma(t)$  converges to the matrix  $\Sigma^* \geq 0$  defined as

$$\lambda_i(\Sigma^*) = \begin{cases} 0 & \text{if } \lambda_i(VA + A^\top V^\top) < 0, \\ \lambda_i(\Sigma_0) & \text{if } \lambda_i(VA + A^\top V^\top) = 0. \end{cases}$$

Hence, under the assumption of Proposition 4.4 and for  $VA + A^{\top}V^{\top} \neq 0$ , any Gaussian solution of the Transformer PDE either i) converges to a degenerate Gaussian measure, i.e., with a low-rank covariance matrix, when  $t \to +\infty$ , or ii) converges in finite time to a probability distribution of zero mass. Case i) corresponds to a collapse of the mass along the axes associated to zero eigenvalues of the limiting covariance  $\Sigma^*$ , and can be seen as the continuous parallel of the clustering for discrete tokens highlighted in [52, 28]. The proof of Proposition 4.4 is in Appendix C.1. We complement this analysis with numerical experiments (Section 4.5), which show that this behavior goes beyond the assumption of Proposition 4.4.

*Rank 1 is preserved.* The case where the initial covariance matrix is of rank 1, which corresponds to a maximally anisotropic distribution of tokens, deserves being analyzed separately, as we point out that rank 1 is preserved along the dynamics.

LEMMA 4.6. Let  $A, V \in \mathbb{R}^{d \times d}$  and  $u_0 \in \mathbb{R}^d$ . Then the maximal solution  $\Sigma(t)$  of the equation

$$\dot{\Sigma} = V\Sigma A\Sigma + \Sigma A^{\top}\Sigma V^{\top}$$

with initial data  $\Sigma_0 := u_0 u_0^{\top}$  is of rank 1 for all times where it is defined. Moreover, denoting  $\Sigma(t) := u(t)u(t)^{\top}$  with  $u(t) \in \mathbb{R}^d$ , we have

$$\dot{u} = (u^{\top} A u) V u.$$

Finding a closed form for u(t) seems a difficult problem. Still, from Lemma 4.6 we can easily characterize rank-1 stationary points of the dynamics.

LEMMA 4.7. Let  $A, V \in \mathbb{R}^{d \times d}$ . The rank-1 matrix  $uu^{\top}$  with  $u \in \mathbb{R}^d$  is a stationary point of Equation (4.1) if and only if  $u \in \ker V$  or  $u^{\top}(A + A^{\top})u = 0$ .

Note that the set  $\{u \in \mathbb{R}^d : u^\top (A + A^\top)u = 0\}$  is the isotropic cone associated to the quadratic form  $A + A^\top$ . Its geometry depends on the signature of  $A + A^\top$ . It is non-trivial if and only if  $A + A^\top$  has zero eigenvalues or eigenvalues of the opposite sign.

Stationary points have low rank when V is identity and A is symmetric. Another interesting case is to look at stationary points of Equation (4.1) when  $V = I_d$  and A is symmetric. We have seen in Proposition 4.4 that when  $A + A^{\top} \leq 0$ , limiting points  $\Sigma^*$  must satisfy

$$\mathrm{rk}\Sigma^* \leq \mathrm{rk}A.$$

We generalize this result to any symmetric matrix A as follows (see Appendix C.1 for the proof).

PROPOSITION 4.8. Let  $A \in \mathbb{R}^{d \times d}$  be a symmetric matrix and assume that  $V = I_d$ . If a symmetric matrix  $\Sigma \in \mathbb{R}^{d \times d}$  is solution of

$$V\Sigma A\Sigma + \Sigma A^{\top}\Sigma V^{\top} = 0,$$

then

 $\operatorname{rk}\Sigma \leq \dim \ker A + \min(\# \{ \text{positive eigenvalues of } A \}, \# \{ \text{negative eigenvalues of } A \} \}.$ 

Therefore, when A is symmetric and  $V = I_d$ , the stationary points of Equation (4.1) have low rank—smaller than d/2 when A is invertible, for instance. Note that when A has a positive eigenvalue, we have seen in Proposition 4.4 that the dynamics

with a positive definite initialization blow up in finite time. In that case, stationary points are never reached.

**4.2.** Multi-Head Softmax Self-Attention. Lemma 4.1 allows us to derive the following result for multi-head Softmax self-attention.

PROPOSITION 4.9. Let  $d \in \mathbb{N}^*$ . Let  $Q, K, V : [0, +\infty) \to \mathbb{R}^{d \times d}$  be three continuous maps, modeling the evolution of parameters Q, K, V across layers of the Transformer. Consider a Gaussian initial condition  $\mu_0 = \mathcal{N}(\alpha_0, \Sigma_0)$  with  $\alpha_0 \in \mathbb{R}^d$  and  $\Sigma_0 \in \mathbb{R}^{d \times d}$ positive definite. Then, the multi-head Transformer PDE

$$\partial_t \mu + \operatorname{div}(\mu \Gamma_{\mu}^{(\mathrm{MH})}) = 0$$

associated to multi-head self-attention with parameters Q(t), K(t), V(t) via Equation (2.9) has a unique maximal solution  $\mu$  on  $[0, t_{\max})$ , such that  $\mu(t)$  is Gaussian for all  $t \in [0, t_{\max})$ . Moreover, denoting  $\mu(t) =: \mathcal{N}(\alpha(t), \Sigma(t))$ , we have

$$\dot{\Sigma} = \sum_{h=1}^{H} W^{(h)} V^{(h)} \Sigma A^{(h)} \Sigma + \Sigma A^{(h)\top} \Sigma (W^{(h)} V^{(h)})^{\top}.$$

and

$$\dot{\alpha} = \sum_{h=1}^{H} W^{(h)} V^{(h)} (I_d + \Sigma A^{(h)}) \alpha.$$

We only analyze the dynamics associated with Softmax multi-head attention numerically (see Section 4.5).

**4.3.** L2 Self-Attention. In the case of L2 self-attention, we have the following closed form for the velocity field when  $\mu$  is Gaussian.

LEMMA 4.10. Let  $\mu = \mathcal{N}(\alpha, \Sigma)$  be a Gaussian measure on  $\mathbb{R}^d$ . Then for all  $x \in \mathbb{R}^d$  it holds

$$\Gamma_{\mu}^{(L2)}(x) = V(\Sigma^{-1} + 2K^{\top}K)^{-1}(\Sigma^{-1}\alpha + 2K^{\top}Qx).$$

As  $\Gamma^{(L2)}_{\mu}$  is an affine function when  $\mu$  is Gaussian, Gaussianity is preserved by the L2 Transformer PDE, so we can rewrite it as two coupled matrix ODEs, with the same method as for Softmax self-attention.

PROPOSITION 4.11. Let  $d \in \mathbb{N}^*$ . Let  $Q, K, V \colon [0, +\infty) \to \mathbb{R}^{d \times d}$  be continuous functions. Consider a Gaussian initial condition  $\mu_0 = \mathcal{N}(\alpha_0, \Sigma_0)$  with  $\alpha_0 \in \mathbb{R}^d$  and  $\Sigma_0 \in \mathbb{R}^{d \times d}$  positive definite. Then, the Transformer equation

$$\partial_t \mu + \operatorname{div}(\mu \Gamma^{(\mathrm{L2})}_{\mu}) = 0$$

associated to L2 self-attention has a unique maximal solution  $\mu$ , such that  $\mu(t)$  is Gaussian for all  $t \in [0, t_{\max})$ . Moreover, denoting  $\mu(t) \eqqcolon \mathcal{N}(\alpha(t), \Sigma(t))$ , we have

(4.3) 
$$\dot{\Sigma} = 2V\Sigma (I_d + 2K^{\top}K\Sigma)^{-1}A\Sigma + 2\Sigma A^{\top} (I_d + 2K^{\top}K\Sigma)^{-1}\Sigma V^{\top}.$$

and

$$\dot{\alpha} = V(\Sigma^{-1} + 2K^{\top}K)^{-1}(\Sigma^{-1} + 2A)\alpha.$$

Contrary to what happens with Softmax self-attention, we show (see Appendix C.1) that the dynamics (4.3) cannot blow up in finite time.

LEMMA 4.12. Let  $\Sigma_0 \succ 0$ . The matrix-valued differential equation

$$\dot{\Sigma} = 2V\Sigma(I_d + 2K^{\top}K\Sigma)^{-1}A\Sigma + 2\Sigma A^{\top}(I_d + 2K^{\top}K\Sigma)^{-1}\Sigma V^{\top}$$

initialized at  $\Sigma_0$  has a unique global solution.

In this sense, L2 self-attention is therefore smoother than Softmax self-attention. This nicely complements—and is connected to—the result in [35] showing that L2 selfattention is globally Lipschitz continuous, contrary to Softmax self-attention. As the equation on  $\Sigma$  is more involved in the L2 case than in the Softmax self-attention case, let us focus only on dimension 1. A clustering phenomenon appears as for Softmax self-attention.

PROPOSITION 4.13. Let  $q, k, v \in \mathbb{R}$  and  $s_0 \in \mathbb{R}^*_+$ . Denote a := qk and consider the differential equation

$$\dot{s} = \frac{4avs^2}{1+2k^2s}$$

with initial condition  $s(0) = s_0$ .

1. If av > 0, then  $s(t) \to +\infty$  when  $t \to +\infty$ .

2. If av < 0, then  $s(t) \to 0$  when  $t \to +\infty$ .

*Proof.* If av > 0, then  $\dot{s} > 0$  so s is increasing. Hence, it converges in  $(s_0, +\infty]$ , and it cannot converge to a finite value as  $0 \notin [s_0, +\infty)$  is the only stationary point of the equation.

It av < 0, then s is decreasing, so it converges in  $[-\infty, s_0)$ . The only stationary point in this interval is 0, and  $s_0 > 0$ , which proves the claim as s(t) is continuous.

In Proposition 4.13, case 1 corresponds, in terms of the Gaussian measure of covariance s(t), to the mass spreading to infinity, while case 2 induces a clustering of the mass into one single Dirac. The 1-dimensional case for L2 self-attention is therefore close to Proposition 4.3 (dimension 1 for Softmax self-attention), where the finite-time blow-up of s is replaced by a divergence when  $t \to +\infty$ . Following this remark, we point out numerically in Section 4.5 that the Softmax and L2 self-attention dynamics are very close except when the former blows up. We also show in Section 4.5 that the clustering behavior observed with Softmax self-attention also occurs with L2 self-attention.

**4.4. Sinkhorn self-Attention.** Let us finally focus on Sinkhorn self-attention. We start by computing a closed form for the velocity field  $\Gamma_{\mu,\varepsilon}^{(\text{sink})}$  when  $\mu$  is a Gaussian probability measure.

LEMMA 4.14. Let  $\mu = \mathcal{N}(\alpha, \Sigma)$  be a Gaussian measure on  $\mathbb{R}^d$ . Let  $Q, K \in \mathbb{R}^{k \times d}$  be two matrices, denote  $A \coloneqq K^\top Q$  and assume that A and  $\Sigma$  are invertible. Then, for all  $x \in \mathbb{R}^d$ , it holds

$$\Gamma_{\mu,\varepsilon}^{(\text{sink})}(x) = \frac{1}{\varepsilon} V(I_d - A^{-\top} \Sigma^{-1} C) \alpha + \frac{1}{\varepsilon} V A^{-\top} \Sigma^{-1} C x,$$

where

$$C \coloneqq \Sigma^{1/2} (\Sigma^{1/2} A^{\top} \Sigma A \Sigma^{1/2} + \frac{\varepsilon^2}{4} I_d)^{1/2} \Sigma^{-1/2} - \frac{\varepsilon}{2} I_d$$

The proof is in Appendix C.1. Building on this result, we obtain with the same method as for Proposition 4.2 two matrix ODEs summarizing the Sinkformer PDE on Gaussian measures.

PROPOSITION 4.15. Let  $d \in \mathbb{N}^*$ . Let  $Q, K, V \colon [0, +\infty) \to \mathbb{R}^{d \times d}$  be continuous functions. Consider a Gaussian initial condition  $\mu_0 = \mathcal{N}(\alpha_0, \Sigma_0)$  with  $\alpha_0 \in \mathbb{R}^d$  and  $\Sigma_0 \in \mathbb{R}^{d \times d}$  positive definite. Then, the Sinkformer PDE

$$\partial_t \mu + \operatorname{div}(\mu \Gamma^{(\operatorname{sink})}_{\mu,\varepsilon}) = 0$$

associated to Sinkhorn self-attention has a unique maximal solution  $\mu$ , such that  $\mu(t)$  is Gaussian for all  $t \in [0, t_{\max})$ . Moreover, denoting  $\mu(t) \eqqcolon \mathcal{N}(\alpha(t), \Sigma(t))$ , we have

$$\dot{\Sigma} = \frac{1}{\varepsilon} V A^{-\top} \Sigma^{-1} C \Sigma + \frac{1}{\varepsilon} \Sigma C^{\top} \Sigma^{-1} A^{-1} V^{\top},$$

with  $C \coloneqq \Sigma^{1/2} (\Sigma^{1/2} A \Sigma A^{\top} \Sigma^{1/2} + \frac{\varepsilon^2}{4} I_d)^{1/2} \Sigma^{-1/2} - \frac{\varepsilon}{2} I_d$ , and

$$\dot{\alpha} = \frac{1}{\varepsilon} V \alpha.$$

Remarkably, in this case, and contrary to L2 and Softmax self-attention, the expectation  $\alpha$  and the covariance matrix  $\Sigma$  evolve independently. We have

$$\alpha(t) = e^{tV/\varepsilon} \alpha_0.$$

Remark 4.16. The fact that the expectation and the covariance evolve independently is not that surprising. Indeed, [47] show that, under certain assumptions on the parameters, the Sinkformer PDE corresponds to the Wasserstein gradient flow of a functional that takes the form  $\phi(\alpha) + \psi(\Sigma)$  on Gaussians, where  $\alpha$  and  $\Sigma$  are respectively the expectation and the covariance matrix of the Gaussian measure (see Equation (5.12)). Then, the Wasserstein gradient flow of  $\phi(\alpha) + \psi(\Sigma)$ , which stays in the space of Gaussian measures, corresponds to following a Euclidean gradient flow on  $\phi(\alpha)$  and a Bures-Wasserstein gradient flow on  $\psi(\Sigma)$  (see Section 5.3).

Contrary to the expectation equation, the covariance equation is much more challenging to be analyzed theoretically. We only consider the 1-dimensional case.

PROPOSITION 4.17. Let  $q, k, v \in \mathbb{R}$  and a := qk. Assume that  $\varepsilon = 1$ , as it can be absorbed in a and v. The covariance equation associated with Sinkhorn self-attention in dimension 1 reads

$$\dot{s} = \frac{v}{a}(\sqrt{4a^2s^2 + 1} - 1).$$

For any initial data  $s_0 \in \mathbb{R}^*_+$ , this equation has a global solution s(t), such that:

- 1. if av > 0, then  $s(t) \to +\infty$  when  $t \to +\infty$ ,
- 2. if av < 0, then  $s(t) \to 0$  when  $t \to +\infty$ .

The proof is the same as for Proposition 4.13. Therefore, in dimension 1, Sinkhorn self-attention is smoother than Softmax self-attention, and induces the same clustering phenomenon when av < 0.

**4.5. Experiments.** Our theoretical results about the Gaussian case typically rely on restrictive hypotheses on the weight matrices. We therefore complement the theoretical analysis with numerical experiments, to understand better the properties of the evolution and of the limiting covariance matrix  $\Sigma$  for each type of self-attention,



FIG. 1. Evolution of the covariance matrix of a 2-dimensional Gaussian measure that goes through the Transformer PDE. The plots (a), (b), and (d) were obtained with Softmax self-attention, respectively with (a) V random and  $A + A^{\top} \prec 0$ , (b)  $V = I_2$  and  $A + A^{\top} \preceq 0$  of rank 1 and (d) V and A chosen specifically to obtain this pattern. The plot (c) corresponds to multi-head self-attention with  $V = I_2$  and  $A + A^{\top} \preceq 0$  of rank 1.



FIG. 2. Projection on the set of trace-1 matrices of the dynamics of the covariance matrix of a Gaussian measure that goes through the Transformer PDE, in cases where curves blow up or diverge. The plots (a), (b), and (c) were obtained with the same parameters ( $V = I_2$  and Q, K fixed so that  $A + A^{\top} > 0$ ), respectively for Softmax, multi-head and L2 self-attention. In (a) and (b), the dynamics explode in finite time, while it is well-posed (but diverging) in (c). Finally, in (d), some of the initializations lead to a finite-time blow-up (purple curves) while others lead to convergence of the covariance matrix (yellow/green curves). (d) was obtained with Softmax self-attention but we observed a very similar behavior with L2 and multi-head self-attention (see Figure 6 in Appendix C.2).

depending on the parameters. In this section, we call a random matrix a matrix whose coordinates are i.i.d. Gaussian. We focus on Softmax, L2 and (Softmax) multi-head attention. Sinkhorn self-attention is investigated in Figure 7.

Qualitative behaviors for d = 2. The 2-dimensional case allows to visualize the different dynamics. Covariance matrices are then symmetric positive semi-definite  $2 \times 2$  matrices, and can be represented in a 3-dimensional space with the following change of coordinates:

$$\begin{pmatrix} a & c \\ c & b \end{pmatrix} \in \mathcal{S}_2^+ \mapsto (x, y, z) \coloneqq (a - b, 2c, a + b) \in \mathbb{R}^3.$$

With this choice of parametrization, the set of symmetric non-negative  $2 \times 2$  matrices becomes the cone of equation  $z \ge \sqrt{x^2 + y^2}$  (see Figure 1), and the evolution of a covariance matrix  $\Sigma(t)$  is represented as a curve inside this cone. Note that the boundary of the cone corresponds to degenerate matrices, i.e., of rank 1 or 0.

We highlight different behaviors which correspond to different choices of parameters and self-attention types. In each case, we set our initial points on a twodimensional grid, orthogonal to the z axis (so all initial matrices have the same trace), and we run a Euler discretization of the ODE satisfied by  $\Sigma$ , with a fixed step-size  $\tau$ . So the equation  $\dot{\Sigma} = g(\Sigma)$  is discretized as

$$\Sigma_{k+1} = \Sigma_k + \tau g(\Sigma_k)$$

starting from some initial point  $\Sigma_0$ . Depending on the parameters and the initial data, we observe a range of different behaviors, illustrated in Figure 1 and 2. Here are some typical evolutions of the covariance matrix.

- 1. Convergence to zero: independently of the initial data, all trajectories converge to the origin. We observe empirically that this is the case when the matrix  $A + A^{\top}$  is negative definite, independently of V, and for Softmax, multi-head and L2 self-attention (see Figure 4). However, this is not the case for Sinkhorn self-attention (see Figure 7). Figure 1 (a) was obtained with Softmax self-attention for a random choice of A provided that  $A + A^{\top} \prec 0$ , and a random choice of V.
- Convergence to a line: all trajectories converge to a line, which is the case for Softmax and L2 self-attention when A+A<sup>T</sup> is negative and of rank 1. Figure 1 (b) corresponds to this case with V random and Softmax self-attention.
- 3. Convergence to a plane: we observe this behavior with MH self-attention when  $A + A^{\top}$  is negative and of rank 1. Figure 1 (c) corresponds to this case for  $V = I_2$ .
- 4. Divergence in finite or infinite time: some initializations lead to a divergence of at least one of the eigenvalues of  $\Sigma$ , in finite time or when  $t \to +\infty$ . In that case, we stop the evolution at some fixed threshold for  $\|\Sigma\|$ , and then we plot in 2D the trajectory of the rescaled covariance matrices  $\frac{\Sigma}{\text{Tr}(\Sigma)}$ , which belong to a horizontal slice of the cone corresponding to matrices of trace 1 (see Figure 2). The plots (a), (b), and (c) of Figure 2 correspond to the same set of parameters, respectively for Softmax, multi-head, and L2 self-attention.

Therefore, when the dynamics is well-posed (no finite-time explosion of an eigenvalue), it appears to converge generically to a low-dimensional subspace, independently of the attention type. This parallels the clustering phenomenon that occurs with a finite number of tokens. We investigate this aspect in higher dimension in Paragraph 4.5.

Our figures also allow for an empirical comparison of the behaviors with the different types of self-attention. In particular, we observe that Softmax and L2 self-attention induce very similar behaviors (see Appendix C.2, Figure 5), except when the former blows up—in that case, L2 diverges but stays well-posed, according to Lemma 4.12, which induces very different trajectories (see for instance Figure 2 (a) and (c)).

*Clustering in higher dimension.* Proposition 4.4 shows that the limiting points of the covariance equation associated with Softmax self-attention have a low rank under some assumptions on the parameters. To complement this analysis, we highlight numerically that this holds generally, as soon as the dynamics converge, and not only for Softmax self-attention but also for L2 self-attention.

Figure 3 plots, in dimensions d = 3, 4, 5, the histogram of the rank of limiting points of the covariance equation for Softmax self-attention, with  $Q \in \mathbb{R}^{\lfloor d/2 \rfloor}$  random (different for each point), K = -Q so that  $A \coloneqq K^{\top}Q$  is symmetric negative and of rank  $\lfloor d/2 \rfloor$ , and  $V = I_d$  (upper row) or V random and different for each point (lower row). The case  $V = I_d$  is covered by Proposition 4.4, which tells us that limiting



FIG. 3. Histogram of the rank of limiting points of the covariance equation for Softmax selfattention, in dimensions 3, 4, and 5. The matrix V has full rank ( $V = I_d$  in the upper row and V random and different for each point in the lower row) and the matrix A has rank  $\lfloor d/2 \rfloor$ , is random (constrained to be negative symmetric) and different for each point. Limiting points have low rank (smaller than  $\lfloor d/2 \rfloor$ ), which parallels the clustering phenomenon observed for discrete tokens.

points have a rank smaller than  $\lceil d/2 \rceil$ : our numerical results confirm this result. Although no theoretical result covers the case V random, Figure 3 highlights that the bound  $\lceil d/2 \rceil$  still holds in that case. We display a similar figure for L2 self-attention in Appendix C.2.

5. Viewing the Dynamics as a Gradient Flow. This Section aims to connect the Transformer PDE with gradient flows in measure spaces. Let us first stress that the derivations in this Section are only formal, the goal being to gain some intuition on the gradient flow structure of the different Transformer evolutions considered so far, and to highlight, in particular, the structure of the Gaussian case. Making these calculations rigorous is beyond the scope of this paper. We start with some background on gradient flows with respect to a geodesic distance, which generalize Wasserstein gradient flows.

5.1. Limiting Minimizing Schemes for Geodesic Distances. We consider a distance  $\mathcal{D}(\mu,\nu)$  on the space  $\mathcal{P}(\mathbb{R}^d)$  of probability distributions on  $\mathbb{R}^d$ . For some function  $\mathcal{F}: \mathcal{P}(\mathbb{R}^d) \to \mathbb{R}$ , one can define minimizing evolutions on the metric space  $\mathcal{P}(\mathbb{R}^d)$  starting from  $\mu_0 \in \mathcal{P}(\mathbb{R}^d)$ , following [1], by considering implicit stepping with step-size  $\tau > 0$ :

(5.1) 
$$\mu_{t+\tau} \in \operatorname*{argmin}_{\mu \in \mathcal{P}(\mathbb{R}^d)} \left\{ \frac{1}{2\tau} \mathcal{D}(\mu_t, \mu)^2 + \mathcal{F}(\mu) \right\}.$$

Under suitable regularity properties on  $\mathcal{F}$ , it is possible to take the limit  $\tau \to 0$  and consider the continuous-time evolution  $t \mapsto \mu_t$ , which solves a PDE. The aim of this Subsection is to formally derive this PDE, called gradient flow of  $\mathcal{F}$  for the distance  $\mathcal{D}$ , and which takes the form of a continuity equation for some velocity field that depends on  $\mathcal{F}$ . Note that if  $\mathcal{D}$  is equal to the Wasserstein distance  $W_2$ , this is by now a textbook computation which corresponds to the JKO flow construction [33, 1, 49] of Wasserstein gradient flows. We consider the more general case where  $\mathcal{D}$  is a geodesic distance with the following dynamical formulation:

(5.2) 
$$\mathcal{D}(\mu,\nu)^2 = \inf_{\rho,\nu} \left\{ \int_0^1 \int \mathcal{G}_{\rho_s}(v_s) \cdot v_s \,\mathrm{d}\rho_s \,\mathrm{d}s : \partial_s \rho + \operatorname{div}(\rho \,\mathcal{G}_{\rho}(v_s)) = 0, \quad \begin{array}{c} \rho_0 = \mu \\ \rho_1 = \nu \end{array} \right\},$$

where  $\mathcal{G}_{\rho}$  is a linear invertible  $L_{\rho}^2$ -self-adjoint operator on vector fields. The classical distance  $\mathcal{D} = W_2$  corresponds to the choice  $\mathcal{G}_{\rho} = \text{Id}$ . We here follow a similar duality strategy to [5, 16] to formally rewrite the distance as a saddle point problem for which optimality conditions are easier to obtain. We first observe that we can dilate the time variable s and scale v such that

(5.3) 
$$\frac{1}{\tau}\mathcal{D}(\mu,\nu)^2 = \inf_{\rho,\nu} \left\{ \int_0^\tau \int \mathcal{G}_{\rho_s}(v_s) \cdot v_s \,\mathrm{d}\rho_s \,\mathrm{d}s : \partial_s \rho + \operatorname{div}(\rho \,\mathcal{G}_{\rho}(v_s)) = 0, \begin{array}{l} \rho_0 = \mu \\ \rho_\tau = \nu \end{array} \right\},$$

Using a Lagrangian dual formulation of  $\mathcal{D}$ , we rewrite

$$\mu_{t+\tau} \in \underset{\mu}{\operatorname{argmin}} \ 2\mathcal{F}(\mu) + \underset{\rho,v}{\inf} \underset{\psi}{\sup} \int_{0}^{\tau} \int \mathcal{G}_{\rho}(v) \cdot v \, \mathrm{d}\rho \, \mathrm{d}s \\ - \int_{0}^{\tau} \int (\partial_{s}\psi + \mathcal{G}_{\rho}(v) \cdot \nabla_{x}\psi) \, \mathrm{d}\rho \, \mathrm{d}s + \int \psi(\tau, \cdot) \, \mathrm{d}\mu - \int \psi(0, \cdot) \, \mathrm{d}\mu_{t}.$$

The optimality condition on  $\mu_{t+\tau}$  is

(5.4) 
$$2\delta \mathcal{F}(\mu_{t+\tau}) + \psi(\tau, \cdot) = 0,$$

where  $\delta \mathcal{F}$  is the first variation (Fréchet derivative) of  $\mathcal{F}$ , i.e.,

$$\mathcal{F}(\mu + h\nu) = \mathcal{F}(\mu) + h \int \delta \mathcal{F}(\mu) \,\mathrm{d}\mu + o(h).$$

Equation (5.4) formally gives

(5.5) 
$$\frac{1}{2}\nabla_x\psi = -\nabla_x\delta\mathcal{F}(\mu),$$

when  $\tau \to 0^+$ . Moreover, we find the following optimality condition for v given by

(5.6) 
$$v = \frac{1}{2} \nabla_x \psi,$$

by taking Fréchet derivatives with respect to v in (5.3) and using that  $\mathcal{G}_{\rho}$  is a linear invertible  $L^2_{\rho}$ -self-adjoint operator on vector fields. Combining Equations (5.5) and (5.6), one can rewrite  $\partial_t \rho + \operatorname{div}(\rho \mathcal{G}_{\rho}(v)) = 0$  as the following non-linear advection equation

(5.7) 
$$\partial_t \rho + \operatorname{div}(\rho \, \Gamma_\rho) = 0.$$

with  $\Gamma_{\rho} := -\mathcal{G}_{\rho}(\nabla_x[\delta \mathcal{F}(\mu)])$ . In this sense, we say that the PDE (5.7) is the gradient flow of  $\mathcal{F}$  for the distance  $\mathcal{D}$ . Notice that for the classical case with  $\mathcal{D} = W_2$  so that  $\mathcal{G}_{\mu} = \text{Id}$ , then  $\Gamma_{\rho}$  is the standard Wasserstein gradient.

Remark 5.1. Classical free energy functionals are of the form

(5.8) 
$$\mathcal{F}(\rho) = \int U(\rho) \, \mathrm{d}x + \int V(x) \, \rho(x) \, \mathrm{d}x + \frac{1}{2} \int W(x-y) \, \rho(x) \, \rho(y) \, \mathrm{d}x \, \mathrm{d}y.$$

where  $U: \mathbb{R}^+ \to \mathbb{R}$  is a density of internal energy,  $V: \mathbb{R}^d \to \mathbb{R}$  is a confinement potential and  $W: \mathbb{R}^d \to \mathbb{R}$  is an interaction potential, see for instance [17, 54]. The corresponding variation is given by  $\delta \mathcal{F}(\rho) = U'(\rho) + V + W * \rho$ . Without interaction potential W = 0, this general family of PDEs, with  $\mathcal{D} = W_2$  so that  $\mathcal{G}_{\mu} = \text{Id}$ , contains well-known models in mathematical physics such as the heat equation,  $U(s) = s \log s$  and V = 0, the porous-medium and fast-diffusion equations [53],  $U(s) = s^m/(m-1)$ , m > 0 and V = 0 and their Fokker-Planck counterparts,  $U(s) = s \log s$  and  $V(x) = |x|^2/2$ , and  $U(s) = s^m/(m-1)$  and  $V(x) = |x|^2/2$  respectively, see [44] for instance. With non trivial interactions potentials, it includes many important PDEs in mathematical biology and mathematical physics, such as the Keller-Segel model, see [12] and the references therein, or nonlocal McKean-Vlasov equations, see [14] and the references therein.

5.2. Restriction to the Subspace of Gaussians. In Section 4, we have seen that the Transformer PDE with a Gaussian measure as initial data stays in the space of Gaussians over time, for several variants of self-attention. For Transformer PDEs that have a gradient flow structure as introduced in Subsection 5.1, the evolution of the Gaussian case can be inferred from the gradient flow structure, and follows a Riemannian gradient flow. This Subsection details that connection, which allows us to check the Gaussian evolutions obtained in Section 4.

Let  $\mathcal{D}$  be a geodesic distance. We assume that  $\mathcal{D}$  is translation invariant, in the sense that:

$$\mathcal{D}(\mu, \nu) = \mathcal{D}(\mu_0, \nu_0)^2 + |m(\mu) - m(\nu)|^2$$

where  $|\cdot|$  is the Euclidean norm,  $m(\mu) \coloneqq \int x d\mu(x)$  is the mean of  $\mu$ , and  $\mu_0 = T_{\#}\mu$ with  $T(x) = x - m(\mu)$  (so that  $\mu_0$  and  $\nu_0$  have zero mean). This is the case for the Wasserstein distance and the twisted distance (5.14) considered below. The metric  $\mathcal{D}$ descends to a finite-dimensional metric D on the cone  $\mathcal{S}_d^{++}$  of covariances:

$$D(\Sigma, \Sigma') \coloneqq \mathcal{D}(\mathcal{N}(0, \Sigma), \mathcal{N}(0, \Sigma')),$$

where  $\mathcal{N}(\alpha, \Sigma)$  is the Gaussian measure with covariance  $\Sigma$  and mean  $\alpha$ . If the initial data  $\mu_{t=0}$  is Gaussian and the vector field  $\Gamma_{\mathcal{N}(\alpha,\Sigma)}$  is affine, solutions of Equation (5.7) are Gaussian for all time, as seen in Section 4: we can write  $\mu_t =: \mathcal{N}(\alpha(t), \Sigma(t))$ , and  $t \mapsto (\alpha(t), \Sigma(t))$  follows a Riemannian gradient flow of the finite-dimensional function

$$F(\alpha, \Sigma) \coloneqq \mathcal{F}(\mathcal{N}(\alpha, \Sigma)),$$

where the Riemannian structure is induced by D as follows.

Similarly to Equation (5.1), the Riemannian flow induced by the gradient flow on the space of Gaussians can be defined via an implicit stepping:

$$(\alpha(t+\tau), \Sigma(t+\tau)) \in \operatorname*{argmin}_{\alpha, \Sigma} D(\Sigma, \Sigma(t))^2 + |\alpha - \alpha(t)|^2 + 2\tau F(\alpha, \Sigma),$$

where the minimization is restricted to semi-definite positive matrices. Taking the limit  $\tau \to 0$  yields a continuous-time trajectory  $t \mapsto (\alpha(t), \Sigma(t))$ . This flow satisfies:

(5.9) 
$$\frac{\mathrm{d}\alpha}{\mathrm{d}t} = \nabla_{\alpha} F(\alpha(t), \Sigma(t)),$$
$$\frac{\mathrm{d}\Sigma}{\mathrm{d}t} = M_{\Sigma}^{-1} \nabla_{\Sigma} F(\alpha(t), \Sigma(t))$$

where the Riemannian metric  $M_{\Sigma} : \mathbb{R}^{d \times d} \to \mathbb{R}^{d \times d}$  is defined as follows:

(5.10) 
$$D(\Sigma, \Sigma')^2 = \langle M_{\Sigma}(\Sigma - \Sigma'), \Sigma - \Sigma' \rangle_{\mathbb{R}^{d \times d}} + o(\|\Sigma - \Sigma'\|_F^2).$$

The following two Subsections detail the gradient flow structure of the Transformer PDE respectively for Softmax and Sinkhorn self-attention, in the general case and the Gaussian case, building on what has been introduced above.

**5.3. Sinkformer: Wasserstein Flow.** As shown in [47], for the specific case of Sinkhorn attention layers, and under the assumption that  $A = A^{\top} = -V$  (with  $A := K^{\top}Q$ ), the Sinkformer PDE is a Wasserstein gradient flow, i.e., it satisfies Equation (5.7) for  $\mathcal{D} = W_2$  (i.e.,  $\mathcal{G}_{\mu} = \text{Id}$ ), and for the functional:

(5.11) 
$$\mathcal{F}_{\varepsilon}(\mu) \coloneqq -\frac{1}{2} \int \kappa_{\mu,\varepsilon}^{\infty} \log\left(\frac{\kappa_{\mu,\varepsilon}^{\infty}}{\kappa_{\mu,\varepsilon}^{0}}\right) \,\mathrm{d}(\mu \otimes \mu) + \frac{1}{4\varepsilon} \int (|Qx|^{2} + |Kx|^{2}) \,\mathrm{d}\mu(x)$$

Remark 5.2. The second term in Equation (5.11) does not appear in [47], because the authors define  $\kappa^0(x,y) \coloneqq \exp(Qx \cdot Ky/\varepsilon)$ , while we choose  $\kappa^0(x,y) \coloneqq \exp(-|Qx - Ky|^2/2\varepsilon)$ . These two choices lead to the same  $\kappa^{\infty}$ .

As seen in Section 4.4, in this case, for a Gaussian  $\mu$ , the vector field  $\Gamma_{\mu}$  is an affine function. Consequently, one can also consider this evolution as a Riemannian gradient flow over the mean and covariance space, according to Subsection 5.2. The corresponding Riemannian metric is the so-called Bures-Wasserstein metric, whose inverse reads [41]:

$$M_{\Sigma}^{-1}: A \in \mathbb{R}^{d \times d} \mapsto A\Sigma + \Sigma A^{\top}.$$

When restricted to the space of Gaussian distributions, we show in Appendix D.1 that the Sinkformer evolution (1.2) corresponds to the Riemannian flow ODE (5.9) for the function:

(5.12)

$$F_{\varepsilon}(\alpha, \Sigma) = \frac{1}{4\varepsilon} \left( -\mathfrak{B}_{\varepsilon}^{2}(\Sigma, \Sigma) + \operatorname{Tr}(Q\Sigma Q^{\top}) + \operatorname{Tr}(K\Sigma K^{\top}) + \alpha^{\top}(Q^{\top}Q + K^{\top}K)\alpha \right),$$

where  $\mathfrak{B}^2_{\varepsilon}(\Sigma_1, \Sigma_2)$  is the entropy-regularized Bures distance, defined as

(5.13) 
$$\mathfrak{B}^2_{\varepsilon}(\Sigma_1, \Sigma_2) \coloneqq 2\varepsilon OT_{\varepsilon}(\mathcal{N}(\alpha_1, \Sigma_1), \mathcal{N}(\alpha_2, \Sigma_2)) - |\alpha_1 - \alpha_2|^2$$

where  $OT_{\varepsilon}$  is defined in Equation (2.5). Note that these Bures flows have several favorable properties. In particular, the low-rank manifold is closed, meaning that the rank cannot increase during evolution. However, it is possible for the rank to decrease, and even if the initial covariance is full-rank, it typically becomes rank-deficient at the final time, as illustrated numerically in Section 4.5.

5.4. Softmax Transformer: A Twisted Wasserstein Flow. For a finite number of tokens, the Transformer evolution can be recasted as a gradient flow for a particular non-Euclidean metric [28]. We generalize this construction to probability measures by defining a twisted Wasserstein distance denoted  $d_{A,V}$ , associated with:

(5.14) 
$$\mathcal{G}_{\mu}(v) \colon x \mapsto \frac{Bv(x)}{\int e^{Ax \cdot y} \,\mathrm{d}\mu(y)},$$

with the notations of Equation (5.2), and where  $B := -VA^{-\top}$ . The twisted Wasserstein distance  $d_{A,V}$  therefore mimics the dynamical formulation of the Wasserstein distance, but changing the mobility, in the spirit of [16, 38].

Using this metric, we obtain with the computation of Subsection 5.1 that when B is symmetric and positive definite, the Transformer PDE becomes the gradient flow PDE (5.7) for the quadratic interaction functional:

$$\mathcal{F}(\mu) \coloneqq \frac{1}{2} \int e^{Ax \cdot y} \,\mathrm{d}\mu(x) \,\mathrm{d}\mu(y).$$

When restricted to the space of Gaussian distributions, the Transformer evolution (1.2) corresponds to the Riemannian flow ODE (5.9) for the function:

$$F(\alpha, \Sigma) \coloneqq \frac{e^{\frac{1}{2}\alpha^{\top}((A+\Sigma^{-1})^{\top}(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})-\Sigma^{-1})\alpha}}{2|\det(I_d - A\Sigma A^{\top}\Sigma)|^{1/2}}$$

and the Riemannian metric induced by  $d_{A,V}$  on the space of Gaussians via Equation (5.10). The computation is in Appendix D.2.

An important open question is to know whether the energy functional  $\mathcal{F}$  is geodesically convex for the twisted Wasserstein distance. We provide a first answer by showing that this is not the case *for any choice of parameters*. In future work, it could be interesting to investigate the geodesic convexity of  $\mathcal{E}$  restricted to smaller classes of measures, or subject to some assumptions on the parameters.

We have the following characterization of the geodesics of  $d_{A,V}$  (see Appendix D.2).

LEMMA 5.3. Let A, V such that  $B \coloneqq -VA^{-\top}$  is symmetric and positive definite. The geodesics  $\rho$  of  $d_{A,V}$  are characterized by

$$\begin{cases} \partial_s \rho + \operatorname{div}\left(\frac{B\nabla_x \psi}{2G*\rho}\rho\right) = 0\\ \partial_s \psi + \frac{1}{4} \frac{\nabla_x \psi \cdot B\nabla_x \psi}{G*\rho} - \frac{1}{4} \int_{\mathbb{R}^d} \frac{\nabla_y \psi \cdot B\nabla_y \psi}{(G*\rho(y))^2} G(y, x) \rho_s(y) \, \mathrm{d}y = 0 \end{cases}$$

for some function  $\psi$ .

Building on this computation, we prove the following negative result (see Appendix D.2).

PROPOSITION 5.4. Let  $A, V \in \mathbb{R}^{d \times d}$  such that  $B \coloneqq -VA^{-\top}$  is symmetric and positive definite. Assume that V is symmetric positive definite and commutes with B. Then  $\mathcal{E}$  is not geodesically convex for the distance  $d_{A,V}$ .

6. Conclusion. This work studies the Transformer PDE, which models the evolution of data that goes through the layers of a deep Transformer model. For several variants of self-attention, we show that the Transformer PDE is well-posed when the initial data is compactly supported, and we derive a stability estimate with respect to the initial condition. Our framework includes in particular masked self-attention. We also consider the case of a Gaussian initial condition, which has the nice property to stay Gaussian across the dynamics for several attention variants. Building on this remark, we show both theoretically and numerically that when the Gaussian evolution converges, the covariance of the limiting Gaussian measure is rank-deficient, which parallels the clustering phenomenon observed with discrete tokens. Finally, we draw a connection between our framework and gradient flows, by introducing formally a distance on probability measures that equips the Transformer PDE associated with Softmax self-attention with a gradient flow structure.

Acknowledgments. The research of JAC was supported by the Advanced Grant Nonlocal-CPD (Nonlocal PDEs for Complex Particle Dynamics: Phase Transitions, Patterns and Synchronization) of the European Research Council Executive Agency (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 883363). JAC was also partially supported by EPSRC grant number EP/V051121/1. JAC was also partially supported by the "Maria de Maeztu" Excellence Unit IMAG, funded by MCIN/AEI/10.13039/501100011033/, reference CEX2020-001105-M. The work of G. Peyré was supported by the French government under the management of Agence Nationale de la Recherche as part of the "France 2030" program, reference ANR-23-IACL-0008 (PRAIRIE-PSAI). The work of G. Peyré and V. Castin was supported by the ERC project WOLF.

## Appendix A. Background on Entropic Optimal Transport.

A.1. Dual Formulation of EOT. Let  $\mu$  and  $\nu$  be two compactly supported or Gaussian probability measures on  $\mathbb{R}^d$ . Recall the entropic optimal transport problem

(A.1) 
$$OT_{\varepsilon}(\mu,\nu) \coloneqq \min_{\pi \in \Pi(\mu,\nu)} \int c_{\varepsilon}(x,y) \, \mathrm{d}\pi(x,y) + \mathrm{KL}(\pi \| \mu \otimes \nu),$$

introduced in Section 2, where  $c_{\varepsilon}(x,y) \coloneqq \frac{1}{2\varepsilon} |Qx - Ky|^2$ . It is well-known [32] that problem (A.1) admits the following dual formulation

$$OT_{\varepsilon}(\mu,\nu) = \max_{\substack{f \in \mathcal{L}^{1}(\mu) \\ g \in \mathcal{L}^{1}(\nu)}} \int_{\mathbb{R}^{d}} f \,\mathrm{d}\mu + \int_{\mathbb{R}^{d}} g \,\mathrm{d}\nu + 1 - \int_{\mathbb{R}^{d} \times \mathbb{R}^{d}} e^{f(x) + g(y) - c_{\varepsilon}(x,y)} \,\mathrm{d}\mu(x) \,\mathrm{d}\nu(y),$$

and that a pair of dual potentials (f, g) is optimal if and only if it satisfies the following optimality conditions respectively  $\nu$ - and  $\mu$ -almost everywhere:

(A.2) 
$$\begin{cases} f(x) = -\log \int e^{g(y) - c_{\varepsilon}(x,y)} d\nu(y) \\ g(y) = -\log \int e^{f(x) - c_{\varepsilon}(x,y)} d\mu(x). \end{cases}$$

There always exist solutions f and g that satisfy Equation (A.2) for every  $(x, y) \in \mathcal{X} \times \mathcal{Y}$  [43]. Let us focus on such pairs. Then, (f, g) is unique modulo the equivalence relation

$$(f_1, g_1) \sim (f_2, g_2) \Leftrightarrow \exists \eta \text{ s.t. } f_1 = f_2 + \eta \text{ and } g_1 = g_2 - \eta.$$

Let then (f, g) be any couple in the equivalence class. We call f and g Schrödinger potentials or dual potentials. They inherit the regularity of c, and satisfy the following useful relation:

$$\kappa_{\mu,\varepsilon}^{\infty}(x,y) = e^{f(x) + g(y) - c_{\varepsilon}(x,y)},$$

We can then rewrite  $OT_{\varepsilon}(\mu, \nu)$  as follows: (A.3)

$$OT_{\varepsilon}(\mu,\nu) = \frac{1}{2\varepsilon} \int |Qx - Ky|^2 \,\kappa_{\mu,\varepsilon}^{\infty}(x,y) \,\mathrm{d}\mu(x) \,\mathrm{d}\nu(y) + \int \log(\kappa_{\mu,\varepsilon}^{\infty}) \kappa_{\mu,\varepsilon}^{\infty} \,\mathrm{d}\mu(x) \,\mathrm{d}\nu(y).$$

Assume now that  $\mu$  and  $\nu$  are compactly supported, with a support included in the ball  $B_R$ . The dual potentials are both Lipschitz continuous.

LEMMA A.1. Assume that  $\operatorname{Supp} \mu$ ,  $\operatorname{Supp} \nu \subset B_R$ . Let  $c: (x, y) \in (\mathbb{R}^d)^2 \mapsto \frac{1}{2\varepsilon} |Qx - Ky|^2$  and (f, g) a couple of dual potentials satisfying Equation (A.2) everywhere. Then

(A.4) 
$$\sup_{x,y\in B_R} f(x) + g(y) \le \frac{1}{\varepsilon} (\|Q\|_2 + \|K\|_2)^2 R^2.$$

Moreover, f and g are both Lipschitz continuous, with a Lipschitz constant bounded by  $\frac{1}{\varepsilon}(\|Q\|_2 + \|K\|_2)^{3/2}R$ .

*Proof.* Equation (A.4) derives from [43], Lemma 4.9, noticing that  $c_{\varepsilon}(x, y) \leq \frac{1}{\varepsilon}(\|Q\|_2 + \|K\|_2)^2 R^2$  for  $x, y \in B_R$ . Then, according to [43] Lemma 4.11, the dual potentials are Lipschitz continuous with the same Lipschitz constant as  $c_{\varepsilon}$ . It is then straightforward to check that  $c_{\varepsilon}$  is  $\frac{1}{\varepsilon}(\|Q\|_2 + \|K\|_2)^{3/2}R$ -Lipschitz continuous in both variables.

Finally, we have the following stability result.

LEMMA A.2 ([10]). For any compactly supported distributions  $\mu, \nu$  on  $\mathbb{R}^d$ , consider the entropic optimal transport problems  $OT_{\varepsilon}(\mu, \mu)$  and  $OT_{\varepsilon}(\nu, \nu)$  for the cost  $c(x, y) \coloneqq \frac{1}{2\varepsilon} |Qx - Ky|^2$ , and denote  $(f^{\mu}, g^{\mu})$  and  $(f^{\nu}, g^{\nu})$  associated Schrödinger potentials. Then, there exists a function  $C_{\text{stab}}(R) > 0$  depending on  $R, Q, K, \varepsilon$  such that for all compactly supported probability measures  $\mu$  and  $\nu$  on  $\mathbb{R}^d$ , we have

$$\inf_{\eta \in \mathbb{R}} \|f^{\mu} - f^{\nu} - \eta\|_{\infty} + \|g^{\mu} - g^{\nu} + \eta\|_{\infty} \le C_{\text{stab}}(R)W_{2}(\mu, \nu).$$

**A.2. EOT Between Gaussians.** We have the following generalization of Theorem 1 in [32], where we take the cost  $\frac{1}{2\varepsilon} |Qx - Ky|^2$  instead of the classical quadratic cost. Note that a similar result is stated in [4], with a generalization to the case A non-invertible.

THEOREM A.3. Let  $\mu = \mathcal{N}(\alpha, \Sigma)$  and  $\nu = \mathcal{N}(\beta, \Omega)$  be two Gaussian measures. Set two matrices Q and K in  $\mathbb{R}^{d \times d}$ , and  $\varepsilon > 0$ . Denote  $A = K^{\top}Q$  and assume that A is invertible. Denote  $\pi^*$  the minimizer of the following entropy-regularized optimal transport problem:

(A.5) 
$$\min_{\pi \in \Pi(\mu,\nu)} \int \frac{1}{2\varepsilon} |Qx - Ky|^2 d\pi(x,y) + \mathrm{KL}(\pi || \mu \otimes \nu),$$

where  $\Pi(\mu, \nu)$  is the set of couplings between  $\mu$  and  $\nu$ , and

$$\mathrm{KL}(\pi \| \mu \otimes \nu) = \int \log \left( \frac{\mathrm{d}\pi}{\mathrm{d}(\mu \otimes \nu)} \right) \,\mathrm{d}\pi$$

is the Kullback-Leibler divergence between  $\pi$  and  $\mu \otimes \nu$ , set equal to  $+\infty$  if  $\pi$  is not absolutely continuous with respect to  $\mu \otimes \nu$ . Then, the optimal coupling  $\pi^*$  is a Gaussian measure, given by

$$\pi^* = \mathcal{N}\left( \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \begin{pmatrix} \Sigma & A^{-1}C^{\top} \\ CA^{-\top} & \Omega \end{pmatrix} \right),$$

with

$$C \coloneqq \Omega^{1/2} \left( \Omega^{1/2} A \Sigma A^{\top} \Omega^{1/2} + \frac{\varepsilon^2}{4} I_d \right)^{1/2} \Omega^{-1/2} - \frac{\varepsilon}{2} I_d.$$

Proof of Theorem A.3. We follow the steps of the proof of Theorem 1 [32]. First of all, we can assume that  $\mu$  and  $\nu$  are centered (Lemma 1 [32]). Then, using the dual formulation of Problem (A.5) introduced in Section A.1, we have that

(A.6) 
$$\frac{\mathrm{d}\pi^*}{\mathrm{d}\mu\otimes\nu}(x,y) = e^{f(x)+g(y)-\frac{|Qx-Ky|^2}{2\varepsilon}},$$

where  $f, g: \mathbb{R}^d \to \mathbb{R}^d$  are optimal dual potentials, that can be obtained as the limit of the following recursion, starting from a potential  $f_0$ :

(A.7)  
$$g_{n+1}(y) = -\log \int e^{f_n(x) - \frac{|Qx - Ky|^2}{2\varepsilon}} d\mu(x)$$
$$f_{n+1}(x) = -\log \int e^{g_{n+1}(y) - \frac{|Qx - Ky|^2}{2\varepsilon}} d\nu(y)$$

for  $n \ge 0$ . Let us choose  $f_0$  to be a quadratic form, and show that the iterates of the recursion (A.7) stay in the space of quadratic forms. For any symmetric matrix  $X \in \mathbb{R}^{d \times d}$ , denote

$$\mathcal{Q}(X): x \in \mathbb{R}^d \mapsto -\frac{1}{2}x^\top Xx.$$

We have the following Lemma.

LEMMA A.4. Let  $\mu = \mathcal{N}(0, \Sigma)$  and  $\nu = \mathcal{N}(0, \Omega)$ , and let X be a symmetric  $d \times d$  matrix and  $m \in \mathbb{R}$  a constant.

(i) The expression

$$T_{\mu,Q,K}(\mathcal{Q}(X)+m) \coloneqq -\log \int e^{-|Qx-Ky|^2/2\varepsilon + \mathcal{Q}(X)(x)+m} \,\mathrm{d}\mu(x)$$

for  $x \in \mathbb{R}^d$  is well-defined if and only if

$$X' \coloneqq Q^\top Q + \varepsilon \Sigma^{-1} + \varepsilon X \succ 0$$

and in that case,  $T_{\mu,Q,K}(\mathcal{Q}(X) + m)$  is a quadratic form, equal to  $\mathcal{Q}(Y)$  up to a constant term, where

$$Y \coloneqq \frac{1}{\varepsilon} (AX'^{-1}A^{\top} - K^{\top}K).$$

(ii) The expression

$$T_{\nu,K,Q}(\mathcal{Q}(X)+m) \coloneqq -\log \int e^{-|Qx-Ky|^2/2\varepsilon + \mathcal{Q}(X)(y)+m} \,\mathrm{d}\nu(y)$$

for  $x \in \mathbb{R}^d$  is well-defined if and only if

$$X'\coloneqq K^\top K + \varepsilon \Omega^{-1} + \varepsilon X \succ 0$$

and in that case, it is a quadratic form, equal to  $\mathcal{Q}(Y)$  up to a constant term, where

$$Y \coloneqq \frac{1}{\varepsilon} (A^{\top} X'^{-1} A - Q^{\top} Q).$$

With the notations of Lemma A.4, we can rewrite the recursion (A.7) as

$$g_{n+1} = T_{\mu,Q,K} (f_n)$$
  
$$f_{n+1} = T_{\nu,K,Q} (g_{n+1}).$$

Let us initialize it with

$$f_0 = \mathcal{Q}(0).$$

According to Lemma A.4, for all  $n \geq 1$ , we can write  $f_n$  and  $g_n$  as quadratic forms  $\mathcal{Q}(U_n)$  and  $\mathcal{Q}(V_n)$  up to constant terms. Assume that  $f_0 = \mathcal{Q}(0)$ , and define  $f_n$  and  $g_n$  with Equation (A.7). Then, we can write

$$f_n = \mathcal{Q}(U_n) + \operatorname{cst}$$
 and  $g_n = \mathcal{Q}(V_n) + \operatorname{cst}$ 

where

$$V_{n+1} = \frac{1}{\varepsilon} \left( A(Q^{\top}Q + \varepsilon \Sigma^{-1} + \varepsilon U_n)^{-1} A^{\top} - K^{\top} K \right)$$
$$U_{n+1} = \frac{1}{\varepsilon} \left( A^{\top} (K^{\top}K + \varepsilon \Omega^{-1} + \varepsilon V_{n+1})^{-1} A - Q^{\top} Q \right).$$

In particular, for all  $n \ge 0$ , the matrices  $Q^{\top}Q + \varepsilon \Sigma^{-1} + \varepsilon U_n$  and  $K^{\top}K + \varepsilon \Omega^{-1} + \varepsilon V_{n+1}$ are positive definite. Indeed, denote  $F_0 \coloneqq Q^{\top}Q + \varepsilon \Sigma^{-1}$  and  $G_0 \coloneqq K^{\top}K + \varepsilon \Omega^{-1}$ , and

(A.8) 
$$F_n \coloneqq Q^\top Q + \varepsilon \Sigma^{-1} + \varepsilon U_n,$$
$$G_n \coloneqq K^\top K + \varepsilon \Omega^{-1} + \varepsilon V_n$$

for all  $n \ge 1$ . We have the following recursion:

(A.9) 
$$\begin{aligned} F_{n+1} &= \varepsilon \Sigma^{-1} + A^{\top} G_n^{-1} A, \\ G_{n+1} &= \varepsilon \Omega^{-1} + A F_n^{-1} A^{\top}, \end{aligned}$$

which proves that  $F_n$  and  $G_n$  stay positive definite along the iterations.

LEMMA A.5. The sequences of matrices  $(F_n)_{n\geq 0}$  and  $(G_n)_{n\geq 0}$  defined by  $F_0 := Q^{\top}Q + \varepsilon \Sigma^{-1}$  and  $G_0 := K^{\top}K + \varepsilon \Omega^{-1}$ , and

$$F_{n+1} \coloneqq \varepsilon \Sigma^{-1} + A^\top G_n^{-1} A,$$
  
$$G_{n+1} \coloneqq \varepsilon \Omega^{-1} + A F_n^{-1} A^\top$$

for all  $n \ge 0$  converge towards positive definite matrices F and G.

*Proof.* Denoting

$$\phi(M) = \varepsilon \Sigma^{-1} + A^{\top} (\varepsilon \Omega^{-1} + A M^{-1} A^{\top})^{-1} A,$$

for any  $M \succ 0$ , we have  $F_{n+1} = \phi(F_n)$  for all  $n \ge 0$ . A similar computation as in the proof of Proposition 2 [32] shows that the operator norm of the differential of  $\phi$  at M is equal to

$$\|D_M\phi\|_2 = \|A^{\top}(\varepsilon\Omega^{-1} + AM^{-1}A^{\top})^{-1}AM^{-1}\|_2^2.$$

Let us assume for now that A is invertible. Then, it holds

$$A^{\top}(\varepsilon \Omega^{-1} + AM^{-1}A^{\top})^{-1}AM^{-1} = (I_d + \varepsilon MA^{-1}\Omega^{-1}A^{-\top})^{-1}.$$

Denoting  $\lambda_1(X) \geq \cdots \geq \lambda_d(X)$  the eigenvalues of any matrix X, we have

$$\begin{split} \left\| A^{\top} (\varepsilon \Omega^{-1} + A M^{-1} A^{\top})^{-1} A M^{-1} \right\|_{2} &= \frac{1}{\lambda_{d} (I_{d} + \varepsilon M A^{-1} \Omega^{-1} A^{-\top})} \\ &\leq \frac{1}{1 + \varepsilon \lambda_{d}(M) \lambda_{d}(A^{-1}) \lambda_{d}(\Omega^{-1}) \lambda_{d}(A^{-\top})} \end{split}$$

Applying Weyl's inequality to the decomposition

$$F_{n+1} = \varepsilon \Sigma^{-1} + A^\top (\varepsilon \Omega^{-1} + A F_n^{-1} A^\top)^{-1} A$$

yields, noticing that the matrix  $A^{\top}(\varepsilon \Omega^{-1} + AF_n^{-1}A^{\top})^{-1}A$  is positive:

$$\lambda_d(F_{n+1}) \ge \frac{\varepsilon}{\lambda_1(\Sigma)},$$

which is also true for  $\lambda_d(F_0)$ . Moreover, for all  $M \succ 0$  such that

$$\lambda_d(M) \ge \frac{\varepsilon}{\lambda_1(\Sigma)},$$

we have

$$\|D_M\phi\|_2 \le \frac{1}{1 + \frac{\varepsilon^2}{\lambda_1(\Sigma)\lambda_1(A)^2\lambda_1(\Omega)}} < 1.$$

An approximation argument shows that this inequality stays true even for noninvertible matrices A. Therefore, we have bounded the operator norm of the differential of  $\phi$  uniformly away from 1 along the trajectory of  $(F_n)$ , which shows that  $F_n$  converges. The same method applies to prove convergence of  $(G_n)$ .

As  $F_n$  and  $G_n$  are positive matrices for all n, the limits F and G are non-negative matrices. Moreover, taking the limit  $n \to \infty$  in Equation (A.9) gives

(A.10) 
$$F = \varepsilon \Sigma^{-1} + A^{\top} G^{-1} A,$$
$$G = \varepsilon \Omega^{-1} + A F^{-1} A^{\top},$$

which shows that F and G are positive matrices.

By taking the limit  $n \to \infty$  in Equations (A.8) and (A.9), we obtain the following relations:

(A.11) 
$$F = Q^{\top}Q + \varepsilon \Sigma^{-1} + \varepsilon U,$$
$$G = K^{\top}K + \varepsilon \Omega^{-1} + \varepsilon V,$$

where the optimal potentials f and g defined in Equation (A.6) can be written as

$$f = \mathcal{Q}(U) + \operatorname{cst}$$
 and  $g = \mathcal{Q}(V) + \operatorname{cst}$ .

We can now show that the optimal coupling  $\pi^*$  is Gaussian, and write its covariance matrix in terms of F and G.

LEMMA A.6. Let  $\pi^*$  be the optimal coupling defined in Theorem A.3, and F and G defined in Equation (A.11). Then  $\pi^*$  is a Gaussian measure on  $\mathbb{R}^d \times \mathbb{R}^d$ , whose covariance matrix is equal to

$$H \coloneqq \varepsilon \begin{pmatrix} F & -A^\top \\ -A & G \end{pmatrix}^{-1}.$$

*Proof.* Using Equation (A.6) and then Equation (A.11), we have that

$$\begin{split} \frac{\mathrm{d}\pi^*}{\mathrm{d}x\,\mathrm{d}y}(x,y) &= \exp\left(f(x) + g(y) - \frac{|Qx - Ky|^2}{2\varepsilon}\right) \frac{\mathrm{d}\mu}{\mathrm{d}x}(x) \frac{\mathrm{d}\nu}{\mathrm{d}y}(y) \\ &\propto \exp\left(\mathcal{Q}(U + \Sigma^{-1})(x) + \mathcal{Q}(V + \Omega^{-1})(y) + \frac{1}{\varepsilon}\mathcal{Q}\left(\begin{smallmatrix}Q^\top Q & -A^\top\\ -A & K^\top K\end{smallmatrix}\right)(\begin{smallmatrix}x\\y\end{smallmatrix}\right)\right) \\ &\propto \exp\left(\frac{1}{\varepsilon}\mathcal{Q}\left(\begin{smallmatrix}\varepsilon U + \varepsilon\Sigma^{-1} + Q^\top Q & -A^\top\\ -A & \varepsilon V + \varepsilon\Omega^{-1} + K^\top K\end{smallmatrix}\right)(\begin{smallmatrix}x\\y\end{smallmatrix}\right) \\ &\propto \exp\left(\frac{1}{\varepsilon}\mathcal{Q}\left(\begin{smallmatrix}F & -A^\top\\ -A & G\end{smallmatrix}\right)(\begin{smallmatrix}x\\y\end{smallmatrix}\right)\right). \end{split}$$

The matrix

$$H^{-1} \coloneqq \frac{1}{\varepsilon} \begin{pmatrix} F & -A^\top \\ -A & G \end{pmatrix}$$

is positive definite, as  $G \succ 0$  and its Schur complement is equal to  $\varepsilon^{-1}(F - A^{\top}G^{-1}A) = \Sigma^{-1} \succ 0$  with Equation (A.10).

The following result paves the way to finding an explicit expression for H.

LEMMA A.7. Let F and G be two positive definite  $d \times d$  matrices satisfying

$$F = \varepsilon \Sigma^{-1} + A^{\top} G^{-1} A,$$
  
$$G = \varepsilon \Omega^{-1} + A F^{-1} A^{\top}.$$

Denote

(A.12) 
$$C_F \coloneqq \Omega A F^{-1} A^{\top},$$
$$C_G \coloneqq \Sigma A^{\top} G^{-1} A.$$

We have

$$C_F = \Omega^{1/2} \left( \Omega^{1/2} A \Sigma A^{\top} \Omega^{1/2} + \frac{\varepsilon^2}{4} I_d \right)^{1/2} \Omega^{-1/2} - \frac{\varepsilon}{2} I_d,$$
  
$$C_G = \Sigma^{1/2} \left( \Sigma^{1/2} A^{\top} \Omega A \Sigma^{1/2} + \frac{\varepsilon^2}{4} I_d \right)^{1/2} \Sigma^{-1/2} - \frac{\varepsilon}{2} I_d,$$

where the square roots are well-defined, as they are taken on positive definite matrices.

*Proof.* Let us prove the result for  $C_F$  only, by symmetry. A very similar computation as in [32] shows that

$$C_F^2 + \varepsilon C_F = \Omega A \Sigma A^\top.$$

Noticing that

$$C_F = \Omega^{1/2} \left( \Omega^{1/2} A F^{-1} A^{\top} \Omega^{1/2} \right) \Omega^{-1/2}$$

shows that all the eigenvalues of  $C_F$  are positive, and it is easy to check that the matrix  $\Omega^{1/2} \left( \Omega^{1/2} A \Sigma A^\top \Omega^{1/2} + \frac{\varepsilon^2}{4} I_d \right)^{1/2} \Omega^{-1/2} - \frac{\varepsilon}{2} I_d$  is the unique solution of the equation  $C^2 + \varepsilon C = \Omega A \Sigma A^\top$ 

with only positive eigenvalues.

Let us now use the block inversion formula on  $H^{-1}$  to obtain the desired expression for H.

$$H = \varepsilon \begin{pmatrix} F & -A^{\top} \\ -A & G \end{pmatrix}^{-1}$$
$$= \varepsilon \begin{pmatrix} (F - A^{\top}G^{-1}A)^{-1} & (F - A^{\top}G^{-1}A)^{-1}A^{\top}G^{-1} \\ (G - AF^{-1}A^{\top})^{-1}AF^{-1} & (G - AF^{-1}A^{\top})^{-1} \end{pmatrix}$$
$$= \begin{pmatrix} \Sigma & \Sigma A^{\top}G^{-1} \\ \Omega AF^{-1} & \Omega \end{pmatrix}$$

using Equation (A.10). We can conclude noticing that  $\Omega AF^{-1} = (\Sigma A^{\top}G^{-1})^{\top}$  as H is symmetric, so that

$$\Sigma A^{\top} G^{-1} = (C_F A^{-\top})^{\top}$$

by definition of  $C_F$ .

## Appendix B. Proofs of Section 3.

Let us start with the following useful result.

LEMMA B.1. Let R > 0. Let  $\mu$  be a probability measure on  $\mathbb{R}^d$  supported in  $B_R$ . Denote  $\alpha \in \mathbb{R}^d$  the expectation of  $\mu$ . Then

$$\|\operatorname{Var} \mu\|_2 \le R^2,$$

where

$$\operatorname{Var} \mu \coloneqq \mathbb{E}_{X \sim \mu} (X - \alpha) (X - \alpha)^{\top}$$

*Proof.* Let X be a random variable distributed according to  $\mu$ . Using the triangle inequality, we have:

$$\begin{split} \left\| \mathbb{E} (X - \alpha) (X - \alpha)^{\top} \right\|_{2} &\leq \mathbb{E} \left\| (X - \alpha) (X - \alpha)^{\top} \right\|_{2} \\ &= \mathbb{E} \left\| X - \alpha \right\|^{2} \\ &= \mathbb{E} \left\| X \right\|^{2} - \left\| \alpha \right\|^{2} \\ &\leq \mathbb{E} \left\| X \right\|^{2}. \end{split}$$

Now let us pick any  $x \in B_r$  such that |x| = r. We have

$$\mathbb{E}(X-x)^{\top}(X+x) \le 0,$$

as the angle between the vectors X - x and X + x is at least  $\pi/2$  for X in  $B_r$ . By expanding this relationship we get

$$E ||X||^{2} - ||x||^{2} = E ||X||^{2} - R^{2} \le 0,$$

which yields the result.

**B.1. Estimates With Constant Parameters.** We gather in this section the estimates on the velocity field  $\Gamma_{\mu}$  when Q, K, V are constant, for each type of unmasked self-attention. Notations are the same as in Section 2. We start with Softmax self-attention.

LEMMA B.2 (Estimates for Softmax self-attention). Let  $p \ge 1$  and R > 0. Let  $\mu$ and  $\nu$  be two probability measures supported in  $B_R$ . We have the following estimates.

(i) 
$$\sup_{x \in \mathbb{R}^d} |\Gamma_{\mu}^{(SM)}| \le ||V||_2 R$$
,  
(ii)  $\sup_{x \in \mathbb{R}^d} ||D_x \Gamma_{\mu}^{(SM)}||_2 \le ||V||_2 ||A||_2 R^2$ ,  
(iii)  $|\Gamma_{\mu}^{(SM)}(x) - \Gamma_{\nu}^{(SM)}(x)| \le ||V||_2 (1 + 2 ||A||_2 R ||x|) e^{2||A||_2 R ||x||} W_p(\mu, \nu)$ .

*Proof.* Equation (i) is straightforward. Equation (ii) relies on the following remark:

$$D_x \Gamma_{\mu} = V \operatorname{Var} \left( \frac{e^{Ax \cdot y}}{\int e^{Ax \cdot z} \, \mathrm{d}\mu(z)} \, \mathrm{d}\mu(y) \right) A$$

As the probability measure  $\frac{e^{Ax \cdot y}}{\int e^{Ax \cdot z} d\mu(z)} d\mu(y)$  is supported in  $B_R$ , its variance is bounded by  $R^2$ , according to Lemma B.1, which proves (ii). Finally, (iii) derives from the proof of Lemma 6.5 in [28].

We obtain the following L2 estimates with the same method as for Softmax selfattention.

LEMMA B.3 (Estimates for L2 self-attention). Let  $p \ge 1$  and R > 0. Let  $\mu$  and  $\nu$  be two probability measures supported in  $B_R$ . We have the following estimates.

(i) 
$$\sup_{x \in \mathbb{R}^d} |\Gamma_{\mu}^{(L2)}| \leq ||V||_2 R,$$
  
(ii)  $\sup_{x \in \mathbb{R}^d} ||D_x \Gamma_{\mu}^{(L2)}||_2 \leq ||V||_2 ||A||_2 R^2,$   
(iii)  $|\Gamma_{\mu}^{(L2)}(x) - \Gamma_{\nu}^{(L2)}(x)| \leq ||V||_2 (1 + 4R(||A||_2 |x| + ||K^{\top}K||_2 R)) \times e^{(||Q||_2 |x| + ||K||_2 R)^2} W_p(\mu, \nu).$ 

The following multi-head attention estimates are straightforward.

LEMMA B.4 (Estimates for multi-head self-attention). Let  $p \ge 1$  and R > 0. Let  $\mu$  and  $\nu$  be two probability measures supported in  $B_R$ . We have the following estimates.

 $\begin{array}{l} (i) \ \sup_{x \in \mathbb{R}^{d}} \left| \Gamma_{\mu}^{(\mathrm{MH})} \right| \leq \sum_{h=1}^{H} \left\| W^{(h)} V^{(h)} \right\|_{2} R, \\ (ii) \ \sup_{x \in \mathbb{R}^{d}} \left\| D_{x} \Gamma_{\mu}^{(\mathrm{MH})} \right\|_{2} \leq \sum_{h=1}^{H} \left\| W^{(h)} V^{(h)} \right\|_{2} \left\| A^{(h)} \right\|_{2} R^{2}, \\ (iii) \ \left| \Gamma_{\mu}(x)^{(\mathrm{MH})} - \Gamma_{\nu}(x)^{(\mathrm{MH})} \right| \leq \sum_{h=1}^{H} \left\| W^{(h)} V^{(h)} \right\|_{2} (1 + 2 \left\| A^{(h)} \right\|_{2} R \left| x \right|) \\ \times e^{2 \left\| A^{(h)} \right\|_{2} R \left| x \right|} W_{p}(\mu, \nu). \end{array}$ 

The estimates for Sinkhorn self-attention are more involved, as they leverage some background on entropic optimal transport (see Section A.1). We only state them for  $\varepsilon = 1$  for simplicity, as the  $\varepsilon$  can be absorbed in the matrices Q, K and V. Let us start by proving estimates on  $\kappa_{\mu}^{\infty}$ . All the notations are defined in Section A.1.

LEMMA B.5. Let  $\kappa^0: (x,y) \in (\mathbb{R}^d)^2 \mapsto e^{-\frac{1}{2}|Qx-Ky|^2}$  and  $\kappa^{\infty}_{\mu}$  defined by the iterates (2.4). Then

(i)  $\sup_{x,y\in B_R} \kappa_{\mu}^{\infty}(x,y) \leq e^{(\|Q\|_2 + \|K\|_2)^2 R^2},$ (ii)  $\sup_{x,y\in B_R} \|D_y \kappa_{\mu}^{\infty}(x,y)\|_2 \leq 2(\|Q\|_2 + \|K\|_2)^2 R e^{(\|Q\|_2 + \|K\|_2)^2 R^2}.$ *Proof.* Let  $x, y \in B_R$ . We have with Lemma A.1 that

$$\kappa_{\mu}^{\infty}(x,y) = e^{f(x) + g(y) - |Qx - Ky|^2} \le e^{(\|Q\|_2 + \|K\|_2)^2 R^2},$$

which proves (i). For (ii), differentiating  $\kappa^{\infty}_{\mu}$  with respect to the second variable gives

$$D_y \kappa_\mu^\infty(x, \cdot) = \kappa_\mu^\infty(x, y) (D_y g - D_y c(x, \cdot)) = \kappa_\mu^\infty(x, y) (D_y g - (Ky - Qx)^\top K)$$

so that

$$\left\| D_{y} \kappa_{\mu}^{\infty}(x, \cdot) \right\|_{2} \leq \left( \|Q\|_{2} + \|K\|_{2} + \left( \|Q\|_{2} + \|K\|_{2} \right)^{3/2} \right) Re^{\left( \|Q\|_{2} + \|K\|_{2} \right)^{2} R^{2}}$$

with Lemma A.1 and the first estimate on  $\kappa_{\mu}^{\infty}.$ 

We can then state estimates on  $\Gamma$ , as for the other types of attention.

LEMMA B.6 (Estimates for Sinkhorn self-attention). Let R > 0 and  $p \ge 1$ . Let  $\mu$ and  $\nu$  be two probability measures supported in  $B_R$ . We have the following estimates for all  $x \in \mathbb{R}^d$ .

(i) 
$$\sup_{x \in \mathbb{R}^d} \left\| \Gamma_{\mu}^{(\text{sink})}(x) \right\| \le \|V\|_2 R,$$
  
(ii)  $\sup_{x \in \mathbb{R}^d} \left\| D_x \Gamma_{\mu}^{(\text{sink})} \right\|_2 \le \|V\|_2 \|A\|_2 R^2,$   
(iii)  $\left| \Gamma_{\mu}^{(\text{sink})}(x) - \Gamma_{\nu}^{(\text{sink})}(x) \right| \le \|V\|_2 \left( 1 + c(R, Q, K)R + 2(\|Q\|_2 + \|K\|_2)^2 R |x| \right) \times e^{(\|Q\|_2 + \|K\|_2)^2 R |x|} W_2(\mu, \nu)$ 

for some function c(R, Q, K) > 0.

Proof. The first estimate is straightforward. To prove the second estimate, notice that

$$D_x f = -\int e^{g(y) - c(x,y)} (Ky - Qx)^\top Q \,\mathrm{d}\mu(y) / e^{-f(x)}$$
$$= \int \kappa_\mu^\infty(x,y) (Qx - Ky)^\top Q \,\mathrm{d}\mu(y)$$
$$= x^\top Q^\top Q - \int \kappa_\mu^\infty(x,y) y^\top A \,\mathrm{d}\mu(y).$$

Then

$$D_x \kappa^{\infty}_{\mu}(\cdot, y) = \kappa^{\infty}_{\mu}(x, y) (D_x f + (Ky - Qx)^{\top} Q)$$
  
=  $\kappa^{\infty}_{\mu}(x, y) \left( y^{\top} - \int \kappa^{\infty}_{\mu}(x, y') y'^{\top} d\mu(y) \right) A,$ 

and finally

$$D_x \Gamma_\mu = V \left( \int y y^\top \kappa_\mu^\infty(x, y) \, \mathrm{d}\mu(y) - \int y \kappa_\mu^\infty(x, y) \, \mathrm{d}\mu(y) \int y^\top \kappa_\mu^\infty(x, y) \, \mathrm{d}\mu(y) \right) A$$
  
=  $V \operatorname{Var}(\kappa_\mu^\infty(x, y) \, \mathrm{d}\mu(y)) A$ ,

which allows us to conclude with Lemma B.1. The last estimate builds on Lemma A.2. Let us first derive a bound on  $\sup_{x,y\in B_R} |\kappa^{\infty}_{\mu}(x,y) - \kappa^{\infty}_{\nu}(x,y)|$ . For all  $x, y \in B_R$ , we have

$$\begin{aligned} \left|\kappa_{\mu}^{\infty}(x,y) - \kappa_{\nu}^{\infty}(x,y)\right| &\leq \left|e^{f^{\mu}(x) + g^{\mu}(y) - \frac{1}{2}|Qx - Ky|^{2}} - e^{f^{\nu}(x) + g^{\nu}(y) - \frac{1}{2}|Qx - Ky|^{2}} \right| \\ &\leq e^{-\frac{1}{2}|Qx - Ky|^{2}} e^{\max(f^{\mu}(x) + g^{\mu}(y), f^{\nu}(x) + g^{\nu}(y))} \\ &\times \left|f^{\mu}(x) - f^{\nu}(x) + g^{\mu}(y) - g^{\nu}(y)\right|. \end{aligned}$$

We can choose  $f^{\mu}$  and  $g^{\mu}$  such that

$$\|f^{\mu} - f^{\nu}\|_{\infty} + \|g^{\mu} - g^{\nu}\|_{\infty} = \inf_{\eta \in \mathbb{R}} \|f^{\mu} - f^{\nu} - \eta\|_{\infty} + \|g^{\mu} - g^{\nu} + \eta\|_{\infty},$$

in order to have

$$|f^{\mu}(x) - f^{\nu}(x) + g^{\mu}(y) - g^{\nu}(y)| \le ||f^{\mu} - f^{\nu}||_{\infty} + ||g^{\mu} - g^{\nu}||_{\infty} \le C_{\text{stab}}(R)W_{2}(\mu, \nu)$$

with Lemma A.2. Finally, we obtain with a variant of Lemma A.1

$$\left|\kappa_{\mu}^{\infty}(x,y) - \kappa_{\nu}^{\infty}(x,y)\right| \le C(R)e^{(\|Q\|_{2} + \|K\|_{2})^{2}R|x|}W_{2}(\mu,\nu),$$

where we upper-bounded  $e^{-\frac{1}{2}|Qx-Ky|^2}$  by 1 and  $\max(f^{\mu}(x) + g^{\mu}(y), f^{\nu}(x) + g^{\nu}(y))$  by  $(\|Q\|_2 + \|K\|_2)^2 R |x|$ . Note that this is a very rough bound, but we do not need more as we just want to prove the continuity of  $\Gamma$ . Then, for all  $x \in \mathbb{R}^d$  we have

$$\begin{split} |\Gamma_{\mu}(x) - \Gamma_{\nu}(x)| &\leq \|V\|_{2} \left( \left| \int y \kappa_{\mu}^{\infty}(x, y) \, d\mu(y) - \int y \kappa_{\nu}^{\infty}(x, y) \, d\mu(y) \right| \\ &+ \left| \int y \kappa_{\nu}^{\infty}(x, y) \, d\mu(y) - \int y \kappa_{\nu}^{\infty}(x, y) \, d\nu(y) \right| \right) \\ &\leq \|V\|_{2} \left( R \sup_{y \in B_{R}} \left| \kappa_{\mu}^{\infty}(x, y) - \kappa_{\nu}^{\infty}(x, y) \right| \\ &+ \sup_{y \in B_{R}} \|D_{y}(y \mapsto y \kappa_{\nu}^{\infty}(x, y))\|_{2} W_{2}(\mu, \nu) \right) \\ &\leq \|V\|_{2} e^{(\|Q\|_{2} + \|K\|_{2})^{2} R|x|} \left( 1 + C_{\text{stab}}(R) R + \\ &\quad (\|Q\|_{2} + \|K\|_{2} + (\|Q\|_{2} + \|K\|_{2})^{3/2}) R\, |x| \right) W_{2}(\mu, \nu). \end{split}$$

Finally, we have the following estimates for Sigmoid attention.

LEMMA B.7 (Estimates for Sigmoid self-attention). Let R > 0 and  $p \ge 1$ . Let  $\mu$ and  $\nu$  be two probability measures supported in  $B_R$ . We have the following estimates for all  $x \in \mathbb{R}^d$ .

 $\begin{array}{l} (i) \; \sup_{x \in \mathbb{R}^d} \; \left| \Gamma_{\mu}^{(\sigma)}(x) \right| \leq \|V\|_2 \, R, \\ (ii) \; \sup_{x \in \mathbb{R}^d} \; \left\| D_x \Gamma_{\mu}^{(\sigma)} \right\|_2 \leq \frac{1}{4} \, \|V\|_2 \, \|A\|_2 \, R^2, \\ (iii) \; \left| \Gamma_{\mu}^{(\sigma)}(x) - \Gamma_{\nu}^{(\sigma)}(x) \right| \leq \|V\|_2 \, (1 + \|A\|_2 \, R^2) W_2(\mu, \nu). \end{array}$ 

Proof. Equation (i) is straightforward. For Equation (ii), we calculate

$$D_x \Gamma_{\mu}^{(\sigma)} = \int V y y^{\top} A \, \sigma (Ax \cdot y) (1 - \sigma (Ax \cdot y)) \, \mathrm{d}\mu(y),$$

and the bound follows, noticing that  $|\sigma(Ax \cdot y)(1 - s(Ax \cdot y))| \le 1/4$  as  $|\sigma(Ax \cdot y)| \in [0, 1]$ . Finally, we have

$$|\Gamma_{\mu}(x) - \Gamma_{\nu}(x)| = \left| \int V y \sigma(Ax \cdot y) \,\mathrm{d}(\mu - \nu)(y) \right|,$$

and

$$D_y(y \mapsto Vy\sigma(Ax \cdot y)) = \sigma(Ax \cdot y)(I_d + (1 - \sigma(Ax \cdot y))yy^{\top}A),$$

so that the Lipschitz constant of  $y \mapsto Vy\sigma(Ax \cdot y)$  is bounded by  $||V||_2 (1+||A||_2 R^2)$  if  $\mu$  and  $\nu$  are supported in  $B_R$ . We conclude for Equation (iii) with the duality formula for  $W_1$ , and with the inequality  $W_1(\mu,\nu) \leq W_2(\mu,\nu)$ , as  $\mu$  and  $\nu$  are compactly supported.

**B.2. Technical Lemmas.** We list here the technical lemmas used in the proof of Theorem 3.1.

LEMMA B.8. Let  $\mu_0 \in \mathcal{P}_c(\mathbb{R}^d)$  be an initial condition, and assume that the associated problem (3.2) has an equi-compactly supported solution  $\mu$  on the time interval [0,T]. For all  $0 \leq t \leq T$ , denote R(t) > 0 the smallest radius such that

$$\operatorname{Supp} \mu(\tau) \subset B_{R(t)},$$

for all  $0 \le \tau \le t$ . Then, it holds

$$R(t) \le e^{\|V\|_2 t} R(0)$$

for all  $0 \leq t \leq T$ .

*Proof.* Let  $x \in \mathbb{R}^d$  belong to the support of  $\mu_0$ . For any  $t \in [0, T]$  we have by Lemma B.2:

$$|x - \phi_t(\mu)(x)| \le \int_0^t \left| \Gamma_{\mu(s)}(\phi_s(\mu)(x)) \right| \, \mathrm{d}s \le \|V\|_2 \int_0^t R(s) \, \mathrm{d}s.$$

By taking a supremum over x in the support of  $\mu_0$  we get

$$R(t) - R(0) \le ||V||_2 \int_0^t R(s) \,\mathrm{d}s$$

and Grönwall's inequality implies the claim.

Lemma B.8 allows us to prove that  $\mathcal{F}$  preserves X.

LEMMA B.9. We have  $\mathcal{F}(X) \subset X$ .

*Proof.* The proof is the same as for Lemma B.8, replacing  $\mu_0$  with  $\bar{\mu}_0$ .  $\Box$ We have the following estimate on  $\mathcal{F}$ .

 $I = \frac{1}{2} \left( \frac{1}{2} \right) \left($ 

LEMMA B.10. Let T > 0 and  $\bar{\mu}_0 \in \mathcal{P}_c(\mathbb{R}^d)$ . Denote  $R_0$  the smallest radius such that  $\operatorname{Supp} \bar{\mu}_0 \subset B_{R_0}$ , and set

$$R(t) \coloneqq e^{\|V\|_2 t} R_0$$

for all  $0 \leq t \leq T$ . Consider X the set of curves  $\mu \in \mathcal{C}_{co}([0,T], \mathcal{P}_{c}(\mathbb{R}^{d}))$  satisfying

$$\operatorname{Supp} \mu(t) \subset B_{R(t)}$$

for all  $0 \leq t \leq T$ , and define  $\mathcal{F} \colon X \to X$  by

$$\mathcal{F}(\mu) \coloneqq \phi_t(\mu)_{\sharp} \bar{\mu}_0.$$

Then, for all  $\mu, \nu \in X$ , we have

$$\mathcal{D}_{p,T}(\mathcal{F}(\mu), \mathcal{F}(\nu)) \le f(T, R_0) \mathcal{D}_{p,T}(\mu, \nu)$$

with f a positive function such that  $f(T, R_0) \to 0$  when  $T \to 0^+$ .

*Proof.* Let us first prove that

(B.1) 
$$\|\phi_t(\mu) - \phi_t(\nu)\|_{L^{\infty}(\operatorname{Supp}\bar{\mu}_0)} \le C_1(R(T)) \int_0^t e^{C_2 R(T)^2(t-s)} W_p(\mu(s),\nu(s)) \,\mathrm{d}s$$

for all  $0 \le t \le T$ , with

$$C_1(R(T)) \coloneqq c(R(t), R(t))$$

(see (iii)) and

 $C_2 \coloneqq \|V\|_2 \|A\|_2.$ 

For all  $x \in \text{Supp } \bar{\mu}_0$  and  $t \in [0, T]$ , it holds:

$$\begin{aligned} |\phi_t(\mu)(x) - \phi_t(\nu)(x)| &\leq \int_0^t |\Gamma_\mu(s, \phi_s(\mu)(x)) - \Gamma_\nu(s, \phi_s(\nu)(x))| \, \mathrm{d}s \\ &\leq \int_0^t |\Gamma_\mu(s, \phi_s(\mu)(x)) - \Gamma_\mu(s, \phi_s(\nu)(x))| \, \mathrm{d}s \\ &+ \int_0^t |\Gamma_\mu(s, \phi_s(\nu)(x)) - \Gamma_\nu(s, \phi_s(\nu)(x))| \, \mathrm{d}s \\ &\leq C_2 R(T)^2 \int_0^t |\phi_s(\mu)(x) - \phi_s(\nu)(x)| \, \mathrm{d}s \\ &+ C_1(R(T)) \int_0^t W_p(\mu(s), \nu(s)) \, \mathrm{d}s \end{aligned}$$

where we used Equations (i), (ii), (iii) for the last inequality, noticing with Lemma B.8 that  $|\phi_s(\nu)(x)| \leq R(T)$  for all  $s \in [0, T]$ . An application of Grönwall's inequality leads to Equation (B.1). Then, by a standard result in optimal transport:

$$\begin{split} W_{p}(\mathcal{F}(\mu)(t), \mathcal{F}(\nu)(t)) &= W_{p}(\phi_{t}(\mu)_{\sharp}\bar{\mu}_{0}, \phi_{t}(\nu)_{\sharp}\bar{\mu}_{0}) \\ &\leq \|\phi_{t}(\mu) - \phi_{t}(\nu)\|_{L^{\infty}(\mathrm{Supp}\,\bar{\mu}_{0})} \\ &\leq C_{1}(R(T)) \left(\int_{0}^{t} e^{C_{2}R(T)^{2}(t-s)} \,\mathrm{d}s\right) \mathcal{D}_{p,T}(\mu, \nu) \\ &= \frac{C_{1}(R(T))}{C_{2}R(T)^{2}} \left(e^{C_{2}R(T)^{2}t} - 1\right) \mathcal{D}_{p,T}(\mu, \nu), \end{split}$$

for all  $0 \le t \le T$ , thanks to Equation (B.1). Taking the supremum over  $t \in [0, T]$  gives the desired estimate with

$$f(T, R_0) \coloneqq \frac{C_1(R(T))}{C_2 R(T)^2} \left( e^{C_2 R(T)^2 T} - 1 \right)$$

We can see that  $f(T, R_0) \to 0$  when  $T \to 0^+$ , which concludes the proof.

We also need the following result to apply a Banach fixed point argument. LEMMA B.11. X equipped with the distance  $\mathcal{D}_{p,T}$  is a complete metric space.

Proof. Let  $(\mu^n)_{n\in\mathbb{N}}$  be a Cauchy sequence taking values in X. Seeing X as a subspace of the complete metric space  $\mathcal{C}([0,T], \mathcal{P}_p(\mathbb{R}^d))$ , we know that  $\mu^n$  converges to some  $\mu^* \in \mathcal{C}([0,T], \mathcal{P}_p(\mathbb{R}^d))$ . Assume by contradiction that  $\mu^*$  does not belong to X. Then, there exists some  $t \in [0,T]$  such that  $\operatorname{Supp} \mu^*(t)$  is not included in  $B_{R(t)}$ , with R(t) defined in Lemma B.10. Let  $x \in \operatorname{Supp} \mu^*(t) \setminus B_{R(t)}$ . There is a small closed ball  $\mathcal{B}$  centered at x, containing a neighborhood of x, and included in the complement of  $B_{R(t)}$ , such that  $\mu^*(t)(\mathcal{B}) > 0$ . Let  $\delta > 0$  such that for all  $x' \in \mathcal{B}$  and  $y \in B_{R(t)}$ , it holds  $|x' - y| \geq \delta$ . Then, for all  $n \in \mathbb{N}$ , we have

$$W_p(\mu^*(t), \mu^n(t))^p \ge \delta^p \mu^*(t)(\mathcal{B}),$$

which contradicts the fact that  $\mathcal{D}_{p,T}(\mu^n, \mu^*) \to 0$  when  $n \to +\infty$ .

**B.3. Stability estimates.** As for the stability estimate, let  $t \ge 0$ . We have

$$\begin{split} W_{p}(\mu(t),\nu(t)) &= W_{p}(\phi_{t}(\mu)_{\sharp}\mu_{0},\phi_{t}(\nu)_{\sharp}\nu_{0}) \\ &\leq W_{p}(\phi_{t}(\mu)_{\sharp}\mu_{0},\phi_{t}(\mu)_{\sharp}\nu_{0}) \\ &+ W_{p}(\phi_{t}(\mu)_{\sharp}\nu_{0},\phi_{t}(\nu)_{\sharp}\nu_{0}) \\ &\leq \operatorname{Lip}(x\mapsto\phi_{t}(\mu)(x))W_{p}(\mu_{0},\nu_{0}) + \|\phi_{t}(\mu) - \phi_{t}(\nu)\|_{L^{\infty}(\operatorname{Supp}\nu_{0})} \,. \end{split}$$

Let us bound the Lipschitz constant  $\operatorname{Lip}(x \mapsto \phi_t(\mu)(x))$ . For all  $x, y \in \mathbb{R}^d$ :

$$\begin{aligned} |\phi_t(\mu)(x) - \phi_t(\mu)(y)| &\le |x - y| + \int_0^t |\Gamma_\mu(s, \phi_s(\mu)(x)) - \Gamma_\mu(s, \phi_s(\mu)(y))| \, \mathrm{d}s \\ &\le |x - y| + C_2 R(T)^2 \int_0^t |\phi_s(\mu)(x) - \phi_s(\mu)(y)| \, \mathrm{d}s \end{aligned}$$

by Lemma B.2, with  $C_2 = \|V\|_2 \|A\|_2$ . Then, Grönwall's inequality entails that

$$|\phi_t(\mu)(x) - \phi_t(\mu)(y)| \le e^{C_2 R(T)^2 t} |x - y|.$$

Plugging this into our previous bound for  $W_p(\mu(t), \nu(t))$ , together with Equation (B.1), we get:

$$W_p(\mu(t),\nu(t)) \le e^{C_2 R(T)^2 t} W_p(\mu_0,\nu_0) + C_1(R(T)) \int_0^t e^{C_2 R(T)^2 (t-s)} W_p(\mu(s),\nu(s)) \,\mathrm{d}s.$$

We can then apply Grönwall's inequality to  $t \mapsto e^{-C_2 R(T)^2 t} W_p(\mu(t), \nu(t))$  to obtain

$$W_p(\mu(t),\nu(t)) \le e^{\left(C_1(R(T))+C_2R(T)^2\right)t} W_p(\mu_0,\nu_0)$$

which concludes the proof, recalling that  $R(T) = e^{\|V\|_2 T} R_0$ .

**B.4. Estimates With Varying Parameters.** The proof of Theorem 3.1 with time-depending parameters is very similar to the case of constant parameters, with a few additional computations. We only write it for Softmax self-attention, but the same ideas apply to the other types of attention. For any equi-compactly supported curve  $\mu \in C_{co}([0, T], \mathcal{P}_c(\mathbb{R}^d))$ , consider again the Cauchy problem

$$\begin{cases} \dot{r}(\tau) = \Gamma_{\mu}(\tau, r(\tau)) \text{ for } 0 \le \tau \le t \\ r(0) = x \in \mathbb{R}^d \end{cases}$$

.

To be able to define the associated flow, we need the same estimates as before: time continuity, local Lipschitz continuity uniformly on [0, T], and at most linear spatial growth of the vector field. We just check time continuity, the other two estimates being straightforward. In what follows, we will write  $\Gamma_{\mu}$  as  $\Gamma_{\mu,A,V}$ , to stress the dependency in A and V. For any measures  $\mu_1, \mu_2$  supported in  $B_R$ , and any matrices  $A_1, A_2, V_1, V_2 \in \mathbb{R}^{d \times d}$ , we have

$$\begin{split} |\Gamma_{\mu_1,A_1,V_1}(x) - \Gamma_{\mu_2,A_2,V_2}(x)| &\leq |\Gamma_{\mu_1,A_1,V_1}(x) - \Gamma_{\mu_2,A_1,V_1}(x)| \\ &+ |\Gamma_{\mu_2,A_1,V_1}(x) - \Gamma_{\mu_2,A_2,V_1}(x)| \\ &+ |\Gamma_{\mu_2,A_2,V_1}(x) - \Gamma_{\mu_2,A_2,V_2}(x)| \\ &\leq \|V_1\|_2 \left(1 + 2 \|A_1\|_2 R \|x\|) e^{2\|A_1\|_2 R \|x\|} W_p(\mu_1,\mu_2) \\ &+ 2 \|V_1\|_2 \|A_1 - A_2\|_2 R^2 \|x\| e^{(\|A_1\|_2 + \|A_2\|_2)R \|x\|} \\ &+ \|V_1 - V_2\|_2 R, \end{split}$$

where we used Lemma B.2 to bound the first term, the bound for the second term relies on the following inequalities:

$$\left| e^{x^{\top} A_1^{\top} y} - e^{x^{\top} A_2^{\top} y} \right| \le \left\| A_1 - A_2 \right\|_2 R \left| x \right| e^{\max(\|A_1\|_2, \|A_2\|_2) R |x|},$$

that derives from the mean-value inequality, and

$$\left|e^{x^{\top}A_i^{\top}y}\right| \ge e^{\|A_i\|_2 R|x|}$$

for i = 1, 2, and the last bound is straightforward. We can therefore prove the continuity of the map  $(t, x) \mapsto \Gamma_{\mu(t)}(x)$  as before, and apply Cauchy-Lipschitz theorem to define the associated flow  $\phi_t(\mu)(x)$ .

Set a compactly supported initial condition  $\bar{\mu}_0$  supported in  $B_{R_0}$  with  $R_0$  minimal, define X the set of curves  $\mu \in \mathcal{C}_{co}([0,T], \mathcal{P}_c(\mathbb{R}^d))$  such that for all  $t \in [0,T]$ , we have

Supp 
$$\mu(t) \subset \overline{B}\left(0, e^{\int_0^t \|V(s)\|_2 \,\mathrm{d}s} R_0\right)$$
,

and consider the map

$$\mathcal{F}\colon \mu\in X\mapsto \phi_t(\mu)_{\sharp}\bar{\mu}_0$$

LEMMA B.12. We have  $\mathcal{F}(X) \subset X$ .

*Proof.* Let  $\mu \in X$  and x in the support of  $\overline{\mu}_0$ . For all  $t \in [0, T]$ , denote R(t) the smallest radius such that  $\mu(s)$  is supported in  $B_{R(t)}$  for all  $s \in [0, t]$ . Then

$$\begin{aligned} |x - \phi_t(\mu)(x)| &\leq \int_0^t |\Gamma_\mu(s, \phi_s(\mu)(x))| \, \mathrm{d}s \\ &\leq \int_0^t \|V(s)\|_2 \, R(s) \, \mathrm{d}s, \end{aligned}$$

using Lemma B.2. Taking the supremum over x in the support of  $\bar{\mu}_0$ , we get

$$R(t) - R(0) \le \int_0^t \|V(s)\|_2 R(s) \,\mathrm{d}s$$

Grönwall's inequality then proves the claim.

A computation that is similar to the proof of Lemma B.10 then gives the following result.

LEMMA B.13. Let T > 0 and  $\bar{\mu}_0 \in \mathcal{P}_c(\mathbb{R}^d)$ . Denote  $R_0$  the smallest radius such that  $\operatorname{Supp} \bar{\mu}_0 \subset B_{R_0}$ , and set

$$R(t) \coloneqq e^{\int_0^t \|V(s)\|_2 \,\mathrm{d}s} R_0$$

for all  $0 \leq t \leq T$ . Consider X the set of curves  $\mu \in \mathcal{C}_{co}([0,T],\mathcal{P}_{c}(\mathbb{R}^{d}))$  satisfying

$$\operatorname{Supp} \mu(t) \subset B_{R(t)}$$

for all  $0 \leq t \leq T$ , and define  $\mathcal{F} \colon X \to X$  by

$$\mathcal{F}(\mu) \coloneqq \phi_t(\mu)_{\sharp} \bar{\mu}_0.$$

Then, for all  $\mu, \nu \in X$ , we have

$$\mathcal{D}_{p,T}(\mathcal{F}(\mu), \mathcal{F}(\nu)) \le f(T, R_0)\mathcal{D}_{p,T}(\mu, \nu)$$

with f a positive function such that  $f(T, R_0) \to 0$  when  $T \to 0^+$ .

As X is a complete metric space, which derives from the proof of Lemma B.11, we can follow the exact same steps as in the previous subsection, and conclude that problem (3.2) is well-posed even for continuous time-depending parameters A and V.

**B.5.** Masked Attention and Wasserstein Distance. The use of the conditional Wasserstein distance for the masked Transformer PDE circumvents the difficulty of proving an estimate of the form

$$\left|\Gamma_{\bar{\mu}}^{(m)}(s,x) - \Gamma_{\bar{\nu}}^{(m)}(s,x)\right| \le C(x)W_2(\bar{\mu},\bar{\nu}).$$

This difficulty is suggested by the following negative result.

**PROPOSITION B.14.** The masked attention map

$$\Gamma \colon \bar{\mu} \in (\mathcal{P}([0,1] \times \mathbb{R}^d), W_2) \mapsto \Gamma_{\bar{\mu}} \in (([0,1] \times \mathbb{R}^d)^{[0,1] \times \mathbb{R}^d}, \|\cdot\|_{\infty})$$

defined in Section 2.6 is not continuous for the Wasserstein 2 distance  $W_2$ .

*Proof.* Let us do the proof for d = 1 for simplicity. For any  $x \in \mathbb{R}$ , denote  $\mathcal{L}_x$  the one-dimensional Lebesgue measure (renormalized to a probability measure) on the segment  $[(0, x), (1, x)] := \{(t, x) : t \in [0, 1]\}$ . Let  $t_0 := 1/2$ . For all  $n \ge 1$ , let  $\mu_n$  and  $\nu_n$  be two probability measures supported in  $[(0, 0), (1, 0)] \cup [(0, n^2), (1, n^2)]$ , defined by their densities w.r.t.  $\mathcal{L}_0$  and  $\mathcal{L}_{n^2}$  as

$$\frac{\mathrm{d}\mu_n}{\mathrm{d}\mathcal{L}_0} \coloneqq \mathbf{1}_{t \notin [t_0, t_0 + 1/n]} \quad \text{and} \quad \frac{\mathrm{d}\mu_n}{\mathrm{d}\mathcal{L}_{n^2}} \coloneqq \mathbf{1}_{t \in [t_0, t_0 + 1/n]}$$

and

$$\frac{\mathrm{d}\nu_n}{\mathrm{d}\mathcal{L}_0} \coloneqq \mathbf{1}_{t \notin [t_0 + 1/2 - 1/n, t_0 + 1/2]} \quad \text{and} \quad \frac{\mathrm{d}\nu_n}{\mathrm{d}\mathcal{L}_{n^2}} \coloneqq \mathbf{1}_{t \in [t_0 + 1/2 - 1/n, t_0 + 1/2]}.$$

One has immediately

$$W_2(\mu_n,\nu_n) \le \sqrt{\frac{1}{4n}},$$

while

$$\Gamma_{\bar{\nu}_n}(t_0 + 1/4, 0) = 0$$

and

$$\Gamma_{\bar{\mu}_n}(t_0 + 1/4, 0) = \frac{n}{t_0 + 1/4} \to_{n \to +\infty} +\infty,$$

which proves the claim.

#### Appendix C. Proofs and Experiments of Section 4.

#### C.1. Proofs.

Proof of Proposition 4.2. Gaussian measures stay Gaussian along the dynamics, as the pushforward of a Gaussian measure by an affine function is still Gaussian. Let us then derive the ODEs on the expectation  $\alpha$  and the covariance  $\Sigma$ . For all  $x \in \mathbb{R}^d$  and  $1 \leq i \leq d$ , denote  $x_i$  the *i*-th coordinate of x. Set two integers  $1 \leq i, j \leq d$ . For all  $t \in [0, T]$ , we have

$$\Sigma(t)_{i,j} = \int (x_i - \alpha(t)_i)(x_j - \alpha(t)_j).$$

In the rest of the proof, we omit the dependency in t of  $\alpha$  and  $\Sigma$  and write  $\Gamma_{\mu}$  for  $\Gamma_{\mu}^{(SM)}$ . Recall that  $\mu$  satisfies

$$\partial_t \mu + \nabla_x \cdot (\mu \Gamma_\mu) = 0.$$

By multiplying this equation by  $(x_i - \alpha_i)(x_j - \alpha_j)$  and then integrating by parts, we get

$$\dot{\Sigma}_{i,j} = \int \sum_{k=1}^{d} \partial_{x_k} \left( (x_i - \alpha_i)(x_j - \alpha_j) \right) \Gamma_{\mu(t)}(x)_k \, \mathrm{d}\mu(t)(x)$$
$$= \int \left( (x_j - \alpha_j) \Gamma_{\mu(t)}(x)_i + (x_i - \alpha_i) \Gamma_{\mu(t)}(x)_j \right) \, \mathrm{d}\mu(t)(x).$$

Replacing  $\Gamma_{\mu(t)}$  with its expression, given in Lemma 4.1, gives

$$\dot{\Sigma}_{i,j} = \int (x_j - \alpha_j) (V\alpha + V\Sigma Ax)_i \, \mathrm{d}\mu(t)(x) + \int (x_i - \alpha_i) (V\alpha + V\Sigma Ax)_j \, \mathrm{d}\mu(t)(x).$$

We compute

$$\int (x_j - \alpha_j) (V\alpha + V\Sigma Ax)_i \, \mathrm{d}\mu(t)(x) = \int (x_j - \alpha_j) \Big( (V\alpha)_i \\ + \sum_{k=1}^d (V\Sigma A)_{i,k} x_k \Big) \, \mathrm{d}\mu(t)(x)$$
$$= \sum_{k=1}^d (V\Sigma A)_{i,k} \int (x_j - \alpha_j) x_k \, \mathrm{d}\mu(t)(x)$$
$$= \sum_{k=1}^d (V\Sigma A)_{i,k} \Sigma_{j,k}$$
$$= (V\Sigma A\Sigma)_{i,j}.$$

Exchanging i and j we get

$$\int (x_i - \alpha_i) (V\alpha + V\Sigma Ax)_j \, \mathrm{d}\mu(t)(x) = (V\Sigma A\Sigma)_{j,i},$$

so that

$$\dot{\Sigma}_{i,j} = (V\Sigma A\Sigma + \Sigma A^{\top}\Sigma V^{\top})_{i,j}.$$

This gives the equation on the covariance. The same strategy, but this time multiplying the PDE by  $x_i - \alpha_i$  before integrating, gives the equation on  $\alpha$ .

Proof of Proposition 4.4. To check the closed form, let us derive an ODE on  $\Omega(t) := \Sigma(t)^{-1}$ . We have that V commutes with  $\Sigma_0$ , so V and  $\Sigma_0$  are co-diagonalizable and V commutes with  $\Omega_0 := \Sigma_0^{-1}$  as well. Using, that  $\dot{\Omega} = -\Omega \dot{\Sigma} \Omega$ , it is easy to check that  $\Omega(t)$  satisfies the following differential equation:

$$\dot{\Omega} = -\Omega V \Omega^{-1} A - A^{\top} \Omega^{-1} V^{\top} \Omega.$$

Thanks to our commutation assumptions, if  $\Omega(t)$  commutes with V, then  $\Omega(t + dt) = \Omega(t) - dt(VA + A^{\top}V^{\top})$  commutes with V as well. Therefore, if  $\Omega_0$  commutes with V, then the matrix  $\Omega$  satisfies the equation

$$\dot{\Omega} = -(VA + A^{\top}V^{\top}).$$

Hence, the solution reads  $\Omega(t) = \Omega_0 - t(VA + A^{\top}V^{\top})$ , which gives the expected formula for  $\Sigma$ .

• Assume first that  $VA + A^{\top}V^{\top}$  has at least one positive eigenvalue. For any symmetric matrix  $M \in \mathbb{R}^{d \times d}$ , denote  $\lambda_1(M) \geq \cdots \geq \lambda_d(M)$  its ordered eigenvalues. According to Weyl's inequality, we have

$$0 \le \lambda_d(\Omega(t)) \le \lambda_1(\Omega_0) - t\lambda_1(VA + A^\top V^\top),$$

so that  $\lambda_d(\Omega(t))$  has to be zero for  $t \geq \lambda_1(\Omega_0)/\lambda_1(VA + A^{\top}V^{\top}) \geq 0$ . Therefore, the matrix  $\Omega$  becomes non-invertible in finite time, and  $\Sigma = \Omega^{-1}$  blows up in finite time.

• Assume now that  $VA + A^{\top}V^{\top} \leq 0$ . Let  $1 \leq i \leq d$ . Again with Weyl's inequality, we have

(C.1) 
$$\lambda_i(\Omega) \ge \lambda_d(\Omega_0) - t\lambda_{d-i}(VA + A^\top V^\top) > 0,$$

so that  $\Omega$  is invertible for all times, and the equation on  $\Sigma = \Omega^{-1}$  has a global solution. Moreover, Equation (C.1) shows that if  $\lambda_{d-i}(VA + A^{\top}V^{\top}) < 0$ , then  $\lambda_i(\Omega) \to +\infty$  when  $t \to +\infty$ . Thus, if  $\lambda_i(VA + A^{\top}V^{\top}) < 0$ , then  $\lambda_i(\Sigma(t)) = \lambda_{d-i}(\Omega(t)) \to 0$  when  $t \to +\infty$ . Finally, if  $\lambda_i(VA + A^{\top}V^{\top}) = 0$ , then  $\lambda_i(\Sigma) = \lambda_i(\Omega)^{-1}$  stays bounded along the dynamics. To prove convergence of  $\Sigma(t)$ , notice that the coefficients of  $\Sigma$  are rational fractions of t, so they converge in  $\mathbb{R} \cup \{+\infty, -\infty\}$  as  $t \to +\infty$ . Moreover, we have seen that they have to be bounded over time, as  $\sum_i \lambda_i(\Sigma(t))^2 = \|\Sigma(t)\|_F^2$  is bounded, which proves convergence of  $\Sigma$  to a positive semi-definite matrix  $\Sigma^*$ .

Proof of Proposition 4.8. Let us first consider the case where A is a diagonal matrix  $\operatorname{diag}(\lambda_1, \ldots, \lambda_d)$  with  $\lambda_1 = \cdots = \lambda_k = 0$ ,  $\lambda_{k+1}, \ldots, \lambda_{k+\ell} = 1$  and  $\lambda_{k+\ell+1} = \cdots = \lambda_d = 0$  where k and  $\ell$  are respectively the multiplicity of 0 and 1 as eigenvalues of A. Then

$$\Sigma A \Sigma = \sum_{i=k+1}^{k+\ell} C_i(\Sigma) C_i(\Sigma)^\top - \sum_{i=k+\ell+1}^d C_i(\Sigma) C_i(\Sigma)^\top,$$

where  $C_i(\Sigma)$  is the *i*-th column of  $\Sigma$ , seen as a column vector. If  $\Sigma A \Sigma = 0$  then

$$\sum_{i=k+1}^{k+\ell} C_i(\Sigma)C_i(\Sigma)^{\top} = \sum_{i=k+\ell+1}^d C_i(\Sigma)C_i(\Sigma)^{\top}$$

so that

$$\Sigma^2 = \sum_{i=1}^d C_i(\Sigma)C_i(\Sigma)^\top = \sum_{i=1}^k C_i(\Sigma)C_i(\Sigma)^\top + 2\sum_{i=k+1}^{k+\ell} C_i(\Sigma)C_i(\Sigma)^\top,$$

which means that the rank of  $\Sigma^2$  cannot exceed  $k + \ell$ . If  $\ell \leq d - (k + \ell)$ , this proves the result. If  $\ell > d - (k + \ell)$ , writing

$$\Sigma^2 = \sum_{i=1}^d C_i(\Sigma)C_i(\Sigma)^\top = \sum_{i=1}^k C_i(\Sigma)C_i(\Sigma)^\top + 2\sum_{i=k+\ell+1}^d C_i(\Sigma)C_i(\Sigma)^\top$$

allows to conclude.

Now if A is a general symmetric matrix, write  $A = ODO^{\top}$  with O an orthogonal matrix and  $D = \text{diag}(\lambda_1, \ldots, \lambda_d)$  with  $\lambda_1 = \cdots = \lambda_k = 0, \lambda_{k+1}, \ldots, \lambda_{k+\ell} > 0$  and

 $\lambda_{k+\ell+1}, \ldots, \lambda_d < 0$  (the case where one these groups of eigenvalues is empty can be solved with the exact same method). Then  $\Sigma A \Sigma = 0$  if and only if  $\tilde{\Sigma} \tilde{D} \tilde{\Sigma} = 0$  with  $\tilde{D} := \operatorname{sign}(D)$  and

$$\tilde{\Sigma} \coloneqq \operatorname{diag}(1,\ldots,1,\sqrt{|\lambda_{k+1}|},\ldots,\sqrt{|\lambda_d|})O^{\top}\Sigma O\operatorname{diag}(1,\ldots,1,\sqrt{|\lambda_{k+1}|},\ldots,\sqrt{|\lambda_d|}).$$

The first part of the proof implies that

 $\operatorname{rk}\tilde{\Sigma} \leq \dim \ker A + \min(\# \{ \text{positive eigenvalues of } A \}, \# \{ \text{negative eigenvalues of } A \} \}).$ 

We conclude by noticing that the matrices  $\Sigma$  and  $\tilde{\Sigma}$  have the same rank, as  $\lambda_i \neq 0$  for  $i \geq k+1$ .

Proof of Lemma 4.12. Consider the map

$$\varphi(M) = (M^{-1} + 2K^{\top}K) M = (I_d + 2K^{\top}KM)^{-1}M^2,$$

defined on the set  $\{M \in \mathbb{R}^{d \times d} : M \text{ symmetric s.t. } M \succeq 0 \text{ and } MK^{\top}K = K^{\top}KM \}$ of non-negative matrices that commute with  $K^{\top}K$ , which is a Banach space when equipped with Frobenius norm. This map is locally Lipschitz continuous, so Cauchy-Lipschitz theorem gives us the local existence and uniqueness of the considered Cauchy problem.

To recover global existence (and uniqueness), let us show that  $\|\varphi(M)\|_F$  grows at most linearly with  $\|M\|_F$ . Using that the squared Frobenius norm of a matrix is equal to the sum of its squared eigenvalues, we get

$$\left\| \left( M^{-1} + 2K^{\top}K \right)^{-1} \right\|_{F}^{2} = \sum_{i=1}^{d} \frac{1}{\lambda_{i}(M^{-1} + 2K^{\top}K)^{2}}$$
$$= \sum_{i=1}^{d} \frac{1}{\left(\lambda_{i}(M^{-1}) + \lambda_{\sigma(i)}(2K^{\top}K)\right)^{2}}$$

for some permutation  $\sigma \in \mathfrak{S}_d$ , where  $\lambda_i$  denotes the *i*-th eigenvalue, as the matrices  $M^{-1}$  and  $2K^{\top}K$  commute. Then

$$\left\| \left( M^{-1} + 2K^{\top}K \right)^{-1} \right\|_{F}^{2} = \sum_{i=1}^{d} \frac{1}{\left( \lambda_{i}(M)^{-1} + \lambda_{\sigma(i)}(2K^{\top}K) \right)^{2}} \\ = \sum_{i=1}^{d} \frac{\lambda_{i}(M)^{2}}{\left( 1 + \lambda_{\sigma(i)}(2K^{\top}K)\lambda_{i}(M) \right)^{2}}.$$

Now, the function

$$x \in \mathbb{R}_+ \mapsto \frac{x^2}{(1 + \lambda_{\sigma(i)}(2K^\top K)x)^2}$$

is bounded on  $\mathbb{R}_+$ , so that there exists a constant C > 0 such that

$$\left\| \left( M^{-1} + 2K^{\top}K \right)^{-1} \right\|_{F}^{2} \le C^{2}$$

for all non-negative symmetric matrices M that commute with  $K^{\top}K$ . Finally, as the Frobenius norm is sub-multiplicative:

$$\|\varphi(M)\|_{F} \leq \left\| \left( M^{-1} + 2K^{\top}K \right)^{-1} \right\|_{F} \|M\|_{F} \leq C \|M\|_{F},$$

which proves the claim and allows us to apply the global Cauchy theorem.

Proof of Lemma 4.14. Let  $c_{\varepsilon}(x,y) := \frac{1}{2\varepsilon} |Qx - Ky|^2$ . Denote  $\pi^*$  the minimizer of the entropic optimal transport problem

$$\min_{\pi \in \Pi(\mu,\mu)} \int c_{\varepsilon}(x,y) \, \mathrm{d}\pi(x,y) + \mathrm{KL}(\pi \| \mu \otimes \mu),$$

with the notations of Section 2. Recall that  $\kappa_{\mu,\varepsilon}^{\infty}$ , simply denoted  $\kappa_{\varepsilon}$  in the rest of the proof, is defined as the density of  $\pi^*$  with respect to the probability measure  $\mu \otimes \mu$ . According to our generalization of [32] (see Section A, Theorem A.3), it holds

$$\pi^* = \mathcal{N}\left(\begin{pmatrix} \alpha \\ \alpha \end{pmatrix}, \begin{pmatrix} \Sigma & A^{-1}C^{\top} \\ CA^{-\top} & \Sigma \end{pmatrix}\right),$$

with

$$C \coloneqq \Sigma^{1/2} \left( \Sigma^{1/2} A \Sigma A^{\top} \Sigma^{1/2} + \frac{\varepsilon^2}{4} I_d \right)^{1/2} \Sigma^{-1/2} - \frac{\varepsilon}{2} I_d$$

Let us compute the density of the probability measure  $\kappa_{\varepsilon}(x, y) d\mu(y)$ , parameterized by  $x \in \mathbb{R}^d$ . This density is proportional to

$$e^{-\frac{1}{2}\begin{pmatrix}x-\alpha\\y-\alpha\end{pmatrix}^{\top}\left(\begin{pmatrix}\Sigma\\CA^{-T}&\Sigma\end{pmatrix}^{-1}-\begin{pmatrix}\Sigma^{-1}&0\\0&0\end{pmatrix}\right)\begin{pmatrix}x-\alpha\\y-\alpha\end{pmatrix}}.$$

With the notations of the proof of Theorem A.3, i.e.,

$$H = \begin{pmatrix} \Sigma & A^{-1}C^{\top} \\ CA^{-\top} & \Sigma \end{pmatrix} \text{ and } \begin{cases} F &= \varepsilon \Sigma^{-1} + A^{\top}G^{-1}A \\ G &= \varepsilon \Sigma^{-1} + A^{\top}F^{-1}A, \end{cases}$$

we have with Lemma A.6

$$\begin{pmatrix} H^{-1} - \begin{pmatrix} \Sigma^{-1} & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix} = \frac{1}{\varepsilon} \begin{pmatrix} F - \varepsilon \Sigma^{-1} & -A^{\top} \\ -A & G \end{pmatrix}$$
$$= \frac{1}{\varepsilon} \begin{pmatrix} A^{\top} G^{-1} A & -A^{\top} \\ -A & G \end{pmatrix}$$

noticing that  $A^{\top}G^{-1}A = F^{-1} - \varepsilon \Sigma^{-1}$  with Equation (A.10). Then, the density of  $\kappa_{\varepsilon}(x, y) d\mu(y)$  is proportional, up to factors that do not depend on y, to

$$\exp\left(-\frac{1}{2\varepsilon}(y-\alpha-G^{-1}A(x-\alpha))^{\top}G(y-\alpha-G^{-1}A(x-\alpha))\right).$$

Finally, we have

$$C_G = \Sigma A^\top G^{-1} A$$

according to Equation (A.12), so that

$$\kappa_{\varepsilon}(x,\cdot) d\mu = \mathcal{N}(\alpha + A^{-\top} \Sigma^{-1} C_G(x-\alpha), \varepsilon G^{-1})$$

and therefore

$$\Gamma_{\mu}(x) = \frac{1}{\varepsilon} V \mathbb{E}[\kappa_{\varepsilon}(x, \cdot) d\mu] = \frac{1}{\varepsilon} V (I_d - A^{-\top} \Sigma^{-1} C_G) \alpha + \frac{1}{\varepsilon} V A^{-\top} \Sigma^{-1} C_G x.$$



FIG. 4. Comparison of the behavior of Softmax, L2 and Multi-head attention in the setting of Figure 1. All plots correspond to the same parameters, with V random and  $A + A^{\top} \prec 0$ . We observe very similar behaviors.



FIG. 5. Evolution of the covariance matrix of a 2-dimensional Gaussian measure that goes through the L2 Transformer PDE. All plots were obtained with L2 self-attention, with the same parameters as in Figure 1 (a, b, d). The behavior looks extremely similar as for Softmax self-attention.

**C.2. Experiments.** Figure 5 is the parallel of Figure 1 for L2 self-attention. Figure 6 complements Figure 2 (d), and highlights that behaviors are very similar for Softmax and multi-head self-attention. The case of L2 self-attention replaces finite-time divergence with infinite-time divergence. Figure 7 plots the behavior of the Sinkformer PDE on Gaussians, in two cases already investigated for Softmax, L2 and multi-head self-attention. We observe a similar behavior as with Softmax self-attention. Figure 8 was obtained with the same procedure as Figure 3 but for L2 self-attention. The conclusion is the same: we observe that limiting points have a low rank.

### Appendix D. Proofs of Section 5.

**D.1. The Bures-Wasserstein Gradient Flow Induced by Sinkformers.** We have seen in Section 5.2 that the Sinkformer PDE initialized with a Gaussian probability measure induces a Bures-Wasserstein gradient flow in the space of Gaussian measures, associated with the energy functional

$$F_{\varepsilon}(\alpha, \Sigma) = \mathcal{F}_{\varepsilon}(\mathcal{N}(\alpha, \Sigma))$$

where  $\mathcal{F}$  is defined in Equation (5.11). Let us compute a more explicit expression for  $F_{\varepsilon}$ . First, for any  $\mu \in \mathcal{P}(\mathbb{R}^d)$  such that everything is well-defined, we have

$$\mathcal{F}_{\varepsilon}(\mu) \coloneqq -\frac{1}{2} \int \kappa_{\mu,\varepsilon}^{\infty} \log \left(\frac{\kappa_{\mu,\varepsilon}^{\infty}}{\kappa_{\mu,\varepsilon}^{0}}\right) \,\mathrm{d}(\mu \otimes \mu) + \frac{1}{4\varepsilon} \int (|Qx|^{2} + |Kx|^{2}) \,\mathrm{d}\mu(x)$$



FIG. 6. Projection on the set of trace-1 matrices of the dynamics of the covariance matrix of a Gaussian measure that goes through the Transformer PDE, in cases where curves blow up or diverge. We obtained the plots (a), (b) and (c) with the same parameters, chosen specifically to observe a division of the behavior between convergence (yellow curves) and blow-up or divergence (purple curves).  $A + A^{\top}$  has one positive and one negative eigenvalue.



FIG. 7. Evolution of the covariance matrix of a 2-dimensional Gaussian measure that goes through the Sinkformer PDE. Figure (a) was obtained with the same parameters as in Figure 1, and Figure (b) with the same parameters as in Figure 2. In both cases, the behavior is very similar to what happens with Softmax self-attention.



FIG. 8. Histogram of the rank of limiting points of the covariance equation for L2 self-attention, in dimensions 3, 4, and 5. The matrix V has full rank ( $V = I_d$  in the upper row and V random and different for each point in the lower row) and the matrix A has rank  $\lfloor d/2 \rfloor$ , is random (constrained to be negative symmetric) and different for each point. Limiting points have a low rank (smaller than  $\lfloor d/2 \rfloor$ .

and, according to Equation (A.3) we have

$$OT_{\varepsilon}(\mu,\mu) = \frac{1}{2\varepsilon} \int |Qx - Ky|^2 \kappa_{\mu,\varepsilon}^{\infty}(x,y) \,\mathrm{d}\mu(x) \,\mathrm{d}\mu(y) + \int \log(\kappa_{\mu,\varepsilon}^{\infty}) \kappa_{\mu,\varepsilon}^{\infty} \,\mathrm{d}\mu(x) \,\mathrm{d}\mu(y).$$

Therefore, it holds, recalling that  $\kappa^0_{\mu,\varepsilon}(x,y) = e^{-\frac{1}{2\varepsilon}|Qx-Ky|^2}$ :

$$\mathcal{F}_{\varepsilon}(\mu) = -\frac{1}{2}OT_{\varepsilon}(\mu,\mu) + \frac{1}{4\varepsilon}\int (|Qx|^2 + |Ky|^2) \,\mathrm{d}\mu(x)$$

Now, by definition of the entropy-regularized Bures distance (5.13) we have

$$OT_{\varepsilon}(\mathcal{N}(\alpha, \Sigma), \mathcal{N}(\alpha, \Sigma)) = \frac{1}{2\varepsilon} \mathfrak{B}_{\varepsilon}^{2}(\Sigma, \Sigma),$$

so that

$$F_{\varepsilon}(\alpha, \Sigma) = -\frac{1}{4\varepsilon} \mathfrak{B}_{\varepsilon}^{2}(\Sigma, \Sigma) + \frac{1}{4\varepsilon} \int (|Qx|^{2} + |Ky|^{2}) \,\mathrm{d}\mu(x).$$

Finally, writing  $|Qx|^2 = \text{Tr}(Qxx^\top Q^\top)$  we compute

$$\int |Qx|^2 \, \mathrm{d}\mathcal{N}(\alpha, \Sigma)(x) = \mathrm{Tr}(Q \int xx^\top \, \mathrm{d}\mathcal{N}(\alpha, \Sigma)(x)Q^\top) = \mathrm{Tr}(Q\Sigma Q^\top) + \alpha^\top Q^\top Q\alpha.$$

The same computation for  $\int |Kx|^2 d\mathcal{N}(\alpha, \Sigma)(x)$  leads to

$$F_{\varepsilon}(\alpha, \Sigma) = \frac{1}{4\varepsilon} \left( -\mathfrak{B}_{\varepsilon}^{2}(\Sigma, \Sigma) + \operatorname{Tr}(Q\Sigma Q^{\top}) + \operatorname{Tr}(K\Sigma K^{\top}) + \alpha^{\top}(Q^{\top}Q + K^{\top}K)\alpha \right).$$

# **D.2.** The Transformer PDE as a $d_{A,V}$ Gradient Flow.

Energy Functional On Gaussians. We have seen in Section 5.4 that the Transformer PDE associated with Softmax self-attention is a gradient flow for the twisted distance  $d_{A,V}$ , the energy functional being

$$\mathcal{F}(\mu) = \frac{1}{2} \int e^{Qx \cdot Ky} \,\mathrm{d}\mu(x) \,\mathrm{d}\mu(y).$$

Let us compute the expression of  $F(\alpha, \Sigma) := \mathcal{F}(\mathcal{N}(\alpha, \Sigma))$  for  $\alpha \in \mathbb{R}^d$  and  $\Sigma$  a  $d \times d$  covariance matrix. First, we have

$$x^{\top}A^{\top}y - \frac{1}{2}(x-\alpha)^{\top}\Sigma^{-1}(x-\alpha) = -\frac{1}{2}(x-\alpha-\Sigma A^{\top}y)^{\top}\Sigma^{-1}(x-\alpha-\Sigma A^{\top}y) + \alpha^{\top}A^{\top}y + \frac{1}{2}y^{\top}A\Sigma A^{\top}y$$

This gives

$$\int e^{x^{\top}A^{\top}y} \, \mathrm{d}\mathcal{N}(\alpha, \Sigma)(x) = \int e^{x^{\top}A^{\top}y - \frac{1}{2}(x-\alpha)^{\top}\Sigma^{-1}(x-\alpha)} \frac{\mathrm{d}x}{(2\pi)^{d/2} \left|\det\Sigma\right|^{1/2}}$$
$$= e^{\alpha^{\top}A^{\top}y + \frac{1}{2}y^{\top}A\Sigma A^{\top}y}.$$

In view of computing the integral of this term in y, we write

$$\begin{aligned} &-\frac{1}{2}(y-\alpha)^{\top}\Sigma^{-1}(y-\alpha) + \alpha^{\top}A^{\top}y + \frac{1}{2}y^{\top}A\Sigma A^{\top}y = \\ &-\frac{1}{2}(y-(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})\alpha)^{\top}(\Sigma^{-1}-A\Sigma A^{\top})(y-(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})\alpha) \\ &+\frac{1}{2}\alpha^{\top}(A^{\top}+\Sigma^{-1})(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})\alpha - \frac{1}{2}\alpha^{\top}\Sigma^{-1}\alpha. \end{aligned}$$

A similar computation as for the integral in x leads to

$$F(\alpha, \Sigma) = \frac{e^{\frac{1}{2}\alpha^{\top}((A+\Sigma^{-1})^{\top}(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})-\Sigma^{-1})\alpha}}{2|\det(\Sigma^{-1}-A\Sigma A^{\top})|^{1/2}|\det\Sigma|^{1/2}}$$
$$= \frac{e^{\frac{1}{2}\alpha^{\top}((A+\Sigma^{-1})^{\top}(\Sigma^{-1}-A\Sigma A^{\top})^{-1}(A+\Sigma^{-1})-\Sigma^{-1})\alpha}}{2|\det(I_d-A\Sigma A^{\top}\Sigma)|^{1/2}}.$$

Geodesics of  $d_{A,V}$ . We look for a characterization of geodesics for  $d_{A,V}$ . We reformulate it via a Lagrange multiplier  $\psi(t, x)$ :

$$d_{A,V}(\mu,\nu)^{2} = \inf_{(\rho,v)} \sup_{\psi} \int_{0}^{1} \int_{\mathbb{R}^{d}} \frac{v \cdot Bv}{G * \rho} \, \mathrm{d}\rho \, \mathrm{d}s$$
$$- \int_{0}^{1} \int_{\mathbb{R}^{d}} \left( \partial_{s}\psi(s,x) + \frac{Bv(s,x) \cdot \nabla_{x}\psi(s,x)}{G * \rho(x)} \right) \, \mathrm{d}\rho(x) \, \mathrm{d}s$$
$$+ \int_{\mathbb{R}^{d}} \psi(1,x) \, \mathrm{d}\nu(x) - \int_{\mathbb{R}^{d}} \psi(0,x) \, \mathrm{d}\mu(x)$$

where  $(\rho, u) \in \mathcal{C}([0, 1], \mathcal{P}_2(\mathbb{R}^d)) \times \mathcal{C}([0, 1], \mathcal{C}(\mathbb{R}^d, \mathbb{R}^d))$  in the infimum is now unconstrained. The optimality condition on v is  $v = \frac{1}{2} \nabla_x \psi$ . Incorporating this in the equation, the problem becomes

$$d_{A,V}(\mu,\nu)^2 = \inf_{\rho} \sup_{\psi} \int_0^1 \int_{\mathbb{R}^d} -\left(\partial_s \psi(s,x) + \frac{1}{4} \frac{B\nabla_x \psi \cdot \nabla_x \psi}{G * \rho}\right) \,\mathrm{d}\rho(x) \,\mathrm{d}s + \int_{\mathbb{R}^d} \psi(1,x) \,\mathrm{d}\nu(x) - \int_{\mathbb{R}^d} \psi(0,x) \,\mathrm{d}\mu(x).$$

Then, the optimality in  $\rho$  gives

$$\partial_s \psi(s,x) + \frac{1}{4} \frac{B\nabla_x \psi \cdot \nabla_x \psi}{G * \rho(x)} = \int \frac{1}{4} \frac{B\nabla_y \psi \cdot \nabla_y \psi}{(G * \rho(y))^2} G(y,x) \,\mathrm{d}\rho(y).$$

We thus obtain the following equations characterizing geodesics for  $d_{A,V}$ :

$$\begin{cases} \partial_s \rho + \operatorname{div}\left(\frac{B\nabla_x \psi}{2G*\rho}\rho\right) = 0\\ \partial_s \psi + \frac{1}{4} \frac{\nabla_x \psi \cdot B\nabla_x \psi}{G*\rho} - \frac{1}{4} \int_{\mathbb{R}^d} \frac{\nabla_y \psi \cdot B\nabla_y \psi}{(G*\rho(y))^2} G(y, x) \rho_s(y) \, \mathrm{d}y = 0. \end{cases}$$

Proof of Proposition 5.4. If  $\rho$  is a geodesic for  $d_{A,V}$ , associated with the test function  $\psi$ , we have

$$\partial_t \mathcal{E}(\rho) = \frac{1}{2} \int \int \nabla_x G(x, y) \cdot (B \nabla_x \psi) \frac{\rho(x) \rho(y)}{G * \rho(x)} \, \mathrm{d}x \, \mathrm{d}y$$
$$= -\frac{1}{2} \int \Gamma_\rho(x) \cdot \nabla_x \psi \, \rho(x) \, \mathrm{d}x.$$

In the whole proof, we write  $d\rho(y) = \rho(y) dy$  for simplicity—this is only a formal

computation. In order to compute  $\partial_t^2 \mathcal{E}(\rho),$  let us compute

$$\begin{split} (\partial_t \Gamma_\rho(x))_i &= \int (Vy)_i G(x,y) \left( \frac{\partial_t \rho(y)}{G * \rho(x)} - \frac{\rho(y)}{(G * \rho(x))^2} \int G(x,z) \partial_t \rho(z) \, \mathrm{d}z \right) \, \mathrm{d}y \\ &= \frac{1}{2} \int \nabla_y ((Vy)_i \cdot G(x,y)) \cdot (B \nabla_y \psi) \frac{\rho(y)}{G * \rho(x) G * \rho(y)} \, \mathrm{d}y \\ &\quad - \frac{1}{2} \int (Vy)_i G(x,y) \frac{\rho(y)}{(G * \rho(x))^2} \int \nabla_z G(x,z) \cdot (B \nabla_z \psi) \frac{\rho(z)}{G * \rho(z)} \, \mathrm{d}z \, \mathrm{d}y \\ &= \frac{1}{2} \int (L_i(V) G(x,y) + (Vy)_i G(x,y) Ax) \cdot (B \nabla_y \psi) \frac{\rho(y)}{G * \rho(y) G * \rho(x)} \, \mathrm{d}y \\ &\quad - \frac{1}{2} \int (Vy)_i G(x,y) \frac{\rho(y)}{(G * \rho(x))^2} \int G(x,z) (Ax) \cdot (B \nabla_z \psi) \frac{\rho(z)}{G * \rho(z)} \, \mathrm{d}z \, \mathrm{d}y, \end{split}$$

where  $L_i(V)$  is the *i*-th row of V, seen as a column vector. Then

$$\begin{split} \partial_t \Gamma_\rho(x) &= \frac{1}{2} \int G(x,y) \left( VB \nabla_y \psi + (Ax) \cdot (B \nabla_y \psi) Vy \right) \frac{\rho(y)}{G * \rho(y) G * \rho(x)} \, \mathrm{d}y \\ &\quad - \frac{1}{2} \int G(x,z) (Ax) \cdot (B \nabla_z \psi) \frac{\rho(z)}{G * \rho(z)} \, \mathrm{d}z \int Vy G(x,y) \frac{\rho(y)}{(G * \rho(x))^2} \, \mathrm{d}y \\ &= \frac{1}{2} \int G(x,y) \left( VB \nabla_y \psi - x^\top V^\top \nabla_y \psi Vy \right) \frac{\rho(y)}{G * \rho(y) G * \rho(x)} \, \mathrm{d}y \\ &\quad + \frac{1}{2} \int G(x,z) x^\top V^\top \nabla_z \psi \frac{\rho(z)}{G * \rho(x) G * \rho(z)} \, \mathrm{d}z \Gamma_\rho(x) \end{split}$$

as  $AB = -V^{\top}$ . Finally,

$$\begin{split} \partial_t \Gamma_\rho(x) &= \frac{1}{2} \int G(x,y) V B \nabla_y \psi \frac{\rho(y)}{G * \rho(y) G * \rho(x)} \, \mathrm{d}y \\ &\quad + \frac{1}{2} \int G(x,y) (Vx) \cdot \nabla_y \psi (\Gamma_\rho(x) - Vy) \frac{\rho(y)}{G * \rho(x) G * \rho(y)} \, \mathrm{d}y \\ &= \frac{1}{2} \int G(x,y) \left( VB + (\Gamma_\rho(x) - Vy) (Vx)^\top \right) \nabla_y \psi \frac{\rho(y)}{G * \rho(x) G * \rho(y)} \, \mathrm{d}y. \end{split}$$

We use this formula to compute  $\partial_t^2 \mathcal{E}(\rho)$ :

$$\partial_t^2 \mathcal{E}(\rho) = -\frac{1}{2} \int (\nabla_x \partial_t \psi) \cdot \Gamma_\rho(x) \rho(x) \, \mathrm{d}x - \frac{1}{2} \int (\nabla_x \psi) \cdot (\Gamma_\rho(x)) \partial_t \rho(x) \, \mathrm{d}x \\ - \frac{1}{2} \int (\nabla_x \psi) \cdot (\partial_t \Gamma_\rho) \rho(x) \, \mathrm{d}x.$$

Mechanical but painful computations, that we do not reproduce here, lead to

$$\begin{split} \partial_t^2 \mathcal{E}(\rho) &= \frac{1}{8} \int \nabla_x \psi \cdot (B \nabla_x \psi) \Gamma_{\rho}(x) \cdot (B^{-1} \Gamma_{\rho}(x)) \frac{\rho(x)}{G * \rho(x)} \, \mathrm{d}x \\ &+ \frac{1}{8} \iint \nabla_y \psi \cdot (B \nabla_y \psi) (Vy) \cdot (B^{-1} \Gamma_{\rho}(x)) G(x, y) \frac{\rho(x) \rho(y)}{(G * \rho(y))^2} \, \mathrm{d}x \, \mathrm{d}y \\ &- \frac{1}{4} \int (\nabla_x \psi \cdot \Gamma_{\rho}(x))^2 \frac{\rho(x)}{G * \rho(x)} \, \mathrm{d}x \\ &+ \frac{1}{4} \iint (\nabla_x \psi \cdot Vy)^2 G(x, y) \frac{\rho(x) \rho(y)}{(G * \rho(x))^2} \, \mathrm{d}x \, \mathrm{d}y \\ &- \frac{1}{4} \iint \nabla_x \psi \cdot (V B \nabla_y \psi) G(x, y) \frac{\rho(x) \rho(y)}{G * \rho(x) G * \rho(y)} \, \mathrm{d}x \, \mathrm{d}y \\ &- \frac{1}{4} \iint \nabla_x \psi \cdot (\Gamma_{\rho}(x) - Vy) (Vx) \cdot \nabla_y \psi \frac{\rho(x) \rho(y)}{G * \rho(x) G * \rho(y)} \, \mathrm{d}x \, \mathrm{d}y. \end{split}$$

Let us now compute  $\partial_t^2 \mathcal{E}(\rho)$  when  $\rho = \delta_z$  is the Dirac measure at  $z \in \mathbb{R}^d$ . We obtain, as  $\Gamma_{\delta_z}(x) = Vz$ , that

$$\partial_t^2 \mathcal{E}(\delta_z) = \frac{1}{4} e^{-Az \cdot z} (\nabla_z \psi)^\top \left( (z^\top V^\top B^{-1} V z) I_d - V \right) B \nabla_z \psi.$$

Assuming that V commutes with  $B \succ 0$ , this can be rewritten as

$$\partial_t^2 \mathcal{E}(\delta_z) = \frac{1}{4} e^{-Az \cdot z} (\nabla_z \psi)^\top B^{1/2} \left( (z^\top V^\top B^{-1} V z) I_d - V \right) B^{1/2} \nabla_z \psi.$$

As  $V \succ 0$ , we can find (infinitely many) vectors  $z \in \mathbb{R}^d$  such that the matrix  $(z^\top V^\top B^{-1} V z) I_d - V$  is negative definite, by imposing that  $z^\top V^\top B^{-1} V z$  is strictly smaller than the smallest eigenvalue of V. Note that we use positive definiteness of V here, as  $z^\top V^\top B^{-1} V z > 0$  for any z. Pick one such  $z \in \mathbb{R}^d$ . Now, consider any (for example compactly supported) measure  $\mu$ , and denote  $(\rho, \psi)$  the geodesic between  $\delta_z$  and  $\mu$  for  $d_{A,V}$ . As  $\partial_{t=0}^2 \mathcal{E}(\delta_z) < 0$ , the functional  $\mathcal{E}$  is not convex along  $\rho$ , which proves the claim.

#### REFERENCES

- L. AMBROSIO, N. GIGLI, AND G. SAVARÉ, Gradient flows: in metric spaces and in the space of probability measures, Springer Science & Business Media, 2008.
- [2] R. BARBONI, G. PEYRÉ, AND F.-X. VIALARD, Understanding the training of infinitely deep and wide resnets with conditional optimal transport, arXiv preprint arXiv:2403.12887, (2024).
- [3] A. L. BERTOZZI, J. A. CARRILLO, AND T. LAURENT, Blow-up in multidimensional aggregation equations with mildly singular interaction kernels, Nonlinearity, 22 (2009), pp. 683– 710, https://doi.org/10.1088/0951-7715/22/3/009, https://doi.org/10.1088/0951-7715/ 22/3/009.
- [4] R. BOJILOV AND A. GALICHON, Matching in closed-form: equilibrium, identification, and comparative statics, Economic Theory, 61 (2016), pp. 587–609.
- [5] Y. BRENIER, Extended Monge-Kantorovich theory, in Optimal transportation and applications (Martina Franca, 2001), vol. 1813 of Lecture Notes in Math., Springer, Berlin, 2003, pp. 91–121, https://doi.org/10.1007/978-3-540-44857-0\_4, https://doi.org/10.1007/ 978-3-540-44857-0\_4.
- [6] D. BURES, An extension of kakutani's theorem on infinite product measures to the tensor product of semifinite w\*-algebras, Transactions of the American Mathematical Society, 135 (1969), pp. 199–212.

- M. BURGER, M. ERBAR, F. HOFFMANN, D. MATTHES, AND A. SCHLICHTING, Covariance-Modulated Optimal Transport and Gradient Flows, Arch. Ration. Mech. Anal., 249 (2025), p. Paper No. 7, https://doi.org/10.1007/s00205-024-02065-w, https://doi.org/10.1007/ s00205-024-02065-w.
- [8] M. BURGER, S. KABRI, Y. KOROLEV, T. ROITH, AND L. WEIGAND, Analysis of mean-field models arising from self-attention dynamics in transformer architectures with layer normalization, arXiv preprint arXiv:2501.03096, (2025).
- [9] J. A. CANIZO, J. A. CARRILLO, AND J. ROSADO, A well-posedness theory in measures for some kinetic models of collective motion, Mathematical Models and Methods in Applied Sciences, 21 (2011), pp. 515–539.
- [10] G. CARLIER, L. CHIZAT, AND M. LABORDE, Lipschitz continuity of the schr\" odinger map in entropic optimal transport, arXiv preprint arXiv:2210.00225, (2022).
- [11] J. A. CARRILLO, Y.-P. CHOI, AND M. HAURAY, The derivation of swarming models: mean-field limit and Wasserstein distances, in Collective dynamics from bacteria to crowds, vol. 553 of CISM Courses and Lect., Springer, Vienna, 2014, pp. 1–46, https://doi.org/10.1007/ 978-3-7091-1785-9\_1, https://doi.org/10.1007/978-3-7091-1785-9\_1.
- [12] J. A. CARRILLO, K. CRAIG, AND Y. YAO, Aggregation-diffusion equations: dynamics, asymptotics, and singular limits, in Active particles. Vol. 2. Advances in theory, models, and applications, Model. Simul. Sci. Eng. Technol., Birkhäuser/Springer, Cham, 2019, pp. 65– 108.
- [13] J. A. CARRILLO, M. G. DELGADINO, L. DESVILLETTES, AND J. S.-H. WU, The Landau equation as a gradient flow, Anal. PDE, 17 (2024), pp. 1331–1375, https://doi.org/10.2140/apde. 2024.17.1331, https://doi.org/10.2140/apde.2024.17.1331.
- [14] J. A. CARRILLO, R. S. GVALANI, AND J. S.-H. WU, An invariance principle for gradient flows in the space of probability measures, J. Differential Equations, 345 (2023), pp. 233–284, https://doi.org/10.1016/j.jde.2022.11.028, https://doi.org/10.1016/j.jde.2022.11.028.
- [15] J. A. CARRILLO, F. HOFFMANN, A. M. STUART, AND U. VAES, Consensus-based sampling, Stud. Appl. Math., 148 (2022), pp. 1069–1140, https://doi.org/10.1111/sapm.12470, https: //doi.org/10.1111/sapm.12470.
- [16] J. A. CARRILLO, S. LISINI, G. SAVARÉ, AND D. SLEPČEV, Nonlinear mobility continuity equations and generalized displacement convexity, J. Funct. Anal., 258 (2010), pp. 1273–1309, https://doi.org/10.1016/j.jfa.2009.10.016, https://doi.org/10.1016/j.jfa.2009.10.016.
- [17] J. A. CARRILLO, R. J. MCCANN, AND C. VILLANI, Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates, Rev. Mat. Iberoamericana, 19 (2003), pp. 971–1018, https://doi.org/10.4171/RMI/376, https: //doi.org/10.4171/RMI/376.
- [18] J. A. CARRILLO AND U. VAES, Wasserstein stability estimates for covariance-preconditioned Fokker-Planck equations, Nonlinearity, 34 (2021), pp. 2275-2295, https://doi.org/10.1088/ 1361-6544/abbe62, https://doi.org/10.1088/1361-6544/abbe62.
- [19] V. CASTIN, P. ABLIN, AND G. PEYRÉ, How smooth is attention?, in ICML 2024, 2024.
- [20] R. T. CHEN, Y. RUBANOVA, J. BETTENCOURT, AND D. K. DUVENAUD, Neural ordinary differential equations, Advances in neural information processing systems, 31 (2018).
- [21] K. CHOROMANSKI, V. LIKHOSHERSTOV, D. DOHAN, X. SONG, A. GANE, T. SARLOS, P. HAWKINS, J. DAVIS, A. MOHIUDDIN, L. KAISER, ET AL., *Rethinking attention with performers*, arXiv preprint arXiv:2009.14794, (2020).
- [22] J. DELON AND A. DESOLNEUX, A wasserstein-type distance in the space of gaussian mixture models, SIAM Journal on Imaging Sciences, 13 (2020), pp. 936–970.
- [23] J. DEVLIN, Bert: Pre-training of deep bidirectional transformers for language understanding, arXiv preprint arXiv:1810.04805, (2018).
- [24] R. L. DOBRUSHIN, Vlasov equations, Functional Analysis and Its Applications, 13 (1979), pp. 115–123.
- [25] Q. FOURNIER, G. M. CARON, AND D. ALOISE, A practical survey on faster and lighter transformers, ACM Computing Surveys, 55 (2023), pp. 1–40.
- [26] A. GARBUNO-INIGO, F. HOFFMANN, W. LI, AND A. M. STUART, Interacting Langevin diffusions: gradient structure and ensemble Kalman sampler, SIAM J. Appl. Dyn. Syst., 19 (2020), pp. 412–441, https://doi.org/10.1137/19M1251655, https://doi.org/10.1137/19M1251655.
- [27] B. GESHKOVSKI, C. LETROUIT, Y. POLYANSKIY, AND P. RIGOLLET, A mathematical perspective on transformers, arXiv preprint arXiv:2312.10794, (2023).
- [28] B. GESHKOVSKI, C. LETROUIT, Y. POLYANSKIY, AND P. RIGOLLET, The emergence of clusters in self-attention dynamics, Advances in Neural Information Processing Systems, 36 (2024).
- [29] F. GOLSE, The mean-field limit for the dynamics of large particle systems, in Journées "Équations aux Dérivées Partielles", Univ. Nantes, Nantes, 2003, pp. Exp. No. IX, 47.

- [30] E. HABER AND L. RUTHOTTO, Stable architectures for deep neural networks, Inverse problems, 34 (2017), p. 014004.
- [31] B. HOSSEINI, A. W. HSU, AND A. TAGHVAEI, Conditional optimal transport on function spaces, arXiv preprint arXiv:2311.05672, (2023).
- [32] H. JANATI, B. MUZELLEC, G. PEYRÉ, AND M. CUTURI, Entropic optimal transport between unbalanced gaussian measures has a closed form, Advances in neural information processing systems, 33 (2020), pp. 10468–10479.
- [33] R. JORDAN, D. KINDERLEHRER, AND F. OTTO, The variational formulation of the fokker-planck equation, SIAM journal on mathematical analysis, 29 (1998), pp. 1–17.
- [34] A. KATHAROPOULOS, A. VYAS, N. PAPPAS, AND F. FLEURET, Transformers are rnns: Fast autoregressive transformers with linear attention, in International conference on machine learning, PMLR, 2020, pp. 5156–5165.
- [35] H. KIM, G. PAPAMAKARIOS, AND A. MNIH, The lipschitz constant of self-attention, 2021, https: //arxiv.org/abs/2006.04710.
- [36] M. LAMBERT, S. CHEWI, F. BACH, S. BONNABEL, AND P. RIGOLLET, Variational inference via wasserstein gradient flows, Advances in Neural Information Processing Systems, 35 (2022), pp. 14434–14447.
- [37] F. LÉGER, G. TODESCHI, AND F.-X. VIALARD, Nonnegative cross-curvature in infinite dimensions: synthetic definition and spaces of measures, arXiv preprint arXiv:2409.18112, (2024).
- [38] W. LI, Hessian metric via transport information geometry, Journal of Mathematical Physics, 62 (2021).
- [39] T. LIN, Y. WANG, X. LIU, AND X. QIU, A survey of transformers, AI open, 3 (2022), pp. 111– 132.
- [40] A. MAHANKALI, T. B. HASHIMOTO, AND T. MA, One step of gradient descent is provably the optimal in-context learner with one layer of linear self-attention, arXiv preprint arXiv:2307.03576, (2023).
- [41] L. MALAGO, L. MONTRUCCHIO, AND G. PISTONE, Wasserstein riemannian geometry of positive definite matrices, arXiv preprint arXiv:1801.09269, (2018).
- [42] S. MOTSCH AND E. TADMOR, Heterophilious dynamics enhances consensus, SIAM review, 56 (2014), pp. 577–621.
- [43] M. NUTZ, Introduction to entropic optimal transport, Lecture notes, Columbia University, (2021).
- [44] F. OTTO, The geometry of dissipative evolution equations: the porous medium equation, Comm. Partial Differential Equations, 26 (2001), pp. 101–174, https://doi.org/10.1081/ PDE-100002243, https://doi.org/10.1081/PDE-100002243.
- [45] H. PENG, N. PAPPAS, D. YOGATAMA, R. SCHWARTZ, N. A. SMITH, AND L. KONG, Random feature attention, arXiv preprint arXiv:2103.02143, (2021).
- [46] J. RAMAPURAM, F. DANIELI, E. DHEKANE, F. WEERS, D. BUSBRIDGE, P. ABLIN, T. LIKHOMA-NENKO, J. DIGANI, Z. GU, A. SHIDANI, ET AL., *Theory, analysis, and best practices for sigmoid self-attention*, arXiv preprint arXiv:2409.04431, (2024).
- [47] M. E. SANDER, P. ABLIN, M. BLONDEL, AND G. PEYRÉ, Sinkformers: Transformers with doubly stochastic attention, in International Conference on Artificial Intelligence and Statistics, PMLR, 2022, pp. 3515–3530.
- [48] M. E. SANDER, R. GIRYES, T. SUZUKI, M. BLONDEL, AND G. PEYRÉ, How do transformers perform in-context autoregressive learning?, arXiv preprint arXiv:2402.05787, (2024).
- [49] F. SANTAMBROGIO, Optimal transport for applied mathematicians, Birkäuser, NY, 55 (2015), p. 94.
- [50] I. SCHLAG, K. IRIE, AND J. SCHMIDHUBER, Linear transformers are secretly fast weight programmers, in International Conference on Machine Learning, PMLR, 2021, pp. 9355–9366.
- [51] Z. SHEN, M. ZHANG, H. ZHAO, S. YI, AND H. LI, Efficient attention: Attention with linear complexities, in Proceedings of the IEEE/CVF winter conference on applications of computer vision, 2021, pp. 3531–3539.
- [52] A. VASWANI, N. SHAZEER, N. PARMAR, J. USZKOREIT, L. JONES, A. N. GOMEZ, L. KAISER, AND I. POLOSUKHIN, Attention is all you need, Advances in neural information processing systems, 30 (2017).
- [53] J. L. VÁZQUEZ, The porous medium equation, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, Oxford, 2007. Mathematical theory.
- [54] C. VILLANI, Topics in optimal transportation, vol. 58 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2003, https://doi.org/10.1090/gsm/058, https://doi.org/10.1090/gsm/058.
- [55] A. A. VLASOV, The vibrational properties of an electron gas, Soviet Physics Uspekhi, 10 (1968),

p. 721.

- [56] J. VON OSWALD, E. NIKLASSON, M. SCHLEGEL, S. KOBAYASHI, N. ZUCCHET, N. SCHERRER, N. MILLER, M. SANDLER, M. VLADYMYROV, R. PASCANU, ET AL., Uncovering mesaoptimization algorithms in transformers, arXiv preprint arXiv:2309.05858, (2023).
- [57] J. VUCKOVIC, A. BARATIN, AND R. T. D. COMBES, A mathematical theory of attention, arXiv preprint arXiv:2007.02876, (2020).
- [58] E. WEINAN, A proposal on machine learning via dynamical systems, Communications in Mathematics and Statistics, 1 (2017), pp. 1–11.
- [59] M. WORTSMAN, J. LEE, J. GILMER, AND S. KORNBLITH, Replacing softmax with relu in vision transformers, arXiv preprint arXiv:2309.08586, (2023).
- [60] Y. D. ZHONG, T. ZHANG, A. CHAKRABORTY, AND B. DEY, A neural ode interpretation of transformer layers, arXiv preprint arXiv:2212.06011, (2022).