

# GRAPH DISCRETIZATION OF LAPLACIAN ON RIEMANNIAN MANIFOLDS WITH BOUNDED RICCI CURVATURE

ANUSHA BHATTACHARYA AND SOMA MAITY

ABSTRACT. We study the approximation of eigenvalues and eigenfunctions for the Laplace-Beltrami operator on compact manifolds without boundary. The analysis is centered on manifolds characterized by specific geometric constraints: lower bounds on Ricci curvature and injectivity radius, and an upper bound on diameter. Using weighted graph techniques, we approximate the eigenvalues for manifolds having a uniform lower bound on the volume of small balls and also investigate uniform bounds of eigenvalues applicable across the entire class.

## 1. INTRODUCTION

The methods of approximating the eigenvalues and eigenfunctions of differential operators using simpler and easily computable objects have been a topic of active research for a long time. Key contributions in this field include Yau and Li's work [12] on eigenvalue behavior in relation to compact manifold geometry and Aubry's work [2] on approximating the spectral data of a compact Riemannian manifold using discrete Laplace operators on immersed graphs. Belkin and Niyogi [3] demonstrate that for random, uniformly distributed points on a manifold, with appropriately chosen edge weights based on distances, the spectrum of the graph Laplacian converges to that of the manifold in a probabilistic sense.

In this paper, we focus on the technique of graph discretization of Riemannian manifolds to approximate the eigenvalues of differential operators like the Laplace-Beltrami operator. This area of research has gained immense popularity in recent decades. Burago, Ivanov and Kurylev [6] have proved the non-probabilistic spectral convergence of the graph Laplacian to the Laplace-Beltrami operator on closed manifolds with certain geometric constraints. They have also studied the spectral and stability properties of an analogue of the Laplace-Beltrami operator called the  $\rho$ -Laplacian for metric measure spaces in [7]. Together with Lu, they have shown that the spectrum of the  $\rho$ -Laplacian approximates the spectrum of a convolution Laplacian acting on vector bundles over metric measure spaces [8]. Lu also generalizes the work of Burago et al. to compact Riemannian manifolds with boundary and metric measure spaces with singularities of certain types in [13].

We build upon the methods implemented by Burago et al. and investigate the convergence of eigenvalues and eigenfunctions on an extended class of manifolds having a lower bound on the Ricci curvature and a lower bound on the volume of balls of small radius instead of considering absolute bounds of sectional curvature as done in [6]. Manifolds with a lower bound on Ricci curvature have many interesting geometric properties that are explained in detailed in a survey article by G. Wei [14]. Throughout this paper,  $M$  is a compact Riemannian manifold of dimension  $n$  without boundary. Let  $\mathcal{M} = \mathcal{M}(n, \lambda, D, i_0)$  be the class of such manifolds with the Ricci curvature and injectivity radius bounded below by  $\lambda$  and  $i_0$ , respectively, and diameter bounded above by  $D$ .

**Definition 1** ( $(\epsilon, \rho)$ -approximation). Given any  $\rho > \epsilon > 0$ , a finite graph  $\Gamma$  with the vertex set  $F = \{x_i\}_{i=1}^N$  and edges  $e_{ij}$  is called an  $(\epsilon, \rho)$ -approximation of  $(M, g)$  if the following conditions hold.

- $F \subset M$ . There exists a partition of  $M$  into measurable subsets  $\{V_i\}_{i=1}^N$  such that  $V_i \subset B_\epsilon(x_i)$  and  $M = \bigcup B_\epsilon(x_i)$  where  $B_\epsilon(x_i)$  is the ball in  $M$  centered at  $x_i$  with radius  $\epsilon$ .

- Two vertices  $x_i$  and  $x_j$  are connected by an edge  $e_{ij}$  if the Riemannian distance between them is less than  $\rho$ .
- *Measure on  $\Gamma$* : Let  $\mu_i = \text{vol}(V_i)$ .  $F$  can then be equipped with a discrete measure  $\mu = \sum_{i=1}^N \mu_i \delta_{x_i}$  where  $\delta_{x_i}$  denotes the Dirac measure at  $x_i$ .
- *Weights on edges*: Let  $\nu_n$  be the volume of the unit ball in Euclidean  $n$ -space. To an edge  $e_{ij}$ , assign the weight

$$w_{ij} = \frac{2(n+2)}{\nu_n \rho^{n+2}} \mu_i \mu_j. \quad (1.1)$$

For any  $M$  from the class  $\mathcal{M}(n, \lambda, D, i_0)$ , there exists a finite set of points  $F$  that is  $\epsilon$ -dense in  $M$  since  $M$  is compact. We can then use a partition obtained by the Voronoi decomposition to define an  $(\epsilon, \rho)$ -approximation of  $(M, g)$ . The weighted graph is denoted by  $\Gamma(F, \rho, \mu)$ . A weighted graph Laplacian  $\Delta_\Gamma$  on  $L^2(F)$  is defined to approximate the Laplace-Beltrami operator  $\Delta_M$  on  $(M, g)$  as follows:

$$\begin{aligned} (\Delta_\Gamma u)(x_i) &= \frac{1}{\mu_i} \sum_{x_i \sim x_j} w_{ij} (u(x_i) - u(x_j)) \\ &= \frac{2(n+2)}{\nu_n \rho^{n+2}} \sum_{x_i \sim x_j} \mu_i (u(x_i) - u(x_j)). \end{aligned} \quad (1.2)$$

The motivation behind the choice of the normalization constant is given in Section 2.3 of [6]. We note that  $-\Delta_\Gamma$  is a self-adjoint non-negative definite operator on  $L^2(F)$  with eigenvalues  $0 \leq \lambda_1(\Gamma) \leq \lambda_2(\Gamma) \leq \dots \leq \lambda_N(\Gamma)$  and the eigenvalues of  $-\Delta_M$  are denoted by  $0 \leq \lambda_1(M) \leq \lambda_2(M) \leq \lambda_3(M) \leq \dots$ . The main goal of this paper is to prove the following theorem.

**Theorem 1.1.** *Consider a manifold  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and an  $(\epsilon, \rho)$ -approximation  $\Gamma$  of  $M$  such that  $\rho < i_0$  and  $\frac{\epsilon}{\rho} < \frac{1}{n}$ . For  $0 < r < \frac{i_0}{2}$  and  $x \in M$ , assume that  $\text{vol}(B_r(x)) \geq \frac{\alpha_n}{2} \left(\frac{2r}{\pi}\right)^n$ . Let  $\lambda_k(M)$ ,  $\lambda_k(\Gamma)$  denote the  $k$ -th eigenvalues of  $-\Delta_M$  and  $-\Delta_\Gamma$  respectively. Then there exists  $\delta(\epsilon, \rho, \lambda, \lambda_k(M)) > 0$  such that for all positive integers  $k$ ,*

$$|\lambda_k(\Gamma) - \lambda_k(M)| \leq \delta(\epsilon, \rho, \lambda, \lambda_k(M)) \lambda_k(M)$$

where  $\delta(\epsilon, \rho, \lambda, \lambda_k(M))$  tends to zero as  $\epsilon, \rho$  and the ratio  $\frac{\epsilon}{\rho}$  approach zero. Moreover, if  $|\text{Ric}| \leq \lambda$ , then there exists a constant  $\mathcal{C}_{k, \mathcal{M}}$  such that

$$|\lambda_k(\Gamma) - \lambda_k(M)| \leq \mathcal{C}_{k, \mathcal{M}} \delta(\epsilon, \rho, \lambda, \mathcal{C}_{k, \mathcal{M}})$$

where  $\delta(\epsilon, \rho, \lambda, \mathcal{C}_{k, \mathcal{M}})$  tends to zero as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

For a detailed value of  $\delta(\epsilon, \rho, \lambda, \mathcal{C}_{k, \mathcal{M}})$ , we refer to equation (5.10). In [1], Micheal Anderson proved that the subclass of  $\mathcal{M}(n, \lambda, D, i_0)$  with  $|\text{Ric}| \leq \lambda$  is compact in the  $C^{1, \alpha}$ -topology. Using this result, we obtain a uniform bound  $\mathcal{C}_{k, \mathcal{M}}$  on the  $k$ -th eigenvalues of  $\Delta_M$  depending only on  $\mathcal{M}$  and  $k$  using. This constant plays a major role in showing uniform convergence of eigenvalues.

The motivation behind the lower bound of  $\text{vol}(B_r(x))$  is that Berger proved in [4] that there exists a uniform constant  $c_n$  such that for small metric balls,

$$\text{vol}(B_r(x)) \geq c_n \left(\frac{2r}{\pi}\right)^n. \quad (1.3)$$

Let  $\alpha_n$  denote the volume of unit  $n$ -sphere. There is a long-standing conjecture asserting that hemispheres minimize the Riemannian volume among small metric balls of a given dimension and intrinsic radius, i.e.  $c_n = \frac{\alpha_n}{2}$  is the optimal constant. Approximate values of  $c_n$  have been derived in [5] and [9]. The conjecture is open even for  $n = 2$ . However, in [10], using isoperimetric inequalities and variational arguments, a new constant  $c_2$  has been derived close to the optimal one. If the sectional curvature is bounded above by 1, then from Gunther's volume comparison it is easy to see that  $c_n = \frac{\alpha_n}{2}$ . If we consider the subclass  $\mathcal{M}(n, \lambda, K, D, i_0)$

of  $\mathcal{M}(n, \lambda, D, i_0)$  where the sectional curvatures of the manifolds are bounded above by  $K$ , we have the following corollary.

**Corollary 1.2.** *Consider a manifold  $M \in \mathcal{M}(n, \lambda, K, D, i_0)$  and an  $(\epsilon, \rho)$ -approximation  $\Gamma$  of  $M$  such that  $\rho < i_0$  and  $\frac{\epsilon}{\rho} < \frac{1}{n}$ . Let  $\lambda_k(M), \lambda_k(\Gamma)$  denote the  $k$ -th eigenvalues of  $-\Delta_M$  and  $-\Delta_\Gamma$  respectively. Then there exists  $\delta(\epsilon, \rho, \lambda, K, \lambda_k(M)) > 0$  such that for all positive integers  $k$ ,*

$$|\lambda_k(\Gamma) - \lambda_k(M)| \leq \delta(\epsilon, \rho, \lambda, K, \lambda_k(M)) \lambda_k(M)$$

where  $\delta(\epsilon, \rho, \lambda, K, \lambda_k(M))$  tends to zero as  $\epsilon, \rho$  and the ratio  $\frac{\epsilon}{\rho}$  approach zero. Moreover, if  $|\text{Ric}| \leq \lambda$ , then there exists a constant  $C_{k, \mathcal{M}}$  such that

$$|\lambda_k(\Gamma) - \lambda_k(M)| \leq C_{k, \mathcal{M}} \delta(\epsilon, \rho, \lambda, K, C_{k, \mathcal{M}})$$

where  $\delta(\epsilon, \rho, \lambda, K, C_{k, \mathcal{M}})$  tends to zero as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . The constant  $C_{k, \mathcal{M}}$  depends only on  $\mathcal{M}$  and  $k$ .

When the absolute value of the sectional curvature is bounded by  $K$ , the Jacobian  $J_x(v)$  of the exponential map at  $x$  in the direction of  $v$  satisfies

$$\frac{1}{1 + CnK|v|^2} \leq J_x(v) \leq 1 + CnK|v|^2$$

within the injectivity radius. This inequality is crucially used in [6]. However, these bounds do not translate to our setting. While they hint at a potential extension to manifolds with Ricci curvature bounds, the generalization was not immediate. We offer a detailed approach by adapting their  $(\epsilon, \rho)$ -approximation construction to our setting.

**Idea of the proof:** We use the Bishop-Gromov theorem and the uniform minimum injectivity radius  $i_0$  of manifolds in  $\mathcal{M}(n, \lambda, D, i_0)$  to derive an upper bound of the Jacobian determinant. While avoiding the reliance on sectional curvature, the methods in [6] can be adapted to our setting for establishing an upper bound of the eigenvalues. By deriving appropriate inequalities to bound the norm of the Dirichlet energy of the image of an  $L^2$  map in  $M$  under the discretization map  $P$ , we obtain an upper bound for the Rayleigh quotient. This can be done by restricting ourselves to the subspace spanned by the first  $k$  eigenvalues in  $H^1(M)$ , thus obtaining an approximation of the  $k$ -th eigenvalue of  $-\Delta_M$ .

To establish a lower bound for the eigenvalues, we regularize  $L^2$  maps on  $(\epsilon, \rho)$ -approximations as elements of  $C^{0,1}(M)$  through the interpolation map  $I$ . Using the lower bound on the volume of small balls, we derive a suitable lower bound of the image of the characteristic function on  $M$  under the associated integral operator and use the upper bound of the Jacobian determinant as obtained before to derive inequalities that constrain the Rayleigh quotient, thus, enabling us to estimate a bound for the  $k$ -th eigenvalue of  $-\Delta_M$ .

## 2. SOME INEQUALITIES AND AVERAGE DISPERSION

Given a Riemannian manifold  $(M, g)$  and a point  $x \in M$ , the exponential map at  $x$  is denoted by  $\exp_x$ . Let  $u$  be a unit vector in  $T_x M$ . We can choose unit vectors  $e_2, e_3, \dots, e_n$  such that  $\{u, e_2, e_3, \dots, e_n\}$  is an orthonormal basis in  $T_x M$ . Let  $\gamma$  be the geodesic defined by  $\gamma(t) = \exp_x(tu)$ . We choose Jacobi fields  $J_2, J_3, \dots, J_n$  along  $\gamma$  such that for all  $i \in \{2, 3, \dots, n\}$ ,

$$J_i(0) = 0, \quad J'_i(0) = e_i.$$

For  $t < i_0$ , using polar co-ordinates  $(u, t)$  in  $T_x M$ , we have,

$$\exp_x^* \text{vol}_g = J(u, t) t^{n-1} dt du$$

where  $J(u, t) = \frac{1}{t^{n-1}} \sqrt{\det(\langle J_i(t), J_k(t) \rangle_g)}$  is called the Jacobian determinant at  $(u, t)$ .

In the following lemma, we derive an upper bound for the Jacobian determinant, relying on the lower bound of Ricci curvature, dimension, and injectivity radius of the manifold in consideration.

**Lemma 2.1.** *Let  $(M, g)$  be an  $n$ -dimensional Riemannian manifold with Ricci curvature bounded below by  $\lambda$ . Then for any  $x \in M$  and a unit tangent vector  $u \in T_x M$ ,*

$$J(u, t) \leq J_\lambda(t), \quad \forall t < i_0(M)$$

where  $J_\lambda(t) = \left(1 + \sum_{j \in \mathbb{N}} \frac{(\sqrt{|\frac{\lambda}{n-1}|}t)^{2j}}{(2j+1)!}\right)^{n-1}$  and  $i_0(M)$  is the injectivity radius of  $M$ .

*Proof.* We will now compute an upper bound of  $J(u, t)$  by using the techniques of Bishop's volume comparison theorem given in Theorem 3.101[11]. For notational convenience, let  $a = \frac{\lambda}{n-1}$ . Then for a fixed  $u \in S^{n-1}$  and  $t < i_0$ ,

$$\frac{1}{J(u, t)} \frac{d}{ds} J(u, s) \Big|_{s=t} \leq \begin{cases} (n-1) \left(\sqrt{a} \cot \sqrt{as} - \frac{1}{s}\right), & \text{if } a > 0 \\ 0 & \text{if } a = 0 \\ (n-1) \left(\sqrt{-a} \coth \sqrt{-as} - \frac{1}{s}\right), & \text{if } a < 0. \end{cases}$$

Integrating the inequalities from  $s$  to  $t$  and taking exponentials, we obtain the following inequalities as limit  $s \rightarrow 0^+$ .

$$J(u, t) \leq \begin{cases} \left(\frac{\sin \sqrt{at}}{\sqrt{at}}\right)^{n-1} & \text{if } a > 0 \\ 1 & \text{if } a = 0 \\ \left(\frac{\sinh \sqrt{-at}}{\sqrt{-at}}\right)^{n-1} & \text{if } a < 0. \end{cases}$$

By using Taylor expansion in each case, we conclude that

$$J(u, t) \leq J_\lambda(t),$$

where  $J_\lambda(t) = \left(1 + \sum_{j \in \mathbb{N}} \frac{(\sqrt{|\frac{\lambda}{n-1}|}t)^{2j}}{(2j+1)!}\right)^{n-1}$ . Clearly,  $J_\lambda$  is a monotone increasing function and  $J_\lambda(t) \rightarrow 1$  as  $t \rightarrow 0$ .  $\square$

**Definition 2.** Let  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and  $B_r(x)$  be a ball in  $M$  centered at  $x$  of radius  $r$ . Let  $f : M \rightarrow \mathbb{R}$  be a smooth function. The average dispersion  $E_r(f) : L^2(M) \rightarrow \mathbb{R}$  is defined as

$$E_r(f) = \int_M \int_{B_r(x)} |f(y) - f(x)|^2 dy dx.$$

We recall that  $H^1(M)$  is the space of all functions in  $L^2(M)$  such that their partial derivatives are also in  $L^2(M)$ . The following lemma gives an upper bound of  $E_r(f)$ .

**Lemma 2.2.** *For  $M \in \mathcal{M}(n, \lambda, D, i_0)$ , let  $0 < r < \frac{i_0}{2}$ ,  $f \in H^1(M)$ . Then*

$$E_r(f) \leq J_\lambda(r) \frac{\nu_n r^{n+2}}{n+2} \|df\|^2.$$

*Proof.* Since smooth functions are dense in  $H^1(M)$ , it is enough to prove the above inequality for smooth functions. For any  $r < \frac{i_0}{2}$ , the exponential map at  $x$  restricted to  $B_r(x)$  is a diffeomorphism onto its image. Using polar co-ordinates  $(u, t)$  on  $T_x M$ ,

$$\int_{B_r(x)} |f(y) - f(x)|^2 dy = \int_{u \in S^{n-1}} \int_{t=0}^r |f(\exp_x(tu)) - f(\exp_x(0))|^2 J(u, t) t^{n-1} dt du. \quad (2.1)$$

Substituting the upper bound of  $J(u, t)$  from Lemma 2.1 in equation (2.1),

$$\int_{B_r(x)} |f(y) - f(x)|^2 dy \leq J_\lambda(r) \int_{u \in S^{n-1}} \int_{t=0}^r |f(\exp_x(tu)) - f(\exp_x(0))|^2 t^{n-1} dt du.$$

Now,

$$\int_{u \in S^{n-1}} \int_{t=0}^r |f(\exp_x(tu)) - f(\exp_x(0))|^2 t^{n-1} dt du = \int_{B_r(0) \subset T_x M} |f(\exp_x(v)) - f(x)|^2 dv.$$

By using the techniques for establishing an upper bound of the same integral in Lemma 3.3 ([6]), we have

$$\int_{B_r(0) \subset T_x M} |f(\exp_x(v)) - f(x)|^2 dv \leq \frac{\nu_n}{n+2} r^{n+2} \|df\|_{L^2}^2.$$

Combining this with the upper bound of the Jacobian determinant, the required inequality is obtained.  $\square$

We establish a Poincaré-type inequality bounding the integral of the difference between a function's squared deviation from its mean by the average dispersion.

**Lemma 2.3.** *For  $0 < \epsilon < r < \frac{i_0}{2}$ , let  $V$  be a measurable subset of  $M$  with  $\text{diam}V \leq \epsilon$  such that  $\text{vol}V = \mu > 0$ . For all  $x \in M$ , let  $\text{vol}(B_r(x)) \geq c_n r^n$ . Let  $f \in L^2(M)$  and  $a = \frac{1}{\mu} \int_V f(x) dx$  be the integral mean of  $f|_V$ . Then*

$$\int_V |f(x) - a|^2 dx \leq \frac{1}{c_n (r - \epsilon)^n} E_r(f, V).$$

*Proof.* The above inequality is obtained in Lemma 3.4 in [6] where an upper bound on the sectional curvature is used to obtain a lower bound on  $\text{vol}(B_r(x))$ . If  $\text{vol}(B_r(x)) \geq c_n r^n$ , then the proof follows using the same techniques.  $\square$

### 3. UPPER BOUND FOR $\lambda_k(\Gamma)$

We find an upper bound of the eigenvalues of  $-\Delta_\Gamma$  in terms of the eigenvalues of corresponding manifolds  $M \in \mathcal{M}(n, \lambda, D, i_0)$  using the inequalities proved in the previous section. consider  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and  $\epsilon > 0$  such that  $0 < \epsilon < \rho < \frac{i_0}{2}$ . Let  $\Gamma$  be an  $(\epsilon, \rho)$ -approximation of  $(M, \text{vol})$  using a decomposition  $\{V_i\}_1^N$ . We define the discretization map that assigns to each vertex of the graph the average value of the function on the partition containing the vertex.

**Definition 3** (Discretization map). Let  $\mu_i = \text{vol}(V_i)$ . Then  $P : L^2(M) \rightarrow L^2(F)$  be a map defined as

$$Pf(x_i) = \frac{1}{\mu_i} \int_{V_i} f d\mu.$$

The map  $P^* : L^2(F) \rightarrow L^2(M)$  is defined as

$$P^*(u) = \sum_{i=1}^N u(x_i) \chi_{V_i}.$$

We see some immediate properties of the discretization map in the following lemma.

**Lemma 3.1.** *Let  $M \in \mathcal{M}(n, \lambda, D, i_0)$ ,  $f \in L^2(M)$  and  $u \in L^2(F)$ . Then*

- (1)  $P : L^2(M) \rightarrow L^2(F)$  is a bounded linear operator :  $\|Pf\| \leq N \|f\|_2$ .
- (2)  $P^*$  is norm-preserving, i.e.  $\|P^*(u)\|_{L^2(M)} = \|u\|_{L^2(F)}$ .
- (3)  $P^*$  is the adjoint of discretization map  $P$ , i.e.  $\langle f, P^*(u) \rangle_{L^2(M)} = \langle P(f), u \rangle_{L^2(F)}$ .

**Lemma 3.2.**

$$\|f - P^*Pf\|_{L^2}^2 \leq C(n) J_\lambda(\rho) \epsilon^2 \|df\|_{L^2}^2.$$

where  $C(n) = \frac{3(n+1)\nu_n}{c_n}$ .

*Proof.* Let  $V$  be a measurable subset of  $M$  and  $\chi_V$  denote the characteristic function on  $V$ . According to definition,

$$\begin{aligned} \|f - P^*Pf\|_{L^2}^2 &= \int_M |f(x) - P^*Pf(x)|^2 dx \\ &= \int_M |f(x) - \sum_{i=1}^N Pf(x_i)\chi_{V_i}(x)|^2 dx \\ &= \sum_{i=1}^N \int_{V_i} |f(x) - Pf(x_i)|^2 dx. \end{aligned}$$

From Lemma 2.3, for all  $r$  such that  $0 < \epsilon < r < 2\rho$  and for all  $1 \leq i \leq N$ ,

$$\int_{V_i} |f(x) - Pf(x_i)|^2 dx \leq \frac{1}{c_n(r-\epsilon)^n} E_r(f, V_i).$$

From Lemma 2.2,

$$\|f - P^*Pf\|_{L^2}^2 \leq \frac{\nu_n}{c_n(r-\epsilon)^n} E_r(f) \leq \frac{J_\lambda(r)\nu_n}{c_n(n+2)} \left(\frac{r}{r-\epsilon}\right)^n r^2 \|df\|_{L^2}^2.$$

Putting  $r = (n+1)\epsilon$ , we have  $(\frac{r}{r-\epsilon})^n = (1 + \frac{1}{n})^n < 3$ . Hence,

$$\|f - P^*Pf\|_{L^2}^2 \leq (n+1)\epsilon^2 \frac{3J_\lambda(\rho)\nu_n}{c_n} \|df\|_{L^2}^2 = C(n)J_\lambda(\rho)\epsilon^2 \|df\|_{L^2}^2.$$

where  $C(n) = \frac{3(n+1)\nu_n}{c_n}$ . □

We emphasize that throughout the paper, we use the notation  $C(n)$  to denote a constant depending on  $n$  and its value can change based on the inequalities derived. The above lemma implies the following straight-forward corollary.

**Corollary 3.3.**

$$|\|Pf\| - \|f\|_{L^2}|^2 \leq C(n)J_\lambda(\rho)\epsilon^2 \|df\|_{L^2}^2.$$

*Proof.* The above statement clearly follows from the triangle inequality and norm-preserving property of  $P^*$ :

$$|\|Pf\| - \|f\|_{L^2}| \leq |\|P^*Pf\| - \|f\|| \leq \sqrt{\frac{3J_\lambda(\rho)\nu_n}{c_n}(n+1)\epsilon} \|df\|_{L^2}.$$

□

The next lemma gives an upper bound of the discrete Dirichlet energy at an image of a point under discretization map using norm of derivative of that point.

**Definition 4** (Discrete Dirichlet energy). Let  $\Gamma = \Gamma(F, \rho, \mu)$  be an  $(\epsilon, \rho)$ - approximation of  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . For  $u \in L^2(F)$ , the Dirichlet energy on  $\Gamma$  is given by

$$\|\delta(u)\|^2 = \frac{n+2}{\nu_n \rho^{n+2}} \sum_i \sum_{j: x_i \sim x_j} \mu_i \mu_j |u(x_j) - u(x_i)|^2. \quad (3.1)$$

**Lemma 3.4.** Let  $\Gamma$  be a  $(\epsilon, \rho)$ -approximation of  $(M, \text{vol})$ . For  $u \in L^2(F)$ ,

$$\|\delta(Pf)\|^2 \leq J_\lambda(\rho + 2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2} \|df\|_{L^2}^2.$$

*Proof.* From Lemma 4.3(2) in [6], we derive

$$\begin{aligned} \|\delta(Pf)\|^2 &\leq \frac{n+2}{\nu_n \rho^{n+2}} \sum_i \sum_{j: x_i \sim x_j} \int_{V_i} \int_{V_j} |f(y) - f(x)|^2 dy dx \\ &= \frac{n+2}{\nu_n \rho^{n+2}} \int_M \int_{U(x)} |f(y) - f(x)|^2 dy dx. \end{aligned} \quad (3.2)$$

where  $U(x) = \bigcup_{j: x_j \sim x_i} V_j$  in (3.2). We also observe that  $U(x) \subseteq B_{\rho+2\epsilon}(x)$ . Hence,

$$\|\delta(Pf)\|^2 \leq \frac{n+2}{\nu_n \rho^{n+2}} E_{\rho+2\epsilon}(f). \quad (3.3)$$

Using Lemma 2.2 and from (3.3), we have

$$\begin{aligned} \|\delta(Pf)\|^2 &\leq J_\lambda(\rho+2\epsilon) \frac{(\rho+2\epsilon)^{n+2}}{\rho^{n+2}} \|df\|_{L^2}^2 \\ &= J_\lambda(\rho+2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2} \|df\|_{L^2}^2. \end{aligned}$$

□

Using the Rayleigh quotient, one can calculate an upper bound of the graph eigenvalue of the graph Laplacian operator using the eigenvalues of Laplace-Beltrami operator as follows.

**Theorem 3.5.** *Let  $0 < \epsilon < \rho < \frac{i_0}{2}$ ,  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and  $\Gamma$  be a  $(\epsilon, \rho)$ -approximation of  $(M, \text{vol})$ . Let  $\lambda_k = \lambda_k(M)$  denote the  $k$ -th eigenvalue of  $-\Delta_M$  and  $\lambda_k(\Gamma)$  denote the  $k$ -th eigenvalue  $-\Delta_\Gamma$ . For sufficiently small  $\epsilon, \rho$  let  $C(n)J_\lambda(\rho)\lambda_k\epsilon^2 < 1$ . Then,*

$$\lambda_k(\Gamma) \leq (1 + \delta_1(\epsilon, \rho, \lambda, \lambda_k))\lambda_k$$

where  $\delta_1(\epsilon, \rho, \lambda, \lambda_k) = \frac{J_\lambda(\rho+2\epsilon)(1+\frac{2\epsilon}{\rho})^{n+2}}{1-\epsilon\sqrt{C(n)J_\lambda(\rho)\lambda_k}} - 1 \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

*Proof.* To prove the theorem we find a linear subspace  $L \subset L^2(X)$  with  $\dim L = k$  such that the Rayleigh quotient restricted to  $L$  satisfies the following inequality,

$$\sup_{u \in L \setminus \{0\}} \frac{\|\delta u\|^2}{\|u\|^2} \leq (1 + \delta_1(\epsilon, \rho, \lambda, \lambda_k))\lambda_k$$

such that  $\delta_1(\epsilon, \rho, \lambda) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . Then the theorem follows from the min-max principle.

Let  $W \subset H^1(M)$  be the linear span of orthonormal eigenfunctions of  $-\Delta_M$  corresponding to eigenvalues  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k$ . For any  $f \in W$ ,

$$\|df\|_{L^2}^2 = \langle -\Delta_M f, f \rangle \leq \lambda_k \|f\|_{L^2}^2.$$

From Corollary 3.3,

$$\|Pf\| \geq \|f\|_{L^2} - \epsilon C(n) \sqrt{J_\lambda(\rho)} \|df\|_{L^2} \geq \left(1 - \epsilon C(n) \sqrt{J_\lambda(\rho)\lambda_k}\right) \|f\|_{L^2}. \quad (3.4)$$

Hence,  $P|_W$  is injective when  $C(n)J_\lambda(\rho)\lambda_k\epsilon^2 < 1$ .

We consider  $\epsilon$  to be small enough such that  $P|_W$  is injective. Let  $L = P(W)$ . Then  $\dim L = k$ . We pick  $u \in L \setminus \{0\}$  and let  $f \in W$  be such that  $u = Pf$ . Then from equation (3.4),

$$\|u\|^2 \geq \left(1 - \epsilon C(n) \sqrt{J_\lambda(\rho)\lambda_k}\right) \|f\|_{L^2}^2. \quad (3.5)$$

By Lemma 3.4,

$$\|\delta(u)\|^2 \leq J_\lambda(\rho+2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2} \lambda_k \|f\|_{L^2}^2. \quad (3.6)$$

From inequalities (3.5) and (3.6), we have

$$\begin{aligned} \frac{\|\delta u\|^2}{\|u\|^2} &\leq \frac{J_\lambda(\rho+2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2} \lambda_k \|f\|_{L^2}^2}{\left(1 - \epsilon C(n) \sqrt{J_\lambda(\rho)\lambda_k}\right) \|f\|_{L^2}^2} \\ &= 1 + \delta_1(\epsilon, \rho, \lambda, \lambda_k) \lambda_k \end{aligned} \quad (3.7)$$

where  $\delta_1(\epsilon, \rho, \lambda, \lambda_k) = \frac{J_\lambda(\rho+2\epsilon)(1+\frac{2\epsilon}{\rho})^{n+2}}{1-\epsilon C(n)\sqrt{J_\lambda(\rho)\lambda_k}} - 1 \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$  which concludes the proof. □

**Remark 3.6.** In [6], the upper bound of the graph eigenvalue was obtained using the upper bound of the absolute value of sectional curvature. But we observe in Theorem 3.7 that the upper bound of the graph eigenvalue can be obtained using only the lower bound on the Ricci curvature.

#### 4. SMOOTHING OPERATOR

In this section, we study the smoothing operator defined in [6] and its properties which will help us to obtain a lower bound for the eigenvalues of  $-\Delta_\Gamma$  using eigenvalues of  $-\Delta_M$ .

**Definition 5.** Fixing a positive  $r < \rho < \frac{i_0}{2}$ , the following functions are defined:

- (1) **Smoothing function.** Let  $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be defined as

$$\psi(t) = \begin{cases} \frac{n+2}{2\nu_n}(1-t^2) & \text{if } 0 \leq t \leq 1 \\ 0 & \text{if } t \geq 1. \end{cases}$$

- (2) **Kernel.** Let  $k_r : M \times M \rightarrow \mathbb{R}_+$  be defined as

$$k_r(x, y) = r^{-n} \psi(r^{-1}d(x, y)).$$

where  $d(x, y)$  denotes the Riemannian distance between  $x$  and  $y$  in  $M$ .

- (3) **Associated integral operator.** Let  $\Lambda_r^0 : L^2(M) \rightarrow C^{0,1}(M)$  be given by

$$\Lambda_r^0 f(x) = \int_M f(y) k_r(x, y) dy.$$

- (4) Let  $\theta \in C^{0,1}(M)$  be defined as

$$\theta = \Lambda_r^0(\mathbf{1}_M).$$

Using the polar co-ordinate in  $\mathbb{R}^n$ , the next lemma follows easily.

**Lemma 4.1.** Let  $p \in \mathbb{R}^n$  and  $x, y \in M$ . Let  $\psi, k_r$  be defined as above. Then,

- (1)  $\int_{\mathbb{R}^n} \psi(|p|) dt = 1$ , and  
(2)  $|k_r(x, y)| \leq \left(\frac{n+2}{2}\right) \frac{1}{\nu_n r^n}$ .

We derive the following approximations for  $\theta$ .

**Lemma 4.2.** For  $0 < r < \frac{i_0}{2}$  and  $M \in \mathcal{M}(n, \lambda, D, i_0)$ , let  $x \in M$  and  $B_r(x)$  be a ball of radius  $r$  centered at  $x$ .

- (1) Then  $\theta(x)$  has the following bounds:

$$C(n) \leq \theta(x) \leq J_\lambda(r), \tag{4.1}$$

where  $C(n)$  is a constant depending on  $n$ .

- (2) If  $\text{vol}(B_r(x)) \geq \frac{\alpha_n}{2} \left(\frac{2r}{\pi}\right)^n$ , then

$$1 - C(n)f_\lambda(r) \leq \theta(x) \leq J_\lambda(r) \tag{4.2}$$

where  $f_\lambda(r) = J_\lambda(r) - 1 \rightarrow 0$  as  $r \rightarrow 0$  and  $C(n)$  is a constant depending on  $n$ .

*Proof.* (1) The upper bound for  $\theta$  follows from the upper bound of the Jacobian calculated in Lemma 2.1 and the value of integral of  $\psi$  obtained from Lemma 4.1(1). To derive the lower limit of  $\theta$ , we only use the uniform lower bound of the injectivity radius of  $\mathcal{M}(n, \lambda, D, i_0)$ .



Let  $x \in M$ . Since  $\left(1 - \frac{t^2}{r^2}\right)$  is a decreasing function, for  $0 < \xi < 1$ ,

$$\begin{aligned} \theta(x) &= \frac{n+2}{2\nu_n r^n} \int_{t=0}^r \int_{u \in S^{n-1}} \left(1 - \frac{t^2}{r^2}\right) J(u, t) t^{n-1} du dt \\ &> \frac{n+2}{2\nu_n r^n} \int_{t=0}^{r\xi} \int_{u \in S^{n-1}} \left(1 - \frac{r^2 \xi^2}{r^2}\right) J(u, t) t^{n-1} du dt \\ &= \frac{n+2}{2\nu_n r^n} (1 - \xi^2) \text{vol}(B_\xi(x)) \\ &\geq C(n)(1 - \xi^2) \xi^n r^n \end{aligned} \quad (4.3)$$

where (4.3) follows from (1.3).

We observe that local maximum of  $(1 - \xi^2) \xi^n$  in  $(0, 1)$  is attained at  $\xi = \sqrt{\frac{n}{n+2}}$ . Substituting in (4.3), we obtain a lower bound  $C(n)$  of  $\theta(x)$  depending only on  $n$ .

Hence,

$$C(n) \leq \theta(x) \leq 1 + J_\lambda(r).$$

(2) The upper bound for  $\theta$  follows from Lemma 2.1 and the Lemma 4.1(1) as above. The lower bound under the volume constraint can be obtained as follows:

$$\begin{aligned} \theta(x) &= \frac{n+2}{2\nu_n r^n} \int_{t=0}^r \int_{u \in S^{n-1}} \left(1 - \frac{t^2}{r^2}\right) J(u, t) t^{n-1} du dt \\ &\geq \frac{n+2}{2\nu_n r^n} \int_{t=0}^r \int_{u \in S^{n-1}} J(u, t) t^{n-1} du dt - J_\lambda(r) \frac{n+2}{2\nu_n r^n} \int_{t=0}^r \int_{u \in S^{n-1}} \frac{t^{n+1}}{r^2} du dt \\ &= \frac{n+2}{2\nu_n r^n} \text{vol}(B_r(x)) - J_\lambda(r) \frac{n}{2} \\ &\geq \frac{n+2}{2\nu_n r^n} \frac{\alpha_n}{2} \left(\frac{2r}{\pi}\right)^n - \frac{n}{2} + \frac{n}{2} (1 - J_\lambda(r)) \\ &= \frac{n+2}{4} \left(\frac{\alpha_n}{\nu_n}\right) \left(\frac{2}{\pi}\right)^n - \frac{n}{2} (1 + f_\lambda(r)) \end{aligned} \quad (4.4)$$

$$\geq 1 - C(n) f_\lambda(r) \quad (4.5)$$

where  $f_\lambda(r) = J_\lambda(r) - 1 \rightarrow 0$  as  $r \rightarrow 0$  in (4.4) and (4.5) follows from the Bishop-Gromov volume comparison theorem.  $\square$

**Lemma 4.3.** *Let  $0 < r < \rho < \frac{i_0}{2}$  and  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . Let  $\nabla\theta$  denote the gradient of  $\theta$ . Then*

$$|\nabla\theta(x)| \leq nr\rho^2 f_\lambda(\rho).$$

where  $f_\lambda(\rho) = J_\lambda(\rho) - 1 \rightarrow 0$ .

*Proof.* One can compute the gradient of  $\theta$  in polar co-ordinates as follows:

$$\nabla\theta(x) = \frac{n+2}{\nu_n r^{n+2}} \int_{t=0}^r \int_{u \in S^{n-1}} tu J(u, t) t^{n-1} du dt.$$

We want to show that  $\int_{u \in S^{n-1}} u du = 0$ .

Let  $A : S^{n-1} \rightarrow S^{n-1}$  be the antipodal map defined as  $A(u) = -u$ . We note that the Jacobian matrix corresponding to  $A$  is  $-I$ , where  $I$  is the identity matrix. By changing co-ordinates using  $A$ , we can show that

$$\int_{u \in S^{n-1}} u du = 0. \quad (4.6)$$

From equation (4.6),

$$\int_{u \in S^{n-1}} \int_{t=0}^r t^n u du dt = \int_{u \in S^{n-1}} u du \int_{t=0}^r t^n dt = 0.$$

Hence,

$$\int_{u \in S^{n-1}} \int_{t=0}^r tJ(u, t)t^{n-1}ududt = \int_{u \in S^{n-1}} \int_{t=0}^r t^n(J(u, t) - 1)ududt$$

Using the upper bound of  $J(u, t)$  from Lemma 2.1, we have

$$\begin{aligned} J(u, t) - 1 &\leq J_\lambda(t) - 1 \\ &= \left(1 + \sum_{j \in \mathbb{N}} \frac{|\frac{\lambda}{n-1}|^j t^{2j}}{(2j+1)!}\right)^{n-1} - 1 \\ &= t^2 f_\lambda(t) \end{aligned}$$

where  $f_\lambda(t) = J_\lambda(t) - 1 \rightarrow 0$  as  $t \rightarrow 0$ .

Let  $w \in T_x M$ ,  $|w| = 1$  be such that  $\max\{\langle \nabla \theta(x), v \rangle | v \in T_x M, |v| = 1\}$  is attained at  $w$ . One can then compute the gradient of  $\theta$  in polar co-ordinates as follows:

$$\begin{aligned} |\nabla \theta(x)| &= \langle \nabla \theta, w \rangle = \frac{n+2}{\nu_n r^{n+2}} \int_{u \in S^{n-1}} \int_{t=0}^r t^n \langle u, w \rangle (J(u, t) - 1) dt du \\ &\leq \rho^2 f_\lambda(\rho) \frac{n+2}{\nu_n r^{n+2}} \int_{u \in S^{n-1}} \int_{t=0}^r t^{n+2} dt du \\ &\leq nr \rho^2 f_\lambda(\rho). \end{aligned}$$

□

Using the associated integral operator and  $\theta$ , we define the following function.

**Definition 6.** Let  $\Lambda_r : L^2(M) \rightarrow C^{0,1}(M)$  be defined as  $\Lambda_r f = \theta^{-1} \Lambda_r^0 f$ .

We observe that  $\Lambda_r f = f$ , for all constant functions  $f$ . In fact,  $\Lambda_r f$  is bounded when  $r$  is less than the injectivity radius as we see in the following lemma.

For uniformity of notation in the following theorems, we denote the lower bound of  $\theta(x)$  by  $C_\theta(r)$  in the subsequent results. The next lemma gives a bound on the  $L^2$  norm of the gradient of  $\theta$ .

**Lemma 4.4.** Let  $0 < r < \frac{i_0}{2}$  and  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . For all  $f \in L^2(M)$ , we have

$$\|\Lambda_r f\|_{L^2} \leq \frac{J_\lambda(r)}{C_\theta(r)} \|f\|_{L^2}$$

where  $C_\theta(r)$  is the lower bound of  $\theta(x)$ .

*Proof.* Let  $x \in M$ . Then

$$\begin{aligned} |\Lambda_r^0 f(x)|^2 &= \left| \int_M f(y) k_r(x, y) dy \right|^2 \\ &\leq \left( \int_M (\sqrt{k_r(x, y)})^2 dy \right) \left( \int_M (f(y) \sqrt{k_r(x, y)})^2 dy \right) \\ &= \theta(x) \int_M |f(y)|^2 k_r(x, y) dy. \end{aligned} \tag{4.7}$$

where (4.7) follows from Cauchy-Schwarz inequality.

So,

$$\begin{aligned} |\Lambda_r f(x)|^2 &= \theta(x)^{-2} |\Lambda_r^0 f(x)|^2 \\ &\leq \theta(x)^{-1} \int_M |f(y)|^2 k_r(x, y) dy \\ &\leq C_\theta(r)^{-1} \int_M |f(y)|^2 k_r(x, y) dy. \end{aligned}$$

Integrating over  $M$ ,

$$\begin{aligned} \|\Lambda_r f\|_{L^2}^2 &= \int_M |\Lambda_r f(x)|^2 dx \leq C_\theta(r)^{-1} \int_M |f(y)|^2 \int_M k_r(x, y) dy dx \\ &\leq C_\theta(r)^{-1} J_\lambda(r) \|f\|_{L^2}^2. \end{aligned} \quad (4.8)$$

where equation (4.8) follows from Lemma 4.2.

Hence,  $\|\Lambda_r f\|_{L^2} \leq C_\theta(r)^{-1} J_\lambda(r) \|f\|_{L^2}$ .  $\square$

When  $r$  is less than the injectivity radius, the following lemma gives an upper bound using average dispersion on how far  $\Lambda_r$  takes a function in  $L^2(M)$ .

**Lemma 4.5.** *Let  $0 < r < \frac{i_0}{2}$  and  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . For all  $f \in L^2(M)$ ,*

$$\|\Lambda_r f - f\|_{L^2}^2 \leq \frac{C(n)}{C_\theta(r)r^n} E_r(f),$$

where  $C(n) = \frac{n+2}{2\nu_n}$  and  $\nu_n$  is the volume of the unit ball in Euclidean  $n$ -space.

*Proof.* We fix a coset representative  $f : M \rightarrow \mathbb{R}$  of an element of  $L^2(M)$ .

Let  $a = f(x)$ . Now,

$$\begin{aligned} \Lambda_r f(x) - f(x) &= \Lambda_r f(x) - a \\ &= \Lambda_r(f - a \mathbf{1}_M)(x) \\ &= \theta^{-1}(x) \int_{B_r(x)} (f(y) - a) k_r(x, y) dy \\ &= \theta^{-1}(x) \int_{B_r(x)} (f(y) - f(x)) k_r(x, y) dy. \end{aligned}$$

By Cauchy-Schwarz inequality,

$$\begin{aligned} |\Lambda_r f(x) - f(x)|^2 &\leq \theta^{-2}(x) \left( \int_{B_r(x)} k_r(x, y) dy \right) \left( \int_{B_r(x)} |f(y) - f(x)|^2 k_r(x, y) dy \right) \\ &= \theta^{-1}(x) \int_{B_r(x)} |f(y) - f(x)|^2 k_r(x, y) dy \\ &\leq \frac{n+2}{2\nu_n C_\theta(r) r^n} \int_{B_r(x)} |f(y) - f(x)|^2 dy \end{aligned} \quad (4.9)$$

where (4.9) follows from Lemma 4.1.

Finally, by integrating on both sides, we get the result.  $\square$

In the following lemma, we obtain an upper bound for the norm of  $d\Lambda_r(f)$  using average dispersion. We are essentially referring to [6] for the proof of the lemma and generalizing it to the class of manifolds with bounded Ricci curvatures instead of sectional curvature.

**Lemma 4.6.** *Let  $0 < r < \rho < \frac{i_0}{2}$  and  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . For every  $f \in L^2(M)$ ,*

$$\|d(\Lambda_r f)\| \leq \sqrt{\frac{n+2}{\nu_n r^{n+2}}} g_\lambda(r, \rho) \sqrt{E_r f}$$

where  $g_\lambda(r, \rho) = \frac{J_\lambda(r)}{C_\theta(r)} + \frac{\rho^2 f_\lambda(\rho) n r^2}{C_\theta(r)^{\frac{3}{2}}}$ .

*Proof.* Fixing a coset representative of  $L^2(M)$  and denoting it by  $f$ , let  $\tilde{f} = \Lambda_r f$ . For any  $a \in \mathbb{R}$ ,

$$\tilde{f}(x) = a + \theta^{-1}(x) \int_{B_r(x)} (f(y) - a) k_r(x, y) dy.$$

Using chain rule and differentiating  $\tilde{f}$  we get,

$$d_x \tilde{f} = \theta^{-1}(x) \int_{B_r(x)} (f(y) - a) d_x k_r(\cdot, y) dy + d_x(\theta^{-1}) \int_{B_r(x)} (f(y) - a) k_r(x, y) dy.$$

Putting  $a = f(x)$ , we have

$$d_x \tilde{f} = \theta(x)^{-1} A_1(x) + A_2(x). \quad (4.10)$$

Here,

$$A_1(x) = \int_{B_r(x)} (f(y) - f(x)) d_x k_r(\cdot, y) dy$$

and

$$A_2(x) = d_x(\theta^{-1}) \int_{B_r(x)} (f(y) - f(x)) k_r(x, y) dy.$$

From Lemma 4.2, and using triangle inequality,

$$\|d\tilde{f}\|_{L^2} \leq C_\theta(r)^{-1} \|A_1\|_{L^2} + \|A_2\|_{L^2}.$$

We evaluate each integral  $A_1$  and  $A_2$  separately to get our desired inequality.

Using Cauchy-Schwarz inequality,

$$\begin{aligned} |A_2(x)|^2 &\leq |d_x(\theta^{-1})|^2 \left( \int_{B_r(x)} k_r(x, y) dy \right) \left( \int_{B_r(x)} |f(y) - f(x)|^2 k_r(x, y) dy \right) \\ &\leq |d_x(\theta^{-1})|^2 \theta(x) \int_{B_r(x)} |f(y) - f(x)|^2 k_r(x, y) dy \\ &\leq \frac{n+2}{2\nu_n r^n} \frac{|d_x \theta|^2}{\theta^3} \int_{B_r(x)} |f(y) - f(x)|^2 dy \\ &\leq \frac{n+2}{2\nu_n r^n} \frac{(\rho^2 f_\lambda(\rho) nr)^2}{C_\theta(r)^3} \int_{B_r(x)} |f(y) - f(x)|^2 dy. \end{aligned}$$

Integrating on  $M$ ,

$$\|A_2\|_{L^2} \leq \frac{\rho^2 f_\lambda(\rho) nr^2}{C_\theta(r)^{\frac{3}{2}}} \sqrt{\frac{n+2}{\nu_n r^{n+2}}} E_r(f). \quad (4.11)$$

On the other hand, fixing  $x \in M$ , let  $w \in T_x M$  be such that  $\max\{|\langle A_1(x), v \rangle| : v \in T_x M, |v| = 1\}$  is attained at  $w$ .

So,

$$\begin{aligned} |A_1(x)| &= \langle A_1(x), w \rangle \\ &= \frac{n+2}{\nu_n r^{n+2}} \int_{B_r(x)} (f(y) - f(x)) \langle \exp^{-1}(y), w \rangle dy \\ &= \frac{n+2}{\nu_n r^{n+2}} \int_{t=0}^r \int_{u \in S^{n-1}} (f(\exp_x(tu)) - f(\exp_x(0))) \langle tu, w \rangle t^{n-1} dudt. \end{aligned}$$

By Cauchy-Schwartz inequality,

$$|A_1(x)|^2 \leq \left( \frac{n+2}{\nu_n r^{n+2}} \right)^2 \left( \int_{B_r(0)} |f(\exp_x(tu)) - f(\exp_x(0))|^2 J(u, t)^2 t^{n-1} dudt \right) \left( \int_{B_r(0)} \langle tu, w \rangle^2 t^{n-1} dudt \right).$$

Now,  $\frac{n+2}{\nu_n r^{n+2}} \int_{B_r(0)} \langle tu, w \rangle^2 t^{n-1} dudt = 1$ . Hence,

$$\begin{aligned} |A_1(x)|^2 &\leq \frac{n+2}{\nu_n r^{n+2}} \int_{B_r(0)} |f(\exp_x(tu)) - f(\exp_x(0))|^2 J(u, t)^2 t^{n-1} dudt \\ &= \frac{n+2}{\nu_n r^{n+2}} \int_{B_r(x)} |f(y) - f(x)|^2 J_x(\exp_x^{-1}(y)) dy \\ &\leq J_\lambda(r) \frac{n+2}{\nu_n r^{n+2}} \int_{B_r(x)} |f(y) - f(x)|^2 dy. \end{aligned}$$

Integrating on  $M$ ,

$$\|A_1\|_{L^2} \leq \sqrt{\frac{n+2}{\nu_n r^{n+2}} J_\lambda(r) E_r f}. \quad (4.12)$$

From (4.10), (4.11) and (4.12),

$$\|df\|_{L^2} \leq \sqrt{\frac{n+2}{\nu_n r^{n+2}} \left( \frac{J_\lambda(r)}{C_\theta(r)} + \frac{\rho^2 f_\lambda(\rho) n r^2}{C_\theta(r)^{\frac{3}{2}}} \right) \sqrt{E_r f}}.$$

We can express  $\left( \frac{J_\lambda(r)}{C_\theta(r)} + \frac{\rho^2 f_\lambda(\rho) n r^2}{C_\theta(r)^{\frac{3}{2}}} \right)$  as  $g_\lambda(r, \rho)$  and observe that for Lemma 4.2(2),  $g_\lambda(r, \rho) \rightarrow 1$  as  $r \rightarrow 0$ .  $\square$

## 5. AN UPPER BOUND FOR $\lambda_k(\Gamma)$

In this section, we compute an appropriate lower bound for the eigenvalues of  $-\Delta_\Gamma$  using eigenvalues of  $-\Delta_M$  for manifolds  $M$  belonging to the class  $\mathcal{M}(n, \lambda, D, i_0)$ .

**Definition 7** (Interpolation map). Let  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and  $F$  be a finite  $\epsilon$ -net in  $M$  such that  $(F, \mu)$   $\epsilon$ -approximates  $(M, vol)$ .

Define  $I : L^2(F) \rightarrow C^{0,1}(M)$  as

$$Iu = \Lambda_{\rho-2\epsilon} P^* u.$$

From Lemma 4.4 and the fact that  $P^*$  preserves norm, we have

$$\|Iu\|_{L^2} \leq \frac{J_\lambda(i_0)}{C_\theta(\rho-2\epsilon)} \|u\|_{L^2}.$$

**Lemma 5.1.** Let  $0 < \epsilon < 2\rho < i_0$ , and  $\frac{\epsilon}{\rho} < \frac{1}{n}$ . Let  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and  $\Gamma = (F, \epsilon, \rho, \mu)$  be a  $(\epsilon, \rho)$ -approximation of  $M$ . For all  $u \in L^2(F)$ , let  $\|\delta(u)\|^2 = \frac{n+2}{\nu_n \rho^{n+2}} \sum_i \sum_{j: x_i \sim x_j} \mu_i \mu_j |u(x_j) - u(x_i)|^2$  denote the discrete Dirichlet energy on  $\Gamma$ . Then

$$(1) \quad \|\|Iu\|_{L^2} - \|u\|\|^2 \leq \frac{C(n)}{C_\theta(\rho-2\epsilon)} \rho^2 \|\delta u\|^2.$$

$$(2) \quad \|dIu\|_{L^2}^2 \leq \left(1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)\right) \|\delta u\|^2$$

where  $C(n) = \frac{3^n}{2}$  and  $g_\lambda(\rho-2\epsilon, \rho) \left(1 + \frac{2\epsilon}{\rho} \left(\frac{n}{n-2}\right)\right)^{n+2} = 1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)$  if

$$vol(B_{\rho-2\epsilon}(x)) \geq \frac{\alpha_n}{2} \left(\frac{2(\rho-2\epsilon)}{\pi}\right)^n.$$

*Proof.* (1) Let  $u \in L^2(F)$ . Using norm preserving property of  $P^*$  and triangle inequality,

$$\|\|Iu\|_{L^2} - \|u\|\| \leq \|Iu - P^*u\|_{L^2}.$$

Using Lemma 4.5, we have

$$\|Iu - P^*u\|_{L^2}^2 = \|\Lambda_{\rho-2\epsilon} P^*u - P^*u\|_{L^2}^2 \leq \frac{n+2}{2\nu_n C_\theta(\rho-2\epsilon)(\rho-2\epsilon)^n} E_{\rho-2\epsilon}(P^*u). \quad (5.1)$$

Now,  $\epsilon < \frac{\rho}{n}$ ,  $\epsilon < \frac{\rho}{3}$ , so  $(\rho-2\epsilon)^{-n} < 3^n \rho^n$ . So from (5.1),

$$\|Iu - P^*u\|_{L^2}^2 \leq \frac{3^n(n+2)}{2\nu_n C_\theta(\rho-2\epsilon)\rho^n} E_{\rho-2\epsilon}(P^*u). \quad (5.2)$$

On the other hand,

$$\|\delta u\|^2 = \frac{n+2}{\nu_n \rho^{n+2}} \int_M \int_{U(x)} |P^*u(y) - P^*u(x)|^2 dy dx,$$

where  $U(x) = \bigcup_{j: x_j \sim x_i} V_j$  such that  $x \in V_i$ .

Since  $U(x) \supset B_{\rho-2\epsilon}(x)$ ,

$$\begin{aligned} \|\delta u\|^2 &\geq \frac{n+2}{\nu_n \rho^{n+2}} \int_M \int_{B_{\rho-2\epsilon}(x)} |P^*u(y) - P^*u(x)|^2 dy dx \\ &= \frac{n+2}{\nu_n \rho^{n+2}} E_{\rho-2\epsilon}(P^*u). \end{aligned} \quad (5.3)$$

Thus, from (5.2) and (5.3),

$$\|Iu - P^*u\|_{L^2}^2 \leq \frac{C(n)}{C_\theta(\rho-2\epsilon)} \rho^2 \|\delta u\|^2.$$

(2) Proceeding with direct computation,

$$\begin{aligned} \|dIu\|_{L^2}^2 &= \|d(\Lambda_{\rho-2\epsilon} P^*u)\|_{L^2}^2 \\ &\leq \frac{n+2}{\nu_n (\rho-2\epsilon)^{n+2}} g_\lambda(\rho-2\epsilon, \rho) E_{\rho-2\epsilon}(P^*u) \end{aligned} \quad (5.4)$$

$$\leq \left( \frac{\rho}{\rho-2\epsilon} \right)^{n+2} g_\lambda(\rho-2\epsilon, \rho) \|\delta u\|^2 \quad (5.5)$$

where (5.4) follows from Lemma 4.6 and (5.5) follows from (5.3). Since  $\frac{\epsilon}{\rho} < \frac{1}{n}$  we have

$$\left( \frac{\rho}{\rho-2\epsilon} \right)^{n+2} \leq \left( 1 + \frac{2\epsilon}{\rho} \left( \frac{n}{n-2} \right) \right)^{n+2}.$$

□

Using the above lemma and the results obtained regarding the interpolation map, we are now ready to compute, under certain constraints, a lower bound for the eigenvalue of the graph Laplacian operator using the eigenvalue of the Laplace-Beltrami operator for all  $M \in \mathcal{M}(n, \lambda, D, i_0)$ .

**Theorem 5.2.** *Let  $M \in \mathcal{M}(n, \lambda, D, i_0)$ , and  $\Gamma = (F, \mu)$  be an  $(\epsilon, \rho)$ -approximation of  $(M, \text{vol})$ . Let  $\Gamma = (F, \mu, \rho)$  denote the weighted graph and  $0 < \epsilon < \rho < \frac{i_0}{2}$ . If for all  $x \in M$  and  $0 < r < \frac{i_0}{2}$ ,  $\text{vol}(B_r(x)) \geq \frac{\alpha_n}{2} \left( \frac{2r}{\pi} \right)^n$  then*

$$\lambda_k(M) \leq (1 - \delta_2(\epsilon, \rho, \lambda_k))^{-1} \lambda_k(\Gamma)$$

where  $\delta_2(\epsilon, \rho, \lambda_k) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

*Proof.* Let  $\lambda_k = \lambda_k(\Gamma)$ . Using the techniques from the proof of Theorem 3.7, it is enough to show there is a subspace  $L \subset H^1(M)$  such that  $\dim(L) = k$  and

$$\sup_{f \in L \setminus \{0\}} \frac{\|df\|_{L^2}^2}{\|f\|_{L^2}^2} \leq (1 - \delta(\epsilon, \rho, \lambda_k))^{-1} \lambda_k.$$

We have  $\|\delta u\|^2 \leq \lambda_k \|u\|^2$ . Using Lemma 5.1(1),

$$\|Iu\|_{L^2} \geq \left( 1 - \frac{C(n)}{\sqrt{C_\theta(\rho-2\epsilon)}} \rho \sqrt{\lambda_k} \right) \|u\|.$$

Let  $f = Iu$ . Then

$$\|f\|_{L^2}^2 \geq \left( 1 - \frac{C(n)}{\sqrt{C_\theta(\rho-2\epsilon)}} \rho \sqrt{\lambda_k} \right) \|u\|^2. \quad (5.6)$$

On the other hand, from Lemma 4.2(2) and Lemma 5.1(2),

$$\begin{aligned} \|df\|_{L^2}^2 &\leq \left(1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)\right) \|\delta u\|^2 \\ &\leq \left(1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)\right) \lambda_k \|u\|^2. \end{aligned} \quad (5.7)$$

From Lemma 5.1(2),  $\sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

Let  $\zeta(\epsilon, \rho) = \frac{\rho}{\sqrt{C_\theta(\rho-2\epsilon)}}$ . Clearly,  $\zeta(\epsilon, \rho) \rightarrow 0$  as  $\epsilon, \rho \rightarrow 0$ .

From equations (5.7) and (5.6), we have

$$\begin{aligned} \frac{\|df\|^2}{\|f\|^2} &\leq \left(\frac{1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)}{1 - \zeta(\epsilon, \rho)C(n)\sqrt{\lambda_k}}\right) \lambda_k \\ &\leq \frac{1}{1 - \delta_2(\epsilon, \rho, \lambda_k)} \lambda_k \end{aligned}$$

where  $\delta_2(\epsilon, \rho, \lambda_k) = \frac{\sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right) + \zeta(\epsilon, \rho)C(n)\sqrt{\lambda_k}}{1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)}$ .

□

**Proof of Theorem 1.1.** From Theorem 3.4, we have

$$\lambda_k(\Gamma) - \lambda_k(M) \leq \delta_1(\epsilon, \rho, \lambda_k(M)) \lambda_k(M). \quad (5.8)$$

where  $\delta_1(\epsilon, \rho, \lambda_k(M)) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

And from Theorem 5.2, we have

$$\lambda_k(M) \leq (1 - \delta_2(\epsilon, \rho, \lambda_k(M)))^{-1} \lambda_k(\Gamma)$$

where  $\delta_2(\epsilon, \rho, \lambda_k) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

Hence we have,

$$\lambda_k(M) - \lambda_k(\Gamma) \leq \delta_2(\epsilon, \rho, \lambda_k(M)) \lambda_k(M). \quad (5.9)$$

Let  $\delta(\epsilon, \rho, \lambda_k(M)) = \max\{\delta_1(\epsilon, \rho, \lambda_k(M)), \delta_2(\epsilon, \rho, \lambda_k(M))\}$ . From (5.8) and (5.9), we have

$$|\lambda_k(M) - \lambda_k(\Gamma)| \leq \delta(\epsilon, \rho, \lambda_k(M)) \lambda_k(M). \quad (5.10)$$

From [1], we obtain a uniform bound  $C_{k, \mathcal{M}}$  on the  $k$ -th eigenvalues of  $\Delta_M$ . Substituting in (5.10),

$$|\lambda_k(M) - \lambda_k(\Gamma)| \leq \delta(\epsilon, \rho, C_{k, \mathcal{M}}) C_{k, \mathcal{M}}.$$

□

**A general case :** We now consider a general case where the condition on the lower bound of a ball with small radius is not assumed. The following theorem provides a uniform bound on the eigenvalues applicable to the entire class  $\mathcal{M}(n, \lambda, D, i_0)$ . The graph eigenvalues are scaled by a constant depending solely on  $n$ , making the resulting expression particularly useful for higher-order eigenvalues.

**Proposition 5.3.** *For a manifold  $M \in \mathcal{M}(n, \lambda, D, i_0)$  with and an  $(\epsilon, \rho)$  approximation  $\Gamma$  of  $M$ , if  $\lambda_k(M)$  and  $\lambda_k(\Gamma)$  denote the  $k$ -th eigenvalues of  $-\Delta_M$  and  $-\Delta_\Gamma$ , respectively. Then there exists  $\delta(\epsilon, \rho, \lambda_k) > 0$  such that for all positive integers  $k$ ,*

$$\lambda_k(M) - C(n) \lambda_k(\Gamma) \leq \delta(\epsilon, \rho, \lambda_k(M)) \lambda_k(M)$$

$C(n)$  is a constant depending on  $n$  and  $\delta(\epsilon, \rho, \lambda_k(M)) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

*Proof.* We use the fact proven by Berger [4] that there exists a uniform constant depending on  $n$  such that for metric balls of radius less than  $\frac{i_0}{2}$ , (1.3) holds. Evaluating the value of  $g_\lambda$  from Lemma 4.6 for Lemma 4.2(1), we observe that  $g_\lambda$  is a constant  $C(n)$  depending on  $n$ . Without loss of generality, we denote all constants depending on  $n$  by  $C(n)$ . Substituting the value of  $g_\lambda$  in Lemma 5.1(2), we get

$$\|df\|^2 \leq C(n)h\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right) \|\delta u\|^2 \quad (5.11)$$

where  $h\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right) - 1 \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . We follow rest of the computations after Lemma 5.1(2) using (5.11). Finally using (5.11) in (5.7),

$$\lambda_k(M) - C(n)\lambda_k(\Gamma) \leq \delta(\epsilon, \rho, \lambda_k(M))\lambda_k(M)$$

for some constant  $C(n)$  depending solely on  $n$ .  $\square$

Using Theorem 1.1 and Proposition 5.3, we have the following corollary.

**Corollary 5.4.** *Consider a manifold  $M \in \mathcal{M}(n, \lambda, D, i_0)$  and an  $(\epsilon, \rho)$ -approximation  $\Gamma$  of  $M$  such that  $\rho < i_0$  and  $\frac{\epsilon}{\rho} < \frac{1}{n}$ . Let  $\lambda_k(M)$  and  $\lambda_k(\Gamma)$  denote the  $k$ -th eigenvalues of  $-\Delta_M$  and  $-\Delta_\Gamma$ , respectively. Then there exists  $\delta(\epsilon, \rho, \lambda_k) > 0$  such that for all positive integers  $k$ ,*

$$\lambda_k(\Gamma) - \lambda_k(M) \leq \delta(\epsilon, \rho, \lambda_k(M))\lambda_k(M)$$

and

$$\lambda_k(M) - C(n)\lambda_k(\Gamma) \leq \delta(\epsilon, \rho, \lambda_k(M))\lambda_k(M)$$

where

$C(n)$  is a constant depending on  $n$  and  $\delta(\epsilon, \rho, \lambda_k(M)) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . Moreover, if  $|\text{Ric}| \leq \lambda$ , then there exists a constant  $C_{k, \mathcal{M}}$  such that  $\lambda_k(M) \leq C_{k, \mathcal{M}}$  for all  $M \in \mathcal{M}(n, \lambda, D, i_0)$ . Then,

$$\lambda_k(\Gamma) - \lambda_k(M) \leq \delta(\epsilon, \rho, C_{k, \mathcal{M}})C_{k, \mathcal{M}}$$

and

$$\lambda_k(M) - C(n)\lambda_k(\Gamma) \leq \delta(\epsilon, \rho, C_{k, \mathcal{M}})C_{k, \mathcal{M}}$$

where  $\delta(\epsilon, \rho, C_{k, \mathcal{M}})$  tends to zero as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

**Remark 5.5.** For a manifold  $(M, g)$ , C. B. Croke [9] provides a lower bound to the volume of balls with radius less than the injectivity radius  $i(M)$ . For any ball  $B_x(r)$  such that  $r < i(M)$ ,

$$\text{vol}(B_x(r)) \geq \frac{2^{n-1}\alpha_{n-1}^n}{n^n\alpha_n^{n-1}}r^n \quad (5.12)$$

where  $\alpha_n$  denotes the volume of  $n$ -sphere with sectional curvature 1. One can follow the above techniques and obtain estimates on the eigenvalues by using (5.12).

## 6. APPROXIMATION OF EIGENFUNCTIONS

In this section, we will investigate the approximation of eigenfunctions of  $M \in \mathcal{M}(n, \lambda, D, i_0)$  such that for all  $0 < r < \frac{i_0}{2}$  and  $x \in M$ ,  $\text{vol}(B_r(x)) \geq \frac{\alpha_n}{2} \left(\frac{2r}{\pi}\right)^n$ . Similar arguments as in [6] hold in our setting. The following lemma shows that the discretization map and the interpolation map are almost inverses of each other.

**Lemma 6.1.** (1) *Let  $f \in H^1(M)$ . Then,*

$$\|IPf - f\| \leq \frac{J_\lambda(i_0)}{C_\theta(r)(i_0)} (1 + C(n)J_\lambda(\rho)) \rho \|df\|.$$

(2) *Let  $u \in L^2(X)$ . Then,*

$$\|PIu - u\| \leq \left( C(n)J_\lambda(\rho) \left( 1 + \sigma(i_0) + C_\theta(r)^{-\frac{1}{2}} \right) \right) \rho \|\delta u\|.$$



*Proof.* (1) By straight forward computation,

$$\begin{aligned} \|IPf - f\| &= \|\Lambda_{\rho-2\epsilon}P^*Pf - f\| \\ &\leq \|\Lambda_{\rho-2\epsilon}(P^*Pf - f)\| + \|\Lambda_{\rho-2\epsilon}f - f\| \end{aligned} \quad (6.1)$$

We aim to find upper bounds of both the terms obtained above using triangle inequality. From Lemma 4.4 and Lemma 3.2,

$$\|\Lambda_{\rho-2\epsilon}(P^*Pf - f)\| \leq \frac{J_\lambda(\rho-2\epsilon)}{C_\theta(r)} \|P^*Pf - f\| \leq \frac{J_\lambda(i_0)}{C_\theta(r)} C(n)J_\lambda(\rho)\epsilon \|df\|_{L^2}. \quad (6.2)$$

Again from Lemma 4.5 and Lemma 2.2,

$$\|\Lambda_{\rho-2\epsilon}f - f\|^2 \leq \frac{n+2}{\nu_n C_\theta(r)(i_0)(\rho-2\epsilon)^n} E_{\rho-2\epsilon}(f) \leq J_\lambda(i_0)C_\theta(r)(i_0)^{-1}\rho^2 \|df\|^2. \quad (6.3)$$

From equations (6.1), (6.2) and (6.3), we conclude that

$$\|IPf - f\| \leq \frac{J_\lambda(i_0)}{C_\theta(r)(i_0)} (1 + C(n)J_\lambda(\rho)) \rho \|df\|.$$

(2) Using Lemma 3.2 and (5.1) and following the similar arguments as above, we obtain the required inequality.  $\square$

Let  $I$  be an interval in  $\mathbb{R}$  and  $H_I(M)$ (or  $H_I(X)$ ) be the subspace of  $H^1(M)$ (or  $H^1(X)$ ) spanned by eigenfunctions having eigenvalues in  $I$ . Also, we denote  $H_{(-\infty, \lambda)}(M)$  by  $H_\lambda(M)$ . Let  $\mathbb{P}_I : L^2(M) \rightarrow H_I(M)$  be the orthogonal projection. We use similar notation for orthogonal projections from  $L^2(X)$  to  $H_I(X)$  and abbreviate  $\mathbb{P}_{(-\infty, \lambda)}$  as  $\mathbb{P}_\lambda$ .

**Lemma 6.2.** (1) Let  $\lambda > 0$  and  $f \in H_\lambda(M)$ . Then

$$\|\delta(Pf)\| \geq (1 - \sigma_1(\rho, \epsilon)) \|df\|$$

where  $\sigma_1$  depends on  $n, \lambda, i_0$  and  $\sigma_1(\rho, \epsilon) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

(2) Let  $\lambda > 0$  and  $u \in H_\lambda(X)$ . Then

$$\|d(Iu)\| \geq (1 - \sigma_2(\rho, \epsilon)) \|\delta u\|$$

where  $\sigma_2$  depends on  $n, \lambda, i_0$  and  $\sigma_2(\rho, \epsilon) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

*Proof.* (1) Since  $\mathbb{P}_\lambda$  does not increase Dirichlet energy,

$$\|d(IPf)\|_{L^2} \geq \|d(\mathbb{P}_\lambda IPf)\|_{L^2} \geq \|df\|_{L^2} - \|d(\mathbb{P}_\lambda IPf - f)\|_{L^2}.$$

Since  $f \in H_\lambda(M)$ , from Lemma 6.1 we obtain

$$\|d(\mathbb{P}_\lambda IPf - f)\|_{L^2} \leq \frac{J_\lambda(i_0)}{C_\theta(r)(i_0)} (1 + C(n)J_\lambda(\rho)) \sqrt{\lambda}\rho \|df\|$$

Hence,

$$\|d(IPf)\|_{L^2} \geq \left(1 - \frac{J_\lambda(i_0)}{C_\theta(r)(i_0)} (1 + C(n)J_\lambda(\rho)) \sqrt{\lambda}\rho\right) \|df\|. \quad (6.4)$$

From Lemma 5.1,

$$\|d(IPf)\| \leq (1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right))(\rho, \epsilon) \|\delta Pf\|. \quad (6.5)$$

Thus, from (6.4) and (6.5),

$$\begin{aligned} \|\delta(Pf)\| &\geq \frac{1 - \frac{J_\lambda(i_0)}{C_\theta(r)(i_0)} (1 + C(n)J_\lambda(\rho)) \sqrt{\lambda}\rho}{\left(1 + \sigma\left(\epsilon, \rho, \frac{\epsilon}{\rho}\right)\right)(\rho, \epsilon)} \|df\| \\ &= (1 - \sigma_1(\rho, \epsilon)) \|df\| \end{aligned}$$

(2) Applying the techniques from the previous part and Lemma 6.1 we obtain

$$\|\delta(PYu)\|_{L^2} \geq \left(1 - \left(C(n)J_\lambda(\rho) \left(1 + \sigma(i_0) + C_\theta(r)^{-\frac{1}{2}}\right)\right) \sqrt{\lambda\rho}\right) \|\delta u\|. \quad (6.6)$$

From Lemma 3.4,

$$\|\delta(PYu)\| \leq \left(J_\lambda(\rho + 2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2}\right)^{\frac{1}{2}} \|d(Yu)\|_{L^2}. \quad (6.7)$$

From (6.6) and (6.7),

$$\begin{aligned} \|d(Yu)\|_{L^2} &\geq \frac{1 - \left(C(n)J_\lambda(\rho) \left(1 + \sigma(i_0) + \frac{1}{\sqrt{C_\theta(r)}}\right)\right) \sqrt{\lambda\rho}}{\sqrt{J_\lambda(\rho + 2\epsilon) \left(1 + \frac{2\epsilon}{\rho}\right)^{n+2}}} \|\delta u\| \\ &= (1 - \sigma_2(\rho, \epsilon)) \|\delta u\| \end{aligned}$$

where  $\sigma_2(\rho, \epsilon) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ .

□

Let  $\{f_k\}_\Gamma^\infty$  and  $\{u_k\}_1^N$  be orthonormal eigenvectors of  $\Delta_M$  and  $\Delta_X$  respectively. The next lemma approximates eigenfunctions of  $\Delta_M$ . It can be proved by using Lemma 7.3 from [6] and analogous inequalities we have obtained so far.

**Lemma 6.3.** *Let  $0 < \epsilon < \rho < \frac{i_0}{2}$ ,  $\mathcal{C}_{\mathcal{M},k}C(n)J_\lambda(\rho)\epsilon < 1$  and  $\mathcal{C}_{\mathcal{M},k} > 1$ .*

(1) *For all  $k \in \mathbb{N}$  and  $a \in \mathbb{R}_+$ ,*

$$\|Pf_k - \mathbb{P}_{\lambda_{k+a}}Pf_k\|^2 \leq \frac{1}{a}\mathcal{C}_{\mathcal{M},k}\sigma(\epsilon, \rho)$$

where  $\sigma_{\mathcal{M}}(\epsilon, \rho) = \max\{\delta(\epsilon, \rho, \mathcal{C}_{\mathcal{M},k}), \mathcal{C}_{\mathcal{M},k}C(n)J_\lambda(\rho)\epsilon\} \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . Also

$$\|\delta(Pf_k - \mathbb{P}_{\lambda_{k+a}}Pf_k)\|^2 \leq \mathcal{C}_{\mathcal{M},k} \left(1 + \frac{1}{a}\right) \sigma_{\mathcal{M}}(\epsilon, \rho).$$

(2) *Let  $\lambda = \lambda_k(X)$ . Then for every  $a \in \mathbb{R}_+$ ,*

$$\|Yu_k - \mathbb{P}_{\lambda_{k+a}}Yu_k\|^2 \leq \frac{1}{a}\mathcal{C}_{\mathcal{M},k}\sigma_{\mathcal{M}}(\epsilon, \rho)$$

where  $\sigma_{\mathcal{M}}(\epsilon, \rho) \rightarrow 0$  as  $\epsilon, \rho, \frac{\epsilon}{\rho} \rightarrow 0$ . Also

$$\|d(Yu_k - \mathbb{P}_{\lambda_{k+a}}Yu_k)\|^2 \leq \mathcal{C}_{\mathcal{M},k} \left(1 + \frac{1}{a}\right) \sigma(\epsilon, \rho).$$

From Lemma 6.1, we consider the maximum of  $\sigma_{\mathcal{M}}(\epsilon, \rho)$ ,  $\sigma_1(\epsilon, \rho)$  and  $\sigma_2(\epsilon, \rho)$  and denote it again by  $\sigma_{\mathcal{M}}(\epsilon, \rho)$ . We use Lemma 7.4[6] to prove the following.

**Lemma 6.4.** *Let  $\mathcal{C}_{\mathcal{M},k}C(n)J_\lambda(\rho)\epsilon < 1$  and  $\mathcal{C}_{\mathcal{M},k} > 1$ .*

(1) *Let  $\lambda = \lambda_k(M)$  and  $\alpha, \beta, \gamma > 0$  such that  $\alpha \leq \beta \leq \gamma \leq 1$  and there are no eigenvalues of  $-\Delta_\Gamma$  in  $(\lambda + \alpha, \lambda + \beta)$ . Then*

$$\|Pf_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\alpha)}Pf_k\|^2 \leq \mathcal{C}_{\mathcal{M},k}\alpha^{-1}\gamma^{-1}\sigma_{\mathcal{M}}(\epsilon, \rho) + 2\beta\gamma^{-1}.$$

(2) *Let  $\lambda = \lambda_k(\Gamma)$  and  $\alpha, \beta, \gamma > 0$  such that  $\alpha \leq \beta \leq \gamma \leq 1$  and there are no eigenvalues of  $-\Delta_M$  in  $(\lambda + \alpha, \lambda + \beta)$ . Then*

$$\|Yu_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\alpha)}Yu_k\|^2 \leq \mathcal{C}_{\mathcal{M},k}\alpha^{-1}\gamma^{-1}\sigma_{\mathcal{M}}(\epsilon, \rho) + 2\beta\gamma^{-1}.$$

**Theorem 6.5.** (1) *Let  $\lambda = \lambda_k(M)$  and let  $f_k$  be corresponding unit-norm eigenfunction of  $\Delta_M$ . Then for every  $\gamma \in (0, 1)$ ,*

$$\|Pf_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\gamma)}Pf_k\|^2 \leq \mathcal{C}_{\mathcal{M},k}\gamma^{-2}\sigma_{\mathcal{M}}(\epsilon, \rho).$$

- (2) Let  $\lambda = \lambda_k(M)$  and let  $f_k$  be corresponding unit-norm eigenfunction of  $\Delta_M$ . Then for every  $\gamma \in (0, 1)$ ,

$$\|Iu_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\gamma)} Iu_k\|^2 \leq \mathcal{C}_{\mathcal{M},k} \gamma^{-2} \sigma_{\mathcal{M}}(\epsilon, \rho).$$

*Proof.* Substituting  $\alpha = \beta = \sqrt{\sigma_{\mathcal{M}}(\epsilon, \rho)} \gamma$  in Lemma 6.4 we get

$$\|Pf_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\gamma)} Pf_k\|^2 \leq C \sqrt{\sigma_{\mathcal{M}}(\epsilon, \rho)} + \mathcal{C}_{\mathcal{M},k} \gamma^{-2} \sqrt{\sigma_{\mathcal{M}}(\epsilon, \rho)}.$$

Now  $\gamma^{-2} \geq 1$ . This implies,

$$\|Pf_k - \mathbb{P}_{(\lambda-\gamma, \lambda+\gamma)} Pf_k\|^2 \leq \mathcal{C}_{\mathcal{M},k} \gamma^{-2} \sqrt{\sigma_{\mathcal{M}}(\epsilon, \rho)}.$$

The second part can also be shown with similar methods.  $\square$

**Theorem 6.6.** Let  $\lambda$  be an eigenvalue of  $\Delta_M$  of multiplicity  $m$ , more precisely,

$$\lambda_{k-1} < \lambda_k = \lambda = \lambda_{k+m-1} < \lambda_{k+m}.$$

where  $\lambda_j = \lambda_j(M)$ . Let  $\delta_\gamma = \min\{1, \lambda_k - \lambda_{k-1}, \lambda_{k+m} - \lambda_{k+m-1}\}$ . Let  $\{u_k, \dots, u_{k+m-1}\}$  be orthonormal eigenvectors of  $-\Delta_\Gamma$  corresponding to eigenvalues  $\lambda_k(\Gamma), \dots, \lambda_{k+m-1}(\Gamma)$ . Then there exists orthonormal eigenfunctions  $f_k, \dots, f_{k+m-1}$  of  $-\Delta_M$  corresponding to eigenvalue  $\lambda$  and such that

$$\|u_j - Pf_j\|^2 \leq \mathcal{C}_{\mathcal{M},k} \delta_\lambda^{-2} \sigma(\epsilon, \rho) \quad (6.8)$$

and

$$\|f_j - Iu_j\|^2 \leq \mathcal{C}_{\mathcal{M},k} \delta_\lambda^{-2} \sigma(\epsilon, \rho) \quad (6.9)$$

for all  $j = k, \dots, k+m-1$ .

*Proof.* The uniform constant in Theorem 5.4, denoted by  $\mathcal{C}_{\mathcal{M},k}$ , can be chosen such that

$$|\lambda_j(\Gamma) - \lambda_j(\Gamma)| < \frac{1}{4} \delta_\lambda,$$

for all  $j = k, \dots, k+m-1$ .

Applying Lemma 6.4 and Lemma 6.2 and using Theorem 4[6] in our setting, we get the desired result.  $\square$

## REFERENCES

- [1] Michael T. Anderson. Convergence and rigidity of manifolds under Ricci curvature bounds. *Invent. Math.*, 102(2):429–445, 1990.
- [2] Erwann Aubry. Approximation of the spectrum of a manifold by discretization. *arXiv preprint arXiv:1301.3663*, 2013.
- [3] Mikhail Belkin and Partha Niyogi. Convergence of laplacian eigenmaps. In B. Schölkopf, J. Platt, and T. Hoffman, editors, *Advances in Neural Information Processing Systems*, volume 19. MIT Press, 2006.
- [4] M. Berger. Some relations between volume, injectivity radius, and convexity radius in Riemannian manifolds. In *Differential geometry and relativity*, volume Vol. 3 of *dath. Phys. Appl. Math.*, pages 33–42. Reidel, Dordrecht-Boston, Mass., 1976.
- [5] Marcel Berger. Volume et rayon d’injectivité dans les variétés riemanniennes de dimension 3. *Osaka Math. J.*, 14(1):191–200, 1977.
- [6] Dmitri Burago, Sergei Ivanov, and Yaroslav Kurylev. A graph discretization of the Laplace-Beltrami operator. *J. Spectr. Theory*, 4(4):675–714, 2014.
- [7] Dmitri Burago, Sergei Ivanov, and Yaroslav Kurylev. Spectral stability of metric-measure Laplacians. *Israel J. Math.*, 232(1):125–158, 2019.
- [8] Dmitri Burago, Sergei Ivanov, Yaroslav Kurylev, and Jinpeng Lu. Approximations of the connection Laplacian spectra. *Math. Z.*, 301(3):3185–3206, 2022.
- [9] Christopher Croke. Some isoperimetric inequalities and eigenvalue estimates. *Annales Scientifiques de l’École Normale Supérieure. Quatrième Série*, 13, 01 1980.
- [10] Christopher B. Croke. Area of small disks. *Bull. Lond. Math. Soc.*, 41(4):701–708, 2009.
- [11] S. Gallot, D. Hulin, and J. Lafontaine. *Riemannian Geometry*. Universitext. Springer Berlin Heidelberg, 2004.
- [12] Peter Li and Shing Tung Yau. Estimates of eigenvalues of a compact Riemannian manifold. In *Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, volume XXXVI of *Proc. Sympos. Pure Math.*, pages 205–239. Amer. Math. Soc., Providence, RI, 1980.
- [13] Jinpeng Lu. Graph approximations to the Laplacian spectra. *J. Topol. Anal.*, 14(1):111–145, 2022.

- [14] Guofang Wei. Manifolds with a lower Ricci curvature bound. In *Surveys in differential geometry. Vol. XI*, volume 11 of *Surv. Differ. Geom.*, pages 203–227. Int. Press, Somerville, MA, 2007.