

COSMIC RAYS MASQUERADING AS HOT CGM GAS: AN INVERSE-COMPTON ORIGIN FOR DIFFUSE X-RAY EMISSION IN THE CIRCUMGALACTIC MEDIUM

PHILIP F. HOPKINS^{1*}, ELIOT QUATAERT^{2**}, SAM B. PONNADA¹, AND EMILY SILICH¹

¹TAPIR, Mailcode 350-17, California Institute of Technology, Pasadena, CA 91125, USA and

²Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA

Version February 3, 2025

ABSTRACT

Observations from *ROSAT* and *eROSITA* have argued that Milky Way (MW), Andromeda, and lower-mass galaxies exhibit extended soft X-ray (1 keV) diffuse halos out to radii $R \gtrsim 100$ kpc in the circumgalactic medium (CGM). If interpreted as thermal emission from hot gas, the surprisingly shallow surface brightness profiles $S_X \propto R^{-1}$ of this emission are difficult to explain, and contradict other observations. We show that such halos instead arise from inverse Compton (IC) scattering of CMB photons with GeV cosmic ray (CR) electrons. GeV electrons have long (\sim Gyr) lifetimes and escape the galaxy, forming a shallow extended radial profile out to $R \gtrsim 100$ kpc, where IC off the CMB dominates their losses and should produce soft, thermal-like X-ray spectra peaked at ~ 1 keV. The observed keV halo luminosities and brightness profiles agree well with those expected for CRs observed in the local interstellar medium (LISM) escaping the galaxy, with energetics consistent with known CRs from SNe and/or AGN, around galaxies with stellar masses $M_* \lesssim 2 \times 10^{11} M_\odot$. At higher masses observed X-ray luminosities are larger than predicted from IC and should be dominated by hot gas. In the MW+M31, the same models of escaping CRs reproduce *Fermi* gamma-ray observations if we assume an LISM-like proton-to-electron ratio and CR-pressure-dominated halo. In all other halos, the associated non-thermal radio and γ -ray brightness is far below detectable limits. If we have indeed detected the expected IC X-ray halos, the observations provide qualitatively new and stringent constraints on the properties of the CGM and CR physics: the observed X-ray brightness *directly* traces the CR lepton energy density $e_{\text{cr},\ell}$ in the CGM (without any degenerate parameters). The implied $e_{\text{cr},\ell}$ agrees well with LISM values at radii $R \lesssim 10$ kpc, while following the profile predicted by simple steady state models of escaping CRs at larger radii. The inferred CR pressure is a major part of the total pressure budget in the CGM of Milky Way-mass galaxies, suggesting that models of thermally dominated halos at Milky Way mass may need to be revised. The measurement of X-ray surface brightness and total luminosity allows one to further determine the effective CGM diffusivity/CR streaming speed at radii $\sim 10 - 1000$ kpc. We show these also agree with LISM values at small radii but the inferred diffusivity increases significantly at larger radii, consistent with independent CGM constraints from UV absorption at ~ 100 kpc.

Subject headings: circumgalactic medium — galaxies: haloes — X-rays — cosmic rays — galaxies: formation

1. INTRODUCTION

Soft X-ray ($\sim 0.5 - 2$ keV) observations from *eROSITA* (Lyskova et al. 2023; Zhang et al. 2024a,b; Bahar et al. 2024; consistent with prior but less-sensitive *ROSAT* observations (Anderson et al. 2013, 2015)) have argued that stacked X-ray images around galaxies with stellar masses $\sim 10^{10} - 10^{11.5} M_\odot$, including Milky Way (MW) and Andromeda (M31)-mass and lower-mass galaxies, exhibit an excess of diffuse emission that appears to be CGM emission from $\gtrsim 10 - 100$ kpc around the central galaxies. Surprisingly, the X-ray surface brightness or flux declines very slowly, $\propto 1/R$ with galacto-centric distance even approaching $\gtrsim R_{500}$ — much shallower than the observed X-ray halos around more massive, individually-detected groups and clusters (Lyskova et al. 2023). The emission is also much more luminous, and declines much more slowly with radius, than predicted in high-resolution numerical simulations of MW-M31-mass systems, if it is assumed to arise from thermal (free-free+metal-line) hot gas emission (Truong et al. 2023; Sultan et al. 2024; Popesso et al. 2024b; Silich et al., in prep.). And regardless of models, reproducing the observed emission via hot gas would require more baryons (uniformly at *super-virial* temperatures) than the universal baryon fraction, and/or more metals than produced by

the sum of all SNe in the history of the galaxy, for MW-mass systems (see §3 below and Zhang et al. 2024a), as well as directly contradicting other observational constraints from both UV (Werk et al. 2014; Faerman et al. 2022; Wijers et al. 2024) and X-ray *eROSITA* (Ponti et al. 2023) and Chandra (Yao et al. 2010) absorption-line studies. Both *eROSITA* and *ROSAT* also find a sharp change in behavior in the total halo luminosity L_X versus stellar mass M_* , where below $M_* \lesssim 2 \times 10^{11} M_\odot$ the correlation is shallow (linear or sub-linear), while it steepens dramatically at higher M_* .

These observations suggest that there is either a dramatic change in halo gas properties in lower mass systems or a change in the nature of the dominant soft X-ray emission process in lower-mass halos. In this paper, we show that inverse Compton (IC) scattering of CMB photons with \sim GeV CR electrons (which are observed to escape the ISM), naturally predicts X-ray halos very similar to those observed in lower-mass systems.

Direct CR observations from experiments like Voyager, AMS-02, and others, show that near the solar circle, \sim GeV CR electrons dominate the CR lepton energy budget, with an energy density ~ 0.02 eV cm^{-3} , and that they have long lifetimes ($\sim 10^9$ yr) compared to their residence/escape timescale from the ISM ($\lesssim 10^7$ yr). This implies that GeV leptons must be escaping from the galaxy into the CGM with an implied galaxy-integrated leptonic rate $\sim 10^{40}$ erg s^{-1} (e.g. Zweibel 2017; Amato & Blasi 2018; Evoli et al. 2019; Maurin 2020; Butsky

* phopkins@caltech.edu
** quataert@princeton.edu

et al. 2023; Di Mauro et al. 2024, and references therein). In this paper, we show that these escaping CR leptons produce X-ray halos with a spectrum peaked in the observed soft X-ray bands, and with a normalization, luminosity, and surface brightness profile very similar to those observed around galaxies with stellar masses $M_* \lesssim 2 \times 10^{11} M_\odot$.

If these observed X-ray halos indeed arise from IC, then they do not trace hot gas, but instead trace the properties of CRs. This enables, for the first time, direct measurement of the CR (leptonic) energy density in the CGM, as well as the CR transport parameters in the CGM – profoundly powerful and unique constraints for CR “feedback” and scattering physics models.

2. INVERSE COMPTON FROM EXTENDED COSMIC RAY HALOS

2.1. Basic Scalings of Emission and Surface Brightness Profiles

Consider the inverse Compton (IC) radiation from a population of CR electrons. A typical CR electron with energy $E_{\text{cr}} \sim E_{\text{cr,GeV}} \text{ GeV}$ (around the peak of the CR spectrum at $E_{\text{cr,GeV}} \sim 1$) will IC scatter a photon to a characteristic energy $h\nu_{\text{IC}} \rightarrow h\nu_{\text{initial}} \gamma_{\text{cr}}^2$, where for CMB photons $\langle h\nu_{\text{initial}} \rangle \sim 3 k_B T_{\text{cmb}} \sim 7 \times 10^{-4} (1+z) \text{ eV}$, so at emission (rest-frame)

$$h\nu_{\text{IC}} \sim 1 - 3 \text{ keV } E_{\text{cr,GeV}}^2 (1+z). \quad (1)$$

The X-ray emissivity will be given by

$$\begin{aligned} \epsilon_X &\sim \frac{4}{3} h\nu_{\text{IC}} n_{\text{cr}, e^\pm} n_{\text{photons}} \sigma_{\text{T}} c \\ &\sim 10^{34.5} \frac{\text{erg}}{\text{s kpc}^3} \left(\frac{e_{\text{cr}, \ell} E_{\text{cr,GeV}}}{0.02 \text{ eV cm}^{-3}} \right) (1+z)^4 \end{aligned} \quad (2)$$

where $e_{\text{cr}, \ell}$ is the total leptonic CR energy density and we have scaled to the Solar-neighborhood value at $\sim \text{GeV}$.¹ A proper calculation of the X-ray spectrum and emissivity are shown in Fig. 1 (and § 2.1.1 below). Integrating through a halo² at some impact parameter $R \equiv R_{100} 100 \text{ kpc}$, this gives a surface brightness³

$$\frac{S_X, \text{keV}}{10^{37} \text{ erg s}^{-1} \text{ kpc}^{-2}} \sim \left(\frac{e_{\text{cr}, \ell} E_{\text{cr,GeV}}}{0.02 \text{ eV cm}^{-3}} \right) \left(\frac{R}{100 \text{ kpc}} \right) (1+z)^4. \quad (3)$$

Now consider the expectation for a CR halo around a galaxy. As discussed and shown in many studies (Su et al. 2020; Hopkins et al. 2020, 2021a,b, 2023b; Ji et al. 2020, 2021; Butsky et al. 2023; Ponnada et al. 2024b,c), the CR energy density e_{cr} in the CGM/ICM/IGM in simulations and more detailed models can usually be represented quite accurately by simply assuming the spherically-symmetric, steady-state solution with weak losses or re-acceleration in the (low-density) CGM (since the $\sim \text{GeV}$ CRs which dominate e_{cr} have small losses except in dense ISM gas and after very long timescales from

¹ We can also define the ratio of leptonic-to-total CR energy, $f_{\text{cr}, \ell} \equiv e_{\text{cr}, \ell} / e_{\text{cr, tot}}$, with $f_{\text{cr}, \ell}^\odot \sim 0.02$ in the LISM.

² For power-law profiles $e_{\text{cr}} \propto r^{-\alpha}$, at impact parameter R , $S_X \approx A_0 R \epsilon_X(r=R)$ with $A_0 \equiv \pi^{1/2} \Gamma[(\alpha-1)/2] / \Gamma[\alpha/2] \sim 3-5$ for the range of slopes of interest here.

³ We follow Zhang et al. 2024b’s convention (since we compare to their observations) for the definition of surface brightness as dL_{iso}/dA in terms of the differential isotropic luminosity and area.




FIG. 1.— Observed-frame continuum spectra predicted by IC from CRs off the CMB (§ 2.1.1). *Top*: $E^2 dN/dE$ over a very broad energy range. *Bottom*: dN/dE focused on standard X-ray energies 0.2 – 8 keV. We calculate the spectrum from IC only, from the CMB (other cosmic backgrounds are totally negligible at these energies), at $z = 0$. For the CR lepton spectrum, we compare (1) the observed LISM spectrum, and (2) that spectrum corrected for IC losses with finite travel time (i.e. finite distance from the ISM) parameterized by $\tau_{\text{loss}}^0 \equiv t_{\text{travel}}/t_{\text{loss}} \sim (R/100 \text{ kpc}) (100 \text{ km s}^{-1}/v_{\text{st, eff}})$ at 1 GeV. For reference, we compare the free-free spectra predicted for different temperatures. IC CRs leaking from the ISM produces a thermal-like spectrum with most of the luminosity at $\sim \text{keV}$ (from $\sim \text{GeV}$ leptons, near the LISM spectral peak, with $\sim \text{Gyr}$ lifetimes), until most of the CR energy is lost ($\tau_{\text{loss}}^0 \gg 1$, expected at $R \gtrsim 0.1 - 1 \text{ Mpc}$ distances from the galaxy).

inverse Compton, which we account for below, with a factor f_{loss} , for some leptonic energy injection rate $\dot{E}_{\text{cr}, \ell}$ from the galaxy (e.g. from SNe shocks, AGN, fast stellar winds, pulsar wind nebulae, etc.). This gives:

$$e_{\text{cr}, \ell} \sim \frac{f_{\text{loss}} \dot{E}_{\text{cr}, \ell}}{4\pi v_{\text{st, eff}} r^2} \approx \frac{f_{\text{loss}} \dot{E}_{\text{cr}, \ell}}{4\pi \kappa_{\text{eff}} [r] r} \quad (4)$$

where κ_{eff} and $v_{\text{st, eff}}$ are some effective, spectrum and isotropically and position-averaged CR diffusion coefficient and/or streaming (+advection) speed in the CGM. In the CGM, at the halo masses of interest, those models and Wiener et al. (2013, 2017); Ruszkowski et al. (2017); Thomas et al. (2023) typically predict that CR transport at $\sim \text{GeV}$ is streaming-dominated with $v_{\text{st, eff}} \sim v_A \sim 30 - 100 \text{ km s}^{-1}$ in a MW-mass halo (one to a few times the Alfvén speed), increasing to a few 100 km s^{-1} in small-group mass halos ($\sim 10^{13} M_\odot$). For the typical SNe

rate in Milky Way to Andromeda-type galaxies (assuming the standard $\sim 10\%$ of the energy per SNe goes into CRs and $\sim 2\%$ of this into leptons; Higdon et al. 1998; Blasi & Amato 2012; Caprioli 2012) plus mean jet power (averaged over the whole population, since the travel time is \sim Gyr so we should average over any AGN outbursts in the last Gyr) for the population (scaled to either radio or X-ray power; see Allen et al. 2006), one expects a population-averaged $\dot{E}_{\text{cr},\ell} \sim 10^{39} - 10^{41} \text{ erg s}^{-1}$ increasing from Milky Way-to-Andromeda mass galaxies (see references in Su et al. 2024, and discussion of energetics in § 2.2.2 below). Inserting these, we obtain:

$$\frac{S_{X,\text{keV}}}{\text{erg s}^{-1} \text{ kpc}^{-2}} \sim 10^{35.3} \left(\frac{\dot{E}_{\text{cr},\ell}}{10^{40} \text{ erg s}^{-1}} \right) \left(\frac{100 \text{ km s}^{-1}}{v_{\text{st,eff}}(R)} \right) \quad (5)$$

$$\times \left(\frac{100 \text{ kpc}}{R} \right) f_{\text{loss}} E_{\text{cr,GeV}} (1+z)^4$$

$$\sim 10^{35.3} (1+z)^4 R_{100}^{-1} f_{\text{loss}} \frac{\dot{E}_{40}}{v_{100}}, \quad (6)$$

where $\dot{E}_{40} \equiv \dot{E}_{\text{cr},\ell}/10^{40} \text{ erg s}^{-1}$, $v_{100} \equiv v_{\text{st,eff}}(R)/100 \text{ km s}^{-1}$, $R_{100} \equiv R/100 \text{ kpc}$. This is remarkably similar to the observed scalings of S_X in Zhang et al. (2024a), both in normalization (for the expected parameters) and (shallow) slope, $S_X \propto R^{-1}$.

This comparison is shown in Fig. 2, where we compare the observed S_X around galaxies of different masses⁴ to the predictions of this toy model, adopting a couple different reasonable choices of the fiducial parameters $\dot{E}_{\text{cr},\ell}$ and $v_{\text{st,eff}}$. For each model prediction we assume the median quoted redshift of the galaxies in that stellar mass interval from Zhang et al. (2024a), namely $\langle z \rangle \approx (0.08, 0.12, 0.15)$ for (MW, M31, 2M31), and calculate f_{loss} self-consistently following § 2.2, for the given $v_{\text{st,eff}}$. We stress that these are not fits; we simply show a couple plausible values to illustrate the similarity of the expected and observed profiles.⁵

In Fig. 2 we also compare the “fiducial” **m12i** simulation from Hopkins et al. (2020, 2021b). This is a cosmological, multi-physics, galaxy+star formation simulation following the evolution of a MW-mass galaxy at high resolution with radiative cooling, magnetohydrodynamics, star formation, and stellar feedback using the Feedback In Realistic Environments (FIRE-2; Hopkins et al. 2014, 2018) methods. Therein, we model CRs as a single ultra-relativistic fluid obeying a simple transport model (which includes advection, streaming at the Alfvén speed, and diffusion with a constant parallel κ fixed to the LISM-inferred value). Since the simulation only follows the total CR energy (dominated by GeV protons), we assume an LISM-like proton-to-electron spectrum and correct for losses using the travel time t_{travel} estimated in-code for the GeV protons. We stress this is just one galaxy, with one (somewhat arbitrary, but empirically-motivated) CR transport model – it is not our intention to make a rigorous comparison here. But it shows that a profile similar to that observed emerges naturally from the simplest CR transport models which have

⁴ Here and throughout, we stress that we always focus on the diffuse CGM emission observed, after subtracting the central galaxy, AGN, XRBs, and other point-source contributions.

⁵ The simple expression in Eq. 5 assumes point-source injection of CRs with constant $v_{\text{st,eff}}$, and so clearly breaks down at small R around the ISM of galaxies. In Fig. 2 we simply truncate the profiles at $\sim 15 \text{ kpc}$ to represent this, but note we obtain a very similar result if we adopt instead a “streaming plus diffusion” model with constant $\kappa_{\text{eff}} \sim 1-3 \times 10^{29} \text{ erg s}^{-1} \text{ cm}^{-2}$ dominant when $\kappa_{\text{eff}} \gtrsim v_{\text{st,eff}} R$, or if we model a finite uniform-density source distribution of size $\sim 10 \text{ kpc}$, as is commonly assumed in codes like GALPROP. In any case, this is only important at ISM radii, not in the CGM.




FIG. 2.— Observed soft X-ray diffuse CGM emission, compared to expected IC from CRs. Shaded range shows *eROSITA* stacked mean CGM emission at 0.5 – 2 keV, around Milky Way (MW; $10^{10.5-11} M_{\odot}$), Andromeda (M31; $10^{11-11.25} M_{\odot}$), and twice M31 (2M31; $10^{11.25-11.5} M_{\odot}$) stellar-mass central, isolated galaxies (after subtracting point sources, AGN, and the XRB; Zhang et al. 2024a). Lines compare the predicted CR-IC emission, using the toy analytic model in § 2.1 (with IC losses accounted for per § 2.2), for values of the injection rate parameter $\dot{E}_{\text{cr},\ell} = \dot{E}_{40} 10^{40} \text{ erg s}^{-1}$ and CR streaming speed $v_{\text{st,eff}} = v_{100} 100 \text{ km s}^{-1}$ (labeled), expected for different mass galaxies (matching colors). We also compare the prediction from a fiducial cosmological CR-MHD simulation (cyan line) of a MW-mass galaxy (which used a simple CR diffusion+streaming model matched to the LISM) from Hopkins et al. (2020, 2021b).

been shown to reproduce a wide variety of LISM observations in the MW (see Chan et al. 2019, 2022; Ji et al. 2020; Hopkins et al. 2021b, 2022a). And we note that the predicted extended thermal emission in X-rays is much smaller (as expected) than the IC emission shown (e.g. van de Voort et al. 2016) – detailed comparison of the IC versus thermal X-ray emission in the CGM (and comparisons with the total X-ray emission including the central galaxy and XRBs) will be shown in Lu et al., in preparation.

Note that for our purposes, the entire ISM/galaxy is the effective CR source region. So e.g. stronger losses from synchrotron or Coulomb/ionization interactions *within* the ISM (e.g. near SNe remnants), or small-scale fluctuations in CR transport physics, are effectively “built into” the initial spectrum we assume (see discussion in Ponnada et al. 2024b). Hence adopting an LISM-like spectrum (appropriate for CRs already at the outskirts of the galaxy in the diffuse ISM, leaking into the CGM), rather than some theoretical CR acceleration spectrum at SNe shocks. This is also demonstrated quantitatively in the simulations above and in Fig. 2: those agree very well with the observed far infrared-radio (synchrotron) correlation of galaxies (with the emission there coming from denser regions in the ISM; Ponnada et al. 2024b,a), while also producing the extended CGM profiles shown.

2.1.1. Spectra and More Detailed Emissivity Calculations

Fig. 1 shows the resulting emissivity as a function of energy, properly integrating over the full CMB spectrum, convolved with the IC emissivity for each CR electron and positron, and then integrating over a full CR spectrum. For the CR spectrum, we show two different choices: first, we directly adopt a best-fit to the observed LISM spectrum, combining Voyager, Pamela, AMS-02, and other data sets as in Cummings et al. (2016); Adriani et al. (2018); Bisschoff et al. (2019), as

fit and parameterized in Hopkins et al. (2022a). The positron correction is also included but is small.⁶ The second choice attempts to correct for IC losses for higher-energy CRs, as described in § 2.2.

As expected the emissivity is large near ~ 1 keV. For a Solar-neighborhood CR spectrum, higher-energy CRs contribute significantly and make the spectrum harder from $\sim 1 - 10$ keV, with a spectral slope $dN/dt dE \propto E^{-\Gamma}$ with $\Gamma \sim 2.5$. But loss corrections likely make this spectrum much softer at higher energies. From this full calculation, we can revise our order-of-magnitude calculation of the emissivity in Eq. 2, integrated specifically in a given band. For $0.5 - 2$ keV (matching the eROSITA data) we obtain $\epsilon_X(0.5 - 2) \sim 2.6 \times 10^{34} \text{ erg s}^{-1} \text{ kpc}^{-3} (e_{\text{cr}, \ell} / 0.02 \text{ eV cm}^{-3}) (1+z)^4$ – nearly identical to equation 2 for $E_{\text{cr}, \text{GeV}} \sim 1$. The pre-factor in this estimate of the emissivity is no smaller than $\sim 1.5 \times 10^{34}$ (given the same *integrated* $e_{\text{cr}, \ell}$) accounting for different estimates of the LISM CR spectrum.

Even though the emissivity in Figure 1 is broadly thermal near the peak, a key difference between IC and thermal emission is that the former is of course pure continuum, while thermal emission at keV temperatures is actually dominated by lines. We do in fact expect some quite weak line emission in the spectrum, even if the emission is IC-dominated and the gas is actually uniformly cool ($< 10^6$ K) and low-density ($n < 10^{-4} \text{ cm}^{-3}$), so that the actual thermal contribution to the observed $\sim 0.5 - 2$ keV emission is negligible. The line emission will come from a mix of fluorescence/recombination/Auger cascades from both photo-excitation by the meta-galactic X-ray background (which still dominates the observed intensity in all but the innermost halo; Zhang et al. 2024b), as well as excitation and ionization by the cosmic rays themselves (not totally negligible for the CR energy densities here, as the integrated CR excitation rate for more-neutral Fe can exceed $\gtrsim 10^{-16} \text{ s}^{-1} (e_{\text{cr}, \text{tot}} / \text{eV cm}^{-3})$; Hsiung Chen & Crasemann 1985; Kaastra & Mewe 1993; Kovaltsov et al. 2001). Detailed calculations of these will depend on the temperature, metallicity, and combination of CR and XRB spectra and cross-sections for arbitrary electron and fluorescence yields, which are quite complicated, so we do not attempt them here, since the X-rays are only detected in broad-band stacks (resolved spectroscopy is only possible at orders-of-magnitude higher surface brightness, in e.g. massive cluster centers).

At present, the only detection of the low-surface-brightness emission around MW-Andromeda mass galaxies is broad-band integrated soft X-ray data, so there is no spectral information to distinguish the different lines in Fig. 1. We show the emissivity as a function of photon energy to highlight that the IC emission is surprisingly thermal-like (especially with IC losses accounted for), and to demonstrate that the emission peaks in the salient X-ray bands. Future measurements of (or non-detections of) diffuse-gas line emission with X-ray microcalorimetry (e.g. XRISM or ATHENA) may be able to place additional useful constraints on the CR-IC scenario. Since the line emission is sensitive to the product $\sim n_{\text{gas}}^2 Z$, upper limits or sufficiently-weak detections would further rule out cooling/line emission as the source of the soft X-ray halos at $\gtrsim 100$ kpc (as already implied by existing X-ray absorption

⁶ There is some factor ~ 2 uncertainty in the exact shape of the LISM spectrum at ~ 1 GeV owing to corrections for Solar modulation and lack of sensitivity of Voyager at these energies; but this is just implicit in $e_{\text{cr}, \ell}$. Likewise for the details of the positron correction.

measurements from Yao et al. 2010; Ponti et al. 2023). Ruling out CR-IC as the source of the halos is more challenging as it would require resolving almost all of the bolometric soft X-ray luminosity into individually detected lines, at very low surface brightness beyond $\gtrsim 100$ kpc around a significant number of MW-mass galaxies – but even if line cooling were the only source of emission, this is not likely to be possible, as most of the cooling luminosity emerges in pseudo-continuum from overlapping lines (e.g. Fe L-shell emission) while these instruments are primarily sensitive to a few spectrally-isolated narrow lines (Kraft et al. 2023; ZuHone et al. 2024).

2.1.2. Redshift Effects & Measuring CR Energy Density

Note that there are two redshift-dependencies which appear in our IC scaling (§ 2.1), both arising from the CMB: (1) the peak of the emissivity in the rest frame scaling $\propto (1+z)$, and (2) the surface brightness scaling $\propto (1+z)^4$. What is worth noting is that in the *observed* frame, these are exactly canceled by cosmological redshift and surface-brightness dimming, respectively. This means that (1) the peak of the *observed* X-ray emission will be at $\sim \text{keV}$:

$$h\nu_{\text{peak}}^{\text{obs}} \sim \text{keV}, \quad (7)$$

and (2) the observed/apparent surface brightness (for the same *physical* parameters in Eq. 3) will be *independent* of the source redshift:

$$\frac{S_{X, 0.5-2\text{keV}}^{\text{obs}}}{10^{37} \text{ erg s}^{-1} \text{ kpc}^{-2}} \sim f_0 \left(\frac{e_{\text{cr}, \ell}}{0.02 \text{ eV cm}^{-3}} \right) \left(\frac{R}{100 \text{ kpc}} \right), \quad (8)$$

where the range of pre-factors $f_0 \sim 0.3 - 0.8$ accounts for the range of more detailed spectral shape calculations in § 2.1.1. The soft X-ray bands are therefore always optimal for observing IC X-ray emission off of the CMB, assuming the LISM CR spectrum is typical.

If the IC interpretation is correct, then, measuring the CGM ~ 1 keV surface brightness *directly* measures the CR lepton energy density around $\sim \text{GeV}$ energies, in a unique manner unlike any other constraint (like synchrotron or γ -ray constraints, which are strictly degenerate with the magnetic field strengths and gas densities and depend qualitatively differently on redshift). In other words, one can directly read off:

$$\frac{e_{\text{cr}, \ell}}{0.02 \text{ eV cm}^{-3}} \sim \frac{S_{X, 0.5-2\text{keV}}^{\text{obs}}}{f_0 10^{37} \text{ erg s}^{-1} \text{ kpc}^{-2}} \frac{100 \text{ kpc}}{R}. \quad (9)$$

This is shown in Fig. 3.⁷ In theoretical *models* for the CR propagation, this in turns constrains the product $f_{\text{loss}} \dot{E}_{\text{cr}, \ell} / v_{\text{st}, \text{eff}}$ (Eq. 5), $f_{\text{loss}} \dot{E}_{40} / v_{100} \sim (S_{X, 0.5-2\text{keV}}^{\text{obs}} / 10^{35.3} \text{ erg s}^{-1} \text{ kpc}^{-2}) R_{100}$.

Note the rest-frame scaling of S_X with $(1+z)^4$ means the IC loss time also scales $\propto (1+z)^{-4}$, which means (§ 2.2) the maximum extent of the IC halo (before being loss-limited) scales will be smaller at higher redshift (modulo possible scaling of varying streaming/diffusion speeds and reacceleration efficiencies, but these should not evolve so rapidly in most

⁷ Our $e_{\text{cr}, \ell} \propto S_X / R$ scaling above assumes a roughly similar power-law decrease of S_X with R , which technically becomes invalid at small R around the ISM ($\ll 20$ kpc) where the observed profiles become flat in Fig. 2. More accurately, in Fig. 3 we de-project assuming spherical symmetry with the standard Abel integral. We obtain a nearly-identical result if we de-project assuming the best-fit spherical β profile model in Zhang et al. (2024a). But in any case this is only important at $\ll 20$ kpc.




FIG. 3.— Leptonic CR energy density around \sim GeV required to explain the observed soft X-ray surface brightness, for each of the three observed stacks in Fig. 2. The central (galactic) values are similar to the observed LISM values (Bischoff et al. 2019) in the MW-mass systems, and those inferred from multi-wavelength modeling of M31’s synchrotron plus γ -ray emission (Lacki et al. 2010), with a power-law falloff and eventual cutoff from IC losses at larger radii.

models). So while the halo *luminosity* remains independent of redshift, the effective maximum extent (in physical units) will be smaller at high redshifts – presenting a potential test of the IC scenario. This redshift-dependence will be quite difficult to detect, however, for two reasons. First, it would require exceptional angular resolution: noting that eROSITA only barely detects the MW-mass halos above their PSF wings at $z \sim 0.08$ (Zhang et al. 2024a), the combination of shrinking physical size and rapidly-rising angular diameter distance means that measuring this predicted trend at $z \gtrsim 0.5$ would require more than an order-of-magnitude better angular resolution (e.g. a $\lesssim 1''$ PSF). Second, the X-ray background also scales $\propto (1+z)^4$, or equivalently surface-brightness dimming scales as $(1+z)^{-4}$; so if one had a halo powered by thermal emission at higher redshift, its *detectable* size above some threshold relative to the backgrounds would also shrink. To properly compare, one would need signal-to-noise scaling as $\propto (1+z)^4$ at higher redshifts. These requirements push the limits even of potential future missions like AXIS.

2.2. (Inverse Compton) Losses

The IC lepton loss timescale $t_{\text{loss}} \equiv p_{\text{cr}}/\dot{p}_{\text{cr}}^{\text{IC}} \sim 1.2(1+z)^{-4} E_{\text{cr, GeV}}^{-1}$ Gyr should eventually become comparable to or shorter than the CR escape/diffusion/travel time to some radius $t_{\text{travel}} \equiv R/v_{\text{st, eff}} \sim 1(R/100 \text{ kpc})(100 \text{ km s}^{-1}/v_{\text{st, eff}})$ Gyr, i.e. $\tau_{\text{loss}} \equiv t_{\text{travel}}/t_{\text{loss}} \sim E_{\text{cr, GeV}} R_{100} (1+z)^4/v_{100}$. This depends on the highly-uncertain CGM streaming/diffusion speeds, so for faster CR transport it could be a small correction, but generically we would expect that at sufficiently large radii and CR energies, $\tau_{\text{loss}} \gg 1$, and the CR spectrum will be suppressed.

It is easy to verify from the scalings above that – for the energies of interest – other CR losses in the CGM/IGM are negligible, including catastrophic/hadronic/pionic, ionization, synchrotron, bremsstrahlung, ionization, and Coulomb (e.g. $t_{\text{loss}}^{\text{Coulomb}} \sim 5000 \text{ Gyr } E_{\text{cr, GeV}} n_{-5}^{-1}$). The “streaming losses” and “adiabatic/ PdV work” terms, which generalize the classic “turbulent reacceleration terms” can be non-zero but in numerical

simulations are found on average to rarely be more than $\mathcal{O}(1)$ corrections to the CR energy density and emissivity, depending on the structure of the turbulence and magnetic fields in the outer halo (Chan et al. 2019; Buck et al. 2020; Hopkins et al. 2022a). But they could modify the details of the outer cutoff and should be studied in future work.

Note in terms of heating the *gas*, the IC radiation is rather inefficient, though a small fraction of the lowest-energy IC photons could go into ionizing higher metal-line transitions. Depending on the details of the CR-gas interactions at the gyro-scale, the CR “streaming heating” term will likely be the most important for heating the gas, though simulations with similar e_{cr} as implied by the X-ray halos generally find this is sub-dominant to gas cooling by factors of $\sim 10 - 100$ in low-mass ($\lesssim 10^{13} M_{\odot}$) halos (Ji et al. 2021, 2022; Martin-Alvarez et al. 2024; Dacunha et al. 2024).

2.2.1. Modification of the CR Spectrum

The evolution of a CR lepton with initial energy $E_{\text{cr}, 0}$ and IC loss time $t_{\text{loss}, 0}$ in a uniform radiation background with only IC losses has the simple solution: $E_{\text{cr}} = E_{\text{cr}, 0}/(1+t/t_{\text{loss}, 0})$. If we assume all the CRs are injected at $r = 0$ and travel together (i.e. pure streaming), then this gives the modified spectrum $dn_{\text{cr}}[E_{\text{cr}}]/d \ln E_{\text{cr}} = dn_{\text{cr}}[E_{\text{cr}}^0]/d \ln E_{\text{cr}}^0 (1+t/t_{\text{loss}, 0})^{-1}$, which depends only on the parameter $\tau \equiv t/t_{\text{loss}, 0}$. The IC-modified spectra are shown in Fig. 1 for different values of the parameter:

$$\tau_{\text{loss}}^0 \equiv \frac{t_{\text{travel}}}{t_{\text{loss}}} \Big|_{\text{GeV}} \sim \frac{0.8}{(1+f_{\text{reacc}})} \frac{R_{100}}{v_{100}} (1+z)^4 \quad (10)$$

Note the spectrum formally depends (1) on how we assume v_{st} depends on CR energy;⁸ (2) the mix of streaming and diffusion and inhomogenous CR transport, which will broaden the “cutoff” in the IC-modified spectrum because of different travel times at a given radius; and (3) the mix of streaming (CR gyroresonant instability) losses, adiabatic CR gains/losses, and turbulent and diffusive re-acceleration (parameterized above with f_{reacc}) during CR transport (see e.g. Gaggero et al. 2015; Hopkins et al. 2021c, 2022b; Korsmeier & Cuoco 2022; Di Mauro et al. 2024). All of these effects introduce only $\mathcal{O}(1)$ changes at most in the total emissivity at ~ 1 keV, since they mostly just modify how rapidly the already-rapid cutoff in the spectrum occurs. Regardless of how these details modify our calculation, we obtain the expected result that when $\tau_{\text{loss}}^0 \gg 1$, the CR spectrum must be strongly modified from its injection spectrum and most of the energy has been lost (so the emissivity at larger R must be small).

Note, as discussed above, that as IC losses modify the spectrum, it becomes softer and more similar to thermal spectra. The measured emissivity spectrum for loss parameter $\tau_{\text{loss}}^0 = 1$, for example, would look similar to gas with $T \approx 1.5 \times 10^7$ K, while at $\tau_{\text{loss}}^0 = 10$, it closely resembles the spectrum of $\sim 10^6$ K gas. At present, observations of the diffuse emission in MW-Andromeda mass galaxies cannot constrain the spectra (just the integrated soft X-ray surface brightness), but in principle this is relevant for future searches, and it informs

⁸ We have compared assuming a constant streaming speed or diffusivity, or allowing v_{st} to depend on energy with some dependence (faster diffusive transport of high-energy CRs) like that inferred in the LISM, $v_{\text{st}} \propto 1+E_{\text{cr, GeV}}^{1/2}$, and find these make only modest quantitative differences to the spectra and do not change any of our qualitative results.




FIG. 4.— Total CGM halo soft X-ray luminosity, $L_{X,\text{keV}}^{\text{tot}} = \int S_X dA$, as a function of central galaxy stellar mass M_* , compared to the expected IC luminosities. Observed values from are shown from stacked ROSAT data (Anderson et al. 2015 CGM only-emission after subtracting XRB+galaxy contributions), as well as eROSITA, using the same brightness profiles from Zhang et al. (2024a) in Fig. 2 and integrating out to radii where any excess emission is detected (i.e. allowing for an IC cutoff; *filled*; “detected”), or using the stacking method in Zhang et al. (2024b) which extrapolates the inferred profiles from smaller radii out to R_{500} in all cases (*open*; “ $R \rightarrow R_{500}$ ”). We compare the predicted time-and-population-averaged SNe \sim GeV leptonic CR injection rate and AGN injection rate (assuming empirical SFR- M_* and $\langle M_{\text{BH}} \rangle - M_*$ relations, and that $\sim 10^{-4} \dot{M}_{\text{BH}} c^2$ goes into leptons, for reference; see § 2.3) from just the central galaxies. These scalings have factor $\sim 2-3$ systematic uncertainties. IC from standard sources (SNe and/or AGN even with very small efficiencies) can naturally explain the halo luminosities at $M_* \lesssim (1-2) \times 10^{11} M_{\odot}$.

our predictions for associated harder X-ray and synchrotron emission below (§ 2.4).

2.2.2. Total Luminosity and Extent of Halos: Estimating the CR Injection Rate and Streaming Speed

Given IC losses, at some radius the CR surface brightness should rapidly truncate, approximately as $\propto \exp(-\tau_{\text{loss}}^0)$. The more robust way of saying this is that once the integrated IC luminosity out to some radius R , $L_{\text{keV}}^{\text{tot}}(R) \equiv \int_0^R S_X(R) \pi R dR$, exceeds the leptonic GeV CR source injection rate $\dot{E}_{\text{cr},\ell}$ (modulo some order-unity correction for other losses and adiabatic/diffusive/turbulent reacceleration, $\sim 1 + f_{\text{reacc}}$), then the CRs must be depleted to lower energies and their IC from the CMB must shift out of the keV band,⁹ so S_X drops rapidly and L_{keV} cannot rise further. The radius where this will occur is just the radius where $\tau_{\text{loss}}^0 \gtrsim$ a couple, i.e. $R \gtrsim R_{\text{crit}} \sim 100 \text{ kpc} (1+z)^{-4} v_{100} (1+f_{\text{reacc}})/0.8$ (Eq. 10). Another way of saying the above is that if we insert R_{crit} into Eq. 5, then we trivially recover $L_{\text{keV}}(R \rightarrow R_{\text{crit}}) \sim \pi R_{\text{crit}}^2 S_X(R_{\text{crit}}) \sim \dot{E}_{\text{cr},\ell}$.

Therefore, if one can *measure* a truncation or “cutoff” radius where S_X exhibits some curvature, one can directly infer the CR streaming speed (since the lifetime is known for IC against the CMB) as $v_{100} \sim R_{\text{crit}} (1+z)^4 (0.8/\tau_{\text{loss}}^0, \text{crit} (1+f_{\text{reacc}}))$. This then, with Eq. 5, allows one to measure $\dot{E}_{\text{cr},\ell}$.

Or equivalently, if one can just measure the asymptotic IC luminosity of the halos, L_{keV} , then one immediately obtains

⁹ There could be some X-ray radiation from lower-energy CRs scattering the cosmic infrared/optical backgrounds, but the photon number density of these backgrounds and corresponding X-ray emissivity is lower than for the CMB by factors of $\sim 10^3 - 10^6$, so we neglect them throughout.

the leptonic injection rate

$$(1 + f_{\text{reacc}}) \dot{E}_{\text{cr},\ell} \sim L_{X,\text{keV}}^{\text{tot}}. \quad (11)$$

This is shown, as a function of galaxy mass, in Fig. 4. Importantly, the observed luminosities are the CGM luminosities *after* subtraction of XRBs, AGN, and any other central source (e.g. hot gas in the galactic ISM). As shown in many studies including those plotted (e.g. Anderson et al. 2015; Zhang et al. 2024a), in low-mass galaxies the total X-ray luminosity *including* the galaxy and all sources is dominated by a combination of HMXBs+AGN in the central $\lesssim 5$ kpc (and simulation comparisons with these are discussed in e.g. van de Voort et al. 2016; Chan et al. 2022 and Lu et al., in prep.).

With this estimate of $\dot{E}_{\text{cr},\ell}$, then from the measured S_X interior to the outer/asymptotic radii ($R \ll R_{\text{crit}}$, where the loss corrections should not be very large) we can infer the CR effective streaming speed or diffusivity:

$$v_{\text{st,eff}}(R < R_{\text{crit}}) \sim \frac{\kappa_{\text{eff}}(R)}{R} \sim 200 \text{ km s}^{-1} \frac{L_{X,\text{keV},40}^{\text{tot}}}{S_{X,35} R_{100}}, \quad (12)$$

$$\kappa_{\text{eff}} \sim R v_{\text{st,eff}}(R) \sim 6 \times 10^{30} \text{ cm}^2 \text{ s}^{-1} \frac{L_{X,\text{keV},40}^{\text{tot}}}{S_{X,35}}, \quad (13)$$

where $S_{X,35} \equiv S_{X,0.5-2\text{keV}}^{\text{obs}}/10^{35} \text{ erg s}^{-1} \text{ kpc}^{-2}$, $L_{X,\text{keV},40}^{\text{tot}} \equiv L_{X,\text{keV}}^{\text{tot}}/10^{40} \text{ erg s}^{-1}$, $R_{100} \equiv R/100 \text{ kpc}$. These are shown in Fig. 5.

Note that for more massive groups and clusters, we should properly account for injection from many galaxies throughout the halo, as the stellar mass and total AGN luminosity becomes increasingly dominated by the sum of galaxies rather than the BCG in clusters (and these, being injected at finite R , will suffer far less IC losses). But in the lower-mass halos of interest here like in the Local Group, the total satellite luminosity/mass is much lower than that of the central (Geha et al. 2024).

2.3. Observational Constraints: CR Energies and Streaming Speeds

Zhang et al. (2024a) measure¹⁰ the stacked 0.5–2 keV CGM surface brightness around three stellar mass-selected samples of galaxies (compared in Fig. 2): their “Milky Way mass” (MW) sample ($M_* \sim 10^{10.5-11} M_{\odot}$), “Andromeda mass” (M31) sample ($M_* \sim 10^{11-11.25} M_{\odot}$), and “twice Andromeda mass” (2M31) sample ($M_* \sim 10^{11.25-11.5} M_{\odot}$). The median central-galaxy stellar mass in these (MW, M31, 2M31) samples is $\sim (0.55, 1.3, 2.2) \times 10^{11} M_{\odot}$.

From the observations, we can estimate $L_{X,\text{keV}}^{\text{tot}}$ in Fig. 4 by integrating along the lower and upper limits on the stacked profiles out to the radii where they become undetectable against the X-ray background (where the lower limits

¹⁰ We compare to the eROSITA data from Zhang et al. (2024a) as the improved sensitivity allows much more detailed model constraints compared to older ROSAT data (which are unable to spatially-resolve the halos around lower-mass systems) in (Anderson et al. 2013, 2015), but note that the integrated CGM/halo luminosities as a function of stellar mass (and halo sizes, where ROSAT can detect and spatially-resolve them) are broadly consistent with one another. We also focus on the stellar mass-selected samples in Zhang et al. (2024a), rather than their halo-mass selected samples, as selection in the latter is model-dependent and can give very different results depending on the scatter in the $M_* - M_{\text{halo}}$ relation (see e.g. Popesso et al. 2024b,a), and exclusively compare their background and point-source subtracted best estimate of the diffuse halo emission. When comparing L_X and M_* , we correct the stellar masses of Anderson et al. (2015) by the same small (0.1 dex) factor as Zhang et al. (2024b) to align their definitions.




Fig. 5.— Implied effective CR diffusivity κ_{eff} , needed to explain the observed S_X profiles given the IC loss timescale and injection rate $\dot{E}_{\text{cr}} \sim L_{X, \text{keV}}^{\text{tot}}$ (§ 2.2.2): $\kappa_{\text{eff}} \sim 6 \times 10^{30} \text{ cm}^2 \text{ s}^{-1} L_{X, \text{keV}, 40}^{\text{tot}} S_{X, 35}^{-1}$. Approaching the ISM the values in MW/M31 are very similar to those inferred from MW LISMs observations from Voyager and AMS-02 (De La Torre Luque et al. 2021; Korsmeier & Cuoco 2022; labeled); at larger radii this rises as expected. We compare the independent constraints on $\sim \text{GeV}$ protons from a completely independent UV absorption-line method applied to cool neutral HI gas (which should not appear in the X-rays) in Butsky et al. (2023), colored by stellar mass (blue matching Z24’s MW-mass sample, grey for masses outside this).

fall below the plot). In principle a lower-level background halo could persist, but this seems to be indicative of some steepening in the profiles as shown in Fig. 2. We obtain $L_{X, \text{keV}, 40}^{\text{tot}} \sim (0.08 - 0.8, 0.8 - 4, 6 - 14)$ for (MW, M31, 2M31).¹¹ For the streaming speeds at $R \lesssim R_{\text{crit}}$, we obtain at $R \sim 10 \text{ kpc}$, $v_{\text{st, eff}} \sim (25 - 80, 80 - 170, 100 - 400) \text{ km s}^{-1}$, or $\kappa_{\text{eff}} \sim (0.8 - 2.4, 2 - 5, 3 - 12) \times 10^{29} \text{ cm}^2 \text{ s}^{-1}$ for (MW, M31, 2M31). At $R \sim 100 \text{ kpc}$, this implies $v_{\text{st, eff}} \sim (80 - 160, 100 - 200, 120 - 160) \text{ km s}^{-1}$ or $\kappa_{\text{eff}} \sim (2 - 5, 3 - 6, 3.5 - 5) \times 10^{30} \text{ cm}^2 \text{ s}^{-1}$, shown in Fig. 5.

2.3.1. Comparison to Other Empirical Constraints and Models

The CR electron injection rates required to explain the X-ray halos of MW-Andromeda mass galaxies via IC emission are quite similar to what is observed and expected in local galaxies from standard sources of CRs. To estimate this, consider CRs accelerated by (1) SNe and (2) AGN (and/or associated jets/bubbles/winds). For (1), we can sum core-collapse and prompt Ia rates (proportional to galactic star formation rates, $\dot{N}_{\text{prompt}} \sim 0.014 (\dot{M}_*/M_\odot)$; Sukhbold et al. 2016; Hopkins et al. 2023a) and delayed Ia rates (proportional to stellar mass $\dot{N}_{\text{late}} \sim 1.5 \times 10^{-13} \text{ yr}^{-1} (M_*/M_\odot)$; Maoz & Graur 2017; Gandhi et al. 2022) using the fact that for stacked samples in these mass ranges, the *mean* (what matters for the stacked result) SFRs and stellar masses are well-known. For stellar mass, $\langle M_* \rangle \sim (0.55, 1.3, 2.2) \times 10^{11} M_\odot$ is defined by the (MW, M31, 2M31) samples above; the mean SFRs are given by the “main sequence” ($\dot{M}_* - M_*$ correlation) average (dominated by the rapidly star-forming population) $\langle \dot{M}_* \rangle \sim (3, 7, 11) M_\odot \text{ yr}^{-1}$ (Pearson et al. 2018; Cooke et al. 2023). If we assume the standard empirically-estimated

¹¹ This is different from the method of estimating $L_{X, \text{CGM}}$ in Zhang et al. (2024b), who extrapolate a fitted or upper limit to S_X out to R_{500} , which leads to factor of a few larger $L_{X, \text{CGM}}$ for the low-mass halos. It is more akin to the estimates in Anderson et al. (2015).

$\sim 10^{50}$ erg of total CR energy accelerated per SNe ($\sim 10\%$ of the ejecta energy), with $\sim 2\%$ of that in $\sim \text{GeV}$ leptons, we obtain $\langle \dot{E}_{\text{SNe}, \ell} \rangle \sim (0.3, 0.8, 1.2) \times 10^{40} \text{ erg s}^{-1}$. So for the lower-mass (MW, M31) samples, the observed halo luminosity is entirely consistent with the expectation from SNe-accelerated leptons. The AGN contribution is more uncertain, but the mean bolometric AGN luminosity and accretion rates are empirically known for galaxies with these stellar masses (and/or SFRs). Here it is more important that we account for the mean, not median, since (given the large range of L_{AGN} at a given M_*) more luminous systems should dominate the injection. From Torbaniuk et al. (2021, 2024), for these mass ranges, we have $\langle \dot{M}_{\text{BH}} \rangle \sim (2.5 - 3, 5.3 - 7.1, 8.8 - 16) \times 0.001 M_\odot \text{ yr}^{-1}$, or an available accretion energy $\langle \dot{E}_{\text{BH}} \rangle \sim (1.4 - 1.7, 3 - 4, 5 - 9) \times 10^{44} \text{ erg s}^{-1}$. So powering the observed halos via AGN injection of CRs would require only a small fraction $\epsilon_{\text{cr}, \ell} \sim (0.4 - 6, 2 - 12, 6 - 28) \times 10^{-5}$ of the accretion energy ultimately goes into accelerating leptons – a completely plausible estimate from previous constraints in radio AGN (Falcke et al. 2004; Bîrzan et al. 2004; Allen et al. 2006; McNamara & Nulsen 2007). In both of these, we could also include the contribution of satellite galaxies, which should grow more important at large radii and more massive halos, but this is expected to be small for MW-M31 mass halos.

Briefly, it is worth noting that the *galaxy* (not diffuse CGM) X-ray luminosity of star-forming (low-mass) galaxies observationally scales as (Mineo et al. 2014)

$$L_X^{\text{galaxy}}(0.5 - 8 \text{ keV}) \approx 4 \times 10^{39} \text{ erg s}^{-1} \left(\frac{\dot{M}_*}{M_\odot \text{ yr}^{-1}} \right), \quad (14)$$

which is expected from the rate of both SNe (creating hot gas) and XRBs (with e.g. Mineo et al. 2012 attributing $\sim 2/3$ of this to resolved HMXBs). Ignoring AGN, the scalings above for CR injection give $L_{X, \text{IC}} \sim \dot{E}_{\text{cr}, \ell} \sim 10^{39} \text{ erg s}^{-1} (\dot{M}_*/M_\odot \text{ yr}^{-1})$. So the expectation is that the extended IC X-ray halo luminosity should scale with (and be within a factor of a few of) the central galaxy/XRB luminosity in star-forming galaxies, although the IC emission is much more diffuse and extended, as indeed appears to be observed.

Regarding AGN, note that we do not expect CR halos powered by BHs/AGN to necessarily correlate with *ongoing* AGN activity. Not only is the stack population-averaged, but the CR travel time to the radii of interest is $\sim \text{Gyr}$ as noted above, much longer than the lifetime of AGN accretion episodes (Martini 2004; Hopkins et al. 2005a,b,c,d; Kelly et al. 2010), so the effective CR injection luminosity is averaged over the whole duty cycle of AGN at low redshifts.

For the implied streaming speeds/diffusivities, the physical uncertainties in theoretical predictions are much larger (Zweibel 2017; Hopkins et al. 2021c, 2022b), but our inferred values are quite reasonable. At $\lesssim 10 \text{ kpc}$, the effective diffusivities/streaming speeds at $\sim \text{GeV}$ are in fact quite close to the empirically-favored LISMs values (from direct calibration to Voyager, AMS, and other CR experiments) $\kappa_{\text{eff}} \sim 1 - 3 \times 10^{29} \text{ cm}^2 \text{ s}^{-1}$ (see e.g. Génolini et al. 2019; De La Torre Luque et al. 2021; Korsmeier & Cuoco 2021, 2022; Hopkins et al. 2022a; Di Mauro et al. 2024). For the outer halo ($\sim 100 \text{ kpc}$), constraints are much more sparse, but Fig. 5 shows that for MW-mass galaxies, the implied κ_{eff} agrees well with the empirical constraints from Butsky et al. (2023) around MW-mass galaxies. What is remarkable is that the latter is derived from an entirely independent technique

and dataset: using indirect arguments about pressure balance in the CGM combined with UV absorption line spectra from background quasars dominated by cool ($< 10^5$ K) gas (indeed primarily HI absorption, so neutral gas at $\sim 10^4$ K) – with completely negligibly thermal X-ray emission – to constrain \sim GeV proton diffusivities. These are also similar to plausible streaming speeds in theoretical models (Wiener et al. 2013, 2017; Ruszkowski et al. 2017; Holguin et al. 2019; Kempster & Quataert 2022; Thomas et al. 2023; Ponnada et al. 2024c).

2.4. Associated Radio and γ -ray Halos

If the observed X-ray emission is indeed non-thermal, it is important to calculate the observed emission in other channels. First consider the radio (synchrotron) emission. This will be doubly-suppressed by weak magnetic (B) fields in the outer halo and IGM (Ponnada et al. 2022, 2024b), $B \sim 30 B_{30}$ nG (Dolag et al. 2011; Vernstrom et al. 2019; Tjemsland et al. 2024).¹² The bolometric synchrotron luminosity will be suppressed relative to the IC by the ratio $e_B/e_{\text{CMB}} \sim 7 \times 10^{-5} B_{30}^2 (1+z)^{-4}$, and the characteristic frequency $\nu_{\text{cr}} = (3/2)\gamma^2 \nu_G \sin \alpha \sim 0.4 \text{ MHz } B_{30} E_{\text{cr, GeV}}^2$. So powering 1.4 GHz emission, for example, requires $E_{\text{cr, GeV}} \sim 60 B_{30}^{-1/2}$ – i.e. extremely high-energy electrons where the CR number density is strongly suppressed even assuming an LISM spectrum (let alone correcting for IC losses in the halo). Neglecting those losses or simply parameterizing them with some term $f_{\text{loss}}^{\sim 100 \text{ GeV}} \ll 1$, we obtain a 1.4 GHz emissivity $\epsilon_{1.4 \text{ GHz}} \sim 10^{29.3} \text{ erg s}^{-1} \text{ kpc}^{-3} B_{30}^{9/4} (f_{\text{loss}}^{\sim 100 \text{ GeV}} e_{\text{cr, } \ell} / 0.02 \text{ eV cm}^{-3})$, and flux $S_{1.4 \text{ GHz}} \sim 10^{31.5} \text{ erg s}^{-1} \text{ kpc}^{-2} B_{30}^{9/4} R_{100} f_{\text{loss}}^{\sim 100 \text{ GeV}} \times (e_{\text{cr, } \ell} / 0.02 \text{ eV cm}^{-3})$, or integrated luminosity

$$\frac{L_{1.4 \text{ GHz}}}{\text{W Hz}^{-1}} \sim 10^{17.8} B_{30}^{9/4} R_{100}^3 \left(\frac{e_{\text{cr, } \ell} f_{\text{loss}}^{\sim 100 \text{ GeV}}}{0.0002 \text{ eV cm}^{-3}} \right), \quad (15)$$

$$\sim 10^{18.2} f_{\text{loss}}^{\sim 100 \text{ GeV}} B_{30}^{9/4} L_{\text{X, keV, 40}}^{\text{tot}},$$

about $\sim 3 - 4$ dex fainter than even the lowest-luminosity detected synchrotron galaxy disks/bulges (Magnelli et al. 2015; Delhaize et al. 2017; Wang et al. 2019) even if $f_{\text{loss}}^{\sim 100 \text{ GeV}} = 1$ (no losses). To be detectable from $R \sim 100 \text{ kpc}$ to $R \sim R_{\text{vir}}$ in terms of either surface brightness or luminosity, even within distances $< 10 \text{ Mpc}$ (let alone the X-ray sample distances which are at cosmological redshifts $z \sim 0.05 - 0.2$, i.e. luminosity distances $\sim 250 - 1000 \text{ Mpc}$), these would require $B \gtrsim \text{few } \mu\text{G}$, $e_{\text{cr, } \ell} \gtrsim 0.02 \text{ eV cm}^{-3}$, and $f_{\text{loss}}^{\sim 100 \text{ GeV}} \sim 1$ i.e. ISM-level magnetic field strengths and CR energy densities without any IC losses. These are orders-of-magnitude larger than any realistic model would predict at $> 100 \text{ kpc}$ distances from the central galaxies, except in massive clusters where $100 \text{ kpc} < 0.1 R_{\text{vir}}$ and these numbers are plausible (precisely where radio synchrotron halos are indeed observed; Gitti 2016). And again, this is an *upper limit* neglecting losses: because 1.4 GHz is so sensitive to high-energy leptons, the

¹² At $R \sim 100 \text{ kpc}$ around MW and M31-mass galaxies, cosmological magnetohydrodynamic+galaxy formation simulations predict a range $\langle |\mathbf{B}|^2 \rangle^{1/2} \sim (10 - 100) \text{ nG}$ (Marinacci et al. 2015; Katz et al. 2019; Ponnada et al. 2022; Mannings et al. 2023), with observational upper limits around such galaxies from Faraday rotation at $R \sim 100 \text{ kpc}$ of $B < 200 - 1000 \text{ nG}$ (Prochaska et al. 2019; Prochaska & Zheng 2019; Lan & Prochaska 2020) and at $R \sim \text{Mpc}$ upper limits are $B < (2 - 20) \text{ nG}$ (Ravi et al. 2016; Vernstrom et al. 2019; O’Sullivan et al. 2020; Padmanabhan & Loeb 2023).

fluxes are reduced (for the same *total* leptonic energy $e_{\text{cr, } \ell}$) by an additional ~ 5 dex ($f_{\text{loss}}^{\sim 100 \text{ GeV}} \sim 10^{-5}$) for even a modest IC correction (unless one could observe at much lower $\lesssim \text{MHz}$ frequencies, tracing the GeV leptons).

Because of this suppression, and the further, additional suppression factor by the scattering rates above and $n_{\text{photons, synch}}/n_{\text{photons, CMB}}$, it follows that synchrotron self-Compton contributes negligibly in the X-rays. And owing to the low gas densities in the outer halos, the non-thermal CR bremsstrahlung emission is also extremely faint.

If we assume an LISM-like proton spectrum and proton-to-electron ratio, the γ -ray emissivity (accounting for all the salient hadronic interactions, appropriate branching ratios, and spectrum at $\sim 0.1 - 10 \text{ GeV}$ energies) is $\approx 1.8 \times 10^{-28} \text{ erg s}^{-1} \text{ cm}^{-3} (e_{\text{cr, tot}} / \text{eV cm}^{-3}) (n_{\text{gas}} / \text{cm}^{-3})$ or $5.3 \times 10^{31} \text{ erg s}^{-1} \text{ kpc}^{-3} (e_{\text{cr, tot}} / \text{eV cm}^{-3}) (n_{\text{gas}} / 10^{-5} \text{ cm}^{-3})$ (Guo & Oh 2008; Lacki et al. 2011; Wiener et al. 2013; Hopkins et al. 2021c), so (assuming the typical profiles above) $S_{\gamma} \sim 8 \times 10^{33} \text{ erg s}^{-1} \text{ kpc}^{-2} (e_{\text{cr, tot}} / \text{eV cm}^{-3}) (n_{\text{gas}} / 10^{-5} \text{ cm}^{-3}) R_{100}$ and

$$\frac{L_{\gamma}}{\text{erg s}^{-1}} \sim 2 \times 10^{36} R_{100}^3 \left(\frac{e_{\text{cr, tot}}}{0.01 \text{ eV cm}^{-3}} \right) \left(\frac{n_{\text{gas}}}{10^{-5} \text{ cm}^{-3}} \right),$$

$$\sim 6 \times 10^{36} \left(\frac{f_{\text{cr, } \ell}^{\circ}}{f_{\text{cr, } \ell}} \right) \left(\frac{n_{\text{gas}}}{10^{-5} \text{ cm}^{-3}} \right) L_{\text{X, keV, 40}}^{\text{tot}}, \quad (16)$$

where we have scaled to a typical gas density seen in simulations at virial radii (Butsky & Quinn 2018; Ji et al. 2020, 2021; Ramesh et al. 2024). For comparison, the sensitivity of *Fermi* is such that outside of $< 1 \text{ Mpc}$, the lowest-luminosity detectable systems (NGC253 and M82, both in the local group) have $L_{\gamma} > 10^{40} \text{ erg s}^{-1}$ (Lacki et al. 2011; Lopez et al. 2018; Zhang et al. 2019), and for systems in the same range of distances/redshifts as the ROSAT or eROSITA samples, detection typically requires $L_{\gamma} \gg 10^{41} - 10^{42} \text{ erg s}^{-1}$ (Ackermann et al. 2014). So the extended γ -ray halo would be completely undetectable unless, again, we assumed an un-realistically large $e_{\text{cr, tot}}$ and n_{gas} for MW-Andromeda like host halos out to even larger radii. Once again, though, the predictions do become interesting for the central regions of massive clusters, where non-trivial constraints from γ -rays do exist (e.g. Ackermann et al. 2014), although these would need to be revisited in an IC+CR-dominated scenario (since the gas densities may be very different from what was assumed).

There are of course two systems of interest *within* $< 1 \text{ Mpc}$, namely the MW and M31, where some diffuse γ -ray emission has been detected. In Fig. 6 we show the profiles measured by Fermi from ring analysis and windows constraining the outer Galaxy (Abdo et al. 2010; Ackermann et al. 2011; Tibaldo 2014; Tibaldo et al. 2015, 2021; Acero et al. 2016; Yang et al. 2016; Pothast et al. 2018),¹³ and the detections of M31 both on ISM scales and the “spherical/outer halo” diffuse emission (Abdo et al. 2010; Karwin et al. 2019; see also Recchia et al. 2021; Do et al. 2021).¹⁴ Note the large range of radii in both cases. We compare these to the predicted γ -ray

¹³ Compiled in Hopkins et al. (2022a), using the median N_{HI} in those maps to convert from γ -ray emissivity to surface brightness where necessary.

¹⁴ The outer radii constraints are unaffected by the central source model discussion in Xing et al. (2023). For the spectra in Fig. 6, we follow Do et al. (2021) and focus on the M31 “S” window which Karwin et al. (2019) argue is less affected by foreground contamination.




FIG. 6.— *Top*: Associated γ -ray emission (*shaded*) from the CRs expected to explain the observed soft X-rays (Fig. 3), assuming (1) an LISM electron-to-proton ratio and proton spectrum (Bischoff et al. 2019); and (2) a gas density profile either (a) in hydrostatic equilibrium with CR pressure alone (*solid*) $\nabla P_{\text{cr}} = \rho \nabla \Phi$ (with Φ given by an NFW profile for the sample median halo mass in Z24), or (b) containing 1/2 of the universal baryon fraction out to R_{vir} with a power-law $\rho \propto r^{-3/2}$ (*dashed*). We compare extended diffuse CGM emission detected at different radii in the MW (Acero et al. 2016; Yang et al. 2016; Pothast et al. 2018) and M31 (Abdo et al. 2010; Karwin et al. 2019). We also compare the prediction from the cosmological MW-mass galaxy simulation in Fig. 2. *Bottom*: Observed γ -ray spectra from *Fermi* in the different MW and M31 regions detected above, compared to the LISM γ -ray spectrum from CRs in diffuse gas (dominated by pion production from LISM CR protons). The observed spectra and profiles are completely consistent with the LISM CR proton spectrum “leaking” out of galaxies at the expected rate. Per § 2.4, these would be undetectable in galaxies beyond M31.

emission (1) from the same simulation as Fig. 2, which features a CR-pressure dominated halo; (2) assuming an LISM proton-to-electron ratio (i.e. same as used to estimate CR pressure below in Fig. 7) and LISM proton spectra and heuristic density profile matching the simulation (assuming the diffuse halo contains 50% of the universal baryon fraction within R_{vir} , with a power-law $n_{\text{gas}} \propto r^{-3/2}$ profile); or (3) assuming again LISM proton spectra but calculating the gas density n_{gas} which would be in quasistatic equilibrium supported by CR pressure alone.¹⁵ We stress these are just assumptions – the X-rays do

¹⁵ Specifically following Butsky et al. (2023), assuming a spherical halo with $\nabla P_{\text{cr}} = \rho_{\text{gas}} \nabla \Phi = -\rho_{\text{gas}} V_C^2/r$, where we assume the circular velocity V_C is given by a dark matter halo with an NFW profile, concentration $c_{\text{vir}} \sim 10$, and halo mass from Zhang et al. (2024a) for each sample, while P_{cr} follows

not directly constrain these ratios. Nonetheless, it is striking that the predicted γ -ray profiles assuming LISM-like CR p/e and spectra and a CR-pressure dominated halo appear to be completely consistent with the claimed *Fermi* detections of diffuse emission from $\sim 10 - 150$ kpc in the MW and M31.

γ -rays can also be produced from IC scattering of higher-energy cosmic IR, optical, UV, and X-ray background photons, but these should be a small correction. Integrating over the observed cosmic background spectrum at $z \approx 0$ from $\sim 100 \mu\text{m}$ to ~ 100 keV (Younger & Hopkins 2011; Ueda et al. 2014; Faucher-Giguère 2020), if we ignore losses (i.e. adopt a strictly Solar-neighborhood CR lepton spectrum) we would obtain a γ -ray emissivity in a broad band around $\sim 0.3 - 3$ GeV of $\sim 1.5 \times 10^{32} \text{erg s}^{-1} \text{kpc}^{-3} (e_{\text{cr}, \ell}/\text{eV cm}^{-3}) f_{\text{loss}}^{\sim 30 \text{ GeV}}$, so this could at most boost the γ -ray luminosities by a factor of a couple (still leaving the halos undetectable), primarily from IC of the cosmic optical background via > 30 GeV electrons. But, like with synchrotron, since this is sensitive to much higher energies, we cannot ignore losses: even the conservative/intermediate loss correction lower the emissivity by a factor of $\sim 1/200$ (with the γ -rays primarily coming from IC of the UVB off few-GeV electrons), leaving this a small correction to the hadronic γ -rays.

Note that, while we expect the radio and γ ray halos to be undetected outside the MW and M31, non-detections are still interesting, as they directly place non-trivial upper limits on magnetic field strengths B and gas densities n_{gas} , respectively.

3. COMPARISON TO HOT GAS EMISSION

We now compare to what we would obtain if we made the common assumption that X-ray halos arise instead from hot gas emission.

3.1. Spectrum

Fig. 1 compares the predicted IC spectra to pure thermal free-free ($d\epsilon_{\text{ff}}/d \ln E_{\text{ph}} \sim 4 \times 10^{30} \text{erg s}^{-1} \text{kpc}^{-3} (n_{\text{gas}}/10^{-5} \text{cm}^{-3})^2 (T/10^6 \text{K})^{1/2} x_{\text{ff}} \exp[-x_{\text{ff}}]$ with $x_{\text{ff}} \equiv E_{\text{ph}}/k_B T$), similar broadly in continuum shape to the observed spectra of cluster cores (even including metal-line emission). Roughly speaking, the predicted IC spectrum looks quite similar to a $\sim 3 \times 10^7$ K free-free spectrum or a line-dominated $\sim 3 \times 10^{6-7}$ K spectrum, peaking at \sim keV.

If we ignore losses and assume an LISM CR spectrum, then the predicted IC spectrum is harder at $\geq 8 - 10$ keV, potentially providing a means to discriminate between these observationally with future facilities. However, once IC losses are accounted for, the spectral shapes are much more similar. There are some differences in detail, but these are (a) difficult to observe without very high-resolution X-ray spectroscopy, and (b) degenerate with having different detailed electron spectral shapes, or a *mix* of temperatures and metallicities and densities emitting thermally. We show this in Fig. 1 for the a typical 0.2–10 keV spectral range, where the actual predicted spectra are almost indistinguishable from thermal, even without losses. More common searches for IC emission in bright systems focus on detection of power-law continua at much harder energies, $\sim 20 - 80$ keV. But there is no predicted detection: even ignoring losses, while the luminosity in this band would only be a factor of a few smaller than at ~ 1 keV, the predicted fluxes and surface brightnesses are far lower than the sensitivity of

instruments at present. For example, given a typical $L_{X,40}$, this predicts a 20 – 80 keV integrated flux for a source at the median redshift of ~ 0.1 of $\sim 10^{-16}$ erg s $^{-1}$ cm $^{-2}$, about ~ 5 orders of magnitude lower than typical upper limits in undetected sources (Bartels et al. 2015a). Including realistic losses suppresses this exponentially (i.e. makes the spectra effectively thermal, as we showed). And because of the scaling of loss rates/lifetimes from IC and synchrotron, almost all of the IC from harder wavelengths is predicted to arise from the central $\lesssim 10$ kpc within/around the galaxies, so any hard X-ray IC which might be detected would appear as exclusively ISM emission, not CGM emission associated with the halo.

3.2. Surface Brightness Profiles

Focusing on the soft band, we can roughly approximate the 0.5 – 2 keV X-ray emissivity of hot gas as the sum of the free-free: $\epsilon_{X,ff}/(\text{erg s}^{-1} \text{kpc}^{-3}) \sim 4 \times 10^{30} (n_{\text{gas}}/10^{-5} \text{cm}^{-3})^2 (T/10^6 \text{K})^{1/2} x_{\text{ff}}^{\text{keV}} \exp[-x_{\text{ff}}^{\text{keV}}]$ (with $x_{\text{ff}}^{\text{keV}} \approx \text{keV}/k_B T$), and line emissivity which can be approximated by $\epsilon_{X,\text{line}}/(\text{erg s}^{-1} \text{kpc}^{-3}) \sim (Z/Z_\odot) 10^{32} (n_{\text{gas}}/10^{-5} \text{cm}^{-3})^2 q^2 \exp(-q)/(q+1)$ (or $\Lambda_X^{0.5-2} \sim 4 (Z/Z_\odot) \times 10^{-23} q^2/(q+1) \exp(-q)$ with $q \equiv 0.3 \text{keV}/k_B T$; Mewe et al. 1985). As expected, free-free will only dominate over line emission (for $Z \gtrsim 0.1 Z_\odot$) at $T \gtrsim 5 \times 10^6$ K. Given that the virial temperature at $z \sim 0$ scales as $T_{\text{vir}} \approx 5 \times 10^5 \text{K} (M_{\text{vir}}/10^{12} M_\odot)^{2/3}$ (Bryan & Norman 1998), the metal-line term should dominate for all $M_{\text{vir}} \lesssim 10^{13.5} M_\odot$. Given the scaling of q , the contribution in these halos is negligible for gas with $\lesssim 10^6$ K, so for low-mass halos, we expect $\epsilon_X \sim 0.8 \times 10^{30} \text{erg s}^{-1} \text{kpc}^{-3} (Z/0.1 Z_\odot) (n_{\text{hot gas}}/10^{-5} \text{cm}^{-3})^2 f_{\text{hot}}$, where f_{hot} is the fraction of hot gas with $T > 10^6$ K (which can be $\ll 1$ in such halos). Note that f_{hot} defined this way means that there should be a very rapid falloff in brightness below $M_{\text{halo}} \lesssim 10^{12.5} M_\odot$, as well.

Both simulations of MW-M31 mass galaxies (Guedes et al. 2011; Kereš et al. 2012; Liang et al. 2016; Butsky & Quinn 2018; Ji et al. 2020; Kim et al. 2022; Truong et al. 2023; Ramesh et al. 2023; and see also Braspenning et al. 2024; Sultan et al. 2024; Lehle et al. 2024 for more massive halos) and observations from UV absorption lines in similar mass halos (Machado et al. 2018; Burchett et al. 2019; Faerman et al. 2022), as well as resolved and individually-detected X-ray profiles in more massive groups/clusters (Cavagnolo et al. 2009; Ettori et al. 2019; Ghirardini et al. 2019; Lyskova et al. 2023) predict at these radii $n_{\text{hot gas}} \sim 4 \times 10^{-5} (r/100 \text{kpc})^{-\alpha_n}$ with α_n between 2 and 3, $T \propto r^{-\alpha_T}$ with $\alpha_T \sim 0.5$, and $Z \sim 0.1 (r/100 \text{kpc})^{-\alpha_Z}$ with α_Z between 0.5 – 1. Combining these, the predicted surface brightness profiles should scale as $S_X \sim 10^{32} \text{erg s}^{-1} \text{kpc}^{-2} (Z/0.1 Z_\odot) (n_{\text{hot}}/10^{-5} \text{cm}^{-3})^2 (R/100 \text{kpc})$ or $\sim 10^{33} \text{erg s}^{-1} \text{kpc}^{-2} f_{\text{hot}} (R/100 \text{kpc})^{-\alpha_R}$ with α_R between 3.5 and 6. And indeed, in said clusters, α_R is typically *observed* between 4 and 5, as expected.

Two problems are evident: (1) the normalization of S_X at $\gtrsim 100$ kpc is predicted to be $\sim 100\times$ lower than observed in the low-mass halos with virial masses $10^{12.5-13} M_\odot$ (the discrepancy increasing by another order-of-magnitude for halos with masses $\sim 10^{12} M_\odot$, owing to the temperature dependence above), and (2) the *slope* of S_X , α_R , is predicted to be qualitatively different, with a very steep $\propto R^{-4.5}$ approaching R_{vir}

as opposed to the observed very shallow $\propto R^{-1}$. If we assume thermal free-free dominates instead (i.e. Z very small), the problems both become more severe.

Explaining the observed S_X via hot gas emission is not just problematic in terms of comparison to simulations or more massive observed systems. If we wish to explain the observed S_X at large radii via hot gas emission, then we require $n_{\text{hot}} \sim 10^{-4} (M_{\text{vir}}/10^{12.5} M_\odot)^{1/2} \text{cm}^{-3} (100 \text{kpc}/r) (Z_\odot/Z)^{1/2} f_{\text{hot}}^{-1/2}$. But integrating out to R_{vir} , this would predict a baryonic mass larger than the Universal baryon fraction times the virial mass, unless f_{hot} and Z are nearly constant with $f_{\text{hot}} \sim 1$ and $Z \sim Z_\odot$ (as shown also in Truong et al. 2023; Zhang et al. 2024a¹⁶). But that, in turn, leads to other contradictions. This requires the unphysical assumption for $M_{\text{vir}} \lesssim 10^{13} M_\odot$ that almost all the gas is super-virial. Moreover, the total metal mass of the halo would be an order of magnitude larger than the maximum possible assuming every SNe had uniformly mixed its metals throughout the halo out to R_{vir} , without losses (this predicts a maximum metallicity¹⁷ of $\sim 0.1 Z_\odot$ at R_{vir} , at these halo masses; see Ma et al. 2016; Muratov et al. 2017). And both $f_{\text{hot}} \sim 1$ and $Z \sim Z_\odot$ also directly contradict what is known from UV absorption-line studies of these halo masses (which imply $Z \sim 0.1 Z_\odot$, as expected, and $f_{\text{hot}} \ll 1$ for $\sim 10^{12} M_\odot$ halos, based on the ratios of HI and cooler-to-hotter lines; Werk et al. 2014; Faerman et al. 2022; Wijers et al. 2024). Very similar constraints ($Z \sim 0.05 Z_\odot$, $f_{\text{hot}} \lesssim 0.1$) are also obtained from eROSITA X-ray absorption studies of the MW halo (Ponti et al. 2023), and from stacked Chandra absorption-line studies of galaxies along quasar sightlines (Yao et al. 2010).

Again we stress these tensions are for low-mass (MW, M31, and smaller) systems ($M_* \ll 2 \times 10^{11} M_\odot$): at the massive group scale and above, the scalings above predict more thermal emission. Also note briefly that the vast majority of historical comparisons to simulations at the lower mass end focus on just the $L_X - M_*$ relation (Popesso et al. 2024b, and references therein). This is in principle much easier to reproduce, as one could conceivably generate the observed total L_X with a relatively small ($\sim 10^8 M_\odot$) mass of Solar-metallicity gas concentrated near the halo center ($R \sim 10$ kpc) at much higher densities $n_{\text{gas}} \sim 10^{-2} \text{cm}^{-3}$, shock-heated to super-virial temperatures $\sim 5 \times 10^6$ K near the peak efficiency of keV emission (which would give $L_{X,40}^{Z=Z_\odot, T=T_{\text{peak}}} \sim (n_{\text{gas}}/10^{-2} \text{cm}^{-3})^2 (R/5 \text{kpc})^3$). Indeed, this appears to be the origin of the “super-virial” hot gas in both Galactic X-ray absorption studies and simulations of MW-mass galaxies (Chan et al. 2022; Ponti et al. 2023; Bhattacharyya et al. 2023; Roy et al. 2024a,b). And in many low-mass galaxies in observations (Anderson et al. 2013; Zhang et al. 2024b) and simulations (van de Voort et al. 2016; Chan et al. 2022) the total X-ray emission *including the galaxy* and CGM together are

¹⁶ Note that Zhang et al. (2024a) use an incorrect expression for the virial temperature of their low-mass halos which is a factor of ~ 5 higher than the standard expressions and those given and tested in adiabatic simulations in Bryan & Norman (1998) which they used to define M_{vir} and R_{vir} . They also do not include the full baryonic mass of the inner galaxy in their available budget. Correcting for this, the baryonic masses they require to explain the observed S_X via thermal emission go up by a factor of $\sim 4 - 5$, making the tension with cosmological constraints much more severe.

¹⁷ For example, at MW mass, assuming the entire Fe yield of every core-collapse and Ia SN (since Fe is what matters for the calculation here) integrated over the history of the galaxy (Maoz & Graur 2017) was retained and uniformly mixed into the halo (with a universal baryon fraction), would predict $[\text{Fe}/\text{H}] \approx -1$ using the most optimistic yields from Leung & Nomoto (2018).




FIG. 7.— CR pressure ($P_{\text{cr}} \approx e_{\text{cr}}/3$) profiles implied by the observed soft X-ray emission from Z24, if we assume an LISM-like electron-to-proton ratio (labeled), motivated by the assumption that GeV electrons and protons both escape the LISM. We compare the directly-measured MW LISM CR pressure (*point*). We also compare CR (*cyan*) and thermal (*pink*) pressure in the same cosmological simulation of a MW-mass galaxy from Fig. 2, as well as a simulation of the same galaxy with no CRs (*pink dashed*) where the pressure is primarily thermal. The observed X-ray halos appear to imply that CR pressure plays an important role in the CGM.

dominated by HMXBs and/or AGN within the central galaxy ($\lesssim 5$ kpc). The challenge, if we ignore the expected CR-IC emission, is how to reproduce the shallow profiles of S_X and emission at $\gtrsim 100$ kpc in MW-mass and smaller halos without violating other observational (and/or cosmological; see Lau et al. 2024) constraints.

4. IMPLICATIONS FOR CR PRESSURE IN THE CGM

Given the inferred $e_{\text{cr}, \ell}(R)$ from the observations in Fig. 3, it is natural to ask what this implies for the role of CR pressure ($P_{\text{cr}} \sim (1/3) e_{\text{cr}, \text{tot}}$, assuming ultra-relativistic particles whose scattering mean-free-path is smaller than the virial radius) in the CGM of MW and M31-mass galaxies. This depends on $e_{\text{cr}, \text{tot}}$, not $e_{\text{cr}, \ell}$, so is not directly constrained by the X-ray data (except to set a lower limit since $e_{\text{cr}, \text{tot}} \geq e_{\text{cr}, \ell}$). But since the GeV electrons (which dominate the X-ray signal) in the halo are consistent with what we expect to escape the ISM, where $e_{\text{cr}, \text{tot}}$ is dominated by GeV protons, it is plausible to assume a Solar-neighborhood-like $f_{\text{cr}, \ell} \equiv e_{\text{cr}, \ell}/e_{\text{cr}, \text{tot}} \sim f_{\text{cr}, \ell}^{\odot} \sim 0.01 - 0.02$. This assumption is also supported by the fact that CR electrons and protons of the same rigidity (and thus the same energy for \gtrsim GeV) have the same basic transport physics (diffusion coefficient and/or streaming speed). Since proton losses are negligible, this also supports $f_{\text{cr}, \ell} \equiv e_{\text{cr}, \ell}/e_{\text{cr}, \text{tot}} \sim f_{\text{cr}, \ell}^{\odot} \sim 0.01 - 0.02$, at least until IC electron losses are important. The resulting estimated $P_{\text{cr}}(R)$ is shown in Fig. 7.

We compare this with the predicted thermal $P_{\text{th}} \equiv nk_B T$ or CR pressure profiles in cosmological MHD+CR simulations of MW-mass halos from Hopkins et al. (2020) with the Feedback In Realistic Environments (FIRE; Hopkins et al. 2014, 2018, 2023a) model for galaxy formation and stellar feedback. Specifically, we compare (a) simulations without CRs, where the pressure is almost entirely thermal (there is some small magnetic+turbulent contribution at large radii, but these are not dominant; Hopkins et al. 2020; Ji et al. 2020), or (b) simulations with CRs using a simple streaming+diffusion model, which form a CR pressure-dominated halo. We see that the

the observationally-inferred total CR pressure in MW-mass halos agrees well with the total pressure in either simulation: this necessarily comes from thermal pressure in the “no CRs” simulation (a), or from CRs in the CR-dominated simulation (b). This, and the agreement with the MW and M31 γ -ray data assuming a CR-pressure dominated equilibrium halo in Fig. 6, strongly suggests that CRs are indeed an important component of the CGM pressure in MW-mass galaxies.

5. CONCLUSIONS

Stacked observations have argued that there is detected soft X-ray (\sim keV) diffuse emission with very shallow profiles $S_X \propto R^{-1}$ out to the virial radius around MW and M31-mass galaxies ($M_* \lesssim 10^{11} M_{\odot}$). Attempting to interpret this emission as thermal free-free or metal-line cooling from in-situ gas is problematic: the profiles are far more shallow than simulations or individually-measured thermal emission in more massive halos, factors $\gtrsim 100$ more luminous than predicted at $\gtrsim 100$ kpc, and attempting to fit to a thermal profile requires almost constant-density, Solar-metallicity, uniformly hot (super-virial-temperature) gas out to the virial radius (contradicting limits from cosmology, the total metal budget of galaxies, and both UV and X-ray absorption-line studies).

We show that the observed X-ray halos instead naturally arise from inverse Compton (IC) scattering of the CMB by extended \sim GeV cosmic ray (CR) halos around galaxies. Measurements of CRs in the ISM of the MW directly show that GeV CRs largely escape the galaxy out into the CGM. Simple models of steady-state CR profiles expected around galaxies (given the GeV CR density observed in the ISM) naturally predict the normalization, shape, and characteristic photon energy of the observed X-ray surface-brightness profiles. We show that the CMB temperature and \sim GeV peak of the CR energy density ensure that the IC X-ray emission peaks in observed soft X-rays (~ 1 keV), with a spectrum that resembles hot gas thermal emission. The total halo X-ray luminosity we find from IC emission also agrees well with that observed around galaxies with stellar masses $M_* \lesssim 2 \times 10^{11} M_{\odot}$. Our models predict associated synchrotron and γ -ray halos but we show that these would be extremely faint, well below present detection limits except in the MW and M31 specifically, where the predicted γ -ray emission assuming a CR-pressure dominated halo with LISM-like proton-to-electron ratios agrees well with Fermi observations.

Our calculations assume that the local ISM CR electron spectrum observed in the MW is representative of the CR spectra in other galaxies with similar total stellar masses; if correct, the surprisingly quasi-thermal spectrum of IC-scattered CMB radiation should be generic in galaxy halos. The assumption of a MW ISM CR electron spectrum is less easily justified in massive galaxies whose CRs likely originate primarily from AGN rather than star formation.

We stress that CR-IC halos *must* be present around MW-mass halos, given what is known about the energy density, lifetimes, and residence times of GeV CRs in the LISM (~ 8 kpc from the Galactic center), which necessarily imply that almost all of the GeV CR electron energy leaks into the CGM. Without invoking unphysically large CGM magnetic fields or gas densities (which are immediately ruled out by existing radio and γ -ray observations), the only way to strongly modify the IC halos would be to either (a) make the CR streaming speed much lower, which would result in the same halo *luminosity*, but a much brighter, more compact halo, which is clearly ruled

out by the observations, or (b) make the streaming speed and diffusivity orders-of-magnitude larger (much larger than any reasonable physical model prediction), so CRs escape the halo much more rapidly and have low energy densities in the CGM.

We predict that CR-IC halos should dominate the X-ray emission for halos less massive than sizeable groups, above which the larger masses and hotter virial temperatures produce larger thermal emission. This appears to be supported by the keV halo luminosity L_X versus stellar-mass M_* correlation, which steepens dramatically above a stellar mass $\sim 1.5 - 2 \times 10^{11} M_\odot$. It is plausible, however, that CR-IC could still contribute significantly to soft X-ray emission at radii $\lesssim 100 - 200$ kpc even at these higher masses (well within the virial radii for the expected large halo masses, so not dominating the total luminosity), especially if strong AGN have been active in the last \sim Gyr, as argued for e.g. some “mini-halos” in massive clusters like Ophiuchus or Abell 3112 (Bartels et al. 2015b; Gitti et al. 2002, 2004; Bonamente et al. 2007; Murgia et al. 2010; Gitti 2016).

If CR-IC is indeed the source of the extended soft X-ray halos around low-mass galaxies, then the keV soft X-ray surface brightness is *directly* proportional to the CR lepton energy density, without any uncertain degenerate factors (unlike e.g. synchrotron and γ -ray emission, which are degenerate with magnetic field strengths and gas densities, respectively, and depend much more sensitively on the CR spectral shape). This is because the seed CMB photons for IC scattering are precisely known. The total halo X-ray luminosity (limited by IC

losses) directly constrains the leptonic energy injection rate from the galaxy; observed values are consistent with expected SNe or AGN (if a fraction of just a few 10^{-5} of $\dot{M}_{\text{BH}} c^2$ accelerates leptons) injection rates in MW and M31 and lower-mass systems. The X-ray surface brightness of the IC halos also yields a direct constraint on the effective diffusion coefficient (or streaming speed) of \sim GeV (rigidity \sim GV) CRs, which we show agrees well with ISM constraints close to galaxies, but κ_{eff} rises by an order of magnitude or more at $\gtrsim 100$ kpc, consistent with a CR streaming speed of $\sim 100 \text{ km s}^{-1}$, in agreement for MW-mass halos with independent (indirect) constraints from UV absorption studies (Butsky et al. 2023). Assuming ISM-like proton-to-electron ratios, our results also imply that the CR pressure is a major component of the total gas pressure in the CGM of MW-mass galaxies, as suggested by previous cosmological zoom-in simulations with CRs (e.g., Ji et al. 2020). Further characterizing the soft X-ray halos of MW-mass galaxies therefore presents a potentially invaluable probe of both CR and CGM physics.

Support for PFH & EQ was provided by Simons Investigator grants. PFH also acknowledges the support of the Institute for Advanced Study where early discussions leading to this work occurred. Numerical calculations were run on allocation AST21010 supported by the NSF. This work was supported in part by NSF AST grant 2107872.

REFERENCES

- Abdo A. A., et al., 2010, *ApJ*, 710, 133
 Acero F., et al., 2016, *ApJS*, 223, 26
 Ackermann M., et al., 2011, *ApJ*, 726, 81
 Ackermann M., et al., 2014, *ApJ*, 787, 18
 Adriani O., et al., 2018, *Phys. Rev. Lett.*, 120, 261102
 Allen S. W., Dunn R. J. H., Fabian A. C., Taylor G. B., Reynolds C. S., 2006, *MNRAS*, 372, 21
 Amato E., Blasi P., 2018, *Advances in Space Research*, 62, 2731
 Anderson M. E., Bregman J. N., Dai X., 2013, *ApJ*, 762, 106
 Anderson M. E., Gaspari M., White S. D. M., Wang W., Dai X., 2015, *MNRAS*, 449, 3806
 Bahar Y. E., et al., 2024, *A&A*, 691, A188
 Bartels R., Zandanel F., Ando S., 2015a, *A&A*, 582, A20
 Bartels R., Zandanel F., Ando S., 2015b, *A&A*, 582, A20
 Bhattacharyya J., Das S., Gupta A., Mathur S., Krongold Y., 2023, *ApJ*, 952, 41
 Birzan L., Rafferty D. A., McNamara B. R., Wise M. W., Nulsen P. E. J., 2004, *ApJ*, 607, 800
 Bisschoff D., Potgieter M. S., Aslam O. P. M., 2019, *ApJ*, 878, 59
 Blasi P., Amato E., 2012, *Journal of Cosmology and Astroparticle Physics*, 1, 010
 Bonamente M., Nevalainen J., Lieu R., 2007, *ApJ*, 668, 796
 Braspenning J., et al., 2024, *MNRAS*, 533, 2656
 Bryan G. L., Norman M. L., 1998, *ApJ*, 495, 80
 Buck T., Pfrommer C., Pakmor R., Grand R. J. J., Springel V., 2020, *MNRAS*, 497, 1712
 Burchett J. N., et al., 2019, *ApJ*, 877, L20
 Butsky I. S., Quinn T. R., 2018, *ApJ*, 868, 108
 Butsky I. S., Nakum S., Ponnada S. B., Hummels C. B., Ji S., Hopkins P. F., 2023, *MNRAS*, 521, 2477
 Caprioli D., 2012, *J. Cosmology Astropart. Phys.*, 2012, 038
 Cavagnolo K. W., Donahue M., Voit G. M., Sun M., 2009, *ApJ*, in press arXiv:0902.1802, 0902, 1802
 Chan T. K., Kereš D., Hopkins P. F., Quataert E., Su K. Y., Hayward C. C., Faucher-Giguère C.-A., 2019, *MNRAS*, 488, 3716
 Chan T. K., Kereš D., Gurvich A. B., Hopkins P. F., Trapp C., Ji S., Faucher-Giguère C.-A., 2022, *MNRAS*, 517, 597
 Cooke K. C., et al., 2023, *ApJ*, 942, 49
 Cummings A. C., et al., 2016, *ApJ*, 831, 18
 Dacunha T., Martin-Alvarez S., Clark S. E., Lopez-Rodriguez E., 2024, arXiv e-prints, p. arXiv:2409.08437
 De La Torre Luque P., Mazziotta M. N., Loparco F., Gargano F., Serini D., 2021, *J. Cosmology Astropart. Phys.*, 2021, 099
 Delhaize J., et al., 2017, *A&A*, 602, A4
 Di Mauro M., Korsmeier M., Cuoco A., 2024, *Phys. Rev. D*, 109, 123003
 Do A., Duong M., McDaniel A., O’Connor C., Profumo S., Rafael J., Sweeney C., Vera W., 2021, *Phys. Rev. D*, 104, 123016
 Dolag K., Kachelriess M., Ostapchenko S., Tomàs R., 2011, *ApJ*, 727, L4
 Etori S., et al., 2019, *A&A*, 621, A39
 Evoli C., Aloisio R., Blasi P., 2019, *Phys. Rev. D*, 99, 103023
 Faerman Y., Pandya V., Somerville R. S., Sternberg A., 2022, *ApJ*, 928, 37
 Falcke H., Körding E., Markoff S., 2004, *A&A*, 414, 895
 Faucher-Giguère C.-A., 2020, *MNRAS*, 493, 1614
 Gaggero D., Urbano A., Valli M., Ullio P., 2015, *Phys. Rev. D*, 91, 083012
 Gandhi P. J., Wetzel A., Hopkins P. F., Shappee B. J., Wheeler C., Faucher-Giguère C.-A., 2022, *MNRAS*, 516, 1941
 Geha M., et al., 2024, *ApJ*, 976, 118
 Génolini Y., et al., 2019, *Phys. Rev. D*, 99, 123028
 Ghirardini V., et al., 2019, *A&A*, 621, A41
 Gitti M., 2016, arXiv e-prints, p. arXiv:1602.08312
 Gitti M., Brunetti G., Setti G., 2002, *A&A*, 386, 456
 Gitti M., Brunetti G., Feretti L., Setti G., 2004, *A&A*, 417, 1
 Guedes J., Callegari S., Madau P., Mayer L., 2011, *ApJ*, 742, 76
 Guo F., Oh S. P., 2008, *MNRAS*, 384, 251
 Higdon J. C., Lingelfelter R. E., Ramaty R., 1998, *ApJ*, 509, L33
 Holguin F., Ruszkowski M., Lazarian A., Farber R., Yang H. Y. K., 2019, *MNRAS*, 490, 1271
 Hopkins P. F., Hernquist L., Martini P., Cox T. J., Robertson B., Di Matteo T., Springel V., 2005a, *ApJ*, 625, L71
 Hopkins P. F., Hernquist L., Cox T. J., Di Matteo T., Martini P., Robertson B., Springel V., 2005b, *ApJ*, 630, 705
 Hopkins P. F., Hernquist L., Cox T. J., Di Matteo T., Robertson B., Springel V., 2005c, *ApJ*, 630, 716
 Hopkins P. F., Hernquist L., Cox T. J., Di Matteo T., Robertson B., Springel V., 2005d, *ApJ*, 632, 81
 Hopkins P. F., Keres D., Onorbe J., Faucher-Giguère C.-A., Quataert E., Murray N., Bullock J. S., 2014, *MNRAS*, 445, 581
 Hopkins P. F., et al., 2018, *MNRAS*, 480, 800
 Hopkins P. F., et al., 2020, *MNRAS*, 492, 3465
 Hopkins P. F., Chan T. K., Ji S., Hummels C. B., Kereš D., Quataert E., Faucher-Giguère C.-A., 2021a, *MNRAS*, 501, 3640
 Hopkins P. F., Chan T. K., Squire J., Quataert E., Ji S., Kereš D., Faucher-Giguère C.-A., 2021b, *MNRAS*, 501, 3663
 Hopkins P. F., Squire J., Chan T. K., Quataert E., Ji S., Kereš D., Faucher-Giguère C.-A., 2021c, *MNRAS*, 501, 4184
 Hopkins P. F., Butsky I. S., Panopoulou G. V., Ji S., Quataert E., Faucher-Giguère C.-A., Kereš D., 2022a, *MNRAS*, 516, 3470
 Hopkins P. F., Squire J., Butsky I. S., Ji S., 2022b, *MNRAS*, 517, 5413
 Hopkins P. F., et al., 2023a, *MNRAS*, 519, 3154
 Hopkins P. F., Butsky I. S., Ji S., Kereš D., 2023b, *MNRAS*, 522, 2936

- Hsiung Chen M., Crasemann B., 1985, *Atomic Data and Nuclear Data Tables*, **33**, 217
- Ji S., et al., 2020, *MNRAS*, **496**, 4221
- Ji S., Kereš D., Chan T. K., Stern J., Hummel C. B., Hopkins P. F., Quataert E., Faucher-Giguère C.-A., 2021, *MNRAS*, **505**, 259
- Ji S., Squire J., Hopkins P. F., 2022, *MNRAS*, **513**, 282
- Kaastra J. S., Mewe R., 1993, *A&AS*, **97**, 443
- Karwin C. M., Murgia S., Campbell S., Moskalenko I. V., 2019, *ApJ*, **880**, 95
- Katz H., Martín-Alvarez S., Devriendt J., Slyz A., Kimm T., 2019, *MNRAS*, **484**, 2620
- Kelly B. C., Vestergaard M., Fan X., Hopkins P., Hernquist L., Siemiginowska A., 2010, *ApJ*, **719**, 1315
- Kempki P., Quataert E., 2022, *MNRAS*, **514**, 657
- Kereš D., Vogelsberger M., Sijacki D., Springel V., Hernquist L., 2012, *MNRAS*, **425**, 2027
- Kim J., et al., 2022, *ApJ*, **926**, 179
- Korsmeier M., Cuoco A., 2021, *Phys. Rev. D*, **103**, 103016
- Korsmeier M., Cuoco A., 2022, *Phys. Rev. D*, **105**, 103033
- Kovaltsov G. A., Barghouty A. F., Kocharov L., Ostryakov V. M., Torsti J., 2001, *A&A*, **375**, 1075
- Kraft R., et al., 2023, Line Emission Mapper (LEM): Probing the physics of cosmic ecosystems ([arXiv:2211.09827](https://arxiv.org/abs/2211.09827)), <https://arxiv.org/abs/2211.09827>
- Lacki B. C., Thompson T. A., Quataert E., 2010, *ApJ*, **717**, 1
- Lacki B. C., Thompson T. A., Quataert E., Loeb A., Waxman E., 2011, *ApJ*, **734**, 107
- Lan T.-W., Prochaska J. X., 2020, *MNRAS*, **496**, 3142
- Lau E. T., et al., 2024, [arXiv e-prints](https://arxiv.org/abs/2412.04559), p. [arXiv:2412.04559](https://arxiv.org/abs/2412.04559)
- Lehle K., Nelson D., Pillepich A., Truong N., Rohr E., 2024, *A&A*, **687**, A129
- Leung S.-C., Nomoto K., 2018, *ApJ*, **861**, 143
- Liang C. J., Kravtsov A. V., Agertz O., 2016, *MNRAS*, **458**, 1164
- Lopez L. A., Auchettl K., Linden T., Bolatto A. D., Thompson T. A., Ramirez-Ruiz E., 2018, *ApJ*, **867**, 44
- Lyskova N., Churazov E., Khabibullin I. I., Burenin R., Starobinsky A. A., Sunyaev R., 2023, *MNRAS*, **525**, 898
- Ma X., Hopkins P. F., Faucher-Giguère C.-A., Zolman N., Muratov A. L., Kereš D., Quataert E., 2016, *MNRAS*, **456**, 2140
- Machado R. E. G., Tissera P. B., Lima Neto G. B., Sodré L., 2018, *A&A*, **609**, A66
- Magnelli B., et al., 2015, *A&A*, **573**, A45
- Mannings A. G., et al., 2023, *ApJ*, **954**, 179
- Maoz D., Graur O., 2017, *ApJ*, **848**, 25
- Marinacci F., Vogelsberger M., Mocz P., Pakmor R., 2015, *MNRAS*, **453**, 3999
- Martin-Alvarez S., et al., 2024, *ApJ*, **966**, 43
- Martini P., 2004, in Ho L. C., ed., *Coevolution of Black Holes and Galaxies* (Cambridge: Cambridge University Press). p. 169
- Maurin D., 2020, *Computer Physics Communications*, **247**, 106942
- McNamara B. R., Nulsen P. E. J., 2007, *ARA&A*, **45**, 117
- Mewe R., Gronenschild E. H. B. M., van den Oord G. H. J., 1985, *A&AS*, **62**, 197
- Mineo S., Gilfanov M., Sunyaev R., 2012, *MNRAS*, **419**, 2095
- Mineo S., Gilfanov M., Lehmer B. D., Morrison G. E., Sunyaev R., 2014, *MNRAS*, **437**, 1698
- Muratov A. L., et al., 2017, *MNRAS*, **468**, 4170
- Murgia M., Eckert D., Govoni F., Ferrari C., Pandey-Pommier M., Nevalainen J., Paltani S., 2010, *A&A*, **514**, A76
- O'Sullivan S. P., et al., 2020, *MNRAS*, **495**, 2607
- Padmanabhan H., Loeb A., 2023, *ApJ*, **946**, L18
- Pearson W. J., et al., 2018, *A&A*, **615**, A146
- Ponnada S. B., et al., 2022, *MNRAS*, **516**, 4417
- Ponnada S. B., et al., 2024a, [arXiv e-prints](https://arxiv.org/abs/2410.02971), p. [arXiv:2410.02971](https://arxiv.org/abs/2410.02971)
- Ponnada S. B., et al., 2024b, *MNRAS*, **527**, 11707
- Ponnada S. B., et al., 2024c, *MNRAS*, **530**, L1
- Ponti G., et al., 2023, *A&A*, **674**, A195
- Popesso P., et al., 2024a, [arXiv e-prints](https://arxiv.org/abs/2411.16546), p. [arXiv:2411.16546](https://arxiv.org/abs/2411.16546)
- Popesso P., et al., 2024b, [arXiv e-prints](https://arxiv.org/abs/2411.17120), p. [arXiv:2411.17120](https://arxiv.org/abs/2411.17120)
- Pothast M., Gaggero D., Storm E., Weniger C., 2018, *J. Cosmology Astropart. Phys.*, **2018**, 045
- Prochaska J. X., Zheng Y., 2019, *MNRAS*, **485**, 648
- Prochaska J. X., et al., 2019, *Science*, **366**, 231
- Ramesh R., Nelson D., Pillepich A., 2023, *MNRAS*, **518**, 5754
- Ramesh R., Nelson D., Girichidis P., 2024, [arXiv e-prints](https://arxiv.org/abs/2409.18238), p. [arXiv:2409.18238](https://arxiv.org/abs/2409.18238)
- Ravi V., et al., 2016, *Science*, **354**, 1249
- Recchia S., Gabici S., Aharonian F. A., Niro V., 2021, *ApJ*, **914**, 135
- Roy M., Su K.-Y., Mathur S., Stern J., 2024a, [arXiv e-prints](https://arxiv.org/abs/2409.17252), p. [arXiv:2409.17252](https://arxiv.org/abs/2409.17252)
- Roy M., Mathur S., Das S., Lara-Di A., Krongold Y., Gupta A., 2024b, [arXiv e-prints](https://arxiv.org/abs/2411.15394), p. [arXiv:2411.15394](https://arxiv.org/abs/2411.15394)
- Ruszkowski M., Yang H.-Y. K., Zweibel E., 2017, *ApJ*, **834**, 208
- Su K.-Y., et al., 2020, *MNRAS*, **491**, 1190
- Su K.-Y., et al., 2024, *MNRAS*, **532**, 2724
- Sukhbold T., Ertl T., Woosley S. E., Brown J. M., Janka H. T., 2016, *ApJ*, **821**, 38
- Sultan I., Faucher-Giguère C.-A., Stern J., Rotshtein S., Byrne L., Wijers N., 2024, [arXiv e-prints](https://arxiv.org/abs/2410.16359), p. [arXiv:2410.16359](https://arxiv.org/abs/2410.16359)
- Thomas T., Pfrommer C., Pakmor R., 2023, *MNRAS*, **521**, 3023
- Tibaldo L., 2014, *Brazilian Journal of Physics*, **44**, 600
- Tibaldo L., et al., 2015, *ApJ*, **807**, 161
- Tibaldo L., Gaggero D., Martin P., 2021, *Universe*, **7**, 141
- Tjemsland J., Meyer M., Vazza F., 2024, *ApJ*, **963**, 135
- Torbaniuk O., Paolillo M., Carrera F., Cavuoti S., Vignali C., Longo G., Aird J., 2021, *MNRAS*, **506**, 2619
- Torbaniuk O., Paolillo M., D'Abrusco R., Vignali C., Georgakakis A., Carrera F. J., Civano F., 2024, *MNRAS*, **527**, 12091
- Truong N., et al., 2023, *MNRAS*, **525**, 1976
- Ueda Y., Akiyama M., Hasinger G., Miyaji T., Watson M. G., 2014, *ApJ*, **786**, 104
- Vernstrom T., Gaensler B. M., Rudnick L., Andernach H., 2019, *ApJ*, **878**, 92
- Wang L., et al., 2019, *A&A*, **631**, A109
- Werk J. K., et al., 2014, *ApJ*, **792**, 8
- Wiener J., Oh S. P., Guo F., 2013, *MNRAS*, **434**, 2209
- Wiener J., Pfrommer C., Oh S. P., 2017, *MNRAS*, **467**, 906
- Wijers N. A., Faucher-Giguère C.-A., Stern J., Byrne L., Sultan I., 2024, *MNRAS*, in press, p. [arXiv:2401.08776](https://arxiv.org/abs/2401.08776)
- Xing Y., Wang Z., Zheng D., Li J., 2023, *ApJ*, **945**, L22
- Yang R., Aharonian F., Evoli C., 2016, *Phys. Rev. D*, **93**, 123007
- Yao Y., Wang Q. D., Penton S. V., Tripp T. M., Shull J. M., Stocke J. T., 2010, *ApJ*, **716**, 1514
- Younger J. D., Hopkins P. F., 2011, *MNRAS*, **410**, 2180
- Zhang Y., Peng F.-K., Wang X.-Y., 2019, *ApJ*, **874**, 173
- Zhang Y., et al., 2024a, *A&A*, **690**, A267
- Zhang Y., et al., 2024b, *A&A*, **690**, A268
- ZuHone J. A., et al., 2024, Properties of the Line-of-Sight Velocity Field in the Hot and X-ray Emitting Circumgalactic Medium of Nearby Simulated Disk Galaxies ([arXiv:2307.01269](https://arxiv.org/abs/2307.01269)), <https://arxiv.org/abs/2307.01269>
- Zweibel E. G., 2017, *Physics of Plasmas*, **24**, 055402
- van de Voort F., Quataert E., Hopkins P. F., Faucher-Giguère C.-A., Feldmann R., Kereš D., Chan T. K., Hafen Z., 2016, *MNRAS*, **463**, 4533

This paper was built using the Open Journal of Astrophysics \LaTeX template. The OJA is a journal which provides fast and easy peer review for new papers in the astro-ph section of the

arXiv, making the reviewing process simpler for authors and referees alike. Learn more at <http://astro.theoj.org>.